

Tobacco Product Use Among U.S. Middle and High School Students — National Youth Tobacco Survey, 2023

Jan Birdsey, MPH¹; Monica Cornelius, PhD¹; Ahmed Jamal, MBBS¹; Eunice Park-Lee, PhD²; Maria R. Cooper, PhD²; Jia Wang, MPH²; Michael D. Sawdey, PhD²; Karen A. Cullen, PhD²; Linda Neff, PhD¹

Abstract

Tobacco product use during adolescence increases the risk for lifelong nicotine addiction and adverse health consequences. CDC and the Food and Drug Administration analyzed data from the 2023 National Youth Tobacco Survey to assess tobacco product use patterns among U.S. middle school (grades 6–8) and high school (grades 9–12) students. In 2023, 10.0% of middle and high school students (2.80 million) reported current (i.e., past 30-day) use of any tobacco product. Current use of any tobacco product by high school students declined by an estimated 540,000, from 2.51 million in 2022 to 1.97 million in 2023. From 2022 to 2023, current e-cigarette use among high school students declined from 14.1% to 10.0%. Among middle and high school students, e-cigarette products were the most used tobacco product in 2023 (7.7%; 2.13 million), followed by cigarettes (1.6%), cigars (1.6%), nicotine pouches (1.5%), smokeless tobacco (1.2%), other oral nicotine products (1.2%), hookahs (1.1%), heated tobacco products (1.0%), and pipe tobacco (0.5%). Among students who had ever used an e-cigarette, 46.7% reported current use. In 2023, among students reporting current e-cigarette use, 89.4% used flavored products and 25.2% used an e-cigarette daily; the most commonly reported brands were Elf Bar, Esco Bars, Vuse, JUUL, and Mr. Fog. Given the number of middle and high school students that use tobacco products, sustained efforts to prevent initiation of tobacco product use among young persons and strategies to help young tobacco users quit are critical to reducing U.S. youth tobacco product use.

Introduction

Commercial tobacco use* among U.S. youths can lead to lifelong nicotine addiction (1) and subsequent disability,

disease, and death (2). This report presents findings from the 2023 National Youth Tobacco Survey (NYTS) and describes the prevalence of ever use (i.e., ever having used, even once or twice) and current use of nine tobacco product types, flavored tobacco products, and e-cigarette use behaviors among U.S. middle and high school students. In addition, 2023 NYTS results are compared with those reported for 2022 NYTS data (3).

Methods

Data Collection

The NYTS is a cross-sectional, school-based, self-administered web-based survey of U.S. middle and high school students. A stratified, three-stage cluster sampling procedure

INSIDE

- 1183 Tuberculosis Testing and Latent Tuberculosis Infection Treatment Practices Among Health Care Providers — United States, 2020–2022
- 1190 Vaccination Coverage by Age 24 Months Among Children Born in 2019 and 2020 — National Immunization Survey–Child, United States, 2020–2022
- 1197 *Vital Signs*: Health Worker–Perceived Working Conditions and Symptoms of Poor Mental Health — Quality of Worklife Survey, United States, 2018–2022
- 1207 QuickStats

Continuing Education examination available at https://www.cdc.gov/mmwr/mmwr_continuingEducation.html

* The term “tobacco” as used in this report refers to commercial tobacco products and not to sacred and traditional use of tobacco by some American Indian communities.



was used to generate a nationally representative sample of U.S. students attending private or public middle (grades 6–8) and high (grades 9–12) schools. In 2023, data were collected during March 9–June 16; a total of 22,069 students from 179 schools participated, with an overall response rate of 30.5%.

Data Analysis

National weighted prevalence estimates, 95% CIs, and population totals[†] were calculated for ever use (i.e., ever having used, even once or twice) and current use (i.e., use on ≥ 1 days during the past 30 days) of nine commercial tobacco products[§] (e-cigarettes, cigarettes, cigars, smokeless tobacco, nicotine pouches,[¶] hookahs, pipe tobacco, heated tobacco

products,^{**} and other oral nicotine products) by student characteristics. Three composite measures were also reported for use of any tobacco product,^{††} any combustible tobacco product,^{§§} and multiple tobacco products.^{¶¶} Current e-cigarette use (i.e., use on ≥ 1 day during the past 30 days) was reported by frequency of use, device type,^{***} brand,^{†††} and flavor.^{§§§}

[†] Data were weighted to account for complex survey design and to adjust for nonresponse. The weighted proportions of students in each grade matched national population proportions for U.S. public and private schools derived from data from Market Data Retrieval Inc. 2021–2022 Common Core of Data and the National Center for Education Statistics 2019–2020 Private School Universe Study. Population total estimates were rounded down to the nearest 10,000 persons.

[§] Products include e-cigarettes, cigarettes, cigars (cigars, cigarillos, or little cigars), smokeless tobacco (chewing tobacco, snuff, dip, or snus), hookahs, heated tobacco products, nicotine pouches, pipe tobacco, bidis (small brown cigarettes wrapped in a leaf), and other oral nicotine products (lozenges, discs, tablets, gums, dissolvable tobacco products, and other products). In 2023, dissolvable tobacco products were reclassified from smokeless tobacco to other oral nicotine products.

[¶] Small, flavored pouches contain nicotine that comes from tobacco. Users place them in their mouth. Nicotine pouches are different from other smokeless tobacco products such as snus, dip, or chewing tobacco, because they do not contain any tobacco leaf.

^{**} Heated tobacco products heat processed tobacco leaf in the form of sticks (“heatsticks”), plugs, or capsules to produce a vapor that the user inhales. They are different from e-cigarettes, which heat a liquid to produce a vapor.

^{††} Any tobacco product use was defined as use of one or more of the following tobacco products: e-cigarettes, cigarettes, cigars, smokeless tobacco, hookahs, heated tobacco products, nicotine pouches, pipe tobacco, bidis, or other oral nicotine products.

^{§§} Any combustible tobacco product use is defined as use of one or more of the following tobacco products: cigarettes, cigars, hookahs, pipe tobacco, or bidis.

^{¶¶} Multiple tobacco product use was defined as use of two or more of the following tobacco products: e-cigarettes, cigarettes, cigars, smokeless tobacco, hookahs, heated tobacco products, nicotine pouches, pipe tobacco, bidis, or other oral nicotine products.

^{***} Categories used are “disposables,” “prefilled or refillable pods or cartridges,” “tanks or mod system,” or “don’t know the type.” Disposable e-cigarettes come prefilled with e-liquid and are designed to be discarded once empty. Some pods or cartridges come prefilled with e-liquid that is replaced after use, although others can be refilled by the user. Tank or mod-type devices can also be refilled but are also usually customizable.

^{†††} Brand response options included blu, Breeze, Elf Bar, Esco Bars, Fume, HQD, JUUL, Kangvape (including Oncee Stick), Logic, Mr. Fog, NJOY, SMOK (including NOVO), Suorin (including Air Bar), Vuse, “some other brand(s) not listed here” with space for a write-in name, and “not sure/I don’t know the brand.” Write-in responses corresponding to an original response option were recoded.

^{§§§} Flavor type was determined by response to the question, “In the past 30 days when you used e-cigarettes, what flavors did you use? (Select one or more).” Those who selected “some other flavor not listed here” could provide a write-in response; write-in responses corresponding to an original response option were recoded.

The *MMWR* series of publications is published by the Office of Science, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30329-4027.

Suggested citation: [Author names; first three, then et al., if more than six.] [Report title]. *MMWR Morb Mortal Wkly Rep* 2023;72:[inclusive page numbers].

Centers for Disease Control and Prevention

Mandy K. Cohen, MD, MPH, *Director*
Debra Houry, MD, MPH, *Chief Medical Officer and Deputy Director for Program and Science*
Paul Muntner, PhD, MHS, *Acting Director, Office of Science*

MMWR Editorial and Production Staff (Weekly)

Charlotte K. Kent, PhD, MPH, *Editor in Chief*
Rachel Gorwitz, MD, MPH, *Acting Executive Editor*
Jacqueline Gindler, MD, *Editor*
Paul Z. Siegel, MD, MPH, *Associate Editor*
Mary Dott, MD, MPH, *Online Editor*
Teresa F. Rutledge, *Managing Editor*
Teresa M. Hood, MS, *Lead Technical Writer-Editor*
Glenn Damon, Jacqueline Farley, MS,
Tiana Garrett, PhD, MPH, Ashley Morici,
Stacy Simon, MA, Morgan Thompson,
Suzanne Webb, PhD, MA,
Technical Writer-Editors

Martha F. Boyd, *Lead Visual Information Specialist*
Alexander J. Gottardy, Maureen A. Leahy,
Stephen R. Spriggs, Armina Velarde, Tong Yang,
Visual Information Specialists
Quang M. Doan, MBA, Phyllis H. King,
Terraye M. Starr, Moua Yang,
Information Technology Specialists

Symone Hairston, MPH,
Acting Lead Health Communication Specialist
Kiana Cohen, MPH,
Leslie Hamlin, Lowery Johnson,
Health Communication Specialists
Dewin Jimenez, Will Yang, MA,
Visual Information Specialists

MMWR Editorial Board

Matthew L. Boulton, MD, MPH
Carolyn Brooks, ScD, MA
Virginia A. Caine, MD
Jonathan E. Fielding, MD, MPH, MBA

Timothy F. Jones, MD, *Chairman*
David W. Fleming, MD
William E. Halperin, MD, DrPH, MPH
Jewel Mullen, MD, MPH, MPA
Jeff Niederdeppe, PhD
Patricia Quinlisk, MD, MPH

Patrick L. Remington, MD, MPH
Carlos Roig, MS, MA
William Schaffner, MD
Morgan Bobb Swanson, MD, PhD

Changes in current-use prevalence since 2022 were estimated using t-tests; details of the 2022 NYTS data collection and estimates have been published previously (3). P-values <0.05 were considered statistically significant. Analyses were conducted using SAS-callable SUDAAN software (version 11.0.4; Research Triangle Institute). Estimates with an unweighted denominator <50 or a relative SE >30% were suppressed. This activity was reviewed by CDC, deemed not research, and was conducted consistent with applicable federal law and CDC policy.^{¶¶¶}

Results

Tobacco Product Use by Population

In 2023, 22.2% of U.S. middle and high school students reported ever using any tobacco product, corresponding to 6.21 million persons (Table 1); 10.0% of students reported current use of any tobacco product, corresponding to 2.80 million persons (Table 2). Overall, current use of any tobacco product was reported by 11.2% of female, 8.9% of male, 12.6% of non-Hispanic multiracial (multiracial), 11.7% of Hispanic or Latino (Hispanic), 9.5% of non-Hispanic White (White), 9.3% of non-Hispanic Black or African American (Black), and 8.0% of non-Hispanic American Indian or Alaska Native (AI/AN) students.^{****} Current use of any combustible tobacco product was reported by 4.7% of Black, 3.9% of Hispanic, 3.7% of multiracial, and 2.7% of White and AI/AN students.

Types of Tobacco Products Used

E-cigarettes were the most commonly reported currently used tobacco product among all students (7.7%) and both middle school (4.6%) and high school students (10.0%). Other currently used tobacco products included cigarettes (1.6%), cigars (1.6%), nicotine pouches (1.5%), smokeless tobacco (1.2%), other oral nicotine products (1.2%), hookahs (1.1%), heated tobacco products (1.0%), and pipe tobacco (0.5%). Among students who had ever used e-cigarettes, 46.7% reported current e-cigarette use.

Characteristics of E-cigarette Use

Among students reporting current e-cigarette use, 25.2% reported using e-cigarettes daily. Frequent use (≥20 of the past 30 days) was reported by 34.7% of current e-cigarette users (Table 3). Disposable e-cigarettes were the most commonly reported device type used (60.7%), followed by prefilled or refillable pods or cartridges (16.1%), and tanks or mod systems

(modifiable devices allowing users to customize the substances in the device) (5.9%). Among students who currently used e-cigarettes, Elf Bar was the most commonly reported brand (56.7%), followed by Esco Bars (21.6%), Vuse (20.7%), JUUL (16.5%), and Mr. Fog (13.6%).

Among students reporting current e-cigarette use, 89.4% reported using a flavored product during the past 30 days, excluding those who only used tobacco-flavored or unflavored e-cigarettes (Table 3). Among students who currently used e-cigarettes, fruit- (63.4%) and candy- (35.0%) flavored categories were reported most commonly; 6.4% of students reported use of tobacco-flavored e-cigarettes. Among those who currently used disposable e-cigarettes, the top reported flavor categories were fruit (70.5%), candy (39.8%), mint (32.0%), menthol (18.7%), unflavored (7.8%), alcoholic drinks (7.2%), and tobacco-flavored (5.4%) (Supplementary Table 1, <https://stacks.cdc.gov/view/cdc/134700>). Among students reporting current use of any tobacco product, 86.9% used a flavored product, ranging from 40.4% of cigarette users (menthol) to 89.4% of e-cigarette users (Supplementary Table 2, <https://stacks.cdc.gov/view/cdc/134701>). Among students currently using tobacco products, use of products with “ice” or “iced”^{††††} included in the flavor name was reported by 57.9% of e-cigarette users, 25.9% of nicotine pouch users, and 22.6% of cigar users; use of concept flavors^{§§§§} was reported by 16.1% of e-cigarette users and 13.4% of cigar users (Supplementary Table 3, <https://stacks.cdc.gov/view/cdc/134702>).

Tobacco Product Use Over Time

From 2022 to 2023, among high school students, statistically significant declines ($p < 0.05$) occurred in current use of any tobacco product (from 16.5% to 12.6%), e-cigarettes (from 14.1% to 10.0%), cigars (from 2.8% to 1.8%), and any combustible tobacco product (from 5.2% to 3.9%). Among middle school students, statistically significant increases ($p < 0.05$) occurred in current use of any tobacco product (from 4.5% to 6.6%) and multiple tobacco products (from 1.5% to 2.5%). Among middle school and high school students combined, no

^{††††} Current users were asked, “Did any of the flavors you used in the past 30 days have names or descriptions that included the word ‘ice’ or ‘iced’ (for example, blueberry ice or strawberry ice)?” Those who reported using only “unflavored” e-cigarettes (n = 60) or nicotine pouches (n = 11) were not asked the question.

^{§§§§} Current users were asked, “Did any of the flavors that you used in the past 30 days have a name that did not describe a specific flavor, such as ‘solar,’ ‘purple,’ ‘jazz,’ ‘island bash,’ ‘fusion,’ or some other word or phrase?” Those who reported using only “unflavored” e-cigarettes (n = 60) or nicotine pouches (n = 11) were not asked the question.

^{¶¶¶} 45 C.F.R. part 46.102(l)(2), 21 C.F.R. part 56; 42 U.S.C. Sect. 241(d); 5 U.S.C. Sect. 552a; 44 U.S.C. Sect. 3501 et seq.

^{****} Estimates among non-Hispanic Asian and non-Hispanic Native Hawaiian or other Pacific Islander students were statistically unreliable for all current measures and are not reported.

TABLE 1. Percentage of middle and high school students who reported ever using tobacco products,* by product, overall and by school level, sex, and race and ethnicity — National Youth Tobacco Survey, United States, 2023

Tobacco product	% (95% CI)									Total estimated weighted no. [§]
	Sex		Race and ethnicity [†]						Total	
	Female	Male	AI/AN, NH	Asian, NH	Black or African American, NH	White, NH	Hispanic or Latino	Multiracial, NH		
Overall										
Any tobacco product [¶]	23.7 (21.5–26.0)	20.8 (18.9–22.8)	22.7 (16.8–30.0)	12.1 (6.5–21.5)	20.1 (17.7–22.6)	23.1 (20.2–26.2)	23.8 (22.2–25.4)	27.9 (22.5–33.9)	22.2 (20.5–23.9)	6,210,000
E-cigarettes	19.4 (17.5–21.5)	14.7 (13.2–16.3)	15.4 (10.7–21.8)	—**	12.9 (11.1–14.8)	18.4 (15.9–21.1)	18.2 (16.3–20.2)	20.8 (15.9–26.8)	17.0 (15.6–18.5)	4,750,000
Cigarettes	7.0 (6.0–8.1)	6.5 (5.4–7.7)	9.5 (5.6–15.5)	—	4.1 (2.9–5.8)	7.5 (6.3–8.9)	7.4 (5.9–9.2)	8.7 (6.0–12.4)	6.7 (6.0–7.6)	1,840,000
Cigars ^{††}	3.8 (2.9–4.8)	5.8 (4.8–7.0)	—	—	4.7 (3.4–6.4)	5.2 (4.1–6.6)	4.7 (4.0–5.5)	6.9 (4.8–9.8)	4.8 (4.0–5.6)	1,300,000
Hookahs	3.4 (2.4–4.8)	2.7 (1.9–3.8)	—	—	4.5 (2.7–7.2)	2.5 (1.7–3.5)	3.5 (2.7–4.5)	3.6 (2.4–5.2)	3.0 (2.4–3.9)	820,000
Smokeless tobacco (composite) ^{††}	2.2 (1.7–2.9)	3.7 (2.8–4.8)	—	—	1.3 (0.8–2.1)	3.4 (2.5–4.6)	2.9 (2.2–3.8)	5.0 (3.3–7.5)	3.0 (2.4–3.6)	800,000
Other oral nicotine products ^{††}	2.7 (2.1–3.4)	3.2 (2.6–4.1)	4.9 (2.8–8.5)	—	1.7 (1.1–2.6)	3.2 (2.4–4.1)	3.5 (2.7–4.6)	4.2 (2.4–7.2)	3.0 (2.5–3.5)	800,000
Nicotine pouches	1.7 (1.2–2.4)	3.0 (2.2–4.1)	—	—	—	3.0 (2.3–3.9)	2.0 (1.2–3.2)	—	2.3 (1.8–3.0)	580,000
Pipe tobacco	1.5 (1.1–2.0)	1.9 (1.4–2.5)	—	—	—	1.8 (1.3–2.5)	2.0 (1.5–2.7)	2.3 (1.3–3.9)	1.7 (1.4–2.0)	440,000
Heated tobacco products	1.5 (1.1–2.0)	1.5 (1.0–2.1)	—	—	1.7 (1.0–2.9)	1.4 (0.9–2.0)	1.8 (1.3–2.4)	1.6 (0.9–3.0)	1.5 (1.1–2.0)	370,000
Any combustible tobacco product ^{§§}	10.9 (9.3–12.8)	11.6 (10.1–13.2)	11.1 (7.0–17.1)	4.4 (2.4–7.8)	11.2 (8.5–14.7)	11.6 (9.7–13.7)	12.0 (10.4–13.8)	14.4 (11.0–18.5)	11.2 (9.9–12.7)	3,090,000
Multiple tobacco products ^{¶¶}	10.1 (8.7–11.8)	9.6 (8.4–10.9)	11.0 (7.2–16.3)	3.6 (2.1–6.0)	7.3 (5.5–9.7)	10.8 (9.1–12.8)	10.3 (8.9–11.8)	13.3 (10.1–17.3)	9.8 (8.7–11.1)	2,750,000
High school students (grades 9–12)										
Any tobacco product [¶]	30.1 (26.9–33.5)	25.9 (23.5–28.5)	29.0 (19.1–41.5)	—	21.8 (18.8–25.2)	31.4 (28.0–34.9)	27.3 (24.8–29.8)	35.1 (27.3–43.7)	27.9 (25.8–30.2)	4,390,000
E-cigarettes	26.0 (23.2–29.0)	19.5 (17.6–21.5)	20.3 (12.5–31.2)	—	14.7 (11.7–18.2)	26.0 (23.0–29.2)	22.3 (20.0–24.9)	27.5 (20.9–35.3)	22.6 (20.9–24.5)	3,550,000
Cigarettes	8.8 (7.3–10.6)	8.3 (7.0–9.7)	—	—	3.0 (1.8–5.0)	10.5 (9.0–12.1)	8.8 (6.9–11.1)	10.5 (6.8–15.7)	8.5 (7.7–9.5)	1,310,000
Cigars ^{††}	4.8 (3.6–6.4)	7.9 (6.3–9.9)	—	—	4.8 (3.2–7.1)	7.8 (6.1–10.0)	5.4 (4.4–6.6)	9.6 (6.4–14.0)	6.4 (5.3–7.7)	980,000
Hookahs	4.0 (2.7–5.9)	3.5 (2.3–5.4)	—	—	—	3.6 (2.5–5.3)	3.9 (2.7–5.5)	3.3 (1.9–5.8)	3.7 (2.8–5.1)	560,000
Smokeless tobacco (composite) ^{††}	2.2 (1.5–3.2)	4.3 (3.3–5.7)	—	—	—	3.8 (2.8–5.1)	2.9 (2.1–4.0)	6.9 (4.1–11.4)	3.3 (2.6–4.1)	500,000
Other oral nicotine products ^{††}	2.8 (2.0–4.0)	4.0 (3.1–5.3)	—	—	1.6 (0.9–2.7)	4.1 (3.0–5.4)	3.8 (3.0–4.8)	—	3.5 (2.8–4.2)	520,000
Nicotine pouches	2.0 (1.4–2.9)	4.1 (3.0–5.6)	—	—	—	4.5 (3.5–5.7)	1.8 (1.1–2.8)	—	3.1 (2.4–4.0)	430,000
Pipe tobacco	1.7 (1.2–2.5)	2.4 (1.8–3.2)	—	—	—	2.7 (2.0–3.5)	2.2 (1.5–3.2)	3.3 (2.0–5.5)	2.1 (1.7–2.5)	310,000
Heated tobacco products	1.7 (1.2–2.5)	1.6 (1.0–2.4)	—	—	—	1.8 (1.2–2.8)	1.5 (0.9–2.3)	—	1.6 (1.2–2.3)	230,000
Any combustible tobacco product ^{§§}	13.6 (11.3–16.2)	14.9 (13.0–16.9)	—	—	10.7 (8.2–14.0)	16.4 (14.1–19.1)	13.8 (11.7–16.3)	17.5 (12.6–23.7)	14.2 (12.6–16.1)	2,190,000
Multiple tobacco products ^{¶¶}	12.8 (10.5–15.4)	12.6 (11.1–14.2)	14.2 (8.0–24.0)	4.6 (2.5–8.3)	7.1 (4.9–10.1)	15.4 (13.1–18.1)	11.7 (10.0–13.6)	17.1 (12.2–23.3)	12.7 (11.1–14.4)	1,990,000

See table footnotes on the next page.

TABLE 1. (Continued) Percentage of middle and high school students who reported ever using tobacco products,* by product, overall and by school level, sex, and race and ethnicity — National Youth Tobacco Survey, United States, 2023

Tobacco product	% (95% CI)									Total estimated weighted no. [§]
	Sex		Race and ethnicity [†]						Total	
	Female	Male	AI/AN, NH	Asian, NH	Black or African American, NH	White, NH	Hispanic or Latino	Multiracial, NH		
Middle school students (grades 6–8)										
Any tobacco product [¶]	15.4 (12.9–18.3)	13.8 (11.3–16.6)	15.3 (9.7–23.2)	—	17.8 (12.9–24.0)	12.3 (10.0–14.9)	18.7 (16.5–21.1)	17.6 (13.0–23.6)	14.7 (12.5–17.1)	1,780,000
E-cigarettes	11.0 (9.1–13.3)	8.2 (6.9–9.8)	—	—	10.6 (8.5–13.1)	8.4 (6.8–10.3)	12.3 (10.5–14.4)	11.3 (6.3–19.5)	9.7 (8.3–11.3)	1,170,000
Cigarettes	4.6 (3.6–5.9)	4.0 (2.7–5.9)	—	—	5.5 (3.9–7.8)	3.5 (2.5–5.1)	5.3 (3.8–7.2)	—	4.3 (3.3–5.5)	510,000
Cigars ^{††}	2.4 (1.6–3.6)	2.9 (2.0–4.2)	—	—	4.6 (2.8–7.4)	1.7 (1.1–2.6)	3.5 (2.3–5.3)	—	2.6 (1.9–3.7)	310,000
Hookahs	—	1.7 (1.2–2.3)	—	—	—	0.9 (0.5–1.6)	2.9 (2.1–4.0)	—	2.1 (1.4–3.2)	240,000
Smokeless tobacco (composite) ^{††}	2.3 (1.6–3.1)	2.7 (1.8–4.0)	—	—	—	2.9 (1.9–4.4)	2.5 (1.6–3.9)	—	2.4 (1.8–3.3)	290,000
Other oral nicotine products ^{††}	2.4 (1.8–3.2)	2.1 (1.6–2.7)	—	—	—	2.0 (1.4–2.9)	2.9 (1.8–4.4)	2.9 (1.6–5.2)	2.2 (1.8–2.7)	260,000
Nicotine pouches	—	—	—	—	—	1.0 (0.6–1.8)	—	—	—	—
Pipe tobacco	1.1 (0.6–2.0)	1.1 (0.6–2.0)	—	—	—	—	1.7 (1.2–2.4)	—	1.1 (0.7–1.6)	120,000
Heated tobacco products	1.2 (0.7–1.9)	—	—	—	—	0.8 (0.5–1.5)	2.1 (1.6–2.8)	—	1.2 (0.8–1.8)	130,000
Any combustible tobacco product ^{§§}	7.5 (5.7–10.0)	7.2 (5.1–9.9)	6.6 (3.6–11.7)	—	11.9 (7.0–19.4)	5.3 (3.8–7.3)	9.3 (7.3–11.7)	9.8 (6.4–14.8)	7.3 (5.6–9.4)	870,000
Multiple tobacco products ^{¶¶}	6.7 (5.3–8.6)	5.5 (4.2–7.2)	—	—	7.6 (4.7–12.2)	4.7 (3.5–6.2)	8.0 (6.0–10.6)	7.9 (5.3–11.6)	6.1 (4.9–7.5)	740,000

Abbreviations: AI/AN = American Indian or Alaska Native; NH = non-Hispanic.

* Ever use is defined as ever having used the product, even once or twice. Because of missing data on the ever use questions, denominators for each tobacco product might be different. For each question, response options were “yes” or “no.”

† Hispanic or Latino persons could be of any race. Estimates among NH Native Hawaiian or other Pacific Islander students, overall and by school level, were statistically unreliable for all measures and are not presented in this table.

§ Estimated weighted total number of ever tobacco product users was rounded down to the nearest 10,000 persons. Overall estimates were reported based on 22,069 U.S. middle and high school students. School level was determined by reported grade level: high school (grades 9–12; n = 10,879) and middle school (grades 6–8; n = 11,067). The sum of subgroup estimates might not sum to overall population estimates because of rounding or exclusion of students who did not report sex, race and ethnicity, or grade level.

¶ Any tobacco product use is defined as ever use of one or more of the following tobacco products: e-cigarettes, cigars, cigarettes, smokeless tobacco (composite), hookahs, nicotine pouches, heated tobacco products, pipe tobacco, bidis (small brown cigarettes wrapped in a leaf), or other oral nicotine products.

** Dashes indicate that data were statistically unreliable because of an unweighted denominator <50 or a relative SE >30%.

†† Cigars were defined as cigars, cigarillos, or little cigars. Smokeless tobacco (composite) was defined as chewing tobacco, snuff, dip, or snus. Other oral nicotine products were defined as lozenges, discs, tablets, gums, dissolvable tobacco products, and other products. In 2023, dissolvable tobacco products were reclassified from smokeless tobacco to other oral nicotine products.

§§ Any combustible tobacco product use was defined as ever use of one or more of the following tobacco products: cigarettes, cigars, hookahs, pipe tobacco, or bidis.

¶¶ Multiple tobacco product use was defined as ever use of two or more of the following tobacco products: e-cigarettes, cigars, cigarettes, smokeless tobacco (composite), hookahs, nicotine pouches, heated tobacco products, pipe tobacco, bidis, or other oral nicotine products.

significant change in current use of any composite measure or individual tobacco product was observed.

Discussion

Current use of any tobacco product by high school students declined by an estimated 540,000 students, from 2.51 million in 2022 (3) to 1.97 million in 2023. In 2023, 22.2% of middle and high school students (representing 6.21 million) reported

ever using any tobacco product, and 10.0% of students (representing 2.80 million) reported current use of any tobacco product. Similar to 2022 (3), ever use of any tobacco product was lowest among non-Hispanic Asian students and did not differ significantly across most racial and ethnic groups.

E-cigarettes have been the most commonly used tobacco product among U.S. youths since 2014 (4). Youth e-cigarette use is a critical public health concern, because approximately

TABLE 2. Percentage of middle and high school students who reported current (past 30-day) tobacco product use, by product,* overall and by school level, sex, and race and ethnicity — National Youth Tobacco Survey, United States, 2023

Tobacco product	% (95% CI)							Total estimated weighted no. [§]	
	Sex		Race and ethnicity [†]						
	Female	Male	AI/AN, NH	Black or African American, NH	White, NH	Hispanic or Latino	Multiracial, NH		Total
Overall									
Any tobacco product [¶]	11.2 (9.5–13.1)	8.9 (7.7–10.3)	8.0 (4.7–13.2)	9.3 (7.5–11.3)	9.5 (7.7–11.6)	11.7 (10.1–13.4)	12.6 (8.8–17.7)	10.0 (8.9–11.2)	2,800,000
E-cigarettes	9.3 (8.1–10.8)	6.1 (5.0–7.4)	5.9 (3.4–10.0)	5.6 (4.5–7.1)	7.7 (6.3–9.4)	8.5 (7.4–9.8)	10.2 (6.8–15.1)	7.7 (6.8–8.6)	2,130,000
Cigarettes	1.4 (1.0–1.9)	1.8 (1.4–2.4)	— ^{**}	—	1.6 (1.1–2.3)	2.1 (1.5–3.1)	1.6 (1.0–2.8)	1.6 (1.2–2.1)	430,000
Cigars	1.3 (0.9–2.0)	1.8 (1.4–2.3)	—	2.3 (1.4–3.8)	1.0 (0.7–1.4)	2.2 (1.7–2.8)	—	1.6 (1.2–2.0)	420,000
Nicotine pouches	0.8 (0.5–1.3)	2.1 (1.5–3.0)	—	—	1.4 (0.9–2.2)	1.9 (1.1–3.3)	—	1.5 (1.0–2.1)	400,000
Smokeless tobacco (composite) ^{††}	—	1.6 (1.1–2.3)	—	—	1.2 (0.7–1.8)	1.6 (1.1–2.4)	—	1.2 (0.9–1.7)	330,000
Other oral nicotine products	1.1 (0.9–1.4)	1.2 (0.9–1.7)	0.5 (0.3–0.8)	—	1.2 (0.9–1.5)	1.5 (1.1–2.0)	—	1.2 (1.0–1.4)	310,000
Hookahs	1.3 (0.8–2.1)	0.9 (0.6–1.3)	—	—	0.7 (0.4–1.1)	1.3 (1.0–1.7)	1.3 (0.7–2.4)	1.1 (0.8–1.5)	290,000
Heated tobacco products	0.7 (0.5–1.0)	1.2 (0.8–1.9)	—	1.0 (0.5–1.7)	0.7 (0.4–1.2)	1.5 (1.0–2.2)	—	1.0 (0.7–1.3)	260,000
Pipe tobacco	0.5 (0.3–0.7)	0.6 (0.4–0.9)	—	—	0.5 (0.3–0.8)	0.9 (0.5–1.4)	—	0.5 (0.4–0.7)	130,000
Any combustible tobacco product ^{§§}	3.3 (2.6–4.1)	3.5 (2.9–4.2)	2.7 (1.5–4.9)	4.7 (3.1–7.0)	2.7 (2.2–3.5)	3.9 (3.1–4.8)	3.7 (2.4–5.5)	3.4 (2.9–4.0)	920,000
Multiple tobacco products ^{¶¶}	3.4 (2.7–4.2)	3.4 (2.7–4.2)	2.0 (1.1–3.5)	3.2 (1.8–5.5)	3.1 (2.4–4.0)	3.9 (3.4–4.5)	4.1 (2.6–6.5)	3.4 (2.9–3.9)	940,000
High school students (grades 9–12)									
Any tobacco product [¶]	14.1 (11.6–17.0)	11.2 (9.4–13.1)	—	9.8 (7.7–12.5)	13.6 (11.2–16.5)	12.4 (10.6–14.4)	17.2 (11.3–25.3)	12.6 (11.1–14.3)	1,970,000
E-cigarettes	12.2 (10.3–14.5)	8.0 (6.3–10.0)	—	5.6 (4.2–7.4)	11.3 (9.2–13.7)	9.7 (8.0–11.8)	14.2 (9.0–21.8)	10.0 (8.8–11.4)	1,560,000
Cigarettes	1.5 (1.0–2.2)	2.3 (1.8–2.9)	—	—	2.2 (1.4–3.4)	2.2 (1.6–3.0)	—	1.9 (1.5–2.4)	290,000
Cigars	1.4 (0.8–2.3)	2.3 (1.7–3.0)	—	1.9 (1.2–3.0)	1.4 (0.9–2.2)	2.3 (1.6–3.3)	—	1.8 (1.4–2.4)	280,000
Nicotine pouches	—	2.6 (1.9–3.6)	—	—	2.2 (1.4–3.4)	1.6 (0.9–2.7)	—	1.7 (1.2–2.5)	260,000
Smokeless tobacco (composite) ^{††}	—	2.1 (1.4–3.0)	—	—	1.7 (1.1–2.6)	1.7 (1.1–2.7)	—	1.5 (1.1–2.2)	230,000
Other oral nicotine products	0.9 (0.7–1.3)	1.5 (1.0–2.2)	—	—	1.3 (1.0–1.8)	1.6 (1.1–2.2)	—	1.2 (1.0–1.6)	180,000
Hookahs	1.4 (0.8–2.4)	0.9 (0.6–1.5)	—	—	—	1.0 (0.6–1.6)	—	1.1 (0.8–1.6)	170,000
Heated tobacco products	0.7 (0.4–1.2)	1.4 (0.8–2.5)	—	—	—	1.6 (0.9–2.7)	—	1.0 (0.7–1.6)	150,000
Pipe tobacco	0.5 (0.3–0.9)	0.7 (0.4–1.2)	—	—	0.6 (0.4–1.0)	—	—	0.6 (0.4–0.9)	90,000
Any combustible tobacco product ^{§§}	3.6 (2.7–4.7)	4.3 (3.6–5.2)	—	4.5 (3.2–6.2)	3.8 (3.0–5.0)	3.8 (2.8–5.0)	5.3 (3.3–8.6)	3.9 (3.4–4.6)	600,000
Multiple tobacco products ^{¶¶}	3.5 (2.7–4.7)	4.3 (3.4–5.5)	—	—	4.3 (3.2–5.7)	3.9 (3.0–5.1)	6.1 (3.6–10.2)	3.9 (3.3–4.7)	610,000

See table footnotes on the next page.

one half of students ever using e-cigarettes reported using them currently, indicating that many young persons who try e-cigarettes remain e-cigarette users. In 2023, 10.0% of high school students and 4.6% of middle school students used e-cigarettes during the past 30 days. From 2022 (3) to 2023, a significant decline in current e-cigarette use occurred among high school students (from 14.1% to 10.0%), while no statistically

significant change occurred among middle school students (from 3.3% in 2022 to 4.6% in 2023). The decline since 2022 in high school student e-cigarette use is likely attributable to multiple factors, such as ongoing efforts at the national, state, and local levels to implement tobacco control strategies, including Food and Drug Administration (FDA) regulatory actions. Continued surveillance is needed to determine the trajectory of

TABLE 2. (Continued) Percentage of middle and high school students who reported current (past 30-day) tobacco product use, by product,* overall and by school level, sex, and race and ethnicity — National Youth Tobacco Survey, United States, 2023

Tobacco product	% (95% CI)							Total	Total estimated weighted no. [§]
	Sex		Race and ethnicity [†]						
	Female	Male	AI/AN, NH	Black or African American, NH	White, NH	Hispanic or Latino	Multiracial, NH		
Middle school students (grades 6–8)									
Any tobacco product [¶]	7.5 (5.9–9.4)	5.7 (4.1–8.0)	—	8.5 (5.8–12.4)	4.1 (3.2–5.2)	10.3 (7.5–14.0)	6.0 (3.4–10.5)	6.6 (5.1–8.5)	800,000
E-cigarettes	5.6 (4.5–7.1)	3.5 (2.5–4.8)	—	5.7 (3.9–8.2)	3.1 (2.2–4.2)	6.6 (5.3–8.2)	—	4.6 (3.6–5.8)	550,000
Cigarettes	1.1 (0.7–1.9)	—	—	—	0.8 (0.4–1.4)	—	—	1.1 (0.6–1.9)	120,000
Cigars	1.2 (0.7–2.2)	1.0 (0.6–1.8)	—	—	—	1.8 (1.1–3.1)	—	1.1 (0.7–1.8)	130,000
Nicotine pouches	—	—	—	—	0.5 (0.3–0.8)	—	—	—	—
Smokeless tobacco (composite) ^{††}	0.6 (0.4–1.0)	—	—	—	—	—	—	0.7 (0.5–1.2)	80,000
Other oral nicotine products	1.3 (0.9–1.8)	0.8 (0.6–1.3)	—	—	1.0 (0.6–1.6)	1.3 (0.8–2.0)	—	1.1 (0.8–1.4)	120,000
Hookahs	—	0.8 (0.5–1.5)	—	—	0.4 (0.2–0.7)	1.8 (1.1–2.9)	—	1.0 (0.6–1.8)	120,000
Heated tobacco products	0.8 (0.4–1.3)	—	—	—	—	1.3 (0.8–2.3)	—	0.8 (0.5–1.4)	90,000
Pipe tobacco	—	—	—	—	—	—	—	0.4 (0.2–0.6)	40,000
Any combustible tobacco product ^{§§}	2.8 (1.8–4.4)	2.3 (1.4–3.6)	—	—	1.3 (0.9–1.9)	3.7 (2.4–5.6)	—	2.5 (1.7–3.8)	300,000
Multiple tobacco products ^{¶¶}	3.1 (2.2–4.4)	2.0 (1.3–3.0)	—	—	1.5 (1.0–2.2)	3.5 (2.5–4.9)	—	2.5 (1.8–3.5)	300,000

Abbreviations: AI/AN = American Indian or Alaska Native; NH = non-Hispanic.

* Current use is defined as use on ≥ 1 days during the past 30 days for each product. Because of missing data on past 30-day use questions, denominators for each tobacco product might be different.

[†] Hispanic or Latino persons could be of any race. Estimates among NH Asian and NH Native Hawaiian or other Pacific Islander students, overall and by school level, were statistically unreliable for all measures and are not presented in this table.

[§] Estimated weighted total number of current tobacco product users was rounded down to the nearest 10,000 persons. Overall estimates were reported based on 22,069 U.S. middle and high school students. School level was determined by reported grade level: high school (grades 9–12; $n = 10,879$) and middle school (grades 6–8; $n = 11,067$). The sum of subgroup estimates might not sum to overall population estimates because of rounding or exclusion of students who did not report sex, race and ethnicity, or grade level.

[¶] Any tobacco product use is defined as current use of one or more of the following tobacco products on ≥ 1 days during the past 30 days: e-cigarettes, cigars, cigarettes, smokeless tobacco (composite), hookahs, nicotine pouches, heated tobacco products, pipe tobacco, bidis (small brown cigarettes wrapped in a leaf), or other oral nicotine products.

** Dashes indicate that data were statistically unreliable because of an unweighted denominator < 50 or a relative SE $> 30\%$.

^{††} Cigars were defined as cigars, cigarillos, or little cigars. Smokeless tobacco (composite) was defined as chewing tobacco, snuff, dip, or snus. Other oral nicotine products were defined as lozenges, discs, tablets, gums, dissolvable tobacco products, and other products. In 2023, dissolvable tobacco products were reclassified from smokeless tobacco to other oral nicotine products.

^{§§} Any combustible tobacco product use was defined as current use of one or more of the following tobacco products: cigarettes, cigars, hookahs, pipe tobacco, or bidis.

^{¶¶} Multiple tobacco product use was defined as current use of two or more of the following tobacco products: e-cigarettes, cigars, cigarettes, smokeless tobacco (composite), hookahs, nicotine pouches, heated tobacco products, pipe tobacco, bidis, or other oral nicotine products.

middle school e-cigarette use. Despite the decline in e-cigarette use among high school students, close to 40% of high school students using e-cigarettes reported frequent use, and 29.9% reported daily use. Furthermore, 550,000 middle school students currently used e-cigarettes, including 20.7% reporting frequent use. Similar patterns were observed in 2022 for both middle school and high school students. These findings are concerning, because adolescents have reported symptoms of nicotine dependence when using tobacco products only 1–3 days per month (1). Efforts aimed at reducing nicotine dependence among adolescents by preventing initiation of tobacco products is important (5).

Among students who reported current e-cigarette use, disposables were the most commonly used device type. Disposable e-cigarettes have been gaining market share; they are relatively inexpensive, have a high nicotine content, and are available in flavors appealing to youths (e.g., fruit and candy) (6). In January 2020, FDA announced that it would prioritize enforcement against prefilled e-cigarettes in flavors other than tobacco and menthol (7). In 2023, NYTS for the first time assessed tobacco-flavored product use, use of flavors that included the word “ice” or “iced” in their name, and use of concept flavors. These results, combined with results of other flavored tobacco

TABLE 3. Percentage of middle and high school students reporting current (past 30-day) e-cigarette use,* overall by selected characteristics and school level — National Youth Tobacco Survey, United States, 2023

Characteristic	Overall		High school		Middle school	
	Estimated weighted no. [†]	% (95% CI)	Estimated weighted no. [†]	% (95% CI)	Estimated weighted no. [†]	% (95% CI)
Among all students	2,130,000	7.7 (6.8–8.6)	1,560,000	10.0 (8.8–11.4)	550,000	4.6 (3.6–5.8)
Among current e-cigarette users						
Frequency of use during past 30 days						
1–5 days	980,000	46.1 (39.8–52.7)	630,000	40.7 (33.1–48.7)	340,000	62.0 (55.7–67.9)
6–19 days	400,000	19.1 (15.0–24.1)	300,000	19.7 (14.1–26.8)	90,000	17.3 (12.2–24.0)
20–30 days	740,000	34.7 (28.4–41.7)	620,000	39.7 (31.3–48.6)	110,000	20.7 (14.6–28.6)
Daily e-cigarette use[§]	530,000	25.2 (19.2–32.3)	460,000	29.9 (22.1–39.1)	60,000	11.4 (7.5–17.0)
Device type most often used[¶]						
Disposables	1,240,000	60.7 (53.3–67.6)	1,000,000	65.2 (56.3–73.1)	240,000	47.9 (39.5–56.5)
Prefilled or refillable pods or cartridges	330,000	16.1 (12.2–21.0)	240,000	16.0 (11.1–22.5)	80,000	16.7 (11.4–23.8)
Tanks or mod system	120,000	5.9 (4.4–7.8)	90,000	6.0 (4.3–8.4)	20,000	4.4 (2.5–7.5)
Don't know the type	350,000	17.3 (12.7–23.1)	190,000	12.8 (8.7–18.4)	150,000	31.1 (22.2–41.5)
Any brand**						
Elf Bar	1,160,000	56.7 (50.6–62.6)	900,000	59.1 (52.9–65.1)	260,000	50.0 (37.5–62.5)
Esco Bars	440,000	21.6 (16.2–28.3)	370,000	24.9 (18.1–33.1)	60,000	12.0 (6.9–20.2)
Vuse	420,000	20.7 (16.4–25.9)	330,000	22.2 (16.9–28.6)	80,000	16.3 (10.8–23.8)
JUUL	330,000	16.5 (12.9–20.9)	240,000	16.3 (11.8–22.1)	80,000	16.8 (11.4–24.1)
Mr. Fog	280,000	13.6 (7.9–22.4)	230,000	15.1 (8.2–26.3)	— ^{††}	—
SMOK (including NOVO)	230,000	11.3 (6.3–19.5)	—	—	30,000	6.7 (3.9–11.1)
Breeze	230,000	11.6 (7.6–17.4)	200,000	13.2 (8.0–21.2)	30,000	6.6 (4.1–10.5)
Kangvape (including Onee Stick)	180,000	8.8 (6.6–11.7)	130,000	8.7 (6.5–11.6)	—	—
Fume	180,000	9.0 (6.4–12.6)	140,000	9.2 (6.0–13.9)	40,000	8.2 (4.7–14.0)
NJOY	150,000	7.5 (5.5–10.3)	120,000	8.1 (5.6–11.7)	20,000	5.4 (3.1–9.2)
blu	120,000	6.0 (4.4–8.3)	70,000	5.2 (3.4–7.8)	40,000	8.1 (5.0–12.9)
HQD	110,000	5.5 (3.4–8.5)	80,000	5.7 (3.3–9.7)	—	—
Logic	80,000	3.9 (2.5–6.1)	50,000	3.7 (2.3–5.8)	—	—
Suorin (including Air Bar)	70,000	3.8 (2.5–5.6)	50,000	3.8 (2.3–6.3)	—	—
Lost Mary ^{§§}	50,000	2.6 (1.4–4.8)	40,000	3.3 (1.8–5.9)	—	—
Some other brand not listed	350,000	17.3 (11.6–24.9)	290,000	19.5 (12.4–29.2)	50,000	10.9 (6.0–19.0)
Not sure or don't know the brand	490,000	23.9 (19.3–29.2)	300,000	19.8 (15.6–24.9)	180,000	35.4 (24.3–48.3)
Usual brand^{¶¶}						
Elf Bar	630,000	31.1 (26.2–36.4)	460,000	30.2 (24.8–36.2)	170,000	33.9 (24.6–44.7)
Vuse	170,000	8.7 (5.8–12.9)	150,000	10.0 (6.4–15.3)	—	—
Esco Bars	120,000	6.0 (3.4–10.4)	110,000	7.7 (4.3–13.5)	—	—
JUUL	70,000	3.4 (1.9–6.1)	—	—	—	—
Mr. Fog	—	—	—	—	—	—
SMOK (including NOVO)	—	—	—	—	—	—
Breeze	—	—	—	—	—	—
Kangvape (including Onee Stick)	—	—	—	—	—	—
Fume	—	—	—	—	—	—
NJOY	—	—	—	—	—	—
blu	—	—	—	—	—	—
HQD	—	—	—	—	—	—
Logic	—	—	—	—	—	—
Suorin (including Air Bar)	—	—	—	—	—	—
Lost Mary	—	—	—	—	—	—
No usual brand	90,000	4.4 (2.8–7.0)	70,000	4.7 (2.8–7.5)	—	—
Some other brand not listed	270,000	13.2 (7.8–21.5)	230,000	15.4 (8.6–25.9)	—	—
Not sure or don't know the brand	400,000	19.8 (16.0–24.1)	240,000	16.0 (12.5–20.4)	150,000	30.5 (22.1–40.4)
Flavored e-cigarette use^{***}						
Any flavor other than tobacco-flavored or unflavored	1,900,000	89.4 (86.2–91.9)	1,410,000	90.3 (86.6–93.1)	480,000	87.1 (79.9–92.0)
Exclusive use of tobacco-flavored or unflavored	110,000	5.6 (3.9–7.9)	80,000	5.4 (3.4–8.4)	30,000	6.2 (3.4–11.0)
Unspecified	100,000	5.0 (3.5–7.2)	60,000	4.3 (2.8–6.6)	—	—

See table footnotes on the next page.

TABLE 3. (Continued) Percentage of middle and high school students reporting current (past 30-day) e-cigarette use,* overall by selected characteristics and school level — National Youth Tobacco Survey, United States, 2023

Characteristic	Overall		High school		Middle school	
	Estimated weighted no. [†]	% (95% CI)	Estimated weighted no. [†]	% (95% CI)	Estimated weighted no. [†]	% (95% CI)
Flavor type used among current e-cigarette users^{†††}						
Fruit	1,280,000	63.4 (59.8–66.9)	930,000	62.6 (57.9–67.0)	340,000	66.3 (59.5–72.5)
Candy, desserts, or other sweets	700,000	35.0 (29.1–41.5)	510,000	34.4 (27.5–42.1)	190,000	37.0 (28.6–46.4)
Mint	560,000	27.8 (22.0–34.4)	470,000	31.6 (24.2–40.1)	80,000	16.5 (11.6–22.9)
Menthol	400,000	20.1 (15.5–25.8)	340,000	23.3 (17.6–30.1)	50,000	10.4 (7.2–14.8)
Unflavored	230,000	11.6 (8.8–15.1)	160,000	10.9 (7.8–15.0)	60,000	13.2 (8.7–19.5)
Non-alcoholic drinks ^{§§§}	220,000	11.3 (6.4–19.1)	—	—	30,000	7.4 (4.1–13.0)
Alcoholic drinks ^{§§§}	170,000	8.4 (5.5–12.7)	130,000	9.0 (5.5–14.4)	—	—
Tobacco-flavored	120,000	6.4 (4.5–9.0)	70,000	5.3 (3.7–7.4)	—	—
Clove or spice ^{§§§}	120,000	6.0 (4.3–8.2)	70,000	5.1 (3.3–7.7)	40,000	7.9 (4.6–13.1)
Chocolate	90,000	4.9 (3.4–7.1)	50,000	3.4 (1.9–6.2)	40,000	8.0 (4.3–14.3)
Some other flavor	120,000	6.0 (4.2–8.5)	60,000	4.6 (3.0–7.0)	50,000	10.0 (5.9–16.4)
Use of any flavors that included the word “ice” or “iced” (such as “blueberry ice” or “strawberry ice”)^{¶¶¶}						
Yes	1,100,000	57.9 (52.5–63.1)	800,000	57.0 (51.3–62.6)	290,000	61.0 (52.8–68.5)
No	560,000	29.5 (24.8–34.8)	440,000	31.6 (26.1–37.7)	110,000	24.1 (18.2–31.1)
Don't know	230,000	12.6 (9.8–16.0)	160,000	11.4 (8.3–15.5)	70,000	15.0 (10.2–21.5)
Use of any concept flavors with a name that did not describe a specific flavor (such as “solar,” “purple,” “jazz,” “island bash,” or “fusion”)^{****}						
Yes	300,000	16.1 (13.5–19.0)	210,000	15.4 (12.2–19.3)	80,000	17.7 (12.3–24.7)
No	1,110,000	58.5 (52.5–64.3)	850,000	60.9 (52.8–68.4)	250,000	52.8 (45.5–60.0)
Don't know	480,000	25.4 (21.6–29.7)	330,000	23.7 (19.0–29.2)	140,000	29.5 (22.6–37.5)

* Current (past 30-day) use of e-cigarettes was determined by asking, “During the past 30 days, on how many days did you use e-cigarettes?” Current use was defined as use on ≥1 days during the past 30 days.

[†] Estimated total number of users was rounded down to the nearest 10,000 persons. The sum of subgroup estimates might not sum to overall population estimates because of rounding or exclusion of students who did not report grade level (n = 102), device type (n = 53), any brand (n = 54), usual brand (n = 61), flavor types used (n = 84), use of flavor including the word “ice” or “iced” (n = 136), or use of flavors without specific flavor descriptor (n = 143).

[§] Daily e-cigarette use was defined as use on all 30 of the past 30 days.

[¶] Device type was determined by the question, “Which of the following best describes the type of e-cigarette you have used in the past 30 days? If you have used more than one type, please think about the one you use most often.”

^{**} All current e-cigarette users were asked, “During the past 30 days, what e-cigarette brands did you use? (Select one or more).” Those who selected “some other brand(s) not listed here” could provide a write-in response. Write-in responses corresponding to an original response option were recoded.

^{††} Data were statistically unreliable because of an unweighted denominator <50 or a relative SE >30%.

^{§§} Lost Mary was not included in the list of prespecified response options, but it was the most commonly provided write-in response for “some other brand(s) not listed here.”

^{¶¶} If a single brand was selected for the question, “During the past 30 days, what e-cigarette brands did you use (Select one or more),” it was reported as the respondent’s usual brand. Those who selected one or more brands were asked, “During the past 30 days, what brand of e-cigarettes did you usually use? (Choose only one answer).” Those who selected “some other brand(s) not listed here” could provide a write-in response. Write-in responses corresponding to an original response option were recoded.

^{***} All current e-cigarette users were asked, “In the past 30 days when you used e-cigarettes, what flavors did you use? (Select one or more)?” Those who provided no valid responses were defined as “Unspecified” flavored users.

^{†††} Flavor type was determined by response to the question, “In the past 30 days when you used e-cigarettes, what flavors did you use? (Select one or more).” Those who selected “some other flavor not listed here” could provide a write-in response; write-in responses corresponding to an original response option were recoded.

^{§§§} These flavor options provided examples: “Alcoholic drinks (such as wine, margarita, or other cocktails);” “Non-alcoholic drinks (such as coffee, soda, lemonade, or other beverage);” and “Spice (such as cinnamon, vanilla, or clove).”

^{¶¶¶} Current e-cigarette users were asked, “Did any of the flavors you used in the past 30 days have names or descriptions that included the word ‘ice’ or ‘iced’ (for example, blueberry ice or strawberry ice)?” Those who reported using only unflavored e-cigarettes (n = 60) did not receive the question.

^{****} Current e-cigarette users were asked, “Did any of the flavors that you used in the past 30 days have a name that did not describe a specific flavor, such as ‘solar,’ ‘purple,’ ‘jazz,’ ‘island bash,’ and ‘fusion,’ or some other word or phrase?” Those who reported using only “unflavored” e-cigarettes (n = 60) did not receive the question.

product use research, continue to demonstrate the high appeal of flavored tobacco products among young persons.

Multiple factors continue to influence tobacco product use and initiation among middle and high school students, including availability of flavored products, marketing, and misperceptions regarding harm. Continued surveillance provides an understanding of the prevalence and frequency of tobacco

product use, the popularity of specific brands and flavors, and how product use behaviors change over time as the tobacco product marketplace continues to diversify.

Limitations

The findings in this report are subject to at least three limitations. First, data were obtained by self-report, which can result in social desirability and recall biases, although previous

Summary**What is already known about this topic?**

Use of tobacco products in any form by youths is unsafe.

What is added by this report?

In 2023, 10.0% of middle and high school students reported current tobacco product use. From 2022 to 2023, current e-cigarette use among high school students declined from 14.1% to 10.0%. E-cigarettes remained the most commonly used tobacco product among youths. Among middle school and high school students who currently use e-cigarettes, 25.2% used e-cigarettes daily, and 89.4% used flavored e-cigarettes.

What are the implications for public health?

Tobacco use declined among high school students; however, sustained public health monitoring with implementation of evidence-based tobacco control strategies, including effective youth interventions, media campaigns, Food and Drug Administration regulations, and other proven tobacco prevention policies might further reduce youth tobacco product use.

research suggests that self-reported measures of tobacco use among persons aged 12–21 years correlate with biomarkers of tobacco use (8). Second, these findings might not be generalizable to youths who are home-schooled, have dropped out of school, are in detention centers, or are enrolled in alternative schools. Finally, the response rate for the 2023 NYTS was lower than that for the 2022 NYTS (30.5% in 2023 versus 45.2% in 2022). The lower response rate can increase the potential for bias and result in higher SEs for some estimates; higher SEs can reduce the power to detect a significant difference, if there is one, when doing year to year comparisons, especially for certain population groups. Adjustments were made to the survey weights to reduce the potential for nonresponse bias. Therefore, 2023 NYTS estimates may be compared with 2022 NYTS estimates for the same population groups.

Implications for Public Health Practice

In 2023, 10.0% (representing 2.80 million) of U.S. middle and high school students reported current tobacco product use. A significant decline in current e-cigarette use occurred among high school students from 2022 to 2023 (from 14.1% to 10.0%). Given the negative health consequences of tobacco use (2) and the unique harms associated with adolescent nicotine exposure (1), prevention of tobacco use by youths is imperative. Thus, a continued comprehensive approach to

tobacco use prevention is needed to further reduce tobacco use among youths, based on knowledge about youth product use behaviors. Further, longstanding and proven tobacco prevention policies, such as price increases, comprehensive smoke-free policies (that include e-cigarettes), counter-marketing campaigns, and health care intervention, will continue to reduce youth initiation and tobacco use (5).

Corresponding author: Jan Birdsey, JBirdsey@cdc.gov.

¹Office on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion, CDC; ²Center for Tobacco Products, Food and Drug Administration, Silver Spring, Maryland.

All authors have completed and submitted the International Committee of Medical Journal Editors form for disclosure of potential conflicts of interest. No potential conflicts of interest were disclosed.

References

1. US Department of Health and Human Services. E-cigarette use among youth and young adults: a report of the Surgeon General. Atlanta, Georgia: US Department of Health and Human Services, CDC, 2016. <https://www.ncbi.nlm.nih.gov/books/NBK538680/>
2. US Department of Health and Human Services. 2014 Surgeon General's report: the health consequences of smoking—50 years of progress. Atlanta, GA: US Department of Health and Human Services, CDC, 2014. <https://www.cdc.gov/tobacco/sgt/50th-anniversary/index.htm>
3. Park-Lee E, Ren C, Cooper M, Cornelius M, Jamal A, Cullen KA. Tobacco product use among middle and high school students—United States, 2022. *MMWR Morb Mortal Wkly Rep* 2022;71:1429–35. PMID:36355596 <https://doi.org/10.15585/mmwr.mm7145a1>
4. Jamal A, Gentzke A, Hu SS, et al. Tobacco use among middle and high school students—United States, 2011–2016. *MMWR Morb Mortal Wkly Rep* 2017;66:597–603. PMID:28617771 <https://doi.org/10.15585/mmwr.mm6623a1>
5. Owens DK, Davidson KW, Krist AH, et al.; US Preventive Services Task Force. Primary care interventions for prevention and cessation of tobacco use in children and adolescents: US Preventive Services Task Force recommendation statement. *JAMA* 2020;323:1590–8. PMID:32343336 <https://doi.org/10.1001/jama.2020.4679>
6. Diaz MC, Silver NA, Bertrand A, Schillo BA. Bigger, stronger and cheaper: growth in e-cigarette market driven by disposable devices with more e-liquid, higher nicotine concentration and declining prices. *Tob Control* 2023;tc-2023-058033. PMID:37536928 <https://doi.org/10.1136/tc-2023-058033>
7. US Department of Health and Human Services. Enforcement priorities for electronic nicotine delivery systems (ENDS) and other deemed products on the market without premarket authorization (revised). Silver Spring, MD: US Department of Health and Human Services, Food and Drug Administration, 2020. <https://www.fda.gov/media/133880/download>
8. Boykan R, Messina CR, Chateau G, Eliscu A, Tolentino J, Goniewicz ML. Self-reported use of tobacco, e-cigarettes, and marijuana versus urinary biomarkers. *Pediatrics* 2019;143:e20183531. PMID:31010908 <https://doi.org/10.1542/peds.2018-3531>