

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,596,993	100.0	821.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	611,105	23.5	193.3
2	Malignant neoplasms (C00-C97)	584,881	22.5	185.0
3	Chronic lower respiratory diseases (J40-J47)	149,205	5.7	47.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130,557	5.0	41.3
5	Cerebrovascular diseases (I60-I69)	128,978	5.0	40.8
6	Alzheimer's disease (G30)	84,767	3.3	26.8
7	Diabetes mellitus (E10-E14)	75,578	2.9	23.9
8	Influenza and pneumonia (J09-J18)	56,979	2.2	18.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47,112	1.8	14.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41,149	1.6	13.0
11	Septicemia (A40-A41)	38,156	1.5	12.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	36,427	1.4	11.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30,770	1.2	9.7
14	Parkinson's disease (G20-G21)	25,196	1.0	8.0
15	Pneumonitis due to solids and liquids (J69)	18,579	0.7	5.9
...	All other causes (Residual)	537,554	20.7	170.0
All races, both sexes, 1-4 years				
...	All causes	4,068	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,316	32.4	8.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	476	11.7	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	337	8.3	2.1
4	Malignant neoplasms (C00-C97)	328	8.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	169	4.2	1.1
6	Influenza and pneumonia (J09-J18)	102	2.5	0.6
7	Chronic lower respiratory diseases (J40-J47)	64	1.6	0.4
8	Septicemia (A40-A41)	53	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	45	1.1	0.3
11	Cerebrovascular diseases (I60-I69)	38	0.9	0.2
12	Meningitis (G00,G03)	23	0.6	0.1
13	Anemias (D50-D64)	20	0.5	0.1
14	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	14	0.3	*
...	All other causes (Residual)	1,021	25.1	6.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-9 years				
...	All causes	2,427	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	746	30.7	3.6
2	Malignant neoplasms (C00-C97)	447	18.4	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	179	7.4	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	5.2	0.6
5	Chronic lower respiratory diseases (J40-J47)	75	3.1	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	3.0	0.4
7	Influenza and pneumonia (J09-J18)	67	2.8	0.3
8	Cerebrovascular diseases (I60-I69)	41	1.7	0.2
9	Septicemia (A40-A41)	35	1.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	1.4	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	18	0.7	*
12	Anemias (D50-D64)	12	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.5	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (Residual)	547	22.5	2.7
All races, both sexes, 10-14 years				
...	All causes	2,913	100.0	14.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	775	26.6	3.8
2	Malignant neoplasms (C00-C97)	448	15.4	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	386	13.3	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	161	5.5	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	152	5.2	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	100	3.4	0.5
7	Chronic lower respiratory diseases (J40-J47)	80	2.7	0.4
8	Influenza and pneumonia (J09-J18)	61	2.1	0.3
9	Cerebrovascular diseases (I60-I69)	48	1.6	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.1	0.2
11	Septicemia (A40-A41)	20	0.7	0.1
11	Diabetes mellitus (E10-E14)	20	0.7	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	15	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.5	*
15	Meningitis (G00,G03)	7	0.2	*
15	Pneumonitis due to solids and liquids (J69)	7	0.2	*
...	All other causes (Residual)	587	20.2	2.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 15-19 years				
...	All causes	9,480	100.0	44.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,652	38.5	17.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,748	18.4	8.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,407	14.8	6.6
4	Malignant neoplasms (C00-C97)	627	6.6	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	297	3.1	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	166	1.8	0.8
7	Influenza and pneumonia (J09-J18)	73	0.8	0.3
8	Chronic lower respiratory diseases (J40-J47)	60	0.6	0.3
9	Cerebrovascular diseases (I60-I69)	56	0.6	0.3
10	Diabetes mellitus (E10-E14)	52	0.5	0.2
11	Septicemia (A40-A41)	42	0.4	0.2
12	Pregnancy, childbirth and the puerperium (O00-O99)	41	0.4	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.4	0.2
14	Anemias (D50-D64)	29	0.3	0.1
15	Legal intervention (Y35,Y89.0)	19	0.2	*
...	All other causes (Residual)	1,173	12.4	5.5
All races, both sexes, 20-24 years				
...	All causes	19,006	100.0	83.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,967	41.9	34.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,130	16.5	13.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,922	15.4	12.8
4	Malignant neoplasms (C00-C97)	869	4.6	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	644	3.4	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	196	1.0	0.9
7	Diabetes mellitus (E10-E14)	141	0.7	0.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	137	0.7	0.6
9	Influenza and pneumonia (J09-J18)	124	0.7	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	106	0.6	0.5
11	Cerebrovascular diseases (I60-I69)	97	0.5	0.4
12	Septicemia (A40-A41)	96	0.5	0.4
13	Chronic lower respiratory diseases (J40-J47)	95	0.5	0.4
14	Legal intervention (Y35,Y89.0)	80	0.4	0.4
15	Anemias (D50-D64)	64	0.3	0.3
...	All other causes (Residual)	2,338	12.3	10.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-29 years				
...	All causes	21,056	100.0	97.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,120	38.6	37.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,265	15.5	15.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,272	10.8	10.5
4	Malignant neoplasms (C00-C97)	1,361	6.5	6.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,155	5.5	5.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	271	1.3	1.3
7	Diabetes mellitus (E10-E14)	265	1.3	1.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	222	1.1	1.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	191	0.9	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	186	0.9	0.9
11	Cerebrovascular diseases (I60-I69)	179	0.9	0.8
12	Influenza and pneumonia (J09-J18)	176	0.8	0.8
13	Septicemia (A40-A41)	133	0.6	0.6
14	Chronic lower respiratory diseases (J40-J47)	130	0.6	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	89	0.4	0.4
...	All other causes (Residual)	3,041	14.4	14.1
All races, both sexes, 30-34 years				
...	All causes	24,407	100.0	114.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,089	33.1	38.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,083	12.6	14.5
3	Malignant neoplasms (C00-C97)	2,312	9.5	10.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,103	8.6	9.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,964	8.0	9.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	485	2.0	2.3
7	Diabetes mellitus (E10-E14)	419	1.7	2.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	360	1.5	1.7
9	Cerebrovascular diseases (I60-I69)	329	1.3	1.5
10	Influenza and pneumonia (J09-J18)	273	1.1	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	237	1.0	1.1
12	Pregnancy, childbirth and the puerperium (O00-O99)	219	0.9	1.0
13	Septicemia (A40-A41)	200	0.8	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	177	0.7	0.8
15	Chronic lower respiratory diseases (J40-J47)	161	0.7	0.8
...	All other causes (Residual)	3,996	16.4	18.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 35-39 years				
...	All causes	27,821	100.0	141.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,240	26.0	36.9
2	Malignant neoplasms (C00-C97)	3,773	13.6	19.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,419	12.3	17.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,029	10.9	15.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,423	5.1	7.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	850	3.1	4.3
7	Diabetes mellitus (E10-E14)	680	2.4	3.5
8	Cerebrovascular diseases (I60-I69)	550	2.0	2.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	494	1.8	2.5
10	Influenza and pneumonia (J09-J18)	341	1.2	1.7
11	Septicemia (A40-A41)	296	1.1	1.5
12	Chronic lower respiratory diseases (J40-J47)	283	1.0	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	230	0.8	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	205	0.7	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	172	0.6	0.9
...	All other causes (Residual)	4,836	17.4	24.7
All races, both sexes, 40-44 years				
...	All causes	41,752	100.0	200.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,114	19.4	38.9
2	Malignant neoplasms (C00-C97)	7,576	18.1	36.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,922	16.6	33.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,522	8.4	16.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,641	3.9	7.9
6	Diabetes mellitus (E10-E14)	1,272	3.0	6.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,158	2.8	5.6
8	Cerebrovascular diseases (I60-I69)	1,137	2.7	5.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	752	1.8	3.6
10	Influenza and pneumonia (J09-J18)	540	1.3	2.6
11	Septicemia (A40-A41)	524	1.3	2.5
12	Chronic lower respiratory diseases (J40-J47)	477	1.1	2.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	397	1.0	1.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	282	0.7	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	236	0.6	1.1
...	All other causes (Residual)	7,202	17.2	34.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-49 years				
...	All causes	66,833	100.0	315.1
1	Malignant neoplasms (C00-C97)	15,256	22.8	71.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,594	18.8	59.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,482	14.2	44.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,011	6.0	18.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,329	5.0	15.7
6	Diabetes mellitus (E10-E14)	2,195	3.3	10.3
7	Cerebrovascular diseases (I60-I69)	1,966	2.9	9.3
8	Chronic lower respiratory diseases (J40-J47)	1,249	1.9	5.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,090	1.6	5.1
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,053	1.6	5.0
11	Septicemia (A40-A41)	903	1.4	4.3
12	Influenza and pneumonia (J09-J18)	817	1.2	3.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	704	1.1	3.3
14	Viral hepatitis (B15-B19)	588	0.9	2.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	483	0.7	2.3
...	All other causes (Residual)	11,113	16.6	52.4
All races, both sexes, 50-54 years				
...	All causes	110,891	100.0	491.6
1	Malignant neoplasms (C00-C97)	30,929	27.9	137.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,573	20.4	100.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,875	9.8	48.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,456	4.9	24.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,610	4.2	20.4
6	Diabetes mellitus (E10-E14)	3,704	3.3	16.4
7	Cerebrovascular diseases (I60-I69)	3,459	3.1	15.3
8	Chronic lower respiratory diseases (J40-J47)	3,370	3.0	14.9
9	Septicemia (A40-A41)	1,542	1.4	6.8
10	Influenza and pneumonia (J09-J18)	1,416	1.3	6.3
11	Viral hepatitis (B15-B19)	1,323	1.2	5.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,321	1.2	5.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,288	1.2	5.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,030	0.9	4.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	936	0.8	4.1
...	All other causes (Residual)	17,059	15.4	75.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 55-59 years				
...	All causes	152,981	100.0	721.8
1	Malignant neoplasms (C00-C97)	49,105	32.1	231.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,210	21.1	152.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,813	6.4	46.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	6,327	4.1	29.9
5	Chronic lower respiratory diseases (J40-J47)	6,241	4.1	29.4
6	Diabetes mellitus (E10-E14)	5,669	3.7	26.7
7	Cerebrovascular diseases (I60-I69)	4,924	3.2	23.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,161	2.7	19.6
9	Septicemia (A40-A41)	2,258	1.5	10.7
10	Viral hepatitis (B15-B19)	2,117	1.4	10.0
11	Influenza and pneumonia (J09-J18)	2,107	1.4	9.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,052	1.3	9.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,444	0.9	6.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,104	0.7	5.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	722	0.5	3.4
...	All other causes (Residual)	22,727	14.9	107.2
All races, both sexes, 60-64 years				
...	All causes	185,146	100.0	1,021.7
1	Malignant neoplasms (C00-C97)	64,219	34.7	354.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,358	21.8	222.7
3	Chronic lower respiratory diseases (J40-J47)	9,701	5.2	53.5
4	Diabetes mellitus (E10-E14)	7,392	4.0	40.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,244	3.9	40.0
6	Cerebrovascular diseases (I60-I69)	6,440	3.5	35.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,624	3.0	31.0
8	Septicemia (A40-A41)	3,087	1.7	17.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,974	1.6	16.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,895	1.6	16.0
11	Influenza and pneumonia (J09-J18)	2,672	1.4	14.7
12	Viral hepatitis (B15-B19)	1,819	1.0	10.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,693	0.9	9.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	866	0.5	4.8
15	Aortic aneurysm and dissection (I71)	740	0.4	4.1
...	All other causes (Residual)	27,422	14.8	151.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-69 years				
...	All causes	214,509	100.0	1,468.4
1	Malignant neoplasms (C00-C97)	76,042	35.4	520.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,702	21.8	319.7
3	Chronic lower respiratory diseases (J40-J47)	15,101	7.0	103.4
4	Diabetes mellitus (E10-E14)	8,510	4.0	58.3
5	Cerebrovascular diseases (I60-I69)	8,163	3.8	55.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,769	2.7	39.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4,074	1.9	27.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,752	1.7	25.7
9	Septicemia (A40-A41)	3,658	1.7	25.0
10	Influenza and pneumonia (J09-J18)	3,319	1.5	22.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,146	1.0	14.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,025	0.9	13.9
13	Alzheimer's disease (G30)	1,289	0.6	8.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,227	0.6	8.4
15	Parkinson's disease (G20-G21)	999	0.5	6.8
...	All other causes (Residual)	31,733	14.8	217.2
All races, both sexes, 70-74 years				
...	All causes	239,920	100.0	2,261.7
1	Malignant neoplasms (C00-C97)	79,510	33.1	749.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,730	21.6	487.6
3	Chronic lower respiratory diseases (J40-J47)	20,502	8.5	193.3
4	Cerebrovascular diseases (I60-I69)	10,559	4.4	99.5
5	Diabetes mellitus (E10-E14)	8,769	3.7	82.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,198	2.2	49.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,772	2.0	45.0
8	Influenza and pneumonia (J09-J18)	4,122	1.7	38.9
9	Septicemia (A40-A41)	4,035	1.7	38.0
10	Alzheimer's disease (G30)	3,272	1.4	30.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	3,013	1.3	28.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,332	1.0	22.0
13	Parkinson's disease (G20-G21)	2,216	0.9	20.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,648	0.7	15.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,546	0.6	14.6
...	All other causes (Residual)	36,696	15.3	345.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 75-79 years				
...	All causes	277,314	100.0	3,611.9
1	Malignant neoplasms (C00-C97)	77,601	28.0	1,010.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62,251	22.4	810.8
3	Chronic lower respiratory diseases (J40-J47)	23,241	8.4	302.7
4	Cerebrovascular diseases (I60-I69)	14,828	5.3	193.1
5	Diabetes mellitus (E10-E14)	9,463	3.4	123.3
6	Alzheimer's disease (G30)	7,594	2.7	98.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,148	2.2	80.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,830	2.1	75.9
9	Influenza and pneumonia (J09-J18)	5,548	2.0	72.3
10	Septicemia (A40-A41)	4,774	1.7	62.2
11	Parkinson's disease (G20-G21)	4,219	1.5	55.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,933	1.1	38.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,287	0.8	29.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,961	0.7	25.5
15	Pneumonitis due to solids and liquids (J69)	1,927	0.7	25.1
...	All other causes (Residual)	46,709	16.8	608.4
All races, both sexes, 80-84 years				
...	All causes	347,699	100.0	6,027.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	85,004	24.4	1,473.6
2	Malignant neoplasms (C00-C97)	75,613	21.7	1,310.8
3	Chronic lower respiratory diseases (J40-J47)	26,105	7.5	452.5
4	Cerebrovascular diseases (I60-I69)	21,323	6.1	369.6
5	Alzheimer's disease (G30)	15,479	4.5	268.3
6	Diabetes mellitus (E10-E14)	10,124	2.9	175.5
7	Influenza and pneumonia (J09-J18)	8,401	2.4	145.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,290	2.4	143.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,487	2.2	129.8
10	Parkinson's disease (G20-G21)	6,330	1.8	109.7
11	Septicemia (A40-A41)	5,500	1.6	95.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,283	1.2	74.2
13	Pneumonitis due to solids and liquids (J69)	3,006	0.9	52.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,593	0.7	44.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,727	0.5	29.9
...	All other causes (Residual)	66,434	19.1	1,151.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85-89 years				
...	All causes	389,258	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	106,258	27.3	---
2	Malignant neoplasms (C00-C97)	60,154	15.5	---
3	Cerebrovascular diseases (I60-I69)	25,546	6.6	---
4	Chronic lower respiratory diseases (J40-J47)	23,832	6.1	---
5	Alzheimer's disease (G30)	23,774	6.1	---
6	Influenza and pneumonia (J09-J18)	11,180	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,747	2.5	---
8	Diabetes mellitus (E10-E14)	9,355	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,436	2.2	---
10	Parkinson's disease (G20-G21)	6,568	1.7	---
11	Septicemia (A40-A41)	5,700	1.5	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,548	1.4	---
13	Pneumonitis due to solids and liquids (J69)	4,142	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,684	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,685	0.4	---
...	All other causes (Residual)	84,649	21.7	---
All races, both sexes, 90-94 years				
...	All causes	294,187	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	89,917	30.6	---
2	Malignant neoplasms (C00-C97)	29,980	10.2	---
3	Alzheimer's disease (G30)	21,221	7.2	---
4	Cerebrovascular diseases (I60-I69)	20,056	6.8	---
5	Chronic lower respiratory diseases (J40-J47)	13,619	4.6	---
6	Influenza and pneumonia (J09-J18)	9,884	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,341	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,172	2.1	---
9	Diabetes mellitus (E10-E14)	5,592	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,216	1.8	---
11	Septicemia (A40-A41)	3,687	1.3	---
12	Pneumonitis due to solids and liquids (J69)	3,354	1.1	---
13	Parkinson's disease (G20-G21)	3,341	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,813	0.6	---
15	Atherosclerosis (I70)	1,402	0.5	---
...	All other causes (Residual)	71,592	24.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 95-99 years				
...	All causes	116,764	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,748	32.3	---
2	Alzheimer's disease (G30)	9,276	7.9	---
3	Malignant neoplasms (C00-C97)	7,715	6.6	---
4	Cerebrovascular diseases (I60-I69)	7,657	6.6	---
5	Influenza and pneumonia (J09-J18)	4,430	3.8	---
6	Chronic lower respiratory diseases (J40-J47)	4,159	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,911	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,486	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,221	1.9	---
10	Diabetes mellitus (E10-E14)	1,649	1.4	---
11	Pneumonitis due to solids and liquids (J69)	1,418	1.2	---
12	Septicemia (A40-A41)	1,219	1.0	---
13	Parkinson's disease (G20-G21)	762	0.7	---
14	Atherosclerosis (I70)	729	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	518	0.4	---
...	All other causes (Residual)	31,866	27.3	---
All races, both sexes, 100 years and over				
...	All causes	24,989	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,546	34.2	---
2	Alzheimer's disease (G30)	1,881	7.5	---
3	Cerebrovascular diseases (I60-I69)	1,470	5.9	---
4	Influenza and pneumonia (J09-J18)	1,147	4.6	---
5	Malignant neoplasms (C00-C97)	943	3.8	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	741	3.0	---
7	Chronic lower respiratory diseases (J40-J47)	635	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	538	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	410	1.6	---
10	Pneumonitis due to solids and liquids (J69)	304	1.2	---
11	Diabetes mellitus (E10-E14)	289	1.2	---
12	Septicemia (A40-A41)	242	1.0	---
13	Atherosclerosis (I70)	233	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	91	0.4	---
14	Parkinson's disease (G20-G21)	91	0.4	---
...	All other causes (Residual)	7,428	29.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, all ages				
...	All causes	1,306,034	100.0	839.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	321,347	24.6	206.5
2	Malignant neoplasms (C00-C97)	307,559	23.5	197.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81,916	6.3	52.6
4	Chronic lower respiratory diseases (J40-J47)	70,317	5.4	45.2
5	Cerebrovascular diseases (I60-I69)	53,691	4.1	34.5
6	Diabetes mellitus (E10-E14)	39,841	3.1	25.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32,055	2.5	20.6
8	Influenza and pneumonia (J09-J18)	26,804	2.1	17.2
9	Alzheimer's disease (G30)	25,836	2.0	16.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	23,709	1.8	15.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,493	1.8	15.1
12	Septicemia (A40-A41)	17,994	1.4	11.6
13	Parkinson's disease (G20-G21)	15,088	1.2	9.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,963	1.0	8.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12,726	1.0	8.2
...	All other causes (Residual)	240,695	18.4	154.6
All races, male, 1-4 years				
...	All causes	2,323	100.0	28.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	831	35.8	10.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	232	10.0	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	184	7.9	2.3
4	Malignant neoplasms (C00-C97)	183	7.9	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	4.1	1.2
6	Influenza and pneumonia (J09-J18)	53	2.3	0.7
7	Chronic lower respiratory diseases (J40-J47)	38	1.6	0.5
8	Septicemia (A40-A41)	32	1.4	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	29	1.2	0.4
10	Cerebrovascular diseases (I60-I69)	22	0.9	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.8	*
12	Meningitis (G00,G03)	13	0.6	*
13	Anemias (D50-D64)	11	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	7	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (Residual)	566	24.4	7.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 5-9 years				
...	All causes	1,383	100.0	13.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	450	32.5	4.3
2	Malignant neoplasms (C00-C97)	241	17.4	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	7.0	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	5.3	0.7
5	Chronic lower respiratory diseases (J40-J47)	50	3.6	0.5
6	Influenza and pneumonia (J09-J18)	37	2.7	0.4
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	2.6	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.7	0.2
9	Septicemia (A40-A41)	21	1.5	0.2
10	Cerebrovascular diseases (I60-I69)	19	1.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	11	0.8	*
12	Anemias (D50-D64)	8	0.6	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.5	*
14	Meningitis (G00,G03)	6	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	299	21.6	2.8
All races, male, 10-14 years				
...	All causes	1,694	100.0	16.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	500	29.5	4.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	245	14.5	2.3
3	Malignant neoplasms (C00-C97)	230	13.6	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	5.6	0.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	4.7	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	3.4	0.5
7	Chronic lower respiratory diseases (J40-J47)	50	3.0	0.5
8	Influenza and pneumonia (J09-J18)	33	1.9	0.3
9	Cerebrovascular diseases (I60-I69)	25	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.9	*
11	Septicemia (A40-A41)	11	0.6	*
11	Diabetes mellitus (E10-E14)	11	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	6	0.4	*
15	Meningitis (G00,G03)	5	0.3	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
...	All other causes (Residual)	315	18.6	3.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-19 years				
...	All causes	6,759	100.0	62.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,554	37.8	23.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,350	20.0	12.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,238	18.3	11.4
4	Malignant neoplasms (C00-C97)	367	5.4	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	192	2.8	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	96	1.4	0.9
7	Influenza and pneumonia (J09-J18)	41	0.6	0.4
8	Chronic lower respiratory diseases (J40-J47)	38	0.6	0.4
9	Diabetes mellitus (E10-E14)	37	0.5	0.3
10	Cerebrovascular diseases (I60-I69)	29	0.4	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.4	0.2
12	Septicemia (A40-A41)	25	0.4	0.2
13	Legal intervention (Y35,Y89.0)	18	0.3	*
14	Anemias (D50-D64)	15	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	12	0.2	*
...	All other causes (Residual)	721	10.7	6.6
All races, male, 20-24 years				
...	All causes	14,105	100.0	120.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,991	42.5	51.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,553	18.1	21.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,527	17.9	21.6
4	Malignant neoplasms (C00-C97)	492	3.5	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	436	3.1	3.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	0.9	1.1
7	Diabetes mellitus (E10-E14)	83	0.6	0.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	78	0.6	0.7
9	Legal intervention (Y35,Y89.0)	77	0.5	0.7
10	Influenza and pneumonia (J09-J18)	73	0.5	0.6
11	Chronic lower respiratory diseases (J40-J47)	63	0.4	0.5
12	Cerebrovascular diseases (I60-I69)	56	0.4	0.5
13	Septicemia (A40-A41)	49	0.3	0.4
14	Anemias (D50-D64)	39	0.3	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.2	0.3
...	All other causes (Residual)	1,430	10.1	12.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 25-29 years				
...	All causes	15,023	100.0	137.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,061	40.3	55.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,664	17.7	24.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,945	12.9	17.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	775	5.2	7.1
5	Malignant neoplasms (C00-C97)	756	5.0	6.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	205	1.4	1.9
7	Diabetes mellitus (E10-E14)	151	1.0	1.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	122	0.8	1.1
9	Cerebrovascular diseases (I60-I69)	101	0.7	0.9
10	Influenza and pneumonia (J09-J18)	97	0.6	0.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	94	0.6	0.9
12	Chronic lower respiratory diseases (J40-J47)	78	0.5	0.7
13	Septicemia (A40-A41)	71	0.5	0.6
14	Legal intervention (Y35,Y89.0)	66	0.4	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	0.3	0.4
...	All other causes (Residual)	1,790	11.9	16.3
All races, male, 30-34 years				
...	All causes	16,439	100.0	153.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,964	36.3	55.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,399	14.6	22.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,610	9.8	15.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,439	8.8	13.5
5	Malignant neoplasms (C00-C97)	1,101	6.7	10.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	313	1.9	2.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	252	1.5	2.4
7	Diabetes mellitus (E10-E14)	252	1.5	2.4
9	Cerebrovascular diseases (I60-I69)	178	1.1	1.7
10	Influenza and pneumonia (J09-J18)	156	0.9	1.5
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	0.9	1.4
12	Septicemia (A40-A41)	107	0.7	1.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	0.6	0.9
14	Chronic lower respiratory diseases (J40-J47)	82	0.5	0.8
15	Legal intervention (Y35,Y89.0)	75	0.5	0.7
...	All other causes (Residual)	2,272	13.8	21.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-39 years				
...	All causes	17,544	100.0	179.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,060	28.8	51.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,351	13.4	24.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,344	13.4	24.0
4	Malignant neoplasms (C00-C97)	1,631	9.3	16.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,143	6.5	11.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	537	3.1	5.5
7	Diabetes mellitus (E10-E14)	403	2.3	4.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	319	1.8	3.3
9	Cerebrovascular diseases (I60-I69)	306	1.7	3.1
10	Influenza and pneumonia (J09-J18)	177	1.0	1.8
11	Chronic lower respiratory diseases (J40-J47)	143	0.8	1.5
12	Septicemia (A40-A41)	138	0.8	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	125	0.7	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	121	0.7	1.2
15	Aortic aneurysm and dissection (I71)	92	0.5	0.9
...	All other causes (Residual)	2,654	15.1	27.1
All races, male, 40-44 years				
...	All causes	25,528	100.0	246.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,506	21.6	53.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,808	18.8	46.4
3	Malignant neoplasms (C00-C97)	3,196	12.5	30.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,652	10.4	25.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,083	4.2	10.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	882	3.5	8.5
7	Diabetes mellitus (E10-E14)	808	3.2	7.8
8	Cerebrovascular diseases (I60-I69)	632	2.5	6.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	523	2.0	5.0
10	Influenza and pneumonia (J09-J18)	287	1.1	2.8
11	Septicemia (A40-A41)	259	1.0	2.5
12	Chronic lower respiratory diseases (J40-J47)	222	0.9	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	217	0.9	2.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	182	0.7	1.8
15	Aortic aneurysm and dissection (I71)	139	0.5	1.3
...	All other causes (Residual)	4,132	16.2	39.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-49 years				
...	All causes	40,531	100.0	386.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,849	21.8	84.3
2	Malignant neoplasms (C00-C97)	7,166	17.7	68.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,353	15.7	60.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,945	7.3	28.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,202	5.4	21.0
6	Diabetes mellitus (E10-E14)	1,351	3.3	12.9
7	Cerebrovascular diseases (I60-I69)	1,129	2.8	10.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	801	2.0	7.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	773	1.9	7.4
10	Chronic lower respiratory diseases (J40-J47)	552	1.4	5.3
11	Septicemia (A40-A41)	486	1.2	4.6
12	Influenza and pneumonia (J09-J18)	461	1.1	4.4
13	Viral hepatitis (B15-B19)	416	1.0	4.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	393	1.0	3.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	339	0.8	3.2
...	All other causes (Residual)	6,315	15.6	60.2
All races, male, 50-54 years				
...	All causes	67,466	100.0	609.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,981	23.7	144.4
2	Malignant neoplasms (C00-C97)	15,800	23.4	142.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,238	10.7	65.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,656	5.4	33.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,448	5.1	31.1
6	Diabetes mellitus (E10-E14)	2,282	3.4	20.6
7	Cerebrovascular diseases (I60-I69)	1,938	2.9	17.5
8	Chronic lower respiratory diseases (J40-J47)	1,590	2.4	14.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	959	1.4	8.7
10	Viral hepatitis (B15-B19)	912	1.4	8.2
11	Septicemia (A40-A41)	818	1.2	7.4
12	Influenza and pneumonia (J09-J18)	816	1.2	7.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	797	1.2	7.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	706	1.0	6.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	651	1.0	5.9
...	All other causes (Residual)	9,874	14.6	89.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-59 years				
...	All causes	93,996	100.0	914.1
1	Malignant neoplasms (C00-C97)	26,932	28.7	261.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,752	24.2	221.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,701	7.1	65.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,471	4.8	43.5
5	Diabetes mellitus (E10-E14)	3,507	3.7	34.1
6	Chronic lower respiratory diseases (J40-J47)	3,168	3.4	30.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,095	3.3	30.1
8	Cerebrovascular diseases (I60-I69)	2,869	3.1	27.9
9	Viral hepatitis (B15-B19)	1,475	1.6	14.3
10	Influenza and pneumonia (J09-J18)	1,236	1.3	12.0
11	Septicemia (A40-A41)	1,217	1.3	11.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,136	1.2	11.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	939	1.0	9.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	820	0.9	8.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	519	0.6	5.0
...	All other causes (Residual)	13,159	14.0	128.0
All races, male, 60-64 years				
...	All causes	112,329	100.0	1,295.0
1	Malignant neoplasms (C00-C97)	35,878	31.9	413.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,928	24.9	322.0
3	Chronic lower respiratory diseases (J40-J47)	5,119	4.6	59.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,957	4.4	57.1
5	Diabetes mellitus (E10-E14)	4,426	3.9	51.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,014	3.6	46.3
7	Cerebrovascular diseases (I60-I69)	3,782	3.4	43.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,278	2.0	26.3
9	Septicemia (A40-A41)	1,675	1.5	19.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,664	1.5	19.2
11	Influenza and pneumonia (J09-J18)	1,509	1.3	17.4
12	Viral hepatitis (B15-B19)	1,317	1.2	15.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,051	0.9	12.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	549	0.5	6.3
15	Aortic aneurysm and dissection (I71)	539	0.5	6.2
...	All other causes (Residual)	15,643	13.9	180.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 65-69 years				
...	All causes	124,324	100.0	1,798.4
1	Malignant neoplasms (C00-C97)	42,172	33.9	610.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,697	24.7	444.0
3	Chronic lower respiratory diseases (J40-J47)	7,590	6.1	109.8
4	Diabetes mellitus (E10-E14)	4,977	4.0	72.0
5	Cerebrovascular diseases (I60-I69)	4,458	3.6	64.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,824	3.1	55.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,682	2.2	38.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,007	1.6	29.0
9	Septicemia (A40-A41)	1,912	1.5	27.7
10	Influenza and pneumonia (J09-J18)	1,859	1.5	26.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,684	1.4	24.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,163	0.9	16.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	739	0.6	10.7
14	Parkinson's disease (G20-G21)	663	0.5	9.6
15	Aortic aneurysm and dissection (I71)	592	0.5	8.6
...	All other causes (Residual)	17,305	13.9	250.3
All races, male, 70-74 years				
...	All causes	133,574	100.0	2,734.7
1	Malignant neoplasms (C00-C97)	43,497	32.6	890.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,937	23.9	653.9
3	Chronic lower respiratory diseases (J40-J47)	10,335	7.7	211.6
4	Cerebrovascular diseases (I60-I69)	5,575	4.2	114.1
5	Diabetes mellitus (E10-E14)	4,880	3.7	99.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,190	2.4	65.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,462	1.8	50.4
8	Influenza and pneumonia (J09-J18)	2,272	1.7	46.5
9	Septicemia (A40-A41)	2,090	1.6	42.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,849	1.4	37.9
11	Parkinson's disease (G20-G21)	1,497	1.1	30.6
12	Alzheimer's disease (G30)	1,401	1.0	28.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,388	1.0	28.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,196	0.9	24.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	918	0.7	18.8
...	All other causes (Residual)	19,087	14.3	390.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-79 years				
...	All causes	146,166	100.0	4,311.2
1	Malignant neoplasms (C00-C97)	41,815	28.6	1,233.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,806	24.5	1,056.1
3	Chronic lower respiratory diseases (J40-J47)	11,442	7.8	337.5
4	Cerebrovascular diseases (I60-I69)	7,037	4.8	207.6
5	Diabetes mellitus (E10-E14)	5,011	3.4	147.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,494	2.4	103.1
7	Alzheimer's disease (G30)	3,128	2.1	92.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,024	2.1	89.2
9	Influenza and pneumonia (J09-J18)	2,911	2.0	85.9
10	Parkinson's disease (G20-G21)	2,782	1.9	82.1
11	Septicemia (A40-A41)	2,257	1.5	66.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,368	0.9	40.3
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,264	0.9	37.3
14	Pneumonitis due to solids and liquids (J69)	1,159	0.8	34.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,110	0.8	32.7
...	All other causes (Residual)	22,558	15.4	665.4
All races, male, 80-84 years				
...	All causes	169,174	100.0	7,137.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,828	25.9	1,849.2
2	Malignant neoplasms (C00-C97)	39,665	23.4	1,673.5
3	Chronic lower respiratory diseases (J40-J47)	12,367	7.3	521.8
4	Cerebrovascular diseases (I60-I69)	8,969	5.3	378.4
5	Alzheimer's disease (G30)	5,661	3.3	238.8
6	Diabetes mellitus (E10-E14)	4,941	2.9	208.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,335	2.6	182.9
8	Influenza and pneumonia (J09-J18)	4,186	2.5	176.6
9	Parkinson's disease (G20-G21)	3,935	2.3	166.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,837	2.3	161.9
11	Septicemia (A40-A41)	2,573	1.5	108.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,784	1.1	75.3
13	Pneumonitis due to solids and liquids (J69)	1,697	1.0	71.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,425	0.8	60.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	945	0.6	39.9
...	All other causes (Residual)	29,026	17.2	1,224.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 85-89 years				
...	All causes	165,286	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,468	28.7	---
2	Malignant neoplasms (C00-C97)	29,719	18.0	---
3	Chronic lower respiratory diseases (J40-J47)	10,470	6.3	---
4	Cerebrovascular diseases (I60-I69)	8,866	5.4	---
5	Alzheimer's disease (G30)	7,419	4.5	---
6	Influenza and pneumonia (J09-J18)	5,015	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,480	2.7	---
8	Diabetes mellitus (E10-E14)	4,044	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,979	2.4	---
10	Parkinson's disease (G20-G21)	3,725	2.3	---
11	Septicemia (A40-A41)	2,434	1.5	---
12	Pneumonitis due to solids and liquids (J69)	2,264	1.4	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,979	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,334	0.8	---
15	Aortic aneurysm and dissection (I71)	727	0.4	---
...	All other causes (Residual)	31,363	19.0	---
All races, male, 90-94 years				
...	All causes	104,050	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,525	32.2	---
2	Malignant neoplasms (C00-C97)	13,529	13.0	---
3	Cerebrovascular diseases (I60-I69)	5,750	5.5	---
4	Chronic lower respiratory diseases (J40-J47)	5,412	5.2	---
5	Alzheimer's disease (G30)	5,380	5.2	---
6	Influenza and pneumonia (J09-J18)	3,811	3.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,791	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,714	2.6	---
9	Diabetes mellitus (E10-E14)	2,105	2.0	---
10	Parkinson's disease (G20-G21)	1,690	1.6	---
11	Pneumonitis due to solids and liquids (J69)	1,648	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,528	1.5	---
13	Septicemia (A40-A41)	1,358	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	772	0.7	---
15	Atherosclerosis (I70)	418	0.4	---
...	All other causes (Residual)	21,619	20.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 95-99 years				
...	All causes	30,779	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,624	34.5	---
2	Malignant neoplasms (C00-C97)	2,879	9.4	---
3	Cerebrovascular diseases (I60-I69)	1,660	5.4	---
4	Alzheimer's disease (G30)	1,624	5.3	---
5	Influenza and pneumonia (J09-J18)	1,400	4.5	---
6	Chronic lower respiratory diseases (J40-J47)	1,348	4.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	853	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	801	2.6	---
9	Pneumonitis due to solids and liquids (J69)	593	1.9	---
10	Diabetes mellitus (E10-E14)	507	1.6	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	498	1.6	---
12	Septicemia (A40-A41)	327	1.1	---
13	Parkinson's disease (G20-G21)	318	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	188	0.6	---
15	Atherosclerosis (I70)	174	0.6	---
...	All other causes (Residual)	6,985	22.7	---
All races, male, 100 years and over				
...	All causes	4,347	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,575	36.2	---
2	Malignant neoplasms (C00-C97)	274	6.3	---
3	Influenza and pneumonia (J09-J18)	263	6.1	---
4	Cerebrovascular diseases (I60-I69)	229	5.3	---
5	Alzheimer's disease (G30)	224	5.2	---
6	Chronic lower respiratory diseases (J40-J47)	145	3.3	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	109	2.5	---
8	Pneumonitis due to solids and liquids (J69)	103	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	2.3	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	2.3	---
11	Diabetes mellitus (E10-E14)	55	1.3	---
12	Septicemia (A40-A41)	46	1.1	---
13	Atherosclerosis (I70)	41	0.9	---
14	Parkinson's disease (G20-G21)	30	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.4	---
...	All other causes (Residual)	1,031	23.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,290,959	100.0	804.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	289,758	22.4	180.6
2	Malignant neoplasms (C00-C97)	277,322	21.5	172.8
3	Chronic lower respiratory diseases (J40-J47)	78,888	6.1	49.2
4	Cerebrovascular diseases (I60-I69)	75,287	5.8	46.9
5	Alzheimer's disease (G30)	58,931	4.6	36.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48,641	3.8	30.3
7	Diabetes mellitus (E10-E14)	35,737	2.8	22.3
8	Influenza and pneumonia (J09-J18)	30,175	2.3	18.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23,619	1.8	14.7
10	Septicemia (A40-A41)	20,162	1.6	12.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,807	1.4	11.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12,718	1.0	7.9
13	Parkinson's disease (G20-G21)	10,108	0.8	6.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,094	0.7	5.7
15	Pneumonitis due to solids and liquids (J69)	8,439	0.7	5.3
...	All other causes (Residual)	294,273	22.8	183.4
All races, female, 1-4 years				
...	All causes	1,745	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	485	27.8	6.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	14.0	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	8.8	2.0
4	Malignant neoplasms (C00-C97)	145	8.3	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	4.2	0.9
6	Influenza and pneumonia (J09-J18)	49	2.8	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.6	0.4
8	Chronic lower respiratory diseases (J40-J47)	26	1.5	0.3
9	Septicemia (A40-A41)	21	1.2	0.3
10	Cerebrovascular diseases (I60-I69)	16	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	16	0.9	*
12	Meningitis (G00,G03)	10	0.6	*
13	Anemias (D50-D64)	9	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
...	All other causes (Residual)	455	26.1	5.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-9 years				
...	All causes	1,044	100.0	10.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	296	28.4	2.9
2	Malignant neoplasms (C00-C97)	206	19.7	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	7.9	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	5.0	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.5	0.4
6	Influenza and pneumonia (J09-J18)	30	2.9	0.3
7	Chronic lower respiratory diseases (J40-J47)	25	2.4	0.2
8	Cerebrovascular diseases (I60-I69)	22	2.1	0.2
9	Septicemia (A40-A41)	14	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
14	Anemias (D50-D64)	4	0.4	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
...	All other causes (Residual)	241	23.1	2.4
All races, female, 10-14 years				
...	All causes	1,219	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	275	22.6	2.7
2	Malignant neoplasms (C00-C97)	218	17.9	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	141	11.6	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	6.7	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	4.7	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	3.5	0.4
7	Chronic lower respiratory diseases (J40-J47)	30	2.5	0.3
8	Influenza and pneumonia (J09-J18)	28	2.3	0.3
9	Cerebrovascular diseases (I60-I69)	23	1.9	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.2	*
11	Septicemia (A40-A41)	9	0.7	*
11	Diabetes mellitus (E10-E14)	9	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	9	0.7	*
14	Anemias (D50-D64)	4	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	272	22.3	2.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 15-19 years				
...	All causes	2,721	100.0	26.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,098	40.4	10.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	398	14.6	3.9
3	Malignant neoplasms (C00-C97)	260	9.6	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	169	6.2	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	3.9	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	2.6	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	41	1.5	0.4
8	Influenza and pneumonia (J09-J18)	32	1.2	0.3
9	Cerebrovascular diseases (I60-I69)	27	1.0	0.3
10	Chronic lower respiratory diseases (J40-J47)	22	0.8	0.2
11	Septicemia (A40-A41)	17	0.6	*
12	Diabetes mellitus (E10-E14)	15	0.6	*
13	Anemias (D50-D64)	14	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.3	*
...	All other causes (Residual)	432	15.9	4.2
All races, female, 20-24 years				
...	All causes	4,901	100.0	44.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,976	40.3	17.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	577	11.8	5.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	395	8.1	3.6
4	Malignant neoplasms (C00-C97)	377	7.7	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	208	4.2	1.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	137	2.8	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	1.4	0.6
8	Diabetes mellitus (E10-E14)	58	1.2	0.5
9	Influenza and pneumonia (J09-J18)	51	1.0	0.5
10	Septicemia (A40-A41)	47	1.0	0.4
11	Cerebrovascular diseases (I60-I69)	41	0.8	0.4
12	Chronic lower respiratory diseases (J40-J47)	32	0.7	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.6	0.3
14	Anemias (D50-D64)	25	0.5	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.3	*
...	All other causes (Residual)	863	17.6	7.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-29 years				
...	All causes	6,033	100.0	56.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,059	34.1	19.4
2	Malignant neoplasms (C00-C97)	605	10.0	5.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	601	10.0	5.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	380	6.3	3.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	327	5.4	3.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	222	3.7	2.1
7	Diabetes mellitus (E10-E14)	114	1.9	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	1.5	0.9
9	Influenza and pneumonia (J09-J18)	79	1.3	0.7
10	Cerebrovascular diseases (I60-I69)	78	1.3	0.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	69	1.1	0.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	66	1.1	0.6
13	Septicemia (A40-A41)	62	1.0	0.6
14	Chronic lower respiratory diseases (J40-J47)	52	0.9	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.7	0.4
...	All other causes (Residual)	1,185	19.6	11.2
All races, female, 30-34 years				
...	All causes	7,968	100.0	75.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,125	26.7	20.1
2	Malignant neoplasms (C00-C97)	1,211	15.2	11.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	684	8.6	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	664	8.3	6.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	354	4.4	3.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	219	2.7	2.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	172	2.2	1.6
8	Diabetes mellitus (E10-E14)	167	2.1	1.6
9	Cerebrovascular diseases (I60-I69)	151	1.9	1.4
10	Influenza and pneumonia (J09-J18)	117	1.5	1.1
11	Human immunodeficiency virus (HIV) disease (B20-B24)	108	1.4	1.0
12	Septicemia (A40-A41)	93	1.2	0.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.1	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	1.1	0.8
15	Chronic lower respiratory diseases (J40-J47)	79	1.0	0.7
...	All other causes (Residual)	1,649	20.7	15.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 35-39 years				
...	All causes	10,277	100.0	104.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,180	21.2	22.2
2	Malignant neoplasms (C00-C97)	2,142	20.8	21.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,075	10.5	10.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	678	6.6	6.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	313	3.0	3.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	280	2.7	2.9
7	Diabetes mellitus (E10-E14)	277	2.7	2.8
8	Cerebrovascular diseases (I60-I69)	244	2.4	2.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	175	1.7	1.8
10	Pregnancy, childbirth and the puerperium (O00-O99)	172	1.7	1.8
11	Influenza and pneumonia (J09-J18)	164	1.6	1.7
12	Septicemia (A40-A41)	158	1.5	1.6
13	Chronic lower respiratory diseases (J40-J47)	140	1.4	1.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	1.0	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	0.8	0.9
...	All other causes (Residual)	2,090	20.3	21.3
All races, female, 40-44 years				
...	All causes	16,224	100.0	154.7
1	Malignant neoplasms (C00-C97)	4,380	27.0	41.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,608	16.1	24.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,114	13.0	20.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	870	5.4	8.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	558	3.4	5.3
6	Cerebrovascular diseases (I60-I69)	505	3.1	4.8
7	Diabetes mellitus (E10-E14)	464	2.9	4.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	276	1.7	2.6
9	Septicemia (A40-A41)	265	1.6	2.5
10	Chronic lower respiratory diseases (J40-J47)	255	1.6	2.4
11	Influenza and pneumonia (J09-J18)	253	1.6	2.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	229	1.4	2.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	180	1.1	1.7
14	Pregnancy, childbirth and the puerperium (O00-O99)	129	0.8	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	0.7	1.0
...	All other causes (Residual)	3,032	18.7	28.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-49 years				
...	All causes	26,302	100.0	245.6
1	Malignant neoplasms (C00-C97)	8,090	30.8	75.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,745	14.2	35.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,129	11.9	29.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,127	4.3	10.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,066	4.1	10.0
6	Diabetes mellitus (E10-E14)	844	3.2	7.9
7	Cerebrovascular diseases (I60-I69)	837	3.2	7.8
8	Chronic lower respiratory diseases (J40-J47)	697	2.6	6.5
9	Septicemia (A40-A41)	417	1.6	3.9
10	Influenza and pneumonia (J09-J18)	356	1.4	3.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	311	1.2	2.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	289	1.1	2.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	280	1.1	2.6
14	Viral hepatitis (B15-B19)	172	0.7	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	149	0.6	1.4
...	All other causes (Residual)	4,793	18.2	44.8
All races, female, 50-54 years				
...	All causes	43,425	100.0	378.0
1	Malignant neoplasms (C00-C97)	15,129	34.8	131.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,592	15.2	57.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,637	8.4	31.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,800	4.1	15.7
5	Chronic lower respiratory diseases (J40-J47)	1,780	4.1	15.5
6	Cerebrovascular diseases (I60-I69)	1,521	3.5	13.2
7	Diabetes mellitus (E10-E14)	1,422	3.3	12.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,162	2.7	10.1
9	Septicemia (A40-A41)	724	1.7	6.3
10	Influenza and pneumonia (J09-J18)	600	1.4	5.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	524	1.2	4.6
12	Viral hepatitis (B15-B19)	411	0.9	3.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	379	0.9	3.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	329	0.8	2.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	230	0.5	2.0
...	All other causes (Residual)	7,185	16.5	62.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 55-59 years				
...	All causes	58,985	100.0	540.5
1	Malignant neoplasms (C00-C97)	22,173	37.6	203.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,458	16.0	86.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,112	5.3	28.5
4	Chronic lower respiratory diseases (J40-J47)	3,073	5.2	28.2
5	Diabetes mellitus (E10-E14)	2,162	3.7	19.8
6	Cerebrovascular diseases (I60-I69)	2,055	3.5	18.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,856	3.1	17.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,066	1.8	9.8
9	Septicemia (A40-A41)	1,041	1.8	9.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	916	1.6	8.4
11	Influenza and pneumonia (J09-J18)	871	1.5	8.0
12	Viral hepatitis (B15-B19)	642	1.1	5.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	505	0.9	4.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	284	0.5	2.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	280	0.5	2.6
...	All other causes (Residual)	9,491	16.1	87.0
All races, female, 60-64 years				
...	All causes	72,817	100.0	770.7
1	Malignant neoplasms (C00-C97)	28,341	38.9	300.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,430	17.1	131.6
3	Chronic lower respiratory diseases (J40-J47)	4,582	6.3	48.5
4	Diabetes mellitus (E10-E14)	2,966	4.1	31.4
5	Cerebrovascular diseases (I60-I69)	2,658	3.7	28.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,287	3.1	24.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,610	2.2	17.0
8	Septicemia (A40-A41)	1,412	1.9	14.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,231	1.7	13.0
10	Influenza and pneumonia (J09-J18)	1,163	1.6	12.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	696	1.0	7.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	642	0.9	6.8
13	Viral hepatitis (B15-B19)	502	0.7	5.3
14	Alzheimer's disease (G30)	383	0.5	4.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	361	0.5	3.8
...	All other causes (Residual)	11,553	15.9	122.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-69 years				
...	All causes	90,185	100.0	1,171.9
1	Malignant neoplasms (C00-C97)	33,870	37.6	440.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,005	17.7	208.0
3	Chronic lower respiratory diseases (J40-J47)	7,511	8.3	97.6
4	Cerebrovascular diseases (I60-I69)	3,705	4.1	48.1
5	Diabetes mellitus (E10-E14)	3,533	3.9	45.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,945	2.2	25.3
7	Septicemia (A40-A41)	1,746	1.9	22.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,745	1.9	22.7
9	Influenza and pneumonia (J09-J18)	1,460	1.6	19.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,392	1.5	18.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	862	1.0	11.2
12	Alzheimer's disease (G30)	706	0.8	9.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	488	0.5	6.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	462	0.5	6.0
15	Pneumonitis due to solids and liquids (J69)	367	0.4	4.8
...	All other causes (Residual)	14,388	16.0	187.0
All races, female, 70-74 years				
...	All causes	106,346	100.0	1,858.0
1	Malignant neoplasms (C00-C97)	36,013	33.9	629.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,793	18.6	345.8
3	Chronic lower respiratory diseases (J40-J47)	10,167	9.6	177.6
4	Cerebrovascular diseases (I60-I69)	4,984	4.7	87.1
5	Diabetes mellitus (E10-E14)	3,889	3.7	67.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,310	2.2	40.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,008	1.9	35.1
8	Septicemia (A40-A41)	1,945	1.8	34.0
9	Alzheimer's disease (G30)	1,871	1.8	32.7
10	Influenza and pneumonia (J09-J18)	1,850	1.7	32.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,164	1.1	20.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,136	1.1	19.8
13	Parkinson's disease (G20-G21)	719	0.7	12.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	628	0.6	11.0
15	Pneumonitis due to solids and liquids (J69)	516	0.5	9.0
...	All other causes (Residual)	17,353	16.3	303.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 75-79 years				
...	All causes	131,148	100.0	3,058.8
1	Malignant neoplasms (C00-C97)	35,786	27.3	834.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,445	20.2	616.8
3	Chronic lower respiratory diseases (J40-J47)	11,799	9.0	275.2
4	Cerebrovascular diseases (I60-I69)	7,791	5.9	181.7
5	Alzheimer's disease (G30)	4,466	3.4	104.2
6	Diabetes mellitus (E10-E14)	4,452	3.4	103.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,806	2.1	65.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,654	2.0	61.9
9	Influenza and pneumonia (J09-J18)	2,637	2.0	61.5
10	Septicemia (A40-A41)	2,517	1.9	58.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,565	1.2	36.5
12	Parkinson's disease (G20-G21)	1,437	1.1	33.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,023	0.8	23.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	851	0.6	19.8
15	Pneumonitis due to solids and liquids (J69)	768	0.6	17.9
...	All other causes (Residual)	24,151	18.4	563.3
All races, female, 80-84 years				
...	All causes	178,525	100.0	5,253.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,176	23.1	1,211.6
2	Malignant neoplasms (C00-C97)	35,948	20.1	1,057.8
3	Chronic lower respiratory diseases (J40-J47)	13,738	7.7	404.2
4	Cerebrovascular diseases (I60-I69)	12,354	6.9	363.5
5	Alzheimer's disease (G30)	9,818	5.5	288.9
6	Diabetes mellitus (E10-E14)	5,183	2.9	152.5
7	Influenza and pneumonia (J09-J18)	4,215	2.4	124.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,955	2.2	116.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,650	2.0	107.4
10	Septicemia (A40-A41)	2,927	1.6	86.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,499	1.4	73.5
12	Parkinson's disease (G20-G21)	2,395	1.3	70.5
13	Pneumonitis due to solids and liquids (J69)	1,309	0.7	38.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,168	0.7	34.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	859	0.5	25.3
...	All other causes (Residual)	37,331	20.9	1,098.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85-89 years				
...	All causes	223,972	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	58,790	26.2	---
2	Malignant neoplasms (C00-C97)	30,435	13.6	---
3	Cerebrovascular diseases (I60-I69)	16,680	7.4	---
4	Alzheimer's disease (G30)	16,355	7.3	---
5	Chronic lower respiratory diseases (J40-J47)	13,362	6.0	---
6	Influenza and pneumonia (J09-J18)	6,165	2.8	---
7	Diabetes mellitus (E10-E14)	5,311	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,267	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,457	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,569	1.6	---
11	Septicemia (A40-A41)	3,266	1.5	---
12	Parkinson's disease (G20-G21)	2,843	1.3	---
13	Pneumonitis due to solids and liquids (J69)	1,878	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,350	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,090	0.5	---
...	All other causes (Residual)	53,154	23.7	---
All races, female, 90-94 years				
...	All causes	190,137	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,392	29.7	---
2	Malignant neoplasms (C00-C97)	16,451	8.7	---
3	Alzheimer's disease (G30)	15,841	8.3	---
4	Cerebrovascular diseases (I60-I69)	14,306	7.5	---
5	Chronic lower respiratory diseases (J40-J47)	8,207	4.3	---
6	Influenza and pneumonia (J09-J18)	6,073	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,550	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,688	1.9	---
9	Diabetes mellitus (E10-E14)	3,487	1.8	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,458	1.8	---
11	Septicemia (A40-A41)	2,329	1.2	---
12	Pneumonitis due to solids and liquids (J69)	1,706	0.9	---
13	Parkinson's disease (G20-G21)	1,651	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,041	0.5	---
15	Atherosclerosis (I70)	984	0.5	---
...	All other causes (Residual)	49,973	26.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 95-99 years				
...	All causes	85,985	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,124	31.5	---
2	Alzheimer's disease (G30)	7,652	8.9	---
3	Cerebrovascular diseases (I60-I69)	5,997	7.0	---
4	Malignant neoplasms (C00-C97)	4,836	5.6	---
5	Influenza and pneumonia (J09-J18)	3,030	3.5	---
6	Chronic lower respiratory diseases (J40-J47)	2,811	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,058	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,988	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,420	1.7	---
10	Diabetes mellitus (E10-E14)	1,142	1.3	---
11	Septicemia (A40-A41)	892	1.0	---
12	Pneumonitis due to solids and liquids (J69)	825	1.0	---
13	Atherosclerosis (I70)	555	0.6	---
14	Parkinson's disease (G20-G21)	444	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	330	0.4	---
...	All other causes (Residual)	24,881	28.9	---
All races, female, 100 years and over				
...	All causes	20,642	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,971	33.8	---
2	Alzheimer's disease (G30)	1,657	8.0	---
3	Cerebrovascular diseases (I60-I69)	1,241	6.0	---
4	Influenza and pneumonia (J09-J18)	884	4.3	---
5	Malignant neoplasms (C00-C97)	669	3.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	632	3.1	---
7	Chronic lower respiratory diseases (J40-J47)	490	2.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	436	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	309	1.5	---
10	Diabetes mellitus (E10-E14)	234	1.1	---
11	Pneumonitis due to solids and liquids (J69)	201	1.0	---
12	Septicemia (A40-A41)	196	0.9	---
13	Atherosclerosis (I70)	192	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.3	---
15	Nutritional deficiencies (E40-E64)	68	0.3	---
...	All other causes (Residual)	6,390	31.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, all ages				
...	All causes	2,217,103	100.0	889.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	522,645	23.6	209.6
2	Malignant neoplasms (C00-C97)	498,116	22.5	199.8
3	Chronic lower respiratory diseases (J40-J47)	136,682	6.2	54.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112,803	5.1	45.2
5	Cerebrovascular diseases (I60-I69)	107,909	4.9	43.3
6	Alzheimer's disease (G30)	77,387	3.5	31.0
7	Diabetes mellitus (E10-E14)	58,925	2.7	23.6
8	Influenza and pneumonia (J09-J18)	49,013	2.2	19.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37,270	1.7	14.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37,154	1.7	14.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	31,871	1.4	12.8
12	Septicemia (A40-A41)	30,889	1.4	12.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24,149	1.1	9.7
14	Parkinson's disease (G20-G21)	23,553	1.1	9.4
15	Pneumonitis due to solids and liquids (J69)	16,458	0.7	6.6
...	All other causes (Residual)	452,279	20.4	181.4
White, both sexes, 1-4 years				
...	All causes	2,774	100.0	23.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	929	33.5	7.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	345	12.4	2.9
3	Malignant neoplasms (C00-C97)	246	8.9	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	179	6.5	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	3.7	0.9
6	Influenza and pneumonia (J09-J18)	70	2.5	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	1.3	0.3
8	Chronic lower respiratory diseases (J40-J47)	32	1.2	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	30	1.1	0.3
10	Septicemia (A40-A41)	29	1.0	0.2
11	Cerebrovascular diseases (I60-I69)	20	0.7	0.2
12	Meningitis (G00,G03)	13	0.5	*
13	Anemias (D50-D64)	11	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	11	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
...	All other causes (Residual)	709	25.6	5.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 5-9 years				
...	All causes	1,675	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	513	30.6	3.3
2	Malignant neoplasms (C00-C97)	351	21.0	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	7.6	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	76	4.5	0.5
5	Influenza and pneumonia (J09-J18)	51	3.0	0.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	2.5	0.3
7	Cerebrovascular diseases (I60-I69)	32	1.9	0.2
8	Chronic lower respiratory diseases (J40-J47)	26	1.6	0.2
9	Septicemia (A40-A41)	25	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.2	0.1
11	Certain conditions originating in the perinatal period (P00-P96)	12	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.4	*
13	Anemias (D50-D64)	5	0.3	*
14	Diabetes mellitus (E10-E14)	4	0.2	*
14	Meningitis (G00,G03)	4	0.2	*
14	Pneumonitis due to solids and liquids (J69)	4	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.2	*
...	All other causes (Residual)	369	22.0	2.4
White, both sexes, 10-14 years				
...	All causes	2,166	100.0	13.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	592	27.3	3.8
2	Malignant neoplasms (C00-C97)	366	16.9	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	308	14.2	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	115	5.3	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	4.3	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	63	2.9	0.4
7	Influenza and pneumonia (J09-J18)	41	1.9	0.3
8	Cerebrovascular diseases (I60-I69)	34	1.6	0.2
9	Chronic lower respiratory diseases (J40-J47)	33	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.2	0.2
11	Diabetes mellitus (E10-E14)	16	0.7	*
12	Septicemia (A40-A41)	13	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	12	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.5	*
15	Meningitis (G00,G03)	5	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
...	All other causes (Residual)	433	20.0	2.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-19 years				
...	All causes	6,852	100.0	42.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,995	43.7	18.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,462	21.3	9.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	489	7.1	3.0
4	Malignant neoplasms (C00-C97)	480	7.0	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	2.8	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	1.8	0.8
7	Influenza and pneumonia (J09-J18)	58	0.8	0.4
8	Cerebrovascular diseases (I60-I69)	35	0.5	0.2
9	Septicemia (A40-A41)	30	0.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.4	0.2
9	Diabetes mellitus (E10-E14)	30	0.4	0.2
12	Pregnancy, childbirth and the puerperium (O00-O99)	29	0.4	0.2
13	Chronic lower respiratory diseases (J40-J47)	25	0.4	0.2
14	Certain conditions originating in the perinatal period (P00-P96)	14	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.2	*
...	All other causes (Residual)	843	12.3	5.2
White, both sexes, 20-24 years				
...	All causes	13,766	100.0	80.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,723	48.8	39.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,541	18.5	14.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,014	7.4	5.9
4	Malignant neoplasms (C00-C97)	658	4.8	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	408	3.0	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	155	1.1	0.9
7	Influenza and pneumonia (J09-J18)	101	0.7	0.6
8	Diabetes mellitus (E10-E14)	83	0.6	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	74	0.5	0.4
10	Cerebrovascular diseases (I60-I69)	72	0.5	0.4
11	Septicemia (A40-A41)	62	0.5	0.4
12	Chronic lower respiratory diseases (J40-J47)	43	0.3	0.3
12	Legal intervention (Y35,Y89.0)	43	0.3	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.2	0.2
...	All other causes (Residual)	1,727	12.5	10.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 25-29 years				
...	All causes	15,567	100.0	94.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,865	44.1	41.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,753	17.7	16.7
3	Malignant neoplasms (C00-C97)	1,029	6.6	6.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	813	5.2	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	757	4.9	4.6
6	Diabetes mellitus (E10-E14)	156	1.0	0.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	153	1.0	0.9
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	140	0.9	0.8
9	Influenza and pneumonia (J09-J18)	132	0.8	0.8
10	Cerebrovascular diseases (I60-I69)	127	0.8	0.8
11	Pregnancy, childbirth and the puerperium (O00-O99)	125	0.8	0.8
12	Septicemia (A40-A41)	92	0.6	0.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	85	0.5	0.5
14	Chronic lower respiratory diseases (J40-J47)	81	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.3	0.3
...	All other causes (Residual)	2,209	14.2	13.4
White, both sexes, 30-34 years				
...	All causes	18,210	100.0	111.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,893	37.9	42.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,641	14.5	16.2
3	Malignant neoplasms (C00-C97)	1,754	9.6	10.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,387	7.6	8.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	799	4.4	4.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	386	2.1	2.4
7	Diabetes mellitus (E10-E14)	273	1.5	1.7
8	Cerebrovascular diseases (I60-I69)	211	1.2	1.3
9	Influenza and pneumonia (J09-J18)	203	1.1	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	196	1.1	1.2
11	Pregnancy, childbirth and the puerperium (O00-O99)	137	0.8	0.8
12	Septicemia (A40-A41)	133	0.7	0.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	131	0.7	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	0.6	0.6
15	Chronic lower respiratory diseases (J40-J47)	103	0.6	0.6
...	All other causes (Residual)	2,859	15.7	17.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-39 years				
...	All causes	20,794	100.0	138.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,130	29.5	40.7
2	Malignant neoplasms (C00-C97)	2,894	13.9	19.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,666	12.8	17.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,257	10.9	15.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	696	3.3	4.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	635	3.1	4.2
7	Diabetes mellitus (E10-E14)	457	2.2	3.0
8	Cerebrovascular diseases (I60-I69)	325	1.6	2.2
9	Influenza and pneumonia (J09-J18)	275	1.3	1.8
10	Septicemia (A40-A41)	205	1.0	1.4
11	Human immunodeficiency virus (HIV) disease (B20-B24)	190	0.9	1.3
12	Chronic lower respiratory diseases (J40-J47)	176	0.8	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	166	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	124	0.6	0.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	105	0.5	0.7
...	All other causes (Residual)	3,493	16.8	23.2
White, both sexes, 40-44 years				
...	All causes	31,736	100.0	195.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,893	21.7	42.4
2	Malignant neoplasms (C00-C97)	5,810	18.3	35.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,853	15.3	29.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,158	10.0	19.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,378	4.3	8.5
6	Diabetes mellitus (E10-E14)	870	2.7	5.4
7	Cerebrovascular diseases (I60-I69)	732	2.3	4.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	608	1.9	3.7
9	Influenza and pneumonia (J09-J18)	401	1.3	2.5
10	Septicemia (A40-A41)	355	1.1	2.2
11	Chronic lower respiratory diseases (J40-J47)	350	1.1	2.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	346	1.1	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	229	0.7	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	188	0.6	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	149	0.5	0.9
...	All other causes (Residual)	5,416	17.1	33.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-49 years				
...	All causes	51,599	100.0	307.1
1	Malignant neoplasms (C00-C97)	11,926	23.1	71.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,204	17.8	54.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,092	15.7	48.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,665	7.1	21.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,882	5.6	17.2
6	Diabetes mellitus (E10-E14)	1,544	3.0	9.2
7	Cerebrovascular diseases (I60-I69)	1,284	2.5	7.6
8	Chronic lower respiratory diseases (J40-J47)	998	1.9	5.9
9	Influenza and pneumonia (J09-J18)	620	1.2	3.7
10	Septicemia (A40-A41)	612	1.2	3.6
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	587	1.1	3.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	509	1.0	3.0
13	Viral hepatitis (B15-B19)	492	1.0	2.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	431	0.8	2.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	272	0.5	1.6
...	All other causes (Residual)	8,481	16.4	50.5
White, both sexes, 50-54 years				
...	All causes	87,487	100.0	480.3
1	Malignant neoplasms (C00-C97)	24,618	28.1	135.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,163	19.6	94.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,192	10.5	50.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,750	5.4	26.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,319	4.9	23.7
6	Chronic lower respiratory diseases (J40-J47)	2,874	3.3	15.8
7	Diabetes mellitus (E10-E14)	2,684	3.1	14.7
8	Cerebrovascular diseases (I60-I69)	2,326	2.7	12.8
9	Viral hepatitis (B15-B19)	1,110	1.3	6.1
10	Septicemia (A40-A41)	1,101	1.3	6.0
11	Influenza and pneumonia (J09-J18)	1,083	1.2	5.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	841	1.0	4.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	641	0.7	3.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	632	0.7	3.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	562	0.6	3.1
...	All other causes (Residual)	13,591	15.5	74.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-59 years				
...	All causes	120,916	100.0	697.6
1	Malignant neoplasms (C00-C97)	39,372	32.6	227.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,546	20.3	141.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,224	6.8	47.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,499	4.5	31.7
5	Chronic lower respiratory diseases (J40-J47)	5,401	4.5	31.2
6	Diabetes mellitus (E10-E14)	4,143	3.4	23.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,913	3.2	22.6
8	Cerebrovascular diseases (I60-I69)	3,416	2.8	19.7
9	Viral hepatitis (B15-B19)	1,729	1.4	10.0
10	Septicemia (A40-A41)	1,683	1.4	9.7
11	Influenza and pneumonia (J09-J18)	1,616	1.3	9.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,290	1.1	7.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	956	0.8	5.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	511	0.4	2.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	492	0.4	2.8
...	All other causes (Residual)	18,125	15.0	104.6
White, both sexes, 60-64 years				
...	All causes	148,369	100.0	986.3
1	Malignant neoplasms (C00-C97)	52,338	35.3	347.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,471	21.2	209.2
3	Chronic lower respiratory diseases (J40-J47)	8,501	5.7	56.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,996	4.0	39.9
5	Diabetes mellitus (E10-E14)	5,454	3.7	36.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,811	3.2	32.0
7	Cerebrovascular diseases (I60-I69)	4,597	3.1	30.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,801	1.9	18.6
9	Septicemia (A40-A41)	2,354	1.6	15.6
10	Influenza and pneumonia (J09-J18)	2,112	1.4	14.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,935	1.3	12.9
12	Viral hepatitis (B15-B19)	1,422	1.0	9.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,167	0.8	7.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	728	0.5	4.8
15	Aortic aneurysm and dissection (I71)	613	0.4	4.1
...	All other causes (Residual)	22,069	14.9	146.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 65-69 years				
...	All causes	179,185	100.0	1,444.9
1	Malignant neoplasms (C00-C97)	64,534	36.0	520.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,038	21.2	306.7
3	Chronic lower respiratory diseases (J40-J47)	13,743	7.7	110.8
4	Diabetes mellitus (E10-E14)	6,604	3.7	53.3
5	Cerebrovascular diseases (I60-I69)	6,228	3.5	50.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,978	2.8	40.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,600	2.0	29.0
8	Septicemia (A40-A41)	2,856	1.6	23.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,751	1.5	22.2
10	Influenza and pneumonia (J09-J18)	2,713	1.5	21.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,047	1.1	16.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,415	0.8	11.4
13	Alzheimer's disease (G30)	1,163	0.6	9.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,058	0.6	8.5
15	Parkinson's disease (G20-G21)	908	0.5	7.3
...	All other causes (Residual)	26,549	14.8	214.1
White, both sexes, 70-74 years				
...	All causes	203,937	100.0	2,246.5
1	Malignant neoplasms (C00-C97)	68,530	33.6	754.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	42,965	21.1	473.3
3	Chronic lower respiratory diseases (J40-J47)	18,856	9.2	207.7
4	Cerebrovascular diseases (I60-I69)	8,444	4.1	93.0
5	Diabetes mellitus (E10-E14)	6,781	3.3	74.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,541	2.2	50.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,581	1.8	39.4
8	Influenza and pneumonia (J09-J18)	3,404	1.7	37.5
9	Septicemia (A40-A41)	3,269	1.6	36.0
10	Alzheimer's disease (G30)	2,893	1.4	31.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,729	1.3	30.1
12	Parkinson's disease (G20-G21)	2,034	1.0	22.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,680	0.8	18.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,569	0.8	17.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,378	0.7	15.2
...	All other causes (Residual)	31,283	15.3	344.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-79 years				
...	All causes	239,264	100.0	3,622.8
1	Malignant neoplasms (C00-C97)	67,428	28.2	1,021.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,953	22.1	801.8
3	Chronic lower respiratory diseases (J40-J47)	21,558	9.0	326.4
4	Cerebrovascular diseases (I60-I69)	12,293	5.1	186.1
5	Diabetes mellitus (E10-E14)	7,374	3.1	111.7
6	Alzheimer's disease (G30)	6,780	2.8	102.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,478	2.3	82.9
8	Influenza and pneumonia (J09-J18)	4,703	2.0	71.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,616	1.9	69.9
10	Parkinson's disease (G20-G21)	3,916	1.6	59.3
11	Septicemia (A40-A41)	3,906	1.6	59.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,229	0.9	33.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,084	0.9	31.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,769	0.7	26.8
15	Pneumonitis due to solids and liquids (J69)	1,665	0.7	25.2
...	All other causes (Residual)	40,512	16.9	613.4
White, both sexes, 80-84 years				
...	All causes	308,071	100.0	6,094.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,844	24.3	1,480.6
2	Malignant neoplasms (C00-C97)	66,883	21.7	1,323.2
3	Chronic lower respiratory diseases (J40-J47)	24,377	7.9	482.3
4	Cerebrovascular diseases (I60-I69)	18,429	6.0	364.6
5	Alzheimer's disease (G30)	14,086	4.6	278.7
6	Diabetes mellitus (E10-E14)	8,285	2.7	163.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,682	2.5	152.0
8	Influenza and pneumonia (J09-J18)	7,379	2.4	146.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,182	2.0	122.3
10	Parkinson's disease (G20-G21)	5,957	1.9	117.8
11	Septicemia (A40-A41)	4,665	1.5	92.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,443	1.1	68.1
13	Pneumonitis due to solids and liquids (J69)	2,699	0.9	53.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,368	0.8	46.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,586	0.5	31.4
...	All other causes (Residual)	59,206	19.2	1,171.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 85-89 years				
...	All causes	352,166	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	96,248	27.3	---
2	Malignant neoplasms (C00-C97)	54,120	15.4	---
3	Cerebrovascular diseases (I60-I69)	22,862	6.5	---
4	Chronic lower respiratory diseases (J40-J47)	22,287	6.3	---
5	Alzheimer's disease (G30)	21,889	6.2	---
6	Influenza and pneumonia (J09-J18)	9,953	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,138	2.6	---
8	Diabetes mellitus (E10-E14)	7,813	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,293	2.1	---
10	Parkinson's disease (G20-G21)	6,179	1.8	---
11	Septicemia (A40-A41)	4,946	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,721	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,776	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,498	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,582	0.4	---
...	All other causes (Residual)	76,861	21.8	---
White, both sexes, 90-94 years				
...	All causes	268,958	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,655	30.7	---
2	Malignant neoplasms (C00-C97)	27,003	10.0	---
3	Alzheimer's disease (G30)	19,573	7.3	---
4	Cerebrovascular diseases (I60-I69)	18,164	6.8	---
5	Chronic lower respiratory diseases (J40-J47)	12,764	4.7	---
6	Influenza and pneumonia (J09-J18)	8,927	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,977	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,432	2.0	---
9	Diabetes mellitus (E10-E14)	4,768	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,549	1.7	---
11	Septicemia (A40-A41)	3,232	1.2	---
12	Parkinson's disease (G20-G21)	3,174	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,084	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,689	0.6	---
15	Atherosclerosis (I70)	1,301	0.5	---
...	All other causes (Residual)	65,666	24.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 95-99 years				
...	All causes	106,531	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,667	32.5	---
2	Alzheimer's disease (G30)	8,494	8.0	---
3	Cerebrovascular diseases (I60-I69)	6,969	6.5	---
4	Malignant neoplasms (C00-C97)	6,920	6.5	---
5	Influenza and pneumonia (J09-J18)	4,048	3.8	---
6	Chronic lower respiratory diseases (J40-J47)	3,871	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,767	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,160	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,955	1.8	---
10	Diabetes mellitus (E10-E14)	1,362	1.3	---
11	Pneumonitis due to solids and liquids (J69)	1,313	1.2	---
12	Septicemia (A40-A41)	1,052	1.0	---
13	Parkinson's disease (G20-G21)	713	0.7	---
14	Atherosclerosis (I70)	694	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	470	0.4	---
...	All other causes (Residual)	29,076	27.3	---
White, both sexes, 100 years and over				
...	All causes	21,854	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,615	34.8	---
2	Alzheimer's disease (G30)	1,635	7.5	---
3	Cerebrovascular diseases (I60-I69)	1,244	5.7	---
4	Influenza and pneumonia (J09-J18)	1,022	4.7	---
5	Malignant neoplasms (C00-C97)	796	3.6	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	627	2.9	---
7	Chronic lower respiratory diseases (J40-J47)	570	2.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	492	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	1.6	---
10	Pneumonitis due to solids and liquids (J69)	272	1.2	---
11	Diabetes mellitus (E10-E14)	216	1.0	---
12	Atherosclerosis (I70)	203	0.9	---
13	Septicemia (A40-A41)	177	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	86	0.4	---
15	Parkinson's disease (G20-G21)	85	0.4	---
...	All other causes (Residual)	6,471	29.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,110,956	100.0	899.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	275,101	24.8	222.6
2	Malignant neoplasms (C00-C97)	263,167	23.7	213.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70,161	6.3	56.8
4	Chronic lower respiratory diseases (J40-J47)	63,757	5.7	51.6
5	Cerebrovascular diseases (I60-I69)	44,203	4.0	35.8
6	Diabetes mellitus (E10-E14)	31,745	2.9	25.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,943	2.6	23.4
8	Alzheimer's disease (G30)	23,648	2.1	19.1
9	Influenza and pneumonia (J09-J18)	22,907	2.1	18.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	20,884	1.9	16.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,800	1.7	15.2
12	Septicemia (A40-A41)	14,602	1.3	11.8
13	Parkinson's disease (G20-G21)	14,127	1.3	11.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,940	0.9	8.0
15	Pneumonitis due to solids and liquids (J69)	8,989	0.8	7.3
...	All other causes (Residual)	199,982	18.0	161.9
White, male, 1-4 years				
...	All causes	1,601	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	604	37.7	9.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	10.1	2.7
3	Malignant neoplasms (C00-C97)	142	8.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	6.1	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.4	0.9
6	Influenza and pneumonia (J09-J18)	39	2.4	0.6
7	Certain conditions originating in the perinatal period (P00-P96)	22	1.4	0.4
8	Chronic lower respiratory diseases (J40-J47)	19	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.1	*
10	Septicemia (A40-A41)	16	1.0	*
11	Cerebrovascular diseases (I60-I69)	9	0.6	*
12	Meningitis (G00,G03)	7	0.4	*
13	Anemias (D50-D64)	6	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (Residual)	393	24.5	6.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-9 years				
...	All causes	967	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	313	32.4	3.9
2	Malignant neoplasms (C00-C97)	189	19.5	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	7.4	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	47	4.9	0.6
5	Influenza and pneumonia (J09-J18)	27	2.8	0.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	2.2	0.3
7	Chronic lower respiratory diseases (J40-J47)	18	1.9	*
8	Septicemia (A40-A41)	14	1.4	*
8	Cerebrovascular diseases (I60-I69)	14	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	0.7	*
12	Anemias (D50-D64)	4	0.4	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.4	*
14	Meningitis (G00,G03)	3	0.3	*
15	Aortic aneurysm and dissection (I71)	2	0.2	*
15	Diseases of appendix (K35-K38)	2	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.2	*
...	All other causes (Residual)	216	22.3	2.7
White, male, 10-14 years				
...	All causes	1,252	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	29.6	4.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	199	15.9	2.5
3	Malignant neoplasms (C00-C97)	182	14.5	2.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	4.6	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	4.5	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	2.6	0.4
7	Influenza and pneumonia (J09-J18)	21	1.7	0.3
8	Chronic lower respiratory diseases (J40-J47)	20	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	17	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.1	*
11	Diabetes mellitus (E10-E14)	10	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.6	*
13	Septicemia (A40-A41)	7	0.6	*
14	Meningitis (G00,G03)	5	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
...	All other causes (Residual)	247	19.7	3.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 15-19 years				
...	All causes	4,767	100.0	57.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,084	43.7	25.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,128	23.7	13.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	406	8.5	4.9
4	Malignant neoplasms (C00-C97)	287	6.0	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	2.6	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	1.5	0.9
7	Influenza and pneumonia (J09-J18)	32	0.7	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.4	0.3
9	Septicemia (A40-A41)	19	0.4	*
9	Diabetes mellitus (E10-E14)	19	0.4	*
9	Cerebrovascular diseases (I60-I69)	19	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	16	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	10	0.2	*
14	Aortic aneurysm and dissection (I71)	8	0.2	*
15	Legal intervention (Y35,Y89.0)	5	0.1	*
...	All other causes (Residual)	516	10.8	6.2
White, male, 20-24 years				
...	All causes	10,113	100.0	114.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,088	50.3	57.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,075	20.5	23.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	806	8.0	9.1
4	Malignant neoplasms (C00-C97)	377	3.7	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	272	2.7	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	1.0	1.1
7	Influenza and pneumonia (J09-J18)	62	0.6	0.7
8	Diabetes mellitus (E10-E14)	48	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	42	0.4	0.5
10	Legal intervention (Y35,Y89.0)	40	0.4	0.5
11	Septicemia (A40-A41)	31	0.3	0.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.3	0.3
12	Chronic lower respiratory diseases (J40-J47)	26	0.3	0.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.2	*
...	All other causes (Residual)	1,081	10.7	12.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-29 years				
...	All causes	11,101	100.0	131.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,149	46.4	60.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,241	20.2	26.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	652	5.9	7.7
4	Malignant neoplasms (C00-C97)	573	5.2	6.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	511	4.6	6.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	96	0.9	1.1
7	Diabetes mellitus (E10-E14)	95	0.9	1.1
8	Cerebrovascular diseases (I60-I69)	74	0.7	0.9
8	Influenza and pneumonia (J09-J18)	74	0.7	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	0.7	0.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	66	0.6	0.8
12	Septicemia (A40-A41)	53	0.5	0.6
13	Chronic lower respiratory diseases (J40-J47)	48	0.4	0.6
14	Legal intervention (Y35,Y89.0)	46	0.4	0.5
15	Aortic aneurysm and dissection (I71)	29	0.3	0.3
...	All other causes (Residual)	1,321	11.9	15.6
White, male, 30-34 years				
...	All causes	12,334	100.0	148.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,087	41.2	61.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,052	16.6	24.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	970	7.9	11.7
4	Malignant neoplasms (C00-C97)	855	6.9	10.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	586	4.8	7.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	259	2.1	3.1
7	Diabetes mellitus (E10-E14)	175	1.4	2.1
8	Cerebrovascular diseases (I60-I69)	122	1.0	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	1.0	1.4
10	Influenza and pneumonia (J09-J18)	110	0.9	1.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	105	0.9	1.3
12	Septicemia (A40-A41)	76	0.6	0.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	0.4	0.7
14	Chronic lower respiratory diseases (J40-J47)	54	0.4	0.7
15	Legal intervention (Y35,Y89.0)	48	0.4	0.6
...	All other causes (Residual)	1,660	13.5	20.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 35-39 years				
...	All causes	13,215	100.0	173.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,274	32.3	56.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,055	15.6	26.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,580	12.0	20.7
4	Malignant neoplasms (C00-C97)	1,271	9.6	16.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	458	3.5	6.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	456	3.5	6.0
7	Diabetes mellitus (E10-E14)	273	2.1	3.6
8	Cerebrovascular diseases (I60-I69)	183	1.4	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	150	1.1	2.0
10	Influenza and pneumonia (J09-J18)	148	1.1	1.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	98	0.7	1.3
12	Septicemia (A40-A41)	93	0.7	1.2
13	Chronic lower respiratory diseases (J40-J47)	82	0.6	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	0.5	0.9
15	Aortic aneurysm and dissection (I71)	60	0.5	0.8
...	All other causes (Residual)	1,969	14.9	25.8
White, male, 40-44 years				
...	All causes	19,669	100.0	240.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,655	23.7	56.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,436	17.5	42.0
3	Malignant neoplasms (C00-C97)	2,515	12.8	30.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,383	12.1	29.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	933	4.7	11.4
6	Diabetes mellitus (E10-E14)	557	2.8	6.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	432	2.2	5.3
8	Cerebrovascular diseases (I60-I69)	420	2.1	5.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	282	1.4	3.4
10	Influenza and pneumonia (J09-J18)	214	1.1	2.6
11	Septicemia (A40-A41)	179	0.9	2.2
12	Chronic lower respiratory diseases (J40-J47)	146	0.7	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	125	0.6	1.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	104	0.5	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	0.5	1.2
...	All other causes (Residual)	3,186	16.2	38.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-49 years				
...	All causes	31,717	100.0	376.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,654	21.0	79.1
2	Malignant neoplasms (C00-C97)	5,698	18.0	67.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,434	17.1	64.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,686	8.5	31.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,926	6.1	22.9
6	Diabetes mellitus (E10-E14)	968	3.1	11.5
7	Cerebrovascular diseases (I60-I69)	725	2.3	8.6
8	Chronic lower respiratory diseases (J40-J47)	434	1.4	5.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	414	1.3	4.9
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	409	1.3	4.9
11	Viral hepatitis (B15-B19)	356	1.1	4.2
12	Influenza and pneumonia (J09-J18)	343	1.1	4.1
13	Septicemia (A40-A41)	326	1.0	3.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	239	0.8	2.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	194	0.6	2.3
...	All other causes (Residual)	4,911	15.5	58.4
White, male, 50-54 years				
...	All causes	54,073	100.0	598.7
1	Malignant neoplasms (C00-C97)	12,752	23.6	141.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,509	23.1	138.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,088	11.3	67.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,230	6.0	35.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,230	6.0	35.8
6	Diabetes mellitus (E10-E14)	1,728	3.2	19.1
7	Chronic lower respiratory diseases (J40-J47)	1,355	2.5	15.0
8	Cerebrovascular diseases (I60-I69)	1,308	2.4	14.5
9	Viral hepatitis (B15-B19)	775	1.4	8.6
10	Influenza and pneumonia (J09-J18)	623	1.2	6.9
11	Septicemia (A40-A41)	588	1.1	6.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	521	1.0	5.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	506	0.9	5.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	422	0.8	4.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	411	0.8	4.6
...	All other causes (Residual)	8,027	14.8	88.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 55-59 years				
...	All causes	75,104	100.0	883.7
1	Malignant neoplasms (C00-C97)	21,634	28.8	254.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,823	23.7	209.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,568	7.4	65.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,927	5.2	46.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,918	3.9	34.3
6	Chronic lower respiratory diseases (J40-J47)	2,752	3.7	32.4
7	Diabetes mellitus (E10-E14)	2,619	3.5	30.8
8	Cerebrovascular diseases (I60-I69)	2,025	2.7	23.8
9	Viral hepatitis (B15-B19)	1,226	1.6	14.4
10	Influenza and pneumonia (J09-J18)	960	1.3	11.3
11	Septicemia (A40-A41)	916	1.2	10.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	731	1.0	8.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	641	0.9	7.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	403	0.5	4.7
15	Aortic aneurysm and dissection (I71)	312	0.4	3.7
...	All other causes (Residual)	10,649	14.2	125.3
White, male, 60-64 years				
...	All causes	90,766	100.0	1,245.4
1	Malignant neoplasms (C00-C97)	29,367	32.4	403.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,283	24.5	305.8
3	Chronic lower respiratory diseases (J40-J47)	4,490	4.9	61.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,075	4.5	55.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,456	3.8	47.4
6	Diabetes mellitus (E10-E14)	3,359	3.7	46.1
7	Cerebrovascular diseases (I60-I69)	2,701	3.0	37.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,147	2.4	29.5
9	Septicemia (A40-A41)	1,291	1.4	17.7
10	Influenza and pneumonia (J09-J18)	1,191	1.3	16.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,095	1.2	15.0
12	Viral hepatitis (B15-B19)	1,043	1.1	14.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	755	0.8	10.4
14	Aortic aneurysm and dissection (I71)	455	0.5	6.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	425	0.5	5.8
...	All other causes (Residual)	12,633	13.9	173.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-69 years				
...	All causes	104,448	100.0	1,760.0
1	Malignant neoplasms (C00-C97)	35,915	34.4	605.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,458	24.4	429.0
3	Chronic lower respiratory diseases (J40-J47)	6,843	6.6	115.3
4	Diabetes mellitus (E10-E14)	3,961	3.8	66.7
5	Cerebrovascular diseases (I60-I69)	3,416	3.3	57.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,302	3.2	55.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,356	2.3	39.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,602	1.5	27.0
9	Influenza and pneumonia (J09-J18)	1,528	1.5	25.7
10	Septicemia (A40-A41)	1,504	1.4	25.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,497	1.4	25.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	817	0.8	13.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	639	0.6	10.8
14	Parkinson's disease (G20-G21)	601	0.6	10.1
15	Aortic aneurysm and dissection (I71)	526	0.5	8.9
...	All other causes (Residual)	14,483	13.9	244.0
White, male, 70-74 years				
...	All causes	114,192	100.0	2,702.7
1	Malignant neoplasms (C00-C97)	37,569	32.9	889.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,962	23.6	638.1
3	Chronic lower respiratory diseases (J40-J47)	9,399	8.2	222.5
4	Cerebrovascular diseases (I60-I69)	4,467	3.9	105.7
5	Diabetes mellitus (E10-E14)	3,889	3.4	92.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,777	2.4	65.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,907	1.7	45.1
8	Influenza and pneumonia (J09-J18)	1,868	1.6	44.2
9	Septicemia (A40-A41)	1,698	1.5	40.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,675	1.5	39.6
11	Parkinson's disease (G20-G21)	1,376	1.2	32.6
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,331	1.2	31.5
13	Alzheimer's disease (G30)	1,238	1.1	29.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	873	0.8	20.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	825	0.7	19.5
...	All other causes (Residual)	16,338	14.3	386.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 75-79 years				
...	All causes	127,019	100.0	4,304.0
1	Malignant neoplasms (C00-C97)	36,529	28.8	1,237.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,945	24.4	1,048.6
3	Chronic lower respiratory diseases (J40-J47)	10,526	8.3	356.7
4	Cerebrovascular diseases (I60-I69)	5,861	4.6	198.6
5	Diabetes mellitus (E10-E14)	4,022	3.2	136.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,110	2.4	105.4
7	Alzheimer's disease (G30)	2,804	2.2	95.0
8	Parkinson's disease (G20-G21)	2,585	2.0	87.6
9	Influenza and pneumonia (J09-J18)	2,473	1.9	83.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,450	1.9	83.0
11	Septicemia (A40-A41)	1,871	1.5	63.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1,165	0.9	39.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,039	0.8	35.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,010	0.8	34.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,010	0.8	34.2
...	All other causes (Residual)	19,619	15.4	664.8
White, male, 80-84 years				
...	All causes	151,438	100.0	7,194.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,285	25.9	1,866.3
2	Malignant neoplasms (C00-C97)	35,354	23.3	1,679.5
3	Chronic lower respiratory diseases (J40-J47)	11,456	7.6	544.2
4	Cerebrovascular diseases (I60-I69)	7,846	5.2	372.7
5	Alzheimer's disease (G30)	5,173	3.4	245.8
6	Diabetes mellitus (E10-E14)	4,190	2.8	199.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,002	2.6	190.1
8	Influenza and pneumonia (J09-J18)	3,718	2.5	176.6
9	Parkinson's disease (G20-G21)	3,714	2.5	176.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,275	2.2	155.6
11	Septicemia (A40-A41)	2,215	1.5	105.2
12	Pneumonitis due to solids and liquids (J69)	1,534	1.0	72.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,435	0.9	68.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,320	0.9	62.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	914	0.6	43.4
...	All other causes (Residual)	26,007	17.2	1,235.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85-89 years				
...	All causes	150,826	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,600	28.9	---
2	Malignant neoplasms (C00-C97)	26,923	17.9	---
3	Chronic lower respiratory diseases (J40-J47)	9,676	6.4	---
4	Cerebrovascular diseases (I60-I69)	7,981	5.3	---
5	Alzheimer's disease (G30)	6,876	4.6	---
6	Influenza and pneumonia (J09-J18)	4,461	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,182	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,545	2.4	---
9	Parkinson's disease (G20-G21)	3,518	2.3	---
10	Diabetes mellitus (E10-E14)	3,492	2.3	---
11	Septicemia (A40-A41)	2,135	1.4	---
12	Pneumonitis due to solids and liquids (J69)	2,079	1.4	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,696	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,263	0.8	---
15	Aortic aneurysm and dissection (I71)	681	0.5	---
...	All other causes (Residual)	28,718	19.0	---
White, male, 90-94 years				
...	All causes	95,593	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,127	32.6	---
2	Malignant neoplasms (C00-C97)	12,212	12.8	---
3	Cerebrovascular diseases (I60-I69)	5,218	5.5	---
4	Chronic lower respiratory diseases (J40-J47)	5,012	5.2	---
5	Alzheimer's disease (G30)	4,983	5.2	---
6	Influenza and pneumonia (J09-J18)	3,441	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,662	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,421	2.5	---
9	Diabetes mellitus (E10-E14)	1,849	1.9	---
10	Parkinson's disease (G20-G21)	1,615	1.7	---
11	Pneumonitis due to solids and liquids (J69)	1,527	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,335	1.4	---
13	Septicemia (A40-A41)	1,200	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	717	0.8	---
15	Atherosclerosis (I70)	396	0.4	---
...	All other causes (Residual)	19,878	20.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 95-99 years				
...	All causes	28,292	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,880	34.9	---
2	Malignant neoplasms (C00-C97)	2,564	9.1	---
3	Cerebrovascular diseases (I60-I69)	1,518	5.4	---
4	Alzheimer's disease (G30)	1,496	5.3	---
5	Influenza and pneumonia (J09-J18)	1,289	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	1,254	4.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	811	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	713	2.5	---
9	Pneumonitis due to solids and liquids (J69)	552	2.0	---
10	Diabetes mellitus (E10-E14)	437	1.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	437	1.5	---
12	Parkinson's disease (G20-G21)	300	1.1	---
13	Septicemia (A40-A41)	284	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	167	0.6	---
15	Atherosclerosis (I70)	166	0.6	---
...	All other causes (Residual)	6,424	22.7	---
White, male, 100 years and over				
...	All causes	3,846	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,440	37.4	---
2	Malignant neoplasms (C00-C97)	225	5.9	---
3	Influenza and pneumonia (J09-J18)	220	5.7	---
4	Alzheimer's disease (G30)	202	5.3	---
4	Cerebrovascular diseases (I60-I69)	202	5.3	---
6	Chronic lower respiratory diseases (J40-J47)	123	3.2	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	2.5	---
7	Pneumonitis due to solids and liquids (J69)	97	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	2.4	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	2.2	---
11	Diabetes mellitus (E10-E14)	47	1.2	---
12	Atherosclerosis (I70)	36	0.9	---
13	Septicemia (A40-A41)	33	0.9	---
14	Parkinson's disease (G20-G21)	27	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.4	---
...	All other causes (Residual)	903	23.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, all ages				
...	All causes	1,106,147	100.0	879.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,544	22.4	196.8
2	Malignant neoplasms (C00-C97)	234,949	21.2	186.8
3	Chronic lower respiratory diseases (J40-J47)	72,925	6.6	58.0
4	Cerebrovascular diseases (I60-I69)	63,706	5.8	50.6
5	Alzheimer's disease (G30)	53,739	4.9	42.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42,642	3.9	33.9
7	Diabetes mellitus (E10-E14)	27,180	2.5	21.6
8	Influenza and pneumonia (J09-J18)	26,106	2.4	20.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,470	1.7	14.7
10	Septicemia (A40-A41)	16,287	1.5	12.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,209	1.3	11.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,987	1.0	8.7
13	Parkinson's disease (G20-G21)	9,426	0.9	7.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,211	0.7	6.5
15	Pneumonitis due to solids and liquids (J69)	7,469	0.7	5.9
...	All other causes (Residual)	252,297	22.8	200.6
White, female, 1-4 years				
...	All causes	1,173	100.0	20.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	325	27.7	5.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	15.6	3.1
3	Malignant neoplasms (C00-C97)	104	8.9	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	6.9	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	4.1	0.8
6	Influenza and pneumonia (J09-J18)	31	2.6	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.6	*
8	Septicemia (A40-A41)	13	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.1	*
10	Cerebrovascular diseases (I60-I69)	11	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
12	Meningitis (G00,G03)	6	0.5	*
13	Anemias (D50-D64)	5	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	316	26.9	5.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 5-9 years				
...	All causes	708	100.0	9.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	28.2	2.6
2	Malignant neoplasms (C00-C97)	162	22.9	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	7.8	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	4.1	0.4
5	Influenza and pneumonia (J09-J18)	24	3.4	0.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.0	0.3
7	Cerebrovascular diseases (I60-I69)	18	2.5	*
8	Septicemia (A40-A41)	11	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	8	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	5	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
14	Diabetes mellitus (E10-E14)	3	0.4	*
14	Pneumonitis due to solids and liquids (J69)	3	0.4	*
...	All other causes (Residual)	153	21.6	2.0
White, female, 10-14 years				
...	All causes	914	100.0	11.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	24.2	2.9
2	Malignant neoplasms (C00-C97)	184	20.1	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	109	11.9	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	6.2	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	4.2	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	3.4	0.4
7	Influenza and pneumonia (J09-J18)	20	2.2	0.3
8	Cerebrovascular diseases (I60-I69)	17	1.9	*
9	Chronic lower respiratory diseases (J40-J47)	13	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	0.8	*
12	Septicemia (A40-A41)	6	0.7	*
12	Diabetes mellitus (E10-E14)	6	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Anemias (D50-D64)	1	0.1	*
15	Nutritional deficiencies (E40-E64)	1	0.1	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.1	*
...	All other causes (Residual)	187	20.5	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-19 years				
...	All causes	2,085	100.0	26.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	911	43.7	11.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	334	16.0	4.3
3	Malignant neoplasms (C00-C97)	193	9.3	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	4.0	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	3.3	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	2.6	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	29	1.4	0.4
8	Influenza and pneumonia (J09-J18)	26	1.2	0.3
9	Cerebrovascular diseases (I60-I69)	16	0.8	*
10	Septicemia (A40-A41)	11	0.5	*
10	Diabetes mellitus (E10-E14)	11	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (Residual)	316	15.2	4.0
White, female, 20-24 years				
...	All causes	3,653	100.0	43.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,635	44.8	19.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	466	12.8	5.6
3	Malignant neoplasms (C00-C97)	281	7.7	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	208	5.7	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	136	3.7	1.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	74	2.0	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	1.5	0.6
8	Influenza and pneumonia (J09-J18)	39	1.1	0.5
9	Diabetes mellitus (E10-E14)	35	1.0	0.4
10	Septicemia (A40-A41)	31	0.8	0.4
11	Cerebrovascular diseases (I60-I69)	30	0.8	0.4
12	Chronic lower respiratory diseases (J40-J47)	17	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.2	*
...	All other causes (Residual)	617	16.9	7.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 25-29 years				
...	All causes	4,466	100.0	55.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,716	38.4	21.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	512	11.5	6.4
3	Malignant neoplasms (C00-C97)	456	10.2	5.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	246	5.5	3.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	161	3.6	2.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	125	2.8	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	1.5	0.8
8	Diabetes mellitus (E10-E14)	61	1.4	0.8
9	Influenza and pneumonia (J09-J18)	58	1.3	0.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	57	1.3	0.7
11	Cerebrovascular diseases (I60-I69)	53	1.2	0.7
12	Septicemia (A40-A41)	39	0.9	0.5
13	Chronic lower respiratory diseases (J40-J47)	33	0.7	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.5	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.5	0.3
...	All other causes (Residual)	839	18.8	10.5
White, female, 30-34 years				
...	All causes	5,876	100.0	73.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,806	30.7	22.6
2	Malignant neoplasms (C00-C97)	899	15.3	11.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	589	10.0	7.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	417	7.1	5.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	213	3.6	2.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	137	2.3	1.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	127	2.2	1.6
8	Diabetes mellitus (E10-E14)	98	1.7	1.2
9	Influenza and pneumonia (J09-J18)	93	1.6	1.2
10	Cerebrovascular diseases (I60-I69)	89	1.5	1.1
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	1.3	1.0
12	Septicemia (A40-A41)	57	1.0	0.7
13	Chronic lower respiratory diseases (J40-J47)	49	0.8	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	0.8	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.5	0.4
...	All other causes (Residual)	1,145	19.5	14.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-39 years				
...	All causes	7,579	100.0	102.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,856	24.5	25.0
2	Malignant neoplasms (C00-C97)	1,623	21.4	21.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	677	8.9	9.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	611	8.1	8.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	240	3.2	3.2
6	Diabetes mellitus (E10-E14)	184	2.4	2.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	177	2.3	2.4
8	Cerebrovascular diseases (I60-I69)	142	1.9	1.9
9	Influenza and pneumonia (J09-J18)	127	1.7	1.7
10	Septicemia (A40-A41)	112	1.5	1.5
11	Pregnancy, childbirth and the puerperium (O00-O99)	105	1.4	1.4
12	Chronic lower respiratory diseases (J40-J47)	94	1.2	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	0.9	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	0.8	0.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	40	0.5	0.5
...	All other causes (Residual)	1,464	19.3	19.7
White, female, 40-44 years				
...	All causes	12,067	100.0	149.8
1	Malignant neoplasms (C00-C97)	3,295	27.3	40.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,238	18.5	27.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,417	11.7	17.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	775	6.4	9.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	445	3.7	5.5
6	Diabetes mellitus (E10-E14)	313	2.6	3.9
7	Cerebrovascular diseases (I60-I69)	312	2.6	3.9
8	Chronic lower respiratory diseases (J40-J47)	204	1.7	2.5
9	Influenza and pneumonia (J09-J18)	187	1.5	2.3
10	Septicemia (A40-A41)	176	1.5	2.2
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	176	1.5	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	0.9	1.3
13	Pregnancy, childbirth and the puerperium (O00-O99)	88	0.7	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	0.7	1.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	64	0.5	0.8
...	All other causes (Residual)	2,187	18.1	27.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-49 years				
...	All causes	19,882	100.0	237.1
1	Malignant neoplasms (C00-C97)	6,228	31.3	74.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,658	13.4	31.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,550	12.8	30.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	979	4.9	11.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	956	4.8	11.4
6	Diabetes mellitus (E10-E14)	576	2.9	6.9
7	Chronic lower respiratory diseases (J40-J47)	564	2.8	6.7
8	Cerebrovascular diseases (I60-I69)	559	2.8	6.7
9	Septicemia (A40-A41)	286	1.4	3.4
10	Influenza and pneumonia (J09-J18)	277	1.4	3.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	192	1.0	2.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	178	0.9	2.1
13	Viral hepatitis (B15-B19)	136	0.7	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	0.6	1.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	95	0.5	1.1
...	All other causes (Residual)	3,521	17.7	42.0
White, female, 50-54 years				
...	All causes	33,414	100.0	363.9
1	Malignant neoplasms (C00-C97)	11,866	35.5	129.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,654	13.9	50.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,104	9.3	33.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,520	4.5	16.6
5	Chronic lower respiratory diseases (J40-J47)	1,519	4.5	16.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,089	3.3	11.9
7	Cerebrovascular diseases (I60-I69)	1,018	3.0	11.1
8	Diabetes mellitus (E10-E14)	956	2.9	10.4
9	Septicemia (A40-A41)	513	1.5	5.6
10	Influenza and pneumonia (J09-J18)	460	1.4	5.0
11	Viral hepatitis (B15-B19)	335	1.0	3.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	335	1.0	3.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	219	0.7	2.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	155	0.5	1.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	0.5	1.6
...	All other causes (Residual)	5,520	16.5	60.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-59 years				
...	All causes	45,812	100.0	518.6
1	Malignant neoplasms (C00-C97)	17,738	38.7	200.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,723	14.7	76.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,656	5.8	30.1
4	Chronic lower respiratory diseases (J40-J47)	2,649	5.8	30.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,572	3.4	17.8
6	Diabetes mellitus (E10-E14)	1,524	3.3	17.3
7	Cerebrovascular diseases (I60-I69)	1,391	3.0	15.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	995	2.2	11.3
9	Septicemia (A40-A41)	767	1.7	8.7
10	Influenza and pneumonia (J09-J18)	656	1.4	7.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	559	1.2	6.3
12	Viral hepatitis (B15-B19)	503	1.1	5.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	315	0.7	3.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	214	0.5	2.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	199	0.4	2.3
...	All other causes (Residual)	7,351	16.0	83.2
White, female, 60-64 years				
...	All causes	57,603	100.0	742.8
1	Malignant neoplasms (C00-C97)	22,971	39.9	296.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,188	16.0	118.5
3	Chronic lower respiratory diseases (J40-J47)	4,011	7.0	51.7
4	Diabetes mellitus (E10-E14)	2,095	3.6	27.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,921	3.3	24.8
6	Cerebrovascular diseases (I60-I69)	1,896	3.3	24.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,355	2.4	17.5
8	Septicemia (A40-A41)	1,063	1.8	13.7
9	Influenza and pneumonia (J09-J18)	921	1.6	11.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	840	1.5	10.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	654	1.1	8.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	412	0.7	5.3
13	Viral hepatitis (B15-B19)	379	0.7	4.9
14	Alzheimer's disease (G30)	351	0.6	4.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	303	0.5	3.9
...	All other causes (Residual)	9,243	16.0	119.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 65-69 years				
...	All causes	74,737	100.0	1,155.8
1	Malignant neoplasms (C00-C97)	28,619	38.3	442.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,580	16.8	194.5
3	Chronic lower respiratory diseases (J40-J47)	6,900	9.2	106.7
4	Cerebrovascular diseases (I60-I69)	2,812	3.8	43.5
5	Diabetes mellitus (E10-E14)	2,643	3.5	40.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,676	2.2	25.9
7	Septicemia (A40-A41)	1,352	1.8	20.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,254	1.7	19.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,244	1.7	19.2
10	Influenza and pneumonia (J09-J18)	1,185	1.6	18.3
11	Alzheimer's disease (G30)	642	0.9	9.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	598	0.8	9.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	445	0.6	6.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	419	0.6	6.5
15	Parkinson's disease (G20-G21)	307	0.4	4.7
...	All other causes (Residual)	12,061	16.1	186.5
White, female, 70-74 years				
...	All causes	89,745	100.0	1,849.4
1	Malignant neoplasms (C00-C97)	30,961	34.5	638.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,003	17.8	329.8
3	Chronic lower respiratory diseases (J40-J47)	9,457	10.5	194.9
4	Cerebrovascular diseases (I60-I69)	3,977	4.4	82.0
5	Diabetes mellitus (E10-E14)	2,892	3.2	59.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,764	2.0	36.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,674	1.9	34.5
8	Alzheimer's disease (G30)	1,655	1.8	34.1
9	Septicemia (A40-A41)	1,571	1.8	32.4
10	Influenza and pneumonia (J09-J18)	1,536	1.7	31.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,054	1.2	21.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	807	0.9	16.6
13	Parkinson's disease (G20-G21)	658	0.7	13.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	553	0.6	11.4
15	Pneumonitis due to solids and liquids (J69)	438	0.5	9.0
...	All other causes (Residual)	14,745	16.4	303.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-79 years				
...	All causes	112,245	100.0	3,072.5
1	Malignant neoplasms (C00-C97)	30,899	27.5	845.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,008	19.6	602.4
3	Chronic lower respiratory diseases (J40-J47)	11,032	9.8	302.0
4	Cerebrovascular diseases (I60-I69)	6,432	5.7	176.1
5	Alzheimer's disease (G30)	3,976	3.5	108.8
6	Diabetes mellitus (E10-E14)	3,352	3.0	91.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,368	2.1	64.8
8	Influenza and pneumonia (J09-J18)	2,230	2.0	61.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,166	1.9	59.3
10	Septicemia (A40-A41)	2,035	1.8	55.7
11	Parkinson's disease (G20-G21)	1,331	1.2	36.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,190	1.1	32.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	919	0.8	25.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	759	0.7	20.8
15	Pneumonitis due to solids and liquids (J69)	669	0.6	18.3
...	All other causes (Residual)	20,879	18.6	571.5
White, female, 80-84 years				
...	All causes	156,633	100.0	5,309.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,559	22.7	1,205.5
2	Malignant neoplasms (C00-C97)	31,529	20.1	1,068.8
3	Chronic lower respiratory diseases (J40-J47)	12,921	8.2	438.0
4	Cerebrovascular diseases (I60-I69)	10,583	6.8	358.8
5	Alzheimer's disease (G30)	8,913	5.7	302.2
6	Diabetes mellitus (E10-E14)	4,095	2.6	138.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,680	2.3	124.8
8	Influenza and pneumonia (J09-J18)	3,661	2.3	124.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,907	1.9	98.5
10	Septicemia (A40-A41)	2,450	1.6	83.1
11	Parkinson's disease (G20-G21)	2,243	1.4	76.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,008	1.3	68.1
13	Pneumonitis due to solids and liquids (J69)	1,165	0.7	39.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,048	0.7	35.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	782	0.5	26.5
...	All other causes (Residual)	33,089	21.1	1,121.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 85-89 years				
...	All causes	201,340	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,648	26.1	---
2	Malignant neoplasms (C00-C97)	27,197	13.5	---
3	Alzheimer's disease (G30)	15,013	7.5	---
4	Cerebrovascular diseases (I60-I69)	14,881	7.4	---
5	Chronic lower respiratory diseases (J40-J47)	12,611	6.3	---
6	Influenza and pneumonia (J09-J18)	5,492	2.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,956	2.5	---
8	Diabetes mellitus (E10-E14)	4,321	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,748	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,025	1.5	---
11	Septicemia (A40-A41)	2,811	1.4	---
12	Parkinson's disease (G20-G21)	2,661	1.3	---
13	Pneumonitis due to solids and liquids (J69)	1,697	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,235	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,023	0.5	---
...	All other causes (Residual)	48,021	23.9	---
White, female, 90-94 years				
...	All causes	173,365	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,528	29.7	---
2	Malignant neoplasms (C00-C97)	14,791	8.5	---
3	Alzheimer's disease (G30)	14,590	8.4	---
4	Cerebrovascular diseases (I60-I69)	12,946	7.5	---
5	Chronic lower respiratory diseases (J40-J47)	7,752	4.5	---
6	Influenza and pneumonia (J09-J18)	5,486	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,315	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,214	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,011	1.7	---
10	Diabetes mellitus (E10-E14)	2,919	1.7	---
11	Septicemia (A40-A41)	2,032	1.2	---
12	Parkinson's disease (G20-G21)	1,559	0.9	---
13	Pneumonitis due to solids and liquids (J69)	1,557	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	972	0.6	---
15	Atherosclerosis (I70)	905	0.5	---
...	All other causes (Residual)	45,788	26.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 95-99 years				
...	All causes	78,239	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,787	31.7	---
2	Alzheimer's disease (G30)	6,998	8.9	---
3	Cerebrovascular diseases (I60-I69)	5,451	7.0	---
4	Malignant neoplasms (C00-C97)	4,356	5.6	---
5	Influenza and pneumonia (J09-J18)	2,759	3.5	---
6	Chronic lower respiratory diseases (J40-J47)	2,617	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,956	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,723	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,242	1.6	---
10	Diabetes mellitus (E10-E14)	925	1.2	---
11	Septicemia (A40-A41)	768	1.0	---
12	Pneumonitis due to solids and liquids (J69)	761	1.0	---
13	Atherosclerosis (I70)	528	0.7	---
14	Parkinson's disease (G20-G21)	413	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	303	0.4	---
...	All other causes (Residual)	22,652	29.0	---
White, female, 100 years and over				
...	All causes	18,008	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,175	34.3	---
2	Alzheimer's disease (G30)	1,433	8.0	---
3	Cerebrovascular diseases (I60-I69)	1,042	5.8	---
4	Influenza and pneumonia (J09-J18)	802	4.5	---
5	Malignant neoplasms (C00-C97)	571	3.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	530	2.9	---
7	Chronic lower respiratory diseases (J40-J47)	447	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	401	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	257	1.4	---
10	Pneumonitis due to solids and liquids (J69)	175	1.0	---
11	Diabetes mellitus (E10-E14)	169	0.9	---
12	Atherosclerosis (I70)	167	0.9	---
13	Septicemia (A40-A41)	144	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.4	---
15	Parkinson's disease (G20-G21)	58	0.3	---
...	All other causes (Residual)	5,568	30.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	302,969	100.0	693.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,010	23.8	164.8
2	Malignant neoplasms (C00-C97)	67,953	22.4	155.5
3	Cerebrovascular diseases (I60-I69)	16,269	5.4	37.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,413	4.4	30.7
5	Diabetes mellitus (E10-E14)	13,385	4.4	30.6
6	Chronic lower respiratory diseases (J40-J47)	9,918	3.3	22.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,393	2.8	19.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,059	2.7	18.4
9	Septicemia (A40-A41)	6,250	2.1	14.3
10	Alzheimer's disease (G30)	5,714	1.9	13.1
11	Influenza and pneumonia (J09-J18)	5,567	1.8	12.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,455	1.8	12.5
13	Certain conditions originating in the perinatal period (P00-P96)	4,055	1.3	9.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,742	1.2	8.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,053	1.0	7.0
...	All other causes (Residual)	59,733	19.7	136.7
Black, both sexes, 1-4 years				
...	All causes	1,006	100.0	37.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	299	29.7	11.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	13.5	5.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	9.0	3.4
4	Malignant neoplasms (C00-C97)	64	6.4	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	5.9	2.2
6	Chronic lower respiratory diseases (J40-J47)	27	2.7	1.0
7	Influenza and pneumonia (J09-J18)	23	2.3	0.8
8	Septicemia (A40-A41)	20	2.0	0.7
9	Cerebrovascular diseases (I60-I69)	15	1.5	*
10	Certain conditions originating in the perinatal period (P00-P96)	13	1.3	*
11	Anemias (D50-D64)	8	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
13	Meningitis (G00,G03)	6	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
...	All other causes (Residual)	231	23.0	8.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-9 years				
...	All causes	582	100.0	17.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	183	31.4	5.4
2	Malignant neoplasms (C00-C97)	65	11.2	1.9
3	Chronic lower respiratory diseases (J40-J47)	45	7.7	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	7.6	1.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	6.5	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.8	0.6
7	Influenza and pneumonia (J09-J18)	13	2.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.9	*
9	Septicemia (A40-A41)	7	1.2	*
9	Anemias (D50-D64)	7	1.2	*
9	Cerebrovascular diseases (I60-I69)	7	1.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.9	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
...	All other causes (Residual)	123	21.1	3.6
Black, both sexes, 10-14 years				
...	All causes	587	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	25.0	4.4
2	Malignant neoplasms (C00-C97)	57	9.7	1.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	9.2	1.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	8.3	1.5
5	Chronic lower respiratory diseases (J40-J47)	42	7.2	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	6.8	1.2
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	4.9	0.9
8	Influenza and pneumonia (J09-J18)	15	2.6	*
9	Cerebrovascular diseases (I60-I69)	12	2.0	*
10	Septicemia (A40-A41)	6	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
12	Anemias (D50-D64)	4	0.7	*
12	Diabetes mellitus (E10-E14)	4	0.7	*
14	Meningitis (G00,G03)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
...	All other causes (Residual)	119	20.3	3.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 15-19 years				
...	All causes	2,180	100.0	62.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	884	40.6	25.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	507	23.3	14.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	163	7.5	4.6
4	Malignant neoplasms (C00-C97)	103	4.7	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	88	4.0	2.5
6	Chronic lower respiratory diseases (J40-J47)	34	1.6	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	1.2	0.8
8	Anemias (D50-D64)	22	1.0	0.6
9	Diabetes mellitus (E10-E14)	20	0.9	0.6
10	Cerebrovascular diseases (I60-I69)	18	0.8	*
11	Influenza and pneumonia (J09-J18)	13	0.6	*
11	Legal intervention (Y35,Y89.0)	13	0.6	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.5	*
14	Septicemia (A40-A41)	9	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.3	*
15	Diseases of appendix (K35-K38)	6	0.3	*
...	All other causes (Residual)	256	11.7	7.3
Black, both sexes, 20-24 years				
...	All causes	4,358	100.0	113.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,847	42.4	48.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	908	20.8	23.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	360	8.3	9.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	202	4.6	5.3
5	Malignant neoplasms (C00-C97)	138	3.2	3.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	83	1.9	2.2
7	Pregnancy, childbirth and the puerperium (O00-O99)	58	1.3	1.5
8	Diabetes mellitus (E10-E14)	53	1.2	1.4
9	Anemias (D50-D64)	48	1.1	1.3
10	Chronic lower respiratory diseases (J40-J47)	46	1.1	1.2
11	Legal intervention (Y35,Y89.0)	31	0.7	0.8
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.7	0.8
13	Septicemia (A40-A41)	27	0.6	0.7
14	Cerebrovascular diseases (I60-I69)	20	0.5	0.5
15	Influenza and pneumonia (J09-J18)	19	0.4	*
...	All other causes (Residual)	488	11.2	12.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-29 years				
...	All causes	4,467	100.0	139.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,374	30.8	43.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	903	20.2	28.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	347	7.8	10.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	303	6.8	9.5
5	Malignant neoplasms (C00-C97)	235	5.3	7.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	179	4.0	5.6
7	Diabetes mellitus (E10-E14)	101	2.3	3.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	87	1.9	2.7
9	Anemias (D50-D64)	53	1.2	1.7
10	Chronic lower respiratory diseases (J40-J47)	46	1.0	1.4
11	Cerebrovascular diseases (I60-I69)	42	0.9	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	0.9	1.2
13	Septicemia (A40-A41)	37	0.8	1.2
14	Influenza and pneumonia (J09-J18)	34	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	0.8	1.1
...	All other causes (Residual)	653	14.6	20.4
Black, both sexes, 30-34 years				
...	All causes	5,064	100.0	166.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,109	21.9	36.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	898	17.7	29.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	627	12.4	20.6
4	Malignant neoplasms (C00-C97)	410	8.1	13.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	261	5.2	8.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	218	4.3	7.2
7	Diabetes mellitus (E10-E14)	125	2.5	4.1
8	Cerebrovascular diseases (I60-I69)	103	2.0	3.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	1.3	2.2
10	Influenza and pneumonia (J09-J18)	63	1.2	2.1
10	Pregnancy, childbirth and the puerperium (O00-O99)	63	1.2	2.1
12	Anemias (D50-D64)	61	1.2	2.0
13	Septicemia (A40-A41)	53	1.0	1.7
13	Chronic lower respiratory diseases (J40-J47)	53	1.0	1.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	0.8	1.2
...	All other causes (Residual)	916	18.1	30.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 35-39 years				
...	All causes	5,702	100.0	207.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,024	18.0	37.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	841	14.7	30.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	737	12.9	26.9
4	Malignant neoplasms (C00-C97)	639	11.2	23.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	292	5.1	10.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	213	3.7	7.8
7	Diabetes mellitus (E10-E14)	189	3.3	6.9
8	Cerebrovascular diseases (I60-I69)	187	3.3	6.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	97	1.7	3.5
10	Chronic lower respiratory diseases (J40-J47)	96	1.7	3.5
11	Septicemia (A40-A41)	81	1.4	3.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	64	1.1	2.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	57	1.0	2.1
14	Influenza and pneumonia (J09-J18)	55	1.0	2.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	52	0.9	1.9
...	All other causes (Residual)	1,078	18.9	39.3
Black, both sexes, 40-44 years				
...	All causes	8,057	100.0	284.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,775	22.0	62.6
2	Malignant neoplasms (C00-C97)	1,278	15.9	45.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	934	11.6	32.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	497	6.2	17.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	390	4.8	13.7
6	Diabetes mellitus (E10-E14)	345	4.3	12.2
7	Cerebrovascular diseases (I60-I69)	309	3.8	10.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	218	2.7	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	148	1.8	5.2
10	Septicemia (A40-A41)	147	1.8	5.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	133	1.7	4.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	1.5	4.3
13	Chronic lower respiratory diseases (J40-J47)	112	1.4	3.9
14	Influenza and pneumonia (J09-J18)	105	1.3	3.7
15	Aortic aneurysm and dissection (I71)	64	0.8	2.3
...	All other causes (Residual)	1,481	18.4	52.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-49 years				
...	All causes	12,459	100.0	437.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,918	23.4	102.4
2	Malignant neoplasms (C00-C97)	2,555	20.5	89.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,082	8.7	38.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	555	4.5	19.5
5	Diabetes mellitus (E10-E14)	543	4.4	19.1
5	Cerebrovascular diseases (I60-I69)	543	4.4	19.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	433	3.5	15.2
8	Septicemia (A40-A41)	253	2.0	8.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	249	2.0	8.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	248	2.0	8.7
11	Chronic lower respiratory diseases (J40-J47)	220	1.8	7.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	198	1.6	7.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	196	1.6	6.9
14	Influenza and pneumonia (J09-J18)	164	1.3	5.8
15	Aortic aneurysm and dissection (I71)	79	0.6	2.8
...	All other causes (Residual)	2,223	17.8	78.0
Black, both sexes, 50-54 years				
...	All causes	19,652	100.0	680.6
1	Malignant neoplasms (C00-C97)	5,111	26.0	177.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,750	24.2	164.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,347	6.9	46.6
4	Cerebrovascular diseases (I60-I69)	927	4.7	32.1
5	Diabetes mellitus (E10-E14)	872	4.4	30.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	637	3.2	22.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	497	2.5	17.2
8	Chronic lower respiratory diseases (J40-J47)	439	2.2	15.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	432	2.2	15.0
10	Septicemia (A40-A41)	388	2.0	13.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	352	1.8	12.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	334	1.7	11.6
13	Influenza and pneumonia (J09-J18)	286	1.5	9.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	174	0.9	6.0
15	Viral hepatitis (B15-B19)	150	0.8	5.2
...	All other causes (Residual)	2,956	15.0	102.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 55-59 years				
...	All causes	27,213	100.0	1,058.6
1	Malignant neoplasms (C00-C97)	7,984	29.3	310.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,716	24.7	261.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,297	4.8	50.5
4	Cerebrovascular diseases (I60-I69)	1,277	4.7	49.7
5	Diabetes mellitus (E10-E14)	1,266	4.7	49.3
6	Chronic lower respiratory diseases (J40-J47)	735	2.7	28.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	687	2.5	26.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	623	2.3	24.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	598	2.2	23.3
10	Septicemia (A40-A41)	524	1.9	20.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	442	1.6	17.2
12	Influenza and pneumonia (J09-J18)	418	1.5	16.3
13	Viral hepatitis (B15-B19)	319	1.2	12.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	231	0.8	9.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	154	0.6	6.0
...	All other causes (Residual)	3,942	14.5	153.4
Black, both sexes, 60-64 years				
...	All causes	30,896	100.0	1,529.1
1	Malignant neoplasms (C00-C97)	9,721	31.5	481.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,604	24.6	376.3
3	Diabetes mellitus (E10-E14)	1,627	5.3	80.5
4	Cerebrovascular diseases (I60-I69)	1,549	5.0	76.7
5	Chronic lower respiratory diseases (J40-J47)	1,065	3.4	52.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,048	3.4	51.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	852	2.8	42.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	653	2.1	32.3
9	Septicemia (A40-A41)	652	2.1	32.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	466	1.5	23.1
11	Influenza and pneumonia (J09-J18)	447	1.4	22.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	384	1.2	19.0
13	Viral hepatitis (B15-B19)	325	1.1	16.1
14	Pneumonitis due to solids and liquids (J69)	124	0.4	6.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	0.4	6.0
...	All other causes (Residual)	4,258	13.8	210.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-69 years				
...	All causes	28,988	100.0	2,032.2
1	Malignant neoplasms (C00-C97)	9,261	31.9	649.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,307	25.2	512.3
3	Diabetes mellitus (E10-E14)	1,549	5.3	108.6
4	Cerebrovascular diseases (I60-I69)	1,519	5.2	106.5
5	Chronic lower respiratory diseases (J40-J47)	1,149	4.0	80.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	871	3.0	61.1
7	Septicemia (A40-A41)	690	2.4	48.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	612	2.1	42.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	529	1.8	37.1
10	Influenza and pneumonia (J09-J18)	493	1.7	34.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	336	1.2	23.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	202	0.7	14.2
13	Viral hepatitis (B15-B19)	169	0.6	11.8
14	Pneumonitis due to solids and liquids (J69)	137	0.5	9.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	130	0.4	9.1
...	All other causes (Residual)	4,034	13.9	282.8
Black, both sexes, 70-74 years				
...	All causes	29,142	100.0	2,943.6
1	Malignant neoplasms (C00-C97)	8,692	29.8	878.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,290	25.0	736.4
3	Cerebrovascular diseases (I60-I69)	1,697	5.8	171.4
4	Diabetes mellitus (E10-E14)	1,626	5.6	164.2
5	Chronic lower respiratory diseases (J40-J47)	1,370	4.7	138.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,023	3.5	103.3
7	Septicemia (A40-A41)	653	2.2	66.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	564	1.9	57.0
9	Influenza and pneumonia (J09-J18)	556	1.9	56.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	473	1.6	47.8
11	Alzheimer's disease (G30)	313	1.1	31.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	186	0.6	18.8
13	Pneumonitis due to solids and liquids (J69)	169	0.6	17.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	120	0.4	12.1
15	Parkinson's disease (G20-G21)	116	0.4	11.7
...	All other causes (Residual)	4,294	14.7	433.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 75-79 years				
...	All causes	29,789	100.0	4,270.6
1	Malignant neoplasms (C00-C97)	7,759	26.0	1,112.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,513	25.2	1,077.1
3	Cerebrovascular diseases (I60-I69)	1,943	6.5	278.6
4	Diabetes mellitus (E10-E14)	1,625	5.5	233.0
5	Chronic lower respiratory diseases (J40-J47)	1,314	4.4	188.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,033	3.5	148.1
7	Septicemia (A40-A41)	745	2.5	106.8
8	Alzheimer's disease (G30)	668	2.2	95.8
9	Influenza and pneumonia (J09-J18)	591	2.0	84.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	568	1.9	81.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	433	1.5	62.1
12	Parkinson's disease (G20-G21)	211	0.7	30.2
13	Pneumonitis due to solids and liquids (J69)	210	0.7	30.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	150	0.5	21.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	132	0.4	18.9
...	All other causes (Residual)	4,894	16.4	701.6
Black, both sexes, 80-84 years				
...	All causes	30,084	100.0	6,479.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,888	26.2	1,698.8
2	Malignant neoplasms (C00-C97)	6,528	21.7	1,405.9
3	Cerebrovascular diseases (I60-I69)	2,165	7.2	466.3
4	Diabetes mellitus (E10-E14)	1,402	4.7	301.9
5	Chronic lower respiratory diseases (J40-J47)	1,259	4.2	271.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,099	3.7	236.7
7	Alzheimer's disease (G30)	1,094	3.6	235.6
8	Septicemia (A40-A41)	703	2.3	151.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	660	2.2	142.1
10	Influenza and pneumonia (J09-J18)	647	2.2	139.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	366	1.2	78.8
12	Parkinson's disease (G20-G21)	231	0.8	49.7
13	Pneumonitis due to solids and liquids (J69)	229	0.8	49.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	159	0.5	34.2
15	Aortic aneurysm and dissection (I71)	102	0.3	22.0
...	All other causes (Residual)	5,552	18.5	1,195.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85-89 years				
...	All causes	27,435	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,473	27.2	---
2	Malignant neoplasms (C00-C97)	4,410	16.1	---
3	Cerebrovascular diseases (I60-I69)	1,902	6.9	---
4	Alzheimer's disease (G30)	1,444	5.3	---
5	Diabetes mellitus (E10-E14)	1,157	4.2	---
6	Chronic lower respiratory diseases (J40-J47)	1,082	3.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	918	3.3	---
8	Influenza and pneumonia (J09-J18)	756	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	636	2.3	---
10	Septicemia (A40-A41)	625	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	352	1.3	---
12	Pneumonitis due to solids and liquids (J69)	250	0.9	---
13	Parkinson's disease (G20-G21)	233	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	131	0.5	---
15	Aortic aneurysm and dissection (I71)	85	0.3	---
...	All other causes (Residual)	5,981	21.8	---
Black, both sexes, 90-94 years				
...	All causes	18,220	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,263	28.9	---
2	Malignant neoplasms (C00-C97)	2,215	12.2	---
3	Cerebrovascular diseases (I60-I69)	1,312	7.2	---
4	Alzheimer's disease (G30)	1,219	6.7	---
5	Diabetes mellitus (E10-E14)	601	3.3	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	574	3.2	---
7	Chronic lower respiratory diseases (J40-J47)	561	3.1	---
8	Influenza and pneumonia (J09-J18)	503	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	477	2.6	---
10	Septicemia (A40-A41)	369	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	231	1.3	---
12	Pneumonitis due to solids and liquids (J69)	166	0.9	---
13	Parkinson's disease (G20-G21)	99	0.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	83	0.5	---
15	Atherosclerosis (I70)	77	0.4	---
...	All other causes (Residual)	4,470	24.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 95-99 years				
...	All causes	7,440	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,258	30.3	---
2	Malignant neoplasms (C00-C97)	596	8.0	---
3	Alzheimer's disease (G30)	581	7.8	---
4	Cerebrovascular diseases (I60-I69)	500	6.7	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	232	3.1	---
6	Influenza and pneumonia (J09-J18)	219	2.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	213	2.9	---
8	Diabetes mellitus (E10-E14)	212	2.8	---
9	Chronic lower respiratory diseases (J40-J47)	171	2.3	---
10	Septicemia (A40-A41)	143	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	1.2	---
12	Pneumonitis due to solids and liquids (J69)	64	0.9	---
13	Anemias (D50-D64)	28	0.4	---
14	Parkinson's disease (G20-G21)	27	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.3	---
...	All other causes (Residual)	2,080	28.0	---
Black, both sexes, 100 years and over				
...	All causes	2,499	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	753	30.1	---
2	Alzheimer's disease (G30)	201	8.0	---
3	Cerebrovascular diseases (I60-I69)	182	7.3	---
4	Malignant neoplasms (C00-C97)	122	4.9	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	3.6	---
6	Influenza and pneumonia (J09-J18)	77	3.1	---
7	Septicemia (A40-A41)	63	2.5	---
7	Diabetes mellitus (E10-E14)	63	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	2.1	---
10	Chronic lower respiratory diseases (J40-J47)	41	1.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	1.2	---
12	Pneumonitis due to solids and liquids (J69)	20	0.8	---
13	Atherosclerosis (I70)	19	0.8	---
14	Anemias (D50-D64)	11	0.4	---
15	Nutritional deficiencies (E40-E64)	10	0.4	---
...	All other causes (Residual)	765	30.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, all ages				
...	All causes	154,767	100.0	739.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,096	24.0	177.2
2	Malignant neoplasms (C00-C97)	34,671	22.4	165.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,017	5.8	43.1
4	Cerebrovascular diseases (I60-I69)	7,338	4.7	35.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,937	4.5	33.1
6	Diabetes mellitus (E10-E14)	6,378	4.1	30.5
7	Chronic lower respiratory diseases (J40-J47)	5,073	3.3	24.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,976	2.6	19.0
9	Septicemia (A40-A41)	2,882	1.9	13.8
10	Influenza and pneumonia (J09-J18)	2,696	1.7	12.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,496	1.6	11.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,479	1.6	11.8
13	Certain conditions originating in the perinatal period (P00-P96)	2,215	1.4	10.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,955	1.3	9.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,891	1.2	9.0
...	All other causes (Residual)	27,667	17.9	132.2
Black, male, 1-4 years				
...	All causes	561	100.0	40.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	31.6	12.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	12.5	5.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	8.9	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	36	6.4	2.6
5	Malignant neoplasms (C00-C97)	33	5.9	2.4
6	Chronic lower respiratory diseases (J40-J47)	15	2.7	*
7	Septicemia (A40-A41)	13	2.3	*
7	Influenza and pneumonia (J09-J18)	13	2.3	*
9	Cerebrovascular diseases (I60-I69)	12	2.1	*
10	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
11	Anemias (D50-D64)	4	0.7	*
11	Meningitis (G00,G03)	4	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	122	21.7	8.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 5-9 years				
...	All causes	321	100.0	18.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	108	33.6	6.2
2	Malignant neoplasms (C00-C97)	32	10.0	1.9
3	Chronic lower respiratory diseases (J40-J47)	29	9.0	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	6.9	1.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	6.2	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	3.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	2.8	*
8	Influenza and pneumonia (J09-J18)	8	2.5	*
9	Septicemia (A40-A41)	4	1.2	*
9	Anemias (D50-D64)	4	1.2	*
11	Cerebrovascular diseases (I60-I69)	3	0.9	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.9	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
...	All other causes (Residual)	62	19.3	3.6
Black, male, 10-14 years				
...	All causes	354	100.0	20.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	30.2	6.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	9.9	2.0
3	Malignant neoplasms (C00-C97)	34	9.6	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	9.6	2.0
5	Chronic lower respiratory diseases (J40-J47)	26	7.3	1.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	5.9	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	5.1	*
8	Influenza and pneumonia (J09-J18)	10	2.8	*
9	Cerebrovascular diseases (I60-I69)	7	2.0	*
10	Septicemia (A40-A41)	4	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
12	Anemias (D50-D64)	1	0.3	*
12	Diabetes mellitus (E10-E14)	1	0.3	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	51	14.4	3.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-19 years				
...	All causes	1,685	100.0	94.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	805	47.8	45.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	22.0	20.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	125	7.4	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	3.4	3.2
5	Malignant neoplasms (C00-C97)	57	3.4	3.2
6	Chronic lower respiratory diseases (J40-J47)	22	1.3	1.2
7	Diabetes mellitus (E10-E14)	16	0.9	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	0.9	*
9	Anemias (D50-D64)	13	0.8	*
9	Legal intervention (Y35,Y89.0)	13	0.8	*
11	Cerebrovascular diseases (I60-I69)	10	0.6	*
12	Influenza and pneumonia (J09-J18)	7	0.4	*
13	Septicemia (A40-A41)	4	0.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	*
15	Meningitis (G00,G03)	3	0.2	*
15	Diseases of appendix (K35-K38)	3	0.2	*
...	All other causes (Residual)	158	9.4	8.8
Black, male, 20-24 years				
...	All causes	3,356	100.0	173.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,674	49.9	86.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	652	19.4	33.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	304	9.1	15.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	136	4.1	7.0
5	Malignant neoplasms (C00-C97)	72	2.1	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	59	1.8	3.1
7	Chronic lower respiratory diseases (J40-J47)	33	1.0	1.7
8	Diabetes mellitus (E10-E14)	32	1.0	1.7
9	Anemias (D50-D64)	31	0.9	1.6
9	Legal intervention (Y35,Y89.0)	31	0.9	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.6	*
12	Septicemia (A40-A41)	15	0.4	*
13	Cerebrovascular diseases (I60-I69)	11	0.3	*
14	Influenza and pneumonia (J09-J18)	9	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
...	All other causes (Residual)	271	8.1	14.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 25-29 years				
...	All causes	3,211	100.0	205.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,229	38.3	78.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	666	20.7	42.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	250	7.8	16.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	225	7.0	14.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	133	4.1	8.5
6	Malignant neoplasms (C00-C97)	131	4.1	8.4
7	Diabetes mellitus (E10-E14)	54	1.7	3.5
8	Anemias (D50-D64)	33	1.0	2.1
9	Chronic lower respiratory diseases (J40-J47)	28	0.9	1.8
10	Cerebrovascular diseases (I60-I69)	25	0.8	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.6	*
12	Septicemia (A40-A41)	17	0.5	*
12	Legal intervention (Y35,Y89.0)	17	0.5	*
14	Influenza and pneumonia (J09-J18)	15	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.5	*
...	All other causes (Residual)	355	11.1	22.7
Black, male, 30-34 years				
...	All causes	3,387	100.0	232.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	980	28.9	67.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	669	19.8	46.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	406	12.0	27.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	216	6.4	14.8
5	Malignant neoplasms (C00-C97)	178	5.3	12.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	140	4.1	9.6
7	Diabetes mellitus (E10-E14)	67	2.0	4.6
8	Cerebrovascular diseases (I60-I69)	49	1.4	3.4
9	Influenza and pneumonia (J09-J18)	40	1.2	2.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.0	2.3
11	Anemias (D50-D64)	26	0.8	1.8
11	Legal intervention (Y35,Y89.0)	26	0.8	1.8
13	Septicemia (A40-A41)	25	0.7	1.7
13	Chronic lower respiratory diseases (J40-J47)	25	0.7	1.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.6	1.5
...	All other causes (Residual)	484	14.3	33.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-39 years				
...	All causes	3,523	100.0	272.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	667	18.9	51.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	651	18.5	50.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	596	16.9	46.1
4	Malignant neoplasms (C00-C97)	255	7.2	19.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	185	5.3	14.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	163	4.6	12.6
7	Diabetes mellitus (E10-E14)	110	3.1	8.5
8	Cerebrovascular diseases (I60-I69)	100	2.8	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.6	4.3
10	Chronic lower respiratory diseases (J40-J47)	52	1.5	4.0
11	Septicemia (A40-A41)	40	1.1	3.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	1.0	2.8
13	Anemias (D50-D64)	29	0.8	2.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	29	0.8	2.2
15	Aortic aneurysm and dissection (I71)	27	0.8	2.1
...	All other causes (Residual)	528	15.0	40.8
Black, male, 40-44 years				
...	All causes	4,691	100.0	351.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,146	24.4	85.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	650	13.9	48.7
3	Malignant neoplasms (C00-C97)	462	9.8	34.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	412	8.8	30.9
5	Human immunodeficiency virus (HIV) disease (B20-B24)	227	4.8	17.0
6	Diabetes mellitus (E10-E14)	211	4.5	15.8
7	Cerebrovascular diseases (I60-I69)	169	3.6	12.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	163	3.5	12.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	1.8	6.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	74	1.6	5.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	73	1.6	5.5
12	Septicemia (A40-A41)	72	1.5	5.4
13	Chronic lower respiratory diseases (J40-J47)	67	1.4	5.0
14	Influenza and pneumonia (J09-J18)	59	1.3	4.4
15	Aortic aneurysm and dissection (I71)	50	1.1	3.7
...	All other causes (Residual)	770	16.4	57.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-49 years				
...	All causes	7,142	100.0	532.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,838	25.7	137.0
2	Malignant neoplasms (C00-C97)	1,123	15.7	83.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	709	9.9	52.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	363	5.1	27.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	341	4.8	25.4
6	Cerebrovascular diseases (I60-I69)	313	4.4	23.3
7	Diabetes mellitus (E10-E14)	309	4.3	23.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	150	2.1	11.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	2.0	10.7
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	143	2.0	10.7
11	Septicemia (A40-A41)	139	1.9	10.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	135	1.9	10.1
13	Chronic lower respiratory diseases (J40-J47)	106	1.5	7.9
14	Influenza and pneumonia (J09-J18)	100	1.4	7.5
15	Aortic aneurysm and dissection (I71)	58	0.8	4.3
...	All other causes (Residual)	1,172	16.4	87.4
Black, male, 50-54 years				
...	All causes	11,139	100.0	824.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,964	26.6	219.3
2	Malignant neoplasms (C00-C97)	2,442	21.9	180.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	932	8.4	68.9
4	Cerebrovascular diseases (I60-I69)	514	4.6	38.0
5	Diabetes mellitus (E10-E14)	467	4.2	34.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	424	3.8	31.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	306	2.7	22.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	265	2.4	19.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	263	2.4	19.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	207	1.9	15.3
11	Chronic lower respiratory diseases (J40-J47)	203	1.8	15.0
12	Septicemia (A40-A41)	199	1.8	14.7
13	Influenza and pneumonia (J09-J18)	167	1.5	12.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	141	1.3	10.4
15	Viral hepatitis (B15-B19)	99	0.9	7.3
...	All other causes (Residual)	1,546	13.9	114.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-59 years				
...	All causes	15,988	100.0	1,347.6
1	Malignant neoplasms (C00-C97)	4,358	27.3	367.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,239	26.5	357.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	948	5.9	79.9
4	Diabetes mellitus (E10-E14)	731	4.6	61.6
5	Cerebrovascular diseases (I60-I69)	716	4.5	60.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	422	2.6	35.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	407	2.5	34.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	367	2.3	30.9
9	Chronic lower respiratory diseases (J40-J47)	363	2.3	30.6
10	Septicemia (A40-A41)	269	1.7	22.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	262	1.6	22.1
12	Influenza and pneumonia (J09-J18)	228	1.4	19.2
13	Viral hepatitis (B15-B19)	208	1.3	17.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	190	1.2	16.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	112	0.7	9.4
...	All other causes (Residual)	2,168	13.6	182.7
Black, male, 60-64 years				
...	All causes	18,078	100.0	1,995.1
1	Malignant neoplasms (C00-C97)	5,352	29.6	590.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,758	26.3	525.1
3	Cerebrovascular diseases (I60-I69)	918	5.1	101.3
4	Diabetes mellitus (E10-E14)	884	4.9	97.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	754	4.2	83.2
6	Chronic lower respiratory diseases (J40-J47)	551	3.0	60.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	507	2.8	56.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	457	2.5	50.4
9	Septicemia (A40-A41)	334	1.8	36.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	282	1.6	31.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	260	1.4	28.7
12	Influenza and pneumonia (J09-J18)	257	1.4	28.4
13	Viral hepatitis (B15-B19)	229	1.3	25.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	0.5	9.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	81	0.4	8.9
...	All other causes (Residual)	2,366	13.1	261.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 65-69 years				
...	All causes	16,278	100.0	2,608.4
1	Malignant neoplasms (C00-C97)	5,090	31.3	815.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,341	26.7	695.6
3	Cerebrovascular diseases (I60-I69)	813	5.0	130.3
4	Diabetes mellitus (E10-E14)	801	4.9	128.4
5	Chronic lower respiratory diseases (J40-J47)	627	3.9	100.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	441	2.7	70.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	422	2.6	67.6
8	Septicemia (A40-A41)	348	2.1	55.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	305	1.9	48.9
10	Influenza and pneumonia (J09-J18)	272	1.7	43.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	240	1.5	38.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	141	0.9	22.6
13	Viral hepatitis (B15-B19)	118	0.7	18.9
14	Pneumonitis due to solids and liquids (J69)	87	0.5	13.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	79	0.5	12.7
...	All other causes (Residual)	2,153	13.2	345.0
Black, male, 70-74 years				
...	All causes	15,542	100.0	3,751.2
1	Malignant neoplasms (C00-C97)	4,686	30.2	1,131.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,044	26.0	976.0
3	Cerebrovascular diseases (I60-I69)	900	5.8	217.2
4	Diabetes mellitus (E10-E14)	787	5.1	189.9
5	Chronic lower respiratory diseases (J40-J47)	783	5.0	189.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	470	3.0	113.4
7	Septicemia (A40-A41)	326	2.1	78.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	307	2.0	74.1
9	Influenza and pneumonia (J09-J18)	306	2.0	73.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	278	1.8	67.1
11	Alzheimer's disease (G30)	132	0.8	31.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	123	0.8	29.7
13	Pneumonitis due to solids and liquids (J69)	101	0.6	24.4
14	Parkinson's disease (G20-G21)	80	0.5	19.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	68	0.4	16.4
...	All other causes (Residual)	2,151	13.8	519.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-79 years				
...	All causes	14,844	100.0	5,409.0
1	Malignant neoplasms (C00-C97)	4,002	27.0	1,458.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,867	26.1	1,409.1
3	Cerebrovascular diseases (I60-I69)	890	6.0	324.3
4	Diabetes mellitus (E10-E14)	743	5.0	270.7
5	Chronic lower respiratory diseases (J40-J47)	709	4.8	258.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	486	3.3	177.1
7	Septicemia (A40-A41)	330	2.2	120.2
8	Influenza and pneumonia (J09-J18)	296	2.0	107.9
9	Alzheimer's disease (G30)	271	1.8	98.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	265	1.8	96.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	261	1.8	95.1
12	Parkinson's disease (G20-G21)	140	0.9	51.0
13	Pneumonitis due to solids and liquids (J69)	131	0.9	47.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.5	28.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	67	0.5	24.4
...	All other causes (Residual)	2,309	15.6	841.4
Black, male, 80-84 years				
...	All causes	13,108	100.0	7,949.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,426	26.1	2,077.7
2	Malignant neoplasms (C00-C97)	3,173	24.2	1,924.2
3	Cerebrovascular diseases (I60-I69)	812	6.2	492.4
4	Chronic lower respiratory diseases (J40-J47)	623	4.8	377.8
5	Diabetes mellitus (E10-E14)	552	4.2	334.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	463	3.5	280.8
7	Alzheimer's disease (G30)	383	2.9	232.3
8	Septicemia (A40-A41)	296	2.3	179.5
9	Influenza and pneumonia (J09-J18)	285	2.2	172.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	271	2.1	164.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	1.5	116.4
12	Parkinson's disease (G20-G21)	144	1.1	87.3
13	Pneumonitis due to solids and liquids (J69)	111	0.8	67.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.6	44.3
15	Aortic aneurysm and dissection (I71)	42	0.3	25.5
...	All other causes (Residual)	2,262	17.3	1,371.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 85-89 years				
...	All causes	10,164	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,734	26.9	---
2	Malignant neoplasms (C00-C97)	1,986	19.5	---
3	Cerebrovascular diseases (I60-I69)	606	6.0	---
4	Chronic lower respiratory diseases (J40-J47)	507	5.0	---
5	Alzheimer's disease (G30)	414	4.1	---
6	Diabetes mellitus (E10-E14)	388	3.8	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	336	3.3	---
8	Influenza and pneumonia (J09-J18)	318	3.1	---
9	Septicemia (A40-A41)	243	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	203	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	1.6	---
12	Pneumonitis due to solids and liquids (J69)	125	1.2	---
13	Parkinson's disease (G20-G21)	119	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.5	---
15	Atherosclerosis (I70)	30	0.3	---
...	All other causes (Residual)	1,941	19.1	---
Black, male, 90-94 years				
...	All causes	5,597	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,573	28.1	---
2	Malignant neoplasms (C00-C97)	944	16.9	---
3	Cerebrovascular diseases (I60-I69)	325	5.8	---
4	Alzheimer's disease (G30)	262	4.7	---
5	Chronic lower respiratory diseases (J40-J47)	235	4.2	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	209	3.7	---
7	Diabetes mellitus (E10-E14)	169	3.0	---
8	Influenza and pneumonia (J09-J18)	168	3.0	---
9	Septicemia (A40-A41)	129	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	1.4	---
12	Pneumonitis due to solids and liquids (J69)	67	1.2	---
13	Parkinson's disease (G20-G21)	40	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.6	---
15	Anemias (D50-D64)	19	0.3	---
15	Nutritional deficiencies (E40-E64)	19	0.3	---
...	All other causes (Residual)	1,205	21.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 95-99 years				
...	All causes	1,579	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	465	29.4	---
2	Malignant neoplasms (C00-C97)	220	13.9	---
3	Cerebrovascular diseases (I60-I69)	100	6.3	---
4	Alzheimer's disease (G30)	81	5.1	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	3.9	---
6	Influenza and pneumonia (J09-J18)	56	3.5	---
7	Chronic lower respiratory diseases (J40-J47)	49	3.1	---
8	Diabetes mellitus (E10-E14)	47	3.0	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	2.2	---
10	Septicemia (A40-A41)	33	2.1	---
11	Pneumonitis due to solids and liquids (J69)	22	1.4	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	1.3	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.6	---
14	Parkinson's disease (G20-G21)	9	0.6	---
15	Anemias (D50-D64)	4	0.3	---
15	Atherosclerosis (I70)	4	0.3	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.3	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.3	---
...	All other causes (Residual)	353	22.4	---
Black, male, 100 years and over				
...	All causes	324	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	25.9	---
2	Malignant neoplasms (C00-C97)	39	12.0	---
3	Cerebrovascular diseases (I60-I69)	22	6.8	---
4	Influenza and pneumonia (J09-J18)	19	5.9	---
5	Alzheimer's disease (G30)	15	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	14	4.3	---
7	Septicemia (A40-A41)	11	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	3.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	2.5	---
10	Diabetes mellitus (E10-E14)	6	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	1.9	---
12	Atherosclerosis (I70)	3	0.9	---
13	Anemias (D50-D64)	2	0.6	---
13	Pneumonitis due to solids and liquids (J69)	2	0.6	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.6	---
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.6	---
...	All other causes (Residual)	79	24.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	148,202	100.0	651.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,914	23.6	153.4
2	Malignant neoplasms (C00-C97)	33,282	22.5	146.2
3	Cerebrovascular diseases (I60-I69)	8,931	6.0	39.2
4	Diabetes mellitus (E10-E14)	7,007	4.7	30.8
5	Chronic lower respiratory diseases (J40-J47)	4,845	3.3	21.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,417	3.0	19.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,396	3.0	19.3
8	Alzheimer's disease (G30)	4,050	2.7	17.8
9	Septicemia (A40-A41)	3,368	2.3	14.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,959	2.0	13.0
11	Influenza and pneumonia (J09-J18)	2,871	1.9	12.6
12	Certain conditions originating in the perinatal period (P00-P96)	1,840	1.2	8.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,263	0.9	5.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,122	0.8	4.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,098	0.7	4.8
...	All other causes (Residual)	31,839	21.5	139.9
Black, female, 1-4 years				
...	All causes	445	100.0	33.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	27.4	9.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	14.8	5.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	9.2	3.1
4	Malignant neoplasms (C00-C97)	31	7.0	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	5.2	1.7
6	Chronic lower respiratory diseases (J40-J47)	12	2.7	*
7	Influenza and pneumonia (J09-J18)	10	2.2	*
8	Septicemia (A40-A41)	7	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.6	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.6	*
11	Anemias (D50-D64)	4	0.9	*
12	Cerebrovascular diseases (I60-I69)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Meningitis (G00,G03)	2	0.4	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	103	23.1	7.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-9 years				
...	All causes	261	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75	28.7	4.5
2	Malignant neoplasms (C00-C97)	33	12.6	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	8.4	1.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	6.9	*
5	Chronic lower respiratory diseases (J40-J47)	16	6.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	4.6	*
7	Influenza and pneumonia (J09-J18)	5	1.9	*
8	Cerebrovascular diseases (I60-I69)	4	1.5	*
9	Septicemia (A40-A41)	3	1.1	*
9	Anemias (D50-D64)	3	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.8	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.8	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.8	*
15	Meningococcal infection (A39)	1	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	60	23.0	3.6
Black, female, 10-14 years				
...	All causes	233	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	17.2	2.4
2	Malignant neoplasms (C00-C97)	23	9.9	1.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	9.4	1.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	8.6	1.2
5	Chronic lower respiratory diseases (J40-J47)	16	6.9	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	6.0	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.4	*
8	Cerebrovascular diseases (I60-I69)	5	2.1	*
8	Influenza and pneumonia (J09-J18)	5	2.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
10	Anemias (D50-D64)	3	1.3	*
10	Diabetes mellitus (E10-E14)	3	1.3	*
13	Septicemia (A40-A41)	2	0.9	*
13	Meningitis (G00,G03)	2	0.9	*
13	Pneumonitis due to solids and liquids (J69)	2	0.9	*
...	All other causes (Residual)	65	27.9	3.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 15-19 years				
...	All causes	495	100.0	28.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	27.5	7.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	16.0	4.6
3	Malignant neoplasms (C00-C97)	46	9.3	2.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	7.7	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	6.1	1.7
6	Chronic lower respiratory diseases (J40-J47)	12	2.4	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	11	2.2	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	2.2	*
9	Anemias (D50-D64)	9	1.8	*
10	Cerebrovascular diseases (I60-I69)	8	1.6	*
11	Influenza and pneumonia (J09-J18)	6	1.2	*
12	Septicemia (A40-A41)	5	1.0	*
13	Diabetes mellitus (E10-E14)	4	0.8	*
14	Diseases of appendix (K35-K38)	3	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
...	All other causes (Residual)	93	18.8	5.4
Black, female, 20-24 years				
...	All causes	1,002	100.0	52.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	256	25.5	13.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	173	17.3	9.1
3	Malignant neoplasms (C00-C97)	66	6.6	3.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	6.6	3.5
5	Pregnancy, childbirth and the puerperium (O00-O99)	58	5.8	3.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	5.6	2.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	24	2.4	1.3
8	Diabetes mellitus (E10-E14)	21	2.1	1.1
9	Anemias (D50-D64)	17	1.7	*
10	Chronic lower respiratory diseases (J40-J47)	13	1.3	*
11	Septicemia (A40-A41)	12	1.2	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.1	*
13	Influenza and pneumonia (J09-J18)	10	1.0	*
14	Cerebrovascular diseases (I60-I69)	9	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.6	*
...	All other causes (Residual)	204	20.4	10.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-29 years				
...	All causes	1,256	100.0	76.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	237	18.9	14.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	145	11.5	8.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	9.7	7.5
4	Malignant neoplasms (C00-C97)	104	8.3	6.4
5	Pregnancy, childbirth and the puerperium (O00-O99)	87	6.9	5.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	4.2	3.2
7	Diabetes mellitus (E10-E14)	47	3.7	2.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	46	3.7	2.8
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	1.7	1.3
10	Septicemia (A40-A41)	20	1.6	1.2
10	Anemias (D50-D64)	20	1.6	1.2
12	Influenza and pneumonia (J09-J18)	19	1.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.5	*
14	Chronic lower respiratory diseases (J40-J47)	18	1.4	*
15	Cerebrovascular diseases (I60-I69)	17	1.4	*
...	All other causes (Residual)	281	22.4	17.2
Black, female, 30-34 years				
...	All causes	1,677	100.0	105.4
1	Malignant neoplasms (C00-C97)	232	13.8	14.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	229	13.7	14.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	221	13.2	13.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	7.7	8.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	78	4.7	4.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	63	3.8	4.0
7	Diabetes mellitus (E10-E14)	58	3.5	3.6
8	Cerebrovascular diseases (I60-I69)	54	3.2	3.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	2.7	2.8
10	Anemias (D50-D64)	35	2.1	2.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	1.9	2.0
12	Septicemia (A40-A41)	28	1.7	1.8
12	Chronic lower respiratory diseases (J40-J47)	28	1.7	1.8
14	Influenza and pneumonia (J09-J18)	23	1.4	1.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.0	*
...	All other causes (Residual)	406	24.2	25.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 35-39 years				
...	All causes	2,179	100.0	150.5
1	Malignant neoplasms (C00-C97)	384	17.6	26.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	357	16.4	24.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	245	11.2	16.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	129	5.9	8.9
5	Cerebrovascular diseases (I60-I69)	87	4.0	6.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	86	3.9	5.9
7	Diabetes mellitus (E10-E14)	79	3.6	5.5
8	Pregnancy, childbirth and the puerperium (O00-O99)	52	2.4	3.6
9	Chronic lower respiratory diseases (J40-J47)	44	2.0	3.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.9	2.9
11	Septicemia (A40-A41)	41	1.9	2.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	35	1.6	2.4
13	Influenza and pneumonia (J09-J18)	32	1.5	2.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	1.3	1.9
15	Anemias (D50-D64)	22	1.0	1.5
...	All other causes (Residual)	516	23.7	35.6
Black, female, 40-44 years				
...	All causes	3,366	100.0	224.1
1	Malignant neoplasms (C00-C97)	816	24.2	54.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	629	18.7	41.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	284	8.4	18.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	163	4.8	10.9
5	Cerebrovascular diseases (I60-I69)	140	4.2	9.3
6	Diabetes mellitus (E10-E14)	134	4.0	8.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	85	2.5	5.7
8	Septicemia (A40-A41)	75	2.2	5.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	1.8	4.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	60	1.8	4.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	1.6	3.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	47	1.4	3.1
13	Influenza and pneumonia (J09-J18)	46	1.4	3.1
14	Chronic lower respiratory diseases (J40-J47)	45	1.3	3.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	35	1.0	2.3
...	All other causes (Residual)	690	20.5	45.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-49 years				
...	All causes	5,317	100.0	352.7
1	Malignant neoplasms (C00-C97)	1,432	26.9	95.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,080	20.3	71.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	373	7.0	24.7
4	Diabetes mellitus (E10-E14)	234	4.4	15.5
5	Cerebrovascular diseases (I60-I69)	230	4.3	15.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	192	3.6	12.7
7	Septicemia (A40-A41)	114	2.1	7.6
7	Chronic lower respiratory diseases (J40-J47)	114	2.1	7.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	2.0	7.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	99	1.9	6.6
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	92	1.7	6.1
12	Influenza and pneumonia (J09-J18)	64	1.2	4.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	63	1.2	4.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	1.0	3.5
15	Anemias (D50-D64)	30	0.6	2.0
...	All other causes (Residual)	1,042	19.6	69.1
Black, female, 50-54 years				
...	All causes	8,513	100.0	554.3
1	Malignant neoplasms (C00-C97)	2,669	31.4	173.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,786	21.0	116.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	415	4.9	27.0
4	Cerebrovascular diseases (I60-I69)	413	4.9	26.9
5	Diabetes mellitus (E10-E14)	405	4.8	26.4
6	Chronic lower respiratory diseases (J40-J47)	236	2.8	15.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	213	2.5	13.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	191	2.2	12.4
9	Septicemia (A40-A41)	189	2.2	12.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	167	2.0	10.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	145	1.7	9.4
12	Influenza and pneumonia (J09-J18)	119	1.4	7.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	0.8	4.6
14	Viral hepatitis (B15-B19)	51	0.6	3.3
15	Anemias (D50-D64)	37	0.4	2.4
...	All other causes (Residual)	1,406	16.5	91.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 55-59 years				
...	All causes	11,225	100.0	810.9
1	Malignant neoplasms (C00-C97)	3,626	32.3	262.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,477	22.1	179.0
3	Cerebrovascular diseases (I60-I69)	561	5.0	40.5
4	Diabetes mellitus (E10-E14)	535	4.8	38.7
5	Chronic lower respiratory diseases (J40-J47)	372	3.3	26.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	349	3.1	25.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	320	2.9	23.1
8	Septicemia (A40-A41)	255	2.3	18.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	201	1.8	14.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	191	1.7	13.8
11	Influenza and pneumonia (J09-J18)	190	1.7	13.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	180	1.6	13.0
13	Viral hepatitis (B15-B19)	111	1.0	8.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.5	3.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	0.4	3.0
...	All other causes (Residual)	1,764	15.7	127.4
Black, female, 60-64 years				
...	All causes	12,818	100.0	1,150.1
1	Malignant neoplasms (C00-C97)	4,369	34.1	392.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,846	22.2	255.4
3	Diabetes mellitus (E10-E14)	743	5.8	66.7
4	Cerebrovascular diseases (I60-I69)	631	4.9	56.6
5	Chronic lower respiratory diseases (J40-J47)	514	4.0	46.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	345	2.7	31.0
7	Septicemia (A40-A41)	318	2.5	28.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	294	2.3	26.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	206	1.6	18.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	196	1.5	17.6
11	Influenza and pneumonia (J09-J18)	190	1.5	17.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	102	0.8	9.2
13	Viral hepatitis (B15-B19)	96	0.7	8.6
14	Pneumonitis due to solids and liquids (J69)	49	0.4	4.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.3	3.9
...	All other causes (Residual)	1,876	14.6	168.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-69 years				
...	All causes	12,710	100.0	1,584.1
1	Malignant neoplasms (C00-C97)	4,171	32.8	519.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,966	23.3	369.7
3	Diabetes mellitus (E10-E14)	748	5.9	93.2
4	Cerebrovascular diseases (I60-I69)	706	5.6	88.0
5	Chronic lower respiratory diseases (J40-J47)	522	4.1	65.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	430	3.4	53.6
7	Septicemia (A40-A41)	342	2.7	42.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	224	1.8	27.9
9	Influenza and pneumonia (J09-J18)	221	1.7	27.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	1.5	23.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	96	0.8	12.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	61	0.5	7.6
13	Alzheimer's disease (G30)	52	0.4	6.5
14	Viral hepatitis (B15-B19)	51	0.4	6.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.4	6.4
...	All other causes (Residual)	1,879	14.8	234.2
Black, female, 70-74 years				
...	All causes	13,600	100.0	2,362.4
1	Malignant neoplasms (C00-C97)	4,006	29.5	695.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,246	23.9	563.9
3	Diabetes mellitus (E10-E14)	839	6.2	145.7
4	Cerebrovascular diseases (I60-I69)	797	5.9	138.4
5	Chronic lower respiratory diseases (J40-J47)	587	4.3	102.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	553	4.1	96.1
7	Septicemia (A40-A41)	327	2.4	56.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	286	2.1	49.7
9	Influenza and pneumonia (J09-J18)	250	1.8	43.4
10	Alzheimer's disease (G30)	181	1.3	31.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	1.2	28.8
12	Pneumonitis due to solids and liquids (J69)	68	0.5	11.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	63	0.5	10.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.4	9.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.3	6.6
...	All other causes (Residual)	2,139	15.7	371.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 75-79 years				
...	All causes	14,945	100.0	3,532.3
1	Malignant neoplasms (C00-C97)	3,757	25.1	888.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,646	24.4	861.7
3	Cerebrovascular diseases (I60-I69)	1,053	7.0	248.9
4	Diabetes mellitus (E10-E14)	882	5.9	208.5
5	Chronic lower respiratory diseases (J40-J47)	605	4.0	143.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	547	3.7	129.3
7	Septicemia (A40-A41)	415	2.8	98.1
8	Alzheimer's disease (G30)	397	2.7	93.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	303	2.0	71.6
10	Influenza and pneumonia (J09-J18)	295	2.0	69.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	1.2	40.7
12	Pneumonitis due to solids and liquids (J69)	79	0.5	18.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.5	17.3
14	Parkinson's disease (G20-G21)	71	0.5	16.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	65	0.4	15.4
...	All other causes (Residual)	2,585	17.3	611.0
Black, female, 80-84 years				
...	All causes	16,976	100.0	5,669.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,462	26.3	1,490.2
2	Malignant neoplasms (C00-C97)	3,355	19.8	1,120.5
3	Cerebrovascular diseases (I60-I69)	1,353	8.0	451.9
4	Diabetes mellitus (E10-E14)	850	5.0	283.9
5	Alzheimer's disease (G30)	711	4.2	237.5
6	Chronic lower respiratory diseases (J40-J47)	636	3.7	212.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	636	3.7	212.4
8	Septicemia (A40-A41)	407	2.4	135.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	389	2.3	129.9
10	Influenza and pneumonia (J09-J18)	362	2.1	120.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	1.0	58.1
12	Pneumonitis due to solids and liquids (J69)	118	0.7	39.4
13	Parkinson's disease (G20-G21)	87	0.5	29.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	86	0.5	28.7
15	Aortic aneurysm and dissection (I71)	60	0.4	20.0
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	60	0.4	20.0
...	All other causes (Residual)	3,230	19.0	1,078.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85-89 years				
...	All causes	17,271	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,739	27.4	---
2	Malignant neoplasms (C00-C97)	2,424	14.0	---
3	Cerebrovascular diseases (I60-I69)	1,296	7.5	---
4	Alzheimer's disease (G30)	1,030	6.0	---
5	Diabetes mellitus (E10-E14)	769	4.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	582	3.4	---
7	Chronic lower respiratory diseases (J40-J47)	575	3.3	---
8	Influenza and pneumonia (J09-J18)	438	2.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	433	2.5	---
10	Septicemia (A40-A41)	382	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	1.1	---
12	Pneumonitis due to solids and liquids (J69)	125	0.7	---
13	Parkinson's disease (G20-G21)	114	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.5	---
15	Aortic aneurysm and dissection (I71)	58	0.3	---
...	All other causes (Residual)	4,037	23.4	---
Black, female, 90-94 years				
...	All causes	12,623	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,690	29.2	---
2	Malignant neoplasms (C00-C97)	1,271	10.1	---
3	Cerebrovascular diseases (I60-I69)	987	7.8	---
4	Alzheimer's disease (G30)	957	7.6	---
5	Diabetes mellitus (E10-E14)	432	3.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	365	2.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	356	2.8	---
8	Influenza and pneumonia (J09-J18)	335	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	326	2.6	---
10	Septicemia (A40-A41)	240	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	1.2	---
12	Pneumonitis due to solids and liquids (J69)	99	0.8	---
13	Atherosclerosis (I70)	60	0.5	---
14	Parkinson's disease (G20-G21)	59	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.4	---
...	All other causes (Residual)	3,244	25.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 95-99 years				
...	All causes	5,861	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,793	30.6	---
2	Alzheimer's disease (G30)	500	8.5	---
3	Cerebrovascular diseases (I60-I69)	400	6.8	---
4	Malignant neoplasms (C00-C97)	376	6.4	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	197	3.4	---
6	Diabetes mellitus (E10-E14)	165	2.8	---
7	Influenza and pneumonia (J09-J18)	163	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	151	2.6	---
9	Chronic lower respiratory diseases (J40-J47)	122	2.1	---
10	Septicemia (A40-A41)	110	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	1.2	---
12	Pneumonitis due to solids and liquids (J69)	42	0.7	---
13	Anemias (D50-D64)	24	0.4	---
14	Nutritional deficiencies (E40-E64)	20	0.3	---
15	Parkinson's disease (G20-G21)	18	0.3	---
15	Atherosclerosis (I70)	18	0.3	---
...	All other causes (Residual)	1,693	28.9	---
Black, female, 100 years and over				
...	All causes	2,175	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	669	30.8	---
2	Alzheimer's disease (G30)	186	8.6	---
3	Cerebrovascular diseases (I60-I69)	160	7.4	---
4	Malignant neoplasms (C00-C97)	83	3.8	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	3.7	---
6	Influenza and pneumonia (J09-J18)	58	2.7	---
7	Diabetes mellitus (E10-E14)	57	2.6	---
8	Septicemia (A40-A41)	52	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	2.0	---
10	Chronic lower respiratory diseases (J40-J47)	27	1.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	1.1	---
12	Pneumonitis due to solids and liquids (J69)	18	0.8	---
13	Atherosclerosis (I70)	16	0.7	---
14	Nutritional deficiencies (E40-E64)	10	0.5	---
15	Anemias (D50-D64)	9	0.4	---
...	All other causes (Residual)	682	31.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, all ages				
...	All causes	17,052	100.0	382.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,139	18.4	70.4
2	Malignant neoplasms (C00-C97)	3,109	18.2	69.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,833	10.7	41.1
4	Diabetes mellitus (E10-E14)	959	5.6	21.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	944	5.5	21.2
6	Chronic lower respiratory diseases (J40-J47)	757	4.4	17.0
7	Cerebrovascular diseases (I60-I69)	595	3.5	13.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	521	3.1	11.7
9	Influenza and pneumonia (J09-J18)	375	2.2	8.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	302	1.8	6.8
11	Septicemia (A40-A41)	288	1.7	6.5
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	241	1.4	5.4
13	Alzheimer's disease (G30)	238	1.4	5.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	183	1.1	4.1
15	Viral hepatitis (B15-B19)	133	0.8	3.0
...	All other causes (Residual)	3,435	20.1	77.1
American Indian, both sexes, 1-4 years				
...	All causes	104	100.0	33.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	41.3	13.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	15.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	10.6	*
4	Malignant neoplasms (C00-C97)	3	2.9	*
4	Influenza and pneumonia (J09-J18)	3	2.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.9	*
7	Nutritional deficiencies (E40-E64)	1	1.0	*
7	Cerebrovascular diseases (I60-I69)	1	1.0	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	1.0	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.0	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.0	*
...	All other causes (Residual)	21	20.2	6.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 5-9 years				
...	All causes	50	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	44.0	5.6
2	Malignant neoplasms (C00-C97)	7	14.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	8.0	*
4	Chronic lower respiratory diseases (J40-J47)	2	4.0	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.0	*
6	Viral hepatitis (B15-B19)	1	2.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.0	*
...	All other causes (Residual)	11	22.0	*
American Indian, both sexes, 10-14 years				
...	All causes	38	100.0	10.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	34.2	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	28.9	*
3	Malignant neoplasms (C00-C97)	3	7.9	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	5.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.6	*
5	Influenza and pneumonia (J09-J18)	1	2.6	*
5	Pneumonitis due to solids and liquids (J69)	1	2.6	*
5	Certain conditions originating in the perinatal period (P00-P96)	1	2.6	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.6	*
...	All other causes (Residual)	4	10.5	*
American Indian, both sexes, 15-19 years				
...	All causes	180	100.0	48.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	41.1	19.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	57	31.7	15.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	7.8	*
4	Malignant neoplasms (C00-C97)	6	3.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	2.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	1.7	*
7	Septicemia (A40-A41)	2	1.1	*
7	Diabetes mellitus (E10-E14)	2	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
...	All other causes (Residual)	16	8.9	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 20-24 years				
...	All causes	368	100.0	93.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	46.5	43.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	93	25.3	23.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	7.1	6.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	3.0	*
5	Malignant neoplasms (C00-C97)	9	2.4	*
6	Diabetes mellitus (E10-E14)	3	0.8	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.8	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.8	*
6	Legal intervention (Y35,Y89.0)	3	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
12	Septicemia (A40-A41)	1	0.3	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
12	Influenza and pneumonia (J09-J18)	1	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
...	All other causes (Residual)	36	9.8	9.1
American Indian, both sexes, 25-29 years				
...	All causes	448	100.0	127.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	45.8	58.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	14.3	18.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	9.6	12.2
4	Malignant neoplasms (C00-C97)	19	4.2	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	16	3.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	2.2	*
7	Influenza and pneumonia (J09-J18)	7	1.6	*
8	Diabetes mellitus (E10-E14)	5	1.1	*
9	Cerebrovascular diseases (I60-I69)	3	0.7	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.7	*
12	Septicemia (A40-A41)	2	0.4	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	*
13	Legal intervention (Y35,Y89.0)	1	0.2	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	64	14.3	18.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 30-34 years				
...	All causes	475	100.0	142.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	31.2	44.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	14.3	20.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	56	11.8	16.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	6.5	9.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	6.3	9.0
6	Malignant neoplasms (C00-C97)	24	5.1	7.2
7	Diabetes mellitus (E10-E14)	13	2.7	*
8	Septicemia (A40-A41)	6	1.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.8	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.8	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
13	Cerebrovascular diseases (I60-I69)	2	0.4	*
13	Influenza and pneumonia (J09-J18)	2	0.4	*
...	All other causes (Residual)	78	16.4	23.4
American Indian, both sexes, 35-39 years				
...	All causes	518	100.0	174.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	145	28.0	48.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	67	12.9	22.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	8.9	15.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	8.3	14.4
5	Malignant neoplasms (C00-C97)	33	6.4	11.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	4.1	7.1
7	Diabetes mellitus (E10-E14)	20	3.9	6.7
8	Septicemia (A40-A41)	9	1.7	*
9	Influenza and pneumonia (J09-J18)	7	1.4	*
10	Cerebrovascular diseases (I60-I69)	5	1.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.6	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
15	Viral hepatitis (B15-B19)	2	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
15	Aortic aneurysm and dissection (I71)	2	0.4	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	99	19.1	33.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 40-44 years				
...	All causes	708	100.0	245.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	22.5	55.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	99	14.0	34.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	93	13.1	32.2
4	Malignant neoplasms (C00-C97)	60	8.5	20.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	6.4	15.6
6	Diabetes mellitus (E10-E14)	34	4.8	11.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	4.1	10.0
8	Influenza and pneumonia (J09-J18)	21	3.0	7.3
9	Cerebrovascular diseases (I60-I69)	18	2.5	*
10	Septicemia (A40-A41)	11	1.6	*
11	Viral hepatitis (B15-B19)	7	1.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.0	*
13	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.4	*
...	All other causes (Residual)	112	15.8	38.8
American Indian, both sexes, 45-49 years				
...	All causes	1,002	100.0	362.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	17.2	62.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	168	16.8	60.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	150	15.0	54.2
4	Malignant neoplasms (C00-C97)	129	12.9	46.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	5.4	19.5
6	Diabetes mellitus (E10-E14)	45	4.5	16.3
7	Septicemia (A40-A41)	20	2.0	7.2
7	Cerebrovascular diseases (I60-I69)	20	2.0	7.2
9	Chronic lower respiratory diseases (J40-J47)	18	1.8	*
10	Viral hepatitis (B15-B19)	17	1.7	*
11	Influenza and pneumonia (J09-J18)	14	1.4	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	1.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.1	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.4	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.4	*
...	All other causes (Residual)	151	15.1	54.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 50-54 years				
...	All causes	1,298	100.0	478.3
1	Malignant neoplasms (C00-C97)	230	17.7	84.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	211	16.3	77.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	14.4	68.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	166	12.8	61.2
5	Diabetes mellitus (E10-E14)	62	4.8	22.8
6	Viral hepatitis (B15-B19)	34	2.6	12.5
7	Cerebrovascular diseases (I60-I69)	30	2.3	11.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	2.2	10.7
9	Chronic lower respiratory diseases (J40-J47)	28	2.2	10.3
10	Septicemia (A40-A41)	27	2.1	9.9
11	Influenza and pneumonia (J09-J18)	22	1.7	8.1
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	1.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	1.1	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.8	*
...	All other causes (Residual)	216	16.6	79.6
American Indian, both sexes, 55-59 years				
...	All causes	1,536	100.0	668.7
1	Malignant neoplasms (C00-C97)	349	22.7	151.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	288	18.8	125.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	144	9.4	62.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	131	8.5	57.0
5	Diabetes mellitus (E10-E14)	114	7.4	49.6
6	Chronic lower respiratory diseases (J40-J47)	53	3.5	23.1
7	Cerebrovascular diseases (I60-I69)	34	2.2	14.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	2.1	13.9
9	Viral hepatitis (B15-B19)	29	1.9	12.6
10	Influenza and pneumonia (J09-J18)	28	1.8	12.2
11	Septicemia (A40-A41)	24	1.6	10.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	1.6	10.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	1.1	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.6	*
...	All other causes (Residual)	250	16.3	108.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 60-64 years				
...	All causes	1,567	100.0	907.1
1	Malignant neoplasms (C00-C97)	427	27.2	247.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	321	20.5	185.8
3	Diabetes mellitus (E10-E14)	124	7.9	71.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	92	5.9	53.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	5.3	48.0
6	Chronic lower respiratory diseases (J40-J47)	60	3.8	34.7
7	Cerebrovascular diseases (I60-I69)	52	3.3	30.1
8	Septicemia (A40-A41)	32	2.0	18.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	2.0	17.9
10	Viral hepatitis (B15-B19)	25	1.6	14.5
10	Influenza and pneumonia (J09-J18)	25	1.6	14.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	1.3	12.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.3	*
...	All other causes (Residual)	245	15.6	141.8
American Indian, both sexes, 65-69 years				
...	All causes	1,599	100.0	1,301.4
1	Malignant neoplasms (C00-C97)	432	27.0	351.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	337	21.1	274.3
3	Diabetes mellitus (E10-E14)	121	7.6	98.5
4	Chronic lower respiratory diseases (J40-J47)	117	7.3	95.2
5	Cerebrovascular diseases (I60-I69)	72	4.5	58.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	64	4.0	52.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	3.8	49.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	2.4	31.7
9	Septicemia (A40-A41)	33	2.1	26.9
10	Influenza and pneumonia (J09-J18)	30	1.9	24.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.8	*
12	Aortic aneurysm and dissection (I71)	9	0.6	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	0.6	*
14	Viral hepatitis (B15-B19)	7	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.3	*
...	All other causes (Residual)	250	15.6	203.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 70-74 years				
...	All causes	1,569	100.0	1,970.6
1	Malignant neoplasms (C00-C97)	423	27.0	531.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	342	21.8	429.5
3	Chronic lower respiratory diseases (J40-J47)	121	7.7	152.0
4	Diabetes mellitus (E10-E14)	113	7.2	141.9
5	Cerebrovascular diseases (I60-I69)	62	4.0	77.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	45	2.9	56.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	2.6	51.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	2.1	41.4
9	Influenza and pneumonia (J09-J18)	32	2.0	40.2
10	Septicemia (A40-A41)	29	1.8	36.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.0	*
12	Alzheimer's disease (G30)	15	1.0	*
13	Parkinson's disease (G20-G21)	11	0.7	*
14	Pneumonitis due to solids and liquids (J69)	9	0.6	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	9	0.6	*
...	All other causes (Residual)	268	17.1	336.6
American Indian, both sexes, 75-79 years				
...	All causes	1,572	100.0	3,064.0
1	Malignant neoplasms (C00-C97)	388	24.7	756.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	327	20.8	637.4
3	Chronic lower respiratory diseases (J40-J47)	118	7.5	230.0
4	Diabetes mellitus (E10-E14)	110	7.0	214.4
5	Cerebrovascular diseases (I60-I69)	81	5.2	157.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	3.3	101.4
7	Alzheimer's disease (G30)	35	2.2	68.2
7	Influenza and pneumonia (J09-J18)	35	2.2	68.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	2.2	68.2
10	Septicemia (A40-A41)	26	1.7	50.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	1.7	50.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.6	48.7
13	Parkinson's disease (G20-G21)	12	0.8	*
14	Pneumonitis due to solids and liquids (J69)	11	0.7	*
15	Aortic aneurysm and dissection (I71)	7	0.4	*
...	All other causes (Residual)	284	18.1	553.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 80-84 years				
...	All causes	1,442	100.0	4,603.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	365	25.3	1,165.2
2	Malignant neoplasms (C00-C97)	278	19.3	887.5
3	Chronic lower respiratory diseases (J40-J47)	97	6.7	309.7
4	Diabetes mellitus (E10-E14)	85	5.9	271.3
5	Cerebrovascular diseases (I60-I69)	75	5.2	239.4
6	Alzheimer's disease (G30)	59	4.1	188.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	3.0	137.3
8	Influenza and pneumonia (J09-J18)	39	2.7	124.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	2.4	108.5
10	Septicemia (A40-A41)	25	1.7	79.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	1.2	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	13	0.9	*
13	Parkinson's disease (G20-G21)	12	0.8	*
14	Pneumonitis due to solids and liquids (J69)	10	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.6	*
...	All other causes (Residual)	281	19.5	897.0
American Indian, both sexes, 85-89 years				
...	All causes	1,198	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	302	25.2	---
2	Malignant neoplasms (C00-C97)	201	16.8	---
3	Chronic lower respiratory diseases (J40-J47)	78	6.5	---
4	Diabetes mellitus (E10-E14)	62	5.2	---
5	Cerebrovascular diseases (I60-I69)	59	4.9	---
6	Alzheimer's disease (G30)	46	3.8	---
6	Influenza and pneumonia (J09-J18)	46	3.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	3.4	---
9	Septicemia (A40-A41)	27	2.3	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	2.2	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25	2.1	---
12	Parkinson's disease (G20-G21)	17	1.4	---
13	Pneumonitis due to solids and liquids (J69)	12	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.3	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.3	---
...	All other causes (Residual)	248	20.7	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 90-94 years				
...	All causes	741	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	193	26.0	---
2	Malignant neoplasms (C00-C97)	71	9.6	---
3	Cerebrovascular diseases (I60-I69)	54	7.3	---
4	Alzheimer's disease (G30)	53	7.2	---
5	Chronic lower respiratory diseases (J40-J47)	40	5.4	---
6	Diabetes mellitus (E10-E14)	34	4.6	---
7	Influenza and pneumonia (J09-J18)	33	4.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	2.2	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.9	---
11	Septicemia (A40-A41)	11	1.5	---
12	Parkinson's disease (G20-G21)	9	1.2	---
13	Pneumonitis due to solids and liquids (J69)	7	0.9	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	5	0.7	---
15	Nutritional deficiencies (E40-E64)	4	0.5	---
...	All other causes (Residual)	176	23.8	---
American Indian, both sexes, 95-99 years				
...	All causes	247	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	25.9	---
2	Cerebrovascular diseases (I60-I69)	22	8.9	---
3	Influenza and pneumonia (J09-J18)	17	6.9	---
4	Malignant neoplasms (C00-C97)	15	6.1	---
4	Alzheimer's disease (G30)	15	6.1	---
6	Diabetes mellitus (E10-E14)	9	3.6	---
7	Chronic lower respiratory diseases (J40-J47)	8	3.2	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	2.4	---
9	Nutritional deficiencies (E40-E64)	4	1.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.6	---
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.2	---
11	Parkinson's disease (G20-G21)	3	1.2	---
11	Pneumonitis due to solids and liquids (J69)	3	1.2	---
14	Anemias (D50-D64)	2	0.8	---
14	Atherosclerosis (I70)	2	0.8	---
14	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	0.8	---
...	All other causes (Residual)	68	27.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 100 years and over				
...	All causes	73	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	30.1	---
2	Alzheimer's disease (G30)	7	9.6	---
3	Influenza and pneumonia (J09-J18)	5	6.8	---
4	Pneumonitis due to solids and liquids (J69)	3	4.1	---
5	Malignant neoplasms (C00-C97)	2	2.7	---
5	Diabetes mellitus (E10-E14)	2	2.7	---
5	Cerebrovascular diseases (I60-I69)	2	2.7	---
5	Atherosclerosis (I70)	2	2.7	---
5	Chronic lower respiratory diseases (J40-J47)	2	2.7	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	2.7	---
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	2.7	---
12	Anemias (D50-D64)	1	1.4	---
12	Nutritional deficiencies (E40-E64)	1	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.4	---
12	Pneumoconioses and chemical effects (J60-J66,J68)	1	1.4	---
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	1.4	---
...	All other causes (Residual)	17	23.3	---
American Indian, male, all ages				
...	All causes	9,331	100.0	416.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,843	19.8	82.3
2	Malignant neoplasms (C00-C97)	1,650	17.7	73.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,177	12.6	52.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	511	5.5	22.8
5	Diabetes mellitus (E10-E14)	490	5.3	21.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	401	4.3	17.9
7	Chronic lower respiratory diseases (J40-J47)	370	4.0	16.5
8	Cerebrovascular diseases (I60-I69)	253	2.7	11.3
9	Influenza and pneumonia (J09-J18)	190	2.0	8.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	187	2.0	8.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	1.5	6.4
12	Septicemia (A40-A41)	139	1.5	6.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	94	1.0	4.2
14	Viral hepatitis (B15-B19)	80	0.9	3.6
15	Alzheimer's disease (G30)	77	0.8	3.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	0.8	3.4
...	All other causes (Residual)	1,649	17.7	73.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 1-4 years				
...	All causes	65	100.0	41.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	36.9	15.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	16.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	10.8	*
4	Malignant neoplasms (C00-C97)	2	3.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.1	*
6	Nutritional deficiencies (E40-E64)	1	1.5	*
6	Cerebrovascular diseases (I60-I69)	1	1.5	*
6	Influenza and pneumonia (J09-J18)	1	1.5	*
6	Chronic lower respiratory diseases (J40-J47)	1	1.5	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.5	*
...	All other causes (Residual)	14	21.5	*
American Indian, male, 5-9 years				
...	All causes	26	100.0	13.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	42.3	*
2	Malignant neoplasms (C00-C97)	6	23.1	*
3	Chronic lower respiratory diseases (J40-J47)	2	7.7	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	7.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	3.8	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	3.8	*
...	All other causes (Residual)	3	11.5	*
American Indian, male, 10-14 years				
...	All causes	21	100.0	11.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	33.3	*
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	33.3	*
3	Malignant neoplasms (C00-C97)	2	9.5	*
4	Pneumonitis due to solids and liquids (J69)	1	4.8	*
4	Certain conditions originating in the perinatal period (P00-P96)	1	4.8	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	4.8	*
...	All other causes (Residual)	2	9.5	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 15-19 years				
...	All causes	128	100.0	67.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	39.8	26.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44	34.4	23.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	7.8	*
4	Malignant neoplasms (C00-C97)	4	3.1	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	2.3	*
6	Diabetes mellitus (E10-E14)	2	1.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.6	*
8	Septicemia (A40-A41)	1	0.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
...	All other causes (Residual)	10	7.8	*
American Indian, male, 20-24 years				
...	All causes	260	100.0	126.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	46.5	59.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	27.3	34.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	6.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	2.7	*
5	Malignant neoplasms (C00-C97)	5	1.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.2	*
6	Legal intervention (Y35,Y89.0)	3	1.2	*
8	Diabetes mellitus (E10-E14)	2	0.8	*
8	Chronic lower respiratory diseases (J40-J47)	2	0.8	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.8	*
11	Septicemia (A40-A41)	1	0.4	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (Residual)	22	8.5	10.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 25-29 years				
...	All causes	305	100.0	165.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	44.6	74.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	17.7	29.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	11.8	19.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	12	3.9	*
5	Malignant neoplasms (C00-C97)	10	3.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	2.3	*
7	Influenza and pneumonia (J09-J18)	5	1.6	*
8	Cerebrovascular diseases (I60-I69)	1	0.3	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
8	Legal intervention (Y35,Y89.0)	1	0.3	*
8	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (Residual)	41	13.4	22.3
American Indian, male, 30-34 years				
...	All causes	310	100.0	180.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	33.9	61.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	16.5	29.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	27	8.7	15.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	8.1	14.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	6.8	12.2
6	Malignant neoplasms (C00-C97)	12	3.9	*
7	Diabetes mellitus (E10-E14)	7	2.3	*
8	Septicemia (A40-A41)	3	1.0	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
10	Influenza and pneumonia (J09-J18)	2	0.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
13	Tuberculosis (A16-A19)	1	0.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
13	Cerebrovascular diseases (I60-I69)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	44	14.2	25.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 35-39 years				
...	All causes	305	100.0	200.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	30.8	61.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	36	11.8	23.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	10.5	21.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	8.5	17.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	5.6	*
6	Diabetes mellitus (E10-E14)	12	3.9	*
7	Malignant neoplasms (C00-C97)	11	3.6	*
8	Septicemia (A40-A41)	5	1.6	*
9	Cerebrovascular diseases (I60-I69)	3	1.0	*
9	Influenza and pneumonia (J09-J18)	3	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	3	1.0	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.0	*
13	Viral hepatitis (B15-B19)	2	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.7	*
13	Aortic aneurysm and dissection (I71)	2	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
...	All other causes (Residual)	52	17.0	34.1
American Indian, male, 40-44 years				
...	All causes	425	100.0	290.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	22.6	65.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	16.0	46.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	56	13.2	38.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	8.7	25.3
5	Malignant neoplasms (C00-C97)	28	6.6	19.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	5.4	15.7
7	Diabetes mellitus (E10-E14)	22	5.2	15.0
8	Cerebrovascular diseases (I60-I69)	6	1.4	*
8	Influenza and pneumonia (J09-J18)	6	1.4	*
10	Septicemia (A40-A41)	5	1.2	*
11	Viral hepatitis (B15-B19)	4	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.9	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
...	All other causes (Residual)	63	14.8	43.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 45-49 years				
...	All causes	600	100.0	433.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	19.5	84.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	19.5	84.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	83	13.8	59.9
4	Malignant neoplasms (C00-C97)	52	8.7	37.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	7.5	32.5
6	Diabetes mellitus (E10-E14)	33	5.5	23.8
7	Cerebrovascular diseases (I60-I69)	12	2.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.8	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	1.8	*
10	Influenza and pneumonia (J09-J18)	10	1.7	*
11	Septicemia (A40-A41)	9	1.5	*
11	Viral hepatitis (B15-B19)	9	1.5	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.5	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.5	*
...	All other causes (Residual)	77	12.8	55.6
American Indian, male, 50-54 years				
...	All causes	775	100.0	582.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	159	20.5	119.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	125	16.1	94.0
3	Malignant neoplasms (C00-C97)	116	15.0	87.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	86	11.1	64.7
5	Diabetes mellitus (E10-E14)	33	4.3	24.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	2.7	15.8
7	Viral hepatitis (B15-B19)	20	2.6	15.0
8	Influenza and pneumonia (J09-J18)	14	1.8	*
9	Septicemia (A40-A41)	13	1.7	*
9	Cerebrovascular diseases (I60-I69)	13	1.7	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	1.7	*
12	Chronic lower respiratory diseases (J40-J47)	12	1.5	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.0	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.8	*
...	All other causes (Residual)	127	16.4	95.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 55-59 years				
...	All causes	893	100.0	805.2
1	Malignant neoplasms (C00-C97)	202	22.6	182.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	200	22.4	180.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	8.8	71.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	76	8.5	68.5
5	Diabetes mellitus (E10-E14)	62	6.9	55.9
6	Chronic lower respiratory diseases (J40-J47)	22	2.5	19.8
7	Cerebrovascular diseases (I60-I69)	19	2.1	*
8	Influenza and pneumonia (J09-J18)	18	2.0	*
9	Viral hepatitis (B15-B19)	17	1.9	*
10	Septicemia (A40-A41)	15	1.7	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	1.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.5	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	1.3	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	1.0	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.9	*
...	All other causes (Residual)	127	14.2	114.5
American Indian, male, 60-64 years				
...	All causes	922	100.0	1,115.7
1	Malignant neoplasms (C00-C97)	250	27.1	302.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	199	21.6	240.8
3	Diabetes mellitus (E10-E14)	65	7.0	78.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	60	6.5	72.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	6.2	69.0
6	Chronic lower respiratory diseases (J40-J47)	29	3.1	35.1
7	Cerebrovascular diseases (I60-I69)	27	2.9	32.7
8	Septicemia (A40-A41)	19	2.1	*
9	Viral hepatitis (B15-B19)	17	1.8	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.7	*
12	Influenza and pneumonia (J09-J18)	12	1.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.5	*
15	Atherosclerosis (I70)	4	0.4	*
15	Pneumonitis due to solids and liquids (J69)	4	0.4	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.4	*
...	All other causes (Residual)	133	14.4	160.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 65-69 years				
...	All causes	893	100.0	1,529.2
1	Malignant neoplasms (C00-C97)	222	24.9	380.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	222	24.9	380.2
3	Chronic lower respiratory diseases (J40-J47)	62	6.9	106.2
4	Diabetes mellitus (E10-E14)	61	6.8	104.5
5	Cerebrovascular diseases (I60-I69)	41	4.6	70.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41	4.6	70.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	37	4.1	63.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	2.1	*
9	Septicemia (A40-A41)	15	1.7	*
10	Influenza and pneumonia (J09-J18)	14	1.6	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	1.0	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	0.9	*
13	Viral hepatitis (B15-B19)	5	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.6	*
13	Aortic aneurysm and dissection (I71)	5	0.6	*
...	All other causes (Residual)	127	14.2	217.5
American Indian, male, 70-74 years				
...	All causes	865	100.0	2,347.9
1	Malignant neoplasms (C00-C97)	233	26.9	632.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	221	25.5	599.9
3	Chronic lower respiratory diseases (J40-J47)	58	6.7	157.4
4	Diabetes mellitus (E10-E14)	57	6.6	154.7
5	Cerebrovascular diseases (I60-I69)	28	3.2	76.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	2.7	62.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	2.5	59.7
8	Influenza and pneumonia (J09-J18)	20	2.3	54.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	18	2.1	*
10	Septicemia (A40-A41)	16	1.8	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.2	*
12	Parkinson's disease (G20-G21)	8	0.9	*
12	Alzheimer's disease (G30)	8	0.9	*
14	Viral hepatitis (B15-B19)	6	0.7	*
15	Pneumonitis due to solids and liquids (J69)	5	0.6	*
15	Peptic ulcer (K25-K28)	5	0.6	*
...	All other causes (Residual)	127	14.7	344.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 75-79 years				
...	All causes	783	100.0	3,494.4
1	Malignant neoplasms (C00-C97)	205	26.2	914.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	182	23.2	812.2
3	Diabetes mellitus (E10-E14)	59	7.5	263.3
4	Chronic lower respiratory diseases (J40-J47)	58	7.4	258.8
5	Cerebrovascular diseases (I60-I69)	40	5.1	178.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	3.6	125.0
7	Influenza and pneumonia (J09-J18)	18	2.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	2.3	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.8	*
10	Alzheimer's disease (G30)	11	1.4	*
11	Septicemia (A40-A41)	10	1.3	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.1	*
13	Parkinson's disease (G20-G21)	7	0.9	*
14	Pneumonitis due to solids and liquids (J69)	5	0.6	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.5	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.5	*
...	All other causes (Residual)	111	14.2	495.4
American Indian, male, 80-84 years				
...	All causes	699	100.0	5,515.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	27.8	1,530.7
2	Malignant neoplasms (C00-C97)	149	21.3	1,175.6
3	Chronic lower respiratory diseases (J40-J47)	45	6.4	355.1
4	Diabetes mellitus (E10-E14)	38	5.4	299.8
5	Cerebrovascular diseases (I60-I69)	27	3.9	213.0
6	Alzheimer's disease (G30)	24	3.4	189.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	2.9	157.8
8	Influenza and pneumonia (J09-J18)	19	2.7	*
9	Septicemia (A40-A41)	13	1.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.6	*
11	Pneumonitis due to solids and liquids (J69)	8	1.1	*
12	Parkinson's disease (G20-G21)	7	1.0	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.6	*
...	All other causes (Residual)	128	18.3	1,009.9

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 85-89 years				
...	All causes	519	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	24.5	---
2	Malignant neoplasms (C00-C97)	102	19.7	---
3	Chronic lower respiratory diseases (J40-J47)	44	8.5	---
4	Diabetes mellitus (E10-E14)	25	4.8	---
5	Influenza and pneumonia (J09-J18)	23	4.4	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	4.2	---
7	Cerebrovascular diseases (I60-I69)	17	3.3	---
8	Alzheimer's disease (G30)	12	2.3	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.9	---
11	Septicemia (A40-A41)	9	1.7	---
11	Parkinson's disease (G20-G21)	9	1.7	---
13	Pneumonitis due to solids and liquids (J69)	7	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	---
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	0.6	---
...	All other causes (Residual)	96	18.5	---
American Indian, male, 90-94 years				
...	All causes	261	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	26.8	---
2	Malignant neoplasms (C00-C97)	33	12.6	---
3	Chronic lower respiratory diseases (J40-J47)	20	7.7	---
4	Influenza and pneumonia (J09-J18)	16	6.1	---
5	Alzheimer's disease (G30)	15	5.7	---
6	Cerebrovascular diseases (I60-I69)	13	5.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	4.2	---
8	Diabetes mellitus (E10-E14)	8	3.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	1.9	---
10	Septicemia (A40-A41)	4	1.5	---
10	Parkinson's disease (G20-G21)	4	1.5	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.1	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	---
13	Nutritional deficiencies (E40-E64)	2	0.8	---
13	Pneumonitis due to solids and liquids (J69)	2	0.8	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.8	---
...	All other causes (Residual)	51	19.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 95-99 years				
...	All causes	61	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	23.0	---
2	Malignant neoplasms (C00-C97)	5	8.2	---
3	Diabetes mellitus (E10-E14)	4	6.6	---
4	Cerebrovascular diseases (I60-I69)	3	4.9	---
4	Influenza and pneumonia (J09-J18)	3	4.9	---
4	Chronic lower respiratory diseases (J40-J47)	3	4.9	---
4	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	4.9	---
8	Anemias (D50-D64)	2	3.3	---
8	Nutritional deficiencies (E40-E64)	2	3.3	---
8	Parkinson's disease (G20-G21)	2	3.3	---
8	Alzheimer's disease (G30)	2	3.3	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.6	---
12	Pneumonitis due to solids and liquids (J69)	1	1.6	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	1.6	---
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	1.6	---
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.6	---
...	All other causes (Residual)	12	19.7	---
American Indian, male, 100 years and over				
...	All causes	15	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	26.7	---
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	13.3	---
3	Malignant neoplasms (C00-C97)	1	6.7	---
3	Anemias (D50-D64)	1	6.7	---
3	Nutritional deficiencies (E40-E64)	1	6.7	---
3	Influenza and pneumonia (J09-J18)	1	6.7	---
3	Chronic lower respiratory diseases (J40-J47)	1	6.7	---
3	Pneumoconioses and chemical effects (J60-J66,J68)	1	6.7	---
3	Pneumonitis due to solids and liquids (J69)	1	6.7	---
...	All other causes (Residual)	2	13.3	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, all ages				
...	All causes	7,721	100.0	348.2
1	Malignant neoplasms (C00-C97)	1,459	18.9	65.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,296	16.8	58.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	656	8.5	29.6
4	Diabetes mellitus (E10-E14)	469	6.1	21.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	433	5.6	19.5
6	Chronic lower respiratory diseases (J40-J47)	387	5.0	17.5
7	Cerebrovascular diseases (I60-I69)	342	4.4	15.4
8	Influenza and pneumonia (J09-J18)	185	2.4	8.3
9	Alzheimer's disease (G30)	161	2.1	7.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	2.1	7.2
11	Septicemia (A40-A41)	149	1.9	6.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	120	1.6	5.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	1.2	4.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	0.7	2.4
15	Viral hepatitis (B15-B19)	53	0.7	2.4
...	All other causes (Residual)	1,709	22.1	77.1
American Indian, female, 1-4 years				
...	All causes	39	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	48.7	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	12.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	10.3	*
4	Influenza and pneumonia (J09-J18)	2	5.1	*
5	Malignant neoplasms (C00-C97)	1	2.6	*
5	Acute bronchitis and bronchiolitis (J20-J21)	1	2.6	*
...	All other causes (Residual)	7	17.9	*
American Indian, female, 5-9 years				
...	All causes	24	100.0	12.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	45.8	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	8.3	*
3	Viral hepatitis (B15-B19)	1	4.2	*
3	Malignant neoplasms (C00-C97)	1	4.2	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	4.2	*
...	All other causes (Residual)	8	33.3	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 10-14 years				
...	All causes	17	100.0	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	35.3	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	23.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	11.8	*
4	Malignant neoplasms (C00-C97)	1	5.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	5.9	*
4	Influenza and pneumonia (J09-J18)	1	5.9	*
...	All other causes (Residual)	2	11.8	*
American Indian, female, 15-19 years				
...	All causes	52	100.0	28.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	44.2	12.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	25.0	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	7.7	*
4	Malignant neoplasms (C00-C97)	2	3.8	*
5	Septicemia (A40-A41)	1	1.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.9	*
5	Chronic lower respiratory diseases (J40-J47)	1	1.9	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.9	*
...	All other causes (Residual)	6	11.5	*
American Indian, female, 20-24 years				
...	All causes	108	100.0	56.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	46.3	26.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	20.4	11.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	7.4	*
4	Malignant neoplasms (C00-C97)	4	3.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	3.7	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	3	2.8	*
7	Diabetes mellitus (E10-E14)	1	0.9	*
7	Influenza and pneumonia (J09-J18)	1	0.9	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.9	*
...	All other causes (Residual)	14	13.0	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-29 years				
...	All causes	143	100.0	85.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	48.3	41.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	7.0	*
3	Malignant neoplasms (C00-C97)	9	6.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	4.9	*
5	Diabetes mellitus (E10-E14)	5	3.5	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4	2.8	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	2.1	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	3	2.1	*
7	Septicemia (A40-A41)	2	1.4	*
9	Cerebrovascular diseases (I60-I69)	2	1.4	*
9	Influenza and pneumonia (J09-J18)	2	1.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.4	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.7	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.7	*
...	All other causes (Residual)	23	16.1	13.7
American Indian, female, 30-34 years				
...	All causes	165	100.0	102.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	26.1	26.6
2	Chronic liver disease and cirrhosis (K70,K73-K74)	29	17.6	18.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	10.3	*
4	Malignant neoplasms (C00-C97)	12	7.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	5.5	*
6	Diabetes mellitus (E10-E14)	6	3.6	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	3.6	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	4	2.4	*
9	Septicemia (A40-A41)	3	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.2	*
12	Cerebrovascular diseases (I60-I69)	1	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
...	All other causes (Residual)	29	17.6	18.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 35-39 years				
...	All causes	213	100.0	146.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	23.9	35.1
2	Chronic liver disease and cirrhosis (K70,K73-K74)	31	14.6	21.3
3	Malignant neoplasms (C00-C97)	22	10.3	15.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	9.4	13.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	5.2	*
6	Diabetes mellitus (E10-E14)	8	3.8	*
7	Septicemia (A40-A41)	4	1.9	*
7	Influenza and pneumonia (J09-J18)	4	1.9	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	1.9	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.4	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.9	*
11	Cerebrovascular diseases (I60-I69)	2	0.9	*
13	Tuberculosis (A16-A19)	1	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
13	Pneumonitis due to solids and liquids (J69)	1	0.5	*
13	Diseases of appendix (K35-K38)	1	0.5	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.5	*
13	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	44	20.7	30.3
American Indian, female, 40-44 years				
...	All causes	283	100.0	198.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	22.3	44.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	43	15.2	30.2
3	Malignant neoplasms (C00-C97)	32	11.3	22.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	8.8	17.6
5	Influenza and pneumonia (J09-J18)	15	5.3	*
6	Diabetes mellitus (E10-E14)	12	4.2	*
6	Cerebrovascular diseases (I60-I69)	12	4.2	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	2.8	*
9	Septicemia (A40-A41)	6	2.1	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	2.1	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.8	*
12	Viral hepatitis (B15-B19)	3	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	3	1.1	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.7	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
...	All other causes (Residual)	45	15.9	31.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 45-49 years				
...	All causes	402	100.0	291.0
1	Malignant neoplasms (C00-C97)	77	19.2	55.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	67	16.7	48.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	13.7	39.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	12.7	36.9
5	Chronic lower respiratory diseases (J40-J47)	14	3.5	*
6	Diabetes mellitus (E10-E14)	12	3.0	*
7	Septicemia (A40-A41)	11	2.7	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	2.2	*
9	Viral hepatitis (B15-B19)	8	2.0	*
9	Cerebrovascular diseases (I60-I69)	8	2.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.7	*
12	Influenza and pneumonia (J09-J18)	4	1.0	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.5	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.2	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	67	16.7	48.5
American Indian, female, 50-54 years				
...	All causes	523	100.0	377.9
1	Malignant neoplasms (C00-C97)	114	21.8	82.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	80	15.3	57.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	11.9	44.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	9.9	37.6
5	Diabetes mellitus (E10-E14)	29	5.5	21.0
6	Cerebrovascular diseases (I60-I69)	17	3.3	*
7	Chronic lower respiratory diseases (J40-J47)	16	3.1	*
8	Septicemia (A40-A41)	14	2.7	*
8	Viral hepatitis (B15-B19)	14	2.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.7	*
11	Influenza and pneumonia (J09-J18)	8	1.5	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	1.5	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	0.6	*
...	All other causes (Residual)	82	15.7	59.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 55-59 years				
...	All causes	643	100.0	541.3
1	Malignant neoplasms (C00-C97)	147	22.9	123.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	88	13.7	74.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	68	10.6	57.2
4	Diabetes mellitus (E10-E14)	52	8.1	43.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	8.1	43.8
6	Chronic lower respiratory diseases (J40-J47)	31	4.8	26.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	3.0	*
8	Cerebrovascular diseases (I60-I69)	15	2.3	*
9	Viral hepatitis (B15-B19)	12	1.9	*
10	Influenza and pneumonia (J09-J18)	10	1.6	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	1.6	*
12	Septicemia (A40-A41)	9	1.4	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Atherosclerosis (I70)	2	0.3	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	2	0.3	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.3	*
...	All other causes (Residual)	113	17.6	95.1
American Indian, female, 60-64 years				
...	All causes	645	100.0	715.8
1	Malignant neoplasms (C00-C97)	177	27.4	196.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	122	18.9	135.4
3	Diabetes mellitus (E10-E14)	59	9.1	65.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	32	5.0	35.5
5	Chronic lower respiratory diseases (J40-J47)	31	4.8	34.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	4.0	28.9
7	Cerebrovascular diseases (I60-I69)	25	3.9	27.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	2.3	*
9	Septicemia (A40-A41)	13	2.0	*
9	Influenza and pneumonia (J09-J18)	13	2.0	*
11	Viral hepatitis (B15-B19)	8	1.2	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.8	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
15	Parkinson's disease (G20-G21)	3	0.5	*
...	All other causes (Residual)	107	16.6	118.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65-69 years				
...	All causes	706	100.0	1,095.1
1	Malignant neoplasms (C00-C97)	210	29.7	325.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	115	16.3	178.4
3	Diabetes mellitus (E10-E14)	60	8.5	93.1
4	Chronic lower respiratory diseases (J40-J47)	55	7.8	85.3
5	Cerebrovascular diseases (I60-I69)	31	4.4	48.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	27	3.8	41.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	2.8	31.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	2.8	31.0
7	Septicemia (A40-A41)	18	2.5	*
10	Influenza and pneumonia (J09-J18)	16	2.3	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.7	*
12	Aortic aneurysm and dissection (I71)	4	0.6	*
13	Tuberculosis (A16-A19)	2	0.3	*
13	Viral hepatitis (B15-B19)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Peptic ulcer (K25-K28)	2	0.3	*
...	All other causes (Residual)	117	16.6	181.5
American Indian, female, 70-74 years				
...	All causes	704	100.0	1,645.7
1	Malignant neoplasms (C00-C97)	190	27.0	444.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	17.2	282.9
3	Chronic lower respiratory diseases (J40-J47)	63	8.9	147.3
4	Diabetes mellitus (E10-E14)	56	8.0	130.9
5	Cerebrovascular diseases (I60-I69)	34	4.8	79.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	27	3.8	63.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	2.6	*
8	Septicemia (A40-A41)	13	1.8	*
9	Influenza and pneumonia (J09-J18)	12	1.7	*
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	1.6	*
11	Alzheimer's disease (G30)	7	1.0	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.9	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	6	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
15	Pneumonitis due to solids and liquids (J69)	4	0.6	*
...	All other causes (Residual)	131	18.6	306.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 75-79 years				
...	All causes	789	100.0	2,730.2
1	Malignant neoplasms (C00-C97)	183	23.2	633.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	18.4	501.7
3	Chronic lower respiratory diseases (J40-J47)	60	7.6	207.6
4	Diabetes mellitus (E10-E14)	51	6.5	176.5
5	Cerebrovascular diseases (I60-I69)	41	5.2	141.9
6	Alzheimer's disease (G30)	24	3.0	83.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	3.0	83.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	2.2	*
8	Influenza and pneumonia (J09-J18)	17	2.2	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	2.2	*
11	Septicemia (A40-A41)	16	2.0	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.4	*
13	Pneumonitis due to solids and liquids (J69)	6	0.8	*
14	Parkinson's disease (G20-G21)	5	0.6	*
15	Aortic aneurysm and dissection (I71)	4	0.5	*
...	All other causes (Residual)	168	21.3	581.3
American Indian, female, 80-84 years				
...	All causes	743	100.0	3,983.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	171	23.0	916.8
2	Malignant neoplasms (C00-C97)	129	17.4	691.7
3	Chronic lower respiratory diseases (J40-J47)	52	7.0	278.8
4	Cerebrovascular diseases (I60-I69)	48	6.5	257.4
5	Diabetes mellitus (E10-E14)	47	6.3	252.0
6	Alzheimer's disease (G30)	35	4.7	187.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	3.1	123.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	3.1	123.3
9	Influenza and pneumonia (J09-J18)	20	2.7	107.2
10	Septicemia (A40-A41)	12	1.6	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	1.5	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9	1.2	*
13	Parkinson's disease (G20-G21)	5	0.7	*
13	Atherosclerosis (I70)	5	0.7	*
15	Aortic aneurysm and dissection (I71)	4	0.5	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.5	*
...	All other causes (Residual)	145	19.5	777.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 85-89 years				
...	All causes	679	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	175	25.8	---
2	Malignant neoplasms (C00-C97)	99	14.6	---
3	Cerebrovascular diseases (I60-I69)	42	6.2	---
4	Diabetes mellitus (E10-E14)	37	5.4	---
5	Alzheimer's disease (G30)	34	5.0	---
5	Chronic lower respiratory diseases (J40-J47)	34	5.0	---
7	Influenza and pneumonia (J09-J18)	23	3.4	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	2.8	---
9	Septicemia (A40-A41)	18	2.7	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	2.4	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	2.2	---
12	Parkinson's disease (G20-G21)	8	1.2	---
13	Pneumonitis due to solids and liquids (J69)	5	0.7	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.4	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.3	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.3	---
...	All other causes (Residual)	147	21.6	---
American Indian, female, 90-94 years				
...	All causes	480	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	123	25.6	---
2	Cerebrovascular diseases (I60-I69)	41	8.5	---
3	Malignant neoplasms (C00-C97)	38	7.9	---
3	Alzheimer's disease (G30)	38	7.9	---
5	Diabetes mellitus (E10-E14)	26	5.4	---
6	Chronic lower respiratory diseases (J40-J47)	20	4.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	3.8	---
8	Influenza and pneumonia (J09-J18)	17	3.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	2.3	---
10	Septicemia (A40-A41)	7	1.5	---
11	Parkinson's disease (G20-G21)	5	1.0	---
11	Pneumonitis due to solids and liquids (J69)	5	1.0	---
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	0.6	---
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	---
15	Nutritional deficiencies (E40-E64)	2	0.4	---
15	Atherosclerosis (I70)	2	0.4	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.4	---
...	All other causes (Residual)	119	24.8	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 95-99 years				
...	All causes	186	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	26.9	---
2	Cerebrovascular diseases (I60-I69)	19	10.2	---
3	Influenza and pneumonia (J09-J18)	14	7.5	---
4	Alzheimer's disease (G30)	13	7.0	---
5	Malignant neoplasms (C00-C97)	10	5.4	---
6	Diabetes mellitus (E10-E14)	5	2.7	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	2.7	---
6	Chronic lower respiratory diseases (J40-J47)	5	2.7	---
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.1	---
9	Nutritional deficiencies (E40-E64)	2	1.1	---
9	Atherosclerosis (I70)	2	1.1	---
9	Pneumonitis due to solids and liquids (J69)	2	1.1	---
13	Septicemia (A40-A41)	1	0.5	---
13	Parkinson's disease (G20-G21)	1	0.5	---
13	Aortic aneurysm and dissection (I71)	1	0.5	---
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.5	---
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	0.5	---
...	All other causes (Residual)	51	27.4	---
American Indian, female, 100 years and over				
...	All causes	58	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	31.0	---
2	Alzheimer's disease (G30)	7	12.1	---
3	Influenza and pneumonia (J09-J18)	4	6.9	---
4	Diabetes mellitus (E10-E14)	2	3.4	---
4	Cerebrovascular diseases (I60-I69)	2	3.4	---
4	Atherosclerosis (I70)	2	3.4	---
4	Pneumonitis due to solids and liquids (J69)	2	3.4	---
4	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	3.4	---
9	Malignant neoplasms (C00-C97)	1	1.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.7	---
9	Chronic lower respiratory diseases (J40-J47)	1	1.7	---
9	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	1.7	---
...	All other causes (Residual)	15	25.9	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	59,869	100.0	321.4
1	Malignant neoplasms (C00-C97)	15,703	26.2	84.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,311	22.2	71.4
3	Cerebrovascular diseases (I60-I69)	4,205	7.0	22.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,508	4.2	13.5
5	Diabetes mellitus (E10-E14)	2,309	3.9	12.4
6	Influenza and pneumonia (J09-J18)	2,024	3.4	10.9
7	Chronic lower respiratory diseases (J40-J47)	1,848	3.1	9.9
8	Alzheimer's disease (G30)	1,428	2.4	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,147	1.9	6.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,121	1.9	6.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	983	1.6	5.3
12	Septicemia (A40-A41)	729	1.2	3.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	559	0.9	3.0
14	Parkinson's disease (G20-G21)	528	0.9	2.8
15	Certain conditions originating in the perinatal period (P00-P96)	487	0.8	2.6
...	All other causes (Residual)	10,979	18.3	58.9
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	184	100.0	18.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	24.5	4.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	15.8	3.0
3	Malignant neoplasms (C00-C97)	15	8.2	*
4	Influenza and pneumonia (J09-J18)	6	3.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	3.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.7	*
7	Septicemia (A40-A41)	4	2.2	*
7	Meningitis (G00,G03)	4	2.2	*
7	Chronic lower respiratory diseases (J40-J47)	4	2.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.6	*
11	Cerebrovascular diseases (I60-I69)	2	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	1.1	*
13	Shigellosis and amebiasis (A03,A06)	1	0.5	*
13	Anemias (D50-D64)	1	0.5	*
13	Hernia (K40-K46)	1	0.5	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.5	*
...	All other causes (Residual)	55	29.9	5.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-9 years				
...	All causes	120	100.0	9.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	23.3	2.3
2	Malignant neoplasms (C00-C97)	24	20.0	1.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	6.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	5.8	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	5.0	*
6	Septicemia (A40-A41)	3	2.5	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	2.5	*
6	Influenza and pneumonia (J09-J18)	3	2.5	*
9	Meningitis (G00,G03)	2	1.7	*
9	Cerebrovascular diseases (I60-I69)	2	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	2	1.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.8	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.8	*
...	All other causes (Residual)	29	24.2	2.3
Asian or Pacific Islander, both sexes, 10-14 years				
...	All causes	122	100.0	10.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	18.9	1.9
2	Malignant neoplasms (C00-C97)	22	18.0	1.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	10.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	5.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	5.7	*
6	Chronic lower respiratory diseases (J40-J47)	5	4.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	4.1	*
8	Influenza and pneumonia (J09-J18)	4	3.3	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	3	2.5	*
10	Cerebrovascular diseases (I60-I69)	2	1.6	*
11	Septicemia (A40-A41)	1	0.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
11	Hernia (K40-K46)	1	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	0.8	*
...	All other causes (Residual)	27	22.1	2.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-19 years				
...	All causes	268	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	28.4	6.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	24.6	5.6
3	Malignant neoplasms (C00-C97)	38	14.2	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	7.5	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	4.5	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	3.4	*
7	Cerebrovascular diseases (I60-I69)	3	1.1	*
8	Influenza and pneumonia (J09-J18)	2	0.7	*
9	Tuberculosis (A16-A19)	1	0.4	*
9	Septicemia (A40-A41)	1	0.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
9	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.4	*
...	All other causes (Residual)	36	13.4	3.0
Asian or Pacific Islander, both sexes, 20-24 years				
...	All causes	514	100.0	36.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	32.1	11.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	136	26.5	9.6
3	Malignant neoplasms (C00-C97)	64	12.5	4.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	6.8	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.5	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.6	*
7	Septicemia (A40-A41)	6	1.2	*
8	Cerebrovascular diseases (I60-I69)	5	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	4	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
10	Influenza and pneumonia (J09-J18)	3	0.6	*
10	Legal intervention (Y35,Y89.0)	3	0.6	*
13	Diabetes mellitus (E10-E14)	2	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.4	*
...	All other causes (Residual)	53	10.3	3.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-29 years				
...	All causes	574	100.0	37.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	25.6	9.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	25.3	9.4
3	Malignant neoplasms (C00-C97)	78	13.6	5.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	7.3	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	7.1	2.6
6	Cerebrovascular diseases (I60-I69)	7	1.2	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	7	1.2	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.0	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.0	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.0	*
11	Viral hepatitis (B15-B19)	3	0.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
11	Diabetes mellitus (E10-E14)	3	0.5	*
11	Influenza and pneumonia (J09-J18)	3	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	71	12.4	4.6
Asian or Pacific Islander, both sexes, 30-34 years				
...	All causes	658	100.0	41.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	22.8	9.5
2	Malignant neoplasms (C00-C97)	124	18.8	7.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	113	17.2	7.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	59	9.0	3.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	3.8	1.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	15	2.3	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	15	2.3	*
8	Cerebrovascular diseases (I60-I69)	13	2.0	*
9	Septicemia (A40-A41)	8	1.2	*
9	Diabetes mellitus (E10-E14)	8	1.2	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.1	*
12	Influenza and pneumonia (J09-J18)	5	0.8	*
13	Tuberculosis (A16-A19)	4	0.6	*
13	Aortic aneurysm and dissection (I71)	4	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.6	*
...	All other causes (Residual)	104	15.8	6.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 35-39 years				
...	All causes	807	100.0	53.8
1	Malignant neoplasms (C00-C97)	207	25.7	13.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	124	15.4	8.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	107	13.3	7.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	92	11.4	6.1
5	Cerebrovascular diseases (I60-I69)	33	4.1	2.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	3.7	2.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	23	2.9	1.5
8	Diabetes mellitus (E10-E14)	14	1.7	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	12	1.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	8	1.0	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.9	*
14	Viral hepatitis (B15-B19)	5	0.6	*
14	Legal intervention (Y35,Y89.0)	5	0.6	*
...	All other causes (Residual)	125	15.5	8.3
Asian or Pacific Islander, both sexes, 40-44 years				
...	All causes	1,251	100.0	84.3
1	Malignant neoplasms (C00-C97)	428	34.2	28.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	201	16.1	13.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	128	10.2	8.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	8.1	6.8
5	Cerebrovascular diseases (I60-I69)	78	6.2	5.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	31	2.5	2.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	1.9	1.6
8	Diabetes mellitus (E10-E14)	23	1.8	1.5
9	Influenza and pneumonia (J09-J18)	13	1.0	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.0	*
12	Septicemia (A40-A41)	11	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.7	*
13	Chronic lower respiratory diseases (J40-J47)	9	0.7	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.7	*
...	All other causes (Residual)	152	12.2	10.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-49 years				
...	All causes	1,773	100.0	138.3
1	Malignant neoplasms (C00-C97)	646	36.4	50.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	304	17.1	23.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	7.7	10.6
4	Cerebrovascular diseases (I60-I69)	119	6.7	9.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	96	5.4	7.5
6	Diabetes mellitus (E10-E14)	63	3.6	4.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	48	2.7	3.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	1.1	1.6
9	Influenza and pneumonia (J09-J18)	19	1.1	*
10	Septicemia (A40-A41)	18	1.0	*
11	Viral hepatitis (B15-B19)	16	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	14	0.8	*
12	Aortic aneurysm and dissection (I71)	14	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.8	*
15	Chronic lower respiratory diseases (J40-J47)	13	0.7	*
...	All other causes (Residual)	233	13.1	18.2
Asian or Pacific Islander, both sexes, 50-54 years				
...	All causes	2,454	100.0	207.1
1	Malignant neoplasms (C00-C97)	970	39.5	81.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	449	18.3	37.9
3	Cerebrovascular diseases (I60-I69)	176	7.2	14.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	6.1	12.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	88	3.6	7.4
6	Diabetes mellitus (E10-E14)	86	3.5	7.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	43	1.8	3.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.3	2.8
9	Viral hepatitis (B15-B19)	29	1.2	2.4
9	Chronic lower respiratory diseases (J40-J47)	29	1.2	2.4
11	Septicemia (A40-A41)	26	1.1	2.2
12	Influenza and pneumonia (J09-J18)	25	1.0	2.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	1.0	2.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	0.9	1.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.7	*
...	All other causes (Residual)	288	11.7	24.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 55-59 years				
...	All causes	3,316	100.0	312.3
1	Malignant neoplasms (C00-C97)	1,400	42.2	131.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	660	19.9	62.2
3	Cerebrovascular diseases (I60-I69)	197	5.9	18.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161	4.9	15.2
5	Diabetes mellitus (E10-E14)	146	4.4	13.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	2.1	6.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	61	1.8	5.7
8	Chronic lower respiratory diseases (J40-J47)	52	1.6	4.9
9	Influenza and pneumonia (J09-J18)	45	1.4	4.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	1.3	4.0
11	Viral hepatitis (B15-B19)	40	1.2	3.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	0.9	2.7
13	Septicemia (A40-A41)	27	0.8	2.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.6	1.9
15	Aortic aneurysm and dissection (I71)	17	0.5	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	0.5	*
...	All other causes (Residual)	331	10.0	31.2
Asian or Pacific Islander, both sexes, 60-64 years				
...	All causes	4,314	100.0	486.9
1	Malignant neoplasms (C00-C97)	1,733	40.2	195.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	962	22.3	108.6
3	Cerebrovascular diseases (I60-I69)	242	5.6	27.3
4	Diabetes mellitus (E10-E14)	187	4.3	21.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	2.7	13.2
6	Influenza and pneumonia (J09-J18)	88	2.0	9.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	1.8	8.7
8	Chronic lower respiratory diseases (J40-J47)	75	1.7	8.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	68	1.6	7.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	1.4	7.0
11	Septicemia (A40-A41)	49	1.1	5.5
12	Viral hepatitis (B15-B19)	47	1.1	5.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	39	0.9	4.4
14	Aortic aneurysm and dissection (I71)	33	0.8	3.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.5	2.4
...	All other causes (Residual)	514	11.9	58.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65-69 years				
...	All causes	4,737	100.0	719.2
1	Malignant neoplasms (C00-C97)	1,815	38.3	275.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,020	21.5	154.9
3	Cerebrovascular diseases (I60-I69)	344	7.3	52.2
4	Diabetes mellitus (E10-E14)	236	5.0	35.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	2.5	17.9
6	Chronic lower respiratory diseases (J40-J47)	92	1.9	14.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	91	1.9	13.8
8	Influenza and pneumonia (J09-J18)	83	1.8	12.6
9	Septicemia (A40-A41)	79	1.7	12.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	74	1.6	11.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	68	1.4	10.3
12	Viral hepatitis (B15-B19)	41	0.9	6.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	0.7	5.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.7	5.2
15	Aortic aneurysm and dissection (I71)	25	0.5	3.8
...	All other causes (Residual)	582	12.3	88.4
Asian or Pacific Islander, both sexes, 70-74 years				
...	All causes	5,272	100.0	1,144.6
1	Malignant neoplasms (C00-C97)	1,865	35.4	404.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,133	21.5	246.0
3	Cerebrovascular diseases (I60-I69)	356	6.8	77.3
4	Diabetes mellitus (E10-E14)	249	4.7	54.1
5	Chronic lower respiratory diseases (J40-J47)	155	2.9	33.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	151	2.9	32.8
7	Influenza and pneumonia (J09-J18)	130	2.5	28.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	2.4	27.6
9	Septicemia (A40-A41)	84	1.6	18.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.4	15.6
11	Parkinson's disease (G20-G21)	55	1.0	11.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	53	1.0	11.5
13	Alzheimer's disease (G30)	51	1.0	11.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.8	8.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	0.6	7.4
...	All other causes (Residual)	717	13.6	155.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 75-79 years				
...	All causes	6,689	100.0	2,060.6
1	Malignant neoplasms (C00-C97)	2,026	30.3	624.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,458	21.8	449.2
3	Cerebrovascular diseases (I60-I69)	511	7.6	157.4
4	Diabetes mellitus (E10-E14)	354	5.3	109.1
5	Chronic lower respiratory diseases (J40-J47)	251	3.8	77.3
6	Influenza and pneumonia (J09-J18)	219	3.3	67.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	2.8	57.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	146	2.2	45.0
9	Alzheimer's disease (G30)	111	1.7	34.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	110	1.6	33.9
11	Septicemia (A40-A41)	97	1.5	29.9
12	Parkinson's disease (G20-G21)	80	1.2	24.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	46	0.7	14.2
14	Pneumonitis due to solids and liquids (J69)	41	0.6	12.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.5	11.1
...	All other causes (Residual)	1,018	15.2	313.6
Asian or Pacific Islander, both sexes, 80-84 years				
...	All causes	8,102	100.0	3,713.6
1	Malignant neoplasms (C00-C97)	1,924	23.7	881.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,907	23.5	874.1
3	Cerebrovascular diseases (I60-I69)	654	8.1	299.8
4	Chronic lower respiratory diseases (J40-J47)	372	4.6	170.5
5	Diabetes mellitus (E10-E14)	352	4.3	161.3
6	Influenza and pneumonia (J09-J18)	336	4.1	154.0
7	Alzheimer's disease (G30)	240	3.0	110.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	2.5	91.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	172	2.1	78.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	162	2.0	74.3
11	Parkinson's disease (G20-G21)	130	1.6	59.6
12	Septicemia (A40-A41)	107	1.3	49.0
13	Pneumonitis due to solids and liquids (J69)	68	0.8	31.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.7	26.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	48	0.6	22.0
...	All other causes (Residual)	1,373	16.9	629.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 85-89 years				
...	All causes	8,459	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,235	26.4	---
2	Malignant neoplasms (C00-C97)	1,423	16.8	---
3	Cerebrovascular diseases (I60-I69)	723	8.5	---
4	Influenza and pneumonia (J09-J18)	425	5.0	---
5	Alzheimer's disease (G30)	395	4.7	---
6	Chronic lower respiratory diseases (J40-J47)	385	4.6	---
7	Diabetes mellitus (E10-E14)	323	3.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	2.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	199	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	166	2.0	---
11	Parkinson's disease (G20-G21)	139	1.6	---
12	Pneumonitis due to solids and liquids (J69)	104	1.2	---
13	Septicemia (A40-A41)	102	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.6	---
15	Aortic aneurysm and dissection (I71)	40	0.5	---
...	All other causes (Residual)	1,533	18.1	---
Asian or Pacific Islander, both sexes, 90-94 years				
...	All causes	6,268	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,806	28.8	---
2	Malignant neoplasms (C00-C97)	691	11.0	---
3	Cerebrovascular diseases (I60-I69)	526	8.4	---
4	Influenza and pneumonia (J09-J18)	421	6.7	---
5	Alzheimer's disease (G30)	376	6.0	---
6	Chronic lower respiratory diseases (J40-J47)	254	4.1	---
7	Diabetes mellitus (E10-E14)	189	3.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	174	2.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	2.4	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	1.8	---
11	Pneumonitis due to solids and liquids (J69)	97	1.5	---
12	Septicemia (A40-A41)	75	1.2	---
13	Parkinson's disease (G20-G21)	59	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.6	---
15	Aortic aneurysm and dissection (I71)	28	0.4	---
...	All other causes (Residual)	1,269	20.2	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 95-99 years				
...	All causes	2,546	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	759	29.8	---
2	Alzheimer's disease (G30)	186	7.3	---
3	Malignant neoplasms (C00-C97)	184	7.2	---
4	Cerebrovascular diseases (I60-I69)	166	6.5	---
5	Influenza and pneumonia (J09-J18)	146	5.7	---
6	Chronic lower respiratory diseases (J40-J47)	109	4.3	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	88	3.5	---
8	Diabetes mellitus (E10-E14)	66	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	2.0	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	49	1.9	---
11	Pneumonitis due to solids and liquids (J69)	38	1.5	---
12	Septicemia (A40-A41)	23	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.7	---
13	Parkinson's disease (G20-G21)	19	0.7	---
15	Atherosclerosis (I70)	11	0.4	---
...	All other causes (Residual)	631	24.8	---
Asian or Pacific Islander, both sexes, 100 years and over				
...	All causes	563	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	27.7	---
2	Influenza and pneumonia (J09-J18)	43	7.6	---
3	Cerebrovascular diseases (I60-I69)	42	7.5	---
4	Alzheimer's disease (G30)	38	6.7	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	4.3	---
6	Malignant neoplasms (C00-C97)	23	4.1	---
7	Chronic lower respiratory diseases (J40-J47)	22	3.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	2.1	---
10	Atherosclerosis (I70)	9	1.6	---
10	Pneumonitis due to solids and liquids (J69)	9	1.6	---
12	Diabetes mellitus (E10-E14)	8	1.4	---
13	Nutritional deficiencies (E40-E64)	6	1.1	---
14	Parkinson's disease (G20-G21)	5	0.9	---
15	Septicemia (A40-A41)	2	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	---
15	Anemias (D50-D64)	2	0.4	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.4	---
...	All other causes (Residual)	144	25.6	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	30,980	100.0	347.4
1	Malignant neoplasms (C00-C97)	8,071	26.1	90.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,307	23.6	81.9
3	Cerebrovascular diseases (I60-I69)	1,897	6.1	21.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,561	5.0	17.5
5	Diabetes mellitus (E10-E14)	1,228	4.0	13.8
6	Chronic lower respiratory diseases (J40-J47)	1,117	3.6	12.5
7	Influenza and pneumonia (J09-J18)	1,011	3.3	11.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	820	2.6	9.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	574	1.9	6.4
10	Alzheimer's disease (G30)	447	1.4	5.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	433	1.4	4.9
12	Septicemia (A40-A41)	371	1.2	4.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	359	1.2	4.0
14	Parkinson's disease (G20-G21)	299	1.0	3.4
15	Certain conditions originating in the perinatal period (P00-P96)	277	0.9	3.1
...	All other causes (Residual)	5,208	16.8	58.4
Asian or Pacific Islander, male, 1-4 years				
...	All causes	96	100.0	19.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	27.1	5.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	13.5	*
3	Malignant neoplasms (C00-C97)	6	6.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	5.2	*
5	Septicemia (A40-A41)	3	3.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.1	*
5	Chronic lower respiratory diseases (J40-J47)	3	3.1	*
8	Meningitis (G00,G03)	2	2.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
9	Anemias (D50-D64)	1	1.0	*
9	Hernia (K40-K46)	1	1.0	*
9	Infections of kidney (N10-N12,N13.6,N15.1)	1	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (Residual)	30	31.3	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-9 years				
...	All causes	69	100.0	11.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	26.1	*
2	Malignant neoplasms (C00-C97)	14	20.3	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	5.8	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	5.8	*
5	Septicemia (A40-A41)	3	4.3	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	4.3	*
7	Meningitis (G00,G03)	2	2.9	*
7	Cerebrovascular diseases (I60-I69)	2	2.9	*
7	Influenza and pneumonia (J09-J18)	2	2.9	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	2.9	*
11	Chronic lower respiratory diseases (J40-J47)	1	1.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	1.4	*
...	All other causes (Residual)	12	17.4	*
Asian or Pacific Islander, male, 10-14 years				
...	All causes	67	100.0	11.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	22.4	*
2	Malignant neoplasms (C00-C97)	12	17.9	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	7.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	6.0	*
4	Chronic lower respiratory diseases (J40-J47)	4	6.0	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	6.0	*
7	Influenza and pneumonia (J09-J18)	2	3.0	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.0	*
7	Complications of medical and surgical care (Y40-Y84,Y88)	2	3.0	*
10	Cerebrovascular diseases (I60-I69)	1	1.5	*
10	Hernia (K40-K46)	1	1.5	*
...	All other causes (Residual)	15	22.4	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-19 years				
...	All causes	179	100.0	29.9
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	29.6	8.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	26.8	8.0
3	Malignant neoplasms (C00-C97)	19	10.6	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	9.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	3.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	3.4	*
7	Influenza and pneumonia (J09-J18)	2	1.1	*
8	Tuberculosis (A16-A19)	1	0.6	*
8	Septicemia (A40-A41)	1	0.6	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
8	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.6	*
...	All other causes (Residual)	24	13.4	4.0
Asian or Pacific Islander, male, 20-24 years				
...	All causes	376	100.0	52.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	34.6	18.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	103	27.4	14.4
3	Malignant neoplasms (C00-C97)	38	10.1	5.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	7.7	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	5.6	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	1.1	*
7	Cerebrovascular diseases (I60-I69)	3	0.8	*
7	Legal intervention (Y35,Y89.0)	3	0.8	*
9	Septicemia (A40-A41)	2	0.5	*
9	Influenza and pneumonia (J09-J18)	2	0.5	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
12	Tuberculosis (A16-A19)	1	0.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
12	Diabetes mellitus (E10-E14)	1	0.3	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
...	All other causes (Residual)	33	8.8	4.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-29 years				
...	All causes	406	100.0	53.7
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	119	29.3	15.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110	27.1	14.5
3	Malignant neoplasms (C00-C97)	42	10.3	5.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	7.9	4.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	6.9	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.5	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	4	1.0	*
8	Viral hepatitis (B15-B19)	3	0.7	*
8	Influenza and pneumonia (J09-J18)	3	0.7	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
12	Diabetes mellitus (E10-E14)	2	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
12	Legal intervention (Y35,Y89.0)	2	0.5	*
15	Septicemia (A40-A41)	1	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
15	Parkinson's disease (G20-G21)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Cerebrovascular diseases (I60-I69)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
...	All other causes (Residual)	41	10.1	5.4
Asian or Pacific Islander, male, 30-34 years				
...	All causes	408	100.0	54.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103	25.2	13.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	19.6	10.7
3	Malignant neoplasms (C00-C97)	56	13.7	7.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	10.3	5.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	4.7	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	10	2.5	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.5	*
7	Cerebrovascular diseases (I60-I69)	6	1.5	*
9	Influenza and pneumonia (J09-J18)	4	1.0	*
10	Tuberculosis (A16-A19)	3	0.7	*
10	Septicemia (A40-A41)	3	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
10	Diabetes mellitus (E10-E14)	3	0.7	*
10	Aortic aneurysm and dissection (I71)	3	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
...	All other causes (Residual)	64	15.7	8.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-39 years				
...	All causes	501	100.0	70.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	19.2	13.6
2	Malignant neoplasms (C00-C97)	94	18.8	13.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	79	15.8	11.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	71	14.2	10.0
5	Cerebrovascular diseases (I60-I69)	20	4.0	2.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	3.4	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	16	3.2	*
8	Diabetes mellitus (E10-E14)	8	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.2	*
10	Viral hepatitis (B15-B19)	4	0.8	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.8	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.8	*
10	Legal intervention (Y35,Y89.0)	4	0.8	*
14	Aortic aneurysm and dissection (I71)	3	0.6	*
14	Influenza and pneumonia (J09-J18)	3	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
...	All other causes (Residual)	66	13.2	9.3
Asian or Pacific Islander, male, 40-44 years				
...	All causes	743	100.0	106.5
1	Malignant neoplasms (C00-C97)	191	25.7	27.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	158	21.3	22.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	14.1	15.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	9.3	9.9
5	Cerebrovascular diseases (I60-I69)	37	5.0	5.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	21	2.8	3.0
7	Diabetes mellitus (E10-E14)	18	2.4	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	2.0	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.3	*
10	Influenza and pneumonia (J09-J18)	8	1.1	*
11	Viral hepatitis (B15-B19)	7	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.8	*
14	Tuberculosis (A16-A19)	5	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.7	*
...	All other causes (Residual)	82	11.0	11.8

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-49 years				
...	All causes	1,072	100.0	177.7
1	Malignant neoplasms (C00-C97)	293	27.3	48.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	240	22.4	39.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	8.7	15.4
4	Cerebrovascular diseases (I60-I69)	79	7.4	13.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	71	6.6	11.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	43	4.0	7.1
7	Diabetes mellitus (E10-E14)	41	3.8	6.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.2	*
9	Septicemia (A40-A41)	12	1.1	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	1.1	*
11	Viral hepatitis (B15-B19)	10	0.9	*
11	Aortic aneurysm and dissection (I71)	10	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
14	Influenza and pneumonia (J09-J18)	8	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
...	All other causes (Residual)	130	12.1	21.5
Asian or Pacific Islander, male, 50-54 years				
...	All causes	1,479	100.0	266.9
1	Malignant neoplasms (C00-C97)	490	33.1	88.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	349	23.6	63.0
3	Cerebrovascular diseases (I60-I69)	103	7.0	18.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	6.3	16.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	3.8	10.1
6	Diabetes mellitus (E10-E14)	54	3.7	9.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	34	2.3	6.1
8	Chronic lower respiratory diseases (J40-J47)	20	1.4	3.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.4	3.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	1.3	*
11	Septicemia (A40-A41)	18	1.2	*
11	Viral hepatitis (B15-B19)	18	1.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.9	*
14	Aortic aneurysm and dissection (I71)	12	0.8	*
14	Influenza and pneumonia (J09-J18)	12	0.8	*
...	All other causes (Residual)	168	11.4	30.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-59 years				
...	All causes	2,011	100.0	413.7
1	Malignant neoplasms (C00-C97)	738	36.7	151.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	490	24.4	100.8
3	Cerebrovascular diseases (I60-I69)	109	5.4	22.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	5.3	21.8
5	Diabetes mellitus (E10-E14)	95	4.7	19.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	2.5	10.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	46	2.3	9.5
8	Chronic lower respiratory diseases (J40-J47)	31	1.5	6.4
9	Influenza and pneumonia (J09-J18)	30	1.5	6.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	1.2	5.1
11	Viral hepatitis (B15-B19)	24	1.2	4.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	1.2	4.9
13	Septicemia (A40-A41)	17	0.8	*
14	Aortic aneurysm and dissection (I71)	11	0.5	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	0.4	*
...	All other causes (Residual)	205	10.2	42.2
Asian or Pacific Islander, male, 60-64 years				
...	All causes	2,563	100.0	644.5
1	Malignant neoplasms (C00-C97)	909	35.5	228.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	688	26.8	173.0
3	Cerebrovascular diseases (I60-I69)	136	5.3	34.2
4	Diabetes mellitus (E10-E14)	118	4.6	29.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	2.8	17.9
6	Influenza and pneumonia (J09-J18)	49	1.9	12.3
6	Chronic lower respiratory diseases (J40-J47)	49	1.9	12.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.8	11.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	1.8	11.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	41	1.6	10.3
11	Septicemia (A40-A41)	31	1.2	7.8
12	Viral hepatitis (B15-B19)	28	1.1	7.0
13	Aortic aneurysm and dissection (I71)	24	0.9	6.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.8	5.0
15	Pneumonitis due to solids and liquids (J69)	12	0.5	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.5	*
...	All other causes (Residual)	284	11.1	71.4

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-69 years				
...	All causes	2,705	100.0	913.5
1	Malignant neoplasms (C00-C97)	945	34.9	319.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	676	25.0	228.3
3	Cerebrovascular diseases (I60-I69)	188	7.0	63.5
4	Diabetes mellitus (E10-E14)	154	5.7	52.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	2.2	19.9
6	Chronic lower respiratory diseases (J40-J47)	58	2.1	19.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.8	16.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	49	1.8	16.5
9	Septicemia (A40-A41)	45	1.7	15.2
9	Influenza and pneumonia (J09-J18)	45	1.7	15.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	1.2	11.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	1.0	9.1
13	Viral hepatitis (B15-B19)	23	0.9	7.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.6	*
15	Aortic aneurysm and dissection (I71)	14	0.5	*
...	All other causes (Residual)	323	11.9	109.1
Asian or Pacific Islander, male, 70-74 years				
...	All causes	2,975	100.0	1,429.4
1	Malignant neoplasms (C00-C97)	1,009	33.9	484.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	710	23.9	341.1
3	Cerebrovascular diseases (I60-I69)	180	6.1	86.5
4	Diabetes mellitus (E10-E14)	147	4.9	70.6
5	Chronic lower respiratory diseases (J40-J47)	95	3.2	45.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	84	2.8	40.4
7	Influenza and pneumonia (J09-J18)	78	2.6	37.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	2.1	29.8
9	Septicemia (A40-A41)	50	1.7	24.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	1.2	16.8
11	Parkinson's disease (G20-G21)	33	1.1	15.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	33	1.1	15.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.8	11.5
14	Alzheimer's disease (G30)	23	0.8	11.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	0.7	9.6
...	All other causes (Residual)	392	13.2	188.3

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-79 years				
...	All causes	3,520	100.0	2,473.7
1	Malignant neoplasms (C00-C97)	1,079	30.7	758.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	812	23.1	570.6
3	Cerebrovascular diseases (I60-I69)	246	7.0	172.9
4	Diabetes mellitus (E10-E14)	187	5.3	131.4
5	Chronic lower respiratory diseases (J40-J47)	149	4.2	104.7
6	Influenza and pneumonia (J09-J18)	124	3.5	87.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	2.7	66.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	2.0	49.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	55	1.6	38.7
10	Parkinson's disease (G20-G21)	50	1.4	35.1
11	Septicemia (A40-A41)	46	1.3	32.3
12	Alzheimer's disease (G30)	42	1.2	29.5
13	Pneumonitis due to solids and liquids (J69)	27	0.8	19.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.6	14.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	18	0.5	*
...	All other causes (Residual)	500	14.2	351.4
Asian or Pacific Islander, male, 80-84 years				
...	All causes	3,929	100.0	4,484.4
1	Malignant neoplasms (C00-C97)	989	25.2	1,128.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	923	23.5	1,053.5
3	Cerebrovascular diseases (I60-I69)	284	7.2	324.1
4	Chronic lower respiratory diseases (J40-J47)	243	6.2	277.3
5	Influenza and pneumonia (J09-J18)	164	4.2	187.2
6	Diabetes mellitus (E10-E14)	161	4.1	183.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	3.1	138.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	2.2	100.4
9	Alzheimer's disease (G30)	81	2.1	92.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	71	1.8	81.0
11	Parkinson's disease (G20-G21)	70	1.8	79.9
12	Septicemia (A40-A41)	49	1.2	55.9
13	Pneumonitis due to solids and liquids (J69)	44	1.1	50.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.7	30.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	26	0.7	29.7
...	All other causes (Residual)	588	15.0	671.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85-89 years				
...	All causes	3,777	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,007	26.7	---
2	Malignant neoplasms (C00-C97)	708	18.7	---
3	Cerebrovascular diseases (I60-I69)	262	6.9	---
4	Chronic lower respiratory diseases (J40-J47)	243	6.4	---
5	Influenza and pneumonia (J09-J18)	213	5.6	---
6	Diabetes mellitus (E10-E14)	139	3.7	---
7	Alzheimer's disease (G30)	117	3.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	2.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	2.3	---
10	Parkinson's disease (G20-G21)	79	2.1	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	1.9	---
12	Pneumonitis due to solids and liquids (J69)	53	1.4	---
13	Septicemia (A40-A41)	47	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.5	---
14	Aortic aneurysm and dissection (I71)	19	0.5	---
...	All other causes (Residual)	602	15.9	---
Asian or Pacific Islander, male, 90-94 years				
...	All causes	2,599	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	755	29.0	---
2	Malignant neoplasms (C00-C97)	340	13.1	---
3	Cerebrovascular diseases (I60-I69)	194	7.5	---
4	Influenza and pneumonia (J09-J18)	186	7.2	---
5	Chronic lower respiratory diseases (J40-J47)	145	5.6	---
6	Alzheimer's disease (G30)	120	4.6	---
7	Diabetes mellitus (E10-E14)	79	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	73	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	2.6	---
10	Pneumonitis due to solids and liquids (J69)	52	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	1.9	---
12	Parkinson's disease (G20-G21)	31	1.2	---
13	Septicemia (A40-A41)	25	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.7	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.4	---
...	All other causes (Residual)	455	17.5	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 95-99 years				
...	All causes	847	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	265	31.3	---
2	Malignant neoplasms (C00-C97)	90	10.6	---
3	Influenza and pneumonia (J09-J18)	52	6.1	---
4	Alzheimer's disease (G30)	45	5.3	---
5	Chronic lower respiratory diseases (J40-J47)	42	5.0	---
6	Cerebrovascular diseases (I60-I69)	39	4.6	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	2.7	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	2.4	---
10	Diabetes mellitus (E10-E14)	19	2.2	---
11	Pneumonitis due to solids and liquids (J69)	18	2.1	---
12	Septicemia (A40-A41)	10	1.2	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.2	---
14	Parkinson's disease (G20-G21)	7	0.8	---
15	Atherosclerosis (I70)	4	0.5	---
15	Aortic aneurysm and dissection (I71)	4	0.5	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.5	---
...	All other causes (Residual)	170	20.1	---
Asian or Pacific Islander, male, 100 years and over				
...	All causes	162	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	29.0	---
2	Influenza and pneumonia (J09-J18)	23	14.2	---
3	Malignant neoplasms (C00-C97)	9	5.6	---
4	Alzheimer's disease (G30)	7	4.3	---
4	Chronic lower respiratory diseases (J40-J47)	7	4.3	---
6	Cerebrovascular diseases (I60-I69)	5	3.1	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	2.5	---
9	Parkinson's disease (G20-G21)	3	1.9	---
9	Pneumonitis due to solids and liquids (J69)	3	1.9	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.9	---
12	Septicemia (A40-A41)	2	1.2	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.2	---
12	Anemias (D50-D64)	2	1.2	---
12	Diabetes mellitus (E10-E14)	2	1.2	---
12	Nutritional deficiencies (E40-E64)	2	1.2	---
12	Atherosclerosis (I70)	2	1.2	---
...	All other causes (Residual)	34	21.0	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	28,889	100.0	297.4
1	Malignant neoplasms (C00-C97)	7,632	26.4	78.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,004	20.8	61.8
3	Cerebrovascular diseases (I60-I69)	2,308	8.0	23.8
4	Diabetes mellitus (E10-E14)	1,081	3.7	11.1
5	Influenza and pneumonia (J09-J18)	1,013	3.5	10.4
6	Alzheimer's disease (G30)	981	3.4	10.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	947	3.3	9.7
8	Chronic lower respiratory diseases (J40-J47)	731	2.5	7.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	573	2.0	5.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	550	1.9	5.7
11	Septicemia (A40-A41)	358	1.2	3.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	301	1.0	3.1
13	Parkinson's disease (G20-G21)	229	0.8	2.4
14	Certain conditions originating in the perinatal period (P00-P96)	210	0.7	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	200	0.7	2.1
...	All other causes (Residual)	5,771	20.0	59.4
Asian or Pacific Islander, female, 1-4 years				
...	All causes	88	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	21.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	18.2	*
3	Malignant neoplasms (C00-C97)	9	10.2	*
4	Influenza and pneumonia (J09-J18)	6	6.8	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.3	*
5	Meningitis (G00,G03)	2	2.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.3	*
5	Cerebrovascular diseases (I60-I69)	2	2.3	*
9	Shigellosis and amebiasis (A03,A06)	1	1.1	*
9	Septicemia (A40-A41)	1	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.1	*
...	All other causes (Residual)	25	28.4	5.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-9 years				
...	All causes	51	100.0	8.3
1	Malignant neoplasms (C00-C97)	10	19.6	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	19.6	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	7.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	7.8	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.9	*
6	Influenza and pneumonia (J09-J18)	1	2.0	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.0	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.0	*
...	All other causes (Residual)	17	33.3	*
Asian or Pacific Islander, female, 10-14 years				
...	All causes	55	100.0	9.3
1	Malignant neoplasms (C00-C97)	10	18.2	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	14.5	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	14.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	5.5	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	5.5	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.5	*
7	Influenza and pneumonia (J09-J18)	2	3.6	*
8	Septicemia (A40-A41)	1	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.8	*
8	Cerebrovascular diseases (I60-I69)	1	1.8	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.8	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.8	*
...	All other causes (Residual)	12	21.8	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-19 years				
...	All causes	89	100.0	15.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	31.5	4.8
2	Malignant neoplasms (C00-C97)	19	21.3	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	14.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	6.7	*
5	Cerebrovascular diseases (I60-I69)	3	3.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	3.4	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.1	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	1	1.1	*
...	All other causes (Residual)	12	13.5	*
Asian or Pacific Islander, female, 20-24 years				
...	All causes	138	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	25.4	5.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	23.9	4.7
3	Malignant neoplasms (C00-C97)	26	18.8	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	4.3	*
5	Septicemia (A40-A41)	4	2.9	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	2.9	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.4	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.4	*
7	Cerebrovascular diseases (I60-I69)	2	1.4	*
7	Chronic lower respiratory diseases (J40-J47)	2	1.4	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.4	*
12	Diabetes mellitus (E10-E14)	1	0.7	*
12	Influenza and pneumonia (J09-J18)	1	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
...	All other causes (Residual)	17	12.3	*

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-29 years				
...	All causes	168	100.0	21.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	22.0	4.7
2	Malignant neoplasms (C00-C97)	36	21.4	4.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	15.5	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	8.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	5.4	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	7	4.2	*
7	Cerebrovascular diseases (I60-I69)	6	3.6	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.2	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2	1.2	*
11	Septicemia (A40-A41)	1	0.6	*
11	Diabetes mellitus (E10-E14)	1	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
11	Pneumonitis due to solids and liquids (J69)	1	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	*
...	All other causes (Residual)	21	12.5	2.7
Asian or Pacific Islander, female, 30-34 years				
...	All causes	250	100.0	29.9
1	Malignant neoplasms (C00-C97)	68	27.2	8.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	18.8	5.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	13.2	3.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	6.8	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	15	6.0	*
6	Cerebrovascular diseases (I60-I69)	7	2.8	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	2.4	*
8	Septicemia (A40-A41)	5	2.0	*
8	Diabetes mellitus (E10-E14)	5	2.0	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	5	2.0	*
11	Tuberculosis (A16-A19)	1	0.4	*
11	Viral hepatitis (B15-B19)	1	0.4	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
11	Anemias (D50-D64)	1	0.4	*
11	Nutritional deficiencies (E40-E64)	1	0.4	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
11	Aortic aneurysm and dissection (I71)	1	0.4	*
11	Influenza and pneumonia (J09-J18)	1	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
...	All other causes (Residual)	31	12.4	3.7

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-39 years				
...	All causes	306	100.0	38.6
1	Malignant neoplasms (C00-C97)	113	36.9	14.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	9.2	3.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	9.2	3.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	6.9	2.6
5	Cerebrovascular diseases (I60-I69)	13	4.2	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	4.2	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	12	3.9	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	7	2.3	*
9	Diabetes mellitus (E10-E14)	6	2.0	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.3	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	1.0	*
14	Tuberculosis (A16-A19)	2	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
...	All other causes (Residual)	46	15.0	5.8
Asian or Pacific Islander, female, 40-44 years				
...	All causes	508	100.0	64.5
1	Malignant neoplasms (C00-C97)	237	46.7	30.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	8.5	5.5
3	Cerebrovascular diseases (I60-I69)	41	8.1	5.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	6.3	4.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	4.5	2.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	10	2.0	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.8	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	1.8	*
9	Septicemia (A40-A41)	8	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	1.2	*
12	Diabetes mellitus (E10-E14)	5	1.0	*
12	Influenza and pneumonia (J09-J18)	5	1.0	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	5	1.0	*
15	Aortic aneurysm and dissection (I71)	3	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
...	All other causes (Residual)	60	11.8	7.6

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-49 years				
...	All causes	701	100.0	103.3
1	Malignant neoplasms (C00-C97)	353	50.4	52.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	9.1	9.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	6.1	6.3
4	Cerebrovascular diseases (I60-I69)	40	5.7	5.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	3.6	3.7
6	Diabetes mellitus (E10-E14)	22	3.1	3.2
7	Influenza and pneumonia (J09-J18)	11	1.6	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.0	*
10	Septicemia (A40-A41)	6	0.9	*
10	Viral hepatitis (B15-B19)	6	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.7	*
14	Aortic aneurysm and dissection (I71)	4	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.4	*
...	All other causes (Residual)	96	13.7	14.1
Asian or Pacific Islander, female, 50-54 years				
...	All causes	975	100.0	154.6
1	Malignant neoplasms (C00-C97)	480	49.2	76.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	100	10.3	15.9
3	Cerebrovascular diseases (I60-I69)	73	7.5	11.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56	5.7	8.9
5	Diabetes mellitus (E10-E14)	32	3.3	5.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	3.3	5.1
7	Influenza and pneumonia (J09-J18)	13	1.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.3	*
9	Viral hepatitis (B15-B19)	11	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.9	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.9	*
14	Septicemia (A40-A41)	8	0.8	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.5	*
...	All other causes (Residual)	114	11.7	18.1

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-59 years				
...	All causes	1,305	100.0	226.7
1	Malignant neoplasms (C00-C97)	662	50.7	115.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	13.0	29.5
3	Cerebrovascular diseases (I60-I69)	88	6.7	15.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	4.2	9.6
5	Diabetes mellitus (E10-E14)	51	3.9	8.9
6	Chronic lower respiratory diseases (J40-J47)	21	1.6	3.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	1.5	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.4	*
9	Viral hepatitis (B15-B19)	16	1.2	*
10	Influenza and pneumonia (J09-J18)	15	1.1	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	15	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.0	*
13	Septicemia (A40-A41)	10	0.8	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	0.6	*
15	Aortic aneurysm and dissection (I71)	6	0.5	*
...	All other causes (Residual)	138	10.6	24.0
Asian or Pacific Islander, female, 60-64 years				
...	All causes	1,751	100.0	358.5
1	Malignant neoplasms (C00-C97)	824	47.1	168.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	274	15.6	56.1
3	Cerebrovascular diseases (I60-I69)	106	6.1	21.7
4	Diabetes mellitus (E10-E14)	69	3.9	14.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	2.6	9.4
6	Influenza and pneumonia (J09-J18)	39	2.2	8.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.8	6.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	27	1.5	5.5
9	Chronic lower respiratory diseases (J40-J47)	26	1.5	5.3
10	Viral hepatitis (B15-B19)	19	1.1	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19	1.1	*
12	Septicemia (A40-A41)	18	1.0	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.6	*
15	Aortic aneurysm and dissection (I71)	9	0.5	*
...	All other causes (Residual)	216	12.3	44.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 65-69 years				
...	All causes	2,032	100.0	560.6
1	Malignant neoplasms (C00-C97)	870	42.8	240.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	344	16.9	94.9
3	Cerebrovascular diseases (I60-I69)	156	7.7	43.0
4	Diabetes mellitus (E10-E14)	82	4.0	22.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	2.9	16.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	2.0	11.3
7	Influenza and pneumonia (J09-J18)	38	1.9	10.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	1.7	9.7
9	Septicemia (A40-A41)	34	1.7	9.4
9	Chronic lower respiratory diseases (J40-J47)	34	1.7	9.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	25	1.2	6.9
12	Viral hepatitis (B15-B19)	18	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.9	*
14	Parkinson's disease (G20-G21)	12	0.6	*
15	Alzheimer's disease (G30)	11	0.5	*
15	Aortic aneurysm and dissection (I71)	11	0.5	*
...	All other causes (Residual)	244	12.0	67.3
Asian or Pacific Islander, female, 70-74 years				
...	All causes	2,297	100.0	909.8
1	Malignant neoplasms (C00-C97)	856	37.3	339.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	423	18.4	167.5
3	Cerebrovascular diseases (I60-I69)	176	7.7	69.7
4	Diabetes mellitus (E10-E14)	102	4.4	40.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	2.9	26.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	2.8	25.7
7	Chronic lower respiratory diseases (J40-J47)	60	2.6	23.8
8	Influenza and pneumonia (J09-J18)	52	2.3	20.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	1.6	14.7
10	Septicemia (A40-A41)	34	1.5	13.5
11	Alzheimer's disease (G30)	28	1.2	11.1
12	Parkinson's disease (G20-G21)	22	1.0	8.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	20	0.9	7.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.7	*
15	Aortic aneurysm and dissection (I71)	14	0.6	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	0.6	*
...	All other causes (Residual)	311	13.5	123.2

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-79 years				
...	All causes	3,169	100.0	1,738.2
1	Malignant neoplasms (C00-C97)	947	29.9	519.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	646	20.4	354.3
3	Cerebrovascular diseases (I60-I69)	265	8.4	145.4
4	Diabetes mellitus (E10-E14)	167	5.3	91.6
5	Chronic lower respiratory diseases (J40-J47)	102	3.2	55.9
6	Influenza and pneumonia (J09-J18)	95	3.0	52.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	2.8	49.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	2.4	41.7
9	Alzheimer's disease (G30)	69	2.2	37.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	55	1.7	30.2
11	Septicemia (A40-A41)	51	1.6	28.0
12	Parkinson's disease (G20-G21)	30	0.9	16.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	28	0.9	15.4
14	Viral hepatitis (B15-B19)	19	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.5	*
...	All other causes (Residual)	513	16.2	281.4
Asian or Pacific Islander, female, 80-84 years				
...	All causes	4,173	100.0	3,196.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	984	23.6	753.7
2	Malignant neoplasms (C00-C97)	935	22.4	716.2
3	Cerebrovascular diseases (I60-I69)	370	8.9	283.4
4	Diabetes mellitus (E10-E14)	191	4.6	146.3
5	Influenza and pneumonia (J09-J18)	172	4.1	131.7
6	Alzheimer's disease (G30)	159	3.8	121.8
7	Chronic lower respiratory diseases (J40-J47)	129	3.1	98.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	91	2.2	69.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	2.0	64.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	1.9	59.7
11	Parkinson's disease (G20-G21)	60	1.4	46.0
12	Septicemia (A40-A41)	58	1.4	44.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.7	23.7
14	Pneumonitis due to solids and liquids (J69)	24	0.6	18.4
15	Viral hepatitis (B15-B19)	23	0.6	17.6
...	All other causes (Residual)	784	18.8	600.5

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 85-89 years				
...	All causes	4,682	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,228	26.2	---
2	Malignant neoplasms (C00-C97)	715	15.3	---
3	Cerebrovascular diseases (I60-I69)	461	9.8	---
4	Alzheimer's disease (G30)	278	5.9	---
5	Influenza and pneumonia (J09-J18)	212	4.5	---
6	Diabetes mellitus (E10-E14)	184	3.9	---
7	Chronic lower respiratory diseases (J40-J47)	142	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	105	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	96	2.1	---
11	Parkinson's disease (G20-G21)	60	1.3	---
12	Septicemia (A40-A41)	55	1.2	---
13	Pneumonitis due to solids and liquids (J69)	51	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.7	---
15	Aortic aneurysm and dissection (I71)	21	0.4	---
...	All other causes (Residual)	931	19.9	---
Asian or Pacific Islander, female, 90-94 years				
...	All causes	3,669	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,051	28.6	---
2	Malignant neoplasms (C00-C97)	351	9.6	---
3	Cerebrovascular diseases (I60-I69)	332	9.0	---
4	Alzheimer's disease (G30)	256	7.0	---
5	Influenza and pneumonia (J09-J18)	235	6.4	---
6	Diabetes mellitus (E10-E14)	110	3.0	---
7	Chronic lower respiratory diseases (J40-J47)	109	3.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	107	2.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	2.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	1.7	---
11	Septicemia (A40-A41)	50	1.4	---
12	Pneumonitis due to solids and liquids (J69)	45	1.2	---
13	Parkinson's disease (G20-G21)	28	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.6	---
15	Aortic aneurysm and dissection (I71)	20	0.5	---
...	All other causes (Residual)	812	22.1	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2013

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 95-99 years				
...	All causes	1,699	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	494	29.1	---
2	Alzheimer's disease (G30)	141	8.3	---
3	Cerebrovascular diseases (I60-I69)	127	7.5	---
4	Malignant neoplasms (C00-C97)	94	5.5	---
4	Influenza and pneumonia (J09-J18)	94	5.5	---
6	Chronic lower respiratory diseases (J40-J47)	67	3.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	63	3.7	---
8	Diabetes mellitus (E10-E14)	47	2.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	1.9	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.5	---
11	Pneumonitis due to solids and liquids (J69)	20	1.2	---
12	Septicemia (A40-A41)	13	0.8	---
13	Parkinson's disease (G20-G21)	12	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.5	---
15	Atherosclerosis (I70)	7	0.4	---
...	All other causes (Residual)	453	26.7	---
Asian or Pacific Islander, female, 100 years and over				
...	All causes	401	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	109	27.2	---
2	Cerebrovascular diseases (I60-I69)	37	9.2	---
3	Alzheimer's disease (G30)	31	7.7	---
4	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	5.0	---
4	Influenza and pneumonia (J09-J18)	20	5.0	---
6	Chronic lower respiratory diseases (J40-J47)	15	3.7	---
7	Malignant neoplasms (C00-C97)	14	3.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	2.7	---
9	Atherosclerosis (I70)	7	1.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.7	---
11	Diabetes mellitus (E10-E14)	6	1.5	---
11	Pneumonitis due to solids and liquids (J69)	6	1.5	---
13	Nutritional deficiencies (E40-E64)	4	1.0	---
14	Parkinson's disease (G20-G21)	2	0.5	---
15	Aortic aneurysm and dissection (I71)	1	0.2	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	---
...	All other causes (Residual)	110	27.4	---

... Category not applicable

* Figure does not meet standards of reliability or precision

--- Data not available

¹ Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013