

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	135,519	100.0	297.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,021	21.4	63.8
2	Malignant neoplasms (C00-C97)	27,660	20.4	60.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,723	8.7	25.8
4	Cerebrovascular diseases (I60-I69)	7,078	5.2	15.6
5	Diabetes mellitus (E10-E14)	6,417	4.7	14.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,913	2.9	8.6
7	Chronic lower respiratory diseases (J40-J47)	3,531	2.6	7.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,466	2.6	7.6
9	Certain conditions originating in the perinatal period (P00-P96)	2,946	2.2	6.5
10	Influenza and pneumonia (J09-J18)	2,735	2.0	6.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,691	2.0	5.9
12	Alzheimer's disease (G30)	2,471	1.8	5.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,465	1.8	5.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,991	1.5	4.4
15	Septicemia (A40-A41)	1,894	1.4	4.2
...	All other causes (Residual)	25,517	18.8	56.1
Hispanic, both sexes, 1-4 years				
...	All causes	1,009	100.0	26.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	348	34.5	9.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	132	13.1	3.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	76	7.5	2.0
4	Malignant neoplasms (C00-C97)	68	6.7	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	3.9	1.0
6	Influenza and pneumonia (J09-J18)	24	2.4	0.6
7	Septicemia (A40-A41)	18	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	16	1.6	*
10	Cerebrovascular diseases (I60-I69)	8	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
12	Pneumonitis due to solids and liquids (J69)	5	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
15	Anemias (D50-D64)	2	0.2	*
15	Meningitis (G00,G03)	2	0.2	*
15	Diseases of appendix (K35-K38)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	236	23.4	6.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	566	100.0	13.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	35.2	4.7
2	Malignant neoplasms (C00-C97)	117	20.7	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	7.8	1.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.1	0.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	4.1	0.5
6	Septicemia (A40-A41)	11	1.9	*
6	Chronic lower respiratory diseases (J40-J47)	11	1.9	*
8	Influenza and pneumonia (J09-J18)	10	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.6	*
10	Cerebrovascular diseases (I60-I69)	7	1.2	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.7	*
12	Anemias (D50-D64)	3	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	Shigellosis and amebiasis (A03,A06)	1	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
...	All other causes (Residual)	96	17.0	2.3
Hispanic, both sexes, 10-14 years				
...	All causes	591	100.0	14.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	31.8	4.7
2	Malignant neoplasms (C00-C97)	111	18.8	2.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	9.0	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	4.9	0.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	3.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.9	*
7	Influenza and pneumonia (J09-J18)	10	1.7	*
8	Cerebrovascular diseases (I60-I69)	9	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	6	1.0	*
11	Septicemia (A40-A41)	5	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
13	Meningitis (G00,G03)	3	0.5	*
14	Anemias (D50-D64)	2	0.3	*
14	Diabetes mellitus (E10-E14)	2	0.3	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	121	20.5	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	2,169	100.0	57.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	934	43.1	24.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	524	24.2	14.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	215	9.9	5.7
4	Malignant neoplasms (C00-C97)	138	6.4	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	2.2	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.4	0.8
7	Cerebrovascular diseases (I60-I69)	12	0.6	*
7	Chronic lower respiratory diseases (J40-J47)	12	0.6	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	7	0.3	*
10	Septicemia (A40-A41)	6	0.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.3	*
10	Diabetes mellitus (E10-E14)	6	0.3	*
13	Anemias (D50-D64)	5	0.2	*
13	Legal intervention (Y35,Y89.0)	5	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.2	*
...	All other causes (Residual)	217	10.0	5.8
Hispanic, both sexes, 20-24 years				
...	All causes	3,517	100.0	95.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,577	44.8	42.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	771	21.9	20.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	312	8.9	8.5
4	Malignant neoplasms (C00-C97)	207	5.9	5.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	2.8	2.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.8	0.8
7	Cerebrovascular diseases (I60-I69)	25	0.7	0.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	0.7	0.7
9	Pregnancy, childbirth and the puerperium (O00-O99)	21	0.6	0.6
10	Legal intervention (Y35,Y89.0)	19	0.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
12	Diabetes mellitus (E10-E14)	12	0.3	*
13	Septicemia (A40-A41)	11	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.3	*
15	Influenza and pneumonia (J09-J18)	10	0.3	*
...	All other causes (Residual)	377	10.7	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	3,315	100.0	79.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,285	38.8	30.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	579	17.5	13.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	342	10.3	8.2
4	Malignant neoplasms (C00-C97)	229	6.9	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	169	5.1	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	64	1.9	1.5
7	Cerebrovascular diseases (I60-I69)	38	1.1	0.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	28	0.8	0.7
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.8	0.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	27	0.8	0.6
11	Diabetes mellitus (E10-E14)	21	0.6	0.5
11	Chronic lower respiratory diseases (J40-J47)	21	0.6	0.5
11	Legal intervention (Y35,Y89.0)	21	0.6	0.5
14	Influenza and pneumonia (J09-J18)	15	0.5	*
15	Septicemia (A40-A41)	13	0.4	*
...	All other causes (Residual)	435	13.1	10.4
Hispanic, both sexes, 30-34 years				
...	All causes	3,231	100.0	80.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,036	32.1	25.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	449	13.9	11.2
3	Malignant neoplasms (C00-C97)	380	11.8	9.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	307	9.5	7.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	209	6.5	5.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	118	3.7	3.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	75	2.3	1.9
8	Cerebrovascular diseases (I60-I69)	62	1.9	1.6
9	Diabetes mellitus (E10-E14)	38	1.2	1.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	31	1.0	0.8
11	Septicemia (A40-A41)	28	0.9	0.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.9	0.7
13	Influenza and pneumonia (J09-J18)	23	0.7	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.6	*
15	Legal intervention (Y35,Y89.0)	15	0.5	*
...	All other causes (Residual)	413	12.8	10.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,768	100.0	103.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	970	25.7	26.6
2	Malignant neoplasms (C00-C97)	582	15.4	16.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	369	9.8	10.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	284	7.5	7.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	245	6.5	6.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	181	4.8	5.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	153	4.1	4.2
8	Cerebrovascular diseases (I60-I69)	118	3.1	3.2
9	Diabetes mellitus (E10-E14)	98	2.6	2.7
10	Septicemia (A40-A41)	36	1.0	1.0
11	Influenza and pneumonia (J09-J18)	34	0.9	0.9
12	Pregnancy, childbirth and the puerperium (O00-O99)	25	0.7	0.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.6	0.7
14	Viral hepatitis (B15-B19)	23	0.6	0.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.6	0.6
...	All other causes (Residual)	603	16.0	16.6
Hispanic, both sexes, 40-44 years				
...	All causes	4,927	100.0	154.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	988	20.1	30.9
2	Malignant neoplasms (C00-C97)	903	18.3	28.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	593	12.0	18.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	324	6.6	10.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	309	6.3	9.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	254	5.2	7.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	218	4.4	6.8
8	Cerebrovascular diseases (I60-I69)	183	3.7	5.7
9	Diabetes mellitus (E10-E14)	130	2.6	4.1
10	Viral hepatitis (B15-B19)	78	1.6	2.4
11	Influenza and pneumonia (J09-J18)	51	1.0	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.0	1.6
13	Septicemia (A40-A41)	45	0.9	1.4
14	Chronic lower respiratory diseases (J40-J47)	25	0.5	0.8
15	Aortic aneurysm and dissection (I71)	21	0.4	0.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.4	0.7
...	All other causes (Residual)	734	14.9	22.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	6,552	100.0	245.0
1	Malignant neoplasms (C00-C97)	1,514	23.1	56.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	959	14.6	35.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	915	14.0	34.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	519	7.9	19.4
5	Cerebrovascular diseases (I60-I69)	300	4.6	11.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	290	4.4	10.8
7	Diabetes mellitus (E10-E14)	233	3.6	8.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	204	3.1	7.6
9	Viral hepatitis (B15-B19)	173	2.6	6.5
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	153	2.3	5.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	1.3	3.3
12	Septicemia (A40-A41)	85	1.3	3.2
13	Chronic lower respiratory diseases (J40-J47)	61	0.9	2.3
14	Influenza and pneumonia (J09-J18)	46	0.7	1.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	0.6	1.6
...	All other causes (Residual)	970	14.8	36.3
Hispanic, both sexes, 50-54 years				
...	All causes	7,848	100.0	378.2
1	Malignant neoplasms (C00-C97)	2,040	26.0	98.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,358	17.3	65.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	744	9.5	35.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	648	8.3	31.2
5	Diabetes mellitus (E10-E14)	398	5.1	19.2
6	Cerebrovascular diseases (I60-I69)	356	4.5	17.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	225	2.9	10.8
8	Viral hepatitis (B15-B19)	219	2.8	10.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	2.4	9.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	142	1.8	6.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	1.3	4.8
12	Septicemia (A40-A41)	97	1.2	4.7
13	Influenza and pneumonia (J09-J18)	84	1.1	4.0
14	Chronic lower respiratory diseases (J40-J47)	80	1.0	3.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	70	0.9	3.4
...	All other causes (Residual)	1,100	14.0	53.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	8,464	100.0	539.5
1	Malignant neoplasms (C00-C97)	2,541	30.0	162.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,696	20.0	108.1
3	Diabetes mellitus (E10-E14)	505	6.0	32.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	493	5.8	31.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	478	5.6	30.5
6	Cerebrovascular diseases (I60-I69)	452	5.3	28.8
7	Viral hepatitis (B15-B19)	233	2.8	14.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	172	2.0	11.0
9	Septicemia (A40-A41)	131	1.5	8.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	1.5	7.8
11	Chronic lower respiratory diseases (J40-J47)	122	1.4	7.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	113	1.3	7.2
13	Influenza and pneumonia (J09-J18)	99	1.2	6.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	78	0.9	5.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	0.7	3.8
...	All other causes (Residual)	1,168	13.8	74.5
Hispanic, both sexes, 60-64 years				
...	All causes	8,960	100.0	798.2
1	Malignant neoplasms (C00-C97)	2,785	31.1	248.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,055	22.9	183.1
3	Diabetes mellitus (E10-E14)	626	7.0	55.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	461	5.1	41.1
5	Cerebrovascular diseases (I60-I69)	459	5.1	40.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	317	3.5	28.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	199	2.2	17.7
8	Chronic lower respiratory diseases (J40-J47)	175	2.0	15.6
9	Influenza and pneumonia (J09-J18)	127	1.4	11.3
10	Viral hepatitis (B15-B19)	125	1.4	11.1
11	Septicemia (A40-A41)	122	1.4	10.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	92	1.0	8.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	79	0.9	7.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	0.8	6.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.5	3.7
...	All other causes (Residual)	1,228	13.7	109.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	9,806	100.0	1,211.9
1	Malignant neoplasms (C00-C97)	3,176	32.4	392.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,274	23.2	281.0
3	Diabetes mellitus (E10-E14)	727	7.4	89.8
4	Cerebrovascular diseases (I60-I69)	526	5.4	65.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	363	3.7	44.9
6	Chronic lower respiratory diseases (J40-J47)	271	2.8	33.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	270	2.8	33.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	252	2.6	31.1
9	Septicemia (A40-A41)	163	1.7	20.1
10	Influenza and pneumonia (J09-J18)	152	1.6	18.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	118	1.2	14.6
12	Viral hepatitis (B15-B19)	86	0.9	10.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	0.6	6.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	50	0.5	6.2
15	Alzheimer's disease (G30)	47	0.5	5.8
...	All other causes (Residual)	1,277	13.0	157.8
Hispanic, both sexes, 70-74 years				
...	All causes	11,321	100.0	1,822.6
1	Malignant neoplasms (C00-C97)	3,296	29.1	530.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,759	24.4	444.2
3	Diabetes mellitus (E10-E14)	808	7.1	130.1
4	Cerebrovascular diseases (I60-I69)	646	5.7	104.0
5	Chronic lower respiratory diseases (J40-J47)	374	3.3	60.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	341	3.0	54.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	337	3.0	54.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	275	2.4	44.3
9	Influenza and pneumonia (J09-J18)	241	2.1	38.8
10	Septicemia (A40-A41)	212	1.9	34.1
11	Alzheimer's disease (G30)	134	1.2	21.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	1.1	19.5
13	Parkinson's disease (G20-G21)	78	0.7	12.6
14	Viral hepatitis (B15-B19)	71	0.6	11.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.5	10.0
...	All other causes (Residual)	1,566	13.8	252.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	14,185	100.0	2,959.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,735	26.3	779.3
2	Malignant neoplasms (C00-C97)	3,570	25.2	744.8
3	Diabetes mellitus (E10-E14)	972	6.9	202.8
4	Cerebrovascular diseases (I60-I69)	929	6.5	193.8
5	Chronic lower respiratory diseases (J40-J47)	602	4.2	125.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	406	2.9	84.7
7	Influenza and pneumonia (J09-J18)	336	2.4	70.1
8	Alzheimer's disease (G30)	315	2.2	65.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	296	2.1	61.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	271	1.9	56.5
11	Septicemia (A40-A41)	225	1.6	46.9
12	Parkinson's disease (G20-G21)	181	1.3	37.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	160	1.1	33.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	91	0.6	19.0
15	Pneumonitis due to solids and liquids (J69)	67	0.5	14.0
...	All other causes (Residual)	2,029	14.3	423.3
Hispanic, both sexes, 80-84 years				
...	All causes	15,305	100.0	4,746.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,379	28.6	1,358.2
2	Malignant neoplasms (C00-C97)	3,113	20.3	965.5
3	Cerebrovascular diseases (I60-I69)	1,093	7.1	339.0
4	Diabetes mellitus (E10-E14)	892	5.8	276.7
5	Chronic lower respiratory diseases (J40-J47)	719	4.7	223.0
6	Alzheimer's disease (G30)	552	3.6	171.2
7	Influenza and pneumonia (J09-J18)	470	3.1	145.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	381	2.5	118.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	320	2.1	99.2
10	Septicemia (A40-A41)	273	1.8	84.7
11	Parkinson's disease (G20-G21)	230	1.5	71.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	225	1.5	69.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	149	1.0	46.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	107	0.7	33.2
15	Pneumonitis due to solids and liquids (J69)	88	0.6	27.3
...	All other causes (Residual)	2,314	15.1	717.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	12,498	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,998	32.0	---
2	Malignant neoplasms (C00-C97)	1,877	15.0	---
3	Cerebrovascular diseases (I60-I69)	1,017	8.1	---
4	Alzheimer's disease (G30)	626	5.0	---
5	Diabetes mellitus (E10-E14)	602	4.8	---
6	Chronic lower respiratory diseases (J40-J47)	562	4.5	---
7	Influenza and pneumonia (J09-J18)	437	3.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	289	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	196	1.6	---
11	Septicemia (A40-A41)	187	1.5	---
12	Parkinson's disease (G20-G21)	120	1.0	---
13	Pneumonitis due to solids and liquids (J69)	97	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.7	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	77	0.6	---
...	All other causes (Residual)	2,106	16.9	---
Hispanic, both sexes, 90-94 years				
...	All causes	7,497	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,646	35.3	---
2	Malignant neoplasms (C00-C97)	768	10.2	---
3	Cerebrovascular diseases (I60-I69)	528	7.0	---
4	Alzheimer's disease (G30)	497	6.6	---
5	Chronic lower respiratory diseases (J40-J47)	315	4.2	---
6	Influenza and pneumonia (J09-J18)	308	4.1	---
7	Diabetes mellitus (E10-E14)	246	3.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	161	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	154	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	1.8	---
11	Septicemia (A40-A41)	113	1.5	---
12	Parkinson's disease (G20-G21)	76	1.0	---
13	Pneumonitis due to solids and liquids (J69)	67	0.9	---
14	Atherosclerosis (I70)	49	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.5	---
...	All other causes (Residual)	1,400	18.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	3,164	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,226	38.7	---
2	Cerebrovascular diseases (I60-I69)	231	7.3	---
3	Alzheimer's disease (G30)	209	6.6	---
4	Malignant neoplasms (C00-C97)	200	6.3	---
5	Influenza and pneumonia (J09-J18)	162	5.1	---
6	Chronic lower respiratory diseases (J40-J47)	106	3.4	---
7	Diabetes mellitus (E10-E14)	81	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	75	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	1.7	---
11	Septicemia (A40-A41)	38	1.2	---
12	Pneumonitis due to solids and liquids (J69)	32	1.0	---
13	Atherosclerosis (I70)	22	0.7	---
14	Parkinson's disease (G20-G21)	16	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.3	---
...	All other causes (Residual)	636	20.1	---
Hispanic, both sexes, 100 years and over				
...	All causes	731	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	279	38.2	---
2	Cerebrovascular diseases (I60-I69)	50	6.8	---
3	Influenza and pneumonia (J09-J18)	45	6.2	---
4	Alzheimer's disease (G30)	40	5.5	---
5	Malignant neoplasms (C00-C97)	26	3.6	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	2.3	---
7	Chronic lower respiratory diseases (J40-J47)	16	2.2	---
8	Diabetes mellitus (E10-E14)	15	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	1.9	---
9	Pneumonitis due to solids and liquids (J69)	14	1.9	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	1.9	---
12	Atherosclerosis (I70)	11	1.5	---
13	Septicemia (A40-A41)	9	1.2	---
14	Anemias (D50-D64)	7	1.0	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	5	0.7	---
...	All other causes (Residual)	169	23.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	75,708	100.0	321.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,657	20.7	66.6
2	Malignant neoplasms (C00-C97)	14,493	19.1	61.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,844	11.7	37.6
4	Cerebrovascular diseases (I60-I69)	3,319	4.4	14.1
5	Diabetes mellitus (E10-E14)	3,199	4.2	13.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,926	3.9	12.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,799	3.7	11.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,078	2.7	8.8
9	Chronic lower respiratory diseases (J40-J47)	1,894	2.5	8.1
10	Certain conditions originating in the perinatal period (P00-P96)	1,643	2.2	7.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,363	1.8	5.8
12	Influenza and pneumonia (J09-J18)	1,361	1.8	5.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,180	1.6	5.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,063	1.4	4.5
15	Septicemia (A40-A41)	944	1.2	4.0
...	All other causes (Residual)	12,945	17.1	55.0
Hispanic, male, 1-4 years				
...	All causes	556	100.0	28.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	225	40.5	11.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	76	13.7	3.8
3	Malignant neoplasms (C00-C97)	38	6.8	1.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	5.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.1	*
6	Influenza and pneumonia (J09-J18)	13	2.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.3	*
8	Septicemia (A40-A41)	6	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	0.9	*
10	Cerebrovascular diseases (I60-I69)	4	0.7	*
10	Pneumonitis due to solids and liquids (J69)	4	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
13	Meningococcal infection (A39)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Diseases of appendix (K35-K38)	1	0.2	*
13	Hernia (K40-K46)	1	0.2	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	118	21.2	5.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	314	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	38.9	5.7
2	Malignant neoplasms (C00-C97)	61	19.4	2.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	6.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	4.8	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	2.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	7	2.2	*
8	Septicemia (A40-A41)	5	1.6	*
9	Cerebrovascular diseases (I60-I69)	4	1.3	*
9	Influenza and pneumonia (J09-J18)	4	1.3	*
11	Anemias (D50-D64)	3	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
13	Shigellosis and amebiasis (A03,A06)	1	0.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
13	Diabetes mellitus (E10-E14)	1	0.3	*
13	Pneumonitis due to solids and liquids (J69)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	51	16.2	2.4
Hispanic, male, 10-14 years				
...	All causes	349	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	35.0	6.0
2	Malignant neoplasms (C00-C97)	53	15.2	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	11.2	1.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	4.6	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	3.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.3	*
7	Influenza and pneumonia (J09-J18)	7	2.0	*
8	Septicemia (A40-A41)	5	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.1	*
9	Cerebrovascular diseases (I60-I69)	4	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.9	*
13	Anemias (D50-D64)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Aortic aneurysm and dissection (I71)	1	0.3	*
13	Legal intervention (Y35,Y89.0)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	67	19.2	3.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,674	100.0	86.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	713	42.6	37.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	474	28.3	24.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	181	10.8	9.4
4	Malignant neoplasms (C00-C97)	86	5.1	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	1.8	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	1.3	1.1
7	Chronic lower respiratory diseases (J40-J47)	6	0.4	*
8	Diabetes mellitus (E10-E14)	5	0.3	*
8	Legal intervention (Y35,Y89.0)	5	0.3	*
10	Cerebrovascular diseases (I60-I69)	4	0.2	*
11	Septicemia (A40-A41)	3	0.2	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.2	*
13	Meningococcal infection (A39)	2	0.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.1	*
15	Viral hepatitis (B15-B19)	1	0.1	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.1	*
15	Anemias (D50-D64)	1	0.1	*
15	Aortic aneurysm and dissection (I71)	1	0.1	*
15	Influenza and pneumonia (J09-J18)	1	0.1	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.1	*
15	Pneumonitis due to solids and liquids (J69)	1	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.1	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.1	*
...	All other causes (Residual)	130	7.8	6.7
Hispanic, male, 20-24 years				
...	All causes	2,832	100.0	143.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,322	46.7	66.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	697	24.6	35.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	270	9.5	13.6
4	Malignant neoplasms (C00-C97)	123	4.3	6.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	2.6	3.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.8	1.1
7	Legal intervention (Y35,Y89.0)	19	0.7	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.6	*
9	Cerebrovascular diseases (I60-I69)	16	0.6	*
10	Diabetes mellitus (E10-E14)	9	0.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
12	Septicemia (A40-A41)	7	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.2	*
14	Aortic aneurysm and dissection (I71)	5	0.2	*
14	Influenza and pneumonia (J09-J18)	5	0.2	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.2	*
...	All other causes (Residual)	226	8.0	11.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,575	100.0	110.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,077	41.8	46.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	509	19.8	21.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	280	10.9	12.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	123	4.8	5.3
5	Malignant neoplasms (C00-C97)	122	4.7	5.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	50	1.9	2.1
7	Cerebrovascular diseases (I60-I69)	24	0.9	1.0
8	Legal intervention (Y35,Y89.0)	21	0.8	0.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	20	0.8	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.8	0.9
11	Chronic lower respiratory diseases (J40-J47)	15	0.6	*
12	Diabetes mellitus (E10-E14)	12	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.3	*
14	Influenza and pneumonia (J09-J18)	8	0.3	*
15	Septicemia (A40-A41)	6	0.2	*
...	All other causes (Residual)	279	10.8	12.0
Hispanic, male, 30-34 years				
...	All causes	2,381	100.0	109.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	862	36.2	39.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	389	16.3	17.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	268	11.3	12.3
4	Malignant neoplasms (C00-C97)	160	6.7	7.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	154	6.5	7.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	85	3.6	3.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	61	2.6	2.8
8	Cerebrovascular diseases (I60-I69)	36	1.5	1.7
9	Diabetes mellitus (E10-E14)	27	1.1	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.7	*
11	Legal intervention (Y35,Y89.0)	15	0.6	*
12	Influenza and pneumonia (J09-J18)	14	0.6	*
13	Septicemia (A40-A41)	13	0.5	*
14	Aortic aneurysm and dissection (I71)	10	0.4	*
15	Viral hepatitis (B15-B19)	9	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	9	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.4	*
...	All other causes (Residual)	243	10.2	11.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,624	100.0	134.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	776	29.6	39.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	269	10.3	13.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	236	9.0	12.1
4	Malignant neoplasms (C00-C97)	235	9.0	12.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	203	7.7	10.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	149	5.7	7.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	124	4.7	6.4
8	Cerebrovascular diseases (I60-I69)	74	2.8	3.8
9	Diabetes mellitus (E10-E14)	70	2.7	3.6
10	Influenza and pneumonia (J09-J18)	23	0.9	1.2
11	Septicemia (A40-A41)	22	0.8	1.1
12	Viral hepatitis (B15-B19)	18	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.7	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	0.5	*
...	All other causes (Residual)	379	14.4	19.5
Hispanic, male, 40-44 years				
...	All causes	3,413	100.0	202.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	782	22.9	46.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	474	13.9	28.2
3	Malignant neoplasms (C00-C97)	398	11.7	23.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	258	7.6	15.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	247	7.2	14.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	214	6.3	12.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	170	5.0	10.1
8	Cerebrovascular diseases (I60-I69)	120	3.5	7.1
9	Diabetes mellitus (E10-E14)	82	2.4	4.9
10	Viral hepatitis (B15-B19)	61	1.8	3.6
11	Influenza and pneumonia (J09-J18)	36	1.1	2.1
12	Septicemia (A40-A41)	27	0.8	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.8	1.6
14	Aortic aneurysm and dissection (I71)	19	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	15	0.4	*
...	All other causes (Residual)	483	14.2	28.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,441	100.0	323.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	755	17.0	54.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	721	16.2	52.4
3	Malignant neoplasms (C00-C97)	718	16.2	52.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	420	9.5	30.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	226	5.1	16.4
6	Cerebrovascular diseases (I60-I69)	189	4.3	13.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	165	3.7	12.0
8	Diabetes mellitus (E10-E14)	159	3.6	11.6
9	Viral hepatitis (B15-B19)	136	3.1	9.9
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	128	2.9	9.3
11	Septicemia (A40-A41)	57	1.3	4.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	1.1	3.5
13	Chronic lower respiratory diseases (J40-J47)	36	0.8	2.6
14	Influenza and pneumonia (J09-J18)	27	0.6	2.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	0.5	1.7
...	All other causes (Residual)	632	14.2	46.0
Hispanic, male, 50-54 years				
...	All causes	5,193	100.0	500.0
1	Malignant neoplasms (C00-C97)	1,073	20.7	103.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,015	19.5	97.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	594	11.4	57.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	509	9.8	49.0
5	Diabetes mellitus (E10-E14)	224	4.3	21.6
6	Cerebrovascular diseases (I60-I69)	210	4.0	20.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	182	3.5	17.5
8	Viral hepatitis (B15-B19)	153	2.9	14.7
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	2.9	14.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	1.8	9.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	1.6	7.8
12	Septicemia (A40-A41)	55	1.1	5.3
13	Influenza and pneumonia (J09-J18)	52	1.0	5.0
14	Chronic lower respiratory diseases (J40-J47)	48	0.9	4.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	0.8	4.1
...	All other causes (Residual)	706	13.6	68.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	5,313	100.0	694.7
1	Malignant neoplasms (C00-C97)	1,324	24.9	173.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,225	23.1	160.2
3	Chronic liver disease and cirrhosis (K70,K73-K74)	387	7.3	50.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	6.4	44.6
5	Diabetes mellitus (E10-E14)	303	5.7	39.6
6	Cerebrovascular diseases (I60-I69)	290	5.5	37.9
7	Viral hepatitis (B15-B19)	174	3.3	22.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	103	1.9	13.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	92	1.7	12.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	81	1.5	10.6
11	Septicemia (A40-A41)	71	1.3	9.3
11	Chronic lower respiratory diseases (J40-J47)	71	1.3	9.3
13	Influenza and pneumonia (J09-J18)	67	1.3	8.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	0.9	6.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	0.8	5.8
...	All other causes (Residual)	691	13.0	90.3
Hispanic, male, 60-64 years				
...	All causes	5,451	100.0	1,028.9
1	Malignant neoplasms (C00-C97)	1,549	28.4	292.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,389	25.5	262.2
3	Diabetes mellitus (E10-E14)	358	6.6	67.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	336	6.2	63.4
5	Cerebrovascular diseases (I60-I69)	266	4.9	50.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	3.9	40.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	2.1	21.7
8	Chronic lower respiratory diseases (J40-J47)	102	1.9	19.3
9	Viral hepatitis (B15-B19)	85	1.6	16.0
10	Influenza and pneumonia (J09-J18)	76	1.4	14.3
11	Septicemia (A40-A41)	73	1.3	13.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	64	1.2	12.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	1.1	11.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	57	1.0	10.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.5	5.1
...	All other causes (Residual)	680	12.5	128.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	5,764	100.0	1,556.5
1	Malignant neoplasms (C00-C97)	1,805	31.3	487.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,468	25.5	396.4
3	Diabetes mellitus (E10-E14)	414	7.2	111.8
4	Cerebrovascular diseases (I60-I69)	279	4.8	75.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	238	4.1	64.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	175	3.0	47.3
7	Chronic lower respiratory diseases (J40-J47)	147	2.6	39.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	2.4	37.5
9	Influenza and pneumonia (J09-J18)	90	1.6	24.3
10	Septicemia (A40-A41)	86	1.5	23.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	63	1.1	17.0
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	0.9	13.2
13	Viral hepatitis (B15-B19)	45	0.8	12.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	42	0.7	11.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.5	7.6
...	All other causes (Residual)	696	12.1	187.9
Hispanic, male, 70-74 years				
...	All causes	6,214	100.0	2,278.5
1	Malignant neoplasms (C00-C97)	1,798	28.9	659.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,599	25.7	586.3
3	Diabetes mellitus (E10-E14)	404	6.5	148.1
4	Cerebrovascular diseases (I60-I69)	329	5.3	120.6
5	Chronic lower respiratory diseases (J40-J47)	240	3.9	88.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	203	3.3	74.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	2.8	64.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	172	2.8	63.1
9	Influenza and pneumonia (J09-J18)	126	2.0	46.2
10	Septicemia (A40-A41)	116	1.9	42.5
11	Alzheimer's disease (G30)	67	1.1	24.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	1.1	24.6
13	Parkinson's disease (G20-G21)	46	0.7	16.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	0.6	14.3
15	Viral hepatitis (B15-B19)	37	0.6	13.6
...	All other causes (Residual)	794	12.8	291.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	7,280	100.0	3,609.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,981	27.2	982.1
2	Malignant neoplasms (C00-C97)	1,936	26.6	959.8
3	Diabetes mellitus (E10-E14)	440	6.0	218.1
4	Cerebrovascular diseases (I60-I69)	431	5.9	213.7
5	Chronic lower respiratory diseases (J40-J47)	341	4.7	169.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	198	2.7	98.2
7	Influenza and pneumonia (J09-J18)	176	2.4	87.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	2.3	84.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	140	1.9	69.4
10	Alzheimer's disease (G30)	119	1.6	59.0
11	Septicemia (A40-A41)	113	1.6	56.0
12	Parkinson's disease (G20-G21)	110	1.5	54.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	64	0.9	31.7
14	Pneumonitis due to solids and liquids (J69)	44	0.6	21.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.6	21.3
...	All other causes (Residual)	974	13.4	482.9
Hispanic, male, 80-84 years				
...	All causes	7,158	100.0	5,546.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,099	29.3	1,626.3
2	Malignant neoplasms (C00-C97)	1,612	22.5	1,249.0
3	Cerebrovascular diseases (I60-I69)	441	6.2	341.7
4	Chronic lower respiratory diseases (J40-J47)	377	5.3	292.1
5	Diabetes mellitus (E10-E14)	370	5.2	286.7
6	Influenza and pneumonia (J09-J18)	243	3.4	188.3
7	Alzheimer's disease (G30)	196	2.7	151.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	193	2.7	149.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	2.3	128.6
10	Parkinson's disease (G20-G21)	135	1.9	104.6
11	Septicemia (A40-A41)	131	1.8	101.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	96	1.3	74.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	55	0.8	42.6
14	Pneumonitis due to solids and liquids (J69)	43	0.6	33.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.6	31.8
...	All other causes (Residual)	960	13.4	743.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	5,233	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,646	31.5	---
2	Malignant neoplasms (C00-C97)	960	18.3	---
3	Cerebrovascular diseases (I60-I69)	377	7.2	---
4	Chronic lower respiratory diseases (J40-J47)	274	5.2	---
5	Diabetes mellitus (E10-E14)	225	4.3	---
6	Alzheimer's disease (G30)	203	3.9	---
7	Influenza and pneumonia (J09-J18)	188	3.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	1.5	---
11	Septicemia (A40-A41)	68	1.3	---
12	Parkinson's disease (G20-G21)	59	1.1	---
13	Pneumonitis due to solids and liquids (J69)	54	1.0	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	35	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.6	---
...	All other causes (Residual)	806	15.4	---
Hispanic, male, 90-94 years				
...	All causes	2,499	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	865	34.6	---
2	Malignant neoplasms (C00-C97)	344	13.8	---
3	Cerebrovascular diseases (I60-I69)	147	5.9	---
4	Chronic lower respiratory diseases (J40-J47)	127	5.1	---
5	Alzheimer's disease (G30)	123	4.9	---
6	Influenza and pneumonia (J09-J18)	105	4.2	---
7	Diabetes mellitus (E10-E14)	71	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	48	1.9	---
11	Parkinson's disease (G20-G21)	36	1.4	---
12	Septicemia (A40-A41)	34	1.4	---
13	Pneumonitis due to solids and liquids (J69)	30	1.2	---
14	Aortic aneurysm and dissection (I71)	18	0.7	---
15	Atherosclerosis (I70)	15	0.6	---
...	All other causes (Residual)	410	16.4	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	886	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	340	38.4	---
2	Malignant neoplasms (C00-C97)	80	9.0	---
3	Influenza and pneumonia (J09-J18)	53	6.0	---
4	Alzheimer's disease (G30)	49	5.5	---
4	Cerebrovascular diseases (I60-I69)	49	5.5	---
6	Chronic lower respiratory diseases (J40-J47)	46	5.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	2.6	---
8	Diabetes mellitus (E10-E14)	19	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	2.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	2.0	---
11	Pneumonitis due to solids and liquids (J69)	11	1.2	---
12	Septicemia (A40-A41)	9	1.0	---
13	Parkinson's disease (G20-G21)	8	0.9	---
14	Atherosclerosis (I70)	6	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	---
...	All other causes (Residual)	153	17.3	---
Hispanic, male, 100 years and over				
...	All causes	191	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	32.5	---
2	Influenza and pneumonia (J09-J18)	18	9.4	---
3	Cerebrovascular diseases (I60-I69)	12	6.3	---
4	Malignant neoplasms (C00-C97)	11	5.8	---
5	Alzheimer's disease (G30)	9	4.7	---
6	Chronic lower respiratory diseases (J40-J47)	7	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	3.1	---
8	Anemias (D50-D64)	4	2.1	---
8	Diabetes mellitus (E10-E14)	4	2.1	---
8	Pneumonitis due to solids and liquids (J69)	4	2.1	---
11	Cholelithiasis and other disorders of gallbladder (K80-K82)	3	1.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.6	---
13	Parkinson's disease (G20-G21)	2	1.0	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	1.0	---
13	Atherosclerosis (I70)	2	1.0	---
...	All other causes (Residual)	42	22.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	59,811	100.0	272.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,364	22.3	60.8
2	Malignant neoplasms (C00-C97)	13,167	22.0	59.9
3	Cerebrovascular diseases (I60-I69)	3,759	6.3	17.1
4	Diabetes mellitus (E10-E14)	3,218	5.4	14.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,879	4.8	13.1
6	Alzheimer's disease (G30)	1,670	2.8	7.6
7	Chronic lower respiratory diseases (J40-J47)	1,637	2.7	7.4
8	Influenza and pneumonia (J09-J18)	1,374	2.3	6.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,328	2.2	6.0
10	Certain conditions originating in the perinatal period (P00-P96)	1,303	2.2	5.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,114	1.9	5.1
12	Septicemia (A40-A41)	950	1.6	4.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	928	1.6	4.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	756	1.3	3.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	540	0.9	2.5
...	All other causes (Residual)	11,824	19.8	53.8
Hispanic, female, 1-4 years				
...	All causes	453	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	27.2	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	12.4	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	10.2	2.4
4	Malignant neoplasms (C00-C97)	30	6.6	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.9	1.2
6	Septicemia (A40-A41)	12	2.6	*
7	Influenza and pneumonia (J09-J18)	11	2.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	11	2.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	2.2	*
10	Cerebrovascular diseases (I60-I69)	4	0.9	*
10	Acute bronchitis and bronchiolitis (J20-J21)	4	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Anemias (D50-D64)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
...	All other causes (Residual)	110	24.3	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	252	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	30.6	3.7
2	Malignant neoplasms (C00-C97)	56	22.2	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	9.9	1.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	5.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	3.2	*
6	Septicemia (A40-A41)	6	2.4	*
6	Influenza and pneumonia (J09-J18)	6	2.4	*
8	Chronic lower respiratory diseases (J40-J47)	4	1.6	*
9	Cerebrovascular diseases (I60-I69)	3	1.2	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	3	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
12	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (Residual)	45	17.9	2.2
Hispanic, female, 10-14 years				
...	All causes	242	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	27.3	3.4
2	Malignant neoplasms (C00-C97)	58	24.0	3.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	5.8	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	5.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.7	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	2.5	*
7	Cerebrovascular diseases (I60-I69)	5	2.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.7	*
9	Influenza and pneumonia (J09-J18)	3	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	3	1.2	*
11	Diabetes mellitus (E10-E14)	2	0.8	*
11	Meningitis (G00,G03)	2	0.8	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
13	Anemias (D50-D64)	1	0.4	*
13	Aortic aneurysm and dissection (I71)	1	0.4	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	51	21.1	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	495	100.0	27.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	44.6	12.2
2	Malignant neoplasms (C00-C97)	52	10.5	2.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	10.1	2.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	6.9	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.4	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.8	*
7	Cerebrovascular diseases (I60-I69)	8	1.6	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	7	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
10	Anemias (D50-D64)	4	0.8	*
12	Septicemia (A40-A41)	3	0.6	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.6	*
12	Pneumonitis due to solids and liquids (J69)	3	0.6	*
15	Influenza and pneumonia (J09-J18)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	68	13.7	3.7
Hispanic, female, 20-24 years				
...	All causes	685	100.0	40.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	255	37.2	14.9
2	Malignant neoplasms (C00-C97)	84	12.3	4.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	10.8	4.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	6.1	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	3.5	1.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	21	3.1	1.2
7	Cerebrovascular diseases (I60-I69)	9	1.3	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.0	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.0	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.9	*
11	Influenza and pneumonia (J09-J18)	5	0.7	*
12	Septicemia (A40-A41)	4	0.6	*
13	Viral hepatitis (B15-B19)	3	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
13	Diabetes mellitus (E10-E14)	3	0.4	*
13	Meningitis (G00,G03)	3	0.4	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
...	All other causes (Residual)	129	18.8	7.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	740	100.0	40.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	28.1	11.2
2	Malignant neoplasms (C00-C97)	107	14.5	5.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	9.5	3.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	8.4	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	6.2	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	27	3.6	1.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.9	*
7	Cerebrovascular diseases (I60-I69)	14	1.9	*
9	Diabetes mellitus (E10-E14)	9	1.2	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.1	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.1	*
12	Septicemia (A40-A41)	7	0.9	*
12	Influenza and pneumonia (J09-J18)	7	0.9	*
14	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.8	*
...	All other causes (Residual)	141	19.1	7.6
Hispanic, female, 30-34 years				
...	All causes	850	100.0	46.8
1	Malignant neoplasms (C00-C97)	220	25.9	12.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	20.5	9.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	7.1	3.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	6.5	3.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	4.6	2.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	33	3.9	1.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	31	3.6	1.7
8	Cerebrovascular diseases (I60-I69)	26	3.1	1.4
9	Septicemia (A40-A41)	15	1.8	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.6	*
11	Diabetes mellitus (E10-E14)	11	1.3	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.2	*
14	Influenza and pneumonia (J09-J18)	9	1.1	*
15	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
...	All other causes (Residual)	137	16.1	7.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,144	100.0	67.5
1	Malignant neoplasms (C00-C97)	347	30.3	20.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	17.0	11.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	100	8.7	5.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	4.2	2.8
5	Cerebrovascular diseases (I60-I69)	44	3.8	2.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	3.7	2.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	32	2.8	1.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	29	2.5	1.7
9	Diabetes mellitus (E10-E14)	28	2.4	1.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	25	2.2	1.5
11	Septicemia (A40-A41)	14	1.2	*
12	Influenza and pneumonia (J09-J18)	11	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.9	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.8	*
15	Chronic lower respiratory diseases (J40-J47)	7	0.6	*
...	All other causes (Residual)	204	17.8	12.0
Hispanic, female, 40-44 years				
...	All causes	1,514	100.0	99.8
1	Malignant neoplasms (C00-C97)	505	33.4	33.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	206	13.6	13.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	119	7.9	7.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	77	5.1	5.1
5	Cerebrovascular diseases (I60-I69)	63	4.2	4.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	51	3.4	3.4
7	Diabetes mellitus (E10-E14)	48	3.2	3.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	3.2	3.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	2.6	2.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.5	1.5
11	Septicemia (A40-A41)	18	1.2	*
12	Viral hepatitis (B15-B19)	17	1.1	*
13	Influenza and pneumonia (J09-J18)	15	1.0	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	14	0.9	*
15	Chronic lower respiratory diseases (J40-J47)	10	0.7	*
...	All other causes (Residual)	260	17.2	17.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	2,111	100.0	162.4
1	Malignant neoplasms (C00-C97)	796	37.7	61.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	204	9.7	15.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	9.2	14.9
4	Cerebrovascular diseases (I60-I69)	111	5.3	8.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	99	4.7	7.6
6	Diabetes mellitus (E10-E14)	74	3.5	5.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	64	3.0	4.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	1.9	3.1
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	1.8	3.0
10	Viral hepatitis (B15-B19)	37	1.8	2.8
11	Septicemia (A40-A41)	28	1.3	2.2
12	Chronic lower respiratory diseases (J40-J47)	25	1.2	1.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	1.2	1.9
14	Influenza and pneumonia (J09-J18)	19	0.9	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.9	*
...	All other causes (Residual)	338	16.0	26.0
Hispanic, female, 50-54 years				
...	All causes	2,655	100.0	256.1
1	Malignant neoplasms (C00-C97)	967	36.4	93.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	343	12.9	33.1
3	Diabetes mellitus (E10-E14)	174	6.6	16.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	5.6	14.5
5	Cerebrovascular diseases (I60-I69)	146	5.5	14.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	139	5.2	13.4
7	Viral hepatitis (B15-B19)	66	2.5	6.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.7	4.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	43	1.6	4.1
10	Septicemia (A40-A41)	42	1.6	4.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	1.3	3.4
12	Influenza and pneumonia (J09-J18)	32	1.2	3.1
12	Chronic lower respiratory diseases (J40-J47)	32	1.2	3.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	1.0	2.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	0.7	*
...	All other causes (Residual)	394	14.8	38.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	3,151	100.0	391.9
1	Malignant neoplasms (C00-C97)	1,217	38.6	151.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	471	14.9	58.6
3	Diabetes mellitus (E10-E14)	202	6.4	25.1
4	Cerebrovascular diseases (I60-I69)	162	5.1	20.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	4.3	17.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	106	3.4	13.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	80	2.5	10.0
8	Septicemia (A40-A41)	60	1.9	7.5
9	Viral hepatitis (B15-B19)	59	1.9	7.3
10	Chronic lower respiratory diseases (J40-J47)	51	1.6	6.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	1.1	4.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	32	1.0	4.0
12	Influenza and pneumonia (J09-J18)	32	1.0	4.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.7	2.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	0.6	2.5
...	All other causes (Residual)	467	14.8	58.1
Hispanic, female, 60-64 years				
...	All causes	3,509	100.0	592.1
1	Malignant neoplasms (C00-C97)	1,236	35.2	208.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	666	19.0	112.4
3	Diabetes mellitus (E10-E14)	268	7.6	45.2
4	Cerebrovascular diseases (I60-I69)	193	5.5	32.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	125	3.6	21.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	2.9	17.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	2.4	14.2
8	Chronic lower respiratory diseases (J40-J47)	73	2.1	12.3
9	Influenza and pneumonia (J09-J18)	51	1.5	8.6
10	Septicemia (A40-A41)	49	1.4	8.3
11	Viral hepatitis (B15-B19)	40	1.1	6.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	1.0	5.9
13	Alzheimer's disease (G30)	20	0.6	3.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	15	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.4	*
...	All other causes (Residual)	537	15.3	90.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	4,042	100.0	921.1
1	Malignant neoplasms (C00-C97)	1,371	33.9	312.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	806	19.9	183.7
3	Diabetes mellitus (E10-E14)	313	7.7	71.3
4	Cerebrovascular diseases (I60-I69)	247	6.1	56.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	131	3.2	29.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	125	3.1	28.5
7	Chronic lower respiratory diseases (J40-J47)	124	3.1	28.3
8	Septicemia (A40-A41)	77	1.9	17.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	1.9	17.5
10	Influenza and pneumonia (J09-J18)	62	1.5	14.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	55	1.4	12.5
12	Viral hepatitis (B15-B19)	41	1.0	9.3
13	Alzheimer's disease (G30)	29	0.7	6.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.4	*
14	Parkinson's disease (G20-G21)	16	0.4	*
...	All other causes (Residual)	552	13.7	125.8
Hispanic, female, 70-74 years				
...	All causes	5,107	100.0	1,465.7
1	Malignant neoplasms (C00-C97)	1,498	29.3	429.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,160	22.7	332.9
3	Diabetes mellitus (E10-E14)	404	7.9	115.9
4	Cerebrovascular diseases (I60-I69)	317	6.2	91.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	169	3.3	48.5
6	Chronic lower respiratory diseases (J40-J47)	134	2.6	38.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	134	2.6	38.5
8	Influenza and pneumonia (J09-J18)	115	2.3	33.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	1.9	28.1
10	Septicemia (A40-A41)	96	1.9	27.6
11	Alzheimer's disease (G30)	67	1.3	19.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	54	1.1	15.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.7	10.6
14	Viral hepatitis (B15-B19)	34	0.7	9.8
15	Parkinson's disease (G20-G21)	32	0.6	9.2
...	All other causes (Residual)	758	14.8	217.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	6,905	100.0	2,487.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,754	25.4	631.9
2	Malignant neoplasms (C00-C97)	1,634	23.7	588.7
3	Diabetes mellitus (E10-E14)	532	7.7	191.7
4	Cerebrovascular diseases (I60-I69)	498	7.2	179.4
5	Chronic lower respiratory diseases (J40-J47)	261	3.8	94.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	3.0	74.9
7	Alzheimer's disease (G30)	196	2.8	70.6
8	Influenza and pneumonia (J09-J18)	160	2.3	57.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	131	1.9	47.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	1.8	45.4
11	Septicemia (A40-A41)	112	1.6	40.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	96	1.4	34.6
13	Parkinson's disease (G20-G21)	71	1.0	25.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.7	17.3
15	Aortic aneurysm and dissection (I71)	29	0.4	10.4
...	All other causes (Residual)	1,049	15.2	377.9
Hispanic, female, 80-84 years				
...	All causes	8,147	100.0	4,213.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,280	28.0	1,179.2
2	Malignant neoplasms (C00-C97)	1,501	18.4	776.3
3	Cerebrovascular diseases (I60-I69)	652	8.0	337.2
4	Diabetes mellitus (E10-E14)	522	6.4	270.0
5	Alzheimer's disease (G30)	356	4.4	184.1
6	Chronic lower respiratory diseases (J40-J47)	342	4.2	176.9
7	Influenza and pneumonia (J09-J18)	227	2.8	117.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	188	2.3	97.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	1.9	79.6
10	Septicemia (A40-A41)	142	1.7	73.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	129	1.6	66.7
12	Parkinson's disease (G20-G21)	95	1.2	49.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	94	1.2	48.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.8	34.1
15	Pneumonitis due to solids and liquids (J69)	45	0.6	23.3
...	All other causes (Residual)	1,354	16.6	700.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	7,265	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,352	32.4	---
2	Malignant neoplasms (C00-C97)	917	12.6	---
3	Cerebrovascular diseases (I60-I69)	640	8.8	---
4	Alzheimer's disease (G30)	423	5.8	---
5	Diabetes mellitus (E10-E14)	377	5.2	---
6	Chronic lower respiratory diseases (J40-J47)	288	4.0	---
7	Influenza and pneumonia (J09-J18)	249	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	2.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	124	1.7	---
10	Septicemia (A40-A41)	119	1.6	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	115	1.6	---
12	Parkinson's disease (G20-G21)	61	0.8	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.7	---
14	Pneumonitis due to solids and liquids (J69)	43	0.6	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	42	0.6	---
...	All other causes (Residual)	1,300	17.9	---
Hispanic, female, 90-94 years				
...	All causes	4,998	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,781	35.6	---
2	Malignant neoplasms (C00-C97)	424	8.5	---
3	Cerebrovascular diseases (I60-I69)	381	7.6	---
4	Alzheimer's disease (G30)	374	7.5	---
5	Influenza and pneumonia (J09-J18)	203	4.1	---
6	Chronic lower respiratory diseases (J40-J47)	188	3.8	---
7	Diabetes mellitus (E10-E14)	175	3.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	106	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	1.9	---
10	Septicemia (A40-A41)	79	1.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	1.4	---
12	Parkinson's disease (G20-G21)	40	0.8	---
13	Pneumonitis due to solids and liquids (J69)	37	0.7	---
14	Atherosclerosis (I70)	34	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.5	---
...	All other causes (Residual)	983	19.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	2,278	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	886	38.9	---
2	Cerebrovascular diseases (I60-I69)	182	8.0	---
3	Alzheimer's disease (G30)	160	7.0	---
4	Malignant neoplasms (C00-C97)	120	5.3	---
5	Influenza and pneumonia (J09-J18)	109	4.8	---
6	Diabetes mellitus (E10-E14)	62	2.7	---
7	Chronic lower respiratory diseases (J40-J47)	60	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	57	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.8	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	1.6	---
11	Septicemia (A40-A41)	29	1.3	---
12	Pneumonitis due to solids and liquids (J69)	21	0.9	---
13	Atherosclerosis (I70)	16	0.7	---
14	Anemias (D50-D64)	8	0.4	---
14	Parkinson's disease (G20-G21)	8	0.4	---
...	All other causes (Residual)	482	21.2	---
Hispanic, female, 100 years and over				
...	All causes	540	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	217	40.2	---
2	Cerebrovascular diseases (I60-I69)	38	7.0	---
3	Alzheimer's disease (G30)	31	5.7	---
4	Influenza and pneumonia (J09-J18)	27	5.0	---
5	Malignant neoplasms (C00-C97)	15	2.8	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	2.2	---
7	Diabetes mellitus (E10-E14)	11	2.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	2.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	2.0	---
10	Pneumonitis due to solids and liquids (J69)	10	1.9	---
11	Atherosclerosis (I70)	9	1.7	---
11	Chronic lower respiratory diseases (J40-J47)	9	1.7	---
13	Septicemia (A40-A41)	8	1.5	---
14	Anemias (D50-D64)	3	0.6	---
15	Nutritional deficiencies (E40-E64)	2	0.4	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.4	---
...	All other causes (Residual)	124	23.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, all ages				
...	All causes	1,939,606	100.0	964.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	502,683	25.9	249.9
2	Malignant neoplasms (C00-C97)	456,576	23.5	226.9
3	Chronic lower respiratory diseases (J40-J47)	114,480	5.9	56.9
4	Cerebrovascular diseases (I60-I69)	107,678	5.6	53.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94,584	4.9	47.0
6	Alzheimer's disease (G30)	66,453	3.4	33.0
7	Diabetes mellitus (E10-E14)	50,046	2.6	24.9
8	Influenza and pneumonia (J09-J18)	43,219	2.2	21.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34,219	1.8	17.0
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,897	1.5	14.4
11	Septicemia (A40-A41)	25,856	1.3	12.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	21,598	1.1	10.7
13	Parkinson's disease (G20-G21)	18,098	0.9	9.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,216	0.9	8.6
15	Pneumonitis due to solids and liquids (J69)	14,680	0.8	7.3
...	All other causes (Residual)	343,323	17.7	170.6
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,325	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	835	35.9	9.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	266	11.4	2.9
3	Malignant neoplasms (C00-C97)	224	9.6	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	6.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	3.2	0.8
6	Influenza and pneumonia (J09-J18)	50	2.2	0.5
7	Septicemia (A40-A41)	42	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.4	0.4
9	Cerebrovascular diseases (I60-I69)	26	1.1	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	25	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	18	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.6	*
14	Meningitis (G00,G03)	10	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.2	*
...	All other causes (Residual)	544	23.4	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 5-9 years				
...	All causes	1,461	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	511	35.0	4.4
2	Malignant neoplasms (C00-C97)	284	19.4	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	6.9	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	4.9	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	3.9	0.5
6	Influenza and pneumonia (J09-J18)	27	1.8	0.2
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.6	0.2
8	Cerebrovascular diseases (I60-I69)	21	1.4	0.2
9	Septicemia (A40-A41)	17	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	17	1.2	*
11	Diabetes mellitus (E10-E14)	7	0.5	*
12	Meningitis (G00,G03)	6	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
14	Anemias (D50-D64)	5	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
...	All other causes (Residual)	303	20.7	2.6
Non-Hispanic, white, both sexes, 10-14 years				
...	All causes	1,901	100.0	15.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	739	38.9	6.1
2	Malignant neoplasms (C00-C97)	275	14.5	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	105	5.5	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	5.3	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	3.7	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	2.5	0.4
7	Influenza and pneumonia (J09-J18)	31	1.6	0.3
8	Septicemia (A40-A41)	26	1.4	0.2
8	Cerebrovascular diseases (I60-I69)	26	1.4	0.2
10	Chronic lower respiratory diseases (J40-J47)	22	1.2	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.1	0.2
12	Meningitis (G00,G03)	10	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.5	*
14	Anemias (D50-D64)	8	0.4	*
14	Pneumonitis due to solids and liquids (J69)	8	0.4	*
...	All other causes (Residual)	403	21.2	3.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-19 years				
...	All causes	7,668	100.0	58.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,569	59.6	34.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,030	13.4	7.8
3	Malignant neoplasms (C00-C97)	392	5.1	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	345	4.5	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	2.3	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	1.3	0.8
7	Cerebrovascular diseases (I60-I69)	40	0.5	0.3
8	Influenza and pneumonia (J09-J18)	37	0.5	0.3
9	Septicemia (A40-A41)	31	0.4	0.2
10	Diabetes mellitus (E10-E14)	29	0.4	0.2
11	Chronic lower respiratory diseases (J40-J47)	27	0.4	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.2	*
13	Pneumonitis due to solids and liquids (J69)	12	0.2	*
13	Pregnancy, childbirth and the puerperium (O00-O99)	12	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.1	*
15	Legal intervention (Y35,Y89.0)	9	0.1	*
...	All other causes (Residual)	825	10.8	6.2
Non-Hispanic, white, both sexes, 20-24 years				
...	All causes	11,939	100.0	91.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,463	54.1	49.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,902	15.9	14.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	592	5.0	4.5
4	Malignant neoplasms (C00-C97)	566	4.7	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	385	3.2	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	142	1.2	1.1
7	Influenza and pneumonia (J09-J18)	79	0.7	0.6
8	Diabetes mellitus (E10-E14)	63	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	62	0.5	0.5
10	Septicemia (A40-A41)	58	0.5	0.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	47	0.4	0.4
12	Chronic lower respiratory diseases (J40-J47)	32	0.3	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.3	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	18	0.2	*
...	All other causes (Residual)	1,472	12.3	11.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 25-29 years				
...	All causes	12,138	100.0	96.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,642	46.5	44.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,946	16.0	15.4
3	Malignant neoplasms (C00-C97)	792	6.5	6.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	642	5.3	5.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	566	4.7	4.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	1.2	1.2
7	Diabetes mellitus (E10-E14)	107	0.9	0.8
8	Influenza and pneumonia (J09-J18)	92	0.8	0.7
9	Cerebrovascular diseases (I60-I69)	91	0.7	0.7
10	Septicemia (A40-A41)	64	0.5	0.5
10	Pregnancy, childbirth and the puerperium (O00-O99)	64	0.5	0.5
12	Chronic lower respiratory diseases (J40-J47)	60	0.5	0.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	60	0.5	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	59	0.5	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.4	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.4	0.3
...	All other causes (Residual)	1,721	14.2	13.6
Non-Hispanic, white, both sexes, 30-34 years				
...	All causes	12,619	100.0	109.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,705	37.3	40.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,946	15.4	16.9
3	Malignant neoplasms (C00-C97)	1,268	10.0	11.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,066	8.4	9.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	488	3.9	4.2
6	Diabetes mellitus (E10-E14)	214	1.7	1.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	146	1.2	1.3
8	Cerebrovascular diseases (I60-I69)	140	1.1	1.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	128	1.0	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	121	1.0	1.0
11	Influenza and pneumonia (J09-J18)	97	0.8	0.8
12	Septicemia (A40-A41)	89	0.7	0.8
13	Pregnancy, childbirth and the puerperium (O00-O99)	72	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	0.6	0.6
15	Chronic lower respiratory diseases (J40-J47)	69	0.5	0.6
...	All other causes (Residual)	1,999	15.8	17.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-39 years				
...	All causes	19,087	100.0	142.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,484	28.7	41.0
2	Malignant neoplasms (C00-C97)	2,731	14.3	20.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,423	12.7	18.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,309	12.1	17.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	501	2.6	3.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	486	2.5	3.6
7	Diabetes mellitus (E10-E14)	402	2.1	3.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	366	1.9	2.7
9	Cerebrovascular diseases (I60-I69)	357	1.9	2.7
10	Septicemia (A40-A41)	178	0.9	1.3
11	Influenza and pneumonia (J09-J18)	171	0.9	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	0.8	1.2
13	Chronic lower respiratory diseases (J40-J47)	142	0.7	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	0.6	0.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	89	0.5	0.7
...	All other causes (Residual)	3,172	16.6	23.7
Non-Hispanic, white, both sexes, 40-44 years				
...	All causes	31,879	100.0	216.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,786	21.3	46.2
2	Malignant neoplasms (C00-C97)	6,099	19.1	41.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,149	16.2	35.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,112	9.8	21.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,240	3.9	8.4
6	Diabetes mellitus (E10-E14)	716	2.2	4.9
7	Cerebrovascular diseases (I60-I69)	711	2.2	4.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	653	2.0	4.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	571	1.8	3.9
10	Chronic lower respiratory diseases (J40-J47)	397	1.2	2.7
11	Septicemia (A40-A41)	347	1.1	2.4
12	Influenza and pneumonia (J09-J18)	325	1.0	2.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	0.7	1.5
14	Viral hepatitis (B15-B19)	216	0.7	1.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	216	0.7	1.5
...	All other causes (Residual)	5,123	16.1	34.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-49 years				
...	All causes	53,142	100.0	327.7
1	Malignant neoplasms (C00-C97)	13,192	24.8	81.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,215	19.2	63.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,113	15.3	50.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,573	6.7	22.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,513	4.7	15.5
6	Cerebrovascular diseases (I60-I69)	1,387	2.6	8.6
7	Diabetes mellitus (E10-E14)	1,320	2.5	8.1
8	Chronic lower respiratory diseases (J40-J47)	1,039	2.0	6.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	700	1.3	4.3
10	Viral hepatitis (B15-B19)	688	1.3	4.2
11	Septicemia (A40-A41)	577	1.1	3.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	559	1.1	3.4
13	Influenza and pneumonia (J09-J18)	512	1.0	3.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	428	0.8	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	242	0.5	1.5
...	All other causes (Residual)	8,084	15.2	49.9
Non-Hispanic, white, both sexes, 50-54 years				
...	All causes	74,756	100.0	483.6
1	Malignant neoplasms (C00-C97)	22,860	30.6	147.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,608	20.9	101.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,232	9.7	46.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,338	4.5	21.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,293	4.4	21.3
6	Chronic lower respiratory diseases (J40-J47)	2,207	3.0	14.3
7	Diabetes mellitus (E10-E14)	2,185	2.9	14.1
8	Cerebrovascular diseases (I60-I69)	2,106	2.8	13.6
9	Viral hepatitis (B15-B19)	1,269	1.7	8.2
10	Septicemia (A40-A41)	868	1.2	5.6
11	Influenza and pneumonia (J09-J18)	758	1.0	4.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	692	0.9	4.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	543	0.7	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	459	0.6	3.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	378	0.5	2.4
...	All other causes (Residual)	10,960	14.7	70.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-59 years				
...	All causes	95,806	100.0	693.4
1	Malignant neoplasms (C00-C97)	34,274	35.8	248.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,200	22.1	153.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,380	5.6	38.9
4	Chronic lower respiratory diseases (J40-J47)	3,905	4.1	28.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,291	3.4	23.8
6	Diabetes mellitus (E10-E14)	3,174	3.3	23.0
7	Cerebrovascular diseases (I60-I69)	2,745	2.9	19.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,676	2.8	19.4
9	Septicemia (A40-A41)	1,269	1.3	9.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,098	1.1	7.9
11	Viral hepatitis (B15-B19)	1,067	1.1	7.7
12	Influenza and pneumonia (J09-J18)	967	1.0	7.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	492	0.5	3.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	486	0.5	3.5
15	Aortic aneurysm and dissection (I71)	442	0.5	3.2
...	All other causes (Residual)	13,340	13.9	96.6
Non-Hispanic, white, both sexes, 60-64 years				
...	All causes	118,070	100.0	1,039.6
1	Malignant neoplasms (C00-C97)	45,829	38.8	403.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,603	22.5	234.2
3	Chronic lower respiratory diseases (J40-J47)	7,195	6.1	63.4
4	Diabetes mellitus (E10-E14)	4,098	3.5	36.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,022	3.4	35.4
6	Cerebrovascular diseases (I60-I69)	3,788	3.2	33.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,636	2.2	23.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,917	1.6	16.9
9	Septicemia (A40-A41)	1,599	1.4	14.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,427	1.2	12.6
11	Influenza and pneumonia (J09-J18)	1,286	1.1	11.3
12	Aortic aneurysm and dissection (I71)	699	0.6	6.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	663	0.6	5.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	614	0.5	5.4
15	Viral hepatitis (B15-B19)	472	0.4	4.2
...	All other causes (Residual)	15,222	12.9	134.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 65-69 years				
...	All causes	135,914	100.0	1,608.0
1	Malignant neoplasms (C00-C97)	52,809	38.9	624.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,878	22.0	353.5
3	Chronic lower respiratory diseases (J40-J47)	10,663	7.8	126.2
4	Cerebrovascular diseases (I60-I69)	5,141	3.8	60.8
5	Diabetes mellitus (E10-E14)	4,543	3.3	53.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,299	2.4	39.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,087	1.5	24.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,041	1.5	24.1
9	Septicemia (A40-A41)	1,949	1.4	23.1
10	Influenza and pneumonia (J09-J18)	1,637	1.2	19.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,254	0.9	14.8
12	Alzheimer's disease (G30)	927	0.7	11.0
13	Aortic aneurysm and dissection (I71)	892	0.7	10.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	823	0.6	9.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	811	0.6	9.6
...	All other causes (Residual)	17,160	12.6	203.0
Non-Hispanic, white, both sexes, 70-74 years				
...	All causes	170,428	100.0	2,499.8
1	Malignant neoplasms (C00-C97)	59,622	35.0	874.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,686	22.7	567.4
3	Chronic lower respiratory diseases (J40-J47)	15,057	8.8	220.9
4	Cerebrovascular diseases (I60-I69)	7,720	4.5	113.2
5	Diabetes mellitus (E10-E14)	5,585	3.3	81.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,698	2.2	54.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,013	1.8	44.2
8	Septicemia (A40-A41)	2,526	1.5	37.1
9	Influenza and pneumonia (J09-J18)	2,524	1.5	37.0
10	Alzheimer's disease (G30)	2,453	1.4	36.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,843	1.1	27.0
12	Parkinson's disease (G20-G21)	1,403	0.8	20.6
13	Aortic aneurysm and dissection (I71)	1,316	0.8	19.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,157	0.7	17.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,144	0.7	16.8
...	All other causes (Residual)	22,681	13.3	332.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-79 years				
...	All causes	239,043	100.0	3,995.4
1	Malignant neoplasms (C00-C97)	69,738	29.2	1,165.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	58,336	24.4	975.0
3	Chronic lower respiratory diseases (J40-J47)	20,599	8.6	344.3
4	Cerebrovascular diseases (I60-I69)	13,425	5.6	224.4
5	Diabetes mellitus (E10-E14)	7,177	3.0	120.0
6	Alzheimer's disease (G30)	6,546	2.7	109.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,015	2.1	83.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,772	2.0	79.8
9	Influenza and pneumonia (J09-J18)	4,627	1.9	77.3
10	Septicemia (A40-A41)	3,569	1.5	59.7
11	Parkinson's disease (G20-G21)	3,305	1.4	55.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,807	0.8	30.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,780	0.7	29.8
14	Aortic aneurysm and dissection (I71)	1,745	0.7	29.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,708	0.7	28.5
...	All other causes (Residual)	34,894	14.6	583.2
Non-Hispanic, white, both sexes, 80-84 years				
...	All causes	310,709	100.0	6,515.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	84,631	27.2	1,774.7
2	Malignant neoplasms (C00-C97)	69,216	22.3	1,451.4
3	Chronic lower respiratory diseases (J40-J47)	23,038	7.4	483.1
4	Cerebrovascular diseases (I60-I69)	20,906	6.7	438.4
5	Alzheimer's disease (G30)	13,702	4.4	287.3
6	Diabetes mellitus (E10-E14)	8,390	2.7	175.9
7	Influenza and pneumonia (J09-J18)	7,647	2.5	160.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,939	2.2	145.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,733	2.2	141.2
10	Parkinson's disease (G20-G21)	5,116	1.6	107.3
11	Septicemia (A40-A41)	4,625	1.5	97.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,928	0.9	61.4
13	Pneumonitis due to solids and liquids (J69)	2,899	0.9	60.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,326	0.7	48.8
15	Aortic aneurysm and dissection (I71)	2,086	0.7	43.7
...	All other causes (Residual)	49,527	15.9	1,038.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 85-89 years				
...	All causes	315,107	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	96,636	30.7	---
2	Malignant neoplasms (C00-C97)	48,719	15.5	---
3	Cerebrovascular diseases (I60-I69)	24,397	7.7	---
4	Alzheimer's disease (G30)	19,084	6.1	---
5	Chronic lower respiratory diseases (J40-J47)	18,083	5.7	---
6	Influenza and pneumonia (J09-J18)	9,722	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,377	2.3	---
8	Diabetes mellitus (E10-E14)	7,151	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,021	2.2	---
10	Parkinson's disease (G20-G21)	4,586	1.5	---
11	Septicemia (A40-A41)	4,427	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,801	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,636	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,129	0.7	---
15	Aortic aneurysm and dissection (I71)	1,769	0.6	---
...	All other causes (Residual)	56,569	18.0	---
Non-Hispanic, white, both sexes, 90-94 years				
...	All causes	213,005	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,440	34.5	---
2	Malignant neoplasms (C00-C97)	21,454	10.1	---
3	Cerebrovascular diseases (I60-I69)	17,005	8.0	---
4	Alzheimer's disease (G30)	15,378	7.2	---
5	Chronic lower respiratory diseases (J40-J47)	8,934	4.2	---
6	Influenza and pneumonia (J09-J18)	8,010	3.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,846	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,543	2.1	---
9	Diabetes mellitus (E10-E14)	3,721	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,082	1.4	---
11	Pneumonitis due to solids and liquids (J69)	2,695	1.3	---
12	Septicemia (A40-A41)	2,517	1.2	---
13	Parkinson's disease (G20-G21)	2,187	1.0	---
14	Atherosclerosis (I70)	1,444	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,248	0.6	---
...	All other causes (Residual)	42,501	20.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 95-99 years				
...	All causes	82,997	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,844	37.2	---
2	Alzheimer's disease (G30)	6,494	7.8	---
3	Cerebrovascular diseases (I60-I69)	6,459	7.8	---
4	Malignant neoplasms (C00-C97)	5,509	6.6	---
5	Influenza and pneumonia (J09-J18)	3,592	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	2,629	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,946	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,566	1.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,378	1.7	---
10	Pneumonitis due to solids and liquids (J69)	1,047	1.3	---
11	Diabetes mellitus (E10-E14)	990	1.2	---
12	Septicemia (A40-A41)	841	1.0	---
13	Atherosclerosis (I70)	763	0.9	---
14	Parkinson's disease (G20-G21)	464	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	377	0.5	---
...	All other causes (Residual)	18,098	21.8	---
Non-Hispanic, white, both sexes, 100 years and over				
...	All causes	16,533	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,490	39.3	---
2	Alzheimer's disease (G30)	1,149	6.9	---
3	Cerebrovascular diseases (I60-I69)	1,057	6.4	---
4	Influenza and pneumonia (J09-J18)	938	5.7	---
5	Malignant neoplasms (C00-C97)	679	4.1	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	355	2.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	332	2.0	---
8	Chronic lower respiratory diseases (J40-J47)	327	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	306	1.9	---
10	Atherosclerosis (I70)	243	1.5	---
11	Pneumonitis due to solids and liquids (J69)	201	1.2	---
12	Diabetes mellitus (E10-E14)	163	1.0	---
13	Septicemia (A40-A41)	139	0.8	---
14	Anemias (D50-D64)	52	0.3	---
15	Parkinson's disease (G20-G21)	47	0.3	---
...	All other causes (Residual)	4,055	24.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	948,662	100.0	960.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	251,229	26.5	254.3
2	Malignant neoplasms (C00-C97)	237,672	25.1	240.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59,274	6.2	60.0
4	Chronic lower respiratory diseases (J40-J47)	53,989	5.7	54.7
5	Cerebrovascular diseases (I60-I69)	41,416	4.4	41.9
6	Diabetes mellitus (E10-E14)	25,584	2.7	25.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22,660	2.4	22.9
8	Alzheimer's disease (G30)	19,378	2.0	19.6
9	Influenza and pneumonia (J09-J18)	19,361	2.0	19.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,896	1.8	17.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	14,007	1.5	14.2
12	Septicemia (A40-A41)	11,653	1.2	11.8
13	Parkinson's disease (G20-G21)	10,457	1.1	10.6
14	Pneumonitis due to solids and liquids (J69)	7,713	0.8	7.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,480	0.7	6.6
...	All other causes (Residual)	150,893	15.9	152.8
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,316	100.0	28.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	39.6	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	10.3	2.9
3	Malignant neoplasms (C00-C97)	111	8.4	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	6.6	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	2.3	0.6
6	Septicemia (A40-A41)	27	2.1	0.6
7	Influenza and pneumonia (J09-J18)	21	1.6	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	14	1.1	*
10	Cerebrovascular diseases (I60-I69)	12	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.8	*
12	Meningitis (G00,G03)	6	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	4	0.3	*
...	All other causes (Residual)	307	23.3	6.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-9 years				
...	All causes	810	100.0	13.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	296	36.5	5.0
2	Malignant neoplasms (C00-C97)	150	18.5	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	6.8	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	4.3	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	3.5	0.5
6	Septicemia (A40-A41)	16	2.0	*
7	Influenza and pneumonia (J09-J18)	14	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	11	1.4	*
9	Cerebrovascular diseases (I60-I69)	10	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.0	*
11	Diabetes mellitus (E10-E14)	5	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	5	0.6	*
13	Anemias (D50-D64)	3	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
15	Meningococcal infection (A39)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	167	20.6	2.8
Non-Hispanic, white, male, 10-14 years				
...	All causes	1,135	100.0	18.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	467	41.1	7.5
2	Malignant neoplasms (C00-C97)	143	12.6	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	6.8	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	4.9	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	3.5	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	1.9	0.4
7	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
8	Cerebrovascular diseases (I60-I69)	14	1.2	*
9	Septicemia (A40-A41)	13	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.1	*
11	Influenza and pneumonia (J09-J18)	11	1.0	*
12	Pneumonitis due to solids and liquids (J69)	7	0.6	*
13	Anemias (D50-D64)	6	0.5	*
13	Meningitis (G00,G03)	6	0.5	*
15	Aortic aneurysm and dissection (I71)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	238	21.0	3.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 15-19 years				
...	All causes	5,239	100.0	77.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,113	59.4	45.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	846	16.1	12.5
3	Malignant neoplasms (C00-C97)	231	4.4	3.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	231	4.4	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	115	2.2	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.0	0.8
7	Cerebrovascular diseases (I60-I69)	24	0.5	0.4
8	Diabetes mellitus (E10-E14)	18	0.3	*
8	Influenza and pneumonia (J09-J18)	18	0.3	*
10	Chronic lower respiratory diseases (J40-J47)	17	0.3	*
11	Septicemia (A40-A41)	13	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.2	*
12	Pneumonitis due to solids and liquids (J69)	10	0.2	*
14	Legal intervention (Y35,Y89.0)	9	0.2	*
15	Aortic aneurysm and dissection (I71)	5	0.1	*
...	All other causes (Residual)	524	10.0	7.7
Non-Hispanic, white, male, 20-24 years				
...	All causes	8,890	100.0	132.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,024	56.5	74.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,614	18.2	24.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	427	4.8	6.3
4	Malignant neoplasms (C00-C97)	345	3.9	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	228	2.6	3.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	0.9	1.2
7	Influenza and pneumonia (J09-J18)	50	0.6	0.7
8	Diabetes mellitus (E10-E14)	42	0.5	0.6
9	Septicemia (A40-A41)	36	0.4	0.5
10	Cerebrovascular diseases (I60-I69)	32	0.4	0.5
11	Chronic lower respiratory diseases (J40-J47)	21	0.2	0.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.2	*
13	Legal intervention (Y35,Y89.0)	13	0.1	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.1	*
14	Pneumonitis due to solids and liquids (J69)	12	0.1	*
...	All other causes (Residual)	936	10.5	13.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-29 years				
...	All causes	8,646	100.0	135.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,254	49.2	66.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,601	18.5	25.1
3	Malignant neoplasms (C00-C97)	444	5.1	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	422	4.9	6.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	406	4.7	6.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	1.0	1.3
7	Diabetes mellitus (E10-E14)	67	0.8	1.1
8	Influenza and pneumonia (J09-J18)	53	0.6	0.8
9	Cerebrovascular diseases (I60-I69)	45	0.5	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	40	0.5	0.6
11	Septicemia (A40-A41)	36	0.4	0.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	33	0.4	0.5
13	Chronic lower respiratory diseases (J40-J47)	32	0.4	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.3	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.3	0.4
...	All other causes (Residual)	1,076	12.4	16.9
Non-Hispanic, white, male, 30-34 years				
...	All causes	8,492	100.0	146.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,455	40.7	59.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,542	18.2	26.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	741	8.7	12.8
4	Malignant neoplasms (C00-C97)	608	7.2	10.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	349	4.1	6.0
6	Diabetes mellitus (E10-E14)	129	1.5	2.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	102	1.2	1.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	90	1.1	1.6
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	0.8	1.1
10	Cerebrovascular diseases (I60-I69)	62	0.7	1.1
11	Influenza and pneumonia (J09-J18)	53	0.6	0.9
12	Septicemia (A40-A41)	48	0.6	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	0.5	0.7
14	Chronic lower respiratory diseases (J40-J47)	33	0.4	0.6
15	Legal intervention (Y35,Y89.0)	29	0.3	0.5
...	All other causes (Residual)	1,147	13.5	19.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 35-39 years				
...	All causes	12,228	100.0	181.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,878	31.7	57.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,882	15.4	28.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,629	13.3	24.2
4	Malignant neoplasms (C00-C97)	1,183	9.7	17.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	2.8	5.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	302	2.5	4.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	292	2.4	4.3
8	Diabetes mellitus (E10-E14)	257	2.1	3.8
9	Cerebrovascular diseases (I60-I69)	181	1.5	2.7
10	Septicemia (A40-A41)	93	0.8	1.4
11	Influenza and pneumonia (J09-J18)	90	0.7	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	0.7	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	0.6	1.0
14	Aortic aneurysm and dissection (I71)	65	0.5	1.0
15	Chronic lower respiratory diseases (J40-J47)	64	0.5	1.0
...	All other causes (Residual)	1,822	14.9	27.1
Non-Hispanic, white, male, 40-44 years				
...	All causes	19,911	100.0	271.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,659	23.4	63.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,689	18.5	50.2
3	Malignant neoplasms (C00-C97)	2,714	13.6	37.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,327	11.7	31.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	795	4.0	10.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	560	2.8	7.6
7	Diabetes mellitus (E10-E14)	452	2.3	6.2
8	Cerebrovascular diseases (I60-I69)	381	1.9	5.2
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	372	1.9	5.1
10	Chronic lower respiratory diseases (J40-J47)	192	1.0	2.6
11	Septicemia (A40-A41)	176	0.9	2.4
11	Influenza and pneumonia (J09-J18)	176	0.9	2.4
13	Viral hepatitis (B15-B19)	130	0.7	1.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	0.6	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	107	0.5	1.5
...	All other causes (Residual)	3,052	15.3	41.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-49 years				
...	All causes	33,039	100.0	409.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,508	22.7	93.1
2	Malignant neoplasms (C00-C97)	6,298	19.1	78.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,483	16.6	68.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,677	8.1	33.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,702	5.2	21.1
6	Diabetes mellitus (E10-E14)	856	2.6	10.6
7	Cerebrovascular diseases (I60-I69)	740	2.2	9.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	573	1.7	7.1
9	Chronic lower respiratory diseases (J40-J47)	531	1.6	6.6
10	Viral hepatitis (B15-B19)	469	1.4	5.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	365	1.1	4.5
12	Septicemia (A40-A41)	307	0.9	3.8
13	Influenza and pneumonia (J09-J18)	285	0.9	3.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	252	0.8	3.1
15	Aortic aneurysm and dissection (I71)	141	0.4	1.7
...	All other causes (Residual)	4,852	14.7	60.2
Non-Hispanic, white, male, 50-54 years				
...	All causes	46,892	100.0	613.1
1	Malignant neoplasms (C00-C97)	12,060	25.7	157.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,599	24.7	151.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,915	10.5	64.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,442	5.2	31.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,379	5.1	31.1
6	Diabetes mellitus (E10-E14)	1,358	2.9	17.8
7	Cerebrovascular diseases (I60-I69)	1,196	2.6	15.6
8	Chronic lower respiratory diseases (J40-J47)	1,155	2.5	15.1
9	Viral hepatitis (B15-B19)	927	2.0	12.1
10	Septicemia (A40-A41)	455	1.0	5.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	453	1.0	5.9
12	Influenza and pneumonia (J09-J18)	442	0.9	5.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	395	0.8	5.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	331	0.7	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	241	0.5	3.2
...	All other causes (Residual)	6,544	14.0	85.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 55-59 years				
...	All causes	59,418	100.0	875.5
1	Malignant neoplasms (C00-C97)	18,670	31.4	275.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,498	26.1	228.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,715	6.3	54.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,402	4.0	35.4
5	Chronic lower respiratory diseases (J40-J47)	2,082	3.5	30.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,985	3.3	29.2
7	Diabetes mellitus (E10-E14)	1,968	3.3	29.0
8	Cerebrovascular diseases (I60-I69)	1,565	2.6	23.1
9	Viral hepatitis (B15-B19)	811	1.4	11.9
10	Septicemia (A40-A41)	677	1.1	10.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	648	1.1	9.5
12	Influenza and pneumonia (J09-J18)	583	1.0	8.6
13	Aortic aneurysm and dissection (I71)	338	0.6	5.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	333	0.6	4.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	297	0.5	4.4
...	All other causes (Residual)	7,846	13.2	115.6
Non-Hispanic, white, male, 60-64 years				
...	All causes	70,559	100.0	1,281.9
1	Malignant neoplasms (C00-C97)	25,283	35.8	459.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,503	26.2	336.2
3	Chronic lower respiratory diseases (J40-J47)	3,689	5.2	67.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,695	3.8	49.0
5	Diabetes mellitus (E10-E14)	2,475	3.5	45.0
6	Cerebrovascular diseases (I60-I69)	2,127	3.0	38.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,840	2.6	33.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,465	2.1	26.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	801	1.1	14.6
10	Septicemia (A40-A41)	794	1.1	14.4
11	Influenza and pneumonia (J09-J18)	758	1.1	13.8
12	Aortic aneurysm and dissection (I71)	534	0.8	9.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	398	0.6	7.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	359	0.5	6.5
15	Viral hepatitis (B15-B19)	333	0.5	6.0
...	All other causes (Residual)	8,505	12.1	154.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-69 years				
...	All causes	78,445	100.0	1,962.5
1	Malignant neoplasms (C00-C97)	29,077	37.1	727.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,697	25.1	492.8
3	Chronic lower respiratory diseases (J40-J47)	5,471	7.0	136.9
4	Cerebrovascular diseases (I60-I69)	2,733	3.5	68.4
5	Diabetes mellitus (E10-E14)	2,720	3.5	68.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,139	2.7	53.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,369	1.7	34.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,077	1.4	26.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,013	1.3	25.3
10	Septicemia (A40-A41)	1,004	1.3	25.1
11	Influenza and pneumonia (J09-J18)	899	1.1	22.5
12	Aortic aneurysm and dissection (I71)	628	0.8	15.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	473	0.6	11.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	440	0.6	11.0
15	Alzheimer's disease (G30)	420	0.5	10.5
...	All other causes (Residual)	9,285	11.8	232.3
Non-Hispanic, white, male, 70-74 years				
...	All causes	94,422	100.0	3,037.2
1	Malignant neoplasms (C00-C97)	32,554	34.5	1,047.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,717	25.1	762.9
3	Chronic lower respiratory diseases (J40-J47)	7,527	8.0	242.1
4	Cerebrovascular diseases (I60-I69)	3,935	4.2	126.6
5	Diabetes mellitus (E10-E14)	3,120	3.3	100.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,308	2.4	74.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,641	1.7	52.8
8	Influenza and pneumonia (J09-J18)	1,361	1.4	43.8
9	Septicemia (A40-A41)	1,288	1.4	41.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,066	1.1	34.3
11	Alzheimer's disease (G30)	1,016	1.1	32.7
12	Parkinson's disease (G20-G21)	902	1.0	29.0
13	Aortic aneurysm and dissection (I71)	834	0.9	26.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	795	0.8	25.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	651	0.7	20.9
...	All other causes (Residual)	11,707	12.4	376.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 75-79 years				
...	All causes	124,921	100.0	4,859.8
1	Malignant neoplasms (C00-C97)	37,517	30.0	1,459.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,639	26.1	1,269.8
3	Chronic lower respiratory diseases (J40-J47)	10,070	8.1	391.8
4	Cerebrovascular diseases (I60-I69)	6,246	5.0	243.0
5	Diabetes mellitus (E10-E14)	3,655	2.9	142.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,791	2.2	108.6
7	Alzheimer's disease (G30)	2,563	2.1	99.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,539	2.0	98.8
9	Influenza and pneumonia (J09-J18)	2,367	1.9	92.1
10	Parkinson's disease (G20-G21)	2,091	1.7	81.3
11	Septicemia (A40-A41)	1,722	1.4	67.0
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,023	0.8	39.8
13	Aortic aneurysm and dissection (I71)	987	0.8	38.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	951	0.8	37.0
15	Pneumonitis due to solids and liquids (J69)	949	0.8	36.9
...	All other causes (Residual)	16,811	13.5	654.0
Non-Hispanic, white, male, 80-84 years				
...	All causes	147,477	100.0	7,942.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,961	28.5	2,259.9
2	Malignant neoplasms (C00-C97)	35,658	24.2	1,920.4
3	Chronic lower respiratory diseases (J40-J47)	10,927	7.4	588.5
4	Cerebrovascular diseases (I60-I69)	8,200	5.6	441.6
5	Alzheimer's disease (G30)	4,727	3.2	254.6
6	Diabetes mellitus (E10-E14)	3,970	2.7	213.8
7	Influenza and pneumonia (J09-J18)	3,731	2.5	200.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,526	2.4	189.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,474	2.4	187.1
10	Parkinson's disease (G20-G21)	3,063	2.1	165.0
11	Septicemia (A40-A41)	2,107	1.4	113.5
12	Pneumonitis due to solids and liquids (J69)	1,585	1.1	85.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,262	0.9	68.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,116	0.8	60.1
15	Aortic aneurysm and dissection (I71)	1,097	0.7	59.1
...	All other causes (Residual)	21,073	14.3	1,134.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85-89 years				
...	All causes	128,774	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,448	31.4	---
2	Malignant neoplasms (C00-C97)	23,420	18.2	---
3	Cerebrovascular diseases (I60-I69)	8,186	6.4	---
4	Chronic lower respiratory diseases (J40-J47)	7,921	6.2	---
5	Alzheimer's disease (G30)	5,718	4.4	---
6	Influenza and pneumonia (J09-J18)	4,211	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,329	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,315	2.6	---
9	Diabetes mellitus (E10-E14)	2,961	2.3	---
10	Parkinson's disease (G20-G21)	2,533	2.0	---
11	Pneumonitis due to solids and liquids (J69)	1,911	1.5	---
12	Septicemia (A40-A41)	1,732	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,184	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	956	0.7	---
15	Aortic aneurysm and dissection (I71)	852	0.7	---
...	All other causes (Residual)	20,097	15.6	---
Non-Hispanic, white, male, 90-94 years				
...	All causes	68,929	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,320	35.3	---
2	Malignant neoplasms (C00-C97)	9,123	13.2	---
3	Cerebrovascular diseases (I60-I69)	4,368	6.3	---
4	Alzheimer's disease (G30)	3,538	5.1	---
5	Chronic lower respiratory diseases (J40-J47)	3,340	4.8	---
6	Influenza and pneumonia (J09-J18)	2,943	4.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,844	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,721	2.5	---
9	Diabetes mellitus (E10-E14)	1,246	1.8	---
10	Pneumonitis due to solids and liquids (J69)	1,223	1.8	---
11	Parkinson's disease (G20-G21)	1,027	1.5	---
12	Septicemia (A40-A41)	843	1.2	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	782	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	480	0.7	---
15	Atherosclerosis (I70)	438	0.6	---
...	All other causes (Residual)	11,693	17.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 95-99 years				
...	All causes	19,207	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,279	37.9	---
2	Malignant neoplasms (C00-C97)	1,873	9.8	---
3	Cerebrovascular diseases (I60-I69)	1,202	6.3	---
4	Influenza and pneumonia (J09-J18)	1,036	5.4	---
5	Alzheimer's disease (G30)	982	5.1	---
6	Chronic lower respiratory diseases (J40-J47)	803	4.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	541	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	515	2.7	---
9	Pneumonitis due to solids and liquids (J69)	376	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	253	1.3	---
11	Diabetes mellitus (E10-E14)	247	1.3	---
12	Septicemia (A40-A41)	196	1.0	---
13	Parkinson's disease (G20-G21)	175	0.9	---
14	Atherosclerosis (I70)	148	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	96	0.5	---
...	All other causes (Residual)	3,485	18.1	---
Non-Hispanic, white, male, 100 years and over				
...	All causes	2,509	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,029	41.0	---
2	Influenza and pneumonia (J09-J18)	200	8.0	---
3	Malignant neoplasms (C00-C97)	184	7.3	---
4	Cerebrovascular diseases (I60-I69)	117	4.7	---
5	Alzheimer's disease (G30)	86	3.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	2.8	---
8	Chronic lower respiratory diseases (J40-J47)	67	2.7	---
9	Pneumonitis due to solids and liquids (J69)	53	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	1.4	---
11	Atherosclerosis (I70)	33	1.3	---
12	Diabetes mellitus (E10-E14)	30	1.2	---
13	Septicemia (A40-A41)	18	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	---
14	Parkinson's disease (G20-G21)	10	0.4	---
...	All other causes (Residual)	496	19.8	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, all ages				
...	All causes	990,944	100.0	967.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	251,454	25.4	245.5
2	Malignant neoplasms (C00-C97)	218,904	22.1	213.7
3	Cerebrovascular diseases (I60-I69)	66,262	6.7	64.7
4	Chronic lower respiratory diseases (J40-J47)	60,491	6.1	59.1
5	Alzheimer's disease (G30)	47,075	4.8	46.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,310	3.6	34.5
7	Diabetes mellitus (E10-E14)	24,462	2.5	23.9
8	Influenza and pneumonia (J09-J18)	23,858	2.4	23.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,323	1.7	16.9
10	Septicemia (A40-A41)	14,203	1.4	13.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,736	1.1	10.5
12	Parkinson's disease (G20-G21)	7,641	0.8	7.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,591	0.8	7.4
14	Pneumonitis due to solids and liquids (J69)	6,967	0.7	6.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,237	0.6	6.1
...	All other causes (Residual)	192,430	19.4	187.9
Non-Hispanic, white, female, 1-4 years				
...	All causes	1,009	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	31.1	7.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	12.9	2.9
3	Malignant neoplasms (C00-C97)	113	11.2	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	5.9	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.5	1.0
6	Influenza and pneumonia (J09-J18)	29	2.9	0.7
7	Septicemia (A40-A41)	15	1.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.4	*
8	Cerebrovascular diseases (I60-I69)	14	1.4	*
10	Certain conditions originating in the perinatal period (P00-P96)	11	1.1	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
14	Meningitis (G00,G03)	4	0.4	*
15	Anemias (D50-D64)	2	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.2	*
...	All other causes (Residual)	234	23.2	5.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 5-9 years				
...	All causes	651	100.0	11.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	33.0	3.8
2	Malignant neoplasms (C00-C97)	134	20.6	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	7.1	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	5.5	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	4.5	0.5
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	2.3	*
7	Influenza and pneumonia (J09-J18)	13	2.0	*
8	Cerebrovascular diseases (I60-I69)	11	1.7	*
9	Meningitis (G00,G03)	6	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	6	0.9	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
12	Anemias (D50-D64)	2	0.3	*
12	Diabetes mellitus (E10-E14)	2	0.3	*
12	Pneumonitis due to solids and liquids (J69)	2	0.3	*
12	Diseases of appendix (K35-K38)	2	0.3	*
...	All other causes (Residual)	128	19.7	2.3
Non-Hispanic, white, female, 10-14 years				
...	All causes	766	100.0	13.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	35.5	4.6
2	Malignant neoplasms (C00-C97)	132	17.2	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	5.9	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	3.9	0.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	3.7	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	3.4	0.4
7	Influenza and pneumonia (J09-J18)	20	2.6	0.3
8	Septicemia (A40-A41)	13	1.7	*
9	Cerebrovascular diseases (I60-I69)	12	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
13	Meningitis (G00,G03)	4	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
15	Anemias (D50-D64)	2	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	155	20.2	2.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-19 years				
...	All causes	2,429	100.0	37.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,456	59.9	22.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	184	7.6	2.9
3	Malignant neoplasms (C00-C97)	161	6.6	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	114	4.7	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	2.7	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	2.0	0.7
7	Influenza and pneumonia (J09-J18)	19	0.8	*
8	Septicemia (A40-A41)	18	0.7	*
9	Cerebrovascular diseases (I60-I69)	16	0.7	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	12	0.5	*
11	Diabetes mellitus (E10-E14)	11	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	10	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	4	0.2	*
...	All other causes (Residual)	297	12.2	4.6
Non-Hispanic, white, female, 20-24 years				
...	All causes	3,049	100.0	47.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,439	47.2	22.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	288	9.4	4.5
3	Malignant neoplasms (C00-C97)	221	7.2	3.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	165	5.4	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	157	5.1	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	2.0	1.0
7	Pregnancy, childbirth and the puerperium (O00-O99)	47	1.5	0.7
8	Cerebrovascular diseases (I60-I69)	30	1.0	0.5
9	Influenza and pneumonia (J09-J18)	29	1.0	0.5
10	Septicemia (A40-A41)	22	0.7	0.3
11	Diabetes mellitus (E10-E14)	21	0.7	0.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	11	0.4	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.3	*
...	All other causes (Residual)	519	17.0	8.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 25-29 years				
...	All causes	3,492	100.0	55.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,388	39.7	22.1
2	Malignant neoplasms (C00-C97)	348	10.0	5.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	345	9.9	5.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	220	6.3	3.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	4.6	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	64	1.8	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	1.8	1.0
8	Cerebrovascular diseases (I60-I69)	46	1.3	0.7
9	Diabetes mellitus (E10-E14)	40	1.1	0.6
10	Influenza and pneumonia (J09-J18)	39	1.1	0.6
11	Septicemia (A40-A41)	28	0.8	0.4
11	Chronic lower respiratory diseases (J40-J47)	28	0.8	0.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	27	0.8	0.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.5	*
...	All other causes (Residual)	659	18.9	10.5
Non-Hispanic, white, female, 30-34 years				
...	All causes	4,127	100.0	72.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,250	30.3	21.8
2	Malignant neoplasms (C00-C97)	660	16.0	11.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	404	9.8	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	325	7.9	5.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	3.4	2.4
6	Diabetes mellitus (E10-E14)	85	2.1	1.5
7	Cerebrovascular diseases (I60-I69)	78	1.9	1.4
8	Pregnancy, childbirth and the puerperium (O00-O99)	72	1.7	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	1.4	1.0
10	Influenza and pneumonia (J09-J18)	44	1.1	0.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	44	1.1	0.8
12	Septicemia (A40-A41)	41	1.0	0.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.9	0.7
14	Chronic lower respiratory diseases (J40-J47)	36	0.9	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.8	0.5
...	All other causes (Residual)	823	19.9	14.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-39 years				
...	All causes	6,859	100.0	103.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,606	23.4	24.1
2	Malignant neoplasms (C00-C97)	1,548	22.6	23.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	680	9.9	10.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	541	7.9	8.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	184	2.7	2.8
6	Cerebrovascular diseases (I60-I69)	176	2.6	2.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	162	2.4	2.4
8	Diabetes mellitus (E10-E14)	145	2.1	2.2
9	Septicemia (A40-A41)	85	1.2	1.3
10	Influenza and pneumonia (J09-J18)	81	1.2	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	1.2	1.2
12	Chronic lower respiratory diseases (J40-J47)	78	1.1	1.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	74	1.1	1.1
14	Pregnancy, childbirth and the puerperium (O00-O99)	73	1.1	1.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	0.7	0.7
...	All other causes (Residual)	1,301	19.0	19.5
Non-Hispanic, white, female, 40-44 years				
...	All causes	11,968	100.0	162.7
1	Malignant neoplasms (C00-C97)	3,385	28.3	46.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,127	17.8	28.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,460	12.2	19.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	785	6.6	10.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	445	3.7	6.0
6	Cerebrovascular diseases (I60-I69)	330	2.8	4.5
7	Diabetes mellitus (E10-E14)	264	2.2	3.6
8	Chronic lower respiratory diseases (J40-J47)	205	1.7	2.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	199	1.7	2.7
10	Septicemia (A40-A41)	171	1.4	2.3
11	Influenza and pneumonia (J09-J18)	149	1.2	2.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	0.9	1.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	93	0.8	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	89	0.7	1.2
15	Viral hepatitis (B15-B19)	86	0.7	1.2
...	All other causes (Residual)	2,071	17.3	28.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-49 years				
...	All causes	20,103	100.0	246.7
1	Malignant neoplasms (C00-C97)	6,894	34.3	84.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,707	13.5	33.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,630	13.1	32.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	896	4.5	11.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	811	4.0	10.0
6	Cerebrovascular diseases (I60-I69)	647	3.2	7.9
7	Chronic lower respiratory diseases (J40-J47)	508	2.5	6.2
8	Diabetes mellitus (E10-E14)	464	2.3	5.7
9	Septicemia (A40-A41)	270	1.3	3.3
10	Influenza and pneumonia (J09-J18)	227	1.1	2.8
11	Viral hepatitis (B15-B19)	219	1.1	2.7
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	194	1.0	2.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	176	0.9	2.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	127	0.6	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	0.5	1.3
...	All other causes (Residual)	3,225	16.0	39.6
Non-Hispanic, white, female, 50-54 years				
...	All causes	27,864	100.0	356.7
1	Malignant neoplasms (C00-C97)	10,800	38.8	138.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,009	14.4	51.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,317	8.3	29.7
4	Chronic lower respiratory diseases (J40-J47)	1,052	3.8	13.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	959	3.4	12.3
6	Cerebrovascular diseases (I60-I69)	910	3.3	11.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	851	3.1	10.9
8	Diabetes mellitus (E10-E14)	827	3.0	10.6
9	Septicemia (A40-A41)	413	1.5	5.3
10	Viral hepatitis (B15-B19)	342	1.2	4.4
11	Influenza and pneumonia (J09-J18)	316	1.1	4.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	297	1.1	3.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	0.5	1.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	0.5	1.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	135	0.5	1.7
...	All other causes (Residual)	4,355	15.6	55.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-59 years				
...	All causes	36,388	100.0	517.6
1	Malignant neoplasms (C00-C97)	15,604	42.9	222.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,702	15.7	81.1
3	Chronic lower respiratory diseases (J40-J47)	1,823	5.0	25.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,665	4.6	23.7
5	Diabetes mellitus (E10-E14)	1,206	3.3	17.2
6	Cerebrovascular diseases (I60-I69)	1,180	3.2	16.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	889	2.4	12.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	691	1.9	9.8
9	Septicemia (A40-A41)	592	1.6	8.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	450	1.2	6.4
11	Influenza and pneumonia (J09-J18)	384	1.1	5.5
12	Viral hepatitis (B15-B19)	256	0.7	3.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	195	0.5	2.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	153	0.4	2.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	139	0.4	2.0
...	All other causes (Residual)	5,459	15.0	77.7
Non-Hispanic, white, female, 60-64 years				
...	All causes	47,511	100.0	811.8
1	Malignant neoplasms (C00-C97)	20,546	43.2	351.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,100	17.0	138.4
3	Chronic lower respiratory diseases (J40-J47)	3,506	7.4	59.9
4	Cerebrovascular diseases (I60-I69)	1,661	3.5	28.4
5	Diabetes mellitus (E10-E14)	1,623	3.4	27.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,327	2.8	22.7
7	Septicemia (A40-A41)	805	1.7	13.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	796	1.7	13.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	626	1.3	10.7
10	Influenza and pneumonia (J09-J18)	528	1.1	9.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	452	1.0	7.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	265	0.6	4.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	255	0.5	4.4
14	Alzheimer's disease (G30)	234	0.5	4.0
15	Aortic aneurysm and dissection (I71)	165	0.3	2.8
...	All other causes (Residual)	6,622	13.9	113.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 65-69 years				
...	All causes	57,469	100.0	1,289.9
1	Malignant neoplasms (C00-C97)	23,732	41.3	532.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,181	17.7	228.5
3	Chronic lower respiratory diseases (J40-J47)	5,192	9.0	116.5
4	Cerebrovascular diseases (I60-I69)	2,408	4.2	54.0
5	Diabetes mellitus (E10-E14)	1,823	3.2	40.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,160	2.0	26.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	964	1.7	21.6
8	Septicemia (A40-A41)	945	1.6	21.2
9	Influenza and pneumonia (J09-J18)	738	1.3	16.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	718	1.2	16.1
11	Alzheimer's disease (G30)	507	0.9	11.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	371	0.6	8.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	350	0.6	7.9
14	Aortic aneurysm and dissection (I71)	264	0.5	5.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	241	0.4	5.4
...	All other causes (Residual)	7,875	13.7	176.8
Non-Hispanic, white, female, 70-74 years				
...	All causes	76,006	100.0	2,049.4
1	Malignant neoplasms (C00-C97)	27,068	35.6	729.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,969	19.7	403.6
3	Chronic lower respiratory diseases (J40-J47)	7,530	9.9	203.0
4	Cerebrovascular diseases (I60-I69)	3,785	5.0	102.1
5	Diabetes mellitus (E10-E14)	2,465	3.2	66.5
6	Alzheimer's disease (G30)	1,437	1.9	38.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,390	1.8	37.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,372	1.8	37.0
9	Septicemia (A40-A41)	1,238	1.6	33.4
10	Influenza and pneumonia (J09-J18)	1,163	1.5	31.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	777	1.0	21.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	581	0.8	15.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	506	0.7	13.6
14	Parkinson's disease (G20-G21)	501	0.7	13.5
15	Aortic aneurysm and dissection (I71)	482	0.6	13.0
...	All other causes (Residual)	10,742	14.1	289.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-79 years				
...	All causes	114,122	100.0	3,344.3
1	Malignant neoplasms (C00-C97)	32,221	28.2	944.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,697	22.5	753.0
3	Chronic lower respiratory diseases (J40-J47)	10,529	9.2	308.5
4	Cerebrovascular diseases (I60-I69)	7,179	6.3	210.4
5	Alzheimer's disease (G30)	3,983	3.5	116.7
6	Diabetes mellitus (E10-E14)	3,522	3.1	103.2
7	Influenza and pneumonia (J09-J18)	2,260	2.0	66.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,233	2.0	65.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,224	1.9	65.2
10	Septicemia (A40-A41)	1,847	1.6	54.1
11	Parkinson's disease (G20-G21)	1,214	1.1	35.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	942	0.8	27.6
13	Aortic aneurysm and dissection (I71)	758	0.7	22.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	757	0.7	22.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	757	0.7	22.2
...	All other causes (Residual)	17,999	15.8	527.4
Non-Hispanic, white, female, 80-84 years				
...	All causes	163,232	100.0	5,605.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	42,670	26.1	1,465.3
2	Malignant neoplasms (C00-C97)	33,558	20.6	1,152.4
3	Cerebrovascular diseases (I60-I69)	12,706	7.8	436.3
4	Chronic lower respiratory diseases (J40-J47)	12,111	7.4	415.9
5	Alzheimer's disease (G30)	8,975	5.5	308.2
6	Diabetes mellitus (E10-E14)	4,420	2.7	151.8
7	Influenza and pneumonia (J09-J18)	3,916	2.4	134.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,413	2.1	117.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,259	2.0	111.9
10	Septicemia (A40-A41)	2,518	1.5	86.5
11	Parkinson's disease (G20-G21)	2,053	1.3	70.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,812	1.1	62.2
13	Pneumonitis due to solids and liquids (J69)	1,314	0.8	45.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,064	0.7	36.5
15	Aortic aneurysm and dissection (I71)	989	0.6	34.0
...	All other causes (Residual)	28,454	17.4	977.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 85-89 years				
...	All causes	186,333	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,188	30.2	---
2	Malignant neoplasms (C00-C97)	25,299	13.6	---
3	Cerebrovascular diseases (I60-I69)	16,211	8.7	---
4	Alzheimer's disease (G30)	13,366	7.2	---
5	Chronic lower respiratory diseases (J40-J47)	10,162	5.5	---
6	Influenza and pneumonia (J09-J18)	5,511	3.0	---
7	Diabetes mellitus (E10-E14)	4,190	2.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,048	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,706	2.0	---
10	Septicemia (A40-A41)	2,695	1.4	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,617	1.4	---
12	Parkinson's disease (G20-G21)	2,053	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,725	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,173	0.6	---
15	Atherosclerosis (I70)	1,058	0.6	---
...	All other causes (Residual)	36,331	19.5	---
Non-Hispanic, white, female, 90-94 years				
...	All causes	144,076	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,120	34.1	---
2	Cerebrovascular diseases (I60-I69)	12,637	8.8	---
3	Malignant neoplasms (C00-C97)	12,331	8.6	---
4	Alzheimer's disease (G30)	11,840	8.2	---
5	Chronic lower respiratory diseases (J40-J47)	5,594	3.9	---
6	Influenza and pneumonia (J09-J18)	5,067	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,125	2.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,699	1.9	---
9	Diabetes mellitus (E10-E14)	2,475	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,300	1.6	---
11	Septicemia (A40-A41)	1,674	1.2	---
12	Pneumonitis due to solids and liquids (J69)	1,472	1.0	---
13	Parkinson's disease (G20-G21)	1,160	0.8	---
14	Atherosclerosis (I70)	1,006	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	768	0.5	---
...	All other causes (Residual)	30,808	21.4	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 95-99 years				
...	All causes	63,790	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,565	36.9	---
2	Alzheimer's disease (G30)	5,512	8.6	---
3	Cerebrovascular diseases (I60-I69)	5,257	8.2	---
4	Malignant neoplasms (C00-C97)	3,636	5.7	---
5	Influenza and pneumonia (J09-J18)	2,556	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	1,826	2.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,405	2.2	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,125	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,051	1.6	---
10	Diabetes mellitus (E10-E14)	743	1.2	---
11	Pneumonitis due to solids and liquids (J69)	671	1.1	---
12	Septicemia (A40-A41)	645	1.0	---
13	Atherosclerosis (I70)	615	1.0	---
14	Parkinson's disease (G20-G21)	289	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	281	0.4	---
...	All other causes (Residual)	14,613	22.9	---
Non-Hispanic, white, female, 100 years and over				
...	All causes	14,024	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,461	38.9	---
2	Alzheimer's disease (G30)	1,063	7.6	---
3	Cerebrovascular diseases (I60-I69)	940	6.7	---
4	Influenza and pneumonia (J09-J18)	738	5.3	---
5	Malignant neoplasms (C00-C97)	495	3.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	320	2.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	263	1.9	---
8	Chronic lower respiratory diseases (J40-J47)	260	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	234	1.7	---
10	Atherosclerosis (I70)	210	1.5	---
11	Pneumonitis due to solids and liquids (J69)	148	1.1	---
12	Diabetes mellitus (E10-E14)	133	0.9	---
13	Septicemia (A40-A41)	121	0.9	---
14	Anemias (D50-D64)	46	0.3	---
15	Nutritional deficiencies (E40-E64)	37	0.3	---
15	Parkinson's disease (G20-G21)	37	0.3	---
...	All other causes (Residual)	3,518	25.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	286,366	100.0	750.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,443	24.6	184.7
2	Malignant neoplasms (C00-C97)	63,441	22.2	166.3
3	Cerebrovascular diseases (I60-I69)	16,934	5.9	44.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,332	4.7	34.9
5	Diabetes mellitus (E10-E14)	12,343	4.3	32.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,746	3.1	22.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,318	2.9	21.8
8	Chronic lower respiratory diseases (J40-J47)	7,830	2.7	20.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,398	2.2	16.8
10	Septicemia (A40-A41)	6,241	2.2	16.4
11	Influenza and pneumonia (J09-J18)	5,091	1.8	13.3
12	Certain conditions originating in the perinatal period (P00-P96)	4,868	1.7	12.8
13	Alzheimer's disease (G30)	4,729	1.7	12.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,597	1.6	12.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,525	0.9	6.6
...	All other causes (Residual)	50,530	17.6	132.5
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,102	100.0	43.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	318	28.9	12.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	160	14.5	6.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	116	10.5	4.6
4	Malignant neoplasms (C00-C97)	51	4.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.8	1.7
6	Chronic lower respiratory diseases (J40-J47)	29	2.6	1.2
7	Influenza and pneumonia (J09-J18)	28	2.5	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	26	2.4	1.0
9	Septicemia (A40-A41)	16	1.5	*
9	Cerebrovascular diseases (I60-I69)	16	1.5	*
11	Anemias (D50-D64)	12	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
...	All other causes (Residual)	262	23.8	10.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-9 years				
...	All causes	563	100.0	18.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	216	38.4	7.1
2	Malignant neoplasms (C00-C97)	57	10.1	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	7.3	1.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	6.6	1.2
5	Chronic lower respiratory diseases (J40-J47)	24	4.3	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.1	0.8
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.6	*
8	Influenza and pneumonia (J09-J18)	7	1.2	*
9	Septicemia (A40-A41)	6	1.1	*
9	Cerebrovascular diseases (I60-I69)	6	1.1	*
11	Anemias (D50-D64)	5	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
13	Meningococcal infection (A39)	3	0.5	*
13	Diseases of appendix (K35-K38)	3	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	119	21.1	3.9
Non-Hispanic, black, both sexes, 10-14 years				
...	All causes	772	100.0	24.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	236	30.6	7.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	101	13.1	3.2
3	Malignant neoplasms (C00-C97)	77	10.0	2.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	5.2	1.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	5.1	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	4.7	1.1
7	Chronic lower respiratory diseases (J40-J47)	35	4.5	1.1
8	Influenza and pneumonia (J09-J18)	14	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.0	*
10	Anemias (D50-D64)	8	1.0	*
12	Cerebrovascular diseases (I60-I69)	7	0.9	*
13	Septicemia (A40-A41)	4	0.5	*
13	Diabetes mellitus (E10-E14)	4	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
...	All other causes (Residual)	148	19.2	4.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 15-19 years				
...	All causes	2,893	100.0	85.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,298	44.9	38.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	723	25.0	21.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	139	4.8	4.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	3.8	3.3
5	Malignant neoplasms (C00-C97)	104	3.6	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	1.6	1.3
7	Chronic lower respiratory diseases (J40-J47)	28	1.0	0.8
8	Anemias (D50-D64)	26	0.9	0.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	25	0.9	0.7
10	Diabetes mellitus (E10-E14)	20	0.7	0.6
10	Pregnancy, childbirth and the puerperium (O00-O99)	20	0.7	0.6
12	Cerebrovascular diseases (I60-I69)	16	0.6	*
13	Septicemia (A40-A41)	12	0.4	*
14	Influenza and pneumonia (J09-J18)	8	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
14	Legal intervention (Y35,Y89.0)	8	0.3	*
...	All other causes (Residual)	302	10.4	9.0
Non-Hispanic, black, both sexes, 20-24 years				
...	All causes	4,344	100.0	142.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,864	42.9	61.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	976	22.5	31.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	257	5.9	8.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	233	5.4	7.6
5	Malignant neoplasms (C00-C97)	154	3.5	5.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	84	1.9	2.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	48	1.1	1.6
8	Anemias (D50-D64)	40	0.9	1.3
9	Legal intervention (Y35,Y89.0)	39	0.9	1.3
10	Chronic lower respiratory diseases (J40-J47)	37	0.9	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	0.8	1.2
12	Septicemia (A40-A41)	35	0.8	1.1
12	Diabetes mellitus (E10-E14)	35	0.8	1.1
14	Cerebrovascular diseases (I60-I69)	29	0.7	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.5	0.7
...	All other causes (Residual)	457	10.5	15.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-29 years				
...	All causes	4,672	100.0	161.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,498	32.1	51.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	914	19.6	31.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	395	8.5	13.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	283	6.1	9.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	244	5.2	8.5
6	Malignant neoplasms (C00-C97)	232	5.0	8.0
7	Diabetes mellitus (E10-E14)	83	1.8	2.9
8	Pregnancy, childbirth and the puerperium (O00-O99)	56	1.2	1.9
9	Anemias (D50-D64)	51	1.1	1.8
10	Influenza and pneumonia (J09-J18)	46	1.0	1.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.0	1.6
12	Septicemia (A40-A41)	45	1.0	1.6
13	Cerebrovascular diseases (I60-I69)	43	0.9	1.5
13	Chronic lower respiratory diseases (J40-J47)	43	0.9	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.9	1.4
...	All other causes (Residual)	653	14.0	22.6
Non-Hispanic, black, both sexes, 30-34 years				
...	All causes	4,849	100.0	192.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,026	21.2	40.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	834	17.2	33.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	590	12.2	23.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	418	8.6	16.6
5	Malignant neoplasms (C00-C97)	380	7.8	15.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	227	4.7	9.0
7	Diabetes mellitus (E10-E14)	129	2.7	5.1
8	Cerebrovascular diseases (I60-I69)	105	2.2	4.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	1.3	2.5
10	Anemias (D50-D64)	50	1.0	2.0
10	Chronic lower respiratory diseases (J40-J47)	50	1.0	2.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	50	1.0	2.0
13	Influenza and pneumonia (J09-J18)	46	0.9	1.8
14	Septicemia (A40-A41)	44	0.9	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	0.9	1.7
...	All other causes (Residual)	794	16.4	31.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 35-39 years				
...	All causes	6,751	100.0	254.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,071	15.9	40.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,006	14.9	37.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	809	12.0	30.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	770	11.4	29.0
5	Malignant neoplasms (C00-C97)	703	10.4	26.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	220	3.3	8.3
7	Cerebrovascular diseases (I60-I69)	214	3.2	8.1
8	Diabetes mellitus (E10-E14)	208	3.1	7.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	2.0	5.1
10	Septicemia (A40-A41)	107	1.6	4.0
11	Influenza and pneumonia (J09-J18)	70	1.0	2.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	65	1.0	2.4
12	Chronic lower respiratory diseases (J40-J47)	65	1.0	2.4
14	Anemias (D50-D64)	62	0.9	2.3
15	Aortic aneurysm and dissection (I71)	46	0.7	1.7
...	All other causes (Residual)	1,200	17.8	45.2
Non-Hispanic, black, both sexes, 40-44 years				
...	All causes	10,046	100.0	366.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,891	18.8	68.9
2	Malignant neoplasms (C00-C97)	1,629	16.2	59.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,181	11.8	43.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,112	11.1	40.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	585	5.8	21.3
6	Cerebrovascular diseases (I60-I69)	434	4.3	15.8
7	Diabetes mellitus (E10-E14)	346	3.4	12.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	195	1.9	7.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	194	1.9	7.1
10	Septicemia (A40-A41)	166	1.7	6.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	159	1.6	5.8
12	Chronic lower respiratory diseases (J40-J47)	128	1.3	4.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	109	1.1	4.0
14	Influenza and pneumonia (J09-J18)	105	1.0	3.8
15	Anemias (D50-D64)	77	0.8	2.8
...	All other causes (Residual)	1,735	17.3	63.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-49 years				
...	All causes	15,453	100.0	565.0
1	Malignant neoplasms (C00-C97)	3,339	21.6	122.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,286	21.3	120.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,373	8.9	50.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,234	8.0	45.1
5	Cerebrovascular diseases (I60-I69)	789	5.1	28.8
6	Diabetes mellitus (E10-E14)	533	3.4	19.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	476	3.1	17.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	346	2.2	12.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	338	2.2	12.4
10	Septicemia (A40-A41)	302	2.0	11.0
11	Chronic lower respiratory diseases (J40-J47)	260	1.7	9.5
12	Influenza and pneumonia (J09-J18)	205	1.3	7.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	200	1.3	7.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	174	1.1	6.4
15	Viral hepatitis (B15-B19)	121	0.8	4.4
...	All other causes (Residual)	2,477	16.0	90.6
Non-Hispanic, black, both sexes, 50-54 years				
...	All causes	21,012	100.0	883.6
1	Malignant neoplasms (C00-C97)	5,415	25.8	227.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,999	23.8	210.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,351	6.4	56.8
4	Cerebrovascular diseases (I60-I69)	1,137	5.4	47.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,084	5.2	45.6
6	Diabetes mellitus (E10-E14)	879	4.2	37.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	497	2.4	20.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	477	2.3	20.1
9	Septicemia (A40-A41)	413	2.0	17.4
10	Chronic lower respiratory diseases (J40-J47)	409	1.9	17.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	331	1.6	13.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	320	1.5	13.5
13	Viral hepatitis (B15-B19)	254	1.2	10.7
14	Influenza and pneumonia (J09-J18)	229	1.1	9.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	143	0.7	6.0
...	All other causes (Residual)	3,074	14.6	129.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 55-59 years				
...	All causes	23,951	100.0	1,243.2
1	Malignant neoplasms (C00-C97)	7,146	29.8	370.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,038	25.2	313.4
3	Cerebrovascular diseases (I60-I69)	1,294	5.4	67.2
4	Diabetes mellitus (E10-E14)	1,190	5.0	61.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,001	4.2	52.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	633	2.6	32.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	623	2.6	32.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	538	2.2	27.9
9	Chronic lower respiratory diseases (J40-J47)	519	2.2	26.9
10	Septicemia (A40-A41)	503	2.1	26.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	362	1.5	18.8
12	Influenza and pneumonia (J09-J18)	287	1.2	14.9
13	Viral hepatitis (B15-B19)	279	1.2	14.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	209	0.9	10.8
15	Aortic aneurysm and dissection (I71)	122	0.5	6.3
...	All other causes (Residual)	3,207	13.4	166.5
Non-Hispanic, black, both sexes, 60-64 years				
...	All causes	23,248	100.0	1,738.9
1	Malignant neoplasms (C00-C97)	7,476	32.2	559.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,118	26.3	457.6
3	Cerebrovascular diseases (I60-I69)	1,278	5.5	95.6
4	Diabetes mellitus (E10-E14)	1,270	5.5	95.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	744	3.2	55.6
6	Chronic lower respiratory diseases (J40-J47)	647	2.8	48.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	583	2.5	43.6
8	Septicemia (A40-A41)	493	2.1	36.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	383	1.6	28.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	357	1.5	26.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	319	1.4	23.9
12	Influenza and pneumonia (J09-J18)	280	1.2	20.9
13	Viral hepatitis (B15-B19)	120	0.5	9.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	107	0.5	8.0
15	Aortic aneurysm and dissection (I71)	105	0.5	7.9
...	All other causes (Residual)	2,968	12.8	222.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-69 years				
...	All causes	24,324	100.0	2,433.4
1	Malignant neoplasms (C00-C97)	7,943	32.7	794.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,380	26.2	638.3
3	Cerebrovascular diseases (I60-I69)	1,490	6.1	149.1
4	Diabetes mellitus (E10-E14)	1,325	5.4	132.6
5	Chronic lower respiratory diseases (J40-J47)	864	3.6	86.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	826	3.4	82.6
7	Septicemia (A40-A41)	617	2.5	61.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	469	1.9	46.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	398	1.6	39.8
10	Influenza and pneumonia (J09-J18)	328	1.3	32.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	220	0.9	22.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	146	0.6	14.6
13	Aortic aneurysm and dissection (I71)	127	0.5	12.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	105	0.4	10.5
15	Pneumonitis due to solids and liquids (J69)	83	0.3	8.3
...	All other causes (Residual)	3,003	12.3	300.4
Non-Hispanic, black, both sexes, 70-74 years				
...	All causes	26,566	100.0	3,373.8
1	Malignant neoplasms (C00-C97)	7,944	29.9	1,008.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,127	26.8	905.1
3	Cerebrovascular diseases (I60-I69)	1,734	6.5	220.2
4	Diabetes mellitus (E10-E14)	1,518	5.7	192.8
5	Chronic lower respiratory diseases (J40-J47)	1,022	3.8	129.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	998	3.8	126.7
7	Septicemia (A40-A41)	697	2.6	88.5
8	Influenza and pneumonia (J09-J18)	463	1.7	58.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	431	1.6	54.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	424	1.6	53.8
11	Alzheimer's disease (G30)	281	1.1	35.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	179	0.7	22.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	144	0.5	18.3
14	Pneumonitis due to solids and liquids (J69)	122	0.5	15.5
15	Aortic aneurysm and dissection (I71)	109	0.4	13.8
...	All other causes (Residual)	3,373	12.7	428.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 75-79 years				
...	All causes	28,763	100.0	4,924.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,825	27.2	1,339.6
2	Malignant neoplasms (C00-C97)	7,655	26.6	1,310.5
3	Cerebrovascular diseases (I60-I69)	2,055	7.1	351.8
4	Diabetes mellitus (E10-E14)	1,523	5.3	260.7
5	Chronic lower respiratory diseases (J40-J47)	1,205	4.2	206.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,041	3.6	178.2
7	Septicemia (A40-A41)	732	2.5	125.3
8	Alzheimer's disease (G30)	615	2.1	105.3
9	Influenza and pneumonia (J09-J18)	574	2.0	98.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	519	1.8	88.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	403	1.4	69.0
12	Pneumonitis due to solids and liquids (J69)	178	0.6	30.5
13	Parkinson's disease (G20-G21)	149	0.5	25.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	144	0.5	24.7
15	Aortic aneurysm and dissection (I71)	136	0.5	23.3
...	All other causes (Residual)	4,009	13.9	686.3
Non-Hispanic, black, both sexes, 80-84 years				
...	All causes	29,635	100.0	7,076.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,534	28.8	2,037.8
2	Malignant neoplasms (C00-C97)	6,487	21.9	1,549.0
3	Cerebrovascular diseases (I60-I69)	2,291	7.7	547.1
4	Diabetes mellitus (E10-E14)	1,508	5.1	360.1
5	Chronic lower respiratory diseases (J40-J47)	1,110	3.7	265.1
6	Alzheimer's disease (G30)	1,005	3.4	240.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,005	3.4	240.0
8	Septicemia (A40-A41)	768	2.6	183.4
9	Influenza and pneumonia (J09-J18)	724	2.4	172.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	566	1.9	135.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	353	1.2	84.3
12	Pneumonitis due to solids and liquids (J69)	258	0.9	61.6
13	Parkinson's disease (G20-G21)	209	0.7	49.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	162	0.5	38.7
15	Aortic aneurysm and dissection (I71)	131	0.4	31.3
...	All other causes (Residual)	4,524	15.3	1,080.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85-89 years				
...	All causes	24,656	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,506	30.4	---
2	Malignant neoplasms (C00-C97)	4,101	16.6	---
3	Cerebrovascular diseases (I60-I69)	2,044	8.3	---
4	Alzheimer's disease (G30)	1,134	4.6	---
5	Diabetes mellitus (E10-E14)	1,041	4.2	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	959	3.9	---
7	Chronic lower respiratory diseases (J40-J47)	769	3.1	---
8	Influenza and pneumonia (J09-J18)	741	3.0	---
9	Septicemia (A40-A41)	626	2.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	557	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	1.3	---
12	Pneumonitis due to solids and liquids (J69)	260	1.1	---
13	Parkinson's disease (G20-G21)	149	0.6	---
14	Atherosclerosis (I70)	112	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	107	0.4	---
...	All other causes (Residual)	4,236	17.2	---
Non-Hispanic, black, both sexes, 90-94 years				
...	All causes	15,275	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,929	32.3	---
2	Malignant neoplasms (C00-C97)	1,863	12.2	---
3	Cerebrovascular diseases (I60-I69)	1,292	8.5	---
4	Alzheimer's disease (G30)	972	6.4	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	504	3.3	---
6	Diabetes mellitus (E10-E14)	499	3.3	---
7	Influenza and pneumonia (J09-J18)	493	3.2	---
8	Chronic lower respiratory diseases (J40-J47)	395	2.6	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	374	2.4	---
10	Septicemia (A40-A41)	370	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	1.3	---
12	Pneumonitis due to solids and liquids (J69)	152	1.0	---
13	Atherosclerosis (I70)	90	0.6	---
14	Parkinson's disease (G20-G21)	77	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.4	---
...	All other causes (Residual)	3,005	19.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 95-99 years				
...	All causes	6,731	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,412	35.8	---
2	Malignant neoplasms (C00-C97)	560	8.3	---
3	Cerebrovascular diseases (I60-I69)	489	7.3	---
4	Alzheimer's disease (G30)	454	6.7	---
5	Influenza and pneumonia (J09-J18)	249	3.7	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	3.1	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	189	2.8	---
8	Diabetes mellitus (E10-E14)	182	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	143	2.1	---
10	Septicemia (A40-A41)	140	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81	1.2	---
12	Pneumonitis due to solids and liquids (J69)	64	1.0	---
13	Atherosclerosis (I70)	54	0.8	---
14	Parkinson's disease (G20-G21)	25	0.4	---
15	Nutritional deficiencies (E40-E64)	24	0.4	---
...	All other causes (Residual)	1,457	21.6	---
Non-Hispanic, black, both sexes, 100 years and over				
...	All causes	2,100	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	754	35.9	---
2	Alzheimer's disease (G30)	143	6.8	---
3	Cerebrovascular diseases (I60-I69)	136	6.5	---
4	Malignant neoplasms (C00-C97)	112	5.3	---
5	Influenza and pneumonia (J09-J18)	105	5.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	65	3.1	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	60	2.9	---
8	Diabetes mellitus (E10-E14)	45	2.1	---
9	Septicemia (A40-A41)	35	1.7	---
10	Chronic lower respiratory diseases (J40-J47)	34	1.6	---
11	Pneumonitis due to solids and liquids (J69)	23	1.1	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	1.0	---
13	Atherosclerosis (I70)	21	1.0	---
14	Nutritional deficiencies (E40-E64)	10	0.5	---
15	Parkinson's disease (G20-G21)	8	0.4	---
...	All other causes (Residual)	527	25.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, all ages				
...	All causes	146,474	100.0	805.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,243	24.1	193.7
2	Malignant neoplasms (C00-C97)	32,766	22.4	180.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,093	6.2	50.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,477	5.1	41.1
5	Cerebrovascular diseases (I60-I69)	7,470	5.1	41.1
6	Diabetes mellitus (E10-E14)	5,434	3.7	29.9
7	Chronic lower respiratory diseases (J40-J47)	4,167	2.8	22.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4,127	2.8	22.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,730	2.5	20.5
10	Septicemia (A40-A41)	2,807	1.9	15.4
11	Certain conditions originating in the perinatal period (P00-P96)	2,734	1.9	15.0
12	Influenza and pneumonia (J09-J18)	2,464	1.7	13.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,963	1.3	10.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,662	1.1	9.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,571	1.1	8.6
...	All other causes (Residual)	23,766	16.2	130.6
Non-Hispanic, black, male, 1-4 years				
...	All causes	609	100.0	47.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	30.5	14.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	14.4	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	9.4	4.4
4	Malignant neoplasms (C00-C97)	31	5.1	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.6	2.2
6	Influenza and pneumonia (J09-J18)	15	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	13	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	2.0	*
9	Septicemia (A40-A41)	11	1.8	*
10	Anemias (D50-D64)	4	0.7	*
10	Cerebrovascular diseases (I60-I69)	4	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
13	Meningitis (G00,G03)	3	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (Residual)	144	23.6	11.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 5-9 years				
...	All causes	325	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	40.6	8.6
2	Malignant neoplasms (C00-C97)	29	8.9	1.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	6.5	1.4
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	6.2	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	4.9	*
5	Chronic lower respiratory diseases (J40-J47)	16	4.9	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.8	*
8	Influenza and pneumonia (J09-J18)	4	1.2	*
9	Meningococcal infection (A39)	2	0.6	*
9	Septicemia (A40-A41)	2	0.6	*
9	Anemias (D50-D64)	2	0.6	*
9	Meningitis (G00,G03)	2	0.6	*
9	Cerebrovascular diseases (I60-I69)	2	0.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.6	*
...	All other causes (Residual)	67	20.6	4.4
Non-Hispanic, black, male, 10-14 years				
...	All causes	476	100.0	29.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	30.0	9.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	15.3	4.6
3	Malignant neoplasms (C00-C97)	41	8.6	2.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	5.7	1.7
5	Chronic lower respiratory diseases (J40-J47)	24	5.0	1.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	4.0	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	3.6	*
8	Influenza and pneumonia (J09-J18)	9	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.7	*
10	Anemias (D50-D64)	6	1.3	*
11	Septicemia (A40-A41)	2	0.4	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
11	Diabetes mellitus (E10-E14)	2	0.4	*
11	Meningitis (G00,G03)	2	0.4	*
11	Pneumonitis due to solids and liquids (J69)	2	0.4	*
11	Diseases of appendix (K35-K38)	2	0.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	93	19.5	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-19 years				
...	All causes	2,247	100.0	131.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,181	52.6	69.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	529	23.5	30.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	118	5.3	6.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	3.5	4.6
5	Malignant neoplasms (C00-C97)	54	2.4	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	1.2	1.6
7	Chronic lower respiratory diseases (J40-J47)	17	0.8	*
8	Anemias (D50-D64)	12	0.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.5	*
9	Cerebrovascular diseases (I60-I69)	11	0.5	*
11	Diabetes mellitus (E10-E14)	8	0.4	*
11	Legal intervention (Y35,Y89.0)	8	0.4	*
13	Septicemia (A40-A41)	6	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	*
14	Aortic aneurysm and dissection (I71)	4	0.2	*
...	All other causes (Residual)	178	7.9	10.4
Non-Hispanic, black, male, 20-24 years				
...	All causes	3,388	100.0	219.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,690	49.9	109.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	738	21.8	47.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	225	6.6	14.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	154	4.5	10.0
5	Malignant neoplasms (C00-C97)	88	2.6	5.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	45	1.3	2.9
7	Legal intervention (Y35,Y89.0)	37	1.1	2.4
8	Chronic lower respiratory diseases (J40-J47)	27	0.8	1.7
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	0.7	1.6
10	Anemias (D50-D64)	24	0.7	1.6
11	Cerebrovascular diseases (I60-I69)	19	0.6	*
12	Septicemia (A40-A41)	16	0.5	*
13	Diabetes mellitus (E10-E14)	13	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.3	*
15	Influenza and pneumonia (J09-J18)	9	0.3	*
...	All other causes (Residual)	267	7.9	17.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 25-29 years				
...	All causes	3,341	100.0	236.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,334	39.9	94.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	710	21.3	50.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	253	7.6	17.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	205	6.1	14.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	151	4.5	10.7
6	Malignant neoplasms (C00-C97)	109	3.3	7.7
7	Diabetes mellitus (E10-E14)	42	1.3	3.0
8	Legal intervention (Y35,Y89.0)	30	0.9	2.1
9	Anemias (D50-D64)	29	0.9	2.1
10	Influenza and pneumonia (J09-J18)	27	0.8	1.9
11	Chronic lower respiratory diseases (J40-J47)	26	0.8	1.8
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	0.7	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.7	1.7
14	Septicemia (A40-A41)	21	0.6	1.5
15	Cerebrovascular diseases (I60-I69)	19	0.6	*
...	All other causes (Residual)	336	10.1	23.8
Non-Hispanic, black, male, 30-34 years				
...	All causes	3,197	100.0	267.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	915	28.6	76.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	632	19.8	52.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	370	11.6	30.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	221	6.9	18.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	194	6.1	16.2
6	Malignant neoplasms (C00-C97)	151	4.7	12.6
7	Diabetes mellitus (E10-E14)	74	2.3	6.2
8	Cerebrovascular diseases (I60-I69)	52	1.6	4.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.1	2.9
10	Chronic lower respiratory diseases (J40-J47)	30	0.9	2.5
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	0.9	2.3
12	Anemias (D50-D64)	26	0.8	2.2
13	Influenza and pneumonia (J09-J18)	24	0.8	2.0
14	Septicemia (A40-A41)	19	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.6	*
...	All other causes (Residual)	408	12.8	34.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-39 years				
...	All causes	4,092	100.0	326.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	733	17.9	58.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	691	16.9	55.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	634	15.5	50.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	464	11.3	37.0
5	Malignant neoplasms (C00-C97)	245	6.0	19.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	4.3	14.0
7	Cerebrovascular diseases (I60-I69)	124	3.0	9.9
8	Diabetes mellitus (E10-E14)	121	3.0	9.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	1.5	4.9
10	Septicemia (A40-A41)	49	1.2	3.9
11	Influenza and pneumonia (J09-J18)	35	0.9	2.8
12	Anemias (D50-D64)	34	0.8	2.7
13	Aortic aneurysm and dissection (I71)	33	0.8	2.6
13	Chronic lower respiratory diseases (J40-J47)	33	0.8	2.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	0.8	2.5
...	All other causes (Residual)	629	15.4	50.2
Non-Hispanic, black, male, 40-44 years				
...	All causes	5,889	100.0	458.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,238	21.0	96.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	798	13.6	62.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	753	12.8	58.6
4	Malignant neoplasms (C00-C97)	650	11.0	50.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	456	7.7	35.5
6	Cerebrovascular diseases (I60-I69)	206	3.5	16.0
7	Diabetes mellitus (E10-E14)	188	3.2	14.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	154	2.6	12.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	2.0	9.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	96	1.6	7.5
11	Septicemia (A40-A41)	77	1.3	6.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	69	1.2	5.4
13	Chronic lower respiratory diseases (J40-J47)	66	1.1	5.1
14	Influenza and pneumonia (J09-J18)	65	1.1	5.1
15	Aortic aneurysm and dissection (I71)	53	0.9	4.1
...	All other causes (Residual)	905	15.4	70.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-49 years				
...	All causes	8,858	100.0	696.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,050	23.1	161.3
2	Malignant neoplasms (C00-C97)	1,530	17.3	120.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	969	10.9	76.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	792	8.9	62.3
5	Cerebrovascular diseases (I60-I69)	399	4.5	31.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	357	4.0	28.1
7	Diabetes mellitus (E10-E14)	315	3.6	24.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	204	2.3	16.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	171	1.9	13.5
10	Septicemia (A40-A41)	152	1.7	12.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	134	1.5	10.5
12	Influenza and pneumonia (J09-J18)	129	1.5	10.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	114	1.3	9.0
13	Chronic lower respiratory diseases (J40-J47)	114	1.3	9.0
15	Viral hepatitis (B15-B19)	76	0.9	6.0
...	All other causes (Residual)	1,352	15.3	106.4
Non-Hispanic, black, male, 50-54 years				
...	All causes	12,402	100.0	1,137.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,201	25.8	293.7
2	Malignant neoplasms (C00-C97)	2,804	22.6	257.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	950	7.7	87.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	741	6.0	68.0
5	Cerebrovascular diseases (I60-I69)	595	4.8	54.6
6	Diabetes mellitus (E10-E14)	469	3.8	43.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	348	2.8	31.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	260	2.1	23.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	259	2.1	23.8
10	Septicemia (A40-A41)	224	1.8	20.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	191	1.5	17.5
12	Chronic lower respiratory diseases (J40-J47)	189	1.5	17.3
13	Viral hepatitis (B15-B19)	173	1.4	15.9
14	Influenza and pneumonia (J09-J18)	138	1.1	12.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	108	0.9	9.9
...	All other causes (Residual)	1,752	14.1	160.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-59 years				
...	All causes	14,320	100.0	1,651.2
1	Malignant neoplasms (C00-C97)	3,953	27.6	455.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,876	27.1	446.9
3	Cerebrovascular diseases (I60-I69)	771	5.4	88.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	726	5.1	83.7
5	Diabetes mellitus (E10-E14)	643	4.5	74.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	445	3.1	51.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	380	2.7	43.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	328	2.3	37.8
9	Chronic lower respiratory diseases (J40-J47)	271	1.9	31.2
10	Septicemia (A40-A41)	270	1.9	31.1
11	Viral hepatitis (B15-B19)	201	1.4	23.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	188	1.3	21.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	174	1.2	20.1
14	Influenza and pneumonia (J09-J18)	157	1.1	18.1
15	Aortic aneurysm and dissection (I71)	84	0.6	9.7
...	All other causes (Residual)	1,853	12.9	213.7
Non-Hispanic, black, male, 60-64 years				
...	All causes	13,503	100.0	2,301.6
1	Malignant neoplasms (C00-C97)	4,160	30.8	709.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,797	28.1	647.2
3	Cerebrovascular diseases (I60-I69)	715	5.3	121.9
4	Diabetes mellitus (E10-E14)	633	4.7	107.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	412	3.1	70.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	395	2.9	67.3
7	Chronic lower respiratory diseases (J40-J47)	392	2.9	66.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	286	2.1	48.7
9	Septicemia (A40-A41)	254	1.9	43.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	218	1.6	37.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	214	1.6	36.5
12	Influenza and pneumonia (J09-J18)	171	1.3	29.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	92	0.7	15.7
14	Viral hepatitis (B15-B19)	86	0.6	14.7
15	Aortic aneurysm and dissection (I71)	60	0.4	10.2
...	All other causes (Residual)	1,618	12.0	275.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 65-69 years				
...	All causes	13,452	100.0	3,178.7
1	Malignant neoplasms (C00-C97)	4,351	32.3	1,028.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,679	27.3	869.3
3	Cerebrovascular diseases (I60-I69)	807	6.0	190.7
4	Diabetes mellitus (E10-E14)	633	4.7	149.6
5	Chronic lower respiratory diseases (J40-J47)	485	3.6	114.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	414	3.1	97.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	332	2.5	78.5
8	Septicemia (A40-A41)	313	2.3	74.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	206	1.5	48.7
10	Influenza and pneumonia (J09-J18)	192	1.4	45.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	159	1.2	37.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	111	0.8	26.2
13	Aortic aneurysm and dissection (I71)	70	0.5	16.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.4	13.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	0.4	13.0
...	All other causes (Residual)	1,586	11.8	374.8
Non-Hispanic, black, male, 70-74 years				
...	All causes	13,925	100.0	4,336.7
1	Malignant neoplasms (C00-C97)	4,299	30.9	1,338.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,826	27.5	1,191.6
3	Cerebrovascular diseases (I60-I69)	885	6.4	275.6
4	Diabetes mellitus (E10-E14)	675	4.8	210.2
5	Chronic lower respiratory diseases (J40-J47)	570	4.1	177.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	451	3.2	140.5
7	Septicemia (A40-A41)	345	2.5	107.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	262	1.9	81.6
9	Influenza and pneumonia (J09-J18)	259	1.9	80.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	202	1.5	62.9
11	Alzheimer's disease (G30)	123	0.9	38.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	112	0.8	34.9
13	Pneumonitis due to solids and liquids (J69)	78	0.6	24.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.5	23.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	59	0.4	18.4
...	All other causes (Residual)	1,703	12.2	530.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-79 years				
...	All causes	13,839	100.0	6,308.6
1	Malignant neoplasms (C00-C97)	3,962	28.6	1,806.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,788	27.4	1,726.8
3	Cerebrovascular diseases (I60-I69)	916	6.6	417.6
4	Chronic lower respiratory diseases (J40-J47)	682	4.9	310.9
5	Diabetes mellitus (E10-E14)	603	4.4	274.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	448	3.2	204.2
7	Septicemia (A40-A41)	315	2.3	143.6
8	Influenza and pneumonia (J09-J18)	300	2.2	136.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	230	1.7	104.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	228	1.6	103.9
11	Alzheimer's disease (G30)	227	1.6	103.5
12	Pneumonitis due to solids and liquids (J69)	106	0.8	48.3
13	Parkinson's disease (G20-G21)	85	0.6	38.7
14	Aortic aneurysm and dissection (I71)	72	0.5	32.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	70	0.5	31.9
...	All other causes (Residual)	1,807	13.1	823.7
Non-Hispanic, black, male, 80-84 years				
...	All causes	12,684	100.0	8,795.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,591	28.3	2,490.2
2	Malignant neoplasms (C00-C97)	3,216	25.4	2,230.2
3	Cerebrovascular diseases (I60-I69)	907	7.2	629.0
4	Chronic lower respiratory diseases (J40-J47)	601	4.7	416.8
5	Diabetes mellitus (E10-E14)	550	4.3	381.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	409	3.2	283.6
7	Influenza and pneumonia (J09-J18)	334	2.6	231.6
8	Alzheimer's disease (G30)	310	2.4	215.0
9	Septicemia (A40-A41)	285	2.2	197.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	203	1.6	140.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	1.3	112.3
12	Pneumonitis due to solids and liquids (J69)	136	1.1	94.3
13	Parkinson's disease (G20-G21)	114	0.9	79.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.6	52.7
15	Aortic aneurysm and dissection (I71)	44	0.3	30.5
...	All other causes (Residual)	1,746	13.8	1,210.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 85-89 years				
...	All causes	8,980	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,583	28.8	---
2	Malignant neoplasms (C00-C97)	2,013	22.4	---
3	Cerebrovascular diseases (I60-I69)	597	6.6	---
4	Chronic lower respiratory diseases (J40-J47)	355	4.0	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	335	3.7	---
6	Alzheimer's disease (G30)	314	3.5	---
7	Diabetes mellitus (E10-E14)	312	3.5	---
8	Influenza and pneumonia (J09-J18)	308	3.4	---
9	Septicemia (A40-A41)	224	2.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	161	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	1.7	---
12	Pneumonitis due to solids and liquids (J69)	117	1.3	---
13	Parkinson's disease (G20-G21)	71	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.5	---
15	Atherosclerosis (I70)	41	0.5	---
...	All other causes (Residual)	1,358	15.1	---
Non-Hispanic, black, male, 90-94 years				
...	All causes	4,375	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,328	30.4	---
2	Malignant neoplasms (C00-C97)	836	19.1	---
3	Cerebrovascular diseases (I60-I69)	299	6.8	---
4	Alzheimer's disease (G30)	195	4.5	---
5	Chronic lower respiratory diseases (J40-J47)	179	4.1	---
6	Influenza and pneumonia (J09-J18)	162	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	160	3.7	---
8	Diabetes mellitus (E10-E14)	121	2.8	---
9	Septicemia (A40-A41)	114	2.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	2.0	---
11	Pneumonitis due to solids and liquids (J69)	52	1.2	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	1.2	---
13	Parkinson's disease (G20-G21)	36	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.5	---
15	Atherosclerosis (I70)	22	0.5	---
...	All other causes (Residual)	707	16.2	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 95-99 years				
...	All causes	1,411	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	468	33.2	---
2	Malignant neoplasms (C00-C97)	198	14.0	---
3	Cerebrovascular diseases (I60-I69)	91	6.4	---
4	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	4.4	---
5	Alzheimer's disease (G30)	60	4.3	---
5	Influenza and pneumonia (J09-J18)	60	4.3	---
7	Chronic lower respiratory diseases (J40-J47)	55	3.9	---
8	Septicemia (A40-A41)	41	2.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	2.4	---
10	Diabetes mellitus (E10-E14)	26	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24	1.7	---
12	Pneumonitis due to solids and liquids (J69)	16	1.1	---
13	Atherosclerosis (I70)	9	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	---
14	Nutritional deficiencies (E40-E64)	7	0.5	---
...	All other causes (Residual)	253	17.9	---
Non-Hispanic, black, male, 100 years and over				
...	All causes	349	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	36.4	---
2	Malignant neoplasms (C00-C97)	38	10.9	---
3	Cerebrovascular diseases (I60-I69)	25	7.2	---
4	Influenza and pneumonia (J09-J18)	21	6.0	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	4.9	---
6	Alzheimer's disease (G30)	16	4.6	---
7	Chronic lower respiratory diseases (J40-J47)	10	2.9	---
8	Septicemia (A40-A41)	7	2.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	1.7	---
10	Diabetes mellitus (E10-E14)	4	1.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	1.1	---
12	Parkinson's disease (G20-G21)	2	0.6	---
12	Atherosclerosis (I70)	2	0.6	---
12	Pneumonitis due to solids and liquids (J69)	2	0.6	---
15	Nutritional deficiencies (E40-E64)	1	0.3	---
15	Aortic aneurysm and dissection (I71)	1	0.3	---
15	Hernia (K40-K46)	1	0.3	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	---
...	All other causes (Residual)	64	18.3	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	139,892	100.0	701.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,200	25.2	176.4
2	Malignant neoplasms (C00-C97)	30,675	21.9	153.7
3	Cerebrovascular diseases (I60-I69)	9,464	6.8	47.4
4	Diabetes mellitus (E10-E14)	6,909	4.9	34.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,588	3.3	23.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,239	3.0	21.2
7	Chronic lower respiratory diseases (J40-J47)	3,663	2.6	18.4
8	Alzheimer's disease (G30)	3,436	2.5	17.2
9	Septicemia (A40-A41)	3,434	2.5	17.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,634	1.9	13.2
11	Influenza and pneumonia (J09-J18)	2,627	1.9	13.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,271	1.6	11.4
13	Certain conditions originating in the perinatal period (P00-P96)	2,134	1.5	10.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,269	0.9	6.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	863	0.6	4.3
...	All other causes (Residual)	26,486	18.9	132.7
Non-Hispanic, black, female, 1-4 years				
...	All causes	493	100.0	39.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	26.8	10.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	14.6	5.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	12.0	4.8
4	Malignant neoplasms (C00-C97)	20	4.1	1.6
5	Chronic lower respiratory diseases (J40-J47)	16	3.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	14	2.8	*
8	Influenza and pneumonia (J09-J18)	13	2.6	*
9	Cerebrovascular diseases (I60-I69)	12	2.4	*
10	Anemias (D50-D64)	8	1.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.2	*
12	Septicemia (A40-A41)	5	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	115	23.3	9.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-9 years				
...	All causes	238	100.0	15.9
1	Accidents (unintentional injuries) (V01-X59, Y85-Y86)	84	35.3	5.6
2	Malignant neoplasms (C00-C97)	28	11.8	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	8.8	1.4
4	Assault (homicide) (*U01-*U02, X85-Y09, Y87.1)	16	6.7	*
5	Chronic lower respiratory diseases (J40-J47)	8	3.4	*
6	Diseases of heart (I00-I09, I11, I13, I20-I51)	7	2.9	*
7	Septicemia (A40-A41)	4	1.7	*
7	Cerebrovascular diseases (I60-I69)	4	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
9	Anemias (D50-D64)	3	1.3	*
9	Influenza and pneumonia (J09-J18)	3	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	3	1.3	*
13	Diabetes mellitus (E10-E14)	2	0.8	*
13	Diseases of appendix (K35-K38)	2	0.8	*
13	Complications of medical and surgical care (Y40-Y84, Y88)	2	0.8	*
...	All other causes (Residual)	48	20.2	3.2
Non-Hispanic, black, female, 10-14 years				
...	All causes	296	100.0	19.1
1	Accidents (unintentional injuries) (V01-X59, Y85-Y86)	93	31.4	6.0
2	Malignant neoplasms (C00-C97)	36	12.2	2.3
3	Assault (homicide) (*U01-*U02, X85-Y09, Y87.1)	28	9.5	1.8
4	Diseases of heart (I00-I09, I11, I13, I20-I51)	21	7.1	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	6.4	*
6	Intentional self-harm (suicide) (*U03, X60-X84, Y87.0)	12	4.1	*
7	Chronic lower respiratory diseases (J40-J47)	11	3.7	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	6	2.0	*
8	Cerebrovascular diseases (I60-I69)	6	2.0	*
10	Influenza and pneumonia (J09-J18)	5	1.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.0	*
12	Septicemia (A40-A41)	2	0.7	*
12	Anemias (D50-D64)	2	0.7	*
12	Diabetes mellitus (E10-E14)	2	0.7	*
12	Complications of medical and surgical care (Y40-Y84, Y88)	2	0.7	*
...	All other causes (Residual)	48	16.2	3.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 15-19 years				
...	All causes	646	100.0	38.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	30.0	11.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	117	18.1	7.0
3	Malignant neoplasms (C00-C97)	50	7.7	3.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	5.0	1.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	3.3	1.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	20	3.1	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	2.8	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	14	2.2	*
8	Anemias (D50-D64)	14	2.2	*
10	Diabetes mellitus (E10-E14)	12	1.9	*
11	Chronic lower respiratory diseases (J40-J47)	11	1.7	*
12	Septicemia (A40-A41)	6	0.9	*
13	Cerebrovascular diseases (I60-I69)	5	0.8	*
13	Influenza and pneumonia (J09-J18)	5	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.8	*
...	All other causes (Residual)	122	18.9	7.3
Non-Hispanic, black, female, 20-24 years				
...	All causes	956	100.0	63.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	24.9	15.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	174	18.2	11.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	8.3	5.2
4	Malignant neoplasms (C00-C97)	66	6.9	4.4
5	Pregnancy, childbirth and the puerperium (O00-O99)	48	5.0	3.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	39	4.1	2.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	3.3	2.1
8	Diabetes mellitus (E10-E14)	22	2.3	1.5
9	Septicemia (A40-A41)	19	2.0	*
10	Anemias (D50-D64)	16	1.7	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	1.2	*
12	Cerebrovascular diseases (I60-I69)	10	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	10	1.0	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.9	*
15	Influenza and pneumonia (J09-J18)	6	0.6	*
...	All other causes (Residual)	177	18.5	11.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-29 years				
...	All causes	1,331	100.0	90.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	15.3	13.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	12.3	11.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	142	10.7	9.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	132	9.9	9.0
5	Malignant neoplasms (C00-C97)	123	9.2	8.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	56	4.2	3.8
7	Diabetes mellitus (E10-E14)	41	3.1	2.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	2.9	2.6
9	Septicemia (A40-A41)	24	1.8	1.6
9	Cerebrovascular diseases (I60-I69)	24	1.8	1.6
11	Anemias (D50-D64)	22	1.7	1.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	1.7	1.5
13	Influenza and pneumonia (J09-J18)	19	1.4	*
14	Chronic lower respiratory diseases (J40-J47)	17	1.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	1.1	*
...	All other causes (Residual)	287	21.6	19.5
Non-Hispanic, black, female, 30-34 years				
...	All causes	1,652	100.0	125.1
1	Malignant neoplasms (C00-C97)	229	13.9	17.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	220	13.3	16.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	12.2	15.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	197	11.9	14.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	111	6.7	8.4
6	Diabetes mellitus (E10-E14)	55	3.3	4.2
7	Cerebrovascular diseases (I60-I69)	53	3.2	4.0
8	Pregnancy, childbirth and the puerperium (O00-O99)	50	3.0	3.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	2.0	2.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.8	2.2
11	Septicemia (A40-A41)	25	1.5	1.9
12	Anemias (D50-D64)	24	1.5	1.8
13	Influenza and pneumonia (J09-J18)	22	1.3	1.7
14	Chronic lower respiratory diseases (J40-J47)	20	1.2	1.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.0	*
...	All other causes (Residual)	366	22.2	27.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 35-39 years				
...	All causes	2,659	100.0	189.3
1	Malignant neoplasms (C00-C97)	458	17.2	32.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	380	14.3	27.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	345	13.0	24.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	273	10.3	19.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	5.1	9.7
6	Cerebrovascular diseases (I60-I69)	90	3.4	6.4
7	Diabetes mellitus (E10-E14)	87	3.3	6.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	74	2.8	5.3
9	Septicemia (A40-A41)	58	2.2	4.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	1.7	3.2
11	Pregnancy, childbirth and the puerperium (O00-O99)	43	1.6	3.1
12	Influenza and pneumonia (J09-J18)	35	1.3	2.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	1.3	2.4
14	Chronic lower respiratory diseases (J40-J47)	32	1.2	2.3
15	Anemias (D50-D64)	28	1.1	2.0
...	All other causes (Residual)	541	20.3	38.5
Non-Hispanic, black, female, 40-44 years				
...	All causes	4,157	100.0	284.7
1	Malignant neoplasms (C00-C97)	979	23.6	67.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	653	15.7	44.7
3	Human immunodeficiency virus (HIV) disease (B20-B24)	428	10.3	29.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	7.6	21.5
5	Cerebrovascular diseases (I60-I69)	228	5.5	15.6
6	Diabetes mellitus (E10-E14)	158	3.8	10.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	3.1	8.8
8	Septicemia (A40-A41)	89	2.1	6.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	1.9	5.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	63	1.5	4.3
11	Chronic lower respiratory diseases (J40-J47)	62	1.5	4.2
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	1.0	2.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	40	1.0	2.7
13	Influenza and pneumonia (J09-J18)	40	1.0	2.7
15	Anemias (D50-D64)	36	0.9	2.5
...	All other causes (Residual)	818	19.7	56.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-49 years				
...	All causes	6,595	100.0	450.5
1	Malignant neoplasms (C00-C97)	1,809	27.4	123.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,236	18.7	84.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	442	6.7	30.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	404	6.1	27.6
5	Cerebrovascular diseases (I60-I69)	390	5.9	26.6
6	Diabetes mellitus (E10-E14)	218	3.3	14.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	167	2.5	11.4
8	Septicemia (A40-A41)	150	2.3	10.2
9	Chronic lower respiratory diseases (J40-J47)	146	2.2	10.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	142	2.2	9.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	1.8	8.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	86	1.3	5.9
13	Influenza and pneumonia (J09-J18)	76	1.2	5.2
14	Viral hepatitis (B15-B19)	45	0.7	3.1
15	Anemias (D50-D64)	44	0.7	3.0
...	All other causes (Residual)	1,121	17.0	76.6
Non-Hispanic, black, female, 50-54 years				
...	All causes	8,610	100.0	668.4
1	Malignant neoplasms (C00-C97)	2,611	30.3	202.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,798	20.9	139.6
3	Cerebrovascular diseases (I60-I69)	542	6.3	42.1
4	Diabetes mellitus (E10-E14)	410	4.8	31.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	401	4.7	31.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	343	4.0	26.6
7	Chronic lower respiratory diseases (J40-J47)	220	2.6	17.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	218	2.5	16.9
9	Septicemia (A40-A41)	189	2.2	14.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	149	1.7	11.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	140	1.6	10.9
12	Influenza and pneumonia (J09-J18)	91	1.1	7.1
13	Viral hepatitis (B15-B19)	81	0.9	6.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	0.7	4.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	0.4	2.7
...	All other causes (Residual)	1,322	15.4	102.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 55-59 years				
...	All causes	9,631	100.0	909.1
1	Malignant neoplasms (C00-C97)	3,193	33.2	301.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,162	22.4	204.1
3	Diabetes mellitus (E10-E14)	547	5.7	51.6
4	Cerebrovascular diseases (I60-I69)	523	5.4	49.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	305	3.2	28.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	275	2.9	26.0
7	Chronic lower respiratory diseases (J40-J47)	248	2.6	23.4
8	Septicemia (A40-A41)	233	2.4	22.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	178	1.8	16.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	174	1.8	16.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	158	1.6	14.9
12	Influenza and pneumonia (J09-J18)	130	1.3	12.3
13	Viral hepatitis (B15-B19)	78	0.8	7.4
14	Aortic aneurysm and dissection (I71)	38	0.4	3.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	0.4	3.3
...	All other causes (Residual)	1,354	14.1	127.8
Non-Hispanic, black, female, 60-64 years				
...	All causes	9,745	100.0	1,298.9
1	Malignant neoplasms (C00-C97)	3,316	34.0	442.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,321	23.8	309.4
3	Diabetes mellitus (E10-E14)	637	6.5	84.9
4	Cerebrovascular diseases (I60-I69)	563	5.8	75.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	349	3.6	46.5
6	Chronic lower respiratory diseases (J40-J47)	255	2.6	34.0
7	Septicemia (A40-A41)	239	2.5	31.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	1.8	22.8
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	169	1.7	22.5
10	Influenza and pneumonia (J09-J18)	109	1.1	14.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	101	1.0	13.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	71	0.7	9.5
13	Aortic aneurysm and dissection (I71)	45	0.5	6.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.4	5.1
15	Viral hepatitis (B15-B19)	34	0.3	4.5
...	All other causes (Residual)	1,327	13.6	176.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-69 years				
...	All causes	10,872	100.0	1,886.2
1	Malignant neoplasms (C00-C97)	3,592	33.0	623.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,701	24.8	468.6
3	Diabetes mellitus (E10-E14)	692	6.4	120.1
4	Cerebrovascular diseases (I60-I69)	683	6.3	118.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	3.8	71.5
6	Chronic lower respiratory diseases (J40-J47)	379	3.5	65.8
7	Septicemia (A40-A41)	304	2.8	52.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	192	1.8	33.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	1.3	23.8
10	Influenza and pneumonia (J09-J18)	136	1.3	23.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	61	0.6	10.6
12	Aortic aneurysm and dissection (I71)	57	0.5	9.9
13	Alzheimer's disease (G30)	49	0.5	8.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.4	8.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	35	0.3	6.1
...	All other causes (Residual)	1,396	12.8	242.2
Non-Hispanic, black, female, 70-74 years				
...	All causes	12,641	100.0	2,710.8
1	Malignant neoplasms (C00-C97)	3,645	28.8	781.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,301	26.1	707.9
3	Cerebrovascular diseases (I60-I69)	849	6.7	182.1
4	Diabetes mellitus (E10-E14)	843	6.7	180.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	547	4.3	117.3
6	Chronic lower respiratory diseases (J40-J47)	452	3.6	96.9
7	Septicemia (A40-A41)	352	2.8	75.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	222	1.8	47.6
9	Influenza and pneumonia (J09-J18)	204	1.6	43.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	1.3	36.2
11	Alzheimer's disease (G30)	158	1.2	33.9
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.5	14.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	67	0.5	14.4
14	Aortic aneurysm and dissection (I71)	52	0.4	11.2
15	Pneumonitis due to solids and liquids (J69)	44	0.3	9.4
...	All other causes (Residual)	1,668	13.2	357.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 75-79 years				
...	All causes	14,924	100.0	4,091.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,037	27.1	1,106.8
2	Malignant neoplasms (C00-C97)	3,693	24.7	1,012.5
3	Cerebrovascular diseases (I60-I69)	1,139	7.6	312.3
4	Diabetes mellitus (E10-E14)	920	6.2	252.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	593	4.0	162.6
6	Chronic lower respiratory diseases (J40-J47)	523	3.5	143.4
7	Septicemia (A40-A41)	417	2.8	114.3
8	Alzheimer's disease (G30)	388	2.6	106.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	291	1.9	79.8
10	Influenza and pneumonia (J09-J18)	274	1.8	75.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	1.2	47.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.5	21.9
13	Pneumonitis due to solids and liquids (J69)	72	0.5	19.7
14	Parkinson's disease (G20-G21)	64	0.4	17.5
14	Aortic aneurysm and dissection (I71)	64	0.4	17.5
...	All other causes (Residual)	2,196	14.7	602.1
Non-Hispanic, black, female, 80-84 years				
...	All causes	16,951	100.0	6,173.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,943	29.2	1,800.2
2	Malignant neoplasms (C00-C97)	3,271	19.3	1,191.3
3	Cerebrovascular diseases (I60-I69)	1,384	8.2	504.0
4	Diabetes mellitus (E10-E14)	958	5.7	348.9
5	Alzheimer's disease (G30)	695	4.1	253.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	596	3.5	217.1
7	Chronic lower respiratory diseases (J40-J47)	509	3.0	185.4
8	Septicemia (A40-A41)	483	2.8	175.9
9	Influenza and pneumonia (J09-J18)	390	2.3	142.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	363	2.1	132.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	1.1	69.6
12	Pneumonitis due to solids and liquids (J69)	122	0.7	44.4
13	Parkinson's disease (G20-G21)	95	0.6	34.6
14	Aortic aneurysm and dissection (I71)	87	0.5	31.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	86	0.5	31.3
...	All other causes (Residual)	2,778	16.4	1,011.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85-89 years				
...	All causes	15,676	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,923	31.4	---
2	Malignant neoplasms (C00-C97)	2,088	13.3	---
3	Cerebrovascular diseases (I60-I69)	1,447	9.2	---
4	Alzheimer's disease (G30)	820	5.2	---
5	Diabetes mellitus (E10-E14)	729	4.7	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	624	4.0	---
7	Influenza and pneumonia (J09-J18)	433	2.8	---
8	Chronic lower respiratory diseases (J40-J47)	414	2.6	---
9	Septicemia (A40-A41)	402	2.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	396	2.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	1.1	---
12	Pneumonitis due to solids and liquids (J69)	143	0.9	---
13	Parkinson's disease (G20-G21)	78	0.5	---
14	Atherosclerosis (I70)	71	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	65	0.4	---
...	All other causes (Residual)	2,878	18.4	---
Non-Hispanic, black, female, 90-94 years				
...	All causes	10,900	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,601	33.0	---
2	Malignant neoplasms (C00-C97)	1,027	9.4	---
3	Cerebrovascular diseases (I60-I69)	993	9.1	---
4	Alzheimer's disease (G30)	777	7.1	---
5	Diabetes mellitus (E10-E14)	378	3.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	344	3.2	---
7	Influenza and pneumonia (J09-J18)	331	3.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	285	2.6	---
9	Septicemia (A40-A41)	256	2.3	---
10	Chronic lower respiratory diseases (J40-J47)	216	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	1.3	---
12	Pneumonitis due to solids and liquids (J69)	100	0.9	---
13	Atherosclerosis (I70)	68	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.4	---
15	Parkinson's disease (G20-G21)	41	0.4	---
...	All other causes (Residual)	2,298	21.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2007

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007; there were no deaths due to J09 in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 95-99 years				
...	All causes	5,320	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,944	36.5	---
2	Cerebrovascular diseases (I60-I69)	398	7.5	---
3	Alzheimer's disease (G30)	394	7.4	---
4	Malignant neoplasms (C00-C97)	362	6.8	---
5	Influenza and pneumonia (J09-J18)	189	3.6	---
6	Diabetes mellitus (E10-E14)	156	2.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	155	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	146	2.7	---
9	Septicemia (A40-A41)	99	1.9	---
10	Chronic lower respiratory diseases (J40-J47)	88	1.7	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	1.1	---
12	Pneumonitis due to solids and liquids (J69)	48	0.9	---
13	Atherosclerosis (I70)	45	0.8	---
14	Parkinson's disease (G20-G21)	19	0.4	---
15	Nutritional deficiencies (E40-E64)	17	0.3	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	17	0.3	---
...	All other causes (Residual)	1,186	22.3	---
Non-Hispanic, black, female, 100 years and over				
...	All causes	1,751	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	627	35.8	---
2	Alzheimer's disease (G30)	127	7.3	---
3	Cerebrovascular diseases (I60-I69)	111	6.3	---
4	Influenza and pneumonia (J09-J18)	84	4.8	---
5	Malignant neoplasms (C00-C97)	74	4.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	3.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	2.7	---
8	Diabetes mellitus (E10-E14)	41	2.3	---
9	Septicemia (A40-A41)	28	1.6	---
10	Chronic lower respiratory diseases (J40-J47)	24	1.4	---
11	Pneumonitis due to solids and liquids (J69)	21	1.2	---
12	Atherosclerosis (I70)	19	1.1	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	0.9	---
14	Nutritional deficiencies (E40-E64)	9	0.5	---
15	Parkinson's disease (G20-G21)	6	0.3	---
...	All other causes (Residual)	460	26.3	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available