

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	131,161	100.0	307.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,555	22.5	69.2
2	Malignant neoplasms (C00-C97)	26,156	19.9	61.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,464	8.7	26.9
4	Cerebrovascular diseases (I60-I69)	6,830	5.2	16.0
5	Diabetes mellitus (E10-E14)	6,665	5.1	15.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,555	2.7	8.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,520	2.7	8.2
8	Chronic lower respiratory diseases (J40-J47)	3,457	2.6	8.1
9	Influenza and pneumonia (J10-J18)	3,085	2.4	7.2
10	Certain conditions originating in the perinatal period (P00-P96)	2,816	2.1	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	2,329	1.8	5.5
12	Alzheimer's disease (G30)	2,201	1.7	5.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,188	1.7	5.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,896	1.4	4.4
15	Septicemia (A40-A41)	1,704	1.3	4.0
...	All other causes (Residual)	23,740	18.1	55.6
Hispanic, both sexes, 1-4 years				
...	All causes	1,041	100.0	28.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	380	36.5	10.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	11.4	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	7.9	2.3
4	Malignant neoplasms (C00-C97)	80	7.7	2.2
5	Influenza and pneumonia (J10-J18)	32	3.1	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.2	0.6
7	Cerebrovascular diseases (I60-I69)	15	1.4	*
8	Septicemia (A40-A41)	14	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.2	*
10	Meningitis (G00,G03)	8	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Anemias (D50-D64)	4	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	4	0.4	*
...	All other causes (Residual)	252	24.2	7.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,068	100.0	13.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	370	34.6	4.7
2	Malignant neoplasms (C00-C97)	194	18.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	8.0	1.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	5.8	0.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	3.7	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.2	0.4
7	Cerebrovascular diseases (I60-I69)	19	1.8	*
8	Chronic lower respiratory diseases (J40-J47)	15	1.4	*
9	Septicemia (A40-A41)	12	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.9	*
11	Influenza and pneumonia (J10-J18)	9	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	5	0.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
13	Anemias (D50-D64)	4	0.4	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	194	18.2	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-24 years				
...	All causes	5,837	100.0	81.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,670	45.7	37.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,307	22.4	18.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	553	9.5	7.7
4	Malignant neoplasms (C00-C97)	323	5.5	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	137	2.3	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	1.3	1.0
7	Cerebrovascular diseases (I60-I69)	44	0.8	0.6
8	Pregnancy, childbirth and the puerperium (O00-099)	35	0.6	0.5
9	Legal intervention (Y35,Y89.0)	32	0.5	0.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	31	0.5	0.4
11	Septicemia (A40-A41)	18	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	15	0.3	*
13	Diabetes mellitus (E10-E14)	14	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.2	*
15	Influenza and pneumonia (J10-J18)	11	0.2	*
...	All other causes (Residual)	560	9.6	7.8
Hispanic, both sexes, 25-34 years				
...	All causes	6,411	100.0	81.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,302	35.9	29.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,085	16.9	13.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	534	8.3	6.8
4	Malignant neoplasms (C00-C97)	530	8.3	6.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	354	5.5	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	192	3.0	2.5
7	Cerebrovascular diseases (I60-I69)	82	1.3	1.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	78	1.2	1.0
9	Diabetes mellitus (E10-E14)	69	1.1	0.9
10	Pregnancy, childbirth and the puerperium (O00-099)	60	0.9	0.8
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	0.8	0.7
12	Influenza and pneumonia (J10-J18)	40	0.6	0.5
13	Septicemia (A40-A41)	36	0.6	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	30	0.5	0.4
15	Chronic lower respiratory diseases (J40-J47)	25	0.4	0.3
...	All other causes (Residual)	942	14.7	12.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-44 years				
...	All causes	8,876	100.0	138.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,920	21.6	30.0
2	Malignant neoplasms (C00-C97)	1,415	15.9	22.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	957	10.8	15.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	616	6.9	9.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	520	5.9	8.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	472	5.3	7.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	467	5.3	7.3
8	Cerebrovascular diseases (I60-I69)	296	3.3	4.6
9	Diabetes mellitus (E10-E14)	216	2.4	3.4
10	Viral hepatitis (B15-B19)	114	1.3	1.8
11	Influenza and pneumonia (J10-J18)	93	1.0	1.5
12	Septicemia (A40-A41)	74	0.8	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	68	0.8	1.1
14	Chronic lower respiratory diseases (J40-J47)	53	0.6	0.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	0.5	0.7
...	All other causes (Residual)	1,548	17.4	24.2
Hispanic, both sexes, 45-54 years				
...	All causes	13,569	100.0	317.8
1	Malignant neoplasms (C00-C97)	3,324	24.5	77.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,233	16.5	52.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,455	10.7	34.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,003	7.4	23.5
5	Cerebrovascular diseases (I60-I69)	639	4.7	15.0
6	Diabetes mellitus (E10-E14)	632	4.7	14.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	557	4.1	13.0
8	Viral hepatitis (B15-B19)	293	2.2	6.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	283	2.1	6.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	264	1.9	6.2
11	Influenza and pneumonia (J10-J18)	172	1.3	4.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	165	1.2	3.9
13	Septicemia (A40-A41)	157	1.2	3.7
14	Chronic lower respiratory diseases (J40-J47)	151	1.1	3.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	82	0.6	1.9
...	All other causes (Residual)	2,159	15.9	50.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-64 years				
...	All causes	16,089	100.0	676.4
1	Malignant neoplasms (C00-C97)	4,836	30.1	203.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,642	22.6	153.1
3	Diabetes mellitus (E10-E14)	1,168	7.3	49.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	877	5.5	36.9
5	Cerebrovascular diseases (I60-I69)	795	4.9	33.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	720	4.5	30.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	351	2.2	14.8
8	Chronic lower respiratory diseases (J40-J47)	298	1.9	12.5
9	Influenza and pneumonia (J10-J18)	247	1.5	10.4
10	Septicemia (A40-A41)	230	1.4	9.7
11	Viral hepatitis (B15-B19)	202	1.3	8.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	171	1.1	7.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	150	0.9	6.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	142	0.9	6.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	75	0.5	3.2
...	All other causes (Residual)	2,185	13.6	91.9
Hispanic, both sexes, 65-74 years				
...	All causes	21,422	100.0	1,621.4
1	Malignant neoplasms (C00-C97)	6,359	29.7	481.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,551	25.9	420.1
3	Diabetes mellitus (E10-E14)	1,670	7.8	126.4
4	Cerebrovascular diseases (I60-I69)	1,181	5.5	89.4
5	Chronic lower respiratory diseases (J40-J47)	668	3.1	50.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	645	3.0	48.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	520	2.4	39.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	506	2.4	38.3
9	Influenza and pneumonia (J10-J18)	455	2.1	34.4
10	Septicemia (A40-A41)	337	1.6	25.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	246	1.1	18.6
12	Alzheimer's disease (G30)	172	0.8	13.0
13	Parkinson's disease (G20-G21)	119	0.6	9.0
14	Viral hepatitis (B15-B19)	113	0.5	8.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	107	0.5	8.1
...	All other causes (Residual)	2,773	12.9	209.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-84 years				
...	All causes	29,010	100.0	3,977.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,583	29.6	1,176.9
2	Malignant neoplasms (C00-C97)	6,438	22.2	882.8
3	Cerebrovascular diseases (I60-I69)	2,009	6.9	275.5
4	Diabetes mellitus (E10-E14)	1,947	6.7	267.0
5	Chronic lower respiratory diseases (J40-J47)	1,254	4.3	171.9
6	Influenza and pneumonia (J10-J18)	892	3.1	122.3
7	Alzheimer's disease (G30)	801	2.8	109.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	699	2.4	95.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	581	2.0	79.7
10	Septicemia (A40-A41)	445	1.5	61.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	394	1.4	54.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	386	1.3	52.9
13	Parkinson's disease (G20-G21)	354	1.2	48.5
14	Pneumonitis due to solids and liquids (J69)	177	0.6	24.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	152	0.5	20.8
...	All other causes (Residual)	3,898	13.4	534.5
Hispanic, both sexes, 85 years and over				
...	All causes	22,058	100.0	9,436.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,965	36.1	3,407.5
2	Malignant neoplasms (C00-C97)	2,637	12.0	1,128.1
3	Cerebrovascular diseases (I60-I69)	1,726	7.8	738.4
4	Alzheimer's disease (G30)	1,187	5.4	507.8
5	Influenza and pneumonia (J10-J18)	1,068	4.8	456.9
6	Chronic lower respiratory diseases (J40-J47)	969	4.4	414.6
7	Diabetes mellitus (E10-E14)	946	4.3	404.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	452	2.0	193.4
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	396	1.8	169.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	359	1.6	153.6
11	Septicemia (A40-A41)	315	1.4	134.8
12	Pneumonitis due to solids and liquids (J69)	257	1.2	109.9
13	Parkinson's disease (G20-G21)	222	1.0	95.0
14	Atherosclerosis (I70)	186	0.8	79.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	81	0.4	34.7
...	All other causes (Residual)	3,292	14.9	1,408.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	73,788	100.0	334.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,900	21.5	72.1
2	Malignant neoplasms (C00-C97)	13,896	18.8	63.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,612	11.7	39.0
4	Diabetes mellitus (E10-E14)	3,296	4.5	14.9
5	Cerebrovascular diseases (I60-I69)	3,188	4.3	14.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,008	4.1	13.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,561	3.5	11.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,841	2.5	8.3
9	Chronic lower respiratory diseases (J40-J47)	1,839	2.5	8.3
10	Certain conditions originating in the perinatal period (P00-P96)	1,612	2.2	7.3
11	Influenza and pneumonia (J10-J18)	1,506	2.0	6.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,309	1.8	5.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	1,186	1.6	5.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	996	1.3	4.5
15	Septicemia (A40-A41)	838	1.1	3.8
...	All other causes (Residual)	12,200	16.5	55.3
Hispanic, male, 1-4 years				
...	All causes	610	100.0	33.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	244	40.0	13.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	10.7	3.5
3	Malignant neoplasms (C00-C97)	52	8.5	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	7.5	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.8	*
6	Influenza and pneumonia (J10-J18)	15	2.5	*
7	Cerebrovascular diseases (I60-I69)	8	1.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
9	Septicemia (A40-A41)	6	1.0	*
10	Meningococcal infection (A39)	3	0.5	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	3	0.5	*
13	Tuberculosis (A16-A19)	2	0.3	*
13	Meningitis (G00,G03)	2	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	2	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	131	21.5	7.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-14 years				
...	All causes	612	100.0	15.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	222	36.3	5.6
2	Malignant neoplasms (C00-C97)	99	16.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	7.7	1.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	6.5	1.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	4.4	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.4	0.5
7	Cerebrovascular diseases (I60-I69)	9	1.5	*
7	Chronic lower respiratory diseases (J40-J47)	9	1.5	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
9	Influenza and pneumonia (J10-J18)	7	1.1	*
11	Septicemia (A40-A41)	6	1.0	*
12	Anemias (D50-D64)	4	0.7	*
13	Diabetes mellitus (E10-E14)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
13	Diseases of appendix (K35-K38)	2	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	106	17.3	2.7
Hispanic, male, 15-24 years				
...	All causes	4,604	100.0	120.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,133	46.3	55.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,187	25.8	31.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	462	10.0	12.1
4	Malignant neoplasms (C00-C97)	202	4.4	5.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	2.0	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	0.9	1.1
7	Legal intervention (Y35,Y89.0)	32	0.7	0.8
8	Cerebrovascular diseases (I60-I69)	25	0.5	0.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.5	0.6
10	Septicemia (A40-A41)	10	0.2	*
11	Diabetes mellitus (E10-E14)	9	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.2	*
12	Influenza and pneumonia (J10-J18)	8	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	8	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	7	0.2	*
...	All other causes (Residual)	356	7.7	9.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-34 years				
...	All causes	4,960	100.0	115.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,930	38.9	44.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	956	19.3	22.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	461	9.3	10.7
4	Malignant neoplasms (C00-C97)	280	5.6	6.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	272	5.5	6.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	142	2.9	3.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	62	1.3	1.4
8	Cerebrovascular diseases (I60-I69)	50	1.0	1.2
9	Diabetes mellitus (E10-E14)	40	0.8	0.9
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.6	0.7
11	Influenza and pneumonia (J10-J18)	26	0.5	0.6
12	Legal intervention (Y35,Y89.0)	24	0.5	0.6
13	Septicemia (A40-A41)	23	0.5	0.5
14	Chronic lower respiratory diseases (J40-J47)	16	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	16	0.3	*
...	All other causes (Residual)	632	12.7	14.7
Hispanic, male, 35-44 years				
...	All causes	6,145	100.0	182.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,513	24.6	44.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	733	11.9	21.7
3	Malignant neoplasms (C00-C97)	601	9.8	17.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	496	8.1	14.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	428	7.0	12.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	399	6.5	11.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	369	6.0	10.9
8	Cerebrovascular diseases (I60-I69)	177	2.9	5.2
9	Diabetes mellitus (E10-E14)	153	2.5	4.5
10	Viral hepatitis (B15-B19)	85	1.4	2.5
11	Influenza and pneumonia (J10-J18)	59	1.0	1.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	37	0.6	1.1
13	Septicemia (A40-A41)	35	0.6	1.0
14	Chronic lower respiratory diseases (J40-J47)	34	0.6	1.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	27	0.4	0.8
...	All other causes (Residual)	999	16.3	29.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-54 years				
...	All causes	8,994	100.0	417.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,679	18.7	77.9
2	Malignant neoplasms (C00-C97)	1,635	18.2	75.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,141	12.7	52.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	814	9.1	37.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	448	5.0	20.8
6	Cerebrovascular diseases (I60-I69)	384	4.3	17.8
7	Diabetes mellitus (E10-E14)	378	4.2	17.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	237	2.6	11.0
9	Viral hepatitis (B15-B19)	222	2.5	10.3
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	218	2.4	10.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	112	1.2	5.2
12	Influenza and pneumonia (J10-J18)	110	1.2	5.1
13	Septicemia (A40-A41)	93	1.0	4.3
14	Chronic lower respiratory diseases (J40-J47)	77	0.9	3.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	46	0.5	2.1
...	All other causes (Residual)	1,400	15.6	65.0
Hispanic, male, 55-64 years				
...	All causes	9,954	100.0	875.8
1	Malignant neoplasms (C00-C97)	2,693	27.1	236.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,492	25.0	219.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	664	6.7	58.4
4	Diabetes mellitus (E10-E14)	662	6.7	58.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	530	5.3	46.6
6	Cerebrovascular diseases (I60-I69)	458	4.6	40.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	201	2.0	17.7
8	Chronic lower respiratory diseases (J40-J47)	180	1.8	15.8
9	Influenza and pneumonia (J10-J18)	147	1.5	12.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	143	1.4	12.6
11	Viral hepatitis (B15-B19)	130	1.3	11.4
12	Septicemia (A40-A41)	121	1.2	10.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	1.2	10.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	69	0.7	6.1
15	Aortic aneurysm and dissection (I71)	54	0.5	4.8
...	All other causes (Residual)	1,294	13.0	113.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-74 years				
...	All causes	11,984	100.0	2,029.4
1	Malignant neoplasms (C00-C97)	3,564	29.7	603.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,316	27.7	561.5
3	Diabetes mellitus (E10-E14)	850	7.1	143.9
4	Cerebrovascular diseases (I60-I69)	627	5.2	106.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	408	3.4	69.1
6	Chronic lower respiratory diseases (J40-J47)	381	3.2	64.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	345	2.9	58.4
8	Influenza and pneumonia (J10-J18)	250	2.1	42.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	244	2.0	41.3
10	Septicemia (A40-A41)	175	1.5	29.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	121	1.0	20.5
12	Alzheimer's disease (G30)	82	0.7	13.9
13	Parkinson's disease (G20-G21)	80	0.7	13.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.5	10.7
14	Aortic aneurysm and dissection (I71)	63	0.5	10.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	0.5	10.7
...	All other causes (Residual)	1,352	11.3	228.9
Hispanic, male, 75-84 years				
...	All causes	14,539	100.0	4,856.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,398	30.2	1,469.2
2	Malignant neoplasms (C00-C97)	3,478	23.9	1,161.8
3	Cerebrovascular diseases (I60-I69)	880	6.1	294.0
4	Diabetes mellitus (E10-E14)	877	6.0	293.0
5	Chronic lower respiratory diseases (J40-J47)	701	4.8	234.2
6	Influenza and pneumonia (J10-J18)	448	3.1	149.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	338	2.3	112.9
8	Alzheimer's disease (G30)	337	2.3	112.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	318	2.2	106.2
10	Septicemia (A40-A41)	222	1.5	74.2
11	Parkinson's disease (G20-G21)	213	1.5	71.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	206	1.4	68.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	162	1.1	54.1
14	Pneumonitis due to solids and liquids (J69)	96	0.7	32.1
15	Aortic aneurysm and dissection (I71)	92	0.6	30.7
...	All other causes (Residual)	1,773	12.2	592.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85 years and over				
...	All causes	8,143	100.0	10,140.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,838	34.9	3,534.2
2	Malignant neoplasms (C00-C97)	1,286	15.8	1,601.5
3	Cerebrovascular diseases (I60-I69)	556	6.8	692.4
4	Chronic lower respiratory diseases (J40-J47)	423	5.2	526.8
5	Influenza and pneumonia (J10-J18)	399	4.9	496.9
6	Alzheimer's disease (G30)	331	4.1	412.2
7	Diabetes mellitus (E10-E14)	325	4.0	404.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	208	2.6	259.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	135	1.7	168.1
10	Pneumonitis due to solids and liquids (J69)	124	1.5	154.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	113	1.4	140.7
12	Parkinson's disease (G20-G21)	111	1.4	138.2
13	Septicemia (A40-A41)	110	1.4	137.0
14	Atherosclerosis (I70)	67	0.8	83.4
15	Aortic aneurysm and dissection (I71)	35	0.4	43.6
...	All other causes (Residual)	1,082	13.3	1,347.4
Hispanic, female, all ages				
...	All causes	57,373	100.0	278.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,655	23.8	66.2
2	Malignant neoplasms (C00-C97)	12,260	21.4	59.5
3	Cerebrovascular diseases (I60-I69)	3,642	6.3	17.7
4	Diabetes mellitus (E10-E14)	3,369	5.9	16.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,852	5.0	13.8
6	Chronic lower respiratory diseases (J40-J47)	1,618	2.8	7.8
7	Influenza and pneumonia (J10-J18)	1,579	2.8	7.7
8	Alzheimer's disease (G30)	1,437	2.5	7.0
9	Certain conditions originating in the perinatal period (P00-P96)	1,204	2.1	5.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	1,143	2.0	5.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	994	1.7	4.8
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	900	1.6	4.4
13	Septicemia (A40-A41)	866	1.5	4.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	767	1.3	3.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	512	0.9	2.5
...	All other causes (Residual)	10,575	18.4	51.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Hispanic, female, 1-4 years			
...	All causes	431	100.0	24.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	31.6	7.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	12.5	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	8.4	2.0
4	Malignant neoplasms (C00-C97)	28	6.5	1.6
5	Influenza and pneumonia (J10-J18)	17	3.9	*
6	Septicemia (A40-A41)	8	1.9	*
7	Cerebrovascular diseases (I60-I69)	7	1.6	*
8	Meningitis (G00,G03)	6	1.4	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
12	Anemias (D50-D64)	3	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
15	Shigellosis and amebiasis (A03,A06)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Hernia (K40-K46)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
...	All other causes (Residual)	108	25.1	6.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	456	100.0	12.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	32.5	3.9
2	Malignant neoplasms (C00-C97)	95	20.8	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	8.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.8	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.9	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.6	*
7	Cerebrovascular diseases (I60-I69)	10	2.2	*
8	Septicemia (A40-A41)	6	1.3	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	4	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
13	Meningitis (G00,G03)	2	0.4	*
13	Influenza and pneumonia (J10-J18)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	90	19.7	2.4
Hispanic, female, 15-24 years				
...	All causes	1,233	100.0	36.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	537	43.6	15.9
2	Malignant neoplasms (C00-C97)	121	9.8	3.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	120	9.7	3.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	7.4	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	3.7	1.4
6	Pregnancy, childbirth and the puerperium (O00-099)	35	2.8	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	2.5	0.9
8	Cerebrovascular diseases (I60-I69)	19	1.5	*
9	Septicemia (A40-A41)	8	0.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	7	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.4	*
12	Diabetes mellitus (E10-E14)	5	0.4	*
14	Tuberculosis (A16-A19)	3	0.2	*
14	Aortic aneurysm and dissection (I71)	3	0.2	*
14	Influenza and pneumonia (J10-J18)	3	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.2	*
...	All other causes (Residual)	188	15.2	5.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,451	100.0	41.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	372	25.6	10.5
2	Malignant neoplasms (C00-C97)	250	17.2	7.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	8.9	3.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	5.7	2.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	5.0	2.1
6	Pregnancy, childbirth and the puerperium (O00-099)	60	4.1	1.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	50	3.4	1.4
8	Cerebrovascular diseases (I60-I69)	32	2.2	0.9
9	Diabetes mellitus (E10-E14)	29	2.0	0.8
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	1.5	0.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	16	1.1	*
12	Influenza and pneumonia (J10-J18)	14	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	14	1.0	*
14	Septicemia (A40-A41)	13	0.9	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.7	*
...	All other causes (Residual)	285	19.6	8.1
Hispanic, female, 35-44 years				
...	All causes	2,731	100.0	90.6
1	Malignant neoplasms (C00-C97)	814	29.8	27.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	407	14.9	13.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	224	8.2	7.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	120	4.4	4.0
5	Cerebrovascular diseases (I60-I69)	119	4.4	3.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	98	3.6	3.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	92	3.4	3.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	2.7	2.4
9	Diabetes mellitus (E10-E14)	63	2.3	2.1
10	Septicemia (A40-A41)	39	1.4	1.3
11	Influenza and pneumonia (J10-J18)	34	1.2	1.1
12	Pregnancy, childbirth and the puerperium (O00-099)	32	1.2	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	31	1.1	1.0
14	Viral hepatitis (B15-B19)	29	1.1	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.8	0.7
...	All other causes (Residual)	535	19.6	17.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	4,575	100.0	216.4
1	Malignant neoplasms (C00-C97)	1,689	36.9	79.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	554	12.1	26.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	314	6.9	14.8
4	Cerebrovascular diseases (I60-I69)	255	5.6	12.1
5	Diabetes mellitus (E10-E14)	254	5.6	12.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	189	4.1	8.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	109	2.4	5.2
8	Chronic lower respiratory diseases (J40-J47)	74	1.6	3.5
9	Viral hepatitis (B15-B19)	71	1.6	3.4
10	Septicemia (A40-A41)	64	1.4	3.0
11	Influenza and pneumonia (J10-J18)	62	1.4	2.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	53	1.2	2.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	1.0	2.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	1.0	2.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	36	0.8	1.7
...	All other causes (Residual)	759	16.6	35.9
Hispanic, female, 55-64 years				
...	All causes	6,135	100.0	493.9
1	Malignant neoplasms (C00-C97)	2,143	34.9	172.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,150	18.7	92.6
3	Diabetes mellitus (E10-E14)	506	8.2	40.7
4	Cerebrovascular diseases (I60-I69)	337	5.5	27.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	213	3.5	17.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	3.1	15.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	150	2.4	12.1
8	Chronic lower respiratory diseases (J40-J47)	118	1.9	9.5
9	Septicemia (A40-A41)	109	1.8	8.8
10	Influenza and pneumonia (J10-J18)	100	1.6	8.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	81	1.3	6.5
12	Viral hepatitis (B15-B19)	72	1.2	5.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.5	2.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.5	2.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	0.4	2.1
...	All other causes (Residual)	881	14.4	70.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	9,438	100.0	1,291.6
1	Malignant neoplasms (C00-C97)	2,795	29.6	382.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,235	23.7	305.9
3	Diabetes mellitus (E10-E14)	820	8.7	112.2
4	Cerebrovascular diseases (I60-I69)	554	5.9	75.8
5	Chronic lower respiratory diseases (J40-J47)	287	3.0	39.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	262	2.8	35.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	237	2.5	32.4
8	Influenza and pneumonia (J10-J18)	205	2.2	28.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	175	1.9	23.9
10	Septicemia (A40-A41)	162	1.7	22.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	125	1.3	17.1
12	Alzheimer's disease (G30)	90	1.0	12.3
13	Viral hepatitis (B15-B19)	60	0.6	8.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.5	6.0
15	Parkinson's disease (G20-G21)	39	0.4	5.3
...	All other causes (Residual)	1,348	14.3	184.5
Hispanic, female, 75-84 years				
...	All causes	14,471	100.0	3,365.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,185	28.9	973.4
2	Malignant neoplasms (C00-C97)	2,960	20.5	688.5
3	Cerebrovascular diseases (I60-I69)	1,129	7.8	262.6
4	Diabetes mellitus (E10-E14)	1,070	7.4	248.9
5	Chronic lower respiratory diseases (J40-J47)	553	3.8	128.6
6	Alzheimer's disease (G30)	464	3.2	107.9
7	Influenza and pneumonia (J10-J18)	444	3.1	103.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	361	2.5	84.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	263	1.8	61.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	224	1.5	52.1
11	Septicemia (A40-A41)	223	1.5	51.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	188	1.3	43.7
13	Parkinson's disease (G20-G21)	141	1.0	32.8
14	Pneumonitis due to solids and liquids (J69)	81	0.6	18.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	70	0.5	16.3
...	All other causes (Residual)	2,115	14.6	491.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	13,915	100.0	9,068.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,127	36.8	3,341.3
2	Malignant neoplasms (C00-C97)	1,351	9.7	880.4
3	Cerebrovascular diseases (I60-I69)	1,170	8.4	762.5
4	Alzheimer's disease (G30)	856	6.2	557.9
5	Influenza and pneumonia (J10-J18)	669	4.8	436.0
6	Diabetes mellitus (E10-E14)	621	4.5	404.7
7	Chronic lower respiratory diseases (J40-J47)	546	3.9	355.8
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	283	2.0	184.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	244	1.8	159.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	224	1.6	146.0
11	Septicemia (A40-A41)	205	1.5	133.6
12	Pneumonitis due to solids and liquids (J69)	133	1.0	86.7
13	Atherosclerosis (I70)	119	0.9	77.6
14	Parkinson's disease (G20-G21)	111	0.8	72.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	52	0.4	33.9
...	All other causes (Residual)	2,204	15.8	1,436.3
White, non-Hispanic, both sexes, all ages				
...	All causes	1,967,142	100.0	981.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	535,101	27.2	267.1
2	Malignant neoplasms (C00-C97)	456,133	23.2	227.7
3	Chronic lower respiratory diseases (J40-J47)	117,337	6.0	58.6
4	Cerebrovascular diseases (I60-I69)	115,074	5.8	57.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88,970	4.5	44.4
6	Alzheimer's disease (G30)	63,958	3.3	31.9
7	Diabetes mellitus (E10-E14)	53,159	2.7	26.5
8	Influenza and pneumonia (J10-J18)	52,431	2.7	26.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	32,492	1.7	16.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,305	1.4	13.6
11	Septicemia (A40-A41)	25,496	1.3	12.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,428	1.0	10.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17,949	0.9	9.0
14	Parkinson's disease (G20-G21)	17,767	0.9	8.9
15	Pneumonitis due to solids and liquids (J69)	14,739	0.7	7.4
...	All other causes (Residual)	328,803	16.7	164.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,404	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	898	37.4	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	270	11.2	2.9
3	Malignant neoplasms (C00-C97)	215	8.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	5.7	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	2.9	0.8
6	Influenza and pneumonia (J10-J18)	46	1.9	0.5
7	Septicemia (A40-A41)	44	1.8	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.3	0.3
9	Cerebrovascular diseases (I60-I69)	30	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	29	1.2	0.3
11	Chronic lower respiratory diseases (J40-J47)	20	0.8	0.2
12	Meningitis (G00,G03)	12	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	10	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	9	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.4	*
...	All other causes (Residual)	573	23.8	6.2
White, non-Hispanic, both sexes, 5-14 years				
...	All causes	3,681	100.0	15.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,423	38.7	5.9
2	Malignant neoplasms (C00-C97)	597	16.2	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	207	5.6	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	170	4.6	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	3.5	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	111	3.0	0.5
7	Influenza and pneumonia (J10-J18)	67	1.8	0.3
8	Septicemia (A40-A41)	47	1.3	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	1.3	0.2
10	Cerebrovascular diseases (I60-I69)	45	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	31	0.8	0.1
12	Diabetes mellitus (E10-E14)	22	0.6	0.1
13	Certain conditions originating in the perinatal period (P00-P96)	19	0.5	*
14	Meningitis (G00,G03)	16	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	13	0.4	*
...	All other causes (Residual)	739	20.1	3.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-24 years				
...	All causes	19,683	100.0	74.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,711	54.4	40.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,992	15.2	11.4
3	Malignant neoplasms (C00-C97)	1,048	5.3	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	895	4.5	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	604	3.1	2.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	313	1.6	1.2
7	Influenza and pneumonia (J10-J18)	123	0.6	0.5
8	Diabetes mellitus (E10-E14)	110	0.6	0.4
9	Cerebrovascular diseases (I60-I69)	92	0.5	0.3
10	Septicemia (A40-A41)	77	0.4	0.3
11	Pregnancy, childbirth and the puerperium (O00-099)	76	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	64	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	45	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	31	0.2	0.1
...	All other causes (Residual)	2,439	12.4	9.3
White, non-Hispanic, both sexes, 25-34 years				
...	All causes	23,826	100.0	98.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,347	39.2	38.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,708	15.6	15.3
3	Malignant neoplasms (C00-C97)	2,175	9.1	9.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,742	7.3	7.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	973	4.1	4.0
6	Diabetes mellitus (E10-E14)	306	1.3	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	287	1.2	1.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	277	1.2	1.1
9	Cerebrovascular diseases (I60-I69)	274	1.2	1.1
10	Influenza and pneumonia (J10-J18)	224	0.9	0.9
11	Septicemia (A40-A41)	157	0.7	0.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	156	0.7	0.6
13	Chronic lower respiratory diseases (J40-J47)	129	0.5	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	123	0.5	0.5
15	Pregnancy, childbirth and the puerperium (O00-099)	121	0.5	0.5
...	All other causes (Residual)	3,827	16.1	15.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 35-44 years				
...	All causes	54,786	100.0	186.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,252	22.4	41.7
2	Malignant neoplasms (C00-C97)	9,976	18.2	34.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,153	14.9	27.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,434	9.9	18.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,814	3.3	6.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,352	2.5	4.6
7	Diabetes mellitus (E10-E14)	1,215	2.2	4.1
8	Cerebrovascular diseases (I60-I69)	1,158	2.1	3.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,099	2.0	3.7
10	Influenza and pneumonia (J10-J18)	593	1.1	2.0
10	Chronic lower respiratory diseases (J40-J47)	593	1.1	2.0
12	Septicemia (A40-A41)	480	0.9	1.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	381	0.7	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	333	0.6	1.1
15	Viral hepatitis (B15-B19)	304	0.6	1.0
...	All other causes (Residual)	9,649	17.6	32.9
White, non-Hispanic, both sexes, 45-54 years				
...	All causes	126,669	100.0	407.1
1	Malignant neoplasms (C00-C97)	36,433	28.8	117.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,135	20.6	84.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,657	10.8	43.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,208	4.9	20.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,323	4.2	17.1
6	Cerebrovascular diseases (I60-I69)	3,461	2.7	11.1
7	Diabetes mellitus (E10-E14)	3,454	2.7	11.1
8	Chronic lower respiratory diseases (J40-J47)	3,054	2.4	9.8
9	Viral hepatitis (B15-B19)	1,602	1.3	5.1
10	Influenza and pneumonia (J10-J18)	1,449	1.1	4.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,408	1.1	4.5
12	Septicemia (A40-A41)	1,342	1.1	4.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	1,059	0.8	3.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	920	0.7	3.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	553	0.4	1.8
...	All other causes (Residual)	20,611	16.3	66.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 55-64 years				
...	All causes	206,078	100.0	873.5
1	Malignant neoplasms (C00-C97)	77,617	37.7	329.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,744	23.2	202.4
3	Chronic lower respiratory diseases (J40-J47)	11,058	5.4	46.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,303	4.0	35.2
5	Diabetes mellitus (E10-E14)	7,293	3.5	30.9
6	Cerebrovascular diseases (I60-I69)	6,351	3.1	26.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,258	2.6	22.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,822	1.9	16.2
9	Septicemia (A40-A41)	2,641	1.3	11.2
10	Influenza and pneumonia (J10-J18)	2,457	1.2	10.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	2,317	1.1	9.8
12	Aortic aneurysm and dissection (I71)	1,148	0.6	4.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,059	0.5	4.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,034	0.5	4.4
15	Viral hepatitis (B15-B19)	858	0.4	3.6
...	All other causes (Residual)	27,118	13.2	114.9
White, non-Hispanic, both sexes, 65-74 years				
...	All causes	314,958	100.0	2,126.0
1	Malignant neoplasms (C00-C97)	112,940	35.9	762.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,716	23.7	504.3
3	Chronic lower respiratory diseases (J40-J47)	26,824	8.5	181.1
4	Cerebrovascular diseases (I60-I69)	13,674	4.3	92.3
5	Diabetes mellitus (E10-E14)	10,826	3.4	73.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,890	2.2	46.5
7	Influenza and pneumonia (J10-J18)	5,022	1.6	33.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	4,846	1.5	32.7
9	Septicemia (A40-A41)	4,413	1.4	29.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	3,929	1.2	26.5
11	Alzheimer's disease (G30)	3,208	1.0	21.7
12	Aortic aneurysm and dissection (I71)	2,443	0.8	16.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,137	0.7	14.4
14	Parkinson's disease (G20-G21)	2,105	0.7	14.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,978	0.6	13.4
...	All other causes (Residual)	39,007	12.4	263.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 75-84 years				
...	All causes	581,712	100.0	5,330.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	160,084	27.5	1,466.8
2	Malignant neoplasms (C00-C97)	142,314	24.5	1,304.0
3	Chronic lower respiratory diseases (J40-J47)	45,941	7.9	420.9
4	Cerebrovascular diseases (I60-I69)	38,830	6.7	355.8
5	Alzheimer's disease (G30)	20,469	3.5	187.5
6	Diabetes mellitus (E10-E14)	17,293	3.0	158.4
7	Influenza and pneumonia (J10-J18)	15,638	2.7	143.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,942	2.1	109.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	11,298	1.9	103.5
10	Parkinson's disease (G20-G21)	8,445	1.5	77.4
11	Septicemia (A40-A41)	8,381	1.4	76.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5,415	0.9	49.6
13	Pneumonitis due to solids and liquids (J69)	4,750	0.8	43.5
14	Aortic aneurysm and dissection (I71)	4,390	0.8	40.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,062	0.7	37.2
...	All other causes (Residual)	82,460	14.2	755.5
White, non-Hispanic, both sexes, 85 years and over				
...	All causes	620,246	100.0	14,222.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	215,545	34.8	4,942.5
2	Malignant neoplasms (C00-C97)	72,770	11.7	1,668.6
3	Cerebrovascular diseases (I60-I69)	51,103	8.2	1,171.8
4	Alzheimer's disease (G30)	39,646	6.4	909.1
5	Chronic lower respiratory diseases (J40-J47)	29,611	4.8	679.0
6	Influenza and pneumonia (J10-J18)	26,717	4.3	612.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,962	2.1	297.2
8	Diabetes mellitus (E10-E14)	12,638	2.0	289.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	12,387	2.0	284.0
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8,788	1.4	201.5
11	Septicemia (A40-A41)	7,814	1.3	179.2
12	Pneumonitis due to solids and liquids (J69)	7,370	1.2	169.0
13	Parkinson's disease (G20-G21)	6,781	1.1	155.5
14	Atherosclerosis (I70)	6,162	1.0	141.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,564	0.6	81.7
...	All other causes (Residual)	106,388	17.2	2,439.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	954,402	100.0	970.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	263,311	27.6	267.8
2	Malignant neoplasms (C00-C97)	236,659	24.8	240.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56,021	5.9	57.0
4	Chronic lower respiratory diseases (J40-J47)	55,015	5.8	56.0
5	Cerebrovascular diseases (I60-I69)	44,038	4.6	44.8
6	Diabetes mellitus (E10-E14)	26,365	2.8	26.8
7	Influenza and pneumonia (J10-J18)	22,903	2.4	23.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,605	2.3	22.0
9	Alzheimer's disease (G30)	18,208	1.9	18.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	15,953	1.7	16.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13,088	1.4	13.3
12	Septicemia (A40-A41)	11,337	1.2	11.5
13	Parkinson's disease (G20-G21)	10,232	1.1	10.4
14	Pneumonitis due to solids and liquids (J69)	7,430	0.8	7.6
15	Aortic aneurysm and dissection (I71)	6,877	0.7	7.0
...	All other causes (Residual)	145,360	15.2	147.8
White, non-Hispanic, male, 1-4 years				
...	All causes	1,408	100.0	29.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	557	39.6	11.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	139	9.9	3.0
3	Malignant neoplasms (C00-C97)	118	8.4	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	5.3	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	2.7	0.8
6	Septicemia (A40-A41)	26	1.8	0.6
7	Influenza and pneumonia (J10-J18)	23	1.6	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.4	0.4
8	Cerebrovascular diseases (I60-I69)	20	1.4	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	18	1.3	*
11	Meningitis (G00,G03)	11	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	6	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	6	0.4	*
...	All other causes (Residual)	335	23.8	7.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-14 years				
...	All causes	2,163	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	886	41.0	7.1
2	Malignant neoplasms (C00-C97)	333	15.4	2.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	133	6.1	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	4.7	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	3.2	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	2.8	0.5
7	Septicemia (A40-A41)	29	1.3	0.2
7	Influenza and pneumonia (J10-J18)	29	1.3	0.2
9	Cerebrovascular diseases (I60-I69)	26	1.2	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.0	0.2
11	Chronic lower respiratory diseases (J40-J47)	17	0.8	*
12	Diabetes mellitus (E10-E14)	9	0.4	*
12	Meningitis (G00,G03)	9	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	9	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (Residual)	422	19.5	3.4
White, non-Hispanic, male, 15-24 years				
...	All causes	14,267	100.0	105.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,942	55.7	58.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,487	17.4	18.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	639	4.5	4.7
4	Malignant neoplasms (C00-C97)	628	4.4	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	422	3.0	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	210	1.5	1.6
7	Influenza and pneumonia (J10-J18)	82	0.6	0.6
8	Diabetes mellitus (E10-E14)	64	0.4	0.5
9	Septicemia (A40-A41)	46	0.3	0.3
10	Cerebrovascular diseases (I60-I69)	40	0.3	0.3
10	Chronic lower respiratory diseases (J40-J47)	40	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.3	0.3
13	Legal intervention (Y35,Y89.0)	26	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	24	0.2	0.2
15	Aortic aneurysm and dissection (I71)	17	0.1	*
...	All other causes (Residual)	1,561	10.9	11.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-34 years				
...	All causes	16,366	100.0	134.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,966	42.6	57.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,995	18.3	24.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,206	7.4	9.9
4	Malignant neoplasms (C00-C97)	1,103	6.7	9.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	664	4.1	5.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	195	1.2	1.6
7	Diabetes mellitus (E10-E14)	189	1.2	1.5
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	1.0	1.3
9	Cerebrovascular diseases (I60-I69)	147	0.9	1.2
10	Influenza and pneumonia (J10-J18)	117	0.7	1.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	92	0.6	0.8
12	Septicemia (A40-A41)	74	0.5	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	67	0.4	0.5
14	Chronic lower respiratory diseases (J40-J47)	57	0.3	0.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.3	0.5
...	All other causes (Residual)	2,276	13.9	18.7
White, non-Hispanic, male, 35-44 years				
...	All causes	34,719	100.0	236.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,585	24.7	58.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,920	17.1	40.3
3	Malignant neoplasms (C00-C97)	4,480	12.9	30.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,164	12.0	28.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,167	3.4	7.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,122	3.2	7.6
7	Diabetes mellitus (E10-E14)	760	2.2	5.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	710	2.0	4.8
9	Cerebrovascular diseases (I60-I69)	581	1.7	4.0
10	Influenza and pneumonia (J10-J18)	334	1.0	2.3
11	Chronic lower respiratory diseases (J40-J47)	274	0.8	1.9
12	Septicemia (A40-A41)	241	0.7	1.6
13	Viral hepatitis (B15-B19)	208	0.6	1.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	201	0.6	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	195	0.6	1.3
...	All other causes (Residual)	5,777	16.6	39.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 45-54 years				
...	All causes	79,860	100.0	517.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,485	24.4	126.2
2	Malignant neoplasms (C00-C97)	18,574	23.3	120.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,562	12.0	61.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,656	5.8	30.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,799	4.8	24.6
6	Diabetes mellitus (E10-E14)	2,179	2.7	14.1
7	Cerebrovascular diseases (I60-I69)	1,868	2.3	12.1
8	Chronic lower respiratory diseases (J40-J47)	1,562	2.0	10.1
9	Viral hepatitis (B15-B19)	1,187	1.5	7.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,177	1.5	7.6
11	Influenza and pneumonia (J10-J18)	847	1.1	5.5
12	Septicemia (A40-A41)	708	0.9	4.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	664	0.8	4.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	656	0.8	4.2
15	Aortic aneurysm and dissection (I71)	370	0.5	2.4
...	All other causes (Residual)	12,566	15.7	81.4
White, non-Hispanic, male, 55-64 years				
...	All causes	124,229	100.0	1,079.6
1	Malignant neoplasms (C00-C97)	42,128	33.9	366.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,711	27.1	293.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,649	4.5	49.1
4	Chronic lower respiratory diseases (J40-J47)	5,629	4.5	48.9
5	Diabetes mellitus (E10-E14)	4,384	3.5	38.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,738	3.0	32.5
7	Cerebrovascular diseases (I60-I69)	3,534	2.8	30.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,953	2.4	25.7
9	Septicemia (A40-A41)	1,416	1.1	12.3
10	Influenza and pneumonia (J10-J18)	1,399	1.1	12.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	1,345	1.1	11.7
12	Aortic aneurysm and dissection (I71)	849	0.7	7.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	658	0.5	5.7
14	Viral hepatitis (B15-B19)	630	0.5	5.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	599	0.5	5.2
...	All other causes (Residual)	15,607	12.6	135.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65-74 years				
...	All causes	177,517	100.0	2,584.5
1	Malignant neoplasms (C00-C97)	62,528	35.2	910.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,538	26.2	677.6
3	Chronic lower respiratory diseases (J40-J47)	13,650	7.7	198.7
4	Cerebrovascular diseases (I60-I69)	7,032	4.0	102.4
5	Diabetes mellitus (E10-E14)	5,941	3.3	86.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,232	2.4	61.6
7	Influenza and pneumonia (J10-J18)	2,753	1.6	40.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	2,625	1.5	38.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,348	1.3	34.2
10	Septicemia (A40-A41)	2,249	1.3	32.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,785	1.0	26.0
12	Aortic aneurysm and dissection (I71)	1,659	0.9	24.2
13	Parkinson's disease (G20-G21)	1,392	0.8	20.3
14	Alzheimer's disease (G30)	1,380	0.8	20.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,143	0.6	16.6
...	All other causes (Residual)	20,262	11.4	295.0
White, non-Hispanic, male, 75-84 years				
...	All causes	285,608	100.0	6,420.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,264	28.8	1,849.3
2	Malignant neoplasms (C00-C97)	74,456	26.1	1,673.7
3	Chronic lower respiratory diseases (J40-J47)	21,973	7.7	493.9
4	Cerebrovascular diseases (I60-I69)	16,379	5.7	368.2
5	Diabetes mellitus (E10-E14)	8,302	2.9	186.6
6	Influenza and pneumonia (J10-J18)	7,767	2.7	174.6
7	Alzheimer's disease (G30)	7,326	2.6	164.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,311	2.2	141.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	5,802	2.0	130.4
10	Parkinson's disease (G20-G21)	5,164	1.8	116.1
11	Septicemia (A40-A41)	3,835	1.3	86.2
12	Pneumonitis due to solids and liquids (J69)	2,611	0.9	58.7
13	Aortic aneurysm and dissection (I71)	2,472	0.9	55.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,165	0.8	48.7
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,163	0.8	48.6
...	All other causes (Residual)	36,618	12.8	823.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 85 years and over				
...	All causes	210,795	100.0	15,401.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,555	34.9	5,374.1
2	Malignant neoplasms (C00-C97)	32,278	15.3	2,358.3
3	Cerebrovascular diseases (I60-I69)	14,378	6.8	1,050.5
4	Chronic lower respiratory diseases (J40-J47)	11,797	5.6	861.9
5	Influenza and pneumonia (J10-J18)	9,491	4.5	693.4
6	Alzheimer's disease (G30)	9,219	4.4	673.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	5,189	2.5	379.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,014	2.4	366.3
9	Diabetes mellitus (E10-E14)	4,537	2.2	331.5
10	Parkinson's disease (G20-G21)	3,393	1.6	247.9
11	Pneumonitis due to solids and liquids (J69)	3,255	1.5	237.8
12	Septicemia (A40-A41)	2,650	1.3	193.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,201	1.0	160.8
14	Atherosclerosis (I70)	1,787	0.8	130.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,368	0.6	100.0
...	All other causes (Residual)	30,683	14.6	2,241.8
White, non-Hispanic, female, all ages				
...	All causes	1,012,740	100.0	992.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	271,790	26.8	266.4
2	Malignant neoplasms (C00-C97)	219,474	21.7	215.1
3	Cerebrovascular diseases (I60-I69)	71,036	7.0	69.6
4	Chronic lower respiratory diseases (J40-J47)	62,322	6.2	61.1
5	Alzheimer's disease (G30)	45,750	4.5	44.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32,949	3.3	32.3
7	Influenza and pneumonia (J10-J18)	29,528	2.9	28.9
8	Diabetes mellitus (E10-E14)	26,794	2.6	26.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	16,539	1.6	16.2
10	Septicemia (A40-A41)	14,159	1.4	13.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11,475	1.1	11.2
12	Parkinson's disease (G20-G21)	7,535	0.7	7.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,340	0.7	7.2
14	Pneumonitis due to solids and liquids (J69)	7,309	0.7	7.2
15	Atherosclerosis (I70)	6,523	0.6	6.4
...	All other causes (Residual)	182,217	18.0	178.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 1-4 years				
...	All causes	996	100.0	22.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	34.2	7.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	131	13.2	2.9
3	Malignant neoplasms (C00-C97)	97	9.7	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	6.4	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	3.2	0.7
6	Influenza and pneumonia (J10-J18)	23	2.3	0.5
7	Septicemia (A40-A41)	18	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	11	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	11	1.1	*
11	Cerebrovascular diseases (I60-I69)	10	1.0	*
12	Meningococcal infection (A39)	5	0.5	*
13	Anemias (D50-D64)	4	0.4	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	231	23.2	5.2
White, non-Hispanic, female, 5-14 years				
...	All causes	1,518	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	537	35.4	4.6
2	Malignant neoplasms (C00-C97)	264	17.4	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	7.0	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	3.8	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	3.3	0.4
6	Influenza and pneumonia (J10-J18)	38	2.5	0.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	2.4	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	19	1.3	*
10	Septicemia (A40-A41)	18	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.9	*
12	Diabetes mellitus (E10-E14)	13	0.9	*
13	Certain conditions originating in the perinatal period (P00-P96)	10	0.7	*
14	Meningitis (G00,G03)	7	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	7	0.5	*
...	All other causes (Residual)	316	20.8	2.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-24 years				
...	All causes	5,416	100.0	42.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,769	51.1	21.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	505	9.3	3.9
3	Malignant neoplasms (C00-C97)	420	7.8	3.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	256	4.7	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	182	3.4	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	1.9	0.8
7	Pregnancy, childbirth and the puerperium (O00-099)	76	1.4	0.6
8	Cerebrovascular diseases (I60-I69)	52	1.0	0.4
9	Diabetes mellitus (E10-E14)	46	0.8	0.4
10	Influenza and pneumonia (J10-J18)	41	0.8	0.3
11	Septicemia (A40-A41)	31	0.6	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.4	0.2
12	Chronic lower respiratory diseases (J40-J47)	24	0.4	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	21	0.4	0.2
15	Pneumonitis due to solids and liquids (J69)	15	0.3	*
...	All other causes (Residual)	851	15.7	6.6
White, non-Hispanic, female, 25-34 years				
...	All causes	7,460	100.0	62.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,381	31.9	19.8
2	Malignant neoplasms (C00-C97)	1,072	14.4	8.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	713	9.6	5.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	536	7.2	4.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	309	4.1	2.6
6	Cerebrovascular diseases (I60-I69)	127	1.7	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	1.7	1.0
8	Pregnancy, childbirth and the puerperium (O00-099)	121	1.6	1.0
9	Diabetes mellitus (E10-E14)	117	1.6	1.0
10	Influenza and pneumonia (J10-J18)	107	1.4	0.9
11	Septicemia (A40-A41)	83	1.1	0.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	82	1.1	0.7
13	Chronic lower respiratory diseases (J40-J47)	72	1.0	0.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	64	0.9	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	56	0.8	0.5
...	All other causes (Residual)	1,496	20.1	12.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 35-44 years				
...	All causes	20,067	100.0	137.0
1	Malignant neoplasms (C00-C97)	5,496	27.4	37.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,667	18.3	25.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,233	11.1	15.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,270	6.3	8.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	647	3.2	4.4
6	Cerebrovascular diseases (I60-I69)	577	2.9	3.9
7	Diabetes mellitus (E10-E14)	455	2.3	3.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	389	1.9	2.7
9	Chronic lower respiratory diseases (J40-J47)	319	1.6	2.2
10	Influenza and pneumonia (J10-J18)	259	1.3	1.8
11	Septicemia (A40-A41)	239	1.2	1.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	230	1.1	1.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	186	0.9	1.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	132	0.7	0.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	119	0.6	0.8
...	All other causes (Residual)	3,849	19.2	26.3
White, non-Hispanic, female, 45-54 years				
...	All causes	46,809	100.0	298.7
1	Malignant neoplasms (C00-C97)	17,859	38.2	113.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,650	14.2	42.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,095	8.7	26.1
4	Cerebrovascular diseases (I60-I69)	1,593	3.4	10.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,552	3.3	9.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,524	3.3	9.7
7	Chronic lower respiratory diseases (J40-J47)	1,492	3.2	9.5
8	Diabetes mellitus (E10-E14)	1,275	2.7	8.1
9	Septicemia (A40-A41)	634	1.4	4.0
10	Influenza and pneumonia (J10-J18)	602	1.3	3.8
11	Viral hepatitis (B15-B19)	415	0.9	2.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	403	0.9	2.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	266	0.6	1.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	256	0.5	1.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	231	0.5	1.5
...	All other causes (Residual)	7,962	17.0	50.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 55-64 years				
...	All causes	81,849	100.0	677.2
1	Malignant neoplasms (C00-C97)	35,489	43.4	293.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,033	17.1	116.1
3	Chronic lower respiratory diseases (J40-J47)	5,429	6.6	44.9
4	Diabetes mellitus (E10-E14)	2,909	3.6	24.1
5	Cerebrovascular diseases (I60-I69)	2,817	3.4	23.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,654	3.2	22.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,520	1.9	12.6
8	Septicemia (A40-A41)	1,225	1.5	10.1
9	Influenza and pneumonia (J10-J18)	1,058	1.3	8.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	972	1.2	8.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	869	1.1	7.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	435	0.5	3.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	401	0.5	3.3
14	Alzheimer's disease (G30)	306	0.4	2.5
15	Aortic aneurysm and dissection (I71)	299	0.4	2.5
...	All other causes (Residual)	11,433	14.0	94.6
White, non-Hispanic, female, 65-74 years				
...	All causes	137,441	100.0	1,729.6
1	Malignant neoplasms (C00-C97)	50,412	36.7	634.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,178	20.5	354.6
3	Chronic lower respiratory diseases (J40-J47)	13,174	9.6	165.8
4	Cerebrovascular diseases (I60-I69)	6,642	4.8	83.6
5	Diabetes mellitus (E10-E14)	4,885	3.6	61.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,658	1.9	33.4
7	Influenza and pneumonia (J10-J18)	2,269	1.7	28.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	2,221	1.6	27.9
9	Septicemia (A40-A41)	2,164	1.6	27.2
10	Alzheimer's disease (G30)	1,828	1.3	23.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,581	1.2	19.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	998	0.7	12.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	815	0.6	10.3
14	Aortic aneurysm and dissection (I71)	784	0.6	9.9
15	Parkinson's disease (G20-G21)	713	0.5	9.0
...	All other causes (Residual)	18,119	13.2	228.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 75-84 years				
...	All causes	296,104	100.0	4,579.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,820	26.3	1,203.6
2	Malignant neoplasms (C00-C97)	67,858	22.9	1,049.5
3	Chronic lower respiratory diseases (J40-J47)	23,968	8.1	370.7
4	Cerebrovascular diseases (I60-I69)	22,451	7.6	347.2
5	Alzheimer's disease (G30)	13,143	4.4	203.3
6	Diabetes mellitus (E10-E14)	8,991	3.0	139.1
7	Influenza and pneumonia (J10-J18)	7,871	2.7	121.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,631	1.9	87.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	5,496	1.9	85.0
10	Septicemia (A40-A41)	4,546	1.5	70.3
11	Parkinson's disease (G20-G21)	3,281	1.1	50.7
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,252	1.1	50.3
13	Pneumonitis due to solids and liquids (J69)	2,139	0.7	33.1
14	Aortic aneurysm and dissection (I71)	1,918	0.6	29.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,897	0.6	29.3
...	All other causes (Residual)	45,842	15.5	709.0
White, non-Hispanic, female, 85 years and over				
...	All causes	409,451	100.0	13,683.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	141,990	34.7	4,745.1
2	Malignant neoplasms (C00-C97)	40,492	9.9	1,353.2
3	Cerebrovascular diseases (I60-I69)	36,725	9.0	1,227.3
4	Alzheimer's disease (G30)	30,427	7.4	1,016.8
5	Chronic lower respiratory diseases (J40-J47)	17,814	4.4	595.3
6	Influenza and pneumonia (J10-J18)	17,226	4.2	575.7
7	Diabetes mellitus (E10-E14)	8,101	2.0	270.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,948	1.9	265.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	7,198	1.8	240.5
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6,587	1.6	220.1
11	Septicemia (A40-A41)	5,164	1.3	172.6
12	Atherosclerosis (I70)	4,375	1.1	146.2
13	Pneumonitis due to solids and liquids (J69)	4,115	1.0	137.5
14	Parkinson's disease (G20-G21)	3,388	0.8	113.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,196	0.5	73.4
...	All other causes (Residual)	75,705	18.5	2,529.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	289,163	100.0	774.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,302	25.3	196.3
2	Malignant neoplasms (C00-C97)	62,510	21.6	167.4
3	Cerebrovascular diseases (I60-I69)	17,378	6.0	46.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,393	4.6	35.9
5	Diabetes mellitus (E10-E14)	12,835	4.4	34.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,522	2.9	22.8
7	Chronic lower respiratory diseases (J40-J47)	8,136	2.8	21.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	8,017	2.8	21.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,908	2.4	18.5
10	Septicemia (A40-A41)	6,159	2.1	16.5
11	Influenza and pneumonia (J10-J18)	5,716	2.0	15.3
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4,906	1.7	13.1
13	Certain conditions originating in the perinatal period (P00-P96)	4,837	1.7	13.0
14	Alzheimer's disease (G30)	4,579	1.6	12.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,495	0.9	6.7
...	All other causes (Residual)	49,470	17.1	132.5
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,065	100.0	42.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	316	29.7	12.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	13.0	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	9.8	4.1
4	Malignant neoplasms (C00-C97)	62	5.8	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.2	1.8
6	Chronic lower respiratory diseases (J40-J47)	30	2.8	1.2
7	Influenza and pneumonia (J10-J18)	28	2.6	1.1
8	Certain conditions originating in the perinatal period (P00-P96)	23	2.2	0.9
9	Septicemia (A40-A41)	20	1.9	0.8
10	Cerebrovascular diseases (I60-I69)	14	1.3	*
11	Anemias (D50-D64)	11	1.0	*
12	Meningitis (G00,G03)	6	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	5	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	255	23.9	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-14 years				
...	All causes	1,500	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	497	33.1	7.9
2	Malignant neoplasms (C00-C97)	165	11.0	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	145	9.7	2.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	5.0	1.2
6	Chronic lower respiratory diseases (J40-J47)	51	3.4	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	2.7	0.7
8	Cerebrovascular diseases (I60-I69)	23	1.5	0.4
9	Influenza and pneumonia (J10-J18)	22	1.5	0.4
10	Septicemia (A40-A41)	20	1.3	0.3
11	Anemias (D50-D64)	15	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.9	*
13	Diabetes mellitus (E10-E14)	12	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.5	*
15	Meningitis (G00,G03)	7	0.5	*
...	All other causes (Residual)	325	21.7	5.2
Black, non-Hispanic, both sexes, 15-24 years				
...	All causes	7,244	100.0	115.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,030	41.8	48.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,734	23.9	27.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	428	5.9	6.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	333	4.6	5.3
5	Malignant neoplasms (C00-C97)	259	3.6	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	108	1.5	1.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	1.4	1.6
8	Anemias (D50-D64)	77	1.1	1.2
9	Chronic lower respiratory diseases (J40-J47)	74	1.0	1.2
10	Diabetes mellitus (E10-E14)	73	1.0	1.2
11	Pregnancy, childbirth and the puerperium (O00-099)	69	1.0	1.1
12	Cerebrovascular diseases (I60-I69)	51	0.7	0.8
13	Septicemia (A40-A41)	44	0.6	0.7
14	Influenza and pneumonia (J10-J18)	31	0.4	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	29	0.4	0.5
...	All other causes (Residual)	805	11.1	12.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-34 years				
...	All causes	9,850	100.0	186.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,538	25.8	47.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,754	17.8	33.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,008	10.2	19.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	824	8.4	15.6
5	Malignant neoplasms (C00-C97)	674	6.8	12.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	474	4.8	9.0
7	Diabetes mellitus (E10-E14)	218	2.2	4.1
8	Cerebrovascular diseases (I60-I69)	163	1.7	3.1
9	Anemias (D50-D64)	131	1.3	2.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	120	1.2	2.3
11	Pregnancy, childbirth and the puerperium (O00-099)	115	1.2	2.2
12	Septicemia (A40-A41)	106	1.1	2.0
13	Chronic lower respiratory diseases (J40-J47)	96	1.0	1.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	0.8	1.5
15	Influenza and pneumonia (J10-J18)	77	0.8	1.5
...	All other causes (Residual)	1,471	14.9	27.8
Black, non-Hispanic, both sexes, 35-44 years				
...	All causes	17,930	100.0	328.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,165	17.7	58.0
2	Malignant neoplasms (C00-C97)	2,521	14.1	46.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	2,298	12.8	42.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,161	12.1	39.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,351	7.5	24.8
6	Cerebrovascular diseases (I60-I69)	683	3.8	12.5
7	Diabetes mellitus (E10-E14)	534	3.0	9.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	415	2.3	7.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	312	1.7	5.7
10	Septicemia (A40-A41)	258	1.4	4.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	234	1.3	4.3
12	Chronic lower respiratory diseases (J40-J47)	217	1.2	4.0
13	Influenza and pneumonia (J10-J18)	214	1.2	3.9
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	194	1.1	3.6
15	Anemias (D50-D64)	144	0.8	2.6
...	All other causes (Residual)	3,229	18.0	59.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-54 years				
...	All causes	37,298	100.0	761.4
1	Malignant neoplasms (C00-C97)	8,805	23.6	179.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,613	23.1	175.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,647	7.1	54.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,475	6.6	50.5
5	Cerebrovascular diseases (I60-I69)	1,986	5.3	40.5
6	Diabetes mellitus (E10-E14)	1,396	3.7	28.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	884	2.4	18.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	785	2.1	16.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	745	2.0	15.2
10	Chronic lower respiratory diseases (J40-J47)	697	1.9	14.2
11	Septicemia (A40-A41)	654	1.8	13.4
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	527	1.4	10.8
13	Influenza and pneumonia (J10-J18)	494	1.3	10.1
14	Viral hepatitis (B15-B19)	349	0.9	7.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	309	0.8	6.3
...	All other causes (Residual)	5,932	15.9	121.1
Black, non-Hispanic, both sexes, 55-64 years				
...	All causes	44,937	100.0	1,513.6
1	Malignant neoplasms (C00-C97)	13,835	30.8	466.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,977	26.7	403.4
3	Cerebrovascular diseases (I60-I69)	2,420	5.4	81.5
4	Diabetes mellitus (E10-E14)	2,415	5.4	81.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,441	3.2	48.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	1,340	3.0	45.1
7	Chronic lower respiratory diseases (J40-J47)	1,195	2.7	40.3
8	Septicemia (A40-A41)	934	2.1	31.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	902	2.0	30.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	751	1.7	25.3
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	684	1.5	23.0
12	Influenza and pneumonia (J10-J18)	603	1.3	20.3
13	Viral hepatitis (B15-B19)	253	0.6	8.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	244	0.5	8.2
15	Aortic aneurysm and dissection (I71)	207	0.5	7.0
...	All other causes (Residual)	5,736	12.8	193.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65-74 years				
...	All causes	51,166	100.0	2,979.6
1	Malignant neoplasms (C00-C97)	15,560	30.4	906.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,840	27.0	806.0
3	Cerebrovascular diseases (I60-I69)	3,194	6.2	186.0
4	Diabetes mellitus (E10-E14)	3,101	6.1	180.6
5	Chronic lower respiratory diseases (J40-J47)	1,998	3.9	116.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	1,744	3.4	101.6
7	Septicemia (A40-A41)	1,171	2.3	68.2
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	948	1.9	55.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	921	1.8	53.6
10	Influenza and pneumonia (J10-J18)	914	1.8	53.2
11	Alzheimer's disease (G30)	390	0.8	22.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	352	0.7	20.5
13	Aortic aneurysm and dissection (I71)	254	0.5	14.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	240	0.5	14.0
15	Pneumonitis due to solids and liquids (J69)	232	0.5	13.5
...	All other causes (Residual)	6,307	12.3	367.3
Black, non-Hispanic, both sexes, 75-84 years				
...	All causes	61,319	100.0	6,218.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,118	29.5	1,837.3
2	Malignant neoplasms (C00-C97)	14,156	23.1	1,435.5
3	Cerebrovascular diseases (I60-I69)	4,691	7.7	475.7
4	Diabetes mellitus (E10-E14)	3,205	5.2	325.0
5	Chronic lower respiratory diseases (J40-J47)	2,414	3.9	244.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	2,090	3.4	211.9
7	Alzheimer's disease (G30)	1,579	2.6	160.1
8	Septicemia (A40-A41)	1,577	2.6	159.9
9	Influenza and pneumonia (J10-J18)	1,497	2.4	151.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,226	2.0	124.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	964	1.6	97.8
12	Pneumonitis due to solids and liquids (J69)	525	0.9	53.2
13	Parkinson's disease (G20-G21)	339	0.6	34.4
14	Aortic aneurysm and dissection (I71)	307	0.5	31.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	280	0.5	28.4
...	All other causes (Residual)	8,351	13.6	846.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85 years and over				
...	All causes	48,481	100.0	13,239.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,016	33.0	4,373.6
2	Malignant neoplasms (C00-C97)	6,461	13.3	1,764.4
3	Cerebrovascular diseases (I60-I69)	4,110	8.5	1,122.3
4	Alzheimer's disease (G30)	2,559	5.3	698.8
5	Diabetes mellitus (E10-E14)	1,880	3.9	513.4
6	Influenza and pneumonia (J10-J18)	1,741	3.6	475.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	1,574	3.2	429.8
8	Chronic lower respiratory diseases (J40-J47)	1,352	2.8	369.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,264	2.6	345.2
10	Septicemia (A40-A41)	1,248	2.6	340.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	647	1.3	176.7
12	Pneumonitis due to solids and liquids (J69)	569	1.2	155.4
13	Atherosclerosis (I70)	365	0.8	99.7
14	Parkinson's disease (G20-G21)	210	0.4	57.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	204	0.4	55.7
...	All other causes (Residual)	8,281	17.1	2,261.4
Black, non-Hispanic, male, all ages				
...	All causes	147,010	100.0	825.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,874	24.4	201.5
2	Malignant neoplasms (C00-C97)	32,363	22.0	181.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,130	6.2	51.3
4	Cerebrovascular diseases (I60-I69)	7,448	5.1	41.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,288	5.0	40.9
6	Diabetes mellitus (E10-E14)	5,666	3.9	31.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,600	3.1	25.8
8	Chronic lower respiratory diseases (J40-J47)	4,406	3.0	24.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	3,612	2.5	20.3
10	Septicemia (A40-A41)	2,755	1.9	15.5
11	Certain conditions originating in the perinatal period (P00-P96)	2,712	1.8	15.2
12	Influenza and pneumonia (J10-J18)	2,697	1.8	15.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,108	1.4	11.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,634	1.1	9.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,592	1.1	8.9
...	All other causes (Residual)	23,125	15.7	129.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 1-4 years				
...	All causes	606	100.0	47.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	29.5	14.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	14.5	6.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	7.6	3.6
4	Malignant neoplasms (C00-C97)	36	5.9	2.8
5	Chronic lower respiratory diseases (J40-J47)	21	3.5	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.1	*
7	Influenza and pneumonia (J10-J18)	16	2.6	*
7	Certain conditions originating in the perinatal period (P00-P96)	16	2.6	*
9	Septicemia (A40-A41)	13	2.1	*
10	Cerebrovascular diseases (I60-I69)	10	1.7	*
11	Anemias (D50-D64)	5	0.8	*
12	Meningitis (G00,G03)	3	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	3	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
...	All other causes (Residual)	147	24.3	11.5
Black, non-Hispanic, male, 5-14 years				
...	All causes	885	100.0	27.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	312	35.3	9.8
2	Malignant neoplasms (C00-C97)	98	11.1	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	10.5	2.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	4.2	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.0	1.1
6	Chronic lower respiratory diseases (J40-J47)	33	3.7	1.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	3.1	0.8
8	Cerebrovascular diseases (I60-I69)	14	1.6	*
9	Septicemia (A40-A41)	13	1.5	*
9	Influenza and pneumonia (J10-J18)	13	1.5	*
11	Anemias (D50-D64)	9	1.0	*
12	Meningitis (G00,G03)	6	0.7	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
15	Diabetes mellitus (E10-E14)	3	0.3	*
15	Diseases of appendix (K35-K38)	3	0.3	*
...	All other causes (Residual)	180	20.3	5.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-24 years				
...	All causes	5,615	100.0	177.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,750	49.0	87.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,301	23.2	41.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	375	6.7	11.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	228	4.1	7.2
5	Malignant neoplasms (C00-C97)	151	2.7	4.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	1.1	2.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	57	1.0	1.8
8	Chronic lower respiratory diseases (J40-J47)	47	0.8	1.5
9	Anemias (D50-D64)	45	0.8	1.4
10	Diabetes mellitus (E10-E14)	39	0.7	1.2
11	Septicemia (A40-A41)	24	0.4	0.8
11	Cerebrovascular diseases (I60-I69)	24	0.4	0.8
11	Legal intervention (Y35,Y89.0)	24	0.4	0.8
14	Influenza and pneumonia (J10-J18)	23	0.4	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	11	0.2	*
...	All other causes (Residual)	453	8.1	14.3
Black, non-Hispanic, male, 25-34 years				
...	All causes	6,706	100.0	264.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,270	33.9	89.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,332	19.9	52.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	624	9.3	24.6
4	Human immunodeficiency virus (HIV) disease (B20-B24)	455	6.8	17.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	394	5.9	15.5
6	Malignant neoplasms (C00-C97)	313	4.7	12.3
7	Diabetes mellitus (E10-E14)	127	1.9	5.0
8	Cerebrovascular diseases (I60-I69)	89	1.3	3.5
9	Anemias (D50-D64)	69	1.0	2.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	60	0.9	2.4
11	Chronic lower respiratory diseases (J40-J47)	52	0.8	2.0
12	Septicemia (A40-A41)	49	0.7	1.9
13	Influenza and pneumonia (J10-J18)	44	0.7	1.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	0.6	1.7
15	Legal intervention (Y35,Y89.0)	39	0.6	1.5
...	All other causes (Residual)	746	11.1	29.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-44 years				
...	All causes	10,451	100.0	407.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,969	18.8	76.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,521	14.6	59.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,454	13.9	56.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,075	10.3	41.9
5	Malignant neoplasms (C00-C97)	960	9.2	37.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	333	3.2	13.0
7	Cerebrovascular diseases (I60-I69)	329	3.1	12.8
8	Diabetes mellitus (E10-E14)	307	2.9	12.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	164	1.6	6.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	141	1.3	5.5
11	Influenza and pneumonia (J10-J18)	127	1.2	5.0
12	Septicemia (A40-A41)	121	1.2	4.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	103	1.0	4.0
13	Chronic lower respiratory diseases (J40-J47)	103	1.0	4.0
15	Aortic aneurysm and dissection (I71)	81	0.8	3.2
...	All other causes (Residual)	1,663	15.9	64.8
Black, non-Hispanic, male, 45-54 years				
...	All causes	21,946	100.0	969.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,497	25.0	242.9
2	Malignant neoplasms (C00-C97)	4,309	19.6	190.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,897	8.6	83.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,728	7.9	76.4
5	Cerebrovascular diseases (I60-I69)	1,039	4.7	45.9
6	Diabetes mellitus (E10-E14)	772	3.5	34.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	624	2.8	27.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	599	2.7	26.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	438	2.0	19.4
10	Chronic lower respiratory diseases (J40-J47)	351	1.6	15.5
11	Septicemia (A40-A41)	344	1.6	15.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	307	1.4	13.6
13	Influenza and pneumonia (J10-J18)	299	1.4	13.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	238	1.1	10.5
15	Viral hepatitis (B15-B19)	234	1.1	10.3
...	All other causes (Residual)	3,270	14.9	144.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 55-64 years				
...	All causes	26,396	100.0	1,993.8
1	Malignant neoplasms (C00-C97)	7,695	29.2	581.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,404	28.0	559.3
3	Cerebrovascular diseases (I60-I69)	1,411	5.3	106.6
4	Diabetes mellitus (E10-E14)	1,274	4.8	96.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,071	4.1	80.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	746	2.8	56.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	689	2.6	52.0
8	Chronic lower respiratory diseases (J40-J47)	660	2.5	49.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	528	2.0	39.9
10	Septicemia (A40-A41)	485	1.8	36.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	395	1.5	29.8
12	Influenza and pneumonia (J10-J18)	372	1.4	28.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	193	0.7	14.6
14	Viral hepatitis (B15-B19)	171	0.6	12.9
15	Aortic aneurysm and dissection (I71)	134	0.5	10.1
...	All other causes (Residual)	3,168	12.0	239.3
Black, non-Hispanic, male, 65-74 years				
...	All causes	27,303	100.0	3,814.1
1	Malignant neoplasms (C00-C97)	8,630	31.6	1,205.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,586	27.8	1,059.7
3	Cerebrovascular diseases (I60-I69)	1,636	6.0	228.5
4	Diabetes mellitus (E10-E14)	1,413	5.2	197.4
5	Chronic lower respiratory diseases (J40-J47)	1,146	4.2	160.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	750	2.7	104.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	589	2.2	82.3
8	Septicemia (A40-A41)	586	2.1	81.9
9	Influenza and pneumonia (J10-J18)	510	1.9	71.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	465	1.7	65.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	225	0.8	31.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	179	0.7	25.0
13	Alzheimer's disease (G30)	163	0.6	22.8
14	Pneumonitis due to solids and liquids (J69)	143	0.5	20.0
15	Aortic aneurysm and dissection (I71)	132	0.5	18.4
...	All other causes (Residual)	3,150	11.5	440.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 75-84 years				
...	All causes	27,723	100.0	7,771.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,952	28.7	2,229.0
2	Malignant neoplasms (C00-C97)	7,304	26.3	2,047.4
3	Cerebrovascular diseases (I60-I69)	1,821	6.6	510.4
4	Chronic lower respiratory diseases (J40-J47)	1,360	4.9	381.2
5	Diabetes mellitus (E10-E14)	1,236	4.5	346.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	891	3.2	249.8
7	Influenza and pneumonia (J10-J18)	683	2.5	191.5
8	Septicemia (A40-A41)	654	2.4	183.3
9	Alzheimer's disease (G30)	545	2.0	152.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	518	1.9	145.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	496	1.8	139.0
12	Pneumonitis due to solids and liquids (J69)	289	1.0	81.0
13	Parkinson's disease (G20-G21)	196	0.7	54.9
14	Aortic aneurysm and dissection (I71)	140	0.5	39.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	123	0.4	34.5
...	All other causes (Residual)	3,515	12.7	985.3
Black, non-Hispanic, male, 85 years and over				
...	All causes	14,690	100.0	13,978.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,495	30.6	4,277.2
2	Malignant neoplasms (C00-C97)	2,859	19.5	2,720.4
3	Cerebrovascular diseases (I60-I69)	1,049	7.1	998.2
4	Chronic lower respiratory diseases (J40-J47)	627	4.3	596.6
5	Influenza and pneumonia (J10-J18)	565	3.8	537.6
6	Alzheimer's disease (G30)	554	3.8	527.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	512	3.5	487.2
8	Diabetes mellitus (E10-E14)	495	3.4	471.0
9	Septicemia (A40-A41)	399	2.7	379.7
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	309	2.1	294.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	1.6	227.4
12	Pneumonitis due to solids and liquids (J69)	216	1.5	205.5
13	Parkinson's disease (G20-G21)	104	0.7	99.0
14	Atherosclerosis (I70)	81	0.6	77.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.4	56.1
...	All other causes (Residual)	2,127	14.5	2,023.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	142,153	100.0	727.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,428	26.3	191.6
2	Malignant neoplasms (C00-C97)	30,147	21.2	154.3
3	Cerebrovascular diseases (I60-I69)	9,930	7.0	50.8
4	Diabetes mellitus (E10-E14)	7,169	5.0	36.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	4,405	3.1	22.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,263	3.0	21.8
7	Chronic lower respiratory diseases (J40-J47)	3,730	2.6	19.1
8	Septicemia (A40-A41)	3,404	2.4	17.4
9	Alzheimer's disease (G30)	3,296	2.3	16.9
10	Influenza and pneumonia (J10-J18)	3,019	2.1	15.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,798	2.0	14.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,308	1.6	11.8
13	Certain conditions originating in the perinatal period (P00-P96)	2,125	1.5	10.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,234	0.9	6.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	861	0.6	4.4
...	All other causes (Residual)	26,036	18.3	133.3
Black, non-Hispanic, female, 1-4 years				
...	All causes	459	100.0	37.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	137	29.8	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	12.6	4.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	10.9	4.1
4	Malignant neoplasms (C00-C97)	26	5.7	2.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	5.7	2.1
6	Influenza and pneumonia (J10-J18)	12	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	9	2.0	*
8	Septicemia (A40-A41)	7	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
10	Anemias (D50-D64)	6	1.3	*
11	Cerebrovascular diseases (I60-I69)	4	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
12	Meningitis (G00,G03)	3	0.7	*
12	Pneumonitis due to solids and liquids (J69)	3	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	106	23.1	8.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-14 years				
...	All causes	615	100.0	20.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	30.1	6.0
2	Malignant neoplasms (C00-C97)	67	10.9	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	8.5	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	7.0	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	6.5	1.3
6	Chronic lower respiratory diseases (J40-J47)	18	2.9	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.6	*
9	Diabetes mellitus (E10-E14)	9	1.5	*
9	Cerebrovascular diseases (I60-I69)	9	1.5	*
9	Influenza and pneumonia (J10-J18)	9	1.5	*
12	Septicemia (A40-A41)	7	1.1	*
13	Anemias (D50-D64)	6	1.0	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
...	All other causes (Residual)	140	22.8	4.5
Black, non-Hispanic, female, 15-24 years				
...	All causes	1,629	100.0	52.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	433	26.6	14.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	280	17.2	9.0
3	Malignant neoplasms (C00-C97)	108	6.6	3.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	105	6.4	3.4
5	Pregnancy, childbirth and the puerperium (O00-099)	69	4.2	2.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	53	3.3	1.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	51	3.1	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	2.2	1.2
9	Diabetes mellitus (E10-E14)	34	2.1	1.1
10	Anemias (D50-D64)	32	2.0	1.0
11	Cerebrovascular diseases (I60-I69)	27	1.7	0.9
11	Chronic lower respiratory diseases (J40-J47)	27	1.7	0.9
13	Septicemia (A40-A41)	20	1.2	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	18	1.1	*
15	Influenza and pneumonia (J10-J18)	8	0.5	*
...	All other causes (Residual)	328	20.1	10.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-34 years				
...	All causes	3,144	100.0	114.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	422	13.4	15.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	384	12.2	13.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	369	11.7	13.4
4	Malignant neoplasms (C00-C97)	361	11.5	13.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	268	8.5	9.7
6	Pregnancy, childbirth and the puerperium (O00-099)	115	3.7	4.2
7	Diabetes mellitus (E10-E14)	91	2.9	3.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	2.5	2.9
9	Cerebrovascular diseases (I60-I69)	74	2.4	2.7
10	Anemias (D50-D64)	62	2.0	2.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	60	1.9	2.2
12	Septicemia (A40-A41)	57	1.8	2.1
13	Chronic lower respiratory diseases (J40-J47)	44	1.4	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	1.2	1.4
15	Influenza and pneumonia (J10-J18)	33	1.0	1.2
...	All other causes (Residual)	686	21.8	24.9
Black, non-Hispanic, female, 35-44 years				
...	All causes	7,479	100.0	258.5
1	Malignant neoplasms (C00-C97)	1,561	20.9	54.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,196	16.0	41.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	844	11.3	29.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	640	8.6	22.1
5	Cerebrovascular diseases (I60-I69)	354	4.7	12.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	276	3.7	9.5
7	Diabetes mellitus (E10-E14)	227	3.0	7.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	148	2.0	5.1
9	Septicemia (A40-A41)	137	1.8	4.7
10	Chronic lower respiratory diseases (J40-J47)	114	1.5	3.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	93	1.2	3.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	91	1.2	3.1
13	Influenza and pneumonia (J10-J18)	87	1.2	3.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	1.1	2.8
15	Anemias (D50-D64)	67	0.9	2.3
...	All other causes (Residual)	1,562	20.9	54.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 45-54 years				
...	All causes	15,352	100.0	582.4
1	Malignant neoplasms (C00-C97)	4,496	29.3	170.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,116	20.3	118.2
3	Cerebrovascular diseases (I60-I69)	947	6.2	35.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	750	4.9	28.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	747	4.9	28.3
6	Diabetes mellitus (E10-E14)	624	4.1	23.7
7	Chronic lower respiratory diseases (J40-J47)	346	2.3	13.1
8	Septicemia (A40-A41)	310	2.0	11.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	307	2.0	11.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	285	1.9	10.8
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	220	1.4	8.3
12	Influenza and pneumonia (J10-J18)	195	1.3	7.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	161	1.0	6.1
14	Viral hepatitis (B15-B19)	115	0.7	4.4
15	Anemias (D50-D64)	77	0.5	2.9
...	All other causes (Residual)	2,656	17.3	100.8
Black, non-Hispanic, female, 55-64 years				
...	All causes	18,541	100.0	1,127.1
1	Malignant neoplasms (C00-C97)	6,140	33.1	373.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,573	24.7	278.0
3	Diabetes mellitus (E10-E14)	1,141	6.2	69.4
4	Cerebrovascular diseases (I60-I69)	1,009	5.4	61.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	594	3.2	36.1
6	Chronic lower respiratory diseases (J40-J47)	535	2.9	32.5
7	Septicemia (A40-A41)	449	2.4	27.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	370	2.0	22.5
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	289	1.6	17.6
10	Influenza and pneumonia (J10-J18)	231	1.2	14.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	223	1.2	13.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	213	1.1	12.9
13	Viral hepatitis (B15-B19)	82	0.4	5.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.4	4.9
15	Aortic aneurysm and dissection (I71)	73	0.4	4.4
...	All other causes (Residual)	2,539	13.7	154.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 65-74 years				
...	All causes	23,863	100.0	2,383.1
1	Malignant neoplasms (C00-C97)	6,930	29.0	692.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,254	26.2	624.6
3	Diabetes mellitus (E10-E14)	1,688	7.1	168.6
4	Cerebrovascular diseases (I60-I69)	1,558	6.5	155.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	994	4.2	99.3
6	Chronic lower respiratory diseases (J40-J47)	852	3.6	85.1
7	Septicemia (A40-A41)	585	2.5	58.4
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	483	2.0	48.2
9	Influenza and pneumonia (J10-J18)	404	1.7	40.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	332	1.4	33.2
11	Alzheimer's disease (G30)	227	1.0	22.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	127	0.5	12.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	123	0.5	12.3
14	Aortic aneurysm and dissection (I71)	122	0.5	12.2
15	Pneumonitis due to solids and liquids (J69)	89	0.4	8.9
...	All other causes (Residual)	3,095	13.0	309.1
Black, non-Hispanic, female, 75-84 years				
...	All causes	33,596	100.0	5,338.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,166	30.3	1,615.3
2	Malignant neoplasms (C00-C97)	6,852	20.4	1,088.7
3	Cerebrovascular diseases (I60-I69)	2,870	8.5	456.0
4	Diabetes mellitus (E10-E14)	1,969	5.9	312.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	1,199	3.6	190.5
6	Chronic lower respiratory diseases (J40-J47)	1,054	3.1	167.5
7	Alzheimer's disease (G30)	1,034	3.1	164.3
8	Septicemia (A40-A41)	923	2.7	146.7
9	Influenza and pneumonia (J10-J18)	814	2.4	129.3
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	730	2.2	116.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	446	1.3	70.9
12	Pneumonitis due to solids and liquids (J69)	236	0.7	37.5
13	Aortic aneurysm and dissection (I71)	167	0.5	26.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	157	0.5	24.9
15	Parkinson's disease (G20-G21)	143	0.4	22.7
...	All other causes (Residual)	4,836	14.4	768.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2005

[Rates per 100,000 population in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
	Black, non-Hispanic, female, 85 years and over			
...	All causes	33,791	100.0	12,941.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,521	34.1	4,412.4
2	Malignant neoplasms (C00-C97)	3,602	10.7	1,379.5
3	Cerebrovascular diseases (I60-I69)	3,061	9.1	1,172.3
4	Alzheimer's disease (G30)	2,005	5.9	767.9
5	Diabetes mellitus (E10-E14)	1,385	4.1	530.4
6	Influenza and pneumonia (J10-J18)	1,176	3.5	450.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)	1,062	3.1	406.7
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	955	2.8	365.8
9	Septicemia (A40-A41)	849	2.5	325.2
10	Chronic lower respiratory diseases (J40-J47)	725	2.1	277.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	408	1.2	156.3
12	Pneumonitis due to solids and liquids (J69)	353	1.0	135.2
13	Atherosclerosis (I70)	284	0.8	108.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	145	0.4	55.5
15	Aortic aneurysm and dissection (I71)	133	0.4	50.9
...	All other causes (Residual)	6,127	18.1	2,346.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable