

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	156,419	100.0	295.0
1	Malignant neoplasms (C00-C97)	34,147	21.8	64.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,595	20.2	59.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,447	7.3	21.6
4	Cerebrovascular diseases (I60-I69)	7,767	5.0	14.6
5	Diabetes mellitus (E10-E14)	7,363	4.7	13.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,988	3.2	9.4
7	Chronic lower respiratory diseases (J40-J47)	4,477	2.9	8.4
8	Alzheimer's disease (G30)	3,772	2.4	7.1
9	Influenza and pneumonia (J09-J18)	3,049	1.9	5.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,964	1.9	5.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,837	1.8	5.4
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,740	1.8	5.2
13	Certain conditions originating in the perinatal period (P00-P96)	2,429	1.6	4.6
14	Septicemia (A40-A41)	2,304	1.5	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,859	1.2	3.5
...	All other causes (Residual)	32,681	20.9	61.6
Hispanic, both sexes, 1-4 years				
...	All causes	899	100.0	21.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	269	29.9	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	14.0	3.1
3	Malignant neoplasms (C00-C97)	99	11.0	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	6.7	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.2	0.7
6	Influenza and pneumonia (J09-J18)	22	2.4	0.5
7	Septicemia (A40-A41)	14	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.3	*
10	Cerebrovascular diseases (I60-I69)	11	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	9	1.0	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.6	*
13	Pneumonitis due to solids and liquids (J69)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Anemias (D50-D64)	2	0.2	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Meningitis (G00,G03)	2	0.2	*
...	All other causes (Residual)	216	24.0	5.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,083	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	307	28.3	3.2
2	Malignant neoplasms (C00-C97)	219	20.2	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	7.1	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	5.3	0.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	4.2	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.9	0.3
7	Cerebrovascular diseases (I60-I69)	24	2.2	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.8	0.2
9	Influenza and pneumonia (J09-J18)	18	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	18	1.7	*
11	Septicemia (A40-A41)	16	1.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
13	Meningitis (G00,G03)	5	0.5	*
14	Anemias (D50-D64)	4	0.4	*
14	Diabetes mellitus (E10-E14)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	225	20.8	2.3
Hispanic, both sexes, 15-24 years				
...	All causes	4,773	100.0	52.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,871	39.2	20.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	915	19.2	10.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	675	14.1	7.4
4	Malignant neoplasms (C00-C97)	370	7.8	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	2.9	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	1.4	0.7
7	Cerebrovascular diseases (I60-I69)	36	0.8	0.4
8	Legal intervention (Y35,Y89.0)	30	0.6	0.3
9	Influenza and pneumonia (J09-J18)	26	0.5	0.3
9	Pregnancy, childbirth and the puerperium (O00-O99)	26	0.5	0.3
11	Diabetes mellitus (E10-E14)	23	0.5	0.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	19	0.4	*
14	Septicemia (A40-A41)	15	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.3	*
...	All other causes (Residual)	526	11.0	5.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	6,216	100.0	71.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,160	34.7	24.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	781	12.6	9.0
3	Malignant neoplasms (C00-C97)	679	10.9	7.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	632	10.2	7.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	356	5.7	4.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	121	1.9	1.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	107	1.7	1.2
8	Cerebrovascular diseases (I60-I69)	98	1.6	1.1
9	Diabetes mellitus (E10-E14)	86	1.4	1.0
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	0.9	0.7
11	Pregnancy, childbirth and the puerperium (O00-O99)	56	0.9	0.6
12	Septicemia (A40-A41)	47	0.8	0.5
13	Influenza and pneumonia (J09-J18)	43	0.7	0.5
14	Legal intervention (Y35,Y89.0)	36	0.6	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.5	0.4
...	All other causes (Residual)	925	14.9	10.7
Hispanic, both sexes, 35-44 years				
...	All causes	8,338	100.0	109.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,812	21.7	23.7
2	Malignant neoplasms (C00-C97)	1,590	19.1	20.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	940	11.3	12.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	547	6.6	7.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	473	5.7	6.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	453	5.4	5.9
7	Cerebrovascular diseases (I60-I69)	288	3.5	3.8
8	Diabetes mellitus (E10-E14)	232	2.8	3.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	184	2.2	2.4
10	Influenza and pneumonia (J09-J18)	103	1.2	1.3
11	Septicemia (A40-A41)	93	1.1	1.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	1.1	1.2
13	Viral hepatitis (B15-B19)	66	0.8	0.9
14	Pregnancy, childbirth and the puerperium (O00-O99)	54	0.6	0.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	0.6	0.6
...	All other causes (Residual)	1,366	16.4	17.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	15,778	100.0	267.9
1	Malignant neoplasms (C00-C97)	4,145	26.3	70.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,473	15.7	42.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,745	11.1	29.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,362	8.6	23.1
5	Diabetes mellitus (E10-E14)	680	4.3	11.5
6	Cerebrovascular diseases (I60-I69)	658	4.2	11.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	457	2.9	7.8
8	Viral hepatitis (B15-B19)	374	2.4	6.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	336	2.1	5.7
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	244	1.5	4.1
11	Septicemia (A40-A41)	232	1.5	3.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	221	1.4	3.8
13	Influenza and pneumonia (J09-J18)	163	1.0	2.8
14	Chronic lower respiratory diseases (J40-J47)	160	1.0	2.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	110	0.7	1.9
...	All other causes (Residual)	2,418	15.3	41.1
Hispanic, both sexes, 55-64 years				
...	All causes	22,620	100.0	622.3
1	Malignant neoplasms (C00-C97)	7,045	31.1	193.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,459	19.7	122.7
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,487	6.6	40.9
4	Diabetes mellitus (E10-E14)	1,364	6.0	37.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,123	5.0	30.9
6	Cerebrovascular diseases (I60-I69)	957	4.2	26.3
7	Viral hepatitis (B15-B19)	499	2.2	13.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	471	2.1	13.0
9	Chronic lower respiratory diseases (J40-J47)	407	1.8	11.2
10	Septicemia (A40-A41)	382	1.7	10.5
11	Influenza and pneumonia (J09-J18)	316	1.4	8.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	261	1.2	7.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	217	1.0	6.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	207	0.9	5.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	0.5	3.3
...	All other causes (Residual)	3,304	14.6	90.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	24,868	100.0	1,330.9
1	Malignant neoplasms (C00-C97)	7,929	31.9	424.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,389	21.7	288.4
3	Diabetes mellitus (E10-E14)	1,737	7.0	93.0
4	Cerebrovascular diseases (I60-I69)	1,244	5.0	66.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	880	3.5	47.1
6	Chronic lower respiratory diseases (J40-J47)	786	3.2	42.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	607	2.4	32.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	598	2.4	32.0
9	Influenza and pneumonia (J09-J18)	456	1.8	24.4
10	Septicemia (A40-A41)	449	1.8	24.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	292	1.2	15.6
12	Alzheimer's disease (G30)	229	0.9	12.3
13	Viral hepatitis (B15-B19)	161	0.6	8.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	149	0.6	8.0
15	Parkinson's disease (G20-G21)	143	0.6	7.7
...	All other causes (Residual)	3,819	15.4	204.4
Hispanic, both sexes, 75-84 years				
...	All causes	33,614	100.0	3,552.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,128	24.2	858.9
2	Malignant neoplasms (C00-C97)	7,851	23.4	829.6
3	Cerebrovascular diseases (I60-I69)	2,178	6.5	230.2
4	Diabetes mellitus (E10-E14)	1,985	5.9	209.8
5	Chronic lower respiratory diseases (J40-J47)	1,487	4.4	157.1
6	Alzheimer's disease (G30)	1,185	3.5	125.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	837	2.5	88.4
8	Influenza and pneumonia (J09-J18)	833	2.5	88.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	701	2.1	74.1
10	Septicemia (A40-A41)	579	1.7	61.2
11	Parkinson's disease (G20-G21)	526	1.6	55.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	502	1.5	53.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	491	1.5	51.9
14	Pneumonitis due to solids and liquids (J69)	226	0.7	23.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	205	0.6	21.7
...	All other causes (Residual)	5,900	17.6	623.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	33,410	100.0	10,157.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,574	28.7	2,910.6
2	Malignant neoplasms (C00-C97)	4,205	12.6	1,278.4
3	Alzheimer's disease (G30)	2,291	6.9	696.5
4	Cerebrovascular diseases (I60-I69)	2,251	6.7	684.3
5	Chronic lower respiratory diseases (J40-J47)	1,511	4.5	459.4
6	Diabetes mellitus (E10-E14)	1,250	3.7	380.0
7	Influenza and pneumonia (J09-J18)	1,041	3.1	316.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	705	2.1	214.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	692	2.1	210.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	682	2.0	207.3
11	Septicemia (A40-A41)	450	1.3	136.8
12	Parkinson's disease (G20-G21)	423	1.3	128.6
13	Pneumonitis due to solids and liquids (J69)	379	1.1	115.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	174	0.5	52.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	156	0.5	47.4
...	All other causes (Residual)	7,626	22.8	2,318.4
Hispanic, male, all ages				
...	All causes	85,238	100.0	316.5
1	Malignant neoplasms (C00-C97)	17,849	20.9	66.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,329	20.3	64.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,255	9.7	30.7
4	Diabetes mellitus (E10-E14)	3,763	4.4	14.0
5	Cerebrovascular diseases (I60-I69)	3,636	4.3	13.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,422	4.0	12.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,292	2.7	8.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,277	2.7	8.5
9	Chronic lower respiratory diseases (J40-J47)	2,268	2.7	8.4
10	Influenza and pneumonia (J09-J18)	1,557	1.8	5.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,552	1.8	5.8
12	Certain conditions originating in the perinatal period (P00-P96)	1,373	1.6	5.1
13	Alzheimer's disease (G30)	1,193	1.4	4.4
14	Septicemia (A40-A41)	1,175	1.4	4.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	952	1.1	3.5
...	All other causes (Residual)	16,345	19.2	60.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	494	100.0	23.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	31.2	7.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	15.0	3.5
3	Malignant neoplasms (C00-C97)	56	11.3	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	6.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.6	*
6	Influenza and pneumonia (J09-J18)	11	2.2	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.8	*
8	Septicemia (A40-A41)	7	1.4	*
8	Cerebrovascular diseases (I60-I69)	7	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
12	Pneumonitis due to solids and liquids (J69)	3	0.6	*
14	Anemias (D50-D64)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	114	23.1	5.4
Hispanic, male, 5-14 years				
...	All causes	608	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	32.1	3.9
2	Malignant neoplasms (C00-C97)	115	18.9	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	6.9	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	5.8	0.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	4.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.0	*
7	Cerebrovascular diseases (I60-I69)	14	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	2.0	*
9	Septicemia (A40-A41)	9	1.5	*
9	Influenza and pneumonia (J09-J18)	9	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	9	1.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
13	Anemias (D50-D64)	2	0.3	*
13	Diabetes mellitus (E10-E14)	2	0.3	*
13	Diseases of appendix (K35-K38)	2	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	110	18.1	2.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,655	100.0	76.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,438	39.3	30.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	818	22.4	17.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	548	15.0	11.5
4	Malignant neoplasms (C00-C97)	228	6.2	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	2.7	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	1.1	0.9
7	Legal intervention (Y35,Y89.0)	29	0.8	0.6
8	Cerebrovascular diseases (I60-I69)	23	0.6	0.5
9	Influenza and pneumonia (J09-J18)	17	0.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.3	*
10	Diabetes mellitus (E10-E14)	12	0.3	*
10	Chronic lower respiratory diseases (J40-J47)	12	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.2	*
14	Septicemia (A40-A41)	7	0.2	*
15	Anemias (D50-D64)	6	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.2	*
...	All other causes (Residual)	351	9.6	7.3
Hispanic, male, 25-34 years				
...	All causes	4,578	100.0	100.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,751	38.2	38.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	672	14.7	14.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	520	11.4	11.4
4	Malignant neoplasms (C00-C97)	346	7.6	7.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	278	6.1	6.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	95	2.1	2.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	90	2.0	2.0
8	Cerebrovascular diseases (I60-I69)	59	1.3	1.3
9	Diabetes mellitus (E10-E14)	52	1.1	1.1
10	Legal intervention (Y35,Y89.0)	34	0.7	0.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	0.7	0.7
12	Influenza and pneumonia (J09-J18)	30	0.7	0.7
13	Septicemia (A40-A41)	26	0.6	0.6
14	Aortic aneurysm and dissection (I71)	17	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	16	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
...	All other causes (Residual)	543	11.9	11.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	5,593	100.0	142.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,400	25.0	35.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	706	12.6	18.0
3	Malignant neoplasms (C00-C97)	679	12.1	17.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	447	8.0	11.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	372	6.7	9.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	360	6.4	9.2
7	Cerebrovascular diseases (I60-I69)	191	3.4	4.9
8	Diabetes mellitus (E10-E14)	154	2.8	3.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	136	2.4	3.5
10	Influenza and pneumonia (J09-J18)	70	1.3	1.8
11	Septicemia (A40-A41)	57	1.0	1.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.0	1.4
13	Viral hepatitis (B15-B19)	48	0.9	1.2
14	Aortic aneurysm and dissection (I71)	39	0.7	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	0.6	0.8
...	All other causes (Residual)	845	15.1	21.6
Hispanic, male, 45-54 years				
...	All causes	10,169	100.0	342.7
1	Malignant neoplasms (C00-C97)	2,005	19.7	67.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,832	18.0	61.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,292	12.7	43.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,045	10.3	35.2
5	Diabetes mellitus (E10-E14)	444	4.4	15.0
6	Cerebrovascular diseases (I60-I69)	400	3.9	13.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	351	3.5	11.8
8	Viral hepatitis (B15-B19)	290	2.9	9.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	256	2.5	8.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	189	1.9	6.4
11	Septicemia (A40-A41)	145	1.4	4.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	1.4	4.7
13	Influenza and pneumonia (J09-J18)	104	1.0	3.5
14	Chronic lower respiratory diseases (J40-J47)	82	0.8	2.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	79	0.8	2.7
...	All other causes (Residual)	1,516	14.9	51.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	14,258	100.0	816.3
1	Malignant neoplasms (C00-C97)	3,927	27.5	224.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,101	21.7	177.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	1,107	7.8	63.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	856	6.0	49.0
5	Diabetes mellitus (E10-E14)	796	5.6	45.6
6	Cerebrovascular diseases (I60-I69)	582	4.1	33.3
7	Viral hepatitis (B15-B19)	371	2.6	21.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	280	2.0	16.0
9	Chronic lower respiratory diseases (J40-J47)	238	1.7	13.6
10	Septicemia (A40-A41)	226	1.6	12.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	215	1.5	12.3
12	Influenza and pneumonia (J09-J18)	192	1.3	11.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	165	1.2	9.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	135	0.9	7.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	0.6	5.2
...	All other causes (Residual)	1,977	13.9	113.2
Hispanic, male, 65-74 years				
...	All causes	14,102	100.0	1,679.1
1	Malignant neoplasms (C00-C97)	4,403	31.2	524.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,411	24.2	406.1
3	Diabetes mellitus (E10-E14)	922	6.5	109.8
4	Cerebrovascular diseases (I60-I69)	634	4.5	75.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	523	3.7	62.3
6	Chronic lower respiratory diseases (J40-J47)	460	3.3	54.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	395	2.8	47.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	329	2.3	39.2
9	Influenza and pneumonia (J09-J18)	248	1.8	29.5
10	Septicemia (A40-A41)	235	1.7	28.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	145	1.0	17.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	0.7	11.2
13	Alzheimer's disease (G30)	92	0.7	11.0
14	Parkinson's disease (G20-G21)	86	0.6	10.2
15	Viral hepatitis (B15-B19)	83	0.6	9.9
...	All other causes (Residual)	2,042	14.5	243.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	16,552	100.0	4,250.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,163	25.2	1,069.1
2	Malignant neoplasms (C00-C97)	4,159	25.1	1,068.1
3	Cerebrovascular diseases (I60-I69)	966	5.8	248.1
4	Diabetes mellitus (E10-E14)	921	5.6	236.5
5	Chronic lower respiratory diseases (J40-J47)	779	4.7	200.1
6	Alzheimer's disease (G30)	440	2.7	113.0
7	Influenza and pneumonia (J09-J18)	429	2.6	110.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	420	2.5	107.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	389	2.4	99.9
10	Parkinson's disease (G20-G21)	295	1.8	75.8
11	Septicemia (A40-A41)	277	1.7	71.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	230	1.4	59.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	209	1.3	53.7
14	Pneumonitis due to solids and liquids (J69)	121	0.7	31.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	98	0.6	25.2
...	All other causes (Residual)	2,656	16.0	682.1
Hispanic, male, 85 years and over				
...	All causes	12,560	100.0	10,799.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,660	29.1	3,147.0
2	Malignant neoplasms (C00-C97)	1,922	15.3	1,652.6
3	Cerebrovascular diseases (I60-I69)	750	6.0	644.9
4	Chronic lower respiratory diseases (J40-J47)	636	5.1	546.9
5	Alzheimer's disease (G30)	629	5.0	540.8
6	Diabetes mellitus (E10-E14)	460	3.7	395.5
7	Influenza and pneumonia (J09-J18)	430	3.4	369.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	298	2.4	256.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	295	2.3	253.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	245	2.0	210.7
11	Parkinson's disease (G20-G21)	210	1.7	180.6
12	Pneumonitis due to solids and liquids (J69)	189	1.5	162.5
13	Septicemia (A40-A41)	171	1.4	147.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	85	0.7	73.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	59	0.5	50.7
...	All other causes (Residual)	2,521	20.1	2,167.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	71,181	100.0	272.7
1	Malignant neoplasms (C00-C97)	16,298	22.9	62.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,266	20.0	54.7
3	Cerebrovascular diseases (I60-I69)	4,131	5.8	15.8
4	Diabetes mellitus (E10-E14)	3,600	5.1	13.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,192	4.5	12.2
6	Alzheimer's disease (G30)	2,579	3.6	9.9
7	Chronic lower respiratory diseases (J40-J47)	2,209	3.1	8.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,566	2.2	6.0
9	Influenza and pneumonia (J09-J18)	1,492	2.1	5.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,412	2.0	5.4
11	Septicemia (A40-A41)	1,129	1.6	4.3
12	Certain conditions originating in the perinatal period (P00-P96)	1,056	1.5	4.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,017	1.4	3.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	838	1.2	3.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	545	0.8	2.1
...	All other causes (Residual)	15,851	22.3	60.7
Hispanic, female, 1-4 years				
...	All causes	405	100.0	20.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	28.4	5.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	12.8	2.6
3	Malignant neoplasms (C00-C97)	43	10.6	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	7.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	4.0	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	2.7	*
6	Influenza and pneumonia (J09-J18)	11	2.7	*
8	Septicemia (A40-A41)	7	1.7	*
9	Cerebrovascular diseases (I60-I69)	4	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
12	Diabetes mellitus (E10-E14)	2	0.5	*
12	Meningitis (G00,G03)	2	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
15	Anemias (D50-D64)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Hernia (K40-K46)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
...	All other causes (Residual)	98	24.2	4.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	475	100.0	10.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	23.6	2.4
2	Malignant neoplasms (C00-C97)	104	21.9	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	7.4	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.6	0.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	3.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.7	*
7	Cerebrovascular diseases (I60-I69)	10	2.1	*
8	Influenza and pneumonia (J09-J18)	9	1.9	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.7	*
11	Septicemia (A40-A41)	7	1.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
13	Meningitis (G00,G03)	4	0.8	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Anemias (D50-D64)	2	0.4	*
15	Diabetes mellitus (E10-E14)	2	0.4	*
...	All other causes (Residual)	113	23.8	2.4
Hispanic, female, 15-24 years				
...	All causes	1,118	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	433	38.7	9.9
2	Malignant neoplasms (C00-C97)	142	12.7	3.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	127	11.4	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	8.7	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.7	0.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	26	2.3	0.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	2.3	0.6
8	Cerebrovascular diseases (I60-I69)	13	1.2	*
9	Diabetes mellitus (E10-E14)	11	1.0	*
10	Influenza and pneumonia (J09-J18)	9	0.8	*
11	Septicemia (A40-A41)	8	0.7	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.6	*
15	Anemias (D50-D64)	4	0.4	*
...	All other causes (Residual)	160	14.3	3.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,638	100.0	40.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	409	25.0	10.0
2	Malignant neoplasms (C00-C97)	333	20.3	8.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	112	6.8	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	6.7	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	78	4.8	1.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	56	3.4	1.4
7	Cerebrovascular diseases (I60-I69)	39	2.4	1.0
8	Diabetes mellitus (E10-E14)	34	2.1	0.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	26	1.6	0.6
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	1.5	0.6
11	Septicemia (A40-A41)	21	1.3	0.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.9	*
14	Influenza and pneumonia (J09-J18)	13	0.8	*
14	Chronic lower respiratory diseases (J40-J47)	13	0.8	*
...	All other causes (Residual)	338	20.6	8.3
Hispanic, female, 35-44 years				
...	All causes	2,745	100.0	73.6
1	Malignant neoplasms (C00-C97)	911	33.2	24.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	412	15.0	11.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	234	8.5	6.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	113	4.1	3.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	3.6	2.7
6	Cerebrovascular diseases (I60-I69)	97	3.5	2.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	3.0	2.2
8	Diabetes mellitus (E10-E14)	78	2.8	2.1
9	Pregnancy, childbirth and the puerperium (O00-O99)	54	2.0	1.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	48	1.7	1.3
11	Septicemia (A40-A41)	36	1.3	1.0
12	Influenza and pneumonia (J09-J18)	33	1.2	0.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	1.2	0.9
14	Chronic lower respiratory diseases (J40-J47)	19	0.7	*
15	Viral hepatitis (B15-B19)	18	0.7	*
...	All other causes (Residual)	479	17.4	12.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	5,609	100.0	192.0
1	Malignant neoplasms (C00-C97)	2,140	38.2	73.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	641	11.4	21.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	453	8.1	15.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	317	5.7	10.8
5	Cerebrovascular diseases (I60-I69)	258	4.6	8.8
6	Diabetes mellitus (E10-E14)	236	4.2	8.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	106	1.9	3.6
8	Septicemia (A40-A41)	87	1.6	3.0
9	Viral hepatitis (B15-B19)	84	1.5	2.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	1.5	2.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	80	1.4	2.7
12	Chronic lower respiratory diseases (J40-J47)	78	1.4	2.7
13	Influenza and pneumonia (J09-J18)	59	1.1	2.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	1.0	1.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	0.6	1.1
...	All other causes (Residual)	902	16.1	30.9
Hispanic, female, 55-64 years				
...	All causes	8,362	100.0	442.8
1	Malignant neoplasms (C00-C97)	3,118	37.3	165.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,358	16.2	71.9
3	Diabetes mellitus (E10-E14)	568	6.8	30.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	380	4.5	20.1
5	Cerebrovascular diseases (I60-I69)	375	4.5	19.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	267	3.2	14.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	191	2.3	10.1
8	Chronic lower respiratory diseases (J40-J47)	169	2.0	9.0
9	Septicemia (A40-A41)	156	1.9	8.3
10	Viral hepatitis (B15-B19)	128	1.5	6.8
11	Influenza and pneumonia (J09-J18)	124	1.5	6.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	82	1.0	4.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	0.6	2.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	42	0.5	2.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.4	1.9
...	All other causes (Residual)	1,323	15.8	70.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	10,766	100.0	1,046.7
1	Malignant neoplasms (C00-C97)	3,526	32.8	342.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,978	18.4	192.3
3	Diabetes mellitus (E10-E14)	815	7.6	79.2
4	Cerebrovascular diseases (I60-I69)	610	5.7	59.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	357	3.3	34.7
6	Chronic lower respiratory diseases (J40-J47)	326	3.0	31.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	278	2.6	27.0
8	Septicemia (A40-A41)	214	2.0	20.8
9	Influenza and pneumonia (J09-J18)	208	1.9	20.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	1.9	19.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	147	1.4	14.3
12	Alzheimer's disease (G30)	137	1.3	13.3
13	Viral hepatitis (B15-B19)	78	0.7	7.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.7	7.0
15	Parkinson's disease (G20-G21)	57	0.5	5.5
...	All other causes (Residual)	1,760	16.3	171.1
Hispanic, female, 75-84 years				
...	All causes	17,062	100.0	3,063.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,965	23.2	711.9
2	Malignant neoplasms (C00-C97)	3,692	21.6	662.9
3	Cerebrovascular diseases (I60-I69)	1,212	7.1	217.6
4	Diabetes mellitus (E10-E14)	1,064	6.2	191.0
5	Alzheimer's disease (G30)	745	4.4	133.8
6	Chronic lower respiratory diseases (J40-J47)	708	4.1	127.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	417	2.4	74.9
8	Influenza and pneumonia (J09-J18)	404	2.4	72.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	312	1.8	56.0
10	Septicemia (A40-A41)	302	1.8	54.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	282	1.7	50.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	272	1.6	48.8
13	Parkinson's disease (G20-G21)	231	1.4	41.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	107	0.6	19.2
15	Pneumonitis due to solids and liquids (J69)	105	0.6	18.9
...	All other causes (Residual)	3,244	19.0	582.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	20,850	100.0	9,805.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,914	28.4	2,781.3
2	Malignant neoplasms (C00-C97)	2,283	10.9	1,073.7
3	Alzheimer's disease (G30)	1,662	8.0	781.6
4	Cerebrovascular diseases (I60-I69)	1,501	7.2	705.9
5	Chronic lower respiratory diseases (J40-J47)	875	4.2	411.5
6	Diabetes mellitus (E10-E14)	790	3.8	371.5
7	Influenza and pneumonia (J09-J18)	611	2.9	287.3
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	460	2.2	216.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	394	1.9	185.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	387	1.9	182.0
11	Septicemia (A40-A41)	279	1.3	131.2
12	Parkinson's disease (G20-G21)	213	1.0	100.2
13	Pneumonitis due to solids and liquids (J69)	190	0.9	89.4
14	Chronic liver disease and cirrhosis (K70,K73-K74)	97	0.5	45.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	89	0.4	41.9
...	All other causes (Residual)	5,105	24.5	2,400.8
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,016,896	100.0	1,004.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	481,991	23.9	240.2
2	Malignant neoplasms (C00-C97)	462,499	22.9	230.4
3	Chronic lower respiratory diseases (J40-J47)	127,116	6.3	63.3
4	Cerebrovascular diseases (I60-I69)	100,154	5.0	49.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99,288	4.9	49.5
6	Alzheimer's disease (G30)	72,772	3.6	36.3
7	Diabetes mellitus (E10-E14)	50,443	2.5	25.1
8	Influenza and pneumonia (J09-J18)	40,460	2.0	20.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,727	1.7	16.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33,105	1.6	16.5
11	Septicemia (A40-A41)	26,592	1.3	13.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	25,720	1.3	12.8
13	Parkinson's disease (G20-G21)	21,201	1.1	10.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21,056	1.0	10.5
15	Pneumonitis due to solids and liquids (J69)	15,073	0.7	7.5
...	All other causes (Residual)	405,699	20.1	202.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,094	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	745	35.6	8.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	11.7	2.9
3	Malignant neoplasms (C00-C97)	202	9.6	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	6.5	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	3.5	0.9
6	Influenza and pneumonia (J09-J18)	46	2.2	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.5	0.4
8	Cerebrovascular diseases (I60-I69)	23	1.1	0.3
9	Septicemia (A40-A41)	22	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	11	0.5	*
10	Certain conditions originating in the perinatal period (P00-P96)	11	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
14	Anemias (D50-D64)	4	0.2	*
14	Diabetes mellitus (E10-E14)	4	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.2	*
...	All other causes (Residual)	522	24.9	6.2
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,667	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	798	29.9	3.5
2	Malignant neoplasms (C00-C97)	501	18.8	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	203	7.6	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	151	5.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	5.4	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	3.3	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	2.0	0.2
8	Influenza and pneumonia (J09-J18)	45	1.7	0.2
9	Cerebrovascular diseases (I60-I69)	44	1.6	0.2
10	Septicemia (A40-A41)	27	1.0	0.1
11	Chronic lower respiratory diseases (J40-J47)	25	0.9	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.6	*
13	Diabetes mellitus (E10-E14)	12	0.4	*
14	Meningitis (G00,G03)	9	0.3	*
15	Anemias (D50-D64)	7	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (Residual)	538	20.2	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	16,475	100.0	65.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,173	49.6	32.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,347	20.3	13.3
3	Malignant neoplasms (C00-C97)	858	5.2	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	706	4.3	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	480	2.9	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	254	1.5	1.0
7	Diabetes mellitus (E10-E14)	109	0.7	0.4
8	Cerebrovascular diseases (I60-I69)	97	0.6	0.4
9	Influenza and pneumonia (J09-J18)	85	0.5	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	73	0.4	0.3
11	Septicemia (A40-A41)	66	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	56	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.3	0.2
14	Legal intervention (Y35,Y89.0)	36	0.2	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.2	0.1
...	All other causes (Residual)	2,054	12.5	8.2
Non-Hispanic, white, both sexes, 25-34 years				
...	All causes	27,266	100.0	110.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,358	41.7	45.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,681	17.2	18.9
3	Malignant neoplasms (C00-C97)	2,120	7.8	8.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,822	6.7	7.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	984	3.6	4.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	366	1.3	1.5
7	Diabetes mellitus (E10-E14)	334	1.2	1.3
8	Cerebrovascular diseases (I60-I69)	278	1.0	1.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	269	1.0	1.1
10	Influenza and pneumonia (J09-J18)	216	0.8	0.9
11	Pregnancy, childbirth and the puerperium (O00-O99)	175	0.6	0.7
12	Septicemia (A40-A41)	170	0.6	0.7
13	Chronic lower respiratory diseases (J40-J47)	140	0.5	0.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	114	0.4	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	0.4	0.4
...	All other causes (Residual)	4,138	15.2	16.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-44 years				
...	All causes	43,940	100.0	179.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,963	24.9	44.7
2	Malignant neoplasms (C00-C97)	7,106	16.2	29.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,313	14.4	25.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,432	12.4	22.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,598	3.6	6.5
6	Diabetes mellitus (E10-E14)	1,042	2.4	4.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	872	2.0	3.6
8	Cerebrovascular diseases (I60-I69)	858	2.0	3.5
9	Chronic lower respiratory diseases (J40-J47)	479	1.1	2.0
10	Septicemia (A40-A41)	422	1.0	1.7
11	Influenza and pneumonia (J09-J18)	402	0.9	1.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	367	0.8	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	277	0.6	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	271	0.6	1.1
15	Viral hepatitis (B15-B19)	168	0.4	0.7
...	All other causes (Residual)	7,370	16.8	30.0
Non-Hispanic, white, both sexes, 45-54 years				
...	All causes	124,142	100.0	410.9
1	Malignant neoplasms (C00-C97)	33,672	27.1	111.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,905	19.3	79.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,552	12.5	51.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,753	6.2	25.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,380	5.1	21.1
6	Chronic lower respiratory diseases (J40-J47)	3,601	2.9	11.9
7	Diabetes mellitus (E10-E14)	3,377	2.7	11.2
8	Cerebrovascular diseases (I60-I69)	3,019	2.4	10.0
9	Septicemia (A40-A41)	1,446	1.2	4.8
10	Viral hepatitis (B15-B19)	1,369	1.1	4.5
11	Influenza and pneumonia (J09-J18)	1,203	1.0	4.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,010	0.8	3.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	977	0.8	3.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	905	0.7	3.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	649	0.5	2.1
...	All other causes (Residual)	19,324	15.6	64.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-64 years				
...	All causes	240,590	100.0	842.1
1	Malignant neoplasms (C00-C97)	84,947	35.3	297.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,756	21.1	177.6
3	Chronic lower respiratory diseases (J40-J47)	12,943	5.4	45.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,072	5.0	42.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	8,271	3.4	28.9
6	Diabetes mellitus (E10-E14)	7,930	3.3	27.8
7	Cerebrovascular diseases (I60-I69)	6,845	2.8	24.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,233	2.6	21.8
9	Septicemia (A40-A41)	3,318	1.4	11.6
10	Influenza and pneumonia (J09-J18)	2,746	1.1	9.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,672	1.1	9.4
12	Viral hepatitis (B15-B19)	2,599	1.1	9.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,772	0.7	6.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,187	0.5	4.2
15	Pneumonitis due to solids and liquids (J69)	912	0.4	3.2
...	All other causes (Residual)	35,387	14.7	123.9
Non-Hispanic, white, both sexes, 65-74 years				
...	All causes	339,392	100.0	1,810.1
1	Malignant neoplasms (C00-C97)	121,681	35.9	649.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,270	21.0	380.1
3	Chronic lower respiratory diseases (J40-J47)	29,881	8.8	159.4
4	Cerebrovascular diseases (I60-I69)	13,032	3.8	69.5
5	Diabetes mellitus (E10-E14)	11,127	3.3	59.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,589	2.5	45.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,336	1.6	28.5
8	Septicemia (A40-A41)	5,088	1.5	27.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	5,016	1.5	26.8
10	Influenza and pneumonia (J09-J18)	4,687	1.4	25.0
11	Alzheimer's disease (G30)	3,579	1.1	19.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,043	0.9	16.2
13	Parkinson's disease (G20-G21)	2,546	0.8	13.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,490	0.7	13.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,171	0.6	11.6
...	All other causes (Residual)	49,856	14.7	265.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-84 years				
...	All causes	511,339	100.0	4,792.4
1	Malignant neoplasms (C00-C97)	127,618	25.0	1,196.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	119,415	23.4	1,119.2
3	Chronic lower respiratory diseases (J40-J47)	43,522	8.5	407.9
4	Cerebrovascular diseases (I60-I69)	28,747	5.6	269.4
5	Alzheimer's disease (G30)	19,960	3.9	187.1
6	Diabetes mellitus (E10-E14)	13,543	2.6	126.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,346	2.4	115.7
8	Influenza and pneumonia (J09-J18)	10,457	2.0	98.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,984	2.0	93.6
10	Parkinson's disease (G20-G21)	8,960	1.8	84.0
11	Septicemia (A40-A41)	7,578	1.5	71.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,960	1.0	46.5
13	Pneumonitis due to solids and liquids (J69)	4,073	0.8	38.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,991	0.8	37.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,061	0.6	28.7
...	All other causes (Residual)	93,124	18.2	872.8
Non-Hispanic, white, both sexes, 85 years and over				
...	All causes	698,324	100.0	14,129.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	207,711	29.7	4,202.8
2	Malignant neoplasms (C00-C97)	83,751	12.0	1,694.6
3	Alzheimer's disease (G30)	48,444	6.9	980.2
4	Cerebrovascular diseases (I60-I69)	47,168	6.8	954.4
5	Chronic lower respiratory diseases (J40-J47)	36,446	5.2	737.4
6	Influenza and pneumonia (J09-J18)	20,504	2.9	414.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,126	2.6	366.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,644	2.0	276.1
9	Diabetes mellitus (E10-E14)	12,964	1.9	262.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,997	1.6	222.5
11	Parkinson's disease (G20-G21)	9,179	1.3	185.7
12	Septicemia (A40-A41)	8,384	1.2	169.6
13	Pneumonitis due to solids and liquids (J69)	7,873	1.1	159.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,441	0.6	89.9
15	Atherosclerosis (I70)	3,540	0.5	71.6
...	All other causes (Residual)	165,152	23.6	3,341.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	998,832	100.0	1,011.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250,116	25.0	253.2
2	Malignant neoplasms (C00-C97)	243,925	24.4	247.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60,388	6.0	61.1
4	Chronic lower respiratory diseases (J40-J47)	59,084	5.9	59.8
5	Cerebrovascular diseases (I60-I69)	39,786	4.0	40.3
6	Diabetes mellitus (E10-E14)	27,043	2.7	27.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,356	2.6	26.7
8	Alzheimer's disease (G30)	22,403	2.2	22.7
9	Influenza and pneumonia (J09-J18)	18,960	1.9	19.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,720	1.7	16.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	16,518	1.7	16.7
12	Parkinson's disease (G20-G21)	12,654	1.3	12.8
13	Septicemia (A40-A41)	12,446	1.2	12.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,263	0.8	8.4
15	Pneumonitis due to solids and liquids (J69)	8,172	0.8	8.3
...	All other causes (Residual)	175,998	17.6	178.2
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,202	100.0	27.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	442	36.8	10.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	132	11.0	3.1
3	Malignant neoplasms (C00-C97)	108	9.0	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	6.7	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	2.8	0.8
6	Influenza and pneumonia (J09-J18)	31	2.6	0.7
7	Cerebrovascular diseases (I60-I69)	17	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.2	*
9	Septicemia (A40-A41)	11	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Diabetes mellitus (E10-E14)	4	0.3	*
15	Meningococcal infection (A39)	3	0.2	*
...	All other causes (Residual)	302	25.1	7.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,548	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	494	31.9	4.3
2	Malignant neoplasms (C00-C97)	288	18.6	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	149	9.6	1.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	4.7	0.6
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	4.6	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	2.8	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	1.9	0.2
8	Cerebrovascular diseases (I60-I69)	20	1.3	0.2
8	Influenza and pneumonia (J09-J18)	20	1.3	0.2
10	Septicemia (A40-A41)	19	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	12	0.8	*
12	Diabetes mellitus (E10-E14)	8	0.5	*
13	Meningitis (G00,G03)	6	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
15	Anemias (D50-D64)	3	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
15	Pneumonitis due to solids and liquids (J69)	3	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.2	*
...	All other causes (Residual)	297	19.2	2.6
Non-Hispanic, white, male, 15-24 years				
...	All causes	11,826	100.0	92.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,050	51.2	47.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,723	23.0	21.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	500	4.2	3.9
4	Malignant neoplasms (C00-C97)	485	4.1	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	316	2.7	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	1.2	1.1
7	Diabetes mellitus (E10-E14)	65	0.5	0.5
8	Cerebrovascular diseases (I60-I69)	54	0.5	0.4
9	Influenza and pneumonia (J09-J18)	52	0.4	0.4
10	Chronic lower respiratory diseases (J40-J47)	37	0.3	0.3
11	Legal intervention (Y35,Y89.0)	33	0.3	0.3
12	Septicemia (A40-A41)	31	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.2	0.2
14	Pneumonitis due to solids and liquids (J69)	21	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.1	*
...	All other causes (Residual)	1,269	10.7	9.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-34 years				
...	All causes	18,606	100.0	149.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,208	44.1	65.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,729	20.0	29.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,254	6.7	10.0
4	Malignant neoplasms (C00-C97)	1,067	5.7	8.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	689	3.7	5.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	228	1.2	1.8
7	Diabetes mellitus (E10-E14)	197	1.1	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	157	0.8	1.3
9	Cerebrovascular diseases (I60-I69)	150	0.8	1.2
10	Influenza and pneumonia (J09-J18)	122	0.7	1.0
11	Septicemia (A40-A41)	98	0.5	0.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	91	0.5	0.7
13	Chronic lower respiratory diseases (J40-J47)	76	0.4	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	0.3	0.5
15	Legal intervention (Y35,Y89.0)	59	0.3	0.5
...	All other causes (Residual)	2,421	13.0	19.4
Non-Hispanic, white, male, 35-44 years				
...	All causes	27,336	100.0	221.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,493	27.4	60.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,386	16.0	35.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,142	15.2	33.6
4	Malignant neoplasms (C00-C97)	3,166	11.6	25.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	992	3.6	8.0
6	Diabetes mellitus (E10-E14)	636	2.3	5.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	589	2.2	4.8
8	Cerebrovascular diseases (I60-I69)	455	1.7	3.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	298	1.1	2.4
10	Septicemia (A40-A41)	216	0.8	1.8
11	Chronic lower respiratory diseases (J40-J47)	207	0.8	1.7
12	Influenza and pneumonia (J09-J18)	197	0.7	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	0.6	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	150	0.5	1.2
15	Aortic aneurysm and dissection (I71)	116	0.4	0.9
...	All other causes (Residual)	4,134	15.1	33.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-54 years				
...	All causes	76,207	100.0	508.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,285	22.7	115.2
2	Malignant neoplasms (C00-C97)	17,132	22.5	114.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,216	13.4	68.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,741	7.5	38.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,254	5.6	28.4
6	Diabetes mellitus (E10-E14)	2,161	2.8	14.4
7	Chronic lower respiratory diseases (J40-J47)	1,669	2.2	11.1
8	Cerebrovascular diseases (I60-I69)	1,620	2.1	10.8
9	Viral hepatitis (B15-B19)	958	1.3	6.4
10	Septicemia (A40-A41)	766	1.0	5.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	766	1.0	5.1
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	680	0.9	4.5
13	Influenza and pneumonia (J09-J18)	662	0.9	4.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	614	0.8	4.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	452	0.6	3.0
...	All other causes (Residual)	11,231	14.7	74.9
Non-Hispanic, white, male, 55-64 years				
...	All causes	147,634	100.0	1,058.0
1	Malignant neoplasms (C00-C97)	47,250	32.0	338.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,160	24.5	259.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,087	5.5	58.0
4	Chronic lower respiratory diseases (J40-J47)	6,654	4.5	47.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,801	3.9	41.6
6	Diabetes mellitus (E10-E14)	5,002	3.4	35.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,777	3.2	34.2
8	Cerebrovascular diseases (I60-I69)	3,982	2.7	28.5
9	Viral hepatitis (B15-B19)	1,873	1.3	13.4
10	Septicemia (A40-A41)	1,811	1.2	13.0
11	Influenza and pneumonia (J09-J18)	1,598	1.1	11.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,527	1.0	10.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,132	0.8	8.1
14	Aortic aneurysm and dissection (I71)	684	0.5	4.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	675	0.5	4.8
...	All other causes (Residual)	20,621	14.0	147.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-74 years				
...	All causes	193,452	100.0	2,175.8
1	Malignant neoplasms (C00-C97)	67,343	34.8	757.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,080	23.8	518.3
3	Chronic lower respiratory diseases (J40-J47)	14,797	7.6	166.4
4	Cerebrovascular diseases (I60-I69)	6,907	3.6	77.7
5	Diabetes mellitus (E10-E14)	6,574	3.4	73.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,436	2.8	61.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,128	1.6	35.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,938	1.5	33.0
9	Influenza and pneumonia (J09-J18)	2,633	1.4	29.6
10	Septicemia (A40-A41)	2,613	1.4	29.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,451	1.3	27.6
12	Parkinson's disease (G20-G21)	1,709	0.9	19.2
13	Alzheimer's disease (G30)	1,556	0.8	17.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,359	0.7	15.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,333	0.7	15.0
...	All other causes (Residual)	26,595	13.7	299.1
Non-Hispanic, white, male, 75-84 years				
...	All causes	258,273	100.0	5,599.2
1	Malignant neoplasms (C00-C97)	68,079	26.4	1,475.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	65,019	25.2	1,409.6
3	Chronic lower respiratory diseases (J40-J47)	20,788	8.0	450.7
4	Cerebrovascular diseases (I60-I69)	12,511	4.8	271.2
5	Alzheimer's disease (G30)	7,470	2.9	161.9
6	Diabetes mellitus (E10-E14)	7,126	2.8	154.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,500	2.5	140.9
8	Parkinson's disease (G20-G21)	5,743	2.2	124.5
9	Influenza and pneumonia (J09-J18)	5,365	2.1	116.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,324	2.1	115.4
11	Septicemia (A40-A41)	3,576	1.4	77.5
12	Pneumonitis due to solids and liquids (J69)	2,354	0.9	51.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,250	0.9	48.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,083	0.8	45.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,776	0.7	38.5
...	All other causes (Residual)	42,309	16.4	917.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85 years and over				
...	All causes	256,821	100.0	15,504.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	79,444	30.9	4,796.1
2	Malignant neoplasms (C00-C97)	38,986	15.2	2,353.6
3	Chronic lower respiratory diseases (J40-J47)	14,830	5.8	895.3
4	Cerebrovascular diseases (I60-I69)	14,046	5.5	848.0
5	Alzheimer's disease (G30)	13,032	5.1	786.7
6	Influenza and pneumonia (J09-J18)	8,240	3.2	497.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,130	2.8	430.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,052	2.4	365.4
9	Diabetes mellitus (E10-E14)	5,270	2.1	318.2
10	Parkinson's disease (G20-G21)	4,876	1.9	294.4
11	Pneumonitis due to solids and liquids (J69)	3,928	1.5	237.1
12	Septicemia (A40-A41)	3,267	1.3	197.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,116	1.2	188.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,984	0.8	119.8
15	Atherosclerosis (I70)	1,133	0.4	68.4
...	All other causes (Residual)	51,487	20.0	3,108.3
Non-Hispanic, white, female, all ages				
...	All causes	1,018,064	100.0	998.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	231,875	22.8	227.5
2	Malignant neoplasms (C00-C97)	218,574	21.5	214.4
3	Chronic lower respiratory diseases (J40-J47)	68,032	6.7	66.7
4	Cerebrovascular diseases (I60-I69)	60,368	5.9	59.2
5	Alzheimer's disease (G30)	50,369	4.9	49.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38,900	3.8	38.2
7	Diabetes mellitus (E10-E14)	23,400	2.3	23.0
8	Influenza and pneumonia (J09-J18)	21,500	2.1	21.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,385	1.6	16.1
10	Septicemia (A40-A41)	14,146	1.4	13.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,793	1.3	12.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,202	0.9	9.0
13	Parkinson's disease (G20-G21)	8,547	0.8	8.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,371	0.7	7.2
15	Pneumonitis due to solids and liquids (J69)	6,901	0.7	6.8
...	All other causes (Residual)	229,701	22.6	225.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	892	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	303	34.0	7.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	12.6	2.7
3	Malignant neoplasms (C00-C97)	94	10.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	6.2	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	4.4	0.9
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	2.0	*
7	Influenza and pneumonia (J09-J18)	15	1.7	*
8	Septicemia (A40-A41)	11	1.2	*
9	Cerebrovascular diseases (I60-I69)	6	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	3	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
14	Anemias (D50-D64)	2	0.2	*
14	Pneumonitis due to solids and liquids (J69)	2	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	218	24.4	5.3
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,119	100.0	10.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	304	27.2	2.8
2	Malignant neoplasms (C00-C97)	213	19.0	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	7.1	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	6.3	0.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	4.8	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	3.9	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	2.2	0.2
7	Influenza and pneumonia (J09-J18)	25	2.2	0.2
9	Cerebrovascular diseases (I60-I69)	24	2.1	0.2
10	Chronic lower respiratory diseases (J40-J47)	13	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	11	1.0	*
12	Septicemia (A40-A41)	8	0.7	*
13	Anemias (D50-D64)	4	0.4	*
13	Diabetes mellitus (E10-E14)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	235	21.0	2.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,649	100.0	37.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,123	45.7	17.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	624	13.4	5.1
3	Malignant neoplasms (C00-C97)	373	8.0	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	206	4.4	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	164	3.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	2.3	0.9
7	Pregnancy, childbirth and the puerperium (O00-O99)	73	1.6	0.6
8	Diabetes mellitus (E10-E14)	44	0.9	0.4
9	Cerebrovascular diseases (I60-I69)	43	0.9	0.3
10	Septicemia (A40-A41)	35	0.8	0.3
11	Influenza and pneumonia (J09-J18)	33	0.7	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	19	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.4	*
15	Pneumonitis due to solids and liquids (J69)	8	0.2	*
...	All other causes (Residual)	759	16.3	6.2
Non-Hispanic, white, female, 25-34 years				
...	All causes	8,660	100.0	70.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,150	36.4	25.7
2	Malignant neoplasms (C00-C97)	1,053	12.2	8.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	952	11.0	7.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	568	6.6	4.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	295	3.4	2.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	175	2.0	1.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	138	1.6	1.1
8	Diabetes mellitus (E10-E14)	137	1.6	1.1
9	Cerebrovascular diseases (I60-I69)	128	1.5	1.0
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	1.3	0.9
11	Influenza and pneumonia (J09-J18)	94	1.1	0.8
12	Septicemia (A40-A41)	72	0.8	0.6
13	Chronic lower respiratory diseases (J40-J47)	64	0.7	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.5	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.5	0.3
...	All other causes (Residual)	1,642	19.0	13.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-44 years				
...	All causes	16,604	100.0	136.0
1	Malignant neoplasms (C00-C97)	3,940	23.7	32.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,470	20.9	28.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,927	11.6	15.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,290	7.8	10.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	606	3.6	5.0
6	Diabetes mellitus (E10-E14)	406	2.4	3.3
7	Cerebrovascular diseases (I60-I69)	403	2.4	3.3
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	283	1.7	2.3
9	Chronic lower respiratory diseases (J40-J47)	272	1.6	2.2
10	Septicemia (A40-A41)	206	1.2	1.7
11	Influenza and pneumonia (J09-J18)	205	1.2	1.7
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	121	0.7	1.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	118	0.7	1.0
14	Pregnancy, childbirth and the puerperium (O00-O99)	102	0.6	0.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	69	0.4	0.6
...	All other causes (Residual)	3,186	19.2	26.1
Non-Hispanic, white, female, 45-54 years				
...	All causes	47,935	100.0	315.2
1	Malignant neoplasms (C00-C97)	16,540	34.5	108.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,620	13.8	43.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,336	11.1	35.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,126	4.4	14.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,012	4.2	13.2
6	Chronic lower respiratory diseases (J40-J47)	1,932	4.0	12.7
7	Cerebrovascular diseases (I60-I69)	1,399	2.9	9.2
8	Diabetes mellitus (E10-E14)	1,216	2.5	8.0
9	Septicemia (A40-A41)	680	1.4	4.5
10	Influenza and pneumonia (J09-J18)	541	1.1	3.6
11	Viral hepatitis (B15-B19)	411	0.9	2.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	396	0.8	2.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	297	0.6	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	235	0.5	1.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	218	0.5	1.4
...	All other causes (Residual)	7,976	16.6	52.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-64 years				
...	All causes	92,956	100.0	635.9
1	Malignant neoplasms (C00-C97)	37,697	40.6	257.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,596	15.7	99.9
3	Chronic lower respiratory diseases (J40-J47)	6,289	6.8	43.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,985	4.3	27.3
5	Diabetes mellitus (E10-E14)	2,928	3.1	20.0
6	Cerebrovascular diseases (I60-I69)	2,863	3.1	19.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,470	2.7	16.9
8	Septicemia (A40-A41)	1,507	1.6	10.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,456	1.6	10.0
10	Influenza and pneumonia (J09-J18)	1,148	1.2	7.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,145	1.2	7.8
12	Viral hepatitis (B15-B19)	726	0.8	5.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	640	0.7	4.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	512	0.6	3.5
15	Alzheimer's disease (G30)	392	0.4	2.7
...	All other causes (Residual)	14,602	15.7	99.9
Non-Hispanic, white, female, 65-74 years				
...	All causes	145,940	100.0	1,480.2
1	Malignant neoplasms (C00-C97)	54,338	37.2	551.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,190	17.3	255.5
3	Chronic lower respiratory diseases (J40-J47)	15,084	10.3	153.0
4	Cerebrovascular diseases (I60-I69)	6,125	4.2	62.1
5	Diabetes mellitus (E10-E14)	4,553	3.1	46.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,153	2.2	32.0
7	Septicemia (A40-A41)	2,475	1.7	25.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,398	1.6	24.3
9	Influenza and pneumonia (J09-J18)	2,054	1.4	20.8
10	Alzheimer's disease (G30)	2,023	1.4	20.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,888	1.3	19.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,131	0.8	11.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	838	0.6	8.5
14	Parkinson's disease (G20-G21)	837	0.6	8.5
15	Pneumonitis due to solids and liquids (J69)	648	0.4	6.6
...	All other causes (Residual)	23,205	15.9	235.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-84 years				
...	All causes	253,066	100.0	4,178.1
1	Malignant neoplasms (C00-C97)	59,539	23.5	983.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,396	21.5	898.1
3	Chronic lower respiratory diseases (J40-J47)	22,734	9.0	375.3
4	Cerebrovascular diseases (I60-I69)	16,236	6.4	268.1
5	Alzheimer's disease (G30)	12,490	4.9	206.2
6	Diabetes mellitus (E10-E14)	6,417	2.5	105.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,846	2.3	96.5
8	Influenza and pneumonia (J09-J18)	5,092	2.0	84.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,660	1.8	76.9
10	Septicemia (A40-A41)	4,002	1.6	66.1
11	Parkinson's disease (G20-G21)	3,217	1.3	53.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,877	1.1	47.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,741	0.7	28.7
14	Pneumonitis due to solids and liquids (J69)	1,719	0.7	28.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,430	0.6	23.6
...	All other causes (Residual)	50,670	20.0	836.5
Non-Hispanic, white, female, 85 years and over				
...	All causes	441,503	100.0	13,437.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	128,267	29.1	3,903.8
2	Malignant neoplasms (C00-C97)	44,765	10.1	1,362.4
3	Alzheimer's disease (G30)	35,412	8.0	1,077.7
4	Cerebrovascular diseases (I60-I69)	33,122	7.5	1,008.1
5	Chronic lower respiratory diseases (J40-J47)	21,616	4.9	657.9
6	Influenza and pneumonia (J09-J18)	12,264	2.8	373.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,996	2.5	334.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,881	1.8	239.9
9	Diabetes mellitus (E10-E14)	7,694	1.7	234.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,592	1.7	231.1
11	Septicemia (A40-A41)	5,117	1.2	155.7
12	Parkinson's disease (G20-G21)	4,303	1.0	131.0
13	Pneumonitis due to solids and liquids (J69)	3,945	0.9	120.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,457	0.6	74.8
15	Atherosclerosis (I70)	2,407	0.5	73.3
...	All other causes (Residual)	113,665	25.7	3,459.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	291,179	100.0	720.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,147	23.7	171.2
2	Malignant neoplasms (C00-C97)	66,565	22.9	164.8
3	Cerebrovascular diseases (I60-I69)	15,714	5.4	38.9
4	Diabetes mellitus (E10-E14)	12,836	4.4	31.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,451	4.3	30.8
6	Chronic lower respiratory diseases (J40-J47)	9,264	3.2	22.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,129	2.8	20.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,123	2.8	20.1
9	Septicemia (A40-A41)	5,918	2.0	14.7
10	Alzheimer's disease (G30)	5,392	1.9	13.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,140	1.8	12.7
12	Influenza and pneumonia (J09-J18)	4,923	1.7	12.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3,848	1.3	9.5
14	Certain conditions originating in the perinatal period (P00-P96)	3,767	1.3	9.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,739	0.9	6.8
...	All other causes (Residual)	57,223	19.7	141.7
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	989	100.0	40.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	271	27.4	11.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	13.0	5.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	10.5	4.2
4	Malignant neoplasms (C00-C97)	64	6.5	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	4.3	1.7
6	Chronic lower respiratory diseases (J40-J47)	30	3.0	1.2
7	Influenza and pneumonia (J09-J18)	20	2.0	0.8
8	Septicemia (A40-A41)	17	1.7	*
8	Cerebrovascular diseases (I60-I69)	17	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	17	1.7	*
11	Anemias (D50-D64)	7	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	252	25.5	10.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,155	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	346	30.0	5.6
2	Malignant neoplasms (C00-C97)	140	12.1	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	9.0	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	6.7	1.2
5	Chronic lower respiratory diseases (J40-J47)	74	6.4	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	3.6	0.7
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.5	0.7
8	Diabetes mellitus (E10-E14)	17	1.5	*
9	Influenza and pneumonia (J09-J18)	15	1.3	*
10	Cerebrovascular diseases (I60-I69)	14	1.2	*
11	Anemias (D50-D64)	13	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	11	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
14	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
15	Septicemia (A40-A41)	7	0.6	*
...	All other causes (Residual)	237	20.5	3.8
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	6,581	100.0	96.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,869	43.6	42.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,357	20.6	20.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	510	7.7	7.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	282	4.3	4.1
5	Malignant neoplasms (C00-C97)	244	3.7	3.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	87	1.3	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	1.3	1.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	67	1.0	1.0
9	Anemias (D50-D64)	65	1.0	1.0
10	Chronic lower respiratory diseases (J40-J47)	62	0.9	0.9
11	Diabetes mellitus (E10-E14)	56	0.9	0.8
12	Legal intervention (Y35,Y89.0)	44	0.7	0.6
13	Cerebrovascular diseases (I60-I69)	42	0.6	0.6
14	Septicemia (A40-A41)	33	0.5	0.5
15	Influenza and pneumonia (J09-J18)	32	0.5	0.5
...	All other causes (Residual)	748	11.4	11.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-34 years				
...	All causes	8,948	100.0	158.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,406	26.9	42.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,692	18.9	29.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	882	9.9	15.6
4	Malignant neoplasms (C00-C97)	641	7.2	11.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	546	6.1	9.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	417	4.7	7.4
7	Diabetes mellitus (E10-E14)	200	2.2	3.5
8	Pregnancy, childbirth and the puerperium (O00-O99)	133	1.5	2.4
9	Cerebrovascular diseases (I60-I69)	121	1.4	2.1
10	Anemias (D50-D64)	113	1.3	2.0
11	Chronic lower respiratory diseases (J40-J47)	109	1.2	1.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	91	1.0	1.6
13	Septicemia (A40-A41)	89	1.0	1.6
14	Influenza and pneumonia (J09-J18)	80	0.9	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	0.6	1.0
...	All other causes (Residual)	1,370	15.3	24.3
Non-Hispanic, black, both sexes, 35-44 years				
...	All causes	13,502	100.0	259.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,765	20.5	53.1
2	Malignant neoplasms (C00-C97)	1,916	14.2	36.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,658	12.3	31.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,279	9.5	24.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	744	5.5	14.3
6	Diabetes mellitus (E10-E14)	505	3.7	9.7
7	Cerebrovascular diseases (I60-I69)	464	3.4	8.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	464	3.4	8.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	256	1.9	4.9
10	Septicemia (A40-A41)	212	1.6	4.1
11	Chronic lower respiratory diseases (J40-J47)	194	1.4	3.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	190	1.4	3.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	158	1.2	3.0
14	Influenza and pneumonia (J09-J18)	151	1.1	2.9
15	Anemias (D50-D64)	143	1.1	2.7
...	All other causes (Residual)	2,403	17.8	46.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-54 years				
...	All causes	32,522	100.0	592.0
1	Malignant neoplasms (C00-C97)	8,098	24.9	147.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,682	23.6	139.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,335	7.2	42.5
4	Cerebrovascular diseases (I60-I69)	1,647	5.1	30.0
5	Diabetes mellitus (E10-E14)	1,395	4.3	25.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,282	3.9	23.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	753	2.3	13.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	743	2.3	13.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	710	2.2	12.9
10	Chronic lower respiratory diseases (J40-J47)	681	2.1	12.4
11	Septicemia (A40-A41)	582	1.8	10.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	532	1.6	9.7
13	Influenza and pneumonia (J09-J18)	388	1.2	7.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	367	1.1	6.7
15	Viral hepatitis (B15-B19)	238	0.7	4.3
...	All other causes (Residual)	5,089	15.6	92.6
Non-Hispanic, black, both sexes, 55-64 years				
...	All causes	54,983	100.0	1,285.3
1	Malignant neoplasms (C00-C97)	17,030	31.0	398.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,749	25.0	321.4
3	Cerebrovascular diseases (I60-I69)	2,690	4.9	62.9
4	Diabetes mellitus (E10-E14)	2,662	4.8	62.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,066	3.8	48.3
6	Chronic lower respiratory diseases (J40-J47)	1,554	2.8	36.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,471	2.7	34.4
8	Septicemia (A40-A41)	1,129	2.1	26.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,094	2.0	25.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	949	1.7	22.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	901	1.6	21.1
12	Influenza and pneumonia (J09-J18)	696	1.3	16.3
13	Viral hepatitis (B15-B19)	631	1.1	14.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	341	0.6	8.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	207	0.4	4.8
...	All other causes (Residual)	7,813	14.2	182.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-74 years				
...	All causes	54,481	100.0	2,470.1
1	Malignant neoplasms (C00-C97)	17,362	31.9	787.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,549	24.9	614.3
3	Diabetes mellitus (E10-E14)	3,084	5.7	139.8
4	Cerebrovascular diseases (I60-I69)	3,034	5.6	137.6
5	Chronic lower respiratory diseases (J40-J47)	2,359	4.3	107.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,767	3.2	80.1
7	Septicemia (A40-A41)	1,279	2.3	58.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	974	1.8	44.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	931	1.7	42.2
10	Influenza and pneumonia (J09-J18)	856	1.6	38.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	475	0.9	21.5
12	Alzheimer's disease (G30)	404	0.7	18.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	294	0.5	13.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	245	0.4	11.1
15	Pneumonitis due to solids and liquids (J69)	229	0.4	10.4
...	All other causes (Residual)	7,639	14.0	346.3
Non-Hispanic, black, both sexes, 75-84 years				
...	All causes	57,565	100.0	5,261.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,768	25.7	1,349.8
2	Malignant neoplasms (C00-C97)	13,962	24.3	1,276.1
3	Cerebrovascular diseases (I60-I69)	3,868	6.7	353.5
4	Diabetes mellitus (E10-E14)	2,945	5.1	269.2
5	Chronic lower respiratory diseases (J40-J47)	2,469	4.3	225.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,027	3.5	185.3
7	Alzheimer's disease (G30)	1,682	2.9	153.7
8	Septicemia (A40-A41)	1,387	2.4	126.8
9	Influenza and pneumonia (J09-J18)	1,171	2.0	107.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,164	2.0	106.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	800	1.4	73.1
12	Pneumonitis due to solids and liquids (J69)	418	0.7	38.2
13	Parkinson's disease (G20-G21)	390	0.7	35.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	314	0.5	28.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	179	0.3	16.4
...	All other causes (Residual)	10,021	17.4	915.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85 years and over				
...	All causes	53,657	100.0	12,812.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,279	28.5	3,648.3
2	Malignant neoplasms (C00-C97)	7,097	13.2	1,694.6
3	Cerebrovascular diseases (I60-I69)	3,781	7.0	902.8
4	Alzheimer's disease (G30)	3,236	6.0	772.7
5	Diabetes mellitus (E10-E14)	1,971	3.7	470.6
6	Chronic lower respiratory diseases (J40-J47)	1,725	3.2	411.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,708	3.2	407.8
8	Influenza and pneumonia (J09-J18)	1,460	2.7	348.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,399	2.6	334.1
10	Septicemia (A40-A41)	1,109	2.1	264.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	571	1.1	136.3
12	Pneumonitis due to solids and liquids (J69)	512	1.0	122.3
13	Parkinson's disease (G20-G21)	297	0.6	70.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	244	0.5	58.3
15	Atherosclerosis (I70)	198	0.4	47.3
...	All other causes (Residual)	13,070	24.4	3,120.8
Non-Hispanic, black, male, all ages				
...	All causes	148,344	100.0	768.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,586	24.0	184.3
2	Malignant neoplasms (C00-C97)	34,005	22.9	176.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,402	5.7	43.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,039	4.7	36.5
5	Cerebrovascular diseases (I60-I69)	6,975	4.7	36.1
6	Diabetes mellitus (E10-E14)	6,072	4.1	31.5
7	Chronic lower respiratory diseases (J40-J47)	4,826	3.3	25.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,776	2.5	19.6
9	Septicemia (A40-A41)	2,677	1.8	13.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,491	1.7	12.9
11	Influenza and pneumonia (J09-J18)	2,371	1.6	12.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,288	1.5	11.9
13	Certain conditions originating in the perinatal period (P00-P96)	2,131	1.4	11.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,861	1.3	9.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,770	1.2	9.2
...	All other causes (Residual)	26,074	17.6	135.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	563	100.0	45.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	164	29.1	13.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	13.9	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	9.1	4.1
4	Malignant neoplasms (C00-C97)	39	6.9	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.0	2.2
6	Chronic lower respiratory diseases (J40-J47)	24	4.3	1.9
7	Influenza and pneumonia (J09-J18)	12	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	11	2.0	*
9	Septicemia (A40-A41)	8	1.4	*
9	Cerebrovascular diseases (I60-I69)	8	1.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.9	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
14	Anemias (D50-D64)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	124	22.0	9.9
Non-Hispanic, black, male, 5-14 years				
...	All causes	707	100.0	22.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	32.2	7.3
2	Malignant neoplasms (C00-C97)	80	11.3	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	9.9	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	6.5	1.5
5	Chronic lower respiratory diseases (J40-J47)	40	5.7	1.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	4.7	1.1
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.3	0.7
8	Anemias (D50-D64)	8	1.1	*
8	Diabetes mellitus (E10-E14)	8	1.1	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	8	1.1	*
11	Influenza and pneumonia (J09-J18)	7	1.0	*
12	Cerebrovascular diseases (I60-I69)	6	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
14	Septicemia (A40-A41)	4	0.6	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
...	All other causes (Residual)	138	19.5	4.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,028	100.0	146.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,586	51.4	75.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	986	19.6	28.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	411	8.2	12.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	188	3.7	5.5
5	Malignant neoplasms (C00-C97)	142	2.8	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.2	1.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	1.1	1.6
8	Chronic lower respiratory diseases (J40-J47)	46	0.9	1.3
9	Legal intervention (Y35,Y89.0)	42	0.8	1.2
10	Anemias (D50-D64)	34	0.7	1.0
11	Diabetes mellitus (E10-E14)	26	0.5	0.8
12	Cerebrovascular diseases (I60-I69)	23	0.5	0.7
13	Influenza and pneumonia (J09-J18)	20	0.4	0.6
14	Septicemia (A40-A41)	13	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.1	*
...	All other causes (Residual)	389	7.7	11.3
Non-Hispanic, black, male, 25-34 years				
...	All causes	6,155	100.0	226.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,153	35.0	79.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,246	20.2	45.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	553	9.0	20.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	467	7.6	17.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	264	4.3	9.7
6	Malignant neoplasms (C00-C97)	263	4.3	9.7
7	Diabetes mellitus (E10-E14)	100	1.6	3.7
8	Cerebrovascular diseases (I60-I69)	70	1.1	2.6
9	Chronic lower respiratory diseases (J40-J47)	68	1.1	2.5
10	Anemias (D50-D64)	59	1.0	2.2
11	Septicemia (A40-A41)	48	0.8	1.8
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	0.7	1.5
13	Influenza and pneumonia (J09-J18)	41	0.7	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.7	1.5
15	Legal intervention (Y35,Y89.0)	40	0.6	1.5
...	All other causes (Residual)	700	11.4	25.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-44 years				
...	All causes	7,905	100.0	322.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,776	22.5	72.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,163	14.7	47.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,107	14.0	45.2
4	Malignant neoplasms (C00-C97)	688	8.7	28.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	385	4.9	15.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	356	4.5	14.5
7	Diabetes mellitus (E10-E14)	300	3.8	12.2
8	Cerebrovascular diseases (I60-I69)	255	3.2	10.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	1.8	5.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	116	1.5	4.7
11	Chronic lower respiratory diseases (J40-J47)	99	1.3	4.0
12	Septicemia (A40-A41)	97	1.2	4.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	87	1.1	3.5
14	Influenza and pneumonia (J09-J18)	73	0.9	3.0
15	Anemias (D50-D64)	64	0.8	2.6
...	All other causes (Residual)	1,198	15.2	48.9
Non-Hispanic, black, male, 45-54 years				
...	All causes	18,577	100.0	720.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,880	26.3	189.1
2	Malignant neoplasms (C00-C97)	3,867	20.8	149.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,581	8.5	61.3
4	Cerebrovascular diseases (I60-I69)	890	4.8	34.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	829	4.5	32.1
6	Diabetes mellitus (E10-E14)	806	4.3	31.2
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	589	3.2	22.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	431	2.3	16.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	406	2.2	15.7
10	Chronic lower respiratory diseases (J40-J47)	335	1.8	13.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	306	1.6	11.9
12	Septicemia (A40-A41)	301	1.6	11.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	281	1.5	10.9
14	Influenza and pneumonia (J09-J18)	218	1.2	8.4
15	Viral hepatitis (B15-B19)	167	0.9	6.5
...	All other causes (Residual)	2,690	14.5	104.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-64 years				
...	All causes	32,243	100.0	1,657.2
1	Malignant neoplasms (C00-C97)	9,354	29.0	480.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,702	27.0	447.3
3	Cerebrovascular diseases (I60-I69)	1,521	4.7	78.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,482	4.6	76.2
5	Diabetes mellitus (E10-E14)	1,477	4.6	75.9
6	Chronic lower respiratory diseases (J40-J47)	815	2.5	41.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	807	2.5	41.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	772	2.4	39.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	668	2.1	34.3
10	Septicemia (A40-A41)	579	1.8	29.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	532	1.6	27.3
12	Viral hepatitis (B15-B19)	421	1.3	21.6
13	Influenza and pneumonia (J09-J18)	409	1.3	21.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	288	0.9	14.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	165	0.5	8.5
...	All other causes (Residual)	4,251	13.2	218.5
Non-Hispanic, black, male, 65-74 years				
...	All causes	29,731	100.0	3,146.5
1	Malignant neoplasms (C00-C97)	9,392	31.6	994.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,759	26.1	821.2
3	Cerebrovascular diseases (I60-I69)	1,598	5.4	169.1
4	Diabetes mellitus (E10-E14)	1,593	5.4	168.6
5	Chronic lower respiratory diseases (J40-J47)	1,337	4.5	141.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	893	3.0	94.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	632	2.1	66.9
8	Septicemia (A40-A41)	609	2.0	64.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	482	1.6	51.0
10	Influenza and pneumonia (J09-J18)	463	1.6	49.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	320	1.1	33.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	225	0.8	23.8
13	Alzheimer's disease (G30)	162	0.5	17.1
14	Pneumonitis due to solids and liquids (J69)	152	0.5	16.1
15	Parkinson's disease (G20-G21)	124	0.4	13.1
...	All other causes (Residual)	3,990	13.4	422.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-84 years				
...	All causes	26,856	100.0	6,550.1
1	Malignant neoplasms (C00-C97)	7,073	26.3	1,725.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,993	26.0	1,705.6
3	Cerebrovascular diseases (I60-I69)	1,620	6.0	395.1
4	Chronic lower respiratory diseases (J40-J47)	1,290	4.8	314.6
5	Diabetes mellitus (E10-E14)	1,229	4.6	299.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	899	3.3	219.3
7	Alzheimer's disease (G30)	633	2.4	154.4
8	Septicemia (A40-A41)	585	2.2	142.7
9	Influenza and pneumonia (J09-J18)	568	2.1	138.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	486	1.8	118.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	459	1.7	111.9
12	Parkinson's disease (G20-G21)	248	0.9	60.5
13	Pneumonitis due to solids and liquids (J69)	236	0.9	57.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	148	0.6	36.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	98	0.4	23.9
...	All other causes (Residual)	4,291	16.0	1,046.6
Non-Hispanic, black, male, 85 years and over				
...	All causes	16,766	100.0	14,006.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,628	27.6	3,866.3
2	Malignant neoplasms (C00-C97)	3,101	18.5	2,590.6
3	Cerebrovascular diseases (I60-I69)	965	5.8	806.2
4	Chronic lower respiratory diseases (J40-J47)	768	4.6	641.6
5	Alzheimer's disease (G30)	704	4.2	588.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	566	3.4	472.8
7	Diabetes mellitus (E10-E14)	532	3.2	444.4
8	Influenza and pneumonia (J09-J18)	529	3.2	441.9
9	Septicemia (A40-A41)	383	2.3	320.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	365	2.2	304.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	1.4	196.3
12	Pneumonitis due to solids and liquids (J69)	205	1.2	171.3
13	Parkinson's disease (G20-G21)	144	0.9	120.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	93	0.6	77.7
15	Atherosclerosis (I70)	63	0.4	52.6
...	All other causes (Residual)	3,485	20.8	2,911.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	142,835	100.0	677.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,561	23.5	159.2
2	Malignant neoplasms (C00-C97)	32,560	22.8	154.4
3	Cerebrovascular diseases (I60-I69)	8,739	6.1	41.4
4	Diabetes mellitus (E10-E14)	6,764	4.7	32.1
5	Chronic lower respiratory diseases (J40-J47)	4,438	3.1	21.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,353	3.0	20.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,049	2.8	19.2
8	Alzheimer's disease (G30)	3,861	2.7	18.3
9	Septicemia (A40-A41)	3,241	2.3	15.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,852	2.0	13.5
11	Influenza and pneumonia (J09-J18)	2,552	1.8	12.1
12	Certain conditions originating in the perinatal period (P00-P96)	1,636	1.1	7.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,357	1.0	6.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,084	0.8	5.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	969	0.7	4.6
...	All other causes (Residual)	30,819	21.6	146.1
Non-Hispanic, black, female, 1-4 years				
...	All causes	426	100.0	35.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	25.1	8.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	12.4	4.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	12.0	4.2
4	Malignant neoplasms (C00-C97)	25	5.9	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.5	*
6	Septicemia (A40-A41)	9	2.1	*
6	Cerebrovascular diseases (I60-I69)	9	2.1	*
8	Influenza and pneumonia (J09-J18)	8	1.9	*
9	Anemias (D50-D64)	6	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Meningitis (G00,G03)	2	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
...	All other causes (Residual)	119	27.9	9.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	448	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	26.3	3.9
2	Malignant neoplasms (C00-C97)	60	13.4	2.0
3	Chronic lower respiratory diseases (J40-J47)	34	7.6	1.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	7.6	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	6.9	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	4.0	*
7	Diabetes mellitus (E10-E14)	9	2.0	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	2.0	*
9	Cerebrovascular diseases (I60-I69)	8	1.8	*
9	Influenza and pneumonia (J09-J18)	8	1.8	*
11	Anemias (D50-D64)	5	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
14	Septicemia (A40-A41)	3	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
...	All other causes (Residual)	99	22.1	3.3
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,553	100.0	46.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	23.9	11.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	283	18.2	8.4
3	Malignant neoplasms (C00-C97)	102	6.6	3.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	6.4	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	6.1	2.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	67	4.3	2.0
7	Anemias (D50-D64)	31	2.0	0.9
8	Diabetes mellitus (E10-E14)	30	1.9	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	1.9	0.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.7	0.8
11	Septicemia (A40-A41)	20	1.3	0.6
12	Cerebrovascular diseases (I60-I69)	19	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.2	*
14	Chronic lower respiratory diseases (J40-J47)	16	1.0	*
15	Influenza and pneumonia (J09-J18)	12	0.8	*
...	All other causes (Residual)	335	21.6	9.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-34 years				
...	All causes	2,793	100.0	95.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	446	16.0	15.2
2	Malignant neoplasms (C00-C97)	378	13.5	12.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	329	11.8	11.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	253	9.1	8.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	153	5.5	5.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	133	4.8	4.5
7	Diabetes mellitus (E10-E14)	100	3.6	3.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	79	2.8	2.7
9	Anemias (D50-D64)	54	1.9	1.8
10	Cerebrovascular diseases (I60-I69)	51	1.8	1.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.8	1.7
12	Septicemia (A40-A41)	41	1.5	1.4
12	Chronic lower respiratory diseases (J40-J47)	41	1.5	1.4
14	Influenza and pneumonia (J09-J18)	39	1.4	1.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	24	0.9	0.8
...	All other causes (Residual)	622	22.3	21.2
Non-Hispanic, black, female, 35-44 years				
...	All causes	5,597	100.0	203.2
1	Malignant neoplasms (C00-C97)	1,228	21.9	44.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	989	17.7	35.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	495	8.8	18.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	359	6.4	13.0
5	Cerebrovascular diseases (I60-I69)	209	3.7	7.6
6	Diabetes mellitus (E10-E14)	205	3.7	7.4
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	172	3.1	6.2
8	Septicemia (A40-A41)	115	2.1	4.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	2.1	4.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	108	1.9	3.9
11	Chronic lower respiratory diseases (J40-J47)	95	1.7	3.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	95	1.7	3.4
13	Anemias (D50-D64)	79	1.4	2.9
14	Influenza and pneumonia (J09-J18)	78	1.4	2.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	74	1.3	2.7
...	All other causes (Residual)	1,181	21.1	42.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-54 years				
...	All causes	13,945	100.0	478.6
1	Malignant neoplasms (C00-C97)	4,231	30.3	145.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,802	20.1	96.2
3	Cerebrovascular diseases (I60-I69)	757	5.4	26.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	754	5.4	25.9
5	Diabetes mellitus (E10-E14)	589	4.2	20.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	453	3.2	15.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	347	2.5	11.9
8	Chronic lower respiratory diseases (J40-J47)	346	2.5	11.9
9	Septicemia (A40-A41)	281	2.0	9.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	279	2.0	9.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	226	1.6	7.8
12	Influenza and pneumonia (J09-J18)	170	1.2	5.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	1.1	5.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	0.6	3.0
15	Anemias (D50-D64)	73	0.5	2.5
...	All other causes (Residual)	2,397	17.2	82.3
Non-Hispanic, black, female, 55-64 years				
...	All causes	22,740	100.0	975.0
1	Malignant neoplasms (C00-C97)	7,676	33.8	329.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,047	22.2	216.4
3	Diabetes mellitus (E10-E14)	1,185	5.2	50.8
4	Cerebrovascular diseases (I60-I69)	1,169	5.1	50.1
5	Chronic lower respiratory diseases (J40-J47)	739	3.2	31.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	664	2.9	28.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	584	2.6	25.0
8	Septicemia (A40-A41)	550	2.4	23.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	369	1.6	15.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	322	1.4	13.8
11	Influenza and pneumonia (J09-J18)	287	1.3	12.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	281	1.2	12.0
13	Viral hepatitis (B15-B19)	210	0.9	9.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	91	0.4	3.9
15	Anemias (D50-D64)	82	0.4	3.5
...	All other causes (Residual)	3,484	15.3	149.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-74 years				
...	All causes	24,750	100.0	1,963.1
1	Malignant neoplasms (C00-C97)	7,970	32.2	632.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,790	23.4	459.2
3	Diabetes mellitus (E10-E14)	1,491	6.0	118.3
4	Cerebrovascular diseases (I60-I69)	1,436	5.8	113.9
5	Chronic lower respiratory diseases (J40-J47)	1,022	4.1	81.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	874	3.5	69.3
7	Septicemia (A40-A41)	670	2.7	53.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	449	1.8	35.6
9	Influenza and pneumonia (J09-J18)	393	1.6	31.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	342	1.4	27.1
11	Alzheimer's disease (G30)	242	1.0	19.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	155	0.6	12.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	125	0.5	9.9
14	Anemias (D50-D64)	82	0.3	6.5
15	Aortic aneurysm and dissection (I71)	79	0.3	6.3
...	All other causes (Residual)	3,630	14.7	287.9
Non-Hispanic, black, female, 75-84 years				
...	All causes	30,709	100.0	4,489.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,775	25.3	1,136.6
2	Malignant neoplasms (C00-C97)	6,889	22.4	1,007.0
3	Cerebrovascular diseases (I60-I69)	2,248	7.3	328.6
4	Diabetes mellitus (E10-E14)	1,716	5.6	250.8
5	Chronic lower respiratory diseases (J40-J47)	1,179	3.8	172.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,128	3.7	164.9
7	Alzheimer's disease (G30)	1,049	3.4	153.3
8	Septicemia (A40-A41)	802	2.6	117.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	678	2.2	99.1
10	Influenza and pneumonia (J09-J18)	603	2.0	88.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	1.1	49.8
12	Pneumonitis due to solids and liquids (J69)	182	0.6	26.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	166	0.5	24.3
14	Parkinson's disease (G20-G21)	142	0.5	20.8
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	93	0.3	13.6
...	All other causes (Residual)	5,718	18.6	835.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85 years and over				
...	All causes	36,891	100.0	12,334.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,651	28.9	3,561.1
2	Malignant neoplasms (C00-C97)	3,996	10.8	1,336.0
3	Cerebrovascular diseases (I60-I69)	2,816	7.6	941.5
4	Alzheimer's disease (G30)	2,532	6.9	846.5
5	Diabetes mellitus (E10-E14)	1,439	3.9	481.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,142	3.1	381.8
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,034	2.8	345.7
8	Chronic lower respiratory diseases (J40-J47)	957	2.6	320.0
9	Influenza and pneumonia (J09-J18)	931	2.5	311.3
10	Septicemia (A40-A41)	726	2.0	242.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	336	0.9	112.3
12	Pneumonitis due to solids and liquids (J69)	307	0.8	102.6
13	Parkinson's disease (G20-G21)	153	0.4	51.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	151	0.4	50.5
15	Atherosclerosis (I70)	135	0.4	45.1
...	All other causes (Residual)	9,585	26.0	3,204.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012