

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	5,735	100.0	551.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,466	25.6	141.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	916	16.0	88.2
3	Newborn affected by maternal complications of pregnancy (P01)	299	5.2	28.8
4	Sudden infant death syndrome (R95)	288	5.0	27.7
5	Newborn affected by complications of placenta, cord and membranes (P02)	232	4.0	22.3
6	Accidents (unintentional injuries) (V01-X59)	189	3.3	18.2
7	Respiratory distress of newborn (P22)	166	2.9	16.0
8	Bacterial sepsis of newborn (P36)	156	2.7	15.0
9	Neonatal hemorrhage (P50-P52,P54)	120	2.1	11.5
10	Diseases of the circulatory system (I00-I99)	108	1.9	10.4
11	Necrotizing enterocolitis of newborn (P77)	90	1.6	8.7
12	Atelectasis (P28.0-P28.1)	84	1.5	8.1
13	Assault (homicide) (*U01,X85-Y09)	81	1.4	7.8
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	68	1.2	6.5
15	Septicemia (A40-A41)	67	1.2	6.4
...	All other causes (Residual)	1,405	24.5	135.2
Infants, Hispanic, male				
...	All causes	3,179	100.0	598.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	761	23.9	143.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	514	16.2	96.8
3	Sudden infant death syndrome (R95)	169	5.3	31.8
4	Newborn affected by maternal complications of pregnancy (P01)	159	5.0	30.0
5	Newborn affected by complications of placenta, cord and membranes (P02)	129	4.1	24.3
6	Accidents (unintentional injuries) (V01-X59)	103	3.2	19.4
7	Respiratory distress of newborn (P22)	96	3.0	18.1
8	Bacterial sepsis of newborn (P36)	86	2.7	16.2
9	Neonatal hemorrhage (P50-P52,P54)	71	2.2	13.4
10	Necrotizing enterocolitis of newborn (P77)	59	1.9	11.1
11	Diseases of the circulatory system (I00-I99)	58	1.8	10.9
12	Assault (homicide) (*U01,X85-Y09)	50	1.6	9.4
13	Atelectasis (P28.0-P28.1)	49	1.5	9.2
14	Septicemia (A40-A41)	37	1.2	7.0
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	37	1.2	7.0
...	All other causes (Residual)	801	25.2	150.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,556	100.0	502.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	705	27.6	138.7
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	402	15.7	79.1
3	Newborn affected by maternal complications of pregnancy (P01)	140	5.5	27.5
4	Sudden infant death syndrome (R95)	119	4.7	23.4
5	Newborn affected by complications of placenta, cord and membranes (P02)	103	4.0	20.3
6	Accidents (unintentional injuries) (V01-X59)	86	3.4	16.9
7	Respiratory distress of newborn (P22)	70	2.7	13.8
7	Bacterial sepsis of newborn (P36)	70	2.7	13.8
9	Diseases of the circulatory system (I00-I99)	50	2.0	9.8
10	Neonatal hemorrhage (P50-P52,P54)	49	1.9	9.6
11	Atelectasis (P28.0-P28.1)	35	1.4	6.9
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	31	1.2	6.1
12	Necrotizing enterocolitis of newborn (P77)	31	1.2	6.1
12	Assault (homicide) (*U01,X85-Y09)	31	1.2	6.1
15	Septicemia (A40-A41)	30	1.2	5.9
...	All other causes (Residual)	604	23.6	118.9
Infants, white, non-Hispanic, both sexes				
...	All causes	12,906	100.0	559.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,963	23.0	128.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,796	13.9	77.8
3	Sudden infant death syndrome (R95)	1,246	9.7	54.0
4	Newborn affected by maternal complications of pregnancy (P01)	739	5.7	32.0
5	Accidents (unintentional injuries) (V01-X59)	555	4.3	24.0
6	Newborn affected by complications of placenta, cord and membranes (P02)	484	3.8	21.0
7	Respiratory distress of newborn (P22)	354	2.7	15.3
8	Bacterial sepsis of newborn (P36)	340	2.6	14.7
9	Neonatal hemorrhage (P50-P52,P54)	303	2.3	13.1
10	Diseases of the circulatory system (I00-I99)	237	1.8	10.3
11	Necrotizing enterocolitis of newborn (P77)	212	1.6	9.2
12	Atelectasis (P28.0-P28.1)	192	1.5	8.3
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	175	1.4	7.6
14	Assault (homicide) (*U01,X85-Y09)	145	1.1	6.3
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	108	0.8	4.7
...	All other causes (Residual)	3,057	23.7	132.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, white, non-Hispanic, male				
...	All causes	7,284	100.0	615.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,584	21.7	133.7
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,014	13.9	85.6
3	Sudden infant death syndrome (R95)	742	10.2	62.7
4	Newborn affected by maternal complications of pregnancy (P01)	418	5.7	35.3
5	Accidents (unintentional injuries) (V01-X59)	323	4.4	27.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	276	3.8	23.3
7	Respiratory distress of newborn (P22)	213	2.9	18.0
8	Neonatal hemorrhage (P50-P52,P54)	177	2.4	14.9
9	Bacterial sepsis of newborn (P36)	171	2.3	14.4
10	Necrotizing enterocolitis of newborn (P77)	132	1.8	11.1
11	Diseases of the circulatory system (I00-I99)	130	1.8	11.0
12	Atelectasis (P28.0-P28.1)	126	1.7	10.6
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	93	1.3	7.9
14	Assault (homicide) (*U01,X85-Y09)	76	1.0	6.4
15	Influenza and pneumonia (J10-J18)	68	0.9	5.7
...	All other causes (Residual)	1,741	23.9	147.0
Infants, white, non-Hispanic, female				
...	All causes	5,622	100.0	500.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,379	24.5	122.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	782	13.9	69.6
3	Sudden infant death syndrome (R95)	504	9.0	44.8
4	Newborn affected by maternal complications of pregnancy (P01)	321	5.7	28.6
5	Accidents (unintentional injuries) (V01-X59)	232	4.1	20.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	208	3.7	18.5
7	Bacterial sepsis of newborn (P36)	169	3.0	15.0
8	Respiratory distress of newborn (P22)	141	2.5	12.5
9	Neonatal hemorrhage (P50-P52,P54)	126	2.2	11.2
10	Diseases of the circulatory system (I00-I99)	107	1.9	9.5
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	82	1.5	7.3
12	Necrotizing enterocolitis of newborn (P77)	80	1.4	7.1
13	Assault (homicide) (*U01,X85-Y09)	69	1.2	6.1
14	Atelectasis (P28.0-P28.1)	66	1.2	5.9
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	52	0.9	4.6
...	All other causes (Residual)	1,304	23.2	116.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, both sexes				
...	All causes	8,507	100.0	1,378.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,903	22.4	308.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,107	13.0	179.3
3	Sudden infant death syndrome (R95)	695	8.2	112.6
4	Newborn affected by maternal complications of pregnancy (P01)	567	6.7	91.9
5	Newborn affected by complications of placenta, cord and membranes (P02)	362	4.3	58.6
6	Accidents (unintentional injuries) (V01-X59)	348	4.1	56.4
7	Bacterial sepsis of newborn (P36)	272	3.2	44.1
8	Respiratory distress of newborn (P22)	271	3.2	43.9
9	Necrotizing enterocolitis of newborn (P77)	204	2.4	33.0
10	Diseases of the circulatory system (I00-I99)	171	2.0	27.7
11	Neonatal hemorrhage (P50-P52,P54)	169	2.0	27.4
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	148	1.7	24.0
13	Chronic respiratory disease originating in the perinatal period (P27)	108	1.3	17.5
14	Atelectasis (P28.0-P28.1)	107	1.3	17.3
15	Assault (homicide) (*U01,X85-Y09)	101	1.2	16.4
...	All other causes (Residual)	1,974	23.2	319.8
Infants, black, non-Hispanic, male				
...	All causes	4,714	100.0	1,498.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,031	21.9	327.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	591	12.5	187.9
3	Sudden infant death syndrome (R95)	381	8.1	121.1
4	Newborn affected by maternal complications of pregnancy (P01)	316	6.7	100.4
5	Accidents (unintentional injuries) (V01-X59)	208	4.4	66.1
6	Newborn affected by complications of placenta, cord and membranes (P02)	202	4.3	64.2
7	Bacterial sepsis of newborn (P36)	150	3.2	47.7
8	Respiratory distress of newborn (P22)	139	2.9	44.2
9	Necrotizing enterocolitis of newborn (P77)	114	2.4	36.2
10	Diseases of the circulatory system (I00-I99)	98	2.1	31.2
11	Neonatal hemorrhage (P50-P52,P54)	91	1.9	28.9
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	87	1.8	27.7
13	Atelectasis (P28.0-P28.1)	67	1.4	21.3
14	Assault (homicide) (*U01,X85-Y09)	66	1.4	21.0
15	Chronic respiratory disease originating in the perinatal period (P27)	55	1.2	17.5
...	All other causes (Residual)	1,118	23.7	355.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, black, non-Hispanic, female				
...	All causes	3,793	100.0	1,253.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	872	23.0	288.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	516	13.6	170.5
3	Sudden infant death syndrome (R95)	314	8.3	103.8
4	Newborn affected by maternal complications of pregnancy (P01)	251	6.6	82.9
5	Newborn affected by complications of placenta, cord and membranes (P02)	160	4.2	52.9
6	Accidents (unintentional injuries) (V01-X59)	140	3.7	46.3
7	Respiratory distress of newborn (P22)	132	3.5	43.6
8	Bacterial sepsis of newborn (P36)	122	3.2	40.3
9	Necrotizing enterocolitis of newborn (P77)	90	2.4	29.7
10	Neonatal hemorrhage (P50-P52,P54)	78	2.1	25.8
11	Diseases of the circulatory system (I00-I99)	73	1.9	24.1
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	61	1.6	20.2
13	Chronic respiratory disease originating in the perinatal period (P27)	53	1.4	17.5
14	Septicemia (A40-A41)	46	1.2	15.2
15	Atelectasis (P28.0-P28.1)	40	1.1	13.2
...	All other causes (Residual)	845	22.3	279.2
Neonates, Hispanic, both sexes				
...	All causes	3,938	100.0	379.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,072	27.2	103.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	899	22.8	86.5
3	Newborn affected by maternal complications of pregnancy (P01)	296	7.5	28.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	228	5.8	21.9
5	Respiratory distress of newborn (P22)	165	4.2	15.9
6	Bacterial sepsis of newborn (P36)	146	3.7	14.1
7	Neonatal hemorrhage (P50-P52,P54)	119	3.0	11.5
8	Atelectasis (P28.0-P28.1)	82	2.1	7.9
9	Necrotizing enterocolitis of newborn (P77)	75	1.9	7.2
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	66	1.7	6.4
11	Pulmonary hemorrhage originating in the perinatal period (P26)	40	1.0	3.8
12	Hydrops fetalis not due to hemolytic disease (P83.2)	29	0.7	2.8
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	28	0.7	2.7
14	Diseases of the circulatory system (I00-I99)	25	0.6	2.4
15	Sudden infant death syndrome (R95)	22	0.6	2.1
...	All other causes (Residual)	646	16.4	62.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	2,162	100.0	407.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	565	26.1	106.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	502	23.2	94.6
3	Newborn affected by maternal complications of pregnancy (P01)	158	7.3	29.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	127	5.9	23.9
5	Respiratory distress of newborn (P22)	96	4.4	18.1
6	Bacterial sepsis of newborn (P36)	82	3.8	15.4
7	Neonatal hemorrhage (P50-P52,P54)	70	3.2	13.2
8	Atelectasis (P28.0-P28.1)	48	2.2	9.0
8	Necrotizing enterocolitis of newborn (P77)	48	2.2	9.0
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	36	1.7	6.8
11	Pulmonary hemorrhage originating in the perinatal period (P26)	24	1.1	4.5
12	Hydrops fetalis not due to hemolytic disease (P83.2)	18	0.8	*
13	Diseases of the circulatory system (I00-I99)	16	0.7	*
14	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	13	0.6	*
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	12	0.6	*
15	Sudden infant death syndrome (R95)	12	0.6	*
...	All other causes (Residual)	335	15.5	63.1
Neonates, Hispanic, female				
...	All causes	1,776	100.0	349.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	507	28.5	99.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	397	22.4	78.1
3	Newborn affected by maternal complications of pregnancy (P01)	138	7.8	27.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	101	5.7	19.9
5	Respiratory distress of newborn (P22)	69	3.9	13.6
6	Bacterial sepsis of newborn (P36)	64	3.6	12.6
7	Neonatal hemorrhage (P50-P52,P54)	49	2.8	9.6
8	Atelectasis (P28.0-P28.1)	34	1.9	6.7
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	30	1.7	5.9
10	Necrotizing enterocolitis of newborn (P77)	27	1.5	5.3
11	Interstitial emphysema and related conditions originating in the perinatal period (P25)	16	0.9	*
11	Pulmonary hemorrhage originating in the perinatal period (P26)	16	0.9	*
13	Hydrops fetalis not due to hemolytic disease (P83.2)	11	0.6	*
13	Accidents (unintentional injuries) (V01-X59)	11	0.6	*
15	Sudden infant death syndrome (R95)	10	0.6	*
...	All other causes (Residual)	296	16.7	58.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, both sexes				
...	All causes	8,508	100.0	368.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,188	25.7	94.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,752	20.6	75.9
3	Newborn affected by maternal complications of pregnancy (P01)	733	8.6	31.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	480	5.6	20.8
5	Respiratory distress of newborn (P22)	345	4.1	14.9
6	Bacterial sepsis of newborn (P36)	324	3.8	14.0
7	Neonatal hemorrhage (P50-P52,P54)	294	3.5	12.7
8	Necrotizing enterocolitis of newborn (P77)	194	2.3	8.4
9	Atelectasis (P28.0-P28.1)	184	2.2	8.0
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	166	2.0	7.2
11	Sudden infant death syndrome (R95)	116	1.4	5.0
12	Hydrops fetalis not due to hemolytic disease (P83.2)	89	1.0	3.9
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	76	0.9	3.3
14	Pulmonary hemorrhage originating in the perinatal period (P26)	69	0.8	3.0
15	Diseases of the circulatory system (I00-I99)	68	0.8	2.9
...	All other causes (Residual)	1,430	16.8	61.9
Neonates, white, non-Hispanic, male				
...	All causes	4,780	100.0	403.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,194	25.0	100.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	984	20.6	83.1
3	Newborn affected by maternal complications of pregnancy (P01)	417	8.7	35.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	274	5.7	23.1
5	Respiratory distress of newborn (P22)	207	4.3	17.5
6	Neonatal hemorrhage (P50-P52,P54)	171	3.6	14.4
7	Bacterial sepsis of newborn (P36)	164	3.4	13.8
8	Necrotizing enterocolitis of newborn (P77)	123	2.6	10.4
9	Atelectasis (P28.0-P28.1)	121	2.5	10.2
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	87	1.8	7.3
11	Sudden infant death syndrome (R95)	73	1.5	6.2
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	50	1.0	4.2
13	Hydrops fetalis not due to hemolytic disease (P83.2)	45	0.9	3.8
14	Pulmonary hemorrhage originating in the perinatal period (P26)	42	0.9	3.5
15	Diseases of the circulatory system (I00-I99)	38	0.8	3.2
...	All other causes (Residual)	790	16.5	66.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, non-Hispanic, female				
...	All causes	3,728	100.0	331.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	994	26.7	88.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	768	20.6	68.3
3	Newborn affected by maternal complications of pregnancy (P01)	316	8.5	28.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	206	5.5	18.3
5	Bacterial sepsis of newborn (P36)	160	4.3	14.2
6	Respiratory distress of newborn (P22)	138	3.7	12.3
7	Neonatal hemorrhage (P50-P52,P54)	123	3.3	10.9
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	79	2.1	7.0
9	Necrotizing enterocolitis of newborn (P77)	71	1.9	6.3
10	Atelectasis (P28.0-P28.1)	63	1.7	5.6
11	Hydrops fetalis not due to hemolytic disease (P83.2)	44	1.2	3.9
12	Sudden infant death syndrome (R95)	43	1.2	3.8
13	Accidents (unintentional injuries) (V01-X59)	31	0.8	2.8
14	Diseases of the circulatory system (I00-I99)	30	0.8	2.7
15	Hematological disorders (P60-P61)	28	0.8	2.5
...	All other causes (Residual)	634	17.0	56.4
Neonates, black, non-Hispanic, both sexes				
...	All causes	5,618	100.0	910.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,844	32.8	298.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	738	13.1	119.6
3	Newborn affected by maternal complications of pregnancy (P01)	556	9.9	90.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	355	6.3	57.5
5	Respiratory distress of newborn (P22)	266	4.7	43.1
6	Bacterial sepsis of newborn (P36)	258	4.6	41.8
7	Necrotizing enterocolitis of newborn (P77)	177	3.2	28.7
8	Neonatal hemorrhage (P50-P52,P54)	162	2.9	26.2
9	Atelectasis (P28.0-P28.1)	100	1.8	16.2
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	71	1.3	11.5
11	Sudden infant death syndrome (R95)	63	1.1	10.2
12	Pulmonary hemorrhage originating in the perinatal period (P26)	62	1.1	10.0
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	34	0.6	5.5
14	Newborn affected by other complications of labor and delivery (P03)	32	0.6	5.2
15	Congenital pneumonia (P23)	30	0.5	4.9
...	All other causes (Residual)	870	15.5	140.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, black, non-Hispanic, male				
...	All causes	3,099	100.0	985.1
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,001	32.3	318.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	395	12.7	125.6
3	Newborn affected by maternal complications of pregnancy (P01)	307	9.9	97.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	199	6.4	63.3
5	Bacterial sepsis of newborn (P36)	142	4.6	45.1
6	Respiratory distress of newborn (P22)	138	4.5	43.9
7	Necrotizing enterocolitis of newborn (P77)	101	3.3	32.1
8	Neonatal hemorrhage (P50-P52,P54)	86	2.8	27.3
9	Atelectasis (P28.0-P28.1)	61	2.0	19.4
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	42	1.4	13.4
11	Pulmonary hemorrhage originating in the perinatal period (P26)	36	1.2	11.4
12	Sudden infant death syndrome (R95)	35	1.1	11.1
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	20	0.6	6.4
14	Diseases of the circulatory system (I00-I99)	17	0.5	*
14	Accidents (unintentional injuries) (V01-X59)	17	0.5	*
...	All other causes (Residual)	502	16.2	159.6
Neonates, black, non-Hispanic, female				
...	All causes	2,519	100.0	832.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	843	33.5	278.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	343	13.6	113.3
3	Newborn affected by maternal complications of pregnancy (P01)	249	9.9	82.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	156	6.2	51.5
5	Respiratory distress of newborn (P22)	128	5.1	42.3
6	Bacterial sepsis of newborn (P36)	116	4.6	38.3
7	Neonatal hemorrhage (P50-P52,P54)	76	3.0	25.1
7	Necrotizing enterocolitis of newborn (P77)	76	3.0	25.1
9	Atelectasis (P28.0-P28.1)	39	1.5	12.9
10	Intrauterine hypoxia and birth asphyxia (P20-P21)	29	1.2	9.6
11	Sudden infant death syndrome (R95)	28	1.1	9.3
12	Pulmonary hemorrhage originating in the perinatal period (P26)	26	1.0	8.6
13	Newborn affected by other complications of labor and delivery (P03)	19	0.8	*
14	Newborn affected by maternal hypertensive disorders (P00.0)	14	0.6	*
14	Congenital pneumonia (P23)	14	0.6	*
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	14	0.6	*
...	All other causes (Residual)	349	13.9	115.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,797	100.0	172.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	394	21.9	37.9
2	Sudden infant death syndrome (R95)	266	14.8	25.6
3	Accidents (unintentional injuries) (V01-X59)	168	9.3	16.2
4	Diseases of the circulatory system (I00-I99)	83	4.6	8.0
5	Assault (homicide) (*U01,X85-Y09)	73	4.1	7.0
6	Septicemia (A40-A41)	67	3.7	6.4
7	Influenza and pneumonia (J10-J18)	59	3.3	5.7
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	47	2.6	4.5
9	Chronic respiratory disease originating in the perinatal period (P27)	32	1.8	3.1
10	Renal failure and other disorders of kidney (N17-N19,N25,N27)	20	1.1	1.9
11	Malignant neoplasms (C00-C97)	18	1.0	*
12	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	17	0.9	*
13	Necrotizing enterocolitis of newborn (P77)	15	0.8	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	13	0.7	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	12	0.7	*
...	All other causes (Residual)	513	28.5	49.4
Postneonates, Hispanic, male				
...	All causes	1,017	100.0	191.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	196	19.3	36.9
2	Sudden infant death syndrome (R95)	157	15.4	29.6
3	Accidents (unintentional injuries) (V01-X59)	93	9.1	17.5
4	Assault (homicide) (*U01,X85-Y09)	46	4.5	8.7
5	Diseases of the circulatory system (I00-I99)	42	4.1	7.9
6	Septicemia (A40-A41)	37	3.6	7.0
7	Influenza and pneumonia (J10-J18)	35	3.4	6.6
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	27	2.7	5.1
9	Chronic respiratory disease originating in the perinatal period (P27)	18	1.8	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	12	1.2	*
11	Necrotizing enterocolitis of newborn (P77)	11	1.1	*
12	Malignant neoplasms (C00-C97)	10	1.0	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	1.0	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	7	0.7	*
14	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	7	0.7	*
...	All other causes (Residual)	309	30.4	58.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	780	100.0	153.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	25.4	39.0
2	Sudden infant death syndrome (R95)	109	14.0	21.4
3	Accidents (unintentional injuries) (V01-X59)	75	9.6	14.8
4	Diseases of the circulatory system (I00-I99)	41	5.3	8.1
5	Septicemia (A40-A41)	30	3.8	5.9
6	Assault (homicide) (*U01,X85-Y09)	27	3.5	5.3
7	Influenza and pneumonia (J10-J18)	24	3.1	4.7
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	20	2.6	3.9
9	Chronic respiratory disease originating in the perinatal period (P27)	14	1.8	*
10	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	1.3	*
11	Malignant neoplasms (C00-C97)	8	1.0	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	6	0.8	*
12	Meningitis (G00,G03)	6	0.8	*
12	Anoxic brain damage, not elsewhere classified (G93.1)	6	0.8	*
12	Bacterial sepsis of newborn (P36)	6	0.8	*
...	All other causes (Residual)	200	25.6	39.4
Postneonates, white, non-Hispanic, both sexes				
...	All causes	4,398	100.0	190.5
1	Sudden infant death syndrome (R95)	1,130	25.7	48.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	775	17.6	33.6
3	Accidents (unintentional injuries) (V01-X59)	489	11.1	21.2
4	Diseases of the circulatory system (I00-I99)	169	3.8	7.3
5	Assault (homicide) (*U01,X85-Y09)	125	2.8	5.4
6	Influenza and pneumonia (J10-J18)	101	2.3	4.4
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	100	2.3	4.3
8	Chronic respiratory disease originating in the perinatal period (P27)	99	2.3	4.3
9	Septicemia (A40-A41)	91	2.1	3.9
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	44	1.0	1.9
11	Malignant neoplasms (C00-C97)	42	1.0	1.8
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	37	0.8	1.6
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	36	0.8	1.6
14	Anoxic brain damage, not elsewhere classified (G93.1)	28	0.6	1.2
15	Acute bronchitis and acute bronchiolitis (J20-J21)	25	0.6	1.1
...	All other causes (Residual)	1,107	25.2	48.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, white, non-Hispanic, male				
...	All causes	2,504	100.0	211.4
1	Sudden infant death syndrome (R95)	669	26.7	56.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	390	15.6	32.9
3	Accidents (unintentional injuries) (V01-X59)	288	11.5	24.3
4	Diseases of the circulatory system (I00-I99)	92	3.7	7.8
5	Assault (homicide) (*U01,X85-Y09)	70	2.8	5.9
6	Influenza and pneumonia (J10-J18)	67	2.7	5.7
7	Chronic respiratory disease originating in the perinatal period (P27)	55	2.2	4.6
8	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	51	2.0	4.3
9	Septicemia (A40-A41)	48	1.9	4.1
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	30	1.2	2.5
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	23	0.9	1.9
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	21	0.8	1.8
13	Malignant neoplasms (C00-C97)	20	0.8	1.7
14	Anoxic brain damage, not elsewhere classified (G93.1)	16	0.6	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	12	0.5	*
...	All other causes (Residual)	652	26.0	55.1
Postneonates, white, non-Hispanic, female				
...	All causes	1,894	100.0	168.5
1	Sudden infant death syndrome (R95)	461	24.3	41.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	385	20.3	34.2
3	Accidents (unintentional injuries) (V01-X59)	201	10.6	17.9
4	Diseases of the circulatory system (I00-I99)	77	4.1	6.8
5	Assault (homicide) (*U01,X85-Y09)	55	2.9	4.9
6	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	49	2.6	4.4
7	Chronic respiratory disease originating in the perinatal period (P27)	44	2.3	3.9
8	Septicemia (A40-A41)	43	2.3	3.8
9	Influenza and pneumonia (J10-J18)	34	1.8	3.0
10	Malignant neoplasms (C00-C97)	22	1.2	2.0
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	15	0.8	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	14	0.7	*
12	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	14	0.7	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	13	0.7	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	12	0.6	*
...	All other causes (Residual)	455	24.0	40.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, non-Hispanic, both sexes				
...	All causes	2,889	100.0	468.0
1	Sudden infant death syndrome (R95)	632	21.9	102.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	369	12.8	59.8
3	Accidents (unintentional injuries) (V01-X59)	320	11.1	51.8
4	Diseases of the circulatory system (I00-I99)	143	4.9	23.2
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	139	4.8	22.5
6	Septicemia (A40-A41)	99	3.4	16.0
7	Chronic respiratory disease originating in the perinatal period (P27)	98	3.4	15.9
8	Assault (homicide) (*U01,X85-Y09)	89	3.1	14.4
9	Influenza and pneumonia (J10-J18)	87	3.0	14.1
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	59	2.0	9.6
11	Necrotizing enterocolitis of newborn (P77)	27	0.9	4.4
12	Anoxic brain damage, not elsewhere classified (G93.1)	19	0.7	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	18	0.6	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	16	0.6	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	14	0.5	*
15	Bacterial sepsis of newborn (P36)	14	0.5	*
...	All other causes (Residual)	746	25.8	120.9
Postneonates, black, non-Hispanic, male				
...	All causes	1,615	100.0	513.3
1	Sudden infant death syndrome (R95)	346	21.4	110.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	196	12.1	62.3
3	Accidents (unintentional injuries) (V01-X59)	191	11.8	60.7
4	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	83	5.1	26.4
5	Diseases of the circulatory system (I00-I99)	81	5.0	25.7
6	Assault (homicide) (*U01,X85-Y09)	58	3.6	18.4
7	Septicemia (A40-A41)	53	3.3	16.8
7	Influenza and pneumonia (J10-J18)	53	3.3	16.8
9	Chronic respiratory disease originating in the perinatal period (P27)	51	3.2	16.2
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	30	1.9	9.5
11	Anoxic brain damage, not elsewhere classified (G93.1)	14	0.9	*
12	Necrotizing enterocolitis of newborn (P77)	13	0.8	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	9	0.6	*
13	Newborn affected by maternal complications of pregnancy (P01)	9	0.6	*
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	8	0.5	*
15	Bacterial sepsis of newborn (P36)	8	0.5	*
...	All other causes (Residual)	412	25.5	131.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2006

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, non-Hispanic, female				
...	All causes	1,274	100.0	421.0
1	Sudden infant death syndrome (R95)	286	22.4	94.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	173	13.6	57.2
3	Accidents (unintentional injuries) (V01-X59)	129	10.1	42.6
4	Diseases of the circulatory system (I00-I99)	62	4.9	20.5
5	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	56	4.4	18.5
6	Chronic respiratory disease originating in the perinatal period (P27)	47	3.7	15.5
7	Septicemia (A40-A41)	46	3.6	15.2
8	Influenza and pneumonia (J10-J18)	34	2.7	11.2
9	Assault (homicide) (*U01,X85-Y09)	31	2.4	10.2
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	29	2.3	9.6
11	Necrotizing enterocolitis of newborn (P77)	14	1.1	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	0.8	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	7	0.5	*
13	Acute bronchitis and acute bronchiolitis (J20-J21)	7	0.5	*
15	Meningitis (G00,G03)	6	0.5	*
15	Bacterial sepsis of newborn (P36)	6	0.5	*
...	All other causes (Residual)	331	26.0	109.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable