

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, race for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	4,813	100.0	530.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,249	26.0	137.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	813	16.9	89.6
3	Newborn affected by maternal complications of pregnancy (P01)	311	6.5	34.3
4	Sudden infant death syndrome (R95)	226	4.7	24.9
5	Newborn affected by complications of placenta, cord and membranes (P02)	211	4.4	23.2
6	Accidents (unintentional injuries) (V01-X59)	168	3.5	18.5
7	Bacterial sepsis of newborn (P36)	109	2.3	12.0
8	Diseases of the circulatory system (I00-I99)	108	2.2	11.9
9	Neonatal hemorrhage (P50-P52,P54)	101	2.1	11.1
10	Respiratory distress of newborn (P22)	82	1.7	9.0
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	68	1.4	7.5
12	Necrotizing enterocolitis of newborn (P77)	66	1.4	7.3
13	Diarrhea and gastroenteritis of infectious origin (A09)	59	1.2	6.5
14	Atelectasis (P28.0-P28.1)	57	1.2	6.3
15	Assault (homicide) (*U01,X85-Y09)	49	1.0	5.4
...	All other causes (Residual)	1,136	23.6	125.2
Infants, Hispanic, male				
...	All causes	2,662	100.0	576.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	642	24.1	139.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	474	17.8	102.6
3	Newborn affected by maternal complications of pregnancy (P01)	168	6.3	36.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	131	4.9	28.4
5	Sudden infant death syndrome (R95)	123	4.6	26.6
6	Accidents (unintentional injuries) (V01-X59)	86	3.2	18.6
7	Bacterial sepsis of newborn (P36)	62	2.3	13.4
8	Diseases of the circulatory system (I00-I99)	61	2.3	13.2
9	Neonatal hemorrhage (P50-P52,P54)	58	2.2	12.6
10	Respiratory distress of newborn (P22)	47	1.8	10.2
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	40	1.5	8.7
12	Diarrhea and gastroenteritis of infectious origin (A09)	35	1.3	7.6
12	Necrotizing enterocolitis of newborn (P77)	35	1.3	7.6
14	Atelectasis (P28.0-P28.1)	31	1.2	6.7
15	Assault (homicide) (*U01,X85-Y09)	26	1.0	5.6
...	All other causes (Residual)	643	24.2	139.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,151	100.0	482.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	607	28.2	136.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	339	15.8	76.0
3	Newborn affected by maternal complications of pregnancy (P01)	143	6.6	32.1
4	Sudden infant death syndrome (R95)	103	4.8	23.1
5	Accidents (unintentional injuries) (V01-X59)	82	3.8	18.4
6	Newborn affected by complications of placenta, cord and membranes (P02)	80	3.7	17.9
7	Diseases of the circulatory system (I00-I99)	47	2.2	10.5
7	Bacterial sepsis of newborn (P36)	47	2.2	10.5
9	Neonatal hemorrhage (P50-P52,P54)	43	2.0	9.6
10	Respiratory distress of newborn (P22)	35	1.6	7.9
11	Necrotizing enterocolitis of newborn (P77)	31	1.4	7.0
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	28	1.3	6.3
12	Hydrops fetalis not due to hemolytic disease (P83.2)	28	1.3	6.3
14	Atelectasis (P28.0-P28.1)	26	1.2	5.8
15	Diarrhea and gastroenteritis of infectious origin (A09)	24	1.1	5.4
...	All other causes (Residual)	488	22.7	109.5
Infants, Non-Hispanic, white, both sexes				
...	All causes	10,601	100.0	496.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,487	23.5	116.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,589	15.0	74.5
3	Sudden infant death syndrome (R95)	860	8.1	40.3
4	Newborn affected by maternal complications of pregnancy (P01)	637	6.0	29.8
5	Accidents (unintentional injuries) (V01-X59)	562	5.3	26.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	436	4.1	20.4
7	Bacterial sepsis of newborn (P36)	248	2.3	11.6
8	Respiratory distress of newborn (P22)	221	2.1	10.4
9	Diseases of the circulatory system (I00-I99)	214	2.0	10.0
10	Neonatal hemorrhage (P50-P52,P54)	204	1.9	9.6
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	158	1.5	7.4
12	Necrotizing enterocolitis of newborn (P77)	142	1.3	6.7
13	Assault (homicide) (*U01,X85-Y09)	133	1.3	6.2
14	Atelectasis (P28.0-P28.1)	128	1.2	6.0
15	Hydrops fetalis not due to hemolytic disease (P83.2)	90	0.8	4.2
...	All other causes (Residual)	2,492	23.5	116.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, white, male				
...	All causes	5,886	100.0	537.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,297	22.0	118.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	854	14.5	78.0
3	Sudden infant death syndrome (R95)	496	8.4	45.3
4	Newborn affected by maternal complications of pregnancy (P01)	378	6.4	34.5
5	Accidents (unintentional injuries) (V01-X59)	330	5.6	30.2
6	Newborn affected by complications of placenta, cord and membranes (P02)	226	3.8	20.6
7	Bacterial sepsis of newborn (P36)	140	2.4	12.8
8	Respiratory distress of newborn (P22)	137	2.3	12.5
9	Neonatal hemorrhage (P50-P52,P54)	123	2.1	11.2
10	Diseases of the circulatory system (I00-I99)	122	2.1	11.1
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	86	1.5	7.9
12	Necrotizing enterocolitis of newborn (P77)	77	1.3	7.0
13	Atelectasis (P28.0-P28.1)	74	1.3	6.8
14	Assault (homicide) (*U01,X85-Y09)	72	1.2	6.6
15	Hydrops fetalis not due to hemolytic disease (P83.2)	51	0.9	4.7
...	All other causes (Residual)	1,423	24.2	130.0
Infants, Non-Hispanic, white, female				
...	All causes	4,715	100.0	453.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,190	25.2	114.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	735	15.6	70.7
3	Sudden infant death syndrome (R95)	364	7.7	35.0
4	Newborn affected by maternal complications of pregnancy (P01)	259	5.5	24.9
5	Accidents (unintentional injuries) (V01-X59)	232	4.9	22.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	210	4.5	20.2
7	Bacterial sepsis of newborn (P36)	108	2.3	10.4
8	Diseases of the circulatory system (I00-I99)	92	2.0	8.8
9	Respiratory distress of newborn (P22)	84	1.8	8.1
10	Neonatal hemorrhage (P50-P52,P54)	81	1.7	7.8
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	72	1.5	6.9
12	Necrotizing enterocolitis of newborn (P77)	65	1.4	6.3
13	Assault (homicide) (*U01,X85-Y09)	61	1.3	5.9
14	Atelectasis (P28.0-P28.1)	54	1.1	5.2
15	Hydrops fetalis not due to hemolytic disease (P83.2)	39	0.8	3.8
...	All other causes (Residual)	1,069	22.7	102.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, black, both sexes				
...	All causes	6,765	100.0	1,159.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,537	22.7	263.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	905	13.4	155.1
3	Sudden infant death syndrome (R95)	518	7.7	88.8
4	Newborn affected by maternal complications of pregnancy (P01)	470	6.9	80.5
5	Accidents (unintentional injuries) (V01-X59)	377	5.6	64.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	292	4.3	50.0
7	Bacterial sepsis of newborn (P36)	182	2.7	31.2
8	Respiratory distress of newborn (P22)	167	2.5	28.6
9	Diseases of the circulatory system (I00-I99)	144	2.1	24.7
10	Necrotizing enterocolitis of newborn (P77)	110	1.6	18.9
11	Diarrhea and gastroenteritis of infectious origin (A09)	94	1.4	16.1
12	Assault (homicide) (*U01,X85-Y09)	93	1.4	15.9
13	Neonatal hemorrhage (P50-P52,P54)	89	1.3	15.3
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	78	1.2	13.4
15	Septicemia (A40-A41)	73	1.1	12.5
...	All other causes (Residual)	1,636	24.2	280.4
Infants, Non-Hispanic, black, male				
...	All causes	3,792	100.0	1,279.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	884	23.3	298.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	461	12.2	155.6
3	Sudden infant death syndrome (R95)	301	7.9	101.6
4	Newborn affected by maternal complications of pregnancy (P01)	282	7.4	95.2
5	Accidents (unintentional injuries) (V01-X59)	222	5.9	74.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	153	4.0	51.6
7	Respiratory distress of newborn (P22)	99	2.6	33.4
8	Bacterial sepsis of newborn (P36)	97	2.6	32.7
9	Diseases of the circulatory system (I00-I99)	75	2.0	25.3
10	Necrotizing enterocolitis of newborn (P77)	61	1.6	20.6
11	Assault (homicide) (*U01,X85-Y09)	59	1.6	19.9
12	Neonatal hemorrhage (P50-P52,P54)	53	1.4	17.9
13	Septicemia (A40-A41)	49	1.3	16.5
14	Diarrhea and gastroenteritis of infectious origin (A09)	41	1.1	13.8
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	40	1.1	13.5
...	All other causes (Residual)	915	24.1	308.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, black, female				
...	All causes	2,973	100.0	1,035.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	653	22.0	227.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	444	14.9	154.6
3	Sudden infant death syndrome (R95)	217	7.3	75.6
4	Newborn affected by maternal complications of pregnancy (P01)	188	6.3	65.5
5	Accidents (unintentional injuries) (V01-X59)	155	5.2	54.0
6	Newborn affected by complications of placenta, cord and membranes (P02)	139	4.7	48.4
7	Bacterial sepsis of newborn (P36)	85	2.9	29.6
8	Diseases of the circulatory system (I00-I99)	69	2.3	24.0
9	Respiratory distress of newborn (P22)	68	2.3	23.7
10	Diarrhea and gastroenteritis of infectious origin (A09)	53	1.8	18.5
11	Necrotizing enterocolitis of newborn (P77)	49	1.6	17.1
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	38	1.3	13.2
13	Neonatal hemorrhage (P50-P52,P54)	36	1.2	12.5
14	Assault (homicide) (*U01,X85-Y09)	34	1.1	11.8
15	Pulmonary hemorrhage originating in the perinatal period (P26)	27	0.9	9.4
15	Atelectasis (P28.0-P28.1)	27	0.9	9.4
...	All other causes (Residual)	691	23.2	240.7
Neonates, Hispanic, both sexes				
...	All causes	3,364	100.0	370.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	908	27.0	100.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	799	23.8	88.0
3	Newborn affected by maternal complications of pregnancy (P01)	309	9.2	34.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	207	6.2	22.8
5	Bacterial sepsis of newborn (P36)	105	3.1	11.6
6	Neonatal hemorrhage (P50-P52,P54)	100	3.0	11.0
7	Respiratory distress of newborn (P22)	81	2.4	8.9
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	61	1.8	6.7
9	Necrotizing enterocolitis of newborn (P77)	55	1.6	6.1
10	Atelectasis (P28.0-P28.1)	51	1.5	5.6
11	Hydrops fetalis not due to hemolytic disease (P83.2)	43	1.3	4.7
12	Pulmonary hemorrhage originating in the perinatal period (P26)	40	1.2	4.4
13	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	26	0.8	2.9
14	Newborn affected by other complications of labor and delivery (P03)	24	0.7	2.6
15	Diseases of the circulatory system (I00-I99)	23	0.7	2.5
...	All other causes (Residual)	532	15.8	58.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	1,870	100.0	404.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	481	25.7	104.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	465	24.9	100.7
3	Newborn affected by maternal complications of pregnancy (P01)	167	8.9	36.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	128	6.8	27.7
5	Bacterial sepsis of newborn (P36)	58	3.1	12.6
6	Neonatal hemorrhage (P50-P52,P54)	57	3.0	12.3
7	Respiratory distress of newborn (P22)	46	2.5	10.0
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	34	1.8	7.4
9	Necrotizing enterocolitis of newborn (P77)	30	1.6	6.5
10	Atelectasis (P28.0-P28.1)	26	1.4	5.6
11	Pulmonary hemorrhage originating in the perinatal period (P26)	21	1.1	4.5
12	Hydrops fetalis not due to hemolytic disease (P83.2)	19	1.0	*
13	Diseases of the circulatory system (I00-I99)	16	0.9	*
14	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	15	0.8	*
14	Newborn affected by other complications of labor and delivery (P03)	15	0.8	*
...	All other causes (Residual)	292	15.6	63.2
Neonates, Hispanic, female				
...	All causes	1,494	100.0	335.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	427	28.6	95.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	334	22.4	74.9
3	Newborn affected by maternal complications of pregnancy (P01)	142	9.5	31.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	79	5.3	17.7
5	Bacterial sepsis of newborn (P36)	47	3.1	10.5
6	Neonatal hemorrhage (P50-P52,P54)	43	2.9	9.6
7	Respiratory distress of newborn (P22)	35	2.3	7.9
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	27	1.8	6.1
9	Atelectasis (P28.0-P28.1)	25	1.7	5.6
9	Necrotizing enterocolitis of newborn (P77)	25	1.7	5.6
11	Hydrops fetalis not due to hemolytic disease (P83.2)	24	1.6	5.4
12	Pulmonary hemorrhage originating in the perinatal period (P26)	19	1.3	*
13	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	11	0.7	*
14	Newborn affected by other complications of labor and delivery (P03)	9	0.6	*
15	Sudden infant death syndrome (R95)	8	0.5	*
...	All other causes (Residual)	239	16.0	53.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, white, both sexes				
...	All causes	7,054	100.0	330.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,835	26.0	86.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,556	22.1	72.9
3	Newborn affected by maternal complications of pregnancy (P01)	633	9.0	29.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	432	6.1	20.2
5	Bacterial sepsis of newborn (P36)	240	3.4	11.2
6	Respiratory distress of newborn (P22)	214	3.0	10.0
7	Neonatal hemorrhage (P50-P52,P54)	193	2.7	9.0
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	148	2.1	6.9
9	Necrotizing enterocolitis of newborn (P77)	130	1.8	6.1
10	Atelectasis (P28.0-P28.1)	126	1.8	5.9
11	Hydrops fetalis not due to hemolytic disease (P83.2)	86	1.2	4.0
12	Sudden infant death syndrome (R95)	79	1.1	3.7
13	Diseases of the circulatory system (I00-I99)	59	0.8	2.8
14	Newborn affected by other complications of labor and delivery (P03)	58	0.8	2.7
14	Pulmonary hemorrhage originating in the perinatal period (P26)	58	0.8	2.7
...	All other causes (Residual)	1,207	17.1	56.6
Neonates, Non-Hispanic, white, male				
...	All causes	3,869	100.0	353.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	957	24.7	87.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	838	21.7	76.6
3	Newborn affected by maternal complications of pregnancy (P01)	375	9.7	34.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	222	5.7	20.3
5	Bacterial sepsis of newborn (P36)	136	3.5	12.4
6	Respiratory distress of newborn (P22)	133	3.4	12.2
7	Neonatal hemorrhage (P50-P52,P54)	115	3.0	10.5
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	83	2.1	7.6
9	Atelectasis (P28.0-P28.1)	73	1.9	6.7
10	Necrotizing enterocolitis of newborn (P77)	70	1.8	6.4
11	Hydrops fetalis not due to hemolytic disease (P83.2)	48	1.2	4.4
12	Sudden infant death syndrome (R95)	43	1.1	3.9
13	Pulmonary hemorrhage originating in the perinatal period (P26)	36	0.9	3.3
14	Newborn affected by other complications of labor and delivery (P03)	35	0.9	3.2
15	Diseases of the circulatory system (I00-I99)	32	0.8	2.9
...	All other causes (Residual)	673	17.4	61.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, white, female				
...	All causes	3,185	100.0	306.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	878	27.6	84.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	718	22.5	69.1
3	Newborn affected by maternal complications of pregnancy (P01)	258	8.1	24.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	210	6.6	20.2
5	Bacterial sepsis of newborn (P36)	104	3.3	10.0
6	Respiratory distress of newborn (P22)	81	2.5	7.8
7	Neonatal hemorrhage (P50-P52,P54)	78	2.4	7.5
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	65	2.0	6.3
9	Necrotizing enterocolitis of newborn (P77)	60	1.9	5.8
10	Atelectasis (P28.0-P28.1)	53	1.7	5.1
11	Hydrops fetalis not due to hemolytic disease (P83.2)	38	1.2	3.7
12	Sudden infant death syndrome (R95)	36	1.1	3.5
13	Diseases of the circulatory system (I00-I99)	27	0.8	2.6
13	Accidents (unintentional injuries) (V01-X59)	27	0.8	2.6
15	Newborn affected by other complications of labor and delivery (P03)	23	0.7	2.2
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	23	0.7	2.2
...	All other causes (Residual)	506	15.9	48.7
Neonates, Non-Hispanic, black, both sexes				
...	All causes	4,422	100.0	757.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,501	33.9	257.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	584	13.2	100.1
3	Newborn affected by maternal complications of pregnancy (P01)	468	10.6	80.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	289	6.5	49.5
5	Bacterial sepsis of newborn (P36)	172	3.9	29.5
6	Respiratory distress of newborn (P22)	160	3.6	27.4
7	Necrotizing enterocolitis of newborn (P77)	94	2.1	16.1
8	Neonatal hemorrhage (P50-P52,P54)	85	1.9	14.6
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	75	1.7	12.9
10	Pulmonary hemorrhage originating in the perinatal period (P26)	63	1.4	10.8
11	Atelectasis (P28.0-P28.1)	61	1.4	10.5
12	Accidents (unintentional injuries) (V01-X59)	42	0.9	7.2
13	Sudden infant death syndrome (R95)	39	0.9	6.7
14	Hydrops fetalis not due to hemolytic disease (P83.2)	30	0.7	5.1
15	Newborn affected by other complications of labor and delivery (P03)	27	0.6	4.6
...	All other causes (Residual)	732	16.6	125.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, black, male				
...	All causes	2,461	100.0	830.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	862	35.0	290.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	306	12.4	103.3
3	Newborn affected by maternal complications of pregnancy (P01)	281	11.4	94.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	152	6.2	51.3
5	Respiratory distress of newborn (P22)	93	3.8	31.4
6	Bacterial sepsis of newborn (P36)	90	3.7	30.4
7	Necrotizing enterocolitis of newborn (P77)	52	2.1	17.5
8	Neonatal hemorrhage (P50-P52,P54)	50	2.0	16.9
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	37	1.5	12.5
9	Atelectasis (P28.0-P28.1)	37	1.5	12.5
11	Pulmonary hemorrhage originating in the perinatal period (P26)	36	1.5	12.1
12	Accidents (unintentional injuries) (V01-X59)	24	1.0	8.1
13	Sudden infant death syndrome (R95)	22	0.9	7.4
14	Newborn affected by other complications of labor and delivery (P03)	18	0.7	*
15	Hydrops fetalis not due to hemolytic disease (P83.2)	15	0.6	*
...	All other causes (Residual)	386	15.7	130.2
Neonates, Non-Hispanic, black, female				
...	All causes	1,961	100.0	683.0
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	639	32.6	222.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	278	14.2	96.8
3	Newborn affected by maternal complications of pregnancy (P01)	187	9.5	65.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	137	7.0	47.7
5	Bacterial sepsis of newborn (P36)	82	4.2	28.6
6	Respiratory distress of newborn (P22)	67	3.4	23.3
7	Necrotizing enterocolitis of newborn (P77)	42	2.1	14.6
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	38	1.9	13.2
9	Neonatal hemorrhage (P50-P52,P54)	35	1.8	12.2
10	Pulmonary hemorrhage originating in the perinatal period (P26)	27	1.4	9.4
11	Atelectasis (P28.0-P28.1)	24	1.2	8.4
12	Accidents (unintentional injuries) (V01-X59)	18	0.9	*
13	Sudden infant death syndrome (R95)	17	0.9	*
14	Slow fetal growth and fetal malnutrition (P05)	15	0.8	*
14	Hydrops fetalis not due to hemolytic disease (P83.2)	15	0.8	*
...	All other causes (Residual)	340	17.3	118.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,449	100.0	159.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	341	23.5	37.6
2	Sudden infant death syndrome (R95)	210	14.5	23.1
3	Accidents (unintentional injuries) (V01-X59)	153	10.6	16.9
4	Diseases of the circulatory system (I00-I99)	85	5.9	9.4
5	Diarrhea and gastroenteritis of infectious origin (A09)	58	4.0	6.4
6	Assault (homicide) (*U01,X85-Y09)	47	3.2	5.2
7	Influenza and pneumonia (J09-J18)	28	1.9	3.1
8	Septicemia (A40-A41)	27	1.9	3.0
9	Chronic respiratory disease originating in the perinatal period (P27)	20	1.4	2.2
10	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	15	1.0	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	14	1.0	*
12	Malignant neoplasms (C00-C97)	11	0.8	*
12	Necrotizing enterocolitis of newborn (P77)	11	0.8	*
14	Whooping cough (A37)	8	0.6	*
14	Anoxic brain damage, not elsewhere classified (G93.1)	8	0.6	*
14	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	8	0.6	*
...	All other causes (Residual)	405	28.0	44.6
Postneonates, Hispanic, male				
...	All causes	792	100.0	171.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	161	20.3	34.9
2	Sudden infant death syndrome (R95)	115	14.5	24.9
3	Accidents (unintentional injuries) (V01-X59)	76	9.6	16.5
4	Diseases of the circulatory system (I00-I99)	45	5.7	9.7
5	Diarrhea and gastroenteritis of infectious origin (A09)	34	4.3	7.4
6	Assault (homicide) (*U01,X85-Y09)	24	3.0	5.2
7	Influenza and pneumonia (J09-J18)	17	2.1	*
8	Septicemia (A40-A41)	15	1.9	*
9	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	12	1.5	*
10	Chronic respiratory disease originating in the perinatal period (P27)	11	1.4	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	9	1.1	*
12	Malignant neoplasms (C00-C97)	7	0.9	*
13	Anoxic brain damage, not elsewhere classified (G93.1)	6	0.8	*
13	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	6	0.8	*
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	6	0.8	*
...	All other causes (Residual)	248	31.3	53.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	657	100.0	147.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	180	27.4	40.4
2	Sudden infant death syndrome (R95)	95	14.5	21.3
3	Accidents (unintentional injuries) (V01-X59)	77	11.7	17.3
4	Diseases of the circulatory system (I00-I99)	40	6.1	9.0
5	Diarrhea and gastroenteritis of infectious origin (A09)	24	3.7	5.4
6	Assault (homicide) (*U01,X85-Y09)	23	3.5	5.2
7	Septicemia (A40-A41)	12	1.8	*
8	Influenza and pneumonia (J09-J18)	11	1.7	*
9	Chronic respiratory disease originating in the perinatal period (P27)	9	1.4	*
10	Necrotizing enterocolitis of newborn (P77)	6	0.9	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	5	0.8	*
12	Malignant neoplasms (C00-C97)	4	0.6	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	4	0.6	*
12	Hydrops fetalis not due to hemolytic disease (P83.2)	4	0.6	*
15	Whooping cough (A37)	3	0.5	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	3	0.5	*
...	All other causes (Residual)	157	23.9	35.2
Postneonates, Non-Hispanic, white, both sexes				
...	All causes	3,547	100.0	166.2
1	Sudden infant death syndrome (R95)	781	22.0	36.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	652	18.4	30.6
3	Accidents (unintentional injuries) (V01-X59)	506	14.3	23.7
4	Diseases of the circulatory system (I00-I99)	155	4.4	7.3
5	Assault (homicide) (*U01,X85-Y09)	122	3.4	5.7
6	Diarrhea and gastroenteritis of infectious origin (A09)	72	2.0	3.4
7	Septicemia (A40-A41)	69	1.9	3.2
8	Influenza and pneumonia (J09-J18)	66	1.9	3.1
9	Chronic respiratory disease originating in the perinatal period (P27)	42	1.2	2.0
10	Malignant neoplasms (C00-C97)	33	0.9	1.5
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	33	0.9	1.5
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.5	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	19	0.5	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	17	0.5	*
15	Meningitis (G00,G03)	14	0.4	*
...	All other causes (Residual)	947	26.7	44.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, white, male				
...	All causes	2,017	100.0	184.3
1	Sudden infant death syndrome (R95)	453	22.5	41.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	340	16.9	31.1
3	Accidents (unintentional injuries) (V01-X59)	301	14.9	27.5
4	Diseases of the circulatory system (I00-I99)	90	4.5	8.2
5	Assault (homicide) (*U01,X85-Y09)	66	3.3	6.0
6	Diarrhea and gastroenteritis of infectious origin (A09)	39	1.9	3.6
7	Influenza and pneumonia (J09-J18)	38	1.9	3.5
8	Septicemia (A40-A41)	37	1.8	3.4
9	Chronic respiratory disease originating in the perinatal period (P27)	25	1.2	2.3
10	Malignant neoplasms (C00-C97)	17	0.8	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	16	0.8	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	13	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.6	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	9	0.4	*
15	Meningitis (G00,G03)	8	0.4	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	8	0.4	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	8	0.4	*
15	Neonatal hemorrhage (P50-P52,P54)	8	0.4	*
...	All other causes (Residual)	529	26.2	48.3
Postneonates, Non-Hispanic, white, female				
...	All causes	1,530	100.0	147.2
1	Sudden infant death syndrome (R95)	328	21.4	31.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	312	20.4	30.0
3	Accidents (unintentional injuries) (V01-X59)	205	13.4	19.7
4	Diseases of the circulatory system (I00-I99)	65	4.2	6.3
5	Assault (homicide) (*U01,X85-Y09)	56	3.7	5.4
6	Diarrhea and gastroenteritis of infectious origin (A09)	33	2.2	3.2
7	Septicemia (A40-A41)	32	2.1	3.1
8	Influenza and pneumonia (J09-J18)	28	1.8	2.7
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	17	1.1	*
9	Chronic respiratory disease originating in the perinatal period (P27)	17	1.1	*
11	Malignant neoplasms (C00-C97)	16	1.0	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	8	0.5	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	8	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	*
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	7	0.5	*
...	All other causes (Residual)	391	25.6	37.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, black, both sexes				
...	All causes	2,343	100.0	401.5
1	Sudden infant death syndrome (R95)	479	20.4	82.1
2	Accidents (unintentional injuries) (V01-X59)	335	14.3	57.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	321	13.7	55.0
4	Diseases of the circulatory system (I00-I99)	121	5.2	20.7
5	Diarrhea and gastroenteritis of infectious origin (A09)	92	3.9	15.8
6	Assault (homicide) (*U01,X85-Y09)	85	3.6	14.6
7	Septicemia (A40-A41)	73	3.1	12.5
8	Influenza and pneumonia (J09-J18)	52	2.2	8.9
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	36	1.5	6.2
10	Chronic respiratory disease originating in the perinatal period (P27)	35	1.5	6.0
11	Necrotizing enterocolitis of newborn (P77)	16	0.7	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	13	0.6	*
13	Meningitis (G00,G03)	12	0.5	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	11	0.5	*
15	Bacterial sepsis of newborn (P36)	10	0.4	*
...	All other causes (Residual)	652	27.8	111.7
Postneonates, Non-Hispanic, black, male				
...	All causes	1,331	100.0	449.1
1	Sudden infant death syndrome (R95)	279	21.0	94.1
2	Accidents (unintentional injuries) (V01-X59)	198	14.9	66.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	155	11.6	52.3
4	Diseases of the circulatory system (I00-I99)	65	4.9	21.9
5	Assault (homicide) (*U01,X85-Y09)	54	4.1	18.2
6	Septicemia (A40-A41)	49	3.7	16.5
7	Diarrhea and gastroenteritis of infectious origin (A09)	40	3.0	13.5
8	Influenza and pneumonia (J09-J18)	30	2.3	10.1
9	Chronic respiratory disease originating in the perinatal period (P27)	23	1.7	7.8
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	22	1.7	7.4
11	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	10	0.8	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	9	0.7	*
12	Necrotizing enterocolitis of newborn (P77)	9	0.7	*
14	Bacterial sepsis of newborn (P36)	7	0.5	*
15	Meningitis (G00,G03)	6	0.5	*
15	Respiratory distress of newborn (P22)	6	0.5	*
...	All other causes (Residual)	369	27.7	124.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2012

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, black, female				
...	All causes	1,012	100.0	352.4
1	Sudden infant death syndrome (R95)	200	19.8	69.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	166	16.4	57.8
3	Accidents (unintentional injuries) (V01-X59)	137	13.5	47.7
4	Diseases of the circulatory system (I00-I99)	56	5.5	19.5
5	Diarrhea and gastroenteritis of infectious origin (A09)	52	5.1	18.1
6	Assault (homicide) (*U01,X85-Y09)	31	3.1	10.8
7	Septicemia (A40-A41)	24	2.4	8.4
8	Influenza and pneumonia (J09-J18)	22	2.2	7.7
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	14	1.4	*
10	Chronic respiratory disease originating in the perinatal period (P27)	12	1.2	*
11	Necrotizing enterocolitis of newborn (P77)	7	0.7	*
12	Meningitis (G00,G03)	6	0.6	*
13	Acute bronchitis and acute bronchiolitis (J20-J21)	5	0.5	*
14	Anoxic brain damage, not elsewhere classified (G93.1)	4	0.4	*
14	Acute upper respiratory infections (J00-J06)	4	0.4	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	4	0.4	*
...	All other causes (Residual)	268	26.5	93.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012