

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, both sexes				
...	All causes	4,751	100.0	527.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,203	25.3	133.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	822	17.3	91.2
3	Newborn affected by maternal complications of pregnancy (P01)	340	7.2	37.7
4	Sudden infant death syndrome (R95)	227	4.8	25.2
5	Newborn affected by complications of placenta, cord and membranes (P02)	178	3.7	19.8
6	Accidents (unintentional injuries) (V01-X59)	166	3.5	18.4
7	Bacterial sepsis of newborn (P36)	126	2.7	14.0
8	Respiratory distress of newborn (P22)	105	2.2	11.7
9	Neonatal hemorrhage (P50-P52,P54)	88	1.9	9.8
10	Diseases of the circulatory system (I00-I99)	81	1.7	9.0
11	Necrotizing enterocolitis of newborn (P77)	65	1.4	7.2
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	59	1.2	6.5
13	Assault (homicide) (*U01,X85-Y09)	55	1.2	6.1
14	Atelectasis (P28.0-P28.1)	47	1.0	5.2
15	Diarrhea and gastroenteritis of infectious origin (A09)	46	1.0	5.1
...	All other causes (Residual)	1,143	24.1	126.9
Infants, Hispanic, male				
...	All causes	2,598	100.0	564.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	626	24.1	136.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	432	16.6	93.9
3	Newborn affected by maternal complications of pregnancy (P01)	184	7.1	40.0
4	Sudden infant death syndrome (R95)	131	5.0	28.5
5	Newborn affected by complications of placenta, cord and membranes (P02)	106	4.1	23.0
6	Accidents (unintentional injuries) (V01-X59)	90	3.5	19.6
7	Bacterial sepsis of newborn (P36)	75	2.9	16.3
8	Respiratory distress of newborn (P22)	64	2.5	13.9
9	Neonatal hemorrhage (P50-P52,P54)	48	1.8	10.4
10	Diseases of the circulatory system (I00-I99)	43	1.7	9.3
11	Necrotizing enterocolitis of newborn (P77)	36	1.4	7.8
12	Assault (homicide) (*U01,X85-Y09)	35	1.3	7.6
13	Diarrhea and gastroenteritis of infectious origin (A09)	30	1.2	6.5
14	Intrauterine hypoxia and birth asphyxia (P20-P21)	26	1.0	5.7
15	Atelectasis (P28.0-P28.1)	24	0.9	5.2
...	All other causes (Residual)	648	24.9	140.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Hispanic, female				
...	All causes	2,153	100.0	488.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	577	26.8	130.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	390	18.1	88.4
3	Newborn affected by maternal complications of pregnancy (P01)	156	7.2	35.4
4	Sudden infant death syndrome (R95)	96	4.5	21.8
5	Accidents (unintentional injuries) (V01-X59)	76	3.5	17.2
6	Newborn affected by complications of placenta, cord and membranes (P02)	72	3.3	16.3
7	Bacterial sepsis of newborn (P36)	51	2.4	11.6
8	Respiratory distress of newborn (P22)	41	1.9	9.3
9	Neonatal hemorrhage (P50-P52,P54)	40	1.9	9.1
10	Diseases of the circulatory system (I00-I99)	38	1.8	8.6
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	33	1.5	7.5
12	Necrotizing enterocolitis of newborn (P77)	29	1.3	6.6
13	Atelectasis (P28.0-P28.1)	23	1.1	5.2
14	Hydrops fetalis not due to hemolytic disease (P83.2)	20	0.9	4.5
14	Assault (homicide) (*U01,X85-Y09)	20	0.9	4.5
...	All other causes (Residual)	491	22.8	111.3
Infants, Non-Hispanic, white, both sexes				
...	All causes	10,563	100.0	496.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,403	22.7	112.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,596	15.1	75.0
3	Sudden infant death syndrome (R95)	786	7.4	36.9
4	Newborn affected by maternal complications of pregnancy (P01)	649	6.1	30.5
5	Accidents (unintentional injuries) (V01-X59)	543	5.1	25.5
6	Newborn affected by complications of placenta, cord and membranes (P02)	440	4.2	20.7
7	Respiratory distress of newborn (P22)	245	2.3	11.5
8	Bacterial sepsis of newborn (P36)	228	2.2	10.7
9	Diseases of the circulatory system (I00-I99)	211	2.0	9.9
10	Neonatal hemorrhage (P50-P52,P54)	183	1.7	8.6
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	172	1.6	8.1
12	Necrotizing enterocolitis of newborn (P77)	146	1.4	6.9
13	Atelectasis (P28.0-P28.1)	124	1.2	5.8
14	Assault (homicide) (*U01,X85-Y09)	113	1.1	5.3
15	Hydrops fetalis not due to hemolytic disease (P83.2)	96	0.9	4.5
...	All other causes (Residual)	2,628	24.9	123.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, white, male				
...	All causes	6,045	100.0	553.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,330	22.0	121.7
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	862	14.3	78.9
3	Sudden infant death syndrome (R95)	471	7.8	43.1
4	Newborn affected by maternal complications of pregnancy (P01)	389	6.4	35.6
5	Accidents (unintentional injuries) (V01-X59)	337	5.6	30.8
6	Newborn affected by complications of placenta, cord and membranes (P02)	253	4.2	23.2
7	Respiratory distress of newborn (P22)	136	2.2	12.4
8	Bacterial sepsis of newborn (P36)	133	2.2	12.2
9	Diseases of the circulatory system (I00-I99)	129	2.1	11.8
10	Neonatal hemorrhage (P50-P52,P54)	118	2.0	10.8
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	85	1.4	7.8
12	Necrotizing enterocolitis of newborn (P77)	80	1.3	7.3
13	Atelectasis (P28.0-P28.1)	72	1.2	6.6
14	Assault (homicide) (*U01,X85-Y09)	69	1.1	6.3
15	Hydrops fetalis not due to hemolytic disease (P83.2)	53	0.9	4.9
...	All other causes (Residual)	1,528	25.3	139.8
Infants, Non-Hispanic, white, female				
...	All causes	4,518	100.0	435.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,073	23.7	103.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	734	16.2	70.8
3	Sudden infant death syndrome (R95)	315	7.0	30.4
4	Newborn affected by maternal complications of pregnancy (P01)	260	5.8	25.1
5	Accidents (unintentional injuries) (V01-X59)	206	4.6	19.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	187	4.1	18.0
7	Respiratory distress of newborn (P22)	109	2.4	10.5
8	Bacterial sepsis of newborn (P36)	95	2.1	9.2
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	87	1.9	8.4
10	Diseases of the circulatory system (I00-I99)	82	1.8	7.9
11	Necrotizing enterocolitis of newborn (P77)	66	1.5	6.4
12	Neonatal hemorrhage (P50-P52,P54)	65	1.4	6.3
13	Atelectasis (P28.0-P28.1)	52	1.2	5.0
14	Assault (homicide) (*U01,X85-Y09)	44	1.0	4.2
15	Hydrops fetalis not due to hemolytic disease (P83.2)	43	1.0	4.1
...	All other causes (Residual)	1,100	24.3	106.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, black, both sexes				
...	All causes	6,781	100.0	1,161.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,533	22.6	262.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	888	13.1	152.1
3	Newborn affected by maternal complications of pregnancy (P01)	513	7.6	87.9
4	Sudden infant death syndrome (R95)	472	7.0	80.8
5	Accidents (unintentional injuries) (V01-X59)	410	6.0	70.2
6	Newborn affected by complications of placenta, cord and membranes (P02)	282	4.2	48.3
7	Bacterial sepsis of newborn (P36)	193	2.8	33.1
8	Respiratory distress of newborn (P22)	146	2.2	25.0
9	Diseases of the circulatory system (I00-I99)	140	2.1	24.0
10	Necrotizing enterocolitis of newborn (P77)	125	1.8	21.4
11	Neonatal hemorrhage (P50-P52,P54)	99	1.5	17.0
12	Assault (homicide) (*U01,X85-Y09)	96	1.4	16.4
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	76	1.1	13.0
14	Diarrhea and gastroenteritis of infectious origin (A09)	72	1.1	12.3
15	Influenza and pneumonia (J09-J18)	69	1.0	11.8
15	Atelectasis (P28.0-P28.1)	69	1.0	11.8
...	All other causes (Residual)	1,598	23.6	273.7
Infants, Non-Hispanic, black, male				
...	All causes	3,698	100.0	1,247.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	853	23.1	287.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	459	12.4	154.8
3	Newborn affected by maternal complications of pregnancy (P01)	278	7.5	93.8
4	Sudden infant death syndrome (R95)	271	7.3	91.4
5	Accidents (unintentional injuries) (V01-X59)	212	5.7	71.5
6	Newborn affected by complications of placenta, cord and membranes (P02)	140	3.8	47.2
7	Bacterial sepsis of newborn (P36)	106	2.9	35.8
8	Respiratory distress of newborn (P22)	80	2.2	27.0
9	Diseases of the circulatory system (I00-I99)	78	2.1	26.3
10	Necrotizing enterocolitis of newborn (P77)	68	1.8	22.9
11	Assault (homicide) (*U01,X85-Y09)	58	1.6	19.6
12	Neonatal hemorrhage (P50-P52,P54)	55	1.5	18.6
13	Diarrhea and gastroenteritis of infectious origin (A09)	44	1.2	14.8
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	44	1.2	14.8
15	Influenza and pneumonia (J09-J18)	39	1.1	13.2
...	All other causes (Residual)	913	24.7	308.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Infants, Non-Hispanic, black, female				
...	All causes	3,083	100.0	1,072.7
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	680	22.1	236.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	429	13.9	149.3
3	Newborn affected by maternal complications of pregnancy (P01)	235	7.6	81.8
4	Sudden infant death syndrome (R95)	201	6.5	69.9
5	Accidents (unintentional injuries) (V01-X59)	198	6.4	68.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	142	4.6	49.4
7	Bacterial sepsis of newborn (P36)	87	2.8	30.3
8	Respiratory distress of newborn (P22)	66	2.1	23.0
9	Diseases of the circulatory system (I00-I99)	62	2.0	21.6
10	Necrotizing enterocolitis of newborn (P77)	57	1.8	19.8
11	Neonatal hemorrhage (P50-P52,P54)	44	1.4	15.3
12	Assault (homicide) (*U01,X85-Y09)	38	1.2	13.2
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	32	1.0	11.1
13	Atelectasis (P28.0-P28.1)	32	1.0	11.1
15	Influenza and pneumonia (J09-J18)	30	1.0	10.4
...	All other causes (Residual)	750	24.3	261.0
Neonates, Hispanic, both sexes				
...	All causes	3,359	100.0	372.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	847	25.2	94.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	806	24.0	89.5
3	Newborn affected by maternal complications of pregnancy (P01)	337	10.0	37.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	177	5.3	19.6
5	Bacterial sepsis of newborn (P36)	122	3.6	13.5
6	Respiratory distress of newborn (P22)	102	3.0	11.3
7	Neonatal hemorrhage (P50-P52,P54)	82	2.4	9.1
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	58	1.7	6.4
8	Necrotizing enterocolitis of newborn (P77)	58	1.7	6.4
10	Atelectasis (P28.0-P28.1)	44	1.3	4.9
11	Hydrops fetalis not due to hemolytic disease (P83.2)	41	1.2	4.6
12	Pulmonary hemorrhage originating in the perinatal period (P26)	29	0.9	3.2
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	26	0.8	2.9
14	Diseases of the circulatory system (I00-I99)	21	0.6	2.3
14	Accidents (unintentional injuries) (V01-X59)	21	0.6	2.3
...	All other causes (Residual)	588	17.5	65.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Hispanic, male				
...	All causes	1,835	100.0	399.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	464	25.3	100.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	426	23.2	92.6
3	Newborn affected by maternal complications of pregnancy (P01)	184	10.0	40.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	106	5.8	23.0
5	Bacterial sepsis of newborn (P36)	72	3.9	15.7
6	Respiratory distress of newborn (P22)	63	3.4	13.7
7	Neonatal hemorrhage (P50-P52,P54)	44	2.4	9.6
8	Necrotizing enterocolitis of newborn (P77)	34	1.9	7.4
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	25	1.4	5.4
10	Atelectasis (P28.0-P28.1)	23	1.3	5.0
11	Hydrops fetalis not due to hemolytic disease (P83.2)	22	1.2	4.8
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	19	1.0	*
13	Pulmonary hemorrhage originating in the perinatal period (P26)	18	1.0	*
14	Accidents (unintentional injuries) (V01-X59)	15	0.8	*
15	Hematological disorders (P60-P61)	12	0.7	*
...	All other causes (Residual)	308	16.8	67.0
Neonates, Hispanic, female				
...	All causes	1,524	100.0	345.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	383	25.1	86.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	380	24.9	86.1
3	Newborn affected by maternal complications of pregnancy (P01)	153	10.0	34.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	71	4.7	16.1
5	Bacterial sepsis of newborn (P36)	50	3.3	11.3
6	Respiratory distress of newborn (P22)	39	2.6	8.8
7	Neonatal hemorrhage (P50-P52,P54)	38	2.5	8.6
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	33	2.2	7.5
9	Necrotizing enterocolitis of newborn (P77)	24	1.6	5.4
10	Atelectasis (P28.0-P28.1)	21	1.4	4.8
11	Hydrops fetalis not due to hemolytic disease (P83.2)	19	1.2	*
12	Diseases of the circulatory system (I00-I99)	13	0.9	*
13	Pulmonary hemorrhage originating in the perinatal period (P26)	11	0.7	*
14	Newborn affected by maternal hypertensive disorders (P00.0)	10	0.7	*
14	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	10	0.7	*
14	Newborn affected by other complications of labor and delivery (P03)	10	0.7	*
...	All other causes (Residual)	259	17.0	58.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, white, both sexes				
...	All causes	7,093	100.0	333.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,743	24.6	81.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,566	22.1	73.5
3	Newborn affected by maternal complications of pregnancy (P01)	646	9.1	30.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	437	6.2	20.5
5	Respiratory distress of newborn (P22)	238	3.4	11.2
6	Bacterial sepsis of newborn (P36)	224	3.2	10.5
7	Neonatal hemorrhage (P50-P52,P54)	179	2.5	8.4
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	167	2.4	7.8
9	Necrotizing enterocolitis of newborn (P77)	135	1.9	6.3
10	Atelectasis (P28.0-P28.1)	116	1.6	5.4
11	Hydrops fetalis not due to hemolytic disease (P83.2)	92	1.3	4.3
12	Sudden infant death syndrome (R95)	85	1.2	4.0
13	Pulmonary hemorrhage originating in the perinatal period (P26)	67	0.9	3.1
14	Newborn affected by other complications of labor and delivery (P03)	59	0.8	2.8
15	Accidents (unintentional injuries) (V01-X59)	55	0.8	2.6
...	All other causes (Residual)	1,284	18.1	60.3
Neonates, Non-Hispanic, white, male				
...	All causes	4,010	100.0	367.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	990	24.7	90.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	848	21.1	77.6
3	Newborn affected by maternal complications of pregnancy (P01)	387	9.7	35.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	252	6.3	23.1
5	Respiratory distress of newborn (P22)	135	3.4	12.4
6	Bacterial sepsis of newborn (P36)	131	3.3	12.0
7	Neonatal hemorrhage (P50-P52,P54)	115	2.9	10.5
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	81	2.0	7.4
9	Necrotizing enterocolitis of newborn (P77)	73	1.8	6.7
10	Atelectasis (P28.0-P28.1)	68	1.7	6.2
11	Hydrops fetalis not due to hemolytic disease (P83.2)	49	1.2	4.5
12	Pulmonary hemorrhage originating in the perinatal period (P26)	45	1.1	4.1
12	Sudden infant death syndrome (R95)	45	1.1	4.1
14	Newborn affected by other complications of labor and delivery (P03)	36	0.9	3.3
15	Accidents (unintentional injuries) (V01-X59)	33	0.8	3.0
...	All other causes (Residual)	722	18.0	66.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, white, female				
...	All causes	3,083	100.0	297.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	753	24.4	72.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	718	23.3	69.3
3	Newborn affected by maternal complications of pregnancy (P01)	259	8.4	25.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	185	6.0	17.8
5	Respiratory distress of newborn (P22)	103	3.3	9.9
6	Bacterial sepsis of newborn (P36)	93	3.0	9.0
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	86	2.8	8.3
8	Neonatal hemorrhage (P50-P52,P54)	64	2.1	6.2
9	Necrotizing enterocolitis of newborn (P77)	62	2.0	6.0
10	Atelectasis (P28.0-P28.1)	48	1.6	4.6
11	Hydrops fetalis not due to hemolytic disease (P83.2)	43	1.4	4.1
12	Sudden infant death syndrome (R95)	40	1.3	3.9
13	Slow fetal growth and fetal malnutrition (P05)	29	0.9	2.8
14	Diseases of the circulatory system (I00-I99)	23	0.7	2.2
14	Newborn affected by other complications of labor and delivery (P03)	23	0.7	2.2
...	All other causes (Residual)	554	18.0	53.4
Neonates, Non-Hispanic, black, both sexes				
...	All causes	4,471	100.0	765.8
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,486	33.2	254.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	577	12.9	98.8
3	Newborn affected by maternal complications of pregnancy (P01)	511	11.4	87.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	275	6.2	47.1
5	Bacterial sepsis of newborn (P36)	185	4.1	31.7
6	Respiratory distress of newborn (P22)	144	3.2	24.7
7	Necrotizing enterocolitis of newborn (P77)	105	2.3	18.0
8	Neonatal hemorrhage (P50-P52,P54)	96	2.1	16.4
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	76	1.7	13.0
10	Atelectasis (P28.0-P28.1)	65	1.5	11.1
11	Slow fetal growth and fetal malnutrition (P05)	48	1.1	8.2
12	Pulmonary hemorrhage originating in the perinatal period (P26)	47	1.1	8.1
13	Accidents (unintentional injuries) (V01-X59)	40	0.9	6.9
14	Newborn affected by other complications of labor and delivery (P03)	38	0.8	6.5
15	Sudden infant death syndrome (R95)	31	0.7	5.3
...	All other causes (Residual)	747	16.7	127.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Neonates, Non-Hispanic, black, male				
...	All causes	2,420	100.0	816.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	823	34.0	277.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	292	12.1	98.5
3	Newborn affected by maternal complications of pregnancy (P01)	277	11.4	93.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	134	5.5	45.2
5	Bacterial sepsis of newborn (P36)	103	4.3	34.7
6	Respiratory distress of newborn (P22)	80	3.3	27.0
7	Necrotizing enterocolitis of newborn (P77)	57	2.4	19.2
8	Neonatal hemorrhage (P50-P52,P54)	52	2.1	17.5
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	44	1.8	14.8
10	Atelectasis (P28.0-P28.1)	34	1.4	11.5
11	Pulmonary hemorrhage originating in the perinatal period (P26)	29	1.2	9.8
12	Newborn affected by other complications of labor and delivery (P03)	22	0.9	7.4
13	Slow fetal growth and fetal malnutrition (P05)	20	0.8	6.7
13	Hydrops fetalis not due to hemolytic disease (P83.2)	20	0.8	6.7
13	Accidents (unintentional injuries) (V01-X59)	20	0.8	6.7
...	All other causes (Residual)	413	17.1	139.3
Neonates, Non-Hispanic, black, female				
...	All causes	2,051	100.0	713.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	663	32.3	230.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	285	13.9	99.2
3	Newborn affected by maternal complications of pregnancy (P01)	234	11.4	81.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	141	6.9	49.1
5	Bacterial sepsis of newborn (P36)	82	4.0	28.5
6	Respiratory distress of newborn (P22)	64	3.1	22.3
7	Necrotizing enterocolitis of newborn (P77)	48	2.3	16.7
8	Neonatal hemorrhage (P50-P52,P54)	44	2.1	15.3
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	32	1.6	11.1
10	Atelectasis (P28.0-P28.1)	31	1.5	10.8
11	Slow fetal growth and fetal malnutrition (P05)	28	1.4	9.7
12	Accidents (unintentional injuries) (V01-X59)	20	1.0	7.0
13	Pulmonary hemorrhage originating in the perinatal period (P26)	18	0.9	*
14	Newborn affected by other complications of labor and delivery (P03)	16	0.8	*
15	Sudden infant death syndrome (R95)	14	0.7	*
...	All other causes (Residual)	331	16.1	115.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, both sexes				
...	All causes	1,392	100.0	154.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	356	25.6	39.5
2	Sudden infant death syndrome (R95)	210	15.1	23.3
3	Accidents (unintentional injuries) (V01-X59)	145	10.4	16.1
4	Diseases of the circulatory system (I00-I99)	60	4.3	6.7
5	Assault (homicide) (*U01,X85-Y09)	50	3.6	5.5
6	Diarrhea and gastroenteritis of infectious origin (A09)	44	3.2	4.9
7	Septicemia (A40-A41)	35	2.5	3.9
7	Influenza and pneumonia (J09-J18)	35	2.5	3.9
9	Chronic respiratory disease originating in the perinatal period (P27)	19	1.4	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	16	1.1	*
11	Anoxic brain damage, not elsewhere classified (G93.1)	13	0.9	*
12	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	10	0.7	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	0.7	*
14	Malignant neoplasms (C00-C97)	8	0.6	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	7	0.5	*
15	Meningitis (G00,G03)	7	0.5	*
15	Necrotizing enterocolitis of newborn (P77)	7	0.5	*
...	All other causes (Residual)	360	25.9	40.0
Postneonates, Hispanic, male				
...	All causes	763	100.0	165.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	21.2	35.2
2	Sudden infant death syndrome (R95)	120	15.7	26.1
3	Accidents (unintentional injuries) (V01-X59)	75	9.8	16.3
4	Diseases of the circulatory system (I00-I99)	35	4.6	7.6
5	Assault (homicide) (*U01,X85-Y09)	33	4.3	7.2
6	Diarrhea and gastroenteritis of infectious origin (A09)	29	3.8	6.3
7	Influenza and pneumonia (J09-J18)	23	3.0	5.0
8	Septicemia (A40-A41)	22	2.9	4.8
9	Chronic respiratory disease originating in the perinatal period (P27)	10	1.3	*
10	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	8	1.0	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	7	0.9	*
12	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	6	0.8	*
13	Acute bronchitis and acute bronchiolitis (J20-J21)	5	0.7	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	4	0.5	*
14	Meningitis (G00,G03)	4	0.5	*
14	Anoxic brain damage, not elsewhere classified (G93.1)	4	0.5	*
14	Neonatal hemorrhage (P50-P52,P54)	4	0.5	*
...	All other causes (Residual)	212	27.8	46.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Hispanic, female				
...	All causes	629	100.0	142.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	194	30.8	44.0
2	Sudden infant death syndrome (R95)	90	14.3	20.4
3	Accidents (unintentional injuries) (V01-X59)	70	11.1	15.9
4	Diseases of the circulatory system (I00-I99)	25	4.0	5.7
5	Assault (homicide) (*U01,X85-Y09)	17	2.7	*
6	Diarrhea and gastroenteritis of infectious origin (A09)	15	2.4	*
7	Septicemia (A40-A41)	13	2.1	*
8	Influenza and pneumonia (J09-J18)	12	1.9	*
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	10	1.6	*
10	Anoxic brain damage, not elsewhere classified (G93.1)	9	1.4	*
10	Chronic respiratory disease originating in the perinatal period (P27)	9	1.4	*
12	Malignant neoplasms (C00-C97)	5	0.8	*
12	Necrotizing enterocolitis of newborn (P77)	5	0.8	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	3	0.5	*
14	Meningitis (G00,G03)	3	0.5	*
14	Bronchitis, chronic and unspecified (J40-J42)	3	0.5	*
14	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	3	0.5	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	3	0.5	*
14	Newborn affected by maternal complications of pregnancy (P01)	3	0.5	*
...	All other causes (Residual)	137	21.8	31.1
Postneonates, Non-Hispanic, white, both sexes				
...	All causes	3,470	100.0	163.0
1	Sudden infant death syndrome (R95)	701	20.2	32.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	660	19.0	31.0
3	Accidents (unintentional injuries) (V01-X59)	488	14.1	22.9
4	Diseases of the circulatory system (I00-I99)	159	4.6	7.5
5	Assault (homicide) (*U01,X85-Y09)	106	3.1	5.0
6	Diarrhea and gastroenteritis of infectious origin (A09)	82	2.4	3.9
7	Influenza and pneumonia (J09-J18)	65	1.9	3.1
8	Septicemia (A40-A41)	58	1.7	2.7
9	Chronic respiratory disease originating in the perinatal period (P27)	41	1.2	1.9
10	Malignant neoplasms (C00-C97)	34	1.0	1.6
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	30	0.9	1.4
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	26	0.7	1.2
13	Anoxic brain damage, not elsewhere classified (G93.1)	16	0.5	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	15	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.4	*
...	All other causes (Residual)	975	28.1	45.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, white, male				
...	All causes	2,035	100.0	186.2
1	Sudden infant death syndrome (R95)	426	20.9	39.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	340	16.7	31.1
3	Accidents (unintentional injuries) (V01-X59)	304	14.9	27.8
4	Diseases of the circulatory system (I00-I99)	100	4.9	9.2
5	Assault (homicide) (*U01,X85-Y09)	67	3.3	6.1
6	Diarrhea and gastroenteritis of infectious origin (A09)	42	2.1	3.8
6	Influenza and pneumonia (J09-J18)	42	2.1	3.8
8	Septicemia (A40-A41)	32	1.6	2.9
9	Chronic respiratory disease originating in the perinatal period (P27)	26	1.3	2.4
10	Malignant neoplasms (C00-C97)	21	1.0	1.9
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	16	0.8	*
12	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	14	0.7	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	11	0.5	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	10	0.5	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	9	0.4	*
...	All other causes (Residual)	575	28.3	52.6
Postneonates, Non-Hispanic, white, female				
...	All causes	1,435	100.0	138.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	320	22.3	30.9
2	Sudden infant death syndrome (R95)	275	19.2	26.5
3	Accidents (unintentional injuries) (V01-X59)	184	12.8	17.8
4	Diseases of the circulatory system (I00-I99)	59	4.1	5.7
5	Diarrhea and gastroenteritis of infectious origin (A09)	40	2.8	3.9
6	Assault (homicide) (*U01,X85-Y09)	39	2.7	3.8
7	Septicemia (A40-A41)	26	1.8	2.5
8	Influenza and pneumonia (J09-J18)	23	1.6	2.2
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	16	1.1	*
10	Chronic respiratory disease originating in the perinatal period (P27)	15	1.0	*
11	Malignant neoplasms (C00-C97)	13	0.9	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	10	0.7	*
13	Anoxic brain damage, not elsewhere classified (G93.1)	9	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.4	*
14	Meningitis (G00,G03)	6	0.4	*
14	Respiratory distress of newborn (P22)	6	0.4	*
...	All other causes (Residual)	388	27.0	37.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, black, both sexes				
...	All causes	2,310	100.0	395.7
1	Sudden infant death syndrome (R95)	441	19.1	75.5
2	Accidents (unintentional injuries) (V01-X59)	370	16.0	63.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	311	13.5	53.3
4	Diseases of the circulatory system (I00-I99)	119	5.2	20.4
5	Assault (homicide) (*U01,X85-Y09)	91	3.9	15.6
6	Diarrhea and gastroenteritis of infectious origin (A09)	72	3.1	12.3
7	Influenza and pneumonia (J09-J18)	67	2.9	11.5
8	Septicemia (A40-A41)	52	2.3	8.9
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	47	2.0	8.1
10	Chronic respiratory disease originating in the perinatal period (P27)	22	1.0	3.8
11	Necrotizing enterocolitis of newborn (P77)	20	0.9	3.4
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	14	0.6	*
12	Meningitis (G00,G03)	14	0.6	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	12	0.5	*
15	Bronchitis, chronic and unspecified (J40-J42)	9	0.4	*
...	All other causes (Residual)	649	28.1	111.2
Postneonates, Non-Hispanic, black, male				
...	All causes	1,278	100.0	431.1
1	Sudden infant death syndrome (R95)	254	19.9	85.7
2	Accidents (unintentional injuries) (V01-X59)	192	15.0	64.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	167	13.1	56.3
4	Diseases of the circulatory system (I00-I99)	66	5.2	22.3
5	Assault (homicide) (*U01,X85-Y09)	56	4.4	18.9
6	Diarrhea and gastroenteritis of infectious origin (A09)	44	3.4	14.8
7	Influenza and pneumonia (J09-J18)	37	2.9	12.5
8	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	30	2.3	10.1
9	Septicemia (A40-A41)	29	2.3	9.8
10	Necrotizing enterocolitis of newborn (P77)	11	0.9	*
11	Acute bronchitis and acute bronchiolitis (J20-J21)	8	0.6	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	6	0.5	*
12	Meningitis (G00,G03)	6	0.5	*
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	6	0.5	*
12	Newborn affected by complications of placenta, cord and membranes (P02)	6	0.5	*
12	Chronic respiratory disease originating in the perinatal period (P27)	6	0.5	*
...	All other causes (Residual)	354	27.7	119.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2013

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Non-Hispanic, black, female				
...	All causes	1,032	100.0	359.1
1	Sudden infant death syndrome (R95)	187	18.1	65.1
2	Accidents (unintentional injuries) (V01-X59)	178	17.2	61.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	14.0	50.1
4	Diseases of the circulatory system (I00-I99)	53	5.1	18.4
5	Assault (homicide) (*U01,X85-Y09)	35	3.4	12.2
6	Influenza and pneumonia (J09-J18)	30	2.9	10.4
7	Diarrhea and gastroenteritis of infectious origin (A09)	28	2.7	9.7
8	Septicemia (A40-A41)	23	2.2	8.0
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	17	1.6	*
10	Chronic respiratory disease originating in the perinatal period (P27)	16	1.6	*
11	Necrotizing enterocolitis of newborn (P77)	9	0.9	*
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	8	0.8	*
12	Meningitis (G00,G03)	8	0.8	*
14	Malignant neoplasms (C00-C97)	6	0.6	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	5	0.5	*
15	Bacterial sepsis of newborn (P36)	5	0.5	*
...	All other causes (Residual)	280	27.1	97.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2013