Person with disabilities experiencing problems accessing NCHS birth and death tables should contact **nchsed@cdc.gov**, or call 301-458-4688

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
United States												
All races, both sexes	45	2	-	-	1	1	-	-	-	-	-	3
Male	15	1	-	-	1	1	-	-	-	-	-	1
Female	30	1	-	-	-	-	-	-	-	-	-	2
White, both sexes	38	1	-	-	1	1	-	-	-	-	-	-
Male	13	1	-	-	1	1	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Florida												
All races, both sexes	5	1	-	-	1	-	-	-	-	-	-	-
Male	3	-	-	-	1	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	1	-	-	-	-	-	-	-
Male	3	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Illinois												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Iowa												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Kentucky												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Maine												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Massachusetts												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Minnesota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
New Mexico												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
North Carolina												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Oregon												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Texas												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Washington												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VII. Diseases of the eye and adnexa (H00-H57)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
United States												
All races, both sexes	-	2	4	-	3	7	5	5	1	10	1	-
Male	-	1	1	-	1	3	2	2	-	-	1	-
Female	-	1	3	-	2	4	3	3	1	10	-	-
White, both sexes	-	2	4	-	3	7	5	5	1	8	-	-
Male	-	1	1	-	1	3	2	2	-	-	-	-
Female	-	1	3	-	2	4	3	3	1	8	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	2	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Florida												
All races, both sexes	-	-	-	-	1	-	1	-	-	1	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	1	-	1	-	-	1	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Illinois												
All races, both sexes	-	1	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	2	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Iowa												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Maine												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Massachusetts												
All races, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	1	1	-	1	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	1	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Ohio												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Oregon												
All races, both sexes	-	1	1	-	-	-	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
White, both sexes	-	1	1	-	-	-	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Texas												
All races, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Washington												
All races, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VII. Diseases of the eye and adnexa (H00-H57)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
United States												
All races, both sexes	58	7	3	-	1	-	1	1	3	1	5	4
Male	33	4	1	-	1	-	-	1	2	1	2	3
Female	25	3	2	-	-	-	1	-	1	-	3	1
White, both sexes	43	3	3	-	-	-	1	1	3	1	3	2
Male	26	2	1	-	-	-	-	1	2	1	2	1
Female	17	1	2	-	-	-	1	-	1	-	1	1
Black, both sexes	11	3	-	-	1	-	-	-	-	-	1	2
Male	6	2	-	-	1	-	-	-	-	-	-	2
Female	5	1	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	5	-	-	-	-	-	-	1	-	-	1	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Colorado												
All races, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Delaware												
All races, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Florida												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Illinois												
All races, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Iowa												
All races, both sexes	4	-	-	-	-	-	1	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	1	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
New Mexico												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Ohio												
All races, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	1	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Texas												
All races, both sexes	11	3	-	-	-	-	-	-	-	-	1	-
Male	10	3	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	2	-	-	-	-	-	-	-	-	1	-
Male	9	2	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Washington												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
VIII. Diseases of the ear and mastoid process (H60-H93)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
United States												
All races, both sexes	4	2	1	4	4	4	3	5	4	1	-	-
Male	3	2	1	3	1	1	2	2	3	-	-	-
Female	1	-	-	1	3	3	1	3	1	1	-	-
White, both sexes	3	2	-	3	3	4	3	4	3	1	-	-
Male	3	2	-	3	1	1	2	2	2	-	-	-
Female	-	-	-	-	2	3	1	2	1	1	-	-
Black, both sexes	1	-	1	1	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	-	-	-	1	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Florida												
All races, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Illinois												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Iowa												
All races, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Maine												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Ohio												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Oregon												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Texas												
All races, both sexes	1	1	-	1	-	1	1	1	1	-	-	-
Male	1	1	-	1	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	1	-	1	-	1	1	1	1	-	-	-
Male	1	1	-	1	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Washington												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
VIII. Diseases of the ear and mastoid process (H60-H93)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
United States												
All races, both sexes	860,843	529	223	169	203	500	927	1,517	2,656	5,180	11,004	19,235
Male	407,936	295	129	82	121	323	606	996	1,723	3,430	7,398	13,158
Female	452,907	234	94	87	82	177	321	521	933	1,750	3,606	6,077
White, both sexes	741,184	339	144	108	137	336	611	962	1,699	3,498	7,810	13,833
Male	350,177	188	88	55	85	213	405	658	1,131	2,386	5,481	9,826
Female	391,007	151	56	53	52	123	206	304	568	1,112	2,329	4,007
Black, both sexes	100,895	157	62	50	54	146	274	492	834	1,492	2,819	4,847
Male	47,972	91	30	23	30	98	175	296	501	918	1,650	2,947
Female	52,923	66	32	27	24	48	99	196	333	574	1,169	1,900
American Indian 1/, both sexes	3,561	9	2	-	2	5	10	16	36	58	105	181
Male	1,853	3	2	-	-	4	4	12	25	33	73	115
Female	1,708	6	-	-	2	1	6	4	11	25	32	66
Asian or Pacific Islander 2/, both sexes	15,203	24	15	11	10	13	32	47	87	132	270	374
Male	7,934	13	9	4	6	8	22	30	66	93	194	270
Female	7,269	11	6	7	4	5	10	17	21	39	76	104
Alabama												
All races, both sexes	16,989	16	3	1	4	10	26	45	79	148	295	493
Male	7,835	7	-	1	2	7	19	30	52	95	187	329
Female	9,154	9	3	-	2	3	7	15	27	53	108	164
White, both sexes	12,869	6	2	1	2	4	13	26	39	89	192	291
Male	5,943	3	-	1	1	2	11	19	27	58	123	193
Female	6,926	3	2	-	1	2	2	7	12	31	69	98
Black, both sexes	4,071	10	1	-	2	6	13	19	39	57	103	201
Male	1,871	4	-	-	1	5	8	11	24	37	64	135
Female	2,200	6	1	-	1	1	5	8	15	20	39	66
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	1	1	-	-
Male	11	-	-	-	-	-	-	-	1	-	-	-
Female	15	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Alaska												
All races, both sexes	853	1	-	-	1	1	6	3	3	8	28	30
Male	496	-	-	-	1	1	4	1	1	6	18	19
Female	357	1	-	-	-	-	2	2	2	2	10	11
White, both sexes	624	-	-	-	1	-	3	2	1	5	17	22
Male	380	-	-	-	1	-	2	1	1	4	13	16
Female	244	-	-	-	-	-	1	1	-	1	4	6
Black, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	175	1	-	-	-	-	1	1	2	2	8	6
Male	86	-	-	-	-	-	-	-	-	1	3	2
Female	89	1	-	-	-	-	1	1	2	1	5	4
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	1	2	-	-	1	2	2
Male	15	-	-	-	-	1	2	-	-	1	1	1
Female	16	-	-	-	-	-	-	-	-	-	1	1
Arizona												
All races, both sexes	14,475	11	3	3	2	9	12	26	41	81	181	296
Male	7,421	4	3	1	1	7	6	15	28	53	114	208
Female	7,054	7	-	2	1	2	6	11	13	28	67	88
White, both sexes	13,669	7	3	3	2	9	8	17	31	65	153	253
Male	7,013	2	3	1	1	7	5	11	22	43	100	183
Female	6,656	5	-	2	1	2	3	6	9	22	53	70
Black, both sexes	359	-	-	-	-	-	2	5	5	8	21	18
Male	175	-	-	-	-	-	1	2	3	5	9	7
Female	184	-	-	-	-	-	1	3	2	3	12	11
American Indian 1/, both sexes	334	2	-	-	-	-	2	3	4	7	6	20
Male	179	2	-	-	-	-	-	2	3	4	4	15
Female	155	-	-	-	-	-	2	1	1	3	2	5
Asian or Pacific Islander 2/, both sexes	113	2	-	-	-	-	-	1	1	1	1	5
Male	54	-	-	-	-	-	-	-	-	1	1	3
Female	59	2	-	-	-	-	-	1	1	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Arkansas												
All races, both sexes	10,125	6	2	4	1	7	17	21	41	78	169	256
Male	4,834	3	1	1	-	6	14	14	27	44	112	177
Female	5,291	3	1	3	1	1	3	7	14	34	57	79
White, both sexes	8,635	5	2	3	1	5	8	4	31	53	115	184
Male	4,157	3	1	1	-	5	6	3	22	29	77	129
Female	4,478	2	1	2	1	-	2	1	9	24	38	55
Black, both sexes	1,453	1	-	1	-	2	9	17	10	24	52	69
Male	663	-	-	-	-	1	8	11	5	15	33	45
Female	790	1	-	1	-	1	1	6	5	9	19	24
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	-	2	2
Male	7	-	-	-	-	-	-	-	-	-	2	2
Female	13	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	1	-	-
California												
All races, both sexes	87,527	44	23	14	20	45	89	162	235	524	1,083	1,847
Male	42,105	28	17	5	14	31	58	111	153	365	766	1,305
Female	45,422	16	6	9	6	14	31	51	82	159	317	542
White, both sexes	73,519	35	15	9	14	31	66	108	170	355	786	1,373
Male	35,089	20	11	3	9	22	43	78	112	260	580	986
Female	38,430	15	4	6	5	9	23	30	58	95	206	387
Black, both sexes	7,090	4	3	-	3	8	12	34	42	120	190	331
Male	3,474	3	1	-	3	5	9	19	24	69	109	219
Female	3,616	1	2	-	-	3	3	15	18	51	81	112
American Indian 1/, both sexes	376	-	-	-	-	2	-	2	3	5	10	16
Male	181	-	-	-	-	2	-	2	2	4	5	8
Female	195	-	-	-	-	-	-	-	1	1	5	8
Asian or Pacific Islander 2/, both sexes	6,542	5	5	5	3	4	11	18	20	44	97	127
Male	3,361	5	5	2	2	2	6	12	15	32	72	92
Female	3,181	-	-	3	1	2	5	6	5	12	25	35

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Colorado												
All races, both sexes	8,933	6	3	5	3	7	7	14	30	70	108	193
Male	4,362	1	3	3	1	5	4	9	20	52	70	143
Female	4,571	5	-	2	2	2	3	5	10	18	38	50
White, both sexes	8,485	5	3	5	2	7	6	10	26	59	97	174
Male	4,142	1	3	3	1	5	3	5	17	47	63	133
Female	4,343	4	-	2	1	2	3	5	9	12	34	41
Black, both sexes	329	-	-	-	-	-	1	3	4	6	7	16
Male	158	-	-	-	-	-	1	3	3	2	3	7
Female	171	-	-	-	-	-	-	-	1	4	4	9
American Indian 1/, both sexes	33	-	-	-	1	-	-	-	-	1	2	2
Male	19	-	-	-	-	-	-	-	-	1	2	2
Female	14	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	86	1	-	-	-	-	-	1	-	4	2	1
Male	43	-	-	-	-	-	-	1	-	2	2	1
Female	43	1	-	-	-	-	-	-	-	2	-	-
Connecticut												
All races, both sexes	9,999	10	2	4	3	4	11	11	30	54	89	188
Male	4,567	4	1	3	1	3	6	8	20	42	60	139
Female	5,432	6	1	1	2	1	5	3	10	12	29	49
White, both sexes	9,296	7	2	3	-	3	9	9	20	43	74	159
Male	4,219	3	1	2	-	3	6	6	15	32	53	117
Female	5,077	4	1	1	-	-	3	3	5	11	21	42
Black, both sexes	626	3	-	1	3	1	2	1	9	9	15	24
Male	309	1	-	1	1	-	-	1	4	9	7	17
Female	317	2	-	-	2	1	2	-	5	-	8	7
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	1	1	-	1
Male	6	-	-	-	-	-	-	-	1	-	-	1
Female	8	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	63	-	-	-	-	-	-	1	-	1	-	4
Male	33	-	-	-	-	-	-	1	-	1	-	4
Female	30	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Delaware												
All races, both sexes	2,586	3	2	-	-	1	2	6	7	13	34	55
Male	1,237	2	-	-	-	1	2	4	5	8	23	38
Female	1,349	1	2	-	-	-	-	2	2	5	11	17
White, both sexes	2,141	1	1	-	-	-	2	4	4	7	22	40
Male	1,043	1	-	-	-	-	2	3	3	5	16	29
Female	1,098	-	1	-	-	-	-	1	1	2	6	11
Black, both sexes	419	2	1	-	-	1	-	2	3	6	12	13
Male	178	1	-	-	-	1	-	1	2	3	7	8
Female	241	1	1	-	-	-	-	1	1	3	5	5
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1,900	-	-	-	1	1	3	3	6	15	33	56
Male	890	-	-	-	1	1	1	3	5	9	21	36
Female	1,010	-	-	-	-	-	2	-	1	6	12	20
White, both sexes	361	-	-	-	-	-	-	-	-	-	1	1
Male	172	-	-	-	-	-	-	-	-	-	1	-
Female	189	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1,511	-	-	-	1	1	3	2	6	14	32	55
Male	699	-	-	-	1	1	1	2	5	9	20	36
Female	812	-	-	-	-	-	2	-	1	5	12	19
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	1	-	1	-	-
Male	18	-	-	-	-	-	-	1	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Florida												
All races, both sexes	60,263	30	12	5	8	31	50	69	136	325	660	1,122
Male	30,153	16	8	2	5	19	35	46	82	207	458	763
Female	30,110	14	4	3	3	12	15	23	54	118	202	359
White, both sexes	53,653	20	5	2	6	18	30	36	73	217	478	827
Male	26,974	11	4	1	5	11	18	23	46	150	354	593
Female	26,679	9	1	1	1	7	12	13	27	67	124	234
Black, both sexes	6,234	9	4	3	2	10	19	32	60	103	174	283
Male	2,989	4	2	1	-	6	16	22	34	54	98	162
Female	3,245	5	2	2	2	4	3	10	26	49	76	121
American Indian 1/, both sexes	55	-	1	-	-	-	-	1	1	1	2	1
Male	33	-	1	-	-	-	-	1	1	1	1	1
Female	22	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	321	1	2	-	-	3	1	-	2	4	6	11
Male	157	1	1	-	-	2	1	-	1	2	5	7
Female	164	-	1	-	-	1	-	-	1	2	1	4
Georgia												
All races, both sexes	22,493	26	14	7	11	23	29	73	109	226	417	753
Male	10,735	19	8	2	4	13	17	43	68	143	275	492
Female	11,758	7	6	5	7	10	12	30	41	83	142	261
White, both sexes	16,244	12	4	5	6	8	6	35	48	121	227	440
Male	7,781	7	2	2	3	3	4	22	30	74	154	311
Female	8,463	5	2	3	3	5	2	13	18	47	73	129
Black, both sexes	6,077	13	9	2	5	15	21	37	57	100	180	310
Male	2,863	11	6	-	1	10	11	21	34	68	115	178
Female	3,214	2	3	2	4	5	10	16	23	32	65	132
American Indian 1/, both sexes	38	-	-	-	-	-	-	-	-	2	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	134	1	1	-	-	-	2	1	4	3	10	3
Male	74	1	-	-	-	-	2	-	4	1	6	3
Female	60	-	1	-	-	-	-	1	-	2	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Hawaii												
All races, both sexes	3,258	3	1	2	3	-	1	6	10	21	60	79
Male	1,749	1	1	-	3	-	1	2	9	16	39	50
Female	1,509	2	-	2	-	-	-	4	1	5	21	29
White, both sexes	821	-	-	-	1	-	-	2	1	3	13	21
Male	446	-	-	-	1	-	-	1	1	3	9	14
Female	375	-	-	-	-	-	-	1	-	-	4	7
Black, both sexes	23	-	-	-	1	-	-	-	-	-	2	3
Male	18	-	-	-	1	-	-	-	-	-	2	3
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2,411	3	1	2	1	-	1	4	9	18	45	55
Male	1,284	1	1	-	1	-	1	1	8	13	28	33
Female	1,127	2	-	2	-	-	-	3	1	5	17	22
Idaho												
All races, both sexes	3,401	-	2	-	1	1	1	4	8	23	38	48
Male	1,700	-	2	-	-	1	-	3	6	14	26	36
Female	1,701	-	-	-	1	-	1	1	2	9	12	12
White, both sexes	3,358	-	2	-	1	1	1	4	8	22	37	48
Male	1,675	-	2	-	-	1	-	3	6	13	26	36
Female	1,683	-	-	-	1	-	1	1	2	9	11	12
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	24	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Illinois												
All races, both sexes	37,198	31	11	9	12	28	49	79	130	235	515	883
Male	17,368	19	4	2	8	14	30	56	87	170	369	622
Female	19,830	12	7	7	4	14	19	23	43	65	146	261
White, both sexes	31,137	22	7	6	6	18	31	48	76	142	348	565
Male	14,382	11	3	1	4	8	18	36	55	110	262	426
Female	16,755	11	4	5	2	10	13	12	21	32	86	139
Black, both sexes	5,598	9	4	3	5	10	16	31	51	89	160	310
Male	2,749	8	1	1	4	6	11	20	29	58	100	189
Female	2,849	1	3	2	1	4	5	11	22	31	60	121
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	-	1	-	-
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	438	-	-	-	1	-	2	-	3	3	7	8
Male	228	-	-	-	-	-	1	-	3	1	7	7
Female	210	-	-	-	1	-	1	-	-	2	-	1
Indiana												
All races, both sexes	19,408	14	7	4	4	18	29	40	85	125	265	479
Male	9,073	5	3	3	2	11	17	26	56	71	169	296
Female	10,335	9	4	1	2	7	12	14	29	54	96	183
White, both sexes	17,909	11	5	2	3	15	26	31	70	100	211	389
Male	8,347	4	3	2	1	10	16	21	43	57	138	248
Female	9,562	7	2	-	2	5	10	10	27	43	73	141
Black, both sexes	1,436	3	1	2	1	3	3	9	14	24	52	88
Male	695	1	-	1	1	1	1	5	12	14	30	47
Female	741	2	1	1	-	2	2	4	2	10	22	41
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	57	-	1	-	-	-	-	-	1	1	2	1
Male	31	-	-	-	-	-	-	-	1	-	1	1
Female	26	-	1	-	-	-	-	-	-	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Iowa												
All races, both sexes	10,263	1	2	3	1	7	9	12	18	45	100	179
Male	4,773	1	2	1	1	3	2	8	8	29	59	121
Female	5,490	-	-	2	-	4	7	4	10	16	41	58
White, both sexes	10,114	1	2	3	1	7	8	11	16	41	90	169
Male	4,692	1	2	1	1	3	2	7	8	27	55	115
Female	5,422	-	-	2	-	4	6	4	8	14	35	54
Black, both sexes	124	-	-	-	-	-	1	1	1	4	10	8
Male	64	-	-	-	-	-	-	1	-	2	4	5
Female	60	-	-	-	-	-	1	-	1	2	6	3
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	1	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	1	-	-	-
Kansas												
All races, both sexes	8,369	7	-	2	-	4	10	15	20	33	86	179
Male	3,904	6	-	1	-	3	6	8	16	24	56	125
Female	4,465	1	-	1	-	1	4	7	4	9	30	54
White, both sexes	7,866	5	-	2	-	2	7	13	16	26	77	147
Male	3,628	4	-	1	-	1	5	7	14	19	50	101
Female	4,238	1	-	1	-	1	2	6	2	7	27	46
Black, both sexes	401	2	-	-	-	2	3	1	4	5	6	25
Male	222	2	-	-	-	2	1	-	2	4	3	21
Female	179	-	-	-	-	-	2	1	2	1	3	4
American Indian 1/, both sexes	75	-	-	-	-	-	-	-	-	1	3	5
Male	44	-	-	-	-	-	-	-	-	-	3	2
Female	31	-	-	-	-	-	-	-	-	1	-	3
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	1	-	1	-	2
Male	10	-	-	-	-	-	-	1	-	1	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Kentucky												
All races, both sexes	13,879	15	3	1	7	4	17	32	54	98	239	368
Male	6,538	10	1	-	6	4	12	21	38	66	155	268
Female	7,341	5	2	1	1	-	5	11	16	32	84	100
White, both sexes	12,936	7	3	1	6	4	15	27	48	85	210	319
Male	6,116	5	1	-	5	4	11	18	34	57	137	240
Female	6,820	2	2	1	1	-	4	9	14	28	73	79
Black, both sexes	909	6	-	-	1	-	2	5	5	13	29	48
Male	400	4	-	-	1	-	1	3	3	9	18	27
Female	509	2	-	-	-	-	1	2	2	4	11	21
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	2	-	-	-	-	-	-	1	-	-	-
Male	17	1	-	-	-	-	-	-	1	-	-	-
Female	11	1	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	14,512	11	6	5	7	13	25	43	53	124	260	468
Male	6,958	6	4	2	3	8	14	29	33	79	160	291
Female	7,554	5	2	3	4	5	11	14	20	45	100	177
White, both sexes	10,113	7	5	1	4	6	16	14	32	66	125	212
Male	4,856	3	3	-	2	3	7	12	21	45	86	143
Female	5,257	4	2	1	2	3	9	2	11	21	39	69
Black, both sexes	4,318	4	1	4	3	7	8	28	21	57	133	253
Male	2,056	3	1	2	1	5	6	16	12	33	73	146
Female	2,262	1	-	2	2	2	2	12	9	24	60	107
American Indian 1/, both sexes	24	-	-	-	-	-	-	1	-	-	1	1
Male	13	-	-	-	-	-	-	1	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	57	-	-	-	-	-	1	-	-	1	1	2
Male	33	-	-	-	-	-	1	-	-	1	1	1
Female	24	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Maine												
All races, both sexes	3,931	1	2	-	-	3	4	5	8	16	41	69
Male	1,863	-	-	-	-	2	3	3	6	11	26	49
Female	2,068	1	2	-	-	1	1	2	2	5	15	20
White, both sexes	3,912	1	1	-	-	3	4	5	8	16	40	67
Male	1,854	-	-	-	-	2	3	3	6	11	25	47
Female	2,058	1	1	-	-	1	1	2	2	5	15	20
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	1	2
Male	5	-	-	-	-	-	-	-	-	-	1	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	15,170	13	8	2	2	14	19	32	66	110	252	380
Male	7,092	7	2	-	1	12	12	26	38	73	163	266
Female	8,078	6	6	2	1	2	7	6	28	37	89	114
White, both sexes	11,078	6	2	1	2	6	13	15	28	48	130	200
Male	5,131	4	1	-	1	5	8	11	16	34	91	149
Female	5,947	2	1	1	1	1	5	4	12	14	39	51
Black, both sexes	3,861	5	5	1	-	8	5	17	37	59	117	176
Male	1,844	2	-	-	-	7	4	15	22	37	69	114
Female	2,017	3	5	1	-	1	1	2	15	22	48	62
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	2	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	215	2	1	-	-	-	1	-	1	1	5	4
Male	109	1	1	-	-	-	-	-	-	1	3	3
Female	106	1	-	-	-	-	1	-	1	-	2	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Massachusetts												
All races, both sexes	17,518	8	4	2	3	4	19	24	42	81	169	281
Male	7,933	5	2	1	3	4	15	19	24	54	120	206
Female	9,585	3	2	1	-	-	4	5	18	27	49	75
White, both sexes	16,667	7	3	2	1	4	16	20	33	66	144	246
Male	7,527	5	2	1	1	4	12	17	18	43	104	183
Female	9,140	2	1	1	-	-	4	3	15	23	40	63
Black, both sexes	686	1	1	-	2	-	2	4	7	13	22	31
Male	316	-	-	-	2	-	2	2	4	9	13	19
Female	370	1	1	-	-	-	-	2	3	4	9	12
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	144	-	-	-	-	-	1	-	2	2	2	3
Male	80	-	-	-	-	-	1	-	2	2	2	3
Female	64	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	32,576	16	5	11	7	21	29	55	94	209	375	713
Male	15,310	9	3	5	4	15	18	36	67	137	254	494
Female	17,266	7	2	6	3	6	11	19	27	72	121	219
White, both sexes	27,622	12	3	6	5	16	18	39	54	131	270	488
Male	12,880	8	2	5	3	11	12	28	42	91	199	364
Female	14,742	4	1	1	2	5	6	11	12	40	71	124
Black, both sexes	4,690	4	2	5	2	5	11	15	37	73	97	216
Male	2,286	1	1	-	1	4	6	7	23	43	50	124
Female	2,404	3	1	5	1	1	5	8	14	30	47	92
American Indian 1/, both sexes	123	-	-	-	-	-	-	1	1	3	3	5
Male	61	-	-	-	-	-	-	1	1	1	2	3
Female	62	-	-	-	-	-	-	-	-	2	1	2
Asian or Pacific Islander 2/, both sexes	141	-	-	-	-	-	-	-	2	2	5	4
Male	83	-	-	-	-	-	-	-	1	2	3	3
Female	58	-	-	-	-	-	-	-	1	-	2	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Minnesota												
All races, both sexes	11,503	4	1	2	3	3	14	19	25	48	131	227
Male	5,546	-	1	1	2	2	9	13	16	34	88	156
Female	5,957	4	-	1	1	1	5	6	9	14	43	71
White, both sexes	11,157	4	1	2	2	3	14	14	18	39	106	193
Male	5,347	-	1	1	1	2	9	10	12	30	73	133
Female	5,810	4	-	1	1	1	5	4	6	9	33	60
Black, both sexes	185	-	-	-	-	-	-	3	5	6	14	27
Male	112	-	-	-	-	-	-	3	3	3	7	18
Female	73	-	-	-	-	-	-	-	2	3	7	9
American Indian 1/, both sexes	71	-	-	-	-	-	-	-	1	2	4	6
Male	39	-	-	-	-	-	-	-	-	-	4	4
Female	32	-	-	-	-	-	-	-	1	2	-	2
Asian or Pacific Islander 2/, both sexes	90	-	-	-	1	-	-	2	1	1	7	1
Male	48	-	-	-	1	-	-	-	1	1	4	1
Female	42	-	-	-	-	-	-	2	-	-	3	-
Mississippi												
All races, both sexes	10,979	5	3	3	3	8	23	29	72	103	207	372
Male	5,178	4	1	3	2	5	10	16	44	60	128	230
Female	5,801	1	2	-	1	3	13	13	28	43	79	142
White, both sexes	7,361	1	2	-	2	-	5	12	27	37	100	171
Male	3,442	1	1	-	2	-	3	8	17	21	63	108
Female	3,919	-	1	-	-	-	2	4	10	16	37	63
Black, both sexes	3,570	4	1	3	1	8	18	16	45	66	105	199
Male	1,713	3	-	3	-	5	7	7	27	39	64	120
Female	1,857	1	1	-	1	3	11	9	18	27	41	79
American Indian 1/, both sexes	22	-	-	-	-	-	-	1	-	-	1	1
Male	13	-	-	-	-	-	-	1	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	16	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Missouri												
All races, both sexes	19,700	13	9	6	4	13	16	43	62	129	264	481
Male	9,154	7	5	2	2	7	13	27	36	85	181	317
Female	10,546	6	4	4	2	6	3	16	26	44	83	164
White, both sexes	17,665	10	7	5	4	8	12	29	45	96	201	380
Male	8,191	6	4	1	2	2	10	21	27	64	147	259
Female	9,474	4	3	4	2	6	2	8	18	32	54	121
Black, both sexes	1,957	3	1	1	-	5	4	14	17	32	63	99
Male	930	1	1	1	-	5	3	6	9	20	34	57
Female	1,027	2	-	-	-	-	1	8	8	12	29	42
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	52	-	1	-	-	-	-	-	-	1	-	2
Male	20	-	-	-	-	-	-	-	-	1	-	1
Female	32	-	1	-	-	-	-	-	-	-	-	1
Montana												
All races, both sexes	2,569	2	2	-	2	2	-	4	10	14	21	47
Male	1,292	1	2	-	2	-	-	4	5	9	16	36
Female	1,277	1	-	-	-	2	-	-	5	5	5	11
White, both sexes	2,458	2	2	-	2	2	-	3	9	9	17	40
Male	1,240	1	2	-	2	-	-	3	4	7	13	29
Female	1,218	1	-	-	-	2	-	-	5	2	4	11
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	100	-	-	-	-	-	-	1	1	4	4	7
Male	45	-	-	-	-	-	-	1	1	1	3	7
Female	55	-	-	-	-	-	-	-	-	3	1	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Nebraska												
All races, both sexes	5,102	2	-	-	-	1	6	5	16	20	51	80
Male	2,393	1	-	-	-	1	2	4	10	16	38	55
Female	2,709	1	-	-	-	-	4	1	6	4	13	25
White, both sexes	4,904	1	-	-	-	1	4	4	11	16	46	72
Male	2,288	-	-	-	-	1	1	3	8	12	34	52
Female	2,616	1	-	-	-	-	3	1	3	4	12	20
Black, both sexes	146	1	-	-	-	-	1	1	4	3	4	8
Male	84	1	-	-	-	-	1	1	2	3	3	3
Female	62	-	-	-	-	-	-	-	2	-	1	5
American Indian 1/, both sexes	31	-	-	-	-	-	1	-	1	1	1	-
Male	11	-	-	-	-	-	-	-	-	1	1	-
Female	20	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	6,491	3	2	1	2	6	11	27	25	55	113	206
Male	3,573	1	1	1	2	4	9	23	18	39	75	142
Female	2,918	2	1	-	-	2	2	4	7	16	38	64
White, both sexes	5,737	2	2	1	1	4	7	18	12	40	87	152
Male	3,161	1	1	1	1	3	5	17	9	27	58	110
Female	2,576	1	1	-	-	1	2	1	3	13	29	42
Black, both sexes	474	1	-	-	1	-	2	8	8	11	22	41
Male	253	-	-	-	1	-	2	5	6	8	14	22
Female	221	1	-	-	-	-	-	3	2	3	8	19
American Indian 1/, both sexes	64	-	-	-	-	-	1	-	3	1	1	3
Male	36	-	-	-	-	-	1	-	2	1	1	2
Female	28	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	216	-	-	-	-	2	1	1	2	3	3	10
Male	123	-	-	-	-	1	1	1	1	3	2	8
Female	93	-	-	-	-	1	-	-	1	-	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
New Hampshire												
All races, both sexes	3,302	1	-	-	-	3	2	6	7	18	32	56
Male	1,535	-	-	-	-	1	2	5	6	14	24	46
Female	1,767	1	-	-	-	2	-	1	1	4	8	10
White, both sexes	3,286	1	-	-	-	3	2	5	7	18	32	55
Male	1,525	-	-	-	-	1	2	4	6	14	24	46
Female	1,761	1	-	-	-	2	-	1	1	4	8	9
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	1	-	-	-	1
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
New Jersey												
All races, both sexes	26,035	11	7	-	7	15	25	39	60	116	263	484
Male	11,854	7	5	-	5	9	18	26	45	80	172	337
Female	14,181	4	2	-	2	6	7	13	15	36	91	147
White, both sexes	22,534	9	5	-	3	12	18	22	30	71	176	329
Male	10,217	6	3	-	2	8	13	15	26	51	123	244
Female	12,317	3	2	-	1	4	5	7	4	20	53	85
Black, both sexes	3,017	1	2	-	3	3	7	14	26	43	80	139
Male	1,374	1	2	-	2	1	5	8	15	28	42	82
Female	1,643	-	-	-	1	2	2	6	11	15	38	57
American Indian 1/, both sexes	36	-	-	-	-	-	-	-	-	-	-	3
Male	15	-	-	-	-	-	-	-	-	-	-	2
Female	21	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	448	1	-	-	1	-	-	3	4	2	7	13
Male	248	-	-	-	1	-	-	3	4	1	7	9
Female	200	1	-	-	-	-	-	-	-	1	-	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
New Mexico												
All races, both sexes	4,511	7	-	1	1	-	-	8	14	16	58	104
Male	2,263	2	-	-	1	-	-	2	8	10	39	72
Female	2,248	5	-	1	-	-	-	6	6	6	19	32
White, both sexes	4,194	3	-	1	1	-	-	7	9	14	45	89
Male	2,088	1	-	-	1	-	-	2	4	8	28	65
Female	2,106	2	-	1	-	-	-	5	5	6	17	24
Black, both sexes	86	-	-	-	-	-	-	1	-	-	3	3
Male	43	-	-	-	-	-	-	-	-	-	2	1
Female	43	-	-	-	-	-	-	1	-	-	1	2
American Indian 1/, both sexes	215	4	-	-	-	-	-	-	5	2	10	11
Male	124	1	-	-	-	-	-	-	4	2	9	5
Female	91	3	-	-	-	-	-	-	1	-	1	6
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	62,512	28	11	9	8	18	42	81	114	235	570	977
Male	27,692	18	6	6	4	12	31	56	70	152	364	698
Female	34,820	10	5	3	4	6	11	25	44	83	206	279
White, both sexes	52,715	15	5	5	6	11	27	43	72	158	364	653
Male	23,291	10	2	4	3	8	19	32	46	101	245	493
Female	29,424	5	3	1	3	3	8	11	26	57	119	160
Black, both sexes	8,418	12	4	2	1	6	15	35	36	72	181	286
Male	3,672	8	3	1	1	4	12	23	20	48	102	172
Female	4,746	4	1	1	-	2	3	12	16	24	79	114
American Indian 1/, both sexes	58	-	-	-	-	-	-	-	-	-	2	-
Male	31	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1,321	1	2	2	1	1	-	3	6	5	23	38
Male	698	-	1	1	-	-	-	1	4	3	16	33
Female	623	1	1	1	1	1	-	2	2	2	7	5

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
North Carolina												
All races, both sexes	24,531	20	13	9	4	17	34	40	107	174	369	655
Male	11,691	9	9	6	3	7	24	21	62	105	245	454
Female	12,840	11	4	3	1	10	10	19	45	69	124	201
White, both sexes	18,979	6	7	3	2	14	17	21	60	99	212	404
Male	9,108	1	5	2	2	6	14	11	37	59	147	292
Female	9,871	5	2	1	-	8	3	10	23	40	65	112
Black, both sexes	5,234	14	6	6	2	3	15	17	43	73	146	241
Male	2,419	8	4	4	1	1	9	9	22	44	91	155
Female	2,815	6	2	2	1	2	6	8	21	29	55	86
American Indian 1/, both sexes	221	-	-	-	-	-	-	1	-	-	7	8
Male	110	-	-	-	-	-	-	-	-	-	4	6
Female	111	-	-	-	-	-	-	1	-	-	3	2
Asian or Pacific Islander 2/, both sexes	97	-	-	-	-	-	2	1	4	2	4	2
Male	54	-	-	-	-	-	1	1	3	2	3	1
Female	43	-	-	-	-	-	1	-	1	-	1	1
North Dakota												
All races, both sexes	2,057	-	-	-	-	1	4	1	6	8	19	22
Male	993	-	-	-	-	1	2	1	4	7	15	16
Female	1,064	-	-	-	-	-	2	-	2	1	4	6
White, both sexes	2,006	-	-	-	-	1	4	1	6	7	16	16
Male	963	-	-	-	-	1	2	1	4	6	13	13
Female	1,043	-	-	-	-	-	2	-	2	1	3	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	47	-	-	-	-	-	-	-	-	1	2	6
Male	28	-	-	-	-	-	-	-	-	1	1	3
Female	19	-	-	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Ohio												
All races, both sexes	38,387	25	11	9	8	13	36	56	123	239	479	902
Male	17,952	16	9	5	7	11	23	35	84	150	326	595
Female	20,435	9	2	4	1	2	13	21	39	89	153	307
White, both sexes	34,220	16	9	6	6	11	22	39	83	171	370	733
Male	15,944	12	8	4	6	9	16	25	60	111	267	493
Female	18,276	4	1	2	-	2	6	14	23	60	103	240
Black, both sexes	4,051	9	2	2	2	2	13	17	39	67	107	165
Male	1,953	4	1	1	1	2	6	10	23	38	58	99
Female	2,098	5	1	1	1	-	7	7	16	29	49	66
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	102	-	-	1	-	-	1	-	1	1	2	2
Male	50	-	-	-	-	-	1	-	1	1	1	2
Female	52	-	-	1	-	-	-	-	-	-	1	-
Oklahoma												
All races, both sexes	13,235	8	6	2	2	14	18	19	37	58	180	304
Male	6,196	4	3	1	1	7	11	12	21	39	127	197
Female	7,039	4	3	1	1	7	7	7	16	19	53	107
White, both sexes	11,707	3	5	2	2	10	13	13	29	35	136	226
Male	5,446	2	2	1	1	5	8	7	16	23	94	153
Female	6,261	1	3	1	1	5	5	6	13	12	42	73
Black, both sexes	828	3	1	-	-	3	2	3	4	12	24	40
Male	384	2	1	-	-	1	1	2	2	9	16	24
Female	444	1	-	-	-	2	1	1	2	3	8	16
American Indian 1/, both sexes	650	1	-	-	-	1	3	3	4	11	19	34
Male	342	-	-	-	-	1	2	3	3	7	16	18
Female	308	1	-	-	-	-	1	-	1	4	3	16
Asian or Pacific Islander 2/, both sexes	50	1	-	-	-	-	-	-	-	-	1	4
Male	24	-	-	-	-	-	-	-	-	-	1	2
Female	26	1	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Oregon												
All races, both sexes	10,088	3	-	2	4	5	12	8	20	50	89	158
Male	4,834	-	-	1	-	3	11	5	15	36	59	108
Female	5,254	3	-	1	4	2	1	3	5	14	30	50
White, both sexes	9,763	3	-	2	2	4	11	6	18	42	85	146
Male	4,663	-	-	1	-	2	10	3	14	31	56	99
Female	5,100	3	-	1	2	2	1	3	4	11	29	47
Black, both sexes	135	-	-	-	1	1	1	2	2	2	3	3
Male	71	-	-	-	-	1	1	2	1	2	2	3
Female	64	-	-	-	1	-	-	-	1	-	1	-
American Indian 1/, both sexes	70	-	-	-	-	-	-	-	-	2	1	5
Male	39	-	-	-	-	-	-	-	-	1	1	4
Female	31	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	120	-	-	-	1	-	-	-	-	4	-	4
Male	61	-	-	-	-	-	-	-	-	2	-	2
Female	59	-	-	-	1	-	-	-	-	2	-	2
Pennsylvania												
All races, both sexes	47,029	23	6	8	5	21	35	52	90	198	440	862
Male	21,403	12	4	5	3	15	22	34	56	143	285	582
Female	25,626	11	2	3	2	6	13	18	34	55	155	280
White, both sexes	42,887	19	3	5	5	14	25	42	69	145	338	669
Male	19,424	9	3	3	3	10	17	28	46	111	227	477
Female	23,463	10	-	2	2	4	8	14	23	34	111	192
Black, both sexes	3,940	3	3	3	-	7	10	9	18	51	98	187
Male	1,871	2	1	2	-	5	5	5	7	30	55	102
Female	2,069	1	2	1	-	2	5	4	11	21	43	85
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	1	1	-	-
Male	7	-	-	-	-	-	-	-	1	1	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	184	1	-	-	-	-	-	1	2	1	4	6
Male	101	1	-	-	-	-	-	1	2	1	3	3
Female	83	-	-	-	-	-	-	-	-	-	1	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Rhode Island												
All races, both sexes	3,773	3	2	-	3	1	1	6	9	19	40	64
Male	1,655	1	1	-	2	1	-	5	4	11	26	42
Female	2,118	2	1	-	1	-	1	1	5	8	14	22
White, both sexes	3,627	2	2	-	3	1	1	4	7	17	36	54
Male	1,575	1	1	-	2	1	-	4	3	10	24	36
Female	2,052	1	1	-	1	-	1	-	4	7	12	18
Black, both sexes	113	1	-	-	-	-	-	2	2	2	3	6
Male	64	-	-	-	-	-	-	1	1	1	2	5
Female	49	1	-	-	-	-	-	1	1	1	1	1
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	1
South Carolina												
All races, both sexes	12,809	11	3	2	5	12	15	39	64	104	240	406
Male	6,363	4	1	1	3	8	10	25	44	68	153	281
Female	6,446	7	2	1	2	4	5	14	20	36	87	125
White, both sexes	9,124	6	1	1	2	5	8	17	25	60	136	234
Male	4,537	2	-	-	1	4	6	10	16	39	94	160
Female	4,587	4	1	1	1	1	2	7	9	21	42	74
Black, both sexes	3,642	5	2	1	3	7	7	21	38	44	103	172
Male	1,803	2	1	1	2	4	4	14	27	29	58	121
Female	1,839	3	1	-	1	3	3	7	11	15	45	51
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	-	-	-	-	-	-	1	-	-	1	-
Male	17	-	-	-	-	-	-	1	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
South Dakota												
All races, both sexes	2,500	1	-	-	1	1	-	1	8	7	19	37
Male	1,237	-	-	-	-	-	-	1	3	4	13	26
Female	1,263	1	-	-	1	1	-	-	5	3	6	11
White, both sexes	2,379	1	-	-	-	1	-	1	7	7	15	30
Male	1,166	-	-	-	-	-	-	1	3	4	10	20
Female	1,213	1	-	-	-	1	-	-	4	3	5	10
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	111	-	-	-	1	-	-	-	1	-	3	6
Male	65	-	-	-	-	-	-	-	-	-	2	5
Female	46	-	-	-	1	-	-	-	1	-	1	1
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	20,091	12	4	4	2	13	26	51	88	162	358	607
Male	9,498	8	1	2	1	7	17	36	64	107	240	400
Female	10,593	4	3	2	1	6	9	15	24	55	118	207
White, both sexes	16,997	8	4	1	1	7	16	31	57	121	262	444
Male	8,005	6	1	1	-	4	10	22	41	85	181	295
Female	8,992	2	3	-	1	3	6	9	16	36	81	149
Black, both sexes	3,021	4	-	3	1	6	10	20	29	41	95	161
Male	1,455	2	-	1	1	3	7	14	22	22	58	104
Female	1,566	2	-	2	-	3	3	6	7	19	37	57
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	58	-	-	-	-	-	-	-	2	-	1	2
Male	29	-	-	-	-	-	-	-	1	-	1	1
Female	29	-	-	-	-	-	-	-	1	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Texas												
All races, both sexes	53,277	42	17	12	22	45	74	115	213	387	846	1,415
Male	25,843	25	10	7	13	30	49	78	138	261	578	984
Female	27,434	17	7	5	9	15	25	37	75	126	268	431
White, both sexes	46,001	29	13	11	16	34	51	84	142	269	614	1,042
Male	22,293	16	8	6	11	20	33	62	91	186	435	756
Female	23,708	13	5	5	5	14	18	22	51	83	179	286
Black, both sexes	6,693	12	4	-	6	11	21	29	62	108	218	348
Male	3,253	8	2	-	2	10	15	15	40	66	133	210
Female	3,440	4	2	-	4	1	6	14	22	42	85	138
American Indian 1/, both sexes	47	-	-	-	-	-	1	1	1	1	1	5
Male	25	-	-	-	-	-	-	-	1	1	1	4
Female	22	-	-	-	-	-	1	1	-	-	-	1
Asian or Pacific Islander 2/, both sexes	536	1	-	1	-	-	1	1	8	9	13	20
Male	272	1	-	1	-	-	1	1	6	8	9	14
Female	264	-	-	-	-	-	-	-	2	1	4	6
Utah												
All races, both sexes	3,956	3	3	2	-	3	6	8	11	27	38	81
Male	1,876	3	2	1	-	3	5	2	6	20	26	59
Female	2,080	-	1	1	-	-	1	6	5	7	12	22
White, both sexes	3,845	3	2	2	-	2	6	8	11	25	33	75
Male	1,810	3	1	1	-	2	5	2	6	19	22	53
Female	2,035	-	1	1	-	-	1	6	5	6	11	22
Black, both sexes	26	-	-	-	-	-	-	-	-	-	3	-
Male	12	-	-	-	-	-	-	-	-	-	2	-
Female	14	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	26	-	1	-	-	-	-	-	-	-	-	3
Male	16	-	1	-	-	-	-	-	-	-	-	3
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	59	-	-	-	-	1	-	-	-	2	2	3
Male	38	-	-	-	-	1	-	-	-	1	2	3
Female	21	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Vermont												
All races, both sexes	1,632	1	-	-	-	1	2	1	2	7	16	35
Male	785	1	-	-	-	1	1	1	2	3	12	25
Female	847	-	-	-	-	-	1	-	-	4	4	10
White, both sexes	1,629	1	-	-	-	1	2	1	2	7	16	35
Male	784	1	-	-	-	1	1	1	2	3	12	25
Female	845	-	-	-	-	-	1	-	-	4	4	10
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	19,267	13	5	9	5	10	29	39	79	139	278	494
Male	9,095	7	2	6	2	4	22	20	52	87	193	324
Female	10,172	6	3	3	3	6	7	19	27	52	85	170
White, both sexes	15,022	6	1	3	4	7	16	22	42	80	176	301
Male	7,091	4	-	3	2	3	12	12	27	51	134	215
Female	7,931	2	1	-	2	4	4	10	15	29	42	86
Black, both sexes	3,963	7	4	6	1	3	11	14	31	55	97	182
Male	1,856	3	2	3	-	1	8	7	20	33	55	101
Female	2,107	4	2	3	1	2	3	7	11	22	42	81
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	1	-	1	1
Male	7	-	-	-	-	-	-	-	1	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	269	-	-	-	-	-	2	3	5	4	4	10
Male	141	-	-	-	-	-	2	1	4	3	3	8
Female	128	-	-	-	-	-	-	2	1	1	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Washington												
All races, both sexes	15,273	7	1	2	5	9	14	17	40	79	143	274
Male	7,502	3	-	-	3	6	10	9	26	51	99	178
Female	7,771	4	1	2	2	3	4	8	14	28	44	96
White, both sexes	14,310	4	1	1	3	5	8	15	33	61	120	237
Male	6,980	2	-	-	1	4	7	7	21	38	85	154
Female	7,330	2	1	1	2	1	1	8	12	23	35	83
Black, both sexes	359	-	-	1	1	1	2	1	1	8	11	18
Male	201	-	-	-	1	-	1	1	1	7	7	10
Female	158	-	-	1	-	1	1	-	-	1	4	8
American Indian 1/, both sexes	151	1	-	-	-	2	1	-	1	4	4	7
Male	84	-	-	-	-	1	1	-	1	2	2	5
Female	67	1	-	-	-	1	-	-	-	2	2	2
Asian or Pacific Islander 2/, both sexes	453	2	-	-	1	1	3	1	5	6	8	12
Male	237	1	-	-	1	1	1	1	3	4	5	9
Female	216	1	-	-	-	-	2	-	2	2	3	3
West Virginia												
All races, both sexes	7,300	1	2	-	1	2	5	7	18	33	79	166
Male	3,409	1	1	-	-	2	2	3	15	21	61	112
Female	3,891	-	1	-	1	-	3	4	3	12	18	54
White, both sexes	7,080	1	2	-	1	2	5	6	15	32	76	156
Male	3,321	1	1	-	-	2	2	2	12	20	58	106
Female	3,759	-	1	-	1	-	3	4	3	12	18	50
Black, both sexes	216	-	-	-	-	-	-	1	3	1	3	10
Male	87	-	-	-	-	-	-	1	3	1	3	6
Female	129	-	-	-	-	-	-	-	-	-	-	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
IX. Diseases of the circulatory system (I00-I99)												
Wisconsin												
All races, both sexes	16,048	6	-	2	3	7	13	16	28	63	149	268
Male	7,601	6	-	1	1	5	7	11	19	44	106	190
Female	8,447	-	-	1	2	2	6	5	9	19	43	78
White, both sexes	15,331	5	-	2	3	6	11	14	17	53	124	228
Male	7,212	5	-	1	1	4	6	10	13	38	90	163
Female	8,119	-	-	1	2	2	5	4	4	15	34	65
Black, both sexes	551	1	-	-	-	1	2	1	9	10	19	33
Male	298	1	-	-	-	1	1	-	5	6	12	22
Female	253	-	-	-	-	-	1	1	4	4	7	11
American Indian 1/, both sexes	93	-	-	-	-	-	-	-	2	-	3	4
Male	54	-	-	-	-	-	-	-	1	-	1	3
Female	39	-	-	-	-	-	-	-	1	-	2	1
Asian or Pacific Islander 2/, both sexes	73	-	-	-	-	-	-	1	-	-	3	3
Male	37	-	-	-	-	-	-	1	-	-	3	2
Female	36	-	-	-	-	-	-	-	-	-	-	1
Wyoming												
All races, both sexes	1,260	1	-	-	2	-	-	-	3	10	15	36
Male	627	1	-	-	1	-	-	-	1	6	9	23
Female	633	-	-	-	1	-	-	-	2	4	6	13
White, both sexes	1,237	1	-	-	2	-	-	-	3	9	14	34
Male	618	1	-	-	1	-	-	-	1	5	8	21
Female	619	-	-	-	1	-	-	-	2	4	6	13
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	1	1	1
Male	7	-	-	-	-	-	-	-	-	1	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
United States												
All races, both sexes	28,313	36,470	44,067	52,531	72,548	108,187	149,303	154,355	115,829	46,857	10,199	41
Male	19,772	25,032	28,833	32,291	41,821	56,957	68,526	59,901	34,891	10,111	1,511	29
Female	8,541	11,438	15,234	20,240	30,727	51,230	80,777	94,454	80,938	36,746	8,688	12
White, both sexes	20,596	27,730	34,731	42,205	60,423	93,465	133,226	141,110	106,452	42,813	8,929	27
Male	14,907	19,500	23,243	26,440	35,443	50,113	61,866	55,309	32,299	9,221	1,301	19
Female	5,689	8,230	11,488	15,765	24,980	43,352	71,360	85,801	74,153	33,592	7,628	8
Black, both sexes	6,870	7,698	8,110	8,771	10,187	12,290	13,125	10,646	7,548	3,305	1,105	13
Male	4,278	4,823	4,843	4,900	5,304	5,639	5,250	3,486	1,897	625	159	9
Female	2,592	2,875	3,267	3,871	4,883	6,651	7,875	7,160	5,651	2,680	946	4
American Indian 1/, both sexes	240	289	317	354	407	435	410	337	230	87	31	-
Male	151	192	197	195	226	213	178	132	72	20	6	-
Female	89	97	120	159	181	222	232	205	158	67	25	-
Asian or Pacific Islander 2/, both sexes	607	753	909	1,201	1,531	1,997	2,542	2,262	1,599	652	134	1
Male	436	517	550	756	848	992	1,232	974	623	245	45	1
Female	171	236	359	445	683	1,005	1,310	1,288	976	407	89	-
Alabama												
All races, both sexes	707	938	1,118	1,205	1,597	2,220	2,807	2,537	1,917	681	142	-
Male	432	607	697	675	886	1,096	1,153	911	489	138	22	-
Female	275	331	421	530	711	1,124	1,654	1,626	1,428	543	120	-
White, both sexes	457	608	775	866	1,213	1,698	2,257	2,137	1,589	520	84	-
Male	297	414	499	501	680	872	957	759	410	105	11	-
Female	160	194	276	365	533	826	1,300	1,378	1,179	415	73	-
Black, both sexes	246	325	341	336	375	513	543	396	327	160	58	-
Male	133	191	196	172	204	220	193	151	79	32	11	-
Female	113	134	145	164	171	293	350	245	248	128	47	-
American Indian 1/, both sexes	2	4	2	1	4	4	6	1	-	-	-	-
Male	-	2	2	1	-	3	2	-	-	-	-	-
Female	2	2	-	-	4	1	4	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	2	5	5	1	3	1	1	-	-
Male	2	-	-	1	2	1	1	1	-	1	-	-
Female	-	1	-	1	3	4	-	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Alaska												
All races, both sexes	53	62	71	65	80	119	129	107	54	24	8	-
Male	37	50	50	43	57	68	63	52	16	6	3	-
Female	16	12	21	22	23	51	66	55	38	18	5	-
White, both sexes	38	51	52	46	47	86	107	79	42	17	8	-
Male	26	42	39	32	35	51	56	41	12	5	3	-
Female	12	9	13	14	12	35	51	38	30	12	5	-
Black, both sexes	3	3	4	2	1	2	2	4	1	-	-	-
Male	3	2	1	2	1	1	1	3	-	-	-	-
Female	-	1	3	-	-	1	1	1	1	-	-	-
American Indian 1/, both sexes	12	7	13	16	27	29	13	20	10	7	-	-
Male	8	6	8	9	19	15	4	6	4	1	-	-
Female	4	1	5	7	8	14	9	14	6	6	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	5	2	7	4	1	-	-	-
Male	-	-	2	-	2	1	2	2	-	-	-	-
Female	-	1	-	1	3	1	5	2	1	-	-	-
Arizona												
All races, both sexes	467	604	818	952	1,350	1,941	2,548	2,569	1,797	643	117	4
Male	339	444	548	588	824	1,060	1,235	1,115	630	174	22	2
Female	128	160	270	364	526	881	1,313	1,454	1,167	469	95	2
White, both sexes	406	541	769	862	1,266	1,837	2,456	2,494	1,750	623	110	4
Male	299	402	524	527	782	1,017	1,196	1,082	616	169	19	2
Female	107	139	245	335	484	820	1,260	1,412	1,134	454	91	2
Black, both sexes	29	25	21	38	37	42	43	33	21	8	3	-
Male	21	17	10	30	18	17	18	11	4	1	1	-
Female	8	8	11	8	19	25	25	22	17	7	2	-
American Indian 1/, both sexes	27	32	22	43	40	41	33	23	19	7	3	-
Male	15	23	12	25	20	18	13	12	7	2	2	-
Female	12	9	10	18	20	23	20	11	12	5	1	-
Asian or Pacific Islander 2/, both sexes	5	6	6	9	7	21	16	19	7	5	1	-
Male	4	2	2	6	4	8	8	10	3	2	-	-
Female	1	4	4	3	3	13	8	9	4	3	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Arkansas												
All races, both sexes	388	502	588	732	887	1,255	1,620	1,673	1,267	517	94	-
Male	273	343	370	446	546	625	747	602	363	104	16	-
Female	115	159	218	286	341	630	873	1,071	904	413	78	-
White, both sexes	289	396	466	619	772	1,100	1,429	1,517	1,120	448	68	-
Male	212	276	309	384	485	570	671	549	323	91	11	-
Female	77	120	157	235	287	530	758	968	797	357	57	-
Black, both sexes	97	105	120	112	111	150	183	152	145	67	26	-
Male	60	66	61	62	59	55	71	53	40	13	5	-
Female	37	39	59	50	52	95	112	99	105	54	21	-
American Indian 1/, both sexes	2	-	2	1	1	2	4	2	1	1	-	-
Male	1	-	-	-	-	-	2	-	-	-	-	-
Female	1	-	2	1	1	2	2	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	3	3	4	2	1	1	-	-
Male	-	1	-	-	2	-	3	-	-	-	-	-
Female	-	-	-	-	1	3	1	2	1	1	-	-
California												
All races, both sexes	2,717	3,594	4,217	5,035	6,947	10,573	15,614	16,115	12,342	5,083	1,202	2
Male	1,926	2,480	2,765	3,050	4,028	5,546	7,354	6,582	4,105	1,214	200	2
Female	791	1,114	1,452	1,985	2,919	5,027	8,260	9,533	8,237	3,869	1,002	-
White, both sexes	2,003	2,706	3,305	3,911	5,521	8,804	13,429	14,254	11,034	4,533	1,056	1
Male	1,457	1,907	2,197	2,392	3,254	4,701	6,351	5,847	3,629	1,062	167	1
Female	546	799	1,108	1,519	2,267	4,103	7,078	8,407	7,405	3,471	889	-
Black, both sexes	467	575	548	634	727	839	946	759	520	242	85	1
Male	296	364	348	350	402	392	413	253	145	38	11	1
Female	171	211	200	284	325	447	533	506	375	204	74	-
American Indian 1/, both sexes	17	40	33	45	45	47	38	41	23	8	1	-
Male	8	25	18	26	18	20	17	18	8	-	-	-
Female	9	15	15	19	27	27	21	23	15	8	1	-
Asian or Pacific Islander 2/, both sexes	230	273	331	445	654	883	1,201	1,061	765	300	60	-
Male	165	184	202	282	354	433	573	464	323	114	22	-
Female	65	89	129	163	300	450	628	597	442	186	38	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Colorado												
All races, both sexes	310	354	441	520	709	1,118	1,536	1,690	1,190	515	104	-
Male	225	262	284	340	424	591	743	703	365	97	17	-
Female	85	92	157	180	285	527	793	987	825	418	87	-
White, both sexes	284	322	403	482	655	1,051	1,484	1,646	1,161	500	103	-
Male	210	238	258	315	398	559	718	691	361	97	16	-
Female	74	84	145	167	257	492	766	955	800	403	87	-
Black, both sexes	24	23	29	29	37	46	32	35	24	12	1	-
Male	14	18	20	19	18	21	16	8	4	-	1	-
Female	10	5	9	10	19	25	16	27	20	12	-	-
American Indian 1/, both sexes	-	4	2	2	8	4	3	1	-	3	-	-
Male	-	3	1	1	5	1	3	-	-	-	-	-
Female	-	1	1	1	3	3	-	1	-	3	-	-
Asian or Pacific Islander 2/, both sexes	2	5	7	7	9	17	17	8	5	-	-	-
Male	1	3	5	5	3	10	6	4	-	-	-	-
Female	1	2	2	2	6	7	11	4	5	-	-	-
Connecticut												
All races, both sexes	266	308	393	461	709	1,119	1,729	2,052	1,675	725	154	2
Male	196	224	265	294	427	591	829	786	511	137	18	2
Female	70	84	128	167	282	528	900	1,266	1,164	588	136	-
White, both sexes	219	255	331	385	634	1,041	1,645	1,978	1,636	698	143	2
Male	167	187	227	252	386	556	795	761	500	130	18	2
Female	52	68	104	133	248	485	850	1,217	1,136	568	125	-
Black, both sexes	45	47	53	70	65	70	73	67	38	22	8	-
Male	27	33	34	36	38	31	31	24	11	3	-	-
Female	18	14	19	34	27	39	42	43	27	19	8	-
American Indian 1/, both sexes	-	-	1	1	2	2	1	2	-	1	1	-
Male	-	-	1	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	2	-	2	-	1	1	-
Asian or Pacific Islander 2/, both sexes	2	6	8	5	8	6	10	5	1	4	2	-
Male	2	4	3	5	2	4	2	1	-	4	-	-
Female	-	2	5	-	6	2	8	4	1	-	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Delaware												
All races, both sexes	75	112	153	176	200	348	452	465	325	135	22	-
Male	51	74	102	103	113	179	211	195	94	29	3	-
Female	24	38	51	73	87	169	241	270	231	106	19	-
White, both sexes	50	85	107	133	158	298	400	408	286	118	17	-
Male	38	60	78	81	89	159	189	177	83	27	3	-
Female	12	25	29	52	69	139	211	231	203	91	14	-
Black, both sexes	24	26	44	39	38	47	50	53	36	17	5	-
Male	13	14	22	19	21	18	21	16	9	2	-	-
Female	11	12	22	20	17	29	29	37	27	15	5	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	2	1	-	-	-
Male	-	-	-	-	1	-	-	1	1	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	4	3	3	2	2	2	-	-	-
Male	-	-	2	3	2	2	1	1	1	-	-	-
Female	-	1	-	1	1	1	1	1	1	-	-	-
Dist. of Columbia												
All races, both sexes	109	100	133	140	164	232	300	294	188	89	31	2
Male	77	58	93	84	93	113	119	101	49	19	6	1
Female	32	42	40	56	71	119	181	193	139	70	25	1
White, both sexes	10	9	19	8	26	41	70	73	54	37	11	1
Male	7	6	15	5	14	24	33	35	21	9	2	-
Female	3	3	4	3	12	17	37	38	33	28	9	1
Black, both sexes	97	89	114	130	136	189	223	217	131	50	20	1
Male	68	50	78	78	77	87	81	63	28	9	4	1
Female	29	39	36	52	59	102	142	154	103	41	16	-
American Indian 1/, both sexes	-	-	-	1	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	-	1	2	2	5	4	3	2	-	-
Male	2	2	-	1	2	2	4	3	-	1	-	-
Female	-	-	-	-	-	-	1	1	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Florida												
All races, both sexes	1,708	2,094	2,871	3,552	4,901	7,759	10,922	11,281	8,591	3,443	689	4
Male	1,211	1,431	1,977	2,269	2,936	4,307	5,376	4,951	3,001	913	137	3
Female	497	663	894	1,283	1,965	3,452	5,546	6,330	5,590	2,530	552	1
White, both sexes	1,265	1,650	2,343	2,924	4,197	6,973	10,056	10,584	8,085	3,236	628	-
Male	934	1,166	1,655	1,924	2,560	3,932	5,020	4,719	2,864	860	124	-
Female	331	484	688	1,000	1,637	3,041	5,036	5,865	5,221	2,376	504	-
Black, both sexes	420	420	499	588	663	742	808	648	485	199	59	4
Male	263	250	304	323	359	354	327	218	125	51	13	3
Female	157	170	195	265	304	388	481	430	360	148	46	1
American Indian 1/, both sexes	5	3	8	4	2	5	8	8	4	-	1	-
Male	3	2	6	1	1	2	6	4	2	-	-	-
Female	2	1	2	3	1	3	2	4	2	-	1	-
Asian or Pacific Islander 2/, both sexes	18	21	21	36	39	39	50	41	17	8	1	-
Male	11	13	12	21	16	19	23	10	10	2	-	-
Female	7	8	9	15	23	20	27	31	7	6	1	-
Georgia												
All races, both sexes	1,076	1,310	1,553	1,729	2,159	2,815	3,456	3,295	2,358	847	205	2
Male	751	876	992	1,064	1,214	1,404	1,413	1,157	582	168	28	2
Female	325	434	561	665	945	1,411	2,043	2,138	1,776	679	177	-
White, both sexes	591	788	1,028	1,117	1,545	2,152	2,741	2,688	1,902	646	132	2
Male	427	545	691	716	897	1,130	1,156	991	472	123	19	2
Female	164	243	337	401	648	1,022	1,585	1,697	1,430	523	113	-
Black, both sexes	478	507	514	587	590	644	695	600	442	198	73	-
Male	319	322	295	331	305	264	249	164	106	44	9	-
Female	159	185	219	256	285	380	446	436	336	154	64	-
American Indian 1/, both sexes	1	2	4	5	6	4	4	1	8	1	-	-
Male	1	2	1	4	2	3	1	-	3	-	-	-
Female	-	-	3	1	4	1	3	1	5	1	-	-
Asian or Pacific Islander 2/, both sexes	6	13	7	20	18	15	16	6	6	2	-	-
Male	4	7	5	13	10	7	7	2	1	1	-	-
Female	2	6	2	7	8	8	9	4	5	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Hawaii												
All races, both sexes	127	144	194	208	260	390	535	540	426	218	30	-
Male	92	114	125	132	154	204	304	250	165	80	7	-
Female	35	30	69	76	106	186	231	290	261	138	23	-
White, both sexes	29	34	60	46	71	103	134	138	105	55	5	-
Male	24	30	45	31	40	53	77	63	39	13	2	-
Female	5	4	15	15	31	50	57	75	66	42	3	-
Black, both sexes	3	1	2	1	1	2	5	1	1	-	-	-
Male	3	1	2	-	1	1	3	-	1	-	-	-
Female	-	-	-	1	-	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	95	109	132	161	188	285	396	398	320	163	25	-
Male	65	83	78	101	113	150	224	186	125	67	5	-
Female	30	26	54	60	75	135	172	212	195	96	20	-
Idaho												
All races, both sexes	99	114	167	216	262	419	599	672	518	185	24	-
Male	71	81	112	139	160	237	302	281	182	42	5	-
Female	28	33	55	77	102	182	297	391	336	143	19	-
White, both sexes	94	112	163	210	260	412	591	667	516	185	24	-
Male	68	79	110	136	158	233	297	279	181	42	5	-
Female	26	33	53	74	102	179	294	388	335	143	19	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	2	3	1	1	6	5	2	-	-	-	-
Male	1	2	2	1	1	3	4	-	-	-	-	-
Female	2	-	1	-	-	3	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	5	1	1	3	3	2	-	-	-
Male	2	-	-	2	1	1	1	2	1	-	-	-
Female	-	-	1	3	-	-	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Illinois												
All races, both sexes	1,277	1,527	1,872	2,193	2,974	4,506	6,300	6,806	5,139	2,165	457	-
Male	910	1,053	1,249	1,349	1,661	2,417	2,903	2,546	1,436	401	62	-
Female	367	474	623	844	1,313	2,089	3,397	4,260	3,703	1,764	395	-
White, both sexes	852	1,065	1,387	1,653	2,354	3,734	5,517	6,178	4,738	1,988	402	-
Male	632	756	947	1,028	1,356	2,052	2,607	2,317	1,335	366	52	-
Female	220	309	440	625	998	1,682	2,910	3,861	3,403	1,622	350	-
Black, both sexes	413	437	450	491	572	713	708	550	365	159	52	-
Male	268	278	280	289	275	341	269	191	92	30	9	-
Female	145	159	170	202	297	372	439	359	273	129	43	-
American Indian 1/, both sexes	-	1	6	2	4	2	3	4	1	1	-	-
Male	-	1	2	1	1	1	-	2	-	-	-	-
Female	-	-	4	1	3	1	3	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes	12	24	29	47	44	57	72	74	35	17	3	-
Male	10	18	20	31	29	23	27	36	9	5	1	-
Female	2	6	9	16	15	34	45	38	26	12	2	-
Indiana												
All races, both sexes	645	798	1,023	1,247	1,725	2,479	3,275	3,448	2,489	1,007	202	-
Male	460	557	666	741	931	1,294	1,519	1,301	719	203	23	-
Female	185	241	357	506	794	1,185	1,756	2,147	1,770	804	179	-
White, both sexes	545	705	914	1,099	1,551	2,308	3,099	3,296	2,386	968	175	-
Male	396	497	599	659	852	1,213	1,448	1,237	689	194	20	-
Female	149	208	315	440	699	1,095	1,651	2,059	1,697	774	155	-
Black, both sexes	98	90	106	136	167	164	167	146	99	37	26	-
Male	64	58	66	74	77	77	67	62	27	8	2	-
Female	34	32	40	62	90	87	100	84	72	29	24	-
American Indian 1/, both sexes	1	-	1	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	1	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	2	11	7	7	8	5	4	2	1	-
Male	-	2	1	8	2	4	4	2	3	1	1	-
Female	1	1	1	3	5	3	4	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Iowa												
All races, both sexes	248	281	406	516	689	1,202	1,871	2,027	1,696	737	213	-
Male	174	201	285	327	415	653	908	850	539	157	29	-
Female	74	80	121	189	274	549	963	1,177	1,157	580	184	-
White, both sexes	232	268	391	505	676	1,191	1,852	2,012	1,692	734	212	-
Male	165	193	275	321	405	647	897	843	538	157	29	-
Female	67	75	116	184	271	544	955	1,169	1,154	577	183	-
Black, both sexes	15	11	14	7	8	9	15	12	4	3	1	-
Male	8	7	9	3	6	4	9	5	1	-	-	-
Female	7	4	5	4	2	5	6	7	3	3	1	-
American Indian 1/, both sexes	1	1	1	-	3	-	1	1	-	-	-	-
Male	1	1	1	-	2	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	4	2	2	3	2	-	-	-	-
Male	-	-	-	3	2	2	2	1	-	-	-	-
Female	-	1	-	1	-	-	1	1	-	-	-	-
Kansas												
All races, both sexes	227	306	356	401	633	951	1,412	1,678	1,311	599	139	-
Male	161	207	244	252	364	525	665	660	413	143	25	-
Female	66	99	112	149	269	426	747	1,018	898	456	114	-
White, both sexes	195	277	312	368	586	892	1,338	1,616	1,269	582	136	-
Male	140	183	214	233	342	490	631	639	392	137	25	-
Female	55	94	98	135	244	402	707	977	877	445	111	-
Black, both sexes	27	23	38	26	29	47	57	56	33	16	1	-
Male	19	18	26	14	13	28	28	20	16	5	-	-
Female	8	5	12	12	16	19	29	36	17	11	1	-
American Indian 1/, both sexes	4	5	5	4	12	9	11	5	8	1	2	-
Male	1	5	4	4	7	6	5	1	5	1	-	-
Female	3	-	1	-	5	3	6	4	3	-	2	-
Asian or Pacific Islander 2/, both sexes	1	1	1	3	6	3	6	1	1	-	-	-
Male	1	1	-	1	2	1	1	-	-	-	-	-
Female	-	-	1	2	4	2	5	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Kentucky												
All races, both sexes	550	689	815	979	1,385	1,813	2,284	2,252	1,576	580	118	-
Male	388	448	517	572	801	979	952	767	417	97	19	-
Female	162	241	298	407	584	834	1,332	1,485	1,159	483	99	-
White, both sexes	475	626	753	907	1,289	1,694	2,170	2,143	1,498	549	107	-
Male	340	412	483	535	752	914	919	736	402	94	17	-
Female	135	214	270	372	537	780	1,251	1,407	1,096	455	90	-
Black, both sexes	72	55	59	68	95	115	114	105	76	31	10	-
Male	46	30	31	34	49	62	33	30	15	3	1	-
Female	26	25	28	34	46	53	81	75	61	28	9	-
American Indian 1/, both sexes	2	2	-	1	-	-	-	-	-	-	-	-
Male	1	2	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	6	3	3	1	4	-	4	2	-	1	-
Male	1	4	3	2	-	3	-	1	-	-	1	-
Female	-	2	-	1	1	1	-	3	2	-	-	-
Louisiana												
All races, both sexes	659	859	947	1,077	1,381	1,895	2,287	2,173	1,517	565	136	1
Male	427	571	588	643	777	944	996	821	417	116	28	1
Female	232	288	359	434	604	951	1,291	1,352	1,100	449	108	-
White, both sexes	369	458	593	684	905	1,396	1,749	1,751	1,204	419	97	-
Male	254	319	392	425	517	737	775	690	330	74	18	-
Female	115	139	201	259	388	659	974	1,061	874	345	79	-
Black, both sexes	285	393	349	383	467	487	527	415	309	144	39	1
Male	172	246	192	210	256	200	216	126	87	42	10	1
Female	113	147	157	173	211	287	311	289	222	102	29	-
American Indian 1/, both sexes	2	2	1	3	1	4	3	2	2	1	-	-
Male	1	2	1	2	-	1	2	2	-	-	-	-
Female	1	-	-	1	1	3	1	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	3	6	4	7	8	8	8	5	2	1	-	-
Male	-	4	3	6	4	6	3	3	-	-	-	-
Female	3	2	1	1	4	2	5	2	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Maine												
All races, both sexes	116	148	198	227	346	506	673	717	575	231	45	-
Male	86	113	126	148	210	258	309	298	169	43	3	-
Female	30	35	72	79	136	248	364	419	406	188	42	-
White, both sexes	114	147	197	226	344	503	670	715	575	231	45	-
Male	85	112	125	148	209	258	308	297	169	43	3	-
Female	29	35	72	78	135	245	362	418	406	188	42	-
Black, both sexes	-	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	1	-	3	3	1	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	1	-	3	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	606	714	852	945	1,250	1,893	2,630	2,575	1,857	767	179	4
Male	405	479	552	579	683	924	1,191	986	499	165	27	2
Female	201	235	300	366	567	969	1,439	1,589	1,358	602	152	2
White, both sexes	305	427	505	570	839	1,420	2,077	2,133	1,563	647	139	2
Male	218	307	344	366	461	714	964	837	432	143	24	1
Female	87	120	161	204	378	706	1,113	1,296	1,131	504	115	1
Black, both sexes	289	270	326	351	388	453	517	410	275	111	39	2
Male	178	164	196	194	212	198	208	139	62	19	3	1
Female	111	106	130	157	176	255	309	271	213	92	36	1
American Indian 1/, both sexes	2	2	1	1	3	1	1	2	-	1	-	-
Male	2	2	1	1	-	-	-	-	-	1	-	-
Female	-	-	-	-	3	1	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	15	20	23	20	19	35	30	19	8	1	-
Male	7	6	11	18	10	12	19	10	5	2	-	-
Female	3	9	9	5	10	7	16	20	14	6	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Massachusetts												
All races, both sexes	411	577	687	906	1,327	2,076	3,059	3,592	2,793	1,178	275	-
Male	310	412	475	575	791	1,128	1,385	1,371	761	237	35	-
Female	101	165	212	331	536	948	1,674	2,221	2,032	941	240	-
White, both sexes	368	525	636	828	1,234	1,955	2,927	3,503	2,730	1,154	265	-
Male	281	381	443	536	748	1,067	1,331	1,335	749	231	35	-
Female	87	144	193	292	486	888	1,596	2,168	1,981	923	230	-
Black, both sexes	37	43	43	66	76	92	96	75	51	17	7	-
Male	24	25	28	35	33	46	33	30	8	3	-	-
Female	13	18	15	31	43	46	63	45	43	14	7	-
American Indian 1/, both sexes	1	-	2	2	4	4	2	-	3	-	1	-
Male	1	-	1	-	2	3	1	-	-	-	-	-
Female	-	-	1	2	2	1	1	-	3	-	1	-
Asian or Pacific Islander 2/, both sexes	5	9	6	10	13	25	34	14	9	7	2	-
Male	4	6	3	4	8	12	20	6	4	3	-	-
Female	1	3	3	6	5	13	14	8	5	4	2	-
Michigan												
All races, both sexes	1,189	1,548	1,698	1,922	2,680	4,159	5,643	5,905	4,268	1,687	342	-
Male	786	1,037	1,118	1,134	1,538	2,162	2,576	2,202	1,321	352	42	-
Female	403	511	580	788	1,142	1,997	3,067	3,703	2,947	1,335	300	-
White, both sexes	815	1,087	1,266	1,544	2,189	3,523	5,023	5,370	3,922	1,540	301	-
Male	570	755	843	913	1,275	1,864	2,314	2,006	1,220	317	38	-
Female	245	332	423	631	914	1,659	2,709	3,364	2,702	1,223	263	-
Black, both sexes	362	439	407	349	456	603	591	504	333	139	40	-
Male	209	267	260	202	242	281	245	187	96	33	4	-
Female	153	172	147	147	214	322	346	317	237	106	36	-
American Indian 1/, both sexes	6	15	12	15	20	15	7	13	4	3	-	-
Male	3	9	6	9	13	7	2	3	1	-	-	-
Female	3	6	6	6	7	8	5	10	3	3	-	-
Asian or Pacific Islander 2/, both sexes	6	7	13	14	15	18	22	18	9	5	1	-
Male	4	6	9	10	8	10	15	6	4	2	-	-
Female	2	1	4	4	7	8	7	12	5	3	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Minnesota												
All races, both sexes	309	416	429	585	815	1,297	1,947	2,338	1,930	810	150	-
Male	223	296	302	394	488	761	955	977	616	189	23	-
Female	86	120	127	191	327	536	992	1,361	1,314	621	127	-
White, both sexes	276	389	401	556	781	1,262	1,919	2,308	1,914	807	148	-
Male	198	276	284	379	468	739	945	965	610	189	22	-
Female	78	113	117	177	313	523	974	1,343	1,304	618	126	-
Black, both sexes	22	16	17	9	14	19	14	12	4	2	1	-
Male	18	9	13	4	10	12	3	7	2	-	-	-
Female	4	7	4	5	4	7	11	5	2	2	1	-
American Indian 1/, both sexes	5	6	6	13	10	4	5	4	5	-	-	-
Male	4	6	3	6	7	2	2	-	1	-	-	-
Female	1	-	3	7	3	2	3	4	4	-	-	-
Asian or Pacific Islander 2/, both sexes	6	5	5	7	10	12	9	14	7	1	1	-
Male	3	5	2	5	3	8	5	5	3	-	1	-
Female	3	-	3	2	7	4	4	9	4	1	-	-
Mississippi												
All races, both sexes	475	622	734	816	1,010	1,434	1,640	1,586	1,185	520	127	2
Male	309	424	472	492	579	719	716	541	306	94	21	2
Female	166	198	262	324	431	715	924	1,045	879	426	106	-
White, both sexes	244	339	449	511	661	1,016	1,237	1,219	886	377	63	2
Male	167	243	300	324	399	537	552	411	221	55	7	2
Female	77	96	149	187	262	479	685	808	665	322	56	-
Black, both sexes	229	278	280	300	343	411	399	362	297	141	64	-
Male	141	177	170	165	179	179	162	128	84	39	14	-
Female	88	101	110	135	164	232	237	234	213	102	50	-
American Indian 1/, both sexes	1	4	4	2	3	3	1	-	1	-	-	-
Male	1	3	2	1	1	2	-	-	1	-	-	-
Female	-	1	2	1	2	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	3	3	4	3	5	1	2	-	-
Male	-	1	-	2	-	1	2	2	-	-	-	-
Female	1	-	1	1	3	3	1	3	1	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Missouri												
All races, both sexes	647	784	1,015	1,189	1,699	2,441	3,334	3,484	2,673	1,141	253	-
Male	442	549	650	738	971	1,306	1,503	1,266	754	257	36	-
Female	205	235	365	451	728	1,135	1,831	2,218	1,919	884	217	-
White, both sexes	502	650	835	1,025	1,496	2,200	3,075	3,273	2,524	1,061	227	-
Male	347	463	550	650	877	1,189	1,392	1,207	704	239	30	-
Female	155	187	285	375	619	1,011	1,683	2,066	1,820	822	197	-
Black, both sexes	142	128	172	156	193	226	251	204	144	76	26	-
Male	94	85	95	84	89	110	107	57	48	18	6	-
Female	48	43	77	72	104	116	144	147	96	58	20	-
American Indian 1/, both sexes	2	1	3	4	3	3	4	3	2	1	-	-
Male	1	1	2	1	1	2	3	1	1	-	-	-
Female	1	-	1	3	2	1	1	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes	1	5	5	4	7	12	4	4	3	3	-	-
Male	-	-	3	3	4	5	1	1	1	-	-	-
Female	1	5	2	1	3	7	3	3	2	3	-	-
Montana												
All races, both sexes	83	103	123	158	209	286	434	509	386	142	32	-
Male	59	69	84	100	120	176	227	215	124	40	3	-
Female	24	34	39	58	89	110	207	294	262	102	29	-
White, both sexes	73	94	113	148	200	273	420	501	380	139	31	-
Male	53	65	76	95	118	170	224	213	123	39	3	-
Female	20	29	37	53	82	103	196	288	257	100	28	-
Black, both sexes	-	-	1	1	-	-	1	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	10	7	9	9	9	11	12	7	5	3	1	-
Male	6	3	7	5	2	5	2	1	-	1	-	-
Female	4	4	2	4	7	6	10	6	5	2	1	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	2	1	-	1	-	-	-
Male	-	1	-	-	-	1	1	-	1	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Nebraska												
All races, both sexes	123	126	186	244	382	570	881	1,035	899	379	96	-
Male	87	95	124	152	233	328	427	421	285	96	18	-
Female	36	31	62	92	149	242	454	614	614	283	78	-
White, both sexes	107	111	169	225	358	545	869	1,010	888	373	94	-
Male	75	86	113	142	223	314	421	411	280	94	18	-
Female	32	25	56	83	135	231	448	599	608	279	76	-
Black, both sexes	12	12	13	13	18	16	11	14	8	6	1	-
Male	10	8	9	8	8	11	6	5	3	2	-	-
Female	2	4	4	5	10	5	5	9	5	4	1	-
American Indian 1/, both sexes	4	2	2	3	3	4	1	5	2	-	1	-
Male	2	-	1	1	1	1	-	2	1	-	-	-
Female	2	2	1	2	2	3	1	3	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	2	3	3	5	-	6	1	-	-	-
Male	-	1	1	1	1	2	-	3	1	-	-	-
Female	-	-	1	2	2	3	-	3	-	-	-	-
Nevada												
All races, both sexes	309	409	539	605	704	891	1,017	861	527	153	25	-
Male	213	303	335	388	435	508	517	348	178	28	5	-
Female	96	106	204	217	269	383	500	513	349	125	20	-
White, both sexes	255	341	464	527	604	808	945	814	486	143	24	-
Male	180	260	295	337	377	471	480	331	164	28	5	-
Female	75	81	169	190	227	337	465	483	322	115	19	-
Black, both sexes	38	45	50	43	63	48	42	25	20	5	1	-
Male	19	28	24	28	36	22	23	8	7	-	-	-
Female	19	17	26	15	27	26	19	17	13	5	1	-
American Indian 1/, both sexes	2	8	5	9	9	5	6	3	6	2	-	-
Male	1	5	4	6	5	3	2	2	1	-	-	-
Female	1	3	1	3	4	2	4	1	5	2	-	-
Asian or Pacific Islander 2/, both sexes	14	15	20	26	28	30	24	19	15	3	-	-
Male	13	10	12	17	17	12	12	7	6	-	-	-
Female	1	5	8	9	11	18	12	12	9	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
New Hampshire												
All races, both sexes	92	118	140	198	246	380	558	631	513	233	68	-
Male	71	91	108	128	147	200	244	233	156	55	4	-
Female	21	27	32	70	99	180	314	398	357	178	64	-
White, both sexes	91	118	139	196	244	376	556	631	511	233	68	-
Male	71	91	107	127	146	196	243	233	155	55	4	-
Female	20	27	32	69	98	180	313	398	356	178	64	-
Black, both sexes	1	-	1	1	-	3	1	-	1	-	-	-
Male	-	-	1	1	-	3	1	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	1	-	1	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	1	-	-	-
New Jersey												
All races, both sexes	689	925	1,128	1,310	2,017	3,272	4,717	5,118	3,855	1,610	362	5
Male	476	637	737	827	1,137	1,694	2,146	1,983	1,116	335	58	4
Female	213	288	391	483	880	1,578	2,571	3,135	2,739	1,275	304	1
White, both sexes	476	689	842	1,011	1,639	2,797	4,232	4,786	3,587	1,480	318	2
Male	344	490	571	633	945	1,472	1,976	1,878	1,053	309	53	2
Female	132	199	271	378	694	1,325	2,256	2,908	2,534	1,171	265	-
Black, both sexes	187	204	245	252	322	412	420	275	227	113	40	2
Male	110	122	137	159	157	193	150	83	55	18	3	1
Female	77	82	108	93	165	219	270	192	172	95	37	1
American Indian 1/, both sexes	-	1	1	7	5	4	3	3	5	4	-	-
Male	-	-	-	3	4	1	1	1	1	2	-	-
Female	-	1	1	4	1	3	2	2	4	2	-	-
Asian or Pacific Islander 2/, both sexes	26	31	40	40	51	59	62	54	36	13	4	1
Male	22	25	29	32	31	28	19	21	7	6	2	1
Female	4	6	11	8	20	31	43	33	29	7	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
New Mexico												
All races, both sexes	157	175	244	290	432	583	804	730	572	257	57	1
Male	111	121	167	180	267	301	392	318	193	67	11	1
Female	46	54	77	110	165	282	412	412	379	190	46	-
White, both sexes	136	158	216	273	399	544	767	695	542	239	55	1
Male	96	111	152	169	248	278	370	297	184	63	10	1
Female	40	47	64	104	151	266	397	398	358	176	45	-
Black, both sexes	5	5	6	5	11	14	12	7	8	6	-	-
Male	3	3	4	4	7	9	6	2	1	1	-	-
Female	2	2	2	1	4	5	6	5	7	5	-	-
American Indian 1/, both sexes	16	12	22	11	18	22	24	27	18	11	2	-
Male	12	7	11	7	11	12	15	18	6	3	1	-
Female	4	5	11	4	7	10	9	9	12	8	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	4	3	1	1	4	1	-	-
Male	-	-	-	-	1	2	1	1	2	-	-	-
Female	-	-	-	1	3	1	-	-	2	1	-	-
New York												
All races, both sexes	1,542	2,222	2,726	3,390	5,131	7,751	10,939	11,849	9,754	4,143	971	1
Male	1,061	1,514	1,709	1,989	2,876	4,004	4,871	4,387	2,877	863	124	-
Female	481	708	1,017	1,401	2,255	3,747	6,068	7,462	6,877	3,280	847	1
White, both sexes	1,058	1,595	1,977	2,526	4,084	6,461	9,451	10,651	8,929	3,786	837	1
Male	771	1,117	1,286	1,504	2,346	3,417	4,274	4,010	2,685	811	107	-
Female	287	478	691	1,022	1,738	3,044	5,177	6,641	6,244	2,975	730	1
Black, both sexes	425	560	669	754	902	1,113	1,265	998	657	310	115	-
Male	248	346	371	416	440	495	486	297	129	39	11	-
Female	177	214	298	338	462	618	779	701	528	271	104	-
American Indian 1/, both sexes	3	7	6	4	6	7	2	12	8	1	-	-
Male	3	6	5	2	1	5	-	6	2	-	-	-
Female	-	1	1	2	5	2	2	6	6	1	-	-
Asian or Pacific Islander 2/, both sexes	56	60	74	106	139	170	221	188	160	46	19	-
Male	39	45	47	67	89	87	111	74	61	13	6	-
Female	17	15	27	39	50	83	110	114	99	33	13	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
North Carolina												
All races, both sexes	967	1,208	1,405	1,714	2,344	3,159	4,173	4,040	2,827	1,035	215	2
Male	673	835	929	1,077	1,315	1,638	1,823	1,447	795	184	28	2
Female	294	373	476	637	1,029	1,521	2,350	2,593	2,032	851	187	-
White, both sexes	619	799	971	1,238	1,796	2,462	3,389	3,418	2,415	855	170	2
Male	467	568	661	805	1,032	1,324	1,525	1,267	700	157	24	2
Female	152	231	310	433	764	1,138	1,864	2,151	1,715	698	146	-
Black, both sexes	334	378	400	444	510	657	739	593	395	174	44	-
Male	196	246	249	252	265	296	280	168	89	26	4	-
Female	138	132	151	192	245	361	459	425	306	148	40	-
American Indian 1/, both sexes	8	20	25	25	27	31	32	20	12	4	1	-
Male	5	12	15	16	15	15	11	9	2	-	-	-
Female	3	8	10	9	12	16	21	11	10	4	1	-
Asian or Pacific Islander 2/, both sexes	6	11	9	7	11	9	13	9	5	2	-	-
Male	5	9	4	4	3	3	7	3	4	1	-	-
Female	1	2	5	3	8	6	6	6	1	1	-	-
North Dakota												
All races, both sexes	44	65	64	91	129	254	355	432	352	182	28	-
Male	33	47	47	60	82	148	198	184	101	45	2	-
Female	11	18	17	31	47	106	157	248	251	137	26	-
White, both sexes	40	57	63	85	125	246	353	427	351	180	28	-
Male	30	44	46	56	78	144	197	182	100	44	2	-
Female	10	13	17	29	47	102	156	245	251	136	26	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	8	1	5	4	6	2	5	1	2	-	-
Male	3	3	1	3	4	4	1	2	1	1	-	-
Female	1	5	-	2	-	2	1	3	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Ohio												
All races, both sexes	1,244	1,544	1,924	2,374	3,163	4,991	6,952	6,984	4,961	1,948	401	-
Male	888	1,042	1,239	1,479	1,817	2,566	3,139	2,657	1,421	388	55	-
Female	356	502	685	895	1,346	2,425	3,813	4,327	3,540	1,560	346	-
White, both sexes	965	1,242	1,600	1,989	2,739	4,449	6,381	6,522	4,679	1,829	359	-
Male	709	859	1,051	1,258	1,599	2,328	2,884	2,489	1,343	368	45	-
Female	256	383	549	731	1,140	2,121	3,497	4,033	3,336	1,461	314	-
Black, both sexes	273	299	315	367	412	529	545	450	276	118	42	-
Male	173	182	185	212	214	233	243	162	76	20	10	-
Female	100	117	130	155	198	296	302	288	200	98	32	-
American Indian 1/, both sexes	1	1	-	-	1	3	2	2	2	-	-	-
Male	1	-	-	-	-	1	2	-	-	-	-	-
Female	-	1	-	-	1	2	-	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	5	2	9	18	11	10	24	10	4	1	-	-
Male	5	1	3	9	4	4	10	6	2	-	-	-
Female	-	1	6	9	7	6	14	4	2	1	-	-
Oklahoma												
All races, both sexes	456	629	738	869	1,216	1,740	2,253	2,195	1,666	664	159	2
Male	316	407	475	520	694	863	1,042	817	466	147	25	1
Female	140	222	263	349	522	877	1,211	1,378	1,200	517	134	1
White, both sexes	350	515	633	740	1,066	1,557	2,052	2,021	1,550	614	133	2
Male	255	346	407	442	611	781	950	751	435	134	21	1
Female	95	169	226	298	455	776	1,102	1,270	1,115	480	112	1
Black, both sexes	60	77	52	65	69	87	98	112	71	31	14	-
Male	29	40	31	42	34	39	45	41	17	6	2	-
Female	31	37	21	23	35	48	53	71	54	25	12	-
American Indian 1/, both sexes	44	34	48	59	73	87	97	58	44	19	11	-
Male	31	21	33	32	46	38	44	24	14	7	2	-
Female	13	13	15	27	27	49	53	34	30	12	9	-
Asian or Pacific Islander 2/, both sexes	2	3	5	5	8	9	6	4	1	-	1	-
Male	1	-	4	4	3	5	3	1	-	-	-	-
Female	1	3	1	1	5	4	3	3	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Oregon												
All races, both sexes	278	363	429	527	718	1,193	1,831	2,042	1,603	639	114	-
Male	207	265	300	309	410	624	901	859	546	155	20	-
Female	71	98	129	218	308	569	930	1,183	1,057	484	94	-
White, both sexes	252	336	402	504	690	1,154	1,790	1,998	1,576	631	111	-
Male	194	244	283	300	390	603	880	842	541	152	18	-
Female	58	92	119	204	300	551	910	1,156	1,035	479	93	-
Black, both sexes	14	13	12	9	11	11	20	13	13	3	1	-
Male	6	10	7	5	8	6	11	4	1	-	1	-
Female	8	3	5	4	3	5	9	9	12	3	-	-
American Indian 1/, both sexes	8	6	6	5	7	9	8	8	3	1	1	-
Male	4	3	5	1	5	6	4	1	2	1	1	-
Female	4	3	1	4	2	3	4	7	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	8	9	9	10	19	13	23	11	4	1	-
Male	3	8	5	3	7	9	6	12	2	2	-	-
Female	1	-	4	6	3	10	7	11	9	2	1	-
Pennsylvania												
All races, both sexes	1,236	1,629	2,024	2,496	3,834	6,058	8,843	9,321	6,701	2,612	532	3
Male	879	1,136	1,311	1,545	2,192	3,199	3,948	3,516	1,920	526	69	1
Female	357	493	713	951	1,642	2,859	4,895	5,805	4,781	2,086	463	2
White, both sexes	971	1,333	1,698	2,153	3,402	5,517	8,236	8,869	6,403	2,478	491	2
Male	707	938	1,109	1,366	1,954	2,928	3,683	3,376	1,854	506	68	1
Female	264	395	589	787	1,448	2,589	4,553	5,493	4,549	1,972	423	1
Black, both sexes	249	282	312	322	408	520	581	431	281	124	40	1
Male	162	186	192	168	224	259	254	134	61	17	-	-
Female	87	96	120	154	184	261	327	297	220	107	40	1
American Indian 1/, both sexes	-	2	1	1	1	2	3	5	-	1	-	-
Male	-	1	-	-	1	-	1	2	-	-	-	-
Female	-	1	1	1	-	2	2	3	-	1	-	-
Asian or Pacific Islander 2/, both sexes	16	12	13	20	23	19	23	16	17	9	1	-
Male	10	11	10	11	13	12	10	4	5	3	1	-
Female	6	1	3	9	10	7	13	12	12	6	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Rhode Island												
All races, both sexes	90	132	149	165	264	434	661	787	615	274	54	-
Male	72	94	105	103	154	231	296	289	166	46	6	-
Female	18	38	44	62	110	203	365	498	449	228	48	-
White, both sexes	81	126	136	152	249	413	644	768	608	271	52	-
Male	65	89	95	94	149	217	288	281	165	46	4	-
Female	16	37	41	58	100	196	356	487	443	225	48	-
Black, both sexes	6	6	12	10	12	17	12	14	6	1	1	-
Male	6	5	9	7	4	11	4	6	1	-	1	-
Female	-	1	3	3	8	6	8	8	5	1	-	-
American Indian 1/, both sexes	2	-	-	1	1	1	2	2	1	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	2	-	-	1	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	2	3	3	3	-	2	1	-
Male	1	-	1	2	1	2	3	1	-	-	1	-
Female	-	-	-	-	1	1	-	2	-	2	-	-
South Carolina												
All races, both sexes	571	770	844	973	1,299	1,726	2,094	1,854	1,234	442	99	2
Male	391	525	554	621	747	879	902	699	345	85	15	2
Female	180	245	290	352	552	847	1,192	1,155	889	357	84	-
White, both sexes	309	466	553	648	909	1,283	1,603	1,474	979	340	63	2
Male	223	324	376	429	540	674	716	570	279	61	11	2
Female	86	142	177	219	369	609	887	904	700	279	52	-
Black, both sexes	258	301	287	323	384	436	487	374	253	101	35	-
Male	164	198	175	190	203	203	186	127	66	24	4	-
Female	94	103	112	133	181	233	301	247	187	77	31	-
American Indian 1/, both sexes	2	1	2	-	1	-	-	1	1	-	1	-
Male	2	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	2	2	2	2	5	7	4	5	1	1	-	-
Male	2	2	2	2	3	2	-	2	-	-	-	-
Female	-	-	-	-	2	5	4	3	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
South Dakota												
All races, both sexes	70	72	111	114	179	293	451	476	411	197	51	-
Male	52	52	76	79	111	182	235	207	138	49	9	-
Female	18	20	35	35	68	111	216	269	273	148	42	-
White, both sexes	60	58	98	98	170	273	444	466	404	195	51	-
Male	43	43	66	70	104	170	233	205	136	49	9	-
Female	17	15	32	28	66	103	211	261	268	146	42	-
Black, both sexes	-	-	-	-	-	2	-	1	-	1	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	9	13	13	16	9	17	6	9	7	1	-	-
Male	8	8	10	9	7	10	2	2	2	-	-	-
Female	1	5	3	7	2	7	4	7	5	1	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	1	1	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Tennessee												
All races, both sexes	815	1,108	1,134	1,488	1,860	2,439	3,393	3,255	2,224	881	167	-
Male	555	738	716	919	1,043	1,267	1,442	1,139	607	166	23	-
Female	260	370	418	569	817	1,172	1,951	2,116	1,617	715	144	-
White, both sexes	597	861	894	1,221	1,552	2,083	3,030	2,914	1,969	781	143	-
Male	416	581	570	786	879	1,090	1,297	1,031	541	152	16	-
Female	181	280	324	435	673	993	1,733	1,883	1,428	629	127	-
Black, both sexes	216	241	234	258	293	344	357	336	248	100	24	-
Male	138	153	141	128	156	172	144	107	61	14	7	-
Female	78	88	93	130	137	172	213	229	187	86	17	-
American Indian 1/, both sexes	1	2	1	3	2	2	2	1	1	-	-	-
Male	-	2	1	2	1	1	1	-	1	-	-	-
Female	1	-	-	1	1	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	5	6	13	10	4	4	6	-	-	-
Male	1	2	4	3	7	4	-	1	4	-	-	-
Female	-	2	1	3	6	6	4	3	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Texas												
All races, both sexes	2,093	2,655	3,154	3,609	4,963	6,879	9,071	8,398	6,268	2,480	519	-
Male	1,466	1,765	2,035	2,190	2,886	3,564	4,148	3,160	1,868	526	62	-
Female	627	890	1,119	1,419	2,077	3,315	4,923	5,238	4,400	1,954	457	-
White, both sexes	1,557	2,084	2,549	2,993	4,242	5,991	8,170	7,691	5,721	2,239	459	-
Male	1,123	1,414	1,671	1,862	2,510	3,141	3,784	2,917	1,719	474	54	-
Female	434	670	878	1,131	1,732	2,850	4,386	4,774	4,002	1,765	405	-
Black, both sexes	506	539	548	558	663	819	821	627	512	223	58	-
Male	319	331	332	298	344	382	337	214	142	45	8	-
Female	187	208	216	260	319	437	484	413	370	178	50	-
American Indian 1/, both sexes	5	2	4	3	2	2	11	4	3	-	1	-
Male	5	1	4	1	-	1	3	2	1	-	-	-
Female	-	1	-	2	2	1	8	2	2	-	1	-
Asian or Pacific Islander 2/, both sexes	25	30	53	55	56	67	69	76	32	18	1	-
Male	19	19	28	29	32	40	24	27	6	7	-	-
Female	6	11	25	26	24	27	45	49	26	11	1	-
Utah												
All races, both sexes	111	135	184	221	281	518	716	819	559	208	22	-
Male	71	93	112	142	150	266	334	328	192	58	3	-
Female	40	42	72	79	131	252	382	491	367	150	19	-
White, both sexes	108	125	174	213	272	504	698	809	550	203	22	-
Male	69	86	105	136	147	260	324	322	187	57	3	-
Female	39	39	69	77	125	244	374	487	363	146	19	-
Black, both sexes	2	3	2	2	1	2	7	1	-	3	-	-
Male	2	1	2	1	-	1	2	-	-	1	-	-
Female	-	2	-	1	1	1	5	1	-	2	-	-
American Indian 1/, both sexes	-	3	3	1	3	2	5	1	3	1	-	-
Male	-	3	-	-	1	1	3	1	3	-	-	-
Female	-	-	3	1	2	1	2	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	4	5	5	5	10	6	8	6	1	-	-
Male	-	3	5	5	2	4	5	5	2	-	-	-
Female	1	1	-	-	3	6	1	3	4	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Vermont												
All races, both sexes	31	56	64	72	137	198	268	340	262	110	29	-
Male	16	43	41	48	76	117	127	153	89	27	2	-
Female	15	13	23	24	61	81	141	187	173	83	27	-
White, both sexes	31	56	64	71	136	198	268	339	262	110	29	-
Male	16	43	41	47	76	117	127	153	89	27	2	-
Female	15	13	23	24	60	81	141	186	173	83	27	-
Black, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	717	957	1,135	1,326	1,708	2,513	3,236	3,231	2,290	857	195	2
Male	498	669	713	808	948	1,307	1,402	1,209	641	159	21	1
Female	219	288	422	518	760	1,206	1,834	2,022	1,649	698	174	1
White, both sexes	454	641	804	966	1,258	1,975	2,671	2,753	1,943	741	157	1
Male	321	463	522	603	723	1,052	1,170	1,059	555	144	16	-
Female	133	178	282	363	535	923	1,501	1,694	1,388	597	141	1
Black, both sexes	252	298	305	336	419	495	522	444	332	112	36	1
Male	169	196	177	190	206	230	212	141	82	15	4	1
Female	83	102	128	146	213	265	310	303	250	97	32	-
American Indian 1/, both sexes	-	1	2	-	1	1	2	2	1	-	-	-
Male	-	1	2	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	11	17	24	24	30	42	41	32	14	4	2	-
Male	8	9	12	15	18	25	19	9	4	-	1	-
Female	3	8	12	9	12	17	22	23	10	4	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Washington												
All races, both sexes	468	573	744	899	1,204	1,898	2,737	2,892	2,178	866	223	-
Male	333	415	539	572	741	1,042	1,333	1,212	697	198	35	-
Female	135	158	205	327	463	856	1,404	1,680	1,481	668	188	-
White, both sexes	404	490	679	826	1,113	1,779	2,588	2,782	2,097	848	216	-
Male	290	355	499	534	683	982	1,262	1,164	670	190	32	-
Female	114	135	180	292	430	797	1,326	1,618	1,427	658	184	-
Black, both sexes	31	43	25	28	36	42	48	40	18	4	-	-
Male	18	34	17	15	22	23	24	14	5	1	-	-
Female	13	9	8	13	14	19	24	26	13	3	-	-
American Indian 1/, both sexes	10	17	17	12	16	19	20	9	7	1	3	-
Male	8	11	10	7	12	14	7	3	-	-	-	-
Female	2	6	7	5	4	5	13	6	7	1	3	-
Asian or Pacific Islander 2/, both sexes	23	23	23	33	39	58	81	61	56	13	4	-
Male	17	15	13	16	24	23	40	31	22	7	3	-
Female	6	8	10	17	15	35	41	30	34	6	1	-
West Virginia												
All races, both sexes	228	347	384	480	654	1,015	1,369	1,185	902	342	80	-
Male	155	228	258	300	365	529	585	442	230	81	18	-
Female	73	119	126	180	289	486	784	743	672	261	62	-
White, both sexes	212	342	367	464	640	988	1,341	1,152	869	331	78	-
Male	147	226	246	291	358	518	575	433	225	80	18	-
Female	65	116	121	173	282	470	766	719	644	251	60	-
Black, both sexes	16	5	16	16	14	26	28	31	33	11	2	-
Male	8	2	11	9	7	11	10	9	5	1	-	-
Female	8	3	5	7	7	15	18	22	28	10	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
IX. Diseases of the circulatory system (I00-I99)												
Wisconsin												
All races, both sexes	397	585	674	836	1,215	2,011	2,728	3,296	2,478	1,051	222	-
Male	301	420	448	530	743	1,129	1,305	1,328	769	203	35	-
Female	96	165	226	306	472	882	1,423	1,968	1,709	848	187	-
White, both sexes	326	517	597	765	1,150	1,926	2,658	3,236	2,438	1,041	214	-
Male	259	371	403	487	707	1,085	1,270	1,302	762	200	35	-
Female	67	146	194	278	443	841	1,388	1,934	1,676	841	179	-
Black, both sexes	60	57	54	54	47	71	49	41	27	8	7	-
Male	36	40	30	35	25	39	22	16	5	2	-	-
Female	24	17	24	19	22	32	27	25	22	6	7	-
American Indian 1/, both sexes	9	8	16	11	7	7	10	8	8	-	-	-
Male	5	7	12	5	6	4	6	3	1	-	-	-
Female	4	1	4	6	1	3	4	5	7	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	7	6	11	7	11	11	5	2	1	-
Male	1	2	3	3	5	1	7	7	1	1	-	-
Female	1	1	4	3	6	6	4	4	4	1	1	-
Wyoming												
All races, both sexes	41	59	71	88	120	175	220	204	135	60	20	-
Male	26	37	47	54	61	104	115	78	52	9	3	-
Female	15	22	24	34	59	71	105	126	83	51	17	-
White, both sexes	37	56	69	88	116	173	218	203	134	60	20	-
Male	24	37	46	54	61	103	114	78	52	9	3	-
Female	13	19	23	34	55	70	104	125	82	51	17	-
Black, both sexes	1	1	-	-	1	1	-	1	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	1	1	-	-	-
American Indian 1/, both sexes	3	1	1	-	2	1	1	-	-	-	-	-
Male	1	-	1	-	-	1	1	-	-	-	-	-
Female	2	1	-	-	2	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
United States												
All races, both sexes	3,365	1	1	1	3	3	6	11	16	26	56	87
Male	1,044	1	-	-	2	1	6	7	7	13	25	35
Female	2,321	-	1	1	1	2	-	4	9	13	31	52
White, both sexes	3,029	1	1	1	3	2	2	7	10	20	37	62
Male	926	1	-	-	2	-	2	5	6	9	19	26
Female	2,103	-	1	1	1	2	-	2	4	11	18	36
Black, both sexes	227	-	-	-	-	1	1	2	6	3	13	19
Male	81	-	-	-	-	1	1	1	1	3	5	7
Female	146	-	-	-	-	-	-	1	5	-	8	12
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	-	4	3
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	3	2
Asian or Pacific Islander 2/, both sexes	89	-	-	-	-	-	3	2	-	3	2	3
Male	30	-	-	-	-	-	3	1	-	1	-	1
Female	59	-	-	-	-	-	-	1	-	2	2	2
Alabama												
All races, both sexes	45	-	-	-	-	-	-	-	2	1	2	1
Male	15	-	-	-	-	-	-	-	-	1	1	-
Female	30	-	-	-	-	-	-	-	2	-	1	1
White, both sexes	30	-	-	-	-	-	-	-	1	-	2	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	13	-	-	-	-	-	-	-	1	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Alaska												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	57	-	-	-	-	-	-	-	1	-	-	2
Male	17	-	-	-	-	-	-	-	1	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	52	-	-	-	-	-	-	-	1	-	-	1
Male	15	-	-	-	-	-	-	-	1	-	-	1
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Arkansas												
All races, both sexes	22	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	403	-	-	-	1	-	-	1	2	6	5	14
Male	138	-	-	-	1	-	-	1	1	3	2	5
Female	265	-	-	-	-	-	-	-	1	3	3	9
White, both sexes	345	-	-	-	1	-	-	-	1	4	4	10
Male	119	-	-	-	1	-	-	-	1	2	2	4
Female	226	-	-	-	-	-	-	-	-	2	2	6
Black, both sexes	20	-	-	-	-	-	-	1	1	-	1	3
Male	7	-	-	-	-	-	-	1	-	-	-	1
Female	13	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	-	-	2	-	1
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	25	-	-	-	-	-	-	-	-	1	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Colorado												
All races, both sexes	60	-	-	-	1	-	-	1	1	-	-	1
Male	15	-	-	-	-	-	-	1	1	-	-	-
Female	45	-	-	-	1	-	-	-	-	-	-	1
White, both sexes	57	-	-	-	1	-	-	-	1	-	-	1
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	43	-	-	-	1	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Delaware												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Florida												
All races, both sexes	216	-	-	-	-	-	1	3	1	1	2	5
Male	92	-	-	-	-	-	1	2	-	1	1	3
Female	124	-	-	-	-	-	-	1	1	-	1	2
White, both sexes	202	-	-	-	-	-	-	2	1	1	1	5
Male	85	-	-	-	-	-	-	2	-	1	1	3
Female	117	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	11	-	-	-	-	-	1	1	-	-	1	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	58	-	1	-	-	1	-	1	-	2	1	3
Male	19	-	-	-	-	-	-	1	-	2	1	-
Female	39	-	1	-	-	1	-	-	-	-	-	3
White, both sexes	48	-	1	-	-	1	-	1	-	2	-	1
Male	16	-	-	-	-	-	-	1	-	2	-	-
Female	32	-	1	-	-	1	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	2
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Hawaii												
All races, both sexes	19	-	-	-	-	-	-	1	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	1	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	1	-	-	1	-
Idaho												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Illinois												
All races, both sexes	128	-	-	1	-	-	-	-	1	-	4	5
Male	37	-	-	-	-	-	-	-	-	-	2	3
Female	91	-	-	1	-	-	-	-	1	-	2	2
White, both sexes	112	-	-	1	-	-	-	-	-	-	4	4
Male	35	-	-	-	-	-	-	-	-	-	2	2
Female	77	-	-	1	-	-	-	-	-	-	2	2
Black, both sexes	13	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	66	-	-	-	-	-	1	-	-	-	2	1
Male	14	-	-	-	-	-	1	-	-	-	1	1
Female	52	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	63	-	-	-	-	-	1	-	-	-	1	1
Male	14	-	-	-	-	-	1	-	-	-	1	1
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Iowa												
All races, both sexes	72	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	55	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	55	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Kentucky												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	34	-	-	-	-	-	1	-	-	-	2	4
Male	16	-	-	-	-	-	1	-	-	-	-	2
Female	18	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	24	-	-	-	-	-	-	-	-	-	2	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Maine												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	33	-	-	-	-	1	-	-	1	1	-	1
Male	8	-	-	-	-	1	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	11	-	-	-	-	1	-	-	1	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Massachusetts												
All races, both sexes	66	-	-	-	-	-	-	-	1	2	1	1
Male	13	-	-	-	-	-	-	-	-	1	1	-
Female	53	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	65	-	-	-	-	-	-	-	1	1	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	53	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	97	-	-	-	-	-	-	-	-	1	1	2
Male	27	-	-	-	-	-	-	-	-	1	1	2
Female	70	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	91	-	-	-	-	-	-	-	-	1	1	1
Male	25	-	-	-	-	-	-	-	-	1	1	1
Female	66	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Minnesota												
All races, both sexes	58	-	-	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	39	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	56	-	-	-	-	-	-	-	-	-	1	1
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	38	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Missouri												
All races, both sexes	90	-	-	-	-	-	-	-	-	-	3	2
Male	37	-	-	-	-	-	-	-	-	-	1	1
Female	53	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	82	-	-	-	-	-	-	-	-	-	2	2
Male	34	-	-	-	-	-	-	-	-	-	1	1
Female	48	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Nebraska												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	14	-	-	-	-	-	1	-	-	-	1	1
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
New Hampshire												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	123	-	-	-	-	-	-	-	-	1	-	2
Male	35	-	-	-	-	-	-	-	-	1	-	1
Female	88	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	108	-	-	-	-	-	-	-	-	1	-	-
Male	29	-	-	-	-	-	-	-	-	1	-	-
Female	79	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
New Mexico												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	203	-	-	-	-	1	-	-	-	1	7	7
Male	61	-	-	-	-	-	-	-	-	-	5	2
Female	142	-	-	-	-	1	-	-	-	1	2	5
White, both sexes	185	-	-	-	-	1	-	-	-	1	5	4
Male	56	-	-	-	-	-	-	-	-	-	4	2
Female	129	-	-	-	-	1	-	-	-	1	1	2
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	3
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
North Carolina												
All races, both sexes	92	-	-	-	-	-	-	-	1	-	-	2
Male	29	-	-	-	-	-	-	-	-	-	-	1
Female	63	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	79	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	54	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	1	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Ohio												
All races, both sexes	192	1	-	-	1	-	-	-	-	1	2	2
Male	65	1	-	-	1	-	-	-	-	1	1	1
Female	127	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	175	1	-	-	1	-	-	-	-	1	1	2
Male	61	1	-	-	1	-	-	-	-	1	1	1
Female	114	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Oklahoma												
All races, both sexes	40	-	-	-	-	-	-	1	1	-	1	1
Male	16	-	-	-	-	-	-	1	1	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	34	-	-	-	-	-	-	1	1	-	-	-
Male	15	-	-	-	-	-	-	1	1	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Oregon												
All races, both sexes	88	-	-	-	-	-	-	-	-	1	1	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	60	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	82	-	-	-	-	-	-	-	-	-	1	1
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	55	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Pennsylvania												
All races, both sexes	175	-	-	-	-	-	-	-	-	1	1	2
Male	48	-	-	-	-	-	-	-	-	1	-	-
Female	127	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	171	-	-	-	-	-	-	-	-	-	1	2
Male	46	-	-	-	-	-	-	-	-	-	-	-
Female	125	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Rhode Island												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	58	-	-	-	-	-	-	1	1	-	3	3
Male	14	-	-	-	-	-	-	-	1	-	1	1
Female	44	-	-	-	-	-	-	1	-	-	2	2
White, both sexes	42	-	-	-	-	-	-	1	-	-	2	2
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	33	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	16	-	-	-	-	-	-	-	1	-	1	1
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
South Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	87	-	-	-	-	-	-	-	1	2	4	1
Male	30	-	-	-	-	-	-	-	1	-	3	-
Female	57	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	79	-	-	-	-	-	-	-	1	2	3	1
Male	25	-	-	-	-	-	-	-	1	-	2	-
Female	54	-	-	-	-	-	-	-	-	2	1	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Texas												
All races, both sexes	148	-	-	-	-	-	1	1	-	2	3	4
Male	43	-	-	-	-	-	1	1	-	-	-	2
Female	105	-	-	-	-	-	-	-	-	2	3	2
White, both sexes	136	-	-	-	-	-	1	1	-	2	2	3
Male	39	-	-	-	-	-	1	1	-	-	-	1
Female	97	-	-	-	-	-	-	-	-	2	2	2
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Vermont												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	56	-	-	-	-	-	1	-	1	1	2	4
Male	14	-	-	-	-	-	1	-	-	1	1	1
Female	42	-	-	-	-	-	-	-	1	-	1	3
White, both sexes	45	-	-	-	-	-	-	-	1	1	-	3
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	37	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	10	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Washington												
All races, both sexes	87	-	-	-	-	-	-	1	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	68	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	80	-	-	-	-	-	-	1	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	63	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	32	-	-	-	-	-	-	-	-	1	-	6
Male	13	-	-	-	-	-	-	-	-	-	-	3
Female	19	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	32	-	-	-	-	-	-	-	-	1	-	6
Male	13	-	-	-	-	-	-	-	-	-	-	3
Female	19	-	-	-	-	-	-	-	-	1	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Wisconsin												
All races, both sexes	77	-	-	-	-	-	-	-	-	-	2	2
Male	24	-	-	-	-	-	-	-	-	-	1	1
Female	53	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	75	-	-	-	-	-	-	-	-	-	1	2
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	13	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
United States												
All races, both sexes	104	160	194	252	361	497	617	530	327	103	9	-
Male	46	74	67	91	119	152	174	131	79	14	-	-
Female	58	86	127	161	242	345	443	399	248	89	9	-
White, both sexes	72	124	160	220	326	460	584	513	314	101	9	-
Male	34	58	58	77	111	144	161	124	75	14	-	-
Female	38	66	102	143	215	316	423	389	239	87	9	-
Black, both sexes	29	22	22	25	17	25	19	12	9	2	-	-
Male	11	10	7	11	2	5	10	4	2	-	-	-
Female	18	12	15	14	15	20	9	8	7	2	-	-
American Indian 1/, both sexes	-	3	-	2	1	1	4	1	1	-	-	-
Male	-	1	-	1	1	1	-	-	1	-	-	-
Female	-	2	-	1	-	-	4	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	11	12	5	17	11	10	4	3	-	-	-
Male	1	5	2	2	5	2	3	3	1	-	-	-
Female	2	6	10	3	12	9	7	1	2	-	-	-
Alabama												
All races, both sexes	3	1	2	2	7	7	9	5	3	-	-	-
Male	-	1	-	1	2	2	5	1	1	-	-	-
Female	3	-	2	1	5	5	4	4	2	-	-	-
White, both sexes	1	-	2	1	5	5	7	4	1	-	-	-
Male	-	-	-	-	1	1	4	-	-	-	-	-
Female	1	-	2	1	4	4	3	4	1	-	-	-
Black, both sexes	2	1	-	1	1	2	1	1	2	-	-	-
Male	-	1	-	1	-	1	1	1	1	-	-	-
Female	2	-	-	-	1	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Alaska												
All races, both sexes	-	-	1	2	1	2	1	-	1	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	1	-	-	-
White, both sexes	-	-	1	2	-	1	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	3	3	5	6	10	13	10	2	1	-	-
Male	1	1	1	3	1	2	4	-	1	1	-	-
Female	-	2	2	2	5	8	9	10	1	-	-	-
White, both sexes	1	2	3	4	6	10	11	10	2	1	-	-
Male	1	-	1	2	1	2	4	-	1	1	-	-
Female	-	2	2	2	5	8	7	10	1	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Arkansas												
All races, both sexes	-	1	3	2	3	3	2	4	2	1	-	-
Male	-	1	-	1	1	2	-	1	-	-	-	-
Female	-	-	3	1	2	1	2	3	2	1	-	-
White, both sexes	-	1	3	2	3	3	2	4	1	1	-	-
Male	-	1	-	1	1	2	-	1	-	-	-	-
Female	-	-	3	1	2	1	2	3	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	10	24	30	28	37	56	71	63	38	16	1	-
Male	6	12	9	12	11	16	18	24	13	4	-	-
Female	4	12	21	16	26	40	53	39	25	12	1	-
White, both sexes	9	16	24	24	27	49	64	61	34	16	1	-
Male	6	8	7	10	10	14	16	23	11	4	-	-
Female	3	8	17	14	17	35	48	38	23	12	1	-
Black, both sexes	1	3	-	1	3	2	3	-	1	-	-	-
Male	-	2	-	-	1	1	1	-	-	-	-	-
Female	1	1	-	1	2	1	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	5	6	3	7	5	3	2	2	-	-	-
Male	-	2	2	2	-	1	1	1	1	-	-	-
Female	-	3	4	1	7	4	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Colorado												
All races, both sexes	4	3	3	2	7	11	11	9	6	-	-	-
Male	1	2	2	1	1	-	-	4	2	-	-	-
Female	3	1	1	1	6	11	11	5	4	-	-	-
White, both sexes	3	3	3	2	6	11	11	9	6	-	-	-
Male	1	2	2	1	1	-	-	4	2	-	-	-
Female	2	1	1	1	5	11	11	5	4	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	1	3	7	3	8	11	6	2	-	-
Male	1	-	-	1	2	1	3	3	1	-	-	-
Female	-	-	1	2	5	2	5	8	5	2	-	-
White, both sexes	1	-	1	2	7	3	8	11	6	2	-	-
Male	1	-	-	1	2	1	3	3	1	-	-	-
Female	-	-	1	1	5	2	5	8	5	2	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Delaware												
All races, both sexes	-	-	2	1	1	1	2	-	2	-	-	-
Male	-	-	1	1	-	-	2	-	1	-	-	-
Female	-	-	1	-	1	1	-	-	1	-	-	-
White, both sexes	-	-	1	1	1	1	2	-	2	-	-	-
Male	-	-	1	1	-	-	2	-	1	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Florida												
All races, both sexes	8	14	9	19	28	26	42	27	24	6	-	-
Male	3	11	5	10	12	11	17	7	7	1	-	-
Female	5	3	4	9	16	15	25	20	17	5	-	-
White, both sexes	6	13	8	16	26	25	42	27	23	6	-	-
Male	3	10	5	8	10	11	17	7	6	1	-	-
Female	3	3	3	8	16	14	25	20	17	5	-	-
Black, both sexes	2	1	1	2	-	1	-	-	1	-	-	-
Male	-	1	-	2	-	-	-	-	1	-	-	-
Female	2	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	-	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	8	3	3	6	9	5	9	4	-	-	-
Male	1	3	1	1	-	6	2	-	1	-	-	-
Female	1	5	2	2	6	3	3	9	3	-	-	-
White, both sexes	1	7	3	3	5	8	4	7	4	-	-	-
Male	-	2	1	1	-	6	2	-	1	-	-	-
Female	1	5	2	2	5	2	2	7	3	-	-	-
Black, both sexes	1	1	-	-	1	1	-	2	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Hawaii												
All races, both sexes	2	2	3	-	2	3	1	3	-	1	-	-
Male	2	1	-	-	-	1	1	2	-	-	-	-
Female	-	1	3	-	2	2	-	1	-	1	-	-
White, both sexes	1	-	-	-	-	1	-	2	-	1	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	3	-	2	2	1	1	-	-	-	-
Male	1	1	-	-	-	1	1	1	-	-	-	-
Female	-	1	3	-	2	1	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	4	3	1	2	4	8	1	-	-	-
Male	-	-	2	1	-	-	1	2	-	-	-	-
Female	-	-	2	2	1	2	3	6	1	-	-	-
White, both sexes	-	-	4	3	1	2	4	8	1	-	-	-
Male	-	-	2	1	-	-	1	2	-	-	-	-
Female	-	-	2	2	1	2	3	6	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Illinois												
All races, both sexes	5	6	8	10	12	19	26	12	13	6	-	-
Male	1	2	2	4	4	3	7	4	5	-	-	-
Female	4	4	6	6	8	16	19	8	8	6	-	-
White, both sexes	2	6	6	7	11	15	25	12	13	6	-	-
Male	1	2	2	3	4	3	7	4	5	-	-	-
Female	1	4	4	4	7	12	18	8	8	6	-	-
Black, both sexes	3	-	2	2	1	4	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	3	-	2	1	1	4	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Indiana												
All races, both sexes	4	3	4	4	5	14	10	7	8	3	-	-
Male	2	-	-	-	1	3	2	-	2	1	-	-
Female	2	3	4	4	4	11	8	7	6	2	-	-
White, both sexes	4	2	4	4	4	14	10	7	8	3	-	-
Male	2	-	-	-	1	3	2	-	2	1	-	-
Female	2	2	4	4	3	11	8	7	6	2	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Iowa												
All races, both sexes	1	2	2	4	7	11	9	17	13	5	-	-
Male	1	1	-	1	1	4	3	2	3	-	-	-
Female	-	1	2	3	6	7	6	15	10	5	-	-
White, both sexes	1	2	2	3	7	11	9	16	13	5	-	-
Male	1	1	-	-	1	4	3	1	3	-	-	-
Female	-	1	2	3	6	7	6	15	10	5	-	-
Black, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	2	-	2	2	8	5	5	1	1	1	-
Male	-	2	-	-	1	2	1	1	-	-	-	-
Female	-	-	-	2	1	6	4	4	1	1	1	-
White, both sexes	-	2	-	2	1	8	5	5	1	1	1	-
Male	-	2	-	-	-	2	1	1	-	-	-	-
Female	-	-	-	2	1	6	4	4	1	1	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Kentucky												
All races, both sexes	1	6	1	6	4	10	9	3	4	2	-	-
Male	1	4	-	2	1	5	2	1	-	-	-	-
Female	-	2	1	4	3	5	7	2	4	2	-	-
White, both sexes	1	6	1	6	4	10	9	3	4	2	-	-
Male	1	4	-	2	1	5	2	1	-	-	-	-
Female	-	2	1	4	3	5	7	2	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	3	4	2	3	5	5	3	1	-	-	-
Male	-	1	3	1	1	2	3	2	-	-	-	-
Female	1	2	1	1	2	3	2	1	1	-	-	-
White, both sexes	1	2	2	1	3	4	3	3	1	-	-	-
Male	-	1	2	-	1	2	1	2	-	-	-	-
Female	1	1	-	1	2	2	2	1	1	-	-	-
Black, both sexes	-	1	2	1	-	1	2	-	-	-	-	-
Male	-	-	1	1	-	-	2	-	-	-	-	-
Female	-	1	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Maine												
All races, both sexes	-	1	4	1	6	2	4	3	1	-	-	-
Male	-	1	2	-	2	1	1	1	-	-	-	-
Female	-	-	2	1	4	1	3	2	1	-	-	-
White, both sexes	-	1	4	1	6	2	4	3	1	-	-	-
Male	-	1	2	-	2	1	1	1	-	-	-	-
Female	-	-	2	1	4	1	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	3	3	2	5	6	4	3	2	-	-	-
Male	-	-	2	1	1	2	-	-	-	-	-	-
Female	1	3	1	1	4	4	4	3	2	-	-	-
White, both sexes	-	1	2	2	4	3	3	3	2	-	-	-
Male	-	-	2	1	1	2	-	-	-	-	-	-
Female	-	1	-	1	3	1	3	3	2	-	-	-
Black, both sexes	1	2	1	-	1	3	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	2	1	-	1	3	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Massachusetts												
All races, both sexes	2	1	5	3	8	13	9	5	12	2	1	-
Male	1	1	1	1	3	3	-	1	-	-	-	-
Female	1	-	4	2	5	10	9	4	12	2	1	-
White, both sexes	2	1	5	3	8	13	9	5	12	2	1	-
Male	1	1	1	1	3	3	-	1	-	-	-	-
Female	1	-	4	2	5	10	9	4	12	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3	4	5	5	12	21	19	14	8	2	-	-
Male	-	1	2	4	3	4	5	2	2	-	-	-
Female	3	3	3	1	9	17	14	12	6	2	-	-
White, both sexes	2	3	5	5	12	20	17	14	8	2	-	-
Male	-	1	2	4	3	4	4	2	2	-	-	-
Female	2	2	3	1	9	16	13	12	6	2	-	-
Black, both sexes	1	1	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Minnesota												
All races, both sexes	2	3	3	-	3	13	16	8	6	2	-	-
Male	1	2	2	-	1	2	3	5	2	-	-	-
Female	1	1	1	-	2	11	13	3	4	2	-	-
White, both sexes	-	3	3	-	3	13	16	8	6	2	-	-
Male	-	2	2	-	1	2	3	5	2	-	-	-
Female	-	1	1	-	2	11	13	3	4	2	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	2	1	5	1	3	6	2	2	-	-	-
Male	-	2	-	2	-	1	1	1	-	-	-	-
Female	-	-	1	3	1	2	5	1	2	-	-	-
White, both sexes	-	1	-	4	1	3	6	1	2	-	-	-
Male	-	1	-	2	-	1	1	-	-	-	-	-
Female	-	-	-	2	1	2	5	1	2	-	-	-
Black, both sexes	-	1	1	1	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Missouri												
All races, both sexes	2	3	1	7	11	12	17	16	11	5	-	-
Male	2	2	-	3	6	6	7	5	4	-	-	-
Female	-	1	1	4	5	6	10	11	7	5	-	-
White, both sexes	1	2	-	7	11	11	16	15	11	4	-	-
Male	1	1	-	3	6	6	6	5	4	-	-	-
Female	-	1	-	4	5	5	10	10	7	4	-	-
Black, both sexes	1	1	1	-	-	1	1	1	-	1	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	2	5	2	2	3	1	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	1	2	4	2	1	2	1	-	-	-
White, both sexes	-	-	1	2	5	2	2	3	1	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	1	2	4	2	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Nebraska												
All races, both sexes	-	-	1	2	2	3	3	4	4	-	-	-
Male	-	-	1	-	-	1	1	-	1	-	-	-
Female	-	-	-	2	2	2	2	4	3	-	-	-
White, both sexes	-	-	1	2	2	3	3	4	4	-	-	-
Male	-	-	1	-	-	1	1	-	1	-	-	-
Female	-	-	-	2	2	2	2	4	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	2	2	-	4	1	1	1	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	2	2	-	2	-	1	1	-	-	-
White, both sexes	-	-	2	2	-	4	1	1	1	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	2	2	-	2	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
New Hampshire												
All races, both sexes	-	-	-	1	-	5	-	3	3	-	1	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	5	-	3	2	-	1	-
White, both sexes	-	-	-	1	-	5	-	3	3	-	1	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	5	-	3	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	3	3	12	12	18	32	23	11	3	-	-
Male	2	-	1	4	6	4	6	6	4	-	-	-
Female	1	3	2	8	6	14	26	17	7	3	-	-
White, both sexes	1	3	3	9	8	17	30	22	11	3	-	-
Male	1	-	1	3	4	4	6	5	4	-	-	-
Female	-	3	2	6	4	13	24	17	7	3	-	-
Black, both sexes	1	-	-	3	1	-	1	1	-	-	-	-
Male	1	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	2	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	3	1	1	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	1	-	-	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
New Mexico												
All races, both sexes	2	1	2	3	6	2	5	2	1	-	-	-
Male	-	1	1	1	2	-	2	-	-	-	-	-
Female	2	-	1	2	4	2	3	2	1	-	-	-
White, both sexes	2	1	2	2	6	2	5	2	1	-	-	-
Male	-	1	1	1	2	-	2	-	-	-	-	-
Female	2	-	1	1	4	2	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	5	14	10	12	25	22	44	31	19	5	-	-
Male	-	8	5	3	11	6	15	3	3	-	-	-
Female	5	6	5	9	14	16	29	28	16	5	-	-
White, both sexes	4	8	9	12	25	21	43	30	17	5	-	-
Male	-	5	4	3	11	6	15	3	3	-	-	-
Female	4	3	5	9	14	15	28	27	14	5	-	-
Black, both sexes	1	5	1	-	-	1	-	1	1	-	-	-
Male	-	2	1	-	-	-	-	-	-	-	-	-
Female	1	3	-	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
North Carolina												
All races, both sexes	4	4	5	12	10	16	17	12	9	-	-	-
Male	1	1	3	4	4	6	4	3	2	-	-	-
Female	3	3	2	8	6	10	13	9	7	-	-	-
White, both sexes	2	4	3	11	9	15	15	11	9	-	-	-
Male	1	1	2	3	4	5	4	3	2	-	-	-
Female	1	3	1	8	5	10	11	8	7	-	-	-
Black, both sexes	2	-	2	1	1	1	2	1	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	2	-	1	-	1	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	-	1	2	1	2	2	1	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	2	2	1	-	-
White, both sexes	-	-	1	-	1	2	1	2	2	1	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Ohio												
All races, both sexes	4	6	10	21	24	27	34	33	15	11	-	-
Male	3	2	4	6	10	8	9	10	5	3	-	-
Female	1	4	6	15	14	19	25	23	10	8	-	-
White, both sexes	2	6	9	18	23	24	31	32	13	11	-	-
Male	2	2	3	6	10	8	7	10	5	3	-	-
Female	-	4	6	12	13	16	24	22	8	8	-	-
Black, both sexes	2	-	1	3	1	3	3	1	2	-	-	-
Male	1	-	1	-	-	-	2	-	-	-	-	-
Female	1	-	-	3	1	3	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	2	4	6	1	4	5	7	3	1	1	-
Male	1	1	1	4	-	3	1	2	-	1	-	-
Female	1	1	3	2	1	1	4	5	3	-	1	-
White, both sexes	2	1	3	6	1	3	5	6	3	1	1	-
Male	1	1	-	4	-	3	1	2	-	1	-	-
Female	1	-	3	2	1	-	4	4	3	-	1	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Oregon												
All races, both sexes	2	1	4	5	10	15	11	27	7	3	-	-
Male	1	-	3	2	1	4	7	9	-	-	-	-
Female	1	1	1	3	9	11	4	18	7	3	-	-
White, both sexes	1	-	3	5	10	14	11	26	7	3	-	-
Male	1	-	3	2	1	4	7	8	-	-	-	-
Female	-	-	-	3	9	10	4	18	7	3	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	5	6	1	10	19	28	46	33	16	6	1	-
Male	3	1	-	3	7	14	9	4	5	1	-	-
Female	2	5	1	7	12	14	37	29	11	5	1	-
White, both sexes	4	6	1	10	17	28	46	33	16	6	1	-
Male	2	1	-	3	7	14	9	4	5	1	-	-
Female	2	5	1	7	10	14	37	29	11	5	1	-
Black, both sexes	1	-	-	-	2	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Rhode Island												
All races, both sexes	-	1	-	-	1	3	2	4	3	1	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	1	-	-	1	3	1	2	3	1	-	-
White, both sexes	-	1	-	-	1	2	2	4	3	1	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	1	-	-	1	2	1	2	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	4	6	4	4	7	11	6	4	1	-	-
Male	1	1	1	1	-	3	3	-	1	-	-	-
Female	2	3	5	3	4	4	8	6	3	1	-	-
White, both sexes	2	2	4	1	3	7	8	6	4	-	-	-
Male	-	1	1	-	-	3	1	-	1	-	-	-
Female	2	1	3	1	3	4	7	6	3	-	-	-
Black, both sexes	1	2	2	3	1	-	3	-	-	1	-	-
Male	1	-	-	1	-	-	2	-	-	-	-	-
Female	-	2	2	2	1	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
South Dakota												
All races, both sexes	-	1	-	2	-	-	2	3	2	-	-	-
Male	-	-	-	1	-	-	-	2	1	-	-	-
Female	-	1	-	1	-	-	2	1	1	-	-	-
White, both sexes	-	1	-	1	-	-	2	3	2	-	-	-
Male	-	-	-	-	-	-	-	2	1	-	-	-
Female	-	1	-	1	-	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	3	6	8	12	12	18	8	8	4	-	-
Male	-	1	2	2	5	7	4	3	2	-	-	-
Female	-	2	4	6	7	5	14	5	6	4	-	-
White, both sexes	-	2	4	7	11	10	18	8	8	4	-	-
Male	-	1	1	1	5	5	4	3	2	-	-	-
Female	-	1	3	6	6	5	14	5	6	4	-	-
Black, both sexes	-	1	2	1	-	2	-	-	-	-	-	-
Male	-	-	1	1	-	2	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Texas												
All races, both sexes	11	8	16	6	20	17	24	20	11	3	1	-
Male	5	2	5	1	5	4	9	6	2	-	-	-
Female	6	6	11	5	15	13	15	14	9	3	1	-
White, both sexes	8	8	13	5	18	17	23	20	11	3	1	-
Male	3	2	5	1	5	4	8	6	2	-	-	-
Female	5	6	8	4	13	13	15	14	9	3	1	-
Black, both sexes	2	-	2	1	2	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	1	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	2	5	1	4	1	7	3	-	-	-
Male	-	-	-	2	-	-	1	-	2	-	-	-
Female	1	-	2	3	1	4	-	7	1	-	-	-
White, both sexes	1	-	2	5	1	4	1	7	3	-	-	-
Male	-	-	-	2	-	-	1	-	2	-	-	-
Female	1	-	2	3	1	4	-	7	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Vermont												
All races, both sexes	-	-	-	1	1	3	1	5	2	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	1	2	-	4	2	-	-	-
White, both sexes	-	-	-	1	1	3	1	5	2	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	1	2	-	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	1	4	3	5	8	10	8	4	-	1	-
Male	1	1	1	1	1	1	2	2	-	-	-	-
Female	2	-	3	2	4	7	8	6	4	-	1	-
White, both sexes	2	-	3	2	5	7	10	7	3	-	1	-
Male	-	-	-	1	1	1	2	2	-	-	-	-
Female	2	-	3	1	4	6	8	5	3	-	1	-
Black, both sexes	1	1	1	1	-	1	-	1	1	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Washington												
All races, both sexes	2	5	5	5	7	11	17	19	8	6	1	-
Male	2	1	1	2	2	2	3	5	-	1	-	-
Female	-	4	4	3	5	9	14	14	8	5	1	-
White, both sexes	1	2	4	5	7	10	16	19	8	6	1	-
Male	1	-	1	2	2	2	3	5	-	1	-	-
Female	-	2	3	3	5	8	13	14	8	5	1	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	3	-	5	3	4	4	2	3	-	-	-
Male	-	1	-	2	2	2	1	1	1	-	-	-
Female	1	2	-	3	1	2	3	1	2	-	-	-
White, both sexes	1	3	-	5	3	4	4	2	3	-	-	-
Male	-	1	-	2	2	2	1	1	1	-	-	-
Female	1	2	-	3	1	2	3	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)												
Wisconsin												
All races, both sexes	2	1	5	3	7	8	15	15	14	3	-	-
Male	2	1	2	-	5	2	3	2	4	1	-	-
Female	-	-	3	3	2	6	12	13	10	2	-	-
White, both sexes	2	1	4	3	7	8	15	15	14	3	-	-
Male	2	1	2	-	5	2	3	2	4	1	-	-
Female	-	-	2	3	2	6	12	13	10	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	1	-	-	2	2	4	2	-	-	-
Male	-	1	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	4	2	-	-	-
White, both sexes	-	1	1	-	-	2	2	4	2	-	-	-
Male	-	1	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
United States												
All races, both sexes	29	-	-	-	-	-	1	-	-	1	1	-
Male	10	-	-	-	-	-	1	-	-	1	-	-
Female	19	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	24	-	-	-	-	-	-	-	-	1	1	-
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Arkansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Illinois												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Maine												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Massachusetts												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
New Mexico												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
North Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Ohio												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Oregon												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Texas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute rheumatic fever (I00-I02)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
United States												
All races, both sexes	2	2	1	4	6	4	2	2	2	1	-	-
Male	1	-	-	-	4	1	1	1	-	-	-	-
Female	1	2	1	4	2	3	1	1	2	1	-	-
White, both sexes	2	-	1	4	5	3	2	2	2	1	-	-
Male	1	-	-	-	4	1	1	1	-	-	-	-
Female	1	-	1	4	1	2	1	1	2	1	-	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	-	1	-	1	1	-	1	1	1	1	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	1	-	1	-	-	-	-	1	1	-	-
White, both sexes	-	-	-	1	1	-	1	1	1	1	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Colorado												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Florida												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Illinois												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Maine												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Massachusetts												
All races, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Missouri												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
New Mexico												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
North Carolina												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Ohio												
All races, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Oregon												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Texas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Washington												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute rheumatic fever (I00-I02)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
United States												
All races, both sexes	3,336	1	1	1	3	3	5	11	16	25	55	87
Male	1,034	1	-	-	2	1	5	7	7	12	25	35
Female	2,302	-	1	1	1	2	-	4	9	13	30	52
White, both sexes	3,005	1	1	1	3	2	2	7	10	19	36	62
Male	917	1	-	-	2	-	2	5	6	8	19	26
Female	2,088	-	1	1	1	2	-	2	4	11	17	36
Black, both sexes	225	-	-	-	-	1	1	2	6	3	13	19
Male	81	-	-	-	-	1	1	1	1	3	5	7
Female	144	-	-	-	-	-	-	1	5	-	8	12
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	4	3
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	3	2
Asian or Pacific Islander 2/, both sexes	87	-	-	-	-	-	2	2	-	3	2	3
Male	29	-	-	-	-	-	2	1	-	1	-	1
Female	58	-	-	-	-	-	-	1	-	2	2	2
Alabama												
All races, both sexes	45	-	-	-	-	-	-	-	2	1	2	1
Male	15	-	-	-	-	-	-	-	-	1	1	-
Female	30	-	-	-	-	-	-	-	2	-	1	1
White, both sexes	30	-	-	-	-	-	-	-	1	-	2	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	13	-	-	-	-	-	-	-	1	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Alaska												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	56	-	-	-	-	-	-	-	1	-	-	2
Male	16	-	-	-	-	-	-	-	1	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	51	-	-	-	-	-	-	-	1	-	-	1
Male	14	-	-	-	-	-	-	-	1	-	-	1
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Arkansas												
All races, both sexes	21	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	395	-	-	-	1	-	-	1	2	5	5	14
Male	134	-	-	-	1	-	-	1	1	2	2	5
Female	261	-	-	-	-	-	-	-	1	3	3	9
White, both sexes	338	-	-	-	1	-	-	-	1	3	4	10
Male	115	-	-	-	1	-	-	-	1	1	2	4
Female	223	-	-	-	-	-	-	-	-	2	2	6
Black, both sexes	20	-	-	-	-	-	-	1	1	-	1	3
Male	7	-	-	-	-	-	-	1	-	-	-	1
Female	13	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	-	2	-	1
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	24	-	-	-	-	-	-	-	-	1	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Colorado												
All races, both sexes	60	-	-	-	1	-	-	1	1	-	-	1
Male	15	-	-	-	-	-	-	1	1	-	-	-
Female	45	-	-	-	1	-	-	-	-	-	-	1
White, both sexes	57	-	-	-	1	-	-	-	1	-	-	1
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	43	-	-	-	1	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Delaware												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Florida												
All races, both sexes	216	-	-	-	-	-	1	3	1	1	2	5
Male	92	-	-	-	-	-	1	2	-	1	1	3
Female	124	-	-	-	-	-	-	1	1	-	1	2
White, both sexes	202	-	-	-	-	-	-	2	1	1	1	5
Male	85	-	-	-	-	-	-	2	-	1	1	3
Female	117	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	11	-	-	-	-	-	1	1	-	-	1	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	57	-	1	-	-	1	-	1	-	2	1	3
Male	19	-	-	-	-	-	-	1	-	2	1	-
Female	38	-	1	-	-	1	-	-	-	-	-	3
White, both sexes	47	-	1	-	-	1	-	1	-	2	-	1
Male	16	-	-	-	-	-	-	1	-	2	-	-
Female	31	-	1	-	-	1	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	2
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Hawaii												
All races, both sexes	19	-	-	-	-	-	-	1	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	1	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	1	-	-	1	-
Idaho												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Illinois												
All races, both sexes	127	-	-	1	-	-	-	-	1	-	4	5
Male	37	-	-	-	-	-	-	-	-	-	2	3
Female	90	-	-	1	-	-	-	-	1	-	2	2
White, both sexes	111	-	-	1	-	-	-	-	-	-	4	4
Male	35	-	-	-	-	-	-	-	-	-	2	2
Female	76	-	-	1	-	-	-	-	-	-	2	2
Black, both sexes	13	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	66	-	-	-	-	-	1	-	-	-	2	1
Male	14	-	-	-	-	-	1	-	-	-	1	1
Female	52	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	63	-	-	-	-	-	1	-	-	-	1	1
Male	14	-	-	-	-	-	1	-	-	-	1	1
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Iowa												
All races, both sexes	72	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	55	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	55	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Kentucky												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	34	-	-	-	-	-	1	-	-	-	2	4
Male	16	-	-	-	-	-	1	-	-	-	-	2
Female	18	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	24	-	-	-	-	-	-	-	-	-	2	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Maine												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	32	-	-	-	-	1	-	-	1	1	-	1
Male	8	-	-	-	-	1	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	10	-	-	-	-	1	-	-	1	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Massachusetts												
All races, both sexes	64	-	-	-	-	-	-	-	1	2	1	1
Male	12	-	-	-	-	-	-	-	-	1	1	-
Female	52	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	63	-	-	-	-	-	-	-	1	1	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	52	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	97	-	-	-	-	-	-	-	-	1	1	2
Male	27	-	-	-	-	-	-	-	-	1	1	2
Female	70	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	91	-	-	-	-	-	-	-	-	1	1	1
Male	25	-	-	-	-	-	-	-	-	1	1	1
Female	66	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Minnesota												
All races, both sexes	58	-	-	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	39	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	56	-	-	-	-	-	-	-	-	-	1	1
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	38	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Missouri												
All races, both sexes	90	-	-	-	-	-	-	-	-	-	3	2
Male	37	-	-	-	-	-	-	-	-	-	1	1
Female	53	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	82	-	-	-	-	-	-	-	-	-	2	2
Male	34	-	-	-	-	-	-	-	-	-	1	1
Female	48	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Nebraska												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	14	-	-	-	-	-	1	-	-	-	1	1
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
New Hampshire												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	123	-	-	-	-	-	-	-	-	1	-	2
Male	35	-	-	-	-	-	-	-	-	1	-	1
Female	88	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	108	-	-	-	-	-	-	-	-	1	-	-
Male	29	-	-	-	-	-	-	-	-	1	-	-
Female	79	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
New Mexico												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	203	-	-	-	-	1	-	-	-	1	7	7
Male	61	-	-	-	-	-	-	-	-	-	5	2
Female	142	-	-	-	-	1	-	-	-	1	2	5
White, both sexes	185	-	-	-	-	1	-	-	-	1	5	4
Male	56	-	-	-	-	-	-	-	-	-	4	2
Female	129	-	-	-	-	1	-	-	-	1	1	2
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	3
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
North Carolina												
All races, both sexes	91	-	-	-	-	-	-	-	1	-	-	2
Male	29	-	-	-	-	-	-	-	-	-	-	1
Female	62	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	78	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	1	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Ohio												
All races, both sexes	190	1	-	-	1	-	-	-	-	1	2	2
Male	64	1	-	-	1	-	-	-	-	1	1	1
Female	126	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	173	1	-	-	1	-	-	-	-	1	1	2
Male	60	1	-	-	1	-	-	-	-	1	1	1
Female	113	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Oklahoma												
All races, both sexes	38	-	-	-	-	-	-	1	1	-	1	1
Male	16	-	-	-	-	-	-	1	1	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	34	-	-	-	-	-	-	1	1	-	-	-
Male	15	-	-	-	-	-	-	1	1	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Oregon												
All races, both sexes	87	-	-	-	-	-	-	-	-	1	1	1
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	60	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	81	-	-	-	-	-	-	-	-	-	1	1
Male	26	-	-	-	-	-	-	-	-	-	-	1
Female	55	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Pennsylvania												
All races, both sexes	173	-	-	-	-	-	-	-	-	1	1	2
Male	48	-	-	-	-	-	-	-	-	1	-	-
Female	125	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	169	-	-	-	-	-	-	-	-	-	1	2
Male	46	-	-	-	-	-	-	-	-	-	-	-
Female	123	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Rhode Island												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	57	-	-	-	-	-	-	1	1	-	3	3
Male	14	-	-	-	-	-	-	-	1	-	1	1
Female	43	-	-	-	-	-	-	1	-	-	2	2
White, both sexes	41	-	-	-	-	-	-	1	-	-	2	2
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	32	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	16	-	-	-	-	-	-	-	1	-	1	1
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
South Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	87	-	-	-	-	-	-	-	1	2	4	1
Male	30	-	-	-	-	-	-	-	1	-	3	-
Female	57	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	79	-	-	-	-	-	-	-	1	2	3	1
Male	25	-	-	-	-	-	-	-	1	-	2	-
Female	54	-	-	-	-	-	-	-	-	2	1	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Texas												
All races, both sexes	147	-	-	-	-	-	1	1	-	2	2	4
Male	43	-	-	-	-	-	1	1	-	-	-	2
Female	104	-	-	-	-	-	-	-	-	2	2	2
White, both sexes	135	-	-	-	-	-	1	1	-	2	1	3
Male	39	-	-	-	-	-	1	1	-	-	-	1
Female	96	-	-	-	-	-	-	-	-	2	1	2
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Vermont												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	55	-	-	-	-	-	-	-	1	1	2	4
Male	13	-	-	-	-	-	-	-	-	1	1	1
Female	42	-	-	-	-	-	-	-	1	-	1	3
White, both sexes	45	-	-	-	-	-	-	-	1	1	-	3
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	37	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	10	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Washington												
All races, both sexes	87	-	-	-	-	-	-	1	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	68	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	80	-	-	-	-	-	-	1	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	63	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	31	-	-	-	-	-	-	-	-	1	-	6
Male	13	-	-	-	-	-	-	-	-	-	-	3
Female	18	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	31	-	-	-	-	-	-	-	-	1	-	6
Male	13	-	-	-	-	-	-	-	-	-	-	3
Female	18	-	-	-	-	-	-	-	-	1	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Chronic rheumatic heart diseases (I05-I09)												
Wisconsin												
All races, both sexes	77	-	-	-	-	-	-	-	-	-	2	2
Male	24	-	-	-	-	-	-	-	-	-	1	1
Female	53	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	75	-	-	-	-	-	-	-	-	-	1	2
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	13	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
United States												
All races, both sexes	102	158	193	248	355	493	615	528	325	102	9	-
Male	45	74	67	91	115	151	173	130	79	14	-	-
Female	57	84	126	157	240	342	442	398	246	88	9	-
White, both sexes	70	124	159	216	321	457	582	511	312	100	9	-
Male	33	58	58	77	107	143	160	123	75	14	-	-
Female	37	66	101	139	214	314	422	388	237	86	9	-
Black, both sexes	29	22	22	25	16	24	19	12	9	2	-	-
Male	11	10	7	11	2	5	10	4	2	-	-	-
Female	18	12	15	14	14	19	9	8	7	2	-	-
American Indian 1/, both sexes	-	2	-	2	1	1	4	1	1	-	-	-
Male	-	1	-	1	1	1	-	-	1	-	-	-
Female	-	1	-	1	-	-	4	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	10	12	5	17	11	10	4	3	-	-	-
Male	1	5	2	2	5	2	3	3	1	-	-	-
Female	2	5	10	3	12	9	7	1	2	-	-	-
Alabama												
All races, both sexes	3	1	2	2	7	7	9	5	3	-	-	-
Male	-	1	-	1	2	2	5	1	1	-	-	-
Female	3	-	2	1	5	5	4	4	2	-	-	-
White, both sexes	1	-	2	1	5	5	7	4	1	-	-	-
Male	-	-	-	-	1	1	4	-	-	-	-	-
Female	1	-	2	1	4	4	3	4	1	-	-	-
Black, both sexes	2	1	-	1	1	2	1	1	2	-	-	-
Male	-	1	-	1	-	1	1	1	1	-	-	-
Female	2	-	-	-	1	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Alaska												
All races, both sexes	-	-	1	2	1	2	1	-	1	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	1	-	-	-
White, both sexes	-	-	1	2	-	1	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	3	3	5	6	9	13	10	2	1	-	-
Male	1	1	1	3	1	1	4	-	1	1	-	-
Female	-	2	2	2	5	8	9	10	1	-	-	-
White, both sexes	1	2	3	4	6	9	11	10	2	1	-	-
Male	1	-	1	2	1	1	4	-	1	1	-	-
Female	-	2	2	2	5	8	7	10	1	-	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Arkansas												
All races, both sexes	-	1	3	2	3	3	2	4	1	1	-	-
Male	-	1	-	1	1	2	-	1	-	-	-	-
Female	-	-	3	1	2	1	2	3	1	1	-	-
White, both sexes	-	1	3	2	3	3	2	4	-	1	-	-
Male	-	1	-	1	1	2	-	1	-	-	-	-
Female	-	-	3	1	2	1	2	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	10	23	30	27	36	56	70	62	37	15	1	-
Male	6	12	9	12	10	16	17	23	13	4	-	-
Female	4	11	21	15	26	40	53	39	24	11	1	-
White, both sexes	9	16	24	23	26	49	63	60	33	15	1	-
Male	6	8	7	10	9	14	15	22	11	4	-	-
Female	3	8	17	13	17	35	48	38	22	11	1	-
Black, both sexes	1	3	-	1	3	2	3	-	1	-	-	-
Male	-	2	-	-	1	1	1	-	-	-	-	-
Female	1	1	-	1	2	1	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	6	3	7	5	3	2	2	-	-	-
Male	-	2	2	2	-	1	1	1	1	-	-	-
Female	-	2	4	1	7	4	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Colorado												
All races, both sexes	4	3	3	2	7	11	11	9	6	-	-	-
Male	1	2	2	1	1	-	-	4	2	-	-	-
Female	3	1	1	1	6	11	11	5	4	-	-	-
White, both sexes	3	3	3	2	6	11	11	9	6	-	-	-
Male	1	2	2	1	1	-	-	4	2	-	-	-
Female	2	1	1	1	5	11	11	5	4	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	1	3	7	3	8	11	6	2	-	-
Male	1	-	-	1	2	1	3	3	1	-	-	-
Female	-	-	1	2	5	2	5	8	5	2	-	-
White, both sexes	1	-	1	2	7	3	8	11	6	2	-	-
Male	1	-	-	1	2	1	3	3	1	-	-	-
Female	-	-	1	1	5	2	5	8	5	2	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Delaware												
All races, both sexes	-	-	2	1	1	1	2	-	2	-	-	-
Male	-	-	1	1	-	-	2	-	1	-	-	-
Female	-	-	1	-	1	1	-	-	1	-	-	-
White, both sexes	-	-	1	1	1	1	2	-	2	-	-	-
Male	-	-	1	1	-	-	2	-	1	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Florida												
All races, both sexes	8	14	9	19	28	26	42	27	24	6	-	-
Male	3	11	5	10	12	11	17	7	7	1	-	-
Female	5	3	4	9	16	15	25	20	17	5	-	-
White, both sexes	6	13	8	16	26	25	42	27	23	6	-	-
Male	3	10	5	8	10	11	17	7	6	1	-	-
Female	3	3	3	8	16	14	25	20	17	5	-	-
Black, both sexes	2	1	1	2	-	1	-	-	1	-	-	-
Male	-	1	-	2	-	-	-	-	1	-	-	-
Female	2	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	-	-	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	8	3	3	6	9	5	8	4	-	-	-
Male	1	3	1	1	-	6	2	-	1	-	-	-
Female	1	5	2	2	6	3	3	8	3	-	-	-
White, both sexes	1	7	3	3	5	8	4	6	4	-	-	-
Male	-	2	1	1	-	6	2	-	1	-	-	-
Female	1	5	2	2	5	2	2	6	3	-	-	-
Black, both sexes	1	1	-	-	1	1	-	2	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Hawaii												
All races, both sexes	2	2	3	-	2	3	1	3	-	1	-	-
Male	2	1	-	-	-	1	1	2	-	-	-	-
Female	-	1	3	-	2	2	-	1	-	1	-	-
White, both sexes	1	-	-	-	-	1	-	2	-	1	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	3	-	2	2	1	1	-	-	-	-
Male	1	1	-	-	-	1	1	1	-	-	-	-
Female	-	1	3	-	2	1	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	4	3	1	2	4	8	1	-	-	-
Male	-	-	2	1	-	-	1	2	-	-	-	-
Female	-	-	2	2	1	2	3	6	1	-	-	-
White, both sexes	-	-	4	3	1	2	4	8	1	-	-	-
Male	-	-	2	1	-	-	1	2	-	-	-	-
Female	-	-	2	2	1	2	3	6	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Illinois												
All races, both sexes	5	6	7	10	12	19	26	12	13	6	-	-
Male	1	2	2	4	4	3	7	4	5	-	-	-
Female	4	4	5	6	8	16	19	8	8	6	-	-
White, both sexes	2	6	5	7	11	15	25	12	13	6	-	-
Male	1	2	2	3	4	3	7	4	5	-	-	-
Female	1	4	3	4	7	12	18	8	8	6	-	-
Black, both sexes	3	-	2	2	1	4	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	3	-	2	1	1	4	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Indiana												
All races, both sexes	4	3	4	4	5	14	10	7	8	3	-	-
Male	2	-	-	-	1	3	2	-	2	1	-	-
Female	2	3	4	4	4	11	8	7	6	2	-	-
White, both sexes	4	2	4	4	4	14	10	7	8	3	-	-
Male	2	-	-	-	1	3	2	-	2	1	-	-
Female	2	2	4	4	3	11	8	7	6	2	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Iowa												
All races, both sexes	1	2	2	4	7	11	9	17	13	5	-	-
Male	1	1	-	1	1	4	3	2	3	-	-	-
Female	-	1	2	3	6	7	6	15	10	5	-	-
White, both sexes	1	2	2	3	7	11	9	16	13	5	-	-
Male	1	1	-	-	1	4	3	1	3	-	-	-
Female	-	1	2	3	6	7	6	15	10	5	-	-
Black, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	2	-	2	2	8	5	5	1	1	1	-
Male	-	2	-	-	1	2	1	1	-	-	-	-
Female	-	-	-	2	1	6	4	4	1	1	1	-
White, both sexes	-	2	-	2	1	8	5	5	1	1	1	-
Male	-	2	-	-	-	2	1	1	-	-	-	-
Female	-	-	-	2	1	6	4	4	1	1	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Kentucky												
All races, both sexes	1	6	1	6	4	10	9	3	4	2	-	-
Male	1	4	-	2	1	5	2	1	-	-	-	-
Female	-	2	1	4	3	5	7	2	4	2	-	-
White, both sexes	1	6	1	6	4	10	9	3	4	2	-	-
Male	1	4	-	2	1	5	2	1	-	-	-	-
Female	-	2	1	4	3	5	7	2	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	3	4	2	3	5	5	3	1	-	-	-
Male	-	1	3	1	1	2	3	2	-	-	-	-
Female	1	2	1	1	2	3	2	1	1	-	-	-
White, both sexes	1	2	2	1	3	4	3	3	1	-	-	-
Male	-	1	2	-	1	2	1	2	-	-	-	-
Female	1	1	-	1	2	2	2	1	1	-	-	-
Black, both sexes	-	1	2	1	-	1	2	-	-	-	-	-
Male	-	-	1	1	-	-	2	-	-	-	-	-
Female	-	1	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Maine												
All races, both sexes	-	1	4	1	5	2	4	3	1	-	-	-
Male	-	1	2	-	2	1	1	1	-	-	-	-
Female	-	-	2	1	3	1	3	2	1	-	-	-
White, both sexes	-	1	4	1	5	2	4	3	1	-	-	-
Male	-	1	2	-	2	1	1	1	-	-	-	-
Female	-	-	2	1	3	1	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	3	3	2	4	6	4	3	2	-	-	-
Male	-	-	2	1	1	2	-	-	-	-	-	-
Female	1	3	1	1	3	4	4	3	2	-	-	-
White, both sexes	-	1	2	2	4	3	3	3	2	-	-	-
Male	-	-	2	1	1	2	-	-	-	-	-	-
Female	-	1	-	1	3	1	3	3	2	-	-	-
Black, both sexes	1	2	1	-	-	3	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	2	1	-	-	3	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Massachusetts												
All races, both sexes	1	1	5	3	7	13	9	5	12	2	1	-
Male	1	1	1	1	2	3	-	1	-	-	-	-
Female	-	-	4	2	5	10	9	4	12	2	1	-
White, both sexes	1	1	5	3	7	13	9	5	12	2	1	-
Male	1	1	1	1	2	3	-	1	-	-	-	-
Female	-	-	4	2	5	10	9	4	12	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3	4	5	5	12	21	19	14	8	2	-	-
Male	-	1	2	4	3	4	5	2	2	-	-	-
Female	3	3	3	1	9	17	14	12	6	2	-	-
White, both sexes	2	3	5	5	12	20	17	14	8	2	-	-
Male	-	1	2	4	3	4	4	2	2	-	-	-
Female	2	2	3	1	9	16	13	12	6	2	-	-
Black, both sexes	1	1	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Minnesota												
All races, both sexes	2	3	3	-	3	13	16	8	6	2	-	-
Male	1	2	2	-	1	2	3	5	2	-	-	-
Female	1	1	1	-	2	11	13	3	4	2	-	-
White, both sexes	-	3	3	-	3	13	16	8	6	2	-	-
Male	-	2	2	-	1	2	3	5	2	-	-	-
Female	-	1	1	-	2	11	13	3	4	2	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	2	1	5	1	3	6	2	2	-	-	-
Male	-	2	-	2	-	1	1	1	-	-	-	-
Female	-	-	1	3	1	2	5	1	2	-	-	-
White, both sexes	-	1	-	4	1	3	6	1	2	-	-	-
Male	-	1	-	2	-	1	1	-	-	-	-	-
Female	-	-	-	2	1	2	5	1	2	-	-	-
Black, both sexes	-	1	1	1	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Missouri												
All races, both sexes	2	3	1	7	11	12	17	16	11	5	-	-
Male	2	2	-	3	6	6	7	5	4	-	-	-
Female	-	1	1	4	5	6	10	11	7	5	-	-
White, both sexes	1	2	-	7	11	11	16	15	11	4	-	-
Male	1	1	-	3	6	6	6	5	4	-	-	-
Female	-	1	-	4	5	5	10	10	7	4	-	-
Black, both sexes	1	1	1	-	-	1	1	1	-	1	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	2	5	2	2	3	1	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	1	2	4	2	1	2	1	-	-	-
White, both sexes	-	-	1	2	5	2	2	3	1	-	-	-
Male	-	-	-	-	1	-	1	1	-	-	-	-
Female	-	-	1	2	4	2	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Nebraska												
All races, both sexes	-	-	1	2	2	3	3	4	4	-	-	-
Male	-	-	1	-	-	1	1	-	1	-	-	-
Female	-	-	-	2	2	2	2	4	3	-	-	-
White, both sexes	-	-	1	2	2	3	3	4	4	-	-	-
Male	-	-	1	-	-	1	1	-	1	-	-	-
Female	-	-	-	2	2	2	2	4	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	2	2	-	4	1	1	1	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	2	2	-	2	-	1	1	-	-	-
White, both sexes	-	-	2	2	-	4	1	1	1	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	2	2	-	2	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
New Hampshire												
All races, both sexes	-	-	-	1	-	5	-	3	3	-	1	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	5	-	3	2	-	1	-
White, both sexes	-	-	-	1	-	5	-	3	3	-	1	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	5	-	3	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	3	3	12	12	18	32	23	11	3	-	-
Male	2	-	1	4	6	4	6	6	4	-	-	-
Female	1	3	2	8	6	14	26	17	7	3	-	-
White, both sexes	1	3	3	9	8	17	30	22	11	3	-	-
Male	1	-	1	3	4	4	6	5	4	-	-	-
Female	-	3	2	6	4	13	24	17	7	3	-	-
Black, both sexes	1	-	-	3	1	-	1	1	-	-	-	-
Male	1	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	2	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	3	1	1	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	1	-	-	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
New Mexico												
All races, both sexes	2	1	2	3	5	2	5	2	1	-	-	-
Male	-	1	1	1	1	-	2	-	-	-	-	-
Female	2	-	1	2	4	2	3	2	1	-	-	-
White, both sexes	2	1	2	2	5	2	5	2	1	-	-	-
Male	-	1	1	1	1	-	2	-	-	-	-	-
Female	2	-	1	1	4	2	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	5	14	10	12	25	22	44	31	19	5	-	-
Male	-	8	5	3	11	6	15	3	3	-	-	-
Female	5	6	5	9	14	16	29	28	16	5	-	-
White, both sexes	4	8	9	12	25	21	43	30	17	5	-	-
Male	-	5	4	3	11	6	15	3	3	-	-	-
Female	4	3	5	9	14	15	28	27	14	5	-	-
Black, both sexes	1	5	1	-	-	1	-	1	1	-	-	-
Male	-	2	1	-	-	-	-	-	-	-	-	-
Female	1	3	-	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
North Carolina												
All races, both sexes	4	4	5	11	10	16	17	12	9	-	-	-
Male	1	1	3	4	4	6	4	3	2	-	-	-
Female	3	3	2	7	6	10	13	9	7	-	-	-
White, both sexes	2	4	3	10	9	15	15	11	9	-	-	-
Male	1	1	2	3	4	5	4	3	2	-	-	-
Female	1	3	1	7	5	10	11	8	7	-	-	-
Black, both sexes	2	-	2	1	1	1	2	1	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	2	-	1	-	1	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	-	1	2	1	2	2	1	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	2	2	1	-	-
White, both sexes	-	-	1	-	1	2	1	2	2	1	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Ohio												
All races, both sexes	4	6	10	21	23	26	34	33	15	11	-	-
Male	3	2	4	6	9	8	9	10	5	3	-	-
Female	1	4	6	15	14	18	25	23	10	8	-	-
White, both sexes	2	6	9	18	22	23	31	32	13	11	-	-
Male	2	2	3	6	9	8	7	10	5	3	-	-
Female	-	4	6	12	13	15	24	22	8	8	-	-
Black, both sexes	2	-	1	3	1	3	3	1	2	-	-	-
Male	1	-	1	-	-	-	2	-	-	-	-	-
Female	1	-	-	3	1	3	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	1	4	6	1	3	5	7	3	1	1	-
Male	1	1	1	4	-	3	1	2	-	1	-	-
Female	1	-	3	2	1	-	4	5	3	-	1	-
White, both sexes	2	1	3	6	1	3	5	6	3	1	1	-
Male	1	1	-	4	-	3	1	2	-	1	-	-
Female	1	-	3	2	1	-	4	4	3	-	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Oregon												
All races, both sexes	1	1	4	5	10	15	11	27	7	3	-	-
Male	-	-	3	2	1	4	7	9	-	-	-	-
Female	1	1	1	3	9	11	4	18	7	3	-	-
White, both sexes	-	-	3	5	10	14	11	26	7	3	-	-
Male	-	-	3	2	1	4	7	8	-	-	-	-
Female	-	-	-	3	9	10	4	18	7	3	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	5	6	1	9	19	28	45	33	16	6	1	-
Male	3	1	-	3	7	14	9	4	5	1	-	-
Female	2	5	1	6	12	14	36	29	11	5	1	-
White, both sexes	4	6	1	9	17	28	45	33	16	6	1	-
Male	2	1	-	3	7	14	9	4	5	1	-	-
Female	2	5	1	6	10	14	36	29	11	5	1	-
Black, both sexes	1	-	-	-	2	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Rhode Island												
All races, both sexes	-	1	-	-	1	3	2	4	3	1	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	1	-	-	1	3	1	2	3	1	-	-
White, both sexes	-	1	-	-	1	2	2	4	3	1	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	1	-	-	1	2	1	2	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
South Carolina												
All races, both sexes	3	4	6	4	4	6	11	6	4	1	-	-
Male	1	1	1	1	-	3	3	-	1	-	-	-
Female	2	3	5	3	4	3	8	6	3	1	-	-
White, both sexes	2	2	4	1	3	6	8	6	4	-	-	-
Male	-	1	1	-	-	3	1	-	1	-	-	-
Female	2	1	3	1	3	3	7	6	3	-	-	-
Black, both sexes	1	2	2	3	1	-	3	-	-	1	-	-
Male	1	-	-	1	-	-	2	-	-	-	-	-
Female	-	2	2	2	1	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
South Dakota												
All races, both sexes	-	1	-	2	-	-	2	3	2	-	-	-
Male	-	-	-	1	-	-	-	2	1	-	-	-
Female	-	1	-	1	-	-	2	1	1	-	-	-
White, both sexes	-	1	-	1	-	-	2	3	2	-	-	-
Male	-	-	-	-	-	-	-	2	1	-	-	-
Female	-	1	-	1	-	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	3	6	8	12	12	18	8	8	4	-	-
Male	-	1	2	2	5	7	4	3	2	-	-	-
Female	-	2	4	6	7	5	14	5	6	4	-	-
White, both sexes	-	2	4	7	11	10	18	8	8	4	-	-
Male	-	1	1	1	5	5	4	3	2	-	-	-
Female	-	1	3	6	6	5	14	5	6	4	-	-
Black, both sexes	-	1	2	1	-	2	-	-	-	-	-	-
Male	-	-	1	1	-	2	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Texas												
All races, both sexes	11	8	16	6	20	17	24	20	11	3	1	-
Male	5	2	5	1	5	4	9	6	2	-	-	-
Female	6	6	11	5	15	13	15	14	9	3	1	-
White, both sexes	8	8	13	5	18	17	23	20	11	3	1	-
Male	3	2	5	1	5	4	8	6	2	-	-	-
Female	5	6	8	4	13	13	15	14	9	3	1	-
Black, both sexes	2	-	2	1	2	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	1	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	2	5	1	4	1	7	3	-	-	-
Male	-	-	-	2	-	-	1	-	2	-	-	-
Female	1	-	2	3	1	4	-	7	1	-	-	-
White, both sexes	1	-	2	5	1	4	1	7	3	-	-	-
Male	-	-	-	2	-	-	1	-	2	-	-	-
Female	1	-	2	3	1	4	-	7	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Vermont												
All races, both sexes	-	-	-	1	1	3	1	5	2	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	1	2	-	4	2	-	-	-
White, both sexes	-	-	-	1	1	3	1	5	2	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	1	1	2	-	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3	1	4	3	5	8	10	8	4	-	1	-
Male	1	1	1	1	1	1	2	2	-	-	-	-
Female	2	-	3	2	4	7	8	6	4	-	1	-
White, both sexes	2	-	3	2	5	7	10	7	3	-	1	-
Male	-	-	-	1	1	1	2	2	-	-	-	-
Female	2	-	3	1	4	6	8	5	3	-	1	-
Black, both sexes	1	1	1	1	-	1	-	1	1	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Washington												
All races, both sexes	2	5	5	5	7	11	17	19	8	6	1	-
Male	2	1	1	2	2	2	3	5	-	1	-	-
Female	-	4	4	3	5	9	14	14	8	5	1	-
White, both sexes	1	2	4	5	7	10	16	19	8	6	1	-
Male	1	-	1	2	2	2	3	5	-	1	-	-
Female	-	2	3	3	5	8	13	14	8	5	1	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	3	-	4	3	4	4	2	3	-	-	-
Male	-	1	-	2	2	2	1	1	1	-	-	-
Female	1	2	-	2	1	2	3	1	2	-	-	-
White, both sexes	1	3	-	4	3	4	4	2	3	-	-	-
Male	-	1	-	2	2	2	1	1	1	-	-	-
Female	1	2	-	2	1	2	3	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Chronic rheumatic heart diseases (I05-I09)												
Wisconsin												
All races, both sexes	2	1	5	3	7	8	15	15	14	3	-	-
Male	2	1	2	-	5	2	3	2	4	1	-	-
Female	-	-	3	3	2	6	12	13	10	2	-	-
White, both sexes	2	1	4	3	7	8	15	15	14	3	-	-
Male	2	1	2	-	5	2	3	2	4	1	-	-
Female	-	-	2	3	2	6	12	13	10	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	1	-	-	2	2	4	2	-	-	-
Male	-	1	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	4	2	-	-	-
White, both sexes	-	1	1	-	-	2	2	4	2	-	-	-
Male	-	1	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
United States												
All races, both sexes	1,551	1	1	-	1	2	2	6	12	16	33	55
Male	490	1	-	-	1	1	2	4	4	7	19	22
Female	1,061	-	1	-	-	1	-	2	8	9	14	33
White, both sexes	1,383	1	1	-	1	1	1	4	7	12	19	39
Male	433	1	-	-	1	-	1	3	3	4	13	15
Female	950	-	1	-	-	1	-	1	4	8	6	24
Black, both sexes	112	-	-	-	-	1	1	1	5	3	10	12
Male	41	-	-	-	-	1	1	1	1	3	5	5
Female	71	-	-	-	-	-	-	-	4	-	5	7
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	47	-	-	-	-	-	-	1	-	1	2	3
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	36	-	-	-	-	-	-	1	-	1	2	2
Alabama												
All races, both sexes	22	-	-	-	-	-	-	-	1	1	2	-
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	15	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	18	-	-	-	-	-	-	-	1	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Alaska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Arkansas												
All races, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	195	-	-	-	1	-	-	1	2	3	3	9
Male	57	-	-	-	1	-	-	1	1	-	2	4
Female	138	-	-	-	-	-	-	-	1	3	1	5
White, both sexes	164	-	-	-	1	-	-	-	1	2	3	7
Male	47	-	-	-	1	-	-	-	1	-	2	3
Female	117	-	-	-	-	-	-	-	-	2	1	4
Black, both sexes	10	-	-	-	-	-	-	1	1	-	-	1
Male	4	-	-	-	-	-	-	1	-	-	-	1
Female	6	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	1	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Colorado												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Delaware												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Florida												
All races, both sexes	95	-	-	-	-	-	1	1	1	1	2	1
Male	42	-	-	-	-	-	1	1	-	1	1	1
Female	53	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	88	-	-	-	-	-	-	1	1	1	1	1
Male	39	-	-	-	-	-	-	1	-	1	1	1
Female	49	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	6	-	-	-	-	-	1	-	-	-	1	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	31	-	1	-	-	1	-	1	-	1	1	3
Male	10	-	-	-	-	-	-	1	-	1	1	-
Female	21	-	1	-	-	1	-	-	-	-	-	3
White, both sexes	25	-	1	-	-	1	-	1	-	1	-	1
Male	9	-	-	-	-	-	-	1	-	1	-	-
Female	16	-	1	-	-	1	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	2
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Hawaii												
All races, both sexes	9	-	-	-	-	-	-	1	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	1	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	1	-	-	1	-
Idaho												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Illinois												
All races, both sexes	64	-	-	-	-	-	-	-	1	-	2	3
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	43	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	57	-	-	-	-	-	-	-	-	-	2	2
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	37	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Iowa												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Kentucky												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Maine												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	21	-	-	-	-	1	-	-	1	-	-	1
Male	7	-	-	-	-	1	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	1	-	-	1	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Massachusetts												
All races, both sexes	34	-	-	-	-	-	-	-	1	1	-	1
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	25	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	33	-	-	-	-	-	-	-	1	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	47	-	-	-	-	-	-	-	-	1	1	1
Male	15	-	-	-	-	-	-	-	-	1	1	1
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	45	-	-	-	-	-	-	-	-	1	1	1
Male	15	-	-	-	-	-	-	-	-	1	1	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Minnesota												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Missouri												
All races, both sexes	41	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Nebraska												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
New Hampshire												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	63	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	55	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
New Mexico												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	101	-	-	-	-	-	-	-	-	1	4	5
Male	33	-	-	-	-	-	-	-	-	-	4	2
Female	68	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	92	-	-	-	-	-	-	-	-	1	3	3
Male	29	-	-	-	-	-	-	-	-	-	3	2
Female	63	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	2
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
North Carolina												
All races, both sexes	40	-	-	-	-	-	-	-	1	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Ohio												
All races, both sexes	78	1	-	-	-	-	-	-	-	-	1	2
Male	24	1	-	-	-	-	-	-	-	-	-	1
Female	54	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	70	1	-	-	-	-	-	-	-	-	-	2
Male	22	1	-	-	-	-	-	-	-	-	-	1
Female	48	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Oklahoma												
All races, both sexes	17	-	-	-	-	-	-	1	-	-	-	1
Male	7	-	-	-	-	-	-	1	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	16	-	-	-	-	-	-	1	-	-	-	-
Male	7	-	-	-	-	-	-	1	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Oregon												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	40	-	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	59	-	-	-	-	-	-	-	-	1	-	1
Male	17	-	-	-	-	-	-	-	-	1	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	58	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Rhode Island												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	22	-	-	-	-	-	-	-	1	-	2	-
Male	5	-	-	-	-	-	-	-	1	-	1	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	1	-	1	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
South Dakota												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	45	-	-	-	-	-	-	-	1	2	3	1
Male	17	-	-	-	-	-	-	-	1	-	3	-
Female	28	-	-	-	-	-	-	-	-	2	-	1
White, both sexes	39	-	-	-	-	-	-	-	1	2	2	1
Male	14	-	-	-	-	-	-	-	1	-	2	-
Female	25	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Texas												
All races, both sexes	81	-	-	-	-	-	1	-	-	2	2	4
Male	27	-	-	-	-	-	1	-	-	-	-	2
Female	54	-	-	-	-	-	-	-	-	2	2	2
White, both sexes	74	-	-	-	-	-	1	-	-	2	1	3
Male	24	-	-	-	-	-	1	-	-	-	-	1
Female	50	-	-	-	-	-	-	-	-	2	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Vermont												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	32	-	-	-	-	-	-	-	1	1	2	3
Male	7	-	-	-	-	-	-	-	-	1	1	1
Female	25	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	24	-	-	-	-	-	-	-	1	1	-	2
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	20	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	2	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Washington												
All races, both sexes	40	-	-	-	-	-	-	1	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	37	-	-	-	-	-	-	1	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	18	-	-	-	-	-	-	-	-	1	-	3
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	18	-	-	-	-	-	-	-	-	1	-	3
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic mitral valve diseases (I05)												
Wisconsin												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	2	2
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	24	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	30	-	-	-	-	-	-	-	-	-	1	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
United States												
All races, both sexes	48	91	108	121	165	245	282	218	106	31	7	-
Male	18	41	37	45	53	73	91	40	26	5	-	-
Female	30	50	71	76	112	172	191	178	80	26	7	-
White, both sexes	34	75	87	103	153	230	263	214	101	30	7	-
Male	13	33	32	38	52	71	84	39	25	5	-	-
Female	21	42	55	65	101	159	179	175	76	25	7	-
Black, both sexes	13	10	12	14	5	7	10	3	4	1	-	-
Male	5	3	4	5	-	1	5	1	-	-	-	-
Female	8	7	8	9	5	6	5	2	4	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	3	-	1	-	-	-
Male	-	1	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	5	9	4	7	7	6	1	-	-	-	-
Male	-	4	1	2	1	-	2	-	-	-	-	-
Female	1	1	8	2	6	7	4	1	-	-	-	-
Alabama												
All races, both sexes	1	-	1	1	3	3	6	3	-	-	-	-
Male	-	-	-	1	1	1	2	-	-	-	-	-
Female	1	-	1	-	2	2	4	3	-	-	-	-
White, both sexes	-	-	1	-	3	3	5	3	-	-	-	-
Male	-	-	-	-	1	1	2	-	-	-	-	-
Female	-	-	1	-	2	2	3	3	-	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Alaska												
All races, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	3	1	2	3	5	7	3	1	1	-	-
Male	1	1	1	-	1	-	3	-	1	1	-	-
Female	-	2	-	2	2	5	4	3	-	-	-	-
White, both sexes	1	2	1	2	3	5	5	3	1	1	-	-
Male	1	-	1	-	1	-	3	-	1	1	-	-
Female	-	2	-	2	2	5	2	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Arkansas												
All races, both sexes	-	-	1	1	2	1	2	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	1	1	1	2	1	1	-	-	-
White, both sexes	-	-	1	1	2	1	2	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	1	1	1	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	5	13	15	11	18	26	34	28	20	5	1	-
Male	2	7	3	6	3	4	9	7	5	2	-	-
Female	3	6	12	5	15	22	25	21	15	3	1	-
White, both sexes	4	9	11	8	15	21	30	27	19	5	1	-
Male	2	5	2	4	3	4	7	7	4	2	-	-
Female	2	4	9	4	12	17	23	20	15	3	1	-
Black, both sexes	1	2	-	1	1	1	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	4	2	2	4	2	1	-	-	-	-
Male	-	1	1	2	-	-	1	-	-	-	-	-
Female	-	1	3	-	2	4	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Colorado												
All races, both sexes	2	2	1	1	3	6	2	3	4	-	-	-
Male	1	1	1	1	1	-	-	2	2	-	-	-
Female	1	1	-	-	2	6	2	1	2	-	-	-
White, both sexes	2	2	1	1	3	6	2	3	4	-	-	-
Male	1	1	1	1	1	-	-	2	2	-	-	-
Female	1	1	-	-	2	6	2	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	1	1	2	2	4	2	1	-	-
Male	1	-	-	-	-	1	1	1	1	-	-	-
Female	-	-	-	1	1	1	1	3	1	1	-	-
White, both sexes	1	-	-	-	1	2	2	4	2	1	-	-
Male	1	-	-	-	-	1	1	1	1	-	-	-
Female	-	-	-	-	1	1	1	3	1	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Delaware												
All races, both sexes	-	-	2	1	-	-	1	-	-	-	-	-
Male	-	-	1	1	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	1	-	-	1	-	-	-	-	-
Male	-	-	1	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Florida												
All races, both sexes	4	8	4	12	12	15	15	13	4	1	-	-
Male	1	6	2	6	5	6	6	4	1	-	-	-
Female	3	2	2	6	7	9	9	9	3	1	-	-
White, both sexes	3	8	3	9	12	15	15	13	4	1	-	-
Male	1	6	2	4	5	6	6	4	1	-	-	-
Female	2	2	1	5	7	9	9	9	3	1	-	-
Black, both sexes	1	-	1	2	-	-	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	2	1	2	4	6	5	3	-	-	-	-
Male	-	-	1	1	-	3	2	-	-	-	-	-
Female	-	2	-	1	4	3	3	3	-	-	-	-
White, both sexes	-	2	1	2	3	5	4	3	-	-	-	-
Male	-	-	1	1	-	3	2	-	-	-	-	-
Female	-	2	-	1	3	2	2	3	-	-	-	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Hawaii												
All races, both sexes	-	1	2	-	2	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	2	-	2	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	2	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	2	-	2	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	2	1	1	-	1	5	-	-	-	-
Male	-	-	1	-	-	-	-	2	-	-	-	-
Female	-	-	1	1	1	-	1	3	-	-	-	-
White, both sexes	-	-	2	1	1	-	1	5	-	-	-	-
Male	-	-	1	-	-	-	-	2	-	-	-	-
Female	-	-	1	1	1	-	1	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Illinois												
All races, both sexes	2	5	5	8	2	11	13	4	5	3	-	-
Male	-	1	2	3	2	2	5	1	3	-	-	-
Female	2	4	3	5	-	9	8	3	2	3	-	-
White, both sexes	1	5	4	6	2	10	13	4	5	3	-	-
Male	-	1	2	3	2	2	5	1	3	-	-	-
Female	1	4	2	3	-	8	8	3	2	3	-	-
Black, both sexes	1	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	1	2	1	1	6	5	2	1	1	-	-
Male	1	-	-	-	1	2	1	-	-	1	-	-
Female	1	1	2	1	-	4	4	2	1	-	-	-
White, both sexes	2	1	2	1	1	6	5	2	1	1	-	-
Male	1	-	-	-	1	2	1	-	-	1	-	-
Female	1	1	2	1	-	4	4	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Iowa												
All races, both sexes	-	2	2	2	2	4	5	6	4	-	-	-
Male	-	1	-	1	-	2	2	1	1	-	-	-
Female	-	1	2	1	2	2	3	5	3	-	-	-
White, both sexes	-	2	2	1	2	4	5	5	4	-	-	-
Male	-	1	-	-	-	2	2	-	1	-	-	-
Female	-	1	2	1	2	2	3	5	3	-	-	-
Black, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	-	1	1	4	2	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	3	2	-	-	-	-	-
White, both sexes	-	1	-	1	1	4	2	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	3	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Kentucky												
All races, both sexes	-	2	1	3	-	3	6	-	-	-	-	-
Male	-	1	-	1	-	2	1	-	-	-	-	-
Female	-	1	1	2	-	1	5	-	-	-	-	-
White, both sexes	-	2	1	3	-	3	6	-	-	-	-	-
Male	-	1	-	1	-	2	1	-	-	-	-	-
Female	-	1	1	2	-	1	5	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	2	3	1	1	4	2	1	-	-	-	-
Male	-	1	2	-	-	1	2	1	-	-	-	-
Female	1	1	1	1	1	3	-	-	-	-	-	-
White, both sexes	1	2	1	1	1	3	-	1	-	-	-	-
Male	-	1	1	-	-	1	-	1	-	-	-	-
Female	1	1	-	1	1	2	-	-	-	-	-	-
Black, both sexes	-	-	2	-	-	1	2	-	-	-	-	-
Male	-	-	1	-	-	-	2	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Maine												
All races, both sexes	-	1	4	1	3	-	1	1	-	-	-	-
Male	-	1	2	-	2	-	-	-	-	-	-	-
Female	-	-	2	1	1	-	1	1	-	-	-	-
White, both sexes	-	1	4	1	3	-	1	1	-	-	-	-
Male	-	1	2	-	2	-	-	-	-	-	-	-
Female	-	-	2	1	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	2	2	2	4	4	3	-	1	-	-	-
Male	-	-	1	1	1	2	-	-	-	-	-	-
Female	-	2	1	1	3	2	3	-	1	-	-	-
White, both sexes	-	-	1	2	4	3	2	-	1	-	-	-
Male	-	-	1	1	1	2	-	-	-	-	-	-
Female	-	-	-	1	3	1	2	-	1	-	-	-
Black, both sexes	-	2	1	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	1	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Massachusetts												
All races, both sexes	1	1	4	2	4	7	3	5	3	-	1	-
Male	1	1	1	1	2	1	-	1	-	-	-	-
Female	-	-	3	1	2	6	3	4	3	-	1	-
White, both sexes	1	1	4	2	4	7	3	5	3	-	1	-
Male	1	1	1	1	2	1	-	1	-	-	-	-
Female	-	-	3	1	2	6	3	4	3	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	2	4	4	6	8	10	7	2	-	-	-
Male	-	-	1	4	2	1	3	-	1	-	-	-
Female	1	2	3	-	4	7	7	7	1	-	-	-
White, both sexes	1	1	4	4	6	8	9	7	2	-	-	-
Male	-	-	1	4	2	1	3	-	1	-	-	-
Female	1	1	3	-	4	7	6	7	1	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Minnesota												
All races, both sexes	1	3	2	-	-	6	6	1	2	1	-	-
Male	-	2	1	-	-	1	2	1	-	-	-	-
Female	1	1	1	-	-	5	4	-	2	1	-	-
White, both sexes	-	3	2	-	-	6	6	1	2	1	-	-
Male	-	2	1	-	-	1	2	1	-	-	-	-
Female	-	1	1	-	-	5	4	-	2	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	2	1	2	1	3	4	1	1	-	-	-
Male	-	2	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	2	1	2	3	1	1	-	-	-
White, both sexes	-	1	-	1	1	3	4	1	1	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	1	2	3	1	1	-	-	-
Black, both sexes	-	1	1	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Missouri												
All races, both sexes	1	2	1	3	5	8	10	5	3	2	-	-
Male	1	1	-	1	3	5	4	2	2	-	-	-
Female	-	1	1	2	2	3	6	3	1	2	-	-
White, both sexes	-	2	-	3	5	8	9	5	3	1	-	-
Male	-	1	-	1	3	5	3	2	2	-	-	-
Female	-	1	-	2	2	3	6	3	1	1	-	-
Black, both sexes	1	-	1	-	-	-	1	-	-	1	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	1	3	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	2	1	-	1	-	-	-	-
White, both sexes	-	-	-	1	3	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	2	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Nebraska												
All races, both sexes	-	-	-	1	2	2	1	1	2	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	2	1	-	1	1	-	-	-
White, both sexes	-	-	-	1	2	2	1	1	2	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	2	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	1	2	-	3	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	2	-	2	-	-	-	-	-	-
White, both sexes	-	-	1	2	-	3	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	2	-	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
New Hampshire												
All races, both sexes	-	-	-	1	-	2	-	-	2	-	1	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	2	-	-	1	-	1	-
White, both sexes	-	-	-	1	-	2	-	-	2	-	1	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	2	-	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	3	2	3	5	12	17	12	5	2	-	-
Male	1	-	1	1	2	3	4	2	2	-	-	-
Female	-	3	1	2	3	9	13	10	3	2	-	-
White, both sexes	-	3	2	2	3	11	15	12	5	2	-	-
Male	-	-	1	1	1	3	4	2	2	-	-	-
Female	-	3	1	1	2	8	11	10	3	2	-	-
Black, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
New Mexico												
All races, both sexes	-	1	-	1	2	2	3	1	-	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	2	2	2	1	-	-	-	-
White, both sexes	-	1	-	1	2	2	3	1	-	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	2	2	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	8	6	5	12	9	22	18	6	2	-	-
Male	-	4	4	2	4	3	7	2	1	-	-	-
Female	3	4	2	3	8	6	15	16	5	2	-	-
White, both sexes	3	5	5	5	12	9	22	17	5	2	-	-
Male	-	2	3	2	4	3	7	2	1	-	-	-
Female	3	3	2	3	8	6	15	15	4	2	-	-
Black, both sexes	-	2	1	-	-	-	-	1	1	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
North Carolina												
All races, both sexes	2	3	1	6	4	4	11	3	3	-	-	-
Male	-	1	-	1	-	1	2	1	-	-	-	-
Female	2	2	1	5	4	3	9	2	3	-	-	-
White, both sexes	1	3	1	6	3	4	9	3	3	-	-	-
Male	-	1	-	1	-	1	2	1	-	-	-	-
Female	1	2	1	5	3	3	7	2	3	-	-	-
Black, both sexes	1	-	-	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	-	1	1	1	1	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	1	-	-
White, both sexes	-	-	1	-	1	1	1	1	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Ohio												
All races, both sexes	2	2	6	10	11	9	12	15	1	6	-	-
Male	1	-	3	2	4	4	5	2	-	1	-	-
Female	1	2	3	8	7	5	7	13	1	5	-	-
White, both sexes	1	2	5	8	10	9	11	15	-	6	-	-
Male	1	-	2	2	4	4	4	2	-	1	-	-
Female	-	2	3	6	6	5	7	13	-	5	-	-
Black, both sexes	1	-	1	2	1	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	-	2	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	1	2	3	-	2	2	2	1	-	1	-
Male	-	1	-	3	-	2	-	-	-	-	-	-
Female	1	-	2	-	-	-	2	2	1	-	1	-
White, both sexes	1	1	2	3	-	2	2	2	1	-	1	-
Male	-	1	-	3	-	2	-	-	-	-	-	-
Female	1	-	2	-	-	-	2	2	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Oregon												
All races, both sexes	1	-	3	2	5	8	6	10	4	1	-	-
Male	-	-	2	-	1	3	4	3	-	-	-	-
Female	1	-	1	2	4	5	2	7	4	1	-	-
White, both sexes	-	-	2	2	5	8	6	10	4	1	-	-
Male	-	-	2	-	1	3	4	3	-	-	-	-
Female	-	-	-	2	4	5	2	7	4	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	3	-	3	7	11	10	13	5	1	1	-
Male	1	-	-	2	5	4	2	-	2	-	-	-
Female	2	3	-	1	2	7	8	13	3	1	1	-
White, both sexes	3	3	-	3	7	11	10	13	5	1	1	-
Male	1	-	-	2	5	4	2	-	2	-	-	-
Female	2	3	-	1	2	7	8	13	3	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Rhode Island												
All races, both sexes	-	-	-	-	1	2	2	-	3	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	2	1	-	3	-	-	-
White, both sexes	-	-	-	-	1	1	2	-	3	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	1	4	1	2	3	3	3	1	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	1	1	3	1	2	2	2	3	1	-	-	-
White, both sexes	1	-	4	-	2	3	3	3	1	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	1	-	3	-	2	2	2	3	1	-	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
South Dakota												
All races, both sexes	-	1	-	-	-	-	2	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	2	1	1	-	-	-
White, both sexes	-	1	-	-	-	-	2	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	2	4	6	8	9	6	2	-	1	-	-
Male	-	1	-	2	3	5	2	-	-	-	-	-
Female	-	1	4	4	5	4	4	2	-	1	-	-
White, both sexes	-	1	3	5	7	8	6	2	-	1	-	-
Male	-	1	-	1	3	4	2	-	-	-	-	-
Female	-	-	3	4	4	4	4	2	-	1	-	-
Black, both sexes	-	1	1	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Texas												
All races, both sexes	7	5	11	2	12	9	15	6	4	1	-	-
Male	4	1	4	-	3	1	8	2	1	-	-	-
Female	3	4	7	2	9	8	7	4	3	1	-	-
White, both sexes	4	5	10	2	11	9	15	6	4	1	-	-
Male	2	1	4	-	3	1	8	2	1	-	-	-
Female	2	4	6	2	8	8	7	4	3	1	-	-
Black, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	1	1	-	2	1	5	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	1	-	1	-	-	2	-	5	-	-	-	-
White, both sexes	1	-	1	1	-	2	1	5	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	1	-	1	-	-	2	-	5	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Vermont												
All races, both sexes	-	-	-	1	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	1	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	1	3	2	6	6	3	2	-	1	-
Male	-	-	1	1	1	1	-	-	-	-	-	-
Female	1	-	-	2	1	5	6	3	2	-	1	-
White, both sexes	1	-	-	2	2	5	6	2	1	-	1	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	1	-	-	1	1	4	6	2	1	-	1	-
Black, both sexes	-	-	1	1	-	1	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Washington												
All races, both sexes	2	1	1	3	5	5	5	11	4	1	1	-
Male	2	1	-	1	1	-	1	2	-	-	-	-
Female	-	-	1	2	4	5	4	9	4	1	1	-
White, both sexes	1	-	1	3	5	4	5	11	4	1	1	-
Male	1	-	-	1	1	-	1	2	-	-	-	-
Female	-	-	1	2	4	4	4	9	4	1	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	3	-	1	2	3	1	2	2	-	-	-
Male	-	1	-	-	1	2	-	1	1	-	-	-
Female	-	2	-	1	1	1	1	1	1	-	-	-
White, both sexes	-	3	-	1	2	3	1	2	2	-	-	-
Male	-	1	-	-	1	2	-	1	1	-	-	-
Female	-	2	-	1	1	1	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic mitral valve diseases (I05)												
Wisconsin												
All races, both sexes	-	1	2	2	2	4	6	7	4	-	-	-
Male	-	1	1	-	1	1	1	1	-	-	-	-
Female	-	-	1	2	1	3	5	6	4	-	-	-
White, both sexes	-	1	1	2	2	4	6	7	4	-	-	-
Male	-	1	1	-	1	1	1	1	-	-	-	-
Female	-	-	-	2	1	3	5	6	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	-	-	-	2	1	2	1	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-
White, both sexes	-	1	-	-	-	2	1	2	1	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
United States												
All races, both sexes	96	-	-	-	-	-	-	-	1	-	3	1
Male	31	-	-	-	-	-	-	-	-	-	2	1
Female	65	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	91	-	-	-	-	-	-	-	-	-	3	1
Male	30	-	-	-	-	-	-	-	-	-	2	1
Female	61	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Colorado												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Florida												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Illinois												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Iowa												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Maine												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Missouri												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Nebraska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
New Mexico												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
North Carolina												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Ohio												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Oregon												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Texas												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Washington												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Rheumatic aortic valve diseases (I06)												
Wisconsin												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
United States												
All races, both sexes	3	3	8	5	9	6	24	18	10	5	-	-
Male	2	-	4	2	3	3	7	4	2	1	-	-
Female	1	3	4	3	6	3	17	14	8	4	-	-
White, both sexes	2	2	8	5	9	5	23	18	10	5	-	-
Male	1	-	4	2	3	3	7	4	2	1	-	-
Female	1	2	4	3	6	2	16	14	8	4	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Alabama												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	1	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Arkansas												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1	-	2	3	-	2	4	3	1	1	-	-
Male	1	-	2	1	-	1	2	1	-	1	-	-
Female	-	-	-	2	-	1	2	2	1	-	-	-
White, both sexes	1	-	2	3	-	2	3	3	1	1	-	-
Male	1	-	2	1	-	1	2	1	-	1	-	-
Female	-	-	-	2	-	1	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Colorado												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Delaware												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Florida												
All races, both sexes	-	1	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	1	-	-	-
White, both sexes	-	1	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	-	1	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Illinois												
All races, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Iowa												
All races, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Kentucky												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Maine												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Massachusetts												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Minnesota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Missouri												
All races, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Nebraska												
All races, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	1	-	-	2	-	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	2	-	-	-	-
White, both sexes	-	-	-	-	2	-	1	2	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	2	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
North Carolina												
All races, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Ohio												
All races, both sexes	-	-	-	-	1	-	2	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	2	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Oregon												
All races, both sexes	-	-	-	-	-	1	-	3	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	3	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	3	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	-	-	-	1	-	3	2	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	2	2	-	-	-	-
White, both sexes	-	-	-	-	1	-	3	2	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	2	2	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Texas												
All races, both sexes	-	-	-	-	-	-	2	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	1	-	-
White, both sexes	-	-	-	-	-	-	2	1	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	-	-	1	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	1	-	-	-	-
White, both sexes	-	-	1	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Washington												
All races, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Rheumatic aortic valve diseases (I06)												
Wisconsin												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
United States												
All races, both sexes	767	-	-	1	1	-	-	1	1	2	4	11
Male	265	-	-	-	-	-	-	1	1	-	-	8
Female	502	-	-	1	1	-	-	-	-	2	4	3
White, both sexes	723	-	-	1	1	-	-	1	1	2	3	9
Male	245	-	-	-	-	-	-	1	1	-	-	6
Female	478	-	-	1	1	-	-	-	-	2	3	3
Black, both sexes	35	-	-	-	-	-	-	-	-	-	1	2
Male	17	-	-	-	-	-	-	-	-	-	-	2
Female	18	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Alaska												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	13	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	1	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	1	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Arkansas												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	82	-	-	-	-	-	-	-	-	-	2	3
Male	34	-	-	-	-	-	-	-	-	-	-	1
Female	48	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	75	-	-	-	-	-	-	-	-	-	1	3
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Colorado												
All races, both sexes	16	-	-	-	1	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	1	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Delaware												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Florida												
All races, both sexes	45	-	-	-	-	-	-	-	-	-	-	2
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	44	-	-	-	-	-	-	-	-	-	-	2
Male	23	-	-	-	-	-	-	-	-	-	-	2
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Hawaii												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Illinois												
All races, both sexes	31	-	-	1	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	1	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Iowa												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Kentucky												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Maine												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Massachusetts												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Minnesota												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Missouri												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Nebraska												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
New Hampshire												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
North Carolina												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Ohio												
All races, both sexes	57	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	53	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Oregon												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	61	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	61	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Rhode Island												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Texas												
All races, both sexes	19	-	-	-	-	-	-	1	-	-	-	-
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	1	-	-	-	-
Male	8	-	-	-	-	-	-	1	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Vermont												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Washington												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Disorders of both mitral and aortic valves (I08.0)												
Wisconsin												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
United States												
All races, both sexes	22	22	27	43	64	116	147	151	118	34	2	-
Male	12	13	14	20	20	45	44	56	27	4	-	-
Female	10	9	13	23	44	71	103	95	91	30	2	-
White, both sexes	16	18	21	40	58	109	142	148	118	33	2	-
Male	10	11	11	19	18	42	40	55	27	4	-	-
Female	6	7	10	21	40	67	102	93	91	29	2	-
Black, both sexes	5	2	6	2	3	6	4	3	-	1	-	-
Male	1	2	3	1	1	2	4	1	-	-	-	-
Female	4	-	3	1	2	4	-	2	-	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	3	1	1	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	1	-	1	2	-	1	-	-	-	-	-
Alabama												
All races, both sexes	1	-	1	-	2	2	3	-	1	-	-	-
Male	-	-	-	-	-	-	3	-	-	-	-	-
Female	1	-	1	-	2	2	-	-	1	-	-	-
White, both sexes	1	-	1	-	2	2	2	-	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	1	-	1	-	2	2	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Alaska												
All races, both sexes	-	-	-	1	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	1	-	-	-
White, both sexes	-	-	-	1	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	-	1	2	-	2	5	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	2	5	1	-	-	-
White, both sexes	-	-	-	1	2	-	2	5	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	2	5	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Arkansas												
All races, both sexes	-	1	-	1	1	1	-	1	-	1	-	-
Male	-	1	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
White, both sexes	-	1	-	1	1	1	-	1	-	1	-	-
Male	-	1	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2	3	6	3	6	14	14	16	10	3	-	-
Male	1	2	2	2	2	6	3	9	6	-	-	-
Female	1	1	4	1	4	8	11	7	4	3	-	-
White, both sexes	2	2	6	2	3	13	14	16	10	3	-	-
Male	1	2	2	2	2	5	3	9	6	-	-	-
Female	1	-	4	-	1	8	11	7	4	3	-	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	2	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Colorado												
All races, both sexes	-	1	1	1	2	2	3	3	2	-	-	-
Male	-	1	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	2	2	3	2	2	-	-	-
White, both sexes	-	1	1	1	2	2	3	3	2	-	-	-
Male	-	1	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	2	2	3	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	1	-	3	1	3	4	4	1	-	-
Male	-	-	-	-	2	-	1	1	-	-	-	-
Female	-	-	1	-	1	1	2	3	4	1	-	-
White, both sexes	-	-	1	-	3	1	3	4	4	1	-	-
Male	-	-	-	-	2	-	1	1	-	-	-	-
Female	-	-	1	-	1	1	2	3	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Delaware												
All races, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Florida												
All races, both sexes	1	2	1	2	4	4	12	7	7	3	-	-
Male	1	2	1	2	2	3	5	3	2	1	-	-
Female	-	-	-	-	2	1	7	4	5	2	-	-
White, both sexes	1	2	1	2	3	4	12	7	7	3	-	-
Male	1	2	1	2	1	3	5	3	2	1	-	-
Female	-	-	-	-	2	1	7	4	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	1	-	1	1	1	-	2	4	-	-	-
Male	-	1	-	-	-	1	-	-	1	-	-	-
Female	1	-	-	1	1	-	-	2	3	-	-	-
White, both sexes	1	-	-	1	1	1	-	2	4	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	1	-	-	1	1	-	-	2	3	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Hawaii												
All races, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Illinois												
All races, both sexes	3	1	1	1	2	4	7	4	5	2	-	-
Male	1	1	-	-	1	1	2	1	1	-	-	-
Female	2	-	1	1	1	3	5	3	4	2	-	-
White, both sexes	1	1	-	1	2	3	7	4	5	2	-	-
Male	1	1	-	-	1	1	2	1	1	-	-	-
Female	-	-	-	1	1	2	5	3	4	2	-	-
Black, both sexes	2	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	1	-	-	1	4	2	4	3	-	-	-
Male	1	-	-	-	-	1	-	-	1	-	-	-
Female	-	1	-	-	1	3	2	4	2	-	-	-
White, both sexes	1	1	-	-	1	4	2	4	3	-	-	-
Male	1	-	-	-	-	1	-	-	1	-	-	-
Female	-	1	-	-	1	3	2	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Iowa												
All races, both sexes	-	-	-	2	2	4	1	5	5	2	-	-
Male	-	-	-	-	-	1	-	-	2	-	-	-
Female	-	-	-	2	2	3	1	5	3	2	-	-
White, both sexes	-	-	-	2	2	4	1	5	5	2	-	-
Male	-	-	-	-	-	1	-	-	2	-	-	-
Female	-	-	-	2	2	3	1	5	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	-	-	-	1	1	1	2	1	-	1	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	1	-
White, both sexes	-	-	-	-	-	1	1	2	1	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Kentucky												
All races, both sexes	-	1	-	1	-	4	-	1	3	1	-	-
Male	-	1	-	1	-	3	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	3	1	-	-
White, both sexes	-	1	-	1	-	4	-	1	3	1	-	-
Male	-	1	-	1	-	3	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	-	-	-	1	1	2	2	-	-	-	-
Male	-	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	1	1	2	2	-	-	-	-
Male	-	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Maine												
All races, both sexes	-	-	-	-	-	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Massachusetts												
All races, both sexes	-	-	-	-	-	2	2	-	5	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	5	-	-	-
White, both sexes	-	-	-	-	-	2	2	-	5	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	1	-	1	3	10	6	3	4	1	-	-
Male	-	-	-	-	1	3	1	-	1	-	-	-
Female	1	1	-	1	2	7	5	3	3	1	-	-
White, both sexes	1	1	-	1	3	10	5	3	4	1	-	-
Male	-	-	-	-	1	3	-	-	1	-	-	-
Female	1	1	-	1	2	7	5	3	3	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Minnesota												
All races, both sexes	-	-	1	-	1	3	2	2	1	1	-	-
Male	-	-	1	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	1	3	2	-	1	1	-	-
White, both sexes	-	-	1	-	1	3	2	2	1	1	-	-
Male	-	-	1	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	1	3	2	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	2	-	-	1	1	1	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	2	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Missouri												
All races, both sexes	1	-	-	3	2	2	7	7	4	3	-	-
Male	1	-	-	1	1	1	3	2	1	-	-	-
Female	-	-	-	2	1	1	4	5	3	3	-	-
White, both sexes	1	-	-	3	2	2	7	7	4	3	-	-
Male	1	-	-	1	1	1	3	2	1	-	-	-
Female	-	-	-	2	1	1	4	5	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Nebraska												
All races, both sexes	-	-	-	-	-	1	1	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	2	-	-	-
White, both sexes	-	-	-	-	-	1	1	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	-	-	-	1	3	1	3	7	2	1	-	-
Male	-	-	-	-	2	1	-	3	1	-	-	-
Female	-	-	-	1	1	-	3	4	1	1	-	-
White, both sexes	-	-	-	-	3	1	3	7	2	1	-	-
Male	-	-	-	-	2	1	-	3	1	-	-	-
Female	-	-	-	-	1	-	3	4	1	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
New Mexico												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	1	1	2	2	6	9	5	4	2	-	-
Male	-	1	1	-	1	2	3	1	1	-	-	-
Female	2	-	-	2	1	4	6	4	3	2	-	-
White, both sexes	1	1	1	2	2	6	8	5	4	2	-	-
Male	-	1	1	-	1	2	3	1	1	-	-	-
Female	1	-	-	2	1	4	5	4	3	2	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
North Carolina												
All races, both sexes	1	1	4	2	2	8	2	5	4	-	-	-
Male	1	-	3	2	1	3	1	1	1	-	-	-
Female	-	1	1	-	1	5	1	4	3	-	-	-
White, both sexes	1	1	2	1	2	8	2	4	4	-	-	-
Male	1	-	2	1	1	3	1	1	1	-	-	-
Female	-	1	-	-	1	5	1	3	3	-	-	-
Black, both sexes	-	-	2	1	-	-	-	1	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	1	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	2	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Ohio												
All races, both sexes	1	1	-	7	5	9	11	13	8	2	-	-
Male	1	-	-	3	2	3	4	5	3	1	-	-
Female	-	1	-	4	3	6	7	8	5	1	-	-
White, both sexes	1	1	-	7	5	7	10	12	8	2	-	-
Male	1	-	-	3	2	3	3	5	3	1	-	-
Female	-	1	-	4	3	4	7	7	5	1	-	-
Black, both sexes	-	-	-	-	-	2	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	-	2	1	-	1	2	2	1	1	-	-
Male	-	-	1	-	-	1	1	2	-	1	-	-
Female	-	-	1	1	-	-	1	-	1	-	-	-
White, both sexes	-	-	1	1	-	1	2	2	1	1	-	-
Male	-	-	-	-	-	1	1	2	-	1	-	-
Female	-	-	1	1	-	-	1	-	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Oregon												
All races, both sexes	-	-	1	2	3	1	4	8	-	1	-	-
Male	-	-	1	1	-	-	3	4	-	-	-	-
Female	-	-	-	1	3	1	1	4	-	1	-	-
White, both sexes	-	-	1	2	3	1	4	8	-	1	-	-
Male	-	-	1	1	-	-	3	4	-	-	-	-
Female	-	-	-	1	3	1	1	4	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	3	-	3	4	11	19	9	5	5	-	-
Male	1	1	-	1	-	6	4	2	1	1	-	-
Female	-	2	-	2	4	5	15	7	4	4	-	-
White, both sexes	1	3	-	3	4	11	19	9	5	5	-	-
Male	1	1	-	1	-	6	4	2	1	1	-	-
Female	-	2	-	2	4	5	15	7	4	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	4	-	1	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	4	-	1	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	1	1	-	1	1	3	1	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	1	-	1	1	2	1	1	1	-	-
White, both sexes	-	1	-	-	-	1	2	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	2	1	1	-	-	-
Black, both sexes	-	-	1	-	1	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	-	1	-	1	3	5	3	6	1	-	-
Male	-	-	1	-	1	2	2	1	1	-	-	-
Female	-	-	-	-	-	1	3	2	5	1	-	-
White, both sexes	-	-	-	-	1	2	5	3	6	1	-	-
Male	-	-	-	-	1	1	2	1	1	-	-	-
Female	-	-	-	-	-	1	3	2	5	1	-	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Texas												
All races, both sexes	1	1	1	1	2	1	1	4	5	-	1	-
Male	-	1	-	1	-	1	-	3	1	-	-	-
Female	1	-	1	-	2	-	1	1	4	-	1	-
White, both sexes	1	1	1	1	2	1	1	4	5	-	1	-
Male	-	1	-	1	-	1	-	3	1	-	-	-
Female	1	-	1	-	2	-	1	1	4	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	1	1	-	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-
White, both sexes	-	-	-	1	1	-	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Vermont												
All races, both sexes	-	-	-	-	-	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	-	-	1	1	2	3	2	-	-	-
Male	1	1	-	-	-	-	2	2	-	-	-	-
Female	-	-	-	-	1	1	-	1	2	-	-	-
White, both sexes	-	-	-	-	1	1	2	3	2	-	-	-
Male	-	-	-	-	-	-	2	2	-	-	-	-
Female	-	-	-	-	1	1	-	1	2	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Washington												
All races, both sexes	-	1	2	-	2	3	4	3	3	1	-	-
Male	-	-	1	-	1	1	1	2	-	-	-	-
Female	-	1	1	-	1	2	3	1	3	1	-	-
White, both sexes	-	-	2	-	2	3	4	3	3	1	-	-
Male	-	-	1	-	1	1	1	2	-	-	-	-
Female	-	-	1	-	1	2	3	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	2	-	1	3	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	2	-	1	-	-	-
White, both sexes	-	-	-	2	-	1	3	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	2	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Disorders of both mitral and aortic valves (I08.0)												
Wisconsin												
All races, both sexes	1	-	-	-	1	2	6	4	6	-	-	-
Male	1	-	-	-	1	-	1	1	2	-	-	-
Female	-	-	-	-	-	2	5	3	4	-	-	-
White, both sexes	1	-	-	-	1	2	6	4	6	-	-	-
Male	1	-	-	-	1	-	1	1	2	-	-	-
Female	-	-	-	-	-	2	5	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
United States												
All races, both sexes	922	-	-	-	1	1	3	4	2	7	15	20
Male	248	-	-	-	1	-	3	2	2	5	4	4
Female	674	-	-	-	-	1	-	2	-	2	11	16
White, both sexes	808	-	-	-	1	1	1	2	2	5	11	13
Male	209	-	-	-	1	-	1	1	2	4	4	4
Female	599	-	-	-	-	1	-	1	-	1	7	9
Black, both sexes	75	-	-	-	-	-	-	1	-	-	2	5
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	1	-	-	2	5
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	2	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	2	2
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	2	1	-	2	-	-
Male	15	-	-	-	-	-	2	1	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	1	-	-
Alabama												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Alaska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Arkansas												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	101	-	-	-	-	-	-	-	-	2	-	2
Male	34	-	-	-	-	-	-	-	-	2	-	-
Female	67	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	83	-	-	-	-	-	-	-	-	1	-	-
Male	26	-	-	-	-	-	-	-	-	1	-	-
Female	57	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Colorado												
All races, both sexes	19	-	-	-	-	-	-	1	1	-	-	1
Male	3	-	-	-	-	-	-	1	1	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	16	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Delaware												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Florida												
All races, both sexes	72	-	-	-	-	-	-	2	-	-	-	2
Male	25	-	-	-	-	-	-	1	-	-	-	-
Female	47	-	-	-	-	-	-	1	-	-	-	2
White, both sexes	66	-	-	-	-	-	-	1	-	-	-	2
Male	22	-	-	-	-	-	-	1	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Hawaii												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Illinois												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	2	2
Male	6	-	-	-	-	-	-	-	-	-	1	2
Female	24	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	2	2
Male	5	-	-	-	-	-	-	-	-	-	1	2
Female	20	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	26	-	-	-	-	-	1	-	-	-	2	-
Male	3	-	-	-	-	-	1	-	-	-	1	-
Female	23	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	23	-	-	-	-	-	1	-	-	-	1	-
Male	3	-	-	-	-	-	1	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Iowa												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Kentucky												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	10	-	-	-	-	-	1	-	-	-	2	1
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	6	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Maine												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Massachusetts												
All races, both sexes	21	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Minnesota												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Missouri												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Nebraska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	-	-	-	-	-	1	-	-	-	1	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
New Hampshire												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	39	-	-	-	-	-	-	-	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
New Mexico												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	61	-	-	-	-	1	-	-	-	-	2	2
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	1	-	-	-	-	2	2
White, both sexes	55	-	-	-	-	1	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	1	-	-	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
North Carolina												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Ohio												
All races, both sexes	51	-	-	-	1	-	-	-	-	1	1	-
Male	16	-	-	-	1	-	-	-	-	1	1	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	1	-	-	-	-	1	1	-
Male	15	-	-	-	1	-	-	-	-	1	1	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	11	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Oregon												
All races, both sexes	20	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Pennsylvania												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	44	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Rhode Island												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	23	-	-	-	-	-	-	1	-	-	-	3
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	1	-	-	-	2
White, both sexes	15	-	-	-	-	-	-	1	-	-	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
South Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Texas												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	37	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Vermont												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Washington												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Wisconsin												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
United States												
All races, both sexes	29	42	50	79	117	126	162	141	91	32	-	-
Male	13	20	12	24	39	30	31	30	24	4	-	-
Female	16	22	38	55	78	96	131	111	67	28	-	-
White, both sexes	18	29	43	68	101	113	154	131	83	32	-	-
Male	9	14	11	18	34	27	29	25	21	4	-	-
Female	9	15	32	50	67	86	125	106	62	28	-	-
Black, both sexes	10	9	4	9	8	11	5	6	5	-	-	-
Male	4	5	-	5	1	2	1	2	2	-	-	-
Female	6	4	4	4	7	9	4	4	3	-	-	-
American Indian 1/, both sexes	-	-	-	2	1	-	1	1	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	3	-	7	2	2	3	3	-	-	-
Male	-	1	1	-	3	1	1	3	1	-	-	-
Female	1	3	2	-	4	1	1	-	2	-	-	-
Alabama												
All races, both sexes	1	1	-	1	2	2	-	2	2	-	-	-
Male	-	1	-	-	1	1	-	1	1	-	-	-
Female	1	-	-	1	1	1	-	1	1	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Black, both sexes	1	1	-	-	1	2	-	1	2	-	-	-
Male	-	1	-	-	-	1	-	1	1	-	-	-
Female	1	-	-	-	1	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Alaska												
All races, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	1	2	1	3	2	2	-	-	-	-
Male	-	-	-	2	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	2	2	2	-	-	-	-
White, both sexes	-	-	1	1	1	3	2	2	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	2	2	2	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Arkansas												
All races, both sexes	-	-	2	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	2	-	-	-	-
White, both sexes	-	-	2	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2	7	7	10	12	14	18	15	6	6	-	-
Male	2	3	2	3	5	5	3	6	2	1	-	-
Female	-	4	5	7	7	9	15	9	4	5	-	-
White, both sexes	2	5	5	10	8	13	16	14	3	6	-	-
Male	2	1	1	3	4	4	3	5	1	1	-	-
Female	-	4	4	7	4	9	13	9	2	5	-	-
Black, both sexes	-	1	-	-	1	-	2	-	1	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	3	1	-	1	2	-	-	-
Male	-	1	1	-	-	1	-	1	1	-	-	-
Female	-	-	1	-	3	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Colorado												
All races, both sexes	2	-	1	-	2	3	5	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	1	-	2	3	5	2	-	-	-	-
White, both sexes	1	-	1	-	1	3	5	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	1	-	1	3	5	2	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	1	3	-	3	3	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	3	-	2	2	-	-	-	-
White, both sexes	-	-	-	1	3	-	3	3	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	3	-	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Delaware												
All races, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Florida												
All races, both sexes	3	3	4	5	11	7	14	7	12	2	-	-
Male	1	3	2	2	5	2	5	-	4	-	-	-
Female	2	-	2	3	6	5	9	7	8	2	-	-
White, both sexes	2	2	4	5	10	6	14	7	11	2	-	-
Male	1	2	2	2	4	2	5	-	3	-	-	-
Female	1	-	2	3	6	4	9	7	8	2	-	-
Black, both sexes	1	1	-	-	-	1	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1	4	1	-	1	1	-	3	-	-	-	-
Male	1	2	-	-	-	1	-	-	-	-	-	-
Female	-	2	1	-	1	-	-	3	-	-	-	-
White, both sexes	-	4	1	-	1	1	-	1	-	-	-	-
Male	-	2	-	-	-	1	-	-	-	-	-	-
Female	-	2	1	-	1	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	2	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Hawaii												
All races, both sexes	-	1	1	-	-	2	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	1	1	-	-	2	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	1	-	-	1	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	2	1	-	2	1	3	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	2	-	3	1	-	-	-
White, both sexes	-	-	2	1	-	2	1	3	1	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	2	-	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Illinois												
All races, both sexes	-	-	1	1	7	4	6	4	2	1	-	-
Male	-	-	-	1	-	-	-	2	-	-	-	-
Female	-	-	1	-	7	4	6	2	2	1	-	-
White, both sexes	-	-	1	-	6	2	5	4	2	1	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	1	-	6	2	5	2	2	1	-	-
Black, both sexes	-	-	-	1	1	2	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Indiana												
All races, both sexes	1	1	2	3	3	4	2	1	4	2	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	1	1	2	3	3	4	2	1	3	2	-	-
White, both sexes	1	-	2	3	2	4	2	1	4	2	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	2	3	2	4	2	1	3	2	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Iowa												
All races, both sexes	1	-	-	-	3	3	3	5	3	3	-	-
Male	1	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	-	2	2	2	4	3	3	-	-
White, both sexes	1	-	-	-	3	3	3	5	3	3	-	-
Male	1	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	-	2	2	2	4	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	-	1	-	2	2	3	-	1	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	2	1	3	-	1	-	-
White, both sexes	-	1	-	1	-	2	2	3	-	1	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	2	1	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Kentucky												
All races, both sexes	1	3	-	2	4	3	3	2	1	1	-	-
Male	1	2	-	-	1	-	1	-	-	-	-	-
Female	-	1	-	2	3	3	2	2	1	1	-	-
White, both sexes	1	3	-	2	4	3	3	2	1	1	-	-
Male	1	2	-	-	1	-	1	-	-	-	-	-
Female	-	1	-	2	3	3	2	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	-	1	1	1	1	-	1	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	1	-	-	-
White, both sexes	-	-	1	-	1	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Maine												
All races, both sexes	-	-	-	-	1	2	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	2	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	1	-	-	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	1	2	1	-	-	-
White, both sexes	-	1	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	2	1	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Massachusetts												
All races, both sexes	-	-	1	1	3	4	4	-	4	2	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	3	3	4	-	4	2	-	-
White, both sexes	-	-	1	1	3	4	4	-	4	2	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	1	3	3	4	-	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	1	1	-	3	3	3	4	2	1	-	-
Male	-	1	1	-	-	-	1	2	-	-	-	-
Female	1	-	-	-	3	3	2	2	2	1	-	-
White, both sexes	-	1	1	-	3	2	3	4	2	1	-	-
Male	-	1	1	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	3	2	2	2	2	1	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Minnesota												
All races, both sexes	1	-	-	-	2	4	8	5	3	-	-	-
Male	1	-	-	-	1	1	1	2	2	-	-	-
Female	-	-	-	-	1	3	7	3	1	-	-	-
White, both sexes	-	-	-	-	2	4	8	5	3	-	-	-
Male	-	-	-	-	1	1	1	2	2	-	-	-
Female	-	-	-	-	1	3	7	3	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Missouri												
All races, both sexes	-	1	-	1	3	2	-	3	4	-	-	-
Male	-	1	-	1	2	-	-	-	1	-	-	-
Female	-	-	-	-	1	2	-	3	3	-	-	-
White, both sexes	-	-	-	1	3	1	-	2	4	-	-	-
Male	-	-	-	1	2	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	2	3	-	-	-
Black, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	1	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Nebraska												
All races, both sexes	-	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	1	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	1	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
New Hampshire												
All races, both sexes	-	-	-	-	-	3	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	3	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	3	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	3	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	-	1	7	4	5	11	4	4	-	-	-
Male	1	-	-	3	2	-	2	1	1	-	-	-
Female	1	-	1	4	2	5	9	3	3	-	-	-
White, both sexes	1	-	1	6	2	5	11	3	4	-	-	-
Male	1	-	-	2	1	-	2	-	1	-	-	-
Female	-	-	1	4	1	5	9	3	3	-	-	-
Black, both sexes	-	-	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
New Mexico												
All races, both sexes	2	-	2	2	3	-	2	1	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	2	-	1	2	2	-	1	1	-	-	-	-
White, both sexes	2	-	2	1	3	-	2	1	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	2	-	1	1	2	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	-	4	3	5	9	7	12	6	9	1	-	-
Male	-	3	-	1	5	1	5	-	1	-	-	-
Female	-	1	3	4	4	6	7	6	8	1	-	-
White, both sexes	-	2	3	5	9	6	12	6	8	1	-	-
Male	-	2	-	1	5	1	5	-	1	-	-	-
Female	-	-	3	4	4	5	7	6	7	1	-	-
Black, both sexes	-	2	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
North Carolina												
All races, both sexes	1	-	-	3	4	4	4	3	-	-	-	-
Male	-	-	-	1	3	2	1	1	-	-	-	-
Female	1	-	-	2	1	2	3	2	-	-	-	-
White, both sexes	-	-	-	3	4	3	4	3	-	-	-	-
Male	-	-	-	1	3	1	1	1	-	-	-	-
Female	-	-	-	2	1	2	3	2	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Ohio												
All races, both sexes	1	3	4	4	6	8	9	4	6	3	-	-
Male	1	2	1	1	2	1	-	2	2	1	-	-
Female	-	1	3	3	4	7	9	2	4	2	-	-
White, both sexes	-	3	4	3	6	7	8	4	5	3	-	-
Male	-	2	1	1	2	1	-	2	2	1	-	-
Female	-	1	3	2	4	6	8	2	3	2	-	-
Black, both sexes	1	-	-	1	-	1	1	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	2	1	-	1	3	1	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	3	1	-	-	-
White, both sexes	1	-	-	2	1	-	1	2	1	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Oregon												
All races, both sexes	-	1	-	1	2	5	1	6	3	-	-	-
Male	-	-	-	1	-	1	-	2	-	-	-	-
Female	-	1	-	-	2	4	1	4	3	-	-	-
White, both sexes	-	-	-	1	2	5	1	5	3	-	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	2	4	1	4	3	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	-	1	3	7	6	13	9	6	-	-	-
Male	-	-	-	-	2	4	2	2	2	-	-	-
Female	-	-	1	3	5	2	11	7	4	-	-	-
White, both sexes	-	-	1	3	5	6	13	9	6	-	-	-
Male	-	-	-	-	2	4	2	2	2	-	-	-
Female	-	-	1	3	3	2	11	7	4	-	-	-
Black, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Rhode Island												
All races, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	2	1	3	1	2	5	2	1	-	-	-
Male	1	1	-	1	-	2	1	-	1	-	-	-
Female	1	1	1	2	1	-	4	2	-	-	-	-
White, both sexes	1	1	-	1	1	2	3	2	1	-	-	-
Male	-	1	-	-	-	2	-	-	1	-	-	-
Female	1	-	-	1	1	-	3	2	-	-	-	-
Black, both sexes	1	1	1	2	-	-	2	-	-	-	-	-
Male	1	-	-	1	-	-	1	-	-	-	-	-
Female	-	1	1	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
South Dakota												
All races, both sexes	-	-	-	2	-	-	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	1	1	2	3	-	6	3	2	1	-	-
Male	-	-	1	-	1	-	-	2	1	-	-	-
Female	-	1	-	2	2	-	6	1	1	1	-	-
White, both sexes	-	1	1	2	3	-	6	3	2	1	-	-
Male	-	-	1	-	1	-	-	2	1	-	-	-
Female	-	1	-	2	2	-	6	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Texas												
All races, both sexes	3	2	4	3	6	7	6	9	1	1	-	-
Male	1	-	1	-	2	2	1	-	-	-	-	-
Female	2	2	3	3	4	5	5	9	1	1	-	-
White, both sexes	3	2	2	2	5	7	5	9	1	1	-	-
Male	1	-	1	-	2	2	-	-	-	-	-	-
Female	2	2	1	2	3	5	5	9	1	1	-	-
Black, both sexes	-	-	2	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	3	-	2	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	1	3	-	2	-	1	-	-	-	-
White, both sexes	-	-	1	3	-	2	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	1	3	-	2	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Vermont												
All races, both sexes	-	-	-	-	1	2	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	2	-	-	-	-
White, both sexes	-	-	-	-	1	2	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	-	2	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	2	-	1	1	1	1	-	-	-	-
White, both sexes	1	-	2	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	2	-	1	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Washington												
All races, both sexes	-	3	1	2	-	3	7	5	1	4	-	-
Male	-	-	-	1	-	1	1	1	-	1	-	-
Female	-	3	1	1	-	2	6	4	1	3	-	-
White, both sexes	-	2	-	2	-	3	6	5	1	4	-	-
Male	-	-	-	1	-	1	1	1	-	1	-	-
Female	-	2	-	1	-	2	5	4	1	3	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)												
Wisconsin												
All races, both sexes	1	-	3	1	4	2	3	3	4	3	-	-
Male	1	-	1	-	3	1	1	-	2	1	-	-
Female	-	-	2	1	1	1	2	3	2	2	-	-
White, both sexes	1	-	3	1	4	2	3	3	4	3	-	-
Male	1	-	1	-	3	1	1	-	2	1	-	-
Female	-	-	2	1	1	1	2	3	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
United States												
All races, both sexes	57,356	4	-	4	2	7	56	148	297	611	1,246	2,036
Male	24,046	2	-	2	1	3	37	93	194	429	824	1,371
Female	33,310	2	-	2	1	4	19	55	103	182	422	665
White, both sexes	43,126	1	-	4	1	3	21	70	137	325	671	1,127
Male	17,312	1	-	2	1	2	15	53	96	246	480	820
Female	25,814	-	-	2	-	1	6	17	41	79	191	307
Black, both sexes	12,765	3	-	-	1	3	34	68	149	268	528	860
Male	6,019	1	-	-	-	1	21	32	90	167	315	513
Female	6,746	2	-	-	1	2	13	36	59	101	213	347
American Indian 1/, both sexes	252	-	-	-	-	-	-	6	4	3	12	12
Male	131	-	-	-	-	-	-	5	1	2	6	10
Female	121	-	-	-	-	-	-	1	3	1	6	2
Asian or Pacific Islander 2/, both sexes	1,213	-	-	-	-	1	1	4	7	15	35	37
Male	584	-	-	-	-	-	1	3	7	14	23	28
Female	629	-	-	-	-	1	-	1	-	1	12	9
Alabama												
All races, both sexes	931	-	-	-	-	1	1	2	2	9	21	33
Male	380	-	-	-	-	-	-	1	1	6	13	24
Female	551	-	-	-	-	1	1	1	1	3	8	9
White, both sexes	530	-	-	-	-	-	-	-	-	2	9	9
Male	197	-	-	-	-	-	-	-	-	2	4	8
Female	333	-	-	-	-	-	-	-	-	-	5	1
Black, both sexes	400	-	-	-	-	1	1	2	2	7	12	24
Male	183	-	-	-	-	-	-	1	1	4	9	16
Female	217	-	-	-	-	1	1	1	1	3	3	8
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Alaska												
All races, both sexes	67	-	-	-	-	-	-	1	-	1	2	4
Male	40	-	-	-	-	-	-	-	-	1	1	4
Female	27	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	46	-	-	-	-	-	-	1	-	1	-	3
Male	29	-	-	-	-	-	-	-	-	1	-	3
Female	17	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Arizona												
All races, both sexes	920	-	-	-	-	-	1	3	6	7	21	27
Male	411	-	-	-	-	-	1	2	4	4	11	20
Female	509	-	-	-	-	-	-	1	2	3	10	7
White, both sexes	837	-	-	-	-	-	1	1	5	6	13	23
Male	373	-	-	-	-	-	1	1	4	3	6	18
Female	464	-	-	-	-	-	-	-	1	3	7	5
Black, both sexes	38	-	-	-	-	-	-	1	-	-	4	3
Male	22	-	-	-	-	-	-	-	-	-	3	1
Female	16	-	-	-	-	-	-	1	-	-	1	2
American Indian 1/, both sexes	36	-	-	-	-	-	-	1	1	-	3	-
Male	11	-	-	-	-	-	-	1	-	-	1	-
Female	25	-	-	-	-	-	-	-	1	-	2	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	1	1	1
Male	5	-	-	-	-	-	-	-	-	1	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Arkansas												
All races, both sexes	543	-	-	-	-	-	-	3	4	3	18	20
Male	216	-	-	-	-	-	-	2	4	3	11	12
Female	327	-	-	-	-	-	-	1	-	-	7	8
White, both sexes	413	-	-	-	-	-	-	-	3	1	11	8
Male	159	-	-	-	-	-	-	-	3	1	8	7
Female	254	-	-	-	-	-	-	-	-	-	3	1
Black, both sexes	127	-	-	-	-	-	-	3	1	2	7	11
Male	56	-	-	-	-	-	-	2	1	2	3	4
Female	71	-	-	-	-	-	-	1	-	-	4	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7,414	-	-	-	-	1	4	17	28	79	157	254
Male	3,167	-	-	-	-	-	4	10	16	61	108	180
Female	4,247	-	-	-	-	1	-	7	12	18	49	74
White, both sexes	5,760	-	-	-	-	-	2	11	16	45	105	167
Male	2,384	-	-	-	-	-	2	6	9	37	77	123
Female	3,376	-	-	-	-	-	-	5	7	8	28	44
Black, both sexes	987	-	-	-	-	-	2	1	10	29	33	69
Male	474	-	-	-	-	-	2	-	5	19	21	45
Female	513	-	-	-	-	-	-	1	5	10	12	24
American Indian 1/, both sexes	31	-	-	-	-	-	-	2	-	-	2	2
Male	15	-	-	-	-	-	-	2	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	2	1
Asian or Pacific Islander 2/, both sexes	636	-	-	-	-	1	-	3	2	5	17	16
Male	294	-	-	-	-	-	-	2	2	5	10	11
Female	342	-	-	-	-	1	-	1	-	-	7	5

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Colorado												
All races, both sexes	509	-	-	-	-	-	-	2	2	4	10	11
Male	200	-	-	-	-	-	-	2	1	1	8	6
Female	309	-	-	-	-	-	-	-	1	3	2	5
White, both sexes	467	-	-	-	-	-	-	-	1	3	9	10
Male	177	-	-	-	-	-	-	-	-	1	7	6
Female	290	-	-	-	-	-	-	-	1	2	2	4
Black, both sexes	36	-	-	-	-	-	-	2	1	-	1	1
Male	21	-	-	-	-	-	-	2	1	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Connecticut												
All races, both sexes	549	1	-	-	-	-	1	1	2	4	8	15
Male	209	-	-	-	-	-	-	1	1	3	7	10
Female	340	1	-	-	-	-	1	-	1	1	1	5
White, both sexes	492	-	-	-	-	-	-	1	1	2	8	12
Male	184	-	-	-	-	-	-	1	1	1	7	8
Female	308	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	49	1	-	-	-	-	1	-	1	1	-	2
Male	21	-	-	-	-	-	-	-	-	1	-	1
Female	28	1	-	-	-	-	1	-	1	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	1	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Delaware												
All races, both sexes	167	-	-	-	-	-	1	1	1	2	4	5
Male	67	-	-	-	-	-	1	-	-	1	2	5
Female	100	-	-	-	-	-	-	1	1	1	2	-
White, both sexes	126	-	-	-	-	-	1	-	-	1	1	3
Male	49	-	-	-	-	-	1	-	-	1	-	3
Female	77	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	41	-	-	-	-	-	-	1	1	1	3	2
Male	18	-	-	-	-	-	-	-	-	-	2	2
Female	23	-	-	-	-	-	-	1	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	281	-	-	-	-	-	1	2	1	1	10	9
Male	138	-	-	-	-	-	-	2	1	-	6	6
Female	143	-	-	-	-	-	1	-	-	1	4	3
White, both sexes	32	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	244	-	-	-	-	-	1	2	1	1	9	8
Male	119	-	-	-	-	-	-	2	1	-	5	6
Female	125	-	-	-	-	-	1	-	-	1	4	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Florida												
All races, both sexes	4,361	1	-	-	-	-	5	12	17	55	101	147
Male	1,969	-	-	-	-	-	5	7	10	38	73	97
Female	2,392	1	-	-	-	-	-	5	7	17	28	50
White, both sexes	3,461	-	-	-	-	-	3	6	6	39	61	92
Male	1,536	-	-	-	-	-	3	5	5	28	48	67
Female	1,925	-	-	-	-	-	-	1	1	11	13	25
Black, both sexes	871	1	-	-	-	-	2	6	11	15	39	54
Male	419	-	-	-	-	-	2	2	5	9	25	29
Female	452	1	-	-	-	-	-	4	6	6	14	25
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	1	-	1
Male	12	-	-	-	-	-	-	-	-	1	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,075	-	-	-	1	1	3	2	15	36	53	109
Male	947	-	-	-	-	1	2	-	9	24	32	71
Female	1,128	-	-	-	1	-	1	2	6	12	21	38
White, both sexes	1,097	-	-	-	-	-	-	1	4	16	15	45
Male	482	-	-	-	-	-	-	-	2	8	10	34
Female	615	-	-	-	-	-	-	1	2	8	5	11
Black, both sexes	968	-	-	-	1	1	3	1	11	20	38	64
Male	459	-	-	-	-	1	2	-	7	16	22	37
Female	509	-	-	-	1	-	1	1	4	4	16	27
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Hawaii												
All races, both sexes	155	-	-	-	-	-	-	-	1	2	4	2
Male	66	-	-	-	-	-	-	-	1	2	2	1
Female	89	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	130	-	-	-	-	-	-	-	1	2	4	1
Male	58	-	-	-	-	-	-	-	1	2	2	1
Female	72	-	-	-	-	-	-	-	-	-	2	-
Idaho												
All races, both sexes	178	-	-	-	-	-	-	-	-	2	2	3
Male	71	-	-	-	-	-	-	-	-	1	1	3
Female	107	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	177	-	-	-	-	-	-	-	-	2	2	3
Male	71	-	-	-	-	-	-	-	-	1	1	3
Female	106	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Illinois												
All races, both sexes	2,589	-	-	-	-	-	4	11	18	34	74	106
Male	1,145	-	-	-	-	-	3	6	14	23	51	76
Female	1,444	-	-	-	-	-	1	5	4	11	23	30
White, both sexes	1,814	-	-	-	-	-	-	2	9	13	27	49
Male	737	-	-	-	-	-	-	2	8	12	20	38
Female	1,077	-	-	-	-	-	-	-	1	1	7	11
Black, both sexes	741	-	-	-	-	-	3	9	8	21	46	55
Male	390	-	-	-	-	-	2	4	5	11	30	37
Female	351	-	-	-	-	-	1	5	3	10	16	18
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	1	-	1	-	1	2
Male	18	-	-	-	-	-	1	-	1	-	1	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
Indiana												
All races, both sexes	1,088	-	-	-	-	-	1	6	4	8	21	25
Male	395	-	-	-	-	-	-	4	2	6	16	17
Female	693	-	-	-	-	-	1	2	2	2	5	8
White, both sexes	935	-	-	-	-	-	-	3	3	3	13	15
Male	324	-	-	-	-	-	-	3	2	3	11	13
Female	611	-	-	-	-	-	-	-	1	-	2	2
Black, both sexes	149	-	-	-	-	-	1	3	1	5	8	10
Male	70	-	-	-	-	-	-	1	-	3	5	4
Female	79	-	-	-	-	-	1	2	1	2	3	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Iowa												
All races, both sexes	600	-	-	-	-	-	-	1	-	3	8	8
Male	223	-	-	-	-	-	-	-	-	3	5	6
Female	377	-	-	-	-	-	-	1	-	-	3	2
White, both sexes	586	-	-	-	-	-	-	1	-	3	4	8
Male	216	-	-	-	-	-	-	-	-	3	3	6
Female	370	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	14	-	-	-	-	-	-	-	-	-	4	-
Male	7	-	-	-	-	-	-	-	-	-	2	-
Female	7	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	345	-	-	-	-	-	1	1	1	1	1	9
Male	125	-	-	-	-	-	1	1	1	1	1	8
Female	220	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	307	-	-	-	-	-	-	1	-	-	-	5
Male	100	-	-	-	-	-	-	1	-	-	-	5
Female	207	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	36	-	-	-	-	-	1	-	1	1	1	4
Male	23	-	-	-	-	-	1	-	1	1	1	3
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Kentucky												
All races, both sexes	767	-	-	-	-	-	2	5	4	16	24	36
Male	308	-	-	-	-	-	2	2	1	13	16	26
Female	459	-	-	-	-	-	-	3	3	3	8	10
White, both sexes	660	-	-	-	-	-	1	4	2	14	20	27
Male	273	-	-	-	-	-	1	1	1	12	15	21
Female	387	-	-	-	-	-	-	3	1	2	5	6
Black, both sexes	104	-	-	-	-	-	1	1	2	2	4	9
Male	33	-	-	-	-	-	1	1	-	1	1	5
Female	71	-	-	-	-	-	-	-	2	1	3	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,166	-	-	-	-	-	1	2	9	17	31	64
Male	506	-	-	-	-	-	-	-	7	12	18	36
Female	660	-	-	-	-	-	1	2	2	5	13	28
White, both sexes	644	-	-	-	-	-	1	-	2	7	8	21
Male	265	-	-	-	-	-	-	-	1	6	5	15
Female	379	-	-	-	-	-	1	-	1	1	3	6
Black, both sexes	519	-	-	-	-	-	-	2	7	10	23	42
Male	239	-	-	-	-	-	-	-	6	6	13	20
Female	280	-	-	-	-	-	-	2	1	4	10	22
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Maine												
All races, both sexes	139	-	-	-	-	-	-	-	1	-	3	-
Male	60	-	-	-	-	-	-	-	1	-	2	-
Female	79	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	137	-	-	-	-	-	-	-	1	-	3	-
Male	60	-	-	-	-	-	-	-	1	-	2	-
Female	77	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,301	-	-	1	-	-	-	2	12	31	50	78
Male	609	-	-	-	-	-	-	2	7	26	35	58
Female	692	-	-	1	-	-	-	-	5	5	15	20
White, both sexes	756	-	-	1	-	-	-	-	4	14	23	40
Male	331	-	-	-	-	-	-	-	2	14	14	29
Female	425	-	-	1	-	-	-	-	2	-	9	11
Black, both sexes	531	-	-	-	-	-	-	2	8	17	27	38
Male	271	-	-	-	-	-	-	2	5	12	21	29
Female	260	-	-	-	-	-	-	-	3	5	6	9
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Massachusetts												
All races, both sexes	808	-	-	-	-	-	2	5	7	13	24	39
Male	344	-	-	-	-	-	2	5	5	7	18	30
Female	464	-	-	-	-	-	-	-	2	6	6	9
White, both sexes	719	-	-	-	-	-	2	4	6	9	16	29
Male	301	-	-	-	-	-	2	4	4	4	13	21
Female	418	-	-	-	-	-	-	-	2	5	3	8
Black, both sexes	77	-	-	-	-	-	-	1	1	4	7	9
Male	35	-	-	-	-	-	-	1	1	3	4	8
Female	42	-	-	-	-	-	-	-	-	1	3	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2,214	-	-	-	-	1	2	3	8	17	38	84
Male	879	-	-	-	-	-	1	1	7	13	25	57
Female	1,335	-	-	-	-	1	1	2	1	4	13	27
White, both sexes	1,668	-	-	-	-	1	1	-	2	8	18	37
Male	626	-	-	-	-	-	-	-	2	6	13	31
Female	1,042	-	-	-	-	1	1	-	-	2	5	6
Black, both sexes	537	-	-	-	-	-	1	3	6	9	20	45
Male	251	-	-	-	-	-	1	1	5	7	12	24
Female	286	-	-	-	-	-	-	2	1	2	8	21
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Minnesota												
All races, both sexes	749	-	-	1	-	-	-	-	2	1	5	12
Male	269	-	-	1	-	-	-	-	1	-	4	6
Female	480	-	-	-	-	-	-	-	1	1	1	6
White, both sexes	712	-	-	1	-	-	-	-	-	-	2	10
Male	253	-	-	1	-	-	-	-	-	-	2	6
Female	459	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	29	-	-	-	-	-	-	-	1	1	2	2
Male	12	-	-	-	-	-	-	-	1	-	1	-
Female	17	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,136	-	-	-	-	1	2	6	9	16	18	49
Male	445	-	-	-	-	-	1	3	4	6	9	30
Female	691	-	-	-	-	1	1	3	5	10	9	19
White, both sexes	570	-	-	-	-	-	-	1	2	5	7	13
Male	217	-	-	-	-	-	-	1	2	4	6	9
Female	353	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	561	-	-	-	-	1	2	4	7	11	11	36
Male	224	-	-	-	-	-	1	1	2	2	3	21
Female	337	-	-	-	-	1	1	3	5	9	8	15
American Indian 1/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Missouri												
All races, both sexes	974	-	-	-	-	-	2	7	12	13	29	52
Male	407	-	-	-	-	-	1	4	6	9	19	33
Female	567	-	-	-	-	-	1	3	6	4	10	19
White, both sexes	722	-	-	-	-	-	1	4	5	5	16	31
Male	290	-	-	-	-	-	1	4	2	4	13	20
Female	432	-	-	-	-	-	-	-	3	1	3	11
Black, both sexes	249	-	-	-	-	-	1	3	7	8	13	21
Male	114	-	-	-	-	-	-	-	4	5	6	13
Female	135	-	-	-	-	-	1	3	3	3	7	8
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	120	-	-	-	-	-	-	2	-	-	1	4
Male	35	-	-	-	-	-	-	2	-	-	-	3
Female	85	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	115	-	-	-	-	-	-	1	-	-	1	4
Male	33	-	-	-	-	-	-	1	-	-	-	3
Female	82	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Nebraska												
All races, both sexes	302	-	-	-	-	-	-	1	-	4	6	3
Male	114	-	-	-	-	-	-	1	-	4	3	3
Female	188	-	-	-	-	-	-	-	-	-	3	-
White, both sexes	274	-	-	-	-	-	-	1	-	3	4	3
Male	98	-	-	-	-	-	-	1	-	3	2	3
Female	176	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	639	-	-	-	-	-	1	2	6	9	15	23
Male	370	-	-	-	-	-	1	2	4	6	11	16
Female	269	-	-	-	-	-	-	-	2	3	4	7
White, both sexes	545	-	-	-	-	-	-	2	2	7	12	12
Male	313	-	-	-	-	-	-	2	2	5	10	10
Female	232	-	-	-	-	-	-	-	-	2	2	2
Black, both sexes	67	-	-	-	-	-	1	-	3	2	3	10
Male	38	-	-	-	-	-	1	-	2	1	1	5
Female	29	-	-	-	-	-	-	-	1	1	2	5
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
New Hampshire												
All races, both sexes	167	-	-	-	-	-	-	-	1	3	-	5
Male	71	-	-	-	-	-	-	-	1	2	-	5
Female	96	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	167	-	-	-	-	-	-	-	1	3	-	5
Male	71	-	-	-	-	-	-	-	1	2	-	5
Female	96	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,454	-	-	-	-	-	1	1	5	12	25	59
Male	607	-	-	-	-	-	-	-	5	8	14	36
Female	847	-	-	-	-	-	1	1	-	4	11	23
White, both sexes	1,096	-	-	-	-	-	-	1	2	5	9	23
Male	428	-	-	-	-	-	-	-	2	4	6	15
Female	668	-	-	-	-	-	-	1	-	1	3	8
Black, both sexes	328	-	-	-	-	-	1	-	3	7	16	33
Male	164	-	-	-	-	-	-	-	3	4	8	19
Female	164	-	-	-	-	-	1	-	-	3	8	14
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	-	-	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
New Mexico												
All races, both sexes	293	-	-	-	-	-	-	-	1	1	13	9
Male	145	-	-	-	-	-	-	-	1	1	9	8
Female	148	-	-	-	-	-	-	-	-	-	4	1
White, both sexes	262	-	-	-	-	-	-	-	1	1	10	9
Male	125	-	-	-	-	-	-	-	1	1	7	8
Female	137	-	-	-	-	-	-	-	-	-	3	1
Black, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	2	-
Male	10	-	-	-	-	-	-	-	-	-	2	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4,275	1	-	-	-	-	4	13	17	30	93	131
Male	1,764	1	-	-	-	-	4	8	15	21	58	88
Female	2,511	-	-	-	-	-	-	5	2	9	35	43
White, both sexes	2,914	-	-	-	-	-	2	4	7	14	36	53
Male	1,146	-	-	-	-	-	2	3	7	9	23	40
Female	1,768	-	-	-	-	-	-	1	-	5	13	13
Black, both sexes	1,239	1	-	-	-	-	2	9	8	15	52	72
Male	555	1	-	-	-	-	2	5	6	11	31	43
Female	684	-	-	-	-	-	-	4	2	4	21	29
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	119	-	-	-	-	-	-	-	2	1	5	6
Male	61	-	-	-	-	-	-	-	2	1	4	5
Female	58	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
North Carolina												
All races, both sexes	1,544	-	-	-	-	-	3	4	14	14	41	66
Male	670	-	-	-	-	-	1	1	9	9	27	40
Female	874	-	-	-	-	-	2	3	5	5	14	26
White, both sexes	924	-	-	-	-	-	1	2	7	5	23	26
Male	379	-	-	-	-	-	-	-	5	3	16	18
Female	545	-	-	-	-	-	1	2	2	2	7	8
Black, both sexes	602	-	-	-	-	-	2	2	7	9	18	39
Male	281	-	-	-	-	-	1	1	4	6	11	21
Female	321	-	-	-	-	-	1	1	3	3	7	18
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	126	-	-	-	-	-	-	-	-	-	-	2
Male	58	-	-	-	-	-	-	-	-	-	-	2
Female	68	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	124	-	-	-	-	-	-	-	-	-	-	2
Male	57	-	-	-	-	-	-	-	-	-	-	2
Female	67	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Ohio												
All races, both sexes	2,860	-	-	-	-	-	3	3	13	31	60	101
Male	1,212	-	-	-	-	-	1	3	8	22	46	66
Female	1,648	-	-	-	-	-	2	-	5	9	14	35
White, both sexes	2,297	-	-	-	-	-	1	2	5	18	38	74
Male	922	-	-	-	-	-	-	2	4	14	32	51
Female	1,375	-	-	-	-	-	1	-	1	4	6	23
Black, both sexes	550	-	-	-	-	-	2	1	8	13	21	27
Male	285	-	-	-	-	-	1	1	4	8	13	15
Female	265	-	-	-	-	-	1	-	4	5	8	12
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	647	-	-	-	-	-	-	1	4	2	6	21
Male	250	-	-	-	-	-	-	1	2	-	3	12
Female	397	-	-	-	-	-	-	-	2	2	3	9
White, both sexes	511	-	-	-	-	-	-	1	2	1	3	12
Male	192	-	-	-	-	-	-	1	1	-	2	6
Female	319	-	-	-	-	-	-	-	1	1	1	6
Black, both sexes	90	-	-	-	-	-	-	-	1	-	2	7
Male	31	-	-	-	-	-	-	-	-	-	1	5
Female	59	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	44	-	-	-	-	-	-	-	1	1	1	2
Male	27	-	-	-	-	-	-	-	1	-	-	1
Female	17	-	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Oregon												
All races, both sexes	700	-	-	-	-	-	1	-	-	1	6	9
Male	259	-	-	-	-	-	1	-	-	1	5	9
Female	441	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	663	-	-	-	-	-	1	-	-	-	5	8
Male	241	-	-	-	-	-	1	-	-	-	4	8
Female	422	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	22	-	-	-	-	-	-	-	-	1	1	-
Male	13	-	-	-	-	-	-	-	-	1	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	2,042	1	-	-	-	-	1	2	7	19	23	81
Male	844	1	-	-	-	-	1	1	4	12	14	53
Female	1,198	-	-	-	-	-	-	1	3	7	9	28
White, both sexes	1,704	1	-	-	-	-	-	1	4	9	12	55
Male	670	1	-	-	-	-	-	1	3	7	7	40
Female	1,034	-	-	-	-	-	-	-	1	2	5	15
Black, both sexes	331	-	-	-	-	-	1	1	3	10	11	26
Male	170	-	-	-	-	-	1	-	1	5	7	13
Female	161	-	-	-	-	-	-	1	2	5	4	13
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Rhode Island												
All races, both sexes	125	-	-	-	-	-	-	-	-	-	2	4
Male	48	-	-	-	-	-	-	-	-	-	1	4
Female	77	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	118	-	-	-	-	-	-	-	-	-	2	4
Male	43	-	-	-	-	-	-	-	-	-	1	4
Female	75	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	835	-	-	-	-	-	-	2	1	8	15	29
Male	350	-	-	-	-	-	-	2	1	7	9	20
Female	485	-	-	-	-	-	-	-	-	1	6	9
White, both sexes	443	-	-	-	-	-	-	-	-	3	8	8
Male	170	-	-	-	-	-	-	-	-	3	4	6
Female	273	-	-	-	-	-	-	-	-	-	4	2
Black, both sexes	389	-	-	-	-	-	-	2	1	5	7	21
Male	178	-	-	-	-	-	-	2	1	4	5	14
Female	211	-	-	-	-	-	-	-	-	1	2	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
South Dakota												
All races, both sexes	144	-	-	-	-	-	-	-	-	-	-	1
Male	56	-	-	-	-	-	-	-	-	-	-	-
Female	88	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	137	-	-	-	-	-	-	-	-	-	-	1
Male	50	-	-	-	-	-	-	-	-	-	-	-
Female	87	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,531	-	-	-	-	-	3	8	16	20	32	53
Male	603	-	-	-	-	-	1	8	10	13	16	31
Female	928	-	-	-	-	-	2	-	6	7	16	22
White, both sexes	1,061	-	-	-	-	-	1	6	8	9	17	29
Male	393	-	-	-	-	-	-	6	5	7	6	14
Female	668	-	-	-	-	-	1	-	3	2	11	15
Black, both sexes	463	-	-	-	-	-	2	2	8	11	15	24
Male	206	-	-	-	-	-	1	2	5	6	10	17
Female	257	-	-	-	-	-	1	-	3	5	5	7
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Texas												
All races, both sexes	3,620	-	-	2	1	-	2	8	29	43	108	140
Male	1,519	-	-	1	1	-	1	5	20	28	64	94
Female	2,101	-	-	1	-	-	1	3	9	15	44	46
White, both sexes	2,799	-	-	2	1	-	1	5	17	25	58	82
Male	1,145	-	-	1	1	-	1	5	10	18	39	60
Female	1,654	-	-	1	-	-	-	-	7	7	19	22
Black, both sexes	782	-	-	-	-	-	1	2	11	16	48	55
Male	354	-	-	-	-	-	-	-	9	8	23	31
Female	428	-	-	-	-	-	1	2	2	8	25	24
American Indian 1/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	-	1	2	2	3
Male	20	-	-	-	-	-	-	-	1	2	2	3
Female	16	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	221	-	-	-	-	-	-	-	-	2	1	2
Male	76	-	-	-	-	-	-	-	-	2	-	1
Female	145	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	217	-	-	-	-	-	-	-	-	2	1	2
Male	74	-	-	-	-	-	-	-	-	2	-	1
Female	143	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Vermont												
All races, both sexes	99	-	-	-	-	-	-	1	-	-	4	5
Male	41	-	-	-	-	-	-	1	-	-	3	2
Female	58	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	99	-	-	-	-	-	-	1	-	-	4	5
Male	41	-	-	-	-	-	-	1	-	-	3	2
Female	58	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,056	-	-	-	-	-	2	5	9	20	23	42
Male	432	-	-	-	-	-	2	3	6	14	21	28
Female	624	-	-	-	-	-	-	2	3	6	2	14
White, both sexes	635	-	-	-	-	-	-	2	4	8	12	14
Male	235	-	-	-	-	-	-	1	3	5	12	10
Female	400	-	-	-	-	-	-	1	1	3	-	4
Black, both sexes	403	-	-	-	-	-	2	2	5	12	10	27
Male	189	-	-	-	-	-	2	1	3	9	8	17
Female	214	-	-	-	-	-	-	1	2	3	2	10
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	1	-	-	-	1
Male	7	-	-	-	-	-	-	1	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Washington												
All races, both sexes	1,089	-	-	-	-	1	-	-	4	10	16	18
Male	422	-	-	-	-	1	-	-	2	9	12	10
Female	667	-	-	-	-	-	-	-	2	1	4	8
White, both sexes	992	-	-	-	-	1	-	-	4	8	11	14
Male	370	-	-	-	-	1	-	-	2	7	9	7
Female	622	-	-	-	-	-	-	-	2	1	2	7
Black, both sexes	43	-	-	-	-	-	-	-	-	-	4	3
Male	21	-	-	-	-	-	-	-	-	-	3	3
Female	22	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	-	-	-	-	-	-	-	-	1	1	1
Male	24	-	-	-	-	-	-	-	-	1	-	-
Female	19	-	-	-	-	-	-	-	-	-	1	1
West Virginia												
All races, both sexes	446	-	-	-	-	1	-	-	-	4	3	7
Male	159	-	-	-	-	1	-	-	-	3	3	4
Female	287	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	424	-	-	-	-	1	-	-	-	4	3	6
Male	152	-	-	-	-	1	-	-	-	3	3	4
Female	272	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive diseases (I10-I13)												
Wisconsin												
All races, both sexes	949	-	-	-	-	-	1	-	4	3	16	16
Male	325	-	-	-	-	-	-	-	2	3	11	10
Female	624	-	-	-	-	-	1	-	2	-	5	6
White, both sexes	870	-	-	-	-	-	1	-	1	1	10	11
Male	290	-	-	-	-	-	-	-	1	1	8	6
Female	580	-	-	-	-	-	1	-	-	-	2	5
Black, both sexes	72	-	-	-	-	-	-	-	3	2	6	5
Male	33	-	-	-	-	-	-	-	1	2	3	4
Female	39	-	-	-	-	-	-	-	2	-	3	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	46	-	-	-	-	-	-	-	-	-	-	4
Male	20	-	-	-	-	-	-	-	-	-	-	4
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	45	-	-	-	-	-	-	-	-	-	-	3
Male	19	-	-	-	-	-	-	-	-	-	-	3
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
United States												
All races, both sexes	2,612	3,021	3,127	3,376	4,162	6,026	8,508	9,422	7,987	3,856	845	3
Male	1,765	1,994	1,948	1,910	2,085	2,699	3,166	2,963	1,817	628	112	3
Female	847	1,027	1,179	1,466	2,077	3,327	5,342	6,459	6,170	3,228	733	-
White, both sexes	1,472	1,868	2,046	2,182	2,878	4,536	6,849	7,998	6,925	3,320	689	3
Male	1,042	1,276	1,309	1,266	1,437	2,037	2,566	2,500	1,566	505	89	3
Female	430	592	737	916	1,441	2,499	4,283	5,498	5,359	2,815	600	-
Black, both sexes	1,070	1,082	1,008	1,098	1,144	1,306	1,442	1,219	895	448	139	-
Male	671	674	602	583	571	581	508	380	198	92	19	-
Female	399	408	406	515	573	725	934	839	697	356	120	-
American Indian 1/, both sexes	21	15	19	19	30	34	26	27	17	4	3	-
Male	15	10	12	11	16	16	11	8	7	1	-	-
Female	6	5	7	8	14	18	15	19	10	3	3	-
Asian or Pacific Islander 2/, both sexes	49	56	54	77	110	150	191	178	150	84	14	-
Male	37	34	25	50	61	65	81	75	46	30	4	-
Female	12	22	29	27	49	85	110	103	104	54	10	-
Alabama												
All races, both sexes	47	53	46	56	84	100	132	164	105	63	12	-
Male	29	29	26	32	45	46	39	58	17	11	3	-
Female	18	24	20	24	39	54	93	106	88	52	9	-
White, both sexes	15	21	23	32	39	65	78	111	77	41	8	-
Male	12	11	12	20	19	28	24	37	12	6	2	-
Female	3	10	11	12	20	37	54	74	65	35	6	-
Black, both sexes	32	32	23	24	45	34	54	53	28	22	4	-
Male	17	18	14	12	26	18	15	21	5	5	1	-
Female	15	14	9	12	19	16	39	32	23	17	3	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Alaska												
All races, both sexes	4	3	9	7	9	7	8	6	3	3	-	-
Male	4	3	6	5	5	4	4	1	1	1	-	-
Female	-	-	3	2	4	3	4	5	2	2	-	-
White, both sexes	3	3	7	5	4	4	8	4	1	2	-	-
Male	3	3	5	4	2	2	4	1	-	1	-	-
Female	-	-	2	1	2	2	4	3	1	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	2	2	5	2	-	2	2	1	-	-
Male	1	-	1	1	3	1	-	-	1	-	-	-
Female	-	-	1	1	2	1	-	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	41	56	52	59	57	107	138	163	123	48	9	2
Male	25	36	27	34	31	59	52	57	37	7	2	2
Female	16	20	25	25	26	48	86	106	86	41	7	-
White, both sexes	34	48	46	51	53	99	131	154	114	47	9	2
Male	21	31	23	29	30	56	50	54	36	6	2	2
Female	13	17	23	22	23	43	81	100	78	41	7	-
Black, both sexes	2	5	3	4	2	1	3	4	6	-	-	-
Male	2	4	2	4	1	-	2	2	1	-	-	-
Female	-	1	1	-	1	1	1	2	5	-	-	-
American Indian 1/, both sexes	4	3	3	3	2	6	4	4	2	-	-	-
Male	2	1	2	-	-	3	-	1	-	-	-	-
Female	2	2	1	3	2	3	4	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	1	-	1	1	1	-	-
Male	-	-	-	1	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	1	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Arkansas												
All races, both sexes	24	30	34	31	37	54	84	73	82	38	8	-
Male	13	22	24	17	17	21	31	13	16	8	2	-
Female	11	8	10	14	20	33	53	60	66	30	6	-
White, both sexes	11	17	19	25	29	43	73	67	69	33	4	-
Male	7	13	16	15	13	19	25	11	13	7	1	-
Female	4	4	3	10	16	24	48	56	56	26	3	-
Black, both sexes	13	13	15	6	8	11	11	6	12	4	4	-
Male	6	9	8	2	4	2	6	2	3	1	1	-
Female	7	4	7	4	4	9	5	4	9	3	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
California												
All races, both sexes	304	376	399	424	515	774	1,107	1,245	1,089	518	123	-
Male	208	233	244	261	253	353	413	446	267	90	20	-
Female	96	143	155	163	262	421	694	799	822	428	103	-
White, both sexes	211	254	286	300	364	601	890	1,033	933	436	106	-
Male	144	167	178	185	178	279	332	364	220	65	18	-
Female	67	87	108	115	186	322	558	669	713	371	88	-
Black, both sexes	76	94	82	86	83	95	109	102	75	31	10	-
Male	50	51	49	50	41	40	39	34	20	7	1	-
Female	26	43	33	36	42	55	70	68	55	24	9	-
American Indian 1/, both sexes	1	6	4	1	3	5	-	3	2	-	-	-
Male	1	3	3	-	1	2	-	2	-	-	-	-
Female	-	3	1	1	2	3	-	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	16	22	27	37	65	73	108	107	79	51	7	-
Male	13	12	14	26	33	32	42	46	27	18	1	-
Female	3	10	13	11	32	41	66	61	52	33	6	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Colorado												
All races, both sexes	21	20	31	24	34	54	84	99	55	49	9	-
Male	11	15	16	18	23	25	29	27	10	8	-	-
Female	10	5	15	6	11	29	55	72	45	41	9	-
White, both sexes	16	17	28	22	29	51	81	91	54	46	9	-
Male	7	12	14	16	21	23	26	26	10	8	-	-
Female	9	5	14	6	8	28	55	65	44	38	9	-
Black, both sexes	4	2	3	2	4	3	3	7	-	3	-	-
Male	3	2	2	2	2	2	3	1	-	-	-	-
Female	1	-	1	-	2	1	-	6	-	3	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	1	1	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Connecticut												
All races, both sexes	15	22	27	33	27	56	73	102	106	47	9	-
Male	13	18	19	13	13	18	32	28	24	9	-	-
Female	2	4	8	20	14	38	41	74	82	38	9	-
White, both sexes	11	17	22	25	23	52	65	97	101	46	9	-
Male	9	16	16	10	13	17	29	25	23	8	-	-
Female	2	1	6	15	10	35	36	72	78	38	9	-
Black, both sexes	4	5	3	7	3	4	7	4	5	1	-	-
Male	4	2	2	2	-	1	3	3	1	1	-	-
Female	-	3	1	5	3	3	4	1	4	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Delaware												
All races, both sexes	3	5	14	6	9	22	29	24	23	15	3	-
Male	-	2	8	5	4	7	11	12	4	4	1	-
Female	3	3	6	1	5	15	18	12	19	11	2	-
White, both sexes	2	3	7	3	7	16	27	21	20	11	3	-
Male	-	1	5	2	3	5	9	10	4	4	1	-
Female	2	2	2	1	4	11	18	11	16	7	2	-
Black, both sexes	1	2	7	3	2	6	2	3	3	4	-	-
Male	-	1	3	3	1	2	2	2	-	-	-	-
Female	1	1	4	-	1	4	-	1	3	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	20	16	21	21	17	29	46	47	23	13	4	-
Male	16	7	15	14	7	13	19	18	7	6	1	-
Female	4	9	6	7	10	16	27	29	16	7	3	-
White, both sexes	1	-	2	1	1	3	7	3	5	6	1	-
Male	1	-	2	1	-	2	3	2	2	1	-	-
Female	-	-	-	-	1	1	4	1	3	5	1	-
Black, both sexes	18	16	19	20	16	25	37	44	17	7	3	-
Male	14	7	13	13	7	10	14	16	5	5	1	-
Female	4	9	6	7	9	15	23	28	12	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	1	-	1	-	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Florida												
All races, both sexes	205	243	248	280	320	462	679	730	548	261	47	-
Male	138	156	170	146	173	220	282	253	138	53	10	-
Female	67	87	78	134	147	242	397	477	410	208	37	-
White, both sexes	120	160	172	188	242	377	585	639	495	234	42	-
Male	87	109	122	105	127	178	248	220	131	45	8	-
Female	33	51	50	83	115	199	337	419	364	189	34	-
Black, both sexes	82	82	73	86	77	83	88	89	51	27	5	-
Male	49	47	47	38	46	40	30	33	7	8	2	-
Female	33	35	26	48	31	43	58	56	44	19	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	3	6	1	2	5	2	2	-	-	-
Male	1	-	1	3	-	2	3	-	-	-	-	-
Female	1	1	2	3	1	-	2	2	2	-	-	-
Georgia												
All races, both sexes	140	167	158	193	179	203	257	248	202	84	24	-
Male	104	113	98	99	98	91	90	60	43	9	3	-
Female	36	54	60	94	81	112	167	188	159	75	21	-
White, both sexes	47	72	84	90	84	109	166	161	140	53	10	-
Male	42	49	56	51	48	51	55	43	27	5	1	-
Female	5	23	28	39	36	58	111	118	113	48	9	-
Black, both sexes	93	93	72	102	93	93	90	86	62	31	14	-
Male	62	62	41	48	49	39	34	17	16	4	2	-
Female	31	31	31	54	44	54	56	69	46	27	12	-
American Indian 1/, both sexes	-	-	1	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	1	1	-	1	-	-	-	-	-
Male	-	2	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Hawaii												
All races, both sexes	10	5	6	8	11	17	17	27	29	14	2	-
Male	6	5	2	3	4	10	5	12	9	4	-	-
Female	4	-	4	5	7	7	12	15	20	10	2	-
White, both sexes	2	1	1	1	4	1	4	1	5	1	-	-
Male	1	1	-	-	2	-	1	1	-	-	-	-
Female	1	-	1	1	2	1	3	-	5	1	-	-
Black, both sexes	1	-	1	-	-	1	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	4	4	7	7	15	13	26	24	13	2	-
Male	4	4	1	3	2	10	4	11	9	4	-	-
Female	3	-	3	4	5	5	9	15	15	9	2	-
Idaho												
All races, both sexes	5	5	5	9	11	19	32	35	32	17	1	-
Male	3	3	4	7	6	4	15	14	9	1	-	-
Female	2	2	1	2	5	15	17	21	23	16	1	-
White, both sexes	5	5	5	9	11	18	32	35	32	17	1	-
Male	3	3	4	7	6	4	15	14	9	1	-	-
Female	2	2	1	2	5	14	17	21	23	16	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Illinois												
All races, both sexes	129	143	178	134	197	267	337	410	326	181	40	-
Male	90	98	127	75	104	127	124	130	65	26	6	-
Female	39	45	51	59	93	140	213	280	261	155	34	-
White, both sexes	65	61	104	74	122	193	262	347	296	154	36	-
Male	49	45	77	38	64	91	99	109	59	21	5	-
Female	16	16	27	36	58	102	163	238	237	133	31	-
Black, both sexes	62	80	74	56	71	71	70	61	26	24	4	-
Male	39	51	50	34	37	36	22	21	6	4	1	-
Female	23	29	24	22	34	35	48	40	20	20	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	2	-	4	4	3	4	2	4	2	-	-
Male	2	2	-	3	3	-	3	-	-	1	-	-
Female	-	-	-	1	1	3	1	2	4	1	-	-
Indiana												
All races, both sexes	33	38	42	53	74	111	173	223	162	92	22	-
Male	24	25	29	24	35	41	60	67	33	11	1	-
Female	9	13	13	29	39	70	113	156	129	81	21	-
White, both sexes	22	31	36	39	57	97	154	203	154	88	17	-
Male	17	21	26	17	30	33	50	57	30	10	1	-
Female	5	10	10	22	27	64	104	146	124	78	16	-
Black, both sexes	10	7	6	14	17	13	19	18	8	4	5	-
Male	7	4	3	7	5	8	10	9	3	1	-	-
Female	3	3	3	7	12	5	9	9	5	3	5	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Iowa												
All races, both sexes	14	6	14	22	21	58	106	127	129	67	16	-
Male	7	4	6	16	9	29	42	45	36	14	1	-
Female	7	2	8	6	12	29	64	82	93	53	15	-
White, both sexes	11	5	12	22	20	58	105	126	129	66	16	-
Male	5	3	5	16	8	29	42	45	36	14	1	-
Female	6	2	7	6	12	29	63	81	93	52	15	-
Black, both sexes	3	1	2	-	1	-	1	1	-	1	-	-
Male	2	1	1	-	1	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	12	8	11	10	15	35	46	77	86	24	7	-
Male	9	6	8	4	8	17	14	18	20	7	1	-
Female	3	2	3	6	7	18	32	59	66	17	6	-
White, both sexes	7	7	8	10	15	33	41	71	79	24	6	-
Male	5	5	6	4	8	15	10	16	17	7	1	-
Female	2	2	2	6	7	18	31	55	62	17	5	-
Black, both sexes	5	1	3	-	-	2	4	6	6	-	1	-
Male	4	1	2	-	-	2	3	2	2	-	-	-
Female	1	-	1	-	-	-	1	4	4	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Kentucky												
All races, both sexes	29	47	50	40	53	84	114	118	92	45	8	-
Male	20	25	29	18	24	37	36	28	21	8	2	-
Female	9	22	21	22	29	47	78	90	71	37	6	-
White, both sexes	20	35	40	36	50	73	101	105	85	40	7	-
Male	15	19	26	18	22	34	34	25	20	8	1	-
Female	5	16	14	18	28	39	67	80	65	32	6	-
Black, both sexes	9	11	10	4	3	10	13	13	7	5	-	-
Male	5	6	3	-	2	2	2	3	1	-	-	-
Female	4	5	7	4	1	8	11	10	6	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	1	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	65	79	70	72	98	139	176	154	125	54	10	-
Male	47	55	42	40	52	56	69	46	18	7	1	-
Female	18	24	28	32	46	83	107	108	107	47	9	-
White, both sexes	33	36	30	30	53	82	109	109	81	34	8	-
Male	24	24	18	20	29	30	43	33	13	3	1	-
Female	9	12	12	10	24	52	66	76	68	31	7	-
Black, both sexes	32	43	40	42	45	56	67	45	44	19	2	-
Male	23	31	24	20	23	25	26	13	5	4	-	-
Female	9	12	16	22	22	31	41	32	39	15	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Maine												
All races, both sexes	5	10	8	5	13	14	27	23	17	11	2	-
Male	4	7	7	5	5	6	12	5	4	2	-	-
Female	1	3	1	-	8	8	15	18	13	9	2	-
White, both sexes	5	10	8	5	12	14	27	22	17	11	2	-
Male	4	7	7	5	5	6	12	5	4	2	-	-
Female	1	3	1	-	7	8	15	17	13	9	2	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	94	87	71	74	89	127	170	178	141	82	14	-
Male	65	65	41	47	56	51	57	60	24	15	-	-
Female	29	22	30	27	33	76	113	118	117	67	14	-
White, both sexes	24	40	25	31	49	72	111	135	113	63	11	-
Male	17	33	13	24	31	33	43	47	21	10	-	-
Female	7	7	12	7	18	39	68	88	92	53	11	-
Black, both sexes	68	46	45	42	40	53	58	40	26	18	3	-
Male	47	31	28	22	25	17	13	12	3	4	-	-
Female	21	15	17	20	15	36	45	28	23	14	3	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	1	-	1	1	3	2	-	-	-
Male	1	-	-	1	-	1	1	1	-	-	-	-
Female	1	-	1	-	-	-	-	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Massachusetts												
All races, both sexes	40	55	27	27	44	82	105	151	129	47	11	-
Male	27	42	20	15	16	39	40	45	23	9	1	-
Female	13	13	7	12	28	43	65	106	106	38	10	-
White, both sexes	35	49	25	22	35	70	93	143	124	47	10	-
Male	25	39	19	13	14	32	37	42	22	9	1	-
Female	10	10	6	9	21	38	56	101	102	38	9	-
Black, both sexes	5	6	2	5	8	8	10	6	4	-	1	-
Male	2	3	1	2	2	4	2	1	1	-	-	-
Female	3	3	1	3	6	4	8	5	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	4	2	2	1	-	-	-
Male	-	-	-	-	-	3	1	2	-	-	-	-
Female	-	-	-	-	1	1	1	-	1	-	-	-
Michigan												
All races, both sexes	91	112	123	153	156	227	319	384	303	162	31	-
Male	48	69	83	74	75	102	119	114	66	20	5	-
Female	43	43	40	79	81	125	200	270	237	142	26	-
White, both sexes	46	75	80	100	104	177	252	335	269	132	31	-
Male	24	49	55	49	48	77	97	97	58	15	5	-
Female	22	26	25	51	56	100	155	238	211	117	26	-
Black, both sexes	45	37	41	52	51	50	67	49	34	27	-	-
Male	24	20	28	25	27	25	22	17	8	5	-	-
Female	21	17	13	27	24	25	45	32	26	22	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	-	-	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	1	-	-	-	-	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Minnesota												
All races, both sexes	19	23	33	37	47	78	102	138	158	74	19	-
Male	14	11	18	18	27	40	46	31	37	13	2	-
Female	5	12	15	19	20	38	56	107	121	61	17	-
White, both sexes	18	18	29	36	44	69	99	137	157	74	18	-
Male	13	9	16	17	25	36	45	31	37	13	2	-
Female	5	9	13	19	19	33	54	106	120	61	16	-
Black, both sexes	1	5	4	-	3	5	2	1	1	-	1	-
Male	1	2	2	-	2	3	-	-	-	-	-	-
Female	-	3	2	-	1	2	2	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	3	1	-	-	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Mississippi												
All races, both sexes	52	67	66	84	97	122	140	185	129	72	21	-
Male	38	42	41	37	47	42	50	50	26	14	5	-
Female	14	25	25	47	50	80	90	135	103	58	16	-
White, both sexes	22	29	31	29	38	54	84	119	84	47	5	-
Male	17	21	18	16	22	20	30	31	13	6	1	-
Female	5	8	13	13	16	34	54	88	71	41	4	-
Black, both sexes	30	37	35	55	59	67	55	66	45	24	16	-
Male	21	20	23	21	25	21	19	19	13	8	4	-
Female	9	17	12	34	34	46	36	47	32	16	12	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Missouri												
All races, both sexes	43	47	64	54	71	86	139	159	126	60	10	-
Male	27	30	39	31	40	35	46	52	23	11	1	-
Female	16	17	25	23	31	51	93	107	103	49	9	-
White, both sexes	20	30	38	30	57	62	112	146	108	48	9	-
Male	13	18	28	18	31	23	39	49	17	9	1	-
Female	7	12	10	12	26	39	73	97	91	39	8	-
Black, both sexes	22	17	25	23	14	24	27	13	18	12	1	-
Male	13	12	10	12	9	12	7	3	6	2	-	-
Female	9	5	15	11	5	12	20	10	12	10	1	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	4	2	10	1	8	10	21	27	15	12	3	-
Male	2	2	4	-	2	6	4	7	2	1	-	-
Female	2	-	6	1	6	4	17	20	13	11	3	-
White, both sexes	4	2	9	1	7	10	19	27	15	12	3	-
Male	2	2	3	-	2	6	4	7	2	1	-	-
Female	2	-	6	1	5	4	15	20	13	11	3	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Nebraska												
All races, both sexes	7	7	12	14	19	21	45	67	63	27	6	-
Male	5	6	8	10	12	6	18	18	13	6	1	-
Female	2	1	4	4	7	15	27	49	50	21	5	-
White, both sexes	5	7	6	13	19	17	40	65	60	25	6	-
Male	4	6	5	9	12	3	14	18	12	5	1	-
Female	1	1	1	4	7	14	26	47	48	20	5	-
Black, both sexes	1	-	6	1	-	4	5	1	2	2	-	-
Male	1	-	3	1	-	3	4	-	-	1	-	-
Female	-	-	3	-	-	1	1	1	2	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	52	58	55	64	64	79	88	63	44	16	-	-
Male	39	42	37	41	36	46	48	21	17	3	-	-
Female	13	16	18	23	28	33	40	42	27	13	-	-
White, both sexes	42	44	46	55	53	74	81	59	41	15	-	-
Male	30	32	32	35	29	44	44	19	16	3	-	-
Female	12	12	14	20	24	30	37	40	25	12	-	-
Black, both sexes	5	12	7	5	8	5	5	-	1	-	-	-
Male	4	8	3	3	5	2	3	-	-	-	-	-
Female	1	4	4	2	3	3	2	-	1	-	-	-
American Indian 1/, both sexes	1	1	1	3	1	-	1	1	-	-	-	-
Male	1	1	1	2	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	1	1	2	-	1	3	2	1	-	-
Male	4	1	1	1	2	-	1	1	1	-	-	-
Female	-	-	-	-	-	-	-	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
New Hampshire												
All races, both sexes	6	10	11	10	10	11	20	31	29	14	6	-
Male	6	6	6	7	5	8	7	9	8	-	1	-
Female	-	4	5	3	5	3	13	22	21	14	5	-
White, both sexes	6	10	11	10	10	11	20	31	29	14	6	-
Male	6	6	6	7	5	8	7	9	8	-	1	-
Female	-	4	5	3	5	3	13	22	21	14	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	62	73	92	72	99	151	202	252	207	111	30	-
Male	39	52	56	52	51	70	79	78	44	19	4	-
Female	23	21	36	20	48	81	123	174	163	92	26	-
White, both sexes	31	42	56	42	57	111	168	236	186	102	25	-
Male	22	34	38	28	29	51	67	73	39	16	4	-
Female	9	8	18	14	28	60	101	163	147	86	21	-
Black, both sexes	30	28	33	29	39	33	29	14	20	8	5	-
Male	16	17	16	23	19	17	11	4	5	2	-	-
Female	14	11	17	6	20	16	18	10	15	6	5	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	3	1	2	6	5	2	1	1	-	-
Male	1	1	2	1	2	2	1	1	-	1	-	-
Female	-	2	1	-	-	4	4	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
New Mexico												
All races, both sexes	22	15	18	15	29	36	34	44	37	17	2	-
Male	16	9	14	10	19	15	15	17	5	5	1	-
Female	6	6	4	5	10	21	19	27	32	12	1	-
White, both sexes	18	15	17	13	24	32	31	39	34	16	2	-
Male	13	9	13	8	16	12	13	14	4	5	1	-
Female	5	6	4	5	8	20	18	25	30	11	1	-
Black, both sexes	2	-	1	2	2	2	1	3	1	1	-	-
Male	1	-	1	2	1	2	1	1	-	-	-	-
Female	1	-	-	-	1	-	-	2	1	1	-	-
American Indian 1/, both sexes	2	-	-	-	2	1	2	2	2	-	-	-
Male	2	-	-	-	1	1	1	2	1	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
New York												
All races, both sexes	205	219	222	283	381	455	640	672	575	273	61	-
Male	128	142	113	157	168	213	247	212	135	46	8	-
Female	77	77	109	126	213	242	393	460	440	227	53	-
White, both sexes	100	110	119	158	255	308	458	534	472	237	47	-
Male	62	71	64	86	109	150	187	168	116	42	7	-
Female	38	39	55	72	146	158	271	366	356	195	40	-
Black, both sexes	101	101	96	121	114	131	158	123	89	32	14	-
Male	62	67	47	68	48	56	51	38	15	3	1	-
Female	39	34	49	53	66	75	107	85	74	29	13	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	8	6	4	12	15	24	14	14	4	-	-
Male	4	4	1	3	11	6	9	6	4	1	-	-
Female	-	4	5	1	1	9	15	8	10	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
North Carolina												
All races, both sexes	76	77	82	109	127	169	241	204	215	82	19	1
Male	61	59	55	67	60	69	89	60	49	11	2	1
Female	15	18	27	42	67	100	152	144	166	71	17	-
White, both sexes	30	38	49	56	66	95	153	138	162	58	14	1
Male	28	29	32	35	28	38	55	44	37	8	2	1
Female	2	9	17	21	38	57	98	94	125	50	12	-
Black, both sexes	43	38	33	51	58	73	86	65	50	24	4	-
Male	31	29	23	31	29	30	34	16	11	3	-	-
Female	12	9	10	20	29	43	52	49	39	21	4	-
American Indian 1/, both sexes	3	1	-	2	3	1	-	1	3	-	1	-
Male	2	1	-	1	3	1	-	-	1	-	-	-
Female	1	-	-	1	-	-	-	1	2	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
North Dakota												
All races, both sexes	1	2	6	3	5	14	12	26	33	21	1	-
Male	1	1	5	1	4	9	6	12	14	3	-	-
Female	-	1	1	2	1	5	6	14	19	18	1	-
White, both sexes	1	2	5	3	5	13	12	26	33	21	1	-
Male	1	1	4	1	4	9	6	12	14	3	-	-
Female	-	1	1	2	1	4	6	14	19	18	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Ohio												
All races, both sexes	142	145	148	150	178	293	445	480	422	201	45	-
Male	99	92	96	84	100	130	169	163	99	28	6	-
Female	43	53	52	66	78	163	276	317	323	173	39	-
White, both sexes	84	100	104	100	132	231	378	429	384	179	38	-
Male	61	68	65	55	71	100	142	142	86	25	4	-
Female	23	32	39	45	61	131	236	287	298	154	34	-
Black, both sexes	56	44	43	47	45	60	67	50	37	22	7	-
Male	36	24	31	28	29	30	27	20	13	3	2	-
Female	20	20	12	19	16	30	40	30	24	19	5	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	3	-	2	-	1	1	-	-	-
Male	2	-	-	1	-	-	-	1	-	-	-	-
Female	-	1	1	2	-	2	-	-	1	-	-	-
Oklahoma												
All races, both sexes	16	28	24	34	47	78	121	120	101	25	19	-
Male	13	18	12	20	24	35	46	30	28	5	1	-
Female	3	10	12	14	23	43	75	90	73	20	18	-
White, both sexes	11	17	21	26	34	55	99	100	91	23	15	-
Male	9	12	11	16	18	24	37	25	25	4	1	-
Female	2	5	10	10	16	31	62	75	66	19	14	-
Black, both sexes	2	9	2	5	9	13	15	13	7	2	3	-
Male	1	5	-	2	3	7	3	3	-	1	-	-
Female	1	4	2	3	6	6	12	10	7	1	3	-
American Indian 1/, both sexes	3	1	1	3	4	9	7	7	3	-	1	-
Male	3	1	1	2	3	4	6	2	3	-	-	-
Female	-	-	-	1	1	5	1	5	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Oregon												
All races, both sexes	18	20	21	21	33	65	115	146	155	72	17	-
Male	14	16	12	11	16	29	47	53	28	14	3	-
Female	4	4	9	10	17	36	68	93	127	58	14	-
White, both sexes	16	17	19	20	29	61	108	142	151	70	16	-
Male	13	13	12	10	14	27	44	52	28	13	2	-
Female	3	4	7	10	15	34	64	90	123	57	14	-
Black, both sexes	1	3	1	-	4	2	5	-	2	1	1	-
Male	-	3	-	-	2	2	3	-	-	-	1	-
Female	1	-	1	-	2	-	2	-	2	1	-	-
American Indian 1/, both sexes	-	-	1	1	-	1	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	1	3	2	1	-	-
Male	1	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	1	1	2	2	-	-	-
Pennsylvania												
All races, both sexes	70	92	100	113	140	228	321	352	316	145	31	-
Male	48	70	66	74	72	99	111	122	71	22	3	-
Female	22	22	34	39	68	129	210	230	245	123	28	-
White, both sexes	42	63	68	83	114	194	285	322	293	133	25	-
Male	33	47	45	54	58	83	93	111	66	18	3	-
Female	9	16	23	29	56	111	192	211	227	115	22	-
Black, both sexes	28	29	32	28	26	34	34	29	21	12	6	-
Male	15	23	21	18	14	16	16	11	5	4	-	-
Female	13	6	11	10	12	18	18	18	16	8	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	2	1	2	-	-	-
Male	-	-	-	2	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Rhode Island												
All races, both sexes	2	4	6	4	9	13	25	24	19	11	2	-
Male	1	3	4	2	2	5	13	7	4	2	-	-
Female	1	1	2	2	7	8	12	17	15	9	2	-
White, both sexes	2	3	5	4	8	12	24	22	19	11	2	-
Male	1	2	3	2	1	4	12	7	4	2	-	-
Female	1	1	2	2	7	8	12	15	15	9	2	-
Black, both sexes	-	1	1	-	1	1	-	2	-	-	-	-
Male	-	1	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	46	54	44	58	86	100	136	118	85	44	9	-
Male	29	33	26	32	40	44	43	33	20	9	2	-
Female	17	21	18	26	46	56	93	85	65	35	7	-
White, both sexes	15	16	22	27	44	51	85	78	55	26	5	-
Male	11	8	14	14	21	23	30	20	13	2	1	-
Female	4	8	8	13	23	28	55	58	42	24	4	-
Black, both sexes	31	37	22	31	42	48	50	40	30	18	4	-
Male	18	24	12	18	19	20	13	13	7	7	1	-
Female	13	13	10	13	23	28	37	27	23	11	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
South Dakota												
All races, both sexes	-	1	3	7	6	6	18	36	35	26	5	-
Male	-	1	1	6	6	3	8	15	10	6	-	-
Female	-	-	2	1	-	3	10	21	25	20	5	-
White, both sexes	-	1	2	5	4	5	18	36	34	26	5	-
Male	-	1	-	4	4	2	8	15	10	6	-	-
Female	-	-	2	1	-	3	10	21	24	20	5	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	2	2	-	-	-	1	-	-	-
Male	-	-	1	2	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	79	80	58	93	118	146	254	255	202	98	16	-
Male	53	52	36	48	55	75	74	64	50	16	1	-
Female	26	28	22	45	63	71	180	191	152	82	15	-
White, both sexes	41	50	23	50	69	94	209	206	158	76	15	-
Male	29	32	14	31	35	47	59	56	37	14	1	-
Female	12	18	9	19	34	47	150	150	121	62	14	-
Black, both sexes	37	30	33	42	47	51	45	49	44	22	1	-
Male	24	20	21	16	18	28	15	8	13	2	-	-
Female	13	10	12	26	29	23	30	41	31	20	1	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	2	1	-	-	-	-	-	-
Male	-	-	1	-	2	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Texas												
All races, both sexes	201	238	229	211	272	388	527	507	445	222	47	-
Male	126	150	145	127	125	158	186	144	109	34	1	-
Female	75	88	84	84	147	230	341	363	336	188	46	-
White, both sexes	131	158	161	155	205	310	434	439	386	193	36	-
Male	88	103	102	94	92	125	157	124	96	28	1	-
Female	43	55	59	61	113	185	277	315	290	165	35	-
Black, both sexes	68	75	67	53	64	76	88	67	55	27	9	-
Male	38	42	42	32	32	32	27	20	12	6	-	-
Female	30	33	25	21	32	44	61	47	43	21	9	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	5	1	3	3	2	4	1	4	2	1	-
Male	-	5	1	1	1	1	2	-	1	-	-	-
Female	2	-	-	2	2	1	2	1	3	2	1	-
Utah												
All races, both sexes	7	9	10	9	11	22	43	50	38	16	1	-
Male	2	7	6	8	5	5	14	16	8	1	1	-
Female	5	2	4	1	6	17	29	34	30	15	-	-
White, both sexes	6	8	10	8	11	22	42	50	38	16	1	-
Male	2	6	6	7	5	5	14	16	8	1	1	-
Female	4	2	4	1	6	17	28	34	30	15	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Vermont												
All races, both sexes	3	7	2	2	7	10	10	20	21	6	1	-
Male	2	5	1	2	2	3	4	6	9	1	-	-
Female	1	2	1	-	5	7	6	14	12	5	1	-
White, both sexes	3	7	2	2	7	10	10	20	21	6	1	-
Male	2	5	1	2	2	3	4	6	9	1	-	-
Female	1	2	1	-	5	7	6	14	12	5	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	51	62	65	69	78	121	146	157	140	49	17	-
Male	33	41	36	37	29	57	49	45	24	5	2	-
Female	18	21	29	32	49	64	97	112	116	44	15	-
White, both sexes	22	32	32	35	43	72	100	120	92	42	5	-
Male	16	20	19	19	16	34	27	34	14	5	-	-
Female	6	12	13	16	27	38	73	86	78	37	5	-
Black, both sexes	29	28	33	34	33	44	44	34	48	7	11	-
Male	17	20	17	18	13	22	21	10	10	-	1	-
Female	12	8	16	16	20	22	23	24	38	7	10	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	2	5	2	3	-	-	1	-
Male	-	1	-	-	-	1	1	1	-	-	1	-
Female	-	1	-	-	2	4	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Washington												
All races, both sexes	40	41	41	41	64	111	195	211	186	90	20	-
Male	31	35	19	19	31	53	81	67	39	10	3	-
Female	9	6	22	22	33	58	114	144	147	80	17	-
White, both sexes	29	34	38	36	52	100	180	200	179	87	19	-
Male	24	31	17	17	24	49	73	62	37	8	2	-
Female	5	3	21	19	28	51	107	138	142	79	17	-
Black, both sexes	8	5	2	4	4	3	3	6	-	1	-	-
Male	4	3	1	1	2	1	1	2	-	-	-	-
Female	4	2	1	3	2	2	2	4	-	1	-	-
American Indian 1/, both sexes	1	1	-	-	3	1	3	-	1	-	-	-
Male	1	1	-	-	2	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	1	5	7	9	5	6	2	1	-
Male	2	-	1	1	3	2	6	3	2	2	1	-
Female	-	1	-	-	2	5	3	2	4	-	-	-
West Virginia												
All races, both sexes	10	21	25	27	30	58	73	71	73	35	8	-
Male	6	14	14	14	14	27	25	17	11	5	1	-
Female	4	7	11	13	16	31	48	54	62	30	7	-
White, both sexes	8	21	21	25	30	53	70	70	70	34	8	-
Male	5	14	11	14	14	26	24	16	11	5	1	-
Female	3	7	10	11	16	27	46	54	59	29	7	-
Black, both sexes	2	-	4	2	-	5	3	1	3	1	-	-
Male	1	-	3	-	-	1	1	1	-	-	-	-
Female	1	-	1	2	-	4	2	-	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive diseases (I10-I13)												
Wisconsin												
All races, both sexes	24	29	33	48	52	104	131	191	181	99	17	-
Male	20	15	25	22	29	38	44	55	40	8	3	-
Female	4	14	8	26	23	66	87	136	141	91	14	-
White, both sexes	16	23	29	37	50	99	123	186	173	94	16	-
Male	14	13	23	17	28	38	41	53	38	6	3	-
Female	2	10	6	20	22	61	82	133	135	88	13	-
Black, both sexes	8	6	3	10	2	4	7	5	7	4	-	-
Male	6	2	2	5	1	-	3	2	1	1	-	-
Female	2	4	1	5	1	4	4	3	6	3	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	-	1	1	1	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	1	-	1	1	-	-	-	1	-
Wyoming												
All races, both sexes	3	4	3	2	4	3	5	8	7	3	-	-
Male	1	2	2	1	1	3	2	2	2	-	-	-
Female	2	2	1	1	3	-	3	6	5	3	-	-
White, both sexes	3	4	3	2	4	3	5	8	7	3	-	-
Male	1	2	2	1	1	3	2	2	2	-	-	-
Female	2	2	1	1	3	-	3	6	5	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
United States												
All races, both sexes	14,857	2	-	1	1	2	5	1	22	47	144	256
Male	5,084	1	-	-	1	1	4	1	10	33	80	159
Female	9,773	1	-	1	-	1	1	-	12	14	64	97
White, both sexes	12,219	-	-	1	1	1	2	-	10	30	86	161
Male	3,996	-	-	-	1	1	2	-	3	22	56	112
Female	8,223	-	-	1	-	-	-	-	7	8	30	49
Black, both sexes	2,278	2	-	-	-	1	3	1	12	16	54	87
Male	924	1	-	-	-	-	2	1	7	10	23	41
Female	1,354	1	-	-	-	1	1	-	5	6	31	46
American Indian 1/, both sexes	59	-	-	-	-	-	-	-	-	-	1	2
Male	30	-	-	-	-	-	-	-	-	-	-	2
Female	29	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	301	-	-	-	-	-	-	-	-	1	3	6
Male	134	-	-	-	-	-	-	-	-	1	1	4
Female	167	-	-	-	-	-	-	-	-	-	2	2
Alabama												
All races, both sexes	272	-	-	-	-	1	-	-	1	1	7	7
Male	104	-	-	-	-	-	-	-	1	1	1	5
Female	168	-	-	-	-	1	-	-	-	-	6	2
White, both sexes	165	-	-	-	-	-	-	-	-	1	3	1
Male	55	-	-	-	-	-	-	-	-	1	-	1
Female	110	-	-	-	-	-	-	-	-	-	3	-
Black, both sexes	106	-	-	-	-	1	-	-	1	-	4	6
Male	49	-	-	-	-	-	-	-	1	-	1	4
Female	57	-	-	-	-	1	-	-	-	-	3	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Alaska												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	190	-	-	-	-	-	-	-	1	-	1	3
Male	60	-	-	-	-	-	-	-	-	-	-	1
Female	130	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	171	-	-	-	-	-	-	-	1	-	-	3
Male	55	-	-	-	-	-	-	-	-	-	-	1
Female	116	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Arkansas												
All races, both sexes	165	-	-	-	-	-	-	-	1	-	6	2
Male	65	-	-	-	-	-	-	-	1	-	4	1
Female	100	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	134	-	-	-	-	-	-	-	1	-	5	1
Male	56	-	-	-	-	-	-	-	1	-	4	1
Female	78	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	30	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,841	-	-	-	-	-	-	-	-	5	22	24
Male	664	-	-	-	-	-	-	-	-	4	11	14
Female	1,177	-	-	-	-	-	-	-	-	1	11	10
White, both sexes	1,498	-	-	-	-	-	-	-	-	1	15	17
Male	518	-	-	-	-	-	-	-	-	1	10	10
Female	980	-	-	-	-	-	-	-	-	-	5	7
Black, both sexes	184	-	-	-	-	-	-	-	-	4	4	5
Male	75	-	-	-	-	-	-	-	-	3	1	3
Female	109	-	-	-	-	-	-	-	-	1	3	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	154	-	-	-	-	-	-	-	-	-	2	2
Male	69	-	-	-	-	-	-	-	-	-	-	1
Female	85	-	-	-	-	-	-	-	-	-	2	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Colorado												
All races, both sexes	136	-	-	-	-	-	-	-	-	-	-	1
Male	44	-	-	-	-	-	-	-	-	-	-	1
Female	92	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	130	-	-	-	-	-	-	-	-	-	-	1
Male	40	-	-	-	-	-	-	-	-	-	-	1
Female	90	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	262	-	-	-	-	-	-	-	-	1	3	10
Male	84	-	-	-	-	-	-	-	-	1	3	5
Female	178	-	-	-	-	-	-	-	-	-	-	5
White, both sexes	239	-	-	-	-	-	-	-	-	-	3	8
Male	76	-	-	-	-	-	-	-	-	-	3	4
Female	163	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Delaware												
All races, both sexes	45	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	37	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Florida												
All races, both sexes	1,002	1	-	-	-	-	-	-	-	5	11	17
Male	365	-	-	-	-	-	-	-	-	4	6	12
Female	637	1	-	-	-	-	-	-	-	1	5	5
White, both sexes	835	-	-	-	-	-	-	-	-	4	5	14
Male	288	-	-	-	-	-	-	-	-	3	3	10
Female	547	-	-	-	-	-	-	-	-	1	2	4
Black, both sexes	162	1	-	-	-	-	-	-	-	1	6	3
Male	74	-	-	-	-	-	-	-	-	1	3	2
Female	88	1	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	607	-	-	-	-	-	1	-	1	7	11	25
Male	237	-	-	-	-	-	-	-	-	4	4	17
Female	370	-	-	-	-	-	1	-	1	3	7	8
White, both sexes	394	-	-	-	-	-	-	-	-	4	5	11
Male	143	-	-	-	-	-	-	-	-	2	3	9
Female	251	-	-	-	-	-	-	-	-	2	2	2
Black, both sexes	213	-	-	-	-	-	1	-	1	3	6	14
Male	94	-	-	-	-	-	-	-	-	2	1	8
Female	119	-	-	-	-	-	1	-	1	1	5	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Hawaii												
All races, both sexes	63	-	-	-	-	-	-	-	-	-	1	-
Male	28	-	-	-	-	-	-	-	-	-	1	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	54	-	-	-	-	-	-	-	-	-	1	-
Male	27	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Illinois												
All races, both sexes	621	-	-	-	-	-	-	-	-	-	7	9
Male	209	-	-	-	-	-	-	-	-	-	6	3
Female	412	-	-	-	-	-	-	-	-	-	1	6
White, both sexes	523	-	-	-	-	-	-	-	-	-	4	4
Male	168	-	-	-	-	-	-	-	-	-	3	1
Female	355	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	92	-	-	-	-	-	-	-	-	-	3	5
Male	39	-	-	-	-	-	-	-	-	-	3	2
Female	53	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	380	-	-	-	-	-	-	-	1	-	3	2
Male	121	-	-	-	-	-	-	-	-	-	3	2
Female	259	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	339	-	-	-	-	-	-	-	1	-	2	2
Male	102	-	-	-	-	-	-	-	-	-	2	2
Female	237	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	40	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Iowa												
All races, both sexes	129	-	-	-	-	-	-	-	-	-	-	2
Male	36	-	-	-	-	-	-	-	-	-	-	1
Female	93	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	127	-	-	-	-	-	-	-	-	-	-	2
Male	36	-	-	-	-	-	-	-	-	-	-	1
Female	91	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	107	-	-	-	-	-	-	-	-	-	-	2
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	75	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	98	-	-	-	-	-	-	-	-	-	-	1
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	71	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Kentucky												
All races, both sexes	218	-	-	-	-	-	-	-	1	1	3	2
Male	65	-	-	-	-	-	-	-	-	1	2	2
Female	153	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	192	-	-	-	-	-	-	-	-	1	3	1
Male	58	-	-	-	-	-	-	-	-	1	2	1
Female	134	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	25	-	-	-	-	-	-	-	1	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	202	-	-	-	-	-	-	-	2	1	2	1
Male	60	-	-	-	-	-	-	-	2	1	-	-
Female	142	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	126	-	-	-	-	-	-	-	-	1	1	-
Male	34	-	-	-	-	-	-	-	-	1	-	-
Female	92	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	76	-	-	-	-	-	-	-	2	-	1	1
Male	26	-	-	-	-	-	-	-	2	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Maine												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	230	-	-	1	-	-	-	-	-	-	-	4
Male	76	-	-	-	-	-	-	-	-	-	-	3
Female	154	-	-	1	-	-	-	-	-	-	-	1
White, both sexes	159	-	-	1	-	-	-	-	-	-	-	2
Male	50	-	-	-	-	-	-	-	-	-	-	1
Female	109	-	-	1	-	-	-	-	-	-	-	1
Black, both sexes	66	-	-	-	-	-	-	-	-	-	-	2
Male	23	-	-	-	-	-	-	-	-	-	-	2
Female	43	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Massachusetts												
All races, both sexes	306	-	-	-	-	-	-	-	-	1	2	6
Male	109	-	-	-	-	-	-	-	-	-	-	2
Female	197	-	-	-	-	-	-	-	-	1	2	4
White, both sexes	286	-	-	-	-	-	-	-	-	1	1	5
Male	101	-	-	-	-	-	-	-	-	-	-	2
Female	185	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	18	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	452	-	-	-	-	-	-	-	1	-	1	6
Male	129	-	-	-	-	-	-	-	1	-	1	3
Female	323	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	391	-	-	-	-	-	-	-	-	-	1	2
Male	106	-	-	-	-	-	-	-	-	-	1	1
Female	285	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	60	-	-	-	-	-	-	-	1	-	-	4
Male	23	-	-	-	-	-	-	-	1	-	-	2
Female	37	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Minnesota												
All races, both sexes	377	-	-	-	-	-	-	-	-	1	-	4
Male	118	-	-	-	-	-	-	-	-	-	-	2
Female	259	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	362	-	-	-	-	-	-	-	-	-	-	3
Male	112	-	-	-	-	-	-	-	-	-	-	2
Female	250	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	12	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	245	-	-	-	-	-	1	-	1	4	4	6
Male	80	-	-	-	-	-	1	-	-	2	2	4
Female	165	-	-	-	-	-	-	-	1	2	2	2
White, both sexes	136	-	-	-	-	-	-	-	-	1	2	3
Male	41	-	-	-	-	-	-	-	-	1	2	2
Female	95	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	106	-	-	-	-	-	1	-	1	3	2	3
Male	37	-	-	-	-	-	1	-	-	1	-	2
Female	69	-	-	-	-	-	-	-	1	2	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Missouri												
All races, both sexes	276	-	-	-	-	-	1	-	-	2	2	6
Male	95	-	-	-	-	-	1	-	-	2	1	1
Female	181	-	-	-	-	-	-	-	-	-	1	5
White, both sexes	235	-	-	-	-	-	1	-	-	1	2	5
Male	81	-	-	-	-	-	1	-	-	1	1	1
Female	154	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	41	-	-	-	-	-	-	-	-	1	-	1
Male	14	-	-	-	-	-	-	-	-	1	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Nebraska												
All races, both sexes	130	-	-	-	-	-	-	-	-	2	1	-
Male	42	-	-	-	-	-	-	-	-	2	-	-
Female	88	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	124	-	-	-	-	-	-	-	-	2	-	-
Male	38	-	-	-	-	-	-	-	-	2	-	-
Female	86	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	78	-	-	-	-	-	-	-	-	1	1	-
Male	29	-	-	-	-	-	-	-	-	1	-	-
Female	49	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	73	-	-	-	-	-	-	-	-	1	1	-
Male	26	-	-	-	-	-	-	-	-	1	-	-
Female	47	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
New Hampshire												
All races, both sexes	51	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	51	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	367	-	-	-	-	-	-	-	-	-	4	2
Male	115	-	-	-	-	-	-	-	-	-	1	1
Female	252	-	-	-	-	-	-	-	-	-	3	1
White, both sexes	296	-	-	-	-	-	-	-	-	-	-	1
Male	86	-	-	-	-	-	-	-	-	-	-	1
Female	210	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	66	-	-	-	-	-	-	-	-	-	4	1
Male	27	-	-	-	-	-	-	-	-	-	1	-
Female	39	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
New Mexico												
All races, both sexes	62	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	59	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,062	1	-	-	-	-	1	1	-	2	9	14
Male	374	1	-	-	-	-	1	1	-	1	4	10
Female	688	-	-	-	-	-	-	-	-	1	5	4
White, both sexes	831	-	-	-	-	-	1	-	-	2	6	7
Male	286	-	-	-	-	-	1	-	-	1	2	6
Female	545	-	-	-	-	-	-	-	-	1	4	1
Black, both sexes	209	1	-	-	-	-	-	1	-	-	3	7
Male	80	1	-	-	-	-	-	1	-	-	2	4
Female	129	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
North Carolina												
All races, both sexes	406	-	-	-	-	-	-	-	3	1	6	7
Male	139	-	-	-	-	-	-	-	2	1	5	5
Female	267	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	278	-	-	-	-	-	-	-	1	1	1	3
Male	90	-	-	-	-	-	-	-	-	1	1	3
Female	188	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	122	-	-	-	-	-	-	-	2	-	5	3
Male	44	-	-	-	-	-	-	-	2	-	4	1
Female	78	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Ohio												
All races, both sexes	673	-	-	-	-	-	1	-	-	-	5	17
Male	220	-	-	-	-	-	1	-	-	-	4	11
Female	453	-	-	-	-	-	-	-	-	-	1	6
White, both sexes	592	-	-	-	-	-	-	-	-	-	4	13
Male	183	-	-	-	-	-	-	-	-	-	4	10
Female	409	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	79	-	-	-	-	-	1	-	-	-	1	4
Male	37	-	-	-	-	-	1	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	196	-	-	-	-	-	-	-	1	-	1	5
Male	66	-	-	-	-	-	-	-	-	-	1	3
Female	130	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	166	-	-	-	-	-	-	-	1	-	1	4
Male	53	-	-	-	-	-	-	-	-	-	1	2
Female	113	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Oregon												
All races, both sexes	311	-	-	-	-	-	-	-	-	-	1	3
Male	111	-	-	-	-	-	-	-	-	-	1	3
Female	200	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	297	-	-	-	-	-	-	-	-	-	1	3
Male	105	-	-	-	-	-	-	-	-	-	1	3
Female	192	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	711	-	-	-	-	-	-	-	3	-	2	18
Male	259	-	-	-	-	-	-	-	1	-	-	12
Female	452	-	-	-	-	-	-	-	2	-	2	6
White, both sexes	629	-	-	-	-	-	-	-	2	-	1	13
Male	220	-	-	-	-	-	-	-	1	-	-	10
Female	409	-	-	-	-	-	-	-	1	-	1	3
Black, both sexes	81	-	-	-	-	-	-	-	1	-	1	5
Male	38	-	-	-	-	-	-	-	-	-	-	2
Female	43	-	-	-	-	-	-	-	1	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Rhode Island												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	228	-	-	-	-	-	-	-	-	1	5	6
Male	92	-	-	-	-	-	-	-	-	1	3	4
Female	136	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	137	-	-	-	-	-	-	-	-	-	3	1
Male	44	-	-	-	-	-	-	-	-	-	1	1
Female	93	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	90	-	-	-	-	-	-	-	-	1	2	5
Male	48	-	-	-	-	-	-	-	-	1	2	3
Female	42	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
South Dakota												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	276	-	-	-	-	-	-	-	-	2	1	6
Male	88	-	-	-	-	-	-	-	-	1	-	2
Female	188	-	-	-	-	-	-	-	-	1	1	4
White, both sexes	211	-	-	-	-	-	-	-	-	-	1	5
Male	69	-	-	-	-	-	-	-	-	-	-	2
Female	142	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	65	-	-	-	-	-	-	-	-	2	-	1
Male	19	-	-	-	-	-	-	-	-	1	-	-
Female	46	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Texas												
All races, both sexes	795	-	-	-	1	-	-	-	2	5	8	20
Male	279	-	-	-	1	-	-	-	1	4	5	14
Female	516	-	-	-	-	-	-	-	1	1	3	6
White, both sexes	678	-	-	-	1	-	-	-	2	5	6	11
Male	240	-	-	-	1	-	-	-	1	4	4	11
Female	438	-	-	-	-	-	-	-	1	1	2	-
Black, both sexes	108	-	-	-	-	-	-	-	-	-	2	7
Male	36	-	-	-	-	-	-	-	-	-	1	1
Female	72	-	-	-	-	-	-	-	-	-	1	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	75	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	55	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	54	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Vermont												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	327	-	-	-	-	-	-	-	2	2	8	6
Male	111	-	-	-	-	-	-	-	1	-	6	5
Female	216	-	-	-	-	-	-	-	1	2	2	1
White, both sexes	223	-	-	-	-	-	-	-	-	2	3	3
Male	70	-	-	-	-	-	-	-	-	-	3	3
Female	153	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	99	-	-	-	-	-	-	-	2	-	5	3
Male	38	-	-	-	-	-	-	-	1	-	3	2
Female	61	-	-	-	-	-	-	-	1	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Washington												
All races, both sexes	279	-	-	-	-	1	-	-	1	1	3	2
Male	84	-	-	-	-	1	-	-	-	1	2	1
Female	195	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	255	-	-	-	-	1	-	-	1	1	3	-
Male	73	-	-	-	-	1	-	-	-	1	2	-
Female	182	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
West Virginia												
All races, both sexes	179	-	-	-	-	-	-	-	-	1	2	4
Male	57	-	-	-	-	-	-	-	-	1	2	3
Female	122	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	170	-	-	-	-	-	-	-	-	1	2	4
Male	54	-	-	-	-	-	-	-	-	1	2	3
Female	116	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Essential (primary) hypertension (I10)												
Wisconsin												
All races, both sexes	261	-	-	-	-	-	-	-	-	-	-	1
Male	69	-	-	-	-	-	-	-	-	-	-	-
Female	192	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	250	-	-	-	-	-	-	-	-	-	-	1
Male	64	-	-	-	-	-	-	-	-	-	-	-
Female	186	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
United States												
All races, both sexes	370	514	629	774	1,020	1,639	2,424	2,756	2,603	1,323	324	-
Male	246	319	389	432	494	686	817	716	476	179	40	-
Female	124	195	240	342	526	953	1,607	2,040	2,127	1,144	284	-
White, both sexes	237	345	467	569	778	1,306	2,069	2,440	2,295	1,162	259	-
Male	165	223	289	321	372	531	687	629	409	143	30	-
Female	72	122	178	248	406	775	1,382	1,811	1,886	1,019	229	-
Black, both sexes	119	157	151	184	214	275	297	260	250	139	56	-
Male	69	91	94	94	108	133	104	66	47	26	7	-
Female	50	66	57	90	106	142	193	194	203	113	49	-
American Indian 1/, both sexes	6	-	4	3	5	13	9	9	4	2	1	-
Male	5	-	3	2	3	5	5	2	2	1	-	-
Female	1	-	1	1	2	8	4	7	2	1	1	-
Asian or Pacific Islander 2/, both sexes	8	12	7	18	23	45	49	47	54	20	8	-
Male	7	5	3	15	11	17	21	19	18	9	3	-
Female	1	7	4	3	12	28	28	28	36	11	5	-
Alabama												
All races, both sexes	9	12	15	18	25	37	44	38	31	20	6	-
Male	5	7	9	13	13	15	11	13	5	4	1	-
Female	4	5	6	5	12	22	33	25	26	16	5	-
White, both sexes	2	5	7	13	11	26	29	27	22	14	4	-
Male	1	3	3	9	5	9	7	9	3	3	1	-
Female	1	2	4	4	6	17	22	18	19	11	3	-
Black, both sexes	7	7	8	5	14	10	15	11	9	6	2	-
Male	4	4	6	4	8	6	4	4	2	1	-	-
Female	3	3	2	1	6	4	11	7	7	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Alaska												
All races, both sexes	-	-	-	2	-	1	1	1	1	-	-	-
Male	-	-	-	1	-	1	-	1	1	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	1	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	4	5	13	7	28	29	37	40	16	4	-
Male	1	2	2	6	6	9	12	11	6	3	1	-
Female	1	2	3	7	1	19	17	26	34	13	3	-
White, both sexes	2	3	4	11	7	25	26	34	35	16	4	-
Male	1	1	1	5	6	8	12	10	6	3	1	-
Female	1	2	3	6	1	17	14	24	29	13	3	-
Black, both sexes	-	1	-	1	-	-	1	-	4	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	4	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	3	2	2	1	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	1	-	2	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Arkansas												
All races, both sexes	8	11	6	10	14	16	25	19	28	15	4	-
Male	5	7	5	4	8	7	8	5	5	5	-	-
Female	3	4	1	6	6	9	17	14	23	10	4	-
White, both sexes	5	8	5	8	12	11	22	18	24	12	2	-
Male	4	5	5	3	8	6	7	4	4	4	-	-
Female	1	3	-	5	4	5	15	14	20	8	2	-
Black, both sexes	3	3	1	2	2	5	3	1	3	3	2	-
Male	1	2	-	1	-	1	1	1	1	1	-	-
Female	2	1	1	1	2	4	2	-	2	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
California												
All races, both sexes	42	44	74	94	126	201	306	359	334	159	51	-
Male	27	30	46	56	63	87	110	114	68	26	8	-
Female	15	14	28	38	63	114	196	245	266	133	43	-
White, both sexes	32	30	56	65	92	160	257	296	296	137	44	-
Male	22	22	34	39	48	70	92	91	55	17	7	-
Female	10	8	22	26	44	90	165	205	241	120	37	-
Black, both sexes	8	13	12	18	19	16	20	32	18	11	4	-
Male	3	7	8	8	9	10	6	9	6	2	-	-
Female	5	6	4	10	10	6	14	23	12	9	4	-
American Indian 1/, both sexes	-	-	1	-	1	2	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	5	11	14	23	29	31	20	11	3	-
Male	2	1	3	9	6	6	12	14	7	7	1	-
Female	-	-	2	2	8	17	17	17	13	4	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Colorado												
All races, both sexes	5	3	5	7	10	18	21	26	18	17	5	-
Male	2	2	3	6	6	7	6	9	1	1	-	-
Female	3	1	2	1	4	11	15	17	17	16	5	-
White, both sexes	4	3	4	7	9	17	21	25	17	17	5	-
Male	1	2	2	6	5	6	6	9	1	1	-	-
Female	3	1	2	1	4	11	15	16	16	16	5	-
Black, both sexes	1	-	1	-	1	1	-	1	-	-	-	-
Male	1	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Connecticut												
All races, both sexes	5	11	8	15	12	26	40	55	52	22	2	-
Male	3	7	6	6	6	6	13	14	9	5	-	-
Female	2	4	2	9	6	20	27	41	43	17	2	-
White, both sexes	5	8	8	9	11	24	37	53	50	21	2	-
Male	3	7	6	3	6	6	13	13	8	4	-	-
Female	2	1	2	6	5	18	24	40	42	17	2	-
Black, both sexes	-	3	-	5	-	2	2	2	2	1	-	-
Male	-	-	-	2	-	-	-	1	1	1	-	-
Female	-	3	-	3	-	2	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Delaware												
All races, both sexes	1	1	3	1	1	9	4	9	11	3	1	-
Male	-	-	2	1	-	4	2	1	1	1	1	-
Female	1	1	1	-	1	5	2	8	10	2	-	-
White, both sexes	-	1	2	-	1	7	3	9	9	3	1	-
Male	-	-	1	-	-	3	1	1	1	1	1	-
Female	-	1	1	-	1	4	2	8	8	2	-	-
Black, both sexes	1	-	1	1	-	2	1	-	2	-	-	-
Male	-	-	1	1	-	1	1	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	1	2	6	4	5	-	1	-
Male	1	-	-	-	-	2	4	3	3	-	-	-
Female	-	-	-	-	1	-	2	1	2	-	1	-
White, both sexes	1	-	-	-	-	-	3	-	1	-	-	-
Male	1	-	-	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	1	1	3	4	3	-	1	-
Male	-	-	-	-	-	1	2	3	2	-	-	-
Female	-	-	-	-	1	-	1	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Florida												
All races, both sexes	25	35	35	46	76	121	179	185	174	76	16	-
Male	18	24	20	20	42	51	66	53	33	13	3	-
Female	7	11	15	26	34	70	113	132	141	63	13	-
White, both sexes	13	24	27	32	60	100	155	163	156	69	13	-
Male	8	18	16	16	29	43	56	44	31	10	1	-
Female	5	6	11	16	31	57	99	119	125	59	12	-
Black, both sexes	11	11	8	13	16	21	22	22	17	7	3	-
Male	9	6	4	3	13	8	9	9	2	3	2	-
Female	2	5	4	10	3	13	13	13	15	4	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Georgia												
All races, both sexes	28	35	45	52	43	63	86	91	79	29	11	-
Male	23	22	28	24	26	29	32	12	11	3	2	-
Female	5	13	17	28	17	34	54	79	68	26	9	-
White, both sexes	18	19	27	31	20	41	63	68	62	20	5	-
Male	16	11	14	15	12	19	20	11	8	2	1	-
Female	2	8	13	16	8	22	43	57	54	18	4	-
Black, both sexes	10	16	18	21	23	22	23	23	17	9	6	-
Male	7	11	14	9	14	10	12	1	3	1	1	-
Female	3	5	4	12	9	12	11	22	14	8	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Hawaii												
All races, both sexes	4	2	1	3	2	9	7	8	20	4	2	-
Male	3	2	-	2	-	7	2	3	7	1	-	-
Female	1	-	1	1	2	2	5	5	13	3	2	-
White, both sexes	1	-	-	-	1	-	3	-	3	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	3	-	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	1	3	1	9	4	8	17	3	2	-
Male	2	2	-	2	-	7	2	3	7	1	-	-
Female	1	-	1	1	1	2	2	5	10	2	2	-
Idaho												
All races, both sexes	1	2	-	4	1	4	5	6	6	5	-	-
Male	-	1	-	3	1	1	1	-	2	1	-	-
Female	1	1	-	1	-	3	4	6	4	4	-	-
White, both sexes	1	2	-	4	1	4	5	6	6	5	-	-
Male	-	1	-	3	1	1	1	-	2	1	-	-
Female	1	1	-	1	-	3	4	6	4	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Illinois												
All races, both sexes	13	12	31	20	39	67	98	127	119	65	14	-
Male	11	8	23	14	21	32	31	31	23	5	1	-
Female	2	4	8	6	18	35	67	96	96	60	13	-
White, both sexes	8	5	23	18	25	52	82	119	109	61	13	-
Male	7	4	20	13	13	23	28	29	22	4	1	-
Female	1	1	3	5	12	29	54	90	87	57	12	-
Black, both sexes	5	7	8	2	13	15	15	7	8	3	1	-
Male	4	4	3	1	7	9	2	2	1	1	-	-
Female	1	3	5	1	6	6	13	5	7	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	1	2	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
Indiana												
All races, both sexes	8	7	17	17	15	50	71	88	53	36	12	-
Male	7	4	11	9	8	21	20	23	9	3	1	-
Female	1	3	6	8	7	29	51	65	44	33	11	-
White, both sexes	7	4	15	13	13	44	65	81	48	35	9	-
Male	6	3	10	7	7	17	16	20	8	3	1	-
Female	1	1	5	6	6	27	49	61	40	32	8	-
Black, both sexes	1	3	2	4	2	5	6	7	5	1	3	-
Male	1	1	1	2	1	4	4	3	1	-	-	-
Female	-	2	1	2	1	1	2	4	4	1	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Iowa												
All races, both sexes	1	1	3	4	7	15	23	20	24	25	4	-
Male	-	-	1	2	1	8	6	10	4	2	1	-
Female	1	1	2	2	6	7	17	10	20	23	3	-
White, both sexes	-	1	3	4	7	15	23	20	24	24	4	-
Male	-	-	1	2	1	8	6	10	4	2	1	-
Female	-	1	2	2	6	7	17	10	20	22	3	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	3	3	5	7	13	12	27	20	9	3	-
Male	2	2	2	1	3	6	2	6	3	3	1	-
Female	1	1	1	4	4	7	10	21	17	6	2	-
White, both sexes	1	2	1	5	7	13	12	26	19	9	2	-
Male	-	1	1	1	3	6	2	6	2	3	1	-
Female	1	1	-	4	4	7	10	20	17	6	1	-
Black, both sexes	2	1	2	-	-	-	-	1	1	-	1	-
Male	2	1	1	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Kentucky												
All races, both sexes	5	9	11	11	14	21	37	47	32	22	2	-
Male	5	5	7	4	6	7	10	8	7	1	-	-
Female	-	4	4	7	8	14	27	39	25	21	2	-
White, both sexes	4	7	9	10	12	20	33	44	29	17	2	-
Male	4	4	6	4	5	7	9	8	6	1	-	-
Female	-	3	3	6	7	13	24	36	23	16	2	-
Black, both sexes	1	1	2	1	2	1	4	3	3	5	-	-
Male	1	1	1	-	1	-	1	-	1	-	-	-
Female	-	-	1	1	1	1	3	3	2	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	5	5	9	6	16	32	42	38	22	16	5	-
Male	4	3	6	4	5	13	11	6	4	1	-	-
Female	1	2	3	2	11	19	31	32	18	15	5	-
White, both sexes	3	3	5	2	9	20	28	29	11	9	5	-
Male	3	2	3	1	3	6	8	4	3	-	-	-
Female	-	1	2	1	6	14	20	25	8	9	5	-
Black, both sexes	2	2	4	4	7	12	14	9	11	7	-	-
Male	1	1	3	3	2	7	3	2	1	1	-	-
Female	1	1	1	1	5	5	11	7	10	6	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Maine												
All races, both sexes	1	4	4	1	2	3	8	7	9	1	-	-
Male	1	3	4	1	-	-	2	1	3	-	-	-
Female	-	1	-	-	2	3	6	6	6	1	-	-
White, both sexes	1	4	4	1	2	3	8	6	9	1	-	-
Male	1	3	4	1	-	-	2	1	3	-	-	-
Female	-	1	-	-	2	3	6	5	6	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5	7	9	12	11	19	39	50	45	24	4	-
Male	4	5	4	7	6	6	11	18	6	6	-	-
Female	1	2	5	5	5	13	28	32	39	18	4	-
White, both sexes	2	3	3	5	8	11	26	43	37	15	3	-
Male	1	2	1	5	5	5	8	14	5	3	-	-
Female	1	1	2	-	3	6	18	29	32	12	3	-
Black, both sexes	2	4	5	7	3	8	13	6	7	8	1	-
Male	2	3	3	2	1	1	3	3	1	2	-	-
Female	-	1	2	5	2	7	10	3	6	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	1	1	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Massachusetts												
All races, both sexes	9	10	11	12	14	37	44	76	56	22	6	-
Male	6	8	9	7	5	20	17	21	10	3	1	-
Female	3	2	2	5	9	17	27	55	46	19	5	-
White, both sexes	9	8	11	10	14	33	39	72	55	22	6	-
Male	6	7	9	7	5	16	15	20	10	3	1	-
Female	3	1	2	3	9	17	24	52	45	19	5	-
Black, both sexes	-	2	-	2	-	3	5	4	-	-	-	-
Male	-	1	-	-	-	3	2	1	-	-	-	-
Female	-	1	-	2	-	-	3	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Michigan												
All races, both sexes	8	9	14	25	29	48	72	82	90	60	7	-
Male	3	6	9	6	14	19	24	23	13	6	1	-
Female	5	3	5	19	15	29	48	59	77	54	6	-
White, both sexes	5	7	9	20	24	42	66	79	80	49	7	-
Male	2	5	5	4	11	16	22	22	12	4	1	-
Female	3	2	4	16	13	26	44	57	68	45	6	-
Black, both sexes	3	2	5	5	5	6	6	3	10	10	-	-
Male	1	1	4	2	3	3	2	1	1	2	-	-
Female	2	1	1	3	2	3	4	2	9	8	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Minnesota												
All races, both sexes	7	9	24	25	28	38	46	67	81	38	9	-
Male	5	5	15	13	17	18	20	10	8	4	1	-
Female	2	4	9	12	11	20	26	57	73	34	8	-
White, both sexes	7	9	21	24	27	33	46	66	80	38	8	-
Male	5	5	13	12	16	16	20	10	8	4	1	-
Female	2	4	8	12	11	17	26	56	72	34	7	-
Black, both sexes	-	-	3	-	1	3	-	1	1	-	1	-
Male	-	-	2	-	1	2	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	7	11	11	13	19	24	31	49	38	19	7	-
Male	6	4	7	8	7	4	11	9	8	6	1	-
Female	1	7	4	5	12	20	20	40	30	13	6	-
White, both sexes	3	6	3	5	7	9	19	38	28	11	1	-
Male	2	4	1	3	3	2	7	8	5	1	-	-
Female	1	2	2	2	4	7	12	30	23	10	1	-
Black, both sexes	4	5	8	8	12	14	11	11	10	7	6	-
Male	4	-	6	5	4	1	3	1	3	5	1	-
Female	-	5	2	3	8	13	8	10	7	2	5	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Missouri												
All races, both sexes	3	14	18	14	24	25	43	49	47	25	3	-
Male	2	3	13	9	10	9	17	15	8	4	-	-
Female	1	11	5	5	14	16	26	34	39	21	3	-
White, both sexes	1	12	11	7	20	23	39	46	45	19	3	-
Male	1	3	11	5	9	8	16	14	8	2	-	-
Female	-	9	-	2	11	15	23	32	37	17	3	-
Black, both sexes	2	2	7	7	4	2	4	3	2	6	-	-
Male	1	-	2	4	1	1	1	1	-	2	-	-
Female	1	2	5	3	3	1	3	2	2	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	2	-	1	3	7	7	5	5	3	-
Male	-	-	1	-	1	1	1	-	-	-	-	-
Female	1	-	1	-	-	2	6	7	5	5	3	-
White, both sexes	1	-	1	-	1	3	6	7	5	5	3	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	1	-	1	-	-	2	5	7	5	5	3	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Nebraska												
All races, both sexes	-	3	6	8	12	10	16	22	34	12	4	-
Male	-	2	5	5	9	2	4	4	7	2	-	-
Female	-	1	1	3	3	8	12	18	27	10	4	-
White, both sexes	-	3	5	8	12	9	15	21	33	12	4	-
Male	-	2	4	5	9	1	3	4	6	2	-	-
Female	-	1	1	3	3	8	12	17	27	10	4	-
Black, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	3	4	3	7	14	17	8	14	4	-	-
Male	1	2	1	1	2	6	9	2	3	1	-	-
Female	1	1	3	2	5	8	8	6	11	3	-	-
White, both sexes	2	1	4	2	6	14	16	8	14	4	-	-
Male	1	1	1	1	1	6	8	2	3	1	-	-
Female	1	-	3	1	5	8	8	6	11	3	-	-
Black, both sexes	-	2	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
New Hampshire												
All races, both sexes	1	6	2	3	4	3	6	11	8	4	2	-
Male	1	2	2	1	3	2	3	3	2	-	-	-
Female	-	4	-	2	1	1	3	8	6	4	2	-
White, both sexes	1	6	2	3	4	3	6	11	8	4	2	-
Male	1	2	2	1	3	2	3	3	2	-	-	-
Female	-	4	-	2	1	1	3	8	6	4	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	8	20	17	16	19	45	54	64	60	50	8	-
Male	5	10	10	11	10	20	14	18	7	8	-	-
Female	3	10	7	5	9	25	40	46	53	42	8	-
White, both sexes	4	10	10	12	11	30	50	60	56	46	6	-
Male	4	6	7	7	5	12	13	17	6	8	-	-
Female	-	4	3	5	6	18	37	43	50	38	6	-
Black, both sexes	3	8	7	4	8	14	4	3	4	4	2	-
Male	-	4	3	4	5	8	1	-	1	-	-	-
Female	3	4	4	-	3	6	3	3	3	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	2	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
New Mexico												
All races, both sexes	3	-	-	4	6	8	11	13	13	4	-	-
Male	-	-	-	2	4	4	4	4	-	2	-	-
Female	3	-	-	2	2	4	7	9	13	2	-	-
White, both sexes	3	-	-	4	4	7	11	13	13	4	-	-
Male	-	-	-	2	3	3	4	4	-	2	-	-
Female	3	-	-	2	1	4	7	9	13	2	-	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	23	36	38	55	99	120	165	205	182	90	21	-
Male	16	23	18	35	38	53	62	46	44	16	5	-
Female	7	13	20	20	61	67	103	159	138	74	16	-
White, both sexes	15	21	28	37	79	91	132	166	149	81	16	-
Male	10	14	14	24	29	39	49	40	37	16	4	-
Female	5	7	14	13	50	52	83	126	112	65	12	-
Black, both sexes	8	11	10	18	18	24	29	38	28	8	5	-
Male	6	8	4	11	7	12	12	6	5	-	1	-
Female	2	3	6	7	11	12	17	32	23	8	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	4	-	-	2	5	4	1	5	1	-	-
Male	-	1	-	-	2	2	1	-	2	-	-	-
Female	-	3	-	-	-	3	3	1	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
North Carolina												
All races, both sexes	8	9	17	23	33	51	74	55	77	32	10	-
Male	7	7	10	15	15	21	21	12	15	2	1	-
Female	1	2	7	8	18	30	53	43	62	30	9	-
White, both sexes	2	4	13	12	18	37	57	40	59	22	8	-
Male	2	3	8	7	7	17	16	10	12	2	1	-
Female	-	1	5	5	11	20	41	30	47	20	7	-
Black, both sexes	3	5	4	10	14	14	17	15	18	10	2	-
Male	3	4	2	7	7	4	5	2	3	-	-	-
Female	-	1	2	3	7	10	12	13	15	10	2	-
American Indian 1/, both sexes	3	-	-	1	1	-	-	-	-	-	-	-
Male	2	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	1	2	1	3	1	3	5	5	4	1	-
Male	1	-	1	1	3	1	-	1	3	-	-	-
Female	-	1	1	-	-	-	3	4	2	4	1	-
White, both sexes	1	1	2	1	3	1	3	5	5	4	1	-
Male	1	-	1	1	3	1	-	1	3	-	-	-
Female	-	1	1	-	-	-	3	4	2	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Ohio												
All races, both sexes	13	20	24	28	33	74	115	140	131	57	15	-
Male	9	10	12	15	19	30	40	39	21	5	4	-
Female	4	10	12	13	14	44	75	101	110	52	11	-
White, both sexes	8	14	21	24	29	61	104	125	123	52	14	-
Male	6	7	9	11	16	23	35	35	19	5	3	-
Female	2	7	12	13	13	38	69	90	104	47	11	-
Black, both sexes	5	5	3	4	3	13	11	15	8	5	1	-
Male	3	3	3	4	3	7	5	4	2	-	1	-
Female	2	2	-	-	-	6	6	11	6	5	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	7	9	6	5	16	20	31	40	39	10	6	-
Male	5	5	3	4	8	5	14	10	6	2	-	-
Female	2	4	3	1	8	15	17	30	33	8	6	-
White, both sexes	4	6	6	4	13	15	25	35	37	10	5	-
Male	3	4	3	4	7	4	10	8	5	2	-	-
Female	1	2	3	-	6	11	15	27	32	8	5	-
Black, both sexes	1	3	-	1	2	2	2	1	1	-	1	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	1	2	-	1	2	1	2	-	1	-	1	-
American Indian 1/, both sexes	2	-	-	-	1	3	4	4	1	-	-	-
Male	2	-	-	-	1	-	4	1	1	-	-	-
Female	-	-	-	-	-	3	-	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Oregon												
All races, both sexes	10	10	14	11	12	31	57	57	65	34	6	-
Male	6	8	9	6	6	12	27	21	7	5	-	-
Female	4	2	5	5	6	19	30	36	58	29	6	-
White, both sexes	8	9	13	11	10	30	55	54	63	34	6	-
Male	5	7	9	6	5	11	25	21	7	5	-	-
Female	3	2	4	5	5	19	30	33	56	29	6	-
Black, both sexes	1	1	-	-	2	1	2	-	1	-	-	-
Male	-	1	-	-	1	1	2	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	2	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
Pennsylvania												
All races, both sexes	16	32	30	41	58	68	118	128	123	61	13	-
Male	11	23	19	30	30	31	37	31	27	6	1	-
Female	5	9	11	11	28	37	81	97	96	55	12	-
White, both sexes	11	21	26	33	53	59	106	122	118	56	8	-
Male	8	15	16	25	27	25	32	29	26	5	1	-
Female	3	6	10	8	26	34	74	93	92	51	7	-
Black, both sexes	5	11	4	8	5	9	11	6	5	5	5	-
Male	3	8	3	5	3	6	4	2	1	1	-	-
Female	2	3	1	3	2	3	7	4	4	4	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Rhode Island												
All races, both sexes	1	-	-	1	1	3	10	6	3	3	-	-
Male	-	-	-	1	-	-	3	2	-	-	-	-
Female	1	-	-	-	1	3	7	4	3	3	-	-
White, both sexes	1	-	-	1	1	3	10	5	3	3	-	-
Male	-	-	-	1	-	-	3	2	-	-	-	-
Female	1	-	-	-	1	3	7	3	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	14	13	16	15	28	30	34	32	19	13	2	-
Male	9	10	10	7	11	12	13	10	1	1	-	-
Female	5	3	6	8	17	18	21	22	18	12	2	-
White, both sexes	4	2	9	11	16	20	27	23	11	9	1	-
Male	3	1	5	5	5	7	10	6	-	-	-	-
Female	1	1	4	6	11	13	17	17	11	9	1	-
Black, both sexes	10	11	7	4	12	10	6	9	8	4	1	-
Male	6	9	5	2	6	5	3	4	1	1	-	-
Female	4	2	2	2	6	5	3	5	7	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
South Dakota												
All races, both sexes	-	1	2	2	1	-	3	9	10	6	2	-
Male	-	1	1	2	1	-	1	5	2	2	-	-
Female	-	-	1	-	-	-	2	4	8	4	2	-
White, both sexes	-	1	1	2	1	-	3	9	9	6	2	-
Male	-	1	-	2	1	-	1	5	2	2	-	-
Female	-	-	1	-	-	-	2	4	7	4	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	10	7	8	14	24	22	56	56	43	20	7	-
Male	5	5	6	7	12	11	16	12	10	1	-	-
Female	5	2	2	7	12	11	40	44	33	19	7	-
White, both sexes	7	6	6	10	17	15	47	46	33	12	6	-
Male	5	4	5	5	8	7	13	10	9	1	-	-
Female	2	2	1	5	9	8	34	36	24	11	6	-
Black, both sexes	3	1	2	4	7	7	9	10	10	8	1	-
Male	-	1	1	2	4	4	3	2	1	-	-	-
Female	3	-	1	2	3	3	6	8	9	8	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Texas												
All races, both sexes	21	28	36	40	54	89	131	134	133	78	15	-
Male	10	15	24	25	23	35	46	34	31	11	-	-
Female	11	13	12	15	31	54	85	100	102	67	15	-
White, both sexes	12	21	26	32	46	77	112	129	116	70	12	-
Male	6	13	19	22	19	30	41	33	27	9	-	-
Female	6	8	7	10	27	47	71	96	89	61	12	-
Black, both sexes	9	7	10	8	8	12	17	4	16	6	2	-
Male	4	2	5	3	4	5	5	1	3	2	-	-
Female	5	5	5	5	4	7	12	3	13	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	1	1	2	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	2	1	-	2	-	-
Utah												
All races, both sexes	2	2	3	3	5	4	19	15	12	10	-	-
Male	1	2	1	3	2	1	4	4	1	1	-	-
Female	1	-	2	-	3	3	15	11	11	9	-	-
White, both sexes	2	2	3	2	5	4	18	15	12	10	-	-
Male	1	2	1	2	2	1	4	4	1	1	-	-
Female	1	-	2	-	3	3	14	11	11	9	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Vermont												
All races, both sexes	-	-	-	2	1	3	2	5	6	3	1	-
Male	-	-	-	2	1	-	1	-	4	-	-	-
Female	-	-	-	-	-	3	1	5	2	3	1	-
White, both sexes	-	-	-	2	1	3	2	5	6	3	1	-
Male	-	-	-	2	1	-	1	-	4	-	-	-
Female	-	-	-	-	-	3	1	5	2	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	7	25	15	28	21	38	54	46	49	16	10	-
Male	3	16	9	15	3	18	12	13	6	2	2	-
Female	4	9	6	13	18	20	42	33	43	14	8	-
White, both sexes	4	18	11	13	14	23	39	42	34	13	4	-
Male	3	11	5	7	2	11	7	12	4	2	-	-
Female	1	7	6	6	12	12	32	30	30	11	4	-
Black, both sexes	3	6	4	15	6	14	14	4	15	3	5	-
Male	-	4	4	8	1	7	4	1	2	-	1	-
Female	3	2	-	7	5	7	10	3	13	3	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	1	-	-	-	1	-
Male	-	1	-	-	-	-	1	-	-	-	1	-
Female	-	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Washington												
All races, both sexes	9	8	7	14	14	32	51	45	53	33	5	-
Male	7	5	1	5	6	16	18	10	7	2	2	-
Female	2	3	6	9	8	16	33	35	46	31	3	-
White, both sexes	7	6	7	14	10	27	48	42	52	32	4	-
Male	6	5	1	5	4	14	16	9	7	1	1	-
Female	1	1	6	9	6	13	32	33	45	31	3	-
Black, both sexes	2	2	-	-	2	2	1	3	-	-	-	-
Male	1	-	-	-	1	1	-	1	-	-	-	-
Female	1	2	-	-	1	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	2	2	-	1	1	1	-
Male	-	-	-	-	1	-	2	-	-	1	1	-
Female	-	-	-	-	1	2	-	-	1	-	-	-
West Virginia												
All races, both sexes	3	10	10	15	10	23	26	33	25	13	4	-
Male	2	7	6	5	6	10	6	6	2	1	-	-
Female	1	3	4	10	4	13	20	27	23	12	4	-
White, both sexes	2	10	7	14	10	21	25	33	24	13	4	-
Male	2	7	4	5	6	9	6	6	2	1	-	-
Female	-	3	3	9	4	12	19	27	22	12	4	-
Black, both sexes	1	-	3	1	-	2	1	-	1	-	-	-
Male	-	-	2	-	-	1	-	-	-	-	-	-
Female	1	-	1	1	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Essential (primary) hypertension (I10)												
Wisconsin												
All races, both sexes	3	8	8	11	13	20	44	54	57	32	10	-
Male	3	5	8	6	7	6	11	9	11	2	1	-
Female	-	3	-	5	6	14	33	45	46	30	9	-
White, both sexes	2	7	8	11	12	20	40	54	55	31	9	-
Male	2	4	8	6	7	6	9	9	10	2	1	-
Female	-	3	-	5	5	14	31	45	45	29	8	-
Black, both sexes	1	1	-	-	1	-	3	-	2	1	-	-
Male	1	1	-	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	1	-
Wyoming												
All races, both sexes	1	2	-	1	2	-	1	1	2	1	-	-
Male	-	1	-	1	-	-	1	-	1	-	-	-
Female	1	1	-	-	2	-	-	1	1	1	-	-
White, both sexes	1	2	-	1	2	-	1	1	2	1	-	-
Male	-	1	-	1	-	-	1	-	1	-	-	-
Female	1	1	-	-	2	-	-	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
United States												
All races, both sexes	29,282	2	-	2	1	-	36	113	215	462	906	1,459
Male	13,219	1	-	2	-	-	24	77	145	338	620	1,010
Female	16,063	1	-	-	1	-	12	36	70	124	286	449
White, both sexes	21,686	1	-	2	-	-	12	59	112	263	511	847
Male	9,408	1	-	2	-	-	8	47	83	205	372	633
Female	12,278	-	-	-	-	-	4	12	29	58	139	214
Black, both sexes	6,923	1	-	-	1	-	23	47	93	186	359	577
Male	3,459	-	-	-	-	-	15	24	54	121	224	350
Female	3,464	1	-	-	1	-	8	23	39	65	135	227
American Indian 1/, both sexes	127	-	-	-	-	-	-	5	3	3	10	9
Male	68	-	-	-	-	-	-	4	1	2	6	7
Female	59	-	-	-	-	-	-	1	2	1	4	2
Asian or Pacific Islander 2/, both sexes	546	-	-	-	-	-	1	2	7	10	26	26
Male	284	-	-	-	-	-	1	2	7	10	18	20
Female	262	-	-	-	-	-	-	-	-	-	8	6
Alabama												
All races, both sexes	461	-	-	-	-	-	1	2	1	4	11	21
Male	202	-	-	-	-	-	-	1	-	3	9	17
Female	259	-	-	-	-	-	1	1	1	1	2	4
White, both sexes	279	-	-	-	-	-	-	-	-	1	6	7
Male	112	-	-	-	-	-	-	-	-	1	4	6
Female	167	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	182	-	-	-	-	-	1	2	1	3	5	14
Male	90	-	-	-	-	-	-	1	-	2	5	11
Female	92	-	-	-	-	-	1	1	1	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Alaska												
All races, both sexes	47	-	-	-	-	-	-	1	-	1	2	4
Male	31	-	-	-	-	-	-	-	-	1	1	4
Female	16	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	35	-	-	-	-	-	-	1	-	1	-	3
Male	24	-	-	-	-	-	-	-	-	1	-	3
Female	11	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Arizona												
All races, both sexes	493	-	-	-	-	-	1	3	3	5	18	21
Male	229	-	-	-	-	-	1	2	2	4	10	17
Female	264	-	-	-	-	-	-	1	1	1	8	4
White, both sexes	447	-	-	-	-	-	1	1	2	4	13	17
Male	207	-	-	-	-	-	1	1	2	3	6	15
Female	240	-	-	-	-	-	-	-	-	1	7	2
Black, both sexes	22	-	-	-	-	-	-	1	-	-	2	3
Male	12	-	-	-	-	-	-	-	-	-	2	1
Female	10	-	-	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	18	-	-	-	-	-	-	1	1	-	2	-
Male	6	-	-	-	-	-	-	1	-	-	1	-
Female	12	-	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	1	1	1
Male	4	-	-	-	-	-	-	-	-	1	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Arkansas												
All races, both sexes	246	-	-	-	-	-	-	2	3	2	10	15
Male	103	-	-	-	-	-	-	1	3	2	7	10
Female	143	-	-	-	-	-	-	1	-	-	3	5
White, both sexes	191	-	-	-	-	-	-	-	2	1	6	6
Male	73	-	-	-	-	-	-	-	2	1	4	6
Female	118	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	53	-	-	-	-	-	-	2	1	1	4	8
Male	29	-	-	-	-	-	-	1	1	1	3	3
Female	24	-	-	-	-	-	-	1	-	-	1	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3,865	-	-	-	-	-	2	14	21	57	105	193
Male	1,723	-	-	-	-	-	2	8	13	45	75	139
Female	2,142	-	-	-	-	-	-	6	8	12	30	54
White, both sexes	3,039	-	-	-	-	-	1	11	13	39	72	128
Male	1,318	-	-	-	-	-	1	6	8	32	53	96
Female	1,721	-	-	-	-	-	-	5	5	7	19	32
Black, both sexes	528	-	-	-	-	-	1	1	6	16	21	51
Male	261	-	-	-	-	-	1	-	3	11	15	34
Female	267	-	-	-	-	-	-	1	3	5	6	17
American Indian 1/, both sexes	21	-	-	-	-	-	-	1	-	-	1	2
Male	11	-	-	-	-	-	-	1	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	277	-	-	-	-	-	-	1	2	2	11	12
Male	133	-	-	-	-	-	-	1	2	2	7	8
Female	144	-	-	-	-	-	-	-	-	-	4	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Colorado												
All races, both sexes	249	-	-	-	-	-	-	1	1	3	6	10
Male	105	-	-	-	-	-	-	1	-	1	6	5
Female	144	-	-	-	-	-	-	-	1	2	-	5
White, both sexes	226	-	-	-	-	-	-	-	1	3	6	9
Male	92	-	-	-	-	-	-	-	-	1	6	5
Female	134	-	-	-	-	-	-	-	1	2	-	4
Black, both sexes	20	-	-	-	-	-	-	1	-	-	-	1
Male	11	-	-	-	-	-	-	1	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	190	1	-	-	-	-	1	1	1	2	5	3
Male	86	-	-	-	-	-	-	1	1	1	4	3
Female	104	1	-	-	-	-	1	-	-	1	1	-
White, both sexes	171	-	-	-	-	-	-	1	1	2	5	2
Male	77	-	-	-	-	-	-	1	1	1	4	2
Female	94	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	17	1	-	-	-	-	1	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	9	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Delaware												
All races, both sexes	86	-	-	-	-	-	-	1	-	2	4	1
Male	34	-	-	-	-	-	-	-	-	1	2	1
Female	52	-	-	-	-	-	-	1	-	1	2	-
White, both sexes	64	-	-	-	-	-	-	-	-	1	1	1
Male	24	-	-	-	-	-	-	-	-	1	-	1
Female	40	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	22	-	-	-	-	-	-	1	-	1	3	-
Male	10	-	-	-	-	-	-	-	-	-	2	-
Female	12	-	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	198	-	-	-	-	-	1	2	-	1	9	8
Male	89	-	-	-	-	-	-	2	-	-	5	5
Female	109	-	-	-	-	-	1	-	-	1	4	3
White, both sexes	23	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	172	-	-	-	-	-	1	2	-	1	8	7
Male	78	-	-	-	-	-	-	2	-	-	4	5
Female	94	-	-	-	-	-	1	-	-	1	4	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Florida												
All races, both sexes	2,330	-	-	-	-	-	3	10	13	42	77	103
Male	1,139	-	-	-	-	-	3	7	8	29	58	67
Female	1,191	-	-	-	-	-	-	3	5	13	19	36
White, both sexes	1,872	-	-	-	-	-	1	6	6	32	49	63
Male	907	-	-	-	-	-	1	5	5	23	40	45
Female	965	-	-	-	-	-	-	1	1	9	9	18
Black, both sexes	445	-	-	-	-	-	2	4	7	9	27	39
Male	227	-	-	-	-	-	2	2	3	5	18	21
Female	218	-	-	-	-	-	-	2	4	4	9	18
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1,055	-	-	-	1	-	1	-	12	26	37	68
Male	542	-	-	-	-	-	1	-	8	18	26	47
Female	513	-	-	-	1	-	-	-	4	8	11	21
White, both sexes	535	-	-	-	-	-	-	-	4	12	10	30
Male	267	-	-	-	-	-	-	-	2	6	7	24
Female	268	-	-	-	-	-	-	-	2	6	3	6
Black, both sexes	510	-	-	-	1	-	1	-	8	14	27	38
Male	269	-	-	-	-	-	1	-	6	12	19	23
Female	241	-	-	-	1	-	-	-	2	2	8	15
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Hawaii												
All races, both sexes	46	-	-	-	-	-	-	-	1	2	2	-
Male	22	-	-	-	-	-	-	-	1	2	1	-
Female	24	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	38	-	-	-	-	-	-	-	1	2	2	-
Male	17	-	-	-	-	-	-	-	1	2	1	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
Idaho												
All races, both sexes	93	-	-	-	-	-	-	-	-	2	2	1
Male	33	-	-	-	-	-	-	-	-	1	1	1
Female	60	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	92	-	-	-	-	-	-	-	-	2	2	1
Male	33	-	-	-	-	-	-	-	-	1	1	1
Female	59	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Illinois												
All races, both sexes	1,469	-	-	-	-	-	4	9	16	32	61	81
Male	739	-	-	-	-	-	3	5	12	23	44	62
Female	730	-	-	-	-	-	1	4	4	9	17	19
White, both sexes	949	-	-	-	-	-	-	2	9	13	22	41
Male	446	-	-	-	-	-	-	2	8	12	17	35
Female	503	-	-	-	-	-	-	-	1	1	5	6
Black, both sexes	499	-	-	-	-	-	3	7	6	19	38	39
Male	281	-	-	-	-	-	2	3	3	11	26	27
Female	218	-	-	-	-	-	1	4	3	8	12	12
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	1	-	1	-	1	1
Male	12	-	-	-	-	-	1	-	1	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Indiana												
All races, both sexes	438	-	-	-	-	-	1	3	1	8	11	20
Male	176	-	-	-	-	-	-	2	1	6	8	14
Female	262	-	-	-	-	-	1	1	-	2	3	6
White, both sexes	367	-	-	-	-	-	-	1	1	3	8	12
Male	144	-	-	-	-	-	-	1	1	3	7	11
Female	223	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	68	-	-	-	-	-	1	2	-	5	3	8
Male	31	-	-	-	-	-	-	1	-	3	1	3
Female	37	-	-	-	-	-	1	1	-	2	2	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Iowa												
All races, both sexes	278	-	-	-	-	-	-	-	-	3	5	5
Male	104	-	-	-	-	-	-	-	-	3	4	4
Female	174	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	274	-	-	-	-	-	-	-	-	3	4	5
Male	100	-	-	-	-	-	-	-	-	3	3	4
Female	174	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	139	-	-	-	-	-	1	1	1	1	1	5
Male	52	-	-	-	-	-	1	1	1	1	1	5
Female	87	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	121	-	-	-	-	-	-	1	-	-	-	3
Male	39	-	-	-	-	-	-	1	-	-	-	3
Female	82	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	1	-	1	1	1	2
Male	12	-	-	-	-	-	1	-	1	1	1	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Kentucky												
All races, both sexes	378	-	-	-	-	-	2	4	2	13	19	31
Male	176	-	-	-	-	-	2	2	1	10	13	23
Female	202	-	-	-	-	-	-	2	1	3	6	8
White, both sexes	322	-	-	-	-	-	1	3	2	11	15	24
Male	156	-	-	-	-	-	1	1	1	9	12	19
Female	166	-	-	-	-	-	-	2	1	2	3	5
Black, both sexes	56	-	-	-	-	-	1	1	-	2	4	7
Male	20	-	-	-	-	-	1	1	-	1	1	4
Female	36	-	-	-	-	-	-	-	-	1	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	686	-	-	-	-	-	1	2	5	11	19	53
Male	312	-	-	-	-	-	-	-	3	8	12	28
Female	374	-	-	-	-	-	1	2	2	3	7	25
White, both sexes	387	-	-	-	-	-	1	-	1	5	5	21
Male	165	-	-	-	-	-	-	-	-	4	3	15
Female	222	-	-	-	-	-	1	-	1	1	2	6
Black, both sexes	297	-	-	-	-	-	-	2	4	6	14	31
Male	146	-	-	-	-	-	-	-	3	4	9	12
Female	151	-	-	-	-	-	-	2	1	2	5	19
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Maine												
All races, both sexes	56	-	-	-	-	-	-	-	1	-	2	-
Male	28	-	-	-	-	-	-	-	1	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	56	-	-	-	-	-	-	-	1	-	2	-
Male	28	-	-	-	-	-	-	-	1	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	799	-	-	-	-	-	-	2	8	28	42	63
Male	400	-	-	-	-	-	-	2	3	24	28	45
Female	399	-	-	-	-	-	-	-	5	4	14	18
White, both sexes	463	-	-	-	-	-	-	-	4	12	20	37
Male	220	-	-	-	-	-	-	-	2	12	12	27
Female	243	-	-	-	-	-	-	-	2	-	8	10
Black, both sexes	327	-	-	-	-	-	-	2	4	16	22	26
Male	176	-	-	-	-	-	-	2	1	12	16	18
Female	151	-	-	-	-	-	-	-	3	4	6	8
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Massachusetts												
All races, both sexes	352	-	-	-	-	-	2	5	7	9	20	30
Male	174	-	-	-	-	-	2	5	5	5	16	26
Female	178	-	-	-	-	-	-	-	2	4	4	4
White, both sexes	310	-	-	-	-	-	2	4	6	7	15	23
Male	151	-	-	-	-	-	2	4	4	4	13	19
Female	159	-	-	-	-	-	-	-	2	3	2	4
Black, both sexes	38	-	-	-	-	-	-	1	1	2	4	6
Male	20	-	-	-	-	-	-	1	1	1	2	6
Female	18	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,288	-	-	-	-	-	1	1	3	16	32	66
Male	541	-	-	-	-	-	-	1	3	13	20	50
Female	747	-	-	-	-	-	1	-	-	3	12	16
White, both sexes	953	-	-	-	-	-	1	-	2	7	14	32
Male	374	-	-	-	-	-	-	-	2	6	10	28
Female	579	-	-	-	-	-	1	-	-	1	4	4
Black, both sexes	331	-	-	-	-	-	-	1	1	9	18	32
Male	165	-	-	-	-	-	-	1	1	7	10	20
Female	166	-	-	-	-	-	-	-	-	2	8	12
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Minnesota												
All races, both sexes	172	-	-	1	-	-	-	-	1	-	3	7
Male	66	-	-	1	-	-	-	-	1	-	3	4
Female	106	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	164	-	-	1	-	-	-	-	-	-	2	7
Male	60	-	-	1	-	-	-	-	-	-	2	4
Female	104	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	675	-	-	-	-	-	1	4	7	8	12	30
Male	285	-	-	-	-	-	-	3	4	2	6	21
Female	390	-	-	-	-	-	1	1	3	6	6	9
White, both sexes	364	-	-	-	-	-	-	1	2	3	5	8
Male	154	-	-	-	-	-	-	1	2	2	4	6
Female	210	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	309	-	-	-	-	-	1	2	5	5	7	22
Male	129	-	-	-	-	-	-	1	2	-	2	15
Female	180	-	-	-	-	-	1	1	3	5	5	7
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Missouri												
All races, both sexes	456	-	-	-	-	-	-	6	11	8	26	39
Male	216	-	-	-	-	-	-	4	6	5	18	26
Female	240	-	-	-	-	-	-	2	5	3	8	13
White, both sexes	311	-	-	-	-	-	-	4	4	3	14	23
Male	140	-	-	-	-	-	-	4	2	3	12	16
Female	171	-	-	-	-	-	-	-	2	-	2	7
Black, both sexes	142	-	-	-	-	-	-	2	7	5	12	16
Male	73	-	-	-	-	-	-	-	4	2	6	10
Female	69	-	-	-	-	-	-	2	3	3	6	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	51	-	-	-	-	-	-	2	-	-	1	2
Male	19	-	-	-	-	-	-	2	-	-	-	2
Female	32	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	49	-	-	-	-	-	-	1	-	-	1	2
Male	18	-	-	-	-	-	-	1	-	-	-	2
Female	31	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Nebraska												
All races, both sexes	89	-	-	-	-	-	-	1	-	2	5	3
Male	40	-	-	-	-	-	-	1	-	2	3	3
Female	49	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	79	-	-	-	-	-	-	1	-	1	4	3
Male	34	-	-	-	-	-	-	1	-	1	2	3
Female	45	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	1	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	504	-	-	-	-	-	-	1	5	7	13	22
Male	305	-	-	-	-	-	-	1	3	5	11	15
Female	199	-	-	-	-	-	-	-	2	2	2	7
White, both sexes	432	-	-	-	-	-	-	1	2	5	11	12
Male	261	-	-	-	-	-	-	1	2	4	10	10
Female	171	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	52	-	-	-	-	-	-	-	2	2	2	9
Male	29	-	-	-	-	-	-	-	1	1	1	4
Female	23	-	-	-	-	-	-	-	1	1	1	5
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
New Hampshire												
All races, both sexes	83	-	-	-	-	-	-	-	1	3	-	4
Male	37	-	-	-	-	-	-	-	1	2	-	4
Female	46	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	83	-	-	-	-	-	-	-	1	3	-	4
Male	37	-	-	-	-	-	-	-	1	2	-	4
Female	46	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	791	-	-	-	-	-	-	-	4	10	15	50
Male	362	-	-	-	-	-	-	-	4	8	9	33
Female	429	-	-	-	-	-	-	-	-	2	6	17
White, both sexes	604	-	-	-	-	-	-	-	2	5	7	21
Male	263	-	-	-	-	-	-	-	2	4	4	14
Female	341	-	-	-	-	-	-	-	-	1	3	7
Black, both sexes	170	-	-	-	-	-	-	-	2	5	8	26
Male	90	-	-	-	-	-	-	-	2	4	5	17
Female	80	-	-	-	-	-	-	-	-	1	3	9
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
New Mexico												
All races, both sexes	176	-	-	-	-	-	-	-	1	1	11	9
Male	99	-	-	-	-	-	-	-	1	1	7	8
Female	77	-	-	-	-	-	-	-	-	-	4	1
White, both sexes	156	-	-	-	-	-	-	-	1	1	8	9
Male	85	-	-	-	-	-	-	-	1	1	5	8
Female	71	-	-	-	-	-	-	-	-	-	3	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	2	-
Male	7	-	-	-	-	-	-	-	-	-	2	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2,368	-	-	-	-	-	3	8	17	25	69	95
Male	1,013	-	-	-	-	-	3	5	15	19	47	63
Female	1,355	-	-	-	-	-	-	3	2	6	22	32
White, both sexes	1,543	-	-	-	-	-	1	3	7	11	27	39
Male	623	-	-	-	-	-	1	2	7	8	19	30
Female	920	-	-	-	-	-	-	1	-	3	8	9
Black, both sexes	753	-	-	-	-	-	2	5	8	13	37	51
Male	350	-	-	-	-	-	2	3	6	10	24	28
Female	403	-	-	-	-	-	-	2	2	3	13	23
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	69	-	-	-	-	-	-	-	2	1	5	5
Male	38	-	-	-	-	-	-	-	2	1	4	5
Female	31	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
North Carolina												
All races, both sexes	687	-	-	-	-	-	1	3	7	8	32	38
Male	321	-	-	-	-	-	1	1	4	5	19	23
Female	366	-	-	-	-	-	-	2	3	3	13	15
White, both sexes	421	-	-	-	-	-	-	1	5	2	22	21
Male	196	-	-	-	-	-	-	-	4	-	15	15
Female	225	-	-	-	-	-	-	1	1	2	7	6
Black, both sexes	260	-	-	-	-	-	1	2	2	6	10	17
Male	124	-	-	-	-	-	1	1	-	5	4	8
Female	136	-	-	-	-	-	-	1	2	1	6	9
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	-	2
Male	27	-	-	-	-	-	-	-	-	-	-	2
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	54	-	-	-	-	-	-	-	-	-	-	2
Male	27	-	-	-	-	-	-	-	-	-	-	2
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Ohio												
All races, both sexes	1,530	-	-	-	-	-	1	3	10	27	48	75
Male	714	-	-	-	-	-	-	3	6	20	37	48
Female	816	-	-	-	-	-	1	-	4	7	11	27
White, both sexes	1,207	-	-	-	-	-	1	2	5	17	30	56
Male	529	-	-	-	-	-	-	2	4	14	26	38
Female	678	-	-	-	-	-	1	-	1	3	4	18
Black, both sexes	315	-	-	-	-	-	-	1	5	10	17	19
Male	181	-	-	-	-	-	-	1	2	6	10	10
Female	134	-	-	-	-	-	-	-	3	4	7	9
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	284	-	-	-	-	-	-	1	2	2	4	11
Male	99	-	-	-	-	-	-	1	2	-	1	5
Female	185	-	-	-	-	-	-	-	-	2	3	6
White, both sexes	228	-	-	-	-	-	-	1	1	1	2	7
Male	84	-	-	-	-	-	-	1	1	-	1	3
Female	144	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	44	-	-	-	-	-	-	-	-	-	1	3
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	35	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	1	1	1	1
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Oregon												
All races, both sexes	224	-	-	-	-	-	-	-	-	1	5	4
Male	78	-	-	-	-	-	-	-	-	1	4	4
Female	146	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	215	-	-	-	-	-	-	-	-	-	4	4
Male	71	-	-	-	-	-	-	-	-	-	3	4
Female	144	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	877	1	-	-	-	-	1	1	2	15	15	51
Male	388	1	-	-	-	-	1	1	2	10	9	34
Female	489	-	-	-	-	-	-	-	-	5	6	17
White, both sexes	709	1	-	-	-	-	-	1	1	8	10	33
Male	297	1	-	-	-	-	-	1	1	7	6	25
Female	412	-	-	-	-	-	-	-	-	1	4	8
Black, both sexes	165	-	-	-	-	-	1	-	1	7	5	18
Male	89	-	-	-	-	-	1	-	1	3	3	9
Female	76	-	-	-	-	-	-	-	-	4	2	9
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Rhode Island												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	-	3
Male	21	-	-	-	-	-	-	-	-	-	-	3
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	45	-	-	-	-	-	-	-	-	-	-	3
Male	19	-	-	-	-	-	-	-	-	-	-	3
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	366	-	-	-	-	-	-	1	-	2	8	13
Male	150	-	-	-	-	-	-	1	-	2	6	9
Female	216	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	213	-	-	-	-	-	-	-	-	1	4	6
Male	90	-	-	-	-	-	-	-	-	1	3	5
Female	123	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	153	-	-	-	-	-	-	1	-	1	4	7
Male	60	-	-	-	-	-	-	1	-	1	3	4
Female	93	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
South Dakota												
All races, both sexes	53	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	842	-	-	-	-	-	3	5	14	16	25	39
Male	337	-	-	-	-	-	1	5	8	11	11	24
Female	505	-	-	-	-	-	2	-	6	5	14	15
White, both sexes	584	-	-	-	-	-	1	4	8	9	16	20
Male	215	-	-	-	-	-	-	4	5	7	6	10
Female	369	-	-	-	-	-	1	-	3	2	10	10
Black, both sexes	255	-	-	-	-	-	2	1	6	7	9	19
Male	121	-	-	-	-	-	1	1	3	4	5	14
Female	134	-	-	-	-	-	1	-	3	3	4	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Texas												
All races, both sexes	1,896	-	-	1	-	-	2	8	22	28	75	94
Male	868	-	-	1	-	-	1	5	15	20	46	63
Female	1,028	-	-	-	-	-	1	3	7	8	29	31
White, both sexes	1,453	-	-	1	-	-	1	5	12	17	44	61
Male	643	-	-	1	-	-	1	5	7	13	29	42
Female	810	-	-	-	-	-	-	-	5	4	15	19
Black, both sexes	422	-	-	-	-	-	1	2	9	9	30	32
Male	212	-	-	-	-	-	-	-	7	5	16	20
Female	210	-	-	-	-	-	1	2	2	4	14	12
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	1	2	1	1
Male	13	-	-	-	-	-	-	-	1	2	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	112	-	-	-	-	-	-	-	-	2	-	2
Male	40	-	-	-	-	-	-	-	-	2	-	1
Female	72	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	111	-	-	-	-	-	-	-	-	2	-	2
Male	39	-	-	-	-	-	-	-	-	2	-	1
Female	72	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Vermont												
All races, both sexes	52	-	-	-	-	-	-	1	-	-	3	3
Male	25	-	-	-	-	-	-	1	-	-	2	1
Female	27	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	52	-	-	-	-	-	-	1	-	-	3	3
Male	25	-	-	-	-	-	-	1	-	-	2	1
Female	27	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	445	-	-	-	-	-	2	4	7	13	11	29
Male	197	-	-	-	-	-	2	2	5	10	11	21
Female	248	-	-	-	-	-	-	2	2	3	-	8
White, both sexes	279	-	-	-	-	-	-	2	4	5	7	9
Male	113	-	-	-	-	-	-	1	3	4	7	7
Female	166	-	-	-	-	-	-	1	1	1	-	2
Black, both sexes	159	-	-	-	-	-	2	1	3	8	3	19
Male	80	-	-	-	-	-	2	-	2	6	3	13
Female	79	-	-	-	-	-	-	1	1	2	-	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	1	-	-	-	1
Male	3	-	-	-	-	-	-	1	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Washington												
All races, both sexes	566	-	-	-	-	-	-	-	2	9	11	15
Male	230	-	-	-	-	-	-	-	2	8	8	8
Female	336	-	-	-	-	-	-	-	-	1	3	7
White, both sexes	516	-	-	-	-	-	-	-	2	7	6	13
Male	204	-	-	-	-	-	-	-	2	6	5	6
Female	312	-	-	-	-	-	-	-	-	1	1	7
Black, both sexes	21	-	-	-	-	-	-	-	-	-	4	2
Male	9	-	-	-	-	-	-	-	-	-	3	2
Female	12	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	1	1	-
Male	14	-	-	-	-	-	-	-	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
West Virginia												
All races, both sexes	163	-	-	-	-	-	-	-	-	2	1	2
Male	62	-	-	-	-	-	-	-	-	1	1	1
Female	101	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	157	-	-	-	-	-	-	-	-	2	1	2
Male	58	-	-	-	-	-	-	-	-	1	1	1
Female	99	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart disease (I11)												
Wisconsin												
All races, both sexes	450	-	-	-	-	-	-	-	2	3	13	12
Male	155	-	-	-	-	-	-	-	-	3	9	8
Female	295	-	-	-	-	-	-	-	2	-	4	4
White, both sexes	397	-	-	-	-	-	-	-	-	1	7	9
Male	131	-	-	-	-	-	-	-	-	1	6	6
Female	266	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	50	-	-	-	-	-	-	-	2	2	6	3
Male	22	-	-	-	-	-	-	-	-	2	3	2
Female	28	-	-	-	-	-	-	-	2	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	3
Male	12	-	-	-	-	-	-	-	-	-	-	3
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
United States												
All races, both sexes	1,834	2,037	1,917	1,780	2,026	2,653	3,717	4,240	3,634	1,834	411	3
Male	1,268	1,393	1,256	1,062	1,085	1,226	1,344	1,262	775	281	47	3
Female	566	644	661	718	941	1,427	2,373	2,978	2,859	1,553	364	-
White, both sexes	1,089	1,308	1,301	1,169	1,429	1,989	2,933	3,586	3,161	1,565	346	3
Male	787	922	879	723	763	937	1,065	1,049	667	223	39	3
Female	302	386	422	446	666	1,052	1,868	2,537	2,494	1,342	307	-
Black, both sexes	697	683	578	571	544	604	694	571	410	225	59	-
Male	446	437	357	312	288	266	241	179	92	45	8	-
Female	251	246	221	259	256	338	453	392	318	180	51	-
American Indian 1/, both sexes	13	13	12	8	11	12	7	12	7	-	2	-
Male	9	10	7	5	4	4	3	5	1	-	-	-
Female	4	3	5	3	7	8	4	7	6	-	2	-
Asian or Pacific Islander 2/, both sexes	35	33	26	32	42	48	83	71	56	44	4	-
Male	26	24	13	22	30	19	35	29	15	13	-	-
Female	9	9	13	10	12	29	48	42	41	31	4	-
Alabama												
All races, both sexes	33	26	20	25	41	42	57	85	52	35	5	-
Male	20	15	12	14	23	22	18	34	7	6	1	-
Female	13	11	8	11	18	20	39	51	45	29	4	-
White, both sexes	13	12	11	14	24	30	34	62	40	22	3	-
Male	11	7	8	9	11	15	10	22	6	2	-	-
Female	2	5	3	5	13	15	24	40	34	20	3	-
Black, both sexes	20	14	9	11	17	12	23	23	12	13	2	-
Male	9	8	4	5	12	7	8	12	1	4	1	-
Female	11	6	5	6	5	5	15	11	11	9	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Alaska												
All races, both sexes	4	3	9	4	5	4	7	2	-	1	-	-
Male	4	3	6	3	3	2	4	-	-	-	-	-
Female	-	-	3	1	2	2	3	2	-	1	-	-
White, both sexes	3	3	7	3	2	2	7	2	-	1	-	-
Male	3	3	5	3	1	1	4	-	-	-	-	-
Female	-	-	2	-	1	1	3	2	-	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	2	1	3	1	-	-	-	-	-	-
Male	1	-	1	-	2	-	-	-	-	-	-	-
Female	-	-	1	1	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	29	45	36	30	33	43	58	81	54	28	3	2
Male	17	30	19	20	16	27	15	24	18	4	1	2
Female	12	15	17	10	17	16	43	57	36	24	2	-
White, both sexes	24	39	31	28	30	39	57	78	51	27	3	2
Male	15	26	16	18	16	26	15	24	17	3	1	2
Female	9	13	15	10	14	13	42	54	34	24	2	-
Black, both sexes	1	4	3	2	1	1	-	2	2	-	-	-
Male	1	3	2	2	-	-	-	-	1	-	-	-
Female	-	1	1	-	1	1	-	2	1	-	-	-
American Indian 1/, both sexes	3	2	2	-	2	2	1	1	1	-	-	-
Male	1	1	1	-	-	1	-	-	-	-	-	-
Female	2	1	1	-	2	1	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Arkansas												
All races, both sexes	11	9	14	14	18	21	34	35	37	17	4	-
Male	6	8	10	10	7	8	14	5	7	3	2	-
Female	5	1	4	4	11	13	20	30	30	14	2	-
White, both sexes	5	6	9	12	14	17	29	32	34	16	2	-
Male	2	5	8	9	4	7	10	4	7	3	1	-
Female	3	1	1	3	10	10	19	28	27	13	1	-
Black, both sexes	6	3	5	2	4	4	5	3	3	-	2	-
Male	4	3	2	1	3	1	4	1	-	-	1	-
Female	2	-	3	1	1	3	1	2	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
California												
All races, both sexes	216	260	253	236	245	346	501	586	507	271	52	-
Male	149	164	162	150	131	158	174	188	118	42	5	-
Female	67	96	91	86	114	188	327	398	389	229	47	-
White, both sexes	152	176	187	178	187	279	401	503	435	232	45	-
Male	106	115	124	116	97	130	138	160	99	32	5	-
Female	46	61	63	62	90	149	263	343	336	200	40	-
Black, both sexes	51	64	49	42	34	41	56	39	39	14	3	-
Male	33	36	30	25	17	14	18	9	10	5	-	-
Female	18	28	19	17	17	27	38	30	29	9	3	-
American Indian 1/, both sexes	1	5	3	1	1	2	-	3	1	-	-	-
Male	1	3	2	-	-	1	-	2	-	-	-	-
Female	-	2	1	1	1	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	12	15	14	15	23	24	44	41	32	25	4	-
Male	9	10	6	9	17	13	18	17	9	5	-	-
Female	3	5	8	6	6	11	26	24	23	20	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Colorado												
All races, both sexes	15	12	21	12	20	19	39	43	23	21	3	-
Male	8	10	12	8	15	9	15	9	3	3	-	-
Female	7	2	9	4	5	10	24	34	20	18	3	-
White, both sexes	11	10	20	10	17	18	37	40	23	18	3	-
Male	5	8	11	6	14	9	13	8	3	3	-	-
Female	6	2	9	4	3	9	24	32	20	15	3	-
Black, both sexes	3	1	1	2	2	1	2	3	-	3	-	-
Male	2	1	1	2	1	-	2	1	-	-	-	-
Female	1	-	-	-	1	1	-	2	-	3	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	10	8	17	9	8	18	16	29	38	17	6	-
Male	10	8	12	3	5	9	11	7	8	3	-	-
Female	-	-	5	6	3	9	5	22	30	14	6	-
White, both sexes	6	7	14	8	7	17	13	28	37	17	6	-
Male	6	7	10	3	5	9	10	7	8	3	-	-
Female	-	-	4	5	2	8	3	21	29	14	6	-
Black, both sexes	4	1	2	1	1	1	3	-	1	-	-	-
Male	4	1	1	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	1	1	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Delaware												
All races, both sexes	2	3	10	4	7	5	19	8	10	8	2	-
Male	-	2	5	4	3	1	7	5	1	2	-	-
Female	2	1	5	-	4	4	12	3	9	6	2	-
White, both sexes	2	1	4	2	5	5	18	7	9	6	2	-
Male	-	1	3	2	2	1	6	4	1	2	-	-
Female	2	-	1	-	3	4	12	3	8	4	2	-
Black, both sexes	-	2	6	2	2	-	1	1	1	2	-	-
Male	-	1	2	2	1	-	1	1	-	-	-	-
Female	-	1	4	-	1	-	-	-	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	13	13	15	15	11	19	29	33	15	11	3	-
Male	12	5	11	9	4	9	11	9	2	4	1	-
Female	1	8	4	6	7	10	18	24	13	7	2	-
White, both sexes	-	-	2	1	1	2	3	2	4	5	1	-
Male	-	-	2	1	-	2	-	1	1	-	-	-
Female	-	-	-	-	1	-	3	1	3	5	1	-
Black, both sexes	12	13	13	14	10	17	24	31	11	6	2	-
Male	11	5	9	8	4	7	9	8	1	4	1	-
Female	1	8	4	6	6	10	15	23	10	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Florida												
All races, both sexes	148	176	172	161	161	205	300	347	254	132	26	-
Male	104	121	131	94	92	107	120	108	57	28	5	-
Female	44	55	41	67	69	98	180	239	197	104	21	-
White, both sexes	97	122	123	114	129	175	261	312	237	120	25	-
Male	73	88	97	70	76	91	109	99	55	25	5	-
Female	24	34	26	44	53	84	152	213	182	95	20	-
Black, both sexes	49	53	47	46	31	30	37	34	17	12	1	-
Male	30	33	34	23	16	16	10	9	2	3	-	-
Female	19	20	13	23	15	14	27	25	15	9	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	2	1	1	-	2	1	-	-	-	-
Male	1	-	-	1	-	-	1	-	-	-	-	-
Female	1	1	2	-	1	-	1	1	-	-	-	-
Georgia												
All races, both sexes	91	99	96	98	88	95	102	94	92	42	13	-
Male	70	70	62	58	49	43	35	27	24	3	1	-
Female	21	29	34	40	39	52	67	67	68	39	12	-
White, both sexes	28	42	50	48	42	47	66	62	65	24	5	-
Male	25	32	37	32	23	21	22	19	16	1	-	-
Female	3	10	13	16	19	26	44	43	49	23	5	-
Black, both sexes	63	55	44	49	44	47	35	31	27	18	8	-
Male	45	36	24	26	25	21	12	8	8	2	1	-
Female	18	19	20	23	19	26	23	23	19	16	7	-
American Indian 1/, both sexes	-	-	1	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	1	1	-	1	-	-	-	-	-
Male	-	2	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Hawaii												
All races, both sexes	5	1	1	1	5	3	7	7	4	7	-	-
Male	3	1	-	-	4	2	3	2	1	2	-	-
Female	2	-	1	1	1	1	4	5	3	5	-	-
White, both sexes	1	1	-	-	2	1	1	-	1	-	-	-
Male	-	1	-	-	2	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	1	1	3	2	6	7	3	7	-	-
Male	2	-	-	-	2	2	2	2	1	2	-	-
Female	1	-	1	1	1	-	4	5	2	5	-	-
Idaho												
All races, both sexes	4	2	3	4	5	8	19	14	16	12	1	-
Male	3	1	2	3	2	1	9	6	3	-	-	-
Female	1	1	1	1	3	7	10	8	13	12	1	-
White, both sexes	4	2	3	4	5	7	19	14	16	12	1	-
Male	3	1	2	3	2	1	9	6	3	-	-	-
Female	1	1	1	1	3	6	10	8	13	12	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Illinois												
All races, both sexes	101	108	117	81	116	150	165	180	146	86	16	-
Male	74	76	91	45	65	71	65	61	28	14	-	-
Female	27	32	26	36	51	79	100	119	118	72	16	-
White, both sexes	52	52	66	42	72	103	121	142	130	68	14	-
Male	41	37	51	21	41	50	48	48	23	12	-	-
Female	11	15	15	21	31	53	73	94	107	56	14	-
Black, both sexes	47	54	51	36	42	44	41	38	15	17	2	-
Male	31	37	40	21	23	21	16	13	5	2	-	-
Female	16	17	11	15	19	23	25	25	10	15	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	-	3	2	3	2	-	1	1	-	-
Male	2	2	-	3	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	3	1	-	1	1	-	-
Indiana												
All races, both sexes	21	23	15	23	37	30	59	72	70	38	6	-
Male	13	15	11	11	20	11	21	20	16	7	-	-
Female	8	8	4	12	17	19	38	52	54	31	6	-
White, both sexes	14	22	12	17	31	25	50	65	67	35	4	-
Male	10	15	10	7	17	8	18	16	14	6	-	-
Female	4	7	2	10	14	17	32	49	53	29	4	-
Black, both sexes	6	1	3	6	6	5	9	5	3	3	2	-
Male	3	-	1	4	3	3	3	3	2	1	-	-
Female	3	1	2	2	3	2	6	2	1	2	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Iowa												
All races, both sexes	10	4	9	12	8	20	47	58	57	29	11	-
Male	7	4	4	9	3	10	15	19	16	6	-	-
Female	3	-	5	3	5	10	32	39	41	23	11	-
White, both sexes	8	3	9	12	8	20	47	58	57	29	11	-
Male	5	3	4	9	3	10	15	19	16	6	-	-
Female	3	-	5	3	5	10	32	39	41	23	11	-
Black, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	6	4	5	2	3	10	16	33	41	7	2	-
Male	5	3	4	1	1	5	7	7	7	2	-	-
Female	1	1	1	1	2	5	9	26	34	5	2	-
White, both sexes	4	4	5	2	3	10	14	28	38	7	2	-
Male	3	3	4	1	1	5	5	5	6	2	-	-
Female	1	1	1	1	2	5	9	23	32	5	2	-
Black, both sexes	2	-	-	-	-	-	2	5	2	-	-	-
Male	2	-	-	-	-	-	2	2	-	-	-	-
Female	-	-	-	-	-	-	-	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Kentucky												
All races, both sexes	21	34	30	21	32	33	44	43	34	12	3	-
Male	12	18	17	11	15	17	15	10	5	4	1	-
Female	9	16	13	10	17	16	29	33	29	8	2	-
White, both sexes	14	26	25	18	31	29	38	39	31	12	3	-
Male	9	14	17	11	14	16	14	8	5	4	1	-
Female	5	12	8	7	17	13	24	31	26	8	2	-
Black, both sexes	7	8	5	3	1	4	6	4	3	-	-	-
Male	3	4	-	-	1	1	1	2	-	-	-	-
Female	4	4	5	3	-	3	5	2	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	44	65	45	49	50	71	88	85	69	25	4	-
Male	34	46	29	26	29	26	37	24	8	1	1	-
Female	10	19	16	23	21	45	51	61	61	24	3	-
White, both sexes	24	29	20	22	30	43	55	61	50	18	2	-
Male	19	20	11	14	17	16	20	18	6	1	1	-
Female	5	9	9	8	13	27	35	43	44	17	1	-
Black, both sexes	20	36	25	27	20	28	33	24	19	6	2	-
Male	15	26	18	12	12	10	17	6	2	-	-	-
Female	5	10	7	15	8	18	16	18	17	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Maine												
All races, both sexes	4	5	3	3	7	5	6	7	6	5	2	-
Male	3	4	3	3	5	3	3	-	1	1	-	-
Female	1	1	-	-	2	2	3	7	5	4	2	-
White, both sexes	4	5	3	3	7	5	6	7	6	5	2	-
Male	3	4	3	3	5	3	3	-	1	1	-	-
Female	1	1	-	-	2	2	3	7	5	4	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	78	69	50	51	49	69	81	87	72	43	7	-
Male	53	52	28	35	33	24	27	29	11	6	-	-
Female	25	17	22	16	16	45	54	58	61	37	7	-
White, both sexes	22	35	22	24	28	41	55	66	56	35	6	-
Male	16	29	12	17	17	16	20	26	10	4	-	-
Female	6	6	10	7	11	25	35	40	46	31	6	-
Black, both sexes	55	33	28	26	21	26	25	19	15	8	1	-
Male	37	22	16	17	16	7	6	3	1	2	-	-
Female	18	11	12	9	5	19	19	16	14	6	1	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	1	1	2	1	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Massachusetts												
All races, both sexes	28	43	13	10	22	23	26	43	46	21	4	-
Male	19	33	10	5	8	12	10	8	6	4	-	-
Female	9	10	3	5	14	11	16	35	40	17	4	-
White, both sexes	23	40	12	8	16	22	22	42	44	21	3	-
Male	17	31	9	4	6	12	9	8	5	4	-	-
Female	6	9	3	4	10	10	13	34	39	17	3	-
Black, both sexes	5	3	1	2	5	1	3	1	2	-	1	-
Male	2	2	1	1	2	-	-	-	1	-	-	-
Female	3	1	-	1	3	1	3	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Michigan												
All races, both sexes	64	93	89	98	84	121	163	213	148	75	21	-
Male	34	59	61	53	42	55	52	57	27	11	3	-
Female	30	34	28	45	42	66	111	156	121	64	18	-
White, both sexes	38	64	62	62	55	97	122	182	131	63	21	-
Male	21	42	45	35	26	42	39	44	22	9	3	-
Female	17	22	17	27	29	55	83	138	109	54	18	-
Black, both sexes	26	29	26	36	28	24	41	31	17	12	-	-
Male	13	17	16	18	16	13	13	13	5	2	-	-
Female	13	12	10	18	12	11	28	18	12	10	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Minnesota												
All races, both sexes	9	11	4	6	10	13	24	24	34	16	9	-
Male	7	5	2	5	5	9	8	4	9	3	-	-
Female	2	6	2	1	5	4	16	20	25	13	9	-
White, both sexes	8	8	4	6	9	12	24	24	34	16	9	-
Male	6	4	2	5	4	8	8	4	9	3	-	-
Female	2	4	2	1	5	4	16	20	25	13	9	-
Black, both sexes	1	3	-	-	1	1	-	-	-	-	-	-
Male	1	1	-	-	1	1	-	-	-	-	-	-
Female	-	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	33	46	44	46	55	72	76	109	78	46	8	-
Male	25	32	27	19	31	27	30	33	17	5	3	-
Female	8	14	17	27	24	45	46	76	61	41	5	-
White, both sexes	16	22	24	19	27	37	50	69	46	32	3	-
Male	13	16	14	11	19	16	18	20	8	3	1	-
Female	3	6	10	8	8	21	32	49	38	29	2	-
Black, both sexes	17	23	20	27	28	35	26	40	32	14	5	-
Male	12	15	13	8	12	11	12	13	9	2	2	-
Female	5	8	7	19	16	24	14	27	23	12	3	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Missouri												
All races, both sexes	38	26	36	22	27	36	53	59	45	20	4	-
Male	24	21	19	11	20	18	16	17	5	6	-	-
Female	14	5	17	11	7	18	37	42	40	14	4	-
White, both sexes	18	16	20	15	21	21	39	53	38	18	4	-
Male	11	13	11	7	15	9	13	15	3	6	-	-
Female	7	3	9	8	6	12	26	38	35	12	4	-
Black, both sexes	19	10	15	6	6	15	14	6	7	2	-	-
Male	12	8	7	3	5	9	3	2	2	-	-	-
Female	7	2	8	3	1	6	11	4	5	2	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	2	7	1	2	3	10	9	6	4	-	-
Male	2	2	2	-	1	2	3	3	-	-	-	-
Female	-	-	5	1	1	1	7	6	6	4	-	-
White, both sexes	2	2	7	1	2	3	9	9	6	4	-	-
Male	2	2	2	-	1	2	3	3	-	-	-	-
Female	-	-	5	1	1	1	6	6	6	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Nebraska												
All races, both sexes	5	3	1	4	5	7	11	18	13	9	2	-
Male	4	3	1	3	3	3	4	5	2	2	1	-
Female	1	-	-	1	2	4	7	13	11	7	1	-
White, both sexes	4	3	-	4	5	6	9	18	11	8	2	-
Male	3	3	-	3	3	2	3	5	2	2	1	-
Female	1	-	-	1	2	4	6	13	9	6	1	-
Black, both sexes	1	-	1	-	-	1	2	-	2	1	-	-
Male	1	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	47	53	49	54	48	59	63	46	26	11	-	-
Male	35	39	35	35	28	37	34	16	10	1	-	-
Female	12	14	14	19	20	22	29	30	16	10	-	-
White, both sexes	38	42	40	48	43	55	59	43	23	10	-	-
Male	27	30	30	30	25	35	32	15	9	1	-	-
Female	11	12	10	18	18	20	27	28	14	9	-	-
Black, both sexes	5	9	7	3	5	4	3	-	1	-	-	-
Male	4	7	3	2	3	2	1	-	-	-	-	-
Female	1	2	4	1	2	2	2	-	1	-	-	-
American Indian 1/, both sexes	1	1	1	2	-	-	-	-	-	-	-	-
Male	1	1	1	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	1	1	-	-	1	3	2	1	-	-
Male	3	1	1	1	-	-	1	1	1	-	-	-
Female	-	-	-	-	-	-	-	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
New Hampshire												
All races, both sexes	4	4	7	5	5	8	9	11	10	9	3	-
Male	4	4	4	4	2	6	1	3	2	-	-	-
Female	-	-	3	1	3	2	8	8	8	9	3	-
White, both sexes	4	4	7	5	5	8	9	11	10	9	3	-
Male	4	4	4	4	2	6	1	3	2	-	-	-
Female	-	-	3	1	3	2	8	8	8	9	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	40	42	55	39	54	69	104	136	106	46	21	-
Male	24	33	35	31	30	31	43	41	29	8	3	-
Female	16	9	20	8	24	38	61	95	77	38	18	-
White, both sexes	23	26	39	23	35	53	86	129	95	42	18	-
Male	15	23	27	17	19	25	38	40	26	6	3	-
Female	8	3	12	6	16	28	48	89	69	36	15	-
Black, both sexes	17	15	15	15	17	12	16	6	10	3	3	-
Male	9	9	7	13	9	5	5	1	3	1	-	-
Female	8	6	8	2	8	7	11	5	7	2	3	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	2	3	2	1	1	1	-	-
Male	-	1	1	1	2	1	-	-	-	1	-	-
Female	-	-	-	-	-	2	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
New Mexico												
All races, both sexes	17	15	16	9	16	18	15	20	15	11	2	-
Male	15	9	12	7	13	6	8	8	2	1	1	-
Female	2	6	4	2	3	12	7	12	13	10	1	-
White, both sexes	13	15	16	7	15	16	13	16	14	10	2	-
Male	12	9	12	5	12	5	6	5	2	1	1	-
Female	1	6	4	2	3	11	7	11	12	9	1	-
Black, both sexes	2	-	-	2	1	1	1	2	-	1	-	-
Male	1	-	-	2	1	1	1	1	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	1	2	1	-	-	-
Male	2	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
New York												
All races, both sexes	150	144	140	167	190	212	333	339	298	143	35	-
Male	91	94	81	91	94	104	115	107	64	17	3	-
Female	59	50	59	76	96	108	218	232	234	126	32	-
White, both sexes	75	77	78	94	121	129	224	267	244	119	27	-
Male	45	50	46	49	61	69	86	80	54	13	3	-
Female	30	27	32	45	60	60	138	187	190	106	24	-
Black, both sexes	72	64	59	70	64	77	95	61	46	21	8	-
Male	43	42	33	40	28	33	25	21	9	3	-	-
Female	29	22	26	30	36	44	70	40	37	18	8	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	2	3	5	5	14	10	8	3	-	-
Male	3	2	1	2	5	1	4	6	1	1	-	-
Female	-	1	1	1	-	4	10	4	7	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
North Carolina												
All races, both sexes	51	59	44	47	47	60	86	89	74	33	7	1
Male	40	46	31	27	19	28	35	29	7	4	1	1
Female	11	13	13	20	28	32	51	60	67	29	6	-
White, both sexes	26	31	27	29	25	33	49	59	60	24	6	1
Male	24	25	19	18	10	15	20	21	6	2	1	1
Female	2	6	8	11	15	18	29	38	54	22	5	-
Black, both sexes	25	27	17	17	22	27	36	30	12	9	-	-
Male	16	20	12	9	9	13	15	8	1	2	-	-
Female	9	7	5	8	13	14	21	22	11	7	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	2	-	1	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	2	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	3	1	1	7	3	10	16	10	-	-
Male	-	1	3	-	1	6	1	4	6	3	-	-
Female	-	-	-	1	-	1	2	6	10	7	-	-
White, both sexes	-	1	3	1	1	7	3	10	16	10	-	-
Male	-	1	3	-	1	6	1	4	6	3	-	-
Female	-	-	-	1	-	1	2	6	10	7	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Ohio												
All races, both sexes	120	108	104	87	96	128	197	201	205	100	20	-
Male	87	73	71	53	56	53	75	67	51	13	1	-
Female	33	35	33	34	40	75	122	134	154	87	19	-
White, both sexes	74	76	76	56	69	102	166	182	187	91	17	-
Male	55	55	53	33	36	42	59	58	42	11	1	-
Female	19	21	23	23	33	60	107	124	145	80	16	-
Black, both sexes	44	32	28	29	27	24	31	19	17	9	3	-
Male	30	18	18	19	20	11	16	9	9	2	-	-
Female	14	14	10	10	7	13	15	10	8	7	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	2	-	2	-	-	1	-	-	-
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	1	-	-	-
Oklahoma												
All races, both sexes	6	11	12	13	20	34	58	42	43	13	12	-
Male	5	7	5	6	11	14	20	5	13	3	1	-
Female	1	4	7	7	9	20	38	37	30	10	11	-
White, both sexes	5	9	10	12	14	26	50	32	38	11	9	-
Male	4	6	5	6	8	11	18	4	13	2	1	-
Female	1	3	5	6	6	15	32	28	25	9	8	-
Black, both sexes	-	1	2	1	5	7	7	8	5	2	2	-
Male	-	-	-	-	2	3	1	-	-	1	-	-
Female	-	1	2	1	3	4	6	8	5	1	2	-
American Indian 1/, both sexes	1	1	-	-	1	1	1	2	-	-	1	-
Male	1	1	-	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Oregon												
All races, both sexes	8	9	5	5	12	16	26	49	48	25	11	-
Male	8	7	2	2	5	6	9	14	9	4	3	-
Female	-	2	3	3	7	10	17	35	39	21	8	-
White, both sexes	8	8	5	4	11	15	26	49	48	23	10	-
Male	8	6	2	1	4	6	9	14	9	3	2	-
Female	-	2	3	3	7	9	17	35	39	20	8	-
Black, both sexes	-	1	-	-	1	-	-	-	-	1	1	-
Male	-	1	-	-	1	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	37	45	54	54	52	90	111	147	134	51	16	-
Male	28	37	37	35	30	36	40	54	25	6	2	-
Female	9	8	17	19	22	54	71	93	109	45	14	-
White, both sexes	23	33	36	38	39	74	97	132	123	45	15	-
Male	21	26	24	24	23	29	33	47	23	4	2	-
Female	2	7	12	14	16	45	64	85	100	41	13	-
Black, both sexes	14	12	18	14	13	16	14	15	10	6	1	-
Male	7	11	13	9	7	7	7	7	2	2	-	-
Female	7	1	5	5	6	9	7	8	8	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	-	-	1	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Rhode Island												
All races, both sexes	-	3	2	1	2	4	5	9	10	7	2	-
Male	-	2	1	1	-	3	3	3	3	2	-	-
Female	-	1	1	-	2	1	2	6	7	5	2	-
White, both sexes	-	2	2	1	2	3	5	8	10	7	2	-
Male	-	1	1	1	-	2	3	3	3	2	-	-
Female	-	1	1	-	2	1	2	5	7	5	2	-
Black, both sexes	-	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	18	26	20	25	31	37	60	54	46	19	6	-
Male	13	12	13	13	13	19	17	13	14	3	2	-
Female	5	14	7	12	18	18	43	41	32	16	4	-
White, both sexes	7	13	12	11	18	21	36	37	33	11	3	-
Male	7	6	9	5	10	13	11	7	11	1	1	-
Female	-	7	3	6	8	8	25	30	22	10	2	-
Black, both sexes	11	13	8	14	13	16	24	17	13	8	3	-
Male	6	6	4	8	3	6	6	6	3	2	1	-
Female	5	7	4	6	10	10	18	11	10	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
South Dakota												
All races, both sexes	-	-	-	3	1	1	7	13	13	14	1	-
Male	-	-	-	2	1	-	1	3	3	3	-	-
Female	-	-	-	1	-	1	6	10	10	11	1	-
White, both sexes	-	-	-	3	-	1	7	13	13	14	1	-
Male	-	-	-	2	-	-	1	3	3	3	-	-
Female	-	-	-	1	-	1	6	10	10	11	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	56	54	25	42	68	68	114	121	121	62	9	-
Male	40	34	15	25	29	34	28	30	28	13	1	-
Female	16	20	10	17	39	34	86	91	93	49	8	-
White, both sexes	32	33	10	23	39	42	93	99	97	49	9	-
Male	23	19	4	17	19	20	23	25	21	11	1	-
Female	9	14	6	6	20	22	70	74	76	38	8	-
Black, both sexes	23	21	14	18	29	26	21	22	24	13	-	-
Male	17	15	11	7	10	14	5	5	7	2	-	-
Female	6	6	3	11	19	12	16	17	17	11	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Texas												
All races, both sexes	146	176	147	104	127	176	232	226	207	102	23	-
Male	93	116	98	72	66	81	77	55	46	12	1	-
Female	53	60	49	32	61	95	155	171	161	90	22	-
White, both sexes	105	115	106	76	99	137	188	195	184	87	20	-
Male	72	76	71	54	51	61	62	47	40	10	1	-
Female	33	39	35	22	48	76	126	148	144	77	19	-
Black, both sexes	39	56	40	28	27	38	41	31	21	15	3	-
Male	21	35	26	18	15	20	13	8	6	2	-	-
Female	18	21	14	10	12	18	28	23	15	13	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	1	-	1	1	2	-	2	-	-	-
Male	-	5	1	-	-	-	2	-	-	-	-	-
Female	2	-	-	-	1	1	-	-	2	-	-	-
Utah												
All races, both sexes	4	6	7	5	4	11	15	29	23	3	1	-
Male	1	5	5	4	1	1	5	8	6	-	1	-
Female	3	1	2	1	3	10	10	21	17	3	-	-
White, both sexes	4	5	7	5	4	11	15	29	23	3	1	-
Male	1	4	5	4	1	1	5	8	6	-	1	-
Female	3	1	2	1	3	10	10	21	17	3	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Vermont												
All races, both sexes	3	6	2	-	3	4	7	9	9	2	-	-
Male	2	4	1	-	1	2	3	4	4	-	-	-
Female	1	2	1	-	2	2	4	5	5	2	-	-
White, both sexes	3	6	2	-	3	4	7	9	9	2	-	-
Male	2	4	1	-	1	2	3	4	4	-	-	-
Female	1	2	1	-	2	2	4	5	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	31	27	29	28	26	35	53	66	55	24	5	-
Male	21	18	17	14	15	17	17	16	8	3	-	-
Female	10	9	12	14	11	18	36	50	47	21	5	-
White, both sexes	15	13	17	19	11	21	41	53	39	22	1	-
Male	10	9	12	11	6	10	12	12	6	3	-	-
Female	5	4	5	8	5	11	29	41	33	19	1	-
Black, both sexes	16	13	12	9	15	12	12	12	16	2	4	-
Male	11	9	5	3	9	6	5	4	2	-	-	-
Female	5	4	7	6	6	6	7	8	14	2	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	2	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Washington												
All races, both sexes	27	25	28	21	35	47	83	113	98	39	13	-
Male	21	24	16	12	19	19	37	32	20	4	-	-
Female	6	1	12	9	16	28	46	81	78	35	13	-
White, both sexes	20	22	26	17	32	44	76	107	94	37	13	-
Male	16	22	15	11	17	19	34	30	18	3	-	-
Female	4	-	11	6	15	25	42	77	76	34	13	-
Black, both sexes	4	1	1	3	-	1	2	2	-	1	-	-
Male	2	1	-	-	-	-	1	-	-	-	-	-
Female	2	-	1	3	-	1	1	2	-	1	-	-
American Indian 1/, both sexes	1	1	-	-	1	-	-	-	1	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	1	2	2	5	4	3	1	-	-
Male	2	-	1	1	2	-	2	2	2	1	-	-
Female	-	1	-	-	-	2	3	2	1	-	-	-
West Virginia												
All races, both sexes	4	9	9	4	12	24	24	24	30	15	3	-
Male	3	6	4	4	4	13	9	8	4	3	1	-
Female	1	3	5	-	8	11	15	16	26	12	2	-
White, both sexes	3	9	8	4	12	24	23	23	28	15	3	-
Male	2	6	3	4	4	13	8	7	4	3	1	-
Female	1	3	5	-	8	11	15	16	24	12	2	-
Black, both sexes	1	-	1	-	-	-	1	1	2	-	-	-
Male	1	-	1	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart disease (I11)												
Wisconsin												
All races, both sexes	18	19	21	24	20	54	52	77	76	55	4	-
Male	14	10	15	11	11	19	16	19	13	6	1	-
Female	4	9	6	13	9	35	36	58	63	49	3	-
White, both sexes	12	14	19	16	19	50	50	73	72	51	4	-
Male	10	9	13	7	10	19	16	17	12	4	1	-
Female	2	5	6	9	9	31	34	56	60	47	3	-
Black, both sexes	6	5	2	8	1	3	2	4	3	3	-	-
Male	4	1	2	4	1	-	-	2	-	1	-	-
Female	2	4	-	4	-	3	2	2	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	2	3	-	2	-	3	5	4	2	-	-
Male	1	1	2	-	1	-	1	2	1	-	-	-
Female	1	1	1	-	1	-	2	3	3	2	-	-
White, both sexes	2	2	3	-	2	-	3	5	4	2	-	-
Male	1	1	2	-	1	-	1	2	1	-	-	-
Female	1	1	1	-	1	-	2	3	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
United States												
All races, both sexes	10,045	-	-	1	-	5	10	28	33	71	144	248
Male	4,374	-	-	-	-	2	5	13	21	36	94	148
Female	5,671	-	-	1	-	3	5	15	12	35	50	100
White, both sexes	7,035	-	-	1	-	2	5	10	9	21	57	94
Male	3,022	-	-	-	-	1	3	5	7	11	41	58
Female	4,013	-	-	1	-	1	2	5	2	10	16	36
Black, both sexes	2,675	-	-	-	-	2	5	15	24	46	81	150
Male	1,204	-	-	-	-	1	2	6	14	22	49	86
Female	1,471	-	-	-	-	1	3	9	10	24	32	64
American Indian 1/, both sexes	54	-	-	-	-	-	-	1	-	-	1	1
Male	26	-	-	-	-	-	-	1	-	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	281	-	-	-	-	1	-	2	-	4	5	3
Male	122	-	-	-	-	-	-	1	-	3	4	3
Female	159	-	-	-	-	1	-	1	-	1	1	-
Alabama												
All races, both sexes	156	-	-	-	-	-	-	-	-	3	3	5
Male	61	-	-	-	-	-	-	-	-	2	3	2
Female	95	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	63	-	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	39	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	93	-	-	-	-	-	-	-	-	3	3	4
Male	37	-	-	-	-	-	-	-	-	2	3	1
Female	56	-	-	-	-	-	-	-	-	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Alaska												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	198	-	-	-	-	-	-	-	2	2	1	1
Male	104	-	-	-	-	-	-	-	2	-	-	-
Female	94	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	184	-	-	-	-	-	-	-	2	2	-	1
Male	96	-	-	-	-	-	-	-	2	-	-	-
Female	88	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Arkansas												
All races, both sexes	97	-	-	-	-	-	-	1	-	1	1	3
Male	36	-	-	-	-	-	-	1	-	1	-	1
Female	61	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	67	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	30	-	-	-	-	-	-	1	-	1	1	2
Male	14	-	-	-	-	-	-	1	-	1	-	1
Female	16	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,193	-	-	-	-	1	1	3	5	10	23	26
Male	549	-	-	-	-	-	1	2	2	6	19	19
Female	644	-	-	-	-	1	-	1	3	4	4	7
White, both sexes	850	-	-	-	-	-	-	-	1	3	15	13
Male	387	-	-	-	-	-	-	-	-	2	13	10
Female	463	-	-	-	-	-	-	-	1	1	2	3
Black, both sexes	189	-	-	-	-	-	1	-	4	4	5	11
Male	97	-	-	-	-	-	1	-	2	1	3	7
Female	92	-	-	-	-	-	-	-	2	3	2	4
American Indian 1/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	150	-	-	-	-	1	-	2	-	3	3	2
Male	64	-	-	-	-	-	-	1	-	3	3	2
Female	86	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Colorado												
All races, both sexes	99	-	-	-	-	-	-	-	1	1	4	-
Male	40	-	-	-	-	-	-	-	1	-	2	-
Female	59	-	-	-	-	-	-	-	-	1	2	-
White, both sexes	90	-	-	-	-	-	-	-	-	-	3	-
Male	37	-	-	-	-	-	-	-	-	-	1	-
Female	53	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	7	-	-	-	-	-	-	-	1	-	1	-
Male	3	-	-	-	-	-	-	-	1	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Connecticut												
All races, both sexes	68	-	-	-	-	-	-	-	1	-	-	2
Male	28	-	-	-	-	-	-	-	-	-	-	2
Female	40	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	58	-	-	-	-	-	-	-	-	-	-	2
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Delaware												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Florida												
All races, both sexes	811	-	-	-	-	-	2	2	3	6	10	22
Male	371	-	-	-	-	-	2	-	1	3	8	15
Female	440	-	-	-	-	-	-	2	2	3	2	7
White, both sexes	595	-	-	-	-	-	2	-	-	2	5	12
Male	276	-	-	-	-	-	2	-	-	1	4	10
Female	319	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	207	-	-	-	-	-	-	2	3	4	5	10
Male	90	-	-	-	-	-	-	-	1	2	4	5
Female	117	-	-	-	-	-	-	2	2	2	1	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	315	-	-	-	-	1	1	2	1	2	3	14
Male	123	-	-	-	-	1	1	-	-	1	1	6
Female	192	-	-	-	-	-	-	2	1	1	2	8
White, both sexes	127	-	-	-	-	-	-	1	-	-	-	3
Male	55	-	-	-	-	-	-	-	-	-	-	1
Female	72	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	188	-	-	-	-	1	1	1	1	2	3	11
Male	68	-	-	-	-	1	1	-	-	1	1	5
Female	120	-	-	-	-	-	-	1	1	1	2	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Hawaii												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	1	-
Idaho												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Illinois												
All races, both sexes	372	-	-	-	-	-	-	1	-	2	4	13
Male	139	-	-	-	-	-	-	1	-	-	1	8
Female	233	-	-	-	-	-	-	-	-	2	3	5
White, both sexes	262	-	-	-	-	-	-	-	-	-	1	3
Male	87	-	-	-	-	-	-	-	-	-	-	1
Female	175	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	104	-	-	-	-	-	-	1	-	2	3	10
Male	49	-	-	-	-	-	-	1	-	-	1	7
Female	55	-	-	-	-	-	-	-	-	2	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	215	-	-	-	-	-	-	3	2	-	5	3
Male	79	-	-	-	-	-	-	2	1	-	3	1
Female	136	-	-	-	-	-	-	1	1	-	2	2
White, both sexes	182	-	-	-	-	-	-	2	1	-	2	1
Male	63	-	-	-	-	-	-	2	1	-	1	-
Female	119	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	33	-	-	-	-	-	-	1	1	-	3	2
Male	16	-	-	-	-	-	-	-	-	-	2	1
Female	17	-	-	-	-	-	-	1	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Iowa												
All races, both sexes	145	-	-	-	-	-	-	1	-	-	1	1
Male	64	-	-	-	-	-	-	-	-	-	-	1
Female	81	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	141	-	-	-	-	-	-	1	-	-	-	1
Male	64	-	-	-	-	-	-	-	-	-	-	1
Female	77	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	80	-	-	-	-	-	-	-	-	-	-	2
Male	35	-	-	-	-	-	-	-	-	-	-	2
Female	45	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Kentucky												
All races, both sexes	135	-	-	-	-	-	-	1	1	2	1	3
Male	57	-	-	-	-	-	-	-	-	2	1	1
Female	78	-	-	-	-	-	-	1	1	-	-	2
White, both sexes	114	-	-	-	-	-	-	1	-	2	1	2
Male	49	-	-	-	-	-	-	-	-	2	1	1
Female	65	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	19	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	205	-	-	-	-	-	-	-	2	3	7	7
Male	96	-	-	-	-	-	-	-	2	2	4	6
Female	109	-	-	-	-	-	-	-	-	1	3	1
White, both sexes	102	-	-	-	-	-	-	-	1	1	2	-
Male	52	-	-	-	-	-	-	-	1	1	2	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	102	-	-	-	-	-	-	-	1	2	5	7
Male	43	-	-	-	-	-	-	-	1	1	2	6
Female	59	-	-	-	-	-	-	-	-	1	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Maine												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	205	-	-	-	-	-	-	-	1	1	4	7
Male	95	-	-	-	-	-	-	-	1	-	4	6
Female	110	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	104	-	-	-	-	-	-	-	-	-	1	1
Male	46	-	-	-	-	-	-	-	-	-	1	1
Female	58	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	101	-	-	-	-	-	-	-	1	1	3	6
Male	49	-	-	-	-	-	-	-	1	-	3	5
Female	52	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Massachusetts												
All races, both sexes	125	-	-	-	-	-	-	-	-	2	1	2
Male	46	-	-	-	-	-	-	-	-	1	1	1
Female	79	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	102	-	-	-	-	-	-	-	-	1	-	1
Male	38	-	-	-	-	-	-	-	-	-	-	-
Female	64	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	18	-	-	-	-	-	-	-	-	1	1	1
Male	5	-	-	-	-	-	-	-	-	1	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	346	-	-	-	-	1	-	1	1	1	2	6
Male	156	-	-	-	-	-	-	-	1	-	2	3
Female	190	-	-	-	-	1	-	1	-	1	-	3
White, both sexes	235	-	-	-	-	1	-	-	-	1	1	2
Male	109	-	-	-	-	-	-	-	-	-	1	2
Female	126	-	-	-	-	1	-	-	-	1	-	-
Black, both sexes	107	-	-	-	-	-	-	1	1	-	1	4
Male	47	-	-	-	-	-	-	-	1	-	1	1
Female	60	-	-	-	-	-	-	1	-	-	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Minnesota												
All races, both sexes	164	-	-	-	-	-	-	-	-	-	2	1
Male	67	-	-	-	-	-	-	-	-	-	1	-
Female	97	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	151	-	-	-	-	-	-	-	-	-	-	-
Male	63	-	-	-	-	-	-	-	-	-	-	-
Female	88	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	2	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	149	-	-	-	-	1	-	1	1	3	1	11
Male	53	-	-	-	-	-	-	-	-	1	-	4
Female	96	-	-	-	-	1	-	1	1	2	1	7
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	108	-	-	-	-	1	-	1	1	3	1	10
Male	42	-	-	-	-	-	-	-	-	1	-	3
Female	66	-	-	-	-	1	-	1	1	2	1	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Missouri												
All races, both sexes	192	-	-	-	-	-	-	-	-	3	1	6
Male	81	-	-	-	-	-	-	-	-	2	-	5
Female	111	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	142	-	-	-	-	-	-	-	-	1	-	2
Male	57	-	-	-	-	-	-	-	-	-	-	2
Female	85	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	50	-	-	-	-	-	-	-	-	2	1	4
Male	24	-	-	-	-	-	-	-	-	2	-	3
Female	26	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Nebraska												
All races, both sexes	69	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	58	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	39	-	-	-	-	-	-	1	-	1	1	1
Male	27	-	-	-	-	-	-	1	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	28	-	-	-	-	-	-	1	-	1	-	-
Male	22	-	-	-	-	-	-	1	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
New Hampshire												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	234	-	-	-	-	-	1	1	1	2	4	7
Male	103	-	-	-	-	-	-	-	1	-	3	2
Female	131	-	-	-	-	-	1	1	-	2	1	5
White, both sexes	153	-	-	-	-	-	-	1	-	-	1	1
Male	62	-	-	-	-	-	-	-	-	-	1	-
Female	91	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	73	-	-	-	-	-	1	-	1	2	3	6
Male	37	-	-	-	-	-	-	-	1	-	2	2
Female	36	-	-	-	-	-	1	-	-	2	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
New Mexico												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	1	-
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	1	-
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	631	-	-	-	-	-	-	4	-	2	9	16
Male	276	-	-	-	-	-	-	2	-	-	5	10
Female	355	-	-	-	-	-	-	2	-	2	4	6
White, both sexes	414	-	-	-	-	-	-	1	-	1	2	7
Male	180	-	-	-	-	-	-	1	-	-	1	4
Female	234	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	198	-	-	-	-	-	-	3	-	1	7	9
Male	85	-	-	-	-	-	-	1	-	-	4	6
Female	113	-	-	-	-	-	-	2	-	1	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
North Carolina												
All races, both sexes	351	-	-	-	-	-	2	1	-	3	3	17
Male	157	-	-	-	-	-	-	-	-	1	3	9
Female	194	-	-	-	-	-	2	1	-	2	-	8
White, both sexes	172	-	-	-	-	-	1	1	-	1	-	1
Male	70	-	-	-	-	-	-	-	-	1	-	-
Female	102	-	-	-	-	-	1	1	-	-	-	1
Black, both sexes	175	-	-	-	-	-	1	-	-	2	3	16
Male	84	-	-	-	-	-	-	-	-	-	3	9
Female	91	-	-	-	-	-	1	-	-	2	-	7
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Ohio												
All races, both sexes	490	-	-	-	-	-	1	-	3	2	6	4
Male	210	-	-	-	-	-	-	-	2	2	4	3
Female	280	-	-	-	-	-	1	-	1	-	2	1
White, both sexes	367	-	-	-	-	-	-	-	-	-	4	3
Male	158	-	-	-	-	-	-	-	-	-	2	2
Female	209	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	121	-	-	-	-	-	1	-	3	2	2	1
Male	51	-	-	-	-	-	-	-	2	2	2	1
Female	70	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	138	-	-	-	-	-	-	-	-	-	1	4
Male	65	-	-	-	-	-	-	-	-	-	1	3
Female	73	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	97	-	-	-	-	-	-	-	-	-	-	1
Male	42	-	-	-	-	-	-	-	-	-	-	1
Female	55	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	25	-	-	-	-	-	-	-	-	-	1	3
Male	14	-	-	-	-	-	-	-	-	-	1	2
Female	11	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Oregon												
All races, both sexes	128	-	-	-	-	-	1	-	-	-	-	2
Male	58	-	-	-	-	-	1	-	-	-	-	2
Female	70	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	117	-	-	-	-	-	1	-	-	-	-	1
Male	53	-	-	-	-	-	1	-	-	-	-	1
Female	64	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	347	-	-	-	-	-	-	1	1	2	5	11
Male	155	-	-	-	-	-	-	-	1	2	4	6
Female	192	-	-	-	-	-	-	1	-	-	1	5
White, both sexes	278	-	-	-	-	-	-	-	1	-	-	8
Male	121	-	-	-	-	-	-	-	1	-	-	4
Female	157	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	67	-	-	-	-	-	-	1	-	2	5	3
Male	34	-	-	-	-	-	-	-	-	2	4	2
Female	33	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Rhode Island												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	2	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	33	-	-	-	-	-	-	-	-	-	2	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	195	-	-	-	-	-	-	1	-	4	2	9
Male	89	-	-	-	-	-	-	1	-	3	-	6
Female	106	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	71	-	-	-	-	-	-	-	-	1	1	1
Male	27	-	-	-	-	-	-	-	-	1	-	-
Female	44	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	122	-	-	-	-	-	-	1	-	3	1	8
Male	60	-	-	-	-	-	-	1	-	2	-	6
Female	62	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
South Dakota												
All races, both sexes	47	-	-	-	-	-	-	-	-	-	-	1
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	43	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	313	-	-	-	-	-	-	2	2	2	5	7
Male	143	-	-	-	-	-	-	2	2	1	4	4
Female	170	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	199	-	-	-	-	-	-	1	-	-	-	4
Male	87	-	-	-	-	-	-	1	-	-	-	2
Female	112	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	110	-	-	-	-	-	-	1	2	2	5	3
Male	53	-	-	-	-	-	-	1	2	1	4	2
Female	57	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Texas												
All races, both sexes	734	-	-	1	-	-	-	-	4	6	22	19
Male	297	-	-	-	-	-	-	-	3	2	12	12
Female	437	-	-	1	-	-	-	-	1	4	10	7
White, both sexes	534	-	-	1	-	-	-	-	2	2	8	10
Male	213	-	-	-	-	-	-	-	1	1	6	7
Female	321	-	-	1	-	-	-	-	1	1	2	3
Black, both sexes	192	-	-	-	-	-	-	-	2	4	13	9
Male	80	-	-	-	-	-	-	-	2	1	5	5
Female	112	-	-	-	-	-	-	-	-	3	8	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Vermont												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	222	-	-	-	-	-	-	1	-	4	2	5
Male	97	-	-	-	-	-	-	1	-	3	2	1
Female	125	-	-	-	-	-	-	-	-	1	-	4
White, both sexes	106	-	-	-	-	-	-	-	-	1	1	2
Male	45	-	-	-	-	-	-	-	-	1	1	-
Female	61	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	111	-	-	-	-	-	-	1	-	3	1	3
Male	51	-	-	-	-	-	-	1	-	2	1	1
Female	60	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Washington												
All races, both sexes	173	-	-	-	-	-	-	-	-	-	2	1
Male	77	-	-	-	-	-	-	-	-	-	2	1
Female	96	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	157	-	-	-	-	-	-	-	-	-	2	1
Male	68	-	-	-	-	-	-	-	-	-	2	1
Female	89	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	74	-	-	-	-	1	-	-	-	1	-	1
Male	28	-	-	-	-	1	-	-	-	1	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	70	-	-	-	-	1	-	-	-	1	-	-
Male	28	-	-	-	-	1	-	-	-	1	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive renal disease (I12)												
Wisconsin												
All races, both sexes	190	-	-	-	-	-	1	-	1	-	2	2
Male	80	-	-	-	-	-	-	-	1	-	1	1
Female	110	-	-	-	-	-	1	-	-	-	1	1
White, both sexes	181	-	-	-	-	-	1	-	1	-	2	1
Male	77	-	-	-	-	-	-	-	1	-	1	-
Female	104	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
United States												
All races, both sexes	292	359	452	639	873	1,361	1,832	1,845	1,288	488	76	-
Male	180	216	227	319	394	621	793	760	417	111	17	-
Female	112	143	225	320	479	740	1,039	1,085	871	377	59	-
White, both sexes	110	173	226	351	524	985	1,445	1,488	1,064	411	59	-
Male	62	103	112	177	231	453	655	638	360	91	14	-
Female	48	70	114	174	293	532	790	850	704	320	45	-
Black, both sexes	176	175	208	257	303	328	334	307	187	61	16	-
Male	114	108	107	128	142	144	116	103	45	15	2	-
Female	62	67	101	129	161	184	218	204	142	46	14	-
American Indian 1/, both sexes	1	1	2	8	12	8	8	5	5	1	-	-
Male	1	-	1	4	7	6	2	-	3	-	-	-
Female	-	1	1	4	5	2	6	5	2	1	-	-
Asian or Pacific Islander 2/, both sexes	5	10	16	23	34	40	45	45	32	15	1	-
Male	3	5	7	10	14	18	20	19	9	5	1	-
Female	2	5	9	13	20	22	25	26	23	10	-	-
Alabama												
All races, both sexes	4	12	10	12	13	16	22	32	18	5	1	-
Male	3	5	5	5	7	7	8	8	4	1	1	-
Female	1	7	5	7	6	9	14	24	14	4	-	-
White, both sexes	-	2	4	4	3	5	12	16	12	3	1	-
Male	-	-	1	2	2	3	6	4	3	1	1	-
Female	-	2	3	2	1	2	6	12	9	2	-	-
Black, both sexes	4	10	6	8	10	11	10	16	6	2	-	-
Male	3	5	4	3	5	4	2	4	1	-	-	-
Female	1	5	2	5	5	7	8	12	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Alaska												
All races, both sexes	-	-	-	-	3	1	-	3	2	1	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	2	1	-	3	2	-	-	-
White, both sexes	-	-	-	-	2	1	-	1	1	1	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	1	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	10	5	9	13	14	33	47	33	23	4	1	-
Male	7	3	4	6	9	21	24	18	10	-	-	-
Female	3	2	5	7	5	12	23	15	13	4	1	-
White, both sexes	8	5	9	10	13	32	45	30	22	4	1	-
Male	5	3	4	5	8	20	23	16	10	-	-	-
Female	3	2	5	5	5	12	22	14	12	4	1	-
Black, both sexes	1	-	-	1	1	-	1	2	-	-	-	-
Male	1	-	-	1	1	-	1	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	2	-	1	1	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	2	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Arkansas												
All races, both sexes	3	8	12	4	2	13	18	14	13	4	-	-
Male	1	6	8	1	-	5	6	3	3	-	-	-
Female	2	2	4	3	2	8	12	11	10	4	-	-
White, both sexes	-	3	4	2	2	13	16	13	9	4	-	-
Male	-	3	3	1	-	5	5	3	2	-	-	-
Female	-	-	1	1	2	8	11	10	7	4	-	-
Black, both sexes	3	5	8	2	-	-	2	1	4	-	-	-
Male	1	3	5	-	-	-	1	-	1	-	-	-
Female	2	2	3	2	-	-	1	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	29	46	54	68	101	160	216	215	164	60	11	-
Male	18	25	27	42	37	72	97	110	55	12	5	-
Female	11	21	27	26	64	88	119	105	109	48	6	-
White, both sexes	18	29	34	42	62	119	165	162	131	46	10	-
Male	9	17	16	22	22	58	76	83	45	9	5	-
Female	9	12	18	20	40	61	89	79	86	37	5	-
Black, both sexes	10	11	13	17	20	24	26	26	12	4	1	-
Male	8	7	7	13	9	9	11	16	3	-	-	-
Female	2	4	6	4	11	15	15	10	9	4	1	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	5	7	9	19	16	25	27	20	10	-	-
Male	1	1	4	7	6	5	10	11	7	3	-	-
Female	-	4	3	2	13	11	15	16	13	7	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Colorado												
All races, both sexes	1	3	3	3	3	16	21	25	9	9	-	-
Male	1	1	-	2	1	9	7	9	4	3	-	-
Female	-	2	3	1	2	7	14	16	5	6	-	-
White, both sexes	1	3	2	3	2	15	21	22	9	9	-	-
Male	1	1	-	2	1	8	7	9	4	3	-	-
Female	-	2	2	1	1	7	14	13	5	6	-	-
Black, both sexes	-	-	1	-	1	1	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Connecticut												
All races, both sexes	-	2	1	5	5	12	14	12	12	2	-	-
Male	-	2	-	3	2	3	6	5	5	-	-	-
Female	-	-	1	2	3	9	8	7	7	2	-	-
White, both sexes	-	2	-	4	4	11	12	11	10	2	-	-
Male	-	2	-	3	2	2	4	4	5	-	-	-
Female	-	-	-	1	2	9	8	7	5	2	-	-
Black, both sexes	-	-	-	1	1	1	2	1	2	-	-	-
Male	-	-	-	-	-	1	2	1	-	-	-	-
Female	-	-	-	1	1	-	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Delaware												
All races, both sexes	-	-	1	1	-	6	5	7	2	2	-	-
Male	-	-	1	-	-	1	1	6	2	1	-	-
Female	-	-	-	1	-	5	4	1	-	1	-	-
White, both sexes	-	-	1	1	-	2	5	5	2	1	-	-
Male	-	-	1	-	-	-	1	5	2	1	-	-
Female	-	-	-	1	-	2	4	-	-	-	-	-
Black, both sexes	-	-	-	-	-	4	-	2	-	1	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	3	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	2	3	2	3	5	6	6	2	2	-	-
Male	2	2	3	2	2	2	2	4	1	2	-	-
Female	1	-	-	-	1	3	4	2	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	3	2	3	2	3	4	6	5	2	1	-	-
Male	2	2	3	2	2	2	2	3	1	1	-	-
Female	1	-	-	-	1	2	4	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Florida												
All races, both sexes	24	29	32	59	61	105	161	154	100	38	3	-
Male	12	9	13	25	27	51	77	75	43	9	1	-
Female	12	20	19	34	34	54	84	79	57	29	2	-
White, both sexes	9	14	19	34	37	76	135	129	88	31	2	-
Male	5	3	7	15	14	34	66	64	42	8	1	-
Female	4	11	12	19	23	42	69	65	46	23	1	-
Black, both sexes	15	15	13	21	24	27	24	24	12	7	1	-
Male	7	6	6	9	13	15	9	11	1	1	-	-
Female	8	9	7	12	11	12	15	13	11	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	4	-	2	2	1	-	-	-	-
Male	-	-	-	1	-	2	2	-	-	-	-	-
Female	-	-	-	3	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes	15	25	15	34	39	32	50	48	26	7	-	-
Male	10	16	6	13	15	14	17	14	7	1	-	-
Female	5	9	9	21	24	18	33	34	19	6	-	-
White, both sexes	1	10	7	9	16	16	30	20	10	4	-	-
Male	1	6	5	3	8	9	11	8	3	-	-	-
Female	-	4	2	6	8	7	19	12	7	4	-	-
Black, both sexes	14	15	8	25	23	16	20	28	16	3	-	-
Male	9	10	1	10	7	5	6	6	4	1	-	-
Female	5	5	7	15	16	11	14	22	12	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Hawaii												
All races, both sexes	1	2	3	3	4	3	3	8	4	3	-	-
Male	-	2	2	-	-	1	-	4	1	1	-	-
Female	1	-	1	3	4	2	3	4	3	2	-	-
White, both sexes	-	-	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	2	3	2	3	7	4	3	-	-
Male	-	2	1	-	-	1	-	3	1	1	-	-
Female	1	-	1	2	3	1	3	4	3	2	-	-
Idaho												
All races, both sexes	-	1	1	1	5	6	6	9	9	-	-	-
Male	-	1	1	1	3	2	5	3	3	-	-	-
Female	-	-	-	-	2	4	1	6	6	-	-	-
White, both sexes	-	1	1	1	5	6	6	9	9	-	-	-
Male	-	1	1	1	3	2	5	3	3	-	-	-
Female	-	-	-	-	2	4	1	6	6	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Illinois												
All races, both sexes	11	12	25	24	34	41	60	73	43	21	8	-
Male	4	8	10	11	13	20	21	27	8	4	3	-
Female	7	4	15	13	21	21	39	46	35	17	5	-
White, both sexes	4	4	14	12	19	31	48	61	40	17	8	-
Male	1	4	5	4	6	14	17	22	8	2	3	-
Female	3	-	9	8	13	17	31	39	32	15	5	-
Black, both sexes	7	8	11	11	14	10	11	11	2	3	-	-
Male	3	4	5	7	6	6	3	5	-	1	-	-
Female	4	4	6	4	8	4	8	6	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	1	1	1	1	-	-
Male	-	-	-	-	1	-	1	-	-	1	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
Indiana												
All races, both sexes	4	6	7	11	17	26	34	47	31	15	4	-
Male	4	4	5	3	6	6	17	19	7	1	-	-
Female	-	2	2	8	11	20	17	28	24	14	4	-
White, both sexes	1	4	6	7	11	23	33	41	31	15	4	-
Male	1	2	4	2	5	5	16	16	7	1	-	-
Female	-	2	2	5	6	18	17	25	24	14	4	-
Black, both sexes	3	2	1	4	6	3	1	6	-	-	-	-
Male	3	2	1	1	1	1	1	3	-	-	-	-
Female	-	-	-	3	5	2	-	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Iowa												
All races, both sexes	3	-	1	4	5	21	30	39	31	8	-	-
Male	-	-	-	3	4	10	17	15	10	4	-	-
Female	3	-	1	1	1	11	13	24	21	4	-	-
White, both sexes	3	-	-	4	5	21	29	38	31	8	-	-
Male	-	-	-	3	4	10	17	15	10	4	-	-
Female	3	-	-	1	1	11	12	23	21	4	-	-
Black, both sexes	-	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	1	3	3	5	11	15	12	20	6	-	-
Male	2	1	2	2	4	5	5	4	8	-	-	-
Female	-	-	1	1	1	6	10	8	12	6	-	-
White, both sexes	2	1	2	3	5	9	12	12	17	6	-	-
Male	2	1	1	2	4	3	3	4	7	-	-	-
Female	-	-	1	1	1	6	9	8	10	6	-	-
Black, both sexes	-	-	1	-	-	2	2	-	3	-	-	-
Male	-	-	1	-	-	2	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Kentucky												
All races, both sexes	3	2	7	7	5	23	27	22	23	6	2	-
Male	3	2	4	3	2	11	9	9	7	2	1	-
Female	-	-	3	4	3	12	18	13	16	4	1	-
White, both sexes	2	1	4	7	5	19	25	16	22	6	1	-
Male	2	1	2	3	2	9	9	8	7	2	-	-
Female	-	-	2	4	3	10	16	8	15	4	1	-
Black, both sexes	1	1	3	-	-	3	2	6	1	-	-	-
Male	1	1	2	-	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	2	2	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	11	6	13	12	26	23	33	25	27	9	1	-
Male	5	4	6	6	16	12	15	10	5	3	-	-
Female	6	2	7	6	10	11	18	15	22	6	1	-
White, both sexes	4	3	5	4	13	13	20	15	15	5	1	-
Male	1	2	4	3	8	7	12	7	3	1	-	-
Female	3	1	1	1	5	6	8	8	12	4	1	-
Black, both sexes	7	3	8	8	13	9	13	10	12	4	-	-
Male	4	2	2	3	8	4	3	3	2	2	-	-
Female	3	1	6	5	5	5	10	7	10	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Maine												
All races, both sexes	-	1	1	1	2	4	13	9	1	3	-	-
Male	-	-	-	1	-	3	7	4	-	1	-	-
Female	-	1	1	-	2	1	6	5	1	2	-	-
White, both sexes	-	1	1	1	2	4	13	9	1	3	-	-
Male	-	-	-	1	-	3	7	4	-	1	-	-
Female	-	1	1	-	2	1	6	5	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	7	6	10	7	23	33	38	32	20	13	3	-
Male	5	4	7	3	14	17	14	11	7	2	-	-
Female	2	2	3	4	9	16	24	21	13	11	3	-
White, both sexes	-	-	-	1	10	18	21	22	17	11	2	-
Male	-	-	-	1	8	10	10	7	6	2	-	-
Female	-	-	-	-	2	8	11	15	11	9	2	-
Black, both sexes	7	6	10	6	13	15	17	10	3	2	1	-
Male	5	4	7	2	6	7	4	4	1	-	-	-
Female	2	2	3	4	7	8	13	6	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Massachusetts												
All races, both sexes	2	2	3	4	6	22	32	25	20	3	1	-
Male	1	1	1	2	2	7	12	11	5	1	-	-
Female	1	1	2	2	4	15	20	14	15	2	1	-
White, both sexes	2	1	2	3	3	15	29	23	18	3	1	-
Male	1	1	1	1	2	4	12	10	5	1	-	-
Female	1	-	1	2	1	11	17	13	13	2	1	-
Black, both sexes	-	1	1	1	3	4	2	1	2	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	1	1	-	3	3	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	1	1	-	-	-	-
Male	-	-	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Michigan												
All races, both sexes	11	8	16	27	31	51	60	68	44	16	2	-
Male	8	3	10	15	15	22	35	22	17	3	-	-
Female	3	5	6	12	16	29	25	46	27	13	2	-
White, both sexes	2	2	9	16	18	33	45	55	38	10	2	-
Male	1	1	5	10	7	15	29	20	16	2	-	-
Female	1	1	4	6	11	18	16	35	22	8	2	-
Black, both sexes	9	6	6	10	13	18	15	13	6	4	-	-
Male	7	2	5	5	8	7	6	2	1	1	-	-
Female	2	4	1	5	5	11	9	11	5	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Minnesota												
All races, both sexes	2	3	4	6	6	24	26	38	34	17	1	-
Male	1	1	1	-	3	11	15	14	14	5	1	-
Female	1	2	3	6	3	13	11	24	20	12	-	-
White, both sexes	2	1	3	6	5	21	23	38	34	17	1	-
Male	1	-	1	-	3	10	14	14	14	5	1	-
Female	1	1	2	6	2	11	9	24	20	12	-	-
Black, both sexes	-	2	1	-	1	1	2	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	1	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Mississippi												
All races, both sexes	11	9	7	16	19	17	18	19	9	2	4	-
Male	6	5	4	7	9	5	6	5	-	-	1	-
Female	5	4	3	9	10	12	12	14	9	2	3	-
White, both sexes	3	1	3	2	4	4	8	6	7	1	1	-
Male	2	1	2	1	-	-	3	1	-	-	-	-
Female	1	-	1	1	4	4	5	5	7	1	1	-
Black, both sexes	8	8	4	14	15	13	10	13	2	1	3	-
Male	4	4	2	6	9	5	3	4	-	-	1	-
Female	4	4	2	8	6	8	7	9	2	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Missouri												
All races, both sexes	1	5	7	16	19	22	30	41	29	9	3	-
Male	-	5	4	11	10	8	10	17	8	-	1	-
Female	1	-	3	5	9	14	20	24	21	9	2	-
White, both sexes	-	1	4	6	15	16	26	39	23	7	2	-
Male	-	1	3	6	7	6	9	17	5	-	1	-
Female	-	-	1	-	8	10	17	22	18	7	1	-
Black, both sexes	1	4	3	10	4	6	4	2	6	2	1	-
Male	-	4	1	5	3	2	1	-	3	-	-	-
Female	1	-	2	5	1	4	3	2	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	-	4	4	3	8	3	3	-	-
Male	-	-	1	-	-	3	-	3	2	1	-	-
Female	-	-	-	-	4	1	3	5	1	2	-	-
White, both sexes	-	-	1	-	3	4	3	8	3	3	-	-
Male	-	-	1	-	-	3	-	3	2	1	-	-
Female	-	-	-	-	3	1	3	5	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Nebraska												
All races, both sexes	1	1	5	2	1	3	14	24	13	5	-	-
Male	1	1	2	2	-	1	8	7	2	2	-	-
Female	-	-	3	-	1	2	6	17	11	3	-	-
White, both sexes	1	1	1	1	1	1	12	23	13	4	-	-
Male	1	1	1	1	-	-	6	7	2	1	-	-
Female	-	-	-	-	1	1	6	16	11	3	-	-
Black, both sexes	-	-	4	1	-	2	2	1	-	1	-	-
Male	-	-	1	1	-	1	2	-	-	1	-	-
Female	-	-	3	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	2	1	5	5	4	5	6	3	1	-	-
Male	3	1	1	4	3	3	4	2	3	1	-	-
Female	-	1	-	1	2	1	1	4	-	-	-	-
White, both sexes	2	1	1	3	1	3	5	6	3	1	-	-
Male	2	1	1	3	1	3	4	2	3	1	-	-
Female	-	-	-	-	-	-	1	4	-	-	-	-
Black, both sexes	-	1	-	1	2	1	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
New Hampshire												
All races, both sexes	1	-	2	1	1	-	5	9	7	1	1	-
Male	1	-	-	1	-	-	3	3	1	-	1	-
Female	-	-	2	-	1	-	2	6	6	1	-	-
White, both sexes	1	-	2	1	1	-	5	9	7	1	1	-
Male	1	-	-	1	-	-	3	3	1	-	1	-
Female	-	-	2	-	1	-	2	6	6	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	10	11	18	13	21	31	33	40	31	10	-	-
Male	6	9	10	8	11	17	16	14	5	1	-	-
Female	4	2	8	5	10	14	17	26	26	9	-	-
White, both sexes	2	6	6	6	10	23	25	36	26	10	-	-
Male	1	5	4	4	5	12	13	12	4	1	-	-
Female	1	1	2	2	5	11	12	24	22	9	-	-
Black, both sexes	8	5	10	7	10	6	5	4	5	-	-	-
Male	5	4	5	4	5	4	2	2	1	-	-	-
Female	3	1	5	3	5	2	3	2	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	2	3	-	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	1	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
New Mexico												
All races, both sexes	-	-	2	2	5	7	5	8	7	1	-	-
Male	-	-	2	1	1	4	3	5	3	1	-	-
Female	-	-	-	1	4	3	2	3	4	-	-	-
White, both sexes	-	-	1	2	4	6	4	7	5	1	-	-
Male	-	-	1	1	1	3	3	5	2	1	-	-
Female	-	-	-	1	3	3	1	2	3	-	-	-
Black, both sexes	-	-	1	-	-	-	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	21	30	23	46	75	97	110	93	74	27	4	-
Male	13	20	5	23	33	45	50	42	20	8	-	-
Female	8	10	18	23	42	52	60	51	54	19	4	-
White, both sexes	6	11	7	22	43	70	79	76	61	25	3	-
Male	3	6	2	11	18	35	37	36	18	8	-	-
Female	3	5	5	11	25	35	42	40	43	17	3	-
Black, both sexes	14	18	14	23	28	25	26	15	12	2	1	-
Male	9	13	3	11	12	9	10	6	1	-	-	-
Female	5	5	11	12	16	16	16	9	11	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	1	4	2	5	2	1	-	-	-
Male	1	1	-	1	3	1	3	-	1	-	-	-
Female	-	-	2	-	1	1	2	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
North Carolina												
All races, both sexes	12	6	17	32	40	48	68	47	39	14	2	-
Male	11	5	11	19	21	17	26	15	16	3	-	-
Female	1	1	6	13	19	31	42	32	23	11	2	-
White, both sexes	1	2	8	12	19	22	41	29	24	10	-	-
Male	1	1	4	8	9	6	17	11	10	2	-	-
Female	-	1	4	4	10	16	24	18	14	8	-	-
Black, both sexes	11	4	9	20	19	26	27	17	14	4	2	-
Male	10	4	7	11	10	11	9	4	5	1	-	-
Female	1	-	2	9	9	15	18	13	9	3	2	-
American Indian 1/, both sexes	-	-	-	-	2	-	-	1	1	-	-	-
Male	-	-	-	-	2	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	1	1	1	6	4	8	10	7	-	-
Male	-	-	1	-	-	2	3	5	4	-	-	-
Female	-	-	-	1	1	4	1	3	6	7	-	-
White, both sexes	-	-	-	1	1	5	4	8	10	7	-	-
Male	-	-	-	-	-	2	3	5	4	-	-	-
Female	-	-	-	1	1	3	1	3	6	7	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Ohio												
All races, both sexes	7	13	16	27	38	69	108	101	58	29	8	-
Male	2	6	10	13	20	33	45	45	16	8	1	-
Female	5	7	6	14	18	36	63	56	42	21	7	-
White, both sexes	2	9	6	16	28	48	89	86	48	22	6	-
Male	-	5	2	10	16	22	41	37	14	7	-	-
Female	2	4	4	6	12	26	48	49	34	15	6	-
Black, both sexes	5	4	10	10	10	21	19	14	10	7	2	-
Male	2	1	8	3	4	11	4	7	2	1	1	-
Female	3	3	2	7	6	10	15	7	8	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	6	4	13	10	23	28	30	14	1	1	-
Male	3	4	2	8	4	16	10	8	6	-	-	-
Female	-	2	2	5	6	7	18	22	8	1	1	-
White, both sexes	2	2	4	8	6	13	21	26	12	1	1	-
Male	2	2	2	4	2	9	8	7	5	-	-	-
Female	-	-	2	4	4	4	13	19	7	1	1	-
Black, both sexes	1	3	-	2	2	4	5	3	1	-	-	-
Male	1	2	-	2	1	3	1	1	-	-	-	-
Female	-	1	-	-	1	1	4	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	3	2	5	2	1	1	-	-	-
Male	-	-	-	2	1	4	1	-	1	-	-	-
Female	-	-	-	1	1	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Oregon												
All races, both sexes	-	1	1	4	8	10	28	32	32	9	-	-
Male	-	1	1	3	4	6	10	16	11	3	-	-
Female	-	-	-	1	4	4	18	16	21	6	-	-
White, both sexes	-	-	1	4	7	9	24	31	30	9	-	-
Male	-	-	1	3	4	5	9	15	11	3	-	-
Female	-	-	-	1	3	4	15	16	19	6	-	-
Black, both sexes	-	1	-	-	1	1	2	-	1	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Pennsylvania												
All races, both sexes	11	12	15	15	22	56	70	61	39	25	1	-
Male	5	7	9	7	10	26	27	30	13	8	-	-
Female	6	5	6	8	12	30	43	31	26	17	1	-
White, both sexes	6	7	6	10	15	51	64	53	32	24	1	-
Male	3	4	5	4	7	23	24	28	11	7	-	-
Female	3	3	1	6	8	28	40	25	21	17	1	-
Black, both sexes	5	5	9	5	7	5	6	7	6	1	-	-
Male	2	3	4	3	3	3	3	2	2	1	-	-
Female	3	2	5	2	4	2	3	5	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Rhode Island												
All races, both sexes	1	1	3	2	6	4	9	3	5	-	-	-
Male	1	1	2	-	2	1	6	2	1	-	-	-
Female	-	-	1	2	4	3	3	1	4	-	-	-
White, both sexes	1	1	2	2	5	4	8	3	5	-	-	-
Male	1	1	1	-	1	1	5	2	1	-	-	-
Female	-	-	1	2	4	3	3	1	4	-	-	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	12	15	7	14	23	27	33	24	14	9	1	-
Male	6	11	3	8	14	11	11	8	3	4	-	-
Female	6	4	4	6	9	16	22	16	11	5	1	-
White, both sexes	3	1	1	4	8	8	16	13	8	5	1	-
Male	-	1	-	3	4	3	7	6	1	1	-	-
Female	3	-	1	1	4	5	9	7	7	4	1	-
Black, both sexes	9	13	6	10	15	18	17	11	6	4	-	-
Male	6	9	3	5	10	7	4	2	2	3	-	-
Female	3	4	3	5	5	11	13	9	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
South Dakota												
All races, both sexes	-	-	-	2	4	5	6	13	10	5	1	-
Male	-	-	-	2	4	3	4	7	5	1	-	-
Female	-	-	-	-	-	2	2	6	5	4	1	-
White, both sexes	-	-	-	-	3	4	6	13	10	5	1	-
Male	-	-	-	-	3	2	4	7	5	1	-	-
Female	-	-	-	-	-	2	2	6	5	4	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	1	-	-	-	-	-	-	-
Male	-	-	-	2	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	10	14	21	29	20	39	64	59	28	11	-	-
Male	7	11	14	12	10	23	22	18	11	2	-	-
Female	3	3	7	17	10	16	42	41	17	9	-	-
White, both sexes	2	10	6	14	9	28	51	45	19	10	-	-
Male	1	8	5	8	5	16	16	17	6	2	-	-
Female	1	2	1	6	4	12	35	28	13	8	-	-
Black, both sexes	8	4	14	15	9	10	13	14	9	1	-	-
Male	6	3	8	4	3	7	6	1	5	-	-	-
Female	2	1	6	11	6	3	7	13	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	1	-	-	-	-	-	-
Male	-	-	1	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Texas												
All races, both sexes	26	29	36	58	76	98	131	121	74	28	5	-
Male	17	16	15	27	29	32	54	44	27	7	-	-
Female	9	13	21	31	47	66	77	77	47	21	5	-
White, both sexes	11	20	23	39	52	78	107	94	60	24	3	-
Male	7	12	8	15	18	26	46	36	24	6	-	-
Female	4	8	15	24	34	52	61	58	36	18	3	-
Black, both sexes	15	9	13	16	22	19	24	27	13	4	2	-
Male	10	4	7	11	10	5	8	8	3	1	-	-
Female	5	5	6	5	12	14	16	19	10	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	2	1	-	-	1	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	-	2	1	-	-	-	1	-	-	-
Utah												
All races, both sexes	1	1	-	1	1	5	4	5	3	3	-	-
Male	-	-	-	1	1	3	2	3	1	-	-	-
Female	1	1	-	-	-	2	2	2	2	3	-	-
White, both sexes	-	1	-	1	1	5	4	5	3	3	-	-
Male	-	-	-	1	1	3	2	3	1	-	-	-
Female	-	1	-	-	-	2	2	2	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Vermont												
All races, both sexes	-	1	-	-	3	3	-	5	5	1	-	-
Male	-	1	-	-	-	1	-	2	1	1	-	-
Female	-	-	-	-	3	2	-	3	4	-	-	-
White, both sexes	-	1	-	-	3	3	-	5	5	1	-	-
Male	-	1	-	-	-	1	-	2	1	1	-	-
Female	-	-	-	-	3	2	-	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	7	10	17	7	26	39	30	36	30	6	2	-
Male	5	7	7	4	9	19	17	15	7	-	-	-
Female	2	3	10	3	17	20	13	21	23	6	2	-
White, both sexes	2	1	4	2	15	22	15	23	14	4	-	-
Male	2	-	2	1	6	12	8	10	2	-	-	-
Female	-	1	2	1	9	10	7	13	12	4	-	-
Black, both sexes	5	9	13	5	10	15	14	12	16	2	2	-
Male	3	7	5	3	3	7	9	4	5	-	-	-
Female	2	2	8	2	7	8	5	8	11	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	2	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Washington												
All races, both sexes	3	6	5	4	12	25	41	35	26	12	1	-
Male	3	4	1	1	6	13	15	19	9	3	-	-
Female	-	2	4	3	6	12	26	16	17	9	1	-
White, both sexes	2	6	4	3	7	23	38	34	24	12	1	-
Male	2	4	-	-	3	12	14	18	9	3	-	-
Female	-	2	4	3	4	11	24	16	15	9	1	-
Black, both sexes	1	-	1	1	2	-	-	-	-	-	-	-
Male	1	-	1	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	2	-	-	-	-	-
Male	-	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	1	2	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	2	-	-	-
West Virginia												
All races, both sexes	3	2	6	6	3	6	16	11	12	5	1	-
Male	1	1	4	4	1	3	7	1	4	-	-	-
Female	2	1	2	2	2	3	9	10	8	5	1	-
White, both sexes	3	2	6	6	3	5	15	11	12	4	1	-
Male	1	1	4	4	1	3	7	1	4	-	-	-
Female	2	1	2	2	2	2	8	10	8	4	1	-
Black, both sexes	-	-	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive renal disease (I12)												
Wisconsin												
All races, both sexes	2	2	3	11	17	23	31	48	34	10	3	-
Male	2	-	2	4	9	11	17	19	12	-	1	-
Female	-	2	1	7	8	12	14	29	22	10	2	-
White, both sexes	1	2	2	10	17	23	29	47	32	10	3	-
Male	1	-	2	4	9	11	16	19	12	-	1	-
Female	-	2	-	6	8	12	13	28	20	10	2	-
Black, both sexes	1	-	-	-	-	-	2	1	2	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	1	-	3	1	2	1	-	-	-
Male	-	-	-	-	-	3	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	2	1	-	-	-
White, both sexes	-	-	-	1	-	3	1	2	1	-	-	-
Male	-	-	-	-	-	3	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
United States												
All races, both sexes	3,172	-	-	-	-	-	5	6	27	31	52	73
Male	1,369	-	-	-	-	-	4	2	18	22	30	54
Female	1,803	-	-	-	-	-	1	4	9	9	22	19
White, both sexes	2,186	-	-	-	-	-	2	1	6	11	17	25
Male	886	-	-	-	-	-	2	1	3	8	11	17
Female	1,300	-	-	-	-	-	-	-	3	3	6	8
Black, both sexes	889	-	-	-	-	-	3	5	20	20	34	46
Male	432	-	-	-	-	-	2	1	15	14	19	36
Female	457	-	-	-	-	-	1	4	5	6	15	10
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	85	-	-	-	-	-	-	-	-	-	1	2
Male	44	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	1	1
Alabama												
All races, both sexes	42	-	-	-	-	-	-	-	-	1	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	19	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Alaska												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	1	2
Male	18	-	-	-	-	-	-	-	-	-	1	2
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	2
Male	15	-	-	-	-	-	-	-	-	-	-	2
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Arkansas												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	515	-	-	-	-	-	1	-	2	7	7	11
Male	231	-	-	-	-	-	1	-	1	6	3	8
Female	284	-	-	-	-	-	-	-	1	1	4	3
White, both sexes	373	-	-	-	-	-	1	-	2	2	3	9
Male	161	-	-	-	-	-	1	-	1	2	1	7
Female	212	-	-	-	-	-	-	-	1	-	2	2
Black, both sexes	86	-	-	-	-	-	-	-	-	5	3	2
Male	41	-	-	-	-	-	-	-	-	4	2	1
Female	45	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	55	-	-	-	-	-	-	-	-	-	1	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Colorado												
All races, both sexes	25	-	-	-	-	-	-	1	-	-	-	-
Male	11	-	-	-	-	-	-	1	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	29	-	-	-	-	-	-	-	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Delaware												
All races, both sexes	10	-	-	-	-	-	1	-	1	-	-	1
Male	5	-	-	-	-	-	1	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	7	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	29	-	-	-	-	-	-	-	1	-	1	1
Male	14	-	-	-	-	-	-	-	1	-	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	28	-	-	-	-	-	-	-	1	-	1	1
Male	13	-	-	-	-	-	-	-	1	-	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Florida												
All races, both sexes	218	-	-	-	-	-	-	-	1	2	3	5
Male	94	-	-	-	-	-	-	-	1	2	1	3
Female	124	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	159	-	-	-	-	-	-	-	-	1	2	3
Male	65	-	-	-	-	-	-	-	-	1	1	2
Female	94	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	57	-	-	-	-	-	-	-	1	1	1	2
Male	28	-	-	-	-	-	-	-	1	1	-	1
Female	29	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	98	-	-	-	-	-	-	-	1	1	2	2
Male	45	-	-	-	-	-	-	-	1	1	1	1
Female	53	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	57	-	-	-	-	-	-	-	1	1	2	1
Male	28	-	-	-	-	-	-	-	1	1	1	1
Female	29	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Hawaii												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Illinois												
All races, both sexes	127	-	-	-	-	-	-	1	2	-	2	3
Male	58	-	-	-	-	-	-	-	2	-	-	3
Female	69	-	-	-	-	-	-	1	-	-	2	-
White, both sexes	80	-	-	-	-	-	-	-	-	-	-	1
Male	36	-	-	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	46	-	-	-	-	-	-	1	2	-	2	1
Male	21	-	-	-	-	-	-	-	2	-	-	1
Female	25	-	-	-	-	-	-	1	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	55	-	-	-	-	-	-	-	-	-	2	-
Male	19	-	-	-	-	-	-	-	-	-	2	-
Female	36	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	47	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Iowa												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	2	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	29	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	44	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Kentucky												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	32	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	73	-	-	-	-	-	-	-	-	2	3	3
Male	38	-	-	-	-	-	-	-	-	1	2	2
Female	35	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	44	-	-	-	-	-	-	-	-	2	3	3
Male	24	-	-	-	-	-	-	-	-	1	2	2
Female	20	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Maine												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	67	-	-	-	-	-	-	-	3	2	4	4
Male	38	-	-	-	-	-	-	-	3	2	3	4
Female	29	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	30	-	-	-	-	-	-	-	-	2	2	-
Male	15	-	-	-	-	-	-	-	-	2	1	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	37	-	-	-	-	-	-	-	3	-	2	4
Male	23	-	-	-	-	-	-	-	3	-	2	4
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Massachusetts												
All races, both sexes	25	-	-	-	-	-	-	-	-	1	1	1
Male	15	-	-	-	-	-	-	-	-	1	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	1	1	1
Male	3	-	-	-	-	-	-	-	-	1	1	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	128	-	-	-	-	-	1	1	3	-	3	6
Male	53	-	-	-	-	-	1	-	2	-	2	1
Female	75	-	-	-	-	-	-	1	1	-	1	5
White, both sexes	89	-	-	-	-	-	-	-	-	-	2	1
Male	37	-	-	-	-	-	-	-	-	-	1	-
Female	52	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	39	-	-	-	-	-	1	1	3	-	1	5
Male	16	-	-	-	-	-	1	-	2	-	1	1
Female	23	-	-	-	-	-	-	1	1	-	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Minnesota												
All races, both sexes	36	-	-	-	-	-	-	-	1	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	67	-	-	-	-	-	-	1	-	1	1	2
Male	27	-	-	-	-	-	-	-	-	1	1	1
Female	40	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	29	-	-	-	-	-	-	-	-	1	-	1
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	38	-	-	-	-	-	-	1	-	-	1	1
Male	16	-	-	-	-	-	-	-	-	-	1	1
Female	22	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Missouri												
All races, both sexes	50	-	-	-	-	-	1	1	1	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	35	-	-	-	-	-	1	1	1	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	1	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	16	-	-	-	-	-	1	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Nebraska												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	18	-	-	-	-	-	1	-	1	-	-	-
Male	9	-	-	-	-	-	1	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	1	-	1	-	-	-
Male	4	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
New Hampshire												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	62	-	-	-	-	-	-	-	-	-	2	-
Male	27	-	-	-	-	-	-	-	-	-	1	-
Female	35	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	19	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
New Mexico												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	214	-	-	-	-	-	-	-	-	1	6	6
Male	101	-	-	-	-	-	-	-	-	1	2	5
Female	113	-	-	-	-	-	-	-	-	-	4	1
White, both sexes	126	-	-	-	-	-	-	-	-	-	1	-
Male	57	-	-	-	-	-	-	-	-	-	1	-
Female	69	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	79	-	-	-	-	-	-	-	-	1	5	5
Male	40	-	-	-	-	-	-	-	-	1	1	5
Female	39	-	-	-	-	-	-	-	-	-	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
North Carolina												
All races, both sexes	100	-	-	-	-	-	-	-	4	2	-	4
Male	53	-	-	-	-	-	-	-	3	2	-	3
Female	47	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	53	-	-	-	-	-	-	-	1	1	-	1
Male	23	-	-	-	-	-	-	-	1	1	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	45	-	-	-	-	-	-	-	3	1	-	3
Male	29	-	-	-	-	-	-	-	2	1	-	3
Female	16	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Ohio												
All races, both sexes	167	-	-	-	-	-	-	-	-	2	1	5
Male	68	-	-	-	-	-	-	-	-	-	1	4
Female	99	-	-	-	-	-	-	-	-	2	-	1
White, both sexes	131	-	-	-	-	-	-	-	-	1	-	2
Male	52	-	-	-	-	-	-	-	-	-	-	1
Female	79	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	35	-	-	-	-	-	-	-	-	1	1	3
Male	16	-	-	-	-	-	-	-	-	-	1	3
Female	19	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	29	-	-	-	-	-	-	-	1	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Oregon												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	107	-	-	-	-	-	-	-	1	2	1	1
Male	42	-	-	-	-	-	-	-	-	-	1	1
Female	65	-	-	-	-	-	-	-	1	2	-	-
White, both sexes	88	-	-	-	-	-	-	-	-	1	1	1
Male	32	-	-	-	-	-	-	-	-	-	1	1
Female	56	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	18	-	-	-	-	-	-	-	1	1	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Rhode Island												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	46	-	-	-	-	-	-	-	1	1	-	1
Male	19	-	-	-	-	-	-	-	1	1	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	1	-	-
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	24	-	-	-	-	-	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	1	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
South Dakota												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	100	-	-	-	-	-	-	1	-	-	1	1
Male	35	-	-	-	-	-	-	1	-	-	1	1
Female	65	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	67	-	-	-	-	-	-	1	-	-	-	-
Male	22	-	-	-	-	-	-	1	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	33	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Texas												
All races, both sexes	195	-	-	-	-	-	-	-	1	4	3	7
Male	75	-	-	-	-	-	-	-	1	2	1	5
Female	120	-	-	-	-	-	-	-	-	2	2	2
White, both sexes	134	-	-	-	-	-	-	-	1	1	-	-
Male	49	-	-	-	-	-	-	-	1	-	-	-
Female	85	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	60	-	-	-	-	-	-	-	-	3	3	7
Male	26	-	-	-	-	-	-	-	-	2	1	5
Female	34	-	-	-	-	-	-	-	-	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Vermont												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	62	-	-	-	-	-	-	-	-	1	2	2
Male	27	-	-	-	-	-	-	-	-	1	2	1
Female	35	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	34	-	-	-	-	-	-	-	-	1	1	2
Male	20	-	-	-	-	-	-	-	-	1	1	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Washington												
All races, both sexes	71	-	-	-	-	-	-	-	1	-	-	-
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	64	-	-	-	-	-	-	-	1	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Hypertensive heart and renal disease (I13)												
Wisconsin												
All races, both sexes	48	-	-	-	-	-	-	-	1	-	1	1
Male	21	-	-	-	-	-	-	-	1	-	1	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	1	-
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	1	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
United States												
All races, both sexes	116	111	129	183	243	373	535	581	462	211	34	-
Male	71	66	76	97	112	166	212	225	149	57	8	-
Female	45	45	53	86	131	207	323	356	313	154	26	-
White, both sexes	36	42	52	93	147	256	402	484	405	182	25	-
Male	28	28	29	45	71	116	159	184	130	48	6	-
Female	8	14	23	48	76	140	243	300	275	134	19	-
Black, both sexes	78	67	71	86	83	99	117	81	48	23	8	-
Male	42	38	44	49	33	38	47	32	14	6	2	-
Female	36	29	27	37	50	61	70	49	34	17	6	-
American Indian 1/, both sexes	1	1	1	-	2	1	2	1	1	1	-	-
Male	-	-	1	-	2	1	1	1	1	-	-	-
Female	1	1	-	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	1	5	4	11	17	14	15	8	5	1	-
Male	1	-	2	3	6	11	5	8	4	3	-	-
Female	-	1	3	1	5	6	9	7	4	2	1	-
Alabama												
All races, both sexes	1	3	1	1	5	5	9	9	4	3	-	-
Male	1	2	-	-	2	2	2	3	1	-	-	-
Female	-	1	1	1	3	3	7	6	3	3	-	-
White, both sexes	-	2	1	1	1	4	3	6	3	2	-	-
Male	-	1	-	-	1	1	1	2	-	-	-	-
Female	-	1	1	1	-	3	2	4	3	2	-	-
Black, both sexes	1	1	-	-	4	1	6	3	1	1	-	-
Male	1	1	-	-	1	1	1	1	1	-	-	-
Female	-	-	-	-	3	-	5	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Alaska												
All races, both sexes	-	-	-	1	1	1	-	-	-	1	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	2	2	3	3	3	4	12	6	-	1	-
Male	-	1	2	2	-	2	1	4	3	-	-	-
Female	-	1	-	1	3	1	3	8	3	-	1	-
White, both sexes	-	1	2	2	3	3	3	12	6	-	1	-
Male	-	1	2	1	-	2	-	4	3	-	-	-
Female	-	-	-	1	3	1	3	8	3	-	1	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Arkansas												
All races, both sexes	2	2	2	3	3	4	7	5	4	2	-	-
Male	1	1	1	2	2	1	3	-	1	-	-	-
Female	1	1	1	1	1	3	4	5	3	2	-	-
White, both sexes	1	-	1	3	1	2	6	4	2	1	-	-
Male	1	-	-	2	1	1	3	-	-	-	-	-
Female	-	-	1	1	-	1	3	4	2	1	-	-
Black, both sexes	1	2	1	-	2	2	1	1	2	1	-	-
Male	-	1	1	-	1	-	-	-	1	-	-	-
Female	1	1	-	-	1	2	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	17	26	18	26	43	67	84	85	84	28	9	-
Male	14	14	9	13	22	36	32	34	26	10	2	-
Female	3	12	9	13	21	31	52	51	58	18	7	-
White, both sexes	9	19	9	15	23	43	67	72	71	21	7	-
Male	7	13	4	8	11	21	26	30	21	7	1	-
Female	2	6	5	7	12	22	41	42	50	14	6	-
Black, both sexes	7	6	8	9	10	14	7	5	6	2	2	-
Male	6	1	4	4	6	7	4	-	1	-	1	-
Female	1	5	4	5	4	7	3	5	5	2	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	2	9	10	10	8	7	5	-	-
Male	1	-	1	1	4	8	2	4	4	3	-	-
Female	-	1	-	1	5	2	8	4	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Colorado												
All races, both sexes	-	2	2	2	1	1	3	5	5	2	1	-
Male	-	2	1	2	1	-	1	-	2	1	-	-
Female	-	-	1	-	-	1	2	5	3	1	1	-
White, both sexes	-	1	2	2	1	1	2	4	5	2	1	-
Male	-	1	1	2	1	-	-	-	2	1	-	-
Female	-	-	1	-	-	1	2	4	3	1	1	-
Black, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	1	1	4	2	-	3	6	4	6	1	-
Male	-	1	1	1	-	-	2	2	2	1	-	-
Female	-	-	-	3	2	-	1	4	2	5	1	-
White, both sexes	-	-	-	4	1	-	3	5	4	6	1	-
Male	-	-	-	1	-	-	2	1	2	1	-	-
Female	-	-	-	3	1	-	1	4	2	5	1	-
Black, both sexes	-	1	1	-	1	-	-	1	-	-	-	-
Male	-	1	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Delaware												
All races, both sexes	-	1	-	-	1	2	1	-	-	2	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	2	-	-
White, both sexes	-	1	-	-	1	2	1	-	-	1	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	1	3	4	2	3	5	4	1	-	-	-
Male	1	-	1	3	1	-	2	2	1	-	-	-
Female	2	1	2	1	1	3	3	2	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	1	3	4	2	3	4	4	1	-	-	-
Male	1	-	1	3	1	-	1	2	1	-	-	-
Female	2	1	2	1	1	3	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Florida												
All races, both sexes	8	3	9	14	22	31	39	44	20	15	2	-
Male	4	2	6	7	12	11	19	17	5	3	1	-
Female	4	1	3	7	10	20	20	27	15	12	1	-
White, both sexes	1	-	3	8	16	26	34	35	14	14	2	-
Male	1	-	2	4	8	10	17	13	3	2	1	-
Female	-	-	1	4	8	16	17	22	11	12	1	-
Black, both sexes	7	3	5	6	6	5	5	9	5	1	-	-
Male	3	2	3	3	4	1	2	4	2	1	-	-
Female	4	1	2	3	2	4	3	5	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Georgia												
All races, both sexes	6	8	2	9	9	13	19	15	5	6	-	-
Male	1	5	2	4	8	5	6	7	1	2	-	-
Female	5	3	-	5	1	8	13	8	4	4	-	-
White, both sexes	-	1	-	2	6	5	7	11	3	5	-	-
Male	-	-	-	1	5	2	2	5	-	2	-	-
Female	-	1	-	1	1	3	5	6	3	3	-	-
Black, both sexes	6	7	2	7	3	8	12	4	2	1	-	-
Male	1	5	2	3	3	3	4	2	1	-	-	-
Female	5	2	-	4	-	5	8	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Hawaii												
All races, both sexes	-	-	1	1	-	2	-	4	1	-	-	-
Male	-	-	-	1	-	-	-	3	-	-	-	-
Female	-	-	1	-	-	2	-	1	1	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	-	4	-	-	-	-
Male	-	-	-	1	-	-	-	3	-	-	-	-
Female	-	-	-	-	-	2	-	1	-	-	-	-
Idaho												
All races, both sexes	-	-	1	-	-	1	2	6	1	-	-	-
Male	-	-	1	-	-	-	-	5	1	-	-	-
Female	-	-	-	-	-	1	2	1	-	-	-	-
White, both sexes	-	-	1	-	-	1	2	6	1	-	-	-
Male	-	-	1	-	-	-	-	5	1	-	-	-
Female	-	-	-	-	-	1	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Illinois												
All races, both sexes	4	11	5	9	8	9	14	30	18	9	2	-
Male	1	6	3	5	5	4	7	11	6	3	2	-
Female	3	5	2	4	3	5	7	19	12	6	-	-
White, both sexes	1	-	1	2	6	7	11	25	17	8	1	-
Male	-	-	1	-	4	4	6	10	6	3	1	-
Female	1	-	-	2	2	3	5	15	11	5	-	-
Black, both sexes	3	11	4	7	2	2	3	5	1	1	1	-
Male	1	6	2	5	1	-	1	1	-	-	1	-
Female	2	5	2	2	1	2	2	4	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	2	3	2	5	5	9	16	8	3	-	-
Male	-	2	2	1	1	3	2	5	1	-	-	-
Female	-	-	1	1	4	2	7	11	7	3	-	-
White, both sexes	-	1	3	2	2	5	6	16	8	3	-	-
Male	-	1	2	1	1	3	-	5	1	-	-	-
Female	-	-	1	1	1	2	6	11	7	3	-	-
Black, both sexes	-	1	-	-	3	-	3	-	-	-	-	-
Male	-	1	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	3	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Iowa												
All races, both sexes	-	1	1	2	1	2	6	10	17	5	1	-
Male	-	-	1	2	1	1	4	1	6	2	-	-
Female	-	1	-	-	-	1	2	9	11	3	1	-
White, both sexes	-	1	-	2	-	2	6	10	17	5	1	-
Male	-	-	-	2	-	1	4	1	6	2	-	-
Female	-	1	-	-	-	1	2	9	11	3	1	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	-	-	-	-	1	3	5	5	2	2	-
Male	-	-	-	-	-	1	-	1	2	2	-	-
Female	1	-	-	-	-	-	3	4	3	-	2	-
White, both sexes	-	-	-	-	-	1	3	5	5	2	2	-
Male	-	-	-	-	-	1	-	1	2	2	-	-
Female	-	-	-	-	-	-	3	4	3	-	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Kentucky												
All races, both sexes	-	2	2	1	2	7	6	6	3	5	1	-
Male	-	-	1	-	1	2	2	1	2	1	-	-
Female	-	2	1	1	1	5	4	5	1	4	1	-
White, both sexes	-	1	2	1	2	5	5	6	3	5	1	-
Male	-	-	1	-	1	2	2	1	2	1	-	-
Female	-	1	1	1	1	3	3	5	1	4	1	-
Black, both sexes	-	1	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	5	3	3	5	6	13	13	6	7	4	-	-
Male	4	2	1	4	2	5	6	6	1	2	-	-
Female	1	1	2	1	4	8	7	-	6	2	-	-
White, both sexes	2	1	-	2	1	6	6	4	5	2	-	-
Male	1	-	-	2	1	1	3	4	1	1	-	-
Female	1	1	-	-	-	5	3	-	4	1	-	-
Black, both sexes	3	2	3	3	5	7	7	2	2	2	-	-
Male	3	2	1	2	1	4	3	2	-	1	-	-
Female	-	-	2	1	4	3	4	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Maine												
All races, both sexes	-	-	-	-	2	2	-	-	1	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	2	-	-	1	2	-	-
White, both sexes	-	-	-	-	1	2	-	-	1	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	1	2	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	4	5	2	4	6	6	12	9	4	2	-	-
Male	3	4	2	2	3	4	5	2	-	1	-	-
Female	1	1	-	2	3	2	7	7	4	1	-	-
White, both sexes	-	2	-	1	3	2	9	4	3	2	-	-
Male	-	2	-	1	1	2	5	-	-	1	-	-
Female	-	-	-	-	2	-	4	4	3	1	-	-
Black, both sexes	4	3	2	3	3	4	3	5	1	-	-	-
Male	3	2	2	1	2	2	-	2	-	-	-	-
Female	1	1	-	2	1	2	3	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Massachusetts												
All races, both sexes	1	-	-	1	2	-	3	7	7	1	-	-
Male	1	-	-	1	1	-	1	5	2	1	-	-
Female	-	-	-	-	1	-	2	2	5	-	-	-
White, both sexes	1	-	-	1	2	-	3	6	7	1	-	-
Male	1	-	-	1	1	-	1	4	2	1	-	-
Female	-	-	-	-	1	-	2	2	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	8	2	4	3	12	7	24	21	21	11	1	-
Male	3	1	3	-	4	6	8	12	9	-	1	-
Female	5	1	1	3	8	1	16	9	12	11	-	-
White, both sexes	1	2	-	2	7	5	19	19	20	10	1	-
Male	-	1	-	-	4	4	7	11	8	-	1	-
Female	1	1	-	2	3	1	12	8	12	10	-	-
Black, both sexes	7	-	4	1	5	2	5	2	1	1	-	-
Male	3	-	3	-	-	2	1	1	1	-	-	-
Female	4	-	1	1	5	-	4	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Minnesota												
All races, both sexes	1	-	1	-	3	3	6	9	9	3	-	-
Male	1	-	-	-	2	2	3	3	6	1	-	-
Female	-	-	1	-	1	1	3	6	3	2	-	-
White, both sexes	1	-	1	-	3	3	6	9	9	3	-	-
Male	1	-	-	-	2	2	3	3	6	1	-	-
Female	-	-	1	-	1	1	3	6	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	1	4	9	4	9	15	8	4	5	2	-
Male	1	1	3	3	-	6	3	3	1	3	-	-
Female	-	-	1	6	4	3	12	5	3	2	2	-
White, both sexes	-	-	1	3	-	4	7	6	3	3	-	-
Male	-	-	1	1	-	2	2	2	-	2	-	-
Female	-	-	-	2	-	2	5	4	3	1	-	-
Black, both sexes	1	1	3	6	4	5	8	2	1	2	2	-
Male	1	1	2	2	-	4	1	1	1	1	-	-
Female	-	-	1	4	4	1	7	1	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Missouri												
All races, both sexes	1	2	3	2	1	3	13	10	5	6	-	-
Male	1	1	3	-	-	-	3	3	2	1	-	-
Female	-	1	-	2	1	3	10	7	3	5	-	-
White, both sexes	1	1	3	2	1	2	8	8	2	4	-	-
Male	1	1	3	-	-	-	1	3	1	1	-	-
Female	-	-	-	2	1	2	7	5	1	3	-	-
Black, both sexes	-	1	-	-	-	1	5	2	3	2	-	-
Male	-	-	-	-	-	-	2	-	1	-	-	-
Female	-	1	-	-	-	1	3	2	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	1	-	1	3	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	1	-	1	2	1	-	-	-
White, both sexes	1	-	-	-	1	-	1	3	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	1	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Nebraska												
All races, both sexes	1	-	-	-	1	1	4	3	3	1	-	-
Male	-	-	-	-	-	-	2	2	2	-	-	-
Female	1	-	-	-	1	1	2	1	1	1	-	-
White, both sexes	-	-	-	-	1	1	4	3	3	1	-	-
Male	-	-	-	-	-	-	2	2	2	-	-	-
Female	-	-	-	-	1	1	2	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	1	2	4	2	3	3	1	-	-	-
Male	-	-	-	1	3	-	1	1	1	-	-	-
Female	-	-	1	1	1	2	2	2	-	-	-	-
White, both sexes	-	-	1	2	3	2	1	2	1	-	-	-
Male	-	-	-	1	2	-	-	-	1	-	-	-
Female	-	-	1	1	1	2	1	2	-	-	-	-
Black, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
New Hampshire												
All races, both sexes	-	-	-	1	-	-	-	-	4	-	-	-
Male	-	-	-	1	-	-	-	-	3	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	4	-	-	-
Male	-	-	-	1	-	-	-	-	3	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4	-	2	4	5	6	11	12	10	5	1	-
Male	4	-	1	2	-	2	6	5	3	2	1	-
Female	-	-	1	2	5	4	5	7	7	3	-	-
White, both sexes	2	-	1	1	1	5	7	11	9	4	1	-
Male	2	-	-	-	-	2	3	4	3	1	1	-
Female	-	-	1	1	1	3	4	7	6	3	-	-
Black, both sexes	2	-	1	3	4	1	4	1	1	1	-	-
Male	2	-	1	2	-	-	3	1	-	1	-	-
Female	-	-	-	1	4	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
New Mexico												
All races, both sexes	2	-	-	-	2	3	3	3	2	1	-	-
Male	1	-	-	-	1	1	-	-	-	1	-	-
Female	1	-	-	-	1	2	3	3	2	-	-	-
White, both sexes	2	-	-	-	1	3	3	3	2	1	-	-
Male	1	-	-	-	-	1	-	-	-	1	-	-
Female	1	-	-	-	1	2	3	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	11	9	21	15	17	26	32	35	21	13	1	-
Male	8	5	9	8	3	11	20	17	7	5	-	-
Female	3	4	12	7	14	15	12	18	14	8	1	-
White, both sexes	4	1	6	5	12	18	23	25	18	12	1	-
Male	4	1	2	2	1	7	15	12	7	5	-	-
Female	-	-	4	3	11	11	8	13	11	7	1	-
Black, both sexes	7	8	13	10	4	5	8	9	3	1	-	-
Male	4	4	7	6	1	2	4	5	-	-	-	-
Female	3	4	6	4	3	3	4	4	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	3	1	1	-	-	-	-
Male	-	-	-	-	1	2	1	-	-	-	-	-
Female	-	-	2	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
North Carolina												
All races, both sexes	5	3	4	7	7	10	13	13	25	3	-	-
Male	3	1	3	6	5	3	7	4	11	2	-	-
Female	2	2	1	1	2	7	6	9	14	1	-	-
White, both sexes	1	1	1	3	4	3	6	10	19	2	-	-
Male	1	-	1	2	2	-	2	2	9	2	-	-
Female	-	1	-	1	2	3	4	8	10	-	-	-
Black, both sexes	4	2	3	4	3	6	6	3	6	1	-	-
Male	2	1	2	4	3	2	5	2	2	-	-	-
Female	2	1	1	-	-	4	1	1	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	2	3	2	-	-	-
Male	-	-	-	-	-	-	2	2	1	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	2	3	2	-	-	-
Male	-	-	-	-	-	-	2	2	1	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Ohio												
All races, both sexes	2	4	4	8	11	22	25	38	28	15	2	-
Male	1	3	3	3	5	14	9	12	11	2	-	-
Female	1	1	1	5	6	8	16	26	17	13	2	-
White, both sexes	-	1	1	4	6	20	19	36	26	14	1	-
Male	-	1	1	1	3	13	7	12	11	2	-	-
Female	-	-	-	3	3	7	12	24	15	12	1	-
Black, both sexes	2	3	2	4	5	2	6	2	2	1	1	-
Male	1	2	2	2	2	1	2	-	-	-	-	-
Female	1	1	-	2	3	1	4	2	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	-	2	2	3	1	1	4	8	5	1	-	-
Male	-	2	2	2	1	-	2	7	3	-	-	-
Female	-	-	-	1	-	1	2	1	2	1	-	-
White, both sexes	-	-	1	2	1	1	3	7	4	1	-	-
Male	-	-	1	2	1	-	1	6	2	-	-	-
Female	-	-	-	-	-	1	2	1	2	1	-	-
Black, both sexes	-	2	-	1	-	-	1	1	-	-	-	-
Male	-	2	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Oregon												
All races, both sexes	-	-	1	1	1	8	4	8	10	4	-	-
Male	-	-	-	-	1	5	1	2	1	2	-	-
Female	-	-	1	1	-	3	3	6	9	2	-	-
White, both sexes	-	-	-	1	1	7	3	8	10	4	-	-
Male	-	-	-	-	1	5	1	2	1	2	-	-
Female	-	-	-	1	-	2	2	6	9	2	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	6	3	1	3	8	14	22	16	20	8	1	-
Male	4	3	1	2	2	6	7	7	6	2	-	-
Female	2	-	-	1	6	8	15	9	14	6	1	-
White, both sexes	2	2	-	2	7	10	18	15	20	8	1	-
Male	1	2	-	1	1	6	4	7	6	2	-	-
Female	1	-	-	1	6	4	14	8	14	6	1	-
Black, both sexes	4	1	1	1	1	4	3	1	-	-	-	-
Male	3	1	1	1	1	-	2	-	-	-	-	-
Female	1	-	-	-	-	4	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Rhode Island												
All races, both sexes	-	-	1	-	-	2	1	6	1	1	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	6	1	1	-	-
White, both sexes	-	-	1	-	-	2	1	6	1	1	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	6	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	-	1	4	4	6	9	8	6	3	-	-
Male	1	-	-	4	2	2	2	2	2	1	-	-
Female	1	-	1	-	2	4	7	6	4	2	-	-
White, both sexes	1	-	-	1	2	2	6	5	3	1	-	-
Male	1	-	-	1	2	-	2	1	1	-	-	-
Female	-	-	-	-	-	2	4	4	2	1	-	-
Black, both sexes	1	-	1	3	2	4	3	3	3	2	-	-
Male	-	-	-	3	-	2	-	1	1	1	-	-
Female	1	-	1	-	2	2	3	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
South Dakota												
All races, both sexes	-	-	1	-	-	-	2	1	2	1	1	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	2	1	1	-
White, both sexes	-	-	1	-	-	-	2	1	2	1	1	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	5	4	8	6	17	20	19	10	5	-	-
Male	1	2	1	4	4	7	8	4	1	-	-	-
Female	2	3	3	4	2	10	12	15	9	5	-	-
White, both sexes	-	1	1	3	4	9	18	16	9	5	-	-
Male	-	1	-	1	3	4	7	4	1	-	-	-
Female	-	-	1	2	1	5	11	12	8	5	-	-
Black, both sexes	3	4	3	5	2	8	2	3	1	-	-	-
Male	1	1	1	3	1	3	1	-	-	-	-	-
Female	2	3	2	2	1	5	1	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Texas												
All races, both sexes	8	5	10	9	15	25	33	26	31	14	4	-
Male	6	3	8	3	7	10	9	11	5	4	-	-
Female	2	2	2	6	8	15	24	15	26	10	4	-
White, both sexes	3	2	6	8	8	18	27	21	26	12	1	-
Male	3	2	4	3	4	8	8	8	5	3	-	-
Female	-	-	2	5	4	10	19	13	21	9	1	-
Black, both sexes	5	3	4	1	7	7	6	5	5	2	2	-
Male	3	1	4	-	3	2	1	3	-	1	-	-
Female	2	2	-	1	4	5	5	2	5	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Utah												
All races, both sexes	-	-	-	-	1	2	5	1	-	-	-	-
Male	-	-	-	-	1	-	3	1	-	-	-	-
Female	-	-	-	-	-	2	2	-	-	-	-	-
White, both sexes	-	-	-	-	1	2	5	1	-	-	-	-
Male	-	-	-	-	1	-	3	1	-	-	-	-
Female	-	-	-	-	-	2	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Vermont												
All races, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	6	-	4	6	5	9	9	9	6	3	-	-
Male	4	-	3	4	2	3	3	1	3	-	-	-
Female	2	-	1	2	3	6	6	8	3	3	-	-
White, both sexes	1	-	-	1	3	6	5	2	5	3	-	-
Male	1	-	-	-	2	1	-	-	2	-	-	-
Female	-	-	-	1	1	5	5	2	3	3	-	-
Black, both sexes	5	-	4	5	2	3	4	6	1	-	-	-
Male	3	-	3	4	-	2	3	1	1	-	-	-
Female	2	-	1	1	2	1	1	5	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Washington												
All races, both sexes	1	2	1	2	3	7	20	18	9	6	1	-
Male	-	2	1	1	-	5	11	6	3	1	1	-
Female	1	-	-	1	3	2	9	12	6	5	-	-
White, both sexes	-	-	1	2	3	6	18	17	9	6	1	-
Male	-	-	1	1	-	4	9	5	3	1	1	-
Female	-	-	-	1	3	2	9	12	6	5	-	-
Black, both sexes	1	2	-	-	-	-	-	1	-	-	-	-
Male	-	2	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	-	2	5	5	7	3	6	2	-	-
Male	-	-	-	1	3	1	3	2	1	1	-	-
Female	-	-	-	1	2	4	4	1	5	1	-	-
White, both sexes	-	-	-	1	5	3	7	3	6	2	-	-
Male	-	-	-	1	3	1	3	2	1	1	-	-
Female	-	-	-	-	2	2	4	1	5	1	-	-
Black, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Hypertensive heart and renal disease (I13)												
Wisconsin												
All races, both sexes	1	-	1	2	2	7	4	12	14	2	-	-
Male	1	-	-	1	2	2	-	8	4	-	-	-
Female	-	-	1	1	-	5	4	4	10	2	-	-
White, both sexes	1	-	-	-	2	6	4	12	14	2	-	-
Male	1	-	-	-	2	2	-	8	4	-	-	-
Female	-	-	-	-	-	4	4	4	10	2	-	-
Black, both sexes	-	-	1	2	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
United States												
All races, both sexes	445,687	13	7	7	7	29	122	283	731	1,918	4,942	9,653
Male	232,115	7	4	4	5	23	98	229	545	1,439	3,749	7,355
Female	213,572	6	3	3	2	6	24	54	186	479	1,193	2,298
White, both sexes	390,421	7	5	6	6	21	85	194	544	1,428	3,897	7,587
Male	203,924	5	2	4	5	18	71	159	405	1,087	3,049	5,910
Female	186,497	2	3	2	1	3	14	35	139	341	848	1,677
Black, both sexes	46,027	5	1	1	1	8	28	75	145	429	887	1,804
Male	22,933	2	1	-	-	5	21	59	102	301	567	1,232
Female	23,094	3	-	1	1	3	7	16	43	128	320	572
American Indian 1/, both sexes	1,788	-	-	-	-	-	1	-	16	22	38	92
Male	1,028	-	-	-	-	-	-	-	13	17	31	67
Female	760	-	-	-	-	-	1	-	3	5	7	25
Asian or Pacific Islander 2/, both sexes	7,451	1	1	-	-	-	8	14	26	39	120	170
Male	4,230	-	1	-	-	-	6	11	25	34	102	146
Female	3,221	1	-	-	-	-	2	3	1	5	18	24
Alabama												
All races, both sexes	5,985	-	-	-	1	-	4	3	11	43	100	179
Male	3,089	-	-	-	1	-	4	2	8	31	66	128
Female	2,896	-	-	-	-	-	-	1	3	12	34	51
White, both sexes	4,770	-	-	-	1	-	3	-	6	25	65	123
Male	2,503	-	-	-	1	-	3	-	6	20	43	86
Female	2,267	-	-	-	-	-	-	-	-	5	22	37
Black, both sexes	1,194	-	-	-	-	-	1	3	5	18	35	56
Male	580	-	-	-	-	-	1	2	2	11	23	42
Female	614	-	-	-	-	-	-	1	3	7	12	14
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Alaska												
All races, both sexes	373	-	-	-	-	-	1	-	2	2	12	18
Male	251	-	-	-	-	-	1	-	1	2	11	13
Female	122	-	-	-	-	-	-	-	1	-	1	5
White, both sexes	299	-	-	-	-	-	-	-	1	1	10	16
Male	205	-	-	-	-	-	-	-	1	1	9	12
Female	94	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	57	-	-	-	-	-	-	-	1	1	1	2
Male	37	-	-	-	-	-	-	-	-	1	1	1
Female	20	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	1	-	-	-	1	-
Male	5	-	-	-	-	-	1	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	8,198	-	-	-	-	-	2	6	10	36	72	163
Male	4,591	-	-	-	-	-	-	5	10	25	53	130
Female	3,607	-	-	-	-	-	2	1	-	11	19	33
White, both sexes	7,795	-	-	-	-	-	1	5	8	30	63	145
Male	4,368	-	-	-	-	-	-	4	8	22	49	115
Female	3,427	-	-	-	-	-	1	1	-	8	14	30
Black, both sexes	186	-	-	-	-	-	-	1	-	3	8	4
Male	97	-	-	-	-	-	-	1	-	2	3	3
Female	89	-	-	-	-	-	-	-	-	1	5	1
American Indian 1/, both sexes	167	-	-	-	-	-	1	-	2	3	1	12
Male	97	-	-	-	-	-	-	-	2	1	1	10
Female	70	-	-	-	-	-	1	-	-	2	-	2
Asian or Pacific Islander 2/, both sexes	50	-	-	-	-	-	-	-	-	-	-	2
Male	29	-	-	-	-	-	-	-	-	-	-	2
Female	21	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Arkansas												
All races, both sexes	5,304	-	-	-	-	1	5	7	20	37	89	150
Male	2,803	-	-	-	-	1	5	6	14	22	64	113
Female	2,501	-	-	-	-	-	-	1	6	15	25	37
White, both sexes	4,580	-	-	-	-	1	2	2	17	29	66	113
Male	2,466	-	-	-	-	1	2	1	12	19	47	83
Female	2,114	-	-	-	-	-	-	1	5	10	19	30
Black, both sexes	706	-	-	-	-	-	3	5	3	7	21	35
Male	328	-	-	-	-	-	3	5	2	3	15	28
Female	378	-	-	-	-	-	-	-	1	4	6	7
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	2	1
Male	6	-	-	-	-	-	-	-	-	-	2	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	1	-	-
California												
All races, both sexes	47,430	-	1	1	1	1	9	26	57	159	433	849
Male	24,803	-	1	1	1	1	6	19	43	132	339	666
Female	22,627	-	-	-	-	-	3	7	14	27	94	183
White, both sexes	40,214	-	1	1	1	-	5	17	48	122	327	661
Male	20,904	-	1	1	1	-	4	13	34	101	266	522
Female	19,310	-	-	-	-	-	1	4	14	21	61	139
Black, both sexes	3,635	-	-	-	-	1	1	5	4	23	56	120
Male	1,887	-	-	-	-	1	1	3	4	18	32	92
Female	1,748	-	-	-	-	-	-	2	-	5	24	28
American Indian 1/, both sexes	198	-	-	-	-	-	-	-	-	2	2	9
Male	103	-	-	-	-	-	-	-	-	2	1	5
Female	95	-	-	-	-	-	-	-	-	-	1	4
Asian or Pacific Islander 2/, both sexes	3,383	-	-	-	-	-	3	4	5	12	48	59
Male	1,909	-	-	-	-	-	1	3	5	11	40	47
Female	1,474	-	-	-	-	-	2	1	-	1	8	12

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Colorado												
All races, both sexes	4,049	-	-	-	-	-	1	2	5	25	34	102
Male	2,335	-	-	-	-	-	1	2	4	23	29	85
Female	1,714	-	-	-	-	-	-	-	1	2	5	17
White, both sexes	3,856	-	-	-	-	-	1	2	5	24	31	94
Male	2,240	-	-	-	-	-	1	2	4	22	27	81
Female	1,616	-	-	-	-	-	-	-	1	2	4	13
Black, both sexes	140	-	-	-	-	-	-	-	-	1	1	6
Male	66	-	-	-	-	-	-	-	-	1	-	2
Female	74	-	-	-	-	-	-	-	-	-	1	4
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	37	-	-	-	-	-	-	-	-	-	2	1
Male	20	-	-	-	-	-	-	-	-	-	2	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	4,977	-	-	-	-	-	1	4	7	16	36	90
Male	2,440	-	-	-	-	-	1	4	5	13	27	66
Female	2,537	-	-	-	-	-	-	-	2	3	9	24
White, both sexes	4,649	-	-	-	-	-	1	4	6	11	28	77
Male	2,271	-	-	-	-	-	1	4	4	9	24	56
Female	2,378	-	-	-	-	-	-	-	2	2	4	21
Black, both sexes	284	-	-	-	-	-	-	-	-	4	8	10
Male	144	-	-	-	-	-	-	-	-	4	3	7
Female	140	-	-	-	-	-	-	-	-	-	5	3
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	1	1	-	1
Male	5	-	-	-	-	-	-	-	1	-	-	1
Female	5	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	34	-	-	-	-	-	-	-	-	-	-	2
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Delaware												
All races, both sexes	1,449	1	-	-	-	-	1	1	2	7	17	33
Male	733	1	-	-	-	-	1	1	1	4	12	22
Female	716	-	-	-	-	-	-	-	1	3	5	11
White, both sexes	1,224	1	-	-	-	-	1	-	2	4	13	27
Male	641	1	-	-	-	-	1	-	1	2	11	19
Female	583	-	-	-	-	-	-	-	1	2	2	8
Black, both sexes	215	-	-	-	-	-	-	1	-	3	4	6
Male	85	-	-	-	-	-	-	1	-	2	1	3
Female	130	-	-	-	-	-	-	-	-	1	3	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1,085	-	-	-	-	-	-	1	1	4	12	22
Male	536	-	-	-	-	-	-	1	1	4	9	16
Female	549	-	-	-	-	-	-	-	-	-	3	6
White, both sexes	228	-	-	-	-	-	-	-	-	-	-	-
Male	112	-	-	-	-	-	-	-	-	-	-	-
Female	116	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	840	-	-	-	-	-	-	-	1	4	12	22
Male	412	-	-	-	-	-	-	-	1	4	9	16
Female	428	-	-	-	-	-	-	-	-	-	3	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	1	-	-	-	-
Male	12	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Florida												
All races, both sexes	34,510	-	-	-	2	1	5	13	33	117	309	562
Male	18,456	-	-	-	1	1	4	9	24	83	238	427
Female	16,054	-	-	-	1	-	1	4	9	34	71	135
White, both sexes	31,500	-	-	-	1	1	2	10	19	76	252	459
Male	16,924	-	-	-	1	1	2	7	17	58	202	359
Female	14,576	-	-	-	-	-	-	3	2	18	50	100
Black, both sexes	2,829	-	-	-	1	-	3	3	12	39	54	97
Male	1,433	-	-	-	-	-	2	2	5	23	33	63
Female	1,396	-	-	-	1	-	1	1	7	16	21	34
American Indian 1/, both sexes	33	-	-	-	-	-	-	-	1	1	-	1
Male	22	-	-	-	-	-	-	-	1	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	148	-	-	-	-	-	-	-	1	1	3	5
Male	77	-	-	-	-	-	-	-	1	1	3	4
Female	71	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	8,714	2	-	-	-	-	2	17	25	68	155	307
Male	4,748	1	-	-	-	-	-	13	19	53	118	233
Female	3,966	1	-	-	-	-	2	4	6	15	37	74
White, both sexes	6,679	-	-	-	-	-	1	8	15	48	107	222
Male	3,733	-	-	-	-	-	-	6	10	35	82	171
Female	2,946	-	-	-	-	-	1	2	5	13	25	51
Black, both sexes	1,975	2	-	-	-	-	1	9	8	19	43	84
Male	980	1	-	-	-	-	-	7	7	17	32	61
Female	995	1	-	-	-	-	1	2	1	2	11	23
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	50	-	-	-	-	-	-	-	2	1	5	1
Male	28	-	-	-	-	-	-	-	2	1	4	1
Female	22	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Hawaii												
All races, both sexes	1,236	-	-	-	-	-	-	-	3	6	22	32
Male	747	-	-	-	-	-	-	-	3	5	17	26
Female	489	-	-	-	-	-	-	-	-	1	5	6
White, both sexes	364	-	-	-	-	-	-	-	-	3	7	10
Male	236	-	-	-	-	-	-	-	-	3	5	9
Female	128	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	863	-	-	-	-	-	-	-	3	3	15	20
Male	504	-	-	-	-	-	-	-	3	2	12	15
Female	359	-	-	-	-	-	-	-	-	1	3	5
Idaho												
All races, both sexes	1,586	-	-	-	-	-	-	1	2	9	21	23
Male	933	-	-	-	-	-	-	1	1	7	15	19
Female	653	-	-	-	-	-	-	-	1	2	6	4
White, both sexes	1,563	-	-	-	-	-	-	1	2	8	20	23
Male	917	-	-	-	-	-	-	1	1	6	15	19
Female	646	-	-	-	-	-	-	-	1	2	5	4
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Illinois												
All races, both sexes	18,774	1	-	1	-	1	5	14	35	76	222	455
Male	9,662	1	-	-	-	1	2	12	29	59	173	353
Female	9,112	-	-	1	-	-	3	2	6	17	49	102
White, both sexes	15,899	1	-	-	-	1	4	10	27	52	173	332
Male	8,214	1	-	-	-	1	1	9	22	41	143	272
Female	7,685	-	-	-	-	-	3	1	5	11	30	60
Black, both sexes	2,643	-	-	1	-	-	1	4	7	23	45	121
Male	1,324	-	-	-	-	-	1	3	6	17	26	79
Female	1,319	-	-	1	-	-	-	1	1	6	19	42
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	220	-	-	-	-	-	-	-	1	1	4	2
Male	119	-	-	-	-	-	-	-	1	1	4	2
Female	101	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	9,272	-	-	-	-	1	7	7	32	49	120	261
Male	4,915	-	-	-	-	1	5	5	25	30	84	185
Female	4,357	-	-	-	-	-	2	2	7	19	36	76
White, both sexes	8,646	-	-	-	-	1	6	7	28	43	102	223
Male	4,588	-	-	-	-	1	5	5	21	25	74	164
Female	4,058	-	-	-	-	-	1	2	7	18	28	59
Black, both sexes	592	-	-	-	-	-	1	-	4	6	17	37
Male	308	-	-	-	-	-	-	-	4	5	10	20
Female	284	-	-	-	-	-	1	-	-	1	7	17
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Iowa												
All races, both sexes	5,670	-	-	1	-	1	-	3	4	18	58	109
Male	2,931	-	-	-	-	1	-	3	3	12	39	83
Female	2,739	-	-	1	-	-	-	-	1	6	19	26
White, both sexes	5,600	-	-	1	-	1	-	3	4	16	54	103
Male	2,892	-	-	-	-	1	-	3	3	11	39	79
Female	2,708	-	-	1	-	-	-	-	1	5	15	24
Black, both sexes	55	-	-	-	-	-	-	-	-	2	4	5
Male	29	-	-	-	-	-	-	-	-	1	-	3
Female	26	-	-	-	-	-	-	-	-	1	4	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3,799	1	-	-	-	-	1	2	8	13	46	95
Male	2,077	1	-	-	-	-	1	1	8	10	32	75
Female	1,722	-	-	-	-	-	-	1	-	3	14	20
White, both sexes	3,595	1	-	-	-	-	1	1	7	12	43	85
Male	1,953	1	-	-	-	-	1	-	7	9	30	67
Female	1,642	-	-	-	-	-	-	1	-	3	13	18
Black, both sexes	157	-	-	-	-	-	-	-	1	1	2	6
Male	95	-	-	-	-	-	-	-	1	1	1	5
Female	62	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	34	-	-	-	-	-	-	-	-	-	1	3
Male	23	-	-	-	-	-	-	-	-	-	1	2
Female	11	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	1	-	-	-	1
Male	6	-	-	-	-	-	-	1	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Kentucky												
All races, both sexes	6,929	-	1	-	-	1	4	13	29	45	138	215
Male	3,645	-	-	-	-	1	4	12	22	32	97	174
Female	3,284	-	1	-	-	-	-	1	7	13	41	41
White, both sexes	6,541	-	1	-	-	1	4	12	28	40	127	201
Male	3,450	-	-	-	-	1	4	11	21	29	88	164
Female	3,091	-	1	-	-	-	-	1	7	11	39	37
Black, both sexes	372	-	-	-	-	-	-	1	-	5	11	13
Male	185	-	-	-	-	-	-	1	-	3	9	9
Female	187	-	-	-	-	-	-	-	-	2	2	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	6,720	-	-	-	1	2	1	9	19	55	117	217
Male	3,665	-	-	-	-	1	1	7	12	39	81	156
Female	3,055	-	-	-	1	1	-	2	7	16	36	61
White, both sexes	4,898	-	-	-	1	1	1	5	14	34	65	119
Male	2,687	-	-	-	-	1	1	4	8	25	48	85
Female	2,211	-	-	-	1	-	-	1	6	9	17	34
Black, both sexes	1,779	-	-	-	-	1	-	4	5	21	52	97
Male	951	-	-	-	-	-	-	3	4	14	33	70
Female	828	-	-	-	-	1	-	1	1	7	19	27
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Maine												
All races, both sexes	1,951	-	-	-	-	1	1	1	3	7	20	39
Male	1,037	-	-	-	-	1	1	1	2	5	16	30
Female	914	-	-	-	-	-	-	-	1	2	4	9
White, both sexes	1,941	-	-	-	-	1	1	1	3	7	20	38
Male	1,032	-	-	-	-	1	1	1	2	5	16	29
Female	909	-	-	-	-	-	-	-	1	2	4	9
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	8,147	-	-	-	-	-	1	3	14	29	90	140
Male	4,066	-	-	-	-	-	1	1	10	20	65	110
Female	4,081	-	-	-	-	-	-	2	4	9	25	30
White, both sexes	6,117	-	-	-	-	-	1	3	10	14	61	86
Male	3,062	-	-	-	-	-	1	1	7	8	47	71
Female	3,055	-	-	-	-	-	-	2	3	6	14	15
Black, both sexes	1,910	-	-	-	-	-	-	-	4	13	27	54
Male	941	-	-	-	-	-	-	-	3	11	16	39
Female	969	-	-	-	-	-	-	-	1	2	11	15
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	111	-	-	-	-	-	-	-	-	1	2	-
Male	59	-	-	-	-	-	-	-	-	1	2	-
Female	52	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Massachusetts												
All races, both sexes	8,219	-	-	-	-	1	4	1	12	31	70	129
Male	4,194	-	-	-	-	1	4	1	5	25	54	98
Female	4,025	-	-	-	-	-	-	-	7	6	16	31
White, both sexes	7,884	-	-	-	-	1	3	1	9	28	64	118
Male	4,029	-	-	-	-	1	3	1	4	22	51	92
Female	3,855	-	-	-	-	-	-	-	5	6	13	26
Black, both sexes	269	-	-	-	-	-	-	-	3	2	6	10
Male	130	-	-	-	-	-	-	-	1	2	3	5
Female	139	-	-	-	-	-	-	-	2	-	3	5
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	55	-	-	-	-	-	1	-	-	1	-	1
Male	29	-	-	-	-	-	1	-	-	1	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	17,503	1	-	1	-	1	4	10	28	82	176	370
Male	9,047	1	-	1	-	1	4	9	23	59	133	285
Female	8,456	-	-	-	-	-	-	1	5	23	43	85
White, both sexes	14,822	1	-	1	-	1	3	9	21	56	141	281
Male	7,655	1	-	1	-	1	3	8	19	42	112	228
Female	7,167	-	-	-	-	-	-	1	2	14	29	53
Black, both sexes	2,545	-	-	-	-	-	1	1	7	23	31	86
Male	1,301	-	-	-	-	-	1	1	4	15	18	56
Female	1,244	-	-	-	-	-	-	-	3	8	13	30
American Indian 1/, both sexes	58	-	-	-	-	-	-	-	-	2	3	2
Male	31	-	-	-	-	-	-	-	-	1	2	-
Female	27	-	-	-	-	-	-	-	-	1	1	2
Asian or Pacific Islander 2/, both sexes	78	-	-	-	-	-	-	-	-	1	1	1
Male	60	-	-	-	-	-	-	-	-	1	1	1
Female	18	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Minnesota												
All races, both sexes	4,818	-	-	-	-	-	5	5	10	27	60	133
Male	2,777	-	-	-	-	-	3	4	7	21	51	106
Female	2,041	-	-	-	-	-	2	1	3	6	9	27
White, both sexes	4,700	-	-	-	-	-	5	5	8	24	54	117
Male	2,699	-	-	-	-	-	3	4	6	19	46	92
Female	2,001	-	-	-	-	-	2	1	2	5	8	25
Black, both sexes	65	-	-	-	-	-	-	-	2	2	3	13
Male	44	-	-	-	-	-	-	-	1	1	2	11
Female	21	-	-	-	-	-	-	-	1	1	1	2
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	-	-	2	3
Male	18	-	-	-	-	-	-	-	-	-	2	3
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	1	1	-
Male	16	-	-	-	-	-	-	-	-	1	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4,782	-	1	-	1	2	5	7	32	42	105	181
Male	2,556	-	-	-	1	2	4	7	21	31	72	131
Female	2,226	-	1	-	-	-	1	-	11	11	33	50
White, both sexes	3,432	-	1	-	1	-	2	5	14	20	52	97
Male	1,839	-	-	-	1	-	1	5	8	14	37	71
Female	1,593	-	1	-	-	-	1	-	6	6	15	26
Black, both sexes	1,331	-	-	-	-	2	3	2	18	22	51	82
Male	709	-	-	-	-	2	3	2	13	17	34	58
Female	622	-	-	-	-	-	-	-	5	5	17	24
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Missouri												
All races, both sexes	10,250	-	1	-	-	-	5	7	16	56	130	262
Male	5,239	-	-	-	-	-	4	4	11	41	104	195
Female	5,011	-	1	-	-	-	1	3	5	15	26	67
White, both sexes	9,315	-	1	-	-	-	3	5	13	47	111	225
Male	4,764	-	-	-	-	-	2	4	9	35	90	168
Female	4,551	-	1	-	-	-	1	1	4	12	21	57
Black, both sexes	901	-	-	-	-	-	2	2	3	9	19	36
Male	460	-	-	-	-	-	2	-	2	6	14	26
Female	441	-	-	-	-	-	-	2	1	3	5	10
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1,083	-	-	-	-	-	-	-	3	6	10	24
Male	662	-	-	-	-	-	-	-	2	6	9	18
Female	421	-	-	-	-	-	-	-	1	-	1	6
White, both sexes	1,036	-	-	-	-	-	-	-	2	4	10	21
Male	634	-	-	-	-	-	-	-	1	4	9	15
Female	402	-	-	-	-	-	-	-	1	-	1	6
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	44	-	-	-	-	-	-	-	1	1	-	3
Male	25	-	-	-	-	-	-	-	1	1	-	3
Female	19	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Nebraska												
All races, both sexes	2,096	-	-	-	-	-	-	1	6	11	34	41
Male	1,145	-	-	-	-	-	-	1	4	9	30	33
Female	951	-	-	-	-	-	-	-	2	2	4	8
White, both sexes	2,023	-	-	-	-	-	-	1	5	9	31	38
Male	1,105	-	-	-	-	-	-	1	3	7	27	32
Female	918	-	-	-	-	-	-	-	2	2	4	6
Black, both sexes	53	-	-	-	-	-	-	-	1	2	3	3
Male	31	-	-	-	-	-	-	-	1	2	3	1
Female	22	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2,650	-	-	-	-	-	1	3	4	9	25	84
Male	1,613	-	-	-	-	-	1	2	4	7	19	68
Female	1,037	-	-	-	-	-	-	1	-	2	6	16
White, both sexes	2,408	-	-	-	-	-	1	2	1	6	23	68
Male	1,479	-	-	-	-	-	1	2	1	4	18	57
Female	929	-	-	-	-	-	-	-	-	2	5	11
Black, both sexes	145	-	-	-	-	-	-	1	2	1	2	8
Male	78	-	-	-	-	-	-	-	2	1	1	4
Female	67	-	-	-	-	-	-	1	-	-	1	4
American Indian 1/, both sexes	29	-	-	-	-	-	-	-	1	1	-	3
Male	15	-	-	-	-	-	-	-	1	1	-	2
Female	14	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	68	-	-	-	-	-	-	-	-	1	-	5
Male	41	-	-	-	-	-	-	-	-	1	-	5
Female	27	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
New Hampshire												
All races, both sexes	1,789	1	-	-	-	-	1	1	3	5	16	32
Male	948	-	-	-	-	-	1	1	3	5	14	29
Female	841	1	-	-	-	-	-	-	-	-	2	3
White, both sexes	1,781	1	-	-	-	-	1	-	3	5	16	32
Male	941	-	-	-	-	-	1	-	3	5	14	29
Female	840	1	-	-	-	-	-	-	-	-	2	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	14,733	1	-	-	-	1	2	7	14	37	116	217
Male	7,071	1	-	-	-	1	1	7	11	28	90	165
Female	7,662	-	-	-	-	-	1	-	3	9	26	52
White, both sexes	13,027	-	-	-	-	1	1	3	11	26	91	170
Male	6,253	-	-	-	-	1	1	3	9	20	73	134
Female	6,774	-	-	-	-	-	-	-	2	6	18	36
Black, both sexes	1,439	1	-	-	-	-	1	2	2	10	19	39
Male	665	1	-	-	-	-	-	2	1	7	11	24
Female	774	-	-	-	-	-	1	-	1	3	8	15
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	246	-	-	-	-	-	-	2	1	1	6	7
Male	145	-	-	-	-	-	-	2	1	1	6	7
Female	101	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
New Mexico												
All races, both sexes	2,338	1	-	-	-	-	-	1	5	5	18	43
Male	1,298	-	-	-	-	-	-	1	5	4	14	34
Female	1,040	1	-	-	-	-	-	-	-	1	4	9
White, both sexes	2,190	1	-	-	-	-	-	1	2	4	15	36
Male	1,206	-	-	-	-	-	-	1	2	3	11	30
Female	984	1	-	-	-	-	-	-	-	1	4	6
Black, both sexes	37	-	-	-	-	-	-	-	-	-	1	2
Male	23	-	-	-	-	-	-	-	-	-	1	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	102	-	-	-	-	-	-	-	3	1	2	5
Male	66	-	-	-	-	-	-	-	3	1	2	3
Female	36	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	41,195	-	1	-	-	2	3	14	26	97	258	525
Male	18,773	-	1	-	-	2	2	12	15	74	188	409
Female	22,422	-	-	-	-	-	1	2	11	23	70	116
White, both sexes	34,949	-	-	-	-	2	3	9	20	71	189	375
Male	15,964	-	-	-	-	2	2	8	10	54	144	310
Female	18,985	-	-	-	-	-	1	1	10	17	45	65
Black, both sexes	5,286	-	-	-	-	-	-	4	4	24	55	127
Male	2,280	-	-	-	-	-	-	4	4	18	34	78
Female	3,006	-	-	-	-	-	-	-	-	6	21	49
American Indian 1/, both sexes	38	-	-	-	-	-	-	-	-	-	1	-
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	922	-	1	-	-	-	-	1	2	2	13	23
Male	508	-	1	-	-	-	-	-	1	2	9	21
Female	414	-	-	-	-	-	-	1	1	-	4	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
North Carolina												
All races, both sexes	11,801	-	-	-	1	1	4	6	30	66	161	348
Male	6,315	-	-	-	1	1	4	2	20	49	124	256
Female	5,486	-	-	-	-	-	-	4	10	17	37	92
White, both sexes	9,517	-	-	-	1	-	3	5	21	43	107	251
Male	5,185	-	-	-	1	-	3	2	14	33	84	187
Female	4,332	-	-	-	-	-	-	3	7	10	23	64
Black, both sexes	2,129	-	-	-	-	1	1	1	8	22	49	92
Male	1,039	-	-	-	-	1	1	-	5	15	35	66
Female	1,090	-	-	-	-	-	-	1	3	7	14	26
American Indian 1/, both sexes	113	-	-	-	-	-	-	-	-	-	3	5
Male	62	-	-	-	-	-	-	-	-	-	3	3
Female	51	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	42	-	-	-	-	-	-	-	1	1	2	-
Male	29	-	-	-	-	-	-	-	1	1	2	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1,123	-	-	-	-	-	1	-	2	6	12	13
Male	609	-	-	-	-	-	1	-	1	5	10	11
Female	514	-	-	-	-	-	-	-	1	1	2	2
White, both sexes	1,093	-	-	-	-	-	1	-	2	6	12	10
Male	590	-	-	-	-	-	1	-	1	5	10	9
Female	503	-	-	-	-	-	-	-	1	1	2	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	29	-	-	-	-	-	-	-	-	-	-	3
Male	18	-	-	-	-	-	-	-	-	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Ohio												
All races, both sexes	20,376	-	-	-	-	-	4	11	36	88	252	475
Male	10,465	-	-	-	-	-	3	10	32	65	185	349
Female	9,911	-	-	-	-	-	1	1	4	23	67	126
White, both sexes	18,379	-	-	-	-	-	1	9	27	65	214	407
Male	9,454	-	-	-	-	-	1	8	26	49	163	305
Female	8,925	-	-	-	-	-	-	1	1	16	51	102
Black, both sexes	1,943	-	-	-	-	-	2	2	8	22	38	65
Male	979	-	-	-	-	-	1	2	5	15	22	41
Female	964	-	-	-	-	-	1	-	3	7	16	24
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	45	-	-	-	-	-	1	-	1	1	-	2
Male	27	-	-	-	-	-	1	-	1	1	-	2
Female	18	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	7,324	1	-	-	-	2	3	5	4	26	109	170
Male	3,817	-	-	-	-	-	3	3	3	23	83	127
Female	3,507	1	-	-	-	2	-	2	1	3	26	43
White, both sexes	6,528	-	-	-	-	1	3	4	3	15	88	138
Male	3,388	-	-	-	-	-	3	2	2	12	66	106
Female	3,140	-	-	-	-	1	-	2	1	3	22	32
Black, both sexes	407	-	-	-	-	1	-	1	-	5	9	12
Male	213	-	-	-	-	-	-	1	-	5	6	7
Female	194	-	-	-	-	1	-	-	-	-	3	5
American Indian 1/, both sexes	367	-	-	-	-	-	-	-	1	6	12	18
Male	206	-	-	-	-	-	-	-	1	6	11	13
Female	161	-	-	-	-	-	-	-	-	-	1	5
Asian or Pacific Islander 2/, both sexes	22	1	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	12	1	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Oregon												
All races, both sexes	4,182	-	-	-	-	-	2	1	4	22	40	74
Male	2,435	-	-	-	-	-	2	1	4	19	33	57
Female	1,747	-	-	-	-	-	-	-	-	3	7	17
White, both sexes	4,072	-	-	-	-	-	2	-	4	19	40	71
Male	2,361	-	-	-	-	-	2	-	4	16	33	54
Female	1,711	-	-	-	-	-	-	-	-	3	7	17
Black, both sexes	41	-	-	-	-	-	-	1	-	-	-	1
Male	29	-	-	-	-	-	-	1	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	28	-	-	-	-	-	-	-	-	1	-	1
Male	19	-	-	-	-	-	-	-	-	1	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	41	-	-	-	-	-	-	-	-	2	-	1
Male	26	-	-	-	-	-	-	-	-	2	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	24,023	-	-	-	-	-	-	7	21	77	192	426
Male	12,014	-	-	-	-	-	-	6	16	62	151	325
Female	12,009	-	-	-	-	-	-	1	5	15	41	101
White, both sexes	22,123	-	-	-	-	-	-	7	16	67	167	350
Male	11,039	-	-	-	-	-	-	6	12	57	135	277
Female	11,084	-	-	-	-	-	-	1	4	10	32	73
Black, both sexes	1,807	-	-	-	-	-	-	-	3	10	25	74
Male	921	-	-	-	-	-	-	-	2	5	16	46
Female	886	-	-	-	-	-	-	-	1	5	9	28
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	87	-	-	-	-	-	-	-	2	-	-	2
Male	51	-	-	-	-	-	-	-	2	-	-	2
Female	36	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Rhode Island												
All races, both sexes	2,498	-	-	-	-	-	-	2	3	7	20	36
Male	1,135	-	-	-	-	-	-	2	1	6	15	29
Female	1,363	-	-	-	-	-	-	-	2	1	5	7
White, both sexes	2,413	-	-	-	-	-	-	2	3	6	19	32
Male	1,091	-	-	-	-	-	-	2	1	5	14	26
Female	1,322	-	-	-	-	-	-	-	2	1	5	6
Black, both sexes	64	-	-	-	-	-	-	-	-	1	1	2
Male	35	-	-	-	-	-	-	-	-	1	1	2
Female	29	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	5,631	-	-	-	-	2	3	10	13	43	103	205
Male	3,191	-	-	-	-	1	2	7	8	32	75	155
Female	2,440	-	-	-	-	1	1	3	5	11	28	50
White, both sexes	4,233	-	-	-	-	1	1	1	6	31	71	137
Male	2,448	-	-	-	-	-	1	1	3	24	53	99
Female	1,785	-	-	-	-	1	-	-	3	7	18	38
Black, both sexes	1,381	-	-	-	-	1	2	8	6	12	31	68
Male	733	-	-	-	-	1	1	5	4	8	21	56
Female	648	-	-	-	-	-	1	3	2	4	10	12
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	1	-	-	1	-
Male	8	-	-	-	-	-	-	1	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
South Dakota												
All races, both sexes	1,406	-	-	-	-	-	-	-	3	4	10	23
Male	768	-	-	-	-	-	-	-	1	2	10	19
Female	638	-	-	-	-	-	-	-	2	2	-	4
White, both sexes	1,338	-	-	-	-	-	-	-	2	4	9	20
Male	728	-	-	-	-	-	-	-	1	2	9	16
Female	610	-	-	-	-	-	-	-	1	2	-	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	64	-	-	-	-	-	-	-	1	-	1	2
Male	37	-	-	-	-	-	-	-	-	-	1	2
Female	27	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	10,877	-	-	-	-	-	6	8	30	76	181	331
Male	5,648	-	-	-	-	-	5	7	24	58	138	241
Female	5,229	-	-	-	-	-	1	1	6	18	43	90
White, both sexes	9,466	-	-	-	-	-	3	5	21	63	148	259
Male	4,937	-	-	-	-	-	3	4	16	49	116	192
Female	4,529	-	-	-	-	-	-	1	5	14	32	67
Black, both sexes	1,375	-	-	-	-	-	3	3	8	13	32	71
Male	689	-	-	-	-	-	2	3	7	9	21	48
Female	686	-	-	-	-	-	1	-	1	4	11	23
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	1	-	1	1
Male	16	-	-	-	-	-	-	-	1	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Texas												
All races, both sexes	27,809	1	1	2	-	4	10	27	52	156	375	727
Male	14,979	-	1	2	-	3	8	25	36	111	292	563
Female	12,830	1	-	-	-	1	2	2	16	45	83	164
White, both sexes	24,390	-	1	2	-	4	8	18	40	115	302	569
Male	13,204	-	1	2	-	3	6	17	28	83	240	450
Female	11,186	-	-	-	-	1	2	1	12	32	62	119
Black, both sexes	3,126	1	-	-	-	-	1	8	10	39	70	144
Male	1,616	-	-	-	-	-	1	7	6	26	49	101
Female	1,510	1	-	-	-	-	-	1	4	13	21	43
American Indian 1/, both sexes	26	-	-	-	-	-	-	-	-	-	1	4
Male	16	-	-	-	-	-	-	-	-	-	1	3
Female	10	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	267	-	-	-	-	-	1	1	2	2	2	10
Male	143	-	-	-	-	-	1	1	2	2	2	9
Female	124	-	-	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	1,479	1	-	-	-	1	1	2	1	5	20	43
Male	879	1	-	-	-	1	1	1	1	4	17	37
Female	600	-	-	-	-	-	-	1	-	1	3	6
White, both sexes	1,436	1	-	-	-	1	1	2	1	3	18	42
Male	852	1	-	-	-	1	1	1	1	3	15	36
Female	584	-	-	-	-	-	-	1	-	-	3	6
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	2	2	-
Male	16	-	-	-	-	-	-	-	-	1	2	-
Female	10	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Vermont												
All races, both sexes	867	-	-	-	-	-	-	-	1	3	5	23
Male	451	-	-	-	-	-	-	-	1	1	5	20
Female	416	-	-	-	-	-	-	-	-	2	-	3
White, both sexes	864	-	-	-	-	-	-	-	1	3	5	23
Male	450	-	-	-	-	-	-	-	1	1	5	20
Female	414	-	-	-	-	-	-	-	-	2	-	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	8,808	1	1	-	-	-	3	5	24	45	131	256
Male	4,738	-	1	-	-	-	3	3	20	31	105	188
Female	4,070	1	-	-	-	-	-	2	4	14	26	68
White, both sexes	7,062	-	-	-	-	-	3	3	18	33	97	180
Male	3,862	-	-	-	-	-	3	2	14	25	81	144
Female	3,200	-	-	-	-	-	-	1	4	8	16	36
Black, both sexes	1,624	1	1	-	-	-	-	1	4	11	31	69
Male	802	-	1	-	-	-	-	1	4	5	21	38
Female	822	1	-	-	-	-	-	-	-	6	10	31
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	117	-	-	-	-	-	-	1	2	1	3	6
Male	71	-	-	-	-	-	-	-	2	1	3	6
Female	46	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Washington												
All races, both sexes	7,792	-	-	-	-	1	4	4	9	21	68	140
Male	4,367	-	-	-	-	-	4	4	7	14	54	101
Female	3,425	-	-	-	-	1	-	-	2	7	14	39
White, both sexes	7,346	-	-	-	-	-	2	3	7	15	60	123
Male	4,100	-	-	-	-	-	2	3	5	10	48	88
Female	3,246	-	-	-	-	-	-	-	2	5	12	35
Black, both sexes	167	-	-	-	-	1	1	1	-	3	2	9
Male	99	-	-	-	-	-	1	1	-	2	1	6
Female	68	-	-	-	-	1	-	-	-	1	1	3
American Indian 1/, both sexes	71	-	-	-	-	-	-	-	1	1	1	3
Male	46	-	-	-	-	-	-	-	1	1	1	3
Female	25	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	208	-	-	-	-	-	1	-	1	2	5	5
Male	122	-	-	-	-	-	1	-	1	1	4	4
Female	86	-	-	-	-	-	-	-	-	1	1	1
West Virginia												
All races, both sexes	3,820	-	-	-	-	1	-	1	7	18	47	102
Male	2,017	-	-	-	-	1	-	1	6	13	38	74
Female	1,803	-	-	-	-	-	-	-	1	5	9	28
White, both sexes	3,723	-	-	-	-	1	-	-	7	17	44	98
Male	1,972	-	-	-	-	1	-	-	6	12	35	71
Female	1,751	-	-	-	-	-	-	-	1	5	9	27
Black, both sexes	97	-	-	-	-	-	-	1	-	1	3	4
Male	45	-	-	-	-	-	-	1	-	1	3	3
Female	52	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Ischemic heart diseases (I20-I25)												
Wisconsin												
All races, both sexes	7,657	-	-	1	-	-	-	4	10	22	65	141
Male	4,151	-	-	-	-	-	-	3	8	20	45	110
Female	3,506	-	-	1	-	-	-	1	2	2	20	31
White, both sexes	7,322	-	-	1	-	-	-	3	6	20	56	125
Male	3,951	-	-	-	-	-	-	2	6	18	40	98
Female	3,371	-	-	1	-	-	-	1	-	2	16	27
Black, both sexes	248	-	-	-	-	-	-	-	2	2	6	12
Male	143	-	-	-	-	-	-	-	1	2	4	9
Female	105	-	-	-	-	-	-	-	1	-	2	3
American Indian 1/, both sexes	50	-	-	-	-	-	-	-	2	-	2	1
Male	35	-	-	-	-	-	-	-	1	-	-	1
Female	15	-	-	-	-	-	-	-	1	-	2	-
Asian or Pacific Islander 2/, both sexes	37	-	-	-	-	-	-	1	-	-	1	3
Male	22	-	-	-	-	-	-	1	-	-	1	2
Female	15	-	-	-	-	-	-	-	-	-	-	1
Wyoming												
All races, both sexes	600	-	-	-	-	-	-	-	-	4	10	18
Male	350	-	-	-	-	-	-	-	-	3	6	12
Female	250	-	-	-	-	-	-	-	-	1	4	6
White, both sexes	591	-	-	-	-	-	-	-	-	4	9	17
Male	346	-	-	-	-	-	-	-	-	3	5	11
Female	245	-	-	-	-	-	-	-	-	1	4	6
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
United States												
All races, both sexes	15,657	20,979	25,820	29,933	40,188	57,709	76,726	76,489	56,574	22,766	5,106	28
Male	11,878	15,484	18,064	19,607	24,890	32,870	38,482	32,582	18,648	5,361	773	18
Female	3,779	5,495	7,756	10,326	15,298	24,839	38,244	43,907	37,926	17,405	4,333	10
White, both sexes	12,266	16,756	21,156	24,855	34,210	50,491	69,005	70,289	52,216	20,889	4,489	19
Male	9,568	12,653	15,120	16,614	21,594	29,326	35,049	30,332	17,334	4,938	669	12
Female	2,698	4,103	6,036	8,241	12,616	21,165	33,956	39,957	34,882	15,951	3,820	7
Black, both sexes	2,965	3,650	4,030	4,259	4,971	6,016	6,251	4,940	3,483	1,539	531	8
Male	1,980	2,401	2,501	2,461	2,711	2,901	2,687	1,680	946	295	75	5
Female	985	1,249	1,529	1,798	2,260	3,115	3,564	3,260	2,537	1,244	456	3
American Indian 1/, both sexes	124	170	185	184	206	218	201	167	108	39	17	-
Male	87	119	131	110	115	120	100	72	32	10	4	-
Female	37	51	54	74	91	98	101	95	76	29	13	-
Asian or Pacific Islander 2/, both sexes	302	403	449	635	801	984	1,269	1,093	767	299	69	1
Male	243	311	312	422	470	523	646	498	336	118	25	1
Female	59	92	137	213	331	461	623	595	431	181	44	-
Alabama												
All races, both sexes	269	372	443	467	586	835	1,000	820	630	189	33	-
Male	183	271	286	286	347	443	466	334	185	44	4	-
Female	86	101	157	181	239	392	534	486	445	145	29	-
White, both sexes	201	270	342	354	470	668	842	710	531	142	17	-
Male	142	206	227	230	282	371	407	288	158	31	2	-
Female	59	64	115	124	188	297	435	422	373	111	15	-
Black, both sexes	64	99	100	113	112	162	157	108	98	47	16	-
Male	39	64	58	56	64	71	59	46	27	13	2	-
Female	25	35	42	57	48	91	98	62	71	34	14	-
American Indian 1/, both sexes	2	3	1	-	2	1	1	1	-	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	2	2	-	-	2	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	2	4	-	1	1	-	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	4	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Alaska												
All races, both sexes	28	50	32	32	36	49	52	37	14	6	2	-
Male	17	40	26	22	27	36	29	21	4	1	-	-
Female	11	10	6	10	9	13	23	16	10	5	2	-
White, both sexes	24	42	26	25	25	39	47	26	12	3	2	-
Male	14	34	23	17	20	28	26	16	4	-	-	-
Female	10	8	3	8	5	11	21	10	8	3	2	-
Black, both sexes	1	1	1	1	1	1	-	2	-	-	-	-
Male	1	-	-	1	1	-	-	1	-	-	-	-
Female	-	1	1	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	3	7	4	5	9	9	4	6	2	3	-	-
Male	2	6	2	4	5	8	3	3	-	1	-	-
Female	1	1	2	1	4	1	1	3	2	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	1	3	-	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	2	-	-	-	-
Arizona												
All races, both sexes	269	393	535	579	826	1,135	1,444	1,369	959	327	71	2
Male	204	313	375	388	541	668	750	652	369	92	16	-
Female	65	80	160	191	285	467	694	717	590	235	55	2
White, both sexes	244	363	507	528	772	1,075	1,395	1,334	937	320	66	2
Male	184	290	361	352	514	641	728	634	362	90	14	-
Female	60	73	146	176	258	434	667	700	575	230	52	2
Black, both sexes	12	8	14	22	25	31	25	20	8	4	1	-
Male	10	7	6	18	13	13	12	6	2	1	-	-
Female	2	1	8	4	12	18	13	14	6	3	1	-
American Indian 1/, both sexes	9	18	10	24	26	21	17	7	12	1	3	-
Male	6	14	6	15	12	10	6	6	5	1	2	-
Female	3	4	4	9	14	11	11	1	7	-	1	-
Asian or Pacific Islander 2/, both sexes	4	4	4	5	3	8	7	8	2	2	1	-
Male	4	2	2	3	2	4	4	6	-	-	-	-
Female	-	2	2	2	1	4	3	2	2	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Arkansas												
All races, both sexes	235	301	345	432	520	680	797	793	592	260	40	-
Male	178	220	224	273	347	354	418	314	190	55	5	-
Female	57	81	121	159	173	326	379	479	402	205	35	-
White, both sexes	191	247	284	375	464	598	691	724	526	218	32	-
Male	150	186	195	240	316	327	374	290	172	47	4	-
Female	41	61	89	135	148	271	317	434	354	171	28	-
Black, both sexes	43	54	59	57	54	79	102	68	66	42	8	-
Male	27	34	29	33	30	27	41	24	18	8	1	-
Female	16	20	30	24	24	52	61	44	48	34	7	-
American Indian 1/, both sexes	1	-	2	-	1	1	3	1	-	-	-	-
Male	1	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	2	-	1	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
California												
All races, both sexes	1,486	2,125	2,555	2,964	4,075	6,005	8,523	8,416	6,414	2,655	674	1
Male	1,160	1,602	1,792	1,908	2,536	3,390	4,346	3,714	2,331	692	122	1
Female	326	523	763	1,056	1,539	2,615	4,177	4,702	4,083	1,963	552	-
White, both sexes	1,135	1,649	2,052	2,356	3,315	5,036	7,372	7,448	5,715	2,366	586	1
Male	901	1,266	1,451	1,535	2,076	2,893	3,784	3,299	2,044	612	99	1
Female	234	383	601	821	1,239	2,143	3,588	4,149	3,671	1,754	487	-
Black, both sexes	226	308	319	329	398	484	505	389	279	133	55	-
Male	159	211	216	187	243	238	227	144	83	20	8	-
Female	67	97	103	142	155	246	278	245	196	113	47	-
American Indian 1/, both sexes	8	20	23	26	26	23	19	23	13	4	-	-
Male	5	13	13	17	12	11	10	8	6	-	-	-
Female	3	7	10	9	14	12	9	15	7	4	-	-
Asian or Pacific Islander 2/, both sexes	117	148	161	253	336	462	627	556	407	152	33	-
Male	95	112	112	169	205	248	325	263	198	60	15	-
Female	22	36	49	84	131	214	302	293	209	92	18	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Colorado												
All races, both sexes	170	198	243	293	368	536	696	692	467	177	40	-
Male	144	164	182	200	241	327	383	337	172	33	8	-
Female	26	34	61	93	127	209	313	355	295	144	32	-
White, both sexes	164	181	221	275	341	504	673	675	455	170	40	-
Male	143	149	164	190	229	313	372	332	170	33	8	-
Female	21	32	57	85	112	191	301	343	285	137	32	-
Black, both sexes	5	13	19	14	21	21	11	12	11	5	-	-
Male	1	12	15	7	8	10	5	3	2	-	-	-
Female	4	1	4	7	13	11	6	9	9	5	-	-
American Indian 1/, both sexes	-	1	1	2	3	4	2	-	-	2	-	-
Male	-	1	1	1	2	1	2	-	-	-	-	-
Female	-	-	-	1	1	3	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	1	3	2	2	3	7	10	5	1	-	-	-
Male	-	2	2	2	2	3	4	2	-	-	-	-
Female	1	1	-	-	1	4	6	3	1	-	-	-
Connecticut												
All races, both sexes	150	164	216	235	362	593	863	1,014	789	355	80	2
Male	113	129	155	165	230	336	420	428	258	76	12	2
Female	37	35	61	70	132	257	443	586	531	279	68	-
White, both sexes	124	137	188	199	332	553	824	979	771	339	74	2
Male	100	108	136	141	213	321	404	414	251	71	12	2
Female	24	29	52	58	119	232	420	565	520	268	62	-
Black, both sexes	25	22	25	34	21	36	32	33	18	12	4	-
Male	12	17	17	22	14	13	13	13	7	2	-	-
Female	13	5	8	12	7	23	19	20	11	10	4	-
American Indian 1/, both sexes	-	-	-	1	2	1	1	-	-	1	1	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	1	5	3	1	7	3	6	2	-	3	1	-
Male	1	4	2	1	2	2	2	1	-	3	-	-
Female	-	1	1	-	5	1	4	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Delaware												
All races, both sexes	54	69	100	100	104	202	243	255	182	68	10	-
Male	41	48	68	61	67	106	123	107	53	16	1	-
Female	13	21	32	39	37	96	120	148	129	52	9	-
White, both sexes	37	56	76	81	88	179	212	224	156	61	6	-
Male	32	41	56	51	57	98	109	100	45	16	1	-
Female	5	15	20	30	31	81	103	124	111	45	5	-
Black, both sexes	16	12	24	16	16	22	30	30	24	7	4	-
Male	9	7	12	8	10	7	13	6	6	-	-	-
Female	7	5	12	8	6	15	17	24	18	7	4	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	3	-	1	1	-	1	-	-	-
Male	-	-	-	2	-	1	1	-	1	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	55	58	73	84	110	156	174	159	107	50	17	2
Male	43	39	50	53	62	79	76	57	29	11	5	1
Female	12	19	23	31	48	77	98	102	78	39	12	1
White, both sexes	5	8	13	6	21	24	42	47	37	19	5	1
Male	3	6	10	4	11	14	19	22	14	7	2	-
Female	2	2	3	2	10	10	23	25	23	12	3	1
Black, both sexes	49	49	60	77	87	131	129	109	68	29	12	1
Male	39	32	40	48	49	64	55	33	15	3	3	1
Female	10	17	20	29	38	67	74	76	53	26	9	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	2	1	2	3	2	2	-	-
Male	1	1	-	1	2	1	2	2	-	1	-	-
Female	-	-	-	-	-	-	-	1	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Florida												
All races, both sexes	925	1,196	1,746	2,102	2,902	4,546	6,253	6,391	4,973	2,012	420	2
Male	722	882	1,280	1,435	1,833	2,735	3,301	3,020	1,830	547	82	2
Female	203	314	466	667	1,069	1,811	2,952	3,371	3,143	1,465	338	-
White, both sexes	749	1,006	1,477	1,823	2,552	4,167	5,841	6,057	4,717	1,902	389	-
Male	598	755	1,108	1,270	1,642	2,539	3,120	2,898	1,752	519	76	-
Female	151	251	369	553	910	1,628	2,721	3,159	2,965	1,383	313	-
Black, both sexes	169	176	253	261	327	361	383	308	248	103	29	2
Male	120	118	160	153	183	187	167	112	71	26	6	2
Female	49	58	93	108	144	174	216	196	177	77	23	-
American Indian 1/, both sexes	2	1	6	2	1	4	4	7	2	-	1	-
Male	1	1	5	1	-	2	4	3	2	-	-	-
Female	1	-	1	1	1	2	-	4	-	-	1	-
Asian or Pacific Islander 2/, both sexes	5	13	10	16	22	14	25	19	6	7	1	-
Male	3	8	7	11	8	7	10	7	5	2	-	-
Female	2	5	3	5	14	7	15	12	1	5	1	-
Georgia												
All races, both sexes	446	560	716	731	870	1,123	1,340	1,162	823	308	59	-
Male	331	407	484	483	554	629	626	471	255	66	5	-
Female	115	153	232	248	316	494	714	691	568	242	54	-
White, both sexes	294	383	532	509	655	882	1,090	965	675	256	37	-
Male	228	291	387	355	438	528	519	416	208	56	3	-
Female	66	92	145	154	217	354	571	549	467	200	34	-
Black, both sexes	148	173	182	205	207	237	246	195	143	51	22	-
Male	100	114	96	116	112	100	106	54	45	10	2	-
Female	48	59	86	89	95	137	140	141	98	41	20	-
American Indian 1/, both sexes	1	-	1	4	-	-	1	-	3	-	-	-
Male	1	-	-	4	-	-	1	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	3	4	1	13	8	4	3	2	2	1	-	-
Male	2	2	1	8	4	1	-	1	1	-	-	-
Female	1	2	-	5	4	3	3	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Hawaii												
All races, both sexes	56	60	90	89	114	157	228	179	124	65	11	-
Male	45	52	63	59	72	80	140	99	54	28	4	-
Female	11	8	27	30	42	77	88	80	70	37	7	-
White, both sexes	14	18	34	19	34	48	65	53	34	23	2	-
Male	13	17	27	13	23	26	44	31	16	8	1	-
Female	1	1	7	6	11	22	21	22	18	15	1	-
Black, both sexes	1	-	1	1	-	-	3	-	1	-	-	-
Male	1	-	1	-	-	-	2	-	1	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	41	42	55	69	80	109	160	126	89	42	9	-
Male	31	35	35	46	49	54	94	68	37	20	3	-
Female	10	7	20	23	31	55	66	58	52	22	6	-
Idaho												
All races, both sexes	65	70	99	123	142	219	265	270	212	57	8	-
Male	51	54	70	81	95	149	154	128	87	20	1	-
Female	14	16	29	42	47	70	111	142	125	37	7	-
White, both sexes	63	68	96	119	141	218	260	269	210	57	8	-
Male	49	52	68	78	94	148	151	128	86	20	1	-
Female	14	16	28	41	47	70	109	141	124	37	7	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	2	1	-	1	2	1	-	-	-	-
Male	1	2	2	1	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	3	1	-	3	-	2	-	-	-
Male	1	-	-	2	1	-	1	-	1	-	-	-
Female	-	-	1	1	-	-	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Illinois												
All races, both sexes	747	899	1,064	1,243	1,635	2,330	3,069	3,314	2,443	1,002	218	-
Male	565	679	753	811	963	1,342	1,558	1,382	737	211	31	-
Female	182	220	311	432	672	988	1,511	1,932	1,706	791	187	-
White, both sexes	540	675	818	976	1,312	1,934	2,684	3,009	2,239	922	190	-
Male	422	529	591	655	797	1,143	1,408	1,272	688	193	26	-
Female	118	146	227	321	515	791	1,276	1,737	1,551	729	164	-
Black, both sexes	204	210	232	242	295	363	346	263	187	72	27	-
Male	140	138	150	139	150	183	135	90	46	16	5	-
Female	64	72	82	103	145	180	211	173	141	56	22	-
American Indian 1/, both sexes	-	1	3	1	2	2	-	2	1	-	-	-
Male	-	1	2	-	1	1	-	-	-	-	-	-
Female	-	-	1	1	1	1	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	13	11	24	26	31	39	40	16	8	1	-
Male	3	11	10	17	15	15	15	20	3	2	-	-
Female	-	2	1	7	11	16	24	20	13	6	1	-
Indiana												
All races, both sexes	385	460	564	695	879	1,201	1,536	1,483	1,085	436	71	-
Male	300	337	406	442	517	698	799	637	336	102	6	-
Female	85	123	158	253	362	503	737	846	749	334	65	-
White, both sexes	344	416	518	623	793	1,116	1,476	1,424	1,044	419	63	-
Male	270	306	377	400	477	652	765	614	327	99	6	-
Female	74	110	141	223	316	464	711	810	717	320	57	-
Black, both sexes	40	42	45	64	82	81	54	56	38	17	8	-
Male	30	29	29	36	38	44	31	22	7	3	-	-
Female	10	13	16	28	44	37	23	34	31	14	8	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	7	4	4	6	3	3	-	-	-
Male	-	2	-	6	2	2	3	1	2	-	-	-
Female	1	-	-	1	2	2	3	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Iowa												
All races, both sexes	160	194	260	325	417	693	991	1,072	862	380	122	-
Male	118	144	204	209	270	397	535	508	296	92	17	-
Female	42	50	56	116	147	296	456	564	566	288	105	-
White, both sexes	154	186	254	318	411	690	976	1,069	861	378	121	-
Male	114	139	198	205	266	395	527	507	296	92	17	-
Female	40	47	56	113	145	295	449	562	565	286	104	-
Black, both sexes	6	6	5	5	3	2	12	1	1	2	1	-
Male	4	4	5	2	2	1	7	-	-	-	-	-
Female	2	2	-	3	1	1	5	1	1	2	1	-
American Indian 1/, both sexes	-	1	1	-	2	-	1	-	-	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	1	1	2	2	-	-	-	-
Male	-	-	-	2	1	1	1	1	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
Kansas												
All races, both sexes	132	184	207	223	341	441	620	703	499	236	47	-
Male	99	138	156	155	217	267	332	332	180	62	11	-
Female	33	46	51	68	124	174	288	371	319	174	36	-
White, both sexes	119	166	184	207	320	418	586	683	485	231	46	-
Male	91	122	138	144	207	254	316	323	171	61	11	-
Female	28	44	46	63	113	164	270	360	314	170	35	-
Black, both sexes	11	12	18	14	13	19	25	19	11	5	-	-
Male	7	10	14	10	5	11	14	8	7	1	-	-
Female	4	2	4	4	8	8	11	11	4	4	-	-
American Indian 1/, both sexes	2	5	5	-	6	3	5	1	2	-	1	-
Male	1	5	4	-	4	2	1	1	2	-	-	-
Female	1	-	1	-	2	1	4	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	2	1	4	-	1	-	-	-
Male	-	1	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	1	1	1	3	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Kentucky												
All races, both sexes	329	413	475	558	733	912	1,094	1,005	660	253	51	-
Male	241	280	305	353	455	535	501	390	189	44	10	-
Female	88	133	170	205	278	377	593	615	471	209	41	-
White, both sexes	298	390	448	522	690	856	1,051	963	623	240	46	-
Male	221	267	288	331	428	505	484	375	182	42	9	-
Female	77	123	160	191	262	351	567	588	441	198	37	-
Black, both sexes	29	20	25	33	43	55	43	40	36	13	5	-
Male	19	10	15	20	27	30	17	15	7	2	1	-
Female	10	10	10	13	16	25	26	25	29	11	4	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	3	-	1	-	2	1	-	-	-
Male	1	2	2	2	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	2	1	-	-	-
Louisiana												
All races, both sexes	341	464	524	569	710	886	1,020	907	599	224	55	-
Male	239	343	344	377	431	497	497	389	181	57	13	-
Female	102	121	180	192	279	389	523	518	418	167	42	-
White, both sexes	211	281	366	393	484	696	800	740	488	163	36	-
Male	157	214	256	271	302	406	393	326	147	36	7	-
Female	54	67	110	122	182	290	407	414	341	127	29	-
Black, both sexes	128	178	156	170	218	183	214	162	110	61	19	-
Male	82	124	86	101	125	87	101	60	34	21	6	-
Female	46	54	70	69	93	96	113	102	76	40	13	-
American Indian 1/, both sexes	-	2	1	1	-	3	2	2	-	-	-	-
Male	-	2	1	1	-	1	2	2	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	1	5	8	4	4	3	1	-	-	-
Male	-	3	1	4	4	3	1	1	-	-	-	-
Female	2	-	-	1	4	1	3	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Maine												
All races, both sexes	69	75	113	134	178	273	318	335	269	92	23	-
Male	54	60	82	89	110	147	175	161	80	23	-	-
Female	15	15	31	45	68	126	143	174	189	69	23	-
White, both sexes	67	74	112	133	178	271	316	335	269	92	23	-
Male	53	59	81	89	110	147	174	161	80	23	-	-
Female	14	15	31	44	68	124	142	174	189	69	23	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	1	-	2	2	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	1	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	332	410	509	570	733	1,043	1,432	1,402	945	387	103	4
Male	235	284	351	360	421	546	703	577	277	85	18	2
Female	97	126	158	210	312	497	729	825	668	302	85	2
White, both sexes	196	265	328	359	502	793	1,139	1,156	796	323	83	2
Male	148	195	241	233	287	427	573	498	236	73	15	1
Female	48	70	87	126	215	366	566	658	560	250	68	1
Black, both sexes	131	137	168	193	212	238	283	231	139	58	20	2
Male	82	84	102	113	126	113	126	74	37	11	3	1
Female	49	53	66	80	86	125	157	157	102	47	17	1
American Indian 1/, both sexes	2	1	1	-	2	-	1	1	-	-	-	-
Male	2	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	7	12	18	17	12	9	14	10	6	-	-
Male	3	4	7	14	8	6	4	5	4	1	-	-
Female	-	3	5	4	9	6	5	9	6	5	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Massachusetts												
All races, both sexes	218	309	425	502	718	997	1,416	1,606	1,177	482	121	-
Male	186	238	317	334	468	609	713	663	361	106	11	-
Female	32	71	108	168	250	388	703	943	816	376	110	-
White, both sexes	200	291	404	466	673	947	1,359	1,574	1,155	473	118	-
Male	170	227	302	316	445	583	689	651	357	104	11	-
Female	30	64	102	150	228	364	670	923	798	369	107	-
Black, both sexes	14	14	17	32	36	41	39	28	18	6	3	-
Male	12	9	13	17	17	22	14	11	3	1	-	-
Female	2	5	4	15	19	19	25	17	15	5	3	-
American Indian 1/, both sexes	1	-	2	1	4	3	-	-	-	-	-	-
Male	1	-	1	-	2	2	-	-	-	-	-	-
Female	-	-	1	1	2	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	4	2	3	5	6	18	4	4	3	-	-
Male	3	2	1	1	4	2	10	1	1	1	-	-
Female	-	2	1	2	1	4	8	3	3	2	-	-
Michigan												
All races, both sexes	729	995	1,080	1,136	1,580	2,277	3,028	2,953	2,098	795	159	-
Male	520	710	728	737	988	1,277	1,510	1,161	692	187	21	-
Female	209	285	352	399	592	1,000	1,518	1,792	1,406	608	138	-
White, both sexes	527	708	804	941	1,295	1,933	2,680	2,664	1,903	721	132	-
Male	394	525	552	610	832	1,107	1,351	1,055	630	167	17	-
Female	133	183	252	331	463	826	1,329	1,609	1,273	554	115	-
Black, both sexes	194	275	262	183	264	325	334	273	188	72	26	-
Male	120	175	164	119	142	160	147	99	58	18	4	-
Female	74	100	98	64	122	165	187	174	130	54	22	-
American Indian 1/, both sexes	4	6	5	7	11	8	2	6	2	-	-	-
Male	2	4	4	3	8	4	1	2	-	-	-	-
Female	2	2	1	4	3	4	1	4	2	-	-	-
Asian or Pacific Islander 2/, both sexes	4	6	9	5	10	11	12	10	5	2	1	-
Male	4	6	8	5	6	6	11	5	4	2	-	-
Female	-	-	1	-	4	5	1	5	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Minnesota												
All races, both sexes	171	232	234	299	384	573	817	888	640	280	60	-
Male	141	189	188	212	250	362	455	447	247	80	14	-
Female	30	43	46	87	134	211	362	441	393	200	46	-
White, both sexes	158	223	225	286	371	564	807	879	635	279	60	-
Male	130	182	181	205	244	355	451	443	244	80	14	-
Female	28	41	44	81	127	209	356	436	391	199	46	-
Black, both sexes	11	4	5	5	5	4	5	4	1	1	-	-
Male	9	2	5	2	3	3	1	3	1	-	-	-
Female	2	2	-	3	2	1	4	1	-	1	-	-
American Indian 1/, both sexes	1	3	2	4	4	1	4	1	2	-	-	-
Male	1	3	1	2	2	1	2	-	1	-	-	-
Female	-	-	1	2	2	-	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	4	4	4	1	4	2	-	-	-
Male	1	2	1	3	1	3	1	1	1	-	-	-
Female	-	-	1	1	3	1	-	3	1	-	-	-
Mississippi												
All races, both sexes	248	320	383	384	485	640	680	595	436	190	43	2
Male	173	238	251	263	288	343	345	226	117	35	6	2
Female	75	82	132	121	197	297	335	369	319	155	37	-
White, both sexes	144	193	259	271	352	486	542	475	342	148	26	2
Male	105	148	179	194	222	274	279	180	91	25	3	2
Female	39	45	80	77	130	212	263	295	251	123	23	-
Black, both sexes	103	124	123	112	130	151	138	119	93	41	17	-
Male	67	87	72	69	66	68	66	46	26	10	3	-
Female	36	37	51	43	64	83	72	73	67	31	14	-
American Indian 1/, both sexes	1	2	1	-	2	1	-	-	-	-	-	-
Male	1	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	2	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	2	-	1	1	1	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Missouri												
All races, both sexes	378	456	582	699	940	1,293	1,671	1,750	1,287	576	141	-
Male	280	342	409	455	585	743	815	690	400	145	16	-
Female	98	114	173	244	355	550	856	1,060	887	431	125	-
White, both sexes	311	399	496	609	845	1,181	1,542	1,645	1,220	537	125	-
Male	235	303	358	403	536	685	751	659	377	135	14	-
Female	76	96	138	206	309	496	791	986	843	402	111	-
Black, both sexes	66	54	81	85	92	106	125	101	67	37	16	-
Male	45	38	48	50	47	55	62	30	23	10	2	-
Female	21	16	33	35	45	51	63	71	44	27	14	-
American Indian 1/, both sexes	1	1	3	2	-	2	3	2	-	1	-	-
Male	-	1	2	-	-	2	2	-	-	-	-	-
Female	1	-	1	2	-	-	1	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	3	3	4	1	2	-	1	-	-
Male	-	-	1	2	2	1	-	1	-	-	-	-
Female	-	2	1	1	1	3	1	1	-	1	-	-
Montana												
All races, both sexes	46	61	75	85	97	130	173	195	118	48	12	-
Male	36	45	57	61	68	86	104	101	46	23	-	-
Female	10	16	18	24	29	44	69	94	72	25	12	-
White, both sexes	41	56	70	80	93	124	172	190	115	47	11	-
Male	31	42	52	59	67	82	104	100	46	22	-	-
Female	10	14	18	21	26	42	68	90	69	25	11	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	4	5	5	4	5	1	5	3	1	1	-
Male	5	2	5	2	1	3	-	1	-	1	-	-
Female	-	2	-	3	3	2	1	4	3	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Nebraska												
All races, both sexes	64	62	101	116	143	253	336	410	351	138	29	-
Male	47	50	72	79	94	167	180	205	131	39	4	-
Female	17	12	29	37	49	86	156	205	220	99	25	-
White, both sexes	55	55	96	105	134	244	334	404	349	135	28	-
Male	40	45	68	73	92	164	180	201	130	38	4	-
Female	15	10	28	32	42	80	154	203	219	97	24	-
Black, both sexes	8	7	3	8	6	5	1	2	-	3	1	-
Male	6	5	2	5	1	2	-	2	-	1	-	-
Female	2	2	1	3	5	3	1	-	-	2	1	-
American Indian 1/, both sexes	1	-	1	2	2	3	1	2	2	-	-	-
Male	1	-	1	1	1	1	-	1	1	-	-	-
Female	-	-	-	1	1	2	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	1	-	2	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	1	1	-	1	-	-	-	-
Nevada												
All races, both sexes	134	197	258	279	305	389	407	332	170	44	9	-
Male	106	153	177	184	200	233	228	158	61	10	2	-
Female	28	44	81	95	105	156	179	174	109	34	7	-
White, both sexes	122	174	231	248	274	358	373	318	160	40	9	-
Male	96	140	160	166	183	218	210	153	58	10	2	-
Female	26	34	71	82	91	140	163	165	102	30	7	-
Black, both sexes	7	16	18	19	18	18	20	7	6	2	-	-
Male	5	10	11	10	10	8	11	2	3	-	-	-
Female	2	6	7	9	8	10	9	5	3	2	-	-
American Indian 1/, both sexes	-	3	3	2	5	4	4	-	2	1	-	-
Male	-	1	2	2	2	3	1	-	-	-	-	-
Female	-	2	1	-	3	1	3	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	5	4	6	10	8	9	10	7	2	1	-	-
Male	5	2	4	6	5	4	6	3	-	-	-	-
Female	-	2	2	4	3	5	4	4	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
New Hampshire												
All races, both sexes	54	73	97	130	147	200	288	329	262	120	30	-
Male	48	62	77	90	96	117	135	140	93	34	3	-
Female	6	11	20	40	51	83	153	189	169	86	27	-
White, both sexes	54	73	97	129	145	197	288	329	261	120	30	-
Male	48	62	77	89	95	114	135	140	92	34	3	-
Female	6	11	20	40	50	83	153	189	169	86	27	-
Black, both sexes	-	-	-	1	-	2	-	-	1	-	-	-
Male	-	-	-	1	-	2	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	349	549	671	755	1,165	1,875	2,704	2,933	2,198	918	217	4
Male	279	406	458	493	709	1,007	1,315	1,202	663	197	35	3
Female	70	143	213	262	456	868	1,389	1,731	1,535	721	182	1
White, both sexes	262	422	531	605	972	1,616	2,449	2,758	2,073	846	189	1
Male	216	324	374	395	609	885	1,217	1,145	631	184	31	1
Female	46	98	157	210	363	731	1,232	1,613	1,442	662	158	-
Black, both sexes	69	104	119	126	165	228	222	141	102	61	26	2
Male	48	64	67	82	83	104	88	42	28	9	3	1
Female	21	40	52	44	82	124	134	99	74	52	23	1
American Indian 1/, both sexes	-	1	-	4	3	2	1	1	5	3	-	-
Male	-	-	-	1	2	1	1	1	1	1	-	-
Female	-	1	-	3	1	1	-	-	4	2	-	-
Asian or Pacific Islander 2/, both sexes	18	22	21	20	25	29	32	33	18	8	2	1
Male	15	18	17	15	15	17	9	14	3	3	1	1
Female	3	4	4	5	10	12	23	19	15	5	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
New Mexico												
All races, both sexes	89	102	153	177	246	299	440	336	283	113	26	1
Male	68	83	111	117	162	173	228	154	107	31	5	1
Female	21	19	42	60	84	126	212	182	176	82	21	-
White, both sexes	80	92	139	167	230	279	421	321	270	105	26	1
Male	61	75	103	111	152	159	216	143	104	29	5	1
Female	19	17	36	56	78	120	205	178	166	76	21	-
Black, both sexes	-	3	3	1	6	5	6	2	6	2	-	-
Male	-	3	3	1	4	5	3	1	1	-	-	-
Female	-	-	-	-	2	-	3	1	5	2	-	-
American Indian 1/, both sexes	9	7	11	8	8	14	13	13	3	5	-	-
Male	7	5	5	5	6	8	9	10	-	2	-	-
Female	2	2	6	3	2	6	4	3	3	3	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	-	-	4	1	-	-
Male	-	-	-	-	-	1	-	-	2	-	-	-
Female	-	-	-	1	2	-	-	-	2	1	-	-
New York												
All races, both sexes	910	1,398	1,838	2,312	3,407	5,196	7,294	7,802	6,607	2,828	677	-
Male	677	1,006	1,196	1,413	2,005	2,731	3,347	2,976	2,022	613	84	-
Female	233	392	642	899	1,402	2,465	3,947	4,826	4,585	2,215	593	-
White, both sexes	661	1,035	1,350	1,741	2,722	4,318	6,273	6,990	6,043	2,567	580	-
Male	525	767	913	1,097	1,641	2,315	2,930	2,723	1,882	568	73	-
Female	136	268	437	644	1,081	2,003	3,343	4,267	4,161	1,999	507	-
Black, both sexes	209	321	432	497	575	754	862	680	436	224	82	-
Male	122	200	245	270	295	347	334	199	90	33	7	-
Female	87	121	187	227	280	407	528	481	346	191	75	-
American Indian 1/, both sexes	2	6	2	3	3	5	1	8	7	-	-	-
Male	2	5	2	1	1	3	-	4	2	-	-	-
Female	-	1	-	2	2	2	1	4	5	-	-	-
Asian or Pacific Islander 2/, both sexes	38	36	54	71	107	119	158	124	121	37	15	-
Male	28	34	36	45	68	66	83	50	48	12	4	-
Female	10	2	18	26	39	53	75	74	73	25	11	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
North Carolina												
All races, both sexes	510	698	788	929	1,250	1,593	1,916	1,773	1,237	403	86	1
Male	381	507	571	623	751	915	916	727	380	80	6	1
Female	129	191	217	306	499	678	1,000	1,046	857	323	80	-
White, both sexes	384	508	580	716	1,001	1,276	1,613	1,543	1,068	332	64	1
Male	300	389	434	497	617	769	789	654	339	68	4	1
Female	84	119	146	219	384	507	824	889	729	264	60	-
Black, both sexes	120	172	190	199	229	297	281	219	158	68	22	-
Male	76	104	122	115	123	138	117	70	37	12	2	-
Female	44	68	68	84	106	159	164	149	121	56	20	-
American Indian 1/, both sexes	4	11	13	12	16	17	15	9	6	2	-	-
Male	3	8	12	9	9	7	6	2	-	-	-	-
Female	1	3	1	3	7	10	9	7	6	2	-	-
Asian or Pacific Islander 2/, both sexes	2	7	5	2	4	3	7	2	5	1	-	-
Male	2	6	3	2	2	1	4	1	4	-	-	-
Female	-	1	2	-	2	2	3	1	1	1	-	-
North Dakota												
All races, both sexes	28	48	39	60	72	146	209	232	149	89	17	-
Male	21	37	28	46	51	96	120	111	43	27	1	-
Female	7	11	11	14	21	50	89	121	106	62	16	-
White, both sexes	24	43	39	55	69	141	208	229	149	88	17	-
Male	18	35	28	42	48	93	120	109	43	27	1	-
Female	6	8	11	13	21	48	88	120	106	61	16	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	5	-	4	3	5	1	3	-	1	-	-
Male	3	2	-	3	3	3	-	2	-	-	-	-
Female	1	3	-	1	-	2	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Ohio												
All races, both sexes	710	907	1,139	1,418	1,808	2,739	3,632	3,551	2,446	965	195	-
Male	548	636	802	949	1,100	1,518	1,766	1,471	789	212	30	-
Female	162	271	337	469	708	1,221	1,866	2,080	1,657	753	165	-
White, both sexes	573	749	972	1,220	1,597	2,467	3,347	3,321	2,314	915	181	-
Male	457	540	699	830	986	1,390	1,637	1,386	750	200	27	-
Female	116	209	273	390	611	1,077	1,710	1,935	1,564	715	154	-
Black, both sexes	135	157	162	190	209	268	269	224	129	49	14	-
Male	89	95	100	115	112	126	122	81	38	12	3	-
Female	46	62	62	75	97	142	147	143	91	37	11	-
American Indian 1/, both sexes	1	-	-	-	-	2	2	2	1	-	-	-
Male	1	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	5	8	2	2	14	4	2	1	-	-
Male	1	1	3	4	2	1	5	4	1	-	-	-
Female	-	-	2	4	-	1	9	-	1	1	-	-
Oklahoma												
All races, both sexes	306	402	472	529	739	949	1,202	1,145	837	346	75	2
Male	224	277	327	335	468	513	611	479	252	76	12	1
Female	82	125	145	194	271	436	591	666	585	270	63	1
White, both sexes	247	339	404	451	669	859	1,104	1,053	772	317	59	2
Male	190	244	279	286	422	467	559	442	230	69	8	1
Female	57	95	125	165	247	392	545	611	542	248	51	1
Black, both sexes	30	40	33	40	31	42	40	58	41	18	6	-
Male	16	20	21	28	19	20	26	23	15	4	2	-
Female	14	20	12	12	12	22	14	35	26	14	4	-
American Indian 1/, both sexes	28	22	31	36	36	44	57	32	24	11	9	-
Male	18	13	24	19	25	24	26	14	7	3	2	-
Female	10	9	7	17	11	20	31	18	17	8	7	-
Asian or Pacific Islander 2/, both sexes	1	1	4	2	3	4	1	2	-	-	1	-
Male	-	-	3	2	2	2	-	-	-	-	-	-
Female	1	1	1	-	1	2	1	2	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Oregon												
All races, both sexes	166	201	232	297	362	509	734	765	544	198	31	-
Male	133	160	177	191	241	303	425	376	244	65	4	-
Female	33	41	55	106	121	206	309	389	300	133	27	-
White, both sexes	153	193	221	288	351	495	719	751	538	196	31	-
Male	123	153	167	186	232	293	418	369	244	63	4	-
Female	30	40	54	102	119	202	301	382	294	133	27	-
Black, both sexes	6	2	6	5	4	4	4	4	4	-	-	-
Male	5	2	5	4	4	3	2	2	-	-	-	-
Female	1	-	1	1	-	1	2	2	4	-	-	-
American Indian 1/, both sexes	5	1	3	1	4	3	5	2	1	1	-	-
Male	3	-	3	-	3	3	3	1	-	1	-	-
Female	2	1	-	1	1	-	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	2	3	3	7	6	8	1	1	-	-
Male	2	5	2	1	2	4	2	4	-	1	-	-
Female	-	-	-	2	1	3	4	4	1	-	-	-
Pennsylvania												
All races, both sexes	678	929	1,198	1,391	2,112	3,245	4,485	4,562	3,209	1,218	270	3
Male	525	693	823	919	1,287	1,874	2,196	1,840	1,002	259	35	1
Female	153	236	375	472	825	1,371	2,289	2,722	2,207	959	235	2
White, both sexes	576	799	1,038	1,225	1,895	2,969	4,179	4,345	3,079	1,159	250	2
Male	449	607	717	828	1,155	1,726	2,050	1,768	968	249	34	1
Female	127	192	321	397	740	1,243	2,129	2,577	2,111	910	216	1
Black, both sexes	95	122	152	153	207	262	298	206	123	57	19	1
Male	71	78	99	84	129	141	142	68	32	8	-	-
Female	24	44	53	69	78	121	156	138	91	49	19	1
American Indian 1/, both sexes	-	-	-	-	-	1	1	4	-	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	8	8	13	10	13	7	7	7	2	1	-
Male	5	8	7	7	3	7	3	2	2	2	1	-
Female	2	-	1	6	7	6	4	5	5	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Rhode Island												
All races, both sexes	61	90	109	112	192	266	442	508	410	198	42	-
Male	51	70	82	69	119	146	201	194	114	32	4	-
Female	10	20	27	43	73	120	241	314	296	166	38	-
White, both sexes	56	86	99	103	180	255	434	498	404	196	40	-
Male	48	66	74	63	115	139	199	191	114	32	2	-
Female	8	20	25	40	65	116	235	307	290	164	38	-
Black, both sexes	2	4	9	7	9	9	7	7	5	-	1	-
Male	2	4	7	5	3	5	2	2	-	-	1	-
Female	-	-	2	2	6	4	5	5	5	-	-	-
American Indian 1/, both sexes	2	-	-	1	1	1	1	2	1	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	2	-	-	1	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	2	1	-	1	-	2	1	-
Male	1	-	1	1	1	1	-	-	-	-	1	-
Female	-	-	-	-	1	-	-	1	-	2	-	-
South Carolina												
All races, both sexes	261	403	465	506	621	768	859	725	460	149	34	1
Male	197	297	338	348	374	429	421	323	147	31	5	1
Female	64	106	127	158	247	339	438	402	313	118	29	-
White, both sexes	174	285	335	362	465	589	686	583	368	117	20	1
Male	139	213	249	267	292	339	351	276	114	23	3	1
Female	35	72	86	95	173	250	335	307	254	94	17	-
Black, both sexes	84	117	128	144	153	177	172	140	92	32	14	-
Male	55	83	88	81	81	89	70	47	33	8	2	-
Female	29	34	40	63	72	88	102	93	59	24	12	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	1	-	3	2	1	2	-	-	-	-
Male	2	1	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	2	1	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
South Dakota												
All races, both sexes	43	53	67	73	107	188	252	237	224	93	29	-
Male	33	39	49	49	69	127	141	106	84	32	7	-
Female	10	14	18	24	38	61	111	131	140	61	22	-
White, both sexes	38	42	57	65	102	180	248	230	220	92	29	-
Male	29	33	42	44	65	122	140	104	82	32	7	-
Female	9	9	15	21	37	58	108	126	138	60	22	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	11	10	8	5	7	3	7	4	1	-	-
Male	3	6	7	5	4	4	1	2	2	-	-	-
Female	1	5	3	3	1	3	2	5	2	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Tennessee												
All races, both sexes	493	668	707	879	1,054	1,357	1,783	1,680	1,120	433	71	-
Male	359	478	468	564	645	750	852	635	327	86	11	-
Female	134	190	239	315	409	607	931	1,045	793	347	60	-
White, both sexes	391	544	593	759	909	1,197	1,597	1,525	1,001	393	58	-
Male	291	391	394	499	566	670	777	581	302	79	7	-
Female	100	153	199	260	343	527	820	944	699	314	51	-
Black, both sexes	101	118	111	114	138	158	183	153	116	40	13	-
Male	67	83	71	62	75	79	74	54	23	7	4	-
Female	34	35	40	52	63	79	109	99	93	33	9	-
American Indian 1/, both sexes	-	2	1	2	2	-	1	1	-	-	-	-
Male	-	2	1	1	1	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	2	4	5	2	2	1	3	-	-	-
Male	1	2	2	2	3	1	-	-	2	-	-	-
Female	-	2	-	2	2	1	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Texas												
All races, both sexes	1,149	1,552	1,881	2,122	2,799	3,722	4,721	4,184	2,942	1,121	261	-
Male	866	1,116	1,306	1,363	1,729	2,120	2,387	1,787	977	246	41	-
Female	283	436	575	759	1,070	1,602	2,334	2,397	1,965	875	220	-
White, both sexes	895	1,259	1,570	1,804	2,431	3,288	4,306	3,848	2,683	1,009	238	-
Male	689	921	1,103	1,192	1,530	1,905	2,215	1,661	898	223	37	-
Female	206	338	467	612	901	1,383	2,091	2,187	1,785	786	201	-
Black, both sexes	239	275	278	291	329	395	380	294	244	105	23	-
Male	163	184	182	154	180	188	163	114	76	18	4	-
Female	76	91	96	137	149	207	217	180	168	87	19	-
American Indian 1/, both sexes	4	2	4	1	1	1	5	3	-	-	-	-
Male	4	1	4	1	-	-	1	1	-	-	-	-
Female	-	1	-	-	1	1	4	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	16	29	26	38	38	30	39	15	7	-	-
Male	10	10	17	16	19	27	8	11	3	5	-	-
Female	1	6	12	10	19	11	22	28	12	2	-	-
Utah												
All races, both sexes	56	76	100	98	109	213	257	247	176	67	6	-
Male	45	59	69	75	68	131	152	127	64	26	-	-
Female	11	17	31	23	41	82	105	120	112	41	6	-
White, both sexes	55	70	96	96	106	207	248	245	172	66	6	-
Male	44	55	66	73	67	128	146	126	62	26	-	-
Female	11	15	30	23	39	79	102	119	110	40	6	-
Black, both sexes	1	1	1	1	-	1	2	-	-	-	-	-
Male	1	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	3	1	-	1	-	2	1	1	-	-	-
Male	-	3	-	-	-	-	2	1	1	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	1	2	5	5	1	3	1	-	-
Male	-	1	2	1	1	3	4	-	1	-	-	-
Female	-	1	-	-	1	2	1	1	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Vermont												
All races, both sexes	20	38	40	48	81	115	142	158	121	56	16	-
Male	12	30	28	32	49	69	69	78	41	15	1	-
Female	8	8	12	16	32	46	73	80	80	41	15	-
White, both sexes	20	38	40	47	80	115	142	157	121	56	16	-
Male	12	30	28	31	49	69	69	78	41	15	1	-
Female	8	8	12	16	31	46	73	79	80	41	15	-
Black, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	401	506	617	703	859	1,151	1,460	1,339	895	336	74	1
Male	314	377	424	459	534	663	708	565	276	64	3	-
Female	87	129	193	244	325	488	752	774	619	272	71	1
White, both sexes	281	346	462	541	664	948	1,233	1,146	762	285	59	1
Male	223	269	322	367	436	553	615	507	240	59	2	-
Female	58	77	140	174	228	395	618	639	522	226	57	1
Black, both sexes	116	149	140	152	181	187	208	180	130	49	14	-
Male	87	101	92	85	88	99	84	55	35	5	1	-
Female	29	48	48	67	93	88	124	125	95	44	13	-
American Indian 1/, both sexes	-	-	1	-	1	-	2	-	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	11	14	10	13	16	17	13	3	2	1	-
Male	4	7	9	7	9	11	8	3	1	-	-	-
Female	-	4	5	3	4	5	9	10	2	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Washington												
All races, both sexes	261	352	480	543	672	996	1,360	1,345	1,032	405	99	-
Male	197	270	385	381	445	629	745	629	377	108	17	-
Female	64	82	95	162	227	367	615	716	655	297	82	-
White, both sexes	235	301	441	511	638	936	1,299	1,284	996	398	97	-
Male	176	230	358	360	423	598	710	604	363	105	17	-
Female	59	71	83	151	215	338	589	680	633	293	80	-
Black, both sexes	10	24	15	11	17	20	21	23	8	1	-	-
Male	7	20	11	8	10	10	11	8	3	-	-	-
Female	3	4	4	3	7	10	10	15	5	1	-	-
American Indian 1/, both sexes	4	12	12	7	3	10	6	5	4	1	1	-
Male	4	8	7	5	3	8	3	2	-	-	-	-
Female	-	4	5	2	-	2	3	3	4	1	1	-
Asian or Pacific Islander 2/, both sexes	12	15	12	14	14	30	34	33	24	5	1	-
Male	10	12	9	8	9	13	21	15	11	3	-	-
Female	2	3	3	6	5	17	13	18	13	2	1	-
West Virginia												
All races, both sexes	154	225	225	281	348	534	688	580	414	160	35	-
Male	112	155	163	187	222	287	333	249	124	44	8	-
Female	42	70	62	94	126	247	355	331	290	116	27	-
White, both sexes	145	222	218	276	343	523	674	570	397	154	34	-
Male	106	154	158	183	218	282	328	246	121	43	8	-
Female	39	68	60	93	125	241	346	324	276	111	26	-
Black, both sexes	9	3	7	5	5	11	14	10	17	6	1	-
Male	6	1	5	4	4	5	5	3	3	1	-	-
Female	3	2	2	1	1	6	9	7	14	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Ischemic heart diseases (I20-I25)												
Wisconsin												
All races, both sexes	244	332	409	471	680	988	1,280	1,457	1,033	441	79	-
Male	198	255	292	336	452	630	675	661	350	101	15	-
Female	46	77	117	135	228	358	605	796	683	340	64	-
White, both sexes	209	302	367	437	642	941	1,254	1,430	1,015	437	77	-
Male	174	231	265	311	429	605	659	650	348	100	15	-
Female	35	71	102	126	213	336	595	780	667	337	62	-
Black, both sexes	27	24	28	25	26	39	20	19	12	4	2	-
Male	18	18	15	17	15	22	12	7	2	1	-	-
Female	9	6	13	8	11	17	8	12	10	3	2	-
American Indian 1/, both sexes	6	4	11	5	3	4	4	4	4	-	-	-
Male	5	4	10	5	3	3	2	1	-	-	-	-
Female	1	-	1	-	-	1	2	3	4	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	3	4	9	4	2	4	2	-	-	-
Male	1	2	2	3	5	-	2	3	-	-	-	-
Female	1	-	1	1	4	4	-	1	2	-	-	-
Wyoming												
All races, both sexes	23	30	46	51	65	93	92	94	50	17	7	-
Male	18	20	35	33	37	61	57	43	24	-	1	-
Female	5	10	11	18	28	32	35	51	26	17	6	-
White, both sexes	21	28	46	51	62	93	92	94	50	17	7	-
Male	16	20	35	33	37	61	57	43	24	-	1	-
Female	5	8	11	18	25	32	35	51	26	17	6	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	2	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	2	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
United States												
All races, both sexes	151,004	11	2	5	2	14	54	124	268	744	1,990	3,892
Male	80,079	6	2	2	1	11	43	98	199	537	1,488	2,939
Female	70,925	5	-	3	1	3	11	26	69	207	502	953
White, both sexes	132,364	6	-	4	1	10	40	78	200	572	1,588	3,135
Male	70,791	4	-	2	1	9	33	64	150	420	1,226	2,424
Female	61,573	2	-	2	-	1	7	14	50	152	362	711
Black, both sexes	15,536	4	1	1	1	4	12	40	55	150	348	672
Male	7,527	2	1	-	-	2	10	29	38	98	217	445
Female	8,009	2	-	1	1	2	2	11	17	52	131	227
American Indian 1/, both sexes	589	-	-	-	-	-	-	-	6	7	12	30
Male	339	-	-	-	-	-	-	-	4	6	11	21
Female	250	-	-	-	-	-	-	-	2	1	1	9
Asian or Pacific Islander 2/, both sexes	2,515	1	1	-	-	-	2	6	7	15	42	55
Male	1,422	-	1	-	-	-	-	5	7	13	34	49
Female	1,093	1	-	-	-	-	2	1	-	2	8	6
Alabama												
All races, both sexes	2,756	-	-	-	-	-	3	2	6	29	73	115
Male	1,465	-	-	-	-	-	3	1	5	20	48	75
Female	1,291	-	-	-	-	-	-	1	1	9	25	40
White, both sexes	2,155	-	-	-	-	-	2	-	3	17	50	85
Male	1,162	-	-	-	-	-	2	-	3	13	32	56
Female	993	-	-	-	-	-	-	-	-	4	18	29
Black, both sexes	593	-	-	-	-	-	1	2	3	12	23	30
Male	301	-	-	-	-	-	1	1	2	7	16	19
Female	292	-	-	-	-	-	-	1	1	5	7	11
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Alaska												
All races, both sexes	78	-	-	-	-	-	-	-	1	1	1	1
Male	50	-	-	-	-	-	-	-	-	1	1	1
Female	28	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	58	-	-	-	-	-	-	-	-	1	1	1
Male	39	-	-	-	-	-	-	-	-	1	1	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2,070	-	-	-	-	-	1	3	2	9	12	34
Male	1,136	-	-	-	-	-	-	2	2	6	9	22
Female	934	-	-	-	-	-	1	1	-	3	3	12
White, both sexes	1,963	-	-	-	-	-	1	2	2	6	11	31
Male	1,079	-	-	-	-	-	-	1	2	4	9	19
Female	884	-	-	-	-	-	1	1	-	2	2	12
Black, both sexes	48	-	-	-	-	-	-	1	-	2	1	1
Male	25	-	-	-	-	-	-	1	-	1	-	1
Female	23	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	43	-	-	-	-	-	-	-	-	1	-	1
Male	25	-	-	-	-	-	-	-	-	1	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Arkansas												
All races, both sexes	3,113	-	-	-	-	-	5	6	13	32	68	118
Male	1,733	-	-	-	-	-	5	5	11	20	50	90
Female	1,380	-	-	-	-	-	-	1	2	12	18	28
White, both sexes	2,671	-	-	-	-	-	2	1	12	26	52	90
Male	1,517	-	-	-	-	-	2	-	10	17	38	67
Female	1,154	-	-	-	-	-	-	1	2	9	14	23
Black, both sexes	429	-	-	-	-	-	3	5	1	5	15	26
Male	210	-	-	-	-	-	3	5	1	3	11	21
Female	219	-	-	-	-	-	-	-	-	2	4	5
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	1	-	-
California												
All races, both sexes	13,534	-	-	-	-	-	4	8	15	40	135	278
Male	7,073	-	-	-	-	-	1	6	13	29	99	220
Female	6,461	-	-	-	-	-	3	2	2	11	36	58
White, both sexes	11,293	-	-	-	-	-	2	6	14	30	102	225
Male	5,891	-	-	-	-	-	1	4	12	21	75	181
Female	5,402	-	-	-	-	-	1	2	2	9	27	44
Black, both sexes	998	-	-	-	-	-	-	1	1	5	13	31
Male	482	-	-	-	-	-	-	1	1	4	8	23
Female	516	-	-	-	-	-	-	-	-	1	5	8
American Indian 1/, both sexes	60	-	-	-	-	-	-	-	-	-	1	5
Male	29	-	-	-	-	-	-	-	-	-	1	1
Female	31	-	-	-	-	-	-	-	-	-	-	4
Asian or Pacific Islander 2/, both sexes	1,183	-	-	-	-	-	2	1	-	5	19	17
Male	671	-	-	-	-	-	-	1	-	4	15	15
Female	512	-	-	-	-	-	2	-	-	1	4	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Colorado												
All races, both sexes	1,297	-	-	-	-	-	-	-	3	8	8	41
Male	768	-	-	-	-	-	-	-	3	8	6	37
Female	529	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	1,227	-	-	-	-	-	-	-	3	7	6	39
Male	733	-	-	-	-	-	-	-	3	7	5	36
Female	494	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	47	-	-	-	-	-	-	-	-	1	1	2
Male	25	-	-	-	-	-	-	-	-	1	-	1
Female	22	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1,349	-	-	-	-	-	-	1	2	3	10	27
Male	664	-	-	-	-	-	-	1	1	1	6	20
Female	685	-	-	-	-	-	-	-	1	2	4	7
White, both sexes	1,276	-	-	-	-	-	-	1	2	2	8	24
Male	628	-	-	-	-	-	-	1	1	1	6	17
Female	648	-	-	-	-	-	-	-	1	1	2	7
Black, both sexes	61	-	-	-	-	-	-	-	-	-	2	1
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Delaware												
All races, both sexes	416	1	-	-	-	-	1	-	-	-	5	11
Male	199	1	-	-	-	-	1	-	-	-	3	7
Female	217	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	366	1	-	-	-	-	1	-	-	-	4	10
Male	180	1	-	-	-	-	1	-	-	-	3	7
Female	186	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	48	-	-	-	-	-	-	-	-	-	1	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	218	-	-	-	-	-	-	-	1	1	2	5
Male	95	-	-	-	-	-	-	-	1	1	1	4
Female	123	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	47	-	-	-	-	-	-	-	-	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	169	-	-	-	-	-	-	-	1	1	2	5
Male	73	-	-	-	-	-	-	-	1	1	1	4
Female	96	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Florida												
All races, both sexes	9,439	-	-	-	2	1	-	2	10	39	73	155
Male	5,012	-	-	-	1	1	-	-	6	26	58	110
Female	4,427	-	-	-	1	-	-	2	4	13	15	45
White, both sexes	8,545	-	-	-	1	1	-	1	4	23	60	115
Male	4,579	-	-	-	1	1	-	-	4	17	49	85
Female	3,966	-	-	-	-	-	-	1	-	6	11	30
Black, both sexes	832	-	-	-	1	-	-	1	5	15	12	38
Male	399	-	-	-	-	-	-	-	1	8	8	23
Female	433	-	-	-	1	-	-	1	4	7	4	15
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	51	-	-	-	-	-	-	-	-	1	1	2
Male	28	-	-	-	-	-	-	-	-	1	1	2
Female	23	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3,324	2	-	-	-	-	1	7	6	25	63	125
Male	1,784	1	-	-	-	-	-	6	5	17	47	90
Female	1,540	1	-	-	-	-	1	1	1	8	16	35
White, both sexes	2,537	-	-	-	-	-	-	4	3	21	46	96
Male	1,400	-	-	-	-	-	-	3	2	14	37	69
Female	1,137	-	-	-	-	-	-	1	1	7	9	27
Black, both sexes	759	2	-	-	-	-	1	3	3	3	15	28
Male	366	1	-	-	-	-	-	3	3	2	9	20
Female	393	1	-	-	-	-	1	-	-	1	6	8
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	1	2	1
Male	14	-	-	-	-	-	-	-	-	1	1	1
Female	9	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Hawaii												
All races, both sexes	387	-	-	-	-	-	-	-	2	-	9	10
Male	215	-	-	-	-	-	-	-	2	-	9	7
Female	172	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	88	-	-	-	-	-	-	-	-	-	3	2
Male	57	-	-	-	-	-	-	-	-	-	3	1
Female	31	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	298	-	-	-	-	-	-	-	2	-	6	8
Male	158	-	-	-	-	-	-	-	2	-	6	6
Female	140	-	-	-	-	-	-	-	-	-	-	2
Idaho												
All races, both sexes	774	-	-	-	-	-	-	1	1	4	12	17
Male	480	-	-	-	-	-	-	1	-	4	9	15
Female	294	-	-	-	-	-	-	-	1	-	3	2
White, both sexes	764	-	-	-	-	-	-	1	1	3	11	17
Male	473	-	-	-	-	-	-	1	-	3	9	15
Female	291	-	-	-	-	-	-	-	1	-	2	2
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Illinois												
All races, both sexes	7,251	-	-	1	-	-	2	3	22	29	96	165
Male	3,778	-	-	-	-	-	1	2	18	19	73	132
Female	3,473	-	-	1	-	-	1	1	4	10	23	33
White, both sexes	6,195	-	-	-	-	-	2	1	16	22	79	129
Male	3,284	-	-	-	-	-	1	1	13	15	65	109
Female	2,911	-	-	-	-	-	1	-	3	7	14	20
Black, both sexes	959	-	-	1	-	-	-	2	5	6	16	35
Male	440	-	-	-	-	-	-	1	4	3	7	22
Female	519	-	-	1	-	-	-	1	1	3	9	13
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	92	-	-	-	-	-	-	-	1	1	1	1
Male	52	-	-	-	-	-	-	-	1	1	1	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	3,521	-	-	-	-	-	3	3	9	23	59	135
Male	1,918	-	-	-	-	-	2	2	7	13	44	99
Female	1,603	-	-	-	-	-	1	1	2	10	15	36
White, both sexes	3,332	-	-	-	-	-	3	3	9	21	51	120
Male	1,811	-	-	-	-	-	2	2	7	11	39	91
Female	1,521	-	-	-	-	-	1	1	2	10	12	29
Black, both sexes	178	-	-	-	-	-	-	-	-	2	7	14
Male	99	-	-	-	-	-	-	-	-	2	5	7
Female	79	-	-	-	-	-	-	-	-	-	2	7
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Iowa												
All races, both sexes	1,918	-	-	1	-	-	-	2	2	4	29	45
Male	1,033	-	-	-	-	-	-	2	1	3	22	37
Female	885	-	-	1	-	-	-	-	1	1	7	8
White, both sexes	1,902	-	-	1	-	-	-	2	2	4	29	43
Male	1,026	-	-	-	-	-	-	2	1	3	22	36
Female	876	-	-	1	-	-	-	-	1	1	7	7
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1,264	1	-	-	-	-	-	1	3	5	15	35
Male	686	1	-	-	-	-	-	1	3	4	12	28
Female	578	-	-	-	-	-	-	-	-	1	3	7
White, both sexes	1,203	1	-	-	-	-	-	-	2	5	14	33
Male	649	1	-	-	-	-	-	-	2	4	12	26
Female	554	-	-	-	-	-	-	-	-	1	2	7
Black, both sexes	50	-	-	-	-	-	-	-	1	-	1	1
Male	29	-	-	-	-	-	-	-	1	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	-	-	1
Male	2	-	-	-	-	-	-	1	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Kentucky												
All races, both sexes	3,485	-	-	-	-	1	2	6	18	27	92	132
Male	1,943	-	-	-	-	1	2	5	13	20	64	109
Female	1,542	-	-	-	-	-	-	1	5	7	28	23
White, both sexes	3,261	-	-	-	-	1	2	5	18	25	84	125
Male	1,822	-	-	-	-	1	2	4	13	18	57	103
Female	1,439	-	-	-	-	-	-	1	5	7	27	22
Black, both sexes	216	-	-	-	-	-	-	1	-	2	8	7
Male	117	-	-	-	-	-	-	1	-	2	7	6
Female	99	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3,043	-	-	-	-	2	1	8	8	26	65	131
Male	1,691	-	-	-	-	1	1	6	6	15	44	95
Female	1,352	-	-	-	-	1	-	2	2	11	21	36
White, both sexes	2,118	-	-	-	-	1	1	4	6	17	36	68
Male	1,181	-	-	-	-	1	1	3	4	11	25	48
Female	937	-	-	-	-	-	-	1	2	6	11	20
Black, both sexes	902	-	-	-	-	1	-	4	2	9	29	63
Male	496	-	-	-	-	-	-	3	2	4	19	47
Female	406	-	-	-	-	1	-	1	-	5	10	16
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Maine												
All races, both sexes	688	-	-	-	-	-	1	-	-	2	5	12
Male	357	-	-	-	-	-	1	-	-	1	4	6
Female	331	-	-	-	-	-	-	-	-	1	1	6
White, both sexes	684	-	-	-	-	-	1	-	-	2	5	12
Male	357	-	-	-	-	-	1	-	-	1	4	6
Female	327	-	-	-	-	-	-	-	-	1	1	6
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,688	-	-	-	-	-	-	-	3	12	40	40
Male	1,320	-	-	-	-	-	-	-	2	8	29	31
Female	1,368	-	-	-	-	-	-	-	1	4	11	9
White, both sexes	2,030	-	-	-	-	-	-	-	2	5	27	25
Male	995	-	-	-	-	-	-	-	2	2	19	20
Female	1,035	-	-	-	-	-	-	-	-	3	8	5
Black, both sexes	621	-	-	-	-	-	-	-	1	6	12	15
Male	303	-	-	-	-	-	-	-	-	5	9	11
Female	318	-	-	-	-	-	-	-	1	1	3	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	-	-	1	1	-
Male	21	-	-	-	-	-	-	-	-	1	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Massachusetts												
All races, both sexes	3,054	-	-	-	-	1	1	-	1	11	21	41
Male	1,567	-	-	-	-	1	1	-	1	9	15	29
Female	1,487	-	-	-	-	-	-	-	-	2	6	12
White, both sexes	2,935	-	-	-	-	1	1	-	-	11	20	39
Male	1,511	-	-	-	-	1	1	-	-	9	15	27
Female	1,424	-	-	-	-	-	-	-	-	2	5	12
Black, both sexes	92	-	-	-	-	-	-	-	1	-	1	1
Male	41	-	-	-	-	-	-	-	1	-	-	1
Female	51	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	5,089	1	-	1	-	1	-	7	4	27	58	124
Male	2,623	1	-	1	-	1	-	7	4	21	46	106
Female	2,466	-	-	-	-	-	-	-	-	6	12	18
White, both sexes	4,545	1	-	1	-	1	-	6	3	23	54	109
Male	2,382	1	-	1	-	1	-	6	3	18	45	95
Female	2,163	-	-	-	-	-	-	-	-	5	9	14
Black, both sexes	507	-	-	-	-	-	-	1	1	4	4	15
Male	218	-	-	-	-	-	-	1	1	3	1	11
Female	289	-	-	-	-	-	-	-	-	1	3	4
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Minnesota												
All races, both sexes	1,534	-	-	-	-	-	1	2	3	10	14	54
Male	870	-	-	-	-	-	1	2	2	7	10	46
Female	664	-	-	-	-	-	-	-	1	3	4	8
White, both sexes	1,505	-	-	-	-	-	1	2	3	10	14	50
Male	851	-	-	-	-	-	1	2	2	7	10	42
Female	654	-	-	-	-	-	-	-	1	3	4	8
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2,109	-	-	-	-	1	3	3	18	13	46	98
Male	1,229	-	-	-	-	1	2	3	12	9	32	73
Female	880	-	-	-	-	-	1	-	6	4	14	25
White, both sexes	1,430	-	-	-	-	-	1	2	10	6	22	51
Male	860	-	-	-	-	-	-	2	6	3	14	41
Female	570	-	-	-	-	-	1	-	4	3	8	10
Black, both sexes	671	-	-	-	-	1	2	1	8	7	23	45
Male	364	-	-	-	-	1	2	1	6	6	17	30
Female	307	-	-	-	-	-	-	-	2	1	6	15
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Missouri												
All races, both sexes	4,382	-	-	-	-	-	4	5	5	24	61	137
Male	2,323	-	-	-	-	-	3	3	3	15	48	95
Female	2,059	-	-	-	-	-	1	2	2	9	13	42
White, both sexes	3,999	-	-	-	-	-	2	3	4	22	55	122
Male	2,135	-	-	-	-	-	1	3	3	15	44	87
Female	1,864	-	-	-	-	-	1	-	1	7	11	35
Black, both sexes	368	-	-	-	-	-	2	2	1	2	6	15
Male	179	-	-	-	-	-	2	-	-	-	4	8
Female	189	-	-	-	-	-	-	2	1	2	2	7
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	394	-	-	-	-	-	-	-	1	4	4	10
Male	240	-	-	-	-	-	-	-	1	4	4	7
Female	154	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	373	-	-	-	-	-	-	-	-	3	4	8
Male	230	-	-	-	-	-	-	-	-	3	4	5
Female	143	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	1	1	-	2
Male	10	-	-	-	-	-	-	-	1	1	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Nebraska												
All races, both sexes	648	-	-	-	-	-	-	-	3	4	13	16
Male	370	-	-	-	-	-	-	-	2	4	12	13
Female	278	-	-	-	-	-	-	-	1	-	1	3
White, both sexes	629	-	-	-	-	-	-	-	3	4	12	16
Male	360	-	-	-	-	-	-	-	2	4	11	13
Female	269	-	-	-	-	-	-	-	1	-	1	3
Black, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	649	-	-	-	-	-	1	2	1	2	12	27
Male	399	-	-	-	-	-	1	1	1	1	8	23
Female	250	-	-	-	-	-	-	1	-	1	4	4
White, both sexes	593	-	-	-	-	-	1	1	-	1	10	21
Male	367	-	-	-	-	-	1	1	-	-	7	18
Female	226	-	-	-	-	-	-	-	-	1	3	3
Black, both sexes	31	-	-	-	-	-	-	1	1	-	2	4
Male	16	-	-	-	-	-	-	-	1	-	1	3
Female	15	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
New Hampshire												
All races, both sexes	466	1	-	-	-	-	1	-	1	1	3	7
Male	259	-	-	-	-	-	1	-	1	1	2	7
Female	207	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	465	1	-	-	-	-	1	-	1	1	3	7
Male	258	-	-	-	-	-	1	-	1	1	2	7
Female	207	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4,673	1	-	-	-	1	2	5	4	8	50	91
Male	2,286	1	-	-	-	1	1	5	3	6	37	63
Female	2,387	-	-	-	-	-	1	-	1	2	13	28
White, both sexes	4,050	-	-	-	-	1	1	3	3	4	39	72
Male	1,980	-	-	-	-	1	1	3	2	3	30	54
Female	2,070	-	-	-	-	-	-	-	1	1	9	18
Black, both sexes	524	1	-	-	-	-	1	1	-	3	9	17
Male	242	1	-	-	-	-	-	1	-	2	5	7
Female	282	-	-	-	-	-	1	-	-	1	4	10
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	92	-	-	-	-	-	-	1	1	1	2	2
Male	62	-	-	-	-	-	-	1	1	1	2	2
Female	30	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
New Mexico												
All races, both sexes	574	1	-	-	-	-	-	-	2	1	7	14
Male	318	-	-	-	-	-	-	-	2	1	4	10
Female	256	1	-	-	-	-	-	-	-	-	3	4
White, both sexes	534	1	-	-	-	-	-	-	2	-	5	12
Male	292	-	-	-	-	-	-	-	2	-	2	10
Female	242	1	-	-	-	-	-	-	-	-	3	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	33	-	-	-	-	-	-	-	-	1	2	2
Male	24	-	-	-	-	-	-	-	-	1	2	-
Female	9	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	9,982	-	1	-	-	-	1	3	7	33	89	148
Male	4,713	-	1	-	-	-	1	3	2	23	65	105
Female	5,269	-	-	-	-	-	-	-	5	10	24	43
White, both sexes	8,645	-	-	-	-	-	1	3	6	23	70	111
Male	4,151	-	-	-	-	-	1	3	1	16	52	85
Female	4,494	-	-	-	-	-	-	-	5	7	18	26
Black, both sexes	1,121	-	-	-	-	-	-	-	1	9	16	32
Male	436	-	-	-	-	-	-	-	1	6	11	15
Female	685	-	-	-	-	-	-	-	-	3	5	17
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	205	-	1	-	-	-	-	-	-	1	3	5
Male	122	-	1	-	-	-	-	-	-	1	2	5
Female	83	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
North Carolina												
All races, both sexes	4,000	-	-	-	-	-	1	3	11	23	57	116
Male	2,097	-	-	-	-	-	1	2	7	18	39	83
Female	1,903	-	-	-	-	-	-	1	4	5	18	33
White, both sexes	3,244	-	-	-	-	-	1	2	7	16	34	84
Male	1,746	-	-	-	-	-	1	2	5	13	25	60
Female	1,498	-	-	-	-	-	-	-	2	3	9	24
Black, both sexes	701	-	-	-	-	-	-	1	3	6	21	30
Male	321	-	-	-	-	-	-	-	1	4	12	22
Female	380	-	-	-	-	-	-	1	2	2	9	8
American Indian 1/, both sexes	39	-	-	-	-	-	-	-	-	-	2	2
Male	21	-	-	-	-	-	-	-	-	-	2	1
Female	18	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	1	1	-	-
Male	9	-	-	-	-	-	-	-	1	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	445	-	-	-	-	-	-	-	-	3	5	3
Male	245	-	-	-	-	-	-	-	-	3	4	3
Female	200	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	434	-	-	-	-	-	-	-	-	3	5	3
Male	238	-	-	-	-	-	-	-	-	3	4	3
Female	196	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Ohio												
All races, both sexes	6,758	-	-	-	-	-	-	2	14	30	97	205
Male	3,535	-	-	-	-	-	-	2	12	21	67	155
Female	3,223	-	-	-	-	-	-	-	2	9	30	50
White, both sexes	6,198	-	-	-	-	-	-	2	9	23	81	181
Male	3,277	-	-	-	-	-	-	2	9	17	60	140
Female	2,921	-	-	-	-	-	-	-	-	6	21	41
Black, both sexes	541	-	-	-	-	-	-	-	4	7	16	22
Male	247	-	-	-	-	-	-	-	2	4	7	13
Female	294	-	-	-	-	-	-	-	2	3	9	9
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	1	-	-	1
Male	8	-	-	-	-	-	-	-	1	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,931	1	-	-	-	1	1	2	-	5	26	47
Male	1,010	-	-	-	-	-	1	1	-	4	18	37
Female	921	1	-	-	-	1	-	1	-	1	8	10
White, both sexes	1,719	-	-	-	-	-	1	1	-	4	19	38
Male	906	-	-	-	-	-	1	-	-	3	13	30
Female	813	-	-	-	-	-	-	1	-	1	6	8
Black, both sexes	118	-	-	-	-	1	-	1	-	1	4	5
Male	52	-	-	-	-	-	-	1	-	1	2	4
Female	66	-	-	-	-	1	-	-	-	-	2	1
American Indian 1/, both sexes	88	-	-	-	-	-	-	-	-	-	3	3
Male	49	-	-	-	-	-	-	-	-	-	3	2
Female	39	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	6	1	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Oregon												
All races, both sexes	1,437	-	-	-	-	-	1	-	1	6	7	23
Male	818	-	-	-	-	-	1	-	1	5	6	19
Female	619	-	-	-	-	-	-	-	-	1	1	4
White, both sexes	1,391	-	-	-	-	-	1	-	1	5	7	22
Male	790	-	-	-	-	-	1	-	1	4	6	18
Female	601	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	8,535	-	-	-	-	-	-	2	9	29	80	203
Male	4,431	-	-	-	-	-	-	1	7	26	64	153
Female	4,104	-	-	-	-	-	-	1	2	3	16	50
White, both sexes	7,959	-	-	-	-	-	-	2	6	28	72	169
Male	4,150	-	-	-	-	-	-	1	5	25	60	134
Female	3,809	-	-	-	-	-	-	1	1	3	12	35
Black, both sexes	542	-	-	-	-	-	-	-	2	1	8	32
Male	262	-	-	-	-	-	-	-	1	1	4	17
Female	280	-	-	-	-	-	-	-	1	-	4	15
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	34	-	-	-	-	-	-	-	1	-	-	2
Male	19	-	-	-	-	-	-	-	1	-	-	2
Female	15	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Rhode Island												
All races, both sexes	893	-	-	-	-	-	-	-	-	2	3	6
Male	437	-	-	-	-	-	-	-	-	2	3	5
Female	456	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	860	-	-	-	-	-	-	-	-	2	3	6
Male	424	-	-	-	-	-	-	-	-	2	3	5
Female	436	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2,467	-	-	-	-	1	-	5	7	24	60	119
Male	1,399	-	-	-	-	1	-	3	5	19	46	87
Female	1,068	-	-	-	-	-	-	2	2	5	14	32
White, both sexes	1,779	-	-	-	-	-	-	1	3	18	39	88
Male	1,029	-	-	-	-	-	-	1	1	15	31	62
Female	750	-	-	-	-	-	-	-	2	3	8	26
Black, both sexes	677	-	-	-	-	1	-	3	3	6	20	31
Male	362	-	-	-	-	1	-	1	3	4	14	25
Female	315	-	-	-	-	-	-	2	-	2	6	6
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	1	-	-	1	-
Male	6	-	-	-	-	-	-	1	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
South Dakota												
All races, both sexes	612	-	-	-	-	-	-	-	3	2	9	10
Male	345	-	-	-	-	-	-	-	1	1	9	9
Female	267	-	-	-	-	-	-	-	2	1	-	1
White, both sexes	570	-	-	-	-	-	-	-	2	2	8	8
Male	320	-	-	-	-	-	-	-	1	1	8	7
Female	250	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	39	-	-	-	-	-	-	-	1	-	1	2
Male	23	-	-	-	-	-	-	-	-	-	1	2
Female	16	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4,571	-	-	-	-	-	4	7	11	48	87	175
Male	2,453	-	-	-	-	-	4	6	8	38	70	131
Female	2,118	-	-	-	-	-	-	1	3	10	17	44
White, both sexes	3,939	-	-	-	-	-	2	4	9	39	75	137
Male	2,132	-	-	-	-	-	2	3	6	32	62	107
Female	1,807	-	-	-	-	-	-	1	3	7	13	30
Black, both sexes	620	-	-	-	-	-	2	3	2	9	12	38
Male	315	-	-	-	-	-	2	3	2	6	8	24
Female	305	-	-	-	-	-	-	-	-	3	4	14
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Texas												
All races, both sexes	11,996	1	-	1	-	4	6	14	23	70	171	307
Male	6,533	-	-	1	-	3	5	13	14	48	134	237
Female	5,463	1	-	-	-	1	1	1	9	22	37	70
White, both sexes	10,569	-	-	1	-	4	6	9	19	52	142	249
Male	5,808	-	-	1	-	3	5	9	12	36	113	198
Female	4,761	-	-	-	-	1	1	-	7	16	29	51
Black, both sexes	1,314	1	-	-	-	-	-	4	4	18	29	56
Male	661	-	-	-	-	-	-	3	2	12	21	38
Female	653	1	-	-	-	-	-	1	2	6	8	18
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	102	-	-	-	-	-	-	1	-	-	-	1
Male	56	-	-	-	-	-	-	1	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	550	1	-	-	-	-	1	1	-	1	7	13
Male	332	1	-	-	-	-	1	-	-	1	6	11
Female	218	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	536	1	-	-	-	-	1	1	-	-	7	13
Male	324	1	-	-	-	-	1	-	-	-	6	11
Female	212	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Vermont												
All races, both sexes	232	-	-	-	-	-	-	-	-	1	2	3
Male	129	-	-	-	-	-	-	-	-	1	2	3
Female	103	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	231	-	-	-	-	-	-	-	-	1	2	3
Male	129	-	-	-	-	-	-	-	-	1	2	3
Female	102	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3,213	-	1	-	-	-	1	4	7	19	60	88
Male	1,752	-	1	-	-	-	1	2	7	13	47	63
Female	1,461	-	-	-	-	-	-	2	-	6	13	25
White, both sexes	2,597	-	-	-	-	-	1	2	6	13	46	66
Male	1,451	-	-	-	-	-	1	1	6	9	40	50
Female	1,146	-	-	-	-	-	-	1	-	4	6	16
Black, both sexes	573	-	1	-	-	-	-	1	1	5	14	19
Male	280	-	1	-	-	-	-	1	1	3	7	11
Female	293	-	-	-	-	-	-	-	-	2	7	8
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	40	-	-	-	-	-	-	1	-	1	-	2
Male	19	-	-	-	-	-	-	-	-	1	-	2
Female	21	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Washington												
All races, both sexes	2,442	-	-	-	-	-	1	2	3	5	23	57
Male	1,361	-	-	-	-	-	1	2	2	2	18	45
Female	1,081	-	-	-	-	-	-	-	1	3	5	12
White, both sexes	2,302	-	-	-	-	-	1	1	2	4	19	50
Male	1,280	-	-	-	-	-	1	1	1	1	16	39
Female	1,022	-	-	-	-	-	-	-	1	3	3	11
Black, both sexes	45	-	-	-	-	-	-	1	-	-	1	5
Male	27	-	-	-	-	-	-	1	-	-	-	4
Female	18	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	1	1	1	2
Male	15	-	-	-	-	-	-	-	1	1	1	2
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	73	-	-	-	-	-	-	-	-	-	2	-
Male	39	-	-	-	-	-	-	-	-	-	1	-
Female	34	-	-	-	-	-	-	-	-	-	1	-
West Virginia												
All races, both sexes	1,288	-	-	-	-	-	-	-	1	3	17	37
Male	662	-	-	-	-	-	-	-	1	3	13	23
Female	626	-	-	-	-	-	-	-	-	-	4	14
White, both sexes	1,270	-	-	-	-	-	-	-	1	3	16	36
Male	655	-	-	-	-	-	-	-	1	3	12	22
Female	615	-	-	-	-	-	-	-	-	-	4	14
Black, both sexes	18	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute myocardial infarction (I21-I22)												
Wisconsin												
All races, both sexes	3,180	-	-	1	-	-	-	2	1	15	25	68
Male	1,769	-	-	-	-	-	-	2	1	14	19	54
Female	1,411	-	-	1	-	-	-	-	-	1	6	14
White, both sexes	3,066	-	-	1	-	-	-	2	1	13	22	62
Male	1,698	-	-	-	-	-	-	2	1	12	17	50
Female	1,368	-	-	1	-	-	-	-	-	1	5	12
Black, both sexes	68	-	-	-	-	-	-	-	-	2	2	3
Male	39	-	-	-	-	-	-	-	-	2	1	2
Female	29	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	-	-	-	1
Wyoming												
All races, both sexes	287	-	-	-	-	-	-	-	-	1	4	13
Male	174	-	-	-	-	-	-	-	-	1	2	9
Female	113	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	282	-	-	-	-	-	-	-	-	1	4	12
Male	172	-	-	-	-	-	-	-	-	1	2	8
Female	110	-	-	-	-	-	-	-	-	-	2	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
United States												
All races, both sexes	6,178	8,292	10,261	11,652	15,022	20,167	25,282	23,679	16,169	5,977	1,216	3
Male	4,624	6,083	7,068	7,520	9,156	11,281	12,259	9,779	5,352	1,447	181	3
Female	1,554	2,209	3,193	4,132	5,866	8,886	13,023	13,900	10,817	4,530	1,035	-
White, both sexes	5,037	6,801	8,692	9,878	12,974	17,689	22,728	21,635	14,835	5,427	1,032	2
Male	3,884	5,088	6,131	6,541	8,074	10,116	11,160	9,051	4,942	1,321	148	2
Female	1,153	1,713	2,561	3,337	4,900	7,573	11,568	12,584	9,893	4,106	884	-
Black, both sexes	996	1,262	1,354	1,484	1,689	2,068	2,056	1,640	1,071	463	164	1
Male	627	818	799	803	880	952	852	536	299	91	27	1
Female	369	444	555	681	809	1,116	1,204	1,104	772	372	137	-
American Indian 1/, both sexes	42	67	58	64	63	78	78	49	22	9	4	-
Male	33	49	35	35	32	43	34	21	11	1	3	-
Female	9	18	23	29	31	35	44	28	11	8	1	-
Asian or Pacific Islander 2/, both sexes	103	162	157	226	296	332	420	355	241	78	16	-
Male	80	128	103	141	170	170	213	171	100	34	3	-
Female	23	34	54	85	126	162	207	184	141	44	13	-
Alabama												
All races, both sexes	163	202	254	228	253	339	429	336	248	62	14	-
Male	110	147	172	130	147	177	187	150	79	12	2	-
Female	53	55	82	98	106	162	242	186	169	50	12	-
White, both sexes	126	145	204	173	205	267	348	283	202	41	4	-
Male	85	110	141	104	120	144	157	125	62	8	-	-
Female	41	35	63	69	85	123	191	158	140	33	4	-
Black, both sexes	37	56	50	55	46	69	80	53	45	21	10	-
Male	25	37	31	26	26	32	30	25	17	4	2	-
Female	12	19	19	29	20	37	50	28	28	17	8	-
American Indian 1/, both sexes	-	1	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Alaska												
All races, both sexes	4	13	11	8	8	9	5	10	4	1	1	-
Male	3	12	7	5	6	6	2	5	1	-	-	-
Female	1	1	4	3	2	3	3	5	3	1	1	-
White, both sexes	3	11	8	6	6	8	3	5	4	-	1	-
Male	2	10	6	4	5	5	-	3	1	-	-	-
Female	1	1	2	2	1	3	3	2	3	-	1	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	2	2	1	1	2	3	-	1	-	-
Male	1	2	1	1	-	1	2	1	-	-	-	-
Female	-	-	1	1	1	-	-	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Arizona												
All races, both sexes	56	96	139	161	235	316	367	318	235	65	21	-
Male	44	73	77	105	153	191	190	135	95	25	7	-
Female	12	23	62	56	82	125	177	183	140	40	14	-
White, both sexes	52	90	134	148	217	302	346	311	230	61	19	-
Male	40	68	74	98	144	184	181	133	92	25	5	-
Female	12	22	60	50	73	118	165	178	138	36	14	-
Black, both sexes	1	2	3	3	9	7	9	5	1	3	-	-
Male	1	2	1	2	4	4	5	2	1	-	-	-
Female	-	-	2	1	5	3	4	3	-	3	-	-
American Indian 1/, both sexes	2	2	2	8	9	5	8	-	3	-	2	-
Male	2	2	2	4	5	2	2	-	2	-	2	-
Female	-	-	-	4	4	3	6	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	2	-	2	4	2	1	1	-	-
Male	1	1	-	1	-	1	2	-	-	-	-	-
Female	-	1	-	1	-	1	2	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Arkansas												
All races, both sexes	169	219	229	277	314	389	436	402	303	111	22	-
Male	132	164	151	176	213	200	227	160	100	26	3	-
Female	37	55	78	101	101	189	209	242	203	85	19	-
White, both sexes	138	182	194	242	279	350	364	366	263	92	18	-
Male	113	140	134	157	194	188	199	145	90	21	2	-
Female	25	42	60	85	85	162	165	221	173	71	16	-
Black, both sexes	31	37	33	35	34	38	68	35	40	19	4	-
Male	19	24	17	19	19	12	25	15	10	5	1	-
Female	12	13	16	16	15	26	43	20	30	14	3	-
American Indian 1/, both sexes	-	-	2	-	1	1	3	1	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	2	-	1	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	398	599	758	958	1,309	1,849	2,508	2,228	1,680	634	133	-
Male	300	441	513	592	797	984	1,246	990	642	178	22	-
Female	98	158	245	366	512	865	1,262	1,238	1,038	456	111	-
White, both sexes	299	469	622	758	1,053	1,527	2,124	1,935	1,461	554	112	-
Male	234	349	426	479	652	824	1,060	853	546	156	18	-
Female	65	120	196	279	401	703	1,064	1,082	915	398	94	-
Black, both sexes	58	64	75	99	110	146	166	105	79	35	10	-
Male	34	45	47	47	58	69	75	41	24	4	1	-
Female	24	19	28	52	52	77	91	64	55	31	9	-
American Indian 1/, both sexes	2	6	5	4	9	10	7	4	4	3	-	-
Male	1	5	4	2	2	4	4	2	3	-	-	-
Female	1	1	1	2	7	6	3	2	1	3	-	-
Asian or Pacific Islander 2/, both sexes	39	60	56	97	137	166	211	184	136	42	11	-
Male	31	42	36	64	85	87	107	94	69	18	3	-
Female	8	18	20	33	52	79	104	90	67	24	8	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Colorado												
All races, both sexes	64	65	98	110	126	194	228	192	121	32	7	-
Male	55	49	70	67	83	117	126	91	48	5	3	-
Female	9	16	28	43	43	77	102	101	73	27	4	-
White, both sexes	59	63	91	105	120	180	216	187	115	29	7	-
Male	54	47	64	65	79	109	119	90	47	5	3	-
Female	5	16	27	40	41	71	97	97	68	24	4	-
Black, both sexes	4	-	6	4	2	9	7	2	6	3	-	-
Male	1	-	5	2	2	6	5	1	1	-	-	-
Female	3	-	1	2	-	3	2	1	5	3	-	-
American Indian 1/, both sexes	-	1	-	1	2	1	1	-	-	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	-	2	4	4	3	-	-	-	-
Male	-	1	1	-	1	2	1	-	-	-	-	-
Female	1	-	-	-	1	2	3	3	-	-	-	-
Connecticut												
All races, both sexes	46	40	70	59	105	176	238	291	189	75	17	-
Male	34	34	53	38	71	100	96	117	66	25	1	-
Female	12	6	17	21	34	76	142	174	123	50	16	-
White, both sexes	41	36	62	51	99	163	226	283	188	74	16	-
Male	32	30	48	32	66	96	93	113	66	25	1	-
Female	9	6	14	19	33	67	133	170	122	49	15	-
Black, both sexes	5	4	7	7	3	11	11	7	1	1	1	-
Male	2	4	5	5	3	3	2	3	-	-	-	-
Female	3	-	2	2	-	8	9	4	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	3	2	1	1	-	-	-	-
Male	-	-	-	1	2	1	1	1	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Delaware												
All races, both sexes	12	21	26	37	36	59	66	71	49	17	4	-
Male	10	13	21	21	21	25	28	30	15	3	-	-
Female	2	8	5	16	15	34	38	41	34	14	4	-
White, both sexes	11	21	22	31	32	53	56	64	42	16	2	-
Male	10	13	18	18	20	22	24	28	12	3	-	-
Female	1	8	4	13	12	31	32	36	30	13	2	-
Black, both sexes	1	-	4	5	4	6	10	6	7	1	2	-
Male	-	-	3	2	1	3	4	1	3	-	-	-
Female	1	-	1	3	3	3	6	5	4	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	13	11	6	15	21	36	40	34	26	6	1	-
Male	8	6	3	7	13	20	15	10	6	-	-	-
Female	5	5	3	8	8	16	25	24	20	6	1	-
White, both sexes	1	1	2	1	5	10	9	10	6	2	-	-
Male	1	1	2	-	4	5	4	3	2	-	-	-
Female	-	-	-	1	1	5	5	7	4	2	-	-
Black, both sexes	12	10	4	14	16	26	31	23	19	4	1	-
Male	7	5	1	7	9	15	11	7	4	-	-	-
Female	5	5	3	7	7	11	20	16	15	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Florida												
All races, both sexes	255	399	577	674	902	1,252	1,664	1,627	1,239	479	88	1
Male	190	298	405	435	541	742	862	743	454	120	19	1
Female	65	101	172	239	361	510	802	884	785	359	69	-
White, both sexes	216	346	488	581	789	1,142	1,546	1,532	1,173	448	79	-
Male	169	264	355	381	487	682	803	710	441	115	15	-
Female	47	82	133	200	302	460	743	822	732	333	64	-
Black, both sexes	37	49	82	87	101	104	111	84	64	31	9	1
Male	21	31	45	50	50	56	56	29	11	5	4	1
Female	16	18	37	37	51	48	55	55	53	26	5	-
American Indian 1/, both sexes	1	1	2	-	1	-	-	4	1	-	-	-
Male	-	1	2	-	-	-	-	1	1	-	-	-
Female	1	-	-	-	1	-	-	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	5	6	11	6	7	7	1	-	-	-
Male	-	2	3	4	4	4	3	3	1	-	-	-
Female	1	1	2	2	7	2	4	4	-	-	-	-
Georgia												
All races, both sexes	174	228	314	311	368	412	483	425	267	95	18	-
Male	124	156	213	208	226	223	205	150	91	22	-	-
Female	50	72	101	103	142	189	278	275	176	73	18	-
White, both sexes	124	157	237	228	275	309	387	347	210	82	11	-
Male	95	116	177	153	176	180	163	130	68	17	-	-
Female	29	41	60	75	99	129	224	217	142	65	11	-
Black, both sexes	49	70	75	75	87	101	95	77	55	13	7	-
Male	28	39	35	49	47	43	42	19	21	5	-	-
Female	21	31	40	26	40	58	53	58	34	8	7	-
American Indian 1/, both sexes	1	-	1	2	-	-	-	-	1	-	-	-
Male	1	-	-	2	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	6	6	2	1	1	1	-	-	-
Male	-	1	1	4	3	-	-	1	1	-	-	-
Female	-	-	-	2	3	2	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Hawaii												
All races, both sexes	21	17	24	25	26	50	80	57	41	23	2	-
Male	15	16	16	14	13	23	47	31	14	8	-	-
Female	6	1	8	11	13	27	33	26	27	15	2	-
White, both sexes	7	4	7	6	5	12	19	10	5	8	-	-
Male	6	4	5	3	4	6	15	6	3	1	-	-
Female	1	-	2	3	1	6	4	4	2	7	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	13	17	19	21	38	60	47	36	15	2	-
Male	9	12	11	11	9	17	32	25	11	7	-	-
Female	5	1	6	8	12	21	28	22	25	8	2	-
Idaho												
All races, both sexes	44	53	67	77	70	105	114	114	73	20	2	-
Male	36	41	45	49	50	75	64	50	32	9	-	-
Female	8	12	22	28	20	30	50	64	41	11	2	-
White, both sexes	43	52	66	75	70	104	113	114	72	20	2	-
Male	35	40	45	48	50	74	63	50	31	9	-	-
Female	8	12	21	27	20	30	50	64	41	11	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	1	1	-	-	-	-	-
Male	1	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	-	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Illinois												
All races, both sexes	315	379	491	538	692	907	1,197	1,198	818	322	76	-
Male	239	281	347	345	408	493	589	498	242	72	19	-
Female	76	98	144	193	284	414	608	700	576	250	57	-
White, both sexes	251	306	404	448	565	755	1,045	1,071	739	299	63	-
Male	195	236	289	299	344	423	531	456	224	67	16	-
Female	56	70	115	149	221	332	514	615	515	232	47	-
Black, both sexes	62	64	80	80	119	140	136	106	72	22	13	-
Male	42	38	53	39	60	65	49	32	17	5	3	-
Female	20	26	27	41	59	75	87	74	55	17	10	-
American Indian 1/, both sexes	-	1	1	-	1	1	-	1	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	8	6	10	7	11	16	20	7	1	-	-
Male	2	6	5	7	4	4	9	10	1	-	-	-
Female	-	2	1	3	3	7	7	10	6	1	-	-
Indiana												
All races, both sexes	178	207	273	324	373	447	535	484	345	109	14	-
Male	138	152	194	202	222	254	252	205	107	25	-	-
Female	40	55	79	122	151	193	283	279	238	84	14	-
White, both sexes	167	191	258	304	356	421	519	461	331	105	12	-
Male	130	140	184	189	212	239	244	194	104	23	-	-
Female	37	51	74	115	144	182	275	267	227	82	12	-
Black, both sexes	11	14	14	17	16	25	15	23	14	4	2	-
Male	8	10	10	11	9	14	7	11	3	2	-	-
Female	3	4	4	6	7	11	8	12	11	2	2	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	2	1	1	1	-	-	-	-	-
Male	-	2	-	2	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Iowa												
All races, both sexes	62	64	102	129	180	244	310	345	272	98	29	-
Male	45	49	86	81	122	142	160	168	82	29	4	-
Female	17	15	16	48	58	102	150	177	190	69	25	-
White, both sexes	60	63	101	127	179	243	306	344	271	98	29	-
Male	44	48	85	81	121	141	159	168	82	29	4	-
Female	16	15	16	46	58	102	147	176	189	69	25	-
Black, both sexes	2	-	1	2	-	-	3	1	1	-	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	1	-	-	2	-	-	2	1	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Kansas												
All races, both sexes	45	80	75	75	121	145	203	248	146	55	11	-
Male	33	52	63	55	71	83	107	112	44	14	3	-
Female	12	28	12	20	50	62	96	136	102	41	8	-
White, both sexes	41	73	70	72	113	140	195	235	144	54	11	-
Male	31	47	58	53	68	79	104	106	42	13	3	-
Female	10	26	12	19	45	61	91	129	102	41	8	-
Black, both sexes	4	6	5	2	5	3	7	12	2	1	-	-
Male	2	4	5	2	1	2	3	5	2	1	-	-
Female	2	2	-	-	4	1	4	7	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	2	2	1	1	-	-	-	-
Male	-	1	-	-	2	2	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Kentucky												
All races, both sexes	211	261	285	313	404	442	496	444	259	80	12	-
Male	154	176	178	205	246	267	236	171	75	16	5	-
Female	57	85	107	108	158	175	260	273	184	64	7	-
White, both sexes	192	245	268	292	373	413	474	420	240	74	10	-
Male	141	166	171	192	224	251	225	163	71	15	5	-
Female	51	79	97	100	149	162	249	257	169	59	5	-
Black, both sexes	19	14	15	20	31	28	22	22	19	6	2	-
Male	13	8	5	13	22	16	11	8	4	1	-	-
Female	6	6	10	7	9	12	11	14	15	5	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	-	1	-	2	-	-	-	-
Male	-	1	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	2	-	-	-	-
Louisiana												
All races, both sexes	191	276	292	288	309	391	438	348	183	69	17	-
Male	133	215	183	197	168	217	193	145	51	17	4	-
Female	58	61	109	91	141	174	245	203	132	52	13	-
White, both sexes	112	160	207	194	206	298	338	274	145	41	10	-
Male	83	127	137	141	114	176	147	114	42	5	2	-
Female	29	33	70	53	92	122	191	160	103	36	8	-
Black, both sexes	77	112	84	90	99	92	96	72	37	28	7	-
Male	50	84	45	53	52	40	44	30	9	12	2	-
Female	27	28	39	37	47	52	52	42	28	16	5	-
American Indian 1/, both sexes	-	2	-	1	-	1	1	1	-	-	-	-
Male	-	2	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	3	4	-	3	1	1	-	-	-
Male	-	2	1	2	2	-	1	-	-	-	-	-
Female	2	-	-	1	2	-	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Maine												
All races, both sexes	28	33	55	45	58	111	101	109	91	26	11	-
Male	22	27	40	27	30	62	56	50	24	7	-	-
Female	6	6	15	18	28	49	45	59	67	19	11	-
White, both sexes	28	33	55	44	58	109	100	109	91	26	11	-
Male	22	27	40	27	30	62	56	50	24	7	-	-
Female	6	6	15	17	28	47	44	59	67	19	11	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	114	129	157	205	263	394	496	440	271	96	28	-
Male	76	95	106	131	145	199	231	162	77	24	4	-
Female	38	34	51	74	118	195	265	278	194	72	24	-
White, both sexes	81	83	104	131	183	300	390	363	234	80	22	-
Male	56	62	74	89	103	156	182	140	67	21	2	-
Female	25	21	30	42	80	144	208	223	167	59	20	-
Black, both sexes	30	42	48	67	74	91	105	75	33	16	6	-
Male	17	29	29	38	40	41	49	22	8	3	2	-
Female	13	13	19	29	34	50	56	53	25	13	4	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	5	7	6	3	1	2	4	-	-	-
Male	2	4	3	4	2	2	-	-	2	-	-	-
Female	-	-	2	3	4	1	1	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Massachusetts												
All races, both sexes	92	148	179	214	297	397	547	551	369	157	27	-
Male	79	116	126	148	193	228	246	237	109	27	2	-
Female	13	32	53	66	104	169	301	314	260	130	25	-
White, both sexes	88	136	171	204	281	375	528	538	362	154	26	-
Male	75	109	122	144	182	217	240	234	106	27	2	-
Female	13	27	49	60	99	158	288	304	256	127	24	-
Black, both sexes	2	8	7	9	12	16	15	10	7	2	1	-
Male	2	5	4	4	7	8	3	3	3	-	-	-
Female	-	3	3	5	5	8	12	7	4	2	1	-
American Indian 1/, both sexes	1	-	-	-	2	3	-	-	-	-	-	-
Male	1	-	-	-	2	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	1	1	2	3	4	3	-	1	-	-
Male	1	2	-	-	2	1	3	-	-	-	-	-
Female	-	2	1	1	-	2	1	3	-	1	-	-
Michigan												
All races, both sexes	213	269	335	362	495	713	883	851	524	183	38	-
Male	145	181	229	230	299	396	434	313	168	38	3	-
Female	68	88	106	132	196	317	449	538	356	145	35	-
White, both sexes	182	220	275	333	444	643	808	767	477	170	28	-
Male	129	159	193	214	280	364	396	284	154	36	3	-
Female	53	61	82	119	164	279	412	483	323	134	25	-
Black, both sexes	28	44	57	25	47	64	70	79	45	13	10	-
Male	13	18	35	14	16	31	33	25	14	2	-	-
Female	15	26	22	11	31	33	37	54	31	11	10	-
American Indian 1/, both sexes	1	2	2	4	1	2	-	3	1	-	-	-
Male	1	1	1	2	1	-	-	2	-	-	-	-
Female	-	1	1	2	-	2	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	1	-	3	4	5	2	1	-	-	-
Male	2	3	-	-	2	1	5	2	-	-	-	-
Female	-	-	1	-	1	3	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Minnesota												
All races, both sexes	66	81	88	123	146	183	265	252	167	69	10	-
Male	55	63	73	87	89	105	140	105	72	13	-	-
Female	11	18	15	36	57	78	125	147	95	56	10	-
White, both sexes	62	77	86	121	140	182	262	251	166	68	10	-
Male	52	60	72	87	86	104	138	104	71	13	-	-
Female	10	17	14	34	54	78	124	147	95	55	10	-
Black, both sexes	4	1	1	1	2	-	-	-	1	1	-	-
Male	3	-	1	-	2	-	-	-	1	-	-	-
Female	1	1	-	1	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	3	-	1	2	-	2	-	-	-	-	-
Male	-	3	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	1	-	2	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	142	186	218	173	216	283	266	222	145	56	19	1
Male	100	147	147	115	135	156	145	95	37	16	3	1
Female	42	39	71	58	81	127	121	127	108	40	16	-
White, both sexes	81	104	149	116	155	201	211	168	105	34	13	1
Male	64	87	108	84	101	122	121	72	24	8	2	1
Female	17	17	41	32	54	79	90	96	81	26	11	-
Black, both sexes	60	81	68	57	61	82	55	53	39	22	6	-
Male	35	59	39	31	34	34	24	23	13	8	1	-
Female	25	22	29	26	27	48	31	30	26	14	5	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Missouri												
All races, both sexes	204	234	314	348	456	551	696	644	455	202	42	-
Male	154	173	220	236	284	315	326	254	142	46	6	-
Female	50	61	94	112	172	236	370	390	313	156	36	-
White, both sexes	169	216	273	308	421	502	643	604	432	186	37	-
Male	129	162	196	211	269	289	301	242	134	43	6	-
Female	40	54	77	97	152	213	342	362	298	143	31	-
Black, both sexes	34	15	37	38	34	48	51	40	23	15	5	-
Male	25	10	21	23	14	25	24	12	8	3	-	-
Female	9	5	16	15	20	23	27	28	15	12	5	-
American Indian 1/, both sexes	1	1	3	-	-	1	1	-	-	1	-	-
Male	-	1	2	-	-	1	1	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	2	1	-	1	-	-	-	-	-
Male	-	-	1	2	1	-	-	-	-	-	-	-
Female	-	2	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	13	19	34	32	36	56	60	72	40	11	2	-
Male	9	14	26	20	29	38	39	35	11	3	-	-
Female	4	5	8	12	7	18	21	37	29	8	2	-
White, both sexes	12	16	32	29	34	54	59	69	40	11	2	-
Male	8	13	24	19	28	38	39	35	11	3	-	-
Female	4	3	8	10	6	16	20	34	29	8	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	3	2	3	2	2	1	3	-	-	-	-
Male	1	1	2	1	1	-	-	-	-	-	-	-
Female	-	2	-	2	1	2	1	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Nebraska												
All races, both sexes	25	28	46	38	52	83	100	127	86	25	2	-
Male	16	22	32	27	35	56	49	64	28	10	-	-
Female	9	6	14	11	17	27	51	63	58	15	2	-
White, both sexes	24	26	45	36	50	81	99	122	86	24	1	-
Male	16	20	31	26	34	55	49	61	28	10	-	-
Female	8	6	14	10	16	26	50	61	58	14	1	-
Black, both sexes	1	2	1	1	2	-	-	2	-	1	1	-
Male	-	2	1	-	1	-	-	2	-	-	-	-
Female	1	-	-	1	1	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	1	-	2	1	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Nevada												
All races, both sexes	35	56	68	76	78	77	74	91	32	15	2	-
Male	26	46	40	52	49	46	40	49	11	5	-	-
Female	9	10	28	24	29	31	34	42	21	10	2	-
White, both sexes	31	51	61	69	72	70	72	88	30	13	2	-
Male	22	42	36	48	46	43	40	47	11	5	-	-
Female	9	9	25	21	26	27	32	41	19	8	2	-
Black, both sexes	2	2	5	3	3	5	-	1	1	1	-	-
Male	2	2	4	1	1	1	-	-	-	-	-	-
Female	-	-	1	2	2	4	-	1	1	1	-	-
American Indian 1/, both sexes	-	1	1	2	1	1	1	-	1	1	-	-
Male	-	-	-	2	-	1	-	-	-	-	-	-
Female	-	1	1	-	1	-	1	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	2	2	1	1	2	-	-	-	-
Male	2	2	-	1	2	1	-	2	-	-	-	-
Female	-	-	1	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
New Hampshire												
All races, both sexes	12	23	26	38	36	60	91	75	64	20	7	-
Male	10	21	22	27	22	34	49	29	24	8	1	-
Female	2	2	4	11	14	26	42	46	40	12	6	-
White, both sexes	12	23	26	38	36	59	91	75	64	20	7	-
Male	10	21	22	27	22	33	49	29	24	8	1	-
Female	2	2	4	11	14	26	42	46	40	12	6	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	134	239	246	287	405	656	863	852	578	216	35	-
Male	107	173	162	179	231	332	400	350	175	56	4	-
Female	27	66	84	108	174	324	463	502	403	160	31	-
White, both sexes	104	177	197	222	325	550	784	795	543	200	30	-
Male	84	133	130	138	191	285	370	330	167	54	4	-
Female	20	44	67	84	134	265	414	465	376	146	26	-
Black, both sexes	25	51	38	57	66	91	68	49	27	15	5	-
Male	18	29	22	38	30	40	25	16	6	2	-	-
Female	7	22	16	19	36	51	43	33	21	13	5	-
American Indian 1/, both sexes	-	-	-	3	-	2	-	-	2	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	2	-	1	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	5	11	11	5	14	13	11	8	6	1	-	-
Male	5	11	10	2	10	6	5	4	2	-	-	-
Female	-	-	1	3	4	7	6	4	4	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
New Mexico												
All races, both sexes	23	31	38	42	68	73	106	84	55	25	4	-
Male	15	24	27	28	48	39	54	38	24	4	-	-
Female	8	7	11	14	20	34	52	46	31	21	4	-
White, both sexes	21	28	36	37	64	71	100	79	52	22	4	-
Male	13	21	27	25	45	38	49	34	23	3	-	-
Female	8	7	9	12	19	33	51	45	29	19	4	-
Black, both sexes	-	-	-	-	1	-	-	-	1	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	2	3	2	5	2	2	6	5	-	1	-	-
Male	2	3	-	3	2	1	5	4	-	1	-	-
Female	-	-	2	2	-	1	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	2	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	1	1	-	-
New York												
All races, both sexes	261	428	551	646	921	1,262	1,725	1,804	1,405	573	124	-
Male	193	318	368	379	544	649	786	674	450	137	15	-
Female	68	110	183	267	377	613	939	1,130	955	436	109	-
White, both sexes	197	340	454	509	750	1,100	1,525	1,633	1,306	519	98	-
Male	157	258	319	311	458	581	715	628	422	130	14	-
Female	40	82	135	198	292	519	810	1,005	884	389	84	-
Black, both sexes	52	75	85	110	140	141	161	148	79	48	24	-
Male	27	47	40	53	68	55	50	34	24	4	1	-
Female	25	28	45	57	72	86	111	114	55	44	23	-
American Indian 1/, both sexes	1	2	-	1	2	2	1	1	1	-	-	-
Male	1	2	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	2	1	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	11	11	12	26	29	19	38	22	19	6	2	-
Male	8	11	9	15	18	12	21	12	4	3	-	-
Female	3	-	3	11	11	7	17	10	15	3	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
North Carolina												
All races, both sexes	182	242	290	319	454	562	657	570	375	118	20	-
Male	133	176	210	195	263	307	302	207	127	24	3	-
Female	49	66	80	124	191	255	355	363	248	94	17	-
White, both sexes	146	173	221	246	376	445	550	499	329	99	16	-
Male	112	133	168	157	219	264	261	187	114	23	2	-
Female	34	40	53	89	157	181	289	312	215	76	14	-
Black, both sexes	34	62	63	67	70	113	99	68	42	18	4	-
Male	20	37	37	33	41	43	38	19	12	1	1	-
Female	14	25	26	34	29	70	61	49	30	17	3	-
American Indian 1/, both sexes	2	4	5	6	7	2	5	1	2	1	-	-
Male	1	3	5	5	3	-	1	-	-	-	-	-
Female	1	1	-	1	4	2	4	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	3	1	-	1	2	3	2	2	-	-	-
Male	-	3	-	-	-	-	2	1	1	-	-	-
Female	-	-	1	-	1	2	1	1	1	-	-	-
North Dakota												
All races, both sexes	12	24	20	31	34	69	78	78	55	28	5	-
Male	9	18	15	25	23	45	42	38	11	9	-	-
Female	3	6	5	6	11	24	36	40	44	19	5	-
White, both sexes	11	20	20	30	33	66	78	77	55	28	5	-
Male	8	16	15	24	22	44	42	37	11	9	-	-
Female	3	4	5	6	11	22	36	40	44	19	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	4	-	1	1	3	-	1	-	-	-	-
Male	1	2	-	1	1	1	-	1	-	-	-	-
Female	-	2	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Ohio												
All races, both sexes	285	351	466	537	641	914	1,158	1,063	686	258	51	-
Male	218	255	329	366	388	503	527	403	218	65	6	-
Female	67	96	137	171	253	411	631	660	468	193	45	-
White, both sexes	243	306	421	486	586	828	1,080	1,004	653	248	47	-
Male	194	227	300	336	363	469	502	386	208	60	4	-
Female	49	79	121	150	223	359	578	618	445	188	43	-
Black, both sexes	40	45	44	48	55	83	73	57	33	10	4	-
Male	22	28	29	28	25	32	24	16	10	5	2	-
Female	18	17	15	20	30	51	49	41	23	5	2	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	3	-	2	5	1	-	-	-	-
Male	1	-	-	2	-	1	1	1	-	-	-	-
Female	-	-	1	1	-	1	4	-	-	-	-	-
Oklahoma												
All races, both sexes	91	110	126	161	230	269	317	274	179	76	14	1
Male	63	76	82	98	141	142	161	105	59	18	3	1
Female	28	34	44	63	89	127	156	169	120	58	11	-
White, both sexes	74	96	106	144	217	231	287	253	167	70	10	1
Male	54	69	70	90	134	127	147	96	53	17	1	1
Female	20	27	36	54	83	104	140	157	114	53	9	-
Black, both sexes	10	7	11	12	7	18	12	13	9	5	2	-
Male	4	3	6	5	4	3	8	5	4	1	1	-
Female	6	4	5	7	3	15	4	8	5	4	1	-
American Indian 1/, both sexes	7	7	9	4	5	18	18	8	3	1	2	-
Male	5	4	6	2	2	12	6	4	2	-	1	-
Female	2	3	3	2	3	6	12	4	1	1	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	2	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Oregon												
All races, both sexes	66	67	76	114	155	177	265	251	168	52	8	-
Male	52	52	53	72	103	99	145	109	86	14	1	-
Female	14	15	23	42	52	78	120	142	82	38	7	-
White, both sexes	63	62	74	109	152	169	257	244	166	51	8	-
Male	50	48	51	69	100	94	142	106	86	13	1	-
Female	13	14	23	40	52	75	115	138	80	38	7	-
Black, both sexes	1	1	2	2	1	3	3	3	2	-	-	-
Male	-	1	2	2	1	2	2	1	-	-	-	-
Female	1	-	-	-	-	1	1	2	2	-	-	-
American Indian 1/, both sexes	1	1	-	1	2	2	2	-	-	-	-	-
Male	1	-	-	-	2	2	1	-	-	-	-	-
Female	-	1	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	-	2	-	3	3	4	-	1	-	-
Male	1	3	-	1	-	1	-	2	-	1	-	-
Female	-	-	-	1	-	2	3	2	-	-	-	-
Pennsylvania												
All races, both sexes	318	420	551	590	857	1,213	1,520	1,419	934	322	68	-
Male	255	307	372	395	513	684	737	552	287	75	3	-
Female	63	113	179	195	344	529	783	867	647	247	65	-
White, both sexes	285	379	495	548	790	1,125	1,439	1,355	892	311	63	-
Male	230	281	336	376	477	640	696	534	278	74	3	-
Female	55	98	159	172	313	485	743	821	614	237	60	-
Black, both sexes	30	38	53	40	61	82	80	61	39	10	5	-
Male	22	23	33	18	35	42	41	17	8	-	-	-
Female	8	15	20	22	26	40	39	44	31	10	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	3	2	6	6	1	3	3	1	-	-
Male	3	3	3	1	1	2	-	1	1	1	-	-
Female	-	-	-	1	5	4	1	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Rhode Island												
All races, both sexes	29	35	49	57	86	111	162	173	117	54	9	-
Male	23	26	37	34	51	61	74	71	37	12	1	-
Female	6	9	12	23	35	50	88	102	80	42	8	-
White, both sexes	27	34	45	53	78	109	156	170	115	53	9	-
Male	22	25	34	32	48	61	72	70	37	12	1	-
Female	5	9	11	21	30	48	84	100	78	41	8	-
Black, both sexes	1	1	4	2	7	2	6	3	2	-	-	-
Male	1	1	3	1	2	-	2	1	-	-	-	-
Female	-	-	1	1	5	2	4	2	2	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	1	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
South Carolina												
All races, both sexes	134	206	220	249	264	320	344	285	167	56	6	-
Male	105	155	170	169	149	175	149	110	39	16	1	-
Female	29	51	50	80	115	145	195	175	128	40	5	-
White, both sexes	88	153	152	166	188	236	263	213	129	39	3	-
Male	71	115	120	124	108	132	117	91	30	10	1	-
Female	17	38	32	42	80	104	146	122	99	29	2	-
Black, both sexes	43	53	66	83	74	84	81	71	38	17	3	-
Male	31	40	49	45	40	43	32	19	9	6	-	-
Female	12	13	17	38	34	41	49	52	29	11	3	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	-	2	-	-	1	-	-	-	-
Male	2	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
South Dakota												
All races, both sexes	27	33	35	48	55	79	104	102	66	30	9	-
Male	19	27	22	33	37	50	58	42	24	9	4	-
Female	8	6	13	15	18	29	46	60	42	21	5	-
White, both sexes	22	26	30	43	51	75	101	99	64	30	9	-
Male	15	23	19	31	33	47	58	41	23	9	4	-
Female	7	3	11	12	18	28	43	58	41	21	5	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	7	5	5	4	3	2	3	2	-	-	-
Male	3	4	3	2	4	2	-	1	1	-	-	-
Female	1	3	2	3	-	1	2	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Tennessee												
All races, both sexes	240	338	347	408	453	559	703	649	382	141	19	-
Male	178	232	231	275	280	304	316	243	108	24	5	-
Female	62	106	116	133	173	255	387	406	274	117	14	-
White, both sexes	194	280	293	346	388	500	616	584	331	126	15	-
Male	148	195	197	238	242	275	282	224	95	23	1	-
Female	46	85	96	108	146	225	334	360	236	103	14	-
Black, both sexes	45	56	52	61	62	59	86	64	50	15	4	-
Male	29	36	32	37	36	29	34	19	13	1	4	-
Female	16	20	20	24	26	30	52	45	37	14	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	-	2	-	1	-	1	-	-	-
Male	1	1	2	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Texas												
All races, both sexes	549	678	885	975	1,296	1,696	1,978	1,717	1,112	418	95	-
Male	402	485	598	626	799	996	996	723	360	82	11	-
Female	147	193	287	349	497	700	982	994	752	336	84	-
White, both sexes	449	564	751	844	1,136	1,497	1,828	1,560	1,011	364	83	-
Male	333	405	519	564	709	893	934	660	333	71	10	-
Female	116	159	232	280	427	604	894	900	678	293	73	-
Black, both sexes	94	106	117	122	145	182	137	141	97	49	12	-
Male	64	74	69	57	81	90	58	57	26	8	1	-
Female	30	32	48	65	64	92	79	84	71	41	11	-
American Indian 1/, both sexes	1	1	2	1	-	-	3	2	-	-	-	-
Male	1	1	2	1	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	7	15	8	15	17	10	14	4	5	-	-
Male	4	5	8	4	9	13	3	5	1	3	-	-
Female	1	2	7	4	6	4	7	9	3	2	-	-
Utah												
All races, both sexes	20	29	38	46	45	96	91	83	60	17	1	-
Male	16	21	26	38	28	64	44	46	23	6	-	-
Female	4	8	12	8	17	32	47	37	37	11	1	-
White, both sexes	19	27	37	46	44	95	87	82	58	17	1	-
Male	15	20	25	38	28	63	42	46	22	6	-	-
Female	4	7	12	8	16	32	45	36	36	11	1	-
Black, both sexes	1	1	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	2	1	1	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Vermont												
All races, both sexes	1	12	9	16	25	40	45	48	20	7	3	-
Male	1	8	7	9	16	27	20	21	11	3	-	-
Female	-	4	2	7	9	13	25	27	9	4	3	-
White, both sexes	1	12	9	16	25	40	45	47	20	7	3	-
Male	1	8	7	9	16	27	20	21	11	3	-	-
Female	-	4	2	7	9	13	25	26	9	4	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	154	214	226	303	328	419	519	477	295	85	13	-
Male	123	160	158	181	202	234	248	194	96	21	1	-
Female	31	54	68	122	126	185	271	283	199	64	12	-
White, both sexes	119	152	178	228	252	345	437	413	255	75	9	-
Male	97	123	127	141	169	195	211	178	84	19	-	-
Female	22	29	51	87	83	150	226	235	171	56	9	-
Black, both sexes	35	57	45	70	73	69	73	59	38	9	4	-
Male	26	35	29	37	32	35	33	14	12	2	1	-
Female	9	22	16	33	41	34	40	45	26	7	3	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	5	3	5	2	5	8	5	2	1	-	-
Male	-	2	2	3	-	4	3	2	-	-	-	-
Female	-	3	1	2	2	1	5	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Washington												
All races, both sexes	91	133	173	193	218	323	429	370	295	105	21	-
Male	69	94	137	133	134	193	234	159	104	31	3	-
Female	22	39	36	60	84	130	195	211	191	74	18	-
White, both sexes	82	115	163	183	211	302	406	354	285	104	20	-
Male	62	81	132	126	129	181	222	156	99	30	3	-
Female	20	34	31	57	82	121	184	198	186	74	17	-
Black, both sexes	2	8	3	2	2	6	7	6	2	-	-	-
Male	2	5	3	2	2	4	2	1	1	-	-	-
Female	-	3	-	-	-	2	5	5	1	-	-	-
American Indian 1/, both sexes	2	4	3	2	-	3	3	-	-	-	-	-
Male	2	3	-	2	-	2	1	-	-	-	-	-
Female	-	1	3	-	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	6	4	6	5	12	13	10	8	1	1	-
Male	3	5	2	3	3	6	9	2	4	1	-	-
Female	2	1	2	3	2	6	4	8	4	-	1	-
West Virginia												
All races, both sexes	53	74	82	118	136	192	244	171	114	38	8	-
Male	31	45	48	78	80	98	115	79	39	8	1	-
Female	22	29	34	40	56	94	129	92	75	30	7	-
White, both sexes	52	73	79	116	136	188	242	170	114	36	8	-
Male	31	45	47	76	80	97	115	79	39	7	1	-
Female	21	28	32	40	56	91	127	91	75	29	7	-
Black, both sexes	1	1	3	2	-	4	2	1	-	2	-	-
Male	-	-	1	2	-	1	-	-	-	1	-	-
Female	1	1	2	-	-	3	2	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute myocardial infarction (I21-I22)												
Wisconsin												
All races, both sexes	128	173	193	223	332	422	521	569	342	141	24	-
Male	106	139	135	158	224	278	239	247	116	30	7	-
Female	22	34	58	65	108	144	282	322	226	111	17	-
White, both sexes	113	166	180	207	319	399	513	564	340	140	24	-
Male	94	132	128	146	217	265	236	245	116	30	7	-
Female	19	34	52	61	102	134	277	319	224	110	17	-
Black, both sexes	10	3	5	10	7	18	3	3	1	1	-	-
Male	8	3	1	7	3	11	-	1	-	-	-	-
Female	2	-	4	3	4	7	3	2	1	1	-	-
American Indian 1/, both sexes	4	2	6	2	1	3	4	2	-	-	-	-
Male	3	2	5	2	1	2	2	1	-	-	-	-
Female	1	-	1	-	-	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	4	5	2	1	-	1	-	-	-
Male	1	2	1	3	3	-	1	-	-	-	-	-
Female	-	-	1	1	2	2	-	-	1	-	-	-
Wyoming												
All races, both sexes	15	19	29	28	34	45	37	34	22	4	2	-
Male	11	12	23	17	21	27	25	14	11	-	1	-
Female	4	7	6	11	13	18	12	20	11	4	1	-
White, both sexes	14	18	29	28	32	45	37	34	22	4	2	-
Male	10	12	23	17	21	27	25	14	11	-	1	-
Female	4	6	6	11	11	18	12	20	11	4	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
United States												
All races, both sexes	3,565	-	1	-	1	1	2	6	5	24	58	127
Male	1,823	-	-	-	1	-	2	4	3	17	42	85
Female	1,742	-	1	-	-	1	-	2	2	7	16	42
White, both sexes	2,924	-	1	-	1	1	2	3	4	18	42	87
Male	1,480	-	-	-	1	-	2	2	2	15	30	66
Female	1,444	-	1	-	-	1	-	1	2	3	12	21
Black, both sexes	558	-	-	-	-	-	-	3	1	6	13	36
Male	296	-	-	-	-	-	-	2	1	2	10	18
Female	262	-	-	-	-	-	-	1	-	4	3	18
American Indian 1/, both sexes	44	-	-	-	-	-	-	-	-	-	3	4
Male	26	-	-	-	-	-	-	-	-	-	2	1
Female	18	-	-	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	27	-	-	-	-	-	-	-	-	1	1	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	14	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	37	-	-	-	-	-	-	-	-	-	-	5
Male	26	-	-	-	-	-	-	-	-	-	-	4
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	5
Male	22	-	-	-	-	-	-	-	-	-	-	4
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Arkansas												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	160	-	-	-	-	-	2	1	-	2	3	5
Male	78	-	-	-	-	-	2	1	-	1	3	5
Female	82	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	132	-	-	-	-	-	2	1	-	2	3	4
Male	65	-	-	-	-	-	2	1	-	1	3	4
Female	67	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Colorado												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Delaware												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Florida												
All races, both sexes	92	-	-	-	-	-	-	-	-	-	1	-
Male	46	-	-	-	-	-	-	-	-	-	1	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	86	-	-	-	-	-	-	-	-	-	1	-
Male	42	-	-	-	-	-	-	-	-	-	1	-
Female	44	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	61	-	-	-	-	-	-	-	1	1	2	3
Male	29	-	-	-	-	-	-	-	-	-	1	3
Female	32	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	43	-	-	-	-	-	-	-	1	-	1	3
Male	24	-	-	-	-	-	-	-	-	-	-	3
Female	19	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	18	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Hawaii												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Illinois												
All races, both sexes	133	-	-	-	-	-	-	-	-	2	2	6
Male	69	-	-	-	-	-	-	-	-	2	2	5
Female	64	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	112	-	-	-	-	-	-	-	-	1	2	5
Male	60	-	-	-	-	-	-	-	-	1	2	5
Female	52	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	19	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Iowa												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	2	1
Male	14	-	-	-	-	-	-	-	-	-	2	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	2	1
Male	13	-	-	-	-	-	-	-	-	-	2	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Kentucky												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	60	-	-	-	-	-	-	-	-	3	2	1
Male	35	-	-	-	-	-	-	-	-	3	2	-
Female	25	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	40	-	-	-	-	-	-	-	-	2	-	-
Male	23	-	-	-	-	-	-	-	-	2	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	-	-	-	-	1	2	1
Male	12	-	-	-	-	-	-	-	-	1	2	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Maine												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Massachusetts												
All races, both sexes	56	-	-	-	-	-	-	-	-	1	-	1
Male	28	-	-	-	-	-	-	-	-	1	-	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	54	-	-	-	-	-	-	-	-	1	-	1
Male	26	-	-	-	-	-	-	-	-	1	-	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	47	-	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	37	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Minnesota												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	16	-	1	-	1	-	-	1	-	-	2	1
Male	10	-	-	-	1	-	-	1	-	-	1	1
Female	6	-	1	-	-	-	-	-	-	-	1	-
White, both sexes	9	-	1	-	1	-	-	1	-	-	-	-
Male	5	-	-	-	1	-	-	1	-	-	-	-
Female	4	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Missouri												
All races, both sexes	65	-	-	-	-	-	-	-	-	-	2	-
Male	23	-	-	-	-	-	-	-	-	-	2	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	59	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Nebraska												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
New Hampshire												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	118	-	-	-	-	-	-	-	1	-	1	1
Male	48	-	-	-	-	-	-	-	-	-	1	1
Female	70	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	108	-	-	-	-	-	-	-	1	-	-	1
Male	40	-	-	-	-	-	-	-	-	-	-	1
Female	68	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
New Mexico												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	185	-	-	-	-	-	-	-	-	-	1	10
Male	92	-	-	-	-	-	-	-	-	-	-	5
Female	93	-	-	-	-	-	-	-	-	-	1	5
White, both sexes	168	-	-	-	-	-	-	-	-	-	1	9
Male	83	-	-	-	-	-	-	-	-	-	-	5
Female	85	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
North Carolina												
All races, both sexes	80	-	-	-	-	-	-	-	-	1	-	2
Male	48	-	-	-	-	-	-	-	-	1	-	2
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	59	-	-	-	-	-	-	-	-	1	-	1
Male	34	-	-	-	-	-	-	-	-	1	-	1
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Ohio												
All races, both sexes	117	-	-	-	-	-	-	1	-	-	1	6
Male	51	-	-	-	-	-	-	-	-	-	-	4
Female	66	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	106	-	-	-	-	-	-	1	-	-	1	6
Male	46	-	-	-	-	-	-	-	-	-	-	4
Female	60	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	753	-	-	-	-	1	-	-	-	2	14	18
Male	380	-	-	-	-	-	-	-	-	2	12	11
Female	373	-	-	-	-	1	-	-	-	-	2	7
White, both sexes	665	-	-	-	-	1	-	-	-	2	11	13
Male	335	-	-	-	-	-	-	-	-	2	10	10
Female	330	-	-	-	-	1	-	-	-	-	1	3
Black, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	32	-	-	-	-	-	-	-	-	-	3	4
Male	18	-	-	-	-	-	-	-	-	-	2	1
Female	14	-	-	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Oregon												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	150	-	-	-	-	-	-	-	-	-	3	6
Male	74	-	-	-	-	-	-	-	-	-	1	5
Female	76	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	143	-	-	-	-	-	-	-	-	-	3	6
Male	72	-	-	-	-	-	-	-	-	-	1	5
Female	71	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Rhode Island												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	227	-	-	-	-	-	-	3	1	5	7	13
Male	139	-	-	-	-	-	-	2	1	2	5	10
Female	88	-	-	-	-	-	-	1	-	3	2	3
White, both sexes	113	-	-	-	-	-	-	-	-	3	3	4
Male	74	-	-	-	-	-	-	-	-	2	2	3
Female	39	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	114	-	-	-	-	-	-	3	1	2	4	9
Male	65	-	-	-	-	-	-	2	1	-	3	7
Female	49	-	-	-	-	-	-	1	-	2	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
South Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	42	-	-	-	-	-	-	-	-	-	1	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	37	-	-	-	-	-	-	-	-	-	1	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Texas												
All races, both sexes	135	-	-	-	-	-	-	-	2	1	1	3
Male	68	-	-	-	-	-	-	-	2	1	-	1
Female	67	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	114	-	-	-	-	-	-	-	2	1	1	3
Male	57	-	-	-	-	-	-	-	2	1	-	1
Female	57	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Vermont												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	571	-	-	-	-	-	-	-	-	4	10	42
Male	317	-	-	-	-	-	-	-	-	3	7	24
Female	254	-	-	-	-	-	-	-	-	1	3	18
White, both sexes	395	-	-	-	-	-	-	-	-	3	8	22
Male	223	-	-	-	-	-	-	-	-	3	6	16
Female	172	-	-	-	-	-	-	-	-	-	2	6
Black, both sexes	174	-	-	-	-	-	-	-	-	1	2	20
Male	92	-	-	-	-	-	-	-	-	-	1	8
Female	82	-	-	-	-	-	-	-	-	1	1	12
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Washington												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	27	-	-	-	-	-	-	-	-	1	-	1
Male	15	-	-	-	-	-	-	-	-	1	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	1	-	1
Male	15	-	-	-	-	-	-	-	-	1	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other acute ischemic heart diseases (I24)												
Wisconsin												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
United States												
All races, both sexes	224	238	277	247	359	401	534	503	365	156	36	-
Male	160	174	197	161	203	221	229	179	108	30	7	-
Female	64	64	80	86	156	180	305	324	257	126	29	-
White, both sexes	150	173	210	203	290	335	466	449	325	134	30	-
Male	117	128	152	131	165	184	201	159	94	25	6	-
Female	33	45	58	72	125	151	265	290	231	109	24	-
Black, both sexes	65	58	57	41	57	58	61	44	34	20	4	-
Male	38	41	38	27	31	33	25	15	11	4	-	-
Female	27	17	19	14	26	25	36	29	23	16	4	-
American Indian 1/, both sexes	4	5	5	3	8	2	3	4	1	-	2	-
Male	1	3	4	3	5	1	2	3	-	-	1	-
Female	3	2	1	-	3	1	1	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	5	2	5	-	4	6	4	6	5	2	-	-
Male	4	2	3	-	2	3	1	2	3	1	-	-
Female	1	-	2	-	2	3	3	4	2	1	-	-
Alabama												
All races, both sexes	2	-	2	1	2	4	7	2	2	3	-	-
Male	1	-	2	1	1	2	2	1	-	2	-	-
Female	1	-	-	-	1	2	5	1	2	1	-	-
White, both sexes	-	-	-	-	2	2	4	1	-	2	-	-
Male	-	-	-	-	1	1	1	-	-	1	-	-
Female	-	-	-	-	1	1	3	1	-	1	-	-
Black, both sexes	1	-	2	1	-	2	3	1	2	1	-	-
Male	-	-	2	1	-	1	1	1	-	1	-	-
Female	1	-	-	-	-	1	2	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Alaska												
All races, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	5	4	2	6	4	1	4	3	-	-	-
Male	2	5	4	1	3	4	1	2	-	-	-	-
Female	1	-	-	1	3	-	-	2	3	-	-	-
White, both sexes	2	3	4	2	4	4	1	4	3	-	-	-
Male	1	3	4	1	2	4	1	2	-	-	-	-
Female	1	-	-	1	2	-	-	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	2	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Arkansas												
All races, both sexes	-	-	2	1	1	1	3	2	2	3	-	-
Male	-	-	2	-	1	1	1	-	-	-	-	-
Female	-	-	-	1	-	-	2	2	2	3	-	-
White, both sexes	-	-	2	-	1	1	2	2	2	1	-	-
Male	-	-	2	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	2	1	-	-
Black, both sexes	-	-	-	1	-	-	1	-	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7	8	13	16	13	19	26	26	10	9	-	-
Male	6	6	8	11	9	7	9	6	4	-	-	-
Female	1	2	5	5	4	12	17	20	6	9	-	-
White, both sexes	7	7	8	13	11	15	21	24	6	8	-	-
Male	6	6	4	8	8	7	8	5	2	-	-	-
Female	1	1	4	5	3	8	13	19	4	8	-	-
Black, both sexes	-	-	2	3	1	3	2	1	-	-	-	-
Male	-	-	2	3	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	3	2	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	1	3	1	4	1	-	-
Male	-	-	1	-	-	-	1	-	2	-	-	-
Female	-	-	1	-	-	1	2	1	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Colorado												
All races, both sexes	-	3	2	3	2	1	-	5	1	1	-	-
Male	-	2	2	2	-	-	-	1	-	-	-	-
Female	-	1	-	1	2	1	-	4	1	1	-	-
White, both sexes	-	3	2	3	2	1	-	5	1	1	-	-
Male	-	2	2	2	-	-	-	1	-	-	-	-
Female	-	1	-	1	2	1	-	4	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	2	4	3	4	5	3	5	1	-
Male	-	-	-	2	1	1	1	3	1	2	1	-
Female	-	-	-	-	3	2	3	2	2	3	-	-
White, both sexes	-	-	-	2	4	2	4	5	3	5	1	-
Male	-	-	-	2	1	1	1	3	1	2	1	-
Female	-	-	-	-	3	1	3	2	2	3	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Delaware												
All races, both sexes	-	1	-	1	-	2	1	1	-	-	-	-
Male	-	1	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	2	1	1	-	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Florida												
All races, both sexes	3	3	6	6	8	12	16	18	12	6	1	-
Male	3	3	5	4	4	6	6	9	4	1	-	-
Female	-	-	1	2	4	6	10	9	8	5	1	-
White, both sexes	2	3	6	5	7	10	16	18	11	6	1	-
Male	2	3	5	4	4	4	6	9	3	1	-	-
Female	-	-	1	1	3	6	10	9	8	5	1	-
Black, both sexes	1	-	-	1	1	2	-	-	1	-	-	-
Male	1	-	-	-	-	2	-	-	1	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2	2	8	5	4	3	10	10	7	1	2	-
Male	1	1	4	3	2	1	5	5	2	1	-	-
Female	1	1	4	2	2	2	5	5	5	-	2	-
White, both sexes	-	2	4	4	3	1	5	10	6	1	2	-
Male	-	1	3	3	2	1	3	5	2	1	-	-
Female	-	1	1	1	1	-	2	5	4	-	2	-
Black, both sexes	2	-	4	1	1	2	5	-	1	-	-	-
Male	1	-	1	-	-	-	2	-	-	-	-	-
Female	1	-	3	1	1	2	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Hawaii												
All races, both sexes	-	-	-	1	-	2	1	1	2	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	2	-	-	-
White, both sexes	-	-	-	1	-	1	1	-	2	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Idaho												
All races, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Illinois												
All races, both sexes	6	11	9	10	13	22	14	17	16	3	2	-
Male	4	7	6	6	6	14	7	7	3	-	-	-
Female	2	4	3	4	7	8	7	10	13	3	2	-
White, both sexes	4	8	7	7	13	20	9	16	16	3	1	-
Male	3	5	5	5	6	12	6	7	3	-	-	-
Female	1	3	2	2	7	8	3	9	13	3	1	-
Black, both sexes	2	3	2	3	-	1	5	-	-	-	1	-
Male	1	2	1	1	-	1	1	-	-	-	-	-
Female	1	1	1	2	-	-	4	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Indiana												
All races, both sexes	-	4	1	1	-	2	1	2	4	2	1	-
Male	-	3	1	1	-	2	1	1	2	1	-	-
Female	-	1	-	-	-	-	-	1	2	1	1	-
White, both sexes	-	3	1	1	-	2	1	2	4	2	1	-
Male	-	3	1	1	-	2	1	1	2	1	-	-
Female	-	-	-	-	-	-	-	1	2	1	1	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Iowa												
All races, both sexes	1	-	-	1	-	1	3	3	3	1	-	-
Male	1	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	-	-	2	3	2	1	-	-
White, both sexes	1	-	-	1	-	1	3	3	3	1	-	-
Male	1	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	-	-	2	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	1	-	2	2	1	2	2	1	3	-	-
Male	3	1	-	2	2	-	2	1	-	1	-	-
Female	-	-	-	-	-	1	-	1	1	2	-	-
White, both sexes	2	1	-	2	2	1	2	2	1	3	-	-
Male	2	1	-	2	2	-	2	1	-	1	-	-
Female	-	-	-	-	-	1	-	1	1	2	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Kentucky												
All races, both sexes	2	1	1	2	-	2	3	1	-	-	-	-
Male	-	1	1	-	-	-	2	-	-	-	-	-
Female	2	-	-	2	-	2	1	1	-	-	-	-
White, both sexes	1	1	1	2	-	1	3	1	-	-	-	-
Male	-	1	1	-	-	-	2	-	-	-	-	-
Female	1	-	-	2	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	6	4	7	4	4	6	6	7	6	4	-	-
Male	5	3	6	2	1	3	1	4	5	-	-	-
Female	1	1	1	2	3	3	5	3	1	4	-	-
White, both sexes	2	3	6	2	2	4	5	6	6	2	-	-
Male	2	2	5	-	1	2	1	3	5	-	-	-
Female	-	1	1	2	1	2	4	3	1	2	-	-
Black, both sexes	4	1	1	2	2	2	1	1	-	2	-	-
Male	3	1	1	2	-	1	-	1	-	-	-	-
Female	1	-	-	-	2	1	1	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Maine												
All races, both sexes	1	3	2	1	1	2	3	1	1	-	-	-
Male	-	1	2	-	1	2	2	-	-	-	-	-
Female	1	2	-	1	-	-	1	1	1	-	-	-
White, both sexes	1	3	1	1	1	2	3	1	1	-	-	-
Male	-	1	1	-	1	2	2	-	-	-	-	-
Female	1	2	-	1	-	-	1	1	1	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	1	-	3	4	5	3	5	3	1	-	-
Male	1	1	-	1	4	3	2	-	1	1	-	-
Female	-	-	-	2	-	2	1	5	2	-	-	-
White, both sexes	1	1	-	2	1	3	3	4	2	-	-	-
Male	1	1	-	-	1	2	2	-	1	-	-	-
Female	-	-	-	2	-	1	1	4	1	-	-	-
Black, both sexes	-	-	-	1	2	2	-	1	1	1	-	-
Male	-	-	-	1	2	1	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Massachusetts												
All races, both sexes	5	2	3	2	7	8	9	9	6	3	-	-
Male	4	2	2	1	4	4	2	4	3	-	-	-
Female	1	-	1	1	3	4	7	5	3	3	-	-
White, both sexes	4	2	3	2	6	8	9	9	6	3	-	-
Male	3	2	2	1	3	4	2	4	3	-	-	-
Female	1	-	1	1	3	4	7	5	3	3	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	3	3	2	6	5	13	8	3	2	-	-
Male	1	2	-	1	4	4	5	2	2	2	-	-
Female	-	1	3	1	2	1	8	6	1	-	-	-
White, both sexes	1	3	2	-	2	5	11	8	3	1	-	-
Male	1	2	-	-	2	4	5	2	2	1	-	-
Female	-	1	2	-	-	1	6	6	1	-	-	-
Black, both sexes	-	-	1	2	4	-	2	-	-	1	-	-
Male	-	-	-	1	2	-	-	-	-	1	-	-
Female	-	-	1	1	2	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Minnesota												
All races, both sexes	-	2	2	1	4	4	4	5	3	-	-	-
Male	-	1	2	1	2	2	1	2	1	-	-	-
Female	-	1	-	-	2	2	3	3	2	-	-	-
White, both sexes	-	2	2	1	4	4	4	5	3	-	-	-
Male	-	1	2	1	2	2	1	2	1	-	-	-
Female	-	1	-	-	2	2	3	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	2	1	-	-	2	-	2	-	2	-	-
Male	1	1	1	-	-	2	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	2	-	-
White, both sexes	1	1	1	-	-	1	-	-	-	2	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	2	-	-
Black, both sexes	-	1	-	-	-	1	-	2	-	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Missouri												
All races, both sexes	-	1	3	2	4	5	16	15	12	3	2	-
Male	-	1	2	2	3	4	3	4	2	-	-	-
Female	-	-	1	-	1	1	13	11	10	3	2	-
White, both sexes	-	1	2	2	3	4	14	15	12	3	2	-
Male	-	1	2	2	2	3	2	4	2	-	-	-
Female	-	-	-	-	1	1	12	11	10	3	2	-
Black, both sexes	-	-	1	-	1	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	1	1	-	1	-	1	1	-	1	-	-
Male	-	-	1	-	1	-	-	-	-	1	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	1	-	-	1	-	1	1	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Nebraska												
All races, both sexes	-	-	-	1	1	2	4	1	3	-	1	-
Male	-	-	-	1	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	2	3	1	2	-	1	-
White, both sexes	-	-	-	1	1	2	4	1	3	-	1	-
Male	-	-	-	1	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	2	3	1	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	3	-	1	2	1	-	1	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	2	-	1	1	1	-	1	-	-	-
White, both sexes	-	1	3	-	1	2	1	-	1	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	2	-	1	1	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
New Hampshire												
All races, both sexes	1	1	-	-	-	1	6	3	-	1	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	1	1	-	-	-	-	5	2	-	1	-	-
White, both sexes	1	1	-	-	-	1	6	3	-	1	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	1	1	-	-	-	-	5	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	2	5	2	14	10	19	20	28	11	2	-
Male	1	2	4	1	10	5	8	6	6	3	-	-
Female	1	-	1	1	4	5	11	14	22	8	2	-
White, both sexes	2	2	4	2	11	8	18	18	28	11	2	-
Male	1	2	3	1	7	4	8	4	6	3	-	-
Female	1	-	1	1	4	4	10	14	22	8	2	-
Black, both sexes	-	-	1	-	2	1	1	1	-	-	-	-
Male	-	-	1	-	2	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
New Mexico												
All races, both sexes	-	-	1	-	1	4	3	2	1	-	-	-
Male	-	-	-	-	-	3	-	1	1	-	-	-
Female	-	-	1	-	1	1	3	1	-	-	-	-
White, both sexes	-	-	1	-	1	3	3	2	1	-	-	-
Male	-	-	-	-	-	2	-	1	1	-	-	-
Female	-	-	1	-	1	1	3	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4	6	8	13	14	20	36	35	23	10	5	-
Male	4	5	6	9	10	14	17	15	6	-	1	-
Female	-	1	2	4	4	6	19	20	17	10	4	-
White, both sexes	2	4	7	11	13	19	32	34	21	10	5	-
Male	2	3	5	8	9	14	16	15	5	-	1	-
Female	-	1	2	3	4	5	16	19	16	10	4	-
Black, both sexes	1	2	1	2	-	1	4	1	1	-	-	-
Male	1	2	1	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	3	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	1	-	-	-
Male	1	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
North Carolina												
All races, both sexes	8	8	6	6	8	10	17	6	8	-	-	-
Male	5	7	6	5	4	7	5	3	3	-	-	-
Female	3	1	-	1	4	3	12	3	5	-	-	-
White, both sexes	6	4	2	5	7	6	15	5	7	-	-	-
Male	5	3	2	4	4	4	5	3	2	-	-	-
Female	1	1	-	1	3	2	10	2	5	-	-	-
Black, both sexes	2	4	4	1	1	4	2	1	1	-	-	-
Male	-	4	4	1	-	3	-	-	1	-	-	-
Female	2	-	-	-	1	1	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	2	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	-	-	-	-	-	-	2	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Ohio												
All races, both sexes	9	5	8	8	11	12	12	26	15	3	-	-
Male	6	3	7	5	5	5	3	8	4	1	-	-
Female	3	2	1	3	6	7	9	18	11	2	-	-
White, both sexes	8	5	7	7	7	12	10	24	15	3	-	-
Male	6	3	6	4	3	5	2	8	4	1	-	-
Female	2	2	1	3	4	7	8	16	11	2	-	-
Black, both sexes	1	-	1	1	4	-	2	2	-	-	-	-
Male	-	-	1	1	2	-	1	-	-	-	-	-
Female	1	-	-	-	2	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	56	45	58	57	78	77	109	111	81	39	7	-
Male	40	33	42	38	39	32	53	47	22	7	2	-
Female	16	12	16	19	39	45	56	64	59	32	5	-
White, both sexes	45	40	50	49	70	72	100	100	72	36	4	-
Male	36	31	37	31	35	30	46	41	18	7	1	-
Female	9	9	13	18	35	42	54	59	54	29	3	-
Black, both sexes	7	3	4	5	3	4	5	6	8	3	1	-
Male	3	1	3	4	1	2	5	3	4	-	-	-
Female	4	2	1	1	2	2	-	3	4	3	1	-
American Indian 1/, both sexes	3	2	2	3	4	1	3	4	1	-	2	-
Male	1	1	1	3	3	-	2	3	-	-	1	-
Female	2	1	1	-	1	1	1	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	2	-	1	-	1	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	1	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Oregon												
All races, both sexes	4	-	2	5	4	1	7	4	2	-	-	-
Male	2	-	1	3	4	1	4	1	-	-	-	-
Female	2	-	1	2	-	-	3	3	2	-	-	-
White, both sexes	4	-	1	5	4	1	7	4	2	-	-	-
Male	2	-	-	3	4	1	4	1	-	-	-	-
Female	2	-	1	2	-	-	3	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	6	8	17	10	13	20	24	19	16	5	3	-
Male	4	5	11	5	7	13	10	5	7	-	1	-
Female	2	3	6	5	6	7	14	14	9	5	2	-
White, both sexes	4	8	16	10	13	19	24	18	14	5	3	-
Male	3	5	11	5	7	13	10	4	7	-	1	-
Female	1	3	5	5	6	6	14	14	7	5	2	-
Black, both sexes	2	-	1	-	-	1	-	1	2	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	1	-	-	1	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Rhode Island												
All races, both sexes	-	-	1	-	3	-	1	-	1	-	-	-
Male	-	-	1	-	3	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	3	-	1	-	1	-	-	-
Male	-	-	-	-	3	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	15	31	35	16	31	17	25	14	12	2	-	-
Male	9	22	25	12	20	9	15	3	4	-	-	-
Female	6	9	10	4	11	8	10	11	8	2	-	-
White, both sexes	4	14	19	9	15	9	15	9	8	1	-	-
Male	4	11	14	7	12	4	8	3	4	-	-	-
Female	-	3	5	2	3	5	7	6	4	1	-	-
Black, both sexes	11	17	16	7	16	8	10	5	4	1	-	-
Male	5	11	11	5	8	5	7	-	-	-	-	-
Female	6	6	5	2	8	3	3	5	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
South Dakota												
All races, both sexes	-	-	-	-	-	2	3	-	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	2	3	-	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	5	1	3	4	4	13	4	3	2	-	-
Male	1	5	-	2	1	3	4	1	2	1	-	-
Female	1	-	1	1	3	1	9	3	1	1	-	-
White, both sexes	2	4	1	3	3	4	11	3	3	2	-	-
Male	1	4	-	2	1	3	3	1	2	1	-	-
Female	1	-	1	1	2	1	8	2	1	1	-	-
Black, both sexes	-	1	-	-	1	-	2	1	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Texas												
All races, both sexes	10	4	10	12	15	16	17	30	9	5	-	-
Male	5	1	9	9	5	11	11	9	2	2	-	-
Female	5	3	1	3	10	5	6	21	7	3	-	-
White, both sexes	6	4	8	11	12	12	16	27	8	3	-	-
Male	3	1	7	9	5	9	10	7	1	1	-	-
Female	3	3	1	2	7	3	6	20	7	2	-	-
Black, both sexes	4	-	2	1	2	4	1	2	1	1	-	-
Male	2	-	2	-	-	2	1	1	1	-	-	-
Female	2	-	-	1	2	2	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	2	-	1	3	1	5	2	1	-	-
Male	-	-	2	-	1	3	-	2	-	-	-	-
Female	-	-	-	-	-	-	1	3	2	1	-	-
White, both sexes	-	-	2	-	1	3	1	5	2	1	-	-
Male	-	-	2	-	1	3	-	2	-	-	-	-
Female	-	-	-	-	-	-	1	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Vermont												
All races, both sexes	-	1	-	-	-	1	-	2	1	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
White, both sexes	-	1	-	-	-	1	-	2	1	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	57	55	47	41	60	66	68	51	44	20	6	-
Male	46	41	29	28	38	38	31	15	13	3	1	-
Female	11	14	18	13	22	28	37	36	31	17	5	-
White, both sexes	32	31	34	32	46	50	55	34	32	12	4	-
Male	27	24	22	22	28	26	27	10	9	2	1	-
Female	5	7	12	10	18	24	28	24	23	10	3	-
Black, both sexes	24	24	12	9	14	16	13	17	12	8	2	-
Male	18	17	6	6	10	12	4	5	4	1	-	-
Female	6	7	6	3	4	4	9	12	8	7	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Washington												
All races, both sexes	1	2	1	1	4	4	4	9	2	2	1	-
Male	-	2	1	-	3	2	3	4	1	-	-	-
Female	1	-	-	1	1	2	1	5	1	2	1	-
White, both sexes	1	-	1	1	4	3	4	9	2	2	1	-
Male	-	-	1	-	3	2	3	4	1	-	-	-
Female	1	-	-	1	1	1	1	5	1	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	2	-	1	4	3	4	3	4	1	1	-
Male	2	2	-	-	1	1	2	1	2	1	1	-
Female	-	-	-	1	3	2	2	2	2	-	-	-
White, both sexes	2	2	-	1	4	3	4	3	4	1	1	-
Male	2	2	-	-	1	1	2	1	2	1	1	-
Female	-	-	-	1	3	2	2	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other acute ischemic heart diseases (I24)												
Wisconsin												
All races, both sexes	2	2	1	2	3	9	10	6	7	1	-	-
Male	2	-	1	1	1	4	3	3	2	-	-	-
Female	-	2	-	1	2	5	7	3	5	1	-	-
White, both sexes	2	2	1	2	3	8	10	5	7	1	-	-
Male	2	-	1	1	1	4	3	3	2	-	-	-
Female	-	2	-	1	2	4	7	2	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	2	-	-	1	-	-	-	1	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
White, both sexes	-	2	-	-	1	-	-	-	1	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
United States												
All races, both sexes	291,118	2	4	2	4	14	66	153	458	1,150	2,894	5,634
Male	150,213	1	2	2	3	12	53	127	343	885	2,219	4,331
Female	140,905	1	2	-	1	2	13	26	115	265	675	1,303
White, both sexes	255,133	1	4	2	4	10	43	113	340	838	2,267	4,365
Male	131,653	1	2	2	3	9	36	93	253	652	1,793	3,420
Female	123,480	-	2	-	1	1	7	20	87	186	474	945
Black, both sexes	29,933	1	-	-	-	4	16	32	89	273	526	1,096
Male	15,110	-	-	-	-	3	11	28	63	201	340	769
Female	14,823	1	-	-	-	1	5	4	26	72	186	327
American Indian 1/, both sexes	1,155	-	-	-	-	-	1	-	10	15	23	58
Male	663	-	-	-	-	-	-	-	9	11	18	45
Female	492	-	-	-	-	-	1	-	1	4	5	13
Asian or Pacific Islander 2/, both sexes	4,897	-	-	-	-	-	6	8	19	24	78	115
Male	2,787	-	-	-	-	-	6	6	18	21	68	97
Female	2,110	-	-	-	-	-	-	2	1	3	10	18
Alabama												
All races, both sexes	3,202	-	-	-	1	-	1	1	5	13	26	64
Male	1,611	-	-	-	1	-	1	1	3	11	17	53
Female	1,591	-	-	-	-	-	-	-	2	2	9	11
White, both sexes	2,602	-	-	-	1	-	1	-	3	7	14	38
Male	1,336	-	-	-	1	-	1	-	3	7	10	30
Female	1,266	-	-	-	-	-	-	-	-	-	4	8
Black, both sexes	588	-	-	-	-	-	-	1	2	6	12	26
Male	272	-	-	-	-	-	-	1	-	4	7	23
Female	316	-	-	-	-	-	-	-	2	2	5	3
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Alaska												
All races, both sexes	293	-	-	-	-	-	1	-	1	1	11	17
Male	201	-	-	-	-	-	1	-	1	1	10	12
Female	92	-	-	-	-	-	-	-	-	-	1	5
White, both sexes	239	-	-	-	-	-	-	-	1	-	9	15
Male	166	-	-	-	-	-	-	-	1	-	8	11
Female	73	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	41	-	-	-	-	-	-	-	-	1	1	2
Male	28	-	-	-	-	-	-	-	-	1	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	1	-	-	-	1	-
Male	4	-	-	-	-	-	1	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	6,091	-	-	-	-	-	1	3	8	27	60	124
Male	3,429	-	-	-	-	-	-	3	8	19	44	104
Female	2,662	-	-	-	-	-	1	-	-	8	16	20
White, both sexes	5,800	-	-	-	-	-	-	3	6	24	52	109
Male	3,267	-	-	-	-	-	-	3	6	18	40	92
Female	2,533	-	-	-	-	-	-	-	-	6	12	17
Black, both sexes	138	-	-	-	-	-	-	-	-	1	7	3
Male	72	-	-	-	-	-	-	-	-	1	3	2
Female	66	-	-	-	-	-	-	-	-	-	4	1
American Indian 1/, both sexes	121	-	-	-	-	-	1	-	2	2	1	11
Male	70	-	-	-	-	-	-	-	2	-	1	9
Female	51	-	-	-	-	-	1	-	-	2	-	2
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Arkansas												
All races, both sexes	2,176	-	-	-	-	1	-	1	7	5	21	32
Male	1,065	-	-	-	-	1	-	1	3	2	14	23
Female	1,111	-	-	-	-	-	-	-	4	3	7	9
White, both sexes	1,898	-	-	-	-	1	-	1	5	3	14	23
Male	944	-	-	-	-	1	-	1	2	2	9	16
Female	954	-	-	-	-	-	-	-	3	1	5	7
Black, both sexes	273	-	-	-	-	-	-	-	2	2	6	9
Male	118	-	-	-	-	-	-	-	1	-	4	7
Female	155	-	-	-	-	-	-	-	1	2	2	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	33,736	-	1	1	1	1	3	17	42	117	295	566
Male	17,652	-	1	1	1	1	3	12	30	102	237	441
Female	16,084	-	-	-	-	-	-	5	12	15	58	125
White, both sexes	28,789	-	1	1	1	-	1	10	34	90	222	432
Male	14,948	-	1	1	1	-	1	8	22	79	188	337
Female	13,841	-	-	-	-	-	-	2	12	11	34	95
Black, both sexes	2,624	-	-	-	-	1	1	4	3	18	43	88
Male	1,397	-	-	-	-	1	1	2	3	14	24	68
Female	1,227	-	-	-	-	-	-	2	-	4	19	20
American Indian 1/, both sexes	135	-	-	-	-	-	-	-	-	2	1	4
Male	73	-	-	-	-	-	-	-	-	2	-	4
Female	62	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2,188	-	-	-	-	-	1	3	5	7	29	42
Male	1,234	-	-	-	-	-	1	2	5	7	25	32
Female	954	-	-	-	-	-	-	1	-	-	4	10

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Colorado												
All races, both sexes	2,734	-	-	-	-	-	1	2	2	17	26	61
Male	1,560	-	-	-	-	-	1	2	1	15	23	48
Female	1,174	-	-	-	-	-	-	-	1	2	3	13
White, both sexes	2,611	-	-	-	-	-	1	2	2	17	25	55
Male	1,500	-	-	-	-	-	1	2	1	15	22	45
Female	1,111	-	-	-	-	-	-	-	1	2	3	10
Black, both sexes	93	-	-	-	-	-	-	-	-	-	-	4
Male	41	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3,601	-	-	-	-	-	1	3	5	13	26	63
Male	1,764	-	-	-	-	-	1	3	4	12	21	46
Female	1,837	-	-	-	-	-	-	-	1	1	5	17
White, both sexes	3,347	-	-	-	-	-	1	3	4	9	20	53
Male	1,631	-	-	-	-	-	1	3	3	8	18	39
Female	1,716	-	-	-	-	-	-	-	1	1	2	14
Black, both sexes	222	-	-	-	-	-	-	-	-	4	6	9
Male	116	-	-	-	-	-	-	-	-	4	3	6
Female	106	-	-	-	-	-	-	-	-	-	3	3
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Delaware												
All races, both sexes	1,027	-	-	-	-	-	-	1	2	7	12	22
Male	530	-	-	-	-	-	-	1	1	4	9	15
Female	497	-	-	-	-	-	-	-	1	3	3	7
White, both sexes	853	-	-	-	-	-	-	-	2	4	9	17
Male	458	-	-	-	-	-	-	-	1	2	8	12
Female	395	-	-	-	-	-	-	-	1	2	1	5
Black, both sexes	166	-	-	-	-	-	-	1	-	3	3	5
Male	67	-	-	-	-	-	-	1	-	2	1	3
Female	99	-	-	-	-	-	-	-	-	1	2	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	864	-	-	-	-	-	-	1	-	3	10	17
Male	439	-	-	-	-	-	-	1	-	3	8	12
Female	425	-	-	-	-	-	-	-	-	-	2	5
White, both sexes	181	-	-	-	-	-	-	-	-	-	-	-
Male	90	-	-	-	-	-	-	-	-	-	-	-
Female	91	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	668	-	-	-	-	-	-	-	-	3	10	17
Male	337	-	-	-	-	-	-	-	-	3	8	12
Female	331	-	-	-	-	-	-	-	-	-	2	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	1	-	-	-	-
Male	12	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Florida												
All races, both sexes	24,979	-	-	-	-	-	5	11	23	78	235	407
Male	13,398	-	-	-	-	-	4	9	18	57	179	317
Female	11,581	-	-	-	-	-	1	2	5	21	56	90
White, both sexes	22,869	-	-	-	-	-	2	9	15	53	191	344
Male	12,303	-	-	-	-	-	2	7	13	41	152	274
Female	10,566	-	-	-	-	-	-	2	2	12	39	70
Black, both sexes	1,991	-	-	-	-	-	3	2	7	24	42	59
Male	1,030	-	-	-	-	-	2	2	4	15	25	40
Female	961	-	-	-	-	-	1	-	3	9	17	19
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	-	1	-	1
Male	16	-	-	-	-	-	-	-	-	1	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	97	-	-	-	-	-	-	-	1	-	2	3
Male	49	-	-	-	-	-	-	-	1	-	2	2
Female	48	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	5,329	-	-	-	-	-	1	10	18	42	90	179
Male	2,935	-	-	-	-	-	-	7	14	36	70	140
Female	2,394	-	-	-	-	-	1	3	4	6	20	39
White, both sexes	4,099	-	-	-	-	-	1	4	11	27	60	123
Male	2,309	-	-	-	-	-	-	3	8	21	45	99
Female	1,790	-	-	-	-	-	1	1	3	6	15	24
Black, both sexes	1,198	-	-	-	-	-	-	6	5	15	27	56
Male	609	-	-	-	-	-	-	4	4	15	22	41
Female	589	-	-	-	-	-	-	2	1	-	5	15
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	2	-	3	-
Male	14	-	-	-	-	-	-	-	2	-	3	-
Female	13	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Hawaii												
All races, both sexes	842	-	-	-	-	-	-	-	1	6	13	22
Male	530	-	-	-	-	-	-	-	1	5	8	19
Female	312	-	-	-	-	-	-	-	-	1	5	3
White, both sexes	271	-	-	-	-	-	-	-	-	3	4	8
Male	178	-	-	-	-	-	-	-	-	3	2	8
Female	93	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	563	-	-	-	-	-	-	-	1	3	9	12
Male	345	-	-	-	-	-	-	-	1	2	6	9
Female	218	-	-	-	-	-	-	-	-	1	3	3
Idaho												
All races, both sexes	810	-	-	-	-	-	-	-	1	5	9	6
Male	453	-	-	-	-	-	-	-	1	3	6	4
Female	357	-	-	-	-	-	-	-	-	2	3	2
White, both sexes	797	-	-	-	-	-	-	-	1	5	9	6
Male	444	-	-	-	-	-	-	-	1	3	6	4
Female	353	-	-	-	-	-	-	-	-	2	3	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Illinois												
All races, both sexes	11,390	1	-	-	-	1	3	11	13	45	124	284
Male	5,815	1	-	-	-	1	1	10	11	38	98	216
Female	5,575	-	-	-	-	-	2	1	2	7	26	68
White, both sexes	9,592	1	-	-	-	1	2	9	11	29	92	198
Male	4,870	1	-	-	-	1	-	8	9	25	76	158
Female	4,722	-	-	-	-	-	2	1	2	4	16	40
Black, both sexes	1,665	-	-	-	-	-	1	2	2	16	29	85
Male	876	-	-	-	-	-	1	2	2	13	19	57
Female	789	-	-	-	-	-	-	-	-	3	10	28
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	126	-	-	-	-	-	-	-	-	-	3	1
Male	66	-	-	-	-	-	-	-	-	-	3	1
Female	60	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	5,733	-	-	-	-	1	4	4	23	26	61	126
Male	2,985	-	-	-	-	1	3	3	18	17	40	86
Female	2,748	-	-	-	-	-	1	1	5	9	21	40
White, both sexes	5,297	-	-	-	-	1	3	4	19	22	51	103
Male	2,765	-	-	-	-	1	3	3	14	14	35	73
Female	2,532	-	-	-	-	-	-	1	5	8	16	30
Black, both sexes	413	-	-	-	-	-	1	-	4	4	10	23
Male	209	-	-	-	-	-	-	-	4	3	5	13
Female	204	-	-	-	-	-	1	-	-	1	5	10
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Iowa												
All races, both sexes	3,739	-	-	-	-	1	-	1	2	14	29	64
Male	1,894	-	-	-	-	1	-	1	2	9	17	46
Female	1,845	-	-	-	-	-	-	-	-	5	12	18
White, both sexes	3,685	-	-	-	-	1	-	1	2	12	25	60
Male	1,862	-	-	-	-	1	-	1	2	8	17	43
Female	1,823	-	-	-	-	-	-	-	-	4	8	17
Black, both sexes	44	-	-	-	-	-	-	-	-	2	4	4
Male	26	-	-	-	-	-	-	-	-	1	-	3
Female	18	-	-	-	-	-	-	-	-	1	4	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2,515	-	-	-	-	-	1	1	5	8	29	59
Male	1,377	-	-	-	-	-	1	-	5	6	18	47
Female	1,138	-	-	-	-	-	-	1	-	2	11	12
White, both sexes	2,373	-	-	-	-	-	1	1	5	7	27	51
Male	1,291	-	-	-	-	-	1	-	5	5	16	41
Female	1,082	-	-	-	-	-	-	1	-	2	11	10
Black, both sexes	106	-	-	-	-	-	-	-	-	1	1	5
Male	65	-	-	-	-	-	-	-	-	1	1	4
Female	41	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	-	-	1	3
Male	17	-	-	-	-	-	-	-	-	-	1	2
Female	10	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Kentucky												
All races, both sexes	3,431	-	1	-	-	-	2	7	11	18	46	82
Male	1,697	-	-	-	-	-	2	7	9	12	33	64
Female	1,734	-	1	-	-	-	-	-	2	6	13	18
White, both sexes	3,269	-	1	-	-	-	2	7	10	15	43	75
Male	1,623	-	-	-	-	-	2	7	8	11	31	60
Female	1,646	-	1	-	-	-	-	-	2	4	12	15
Black, both sexes	155	-	-	-	-	-	-	-	-	3	3	6
Male	68	-	-	-	-	-	-	-	-	1	2	3
Female	87	-	-	-	-	-	-	-	-	2	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3,617	-	-	-	1	-	-	1	11	26	50	85
Male	1,939	-	-	-	-	-	-	1	6	21	35	61
Female	1,678	-	-	-	1	-	-	-	5	5	15	24
White, both sexes	2,740	-	-	-	1	-	-	1	8	15	29	51
Male	1,483	-	-	-	-	-	-	1	4	12	23	37
Female	1,257	-	-	-	1	-	-	-	4	3	6	14
Black, both sexes	857	-	-	-	-	-	-	-	3	11	21	33
Male	443	-	-	-	-	-	-	-	2	9	12	23
Female	414	-	-	-	-	-	-	-	1	2	9	10
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Maine												
All races, both sexes	1,248	-	-	-	-	1	-	1	3	5	15	27
Male	672	-	-	-	-	1	-	1	2	4	12	24
Female	576	-	-	-	-	-	-	-	1	1	3	3
White, both sexes	1,243	-	-	-	-	1	-	1	3	5	15	26
Male	668	-	-	-	-	1	-	1	2	4	12	23
Female	575	-	-	-	-	-	-	-	1	1	3	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5,433	-	-	-	-	-	1	3	11	17	50	100
Male	2,732	-	-	-	-	-	1	1	8	12	36	79
Female	2,701	-	-	-	-	-	-	2	3	5	14	21
White, both sexes	4,070	-	-	-	-	-	1	3	8	9	34	61
Male	2,059	-	-	-	-	-	1	1	5	6	28	51
Female	2,011	-	-	-	-	-	-	2	3	3	6	10
Black, both sexes	1,281	-	-	-	-	-	-	-	3	7	15	39
Male	633	-	-	-	-	-	-	-	3	6	7	28
Female	648	-	-	-	-	-	-	-	-	1	8	11
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	74	-	-	-	-	-	-	-	-	-	1	-
Male	37	-	-	-	-	-	-	-	-	-	1	-
Female	37	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Massachusetts												
All races, both sexes	5,109	-	-	-	-	-	3	1	11	19	49	87
Male	2,599	-	-	-	-	-	3	1	4	15	39	68
Female	2,510	-	-	-	-	-	-	-	7	4	10	19
White, both sexes	4,895	-	-	-	-	-	2	1	9	16	44	78
Male	2,492	-	-	-	-	-	2	1	4	12	36	64
Female	2,403	-	-	-	-	-	-	-	5	4	8	14
Black, both sexes	175	-	-	-	-	-	-	-	2	2	5	9
Male	87	-	-	-	-	-	-	-	-	2	3	4
Female	88	-	-	-	-	-	-	-	2	-	2	5
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	34	-	-	-	-	-	1	-	-	1	-	-
Male	19	-	-	-	-	-	1	-	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	12,367	-	-	-	-	-	4	3	24	55	118	245
Male	6,400	-	-	-	-	-	4	2	19	38	87	178
Female	5,967	-	-	-	-	-	-	1	5	17	31	67
White, both sexes	10,240	-	-	-	-	-	3	3	18	33	87	171
Male	5,253	-	-	-	-	-	3	2	16	24	67	132
Female	4,987	-	-	-	-	-	-	1	2	9	20	39
Black, both sexes	2,028	-	-	-	-	-	1	-	6	19	27	71
Male	1,079	-	-	-	-	-	1	-	3	12	17	45
Female	949	-	-	-	-	-	-	-	3	7	10	26
American Indian 1/, both sexes	42	-	-	-	-	-	-	-	-	2	3	2
Male	23	-	-	-	-	-	-	-	-	1	2	-
Female	19	-	-	-	-	-	-	-	-	1	1	2
Asian or Pacific Islander 2/, both sexes	57	-	-	-	-	-	-	-	-	1	1	1
Male	45	-	-	-	-	-	-	-	-	1	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Minnesota												
All races, both sexes	3,259	-	-	-	-	-	4	3	7	17	46	79
Male	1,895	-	-	-	-	-	2	2	5	14	41	60
Female	1,364	-	-	-	-	-	2	1	2	3	5	19
White, both sexes	3,170	-	-	-	-	-	4	3	5	14	40	67
Male	1,836	-	-	-	-	-	2	2	4	12	36	50
Female	1,334	-	-	-	-	-	2	1	1	2	4	17
Black, both sexes	52	-	-	-	-	-	-	-	2	2	3	11
Male	35	-	-	-	-	-	-	-	1	1	2	9
Female	17	-	-	-	-	-	-	-	1	1	1	2
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	2	1
Male	11	-	-	-	-	-	-	-	-	-	2	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	1	1	-
Male	13	-	-	-	-	-	-	-	-	1	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2,657	-	-	-	-	1	2	3	14	29	57	82
Male	1,317	-	-	-	-	1	2	3	9	22	39	57
Female	1,340	-	-	-	-	-	-	-	5	7	18	25
White, both sexes	1,993	-	-	-	-	-	1	2	4	14	30	46
Male	974	-	-	-	-	-	1	2	2	11	23	30
Female	1,019	-	-	-	-	-	-	-	2	3	7	16
Black, both sexes	653	-	-	-	-	1	1	1	10	15	26	36
Male	340	-	-	-	-	1	1	1	7	11	16	27
Female	313	-	-	-	-	-	-	-	3	4	10	9
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Missouri												
All races, both sexes	5,803	-	1	-	-	-	1	2	11	32	67	125
Male	2,893	-	-	-	-	-	1	1	8	26	54	100
Female	2,910	-	1	-	-	-	-	1	3	6	13	25
White, both sexes	5,257	-	1	-	-	-	1	2	9	25	55	103
Male	2,610	-	-	-	-	-	1	1	6	20	45	81
Female	2,647	-	1	-	-	-	-	1	3	5	10	22
Black, both sexes	527	-	-	-	-	-	-	-	2	7	12	21
Male	277	-	-	-	-	-	-	-	2	6	9	18
Female	250	-	-	-	-	-	-	-	-	1	3	3
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	683	-	-	-	-	-	-	-	2	2	6	14
Male	419	-	-	-	-	-	-	-	1	2	5	11
Female	264	-	-	-	-	-	-	-	1	-	1	3
White, both sexes	658	-	-	-	-	-	-	-	2	1	6	13
Male	402	-	-	-	-	-	-	-	1	1	5	10
Female	256	-	-	-	-	-	-	-	1	-	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Nebraska												
All races, both sexes	1,435	-	-	-	-	-	-	1	3	7	21	25
Male	771	-	-	-	-	-	-	1	2	5	18	20
Female	664	-	-	-	-	-	-	-	1	2	3	5
White, both sexes	1,381	-	-	-	-	-	-	1	2	5	19	22
Male	741	-	-	-	-	-	-	1	1	3	16	19
Female	640	-	-	-	-	-	-	-	1	2	3	3
Black, both sexes	41	-	-	-	-	-	-	-	1	2	2	3
Male	24	-	-	-	-	-	-	-	1	2	2	1
Female	17	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,992	-	-	-	-	-	-	1	3	7	13	57
Male	1,211	-	-	-	-	-	-	1	3	6	11	45
Female	781	-	-	-	-	-	-	-	-	1	2	12
White, both sexes	1,806	-	-	-	-	-	-	1	1	5	13	47
Male	1,109	-	-	-	-	-	-	1	1	4	11	39
Female	697	-	-	-	-	-	-	-	-	1	2	8
Black, both sexes	114	-	-	-	-	-	-	-	1	1	-	4
Male	62	-	-	-	-	-	-	-	1	1	-	1
Female	52	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	1	-	-	3
Male	11	-	-	-	-	-	-	-	1	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	53	-	-	-	-	-	-	-	-	1	-	3
Male	29	-	-	-	-	-	-	-	-	1	-	3
Female	24	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
New Hampshire												
All races, both sexes	1,310	-	-	-	-	-	-	1	2	4	13	25
Male	686	-	-	-	-	-	-	1	2	4	12	22
Female	624	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	1,303	-	-	-	-	-	-	-	2	4	13	25
Male	680	-	-	-	-	-	-	-	2	4	12	22
Female	623	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	9,942	-	-	-	-	-	-	2	9	29	65	125
Male	4,737	-	-	-	-	-	-	2	8	22	52	101
Female	5,205	-	-	-	-	-	-	-	1	7	13	24
White, both sexes	8,869	-	-	-	-	-	-	-	7	22	52	97
Male	4,233	-	-	-	-	-	-	-	7	17	43	79
Female	4,636	-	-	-	-	-	-	-	-	5	9	18
Black, both sexes	908	-	-	-	-	-	-	1	2	7	9	22
Male	418	-	-	-	-	-	-	1	1	5	5	17
Female	490	-	-	-	-	-	-	-	1	2	4	5
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	152	-	-	-	-	-	-	1	-	-	4	5
Male	81	-	-	-	-	-	-	1	-	-	4	5
Female	71	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
New Mexico												
All races, both sexes	1,752	-	-	-	-	-	-	1	3	4	11	29
Male	975	-	-	-	-	-	-	1	3	3	10	24
Female	777	-	-	-	-	-	-	-	-	1	1	5
White, both sexes	1,645	-	-	-	-	-	-	1	-	4	10	24
Male	910	-	-	-	-	-	-	1	-	3	9	20
Female	735	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	34	-	-	-	-	-	-	-	-	-	1	2
Male	22	-	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	68	-	-	-	-	-	-	-	3	-	-	3
Male	41	-	-	-	-	-	-	-	3	-	-	3
Female	27	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	31,028	-	-	-	-	2	2	11	19	64	168	367
Male	13,968	-	-	-	-	2	1	9	13	51	123	299
Female	17,060	-	-	-	-	-	1	2	6	13	45	68
White, both sexes	26,136	-	-	-	-	2	2	6	14	48	118	255
Male	11,730	-	-	-	-	2	1	5	9	38	92	220
Female	14,406	-	-	-	-	-	1	1	5	10	26	35
Black, both sexes	4,151	-	-	-	-	-	-	4	3	15	39	94
Male	1,838	-	-	-	-	-	-	4	3	12	23	63
Female	2,313	-	-	-	-	-	-	-	-	3	16	31
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	714	-	-	-	-	-	-	1	2	1	10	18
Male	383	-	-	-	-	-	-	-	1	1	7	16
Female	331	-	-	-	-	-	-	1	1	-	3	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
North Carolina												
All races, both sexes	7,721	-	-	-	1	1	3	3	19	42	104	230
Male	4,170	-	-	-	1	1	3	-	13	30	85	171
Female	3,551	-	-	-	-	-	-	3	6	12	19	59
White, both sexes	6,214	-	-	-	1	-	2	3	14	26	73	166
Male	3,405	-	-	-	1	-	2	-	9	19	59	126
Female	2,809	-	-	-	-	-	-	3	5	7	14	40
Black, both sexes	1,407	-	-	-	-	1	1	-	5	16	28	61
Male	704	-	-	-	-	1	1	-	4	11	23	43
Female	703	-	-	-	-	-	-	-	1	5	5	18
American Indian 1/, both sexes	74	-	-	-	-	-	-	-	-	-	1	3
Male	41	-	-	-	-	-	-	-	-	-	1	2
Female	33	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	2	-
Male	20	-	-	-	-	-	-	-	-	-	2	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	675	-	-	-	-	-	1	-	2	3	7	10
Male	363	-	-	-	-	-	1	-	1	2	6	8
Female	312	-	-	-	-	-	-	-	1	1	1	2
White, both sexes	656	-	-	-	-	-	1	-	2	3	7	7
Male	351	-	-	-	-	-	1	-	1	2	6	6
Female	305	-	-	-	-	-	-	-	1	1	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	-	-	-	3
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	7	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Ohio												
All races, both sexes	13,501	-	-	-	-	-	4	8	22	58	154	264
Male	6,879	-	-	-	-	-	3	8	20	44	118	190
Female	6,622	-	-	-	-	-	1	-	2	14	36	74
White, both sexes	12,075	-	-	-	-	-	1	6	18	42	132	220
Male	6,131	-	-	-	-	-	1	6	17	32	103	161
Female	5,944	-	-	-	-	-	-	-	1	10	29	59
Black, both sexes	1,391	-	-	-	-	-	2	2	4	15	22	43
Male	727	-	-	-	-	-	1	2	3	11	15	28
Female	664	-	-	-	-	-	1	-	1	4	7	15
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	1	-	-	1	-	1
Male	19	-	-	-	-	-	1	-	-	1	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4,640	-	-	-	-	-	2	3	4	19	69	105
Male	2,427	-	-	-	-	-	2	2	3	17	53	79
Female	2,213	-	-	-	-	-	-	1	1	2	16	26
White, both sexes	4,144	-	-	-	-	-	2	3	3	9	58	87
Male	2,147	-	-	-	-	-	2	2	2	7	43	66
Female	1,997	-	-	-	-	-	-	1	1	2	15	21
Black, both sexes	239	-	-	-	-	-	-	-	-	4	5	6
Male	135	-	-	-	-	-	-	-	-	4	4	3
Female	104	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	247	-	-	-	-	-	-	-	1	6	6	11
Male	139	-	-	-	-	-	-	-	1	6	6	10
Female	108	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Oregon												
All races, both sexes	2,715	-	-	-	-	-	1	1	3	16	32	51
Male	1,600	-	-	-	-	-	1	1	3	14	26	38
Female	1,115	-	-	-	-	-	-	-	-	2	6	13
White, both sexes	2,652	-	-	-	-	-	1	-	3	14	32	49
Male	1,555	-	-	-	-	-	1	-	3	12	26	36
Female	1,097	-	-	-	-	-	-	-	-	2	6	13
Black, both sexes	23	-	-	-	-	-	-	1	-	-	-	1
Male	18	-	-	-	-	-	-	1	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	2	-	1
Male	17	-	-	-	-	-	-	-	-	2	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	15,338	-	-	-	-	-	-	5	12	48	109	217
Male	7,509	-	-	-	-	-	-	5	9	36	86	167
Female	7,829	-	-	-	-	-	-	-	3	12	23	50
White, both sexes	14,021	-	-	-	-	-	-	5	10	39	92	175
Male	6,817	-	-	-	-	-	-	5	7	32	74	138
Female	7,204	-	-	-	-	-	-	-	3	7	18	37
Black, both sexes	1,258	-	-	-	-	-	-	-	1	9	17	42
Male	657	-	-	-	-	-	-	-	1	4	12	29
Female	601	-	-	-	-	-	-	-	-	5	5	13
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	53	-	-	-	-	-	-	-	1	-	-	-
Male	32	-	-	-	-	-	-	-	1	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Rhode Island												
All races, both sexes	1,599	-	-	-	-	-	-	2	3	5	17	30
Male	693	-	-	-	-	-	-	2	1	4	12	24
Female	906	-	-	-	-	-	-	-	2	1	5	6
White, both sexes	1,548	-	-	-	-	-	-	2	3	4	16	26
Male	663	-	-	-	-	-	-	2	1	3	11	21
Female	885	-	-	-	-	-	-	-	2	1	5	5
Black, both sexes	35	-	-	-	-	-	-	-	-	1	1	2
Male	23	-	-	-	-	-	-	-	-	1	1	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2,937	-	-	-	-	1	3	2	5	14	36	73
Male	1,653	-	-	-	-	-	2	2	2	11	24	58
Female	1,284	-	-	-	-	1	1	-	3	3	12	15
White, both sexes	2,341	-	-	-	-	1	1	-	3	10	29	45
Male	1,345	-	-	-	-	-	1	-	2	7	20	34
Female	996	-	-	-	-	1	-	-	1	3	9	11
Black, both sexes	590	-	-	-	-	-	2	2	2	4	7	28
Male	306	-	-	-	-	-	1	2	-	4	4	24
Female	284	-	-	-	-	-	1	-	2	-	3	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
South Dakota												
All races, both sexes	789	-	-	-	-	-	-	-	-	2	1	13
Male	420	-	-	-	-	-	-	-	-	1	1	10
Female	369	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	763	-	-	-	-	-	-	-	-	2	1	12
Male	405	-	-	-	-	-	-	-	-	1	1	9
Female	358	-	-	-	-	-	-	-	-	1	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6,264	-	-	-	-	-	2	1	19	28	93	156
Male	3,175	-	-	-	-	-	1	1	16	20	68	110
Female	3,089	-	-	-	-	-	1	-	3	8	25	46
White, both sexes	5,490	-	-	-	-	-	1	1	12	24	72	122
Male	2,787	-	-	-	-	-	1	1	10	17	54	85
Female	2,703	-	-	-	-	-	-	-	2	7	18	37
Black, both sexes	750	-	-	-	-	-	1	-	6	4	20	33
Male	372	-	-	-	-	-	-	-	5	3	13	24
Female	378	-	-	-	-	-	1	-	1	1	7	9
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	1	-	1	1
Male	11	-	-	-	-	-	-	-	1	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Texas												
All races, both sexes	15,678	-	1	1	-	-	4	13	27	85	203	417
Male	8,378	-	1	1	-	-	3	12	20	62	158	325
Female	7,300	-	-	-	-	-	1	1	7	23	45	92
White, both sexes	13,707	-	1	1	-	-	2	9	19	62	159	317
Male	7,339	-	1	1	-	-	1	8	14	46	127	251
Female	6,368	-	-	-	-	-	1	1	5	16	32	66
Black, both sexes	1,794	-	-	-	-	-	1	4	6	21	41	88
Male	946	-	-	-	-	-	1	4	4	14	28	63
Female	848	-	-	-	-	-	-	-	2	7	13	25
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	1	3
Male	8	-	-	-	-	-	-	-	-	-	1	2
Female	7	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	162	-	-	-	-	-	1	-	2	2	2	9
Male	85	-	-	-	-	-	1	-	2	2	2	9
Female	77	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	914	-	-	-	-	1	-	1	1	4	13	30
Male	539	-	-	-	-	1	-	1	1	3	11	26
Female	375	-	-	-	-	-	-	-	-	1	2	4
White, both sexes	885	-	-	-	-	1	-	1	1	3	11	29
Male	520	-	-	-	-	1	-	1	1	3	9	25
Female	365	-	-	-	-	-	-	-	-	-	2	4
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	1	2	-
Male	12	-	-	-	-	-	-	-	-	-	2	-
Female	7	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Vermont												
All races, both sexes	630	-	-	-	-	-	-	-	1	2	3	20
Male	320	-	-	-	-	-	-	-	1	-	3	17
Female	310	-	-	-	-	-	-	-	-	2	-	3
White, both sexes	628	-	-	-	-	-	-	-	1	2	3	20
Male	319	-	-	-	-	-	-	-	1	-	3	17
Female	309	-	-	-	-	-	-	-	-	2	-	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5,024	1	-	-	-	-	2	1	17	22	61	126
Male	2,669	-	-	-	-	-	2	1	13	15	51	101
Female	2,355	1	-	-	-	-	-	-	4	7	10	25
White, both sexes	4,070	-	-	-	-	-	2	1	12	17	43	92
Male	2,188	-	-	-	-	-	2	1	8	13	35	78
Female	1,882	-	-	-	-	-	-	-	4	4	8	14
Black, both sexes	877	1	-	-	-	-	-	-	3	5	15	30
Male	430	-	-	-	-	-	-	-	3	2	13	19
Female	447	1	-	-	-	-	-	-	-	3	2	11
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	75	-	-	-	-	-	-	-	2	-	3	4
Male	50	-	-	-	-	-	-	-	2	-	3	4
Female	25	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Washington												
All races, both sexes	5,319	-	-	-	-	1	3	2	6	16	45	83
Male	2,990	-	-	-	-	-	3	2	5	12	36	56
Female	2,329	-	-	-	-	1	-	-	1	4	9	27
White, both sexes	5,016	-	-	-	-	-	1	2	5	11	41	73
Male	2,806	-	-	-	-	-	1	2	4	9	32	49
Female	2,210	-	-	-	-	-	-	-	1	2	9	24
Black, both sexes	122	-	-	-	-	1	1	-	-	3	1	4
Male	72	-	-	-	-	-	1	-	-	2	1	2
Female	50	-	-	-	-	1	-	-	-	1	-	2
American Indian 1/, both sexes	48	-	-	-	-	-	-	-	-	-	-	1
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	133	-	-	-	-	-	1	-	1	2	3	5
Male	82	-	-	-	-	-	1	-	1	1	3	4
Female	51	-	-	-	-	-	-	-	-	1	-	1
West Virginia												
All races, both sexes	2,505	-	-	-	-	1	-	1	6	14	30	64
Male	1,340	-	-	-	-	1	-	1	5	9	25	50
Female	1,165	-	-	-	-	-	-	-	1	5	5	14
White, both sexes	2,426	-	-	-	-	1	-	-	6	13	28	61
Male	1,302	-	-	-	-	1	-	-	5	8	23	48
Female	1,124	-	-	-	-	-	-	-	1	5	5	13
Black, both sexes	79	-	-	-	-	-	-	1	-	1	2	3
Male	38	-	-	-	-	-	-	1	-	1	2	2
Female	41	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other forms of chronic ischemic heart disease (I20,I25)												
Wisconsin												
All races, both sexes	4,433	-	-	-	-	-	-	2	9	7	39	73
Male	2,365	-	-	-	-	-	-	1	7	6	26	56
Female	2,068	-	-	-	-	-	-	1	2	1	13	17
White, both sexes	4,214	-	-	-	-	-	-	1	5	7	33	63
Male	2,236	-	-	-	-	-	-	-	5	6	23	48
Female	1,978	-	-	-	-	-	-	1	-	1	10	15
Black, both sexes	179	-	-	-	-	-	-	-	2	-	4	9
Male	104	-	-	-	-	-	-	-	1	-	3	7
Female	75	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	2	-	2	-
Male	16	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	1	-	2	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	1	-	-	-	1
Male	9	-	-	-	-	-	-	1	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	309	-	-	-	-	-	-	-	-	3	6	5
Male	174	-	-	-	-	-	-	-	-	2	4	3
Female	135	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	305	-	-	-	-	-	-	-	-	3	5	5
Male	172	-	-	-	-	-	-	-	-	2	3	3
Female	133	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
United States												
All races, both sexes	9,255	12,449	15,282	18,034	24,807	37,141	50,910	52,307	40,040	16,633	3,854	25
Male	7,094	9,227	10,799	11,926	15,531	21,368	25,994	22,624	13,188	3,884	585	15
Female	2,161	3,222	4,483	6,108	9,276	15,773	24,916	29,683	26,852	12,749	3,269	10
White, both sexes	7,079	9,782	12,254	14,774	20,946	32,467	45,811	48,205	37,056	15,328	3,427	17
Male	5,567	7,437	8,837	9,942	13,355	19,026	23,688	21,122	12,298	3,592	515	10
Female	1,512	2,345	3,417	4,832	7,591	13,441	22,123	27,083	24,758	11,736	2,912	7
Black, both sexes	1,904	2,330	2,619	2,734	3,225	3,890	4,134	3,256	2,378	1,056	363	7
Male	1,315	1,542	1,664	1,631	1,800	1,916	1,810	1,129	636	200	48	4
Female	589	788	955	1,103	1,425	1,974	2,324	2,127	1,742	856	315	3
American Indian 1/, both sexes	78	98	122	117	135	138	120	114	85	30	11	-
Male	53	67	92	72	78	76	64	48	21	9	-	-
Female	25	31	30	45	57	62	56	66	64	21	11	-
Asian or Pacific Islander 2/, both sexes	194	239	287	409	501	646	845	732	521	219	53	1
Male	159	181	206	281	298	350	432	325	233	83	22	1
Female	35	58	81	128	203	296	413	407	288	136	31	-
Alabama												
All races, both sexes	104	170	187	238	331	492	564	482	380	124	19	-
Male	72	124	112	155	199	264	277	183	106	30	2	-
Female	32	46	75	83	132	228	287	299	274	94	17	-
White, both sexes	75	125	138	181	263	399	490	426	329	99	13	-
Male	57	96	86	126	161	226	249	163	96	22	2	-
Female	18	29	52	55	102	173	241	263	233	77	11	-
Black, both sexes	26	43	48	57	66	91	74	54	51	25	6	-
Male	14	27	25	29	38	38	28	20	10	8	-	-
Female	12	16	23	28	28	53	46	34	41	17	6	-
American Indian 1/, both sexes	2	2	1	-	1	-	-	1	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	2	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Alaska												
All races, both sexes	24	37	21	24	27	40	46	27	10	5	1	-
Male	14	28	19	17	21	30	27	16	3	1	-	-
Female	10	9	2	7	6	10	19	11	7	4	1	-
White, both sexes	21	31	18	19	18	31	43	21	8	3	1	-
Male	12	24	17	13	15	23	26	13	3	-	-	-
Female	9	7	1	6	3	8	17	8	5	3	1	-
Black, both sexes	1	1	-	1	-	1	-	2	-	-	-	-
Male	1	-	-	1	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	2	5	2	3	8	8	2	3	2	2	-	-
Male	1	4	1	3	5	7	1	2	-	1	-	-
Female	1	1	1	-	3	1	1	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	1	1	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Arizona												
All races, both sexes	210	292	392	416	585	815	1,076	1,047	721	262	50	2
Male	158	235	294	282	385	473	559	515	274	67	9	-
Female	52	57	98	134	200	342	517	532	447	195	41	2
White, both sexes	190	270	369	378	551	769	1,048	1,019	704	259	47	2
Male	143	219	283	253	368	453	546	499	270	65	9	-
Female	47	51	86	125	183	316	502	520	434	194	38	2
Black, both sexes	11	6	11	19	16	24	16	15	7	1	1	-
Male	9	5	5	16	9	9	7	4	1	1	-	-
Female	2	1	6	3	7	15	9	11	6	-	1	-
American Indian 1/, both sexes	7	15	8	16	15	16	9	7	9	1	1	-
Male	4	11	4	11	6	8	4	6	3	1	-	-
Female	3	4	4	5	9	8	5	1	6	-	1	-
Asian or Pacific Islander 2/, both sexes	2	1	4	3	3	6	3	6	1	1	1	-
Male	2	-	2	2	2	3	2	6	-	-	-	-
Female	-	1	2	1	1	3	1	-	1	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Arkansas												
All races, both sexes	66	82	114	154	205	290	358	389	287	146	18	-
Male	46	56	71	97	133	153	190	154	90	29	2	-
Female	20	26	43	57	72	137	168	235	197	117	16	-
White, both sexes	53	65	88	133	184	247	325	356	261	125	14	-
Male	37	46	59	83	121	138	174	145	82	26	2	-
Female	16	19	29	50	63	109	151	211	179	99	12	-
Black, both sexes	12	17	26	21	20	41	33	33	26	21	4	-
Male	8	10	12	14	11	15	16	9	8	3	-	-
Female	4	7	14	7	9	26	17	24	18	18	4	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
California												
All races, both sexes	1,081	1,518	1,784	1,990	2,753	4,137	5,989	6,162	4,724	2,012	541	1
Male	854	1,155	1,271	1,305	1,730	2,399	3,091	2,718	1,685	514	100	1
Female	227	363	513	685	1,023	1,738	2,898	3,444	3,039	1,498	441	-
White, both sexes	829	1,173	1,422	1,585	2,251	3,494	5,227	5,489	4,248	1,804	474	1
Male	661	911	1,021	1,048	1,416	2,062	2,716	2,441	1,496	456	81	1
Female	168	262	401	537	835	1,432	2,511	3,048	2,752	1,348	393	-
Black, both sexes	168	244	242	227	287	335	337	283	200	98	45	-
Male	125	166	167	137	184	169	152	102	59	16	7	-
Female	43	78	75	90	103	166	185	181	141	82	38	-
American Indian 1/, both sexes	6	13	17	22	16	13	12	19	9	1	-	-
Male	4	8	8	15	10	7	6	6	3	-	-	-
Female	2	5	9	7	6	6	6	13	6	1	-	-
Asian or Pacific Islander 2/, both sexes	78	88	103	156	199	295	413	371	267	109	22	-
Male	64	70	75	105	120	161	217	169	127	42	12	-
Female	14	18	28	51	79	134	196	202	140	67	10	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Colorado												
All races, both sexes	106	130	143	180	240	341	468	495	345	144	33	-
Male	89	113	110	131	158	210	257	245	124	28	5	-
Female	17	17	33	49	82	131	211	250	221	116	28	-
White, both sexes	105	115	128	167	219	323	457	483	339	140	33	-
Male	89	100	98	123	150	204	253	241	123	28	5	-
Female	16	15	30	44	69	119	204	242	216	112	28	-
Black, both sexes	1	13	13	10	19	12	4	10	5	2	-	-
Male	-	12	10	5	6	4	-	2	1	-	-	-
Female	1	1	3	5	13	8	4	8	4	2	-	-
American Indian 1/, both sexes	-	-	1	1	1	3	1	-	-	2	-	-
Male	-	-	1	1	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	2	1	3	6	2	1	-	-	-
Male	-	1	1	2	1	1	3	2	-	-	-	-
Female	-	1	-	-	-	2	3	-	1	-	-	-
Connecticut												
All races, both sexes	104	124	146	174	253	414	621	718	597	275	62	2
Male	79	95	102	125	158	235	323	308	191	49	10	2
Female	25	29	44	49	95	179	298	410	406	226	52	-
White, both sexes	83	101	126	146	229	388	594	691	580	260	57	2
Male	68	78	88	107	146	224	310	298	184	44	10	2
Female	15	23	38	39	83	164	284	393	396	216	47	-
Black, both sexes	20	18	18	27	18	24	21	26	17	11	3	-
Male	10	13	12	17	11	10	11	10	7	2	-	-
Female	10	5	6	10	7	14	10	16	10	9	3	-
American Indian 1/, both sexes	-	-	-	1	2	1	1	-	-	1	1	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	1	5	2	-	4	1	5	1	-	3	1	-
Male	1	4	2	-	-	1	1	-	-	3	-	-
Female	-	1	-	-	4	-	4	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Delaware												
All races, both sexes	42	47	74	62	68	141	176	183	133	51	6	-
Male	31	34	47	39	46	80	94	77	38	13	1	-
Female	11	13	27	23	22	61	82	106	95	38	5	-
White, both sexes	26	35	54	49	56	124	155	159	114	45	4	-
Male	22	28	38	32	37	75	84	72	33	13	1	-
Female	4	7	16	17	19	49	71	87	81	32	3	-
Black, both sexes	15	11	20	11	12	16	20	24	17	6	2	-
Male	9	6	9	6	9	4	9	5	3	-	-	-
Female	6	5	11	5	3	12	11	19	14	6	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	-	1	1	-	1	-	-	-
Male	-	-	-	1	-	1	1	-	1	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	41	47	67	69	88	119	134	125	81	44	16	2
Male	35	33	47	46	48	58	61	47	23	11	5	1
Female	6	14	20	23	40	61	73	78	58	33	11	1
White, both sexes	4	7	11	5	16	14	33	37	31	17	5	1
Male	2	5	8	4	7	9	15	19	12	7	2	-
Female	2	2	3	1	9	5	18	18	19	10	3	1
Black, both sexes	36	39	56	63	70	104	98	86	49	25	11	1
Male	32	27	39	41	39	48	44	26	11	3	3	1
Female	4	12	17	22	31	56	54	60	38	22	8	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	2	1	2	2	1	2	-	-
Male	1	1	-	1	2	1	2	2	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Florida												
All races, both sexes	667	794	1,163	1,422	1,992	3,282	4,573	4,746	3,722	1,527	331	1
Male	529	581	870	996	1,288	1,987	2,433	2,268	1,372	426	63	1
Female	138	213	293	426	704	1,295	2,140	2,478	2,350	1,101	268	-
White, both sexes	531	657	983	1,237	1,756	3,015	4,279	4,507	3,533	1,448	309	-
Male	427	488	748	885	1,151	1,853	2,311	2,179	1,308	403	61	-
Female	104	169	235	352	605	1,162	1,968	2,328	2,225	1,045	248	-
Black, both sexes	131	127	171	173	225	255	272	224	183	72	20	1
Male	98	87	115	103	133	129	111	83	59	21	2	1
Female	33	40	56	70	92	126	161	141	124	51	18	-
American Indian 1/, both sexes	1	-	4	2	-	4	4	3	1	-	1	-
Male	1	-	3	1	-	2	4	2	1	-	-	-
Female	-	-	1	1	-	2	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	10	5	10	11	8	18	12	5	7	1	-
Male	3	6	4	7	4	3	7	4	4	2	-	-
Female	1	4	1	3	7	5	11	8	1	5	1	-
Georgia												
All races, both sexes	270	330	394	415	498	708	847	727	549	212	39	-
Male	206	250	267	272	326	405	416	316	162	43	5	-
Female	64	80	127	143	172	303	431	411	387	169	34	-
White, both sexes	170	224	291	277	377	572	698	608	459	173	24	-
Male	133	174	207	199	260	347	353	281	138	38	3	-
Female	37	50	84	78	117	225	345	327	321	135	21	-
Black, both sexes	97	103	103	129	119	134	146	118	87	38	15	-
Male	71	75	60	67	65	57	62	35	24	5	2	-
Female	26	28	43	62	54	77	84	83	63	33	13	-
American Indian 1/, both sexes	-	-	-	2	-	-	1	-	2	-	-	-
Male	-	-	-	2	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	-	7	2	2	2	1	1	1	-	-
Male	2	1	-	4	1	1	-	-	-	-	-	-
Female	1	2	-	3	1	1	2	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Hawaii												
All races, both sexes	35	43	66	63	88	105	147	121	81	42	9	-
Male	30	36	47	45	59	55	93	68	40	20	4	-
Female	5	7	19	18	29	50	54	53	41	22	5	-
White, both sexes	7	14	27	12	29	35	45	43	27	15	2	-
Male	7	13	22	10	19	19	29	25	13	7	1	-
Female	-	1	5	2	10	16	16	18	14	8	1	-
Black, both sexes	1	-	1	1	-	-	2	-	1	-	-	-
Male	1	-	1	-	-	-	2	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	29	38	50	59	70	100	78	53	27	7	-
Male	22	23	24	35	40	36	62	43	26	13	3	-
Female	5	6	14	15	19	34	38	35	27	14	4	-
Idaho												
All races, both sexes	21	17	31	46	72	114	151	156	138	37	6	-
Male	15	13	25	32	45	74	90	78	55	11	1	-
Female	6	4	6	14	27	40	61	78	83	26	5	-
White, both sexes	20	16	29	44	71	114	147	155	137	37	6	-
Male	14	12	23	30	44	74	88	78	55	11	1	-
Female	6	4	6	14	27	40	59	77	82	26	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	2	1	-	-	1	1	-	-	-	-
Male	-	1	2	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	-	3	-	1	-	-	-
Male	1	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Illinois												
All races, both sexes	426	509	564	695	930	1,401	1,858	2,099	1,609	677	140	-
Male	322	391	400	460	549	835	962	877	492	139	12	-
Female	104	118	164	235	381	566	896	1,222	1,117	538	128	-
White, both sexes	285	361	407	521	734	1,159	1,630	1,922	1,484	620	126	-
Male	224	288	297	351	447	708	871	809	461	126	10	-
Female	61	73	110	170	287	451	759	1,113	1,023	494	116	-
Black, both sexes	140	143	150	159	176	222	205	157	115	50	13	-
Male	97	98	96	99	90	117	85	58	29	11	2	-
Female	43	45	54	60	86	105	120	99	86	39	11	-
American Indian 1/, both sexes	-	-	2	1	1	1	-	1	1	-	-	-
Male	-	-	2	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	5	5	14	19	19	23	19	9	7	1	-
Male	1	5	5	10	11	10	6	10	2	2	-	-
Female	-	-	-	4	8	9	17	9	7	5	1	-
Indiana												
All races, both sexes	207	249	290	370	506	752	1,000	997	736	325	56	-
Male	162	182	211	239	295	442	546	431	227	76	6	-
Female	45	67	79	131	211	310	454	566	509	249	50	-
White, both sexes	177	222	259	318	437	693	956	961	709	312	50	-
Male	140	163	192	210	265	411	520	419	221	75	6	-
Female	37	59	67	108	172	282	436	542	488	237	44	-
Black, both sexes	29	27	31	47	66	56	39	33	24	13	6	-
Male	22	19	19	25	29	30	24	11	4	1	-	-
Female	7	8	12	22	37	26	15	22	20	12	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	5	3	3	5	3	3	-	-	-
Male	-	-	-	4	1	1	2	1	2	-	-	-
Female	1	-	-	1	2	2	3	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Iowa												
All races, both sexes	97	130	158	195	237	448	678	724	587	281	93	-
Male	72	95	118	128	148	254	374	340	213	63	13	-
Female	25	35	40	67	89	194	304	384	374	218	80	-
White, both sexes	93	123	153	190	232	446	667	722	587	279	92	-
Male	69	91	113	124	145	253	367	339	213	63	13	-
Female	24	32	40	66	87	193	300	383	374	216	79	-
Black, both sexes	4	6	4	3	3	2	9	-	-	2	1	-
Male	3	4	4	2	2	1	6	-	-	-	-	-
Female	1	2	-	1	1	1	3	-	-	2	1	-
American Indian 1/, both sexes	-	-	1	-	2	-	1	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	-	-	1	2	-	-	-	-
Male	-	-	-	2	-	-	1	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Kansas												
All races, both sexes	84	103	132	146	218	295	415	453	352	178	36	-
Male	63	85	93	98	144	184	223	219	136	47	8	-
Female	21	18	39	48	74	111	192	234	216	131	28	-
White, both sexes	76	92	114	133	205	277	389	446	340	174	35	-
Male	58	74	80	89	137	175	210	216	129	47	8	-
Female	18	18	34	44	68	102	179	230	211	127	27	-
Black, both sexes	6	6	13	12	8	16	18	7	9	4	-	-
Male	4	6	9	8	4	9	11	3	5	-	-	-
Female	2	-	4	4	4	7	7	4	4	4	-	-
American Indian 1/, both sexes	2	4	5	-	4	1	4	-	2	-	1	-
Male	1	4	4	-	2	-	1	-	2	-	-	-
Female	1	-	1	-	2	1	3	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	1	4	-	1	-	-	-
Male	-	1	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	3	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Kentucky												
All races, both sexes	116	151	189	243	329	468	595	560	401	173	39	-
Male	87	103	126	148	209	268	263	219	114	28	5	-
Female	29	48	63	95	120	200	332	341	287	145	34	-
White, both sexes	105	144	179	228	317	442	574	542	383	166	36	-
Male	80	100	116	139	204	254	257	212	111	27	4	-
Female	25	44	63	89	113	188	317	330	272	139	32	-
Black, both sexes	10	6	10	13	12	26	21	18	17	7	3	-
Male	6	2	10	7	5	14	6	7	3	1	1	-
Female	4	4	-	6	7	12	15	11	14	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	2	-	-	-	-	1	-	-	-
Male	1	1	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Louisiana												
All races, both sexes	144	184	225	277	397	489	576	552	410	151	38	-
Male	101	125	155	178	262	277	303	240	125	40	9	-
Female	43	59	70	99	135	212	273	312	285	111	29	-
White, both sexes	97	118	153	197	276	394	457	460	337	120	26	-
Male	72	85	114	130	187	228	245	209	100	31	5	-
Female	25	33	39	67	89	166	212	251	237	89	21	-
Black, both sexes	47	65	71	78	117	89	117	89	73	31	12	-
Male	29	39	40	46	73	46	57	29	25	9	4	-
Female	18	26	31	32	44	43	60	60	48	22	8	-
American Indian 1/, both sexes	-	-	1	-	-	2	1	1	-	-	-	-
Male	-	-	1	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	4	4	1	2	-	-	-	-
Male	-	1	-	2	2	3	-	1	-	-	-	-
Female	-	-	-	-	2	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Maine												
All races, both sexes	40	39	56	88	119	160	214	225	177	66	12	-
Male	32	32	40	62	79	83	117	111	56	16	-	-
Female	8	7	16	26	40	77	97	114	121	50	12	-
White, both sexes	38	38	56	88	119	160	213	225	177	66	12	-
Male	31	31	40	62	79	83	116	111	56	16	-	-
Female	7	7	16	26	40	77	97	114	121	50	12	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	217	280	352	362	466	644	933	957	671	290	75	4
Male	158	188	245	228	272	344	470	415	199	60	14	2
Female	59	92	107	134	194	300	463	542	472	230	61	2
White, both sexes	114	181	224	226	318	490	746	789	560	243	61	2
Male	91	132	167	144	183	269	389	358	168	52	13	1
Female	23	49	57	82	135	221	357	431	392	191	48	1
Black, both sexes	101	95	120	125	136	145	178	155	105	41	14	2
Male	65	55	73	74	84	71	77	52	29	7	1	1
Female	36	40	47	51	52	74	101	103	76	34	13	1
American Indian 1/, both sexes	1	1	1	-	2	-	1	1	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	7	11	10	9	8	12	6	6	-	-
Male	1	-	4	10	5	4	4	5	2	1	-	-
Female	-	3	3	1	5	5	4	7	4	5	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Massachusetts												
All races, both sexes	121	159	243	286	414	592	860	1,046	802	322	94	-
Male	103	120	189	185	271	377	465	422	249	79	9	-
Female	18	39	54	101	143	215	395	624	553	243	85	-
White, both sexes	108	153	230	260	386	564	822	1,027	787	316	92	-
Male	92	116	178	171	260	362	447	413	248	77	9	-
Female	16	37	52	89	126	202	375	614	539	239	83	-
Black, both sexes	11	6	10	23	23	25	24	18	11	4	2	-
Male	9	4	9	13	9	14	11	8	-	1	-	-
Female	2	2	1	10	14	11	13	10	11	3	2	-
American Indian 1/, both sexes	-	-	2	1	2	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	2	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	2	3	3	14	1	4	2	-	-
Male	2	-	1	1	2	1	7	1	1	1	-	-
Female	-	-	-	1	1	2	7	-	3	1	-	-
Michigan												
All races, both sexes	515	723	742	772	1,079	1,559	2,132	2,094	1,571	610	121	-
Male	374	527	499	506	685	877	1,071	846	522	147	18	-
Female	141	196	243	266	394	682	1,061	1,248	1,049	463	103	-
White, both sexes	344	485	527	608	849	1,285	1,861	1,889	1,423	550	104	-
Male	264	364	359	396	550	739	950	769	474	130	14	-
Female	80	121	168	212	299	546	911	1,120	949	420	90	-
Black, both sexes	166	231	204	156	213	261	262	194	143	58	16	-
Male	107	157	129	104	124	129	114	74	44	15	4	-
Female	59	74	75	52	89	132	148	120	99	43	12	-
American Indian 1/, both sexes	3	4	3	3	10	6	2	3	1	-	-	-
Male	1	3	3	1	7	4	1	-	-	-	-	-
Female	2	1	-	2	3	2	1	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	8	5	7	7	7	8	4	2	1	-
Male	2	3	8	5	4	5	6	3	4	2	-	-
Female	-	-	-	-	3	2	1	5	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Minnesota												
All races, both sexes	105	149	144	175	234	386	548	631	470	211	50	-
Male	86	125	113	124	159	255	314	340	174	67	14	-
Female	19	24	31	51	75	131	234	291	296	144	36	-
White, both sexes	96	144	137	164	227	378	541	623	466	211	50	-
Male	78	121	107	117	156	249	312	337	172	67	14	-
Female	18	23	30	47	71	129	229	286	294	144	36	-
Black, both sexes	7	3	4	4	3	4	5	4	-	-	-	-
Male	6	2	4	2	1	3	1	3	-	-	-	-
Female	1	1	-	2	2	1	4	1	-	-	-	-
American Indian 1/, both sexes	1	-	2	3	2	1	2	1	2	-	-	-
Male	1	-	1	2	1	1	1	-	1	-	-	-
Female	-	-	1	1	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	4	2	3	-	3	2	-	-	-
Male	1	2	1	3	1	2	-	-	1	-	-	-
Female	-	-	-	1	1	1	-	3	1	-	-	-
Mississippi												
All races, both sexes	105	132	164	211	269	355	414	371	291	132	24	1
Male	72	90	103	148	153	185	200	130	80	19	3	1
Female	33	42	61	63	116	170	214	241	211	113	21	-
White, both sexes	62	88	109	155	197	284	331	307	237	112	13	1
Male	40	61	70	110	121	151	158	108	67	17	1	1
Female	22	27	39	45	76	133	173	199	170	95	12	-
Black, both sexes	43	42	55	55	69	68	83	64	54	19	11	-
Male	32	27	33	38	32	33	42	22	13	2	2	-
Female	11	15	22	17	37	35	41	42	41	17	9	-
American Indian 1/, both sexes	-	2	-	-	2	1	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	2	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Missouri												
All races, both sexes	174	221	265	349	480	737	959	1,091	820	371	97	-
Male	126	168	187	217	298	424	486	432	256	99	10	-
Female	48	53	78	132	182	313	473	659	564	272	87	-
White, both sexes	142	182	221	299	421	675	885	1,026	776	348	86	-
Male	106	140	160	190	265	393	448	413	241	92	8	-
Female	36	42	61	109	156	282	437	613	535	256	78	-
Black, both sexes	32	39	43	47	57	57	72	61	44	22	11	-
Male	20	28	27	27	32	29	37	18	15	7	2	-
Female	12	11	16	20	25	28	35	43	29	15	9	-
American Indian 1/, both sexes	-	-	-	2	-	1	2	2	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	2	-	-	1	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	4	-	2	-	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	1	1	1	3	-	1	-	1	-	-
Montana												
All races, both sexes	33	41	40	53	60	74	112	122	78	36	10	-
Male	27	31	30	41	38	48	65	66	35	19	-	-
Female	6	10	10	12	22	26	47	56	43	17	10	-
White, both sexes	29	39	38	51	58	70	112	120	75	35	9	-
Male	23	29	28	40	38	44	65	65	35	18	-	-
Female	6	10	10	11	20	26	47	55	40	17	9	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	1	2	2	2	3	-	2	3	1	1	-
Male	4	1	2	1	-	3	-	1	-	1	-	-
Female	-	-	-	1	2	-	-	1	3	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Nebraska												
All races, both sexes	39	34	55	77	90	168	232	282	262	113	26	-
Male	31	28	40	51	58	111	130	141	102	29	4	-
Female	8	6	15	26	32	57	102	141	160	84	22	-
White, both sexes	31	29	51	68	83	161	231	281	260	111	26	-
Male	24	25	37	46	57	109	130	140	101	28	4	-
Female	7	4	14	22	26	52	101	141	159	83	22	-
Black, both sexes	7	5	2	7	4	5	1	-	-	2	-	-
Male	6	3	1	5	-	2	-	-	-	1	-	-
Female	1	2	1	2	4	3	1	-	-	1	-	-
American Indian 1/, both sexes	1	-	1	1	2	1	-	1	2	-	-	-
Male	1	-	1	-	1	-	-	1	1	-	-	-
Female	-	-	-	1	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
Nevada												
All races, both sexes	99	140	187	203	226	310	332	241	137	29	7	-
Male	80	106	136	132	151	186	188	109	50	5	2	-
Female	19	34	51	71	75	124	144	132	87	24	5	-
White, both sexes	91	122	167	179	201	286	300	230	129	27	7	-
Male	74	97	123	118	137	174	170	106	47	5	2	-
Female	17	25	44	61	64	112	130	124	82	22	5	-
Black, both sexes	5	14	13	16	15	13	20	6	5	1	-	-
Male	3	8	7	9	9	7	11	2	3	-	-	-
Female	2	6	6	7	6	6	9	4	2	1	-	-
American Indian 1/, both sexes	-	2	2	-	4	3	3	-	1	-	-	-
Male	-	1	2	-	2	2	1	-	-	-	-	-
Female	-	1	-	-	2	1	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	5	8	6	8	9	5	2	1	-	-
Male	3	-	4	5	3	3	6	1	-	-	-	-
Female	-	2	1	3	3	5	3	4	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
New Hampshire												
All races, both sexes	41	49	71	92	111	139	191	251	198	99	23	-
Male	38	41	55	63	74	82	85	110	69	26	2	-
Female	3	8	16	29	37	57	106	141	129	73	21	-
White, both sexes	41	49	71	91	109	137	191	251	197	99	23	-
Male	38	41	55	62	73	80	85	110	68	26	2	-
Female	3	8	16	29	36	57	106	141	129	73	21	-
Black, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	1	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	213	308	420	466	746	1,209	1,822	2,061	1,592	691	180	4
Male	171	231	292	313	468	670	907	846	482	138	31	3
Female	42	77	128	153	278	539	915	1,215	1,110	553	149	1
White, both sexes	156	243	330	381	636	1,058	1,647	1,945	1,502	635	157	1
Male	131	189	241	256	411	596	839	811	458	127	27	1
Female	25	54	89	125	225	462	808	1,134	1,044	508	130	-
Black, both sexes	44	53	80	69	97	136	153	91	75	46	21	2
Male	30	35	44	44	51	64	63	25	22	7	3	1
Female	14	18	36	25	46	72	90	66	53	39	18	1
American Indian 1/, both sexes	-	1	-	1	2	-	1	1	3	3	-	-
Male	-	-	-	-	1	-	1	1	1	1	-	-
Female	-	1	-	1	1	-	-	-	2	2	-	-
Asian or Pacific Islander 2/, both sexes	13	11	10	15	11	15	21	24	12	7	2	1
Male	10	7	7	13	5	10	4	9	1	3	1	1
Female	3	4	3	2	6	5	17	15	11	4	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
New Mexico												
All races, both sexes	66	71	114	135	177	222	331	250	227	88	22	1
Male	53	59	84	89	114	131	174	115	82	27	5	1
Female	13	12	30	46	63	91	157	135	145	61	17	-
White, both sexes	59	64	102	130	165	205	318	240	217	83	22	1
Male	48	54	76	86	107	119	167	108	80	26	5	1
Female	11	10	26	44	58	86	151	132	137	57	17	-
Black, both sexes	-	3	3	1	5	5	6	2	5	1	-	-
Male	-	3	3	1	3	5	3	1	1	-	-	-
Female	-	-	-	-	2	-	3	1	4	1	-	-
American Indian 1/, both sexes	7	4	9	3	6	11	7	8	3	4	-	-
Male	5	2	5	2	4	6	4	6	-	1	-	-
Female	2	2	4	1	2	5	3	2	3	3	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	-	2	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-
New York												
All races, both sexes	645	964	1,279	1,653	2,472	3,914	5,533	5,963	5,179	2,245	548	-
Male	480	683	822	1,025	1,451	2,068	2,544	2,287	1,566	476	68	-
Female	165	281	457	628	1,021	1,846	2,989	3,676	3,613	1,769	480	-
White, both sexes	462	691	889	1,221	1,959	3,199	4,716	5,323	4,716	2,038	477	-
Male	366	506	589	778	1,174	1,720	2,199	2,080	1,455	438	58	-
Female	96	185	300	443	785	1,479	2,517	3,243	3,261	1,600	419	-
Black, both sexes	156	244	346	385	435	612	697	531	356	176	58	-
Male	94	151	204	216	227	292	283	165	66	29	6	-
Female	62	93	142	169	208	320	414	366	290	147	52	-
American Indian 1/, both sexes	1	4	2	2	1	3	-	7	6	-	-	-
Male	1	3	2	1	1	2	-	4	2	-	-	-
Female	-	1	-	1	-	1	-	3	4	-	-	-
Asian or Pacific Islander 2/, both sexes	26	25	42	45	77	100	120	102	101	31	13	-
Male	19	23	27	30	49	54	62	38	43	9	4	-
Female	7	2	15	15	28	46	58	64	58	22	9	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
North Carolina												
All races, both sexes	320	448	492	604	788	1,021	1,242	1,197	854	285	66	1
Male	243	324	355	423	484	601	609	517	250	56	3	1
Female	77	124	137	181	304	420	633	680	604	229	63	-
White, both sexes	232	331	357	465	618	825	1,048	1,039	732	233	48	1
Male	183	253	264	336	394	501	523	464	223	45	2	1
Female	49	78	93	129	224	324	525	575	509	188	46	-
Black, both sexes	84	106	123	131	158	180	180	150	115	50	18	-
Male	56	63	81	81	82	92	79	51	24	11	1	-
Female	28	43	42	50	76	88	101	99	91	39	17	-
American Indian 1/, both sexes	2	7	8	6	9	15	10	8	4	1	-	-
Male	2	5	7	4	6	7	5	2	-	-	-	-
Female	-	2	1	2	3	8	5	6	4	1	-	-
Asian or Pacific Islander 2/, both sexes	2	4	4	2	3	1	4	-	3	1	-	-
Male	2	3	3	2	2	1	2	-	3	-	-	-
Female	-	1	1	-	1	-	2	-	-	1	-	-
North Dakota												
All races, both sexes	16	24	19	29	38	77	129	154	94	61	11	-
Male	12	19	13	21	28	51	77	73	32	18	1	-
Female	4	5	6	8	10	26	52	81	62	43	10	-
White, both sexes	13	23	19	25	36	75	128	152	94	60	11	-
Male	10	19	13	18	26	49	77	72	32	18	1	-
Female	3	4	6	7	10	26	51	80	62	42	10	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	1	-	3	2	2	1	2	-	1	-	-
Male	2	-	-	2	2	2	-	1	-	-	-	-
Female	1	1	-	1	-	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Ohio												
All races, both sexes	416	551	665	873	1,156	1,813	2,462	2,462	1,745	704	144	-
Male	324	378	466	578	707	1,010	1,236	1,060	567	146	24	-
Female	92	173	199	295	449	803	1,226	1,402	1,178	558	120	-
White, both sexes	322	438	544	727	1,004	1,627	2,257	2,293	1,646	664	134	-
Male	257	310	393	490	620	916	1,133	992	538	139	23	-
Female	65	128	151	237	384	711	1,124	1,301	1,108	525	111	-
Black, both sexes	94	112	117	141	150	185	194	165	96	39	10	-
Male	67	67	70	86	85	94	97	65	28	7	1	-
Female	27	45	47	55	65	91	97	100	68	32	9	-
American Indian 1/, both sexes	-	-	-	-	-	1	2	1	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	4	5	2	-	9	3	2	1	-	-
Male	-	1	3	2	2	-	4	3	1	-	-	-
Female	-	-	1	3	-	-	5	-	1	1	-	-
Oklahoma												
All races, both sexes	159	247	288	311	431	603	776	760	577	231	54	1
Male	121	168	203	199	288	339	397	327	171	51	7	-
Female	38	79	85	112	143	264	379	433	406	180	47	1
White, both sexes	128	203	248	258	382	556	717	700	533	211	45	1
Male	100	144	172	165	253	310	366	305	159	45	6	-
Female	28	59	76	93	129	246	351	395	374	166	39	1
Black, both sexes	13	30	18	23	21	20	23	39	24	10	3	-
Male	9	16	12	19	14	15	13	15	7	3	1	-
Female	4	14	6	4	7	5	10	24	17	7	2	-
American Indian 1/, both sexes	18	13	20	29	27	25	36	20	20	10	5	-
Male	12	8	17	14	20	12	18	7	5	3	-	-
Female	6	5	3	15	7	13	18	13	15	7	5	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	1	2	-	1	-	-	1	-
Male	-	-	2	1	1	2	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Oregon												
All races, both sexes	96	134	154	178	203	331	462	510	374	146	23	-
Male	79	108	123	116	134	203	276	266	158	51	3	-
Female	17	26	31	62	69	128	186	244	216	95	20	-
White, both sexes	86	131	146	174	195	325	455	503	370	145	23	-
Male	71	105	116	114	128	198	272	262	158	50	3	-
Female	15	26	30	60	67	127	183	241	212	95	20	-
Black, both sexes	5	1	4	3	3	1	1	1	2	-	-	-
Male	5	1	3	2	3	1	-	1	-	-	-	-
Female	-	-	1	1	-	-	1	-	2	-	-	-
American Indian 1/, both sexes	4	-	2	-	2	1	3	2	1	1	-	-
Male	2	-	2	-	1	1	2	1	-	1	-	-
Female	2	-	-	-	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	1	3	4	3	4	1	-	-	-
Male	1	2	2	-	2	3	2	2	-	-	-	-
Female	-	-	-	1	1	1	1	2	1	-	-	-
Pennsylvania												
All races, both sexes	354	501	630	791	1,242	2,012	2,941	3,124	2,259	891	199	3
Male	266	381	440	519	767	1,177	1,449	1,283	708	184	31	1
Female	88	120	190	272	475	835	1,492	1,841	1,551	707	168	2
White, both sexes	287	412	527	667	1,092	1,825	2,716	2,972	2,173	843	184	2
Male	216	321	370	447	671	1,073	1,344	1,230	683	175	30	1
Female	71	91	157	220	421	752	1,372	1,742	1,490	668	154	1
Black, both sexes	63	84	98	113	146	179	218	144	82	47	14	1
Male	48	55	66	66	94	99	101	50	24	8	-	-
Female	15	29	32	47	52	80	117	94	58	39	14	1
American Indian 1/, both sexes	-	-	-	-	-	1	1	4	-	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	5	5	11	4	7	6	4	4	1	1	-
Male	2	5	4	6	2	5	3	1	1	1	1	-
Female	2	-	1	5	2	2	3	3	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Rhode Island												
All races, both sexes	32	55	59	55	103	155	279	335	292	144	33	-
Male	28	44	44	35	65	85	127	123	76	20	3	-
Female	4	11	15	20	38	70	152	212	216	124	30	-
White, both sexes	29	52	54	50	99	146	277	328	288	143	31	-
Male	26	41	40	31	64	78	127	121	76	20	1	-
Female	3	11	14	19	35	68	150	207	212	123	30	-
Black, both sexes	1	3	4	5	2	7	1	4	3	-	1	-
Male	1	3	3	4	1	5	-	1	-	-	1	-
Female	-	-	1	1	1	2	1	3	3	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	1	1	2	1	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	1	-	-	-	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	1	1	-	1	-	1	1	-
Male	1	-	1	-	-	1	-	-	-	-	1	-
Female	-	-	-	-	1	-	-	1	-	1	-	-
South Carolina												
All races, both sexes	112	166	210	241	326	431	490	426	281	91	28	1
Male	83	120	143	167	205	245	257	210	104	15	4	1
Female	29	46	67	74	121	186	233	216	177	76	24	-
White, both sexes	82	118	164	187	262	344	408	361	231	77	17	1
Male	64	87	115	136	172	203	226	182	80	13	2	1
Female	18	31	49	51	90	141	182	179	151	64	15	-
Black, both sexes	30	47	46	54	63	85	81	64	50	14	11	-
Male	19	32	28	31	33	41	31	28	24	2	2	-
Female	11	15	18	23	30	44	50	36	26	12	9	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	2	1	1	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
South Dakota												
All races, both sexes	16	20	32	25	52	107	145	135	158	63	20	-
Male	14	12	27	16	32	76	81	64	60	23	3	-
Female	2	8	5	9	20	31	64	71	98	40	17	-
White, both sexes	16	16	27	22	51	103	144	131	156	62	20	-
Male	14	10	23	13	32	74	80	63	59	23	3	-
Female	2	6	4	9	19	29	64	68	97	39	17	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	4	5	3	1	4	1	4	2	1	-	-
Male	-	2	4	3	-	2	1	1	1	-	-	-
Female	-	2	1	-	1	2	-	3	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	251	325	359	468	597	794	1,067	1,027	735	290	52	-
Male	180	241	237	287	364	443	532	391	217	61	6	-
Female	71	84	122	181	233	351	535	636	518	229	46	-
White, both sexes	195	260	299	410	518	693	970	938	667	265	43	-
Male	142	192	197	259	323	392	492	356	205	55	6	-
Female	53	68	102	151	195	301	478	582	462	210	37	-
Black, both sexes	56	61	59	53	75	99	95	88	66	25	9	-
Male	38	46	39	25	39	50	39	35	10	6	-	-
Female	18	15	20	28	36	49	56	53	56	19	9	-
American Indian 1/, both sexes	-	2	1	1	1	-	1	-	-	-	-	-
Male	-	2	1	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	4	3	2	1	1	2	-	-	-
Male	-	1	-	2	2	1	-	-	2	-	-	-
Female	-	1	-	2	1	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Texas												
All races, both sexes	590	870	986	1,135	1,488	2,010	2,726	2,437	1,821	698	166	-
Male	459	630	699	728	925	1,113	1,380	1,055	615	162	30	-
Female	131	240	287	407	563	897	1,346	1,382	1,206	536	136	-
White, both sexes	440	691	811	949	1,283	1,779	2,462	2,261	1,664	642	155	-
Male	353	515	577	619	816	1,003	1,271	994	564	151	27	-
Female	87	176	234	330	467	776	1,191	1,267	1,100	491	128	-
Black, both sexes	141	169	159	168	182	209	242	151	146	55	11	-
Male	97	110	111	97	99	96	104	56	49	10	3	-
Female	44	59	48	71	83	113	138	95	97	45	8	-
American Indian 1/, both sexes	3	1	2	-	1	1	2	1	-	-	-	-
Male	3	-	2	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	9	14	18	22	21	20	24	11	1	-	-
Male	6	5	9	12	10	14	5	5	2	1	-	-
Female	-	4	5	6	12	7	15	19	9	-	-	-
Utah												
All races, both sexes	36	47	60	52	63	114	165	159	114	49	5	-
Male	29	38	41	37	39	64	108	79	41	20	-	-
Female	7	9	19	15	24	50	57	80	73	29	5	-
White, both sexes	36	43	57	50	61	109	160	158	112	48	5	-
Male	29	35	39	35	38	62	104	78	40	20	-	-
Female	7	8	18	15	23	47	56	80	72	28	5	-
Black, both sexes	-	-	1	1	-	1	1	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	2	1	-	-	-	1	1	-	-	-	-
Male	-	2	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	1	2	4	3	-	2	1	-	-
Male	-	1	1	1	1	2	3	-	1	-	-	-
Female	-	1	-	-	1	2	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Vermont												
All races, both sexes	19	25	31	32	56	74	97	108	100	49	13	-
Male	11	21	21	23	33	42	49	56	30	12	1	-
Female	8	4	10	9	23	32	48	52	70	37	12	-
White, both sexes	19	25	31	31	55	74	97	108	100	49	13	-
Male	11	21	21	22	33	42	49	56	30	12	1	-
Female	8	4	10	9	22	32	48	52	70	37	12	-
Black, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	190	237	344	359	471	666	873	811	556	231	55	1
Male	145	176	237	250	294	391	429	356	167	40	1	-
Female	45	61	107	109	177	275	444	455	389	191	54	1
White, both sexes	130	163	250	281	366	553	741	699	475	198	46	1
Male	99	122	173	204	239	332	377	319	147	38	1	-
Female	31	41	77	77	127	221	364	380	328	160	45	1
Black, both sexes	57	68	83	73	94	102	122	104	80	32	8	-
Male	43	49	57	42	46	52	47	36	19	2	-	-
Female	14	19	26	31	48	50	75	68	61	30	8	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	6	10	5	11	11	9	8	1	1	1	-
Male	3	5	6	4	9	7	5	1	1	-	-	-
Female	-	1	4	1	2	4	4	7	-	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Washington												
All races, both sexes	169	217	306	349	450	669	927	966	735	298	77	-
Male	128	174	247	248	308	434	508	466	272	77	14	-
Female	41	43	59	101	142	235	419	500	463	221	63	-
White, both sexes	152	186	277	327	423	631	889	921	709	292	76	-
Male	114	149	225	234	291	415	485	444	263	75	14	-
Female	38	37	52	93	132	216	404	477	446	217	62	-
Black, both sexes	8	16	12	9	15	14	14	17	6	1	-	-
Male	5	15	8	6	8	6	9	7	2	-	-	-
Female	3	1	4	3	7	8	5	10	4	1	-	-
American Indian 1/, both sexes	2	7	9	5	3	7	3	5	4	1	1	-
Male	2	4	7	3	3	6	2	2	-	-	-	-
Female	-	3	2	2	-	1	1	3	4	1	1	-
Asian or Pacific Islander 2/, both sexes	7	8	8	8	9	17	21	23	16	4	-	-
Male	7	6	7	5	6	7	12	13	7	2	-	-
Female	-	2	1	3	3	10	9	10	9	2	-	-
West Virginia												
All races, both sexes	99	149	143	162	208	339	440	406	296	121	26	-
Male	79	108	115	109	141	188	216	169	83	35	6	-
Female	20	41	28	53	67	151	224	237	213	86	20	-
White, both sexes	91	147	139	159	203	332	428	397	279	117	25	-
Male	73	107	111	107	137	184	211	166	80	35	6	-
Female	18	40	28	52	66	148	217	231	199	82	19	-
Black, both sexes	8	2	4	3	5	7	12	9	17	4	1	-
Male	6	1	4	2	4	4	5	3	3	-	-	-
Female	2	1	-	1	1	3	7	6	14	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other forms of chronic ischemic heart disease (I20,I25)												
Wisconsin												
All races, both sexes	114	157	215	246	345	557	749	882	684	299	55	-
Male	90	116	156	177	227	348	433	411	232	71	8	-
Female	24	41	59	69	118	209	316	471	452	228	47	-
White, both sexes	94	134	186	228	320	534	731	861	668	296	53	-
Male	78	99	136	164	211	336	420	402	230	70	8	-
Female	16	35	50	64	109	198	311	459	438	226	45	-
Black, both sexes	17	21	23	15	19	21	17	15	11	3	2	-
Male	10	15	14	10	12	11	12	6	2	1	-	-
Female	7	6	9	5	7	10	5	9	9	2	2	-
American Indian 1/, both sexes	2	2	5	3	2	1	-	2	4	-	-	-
Male	2	2	5	3	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	4	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	4	1	1	4	1	-	-	-
Male	-	-	1	-	2	-	1	3	-	-	-	-
Female	1	-	-	-	2	1	-	1	1	-	-	-
Wyoming												
All races, both sexes	8	9	17	23	30	48	55	60	27	13	5	-
Male	7	6	12	16	16	34	32	29	13	-	-	-
Female	1	3	5	7	14	14	23	31	14	13	5	-
White, both sexes	7	8	17	23	29	48	55	60	27	13	5	-
Male	6	6	12	16	16	34	32	29	13	-	-	-
Female	1	2	5	7	13	14	23	31	14	13	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
United States												
All races, both sexes	62,799	1	-	-	-	2	14	45	137	403	1,017	2,159
Male	34,453	-	-	-	-	2	13	39	107	318	802	1,702
Female	28,346	1	-	-	-	-	1	6	30	85	215	457
White, both sexes	51,717	-	-	-	-	2	12	32	92	283	790	1,663
Male	28,155	-	-	-	-	2	11	26	71	227	649	1,345
Female	23,562	-	-	-	-	-	1	6	21	56	141	318
Black, both sexes	9,576	1	-	-	-	-	1	12	30	106	190	431
Male	5,372	-	-	-	-	-	1	12	22	79	123	299
Female	4,204	1	-	-	-	-	-	-	8	27	67	132
American Indian 1/, both sexes	344	-	-	-	-	-	-	-	6	9	10	23
Male	216	-	-	-	-	-	-	-	5	8	6	20
Female	128	-	-	-	-	-	-	-	1	1	4	3
Asian or Pacific Islander 2/, both sexes	1,162	-	-	-	-	-	1	1	9	5	27	42
Male	710	-	-	-	-	-	1	1	9	4	24	38
Female	452	-	-	-	-	-	-	-	-	1	3	4
Alabama												
All races, both sexes	610	-	-	-	-	-	-	1	2	5	9	22
Male	290	-	-	-	-	-	-	1	2	4	6	20
Female	320	-	-	-	-	-	-	-	-	1	3	2
White, both sexes	450	-	-	-	-	-	-	-	2	4	5	11
Male	211	-	-	-	-	-	-	-	2	4	4	9
Female	239	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	159	-	-	-	-	-	-	1	-	1	4	11
Male	79	-	-	-	-	-	-	1	-	-	2	11
Female	80	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Alaska												
All races, both sexes	173	-	-	-	-	-	-	-	1	-	11	15
Male	118	-	-	-	-	-	-	-	1	-	10	11
Female	55	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	147	-	-	-	-	-	-	-	1	-	9	14
Male	101	-	-	-	-	-	-	-	1	-	8	10
Female	46	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2,021	-	-	-	-	-	-	-	2	12	28	69
Male	1,128	-	-	-	-	-	-	-	2	9	23	59
Female	893	-	-	-	-	-	-	-	-	3	5	10
White, both sexes	1,883	-	-	-	-	-	-	-	1	11	25	60
Male	1,046	-	-	-	-	-	-	-	1	8	21	51
Female	837	-	-	-	-	-	-	-	-	3	4	9
Black, both sexes	59	-	-	-	-	-	-	-	-	1	2	1
Male	33	-	-	-	-	-	-	-	-	1	1	-
Female	26	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	65	-	-	-	-	-	-	-	1	-	1	7
Male	40	-	-	-	-	-	-	-	1	-	1	7
Female	25	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Arkansas												
All races, both sexes	292	-	-	-	-	-	-	-	6	2	12	18
Male	145	-	-	-	-	-	-	-	3	1	8	13
Female	147	-	-	-	-	-	-	-	3	1	4	5
White, both sexes	248	-	-	-	-	-	-	-	5	1	8	14
Male	122	-	-	-	-	-	-	-	2	1	4	10
Female	126	-	-	-	-	-	-	-	3	-	4	4
Black, both sexes	42	-	-	-	-	-	-	-	1	1	3	4
Male	21	-	-	-	-	-	-	-	1	-	3	3
Female	21	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7,810	-	-	-	-	-	-	2	9	34	90	183
Male	4,455	-	-	-	-	-	-	2	8	33	76	147
Female	3,355	-	-	-	-	-	-	-	1	1	14	36
White, both sexes	6,389	-	-	-	-	-	-	1	7	24	64	142
Male	3,616	-	-	-	-	-	-	1	6	23	59	115
Female	2,773	-	-	-	-	-	-	-	1	1	5	27
Black, both sexes	884	-	-	-	-	-	-	-	1	5	18	26
Male	527	-	-	-	-	-	-	-	1	5	10	20
Female	357	-	-	-	-	-	-	-	-	-	8	6
American Indian 1/, both sexes	33	-	-	-	-	-	-	-	-	2	1	1
Male	18	-	-	-	-	-	-	-	-	2	-	1
Female	15	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	504	-	-	-	-	-	-	1	1	3	7	14
Male	294	-	-	-	-	-	-	1	1	3	7	11
Female	210	-	-	-	-	-	-	-	-	-	-	3

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Colorado												
All races, both sexes	572	-	-	-	-	-	-	1	-	4	7	20
Male	348	-	-	-	-	-	-	1	-	4	5	15
Female	224	-	-	-	-	-	-	-	-	-	2	5
White, both sexes	533	-	-	-	-	-	-	1	-	4	7	18
Male	328	-	-	-	-	-	-	1	-	4	5	15
Female	205	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	31	-	-	-	-	-	-	-	-	-	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	825	-	-	-	-	-	-	-	-	3	7	32
Male	419	-	-	-	-	-	-	-	-	2	5	22
Female	406	-	-	-	-	-	-	-	-	1	2	10
White, both sexes	749	-	-	-	-	-	-	-	-	1	6	27
Male	381	-	-	-	-	-	-	-	-	-	5	19
Female	368	-	-	-	-	-	-	-	-	1	1	8
Black, both sexes	66	-	-	-	-	-	-	-	-	2	1	4
Male	32	-	-	-	-	-	-	-	-	2	-	2
Female	34	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Delaware												
All races, both sexes	337	-	-	-	-	-	-	1	2	6	5	9
Male	178	-	-	-	-	-	-	1	1	3	3	6
Female	159	-	-	-	-	-	-	-	1	3	2	3
White, both sexes	263	-	-	-	-	-	-	-	2	4	4	7
Male	139	-	-	-	-	-	-	-	1	2	3	5
Female	124	-	-	-	-	-	-	-	1	2	1	2
Black, both sexes	70	-	-	-	-	-	-	1	-	2	1	2
Male	36	-	-	-	-	-	-	1	-	1	-	1
Female	34	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	544	-	-	-	-	-	-	-	-	2	6	13
Male	288	-	-	-	-	-	-	-	-	2	5	10
Female	256	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	121	-	-	-	-	-	-	-	-	-	-	-
Male	60	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	415	-	-	-	-	-	-	-	-	2	6	13
Male	221	-	-	-	-	-	-	-	-	2	5	10
Female	194	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Florida												
All races, both sexes	5,073	-	-	-	-	-	-	4	8	22	76	151
Male	2,779	-	-	-	-	-	-	3	4	19	58	123
Female	2,294	-	-	-	-	-	-	1	4	3	18	28
White, both sexes	4,596	-	-	-	-	-	-	3	4	18	60	131
Male	2,496	-	-	-	-	-	-	2	2	16	49	109
Female	2,100	-	-	-	-	-	-	1	2	2	11	22
Black, both sexes	440	-	-	-	-	-	-	1	3	3	14	18
Male	260	-	-	-	-	-	-	1	1	2	7	12
Female	180	-	-	-	-	-	-	-	2	1	7	6
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	-	1	-	2	1
Male	19	-	-	-	-	-	-	-	1	-	2	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1,114	-	-	-	-	-	-	1	5	12	25	55
Male	625	-	-	-	-	-	-	1	4	12	20	45
Female	489	-	-	-	-	-	-	-	1	-	5	10
White, both sexes	757	-	-	-	-	-	-	-	2	5	17	42
Male	433	-	-	-	-	-	-	-	2	5	13	36
Female	324	-	-	-	-	-	-	-	-	-	4	6
Black, both sexes	351	-	-	-	-	-	-	1	2	7	7	13
Male	187	-	-	-	-	-	-	1	1	7	6	9
Female	164	-	-	-	-	-	-	-	1	-	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	1	-
Male	4	-	-	-	-	-	-	-	1	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Hawaii												
All races, both sexes	190	-	-	-	-	-	-	-	-	2	5	5
Male	126	-	-	-	-	-	-	-	-	2	3	5
Female	64	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	77	-	-	-	-	-	-	-	-	2	3	3
Male	52	-	-	-	-	-	-	-	-	2	2	3
Female	25	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	112	-	-	-	-	-	-	-	-	-	2	2
Male	74	-	-	-	-	-	-	-	-	-	1	2
Female	38	-	-	-	-	-	-	-	-	-	1	-
Idaho												
All races, both sexes	93	-	-	-	-	-	-	-	-	-	-	-
Male	38	-	-	-	-	-	-	-	-	-	-	-
Female	55	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	92	-	-	-	-	-	-	-	-	-	-	-
Male	38	-	-	-	-	-	-	-	-	-	-	-
Female	54	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Illinois												
All races, both sexes	1,978	-	-	-	-	-	-	2	1	3	14	101
Male	1,162	-	-	-	-	-	-	2	-	2	12	79
Female	816	-	-	-	-	-	-	-	1	1	2	22
White, both sexes	1,424	-	-	-	-	-	-	2	1	1	10	65
Male	817	-	-	-	-	-	-	2	-	-	8	54
Female	607	-	-	-	-	-	-	-	1	1	2	11
Black, both sexes	520	-	-	-	-	-	-	-	-	2	3	35
Male	323	-	-	-	-	-	-	-	-	2	3	24
Female	197	-	-	-	-	-	-	-	-	-	-	11
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	-	-	-	-	-	-	-	-	-	1	1
Male	22	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	620	-	-	-	-	-	-	2	2	9	19	29
Male	346	-	-	-	-	-	-	1	2	5	13	18
Female	274	-	-	-	-	-	-	1	-	4	6	11
White, both sexes	552	-	-	-	-	-	-	2	2	7	15	23
Male	301	-	-	-	-	-	-	1	2	4	11	15
Female	251	-	-	-	-	-	-	1	-	3	4	8
Black, both sexes	65	-	-	-	-	-	-	-	-	2	4	6
Male	44	-	-	-	-	-	-	-	-	1	2	3
Female	21	-	-	-	-	-	-	-	-	1	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Iowa												
All races, both sexes	996	-	-	-	-	-	-	-	-	8	14	22
Male	515	-	-	-	-	-	-	-	-	5	11	15
Female	481	-	-	-	-	-	-	-	-	3	3	7
White, both sexes	969	-	-	-	-	-	-	-	-	6	12	20
Male	499	-	-	-	-	-	-	-	-	4	11	13
Female	470	-	-	-	-	-	-	-	-	2	1	7
Black, both sexes	24	-	-	-	-	-	-	-	-	2	2	2
Male	14	-	-	-	-	-	-	-	-	1	-	2
Female	10	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	321	-	-	-	-	-	-	-	2	2	9	16
Male	203	-	-	-	-	-	-	-	2	1	6	13
Female	118	-	-	-	-	-	-	-	-	1	3	3
White, both sexes	298	-	-	-	-	-	-	-	2	2	8	15
Male	186	-	-	-	-	-	-	-	2	1	5	13
Female	112	-	-	-	-	-	-	-	-	1	3	2
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Kentucky												
All races, both sexes	533	-	-	-	-	-	1	2	2	4	7	14
Male	259	-	-	-	-	-	1	2	2	2	5	11
Female	274	-	-	-	-	-	-	-	-	2	2	3
White, both sexes	507	-	-	-	-	-	1	2	2	4	6	13
Male	248	-	-	-	-	-	1	2	2	2	5	11
Female	259	-	-	-	-	-	-	-	-	2	1	2
Black, both sexes	25	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,135	-	-	-	-	-	-	1	3	12	24	39
Male	599	-	-	-	-	-	-	1	2	9	17	27
Female	536	-	-	-	-	-	-	-	1	3	7	12
White, both sexes	818	-	-	-	-	-	-	1	2	6	17	22
Male	440	-	-	-	-	-	-	1	1	4	14	16
Female	378	-	-	-	-	-	-	-	1	2	3	6
Black, both sexes	315	-	-	-	-	-	-	-	1	6	7	17
Male	157	-	-	-	-	-	-	-	1	5	3	11
Female	158	-	-	-	-	-	-	-	-	1	4	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Maine												
All races, both sexes	233	-	-	-	-	1	-	1	2	3	7	11
Male	120	-	-	-	-	1	-	1	1	2	5	11
Female	113	-	-	-	-	-	-	-	1	1	2	-
White, both sexes	231	-	-	-	-	1	-	1	2	3	7	11
Male	119	-	-	-	-	1	-	1	1	2	5	11
Female	112	-	-	-	-	-	-	-	1	1	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,471	-	-	-	-	-	1	1	10	13	43	75
Male	1,219	-	-	-	-	-	1	1	8	9	32	59
Female	1,252	-	-	-	-	-	-	-	2	4	11	16
White, both sexes	1,724	-	-	-	-	-	1	1	7	6	30	47
Male	854	-	-	-	-	-	1	1	5	4	26	40
Female	870	-	-	-	-	-	-	-	2	2	4	7
Black, both sexes	706	-	-	-	-	-	-	-	3	6	12	28
Male	349	-	-	-	-	-	-	-	3	5	5	19
Female	357	-	-	-	-	-	-	-	-	1	7	9
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	21	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Massachusetts												
All races, both sexes	774	-	-	-	-	-	2	-	-	8	19	34
Male	423	-	-	-	-	-	2	-	-	6	17	26
Female	351	-	-	-	-	-	-	-	-	2	2	8
White, both sexes	725	-	-	-	-	-	1	-	-	6	19	32
Male	393	-	-	-	-	-	1	-	-	4	17	25
Female	332	-	-	-	-	-	-	-	-	2	2	7
Black, both sexes	40	-	-	-	-	-	-	-	-	2	-	2
Male	23	-	-	-	-	-	-	-	-	2	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	1	-	-	-	-	-
Male	7	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	3,826	-	-	-	-	-	2	2	11	34	70	148
Male	2,197	-	-	-	-	-	2	2	9	25	55	108
Female	1,629	-	-	-	-	-	-	-	2	9	15	40
White, both sexes	2,525	-	-	-	-	-	2	2	7	20	47	92
Male	1,450	-	-	-	-	-	2	2	7	15	39	73
Female	1,075	-	-	-	-	-	-	-	-	5	8	19
Black, both sexes	1,259	-	-	-	-	-	-	-	4	13	20	54
Male	718	-	-	-	-	-	-	-	2	9	14	34
Female	541	-	-	-	-	-	-	-	2	4	6	20
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	1	2	1
Male	10	-	-	-	-	-	-	-	-	1	1	-
Female	9	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Minnesota												
All races, both sexes	320	-	-	-	-	-	-	1	-	1	2	5
Male	176	-	-	-	-	-	-	-	-	1	1	3
Female	144	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	316	-	-	-	-	-	-	1	-	1	2	4
Male	174	-	-	-	-	-	-	-	-	1	1	2
Female	142	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	654	-	-	-	-	-	-	-	-	-	6	9
Male	266	-	-	-	-	-	-	-	-	-	4	5
Female	388	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	502	-	-	-	-	-	-	-	-	-	2	7
Male	205	-	-	-	-	-	-	-	-	-	2	4
Female	297	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	149	-	-	-	-	-	-	-	-	-	3	2
Male	60	-	-	-	-	-	-	-	-	-	2	1
Female	89	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Missouri												
All races, both sexes	629	-	-	-	-	-	-	1	-	-	3	16
Male	343	-	-	-	-	-	-	-	-	-	2	13
Female	286	-	-	-	-	-	-	1	-	-	1	3
White, both sexes	564	-	-	-	-	-	-	1	-	-	3	15
Male	310	-	-	-	-	-	-	-	-	-	2	13
Female	254	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	62	-	-	-	-	-	-	-	-	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	111	-	-	-	-	-	-	-	-	1	2	6
Male	74	-	-	-	-	-	-	-	-	1	2	5
Female	37	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	107	-	-	-	-	-	-	-	-	1	2	6
Male	71	-	-	-	-	-	-	-	-	1	2	5
Female	36	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Nebraska												
All races, both sexes	124	-	-	-	-	-	-	-	-	-	1	1
Male	44	-	-	-	-	-	-	-	-	-	1	-
Female	80	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	122	-	-	-	-	-	-	-	-	-	1	1
Male	44	-	-	-	-	-	-	-	-	-	1	-
Female	78	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	937	-	-	-	-	-	-	1	-	2	7	31
Male	594	-	-	-	-	-	-	1	-	2	7	26
Female	343	-	-	-	-	-	-	-	-	-	-	5
White, both sexes	847	-	-	-	-	-	-	1	-	2	7	24
Male	539	-	-	-	-	-	-	1	-	2	7	21
Female	308	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	54	-	-	-	-	-	-	-	-	-	-	2
Male	34	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	3
Male	8	-	-	-	-	-	-	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	-	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	2
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
New Hampshire												
All races, both sexes	322	-	-	-	-	-	-	-	1	3	6	12
Male	204	-	-	-	-	-	-	-	1	3	5	10
Female	118	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	321	-	-	-	-	-	-	-	1	3	6	12
Male	203	-	-	-	-	-	-	-	1	3	5	10
Female	118	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2,070	-	-	-	-	-	-	-	1	13	23	58
Male	1,096	-	-	-	-	-	-	-	1	12	18	50
Female	974	-	-	-	-	-	-	-	-	1	5	8
White, both sexes	1,770	-	-	-	-	-	-	-	1	10	18	43
Male	927	-	-	-	-	-	-	-	1	10	15	38
Female	843	-	-	-	-	-	-	-	-	-	3	5
Black, both sexes	253	-	-	-	-	-	-	-	-	3	4	12
Male	137	-	-	-	-	-	-	-	-	2	2	9
Female	116	-	-	-	-	-	-	-	-	1	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	47	-	-	-	-	-	-	-	-	-	1	3
Male	32	-	-	-	-	-	-	-	-	-	1	3
Female	15	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
New Mexico												
All races, both sexes	720	-	-	-	-	-	-	1	3	2	7	22
Male	445	-	-	-	-	-	-	1	3	1	7	18
Female	275	-	-	-	-	-	-	-	-	1	-	4
White, both sexes	677	-	-	-	-	-	-	1	-	2	6	20
Male	418	-	-	-	-	-	-	1	-	1	6	16
Female	259	-	-	-	-	-	-	-	-	1	-	4
Black, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	3	-	-	1
Male	16	-	-	-	-	-	-	-	3	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	4,272	-	-	-	-	1	1	5	8	30	62	146
Male	2,394	-	-	-	-	1	1	5	6	27	46	122
Female	1,878	-	-	-	-	-	-	-	2	3	16	24
White, both sexes	3,146	-	-	-	-	1	1	2	5	22	43	95
Male	1,742	-	-	-	-	1	1	2	3	21	35	83
Female	1,404	-	-	-	-	-	-	-	2	1	8	12
Black, both sexes	1,013	-	-	-	-	-	-	3	2	8	14	44
Male	574	-	-	-	-	-	-	3	2	6	8	32
Female	439	-	-	-	-	-	-	-	-	2	6	12
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	108	-	-	-	-	-	-	-	1	-	5	7
Male	74	-	-	-	-	-	-	-	1	-	3	7
Female	34	-	-	-	-	-	-	-	-	-	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
North Carolina												
All races, both sexes	1,147	-	-	-	-	-	1	1	4	6	12	31
Male	527	-	-	-	-	-	1	-	4	3	11	25
Female	620	-	-	-	-	-	-	1	-	3	1	6
White, both sexes	920	-	-	-	-	-	1	1	2	-	8	24
Male	424	-	-	-	-	-	1	-	2	-	8	19
Female	496	-	-	-	-	-	-	1	-	-	-	5
Black, both sexes	219	-	-	-	-	-	-	-	2	6	4	7
Male	100	-	-	-	-	-	-	-	2	3	3	6
Female	119	-	-	-	-	-	-	-	-	3	1	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	126	-	-	-	-	-	1	-	-	2	5	5
Male	75	-	-	-	-	-	1	-	-	2	4	4
Female	51	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	123	-	-	-	-	-	1	-	-	2	5	5
Male	73	-	-	-	-	-	1	-	-	2	4	4
Female	50	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Ohio												
All races, both sexes	2,074	-	-	-	-	-	2	2	6	21	50	79
Male	1,143	-	-	-	-	-	2	2	6	14	41	60
Female	931	-	-	-	-	-	-	-	-	7	9	19
White, both sexes	1,770	-	-	-	-	-	1	-	5	13	44	69
Male	970	-	-	-	-	-	1	-	5	8	37	53
Female	800	-	-	-	-	-	-	-	-	5	7	16
Black, both sexes	293	-	-	-	-	-	1	2	1	8	6	9
Male	168	-	-	-	-	-	1	2	1	6	4	6
Female	125	-	-	-	-	-	-	-	-	2	2	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,085	-	-	-	-	-	-	1	2	12	36	44
Male	614	-	-	-	-	-	-	1	2	11	28	34
Female	471	-	-	-	-	-	-	-	-	1	8	10
White, both sexes	948	-	-	-	-	-	-	1	1	6	31	36
Male	519	-	-	-	-	-	-	1	1	5	24	29
Female	429	-	-	-	-	-	-	-	-	1	7	7
Black, both sexes	67	-	-	-	-	-	-	-	-	2	4	3
Male	42	-	-	-	-	-	-	-	-	2	3	1
Female	25	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	69	-	-	-	-	-	-	-	1	4	1	5
Male	52	-	-	-	-	-	-	-	1	4	1	4
Female	17	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Oregon												
All races, both sexes	297	-	-	-	-	-	1	-	-	1	8	8
Male	163	-	-	-	-	-	1	-	-	1	8	6
Female	134	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	286	-	-	-	-	-	1	-	-	-	8	8
Male	154	-	-	-	-	-	1	-	-	-	8	6
Female	132	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3,806	-	-	-	-	-	-	3	4	19	55	111
Male	1,886	-	-	-	-	-	-	3	4	15	46	86
Female	1,920	-	-	-	-	-	-	-	-	4	9	25
White, both sexes	3,228	-	-	-	-	-	-	3	3	16	45	89
Male	1,550	-	-	-	-	-	-	3	3	13	38	69
Female	1,678	-	-	-	-	-	-	-	-	3	7	20
Black, both sexes	560	-	-	-	-	-	-	-	-	3	10	22
Male	325	-	-	-	-	-	-	-	-	2	8	17
Female	235	-	-	-	-	-	-	-	-	1	2	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	1	-	-	-
Male	11	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Rhode Island												
All races, both sexes	565	-	-	-	-	-	-	-	1	3	4	13
Male	272	-	-	-	-	-	-	-	-	3	2	11
Female	293	-	-	-	-	-	-	-	1	-	2	2
White, both sexes	550	-	-	-	-	-	-	-	1	3	4	12
Male	263	-	-	-	-	-	-	-	-	3	2	10
Female	287	-	-	-	-	-	-	-	1	-	2	2
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	464	-	-	-	-	-	-	-	1	5	11	17
Male	254	-	-	-	-	-	-	-	1	4	8	13
Female	210	-	-	-	-	-	-	-	-	1	3	4
White, both sexes	350	-	-	-	-	-	-	-	1	3	9	9
Male	190	-	-	-	-	-	-	-	1	2	7	7
Female	160	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	114	-	-	-	-	-	-	-	-	2	2	8
Male	64	-	-	-	-	-	-	-	-	2	1	6
Female	50	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
South Dakota												
All races, both sexes	117	-	-	-	-	-	-	-	-	-	-	-
Male	64	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	115	-	-	-	-	-	-	-	-	-	-	-
Male	62	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,779	-	-	-	-	-	-	1	10	13	36	87
Male	961	-	-	-	-	-	-	1	9	10	25	63
Female	818	-	-	-	-	-	-	-	1	3	11	24
White, both sexes	1,533	-	-	-	-	-	-	1	5	10	31	69
Male	820	-	-	-	-	-	-	1	4	8	21	49
Female	713	-	-	-	-	-	-	-	1	2	10	20
Black, both sexes	238	-	-	-	-	-	-	-	4	3	5	17
Male	134	-	-	-	-	-	-	-	4	2	4	13
Female	104	-	-	-	-	-	-	-	-	1	1	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Texas												
All races, both sexes	4,071	-	-	-	-	-	1	6	16	47	95	223
Male	2,404	-	-	-	-	-	-	6	11	34	74	184
Female	1,667	-	-	-	-	-	1	-	5	13	21	39
White, both sexes	3,350	-	-	-	-	-	1	3	10	35	72	174
Male	1,961	-	-	-	-	-	-	3	7	27	56	147
Female	1,389	-	-	-	-	-	1	-	3	8	16	27
Black, both sexes	661	-	-	-	-	-	-	3	4	12	22	45
Male	412	-	-	-	-	-	-	3	2	7	17	33
Female	249	-	-	-	-	-	-	-	2	5	5	12
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	53	-	-	-	-	-	-	-	2	-	1	3
Male	27	-	-	-	-	-	-	-	2	-	1	3
Female	26	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	83	-	-	-	-	-	-	-	-	-	-	2
Male	46	-	-	-	-	-	-	-	-	-	-	2
Female	37	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	82	-	-	-	-	-	-	-	-	-	-	2
Male	45	-	-	-	-	-	-	-	-	-	-	2
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Vermont												
All races, both sexes	249	-	-	-	-	-	-	-	1	2	2	15
Male	134	-	-	-	-	-	-	-	1	-	2	13
Female	115	-	-	-	-	-	-	-	-	2	-	2
White, both sexes	248	-	-	-	-	-	-	-	1	2	2	15
Male	133	-	-	-	-	-	-	-	1	-	2	13
Female	115	-	-	-	-	-	-	-	-	2	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,164	1	-	-	-	-	-	-	1	8	21	46
Male	590	-	-	-	-	-	-	-	1	8	18	38
Female	574	1	-	-	-	-	-	-	-	-	3	8
White, both sexes	924	-	-	-	-	-	-	-	-	7	13	35
Male	460	-	-	-	-	-	-	-	-	7	11	32
Female	464	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	212	1	-	-	-	-	-	-	-	1	6	10
Male	109	-	-	-	-	-	-	-	-	1	5	5
Female	103	1	-	-	-	-	-	-	-	-	1	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	-	1	-	2	1
Male	21	-	-	-	-	-	-	-	1	-	2	1
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Washington												
All races, both sexes	1,566	-	-	-	-	-	1	-	4	8	30	52
Male	885	-	-	-	-	-	1	-	3	5	26	37
Female	681	-	-	-	-	-	-	-	1	3	4	15
White, both sexes	1,445	-	-	-	-	-	1	-	4	6	28	46
Male	815	-	-	-	-	-	1	-	3	4	24	33
Female	630	-	-	-	-	-	-	-	1	2	4	13
Black, both sexes	55	-	-	-	-	-	-	-	-	1	-	2
Male	30	-	-	-	-	-	-	-	-	1	-	1
Female	25	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	50	-	-	-	-	-	-	-	-	1	2	3
Male	29	-	-	-	-	-	-	-	-	-	2	2
Female	21	-	-	-	-	-	-	-	-	1	-	1
West Virginia												
All races, both sexes	574	-	-	-	-	-	-	-	2	-	8	14
Male	323	-	-	-	-	-	-	-	1	-	7	12
Female	251	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	547	-	-	-	-	-	-	-	2	-	7	13
Male	309	-	-	-	-	-	-	-	1	-	6	11
Female	238	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	27	-	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerotic cardiovascular disease, so described (I25.0)												
Wisconsin												
All races, both sexes	902	-	-	-	-	-	-	1	4	3	17	25
Male	481	-	-	-	-	-	-	-	2	3	13	19
Female	421	-	-	-	-	-	-	1	2	-	4	6
White, both sexes	819	-	-	-	-	-	-	1	1	3	13	21
Male	430	-	-	-	-	-	-	-	1	3	10	16
Female	389	-	-	-	-	-	-	1	-	-	3	5
Black, both sexes	72	-	-	-	-	-	-	-	2	-	3	4
Male	45	-	-	-	-	-	-	-	1	-	3	3
Female	27	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	30	-	-	-	-	-	-	-	-	1	1	-
Male	15	-	-	-	-	-	-	-	-	1	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	1	1	-
Male	14	-	-	-	-	-	-	-	-	1	1	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
United States												
All races, both sexes	3,725	4,665	4,900	4,746	5,500	7,151	9,135	8,651	6,855	2,953	723	17
Male	2,941	3,580	3,580	3,224	3,485	3,961	4,423	3,451	2,059	656	100	10
Female	784	1,085	1,320	1,522	2,015	3,190	4,712	5,200	4,796	2,297	623	7
White, both sexes	2,757	3,493	3,707	3,693	4,333	5,886	7,834	7,694	6,150	2,665	620	11
Male	2,238	2,753	2,750	2,541	2,763	3,307	3,795	3,130	1,862	592	88	5
Female	519	740	957	1,152	1,570	2,579	4,039	4,564	4,288	2,073	532	6
Black, both sexes	857	1,023	1,047	913	1,015	1,096	1,092	818	602	247	90	5
Male	612	715	717	590	635	563	515	268	160	48	9	4
Female	245	308	330	323	380	533	577	550	442	199	81	1
American Indian 1/, both sexes	34	43	37	34	38	32	34	20	14	7	3	-
Male	24	29	28	23	17	18	21	10	4	3	-	-
Female	10	14	9	11	21	14	13	10	10	4	3	-
Asian or Pacific Islander 2/, both sexes	77	106	109	106	114	137	175	119	89	34	10	1
Male	67	83	85	70	70	73	92	43	33	13	3	1
Female	10	23	24	36	44	64	83	76	56	21	7	-
Alabama												
All races, both sexes	27	37	30	32	50	68	98	91	101	31	6	-
Male	16	31	21	22	32	39	41	25	22	7	1	-
Female	11	6	9	10	18	29	57	66	79	24	5	-
White, both sexes	18	28	21	22	31	48	72	82	78	24	4	-
Male	12	25	15	16	19	29	30	22	19	4	1	-
Female	6	3	6	6	12	19	42	60	59	20	3	-
Black, both sexes	9	8	9	10	19	20	26	9	23	7	2	-
Male	4	6	6	6	13	10	11	3	3	3	-	-
Female	5	2	3	4	6	10	15	6	20	4	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Alaska												
All races, both sexes	18	28	18	15	16	14	20	13	3	1	-	-
Male	10	20	16	11	11	11	9	8	-	-	-	-
Female	8	8	2	4	5	3	11	5	3	1	-	-
White, both sexes	16	24	15	11	12	11	19	11	3	1	-	-
Male	9	18	14	8	9	9	8	7	-	-	-	-
Female	7	6	1	3	3	2	11	4	3	1	-	-
Black, both sexes	-	1	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	2	3	2	2	4	3	1	-	-	-	-	-
Male	1	2	1	2	2	2	1	-	-	-	-	-
Female	1	1	1	-	2	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	113	151	167	164	223	235	317	266	183	74	15	2
Male	90	119	126	105	142	130	135	117	57	12	2	-
Female	23	32	41	59	81	105	182	149	126	62	13	2
White, both sexes	99	137	158	145	207	216	302	256	178	72	14	2
Male	79	110	121	90	136	121	129	109	57	11	2	-
Female	20	27	37	55	71	95	173	147	121	61	12	2
Black, both sexes	8	4	6	8	7	11	7	2	2	-	-	-
Male	7	3	3	6	4	5	2	1	-	-	-	-
Female	1	1	3	2	3	6	5	1	2	-	-	-
American Indian 1/, both sexes	4	9	3	8	9	7	6	6	2	1	1	-
Male	2	6	2	7	2	4	2	5	-	1	-	-
Female	2	3	1	1	7	3	4	1	2	-	1	-
Asian or Pacific Islander 2/, both sexes	2	1	-	3	-	1	2	2	1	1	-	-
Male	2	-	-	2	-	-	2	2	-	-	-	-
Female	-	1	-	1	-	1	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Arkansas												
All races, both sexes	21	16	21	22	19	33	23	52	31	16	-	-
Male	15	12	16	14	13	16	9	15	10	-	-	-
Female	6	4	5	8	6	17	14	37	21	16	-	-
White, both sexes	18	13	17	17	17	29	22	44	28	15	-	-
Male	12	9	15	11	12	16	9	12	9	-	-	-
Female	6	4	2	6	5	13	13	32	19	15	-	-
Black, both sexes	2	3	4	5	2	4	1	8	3	1	-	-
Male	2	3	1	3	1	-	-	3	1	-	-	-
Female	-	-	3	2	1	4	1	5	2	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	470	668	683	661	689	884	1,141	1,007	819	369	101	-
Male	403	561	498	454	445	507	561	385	281	77	17	-
Female	67	107	185	207	244	377	580	622	538	292	84	-
White, both sexes	350	519	522	533	542	736	950	868	721	325	85	-
Male	306	440	382	372	344	436	462	344	240	70	16	-
Female	44	79	140	161	198	300	488	524	481	255	69	-
Black, both sexes	88	103	116	82	100	85	99	71	48	31	11	-
Male	70	83	82	51	73	43	48	20	18	3	-	-
Female	18	20	34	31	27	42	51	51	30	28	11	-
American Indian 1/, both sexes	1	6	4	6	4	3	3	1	1	-	-	-
Male	-	4	2	5	2	-	2	-	-	-	-	-
Female	1	2	2	1	2	3	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	31	40	41	40	43	60	89	67	49	13	5	-
Male	27	34	32	26	26	28	49	21	23	4	1	-
Female	4	6	9	14	17	32	40	46	26	9	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Colorado												
All races, both sexes	44	52	45	48	60	69	78	70	44	25	5	-
Male	38	47	36	33	36	36	39	36	14	8	-	-
Female	6	5	9	15	24	33	39	34	30	17	5	-
White, both sexes	44	45	38	45	54	63	75	67	43	24	5	-
Male	38	41	30	31	34	35	37	35	14	8	-	-
Female	6	4	8	14	20	28	38	32	29	16	5	-
Black, both sexes	-	5	7	3	6	5	-	3	-	-	-	-
Male	-	5	6	2	2	1	-	1	-	-	-	-
Female	-	-	1	1	4	4	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	1	2	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	1	1	-	1	-	-	-
Connecticut												
All races, both sexes	44	51	64	64	59	92	116	121	104	53	13	2
Male	32	36	41	44	35	49	67	44	30	10	-	2
Female	12	15	23	20	24	43	49	77	74	43	13	-
White, both sexes	33	42	58	55	52	87	110	115	100	50	11	2
Male	27	31	37	37	32	46	63	43	29	10	-	2
Female	6	11	21	18	20	41	47	72	71	40	11	-
Black, both sexes	10	8	5	9	4	5	5	5	4	3	1	-
Male	4	4	3	7	2	3	3	1	1	-	-	-
Female	6	4	2	2	2	2	2	4	3	3	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	1	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	1	1	-	2	-	-	1	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Delaware												
All races, both sexes	24	25	32	25	23	40	41	44	38	20	2	-
Male	18	20	22	16	18	22	18	17	10	3	-	-
Female	6	5	10	9	5	18	23	27	28	17	2	-
White, both sexes	15	19	23	17	19	34	32	38	31	16	2	-
Male	13	16	17	10	15	19	12	16	7	3	-	-
Female	2	3	6	7	4	15	20	22	24	13	2	-
Black, both sexes	9	6	9	6	4	6	8	6	6	4	-	-
Male	5	4	5	5	3	3	5	1	2	-	-	-
Female	4	2	4	1	1	3	3	5	4	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	28	33	54	43	63	73	81	62	54	23	7	2
Male	24	25	39	27	34	36	36	23	16	8	2	1
Female	4	8	15	16	29	37	45	39	38	15	5	1
White, both sexes	2	5	9	5	10	11	23	18	23	12	2	1
Male	1	4	6	4	3	7	9	10	10	5	1	-
Female	1	1	3	1	7	4	14	8	13	7	1	1
Black, both sexes	26	28	45	37	51	61	56	43	31	10	5	1
Male	23	21	33	22	29	28	25	12	6	3	1	1
Female	3	7	12	15	22	33	31	31	25	7	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	2	1	2	1	-	1	-	-
Male	-	-	-	1	2	1	2	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Florida												
All races, both sexes	206	246	314	303	381	595	816	846	708	324	72	1
Male	176	190	250	216	249	356	412	375	243	87	17	1
Female	30	56	64	87	132	239	404	471	465	237	55	-
White, both sexes	167	200	261	266	335	530	763	810	671	309	68	-
Male	145	161	210	193	217	319	387	360	227	83	16	-
Female	22	39	51	73	118	211	376	450	444	226	52	-
Black, both sexes	37	40	49	35	43	63	48	33	36	13	3	1
Male	30	26	36	22	30	36	23	14	15	3	1	1
Female	7	14	13	13	13	27	25	19	21	10	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	6	4	1	3	2	5	1	1	2	1	-
Male	1	3	4	-	2	1	2	-	1	1	-	-
Female	1	3	-	1	1	1	3	1	-	1	1	-
Georgia												
All races, both sexes	86	84	116	92	83	123	143	133	103	44	9	-
Male	69	66	78	50	54	61	67	56	32	10	-	-
Female	17	18	38	42	29	62	76	77	71	34	9	-
White, both sexes	51	58	74	42	52	82	112	101	81	34	4	-
Male	41	46	52	24	38	45	54	43	26	8	-	-
Female	10	12	22	18	14	37	58	58	55	26	4	-
Black, both sexes	34	26	42	47	31	41	31	32	22	10	5	-
Male	27	20	26	24	16	16	13	13	6	2	-	-
Female	7	6	16	23	15	25	18	19	16	8	5	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Hawaii												
All races, both sexes	6	15	18	18	27	23	32	21	10	7	1	-
Male	6	14	14	12	22	11	16	13	3	4	1	-
Female	-	1	4	6	5	12	16	8	7	3	-	-
White, both sexes	2	8	10	4	9	10	12	7	5	2	-	-
Male	2	8	8	3	7	5	5	4	2	1	-	-
Female	-	-	2	1	2	5	7	3	3	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	7	8	13	18	13	20	14	5	5	1	-
Male	4	6	6	9	15	6	11	9	1	3	1	-
Female	-	1	2	4	3	7	9	5	4	2	-	-
Idaho												
All races, both sexes	5	3	3	4	6	9	19	18	23	3	-	-
Male	4	2	2	4	2	4	8	7	5	-	-	-
Female	1	1	1	-	4	5	11	11	18	3	-	-
White, both sexes	5	3	3	4	6	9	18	18	23	3	-	-
Male	4	2	2	4	2	4	8	7	5	-	-	-
Female	1	1	1	-	4	5	10	11	18	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Illinois												
All races, both sexes	178	212	172	163	208	233	245	204	169	63	10	-
Male	131	168	128	111	119	140	136	87	39	7	1	-
Female	47	44	44	52	89	93	109	117	130	56	9	-
White, both sexes	108	133	103	119	138	166	192	172	146	60	8	-
Male	80	109	78	79	81	102	109	74	33	7	1	-
Female	28	24	25	40	57	64	83	98	113	53	7	-
Black, both sexes	70	76	66	40	62	63	47	30	21	3	2	-
Male	51	56	47	30	34	36	23	11	6	-	-	-
Female	19	20	19	10	28	27	24	19	15	3	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	3	3	8	4	6	2	2	-	-	-
Male	-	3	3	2	4	2	4	2	-	-	-	-
Female	-	-	-	1	4	2	2	-	2	-	-	-
Indiana												
All races, both sexes	44	40	38	40	59	64	96	69	71	31	7	-
Male	31	30	24	35	43	37	51	28	19	9	-	-
Female	13	10	14	5	16	27	45	41	52	22	7	-
White, both sexes	36	34	32	30	49	59	91	67	68	31	6	-
Male	26	25	20	26	36	35	47	27	17	9	-	-
Female	10	9	12	4	13	24	44	40	51	22	6	-
Black, both sexes	7	6	6	9	10	4	5	2	3	-	1	-
Male	5	5	4	8	7	2	4	1	2	-	-	-
Female	2	1	2	1	3	2	1	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Iowa												
All races, both sexes	50	59	58	70	77	116	170	161	106	62	23	-
Male	34	44	44	45	49	61	90	66	33	14	4	-
Female	16	15	14	25	28	55	80	95	73	48	19	-
White, both sexes	47	55	55	68	76	115	164	161	106	62	22	-
Male	32	43	41	44	48	60	86	66	33	14	4	-
Female	15	12	14	24	28	55	78	95	73	48	18	-
Black, both sexes	3	3	2	2	1	1	5	-	-	-	1	-
Male	2	1	2	1	1	1	3	-	-	-	-	-
Female	1	2	-	1	-	-	2	-	-	-	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	26	28	30	19	34	30	45	38	27	12	3	-
Male	21	24	24	15	20	20	22	24	6	3	2	-
Female	5	4	6	4	14	10	23	14	21	9	1	-
White, both sexes	22	24	25	17	32	29	42	38	27	12	3	-
Male	20	20	20	13	18	19	20	24	6	3	2	-
Female	2	4	5	4	14	10	22	14	21	9	1	-
Black, both sexes	3	1	4	1	1	1	1	-	-	-	-	-
Male	1	1	4	1	1	1	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	1	-	1	-	1	-	-	-	-	-
Male	-	2	-	-	1	-	1	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	1	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Kentucky												
All races, both sexes	34	31	38	34	46	59	78	64	83	27	9	-
Male	24	21	28	18	28	28	35	28	22	3	1	-
Female	10	10	10	16	18	31	43	36	61	24	8	-
White, both sexes	33	31	32	32	44	55	78	63	78	25	8	-
Male	23	21	22	16	28	27	35	28	21	3	1	-
Female	10	10	10	16	16	28	43	35	57	22	7	-
Black, both sexes	1	-	6	1	2	4	-	1	5	2	1	-
Male	1	-	6	1	-	1	-	-	1	-	-	-
Female	-	-	-	-	2	3	-	1	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	54	68	67	78	114	144	154	163	144	58	12	-
Male	40	48	44	52	73	83	75	65	42	16	5	-
Female	14	20	23	26	41	61	79	98	102	42	7	-
White, both sexes	38	43	40	53	70	112	119	129	115	43	8	-
Male	30	34	29	36	45	67	60	56	32	12	3	-
Female	8	9	11	17	25	45	59	73	83	31	5	-
Black, both sexes	16	25	26	25	44	31	35	34	29	15	4	-
Male	10	14	14	16	28	15	15	9	10	4	2	-
Female	6	11	12	9	16	16	20	25	19	11	2	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Maine												
All races, both sexes	14	10	6	16	23	30	33	32	27	10	7	-
Male	9	9	4	13	14	13	16	13	5	3	-	-
Female	5	1	2	3	9	17	17	19	22	7	7	-
White, both sexes	13	9	6	16	23	30	33	32	27	10	7	-
Male	9	8	4	13	14	13	16	13	5	3	-	-
Female	4	1	2	3	9	17	17	19	22	7	7	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	158	185	180	171	206	263	344	376	280	127	34	4
Male	118	121	126	102	122	134	146	134	76	26	2	2
Female	40	64	54	69	84	129	198	242	204	101	32	2
White, both sexes	83	119	111	101	127	193	260	290	214	104	28	2
Male	68	84	84	61	72	97	113	112	61	23	1	1
Female	15	35	27	40	55	96	147	178	153	81	27	1
Black, both sexes	74	63	62	66	75	65	79	82	64	19	6	2
Male	49	36	38	38	50	35	31	21	15	2	1	1
Female	25	27	24	28	25	30	48	61	49	17	5	1
American Indian 1/, both sexes	-	1	1	-	2	-	-	1	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	6	4	2	5	5	3	2	4	-	-
Male	1	-	3	3	-	2	2	1	-	1	-	-
Female	-	2	3	1	2	3	3	2	2	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Massachusetts												
All races, both sexes	58	53	53	71	71	73	96	123	70	31	12	-
Male	50	42	41	48	42	43	45	40	16	5	-	-
Female	8	11	12	23	29	30	51	83	54	26	12	-
White, both sexes	53	51	51	60	64	68	88	121	70	29	12	-
Male	46	40	40	44	38	39	40	39	16	4	-	-
Female	7	11	11	16	26	29	48	82	54	25	12	-
Black, both sexes	4	2	1	11	5	4	6	2	-	1	-	-
Male	3	2	-	4	3	4	3	1	-	-	-	-
Female	1	-	1	7	2	-	3	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	2	1	2	-	-	1	-	-
Male	1	-	1	-	1	-	2	-	-	1	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Michigan												
All races, both sexes	315	429	395	299	381	482	480	390	265	105	18	-
Male	223	309	271	191	232	267	226	161	87	25	4	-
Female	92	120	124	108	149	215	254	229	178	80	14	-
White, both sexes	181	244	242	189	242	322	341	300	198	85	11	-
Male	137	182	171	111	152	183	159	127	69	19	2	-
Female	44	62	71	78	90	139	182	173	129	66	9	-
Black, both sexes	131	181	147	105	134	152	136	89	66	20	7	-
Male	84	124	94	76	77	79	65	34	18	6	2	-
Female	47	57	53	29	57	73	71	55	48	14	5	-
American Indian 1/, both sexes	2	2	1	1	4	4	-	-	1	-	-	-
Male	1	1	1	-	2	3	-	-	-	-	-	-
Female	1	1	-	1	2	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	5	4	1	4	3	1	-	-	-	-
Male	1	2	5	4	1	2	2	-	-	-	-	-
Female	-	-	-	-	-	2	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Minnesota												
All races, both sexes	9	15	14	17	23	35	47	66	46	31	8	-
Male	8	10	12	15	19	23	24	33	21	4	2	-
Female	1	5	2	2	4	12	23	33	25	27	6	-
White, both sexes	9	15	13	16	23	35	46	66	46	31	8	-
Male	8	10	11	15	19	23	24	33	21	4	2	-
Female	1	5	2	1	4	12	22	33	25	27	6	-
Black, both sexes	-	-	1	1	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	16	32	27	49	61	74	114	100	103	53	10	-
Male	12	17	15	41	29	34	53	27	19	8	2	-
Female	4	15	12	8	32	40	61	73	84	45	8	-
White, both sexes	10	22	21	37	43	56	90	83	82	45	4	-
Male	6	11	12	31	22	25	44	23	17	7	1	-
Female	4	11	9	6	21	31	46	60	65	38	3	-
Black, both sexes	6	9	6	12	17	18	24	17	21	8	6	-
Male	6	5	3	10	7	9	9	4	2	1	1	-
Female	-	4	3	2	10	9	15	13	19	7	5	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Missouri												
All races, both sexes	22	34	39	55	48	67	105	102	90	41	6	-
Male	18	25	33	37	36	41	50	40	33	15	-	-
Female	4	9	6	18	12	26	55	62	57	26	6	-
White, both sexes	19	28	35	48	42	60	91	96	85	36	5	-
Male	17	20	30	34	33	36	44	37	31	13	-	-
Female	2	8	5	14	9	24	47	59	54	23	5	-
Black, both sexes	3	6	4	5	6	7	14	5	5	5	1	-
Male	1	5	3	3	3	5	6	3	2	2	-	-
Female	2	1	1	2	3	2	8	2	3	3	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	11	5	13	12	10	9	15	13	6	6	2	-
Male	9	4	11	7	8	5	10	6	3	3	-	-
Female	2	1	2	5	2	4	5	7	3	3	2	-
White, both sexes	11	5	13	12	10	7	15	13	5	5	2	-
Male	9	4	11	7	8	3	10	6	3	2	-	-
Female	2	1	2	5	2	4	5	7	2	3	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	1	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Nebraska												
All races, both sexes	-	2	3	7	5	8	12	23	40	17	5	-
Male	-	2	2	4	3	3	6	6	13	4	-	-
Female	-	-	1	3	2	5	6	17	27	13	5	-
White, both sexes	-	2	3	6	5	8	12	23	39	17	5	-
Male	-	2	2	4	3	3	6	6	13	4	-	-
Female	-	-	1	2	2	5	6	17	26	13	5	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	72	87	120	117	88	138	124	88	43	16	3	-
Male	61	62	91	77	57	79	74	37	17	3	-	-
Female	11	25	29	40	31	59	50	51	26	13	3	-
White, both sexes	66	77	105	103	76	129	115	84	40	15	3	-
Male	55	57	80	69	49	73	70	36	16	3	-	-
Female	11	20	25	34	27	56	45	48	24	12	3	-
Black, both sexes	3	7	10	11	7	6	5	1	1	1	-	-
Male	3	4	7	6	4	4	3	1	1	-	-	-
Female	-	3	3	5	3	2	2	-	-	1	-	-
American Indian 1/, both sexes	-	2	2	-	3	-	3	-	-	-	-	-
Male	-	1	2	-	2	-	1	-	-	-	-	-
Female	-	1	-	-	1	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	3	3	2	3	1	3	2	-	-	-
Male	3	-	2	2	2	2	-	-	-	-	-	-
Female	-	1	1	1	-	1	1	3	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
New Hampshire												
All races, both sexes	27	20	26	36	21	35	40	35	39	16	5	-
Male	25	18	19	24	13	23	17	16	19	9	2	-
Female	2	2	7	12	8	12	23	19	20	7	3	-
White, both sexes	27	20	26	36	21	35	40	35	38	16	5	-
Male	25	18	19	24	13	23	17	16	18	9	2	-
Female	2	2	7	12	8	12	23	19	20	7	3	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	99	133	156	133	182	254	306	317	240	111	42	2
Male	82	104	104	97	119	145	150	119	71	18	4	2
Female	17	29	52	36	63	109	156	198	169	93	38	-
White, both sexes	75	105	118	112	152	211	272	296	222	101	34	-
Male	62	84	80	81	100	122	135	117	65	14	3	-
Female	13	21	38	31	52	89	137	179	157	87	31	-
Black, both sexes	16	21	33	17	27	36	31	20	16	9	7	1
Male	14	15	20	12	18	19	14	2	6	3	-	1
Female	2	6	13	5	9	17	17	18	10	6	7	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	7	5	4	3	7	3	1	2	1	1	1
Male	6	5	4	4	1	4	1	-	-	1	1	1
Female	2	2	1	-	2	3	2	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
New Mexico												
All races, both sexes	46	43	75	69	64	90	114	71	72	31	9	1
Male	35	37	56	47	41	56	60	38	31	9	4	1
Female	11	6	19	22	23	34	54	33	41	22	5	-
White, both sexes	40	39	70	67	61	84	110	69	68	30	9	1
Male	31	33	54	47	40	51	57	36	31	9	4	1
Female	9	6	16	20	21	33	53	33	37	21	5	-
Black, both sexes	-	2	1	-	2	3	1	-	2	-	-	-
Male	-	2	1	-	1	3	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	2	-	-	-
American Indian 1/, both sexes	6	2	4	1	1	2	3	2	1	1	-	-
Male	4	2	1	-	-	1	2	2	-	-	-	-
Female	2	-	3	1	1	1	1	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
New York												
All races, both sexes	243	331	368	386	432	541	607	539	387	151	34	-
Male	201	252	266	262	273	286	288	210	114	34	-	-
Female	42	79	102	124	159	255	319	329	273	117	34	-
White, both sexes	158	204	230	253	313	413	465	443	336	133	29	-
Male	137	168	165	175	199	219	222	179	104	28	-	-
Female	21	36	65	78	114	194	243	264	232	105	29	-
Black, both sexes	76	117	127	125	111	113	124	86	43	16	4	-
Male	55	75	93	82	68	54	57	27	7	5	-	-
Female	21	42	34	43	43	59	67	59	36	11	4	-
American Indian 1/, both sexes	-	2	1	-	1	1	-	-	-	-	-	-
Male	-	1	1	-	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	8	10	8	7	14	18	10	8	2	1	-
Male	9	8	7	5	5	12	9	4	3	1	-	-
Female	-	-	3	3	2	2	9	6	5	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
North Carolina												
All races, both sexes	41	47	57	72	100	127	194	196	177	63	18	-
Male	29	33	37	47	59	57	90	73	45	12	1	-
Female	12	14	20	25	41	70	104	123	132	51	17	-
White, both sexes	32	31	40	50	78	104	168	165	151	51	14	-
Male	25	22	28	32	47	48	77	66	40	8	1	-
Female	7	9	12	18	31	56	91	99	111	43	13	-
Black, both sexes	9	13	17	22	22	22	25	30	25	11	4	-
Male	4	9	9	15	12	9	13	7	4	4	-	-
Female	5	4	8	7	10	13	12	23	21	7	4	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	3	-	-	-	-	-	-	1	-	-	-
Male	-	2	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	8	8	6	7	6	16	21	13	18	8	2	-
Male	7	5	3	6	5	9	11	9	7	2	-	-
Female	1	3	3	1	1	7	10	4	11	6	2	-
White, both sexes	7	8	6	6	6	16	20	13	18	8	2	-
Male	6	5	3	5	5	9	11	9	7	2	-	-
Female	1	3	3	1	1	7	9	4	11	6	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Ohio												
All races, both sexes	135	143	137	184	145	219	306	309	233	81	22	-
Male	105	107	101	136	100	117	142	113	74	20	3	-
Female	30	36	36	48	45	102	164	196	159	61	19	-
White, both sexes	106	116	102	148	123	186	264	284	217	75	17	-
Male	87	89	78	114	84	102	117	105	68	19	3	-
Female	19	27	24	34	39	84	147	179	149	56	14	-
Black, both sexes	29	26	34	33	22	33	39	24	15	6	5	-
Male	18	17	23	21	16	15	23	8	6	1	-	-
Female	11	9	11	12	6	18	16	16	9	5	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	3	-	-	2	-	1	-	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	1	2	-	-	1	-	1	-	-	-
Oklahoma												
All races, both sexes	67	91	94	84	84	117	153	133	115	42	9	1
Male	51	66	72	56	52	64	78	57	36	6	-	-
Female	16	25	22	28	32	53	75	76	79	36	9	1
White, both sexes	51	69	76	67	74	108	143	126	109	40	9	1
Male	38	55	59	44	44	58	71	53	32	5	-	-
Female	13	14	17	23	30	50	72	73	77	35	9	1
Black, both sexes	6	15	8	10	5	4	1	7	2	-	-	-
Male	4	6	4	9	4	4	-	4	1	-	-	-
Female	2	9	4	1	1	-	1	3	1	-	-	-
American Indian 1/, both sexes	10	7	9	7	5	5	9	-	4	2	-	-
Male	9	5	8	3	4	2	7	-	3	1	-	-
Female	1	2	1	4	1	3	2	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Oregon												
All races, both sexes	8	17	11	16	13	24	55	56	51	24	4	-
Male	7	13	10	8	9	16	34	26	16	6	2	-
Female	1	4	1	8	4	8	21	30	35	18	2	-
White, both sexes	7	16	8	16	12	24	54	54	50	24	4	-
Male	6	12	7	8	8	16	33	25	16	6	2	-
Female	1	4	1	8	4	8	21	29	34	18	2	-
Black, both sexes	1	1	2	-	1	-	-	1	1	-	-	-
Male	1	1	2	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	167	225	215	221	315	450	624	635	501	209	52	-
Male	133	173	152	138	204	239	296	217	131	40	9	-
Female	34	52	63	83	111	211	328	418	370	169	43	-
White, both sexes	122	168	164	160	245	366	551	582	472	194	48	-
Male	99	132	114	105	151	191	258	200	127	38	9	-
Female	23	36	50	55	94	175	293	382	345	156	39	-
Black, both sexes	42	54	48	60	68	83	72	52	27	15	4	-
Male	33	38	36	32	51	47	38	17	4	2	-	-
Female	9	16	12	28	17	36	34	35	23	13	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	3	1	2	1	1	-	2	-	-	-
Male	1	3	2	1	2	1	-	-	-	-	-	-
Female	2	-	1	-	-	-	1	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Rhode Island												
All races, both sexes	18	28	30	23	41	55	99	107	98	39	6	-
Male	16	24	23	16	28	31	45	38	24	10	1	-
Female	2	4	7	7	13	24	54	69	74	29	5	-
White, both sexes	17	27	28	20	38	53	99	106	98	38	6	-
Male	15	23	22	13	28	29	45	38	24	10	1	-
Female	2	4	6	7	10	24	54	68	74	28	5	-
Black, both sexes	1	1	2	3	1	2	-	1	-	-	-	-
Male	1	1	1	3	-	2	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
South Carolina												
All races, both sexes	24	31	43	28	46	39	84	59	47	20	8	1
Male	17	24	31	20	27	20	40	30	14	3	1	1
Female	7	7	12	8	19	19	44	29	33	17	7	-
White, both sexes	16	22	35	20	34	25	71	46	36	17	5	1
Male	12	17	25	15	19	13	36	23	9	2	1	1
Female	4	5	10	5	15	12	35	23	27	15	4	-
Black, both sexes	8	9	8	8	12	14	13	13	11	3	3	-
Male	5	7	6	5	8	7	4	7	5	1	-	-
Female	3	2	2	3	4	7	9	6	6	2	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
South Dakota												
All races, both sexes	3	2	5	2	6	8	20	22	33	13	3	-
Male	3	2	4	2	3	5	14	12	12	7	-	-
Female	-	-	1	-	3	3	6	10	21	6	3	-
White, both sexes	3	2	4	2	6	8	20	21	33	13	3	-
Male	3	2	3	2	3	5	14	11	12	7	-	-
Female	-	-	1	-	3	3	6	10	21	6	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	105	140	132	127	161	184	264	246	177	80	16	-
Male	80	102	91	86	102	97	137	96	47	15	-	-
Female	25	38	41	41	59	87	127	150	130	65	16	-
White, both sexes	86	111	106	108	144	155	236	221	163	74	13	-
Male	65	80	75	74	91	84	125	86	44	13	-	-
Female	21	31	31	34	53	71	111	135	119	61	13	-
Black, both sexes	19	26	25	18	17	28	28	25	14	6	3	-
Male	15	20	15	11	11	12	12	10	3	2	-	-
Female	4	6	10	7	6	16	16	15	11	4	3	-
American Indian 1/, both sexes	-	1	1	1	-	-	-	-	-	-	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Texas												
All races, both sexes	306	394	413	323	364	407	507	448	348	136	37	-
Male	245	309	310	225	224	214	242	184	103	33	6	-
Female	61	85	103	98	140	193	265	264	245	103	31	-
White, both sexes	229	304	331	262	287	337	425	405	319	121	35	-
Male	188	241	248	186	181	177	203	167	93	32	5	-
Female	41	63	83	76	106	160	222	238	226	89	30	-
Black, both sexes	71	86	76	57	66	62	77	39	25	14	2	-
Male	51	65	57	37	41	33	39	17	9	-	1	-
Female	20	21	19	20	25	29	38	22	16	14	1	-
American Indian 1/, both sexes	3	-	-	-	1	1	1	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	4	6	4	10	7	4	4	4	1	-	-
Male	3	3	5	2	2	4	-	-	1	1	-	-
Female	-	1	1	2	8	3	4	4	3	-	-	-
Utah												
All races, both sexes	1	6	6	4	6	7	13	24	7	7	-	-
Male	1	5	4	4	4	3	8	11	1	3	-	-
Female	-	1	2	-	2	4	5	13	6	4	-	-
White, both sexes	1	6	6	4	5	7	13	24	7	7	-	-
Male	1	5	4	4	3	3	8	11	1	3	-	-
Female	-	1	2	-	2	4	5	13	6	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Vermont												
All races, both sexes	12	17	15	16	22	25	31	33	39	15	4	-
Male	9	13	6	14	15	18	13	16	9	4	1	-
Female	3	4	9	2	7	7	18	17	30	11	3	-
White, both sexes	12	17	15	15	22	25	31	33	39	15	4	-
Male	9	13	6	13	15	18	13	16	9	4	1	-
Female	3	4	9	2	7	7	18	17	30	11	3	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	69	80	76	70	99	127	174	188	125	61	17	1
Male	54	54	56	47	62	68	77	73	24	10	-	-
Female	15	26	20	23	37	59	97	115	101	51	17	1
White, both sexes	49	57	53	59	75	100	147	158	104	51	15	1
Male	37	38	37	38	48	52	66	65	20	9	-	-
Female	12	19	16	21	27	48	81	93	84	42	15	1
Black, both sexes	18	19	18	11	21	25	23	28	21	9	1	-
Male	15	12	15	9	11	14	9	8	4	1	-	-
Female	3	7	3	2	10	11	14	20	17	8	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	5	-	3	2	4	2	-	1	1	-
Male	2	4	4	-	3	2	2	-	-	-	-	-
Female	-	-	1	-	-	-	2	2	-	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Washington												
All races, both sexes	100	109	124	116	132	168	209	231	184	76	22	-
Male	76	88	105	85	88	102	87	105	53	21	3	-
Female	24	21	19	31	44	66	122	126	131	55	19	-
White, both sexes	91	89	111	106	124	152	202	214	175	75	21	-
Male	70	72	94	79	84	95	84	99	49	21	3	-
Female	21	17	17	27	40	57	118	115	126	54	18	-
Black, both sexes	5	11	5	4	5	7	4	8	3	-	-	-
Male	2	11	4	2	2	2	2	2	1	-	-	-
Female	3	-	1	2	3	5	2	6	2	-	-	-
American Indian 1/, both sexes	-	3	3	2	-	3	1	2	-	-	1	-
Male	-	1	3	1	-	3	1	1	-	-	-	-
Female	-	2	-	1	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	4	6	5	4	3	6	2	7	6	1	-	-
Male	4	4	4	3	2	2	-	3	3	-	-	-
Female	-	2	1	1	1	4	2	4	3	1	-	-
West Virginia												
All races, both sexes	42	45	29	49	39	59	87	85	62	46	7	-
Male	35	32	26	38	28	37	46	34	15	12	-	-
Female	7	13	3	11	11	22	41	51	47	34	7	-
White, both sexes	38	43	28	48	39	58	80	83	59	43	6	-
Male	32	31	25	37	28	36	42	33	15	12	-	-
Female	6	12	3	11	11	22	38	50	44	31	6	-
Black, both sexes	4	2	1	1	-	1	7	2	3	3	1	-
Male	3	1	1	1	-	1	4	1	-	-	-	-
Female	1	1	-	-	-	-	3	1	3	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerotic cardiovascular disease, so described (I25.0)												
Wisconsin												
All races, both sexes	49	56	61	69	68	111	136	142	111	43	6	-
Male	39	40	45	45	41	63	67	57	39	8	-	-
Female	10	16	16	24	27	48	69	85	72	35	6	-
White, both sexes	40	46	51	59	63	104	128	137	105	41	6	-
Male	35	32	38	37	37	58	61	55	39	8	-	-
Female	5	14	13	22	26	46	67	82	66	33	6	-
Black, both sexes	8	9	9	9	4	6	8	5	3	2	-	-
Male	3	7	6	7	3	4	6	2	-	-	-	-
Female	5	2	3	2	1	2	2	3	3	2	-	-
American Indian 1/, both sexes	1	1	1	1	-	1	-	-	2	-	-	-
Male	1	1	1	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Wyoming												
All races, both sexes	2	-	3	2	1	5	8	6	-	1	-	-
Male	1	-	2	2	1	2	4	1	-	-	-	-
Female	1	-	1	-	-	3	4	5	-	1	-	-
White, both sexes	1	-	3	2	1	5	8	6	-	1	-	-
Male	-	-	2	2	1	2	4	1	-	-	-	-
Female	1	-	1	-	-	3	4	5	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
United States												
All races, both sexes	228,319	1	4	2	4	12	52	108	321	747	1,877	3,475
Male	115,760	1	2	2	3	10	40	88	236	567	1,417	2,629
Female	112,559	-	2	-	1	2	12	20	85	180	460	846
White, both sexes	203,416	1	4	2	4	8	31	81	248	555	1,477	2,702
Male	103,498	1	2	2	3	7	25	67	182	425	1,144	2,075
Female	99,918	-	2	-	1	1	6	14	66	130	333	627
Black, both sexes	20,357	-	-	-	-	4	15	20	59	167	336	665
Male	9,738	-	-	-	-	3	10	16	41	122	217	470
Female	10,619	-	-	-	-	1	5	4	18	45	119	195
American Indian 1/, both sexes	811	-	-	-	-	-	1	-	4	6	13	35
Male	447	-	-	-	-	-	-	-	4	3	12	25
Female	364	-	-	-	-	-	1	-	-	3	1	10
Asian or Pacific Islander 2/, both sexes	3,735	-	-	-	-	-	5	7	10	19	51	73
Male	2,077	-	-	-	-	-	5	5	9	17	44	59
Female	1,658	-	-	-	-	-	-	2	1	2	7	14
Alabama												
All races, both sexes	2,592	-	-	-	1	-	1	-	3	8	17	42
Male	1,321	-	-	-	1	-	1	-	1	7	11	33
Female	1,271	-	-	-	-	-	-	-	2	1	6	9
White, both sexes	2,152	-	-	-	1	-	1	-	1	3	9	27
Male	1,125	-	-	-	1	-	1	-	1	3	6	21
Female	1,027	-	-	-	-	-	-	-	-	-	3	6
Black, both sexes	429	-	-	-	-	-	-	-	2	5	8	15
Male	193	-	-	-	-	-	-	-	-	4	5	12
Female	236	-	-	-	-	-	-	-	2	1	3	3
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Alaska												
All races, both sexes	120	-	-	-	-	-	1	-	-	1	-	2
Male	83	-	-	-	-	-	1	-	-	1	-	1
Female	37	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	92	-	-	-	-	-	-	-	-	-	-	1
Male	65	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	-	1	-	1
Male	15	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	4,070	-	-	-	-	-	1	3	6	15	32	55
Male	2,301	-	-	-	-	-	-	3	6	10	21	45
Female	1,769	-	-	-	-	-	1	-	-	5	11	10
White, both sexes	3,917	-	-	-	-	-	-	3	5	13	27	49
Male	2,221	-	-	-	-	-	-	3	5	10	19	41
Female	1,696	-	-	-	-	-	-	-	-	3	8	8
Black, both sexes	79	-	-	-	-	-	-	-	-	-	5	2
Male	39	-	-	-	-	-	-	-	-	-	2	2
Female	40	-	-	-	-	-	-	-	-	-	3	-
American Indian 1/, both sexes	56	-	-	-	-	-	1	-	1	2	-	4
Male	30	-	-	-	-	-	-	-	1	-	-	2
Female	26	-	-	-	-	-	1	-	-	2	-	2
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Arkansas												
All races, both sexes	1,884	-	-	-	-	1	-	1	1	3	9	14
Male	920	-	-	-	-	1	-	1	-	1	6	10
Female	964	-	-	-	-	-	-	-	1	2	3	4
White, both sexes	1,650	-	-	-	-	1	-	1	-	2	6	9
Male	822	-	-	-	-	1	-	1	-	1	5	6
Female	828	-	-	-	-	-	-	-	-	1	1	3
Black, both sexes	231	-	-	-	-	-	-	-	1	1	3	5
Male	97	-	-	-	-	-	-	-	-	-	1	4
Female	134	-	-	-	-	-	-	-	1	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	25,926	-	1	1	1	1	3	15	33	83	205	383
Male	13,197	-	1	1	1	1	3	10	22	69	161	294
Female	12,729	-	-	-	-	-	-	5	11	14	44	89
White, both sexes	22,400	-	1	1	1	-	1	9	27	66	158	290
Male	11,332	-	1	1	1	-	1	7	16	56	129	222
Female	11,068	-	-	-	-	-	-	2	11	10	29	68
Black, both sexes	1,740	-	-	-	-	1	1	4	2	13	25	62
Male	870	-	-	-	-	1	1	2	2	9	14	48
Female	870	-	-	-	-	-	-	2	-	4	11	14
American Indian 1/, both sexes	102	-	-	-	-	-	-	-	-	-	-	3
Male	55	-	-	-	-	-	-	-	-	-	-	3
Female	47	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1,684	-	-	-	-	-	1	2	4	4	22	28
Male	940	-	-	-	-	-	1	1	4	4	18	21
Female	744	-	-	-	-	-	-	1	-	-	4	7

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Colorado												
All races, both sexes	2,162	-	-	-	-	-	1	1	2	13	19	41
Male	1,212	-	-	-	-	-	1	1	1	11	18	33
Female	950	-	-	-	-	-	-	-	1	2	1	8
White, both sexes	2,078	-	-	-	-	-	1	1	2	13	18	37
Male	1,172	-	-	-	-	-	1	1	1	11	17	30
Female	906	-	-	-	-	-	-	-	1	2	1	7
Black, both sexes	62	-	-	-	-	-	-	-	-	-	-	2
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	38	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2,776	-	-	-	-	-	1	3	5	10	19	31
Male	1,345	-	-	-	-	-	1	3	4	10	16	24
Female	1,431	-	-	-	-	-	-	-	1	-	3	7
White, both sexes	2,598	-	-	-	-	-	1	3	4	8	14	26
Male	1,250	-	-	-	-	-	1	3	3	8	13	20
Female	1,348	-	-	-	-	-	-	-	1	-	1	6
Black, both sexes	156	-	-	-	-	-	-	-	-	2	5	5
Male	84	-	-	-	-	-	-	-	-	2	3	4
Female	72	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Delaware												
All races, both sexes	690	-	-	-	-	-	-	-	-	1	7	13
Male	352	-	-	-	-	-	-	-	-	1	6	9
Female	338	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	590	-	-	-	-	-	-	-	-	-	5	10
Male	319	-	-	-	-	-	-	-	-	-	5	7
Female	271	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	96	-	-	-	-	-	-	-	-	1	2	3
Male	31	-	-	-	-	-	-	-	-	1	1	2
Female	65	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	320	-	-	-	-	-	-	1	-	1	4	4
Male	151	-	-	-	-	-	-	1	-	1	3	2
Female	169	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	60	-	-	-	-	-	-	-	-	-	-	-
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	253	-	-	-	-	-	-	-	-	1	4	4
Male	116	-	-	-	-	-	-	-	-	1	3	2
Female	137	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Florida												
All races, both sexes	19,906	-	-	-	-	-	5	7	15	56	159	256
Male	10,619	-	-	-	-	-	4	6	14	38	121	194
Female	9,287	-	-	-	-	-	1	1	1	18	38	62
White, both sexes	18,273	-	-	-	-	-	2	6	11	35	131	213
Male	9,807	-	-	-	-	-	2	5	11	25	103	165
Female	8,466	-	-	-	-	-	-	1	-	10	28	48
Black, both sexes	1,551	-	-	-	-	-	3	1	4	21	28	41
Male	770	-	-	-	-	-	2	1	3	13	18	28
Female	781	-	-	-	-	-	1	-	1	8	10	13
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	65	-	-	-	-	-	-	-	-	-	-	2
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	35	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	4,215	-	-	-	-	-	1	9	13	30	65	124
Male	2,310	-	-	-	-	-	-	6	10	24	50	95
Female	1,905	-	-	-	-	-	1	3	3	6	15	29
White, both sexes	3,342	-	-	-	-	-	1	4	9	22	43	81
Male	1,876	-	-	-	-	-	-	3	6	16	32	63
Female	1,466	-	-	-	-	-	1	1	3	6	11	18
Black, both sexes	847	-	-	-	-	-	-	5	3	8	20	43
Male	422	-	-	-	-	-	-	3	3	8	16	32
Female	425	-	-	-	-	-	-	2	-	-	4	11
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	1	-	2	-
Male	10	-	-	-	-	-	-	-	1	-	2	-
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Hawaii												
All races, both sexes	652	-	-	-	-	-	-	-	1	4	8	17
Male	404	-	-	-	-	-	-	-	1	3	5	14
Female	248	-	-	-	-	-	-	-	-	1	3	3
White, both sexes	194	-	-	-	-	-	-	-	-	1	1	5
Male	126	-	-	-	-	-	-	-	-	1	-	5
Female	68	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	2
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	451	-	-	-	-	-	-	-	1	3	7	10
Male	271	-	-	-	-	-	-	-	1	2	5	7
Female	180	-	-	-	-	-	-	-	-	1	2	3
Idaho												
All races, both sexes	717	-	-	-	-	-	-	-	1	5	9	6
Male	415	-	-	-	-	-	-	-	1	3	6	4
Female	302	-	-	-	-	-	-	-	-	2	3	2
White, both sexes	705	-	-	-	-	-	-	-	1	5	9	6
Male	406	-	-	-	-	-	-	-	1	3	6	4
Female	299	-	-	-	-	-	-	-	-	2	3	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Illinois												
All races, both sexes	9,412	1	-	-	-	1	3	9	12	42	110	183
Male	4,653	1	-	-	-	1	1	8	11	36	86	137
Female	4,759	-	-	-	-	-	2	1	1	6	24	46
White, both sexes	8,168	1	-	-	-	1	2	7	10	28	82	133
Male	4,053	1	-	-	-	1	-	6	9	25	68	104
Female	4,115	-	-	-	-	-	2	1	1	3	14	29
Black, both sexes	1,145	-	-	-	-	-	1	2	2	14	26	50
Male	553	-	-	-	-	-	1	2	2	11	16	33
Female	592	-	-	-	-	-	-	-	-	3	10	17
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	93	-	-	-	-	-	-	-	-	-	2	-
Male	44	-	-	-	-	-	-	-	-	-	2	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	5,113	-	-	-	-	1	4	2	21	17	42	97
Male	2,639	-	-	-	-	1	3	2	16	12	27	68
Female	2,474	-	-	-	-	-	1	-	5	5	15	29
White, both sexes	4,745	-	-	-	-	1	3	2	17	15	36	80
Male	2,464	-	-	-	-	1	3	2	12	10	24	58
Female	2,281	-	-	-	-	-	-	-	5	5	12	22
Black, both sexes	348	-	-	-	-	-	1	-	4	2	6	17
Male	165	-	-	-	-	-	-	-	4	2	3	10
Female	183	-	-	-	-	-	1	-	-	-	3	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Iowa												
All races, both sexes	2,743	-	-	-	-	1	-	1	2	6	15	42
Male	1,379	-	-	-	-	1	-	1	2	4	6	31
Female	1,364	-	-	-	-	-	-	-	-	2	9	11
White, both sexes	2,716	-	-	-	-	1	-	1	2	6	13	40
Male	1,363	-	-	-	-	1	-	1	2	4	6	30
Female	1,353	-	-	-	-	-	-	-	-	2	7	10
Black, both sexes	20	-	-	-	-	-	-	-	-	-	2	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2,194	-	-	-	-	-	1	1	3	6	20	43
Male	1,174	-	-	-	-	-	1	-	3	5	12	34
Female	1,020	-	-	-	-	-	-	1	-	1	8	9
White, both sexes	2,075	-	-	-	-	-	1	1	3	5	19	36
Male	1,105	-	-	-	-	-	1	-	3	4	11	28
Female	970	-	-	-	-	-	-	1	-	1	8	8
Black, both sexes	93	-	-	-	-	-	-	-	-	1	1	4
Male	55	-	-	-	-	-	-	-	-	1	1	4
Female	38	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	-	-	3
Male	12	-	-	-	-	-	-	-	-	-	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Kentucky												
All races, both sexes	2,898	-	1	-	-	-	1	5	9	14	39	68
Male	1,438	-	-	-	-	-	1	5	7	10	28	53
Female	1,460	-	1	-	-	-	-	-	2	4	11	15
White, both sexes	2,762	-	1	-	-	-	1	5	8	11	37	62
Male	1,375	-	-	-	-	-	1	5	6	9	26	49
Female	1,387	-	1	-	-	-	-	-	2	2	11	13
Black, both sexes	130	-	-	-	-	-	-	-	-	3	2	5
Male	58	-	-	-	-	-	-	-	-	1	2	3
Female	72	-	-	-	-	-	-	-	-	2	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2,482	-	-	-	1	-	-	-	8	14	26	46
Male	1,340	-	-	-	-	-	-	-	4	12	18	34
Female	1,142	-	-	-	1	-	-	-	4	2	8	12
White, both sexes	1,922	-	-	-	1	-	-	-	6	9	12	29
Male	1,043	-	-	-	-	-	-	-	3	8	9	21
Female	879	-	-	-	1	-	-	-	3	1	3	8
Black, both sexes	542	-	-	-	-	-	-	-	2	5	14	16
Male	286	-	-	-	-	-	-	-	1	4	9	12
Female	256	-	-	-	-	-	-	-	1	1	5	4
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Maine												
All races, both sexes	1,015	-	-	-	-	-	-	-	1	2	8	16
Male	552	-	-	-	-	-	-	-	1	2	7	13
Female	463	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	1,012	-	-	-	-	-	-	-	1	2	8	15
Male	549	-	-	-	-	-	-	-	1	2	7	12
Female	463	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,962	-	-	-	-	-	-	2	1	4	7	25
Male	1,513	-	-	-	-	-	-	-	-	3	4	20
Female	1,449	-	-	-	-	-	-	2	1	1	3	5
White, both sexes	2,346	-	-	-	-	-	-	2	1	3	4	14
Male	1,205	-	-	-	-	-	-	-	-	2	2	11
Female	1,141	-	-	-	-	-	-	2	1	1	2	3
Black, both sexes	575	-	-	-	-	-	-	-	-	1	3	11
Male	284	-	-	-	-	-	-	-	-	1	2	9
Female	291	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Massachusetts												
All races, both sexes	4,335	-	-	-	-	-	1	1	11	11	30	53
Male	2,176	-	-	-	-	-	1	1	4	9	22	42
Female	2,159	-	-	-	-	-	-	-	7	2	8	11
White, both sexes	4,170	-	-	-	-	-	1	1	9	10	25	46
Male	2,099	-	-	-	-	-	1	1	4	8	19	39
Female	2,071	-	-	-	-	-	-	-	5	2	6	7
Black, both sexes	135	-	-	-	-	-	-	-	2	-	5	7
Male	64	-	-	-	-	-	-	-	-	-	3	3
Female	71	-	-	-	-	-	-	-	2	-	2	4
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	1	-	-
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	8,541	-	-	-	-	-	2	1	13	21	48	97
Male	4,203	-	-	-	-	-	2	-	10	13	32	70
Female	4,338	-	-	-	-	-	-	1	3	8	16	27
White, both sexes	7,715	-	-	-	-	-	1	1	11	13	40	79
Male	3,803	-	-	-	-	-	1	-	9	9	28	59
Female	3,912	-	-	-	-	-	-	1	2	4	12	20
Black, both sexes	769	-	-	-	-	-	1	-	2	6	7	17
Male	361	-	-	-	-	-	1	-	1	3	3	11
Female	408	-	-	-	-	-	-	-	1	3	4	6
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	-	1	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	34	-	-	-	-	-	-	-	-	1	-	-
Male	26	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Minnesota												
All races, both sexes	2,939	-	-	-	-	-	4	2	7	16	44	74
Male	1,719	-	-	-	-	-	2	2	5	13	40	57
Female	1,220	-	-	-	-	-	2	-	2	3	4	17
White, both sexes	2,854	-	-	-	-	-	4	2	5	13	38	63
Male	1,662	-	-	-	-	-	2	2	4	11	35	48
Female	1,192	-	-	-	-	-	2	-	1	2	3	15
Black, both sexes	49	-	-	-	-	-	-	-	2	2	3	11
Male	34	-	-	-	-	-	-	-	1	1	2	9
Female	15	-	-	-	-	-	-	-	1	1	1	2
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	-	2	-
Male	10	-	-	-	-	-	-	-	-	-	2	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	1	1	-
Male	13	-	-	-	-	-	-	-	-	1	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2,003	-	-	-	-	1	2	3	14	29	51	73
Male	1,051	-	-	-	-	1	2	3	9	22	35	52
Female	952	-	-	-	-	-	-	-	5	7	16	21
White, both sexes	1,491	-	-	-	-	-	1	2	4	14	28	39
Male	769	-	-	-	-	-	1	2	2	11	21	26
Female	722	-	-	-	-	-	-	-	2	3	7	13
Black, both sexes	504	-	-	-	-	1	1	1	10	15	23	34
Male	280	-	-	-	-	1	1	1	7	11	14	26
Female	224	-	-	-	-	-	-	-	3	4	9	8
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Missouri												
All races, both sexes	5,174	-	1	-	-	-	1	1	11	32	64	109
Male	2,550	-	-	-	-	-	1	1	8	26	52	87
Female	2,624	-	1	-	-	-	-	-	3	6	12	22
White, both sexes	4,693	-	1	-	-	-	1	1	9	25	52	88
Male	2,300	-	-	-	-	-	1	1	6	20	43	68
Female	2,393	-	1	-	-	-	-	-	3	5	9	20
Black, both sexes	465	-	-	-	-	-	-	-	2	7	12	20
Male	244	-	-	-	-	-	-	-	2	6	9	18
Female	221	-	-	-	-	-	-	-	-	1	3	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	572	-	-	-	-	-	-	-	2	1	4	8
Male	345	-	-	-	-	-	-	-	1	1	3	6
Female	227	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	551	-	-	-	-	-	-	-	2	-	4	7
Male	331	-	-	-	-	-	-	-	1	-	3	5
Female	220	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Nebraska												
All races, both sexes	1,311	-	-	-	-	-	-	1	3	7	20	24
Male	727	-	-	-	-	-	-	1	2	5	17	20
Female	584	-	-	-	-	-	-	-	1	2	3	4
White, both sexes	1,259	-	-	-	-	-	-	1	2	5	18	21
Male	697	-	-	-	-	-	-	1	1	3	15	19
Female	562	-	-	-	-	-	-	-	1	2	3	2
Black, both sexes	40	-	-	-	-	-	-	-	1	2	2	3
Male	24	-	-	-	-	-	-	-	1	2	2	1
Female	16	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,055	-	-	-	-	-	-	-	3	5	6	26
Male	617	-	-	-	-	-	-	-	3	4	4	19
Female	438	-	-	-	-	-	-	-	-	1	2	7
White, both sexes	959	-	-	-	-	-	-	-	1	3	6	23
Male	570	-	-	-	-	-	-	-	1	2	4	18
Female	389	-	-	-	-	-	-	-	-	1	2	5
Black, both sexes	60	-	-	-	-	-	-	-	1	1	-	2
Male	28	-	-	-	-	-	-	-	1	1	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	-	-	-	1	-	1
Male	16	-	-	-	-	-	-	-	-	1	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
New Hampshire												
All races, both sexes	988	-	-	-	-	-	-	1	1	1	7	13
Male	482	-	-	-	-	-	-	1	1	1	7	12
Female	506	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	982	-	-	-	-	-	-	-	1	1	7	13
Male	477	-	-	-	-	-	-	-	1	1	7	12
Female	505	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	7,872	-	-	-	-	-	-	2	8	16	42	67
Male	3,641	-	-	-	-	-	-	2	7	10	34	51
Female	4,231	-	-	-	-	-	-	-	1	6	8	16
White, both sexes	7,099	-	-	-	-	-	-	-	6	12	34	54
Male	3,306	-	-	-	-	-	-	-	6	7	28	41
Female	3,793	-	-	-	-	-	-	-	-	5	6	13
Black, both sexes	655	-	-	-	-	-	-	1	2	4	5	10
Male	281	-	-	-	-	-	-	1	1	3	3	8
Female	374	-	-	-	-	-	-	-	1	1	2	2
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	105	-	-	-	-	-	-	1	-	-	3	2
Male	49	-	-	-	-	-	-	1	-	-	3	2
Female	56	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
New Mexico												
All races, both sexes	1,032	-	-	-	-	-	-	-	-	2	4	7
Male	530	-	-	-	-	-	-	-	-	2	3	6
Female	502	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	968	-	-	-	-	-	-	-	-	2	4	4
Male	492	-	-	-	-	-	-	-	-	2	3	4
Female	476	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	41	-	-	-	-	-	-	-	-	-	-	2
Male	25	-	-	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	26,756	-	-	-	-	1	1	6	11	34	106	221
Male	11,574	-	-	-	-	1	-	4	7	24	77	177
Female	15,182	-	-	-	-	-	1	2	4	10	29	44
White, both sexes	22,990	-	-	-	-	1	1	4	9	26	75	160
Male	9,988	-	-	-	-	1	-	3	6	17	57	137
Female	13,002	-	-	-	-	-	1	1	3	9	18	23
Black, both sexes	3,138	-	-	-	-	-	-	1	1	7	25	50
Male	1,264	-	-	-	-	-	-	1	1	6	15	31
Female	1,874	-	-	-	-	-	-	-	-	1	10	19
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	606	-	-	-	-	-	-	1	1	1	5	11
Male	309	-	-	-	-	-	-	-	-	1	4	9
Female	297	-	-	-	-	-	-	1	1	-	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
North Carolina												
All races, both sexes	6,574	-	-	-	1	1	2	2	15	36	92	199
Male	3,643	-	-	-	1	1	2	-	9	27	74	146
Female	2,931	-	-	-	-	-	-	2	6	9	18	53
White, both sexes	5,294	-	-	-	1	-	1	2	12	26	65	142
Male	2,981	-	-	-	1	-	1	-	7	19	51	107
Female	2,313	-	-	-	-	-	-	2	5	7	14	35
Black, both sexes	1,188	-	-	-	-	1	1	-	3	10	24	54
Male	604	-	-	-	-	1	1	-	2	8	20	37
Female	584	-	-	-	-	-	-	-	1	2	4	17
American Indian 1/, both sexes	70	-	-	-	-	-	-	-	-	-	1	3
Male	41	-	-	-	-	-	-	-	-	-	1	2
Female	29	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	2	-
Male	17	-	-	-	-	-	-	-	-	-	2	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	549	-	-	-	-	-	-	-	2	1	2	5
Male	288	-	-	-	-	-	-	-	1	-	2	4
Female	261	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	533	-	-	-	-	-	-	-	2	1	2	2
Male	278	-	-	-	-	-	-	-	1	-	2	2
Female	255	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Ohio												
All races, both sexes	11,427	-	-	-	-	-	2	6	16	37	104	185
Male	5,736	-	-	-	-	-	1	6	14	30	77	130
Female	5,691	-	-	-	-	-	1	-	2	7	27	55
White, both sexes	10,305	-	-	-	-	-	-	6	13	29	88	151
Male	5,161	-	-	-	-	-	-	6	12	24	66	108
Female	5,144	-	-	-	-	-	-	-	1	5	22	43
Black, both sexes	1,098	-	-	-	-	-	1	-	3	7	16	34
Male	559	-	-	-	-	-	-	-	2	5	11	22
Female	539	-	-	-	-	-	1	-	1	2	5	12
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	1	-	-	1	-	-
Male	15	-	-	-	-	-	1	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3,555	-	-	-	-	-	2	2	2	7	33	61
Male	1,813	-	-	-	-	-	2	1	1	6	25	45
Female	1,742	-	-	-	-	-	-	1	1	1	8	16
White, both sexes	3,196	-	-	-	-	-	2	2	2	3	27	51
Male	1,628	-	-	-	-	-	2	1	1	2	19	37
Female	1,568	-	-	-	-	-	-	1	1	1	8	14
Black, both sexes	172	-	-	-	-	-	-	-	-	2	1	3
Male	93	-	-	-	-	-	-	-	-	2	1	2
Female	79	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	178	-	-	-	-	-	-	-	-	2	5	6
Male	87	-	-	-	-	-	-	-	-	2	5	6
Female	91	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Oregon												
All races, both sexes	2,418	-	-	-	-	-	-	1	3	15	24	43
Male	1,437	-	-	-	-	-	-	1	3	13	18	32
Female	981	-	-	-	-	-	-	-	-	2	6	11
White, both sexes	2,366	-	-	-	-	-	-	-	3	14	24	41
Male	1,401	-	-	-	-	-	-	-	3	12	18	30
Female	965	-	-	-	-	-	-	-	-	2	6	11
Black, both sexes	16	-	-	-	-	-	-	1	-	-	-	1
Male	12	-	-	-	-	-	-	1	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	-	1	-	1
Male	16	-	-	-	-	-	-	-	-	1	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	11,532	-	-	-	-	-	-	2	8	29	54	106
Male	5,623	-	-	-	-	-	-	2	5	21	40	81
Female	5,909	-	-	-	-	-	-	-	3	8	14	25
White, both sexes	10,793	-	-	-	-	-	-	2	7	23	47	86
Male	5,267	-	-	-	-	-	-	2	4	19	36	69
Female	5,526	-	-	-	-	-	-	-	3	4	11	17
Black, both sexes	698	-	-	-	-	-	-	-	1	6	7	20
Male	332	-	-	-	-	-	-	-	1	2	4	12
Female	366	-	-	-	-	-	-	-	-	4	3	8
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Rhode Island												
All races, both sexes	1,034	-	-	-	-	-	-	2	2	2	13	17
Male	421	-	-	-	-	-	-	2	1	1	10	13
Female	613	-	-	-	-	-	-	-	1	1	3	4
White, both sexes	998	-	-	-	-	-	-	2	2	1	12	14
Male	400	-	-	-	-	-	-	2	1	-	9	11
Female	598	-	-	-	-	-	-	-	1	1	3	3
Black, both sexes	23	-	-	-	-	-	-	-	-	1	1	1
Male	14	-	-	-	-	-	-	-	-	1	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2,473	-	-	-	-	1	3	2	4	9	25	56
Male	1,399	-	-	-	-	-	2	2	1	7	16	45
Female	1,074	-	-	-	-	1	1	-	3	2	9	11
White, both sexes	1,991	-	-	-	-	1	1	-	2	7	20	36
Male	1,155	-	-	-	-	-	1	-	1	5	13	27
Female	836	-	-	-	-	1	-	-	1	2	7	9
Black, both sexes	476	-	-	-	-	-	2	2	2	2	5	20
Male	242	-	-	-	-	-	1	2	-	2	3	18
Female	234	-	-	-	-	-	1	-	2	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
South Dakota												
All races, both sexes	672	-	-	-	-	-	-	-	-	2	1	13
Male	356	-	-	-	-	-	-	-	-	1	1	10
Female	316	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	648	-	-	-	-	-	-	-	-	2	1	12
Male	343	-	-	-	-	-	-	-	-	1	1	9
Female	305	-	-	-	-	-	-	-	-	1	-	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4,485	-	-	-	-	-	2	-	9	15	57	69
Male	2,214	-	-	-	-	-	1	-	7	10	43	47
Female	2,271	-	-	-	-	-	1	-	2	5	14	22
White, both sexes	3,957	-	-	-	-	-	1	-	7	14	41	53
Male	1,967	-	-	-	-	-	1	-	6	9	33	36
Female	1,990	-	-	-	-	-	-	-	1	5	8	17
Black, both sexes	512	-	-	-	-	-	1	-	2	1	15	16
Male	238	-	-	-	-	-	-	-	1	1	9	11
Female	274	-	-	-	-	-	1	-	1	-	6	5
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Texas												
All races, both sexes	11,607	-	1	1	-	-	3	7	11	38	108	194
Male	5,974	-	1	1	-	-	3	6	9	28	84	141
Female	5,633	-	-	-	-	-	-	1	2	10	24	53
White, both sexes	10,357	-	1	1	-	-	1	6	9	27	87	143
Male	5,378	-	1	1	-	-	1	5	7	19	71	104
Female	4,979	-	-	-	-	-	-	1	2	8	16	39
Black, both sexes	1,133	-	-	-	-	-	1	1	2	9	19	43
Male	534	-	-	-	-	-	1	1	2	7	11	30
Female	599	-	-	-	-	-	-	-	-	2	8	13
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	1	2
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	109	-	-	-	-	-	1	-	-	2	1	6
Male	58	-	-	-	-	-	1	-	-	2	1	6
Female	51	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	831	-	-	-	-	1	-	1	1	4	13	28
Male	493	-	-	-	-	1	-	1	1	3	11	24
Female	338	-	-	-	-	-	-	-	-	1	2	4
White, both sexes	803	-	-	-	-	1	-	1	1	3	11	27
Male	475	-	-	-	-	1	-	1	1	3	9	23
Female	328	-	-	-	-	-	-	-	-	-	2	4
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	1	2	-
Male	11	-	-	-	-	-	-	-	-	-	2	-
Female	7	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Vermont												
All races, both sexes	381	-	-	-	-	-	-	-	-	-	1	5
Male	186	-	-	-	-	-	-	-	-	-	1	4
Female	195	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	380	-	-	-	-	-	-	-	-	-	1	5
Male	186	-	-	-	-	-	-	-	-	-	1	4
Female	194	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3,860	-	-	-	-	-	2	1	16	14	40	80
Male	2,079	-	-	-	-	-	2	1	12	7	33	63
Female	1,781	-	-	-	-	-	-	-	4	7	7	17
White, both sexes	3,146	-	-	-	-	-	2	1	12	10	30	57
Male	1,728	-	-	-	-	-	2	1	8	6	24	46
Female	1,418	-	-	-	-	-	-	-	4	4	6	11
Black, both sexes	665	-	-	-	-	-	-	-	3	4	9	20
Male	321	-	-	-	-	-	-	-	3	1	8	14
Female	344	-	-	-	-	-	-	-	-	3	1	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	47	-	-	-	-	-	-	-	1	-	1	3
Male	29	-	-	-	-	-	-	-	1	-	1	3
Female	18	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Washington												
All races, both sexes	3,753	-	-	-	-	1	2	2	2	8	15	31
Male	2,105	-	-	-	-	-	2	2	2	7	10	19
Female	1,648	-	-	-	-	1	-	-	-	1	5	12
White, both sexes	3,571	-	-	-	-	-	-	2	1	5	13	27
Male	1,991	-	-	-	-	-	-	2	1	5	8	16
Female	1,580	-	-	-	-	-	-	-	-	-	5	11
Black, both sexes	67	-	-	-	-	1	1	-	-	2	1	2
Male	42	-	-	-	-	-	1	-	-	1	1	1
Female	25	-	-	-	-	1	-	-	-	1	-	1
American Indian 1/, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	83	-	-	-	-	-	1	-	1	1	1	2
Male	53	-	-	-	-	-	1	-	1	1	1	2
Female	30	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1,931	-	-	-	-	1	-	1	4	14	22	50
Male	1,017	-	-	-	-	1	-	1	4	9	18	38
Female	914	-	-	-	-	-	-	-	-	5	4	12
White, both sexes	1,879	-	-	-	-	1	-	-	4	13	21	48
Male	993	-	-	-	-	1	-	-	4	8	17	37
Female	886	-	-	-	-	-	-	-	-	5	4	11
Black, both sexes	52	-	-	-	-	-	-	1	-	1	1	2
Male	24	-	-	-	-	-	-	1	-	1	1	1
Female	28	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Wisconsin												
All races, both sexes	3,531	-	-	-	-	-	-	1	5	4	22	48
Male	1,884	-	-	-	-	-	-	1	5	3	13	37
Female	1,647	-	-	-	-	-	-	-	-	1	9	11
White, both sexes	3,395	-	-	-	-	-	-	-	4	4	20	42
Male	1,806	-	-	-	-	-	-	-	4	3	13	32
Female	1,589	-	-	-	-	-	-	-	-	1	7	10
Black, both sexes	107	-	-	-	-	-	-	-	-	-	1	5
Male	59	-	-	-	-	-	-	-	-	-	-	4
Female	48	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	1	-	1	-
Male	11	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	1	-	-	-	1
Male	8	-	-	-	-	-	-	1	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	279	-	-	-	-	-	-	-	-	2	5	5
Male	159	-	-	-	-	-	-	-	-	1	3	3
Female	120	-	-	-	-	-	-	-	-	1	2	2
White, both sexes	276	-	-	-	-	-	-	-	-	2	4	5
Male	158	-	-	-	-	-	-	-	-	1	2	3
Female	118	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
United States												
All races, both sexes	5,530	7,784	10,382	13,288	19,307	29,990	41,775	43,656	33,185	13,680	3,131	8
Male	4,153	5,647	7,219	8,702	12,046	17,407	21,571	19,173	11,129	3,228	485	5
Female	1,377	2,137	3,163	4,586	7,261	12,583	20,204	24,483	22,056	10,452	2,646	3
White, both sexes	4,322	6,289	8,547	11,081	16,613	26,581	37,977	40,511	30,906	12,663	2,807	6
Male	3,329	4,684	6,087	7,401	10,592	15,719	19,893	17,992	10,436	3,000	427	5
Female	993	1,605	2,460	3,680	6,021	10,862	18,084	22,519	20,470	9,663	2,380	1
Black, both sexes	1,047	1,307	1,572	1,821	2,210	2,794	3,042	2,438	1,776	809	273	2
Male	703	827	947	1,041	1,165	1,353	1,295	861	476	152	39	-
Female	344	480	625	780	1,045	1,441	1,747	1,577	1,300	657	234	2
American Indian 1/, both sexes	44	55	85	83	97	106	86	94	71	23	8	-
Male	29	38	64	49	61	58	43	38	17	6	-	-
Female	15	17	21	34	36	48	43	56	54	17	8	-
Asian or Pacific Islander 2/, both sexes	117	133	178	303	387	509	670	613	432	185	43	-
Male	92	98	121	211	228	277	340	282	200	70	19	-
Female	25	35	57	92	159	232	330	331	232	115	24	-
Alabama												
All races, both sexes	77	133	157	206	281	424	466	391	279	93	13	-
Male	56	93	91	133	167	225	236	158	84	23	1	-
Female	21	40	66	73	114	199	230	233	195	70	12	-
White, both sexes	57	97	117	159	232	351	418	344	251	75	9	-
Male	45	71	71	110	142	197	219	141	77	18	1	-
Female	12	26	46	49	90	154	199	203	174	57	8	-
Black, both sexes	17	35	39	47	47	71	48	45	28	18	4	-
Male	10	21	19	23	25	28	17	17	7	5	-	-
Female	7	14	20	24	22	43	31	28	21	13	4	-
American Indian 1/, both sexes	2	1	1	-	1	-	-	1	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	2	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Alaska												
All races, both sexes	6	9	3	9	11	26	26	14	7	4	1	-
Male	4	8	3	6	10	19	18	8	3	1	-	-
Female	2	1	-	3	1	7	8	6	4	3	1	-
White, both sexes	5	7	3	8	6	20	24	10	5	2	1	-
Male	3	6	3	5	6	14	18	6	3	-	-	-
Female	2	1	-	3	-	6	6	4	2	2	1	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	1	4	5	1	3	2	2	-	-
Male	-	2	-	1	3	5	-	2	-	1	-	-
Female	-	-	-	-	1	-	1	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Arizona												
All races, both sexes	97	141	225	252	362	580	759	781	538	188	35	-
Male	68	116	168	177	243	343	424	398	217	55	7	-
Female	29	25	57	75	119	237	335	383	321	133	28	-
White, both sexes	91	133	211	233	344	553	746	763	526	187	33	-
Male	64	109	162	163	232	332	417	390	213	54	7	-
Female	27	24	49	70	112	221	329	373	313	133	26	-
Black, both sexes	3	2	5	11	9	13	9	13	5	1	1	-
Male	2	2	2	10	5	4	5	3	1	1	-	-
Female	1	-	3	1	4	9	4	10	4	-	1	-
American Indian 1/, both sexes	3	6	5	8	6	9	3	1	7	-	-	-
Male	2	5	2	4	4	4	2	1	3	-	-	-
Female	1	1	3	4	2	5	1	-	4	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	4	-	3	5	1	4	-	-	1	-
Male	-	-	2	-	2	3	-	4	-	-	-	-
Female	-	-	2	-	1	2	1	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Arkansas												
All races, both sexes	45	66	93	132	186	257	335	337	256	130	18	-
Male	31	44	55	83	120	137	181	139	80	29	2	-
Female	14	22	38	49	66	120	154	198	176	101	16	-
White, both sexes	35	52	71	116	167	218	303	312	233	110	14	-
Male	25	37	44	72	109	122	165	133	73	26	2	-
Female	10	15	27	44	58	96	138	179	160	84	12	-
Black, both sexes	10	14	22	16	18	37	32	25	23	20	4	-
Male	6	7	11	11	10	15	16	6	7	3	-	-
Female	4	7	11	5	8	22	16	19	16	17	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
California												
All races, both sexes	611	850	1,101	1,329	2,064	3,253	4,848	5,155	3,905	1,643	440	1
Male	451	594	773	851	1,285	1,892	2,530	2,333	1,404	437	83	1
Female	160	256	328	478	779	1,361	2,318	2,822	2,501	1,206	357	-
White, both sexes	479	654	900	1,052	1,709	2,758	4,277	4,621	3,527	1,479	389	1
Male	355	471	639	676	1,072	1,626	2,254	2,097	1,256	386	65	1
Female	124	183	261	376	637	1,132	2,023	2,524	2,271	1,093	324	-
Black, both sexes	80	141	126	145	187	250	238	212	152	67	34	-
Male	55	83	85	86	111	126	104	82	41	13	7	-
Female	25	58	41	59	76	124	134	130	111	54	27	-
American Indian 1/, both sexes	5	7	13	16	12	10	9	18	8	1	-	-
Male	4	4	6	10	8	7	4	6	3	-	-	-
Female	1	3	7	6	4	3	5	12	5	1	-	-
Asian or Pacific Islander 2/, both sexes	47	48	62	116	156	235	324	304	218	96	17	-
Male	37	36	43	79	94	133	168	148	104	38	11	-
Female	10	12	19	37	62	102	156	156	114	58	6	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Colorado												
All races, both sexes	62	78	98	132	180	272	390	425	301	119	28	-
Male	51	66	74	98	122	174	218	209	110	20	5	-
Female	11	12	24	34	58	98	172	216	191	99	23	-
White, both sexes	61	70	90	122	165	260	382	416	296	116	28	-
Male	51	59	68	92	116	169	216	206	109	20	5	-
Female	10	11	22	30	49	91	166	210	187	96	23	-
Black, both sexes	1	8	6	7	13	7	4	7	5	2	-	-
Male	-	7	4	3	4	3	-	1	1	-	-	-
Female	1	1	2	4	9	4	4	6	4	2	-	-
American Indian 1/, both sexes	-	-	1	1	1	3	-	-	-	1	-	-
Male	-	-	1	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	1	2	4	2	-	-	-	-
Male	-	-	1	2	1	1	2	2	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Connecticut												
All races, both sexes	60	73	82	110	194	322	505	597	493	222	49	-
Male	47	59	61	81	123	186	256	264	161	39	10	-
Female	13	14	21	29	71	136	249	333	332	183	39	-
White, both sexes	50	59	68	91	177	301	484	576	480	210	46	-
Male	41	47	51	70	114	178	247	255	155	34	10	-
Female	9	12	17	21	63	123	237	321	325	176	36	-
Black, both sexes	10	10	13	18	14	19	16	21	13	8	2	-
Male	6	9	9	10	9	7	8	9	6	2	-	-
Female	4	1	4	8	5	12	8	12	7	6	2	-
American Indian 1/, both sexes	-	-	-	1	1	1	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	4	1	-	2	1	5	-	-	3	1	-
Male	-	3	1	-	-	1	1	-	-	3	-	-
Female	-	1	-	-	2	-	4	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Delaware												
All races, both sexes	18	22	42	37	45	101	135	139	95	31	4	-
Male	13	14	25	23	28	58	76	60	28	10	1	-
Female	5	8	17	14	17	43	59	79	67	21	3	-
White, both sexes	11	16	31	32	37	90	123	121	83	29	2	-
Male	9	12	21	22	22	56	72	56	26	10	1	-
Female	2	4	10	10	15	34	51	65	57	19	1	-
Black, both sexes	6	5	11	5	8	10	12	18	11	2	2	-
Male	4	2	4	1	6	1	4	4	1	-	-	-
Female	2	3	7	4	2	9	8	14	10	2	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	13	14	13	26	25	46	53	63	27	21	9	-
Male	11	8	8	19	14	22	25	24	7	3	3	-
Female	2	6	5	7	11	24	28	39	20	18	6	-
White, both sexes	2	2	2	-	6	3	10	19	8	5	3	-
Male	1	1	2	-	4	2	6	9	2	2	1	-
Female	1	1	-	-	2	1	4	10	6	3	2	-
Black, both sexes	10	11	11	26	19	43	42	43	18	15	6	-
Male	9	6	6	19	10	20	19	14	5	-	2	-
Female	1	5	5	7	9	23	23	29	13	15	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	1	1	1	-	-
Male	1	1	-	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Florida												
All races, both sexes	461	548	849	1,119	1,611	2,687	3,757	3,900	3,014	1,203	259	-
Male	353	391	620	780	1,039	1,631	2,021	1,893	1,129	339	46	-
Female	108	157	229	339	572	1,056	1,736	2,007	1,885	864	213	-
White, both sexes	364	457	722	971	1,421	2,485	3,516	3,697	2,862	1,139	241	-
Male	282	327	538	692	934	1,534	1,924	1,819	1,081	320	45	-
Female	82	130	184	279	487	951	1,592	1,878	1,781	819	196	-
Black, both sexes	94	87	122	138	182	192	224	191	147	59	17	-
Male	68	61	79	81	103	93	88	69	44	18	1	-
Female	26	26	43	57	79	99	136	122	103	41	16	-
American Indian 1/, both sexes	1	-	4	1	-	4	4	1	1	-	1	-
Male	1	-	3	-	-	2	4	1	1	-	-	-
Female	-	-	1	1	-	2	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	4	1	9	8	6	13	11	4	5	-	-
Male	2	3	-	7	2	2	5	4	3	1	-	-
Female	-	1	1	2	6	4	8	7	1	4	-	-
Georgia												
All races, both sexes	184	246	278	323	415	585	704	594	446	168	30	-
Male	137	184	189	222	272	344	349	260	130	33	5	-
Female	47	62	89	101	143	241	355	334	316	135	25	-
White, both sexes	119	166	217	235	325	490	586	507	378	139	20	-
Male	92	128	155	175	222	302	299	238	112	30	3	-
Female	27	38	62	60	103	188	287	269	266	109	17	-
Black, both sexes	63	77	61	82	88	93	115	86	65	28	10	-
Male	44	55	34	43	49	41	49	22	18	3	2	-
Female	19	22	27	39	39	52	66	64	47	25	8	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	2	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	-	5	2	2	2	1	1	1	-	-
Male	1	1	-	3	1	1	-	-	-	-	-	-
Female	1	2	-	2	1	1	2	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Hawaii												
All races, both sexes	29	28	48	45	61	82	115	100	71	35	8	-
Male	24	22	33	33	37	44	77	55	37	16	3	-
Female	5	6	15	12	24	38	38	45	34	19	5	-
White, both sexes	5	6	17	8	20	25	33	36	22	13	2	-
Male	5	5	14	7	12	14	24	21	11	6	1	-
Female	-	1	3	1	8	11	9	15	11	7	1	-
Black, both sexes	1	-	1	-	-	-	2	-	1	-	-	-
Male	1	-	1	-	-	-	2	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	22	30	37	41	57	80	64	48	22	6	-
Male	18	17	18	26	25	30	51	34	25	10	2	-
Female	5	5	12	11	16	27	29	30	23	12	4	-
Idaho												
All races, both sexes	16	14	28	42	66	105	132	138	115	34	6	-
Male	11	11	23	28	43	70	82	71	50	11	1	-
Female	5	3	5	14	23	35	50	67	65	23	5	-
White, both sexes	15	13	26	40	65	105	129	137	114	34	6	-
Male	10	10	21	26	42	70	80	71	50	11	1	-
Female	5	3	5	14	23	35	49	66	64	23	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	2	1	-	-	1	1	-	-	-	-
Male	-	1	2	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	1	-	2	-	1	-	-	-
Male	1	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Illinois												
All races, both sexes	248	297	392	532	722	1,168	1,613	1,895	1,440	614	130	-
Male	191	223	272	349	430	695	826	790	453	132	11	-
Female	57	74	120	183	292	473	787	1,105	987	482	119	-
White, both sexes	177	228	304	402	596	993	1,438	1,750	1,338	560	118	-
Male	144	179	219	272	366	606	762	735	428	119	9	-
Female	33	49	85	130	230	387	676	1,015	910	441	109	-
Black, both sexes	70	67	84	119	114	159	158	127	94	47	11	-
Male	46	42	49	69	56	81	62	47	23	11	2	-
Female	24	25	35	50	58	78	96	80	71	36	9	-
American Indian 1/, both sexes	-	-	2	-	1	1	-	1	1	-	-	-
Male	-	-	2	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	11	11	15	17	17	7	7	1	-
Male	1	2	2	8	7	8	2	8	2	2	-	-
Female	-	-	-	3	4	7	15	9	5	5	1	-
Indiana												
All races, both sexes	163	209	252	330	447	688	904	928	665	294	49	-
Male	131	152	187	204	252	405	495	403	208	67	6	-
Female	32	57	65	126	195	283	409	525	457	227	43	-
White, both sexes	141	188	227	288	388	634	865	894	641	281	44	-
Male	114	138	172	184	229	376	473	392	204	66	6	-
Female	27	50	55	104	159	258	392	502	437	215	38	-
Black, both sexes	22	21	25	38	56	52	34	31	21	13	5	-
Male	17	14	15	17	22	28	20	10	2	1	-	-
Female	5	7	10	21	34	24	14	21	19	12	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	4	3	2	5	3	3	-	-	-
Male	-	-	-	3	1	1	2	1	2	-	-	-
Female	-	-	-	1	2	1	3	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Iowa												
All races, both sexes	47	71	100	125	160	332	508	563	481	219	70	-
Male	38	51	74	83	99	193	284	274	180	49	9	-
Female	9	20	26	42	61	139	224	289	301	170	61	-
White, both sexes	46	68	98	122	156	331	503	561	481	217	70	-
Male	37	48	72	80	97	193	281	273	180	49	9	-
Female	9	20	26	42	59	138	222	288	301	168	61	-
Black, both sexes	1	3	2	1	2	1	4	-	-	2	-	-
Male	1	3	2	1	1	-	3	-	-	-	-	-
Female	-	-	-	-	1	1	1	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	-	2	-	-	-	-
Male	-	-	-	2	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Kansas												
All races, both sexes	58	75	102	127	184	265	370	415	325	166	33	-
Male	42	61	69	83	124	164	201	195	130	44	6	-
Female	16	14	33	44	60	101	169	220	195	122	27	-
White, both sexes	54	68	89	116	173	248	347	408	313	162	32	-
Male	38	54	60	76	119	156	190	192	123	44	6	-
Female	16	14	29	40	54	92	157	216	190	118	26	-
Black, both sexes	3	5	9	11	7	15	17	7	9	4	-	-
Male	3	5	5	7	3	8	10	3	5	-	-	-
Female	-	-	4	4	4	7	7	4	4	4	-	-
American Indian 1/, both sexes	1	2	4	-	3	1	3	-	2	-	1	-
Male	1	2	4	-	1	-	-	-	2	-	-	-
Female	-	-	-	-	2	1	3	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	3	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Kentucky												
All races, both sexes	82	120	151	209	283	409	517	496	318	146	30	-
Male	63	82	98	130	181	240	228	191	92	25	4	-
Female	19	38	53	79	102	169	289	305	226	121	26	-
White, both sexes	72	113	147	196	273	387	496	479	305	141	28	-
Male	57	79	94	123	176	227	222	184	90	24	3	-
Female	15	34	53	73	97	160	274	295	215	117	25	-
Black, both sexes	9	6	4	12	10	22	21	17	12	5	2	-
Male	5	2	4	6	5	13	6	7	2	1	1	-
Female	4	4	-	6	5	9	15	10	10	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	-	-	-	-	1	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Louisiana												
All races, both sexes	90	116	158	199	283	345	422	389	266	93	26	-
Male	61	77	111	126	189	194	228	175	83	24	4	-
Female	29	39	47	73	94	151	194	214	183	69	22	-
White, both sexes	59	75	113	144	206	282	338	331	222	77	18	-
Male	42	51	85	94	142	161	185	153	68	19	2	-
Female	17	24	28	50	64	121	153	178	154	58	16	-
Black, both sexes	31	40	45	53	73	58	82	55	44	16	8	-
Male	19	25	26	30	45	31	42	20	15	5	2	-
Female	12	15	19	23	28	27	40	35	29	11	6	-
American Indian 1/, both sexes	-	-	-	-	-	2	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	4	3	1	2	-	-	-	-
Male	-	1	-	2	2	2	-	1	-	-	-	-
Female	-	-	-	-	2	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Maine												
All races, both sexes	26	29	50	72	96	130	181	193	150	56	5	-
Male	23	23	36	49	65	70	101	98	51	13	-	-
Female	3	6	14	23	31	60	80	95	99	43	5	-
White, both sexes	25	29	50	72	96	130	180	193	150	56	5	-
Male	22	23	36	49	65	70	100	98	51	13	-	-
Female	3	6	14	23	31	60	80	95	99	43	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	59	95	172	191	260	381	589	581	391	163	41	-
Male	40	67	119	126	150	210	324	281	123	34	12	-
Female	19	28	53	65	110	171	265	300	268	129	29	-
White, both sexes	31	62	113	125	191	297	486	499	346	139	33	-
Male	23	48	83	83	111	172	276	246	107	29	12	-
Female	8	14	30	42	80	125	210	253	239	110	21	-
Black, both sexes	27	32	58	59	61	80	99	73	41	22	8	-
Male	16	19	35	36	34	36	46	31	14	5	-	-
Female	11	13	23	23	27	44	53	42	27	17	8	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	7	8	4	3	9	4	2	-	-
Male	-	-	1	7	5	2	2	4	2	-	-	-
Female	-	1	-	-	3	2	1	5	2	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Massachusetts												
All races, both sexes	63	106	190	215	343	519	764	923	732	291	82	-
Male	53	78	148	137	229	334	420	382	233	74	9	-
Female	10	28	42	78	114	185	344	541	499	217	73	-
White, both sexes	55	102	179	200	322	496	734	906	717	287	80	-
Male	46	76	138	127	222	323	407	374	232	73	9	-
Female	9	26	41	73	100	173	327	532	485	214	71	-
Black, both sexes	7	4	9	12	18	21	18	16	11	3	2	-
Male	6	2	9	9	6	10	8	7	-	1	-	-
Female	1	2	-	3	12	11	10	9	11	2	2	-
American Indian 1/, both sexes	-	-	2	1	2	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	2	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	1	2	12	1	4	1	-	-
Male	1	-	-	1	1	1	5	1	1	-	-	-
Female	-	-	-	1	-	1	7	-	3	1	-	-
Michigan												
All races, both sexes	200	294	347	473	698	1,077	1,652	1,704	1,306	505	103	-
Male	151	218	228	315	453	610	845	685	435	122	14	-
Female	49	76	119	158	245	467	807	1,019	871	383	89	-
White, both sexes	163	241	285	419	607	963	1,520	1,589	1,225	465	93	-
Male	127	182	188	285	398	556	791	642	405	111	12	-
Female	36	59	97	134	209	407	729	947	820	354	81	-
Black, both sexes	35	50	57	51	79	109	126	105	77	38	9	-
Male	23	33	35	28	47	50	49	40	26	9	2	-
Female	12	17	22	23	32	59	77	65	51	29	7	-
American Indian 1/, both sexes	1	2	2	2	6	2	2	3	-	-	-	-
Male	-	2	2	1	5	1	1	-	-	-	-	-
Female	1	-	-	1	1	1	1	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	3	1	6	3	4	7	4	2	1	-
Male	1	1	3	1	3	3	4	3	4	2	-	-
Female	-	-	-	-	3	-	-	4	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Minnesota												
All races, both sexes	96	134	130	158	211	351	501	565	424	180	42	-
Male	78	115	101	109	140	232	290	307	153	63	12	-
Female	18	19	29	49	71	119	211	258	271	117	30	-
White, both sexes	87	129	124	148	204	343	495	557	420	180	42	-
Male	70	111	96	102	137	226	288	304	151	63	12	-
Female	17	18	28	46	67	117	207	253	269	117	30	-
Black, both sexes	7	3	3	3	3	4	4	4	-	-	-	-
Male	6	2	3	2	1	3	1	3	-	-	-	-
Female	1	1	-	1	2	1	3	1	-	-	-	-
American Indian 1/, both sexes	1	-	2	3	2	1	2	1	2	-	-	-
Male	1	-	1	2	1	1	1	-	1	-	-	-
Female	-	-	1	1	1	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	4	2	3	-	3	2	-	-	-
Male	1	2	1	3	1	2	-	-	1	-	-	-
Female	-	-	-	1	1	1	-	3	1	-	-	-
Mississippi												
All races, both sexes	89	100	137	162	208	281	300	271	188	79	14	1
Male	60	73	88	107	124	151	147	103	61	11	1	1
Female	29	27	49	55	84	130	153	168	127	68	13	-
White, both sexes	52	66	88	118	154	228	241	224	155	67	9	1
Male	34	50	58	79	99	126	114	85	50	10	-	1
Female	18	16	30	39	55	102	127	139	105	57	9	-
Black, both sexes	37	33	49	43	52	50	59	47	33	11	5	-
Male	26	22	30	28	25	24	33	18	11	1	1	-
Female	11	11	19	15	27	26	26	29	22	10	4	-
American Indian 1/, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	2	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Missouri												
All races, both sexes	152	187	226	294	432	670	854	989	730	330	91	-
Male	108	143	154	180	262	383	436	392	223	84	10	-
Female	44	44	72	114	170	287	418	597	507	246	81	-
White, both sexes	123	154	186	251	379	615	794	930	691	312	81	-
Male	89	120	130	156	232	357	404	376	210	79	8	-
Female	34	34	56	95	147	258	390	554	481	233	73	-
Black, both sexes	29	33	39	42	51	50	58	56	39	17	10	-
Male	19	23	24	24	29	24	31	15	13	5	2	-
Female	10	10	15	18	22	26	27	41	26	12	8	-
American Indian 1/, both sexes	-	-	-	1	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	4	-	2	-	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	1	-	1	3	-	1	-	1	-	-
Montana												
All races, both sexes	22	36	27	41	50	65	97	109	72	30	8	-
Male	18	27	19	34	30	43	55	60	32	16	-	-
Female	4	9	8	7	20	22	42	49	40	14	8	-
White, both sexes	18	34	25	39	48	63	97	107	70	30	7	-
Male	14	25	17	33	30	41	55	59	32	16	-	-
Female	4	9	8	6	18	22	42	48	38	14	7	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	1	2	2	2	2	-	2	2	-	1	-
Male	4	1	2	1	-	2	-	1	-	-	-	-
Female	-	-	-	1	2	-	-	1	2	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Nebraska												
All races, both sexes	39	32	52	70	85	160	220	259	222	96	21	-
Male	31	26	38	47	55	108	124	135	89	25	4	-
Female	8	6	14	23	30	52	96	124	133	71	17	-
White, both sexes	31	27	48	62	78	153	219	258	221	94	21	-
Male	24	23	35	42	54	106	124	134	88	24	4	-
Female	7	4	13	20	24	47	95	124	133	70	17	-
Black, both sexes	7	5	2	6	4	5	1	-	-	2	-	-
Male	6	3	1	5	-	2	-	-	-	1	-	-
Female	1	2	1	1	4	3	1	-	-	1	-	-
American Indian 1/, both sexes	1	-	1	1	2	1	-	1	1	-	-	-
Male	1	-	1	-	1	-	-	1	1	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	1	-	-	-	-	-	-
Nevada												
All races, both sexes	27	53	67	86	138	172	208	153	94	13	4	-
Male	19	44	45	55	94	107	114	72	33	2	2	-
Female	8	9	22	31	44	65	94	81	61	11	2	-
White, both sexes	25	45	62	76	125	157	185	146	89	12	4	-
Male	19	40	43	49	88	101	100	70	31	2	2	-
Female	6	5	19	27	37	56	85	76	58	10	2	-
Black, both sexes	2	7	3	5	8	7	15	5	4	-	-	-
Male	-	4	-	3	5	3	8	1	2	-	-	-
Female	2	3	3	2	3	4	7	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	3	-	-	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	5	4	5	8	2	-	1	-	-
Male	-	-	2	3	1	1	6	1	-	-	-	-
Female	-	1	-	2	3	4	2	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
New Hampshire												
All races, both sexes	14	29	45	56	90	104	151	216	159	83	18	-
Male	13	23	36	39	61	59	68	94	50	17	-	-
Female	1	6	9	17	29	45	83	122	109	66	18	-
White, both sexes	14	29	45	55	88	102	151	216	159	83	18	-
Male	13	23	36	38	60	57	68	94	50	17	-	-
Female	1	6	9	17	28	45	83	122	109	66	18	-
Black, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	114	175	264	333	564	955	1,516	1,744	1,352	580	138	2
Male	89	127	188	216	349	525	757	727	411	120	27	1
Female	25	48	76	117	215	430	759	1,017	941	460	111	1
White, both sexes	81	138	212	269	484	847	1,375	1,649	1,280	534	123	1
Male	69	105	161	175	311	474	704	694	393	113	24	1
Female	12	33	51	94	173	373	671	955	887	421	99	-
Black, both sexes	28	32	47	52	70	100	122	71	59	37	14	1
Male	16	20	24	32	33	45	49	23	16	4	3	-
Female	12	12	23	20	37	55	73	48	43	33	11	1
American Indian 1/, both sexes	-	1	-	1	2	-	1	1	3	3	-	-
Male	-	-	-	-	1	-	1	1	1	1	-	-
Female	-	1	-	1	1	-	-	-	2	2	-	-
Asian or Pacific Islander 2/, both sexes	5	4	5	11	8	8	18	23	10	6	1	-
Male	4	2	3	9	4	6	3	9	1	2	-	-
Female	1	2	2	2	4	2	15	14	9	4	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
New Mexico												
All races, both sexes	20	28	39	66	113	132	217	179	155	57	13	-
Male	18	22	28	42	73	75	114	77	51	18	1	-
Female	2	6	11	24	40	57	103	102	104	39	12	-
White, both sexes	19	25	32	63	104	121	208	171	149	53	13	-
Male	17	21	22	39	67	68	110	72	49	17	1	-
Female	2	4	10	24	37	53	98	99	100	36	12	-
Black, both sexes	-	1	2	1	3	2	5	2	3	1	-	-
Male	-	1	2	1	2	2	2	1	1	-	-	-
Female	-	-	-	-	1	-	3	1	2	1	-	-
American Indian 1/, both sexes	1	2	5	2	5	9	4	6	2	3	-	-
Male	1	-	4	2	4	5	2	4	-	1	-	-
Female	-	2	1	-	1	4	2	2	2	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
New York												
All races, both sexes	402	633	911	1,267	2,040	3,373	4,926	5,424	4,792	2,094	514	-
Male	279	431	556	763	1,178	1,782	2,256	2,077	1,452	442	68	-
Female	123	202	355	504	862	1,591	2,670	3,347	3,340	1,652	446	-
White, both sexes	304	487	659	968	1,646	2,786	4,251	4,880	4,380	1,905	448	-
Male	229	338	424	603	975	1,501	1,977	1,901	1,351	410	58	-
Female	75	149	235	365	671	1,285	2,274	2,979	3,029	1,495	390	-
Black, both sexes	80	127	219	260	324	499	573	445	313	160	54	-
Male	39	76	111	134	159	238	226	138	59	24	6	-
Female	41	51	108	126	165	261	347	307	254	136	48	-
American Indian 1/, both sexes	1	2	1	2	-	2	-	7	6	-	-	-
Male	1	2	1	1	-	1	-	4	2	-	-	-
Female	-	-	-	1	-	1	-	3	4	-	-	-
Asian or Pacific Islander 2/, both sexes	17	17	32	37	70	86	102	92	93	29	12	-
Male	10	15	20	25	44	42	53	34	40	8	4	-
Female	7	2	12	12	26	44	49	58	53	21	8	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
North Carolina												
All races, both sexes	279	401	435	532	688	894	1,048	1,001	677	222	48	1
Male	214	291	318	376	425	544	519	444	205	44	2	1
Female	65	110	117	156	263	350	529	557	472	178	46	-
White, both sexes	200	300	317	415	540	721	880	874	581	182	34	1
Male	158	231	236	304	347	453	446	398	183	37	1	1
Female	42	69	81	111	193	268	434	476	398	145	33	-
Black, both sexes	75	93	106	109	136	158	155	120	90	39	14	-
Male	52	54	72	66	70	83	66	44	20	7	1	-
Female	23	39	34	43	66	75	89	76	70	32	13	-
American Indian 1/, both sexes	2	7	8	6	9	14	9	7	4	-	-	-
Male	2	5	7	4	6	7	5	2	-	-	-	-
Female	-	2	1	2	3	7	4	5	4	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	4	2	3	1	4	-	2	1	-	-
Male	2	1	3	2	2	1	2	-	2	-	-	-
Female	-	-	1	-	1	-	2	-	-	1	-	-
North Dakota												
All races, both sexes	8	16	13	22	32	61	108	141	76	53	9	-
Male	5	14	10	15	23	42	66	64	25	16	1	-
Female	3	2	3	7	9	19	42	77	51	37	8	-
White, both sexes	6	15	13	19	30	59	108	139	76	52	9	-
Male	4	14	10	13	21	40	66	63	25	16	1	-
Female	2	1	3	6	9	19	42	76	51	36	8	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	2	2	2	-	2	-	1	-	-
Male	1	-	-	1	2	2	-	1	-	-	-	-
Female	1	1	-	1	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Ohio												
All races, both sexes	281	408	528	689	1,011	1,594	2,156	2,153	1,512	623	122	-
Male	219	271	365	442	607	893	1,094	947	493	126	21	-
Female	62	137	163	247	404	701	1,062	1,206	1,019	497	101	-
White, both sexes	216	322	442	579	881	1,441	1,993	2,009	1,429	589	117	-
Male	170	221	315	376	536	814	1,016	887	470	120	20	-
Female	46	101	127	203	345	627	977	1,122	959	469	97	-
Black, both sexes	65	86	83	108	128	152	155	141	81	33	5	-
Male	49	50	47	65	69	79	74	57	22	6	1	-
Female	16	36	36	43	59	73	81	84	59	27	4	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	3	2	2	-	7	3	1	1	-	-
Male	-	-	3	1	2	-	3	3	1	-	-	-
Female	-	-	-	1	-	-	4	-	-	1	-	-
Oklahoma												
All races, both sexes	92	156	194	227	347	486	623	627	462	189	45	-
Male	70	102	131	143	236	275	319	270	135	45	7	-
Female	22	54	63	84	111	211	304	357	327	144	38	-
White, both sexes	77	134	172	191	308	448	574	574	424	171	36	-
Male	62	89	113	121	209	252	295	252	127	40	6	-
Female	15	45	59	70	99	196	279	322	297	131	30	-
Black, both sexes	7	15	10	13	16	16	22	32	22	10	3	-
Male	5	10	8	10	10	11	13	11	6	3	1	-
Female	2	5	2	3	6	5	9	21	16	7	2	-
American Indian 1/, both sexes	8	6	11	22	22	20	27	20	16	8	5	-
Male	3	3	9	11	16	10	11	7	2	2	-	-
Female	5	3	2	11	6	10	16	13	14	6	5	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	1	2	-	1	-	-	1	-
Male	-	-	1	1	1	2	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Oregon												
All races, both sexes	88	117	143	162	190	307	407	454	323	122	19	-
Male	72	95	113	108	125	187	242	240	142	45	1	-
Female	16	22	30	54	65	120	165	214	181	77	18	-
White, both sexes	79	115	138	158	183	301	401	449	320	121	19	-
Male	65	93	109	106	120	182	239	237	142	44	1	-
Female	14	22	29	52	63	119	162	212	178	77	18	-
Black, both sexes	4	-	2	3	2	1	1	-	1	-	-	-
Male	4	-	1	2	2	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	4	-	1	-	2	1	2	1	1	1	-	-
Male	2	-	1	-	1	1	1	1	-	1	-	-
Female	2	-	-	-	1	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	1	3	4	3	4	1	-	-	-
Male	1	2	2	-	2	3	2	2	-	-	-	-
Female	-	-	-	1	1	1	1	2	1	-	-	-
Pennsylvania												
All races, both sexes	187	276	415	570	927	1,562	2,317	2,489	1,758	682	147	3
Male	133	208	288	381	563	938	1,153	1,066	577	144	22	1
Female	54	68	127	189	364	624	1,164	1,423	1,181	538	125	2
White, both sexes	165	244	363	507	847	1,459	2,165	2,390	1,701	649	136	2
Male	117	189	256	342	520	882	1,086	1,030	556	137	21	1
Female	48	55	107	165	327	577	1,079	1,360	1,145	512	115	1
Black, both sexes	21	30	50	53	78	96	146	92	55	32	10	1
Male	15	17	30	34	43	52	63	33	20	6	-	-
Female	6	13	20	19	35	44	83	59	35	26	10	1
American Indian 1/, both sexes	-	-	-	-	-	1	1	3	-	-	-	-
Male	-	-	-	-	-	-	1	2	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	10	2	6	5	4	2	1	1	-
Male	1	2	2	5	-	4	3	1	1	1	1	-
Female	-	-	-	5	2	2	2	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Rhode Island												
All races, both sexes	14	27	29	32	62	100	180	228	194	105	27	-
Male	12	20	21	19	37	54	82	85	52	10	2	-
Female	2	7	8	13	25	46	98	143	142	95	25	-
White, both sexes	12	25	26	30	61	93	178	222	190	105	25	-
Male	11	18	18	18	36	49	82	83	52	10	-	-
Female	1	7	8	12	25	44	96	139	138	95	25	-
Black, both sexes	-	2	2	2	1	5	1	3	3	-	1	-
Male	-	2	2	1	1	3	-	1	-	-	1	-
Female	-	-	-	1	-	2	1	2	3	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	1	-	-	-	-	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	1	-	1	-	-	1	-
Male	1	-	1	-	-	1	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
South Carolina												
All races, both sexes	88	135	167	213	280	392	406	367	234	71	20	-
Male	66	96	112	147	178	225	217	180	90	12	3	-
Female	22	39	55	66	102	167	189	187	144	59	17	-
White, both sexes	66	96	129	167	228	319	337	315	195	60	12	-
Male	52	70	90	121	153	190	190	159	71	11	1	-
Female	14	26	39	46	75	129	147	156	124	49	11	-
Black, both sexes	22	38	38	46	51	71	68	51	39	11	8	-
Male	14	25	22	26	25	34	27	21	19	1	2	-
Female	8	13	16	20	26	37	41	30	20	10	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	2	1	1	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
South Dakota												
All races, both sexes	13	18	27	23	46	99	125	113	125	50	17	-
Male	11	10	23	14	29	71	67	52	48	16	3	-
Female	2	8	4	9	17	28	58	61	77	34	14	-
White, both sexes	13	14	23	20	45	95	124	110	123	49	17	-
Male	11	8	20	11	29	69	66	52	47	16	3	-
Female	2	6	3	9	16	26	58	58	76	33	14	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	4	4	3	1	4	1	3	2	1	-	-
Male	-	2	3	3	-	2	1	-	1	-	-	-
Female	-	2	1	-	1	2	-	3	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	146	185	227	341	436	610	803	781	558	210	36	-
Male	100	139	146	201	262	346	395	295	170	46	6	-
Female	46	46	81	140	174	264	408	486	388	164	30	-
White, both sexes	109	149	193	302	374	538	734	717	504	191	30	-
Male	77	112	122	185	232	308	367	270	161	42	6	-
Female	32	37	71	117	142	230	367	447	343	149	24	-
Black, both sexes	37	35	34	35	58	71	67	63	52	19	6	-
Male	23	26	24	14	28	38	27	25	7	4	-	-
Female	14	9	10	21	30	33	40	38	45	15	6	-
American Indian 1/, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	4	3	1	1	1	2	-	-	-
Male	-	-	-	2	2	-	-	-	2	-	-	-
Female	-	-	-	2	1	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Texas												
All races, both sexes	284	476	573	812	1,124	1,603	2,219	1,989	1,473	562	129	-
Male	214	321	389	503	701	899	1,138	871	512	129	24	-
Female	70	155	184	309	423	704	1,081	1,118	961	433	105	-
White, both sexes	211	387	480	687	996	1,442	2,037	1,856	1,345	521	120	-
Male	165	274	329	433	635	826	1,068	827	471	119	22	-
Female	46	113	151	254	361	616	969	1,029	874	402	98	-
Black, both sexes	70	83	83	111	116	147	165	112	121	41	9	-
Male	46	45	54	60	58	63	65	39	40	10	2	-
Female	24	38	29	51	58	84	100	73	81	31	7	-
American Indian 1/, both sexes	-	1	2	-	-	-	1	1	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	8	14	12	14	16	20	7	-	-	-
Male	3	2	4	10	8	10	5	5	1	-	-	-
Female	-	3	4	4	4	4	11	15	6	-	-	-
Utah												
All races, both sexes	35	41	54	48	57	107	152	135	107	42	5	-
Male	28	33	37	33	35	61	100	68	40	17	-	-
Female	7	8	17	15	22	46	52	67	67	25	5	-
White, both sexes	35	37	51	46	56	102	147	134	105	41	5	-
Male	28	30	35	31	35	59	96	67	39	17	-	-
Female	7	7	16	15	21	43	51	67	66	24	5	-
Black, both sexes	-	-	1	1	-	1	1	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	2	1	-	-	-	1	1	-	-	-	-
Male	-	2	-	-	-	-	1	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	1	1	4	3	-	2	1	-	-
Male	-	1	1	1	-	2	3	-	1	-	-	-
Female	-	1	-	-	1	2	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Vermont												
All races, both sexes	7	8	16	16	34	49	66	75	61	34	9	-
Male	2	8	15	9	18	24	36	40	21	8	-	-
Female	5	-	1	7	16	25	30	35	40	26	9	-
White, both sexes	7	8	16	16	33	49	66	75	61	34	9	-
Male	2	8	15	9	18	24	36	40	21	8	-	-
Female	5	-	1	7	15	25	30	35	40	26	9	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	121	157	268	289	372	539	699	623	431	170	38	-
Male	91	122	181	203	232	323	352	283	143	30	1	-
Female	30	35	87	86	140	216	347	340	288	140	37	-
White, both sexes	81	106	197	222	291	453	594	541	371	147	31	-
Male	62	84	136	166	191	280	311	254	127	29	1	-
Female	19	22	61	56	100	173	283	287	244	118	30	-
Black, both sexes	39	49	65	62	73	77	99	76	59	23	7	-
Male	28	37	42	33	35	38	38	28	15	1	-	-
Female	11	12	23	29	38	39	61	48	44	22	7	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	5	5	8	9	5	6	1	-	-	-
Male	1	1	2	4	6	5	3	1	1	-	-	-
Female	-	1	3	1	2	4	2	5	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Washington												
All races, both sexes	69	108	182	233	318	501	718	735	551	222	55	-
Male	52	86	142	163	220	332	421	361	219	56	11	-
Female	17	22	40	70	98	169	297	374	332	166	44	-
White, both sexes	61	97	166	221	299	479	687	707	534	217	55	-
Male	44	77	131	155	207	320	401	345	214	54	11	-
Female	17	20	35	66	92	159	286	362	320	163	44	-
Black, both sexes	3	5	7	5	10	7	10	9	3	1	-	-
Male	3	4	4	4	6	4	7	5	1	-	-	-
Female	-	1	3	1	4	3	3	4	2	1	-	-
American Indian 1/, both sexes	2	4	6	3	3	4	2	3	4	1	-	-
Male	2	3	4	2	3	3	1	1	-	-	-	-
Female	-	1	2	1	-	1	1	2	4	1	-	-
Asian or Pacific Islander 2/, both sexes	3	2	3	4	6	11	19	16	10	3	-	-
Male	3	2	3	2	4	5	12	10	4	2	-	-
Female	-	-	-	2	2	6	7	6	6	1	-	-
West Virginia												
All races, both sexes	57	104	114	113	169	280	353	321	234	75	19	-
Male	44	76	89	71	113	151	170	135	68	23	6	-
Female	13	28	25	42	56	129	183	186	166	52	13	-
White, both sexes	53	104	111	111	164	274	348	314	220	74	19	-
Male	41	76	86	70	109	148	169	133	65	23	6	-
Female	12	28	25	41	55	126	179	181	155	51	13	-
Black, both sexes	4	-	3	2	5	6	5	7	14	1	-	-
Male	3	-	3	1	4	3	1	2	3	-	-	-
Female	1	-	-	1	1	3	4	5	11	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)												
Wisconsin												
All races, both sexes	65	101	154	177	277	446	613	740	573	256	49	-
Male	51	76	111	132	186	285	366	354	193	63	8	-
Female	14	25	43	45	91	161	247	386	380	193	41	-
White, both sexes	54	88	135	169	257	430	603	724	563	255	47	-
Male	43	67	98	127	174	278	359	347	191	62	8	-
Female	11	21	37	42	83	152	244	377	372	193	39	-
Black, both sexes	9	12	14	6	15	15	9	10	8	1	2	-
Male	7	8	8	3	9	7	6	4	2	1	-	-
Female	2	4	6	3	6	8	3	6	6	-	2	-
American Indian 1/, both sexes	1	1	4	2	2	-	-	2	2	-	-	-
Male	1	1	4	2	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	3	1	1	4	-	-	-	-
Male	-	-	1	-	1	-	1	3	-	-	-	-
Female	1	-	-	-	2	1	-	1	-	-	-	-
Wyoming												
All races, both sexes	6	9	14	21	29	43	47	54	27	12	5	-
Male	6	6	10	14	15	32	28	28	13	-	-	-
Female	-	3	4	7	14	11	19	26	14	12	5	-
White, both sexes	6	8	14	21	28	43	47	54	27	12	5	-
Male	6	6	10	14	15	32	28	28	13	-	-	-
Female	-	2	4	7	13	11	19	26	14	12	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
United States												
All races, both sexes	170,585	342	143	96	135	357	561	767	1,080	1,639	2,656	4,023
Male	75,094	188	77	44	74	244	370	499	677	1,016	1,679	2,513
Female	95,491	154	66	52	61	113	191	268	403	623	977	1,510
White, both sexes	147,474	219	87	54	87	239	373	498	653	1,071	1,846	2,803
Male	64,180	118	53	27	50	158	250	340	420	689	1,195	1,803
Female	83,294	101	34	27	37	81	123	158	233	382	651	1,000
Black, both sexes	20,093	101	44	34	40	107	163	246	383	514	744	1,117
Male	9,438	60	18	14	22	79	105	145	231	297	438	647
Female	10,655	41	26	20	18	28	58	101	152	217	306	470
American Indian 1/, both sexes	712	5	1	-	2	4	8	8	13	18	25	43
Male	361	1	1	-	-	3	4	5	10	9	18	25
Female	351	4	-	-	2	1	4	3	3	9	7	18
Asian or Pacific Islander 2/, both sexes	2,306	17	11	8	6	7	17	15	31	36	41	60
Male	1,115	9	5	3	2	4	11	9	16	21	28	38
Female	1,191	8	6	5	4	3	6	6	15	15	13	22
Alabama												
All races, both sexes	6,336	12	3	1	2	8	18	32	49	69	119	187
Male	2,871	5	-	1	1	6	13	21	33	47	78	127
Female	3,465	7	3	-	1	2	5	11	16	22	41	60
White, both sexes	4,820	6	2	1	-	4	7	20	25	44	82	116
Male	2,166	3	-	1	-	2	6	14	16	31	55	78
Female	2,654	3	2	-	-	2	1	6	9	13	27	38
Black, both sexes	1,503	6	1	-	2	4	11	12	24	23	37	70
Male	697	2	-	-	1	4	7	7	17	16	23	48
Female	806	4	1	-	1	-	4	5	7	7	14	22
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Alaska												
All races, both sexes	193	1	-	-	1	1	4	2	1	2	6	5
Male	102	-	-	-	1	1	3	1	-	2	3	2
Female	91	1	-	-	-	-	1	1	1	-	3	3
White, both sexes	135	-	-	-	1	-	2	1	-	2	4	2
Male	76	-	-	-	1	-	2	1	-	2	2	1
Female	59	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	41	1	-	-	-	-	1	1	1	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	27	1	-	-	-	-	1	1	1	-	1	1
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	1	1	-	-	-	-	2
Male	6	-	-	-	-	1	1	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
Arizona												
All races, both sexes	2,179	8	1	1	2	6	7	12	16	24	52	51
Male	1,047	2	1	1	1	4	5	6	9	17	33	31
Female	1,132	6	-	-	1	2	2	6	7	7	19	20
White, both sexes	2,045	5	1	1	2	6	5	7	10	18	47	41
Male	975	1	1	1	1	4	4	4	6	13	30	25
Female	1,070	4	-	-	1	2	1	3	4	5	17	16
Black, both sexes	46	-	-	-	-	-	1	3	4	3	4	5
Male	22	-	-	-	-	-	1	1	2	1	2	2
Female	24	-	-	-	-	-	-	2	2	2	2	3
American Indian 1/, both sexes	68	1	-	-	-	-	1	2	1	3	1	5
Male	43	1	-	-	-	-	-	1	1	3	1	4
Female	25	-	-	-	-	-	1	1	-	-	-	1
Asian or Pacific Islander 2/, both sexes	20	2	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	2	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Arkansas												
All races, both sexes	1,968	5	1	3	-	5	8	8	8	24	33	41
Male	879	2	-	1	-	5	7	5	5	13	21	24
Female	1,089	3	1	2	-	-	1	3	3	11	12	17
White, both sexes	1,692	4	1	3	-	4	3	2	7	13	20	29
Male	747	2	-	1	-	4	3	2	5	5	12	17
Female	945	2	1	2	-	-	-	-	2	8	8	12
Black, both sexes	273	1	-	-	-	1	5	6	1	11	13	12
Male	131	-	-	-	-	1	4	3	-	8	9	7
Female	142	1	-	-	-	-	1	3	1	3	4	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	12,703	31	15	9	15	31	53	83	99	154	240	336
Male	5,817	19	10	2	9	24	35	60	66	106	170	227
Female	6,886	12	5	7	6	7	18	23	33	48	70	109
White, both sexes	11,032	25	10	4	10	21	39	57	72	102	184	249
Male	5,021	14	7	-	5	17	26	46	50	76	135	174
Female	6,011	11	3	4	5	4	13	11	22	26	49	75
Black, both sexes	969	3	2	-	2	6	8	20	19	38	45	73
Male	481	2	-	-	2	4	5	11	11	21	27	41
Female	488	1	2	-	-	2	3	9	8	17	18	32
American Indian 1/, both sexes	54	-	-	-	-	1	-	-	2	2	2	2
Male	27	-	-	-	-	1	-	-	2	2	1	2
Female	27	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	648	3	3	5	3	3	6	6	6	12	9	12
Male	288	3	3	2	2	2	4	3	3	7	7	10
Female	360	-	-	3	1	1	2	3	3	5	2	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Colorado												
All races, both sexes	1,924	5	-	2	1	7	3	2	15	29	34	46
Male	829	1	-	1	-	5	2	1	12	20	19	30
Female	1,095	4	-	1	1	2	1	1	3	9	15	16
White, both sexes	1,843	4	-	2	-	7	2	2	13	24	30	40
Male	787	1	-	1	-	5	1	1	10	18	17	27
Female	1,056	3	-	1	-	2	1	1	3	6	13	13
Black, both sexes	57	-	-	-	-	-	1	-	2	4	3	6
Male	27	-	-	-	-	-	1	-	2	1	1	3
Female	30	-	-	-	-	-	-	-	-	3	2	3
American Indian 1/, both sexes	10	-	-	-	1	-	-	-	-	1	1	-
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	3	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	1	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2,412	5	2	1	2	3	7	3	15	24	35	58
Male	1,077	1	1	1	1	2	4	2	10	20	22	46
Female	1,335	4	1	-	1	1	3	1	5	4	13	12
White, both sexes	2,242	5	2	-	-	2	7	2	9	23	31	50
Male	992	1	1	-	-	2	4	1	7	19	19	39
Female	1,250	4	1	-	-	-	3	1	2	4	12	11
Black, both sexes	158	-	-	1	2	1	-	1	6	1	4	7
Male	82	-	-	1	1	-	-	1	3	1	3	6
Female	76	-	-	-	1	1	-	-	3	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Delaware												
All races, both sexes	477	2	1	-	-	1	-	2	3	1	12	9
Male	210	1	-	-	-	1	-	1	3	1	9	7
Female	267	1	1	-	-	-	-	1	-	-	3	2
White, both sexes	389	-	-	-	-	-	-	2	1	1	7	5
Male	168	-	-	-	-	-	-	1	1	1	5	4
Female	221	-	-	-	-	-	-	1	-	-	2	1
Black, both sexes	78	2	1	-	-	1	-	-	2	-	5	3
Male	37	1	-	-	-	1	-	-	2	-	4	2
Female	41	1	1	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	202	-	-	-	1	1	1	-	3	2	5	10
Male	77	-	-	-	1	1	-	-	3	1	2	3
Female	125	-	-	-	-	-	1	-	-	1	3	7
White, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	164	-	-	-	1	1	1	-	3	1	5	10
Male	61	-	-	-	1	1	-	-	3	1	2	3
Female	103	-	-	-	-	-	1	-	-	-	3	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Florida												
All races, both sexes	9,005	18	7	2	3	19	28	30	51	88	123	210
Male	4,233	11	4	-	2	16	19	21	28	51	77	124
Female	4,772	7	3	2	1	3	9	9	23	37	46	86
White, both sexes	7,994	11	2	1	2	11	18	15	27	60	86	138
Male	3,759	6	1	-	2	9	10	9	13	38	60	85
Female	4,235	5	1	1	-	2	8	6	14	22	26	53
Black, both sexes	958	6	2	1	1	7	9	14	23	27	36	69
Male	444	4	1	-	-	6	8	11	15	13	16	38
Female	514	2	1	1	1	1	1	3	8	14	20	31
American Indian 1/, both sexes	6	-	1	-	-	-	-	1	-	-	1	-
Male	4	-	1	-	-	-	-	1	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	47	1	2	-	-	1	1	-	1	1	-	3
Male	26	1	1	-	-	1	1	-	-	-	-	1
Female	21	-	1	-	-	-	-	-	1	1	-	2
Georgia												
All races, both sexes	6,856	15	8	3	8	18	21	39	53	80	126	191
Male	3,053	11	5	1	3	11	13	22	32	42	78	114
Female	3,803	4	3	2	5	7	8	17	21	38	48	77
White, both sexes	5,043	6	1	2	5	5	4	19	24	30	61	102
Male	2,209	3	1	1	2	3	3	13	16	15	38	68
Female	2,834	3	-	1	3	2	1	6	8	15	23	34
Black, both sexes	1,754	8	6	1	3	13	15	19	28	48	60	88
Male	812	7	4	-	1	8	8	9	15	27	38	45
Female	942	1	2	1	2	5	7	10	13	21	22	43
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	44	1	1	-	-	-	2	1	1	1	5	1
Male	26	1	-	-	-	-	2	-	1	-	2	1
Female	18	-	1	-	-	-	-	1	-	1	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Hawaii												
All races, both sexes	1,009	2	-	1	1	-	1	4	6	9	22	26
Male	523	1	-	-	1	-	1	1	5	5	16	13
Female	486	1	-	1	-	-	-	3	1	4	6	13
White, both sexes	257	-	-	-	-	-	-	2	1	-	5	4
Male	126	-	-	-	-	-	-	1	1	-	4	1
Female	131	-	-	-	-	-	-	1	-	-	1	3
Black, both sexes	9	-	-	-	1	-	-	-	-	-	1	1
Male	7	-	-	-	1	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	740	2	-	1	-	-	1	2	5	9	16	21
Male	389	1	-	-	-	-	1	-	4	5	11	11
Female	351	1	-	1	-	-	-	2	1	4	5	10
Idaho												
All races, both sexes	737	-	1	-	1	1	1	3	3	9	8	14
Male	309	-	1	-	-	1	-	2	2	4	7	10
Female	428	-	-	-	1	-	1	1	1	5	1	4
White, both sexes	730	-	1	-	1	1	1	3	3	9	8	14
Male	306	-	1	-	-	1	-	2	2	4	7	10
Female	424	-	-	-	1	-	1	1	1	5	1	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Illinois												
All races, both sexes	7,728	15	6	2	8	25	31	46	54	81	137	189
Male	3,311	9	2	-	4	12	18	35	33	56	96	119
Female	4,417	6	4	2	4	13	13	11	21	25	41	70
White, both sexes	6,615	9	2	1	4	16	21	29	30	55	93	113
Male	2,769	4	1	-	2	7	13	23	20	42	66	77
Female	3,846	5	1	1	2	9	8	6	10	13	27	36
Black, both sexes	1,037	6	4	1	3	9	9	17	23	25	43	75
Male	504	5	1	-	2	5	5	12	12	13	29	41
Female	533	1	3	1	1	4	4	5	11	12	14	34
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	72	-	-	-	1	-	1	-	1	-	1	1
Male	37	-	-	-	-	-	-	-	1	-	1	1
Female	35	-	-	-	1	-	1	-	-	-	-	-
Indiana												
All races, both sexes	4,711	10	4	3	3	15	16	17	39	46	66	121
Male	2,005	5	1	2	1	9	8	12	23	25	44	63
Female	2,706	5	3	1	2	6	8	5	16	21	22	58
White, both sexes	4,365	8	2	1	2	13	16	14	32	41	50	93
Male	1,849	4	1	1	-	8	8	10	16	23	34	45
Female	2,516	4	1	-	2	5	8	4	16	18	16	48
Black, both sexes	331	2	1	2	1	2	-	3	6	5	16	27
Male	150	1	-	1	1	1	-	2	6	2	10	18
Female	181	1	1	1	-	1	-	1	-	3	6	9
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	13	-	1	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Iowa												
All races, both sexes	1,369	-	2	1	1	3	5	5	7	18	16	32
Male	565	-	2	1	1	2	2	4	2	12	10	17
Female	804	-	-	-	-	1	3	1	5	6	6	15
White, both sexes	1,338	-	2	1	1	3	5	4	5	16	14	29
Male	551	-	2	1	1	2	2	3	2	11	8	16
Female	787	-	-	-	-	1	3	1	3	5	6	13
Black, both sexes	28	-	-	-	-	-	-	1	1	2	2	2
Male	13	-	-	-	-	-	-	1	-	1	2	1
Female	15	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Kansas												
All races, both sexes	1,975	3	-	2	-	4	7	8	7	15	20	41
Male	813	2	-	1	-	3	4	6	5	9	14	26
Female	1,162	1	-	1	-	1	3	2	2	6	6	15
White, both sexes	1,848	2	-	2	-	2	5	8	5	11	17	29
Male	748	1	-	1	-	1	4	6	5	7	13	17
Female	1,100	1	-	1	-	1	1	2	-	4	4	12
Black, both sexes	103	1	-	-	-	2	2	-	2	2	3	10
Male	52	1	-	-	-	2	-	-	-	1	1	9
Female	51	-	-	-	-	-	2	-	2	1	2	1
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	1	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	2
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Kentucky												
All races, both sexes	3,393	12	2	1	3	2	8	12	17	28	51	68
Male	1,461	7	1	-	2	2	5	6	12	15	33	43
Female	1,932	5	1	1	1	-	3	6	5	13	18	25
White, both sexes	3,156	5	2	1	3	2	7	9	14	23	43	55
Male	1,352	3	1	-	2	2	5	5	9	10	27	37
Female	1,804	2	1	1	1	-	2	4	5	13	16	18
Black, both sexes	227	5	-	-	-	-	1	3	3	5	8	13
Male	102	3	-	-	-	-	-	1	3	5	6	6
Female	125	2	-	-	-	-	1	2	-	-	2	7
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	2	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3,495	7	3	4	3	8	18	22	17	36	60	85
Male	1,533	3	3	2	2	6	11	15	8	19	36	47
Female	1,962	4	-	2	1	2	7	7	9	17	24	38
White, both sexes	2,519	5	2	1	2	4	11	7	9	16	33	43
Male	1,110	2	2	-	2	2	6	6	7	9	22	27
Female	1,409	3	-	1	-	2	5	1	2	7	11	16
Black, both sexes	962	2	1	3	1	4	7	14	8	19	26	42
Male	419	1	1	2	-	4	5	8	1	9	14	20
Female	543	1	-	1	1	-	2	6	7	10	12	22
American Indian 1/, both sexes	9	-	-	-	-	-	-	1	-	-	1	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Maine												
All races, both sexes	905	1	2	-	-	2	3	4	3	6	12	17
Male	397	-	-	-	-	1	2	2	2	5	5	13
Female	508	1	2	-	-	1	1	2	1	1	7	4
White, both sexes	900	1	1	-	-	2	3	4	3	6	11	17
Male	394	-	-	-	-	1	2	2	2	5	4	13
Female	506	1	1	-	-	1	1	2	1	1	7	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,548	9	6	1	2	9	15	20	32	27	65	90
Male	1,148	6	2	-	1	8	10	18	18	15	41	57
Female	1,400	3	4	1	1	1	5	2	14	12	24	33
White, both sexes	1,870	3	1	-	2	5	10	10	10	11	31	49
Male	828	3	1	-	1	4	7	9	5	6	22	32
Female	1,042	-	-	-	1	1	3	1	5	5	9	17
Black, both sexes	640	5	4	1	-	4	4	10	21	15	34	38
Male	301	2	-	-	-	4	3	9	13	8	19	23
Female	339	3	4	1	-	-	1	1	8	7	15	15
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	1	1	-	-	-	1	-	1	-	-	3
Male	18	1	1	-	-	-	-	-	-	-	-	2
Female	18	-	-	-	-	-	1	-	1	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Massachusetts												
All races, both sexes	4,618	6	3	1	2	3	10	15	10	25	49	70
Male	1,923	4	1	1	2	3	8	11	4	16	32	51
Female	2,695	2	2	-	-	-	2	4	6	9	17	19
White, both sexes	4,394	5	2	1	1	3	8	13	7	19	42	64
Male	1,818	4	1	1	1	3	6	10	2	13	26	47
Female	2,576	1	1	-	-	-	2	3	5	6	16	17
Black, both sexes	187	1	1	-	1	-	2	2	2	5	6	5
Male	84	-	-	-	1	-	2	1	1	2	5	3
Female	103	1	1	-	-	-	-	1	1	3	1	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	-	1	1	1	1
Male	21	-	-	-	-	-	-	-	1	1	1	1
Female	11	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	6,112	10	4	9	5	14	18	25	40	60	92	146
Male	2,623	5	2	3	4	11	12	15	29	38	62	91
Female	3,489	5	2	6	1	3	6	10	11	22	30	55
White, both sexes	5,340	7	2	4	3	11	12	17	21	31	64	96
Male	2,252	4	1	3	3	8	8	12	17	21	48	64
Female	3,088	3	1	1	-	3	4	5	4	10	16	32
Black, both sexes	723	3	2	5	2	3	6	8	17	28	26	47
Male	351	1	1	-	1	3	4	3	10	17	13	25
Female	372	2	1	5	1	-	2	5	7	11	13	22
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	1	1	-	1
Male	13	-	-	-	-	-	-	-	1	-	-	1
Female	12	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	1	-	2	2
Male	7	-	-	-	-	-	-	-	1	-	1	1
Female	17	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Minnesota												
All races, both sexes	2,842	1	1	1	1	3	6	9	6	10	28	46
Male	1,244	-	1	-	-	2	5	6	5	6	12	29
Female	1,598	1	-	1	1	1	1	3	1	4	16	17
White, both sexes	2,767	1	1	1	1	3	6	7	6	7	23	39
Male	1,201	-	1	-	-	2	5	4	5	5	9	24
Female	1,566	1	-	1	1	1	1	3	1	2	14	15
Black, both sexes	41	-	-	-	-	-	-	2	-	2	4	5
Male	27	-	-	-	-	-	-	2	-	1	2	4
Female	14	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	22	-	-	-	-	-	-	-	-	1	1	2
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	3,090	2	2	2	1	4	14	11	22	30	37	79
Male	1,287	1	1	2	-	3	4	3	13	15	19	44
Female	1,803	1	1	-	1	1	10	8	9	15	18	35
White, both sexes	2,064	-	1	-	-	-	2	3	7	9	15	35
Male	825	-	1	-	-	-	1	-	3	2	7	18
Female	1,239	-	-	-	-	-	1	3	4	7	8	17
Black, both sexes	1,017	2	1	2	1	4	12	8	15	21	22	44
Male	457	1	-	2	-	3	3	3	10	13	12	26
Female	560	1	1	-	1	1	9	5	5	8	10	18
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Missouri												
All races, both sexes	4,128	10	7	3	4	10	8	17	26	42	51	89
Male	1,790	6	4	1	2	6	7	14	16	25	30	52
Female	2,338	4	3	2	2	4	1	3	10	17	21	37
White, both sexes	3,749	9	6	3	4	5	7	11	21	33	36	67
Male	1,614	6	4	1	2	1	6	10	13	19	22	41
Female	2,135	3	2	2	2	4	1	1	8	14	14	26
Black, both sexes	360	1	-	-	-	5	1	6	5	9	15	22
Male	169	-	-	-	-	5	1	4	3	6	8	11
Female	191	1	-	-	-	-	-	2	2	3	7	11
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	1	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	697	2	2	-	2	1	-	1	5	4	7	12
Male	312	1	2	-	2	-	-	1	1	2	5	9
Female	385	1	-	-	-	1	-	-	4	2	2	3
White, both sexes	667	2	2	-	2	1	-	1	5	3	4	9
Male	295	1	2	-	2	-	-	1	1	2	2	6
Female	372	1	-	-	-	1	-	-	4	1	2	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	-	1	3	3
Male	15	-	-	-	-	-	-	-	-	-	3	3
Female	10	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Nebraska												
All races, both sexes	1,421	2	-	-	-	1	5	1	5	3	8	21
Male	588	1	-	-	-	1	1	-	2	2	5	12
Female	833	1	-	-	-	-	4	1	3	1	3	9
White, both sexes	1,372	1	-	-	-	1	3	1	2	2	8	19
Male	564	-	-	-	-	1	-	-	2	1	5	12
Female	808	1	-	-	-	-	3	1	-	1	3	7
Black, both sexes	37	1	-	-	-	-	1	-	2	1	-	2
Male	17	1	-	-	-	-	1	-	-	1	-	-
Female	20	-	-	-	-	-	-	-	2	-	-	2
American Indian 1/, both sexes	7	-	-	-	-	-	1	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,908	3	-	-	1	4	8	18	7	24	53	64
Male	1,005	1	-	-	1	3	6	15	6	19	31	43
Female	903	2	-	-	-	1	2	3	1	5	22	21
White, both sexes	1,658	2	-	-	1	3	6	11	5	16	37	50
Male	868	1	-	-	1	3	4	10	4	13	20	35
Female	790	1	-	-	-	-	2	1	1	3	17	15
Black, both sexes	168	1	-	-	-	-	1	6	1	6	12	13
Male	90	-	-	-	-	-	1	4	1	4	8	7
Female	78	1	-	-	-	-	-	2	-	2	4	6
American Indian 1/, both sexes	20	-	-	-	-	-	1	-	1	-	1	-
Male	13	-	-	-	-	-	1	-	1	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	62	-	-	-	-	1	-	1	-	2	3	1
Male	34	-	-	-	-	-	-	1	-	2	2	1
Female	28	-	-	-	-	1	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
New Hampshire												
All races, both sexes	639	-	-	-	-	2	1	4	2	7	8	8
Male	267	-	-	-	-	-	1	4	2	5	5	6
Female	372	-	-	-	-	2	-	-	-	2	3	2
White, both sexes	636	-	-	-	-	2	1	4	2	7	8	8
Male	265	-	-	-	-	-	1	4	2	5	5	6
Female	371	-	-	-	-	2	-	-	-	2	3	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4,946	8	3	-	4	14	18	24	30	41	72	123
Male	2,193	5	2	-	3	8	15	17	23	28	35	82
Female	2,753	3	1	-	1	6	3	7	7	13	37	41
White, both sexes	4,283	7	2	-	1	11	15	15	13	24	43	81
Male	1,874	5	1	-	1	7	12	11	13	17	21	57
Female	2,409	2	1	-	-	4	3	4	-	7	22	24
Black, both sexes	595	-	1	-	3	3	3	8	15	16	28	38
Male	282	-	1	-	2	1	3	5	8	11	13	23
Female	313	-	-	-	1	2	-	3	7	5	15	15
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	63	1	-	-	-	-	-	1	2	1	1	3
Male	33	-	-	-	-	-	-	1	2	-	1	1
Female	30	1	-	-	-	-	-	-	-	1	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
New Mexico												
All races, both sexes	878	3	-	-	1	-	-	6	3	7	17	24
Male	397	1	-	-	1	-	-	1	2	5	8	14
Female	481	2	-	-	-	-	-	5	1	2	9	10
White, both sexes	808	1	-	-	1	-	-	5	2	6	13	21
Male	367	1	-	-	1	-	-	1	1	4	5	13
Female	441	-	-	-	-	-	-	4	1	2	8	8
Black, both sexes	13	-	-	-	-	-	-	1	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	56	2	-	-	-	-	-	-	1	1	4	2
Male	27	-	-	-	-	-	-	-	1	1	3	1
Female	29	2	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	8,005	21	8	4	4	11	21	36	42	64	86	157
Male	3,444	12	3	3	2	9	16	25	22	35	52	103
Female	4,561	9	5	1	2	2	5	11	20	29	34	54
White, both sexes	7,148	12	5	2	2	5	13	22	26	41	57	119
Male	3,072	8	2	2	1	5	9	15	14	22	34	80
Female	4,076	4	3	-	1	-	4	7	12	19	23	39
Black, both sexes	779	8	2	1	1	5	8	13	15	23	29	38
Male	341	4	1	1	1	4	7	9	8	13	18	23
Female	438	4	1	-	-	1	1	4	7	10	11	15
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	74	1	1	1	1	1	-	1	1	-	-	-
Male	29	-	-	-	-	-	-	1	-	-	-	-
Female	45	1	1	1	1	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
North Carolina												
All races, both sexes	5,085	11	9	8	3	13	20	18	37	56	80	120
Male	2,198	5	6	6	2	3	17	8	22	31	49	84
Female	2,887	6	3	2	1	10	3	10	15	25	31	36
White, both sexes	3,915	3	5	2	1	11	9	9	15	28	41	58
Male	1,655	-	3	2	1	3	9	4	11	15	25	43
Female	2,260	3	2	-	-	8	-	5	4	13	16	15
Black, both sexes	1,110	8	4	6	2	2	10	7	20	28	37	60
Male	516	5	3	4	1	-	7	3	10	16	23	39
Female	594	3	1	2	1	2	3	4	10	12	14	21
American Indian 1/, both sexes	42	-	-	-	-	-	-	1	-	-	2	2
Male	20	-	-	-	-	-	-	-	-	-	1	2
Female	22	-	-	-	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	1	1	2	-	-	-
Male	7	-	-	-	-	-	1	1	1	-	-	-
Female	11	-	-	-	-	-	-	-	1	-	-	-
North Dakota												
All races, both sexes	317	-	-	-	-	1	2	1	2	1	7	5
Male	121	-	-	-	-	1	1	1	1	1	5	2
Female	196	-	-	-	-	-	1	-	1	-	2	3
White, both sexes	311	-	-	-	-	1	2	1	2	1	4	4
Male	118	-	-	-	-	1	1	1	1	1	3	2
Female	193	-	-	-	-	-	1	-	1	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	2	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Ohio												
All races, both sexes	6,738	16	11	5	7	9	19	31	46	79	100	156
Male	2,925	9	9	2	6	7	12	16	28	45	61	87
Female	3,813	7	2	3	1	2	7	15	18	34	39	69
White, both sexes	6,029	8	9	4	5	7	14	22	31	56	75	126
Male	2,593	6	8	2	5	5	10	12	19	34	49	69
Female	3,436	2	1	2	-	2	4	10	12	22	26	57
Black, both sexes	695	8	2	1	2	2	5	9	15	23	25	30
Male	327	3	1	-	1	2	2	4	9	11	12	18
Female	368	5	1	1	1	-	3	5	6	12	13	12
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2,366	5	4	1	1	10	11	8	18	17	36	58
Male	969	3	2	1	-	5	7	5	10	10	25	34
Female	1,397	2	2	-	1	5	4	3	8	7	11	24
White, both sexes	2,092	2	4	1	1	7	9	5	14	11	27	38
Male	840	1	2	1	-	3	5	2	8	8	19	23
Female	1,252	1	2	-	1	4	4	3	6	3	8	15
Black, both sexes	155	2	-	-	-	2	-	1	2	4	7	11
Male	70	2	-	-	-	1	-	1	1	2	4	8
Female	85	-	-	-	-	1	-	-	1	2	3	3
American Indian 1/, both sexes	115	1	-	-	-	1	2	2	2	2	2	9
Male	58	-	-	-	-	1	2	2	1	-	2	3
Female	57	1	-	-	-	-	-	-	1	2	-	6
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Oregon												
All races, both sexes	2,260	1	-	1	4	4	7	5	10	18	20	31
Male	948	-	-	-	-	3	6	3	6	12	12	17
Female	1,312	1	-	1	4	1	1	2	4	6	8	14
White, both sexes	2,197	1	-	1	2	3	6	4	9	15	19	28
Male	920	-	-	-	-	2	5	2	6	11	11	14
Female	1,277	1	-	1	2	1	1	2	3	4	8	14
Black, both sexes	31	-	-	-	1	1	1	1	1	1	1	1
Male	10	-	-	-	-	1	1	1	-	1	1	1
Female	21	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	1	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	1	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	1	-	-	-	-	1	-	-
Pennsylvania												
All races, both sexes	11,025	16	2	5	4	16	25	27	38	68	140	221
Male	4,714	8	1	3	3	13	16	21	23	42	81	137
Female	6,311	8	1	2	1	3	9	6	15	26	59	84
White, both sexes	10,062	14	-	5	4	11	18	19	28	46	98	175
Male	4,269	6	-	3	3	9	12	15	18	29	58	111
Female	5,793	8	-	2	1	2	6	4	10	17	40	64
Black, both sexes	928	1	2	-	-	5	7	7	9	21	41	45
Male	428	1	1	-	-	4	4	5	4	12	22	26
Female	500	-	1	-	-	1	3	2	5	9	19	19
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	1	-	-	-	-	-	1	-	1	1	1
Male	15	1	-	-	-	-	-	1	-	1	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Rhode Island												
All races, both sexes	431	3	2	-	3	-	1	3	4	8	11	8
Male	179	1	1	-	2	-	-	2	3	4	7	4
Female	252	2	1	-	1	-	1	1	1	4	4	4
White, both sexes	409	2	2	-	3	-	1	2	3	7	9	7
Male	165	1	1	-	2	-	-	2	2	4	6	3
Female	244	1	1	-	1	-	1	-	1	3	3	4
Black, both sexes	21	1	-	-	-	-	-	1	1	1	2	1
Male	13	-	-	-	-	-	-	-	1	-	1	1
Female	8	1	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3,258	9	2	2	5	8	9	19	34	43	74	97
Male	1,517	3	1	1	3	5	5	12	25	23	44	59
Female	1,741	6	1	1	2	3	4	7	9	20	30	38
White, both sexes	2,299	6	-	1	2	3	5	11	11	22	36	57
Male	1,036	2	-	-	1	3	3	6	8	10	23	35
Female	1,263	4	-	1	1	-	2	5	3	12	13	22
Black, both sexes	947	3	2	1	3	5	4	8	23	21	38	40
Male	474	1	1	1	2	2	2	6	17	13	21	24
Female	473	2	1	-	1	3	2	2	6	8	17	16
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
South Dakota												
All races, both sexes	299	1	-	-	1	-	-	1	4	2	4	6
Male	134	-	-	-	-	-	-	1	1	1	-	4
Female	165	1	-	-	1	-	-	-	3	1	4	2
White, both sexes	276	1	-	-	-	-	-	1	4	2	4	3
Male	121	-	-	-	-	-	-	1	1	1	-	2
Female	155	1	-	-	-	-	-	-	3	1	4	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	22	-	-	-	1	-	-	-	-	-	-	3
Male	12	-	-	-	-	-	-	-	-	-	-	2
Female	10	-	-	-	1	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3,040	8	2	2	1	10	12	27	28	44	69	113
Male	1,373	6	-	-	-	5	8	17	20	26	46	69
Female	1,667	2	2	2	1	5	4	10	8	18	23	44
White, both sexes	2,524	6	2	-	1	6	8	15	17	33	51	83
Male	1,111	5	-	-	-	3	5	9	12	22	32	48
Female	1,413	1	2	-	1	3	3	6	5	11	19	35
Black, both sexes	505	2	-	2	-	4	4	12	10	11	18	30
Male	259	1	-	-	-	2	3	8	8	4	14	21
Female	246	1	-	2	-	2	1	4	2	7	4	9
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Texas												
All races, both sexes	10,104	27	10	5	13	32	45	61	88	110	184	268
Male	4,492	18	6	2	8	25	30	37	57	79	118	159
Female	5,612	9	4	3	5	7	15	24	31	31	66	109
White, both sexes	8,743	20	7	4	8	21	29	48	54	71	134	192
Male	3,834	11	5	1	6	15	18	31	36	51	89	118
Female	4,909	9	2	3	2	6	11	17	18	20	45	74
Black, both sexes	1,273	7	3	-	5	11	15	13	32	37	49	73
Male	617	7	1	-	2	10	12	6	20	26	29	40
Female	656	-	2	-	3	1	3	7	12	11	20	33
American Indian 1/, both sexes	8	-	-	-	-	-	1	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	80	-	-	1	-	-	-	-	1	2	1	3
Male	38	-	-	1	-	-	-	-	-	2	-	1
Female	42	-	-	-	-	-	-	-	1	-	1	2
Utah												
All races, both sexes	1,248	-	1	-	-	1	4	4	8	12	13	17
Male	494	-	-	-	-	1	4	-	4	8	7	12
Female	754	-	1	-	-	-	-	4	4	4	6	5
White, both sexes	1,222	-	1	-	-	1	4	4	8	12	11	15
Male	482	-	-	-	-	1	4	-	4	8	5	10
Female	740	-	1	-	-	-	-	4	4	4	6	5
Black, both sexes	12	-	-	-	-	-	-	-	-	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	2	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Vermont												
All races, both sexes	299	-	-	-	-	1	2	-	1	2	3	3
Male	131	-	-	-	-	1	1	-	1	1	2	2
Female	168	-	-	-	-	-	1	-	-	1	1	1
White, both sexes	299	-	-	-	-	1	2	-	1	2	3	3
Male	131	-	-	-	-	1	1	-	1	1	2	2
Female	168	-	-	-	-	-	1	-	-	1	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4,821	10	3	8	4	6	19	22	31	39	71	105
Male	2,130	7	1	5	1	3	13	12	18	22	45	68
Female	2,691	3	2	3	3	3	6	10	13	17	26	37
White, both sexes	3,816	6	1	3	3	6	11	13	12	20	45	58
Male	1,674	4	-	3	1	3	8	8	7	10	30	39
Female	2,142	2	1	-	2	3	3	5	5	10	15	19
Black, both sexes	947	4	2	5	1	-	7	9	16	19	26	46
Male	427	3	1	2	-	-	4	4	9	12	15	28
Female	520	1	1	3	1	-	3	5	7	7	11	18
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	55	-	-	-	-	-	1	-	2	-	-	1
Male	27	-	-	-	-	-	1	-	1	-	-	1
Female	28	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Washington												
All races, both sexes	2,469	1	1	2	3	5	10	9	18	23	38	57
Male	1,104	-	-	-	1	3	6	4	9	15	22	31
Female	1,365	1	1	2	2	2	4	5	9	8	16	26
White, both sexes	2,312	-	1	1	2	3	6	8	15	19	32	50
Male	1,031	-	-	-	-	2	5	3	8	11	19	29
Female	1,281	-	1	1	2	1	1	5	7	8	13	21
Black, both sexes	74	-	-	1	1	-	1	-	1	3	4	4
Male	40	-	-	-	1	-	-	-	1	3	2	-
Female	34	-	-	1	-	-	1	-	-	-	2	4
American Indian 1/, both sexes	24	-	-	-	-	2	1	-	-	-	1	2
Male	12	-	-	-	-	1	1	-	-	-	-	1
Female	12	-	-	-	-	1	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	59	1	-	-	-	-	2	1	2	1	1	1
Male	21	-	-	-	-	-	-	1	-	1	1	1
Female	38	1	-	-	-	-	2	-	2	-	-	-
West Virginia												
All races, both sexes	1,493	-	2	-	1	-	3	4	5	5	16	26
Male	637	-	1	-	-	-	1	2	5	4	14	19
Female	856	-	1	-	1	-	2	2	-	1	2	7
White, both sexes	1,442	-	2	-	1	-	3	4	3	5	16	24
Male	616	-	1	-	-	-	1	2	3	4	14	18
Female	826	-	1	-	1	-	2	2	-	1	2	6
Black, both sexes	48	-	-	-	-	-	-	-	2	-	-	2
Male	20	-	-	-	-	-	-	-	2	-	-	1
Female	28	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other heart diseases (I26-I51)												
Wisconsin												
All races, both sexes	3,610	4	-	-	2	5	8	6	9	19	37	58
Male	1,560	4	-	-	-	3	6	5	7	9	28	42
Female	2,050	-	-	-	2	2	2	1	2	10	9	16
White, both sexes	3,461	3	-	-	2	5	7	5	6	14	31	48
Male	1,487	3	-	-	-	3	5	5	4	8	23	37
Female	1,974	-	-	-	2	2	2	-	2	6	8	11
Black, both sexes	115	1	-	-	-	-	1	1	3	5	5	8
Male	59	1	-	-	-	-	1	-	3	1	4	4
Female	56	-	-	-	-	-	-	1	-	4	1	4
American Indian 1/, both sexes	25	-	-	-	-	-	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	313	1	-	-	1	-	-	-	3	4	3	8
Male	135	1	-	-	-	-	-	-	1	2	3	4
Female	178	-	-	-	1	-	-	-	2	2	-	4
White, both sexes	310	1	-	-	1	-	-	-	3	4	3	8
Male	134	1	-	-	-	-	-	-	1	2	3	4
Female	176	-	-	-	1	-	-	-	2	2	-	4
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
United States												
All races, both sexes	5,097	6,394	7,467	8,936	12,827	19,611	28,255	32,000	25,325	10,543	2,325	6
Male	3,256	3,995	4,420	5,101	6,886	9,674	12,280	11,868	7,540	2,299	390	4
Female	1,841	2,399	3,047	3,835	5,941	9,937	15,975	20,132	17,785	8,244	1,935	2
White, both sexes	3,587	4,754	5,754	7,108	10,716	17,076	25,515	29,616	23,585	9,747	2,081	5
Male	2,367	3,014	3,474	4,105	5,792	8,528	11,215	11,043	7,077	2,118	340	4
Female	1,220	1,740	2,280	3,003	4,924	8,548	14,300	18,573	16,508	7,629	1,741	1
Black, both sexes	1,384	1,480	1,512	1,602	1,847	2,205	2,321	1,969	1,415	649	215	1
Male	818	879	831	860	949	999	875	654	349	129	39	-
Female	566	601	681	742	898	1,206	1,446	1,315	1,066	520	176	1
American Indian 1/, both sexes	45	50	50	66	72	75	76	66	52	28	5	-
Male	25	31	28	35	46	30	34	26	21	7	2	-
Female	20	19	22	31	26	45	42	40	31	21	3	-
Asian or Pacific Islander 2/, both sexes	81	110	151	160	192	255	343	349	273	119	24	-
Male	46	71	87	101	99	117	156	145	93	45	9	-
Female	35	39	64	59	93	138	187	204	180	74	15	-
Alabama												
All races, both sexes	263	335	417	423	557	772	998	945	780	280	66	-
Male	149	207	252	231	305	386	408	326	202	60	13	-
Female	114	128	165	192	252	386	590	619	578	220	53	-
White, both sexes	173	223	290	302	424	571	811	804	647	225	43	-
Male	105	144	181	167	235	298	337	270	168	48	7	-
Female	68	79	109	135	189	273	474	534	479	177	36	-
Black, both sexes	90	111	127	119	131	198	186	141	133	54	23	-
Male	44	62	71	63	70	85	70	56	34	11	6	-
Female	46	49	56	56	61	113	116	85	99	43	17	-
American Indian 1/, both sexes	-	1	-	1	1	2	-	-	-	-	-	-
Male	-	1	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	1	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Alaska												
All races, both sexes	11	3	13	14	18	27	27	27	21	8	1	-
Male	11	2	8	10	13	11	12	12	6	3	1	-
Female	-	1	5	4	5	16	15	15	15	5	-	-
White, both sexes	7	1	10	8	11	18	23	21	17	6	1	-
Male	7	1	5	7	9	9	11	9	5	3	1	-
Female	-	-	5	1	2	9	12	12	12	3	-	-
Black, both sexes	1	1	1	1	-	-	-	1	1	-	-	-
Male	1	1	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	3	-	1	5	5	9	3	4	3	2	-	-
Male	3	-	1	2	3	2	1	1	1	-	-	-
Female	-	-	-	3	2	7	2	3	2	2	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	2	-	1	1	-	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	-	-
Arizona												
All races, both sexes	69	76	100	131	175	254	360	398	304	118	14	-
Male	49	45	68	74	92	121	173	179	103	32	1	-
Female	20	31	32	57	83	133	187	219	201	86	13	-
White, both sexes	56	66	95	117	167	244	349	383	302	109	14	-
Male	42	40	66	66	86	118	165	170	101	30	1	-
Female	14	26	29	51	81	126	184	213	201	79	13	-
Black, both sexes	5	4	-	3	1	1	4	5	-	3	-	-
Male	3	2	-	1	1	-	3	3	-	-	-	-
Female	2	2	-	2	-	1	1	2	-	3	-	-
American Indian 1/, both sexes	8	5	5	10	5	6	5	5	1	4	-	-
Male	4	3	2	6	5	2	5	3	1	1	-	-
Female	4	2	3	4	-	4	-	2	-	3	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	2	3	2	5	1	2	-	-
Male	-	-	-	1	-	1	-	3	1	1	-	-
Female	-	1	-	-	2	2	2	2	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Arkansas												
All races, both sexes	62	77	95	127	149	208	324	371	294	106	19	-
Male	43	49	61	76	82	103	141	124	90	22	5	-
Female	19	28	34	51	67	105	183	247	204	84	14	-
White, both sexes	42	59	76	102	125	186	302	335	265	100	14	-
Male	26	37	52	63	72	94	132	113	82	21	4	-
Female	16	22	24	39	53	92	170	222	183	79	10	-
Black, both sexes	20	18	19	25	22	22	22	35	29	6	5	-
Male	17	12	9	13	9	9	9	11	8	1	1	-
Female	3	6	10	12	13	13	13	24	21	5	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
California												
All races, both sexes	386	455	487	585	809	1,282	2,126	2,459	2,016	841	191	-
Male	246	283	307	325	451	640	967	942	686	212	30	-
Female	140	172	180	260	358	642	1,159	1,517	1,330	629	161	-
White, both sexes	286	342	390	470	663	1,102	1,892	2,276	1,879	779	180	-
Male	189	212	249	263	383	565	863	884	642	193	28	-
Female	97	130	141	207	280	537	1,029	1,392	1,237	586	152	-
Black, both sexes	72	79	59	79	92	99	107	75	58	29	4	-
Male	42	48	36	41	47	46	57	20	15	5	-	-
Female	30	31	23	38	45	53	50	55	43	24	4	-
American Indian 1/, both sexes	3	5	2	6	2	9	9	6	1	2	-	-
Male	1	3	1	4	1	2	4	3	-	-	-	-
Female	2	2	1	2	1	7	5	3	1	2	-	-
Asian or Pacific Islander 2/, both sexes	25	29	36	30	52	72	118	102	78	31	7	-
Male	14	20	21	17	20	27	43	35	29	14	2	-
Female	11	9	15	13	32	45	75	67	49	17	5	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Colorado												
All races, both sexes	53	57	80	89	127	194	313	405	286	146	30	-
Male	33	36	35	56	62	86	147	157	89	29	8	-
Female	20	21	45	33	65	108	166	248	197	117	22	-
White, both sexes	47	53	77	84	118	183	305	397	283	143	29	-
Male	29	33	34	52	55	81	143	155	88	29	7	-
Female	18	20	43	32	63	102	162	242	195	114	22	-
Black, both sexes	6	1	2	4	5	6	5	7	2	2	1	-
Male	4	1	-	3	4	2	2	1	1	-	1	-
Female	2	-	2	1	1	4	3	6	1	2	-	-
American Indian 1/, both sexes	-	3	-	-	2	-	1	-	-	1	-	-
Male	-	2	-	-	2	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	5	2	1	1	-	-	-
Male	-	-	1	1	1	3	1	1	-	-	-	-
Female	-	-	-	-	1	2	1	-	1	-	-	-
Connecticut												
All races, both sexes	63	72	90	113	149	230	393	499	423	182	43	-
Male	48	44	56	63	93	125	190	195	122	28	3	-
Female	15	28	34	50	56	105	203	304	301	154	40	-
White, both sexes	52	58	73	96	125	216	377	484	414	176	40	-
Male	41	35	48	56	77	118	183	191	120	27	3	-
Female	11	23	25	40	48	98	194	293	294	149	37	-
Black, both sexes	11	13	15	17	24	12	15	12	8	5	3	-
Male	7	9	8	7	16	6	7	4	2	-	-	-
Female	4	4	7	10	8	6	8	8	6	5	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	-	2	1	2	1	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	1	2	-	-	1	1	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Delaware												
All races, both sexes	11	21	15	43	34	57	83	90	57	30	5	-
Male	7	14	10	23	17	26	39	30	16	4	1	-
Female	4	7	5	20	17	31	44	60	41	26	4	-
White, both sexes	7	15	10	31	23	46	73	81	55	27	5	-
Male	4	12	7	20	10	20	36	27	16	3	1	-
Female	3	3	3	11	13	26	37	54	39	24	4	-
Black, both sexes	4	6	4	11	10	10	9	6	1	3	-	-
Male	3	2	2	2	6	6	3	2	-	1	-	-
Female	1	4	2	9	4	4	6	4	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	1	1	2	1	-	-	-
Male	-	-	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
Dist. of Columbia												
All races, both sexes	13	6	13	10	13	16	31	40	21	12	4	-
Male	6	5	9	7	10	8	7	11	2	1	-	-
Female	7	1	4	3	3	8	24	29	19	11	4	-
White, both sexes	-	-	1	-	1	5	7	11	3	5	3	-
Male	-	-	1	-	1	3	2	6	1	1	-	-
Female	-	-	-	-	-	2	5	5	2	4	3	-
Black, both sexes	13	5	12	10	12	11	24	29	18	7	1	-
Male	6	4	8	7	9	5	5	5	1	-	-	-
Female	7	1	4	3	3	6	19	24	17	7	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Florida												
All races, both sexes	255	291	353	453	657	1,061	1,502	1,806	1,396	551	100	1
Male	163	178	204	274	373	537	700	769	500	160	22	-
Female	92	113	149	179	284	524	802	1,037	896	391	78	1
White, both sexes	182	221	276	376	555	951	1,380	1,729	1,334	528	91	-
Male	125	144	165	234	318	485	651	750	483	151	20	-
Female	57	77	111	142	237	466	729	979	851	377	71	-
Black, both sexes	67	68	75	75	97	104	115	72	57	23	9	1
Male	35	33	38	39	51	47	45	19	14	9	2	-
Female	32	35	37	36	46	57	70	53	43	14	7	1
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	5	2	1	2	5	6	7	5	4	-	-	-
Male	3	1	-	1	4	5	4	-	3	-	-	-
Female	2	1	1	1	1	1	3	5	1	-	-	-
Georgia												
All races, both sexes	300	345	418	478	637	861	1,047	1,057	781	289	79	2
Male	189	211	261	280	342	410	403	368	180	60	15	2
Female	111	134	157	198	295	451	644	689	601	229	64	-
White, both sexes	176	216	263	305	461	696	862	874	658	216	55	2
Male	111	137	162	176	247	340	346	318	155	42	10	2
Female	65	79	101	129	214	356	516	556	503	174	45	-
Black, both sexes	124	126	151	167	168	159	179	180	116	71	24	-
Male	78	72	96	100	91	66	53	49	23	17	5	-
Female	46	54	55	67	77	93	126	131	93	54	19	-
American Indian 1/, both sexes	-	1	1	1	4	1	1	-	4	1	-	-
Male	-	1	1	-	1	1	-	-	2	-	-	-
Female	-	-	-	1	3	-	1	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	2	3	5	4	5	5	3	3	1	-	-
Male	-	1	2	4	3	3	4	1	-	1	-	-
Female	-	1	1	1	1	2	1	2	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Hawaii												
All races, both sexes	33	47	59	69	71	108	142	174	149	74	11	-
Male	19	39	39	46	39	56	84	76	55	24	3	-
Female	14	8	20	23	32	52	58	98	94	50	8	-
White, both sexes	11	10	13	18	15	29	35	50	42	19	3	-
Male	8	8	11	13	5	14	18	19	17	5	1	-
Female	3	2	2	5	10	15	17	31	25	14	2	-
Black, both sexes	1	1	-	-	1	1	1	1	-	-	-	-
Male	1	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	36	46	51	55	78	106	120	107	55	8	-
Male	10	30	28	33	33	41	66	56	38	19	2	-
Female	11	6	18	18	22	37	40	64	69	36	6	-
Idaho												
All races, both sexes	14	18	26	28	41	79	125	164	131	59	11	-
Male	8	11	17	15	24	35	48	58	52	11	3	-
Female	6	7	9	13	17	44	77	106	79	48	8	-
White, both sexes	14	18	25	28	41	77	123	162	131	59	11	-
Male	8	11	17	15	24	34	47	57	52	11	3	-
Female	6	7	8	13	17	43	76	105	79	48	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	2	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Illinois												
All races, both sexes	224	258	331	393	531	875	1,238	1,447	1,179	535	123	-
Male	147	149	194	230	297	424	547	505	325	91	18	-
Female	77	109	137	163	234	451	691	942	854	444	105	-
White, both sexes	146	179	250	294	435	745	1,110	1,352	1,113	505	113	-
Male	96	105	155	173	245	370	493	467	310	85	15	-
Female	50	74	95	121	190	375	617	885	803	420	98	-
Black, both sexes	76	77	69	91	89	125	120	83	58	25	9	-
Male	49	44	34	52	46	52	52	31	12	5	2	-
Female	27	33	35	39	43	73	68	52	46	20	7	-
American Indian 1/, both sexes	-	-	2	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	10	8	7	5	7	12	8	5	1	-
Male	2	-	5	5	6	2	2	7	3	1	1	-
Female	-	2	5	3	1	3	5	5	5	4	-	-
Indiana												
All races, both sexes	121	177	228	240	393	562	764	872	676	272	66	-
Male	74	112	128	131	191	267	335	309	203	53	9	-
Female	47	65	100	109	202	295	429	563	473	219	57	-
White, both sexes	104	159	200	214	357	534	721	838	644	262	60	-
Male	63	101	111	119	176	261	326	295	191	49	7	-
Female	41	58	89	95	181	273	395	543	453	213	53	-
Black, both sexes	17	18	28	24	33	28	41	33	31	8	5	-
Male	11	11	17	11	15	6	8	14	11	3	1	-
Female	6	7	11	13	18	22	33	19	20	5	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	3	-	1	1	1	2	1	-
Male	-	-	-	1	-	-	1	-	1	1	1	-
Female	-	-	-	1	3	-	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Iowa												
All races, both sexes	34	37	56	60	87	138	250	257	228	101	31	-
Male	25	22	27	36	43	70	106	89	65	21	8	-
Female	9	15	29	24	44	68	144	168	163	80	23	-
White, both sexes	33	34	51	59	83	133	249	256	228	101	31	-
Male	24	21	26	36	40	68	105	89	65	21	8	-
Female	9	13	25	23	43	65	144	167	163	80	23	-
Black, both sexes	1	3	5	1	3	5	1	1	-	-	-	-
Male	1	1	1	-	2	2	1	-	-	-	-	-
Female	-	2	4	1	1	3	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	41	52	65	76	126	221	312	400	361	173	41	-
Male	26	24	37	44	62	102	144	144	106	47	7	-
Female	15	28	28	32	64	119	168	256	255	126	34	-
White, both sexes	36	49	55	72	114	200	297	383	356	165	40	-
Male	24	22	30	41	55	90	138	140	103	43	7	-
Female	12	27	25	31	59	110	159	243	253	122	33	-
Black, both sexes	4	3	10	2	11	14	12	14	4	7	-	-
Male	2	2	7	1	6	7	4	4	2	3	-	-
Female	2	1	3	1	5	7	8	10	2	4	-	-
American Indian 1/, both sexes	1	-	-	2	1	5	2	3	1	1	1	-
Male	-	-	-	2	1	4	2	-	1	1	-	-
Female	1	-	-	-	-	1	-	3	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Kentucky												
All races, both sexes	111	135	159	203	316	414	570	610	460	179	32	-
Male	77	90	107	107	173	204	219	198	126	31	3	-
Female	34	45	52	96	143	210	351	412	334	148	29	-
White, both sexes	95	121	150	186	285	386	539	584	445	171	30	-
Male	66	78	103	100	162	184	213	189	123	30	3	-
Female	29	43	47	86	123	202	326	395	322	141	27	-
Black, both sexes	16	13	8	16	30	26	31	25	14	8	2	-
Male	11	11	3	6	11	18	6	8	3	1	-	-
Female	5	2	5	10	19	8	25	17	11	7	2	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	2	-	1	1	-	-	-
Male	-	-	1	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Louisiana												
All races, both sexes	130	160	173	220	280	441	526	628	445	185	44	-
Male	82	87	96	129	144	213	229	224	129	38	10	-
Female	48	73	77	91	136	228	297	404	316	147	34	-
White, both sexes	74	79	105	144	184	322	422	512	366	144	34	-
Male	49	49	64	86	89	172	187	194	101	27	7	-
Female	25	30	41	58	95	150	235	318	265	117	27	-
Black, both sexes	55	80	67	74	95	119	102	115	78	40	10	-
Male	33	38	32	42	55	41	42	29	28	11	3	-
Female	22	42	35	32	40	78	60	86	50	29	7	-
American Indian 1/, both sexes	1	-	-	2	1	-	1	-	1	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	-	1	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Maine												
All races, both sexes	29	40	40	50	65	101	144	165	137	73	11	-
Male	19	30	22	35	43	46	56	63	41	10	2	-
Female	10	10	18	15	22	55	88	102	96	63	9	-
White, both sexes	29	40	40	50	64	100	144	164	137	73	11	-
Male	19	30	22	35	42	46	56	62	41	10	2	-
Female	10	10	18	15	22	54	88	102	96	63	9	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	92	107	119	133	195	287	437	426	323	120	33	-
Male	61	65	66	79	92	138	176	166	89	34	6	-
Female	31	42	53	54	103	149	261	260	234	86	27	-
White, both sexes	45	67	74	85	135	218	349	364	270	105	26	-
Male	32	42	38	52	63	107	145	144	78	31	6	-
Female	13	25	36	33	72	111	204	220	192	74	20	-
Black, both sexes	44	38	42	46	58	69	79	56	51	14	7	-
Male	27	23	26	25	28	31	25	22	11	2	-	-
Female	17	15	16	21	30	38	54	34	40	12	7	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	3	2	1	-	9	6	2	1	-	-
Male	2	-	2	2	1	-	6	-	-	1	-	-
Female	1	2	1	-	-	-	3	6	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Massachusetts												
All races, both sexes	88	123	132	200	304	501	778	982	841	378	97	-
Male	57	81	83	121	166	255	341	372	220	75	19	-
Female	31	42	49	79	138	246	437	610	621	303	78	-
White, both sexes	79	106	118	184	286	474	742	955	820	372	93	-
Male	52	71	75	113	161	239	324	359	218	73	19	-
Female	27	35	43	71	125	235	418	596	602	299	74	-
Black, both sexes	9	15	13	14	17	20	26	24	16	6	2	-
Male	5	9	7	7	5	11	12	10	1	2	-	-
Female	4	6	6	7	12	9	14	14	15	4	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	3	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	3	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	1	1	7	9	3	2	-	2	-
Male	-	1	1	1	-	5	5	3	1	-	-	-
Female	-	1	-	-	1	2	4	-	1	-	2	-
Michigan												
All races, both sexes	183	222	217	273	400	718	1,039	1,207	952	394	84	-
Male	110	129	139	146	201	338	446	460	287	87	8	-
Female	73	93	78	127	199	380	593	747	665	307	76	-
White, both sexes	125	158	168	220	341	618	953	1,132	906	372	79	-
Male	78	92	105	123	169	292	412	432	270	82	8	-
Female	47	66	63	97	172	326	541	700	636	290	71	-
Black, both sexes	57	58	47	48	55	94	80	69	43	20	5	-
Male	32	33	34	21	30	44	31	27	16	5	-	-
Female	25	25	13	27	25	50	49	42	27	15	5	-
American Indian 1/, both sexes	-	6	1	2	3	3	2	3	1	1	-	-
Male	-	4	-	1	2	1	1	1	1	-	-	-
Female	-	2	1	1	1	2	1	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	3	1	3	4	3	2	1	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	1	-	1	2	1	2	2	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Minnesota												
All races, both sexes	53	84	81	107	171	282	440	629	583	263	37	-
Male	34	42	48	78	101	168	208	241	195	58	5	-
Female	19	42	33	29	70	114	232	388	388	205	32	-
White, both sexes	44	78	74	104	165	277	434	620	579	261	36	-
Male	28	38	43	77	96	165	206	237	194	58	4	-
Female	16	40	31	27	69	112	228	383	385	203	32	-
Black, both sexes	4	4	4	1	3	3	4	2	2	1	-	-
Male	3	2	3	-	3	2	2	2	1	-	-	-
Female	1	2	1	1	-	1	2	-	1	1	-	-
American Indian 1/, both sexes	4	1	3	2	3	1	1	2	1	-	-	-
Male	3	1	2	1	2	-	-	-	-	-	-	-
Female	1	-	1	1	1	1	1	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	1	1	5	1	1	1	-
Male	-	1	-	-	-	1	-	2	-	-	1	-
Female	1	-	-	-	-	-	1	3	1	1	-	-
Mississippi												
All races, both sexes	85	154	169	186	244	413	477	501	426	182	49	-
Male	51	95	98	99	127	204	183	164	121	32	8	-
Female	34	59	71	87	117	209	294	337	305	150	41	-
White, both sexes	36	76	99	107	158	291	358	393	319	132	23	-
Male	22	48	61	57	82	148	140	127	87	19	2	-
Female	14	28	38	50	76	143	218	266	232	113	21	-
Black, both sexes	48	78	67	78	85	121	119	107	106	50	26	-
Male	29	47	36	41	44	55	43	37	33	13	6	-
Female	19	31	31	37	41	66	76	70	73	37	20	-
American Indian 1/, both sexes	-	-	2	-	1	1	-	-	1	-	-	-
Male	-	-	1	-	1	1	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Missouri												
All races, both sexes	119	157	190	213	317	487	695	740	618	272	53	-
Male	75	106	106	119	165	249	296	260	185	55	11	-
Female	44	51	84	94	152	238	399	480	433	217	42	-
White, both sexes	96	128	161	197	274	435	650	706	588	263	49	-
Male	63	88	91	111	148	225	278	249	174	53	9	-
Female	33	40	70	86	126	210	372	457	414	210	40	-
Black, both sexes	22	27	27	15	41	50	42	32	28	8	4	-
Male	12	18	14	7	16	24	16	11	9	2	2	-
Female	10	9	13	8	25	26	26	21	19	6	2	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	1	1	2	2	2	1	1	-	-
Male	-	-	1	1	-	-	1	-	1	-	-	-
Female	1	2	1	-	1	2	1	2	-	1	-	-
Montana												
All races, both sexes	24	25	24	38	44	64	118	127	135	50	12	-
Male	17	14	16	21	24	33	65	45	41	10	3	-
Female	7	11	8	17	20	31	53	82	94	40	9	-
White, both sexes	21	22	22	35	43	61	113	127	133	49	12	-
Male	16	13	14	19	23	32	63	45	40	10	3	-
Female	5	9	8	16	20	29	50	82	93	39	9	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	3	2	2	2	1	3	3	-	1	1	-	-
Male	1	1	2	2	1	1	1	-	-	-	-	-
Female	2	1	-	-	-	2	2	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Nebraska												
All races, both sexes	28	33	42	60	119	135	240	283	255	135	45	-
Male	21	23	27	29	67	76	106	99	77	30	9	-
Female	7	10	15	31	52	59	134	184	178	105	36	-
White, both sexes	26	30	39	57	113	129	236	274	252	135	44	-
Male	19	22	24	29	65	71	105	93	76	30	9	-
Female	7	8	15	28	48	58	131	181	176	105	35	-
Black, both sexes	1	1	3	2	6	4	4	6	3	-	-	-
Male	1	-	3	-	2	3	1	3	1	-	-	-
Female	-	1	-	2	4	1	3	3	2	-	-	-
American Indian 1/, both sexes	1	1	-	1	-	-	-	1	-	-	1	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	2	-	2	-	-	-	-
Male	-	1	-	-	-	2	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	87	104	148	166	193	225	293	247	196	56	11	-
Male	49	75	84	109	122	125	137	98	69	10	2	-
Female	38	29	64	57	71	100	156	149	127	46	9	-
White, both sexes	65	82	123	138	162	201	279	233	182	52	10	-
Male	40	60	70	89	102	116	131	93	64	10	2	-
Female	25	22	53	49	60	85	148	140	118	42	8	-
Black, both sexes	18	13	17	15	22	16	8	9	6	3	1	-
Male	7	8	9	12	13	7	3	4	2	-	-	-
Female	11	5	8	3	9	9	5	5	4	3	1	-
American Indian 1/, both sexes	1	2	1	4	2	-	1	2	3	1	-	-
Male	-	2	1	2	2	-	1	1	1	-	-	-
Female	1	-	-	2	-	-	-	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	3	7	7	9	7	8	5	3	5	-	-	-
Male	2	5	4	6	5	2	2	-	2	-	-	-
Female	1	2	3	3	2	6	3	3	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
New Hampshire												
All races, both sexes	16	16	14	25	47	66	112	125	113	54	19	-
Male	9	13	10	12	27	28	50	47	37	11	-	-
Female	7	3	4	13	20	38	62	78	76	43	19	-
White, both sexes	16	16	14	24	47	65	111	125	113	54	19	-
Male	9	13	10	12	27	27	49	47	37	11	-	-
Female	7	3	4	12	20	38	62	78	76	43	19	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	140	146	184	212	358	587	852	1,003	748	320	58	1
Male	88	88	107	139	188	300	370	378	224	79	13	1
Female	52	58	77	73	170	287	482	625	524	241	45	-
White, both sexes	91	114	130	163	298	515	774	944	696	292	53	1
Male	60	68	76	103	155	264	346	358	214	72	12	1
Female	31	46	54	60	143	251	428	586	482	220	41	-
Black, both sexes	47	31	48	42	53	65	72	52	41	25	4	-
Male	26	19	26	30	28	33	22	18	8	5	-	-
Female	21	12	22	12	25	32	50	34	33	20	4	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	1	-	1	-	-
Male	-	-	-	1	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	6	6	6	7	6	6	11	2	1	-
Male	2	1	5	5	4	3	2	2	2	1	1	-
Female	-	-	1	1	2	4	4	4	9	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
New Mexico												
All races, both sexes	20	31	25	45	69	107	140	163	132	72	13	-
Male	14	15	10	25	39	56	65	74	42	21	4	-
Female	6	16	15	20	30	51	75	89	90	51	9	-
White, both sexes	16	27	19	42	60	102	131	159	124	67	12	-
Male	12	14	9	22	35	54	60	73	39	20	3	-
Female	4	13	10	20	25	48	71	86	85	47	9	-
Black, both sexes	1	1	2	1	2	-	1	1	1	1	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	1	1	2	-	1	-	-	1	1	1	-	-
American Indian 1/, both sexes	3	3	4	2	6	5	8	3	7	4	1	-
Male	2	1	1	2	3	2	4	1	3	1	1	-
Female	1	2	3	-	3	3	4	2	4	3	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
New York												
All races, both sexes	176	259	295	330	566	940	1,335	1,637	1,315	562	135	1
Male	105	159	175	185	302	483	586	632	392	123	20	-
Female	71	100	120	145	264	457	749	1,005	923	439	115	1
White, both sexes	125	205	223	273	481	842	1,238	1,546	1,253	537	120	1
Male	81	131	134	150	263	451	552	600	380	121	17	-
Female	44	74	89	123	218	391	686	946	873	416	103	1
Black, both sexes	49	49	68	48	81	91	91	71	52	24	12	-
Male	24	26	39	28	37	31	32	23	8	2	2	-
Female	25	23	29	20	44	60	59	48	44	22	10	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	1	-	1	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	5	3	8	4	7	6	19	10	-	3	-
Male	-	2	2	6	2	1	2	8	4	-	1	-
Female	2	3	1	2	2	6	4	11	6	-	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
North Carolina												
All races, both sexes	168	209	262	311	447	599	876	875	631	278	54	-
Male	104	133	153	172	241	282	368	281	170	47	14	-
Female	64	76	109	139	206	317	508	594	461	231	40	-
White, both sexes	90	129	163	218	340	474	725	757	553	234	50	-
Male	62	79	97	123	188	220	318	252	146	40	14	-
Female	28	50	66	95	152	254	407	505	407	194	36	-
Black, both sexes	77	75	93	89	99	117	142	113	75	42	4	-
Male	41	52	55	47	50	58	46	27	23	6	-	-
Female	36	23	38	42	49	59	96	86	52	36	4	-
American Indian 1/, both sexes	-	5	5	2	4	6	8	3	3	1	-	-
Male	-	2	1	1	3	4	3	2	1	-	-	-
Female	-	3	4	1	1	2	5	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	4	2	1	2	-	1	-	-
Male	1	-	-	1	-	-	1	-	-	1	-	-
Female	-	-	1	1	4	2	-	2	-	-	-	-
North Dakota												
All races, both sexes	7	8	8	8	18	29	42	74	63	34	7	-
Male	6	5	6	3	9	9	21	22	20	7	1	-
Female	1	3	2	5	9	20	21	52	43	27	6	-
White, both sexes	7	8	8	8	18	29	42	73	62	34	7	-
Male	6	5	6	3	9	9	21	22	19	7	1	-
Female	1	3	2	5	9	20	21	51	43	27	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Ohio												
All races, both sexes	186	258	291	345	512	816	1,169	1,312	933	365	72	-
Male	120	176	155	193	262	398	517	467	265	75	15	-
Female	66	82	136	152	250	418	652	845	668	290	57	-
White, both sexes	147	203	234	274	451	730	1,095	1,236	894	348	60	-
Male	95	137	128	155	235	361	488	434	259	71	11	-
Female	52	66	106	119	216	369	607	802	635	277	49	-
Black, both sexes	38	54	56	68	58	83	73	75	39	17	12	-
Male	24	39	27	36	27	36	29	32	6	4	4	-
Female	14	15	29	32	31	47	44	43	33	13	8	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	3	3	2	1	1	-	-	-	-
Male	1	-	-	2	-	1	-	1	-	-	-	-
Female	-	-	1	1	3	1	1	-	-	-	-	-
Oklahoma												
All races, both sexes	62	80	109	129	188	289	404	403	355	142	36	-
Male	32	46	63	72	86	126	169	134	92	37	10	-
Female	30	34	46	57	102	163	235	269	263	105	26	-
White, both sexes	45	63	90	106	161	266	364	372	341	131	34	-
Male	26	36	50	57	71	121	157	118	89	33	10	-
Female	19	27	40	49	90	145	207	254	252	98	24	-
Black, both sexes	12	12	12	12	11	10	25	17	7	6	2	-
Male	4	7	8	9	6	2	6	8	-	1	-	-
Female	8	5	4	3	5	8	19	9	7	5	2	-
American Indian 1/, both sexes	5	4	7	11	16	12	14	13	7	5	-	-
Male	2	3	5	6	9	2	6	8	3	3	-	-
Female	3	1	2	5	7	10	8	5	4	2	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Oregon												
All races, both sexes	43	61	71	78	126	224	378	489	453	191	45	-
Male	30	42	46	32	64	118	170	193	149	38	7	-
Female	13	19	25	46	62	106	208	296	304	153	38	-
White, both sexes	38	56	65	75	125	222	374	478	443	189	44	-
Male	28	38	43	32	64	117	167	189	147	38	6	-
Female	10	18	22	43	61	105	207	289	296	151	38	-
Black, both sexes	3	2	3	2	1	-	2	6	4	-	-	-
Male	1	1	-	-	-	-	1	1	-	-	-	-
Female	2	1	3	2	1	-	1	5	4	-	-	-
American Indian 1/, both sexes	2	1	2	1	-	1	1	2	1	-	1	-
Male	1	1	2	-	-	1	1	-	1	-	1	-
Female	1	-	-	1	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	-	-	1	1	3	5	2	-	-
Male	-	2	1	-	-	-	1	3	1	-	-	-
Female	-	-	-	-	-	1	-	-	4	2	-	-
Pennsylvania												
All races, both sexes	288	341	421	523	797	1,282	2,005	2,268	1,709	700	129	-
Male	200	215	260	298	433	629	856	853	459	143	20	-
Female	88	126	161	225	364	653	1,149	1,415	1,250	557	109	-
White, both sexes	213	262	351	446	709	1,178	1,888	2,168	1,638	670	121	-
Male	150	162	220	263	393	573	808	826	448	142	20	-
Female	63	100	131	183	316	605	1,080	1,342	1,190	528	101	-
Black, both sexes	72	75	69	75	84	100	113	98	68	28	8	-
Male	47	51	40	35	38	53	47	26	11	1	-	-
Female	25	24	29	40	46	47	66	72	57	27	8	-
American Indian 1/, both sexes	-	2	1	1	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	-	1	4	4	3	2	3	2	-	-
Male	3	1	-	-	2	3	1	1	-	-	-	-
Female	-	1	-	1	2	1	2	1	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Rhode Island												
All races, both sexes	15	18	15	13	26	57	56	91	65	24	8	-
Male	11	10	7	9	11	29	25	32	16	3	2	-
Female	4	8	8	4	15	28	31	59	49	21	6	-
White, both sexes	12	17	13	10	26	55	54	89	65	24	8	-
Male	8	10	6	7	11	27	24	30	16	3	2	-
Female	4	7	7	3	15	28	30	59	49	21	6	-
Black, both sexes	3	1	2	3	-	2	1	2	-	-	-	-
Male	3	-	1	2	-	2	-	2	-	-	-	-
Female	-	1	1	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	151	184	175	212	288	418	502	530	333	132	30	1
Male	100	115	96	121	177	209	214	180	97	22	4	1
Female	51	69	79	91	111	209	288	350	236	110	26	-
White, both sexes	79	104	105	133	197	312	382	431	280	101	20	1
Male	53	66	60	72	121	155	168	142	85	19	3	1
Female	26	38	45	61	76	157	214	289	195	82	17	-
Black, both sexes	71	80	69	78	89	106	119	96	52	30	9	-
Male	46	49	35	48	54	54	46	36	12	3	1	-
Female	25	31	34	30	35	52	73	60	40	27	8	-
American Indian 1/, both sexes	1	-	1	-	1	-	-	1	1	-	1	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	1	2	-	1	-	-
Male	-	-	-	1	1	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
South Dakota												
All races, both sexes	15	5	12	14	18	31	45	58	47	27	8	-
Male	12	3	7	11	10	16	26	31	9	2	-	-
Female	3	2	5	3	8	15	19	27	38	25	8	-
White, both sexes	11	4	11	11	17	28	42	56	46	27	8	-
Male	8	2	6	10	9	14	25	31	9	2	-	-
Female	3	2	5	1	8	14	17	25	37	25	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	1	3	1	3	3	2	1	-	-	-
Male	4	-	1	1	1	2	1	-	-	-	-	-
Female	-	-	-	2	-	1	2	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	114	184	166	215	274	311	484	468	355	118	35	-
Male	74	119	98	122	133	141	193	167	97	28	4	-
Female	40	65	68	93	141	170	291	301	258	90	31	-
White, both sexes	71	131	127	172	229	265	427	422	321	105	32	-
Male	47	87	77	106	106	117	167	150	89	26	3	-
Female	24	44	50	66	123	148	260	272	232	79	29	-
Black, both sexes	43	53	39	42	44	42	55	45	33	13	3	-
Male	27	32	21	16	26	22	26	17	8	2	1	-
Female	16	21	18	26	18	20	29	28	25	11	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	3	2	1	1	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	1	-	2	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Texas												
All races, both sexes	369	406	495	584	835	1,165	1,651	1,684	1,373	584	115	-
Male	245	239	281	327	453	557	697	594	405	142	13	-
Female	124	167	214	257	382	608	954	1,090	968	442	102	-
White, both sexes	253	311	385	474	723	1,030	1,502	1,561	1,276	539	101	-
Male	170	190	229	270	401	490	634	552	375	131	11	-
Female	83	121	156	204	322	540	868	1,009	901	408	90	-
Black, both sexes	111	93	99	101	105	126	137	114	89	39	14	-
Male	70	48	46	52	48	64	60	37	28	9	2	-
Female	41	45	53	49	57	62	77	77	61	30	12	-
American Indian 1/, both sexes	1	-	-	1	1	-	1	-	2	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	1	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	11	8	6	9	11	9	6	6	-	-
Male	4	1	6	5	4	3	3	5	1	2	-	-
Female	-	1	5	3	2	6	8	4	5	4	-	-
Utah												
All races, both sexes	25	27	37	47	77	144	217	301	219	85	9	-
Male	13	14	22	25	27	70	87	102	73	23	2	-
Female	12	13	15	22	50	74	130	199	146	62	7	-
White, both sexes	24	26	35	47	72	141	215	299	216	82	9	-
Male	12	14	20	25	26	69	87	101	71	23	2	-
Female	12	12	15	22	46	72	128	198	145	59	7	-
Black, both sexes	1	1	1	-	1	1	2	1	-	2	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	-	2	1	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	3	1	-	1	2	-	-	-
Male	-	-	1	-	1	-	-	1	1	-	-	-
Female	-	-	-	-	2	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Vermont												
All races, both sexes	4	3	12	8	17	33	47	72	57	27	7	-
Male	1	3	7	7	12	17	19	32	17	7	1	-
Female	3	-	5	1	5	16	28	40	40	20	6	-
White, both sexes	4	3	12	8	17	33	47	72	57	27	7	-
Male	1	3	7	7	12	17	19	32	17	7	1	-
Female	3	-	5	1	5	16	28	40	40	20	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	147	232	238	288	380	594	802	860	649	255	58	-
Male	87	163	135	152	199	288	336	322	189	55	9	-
Female	60	69	103	136	181	306	466	538	460	200	49	-
White, both sexes	89	157	172	202	288	467	671	741	570	227	54	-
Male	52	116	104	105	153	233	287	278	174	50	9	-
Female	37	41	68	97	135	234	384	463	396	177	45	-
Black, both sexes	54	73	61	80	86	121	123	109	74	27	4	-
Male	33	46	29	43	42	51	46	41	13	5	-	-
Female	21	27	32	37	44	70	77	68	61	22	4	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	4	6	6	6	8	9	5	1	-	-
Male	2	1	1	4	4	4	3	3	2	-	-	-
Female	2	1	3	2	2	2	5	6	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Washington												
All races, both sexes	70	71	88	104	169	263	430	517	383	161	46	-
Male	43	46	57	62	93	128	188	225	125	41	5	-
Female	27	25	31	42	76	135	242	292	258	120	41	-
White, both sexes	60	59	82	92	153	252	401	501	371	158	46	-
Male	39	39	54	57	82	121	178	217	122	40	5	-
Female	21	20	28	35	71	131	223	284	249	118	41	-
Black, both sexes	7	8	3	8	8	4	12	6	2	1	-	-
Male	3	5	1	4	5	4	7	3	1	-	-	-
Female	4	3	2	4	3	-	5	3	1	1	-	-
American Indian 1/, both sexes	2	2	1	1	4	1	4	1	2	-	-	-
Male	1	1	1	-	3	1	1	1	-	-	-	-
Female	1	1	-	1	1	-	3	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	3	4	6	13	9	8	2	-	-
Male	-	1	1	1	3	2	2	4	2	1	-	-
Female	1	1	1	2	1	4	11	5	6	1	-	-
West Virginia												
All races, both sexes	30	59	77	90	141	199	287	259	203	61	25	-
Male	15	35	47	50	65	109	112	93	48	12	5	-
Female	15	24	30	40	76	90	175	166	155	49	20	-
White, both sexes	28	59	74	82	138	195	282	246	195	60	25	-
Male	15	35	45	46	64	106	110	89	46	12	5	-
Female	13	24	29	36	74	89	172	157	149	48	20	-
Black, both sexes	2	-	2	8	3	4	5	11	8	1	-	-
Male	-	-	1	4	1	3	2	4	2	-	-	-
Female	2	-	1	4	2	1	3	7	6	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other heart diseases (I26-I51)												
Wisconsin												
All races, both sexes	60	110	118	152	223	410	575	806	676	268	64	-
Male	48	79	65	79	121	209	275	311	213	49	7	-
Female	12	31	53	73	102	201	300	495	463	219	57	-
White, both sexes	52	95	100	143	210	393	561	791	665	268	62	-
Male	44	66	56	76	114	201	271	304	211	49	7	-
Female	8	29	44	67	96	192	290	487	454	219	55	-
Black, both sexes	7	11	13	6	10	15	10	11	6	-	2	-
Male	4	10	6	3	5	8	3	5	1	-	-	-
Female	3	1	7	3	5	7	7	6	5	-	2	-
American Indian 1/, both sexes	1	4	3	2	3	1	1	4	3	-	-	-
Male	-	3	2	-	2	-	-	2	1	-	-	-
Female	1	1	1	2	1	1	1	2	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	1	3	-	2	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	1	2	-	2	-	-	-
Wyoming												
All races, both sexes	9	15	14	22	29	34	54	49	39	19	9	-
Male	3	9	8	12	13	16	23	14	19	6	1	-
Female	6	6	6	10	16	18	31	35	20	13	8	-
White, both sexes	9	15	13	22	28	34	54	49	38	19	9	-
Male	3	9	7	12	13	16	23	14	19	6	1	-
Female	6	6	6	10	15	18	31	35	19	13	8	-
Black, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
United States												
All races, both sexes	13,504	98	20	8	8	47	78	128	192	288	435	612
Male	5,357	55	11	2	4	16	34	44	68	131	188	258
Female	8,147	43	9	6	4	31	44	84	124	157	247	354
White, both sexes	11,085	67	16	4	6	37	53	76	108	198	300	441
Male	4,422	39	9	1	2	11	22	25	40	95	130	191
Female	6,663	28	7	3	4	26	31	51	68	103	170	250
Black, both sexes	2,201	28	4	3	2	9	23	47	77	83	127	158
Male	852	16	2	1	2	4	11	17	26	33	54	63
Female	1,349	12	2	2	-	5	12	30	51	50	73	95
American Indian 1/, both sexes	69	1	-	-	-	-	-	3	1	1	4	5
Male	29	-	-	-	-	-	-	1	-	-	3	1
Female	40	1	-	-	-	-	-	2	1	1	1	4
Asian or Pacific Islander 2/, both sexes	149	2	-	1	-	1	2	2	6	6	4	8
Male	54	-	-	-	-	1	1	1	2	3	1	3
Female	95	2	-	1	-	-	1	1	4	3	3	5
Alabama												
All races, both sexes	328	1	1	1	-	1	-	4	4	5	7	18
Male	125	-	-	1	-	-	-	2	2	4	2	8
Female	203	1	1	-	-	1	-	2	2	1	5	10
White, both sexes	224	1	1	1	-	1	-	1	4	4	4	9
Male	84	-	-	1	-	-	-	-	2	4	1	2
Female	140	1	1	-	-	1	-	1	2	-	3	7
Black, both sexes	104	-	-	-	-	-	-	3	-	1	3	9
Male	41	-	-	-	-	-	-	2	-	-	1	6
Female	63	-	-	-	-	-	-	1	-	1	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Alaska												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Arizona												
All races, both sexes	211	1	-	-	-	-	2	4	1	1	4	12
Male	92	1	-	-	-	-	2	-	-	1	1	5
Female	119	-	-	-	-	-	-	4	1	-	3	7
White, both sexes	199	1	-	-	-	-	2	2	-	1	4	12
Male	88	1	-	-	-	-	2	-	-	1	1	5
Female	111	-	-	-	-	-	-	2	-	-	3	7
Black, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Arkansas												
All races, both sexes	207	2	-	-	-	-	2	1	3	7	7	12
Male	92	1	-	-	-	-	2	-	1	2	3	7
Female	115	1	-	-	-	-	-	1	2	5	4	5
White, both sexes	175	2	-	-	-	-	1	-	2	5	6	10
Male	78	1	-	-	-	-	1	-	1	1	3	6
Female	97	1	-	-	-	-	-	-	1	4	3	4
Black, both sexes	32	-	-	-	-	-	1	1	1	2	1	2
Male	14	-	-	-	-	-	1	-	-	1	-	1
Female	18	-	-	-	-	-	-	1	1	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	776	7	2	2	-	4	5	11	18	19	40	33
Male	283	4	-	-	-	1	1	3	5	6	18	13
Female	493	3	2	2	-	3	4	8	13	13	22	20
White, both sexes	623	5	2	1	-	3	4	7	12	11	30	21
Male	232	3	-	-	-	1	1	1	3	5	14	7
Female	391	2	2	1	-	2	3	6	9	6	16	14
Black, both sexes	101	2	-	-	-	1	-	3	4	7	9	11
Male	33	1	-	-	-	-	-	1	1	1	4	5
Female	68	1	-	-	-	1	-	2	3	6	5	6
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	50	-	-	1	-	-	1	1	2	1	1	1
Male	16	-	-	-	-	-	-	1	1	-	-	1
Female	34	-	-	1	-	-	1	-	1	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Colorado												
All races, both sexes	232	2	-	-	-	-	1	-	4	4	4	9
Male	109	-	-	-	-	-	1	-	4	2	1	6
Female	123	2	-	-	-	-	-	-	-	2	3	3
White, both sexes	213	2	-	-	-	-	-	-	2	3	3	8
Male	99	-	-	-	-	-	-	-	2	2	1	6
Female	114	2	-	-	-	-	-	-	-	1	2	2
Black, both sexes	15	-	-	-	-	-	1	-	2	1	1	1
Male	7	-	-	-	-	-	1	-	2	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	156	-	-	1	1	-	1	1	3	2	8	8
Male	69	-	-	1	1	-	-	-	1	2	5	5
Female	87	-	-	-	-	-	1	1	2	-	3	3
White, both sexes	137	-	-	-	-	-	1	1	1	2	7	6
Male	60	-	-	-	-	-	-	-	1	2	4	4
Female	77	-	-	-	-	-	1	1	-	-	3	2
Black, both sexes	19	-	-	1	1	-	-	-	2	-	1	2
Male	9	-	-	1	1	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	2	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Delaware												
All races, both sexes	52	2	-	-	-	-	-	-	-	-	3	1
Male	22	1	-	-	-	-	-	-	-	-	1	-
Female	30	1	-	-	-	-	-	-	-	-	2	1
White, both sexes	42	-	-	-	-	-	-	-	-	-	2	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	9	2	-	-	-	-	-	-	-	-	1	-
Male	4	1	-	-	-	-	-	-	-	-	1	-
Female	5	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	24	-	-	-	-	-	-	-	1	-	1	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	-	-	-	1	-	1	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Florida												
All races, both sexes	761	6	1	-	-	2	-	4	7	20	20	34
Male	314	4	-	-	-	1	-	3	1	9	7	12
Female	447	2	1	-	-	1	-	1	6	11	13	22
White, both sexes	641	3	-	-	-	1	-	2	4	17	14	19
Male	268	2	-	-	-	-	-	1	-	7	4	8
Female	373	1	-	-	-	1	-	1	4	10	10	11
Black, both sexes	116	3	1	-	-	-	-	2	3	3	6	14
Male	44	2	-	-	-	-	-	2	1	2	3	4
Female	72	1	1	-	-	-	-	-	2	1	3	10
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	441	5	-	-	-	2	4	7	11	21	20	18
Male	174	3	-	-	-	-	1	1	3	10	12	6
Female	267	2	-	-	-	2	3	6	8	11	8	12
White, both sexes	295	2	-	-	-	1	1	2	3	11	10	8
Male	118	-	-	-	-	-	-	-	1	5	5	3
Female	177	2	-	-	-	1	1	2	2	6	5	5
Black, both sexes	139	3	-	-	-	1	3	4	8	9	10	10
Male	54	3	-	-	-	-	1	1	2	5	7	3
Female	85	-	-	-	-	1	2	3	6	4	3	7
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	1	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Hawaii												
All races, both sexes	29	1	-	-	-	-	-	1	-	2	1	3
Male	12	-	-	-	-	-	-	-	-	2	-	1
Female	17	1	-	-	-	-	-	1	-	-	1	2
White, both sexes	8	-	-	-	-	-	-	1	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	1	-	-	-	-	-	-	-	2	1	3
Male	7	-	-	-	-	-	-	-	-	2	-	1
Female	14	1	-	-	-	-	-	-	-	-	1	2
Idaho												
All races, both sexes	82	-	-	-	-	-	-	1	-	4	3	6
Male	33	-	-	-	-	-	-	-	-	3	3	2
Female	49	-	-	-	-	-	-	1	-	1	-	4
White, both sexes	81	-	-	-	-	-	-	1	-	4	3	6
Male	33	-	-	-	-	-	-	-	-	3	3	2
Female	48	-	-	-	-	-	-	1	-	1	-	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Illinois												
All races, both sexes	655	6	1	-	-	1	3	5	11	12	23	34
Male	260	5	1	-	-	-	2	2	4	6	14	14
Female	395	1	-	-	-	1	1	3	7	6	9	20
White, both sexes	517	4	-	-	-	-	3	4	8	7	12	25
Male	210	3	-	-	-	-	2	1	4	4	8	11
Female	307	1	-	-	-	-	1	3	4	3	4	14
Black, both sexes	127	2	1	-	-	1	-	1	3	5	11	9
Male	48	2	1	-	-	-	-	1	-	2	6	3
Female	79	-	-	-	-	1	-	-	3	3	5	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	394	5	1	-	-	3	2	4	12	8	14	24
Male	162	2	-	-	-	1	1	3	3	3	9	8
Female	232	3	1	-	-	2	1	1	9	5	5	16
White, both sexes	356	4	1	-	-	3	2	2	10	7	12	23
Male	144	1	-	-	-	1	1	2	1	3	8	7
Female	212	3	1	-	-	2	1	-	9	4	4	16
Black, both sexes	37	1	-	-	-	-	-	2	1	1	2	1
Male	17	1	-	-	-	-	-	1	1	-	1	1
Female	20	-	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Iowa												
All races, both sexes	192	-	-	-	-	1	2	-	2	4	5	4
Male	77	-	-	-	-	1	-	-	1	3	4	2
Female	115	-	-	-	-	-	2	-	1	1	1	2
White, both sexes	184	-	-	-	-	1	2	-	1	4	5	4
Male	73	-	-	-	-	1	-	-	1	3	4	2
Female	111	-	-	-	-	-	2	-	-	1	1	2
Black, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	152	1	-	-	-	-	3	1	1	2	5	7
Male	54	-	-	-	-	-	1	1	-	1	2	5
Female	98	1	-	-	-	-	2	-	1	1	3	2
White, both sexes	140	1	-	-	-	-	1	1	-	2	3	6
Male	49	-	-	-	-	-	1	1	-	1	2	4
Female	91	1	-	-	-	-	-	-	-	1	1	2
Black, both sexes	12	-	-	-	-	-	2	-	1	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	2	-	1	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Kentucky												
All races, both sexes	251	1	-	1	-	-	1	6	-	5	12	14
Male	99	-	-	-	-	-	1	3	-	2	7	6
Female	152	1	-	1	-	-	-	3	-	3	5	8
White, both sexes	230	1	-	1	-	-	1	4	-	4	8	12
Male	93	-	-	-	-	-	1	2	-	1	5	6
Female	137	1	-	1	-	-	-	2	-	3	3	6
Black, both sexes	21	-	-	-	-	-	-	2	-	1	4	2
Male	6	-	-	-	-	-	-	1	-	1	2	-
Female	15	-	-	-	-	-	-	1	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	270	1	1	-	-	-	6	5	4	4	14	13
Male	104	1	1	-	-	-	2	2	1	2	5	7
Female	166	-	-	-	-	-	4	3	3	2	9	6
White, both sexes	180	1	1	-	-	-	4	2	1	4	8	7
Male	72	1	1	-	-	-	1	1	1	2	4	4
Female	108	-	-	-	-	-	3	1	-	2	4	3
Black, both sexes	87	-	-	-	-	-	2	2	3	-	5	6
Male	31	-	-	-	-	-	1	-	-	-	1	3
Female	56	-	-	-	-	-	1	2	3	-	4	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	1	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Maine												
All races, both sexes	51	-	-	-	-	-	-	1	-	-	3	2
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	32	-	-	-	-	-	-	1	-	-	2	1
White, both sexes	51	-	-	-	-	-	-	1	-	-	3	2
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	32	-	-	-	-	-	-	1	-	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	262	4	-	-	-	-	1	2	6	3	12	15
Male	122	3	-	-	-	-	1	2	2	1	5	7
Female	140	1	-	-	-	-	-	-	4	2	7	8
White, both sexes	164	2	-	-	-	-	1	1	3	1	4	8
Male	79	2	-	-	-	-	1	1	1	-	2	4
Female	85	-	-	-	-	-	-	-	2	1	2	4
Black, both sexes	94	2	-	-	-	-	-	1	3	2	8	6
Male	42	1	-	-	-	-	-	1	1	1	3	3
Female	52	1	-	-	-	-	-	-	2	1	5	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Massachusetts												
All races, both sexes	309	3	-	-	-	1	-	3	3	6	9	9
Male	120	2	-	-	-	1	-	-	1	2	5	6
Female	189	1	-	-	-	-	-	3	2	4	4	3
White, both sexes	291	3	-	-	-	1	-	2	2	5	7	9
Male	114	2	-	-	-	1	-	-	1	2	3	6
Female	177	1	-	-	-	-	-	2	1	3	4	3
Black, both sexes	17	-	-	-	-	-	-	1	1	1	2	-
Male	6	-	-	-	-	-	-	-	-	-	2	-
Female	11	-	-	-	-	-	-	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	464	2	2	-	-	1	1	3	7	5	9	16
Male	183	-	2	-	-	-	1	1	3	1	3	1
Female	281	2	-	-	-	1	-	2	4	4	6	15
White, both sexes	368	1	1	-	-	1	-	1	3	1	4	13
Male	146	-	1	-	-	-	-	-	1	1	1	1
Female	222	1	-	-	-	1	-	1	2	-	3	12
Black, both sexes	88	1	1	-	-	-	1	2	4	3	4	3
Male	34	-	1	-	-	-	1	1	2	-	2	-
Female	54	1	-	-	-	-	-	1	2	3	2	3
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Minnesota												
All races, both sexes	204	-	-	-	-	-	1	1	1	4	4	11
Male	73	-	-	-	-	-	-	-	-	2	-	5
Female	131	-	-	-	-	-	1	1	1	2	4	6
White, both sexes	199	-	-	-	-	-	1	1	1	3	4	10
Male	72	-	-	-	-	-	-	-	-	2	-	5
Female	127	-	-	-	-	-	1	1	1	1	4	5
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	205	1	1	-	-	-	6	4	6	4	6	9
Male	70	-	1	-	-	-	2	1	2	1	1	3
Female	135	1	-	-	-	-	4	3	4	3	5	6
White, both sexes	131	-	1	-	-	-	2	-	2	2	4	3
Male	41	-	1	-	-	-	1	-	1	-	-	1
Female	90	-	-	-	-	-	1	-	1	2	4	2
Black, both sexes	74	1	-	-	-	-	4	4	4	2	2	6
Male	29	-	-	-	-	-	1	1	1	1	1	2
Female	45	1	-	-	-	-	3	3	3	1	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Missouri												
All races, both sexes	354	4	4	1	1	1	-	1	4	9	10	14
Male	146	2	3	-	-	-	-	1	-	5	4	6
Female	208	2	1	1	1	1	-	-	4	4	6	8
White, both sexes	304	3	4	1	1	1	-	1	3	8	7	9
Male	123	2	3	-	-	-	-	1	-	5	3	4
Female	181	1	1	1	1	1	-	-	3	3	4	5
Black, both sexes	49	1	-	-	-	-	-	-	1	1	3	5
Male	22	-	-	-	-	-	-	-	-	-	1	2
Female	27	1	-	-	-	-	-	-	1	1	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	51	1	-	-	-	1	-	1	-	1	1	1
Male	18	1	-	-	-	-	-	1	-	-	1	-
Female	33	-	-	-	-	1	-	-	-	1	-	1
White, both sexes	47	1	-	-	-	1	-	1	-	1	1	1
Male	17	1	-	-	-	-	-	1	-	-	1	-
Female	30	-	-	-	-	1	-	-	-	1	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Nebraska												
All races, both sexes	96	-	-	-	-	-	1	1	1	1	2	3
Male	28	-	-	-	-	-	-	-	-	1	1	-
Female	68	-	-	-	-	-	1	1	1	-	1	3
White, both sexes	89	-	-	-	-	-	1	1	-	-	2	2
Male	26	-	-	-	-	-	-	-	-	-	1	-
Female	63	-	-	-	-	-	1	1	-	-	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	115	-	-	-	-	-	-	2	-	2	2	6
Male	56	-	-	-	-	-	-	2	-	1	1	4
Female	59	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	99	-	-	-	-	-	-	2	-	2	-	6
Male	49	-	-	-	-	-	-	2	-	1	-	4
Female	50	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	13	-	-	-	-	-	-	-	-	-	2	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
New Hampshire												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	2	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	35	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	54	-	-	-	-	-	-	-	-	-	2	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	35	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	375	4	1	-	1	2	3	3	3	7	15	10
Male	137	3	1	-	1	1	1	-	1	4	5	4
Female	238	1	-	-	-	1	2	3	2	3	10	6
White, both sexes	298	3	1	-	-	-	3	1	1	5	7	8
Male	114	3	1	-	-	-	1	-	1	3	2	3
Female	184	-	-	-	-	-	2	1	-	2	5	5
Black, both sexes	73	-	-	-	1	2	-	2	2	2	7	2
Male	22	-	-	-	1	1	-	-	-	1	2	1
Female	51	-	-	-	-	1	-	2	2	1	5	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
New Mexico												
All races, both sexes	74	-	-	-	1	-	-	3	1	2	5	3
Male	28	-	-	-	1	-	-	-	-	1	1	1
Female	46	-	-	-	-	-	-	3	1	1	4	2
White, both sexes	67	-	-	-	1	-	-	2	1	2	4	2
Male	26	-	-	-	1	-	-	-	-	1	-	1
Female	41	-	-	-	-	-	-	2	1	1	4	1
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	583	6	1	-	-	3	2	5	9	16	18	23
Male	238	4	-	-	-	3	2	2	3	6	10	10
Female	345	2	1	-	-	-	-	3	6	10	8	13
White, both sexes	475	5	-	-	-	2	1	3	4	11	14	15
Male	191	3	-	-	-	2	1	1	1	5	8	7
Female	284	2	-	-	-	-	-	2	3	6	6	8
Black, both sexes	104	1	1	-	-	1	1	2	5	5	4	8
Male	46	1	-	-	-	1	1	1	2	1	2	3
Female	58	-	1	-	-	-	-	1	3	4	2	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
North Carolina												
All races, both sexes	532	5	-	-	-	5	3	3	4	18	17	25
Male	193	2	-	-	-	-	2	1	1	10	5	12
Female	339	3	-	-	-	5	1	2	3	8	12	13
White, both sexes	360	1	-	-	-	5	-	2	2	9	7	11
Male	127	-	-	-	-	-	-	1	1	4	2	6
Female	233	1	-	-	-	5	-	1	1	5	5	5
Black, both sexes	163	4	-	-	-	-	2	-	2	9	10	14
Male	62	2	-	-	-	-	1	-	-	6	3	6
Female	101	2	-	-	-	-	1	-	2	3	7	8
American Indian 1/, both sexes	7	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	39	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	39	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Ohio												
All races, both sexes	575	2	-	-	-	1	3	10	12	16	13	35
Male	217	2	-	-	-	-	1	3	3	3	7	11
Female	358	-	-	-	-	1	2	7	9	13	6	24
White, both sexes	491	2	-	-	-	1	2	6	7	13	11	26
Male	193	2	-	-	-	-	1	1	2	2	6	6
Female	298	-	-	-	-	1	1	5	5	11	5	20
Black, both sexes	80	-	-	-	-	-	1	4	5	3	2	9
Male	23	-	-	-	-	-	-	2	1	1	1	5
Female	57	-	-	-	-	-	1	2	4	2	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	173	1	2	-	-	3	3	1	1	2	4	7
Male	69	-	1	-	-	-	-	-	-	1	3	3
Female	104	1	1	-	-	3	3	1	1	1	1	4
White, both sexes	147	-	2	-	-	3	3	1	1	1	3	6
Male	57	-	1	-	-	-	-	-	-	1	2	3
Female	90	-	1	-	-	3	3	1	1	-	1	3
Black, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	10	1	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	3	1	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Oregon												
All races, both sexes	161	-	-	-	1	1	1	2	1	3	4	9
Male	70	-	-	-	-	-	1	1	1	1	2	3
Female	91	-	-	-	1	1	-	1	-	2	2	6
White, both sexes	153	-	-	-	1	1	1	2	1	2	4	9
Male	69	-	-	-	-	-	1	1	1	1	2	3
Female	84	-	-	-	1	1	-	1	-	1	2	6
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Pennsylvania												
All races, both sexes	760	4	-	-	1	5	6	4	6	10	22	31
Male	294	2	-	-	-	2	3	3	4	3	5	16
Female	466	2	-	-	1	3	3	1	2	7	17	15
White, both sexes	650	3	-	-	1	4	5	4	4	7	15	25
Male	256	1	-	-	-	2	3	3	3	1	5	13
Female	394	2	-	-	1	2	2	1	1	6	10	12
Black, both sexes	107	1	-	-	-	1	1	-	2	2	7	6
Male	36	1	-	-	-	-	-	-	1	1	-	3
Female	71	-	-	-	-	1	1	-	1	1	7	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Rhode Island												
All races, both sexes	53	2	-	-	-	-	-	1	-	3	2	1
Male	16	-	-	-	-	-	-	-	-	1	-	1
Female	37	2	-	-	-	-	-	1	-	2	2	-
White, both sexes	47	1	-	-	-	-	-	-	-	2	2	-
Male	14	-	-	-	-	-	-	-	-	1	-	-
Female	33	1	-	-	-	-	-	-	-	1	2	-
Black, both sexes	6	1	-	-	-	-	-	1	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	1	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	244	2	-	-	-	-	-	1	4	7	10	15
Male	88	-	-	-	-	-	-	-	4	4	6	5
Female	156	2	-	-	-	-	-	1	-	3	4	10
White, both sexes	160	1	-	-	-	-	-	1	1	2	6	8
Male	55	-	-	-	-	-	-	-	1	1	3	1
Female	105	1	-	-	-	-	-	1	-	1	3	7
Black, both sexes	84	1	-	-	-	-	-	-	3	5	4	7
Male	33	-	-	-	-	-	-	-	3	3	3	4
Female	51	1	-	-	-	-	-	-	-	2	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
South Dakota												
All races, both sexes	43	-	-	-	-	-	-	-	2	1	1	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	2	1	1	1
White, both sexes	41	-	-	-	-	-	-	-	2	1	1	1
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	2	1	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	355	3	-	1	-	1	2	1	4	9	13	23
Male	140	2	-	-	-	-	1	1	1	5	5	14
Female	215	1	-	1	-	1	1	-	3	4	8	9
White, both sexes	304	2	-	-	-	1	1	1	2	5	12	21
Male	124	2	-	-	-	-	-	1	-	4	5	13
Female	180	-	-	-	-	1	1	-	2	1	7	8
Black, both sexes	51	1	-	1	-	-	1	-	2	4	1	2
Male	16	-	-	-	-	-	1	-	1	1	-	1
Female	35	1	-	1	-	-	-	-	1	3	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Texas												
All races, both sexes	900	7	-	-	2	4	7	12	15	14	24	42
Male	373	4	-	-	1	3	1	5	6	7	7	14
Female	527	3	-	-	1	1	6	7	9	7	17	28
White, both sexes	709	6	-	-	2	2	5	8	6	7	17	33
Male	284	3	-	-	1	1	-	3	2	3	4	14
Female	425	3	-	-	1	1	5	5	4	4	13	19
Black, both sexes	180	1	-	-	-	2	2	4	8	7	7	9
Male	82	1	-	-	-	2	1	2	4	4	3	-
Female	98	-	-	-	-	-	1	2	4	3	4	9
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
Utah												
All races, both sexes	122	-	-	-	-	1	2	2	4	3	3	6
Male	56	-	-	-	-	1	2	-	2	2	1	4
Female	66	-	-	-	-	-	-	2	2	1	2	2
White, both sexes	120	-	-	-	-	1	2	2	4	3	3	5
Male	55	-	-	-	-	1	2	-	2	2	1	3
Female	65	-	-	-	-	-	-	2	2	1	2	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Vermont												
All races, both sexes	36	-	-	-	-	-	1	-	-	1	1	1
Male	16	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	1	-	-	1	-	1
White, both sexes	36	-	-	-	-	-	1	-	-	1	1	1
Male	16	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	1	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	383	3	-	1	-	-	-	3	6	9	12	15
Male	150	3	-	-	-	-	-	-	2	2	3	7
Female	233	-	-	1	-	-	-	3	4	7	9	8
White, both sexes	275	3	-	-	-	-	-	-	1	7	5	7
Male	103	3	-	-	-	-	-	-	-	2	-	4
Female	172	-	-	-	-	-	-	-	1	5	5	3
Black, both sexes	105	-	-	1	-	-	-	3	5	2	7	8
Male	46	-	-	-	-	-	-	-	2	-	3	3
Female	59	-	-	1	-	-	-	3	3	2	4	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Washington												
All races, both sexes	262	-	1	-	-	1	-	2	5	4	9	12
Male	109	-	-	-	-	-	-	-	1	3	4	5
Female	153	-	1	-	-	1	-	2	4	1	5	7
White, both sexes	237	-	1	-	-	1	-	2	4	4	9	10
Male	98	-	-	-	-	-	-	-	1	3	4	4
Female	139	-	1	-	-	1	-	2	3	1	5	6
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	1	-	-	-
West Virginia												
All races, both sexes	103	-	1	-	-	-	1	-	2	1	-	6
Male	43	-	1	-	-	-	-	-	2	1	-	4
Female	60	-	-	-	-	-	1	-	-	-	-	2
White, both sexes	101	-	1	-	-	-	1	-	2	1	-	6
Male	42	-	1	-	-	-	-	-	2	1	-	4
Female	59	-	-	-	-	-	1	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Wisconsin												
All races, both sexes	296	2	-	-	-	2	2	1	1	5	9	8
Male	131	2	-	-	-	1	2	-	1	3	5	4
Female	165	-	-	-	-	1	-	1	-	2	4	4
White, both sexes	277	2	-	-	-	2	1	-	1	4	8	5
Male	123	2	-	-	-	1	1	-	1	3	4	3
Female	154	-	-	-	-	1	-	-	-	1	4	2
Black, both sexes	13	-	-	-	-	-	1	1	-	1	-	3
Male	5	-	-	-	-	-	1	-	-	-	-	1
Female	8	-	-	-	-	-	-	1	-	1	-	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	43	1	-	-	-	-	-	-	2	2	1	2
Male	13	1	-	-	-	-	-	-	1	2	1	-
Female	30	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	42	1	-	-	-	-	-	-	2	2	1	2
Male	13	1	-	-	-	-	-	-	1	2	1	-
Female	29	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
United States												
All races, both sexes	742	903	983	1,160	1,433	1,651	1,944	1,618	855	264	37	-
Male	364	424	438	546	610	683	715	498	210	52	6	-
Female	378	479	545	614	823	968	1,229	1,120	645	212	31	-
White, both sexes	556	706	763	941	1,206	1,414	1,704	1,449	781	227	32	-
Male	282	348	346	451	523	593	637	440	188	43	6	-
Female	274	358	417	490	683	821	1,067	1,009	593	184	26	-
Black, both sexes	174	178	204	203	201	219	213	140	68	35	5	-
Male	73	71	84	90	77	86	69	46	19	8	-	-
Female	101	107	120	113	124	133	144	94	49	27	5	-
American Indian 1/, both sexes	5	10	4	7	10	1	8	5	4	-	-	-
Male	4	3	3	1	5	-	4	3	1	-	-	-
Female	1	7	1	6	5	1	4	2	3	-	-	-
Asian or Pacific Islander 2/, both sexes	7	9	12	9	16	17	19	24	2	2	-	-
Male	5	2	5	4	5	4	5	9	2	1	-	-
Female	2	7	7	5	11	13	14	15	-	1	-	-
Alabama												
All races, both sexes	18	29	36	29	30	52	40	28	13	11	-	-
Male	6	12	15	11	11	23	18	7	3	-	-	-
Female	12	17	21	18	19	29	22	21	10	11	-	-
White, both sexes	13	16	22	20	24	34	29	23	9	8	-	-
Male	5	6	9	7	9	17	12	7	2	-	-	-
Female	8	10	13	13	15	17	17	16	7	8	-	-
Black, both sexes	5	13	14	9	6	18	11	5	4	3	-	-
Male	1	6	6	4	2	6	6	-	1	-	-	-
Female	4	7	8	5	4	12	5	5	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Alaska												
All races, both sexes	1	-	2	2	1	1	4	2	-	-	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	2	2	1	-	3	2	-	-	-	-
White, both sexes	-	-	2	1	1	1	4	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	2	1	1	-	3	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	19	15	11	18	31	31	24	19	14	3	1	-
Male	10	10	7	9	6	17	11	7	4	1	-	-
Female	9	5	4	9	25	14	13	12	10	2	1	-
White, both sexes	16	14	11	15	31	30	24	18	14	3	1	-
Male	10	9	7	7	6	17	11	6	4	1	-	-
Female	6	5	4	8	25	13	13	12	10	2	1	-
Black, both sexes	2	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	2	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Arkansas												
All races, both sexes	13	12	8	19	16	21	38	22	20	3	1	-
Male	8	6	7	11	5	9	16	7	6	1	-	-
Female	5	6	1	8	11	12	22	15	14	2	1	-
White, both sexes	10	11	7	16	14	19	36	16	17	3	-	-
Male	5	6	6	9	5	9	15	5	4	1	-	-
Female	5	5	1	7	9	10	21	11	13	2	-	-
Black, both sexes	3	1	1	3	2	2	2	6	3	-	1	-
Male	3	-	1	2	-	-	1	2	2	-	-	-
Female	-	1	-	1	2	2	1	4	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	45	54	59	55	69	81	103	100	53	14	2	-
Male	26	25	25	23	29	29	34	23	14	3	1	-
Female	19	29	34	32	40	52	69	77	39	11	1	-
White, both sexes	38	47	49	44	56	60	88	84	48	11	2	-
Male	24	22	24	19	26	22	27	18	13	1	1	-
Female	14	25	25	25	30	38	61	66	35	10	1	-
Black, both sexes	3	5	7	9	8	13	7	6	4	2	-	-
Male	-	3	1	3	2	4	4	2	-	1	-	-
Female	3	2	6	6	6	9	3	4	4	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	2	3	2	5	8	7	10	1	1	-	-
Male	1	-	-	1	1	3	2	3	1	1	-	-
Female	2	2	3	1	4	5	5	7	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Colorado												
All races, both sexes	11	19	16	17	19	20	42	41	18	4	1	-
Male	3	10	8	14	12	9	19	16	4	-	-	-
Female	8	9	8	3	7	11	23	25	14	4	1	-
White, both sexes	9	16	15	16	18	19	39	40	18	4	1	-
Male	3	8	8	13	11	9	17	15	4	-	-	-
Female	6	8	7	3	7	10	22	25	14	4	1	-
Black, both sexes	2	1	1	1	-	1	2	1	-	-	-	-
Male	-	1	-	1	-	-	1	1	-	-	-	-
Female	2	-	1	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	6	11	2	10	14	19	25	24	14	3	3	-
Male	4	4	1	5	7	12	9	9	2	1	-	-
Female	2	7	1	5	7	7	16	15	12	2	3	-
White, both sexes	5	8	2	7	11	18	24	24	14	3	3	-
Male	4	2	1	3	6	12	9	9	2	1	-	-
Female	1	6	1	4	5	6	15	15	12	2	3	-
Black, both sexes	1	3	-	3	3	1	1	-	-	-	-	-
Male	-	2	-	2	1	-	-	-	-	-	-	-
Female	1	1	-	1	2	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Delaware												
All races, both sexes	4	3	1	6	4	13	7	5	3	-	-	-
Male	3	2	1	2	1	8	2	-	1	-	-	-
Female	1	1	-	4	3	5	5	5	2	-	-	-
White, both sexes	3	3	-	5	2	12	7	5	2	-	-	-
Male	2	2	-	2	1	7	2	-	1	-	-	-
Female	1	1	-	3	1	5	5	5	1	-	-	-
Black, both sexes	1	-	-	1	2	1	-	-	1	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	2	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	1	-	-	2	3	6	4	4	1	-	-
Male	-	-	-	-	1	2	1	1	1	-	-	-
Female	1	1	-	-	1	1	5	3	3	1	-	-
White, both sexes	-	-	-	-	-	1	-	2	-	1	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	1	1	-	-	2	2	6	2	4	-	-	-
Male	-	-	-	-	1	1	1	-	1	-	-	-
Female	1	1	-	-	1	1	5	2	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Florida												
All races, both sexes	34	45	56	67	89	92	118	94	52	15	5	-
Male	15	16	25	29	36	48	44	41	13	8	2	-
Female	19	29	31	38	53	44	74	53	39	7	3	-
White, both sexes	24	34	42	59	75	85	107	88	50	13	4	-
Male	11	16	20	28	29	44	40	37	13	6	2	-
Female	13	18	22	31	46	41	67	51	37	7	2	-
Black, both sexes	9	10	14	8	14	7	11	6	2	2	1	-
Male	3	-	5	1	7	4	4	4	-	2	-	-
Female	6	10	9	7	7	3	7	2	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	28	26	25	46	44	55	56	39	23	9	2	-
Male	15	14	15	17	22	22	19	8	4	2	-	-
Female	13	12	10	29	22	33	37	31	19	7	2	-
White, both sexes	16	16	18	32	32	45	41	32	18	5	2	-
Male	11	10	11	11	19	18	14	6	4	-	-	-
Female	5	6	7	21	13	27	27	26	14	5	2	-
Black, both sexes	12	10	7	14	11	9	13	6	5	4	-	-
Male	4	4	4	6	2	4	4	2	-	2	-	-
Female	8	6	3	8	9	5	9	4	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Hawaii												
All races, both sexes	1	-	1	1	2	5	5	4	1	1	-	-
Male	-	-	-	1	-	1	4	2	1	-	-	-
Female	1	-	1	-	2	4	1	2	-	1	-	-
White, both sexes	1	-	-	-	-	1	2	1	1	1	-	-
Male	-	-	-	-	-	1	2	1	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	4	3	3	-	-	-	-
Male	-	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	1	-	2	4	1	2	-	-	-	-
Idaho												
All races, both sexes	4	3	8	3	6	9	12	17	3	3	-	-
Male	-	-	4	2	4	2	5	7	1	-	-	-
Female	4	3	4	1	2	7	7	10	2	3	-	-
White, both sexes	4	3	8	3	6	9	12	16	3	3	-	-
Male	-	-	4	2	4	2	5	7	1	-	-	-
Female	4	3	4	1	2	7	7	9	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Illinois												
All races, both sexes	37	45	53	53	61	78	93	80	42	15	2	-
Male	17	23	24	30	26	28	26	23	14	1	-	-
Female	20	22	29	23	35	50	67	57	28	14	2	-
White, both sexes	26	30	39	42	50	63	79	70	39	14	2	-
Male	12	17	20	24	24	21	23	22	13	1	-	-
Female	14	13	19	18	26	42	56	48	26	13	2	-
Black, both sexes	11	14	10	10	10	14	13	10	2	-	-	-
Male	5	6	3	6	2	7	3	1	-	-	-	-
Female	6	8	7	4	8	7	10	9	2	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	3	1	1	1	1	-	1	1	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	1	2	1	1	1	1	-	-	1	-	-
Indiana												
All races, both sexes	21	25	32	35	44	48	58	37	19	2	-	-
Male	12	17	13	16	15	20	20	14	5	-	-	-
Female	9	8	19	19	29	28	38	23	14	2	-	-
White, both sexes	20	24	24	33	37	45	53	37	17	2	-	-
Male	11	16	9	15	13	20	18	14	4	-	-	-
Female	9	8	15	18	24	25	35	23	13	2	-	-
Black, both sexes	1	1	8	2	7	3	5	-	2	-	-	-
Male	1	1	4	1	2	-	2	-	1	-	-	-
Female	-	-	4	1	5	3	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Iowa												
All races, both sexes	9	12	19	19	26	20	21	32	15	-	1	-
Male	8	8	7	14	9	8	6	5	1	-	-	-
Female	1	4	12	5	17	12	15	27	14	-	1	-
White, both sexes	8	11	17	19	24	20	21	31	15	-	1	-
Male	7	7	7	14	7	8	6	5	1	-	-	-
Female	1	4	10	5	17	12	15	26	14	-	1	-
Black, both sexes	1	1	2	-	1	-	-	1	-	-	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	2	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	5	8	11	7	13	16	23	29	12	8	-	-
Male	2	3	4	2	7	4	9	6	3	4	-	-
Female	3	5	7	5	6	12	14	23	9	4	-	-
White, both sexes	5	7	10	6	11	16	23	28	12	8	-	-
Male	2	2	4	1	5	4	9	6	3	4	-	-
Female	3	5	6	5	6	12	14	22	9	4	-	-
Black, both sexes	-	1	1	1	2	-	-	1	-	-	-	-
Male	-	1	-	1	2	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Kentucky												
All races, both sexes	12	16	16	22	37	32	36	25	14	1	-	-
Male	5	8	6	12	13	13	11	9	3	-	-	-
Female	7	8	10	10	24	19	25	16	11	1	-	-
White, both sexes	10	16	15	19	33	32	34	25	14	1	-	-
Male	4	8	6	11	13	13	11	9	3	-	-	-
Female	6	8	9	8	20	19	23	16	11	1	-	-
Black, both sexes	2	-	1	3	4	-	2	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	2	4	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	23	28	18	23	23	37	30	28	11	1	-	-
Male	13	14	6	9	11	13	8	6	3	-	-	-
Female	10	14	12	14	12	24	22	22	8	1	-	-
White, both sexes	17	15	7	15	17	23	25	23	10	-	-	-
Male	9	10	2	7	7	8	7	5	2	-	-	-
Female	8	5	5	8	10	15	18	18	8	-	-	-
Black, both sexes	6	13	10	8	6	14	5	5	1	1	-	-
Male	4	4	4	2	4	5	1	1	1	-	-	-
Female	2	9	6	6	2	9	4	4	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Maine												
All races, both sexes	3	2	3	6	5	6	9	7	4	-	-	-
Male	2	1	1	4	2	1	1	4	1	-	-	-
Female	1	1	2	2	3	5	8	3	3	-	-	-
White, both sexes	3	2	3	6	5	6	9	7	4	-	-	-
Male	2	1	1	4	2	1	1	4	1	-	-	-
Female	1	1	2	2	3	5	8	3	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	22	22	16	24	28	24	42	28	10	3	-	-
Male	13	10	8	11	12	17	18	9	2	1	-	-
Female	9	12	8	13	16	7	24	19	8	2	-	-
White, both sexes	11	16	9	11	18	17	32	21	9	-	-	-
Male	6	8	4	5	11	13	13	6	2	-	-	-
Female	5	8	5	6	7	4	19	15	7	-	-	-
Black, both sexes	10	6	7	13	9	7	10	6	1	3	-	-
Male	6	2	4	6	1	4	5	3	-	1	-	-
Female	4	4	3	7	8	3	5	3	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Massachusetts												
All races, both sexes	17	13	20	24	25	41	56	45	25	9	-	-
Male	12	7	10	12	10	15	19	12	5	1	-	-
Female	5	6	10	12	15	26	37	33	20	8	-	-
White, both sexes	14	12	18	23	24	40	55	44	24	8	-	-
Male	10	7	9	12	10	14	19	12	5	1	-	-
Female	4	5	9	11	14	26	36	32	19	7	-	-
Black, both sexes	3	1	2	1	1	1	-	1	1	1	-	-
Male	2	-	1	-	-	1	-	-	-	-	-	-
Female	1	1	1	1	1	-	-	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Michigan												
All races, both sexes	28	38	40	41	49	63	51	64	32	11	1	-
Male	12	16	20	17	18	26	23	27	10	2	-	-
Female	16	22	20	24	31	37	28	37	22	9	1	-
White, both sexes	23	29	34	31	41	48	45	55	27	9	1	-
Male	9	14	16	13	15	21	21	23	7	2	-	-
Female	14	15	18	18	26	27	24	32	20	7	1	-
Black, both sexes	5	8	6	9	6	15	5	8	5	2	-	-
Male	3	1	4	4	2	5	2	3	3	-	-	-
Female	2	7	2	5	4	10	3	5	2	2	-	-
American Indian 1/, both sexes	-	1	-	1	2	-	1	1	-	-	-	-
Male	-	1	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	1	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Minnesota												
All races, both sexes	9	19	8	15	20	12	42	31	19	7	-	-
Male	5	5	3	8	10	5	13	12	5	-	-	-
Female	4	14	5	7	10	7	29	19	14	7	-	-
White, both sexes	9	19	8	14	19	12	41	31	19	7	-	-
Male	5	5	3	8	9	5	13	12	5	-	-	-
Female	4	14	5	6	10	7	28	19	14	7	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	6	20	10	20	21	27	28	18	15	3	-	-
Male	2	4	4	8	11	11	9	3	7	-	-	-
Female	4	16	6	12	10	16	19	15	8	3	-	-
White, both sexes	2	9	7	14	13	23	23	14	10	2	-	-
Male	-	2	2	5	5	9	7	2	5	-	-	-
Female	2	7	5	9	8	14	16	12	5	2	-	-
Black, both sexes	4	11	3	6	8	4	5	4	5	1	-	-
Male	2	2	2	3	6	2	2	1	2	-	-	-
Female	2	9	1	3	2	2	3	3	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Missouri												
All races, both sexes	27	20	19	38	41	42	43	48	21	5	1	-
Male	16	11	5	19	18	13	20	15	7	1	-	-
Female	11	9	14	19	23	29	23	33	14	4	1	-
White, both sexes	24	17	16	35	33	35	34	45	21	5	1	-
Male	15	9	4	17	13	10	16	13	7	1	-	-
Female	9	8	12	18	20	25	18	32	14	4	1	-
Black, both sexes	3	3	3	3	8	7	8	3	-	-	-	-
Male	1	2	1	2	5	3	3	2	-	-	-	-
Female	2	1	2	1	3	4	5	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	8	1	6	8	1	10	4	4	-	1	-
Male	2	3	-	-	2	1	6	-	1	-	-	-
Female	-	5	1	6	6	-	4	4	3	-	1	-
White, both sexes	2	7	1	5	7	1	9	4	4	-	1	-
Male	2	3	-	-	1	1	6	-	1	-	-	-
Female	-	4	1	5	6	-	3	4	3	-	1	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Nebraska												
All races, both sexes	4	3	2	9	11	8	19	12	11	7	1	-
Male	-	-	1	3	4	2	9	1	3	3	-	-
Female	4	3	1	6	7	6	10	11	8	4	1	-
White, both sexes	4	1	2	8	11	7	19	12	11	7	1	-
Male	-	-	1	3	4	1	9	1	3	3	-	-
Female	4	1	1	5	7	6	10	11	8	4	1	-
Black, both sexes	-	1	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	10	2	15	13	18	11	15	11	6	2	-	-
Male	4	1	9	5	16	4	6	3	-	-	-	-
Female	6	1	6	8	2	7	9	8	6	2	-	-
White, both sexes	9	2	13	9	15	9	14	10	6	2	-	-
Male	4	1	9	2	14	4	5	3	-	-	-	-
Female	5	1	4	7	1	5	9	7	6	2	-	-
Black, both sexes	1	-	2	3	3	1	1	-	-	-	-	-
Male	-	-	-	2	2	-	1	-	-	-	-	-
Female	1	-	2	1	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
New Hampshire												
All races, both sexes	2	2	3	6	4	9	11	5	8	2	-	-
Male	1	-	2	3	2	3	5	-	1	1	-	-
Female	1	2	1	3	2	6	6	5	7	1	-	-
White, both sexes	2	2	3	6	4	9	11	5	8	2	-	-
Male	1	-	2	3	2	3	5	-	1	1	-	-
Female	1	2	1	3	2	6	6	5	7	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	23	22	31	31	36	39	65	49	25	4	1	-
Male	9	7	11	17	15	13	20	15	7	1	1	-
Female	14	15	20	14	21	26	45	34	18	3	-	-
White, both sexes	14	17	22	24	30	36	55	44	23	3	1	-
Male	7	6	7	13	11	13	20	14	7	1	1	-
Female	7	11	15	11	19	23	35	30	16	2	-	-
Black, both sexes	9	5	9	7	6	2	9	5	2	1	-	-
Male	2	1	4	4	4	-	-	1	-	-	-	-
Female	7	4	5	3	2	2	9	4	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
New Mexico												
All races, both sexes	3	8	2	9	8	6	10	5	6	2	-	-
Male	1	4	1	4	5	2	3	2	2	-	-	-
Female	2	4	1	5	3	4	7	3	4	2	-	-
White, both sexes	3	7	2	9	7	6	10	5	4	2	-	-
Male	1	4	1	4	5	2	3	2	1	-	-	-
Female	2	3	1	5	2	4	7	3	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	30	39	43	43	53	73	95	75	34	14	1	-
Male	13	19	21	21	22	31	35	23	9	4	-	-
Female	17	20	22	22	31	42	60	52	25	10	1	-
White, both sexes	21	29	33	32	46	60	84	68	32	14	1	-
Male	9	13	16	15	20	26	32	20	8	4	-	-
Female	12	16	17	17	26	34	52	48	24	10	1	-
Black, both sexes	9	9	10	10	7	13	11	5	2	-	-	-
Male	4	6	5	6	2	5	3	2	1	-	-	-
Female	5	3	5	4	5	8	8	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	1	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
North Carolina												
All races, both sexes	24	25	53	53	57	64	69	59	31	16	1	-
Male	9	12	28	24	22	21	22	18	3	1	-	-
Female	15	13	25	29	35	43	47	41	28	15	1	-
White, both sexes	10	16	35	32	43	52	52	50	22	11	-	-
Male	2	10	17	14	18	19	17	15	1	-	-	-
Female	8	6	18	18	25	33	35	35	21	11	-	-
Black, both sexes	14	8	17	20	12	12	17	8	8	5	1	-
Male	7	2	10	10	3	2	5	2	2	1	-	-
Female	7	6	7	10	9	10	12	6	6	4	1	-
American Indian 1/, both sexes	-	1	1	1	1	-	-	1	1	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	1	-	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	3	1	7	3	7	6	5	3	2	1	-
Male	-	-	1	3	3	2	2	-	-	-	-	-
Female	-	3	-	4	-	5	4	5	3	2	1	-
White, both sexes	-	3	1	7	3	7	6	5	3	2	1	-
Male	-	-	1	3	3	2	2	-	-	-	-	-
Female	-	3	-	4	-	5	4	5	3	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Ohio												
All races, both sexes	25	43	49	43	55	71	89	68	30	9	1	-
Male	9	23	19	20	24	32	35	20	4	1	-	-
Female	16	20	30	23	31	39	54	48	26	8	1	-
White, both sexes	21	38	40	38	48	63	77	63	26	8	1	-
Male	9	21	17	17	23	31	32	18	4	1	-	-
Female	12	17	23	21	25	32	45	45	22	7	1	-
Black, both sexes	4	4	9	5	6	8	11	4	4	1	-	-
Male	-	2	2	3	1	1	3	1	-	-	-	-
Female	4	2	7	2	5	7	8	3	4	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	9	15	20	14	20	19	23	21	7	1	-	-
Male	3	7	8	6	9	7	9	8	4	-	-	-
Female	6	8	12	8	11	12	14	13	3	1	-	-
White, both sexes	5	13	15	14	16	18	19	19	7	1	-	-
Male	2	7	4	6	6	6	8	7	4	-	-	-
Female	3	6	11	8	10	12	11	12	3	1	-	-
Black, both sexes	3	1	3	-	2	-	3	2	-	-	-	-
Male	-	-	2	-	1	-	-	1	-	-	-	-
Female	3	1	1	-	1	-	3	1	-	-	-	-
American Indian 1/, both sexes	1	1	2	-	2	-	1	-	-	-	-	-
Male	1	-	2	-	2	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Oregon												
All races, both sexes	4	11	13	10	20	18	26	24	7	6	-	-
Male	2	9	9	3	9	9	12	6	2	-	-	-
Female	2	2	4	7	11	9	14	18	5	6	-	-
White, both sexes	3	10	12	8	20	18	26	22	7	6	-	-
Male	2	8	9	3	9	9	12	6	2	-	-	-
Female	1	2	3	5	11	9	14	16	5	6	-	-
Black, both sexes	1	-	1	2	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	2	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	38	41	48	73	93	92	108	101	57	18	2	-
Male	20	20	17	35	38	34	46	25	16	4	1	-
Female	18	21	31	38	55	58	62	76	41	14	1	-
White, both sexes	27	34	38	60	73	85	95	95	56	17	2	-
Male	14	15	15	32	32	32	41	24	15	4	1	-
Female	13	19	23	28	41	53	54	71	41	13	1	-
Black, both sexes	10	7	10	12	20	7	13	6	1	1	-	-
Male	5	5	2	3	6	2	5	1	1	-	-	-
Female	5	2	8	9	14	5	8	5	-	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Rhode Island												
All races, both sexes	2	6	2	2	6	8	6	10	2	-	-	-
Male	2	2	-	-	-	5	-	4	1	-	-	-
Female	-	4	2	2	6	3	6	6	1	-	-	-
White, both sexes	1	6	2	2	6	8	5	10	2	-	-	-
Male	1	2	-	-	-	5	-	4	1	-	-	-
Female	-	4	2	2	6	3	5	6	1	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	17	23	26	20	20	33	28	20	14	2	2	-
Male	6	8	8	7	8	13	11	6	2	-	-	-
Female	11	15	18	13	12	20	17	14	12	2	2	-
White, both sexes	10	12	15	14	16	22	20	15	14	2	1	-
Male	5	4	5	6	6	9	8	4	2	-	-	-
Female	5	8	10	8	10	13	12	11	12	2	1	-
Black, both sexes	7	11	11	6	4	11	8	5	-	-	1	-
Male	1	4	3	1	2	4	3	2	-	-	-	-
Female	6	7	8	5	2	7	5	3	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
South Dakota												
All races, both sexes	5	2	3	2	-	9	9	4	1	3	-	-
Male	4	1	1	2	-	3	6	3	-	-	-	-
Female	1	1	2	-	-	6	3	1	1	3	-	-
White, both sexes	4	2	3	2	-	9	9	4	-	3	-	-
Male	3	1	1	2	-	3	6	3	-	-	-	-
Female	1	1	2	-	-	6	3	1	-	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	19	29	26	30	46	45	40	30	28	5	-	-
Male	11	15	8	14	22	18	13	6	3	1	-	-
Female	8	14	18	16	24	27	27	24	25	4	-	-
White, both sexes	15	26	21	28	42	38	34	26	25	4	-	-
Male	10	13	8	13	20	14	13	4	3	1	-	-
Female	5	13	13	15	22	24	21	22	22	3	-	-
Black, both sexes	4	3	5	2	4	7	6	4	3	1	-	-
Male	1	2	-	1	2	4	-	2	-	-	-	-
Female	3	1	5	1	2	3	6	2	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Texas												
All races, both sexes	45	64	65	104	110	114	117	90	44	17	3	-
Male	20	29	33	55	49	52	42	31	11	3	-	-
Female	25	35	32	49	61	62	75	59	33	14	3	-
White, both sexes	34	50	43	83	94	91	100	76	39	11	2	-
Male	12	24	20	44	41	40	37	24	9	2	-	-
Female	22	26	23	39	53	51	63	52	30	9	2	-
Black, both sexes	10	13	20	19	13	23	17	13	5	6	1	-
Male	7	4	11	10	7	12	5	6	2	1	-	-
Female	3	9	9	9	6	11	12	7	3	5	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	2	2	-	-	1	-	-	-	-
Male	1	1	2	1	1	-	-	1	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Utah												
All races, both sexes	8	5	8	8	12	13	15	18	12	1	1	-
Male	3	1	4	4	5	6	10	7	2	1	1	-
Female	5	4	4	4	7	7	5	11	10	-	-	-
White, both sexes	8	5	8	8	11	13	15	18	12	1	1	-
Male	3	1	4	4	5	6	10	7	2	1	1	-
Female	5	4	4	4	6	7	5	11	10	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Vermont												
All races, both sexes	1	1	2	3	2	7	7	6	2	1	-	-
Male	-	1	-	2	1	3	5	3	-	-	-	-
Female	1	-	2	1	1	4	2	3	2	1	-	-
White, both sexes	1	1	2	3	2	7	7	6	2	1	-	-
Male	-	1	-	2	1	3	5	3	-	-	-	-
Female	1	-	2	1	1	4	2	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	28	21	32	32	45	56	55	49	12	3	1	-
Male	12	10	13	13	21	23	19	17	3	2	-	-
Female	16	11	19	19	24	33	36	32	9	1	1	-
White, both sexes	16	13	24	25	38	42	47	33	10	3	1	-
Male	6	5	10	10	17	16	15	11	2	2	-	-
Female	10	8	14	15	21	26	32	22	8	1	1	-
Black, both sexes	12	8	8	7	7	14	7	14	2	-	-	-
Male	6	5	3	3	4	7	4	5	1	-	-	-
Female	6	3	5	4	3	7	3	9	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Washington												
All races, both sexes	19	15	19	11	28	30	43	32	25	6	-	-
Male	11	11	11	7	14	10	14	11	6	1	-	-
Female	8	4	8	4	14	20	29	21	19	5	-	-
White, both sexes	16	10	18	10	24	30	38	31	24	5	-	-
Male	10	8	10	7	12	10	13	10	5	1	-	-
Female	6	2	8	3	12	20	25	21	19	4	-	-
Black, both sexes	2	4	1	1	3	-	1	-	1	1	-	-
Male	-	3	1	-	2	-	1	-	1	-	-	-
Female	2	1	-	1	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	1	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	3	-	-	-	-	-
West Virginia												
All races, both sexes	7	7	7	8	18	16	15	5	9	-	-	-
Male	2	2	6	3	5	7	3	2	5	-	-	-
Female	5	5	1	5	13	9	12	3	4	-	-	-
White, both sexes	7	7	7	7	18	16	15	4	9	-	-	-
Male	2	2	6	3	5	7	3	1	5	-	-	-
Female	5	5	1	4	13	9	12	3	4	-	-	-
Black, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)												
Wisconsin												
All races, both sexes	12	20	19	22	32	47	43	42	18	10	1	-
Male	10	13	7	9	17	22	15	13	5	2	-	-
Female	2	7	12	13	15	25	28	29	13	8	1	-
White, both sexes	11	18	17	20	30	46	42	41	18	10	1	-
Male	10	12	5	8	17	21	15	13	5	2	-	-
Female	1	6	12	12	13	25	27	28	13	8	1	-
Black, both sexes	1	-	1	1	2	1	1	-	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	1	-	-	-	2	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	2	3	1	8	7	8	2	2	1	-	-
Male	-	-	1	-	1	3	1	1	-	1	-	-
Female	1	2	2	1	7	4	7	1	2	-	-	-
White, both sexes	1	2	3	1	7	7	8	2	2	1	-	-
Male	-	-	1	-	1	3	1	1	-	1	-	-
Female	1	2	2	1	6	4	7	1	2	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
United States												
All races, both sexes	8,283	4	-	2	4	28	53	87	129	204	295	382
Male	3,543	3	-	-	2	9	24	35	47	105	145	178
Female	4,740	1	-	2	2	19	29	52	82	99	150	204
White, both sexes	6,778	4	-	1	3	21	34	49	71	143	201	282
Male	2,896	3	-	-	1	6	15	20	30	77	99	130
Female	3,882	1	-	1	2	15	19	29	41	66	102	152
Black, both sexes	1,394	-	-	1	1	7	18	36	53	57	91	95
Male	596	-	-	-	1	3	8	13	15	26	44	46
Female	798	-	-	1	-	4	10	23	38	31	47	49
American Indian 1/, both sexes	38	-	-	-	-	-	-	1	1	1	2	3
Male	16	-	-	-	-	-	-	1	-	-	1	1
Female	22	-	-	-	-	-	-	-	1	1	1	2
Asian or Pacific Islander 2/, both sexes	73	-	-	-	-	-	1	1	4	3	1	2
Male	35	-	-	-	-	-	1	1	2	2	1	1
Female	38	-	-	-	-	-	-	-	2	1	-	1
Alabama												
All races, both sexes	226	-	-	-	-	1	-	3	3	5	4	13
Male	85	-	-	-	-	-	-	2	2	4	2	5
Female	141	-	-	-	-	1	-	1	1	1	2	8
White, both sexes	157	-	-	-	-	1	-	-	3	4	3	6
Male	58	-	-	-	-	-	-	-	2	4	1	1
Female	99	-	-	-	-	1	-	-	1	-	2	5
Black, both sexes	69	-	-	-	-	-	-	3	-	1	1	7
Male	27	-	-	-	-	-	-	2	-	-	1	4
Female	42	-	-	-	-	-	-	1	-	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Alaska												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	126	-	-	-	-	-	2	3	1	1	3	6
Male	56	-	-	-	-	-	2	-	-	1	1	2
Female	70	-	-	-	-	-	-	3	1	-	2	4
White, both sexes	119	-	-	-	-	-	2	2	-	1	3	6
Male	54	-	-	-	-	-	2	-	-	1	1	2
Female	65	-	-	-	-	-	-	2	-	-	2	4
Black, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Arkansas												
All races, both sexes	140	1	-	-	-	-	1	1	3	4	5	8
Male	66	1	-	-	-	-	1	-	1	1	2	6
Female	74	-	-	-	-	-	-	1	2	3	3	2
White, both sexes	119	1	-	-	-	-	1	-	2	3	4	7
Male	57	1	-	-	-	-	1	-	1	1	2	5
Female	62	-	-	-	-	-	-	-	1	2	2	2
Black, both sexes	21	-	-	-	-	-	-	1	1	1	1	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	1	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	355	-	-	-	-	3	4	3	6	6	19	12
Male	154	-	-	-	-	1	1	1	4	4	11	9
Female	201	-	-	-	-	2	3	2	2	2	8	3
White, both sexes	290	-	-	-	-	2	4	1	4	4	14	6
Male	127	-	-	-	-	1	1	-	2	3	8	5
Female	163	-	-	-	-	1	3	1	2	1	6	1
Black, both sexes	42	-	-	-	-	1	-	1	1	2	5	6
Male	16	-	-	-	-	-	-	-	1	1	3	4
Female	26	-	-	-	-	1	-	1	-	1	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	1	1	-	-	-
Male	10	-	-	-	-	-	-	1	1	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Colorado												
All races, both sexes	124	-	-	-	-	-	1	-	2	1	3	6
Male	65	-	-	-	-	-	1	-	2	-	-	5
Female	59	-	-	-	-	-	-	-	-	1	3	1
White, both sexes	111	-	-	-	-	-	-	-	2	-	2	6
Male	58	-	-	-	-	-	-	-	2	-	-	5
Female	53	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	11	-	-	-	-	-	1	-	-	1	1	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	100	-	-	-	-	-	1	1	3	1	4	7
Male	38	-	-	-	-	-	-	-	1	1	3	4
Female	62	-	-	-	-	-	1	1	2	-	1	3
White, both sexes	88	-	-	-	-	-	1	1	1	1	3	5
Male	33	-	-	-	-	-	-	-	1	1	2	3
Female	55	-	-	-	-	-	1	1	-	-	1	2
Black, both sexes	12	-	-	-	-	-	-	-	2	-	1	2
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	2	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Delaware												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	3	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	2	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Florida												
All races, both sexes	458	-	-	-	-	-	-	3	4	15	9	23
Male	193	-	-	-	-	-	-	2	1	7	4	9
Female	265	-	-	-	-	-	-	1	3	8	5	14
White, both sexes	384	-	-	-	-	-	-	2	2	12	6	15
Male	159	-	-	-	-	-	-	1	-	5	2	5
Female	225	-	-	-	-	-	-	1	2	7	4	10
Black, both sexes	73	-	-	-	-	-	-	1	2	3	3	8
Male	34	-	-	-	-	-	-	1	1	2	2	4
Female	39	-	-	-	-	-	-	-	1	1	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	266	-	-	-	-	1	3	6	8	15	16	11
Male	106	-	-	-	-	-	-	1	2	7	11	4
Female	160	-	-	-	-	1	3	5	6	8	5	7
White, both sexes	176	-	-	-	-	-	1	2	2	8	7	6
Male	68	-	-	-	-	-	-	-	-	3	4	2
Female	108	-	-	-	-	-	1	2	2	5	3	4
Black, both sexes	89	-	-	-	-	1	2	4	6	7	9	5
Male	37	-	-	-	-	-	-	1	2	4	7	2
Female	52	-	-	-	-	1	2	3	4	3	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Hawaii												
All races, both sexes	16	-	-	-	-	-	-	1	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	58	-	-	-	-	-	-	1	-	4	3	6
Male	27	-	-	-	-	-	-	-	-	3	3	2
Female	31	-	-	-	-	-	-	1	-	1	-	4
White, both sexes	57	-	-	-	-	-	-	1	-	4	3	6
Male	27	-	-	-	-	-	-	-	-	3	3	2
Female	30	-	-	-	-	-	-	1	-	1	-	4
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Illinois												
All races, both sexes	428	-	-	-	-	1	2	4	8	9	19	21
Male	188	-	-	-	-	-	2	1	4	5	14	11
Female	240	-	-	-	-	1	-	3	4	4	5	10
White, both sexes	339	-	-	-	-	-	2	3	6	6	10	15
Male	151	-	-	-	-	-	2	-	4	4	8	8
Female	188	-	-	-	-	-	-	3	2	2	2	7
Black, both sexes	83	-	-	-	-	1	-	1	2	3	9	6
Male	36	-	-	-	-	-	-	1	-	1	6	3
Female	47	-	-	-	-	1	-	-	2	2	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	255	-	-	-	-	1	2	4	11	8	12	17
Male	116	-	-	-	-	-	1	3	3	3	8	6
Female	139	-	-	-	-	1	1	1	8	5	4	11
White, both sexes	226	-	-	-	-	1	2	2	9	7	10	16
Male	101	-	-	-	-	-	1	2	1	3	7	5
Female	125	-	-	-	-	1	1	-	8	4	3	11
Black, both sexes	28	-	-	-	-	-	-	2	1	1	2	1
Male	14	-	-	-	-	-	-	1	1	-	1	1
Female	14	-	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Iowa												
All races, both sexes	124	-	-	-	-	1	2	-	2	4	4	3
Male	58	-	-	-	-	1	-	-	1	3	4	2
Female	66	-	-	-	-	-	2	-	1	1	-	1
White, both sexes	118	-	-	-	-	1	2	-	1	4	4	3
Male	55	-	-	-	-	1	-	-	1	3	4	2
Female	63	-	-	-	-	-	2	-	-	1	-	1
Black, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	94	-	-	-	-	-	3	1	1	1	1	5
Male	39	-	-	-	-	-	1	1	-	-	-	4
Female	55	-	-	-	-	-	2	-	1	1	1	1
White, both sexes	85	-	-	-	-	-	1	1	-	1	-	4
Male	35	-	-	-	-	-	1	1	-	-	-	3
Female	50	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	9	-	-	-	-	-	2	-	1	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	2	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Kentucky												
All races, both sexes	164	-	-	-	-	-	-	4	-	3	9	9
Male	74	-	-	-	-	-	-	3	-	2	4	6
Female	90	-	-	-	-	-	-	1	-	1	5	3
White, both sexes	151	-	-	-	-	-	-	3	-	2	7	9
Male	71	-	-	-	-	-	-	2	-	1	4	6
Female	80	-	-	-	-	-	-	1	-	1	3	3
Black, both sexes	13	-	-	-	-	-	-	1	-	1	2	-
Male	3	-	-	-	-	-	-	1	-	1	-	-
Female	10	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	187	-	-	-	-	-	3	4	2	3	12	9
Male	79	-	-	-	-	-	-	2	-	2	5	6
Female	108	-	-	-	-	-	3	2	2	1	7	3
White, both sexes	125	-	-	-	-	-	2	2	-	3	6	6
Male	55	-	-	-	-	-	-	1	-	2	4	4
Female	70	-	-	-	-	-	2	1	-	1	2	2
Black, both sexes	60	-	-	-	-	-	1	1	2	-	5	3
Male	23	-	-	-	-	-	-	-	-	-	1	2
Female	37	-	-	-	-	-	1	1	2	-	4	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	1	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Maine												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	2	1
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	27	-	-	-	-	-	-	-	-	-	2	1
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	181	-	-	-	-	-	1	2	5	1	11	11
Male	84	-	-	-	-	-	1	2	1	-	5	5
Female	97	-	-	-	-	-	-	-	4	1	6	6
White, both sexes	115	-	-	-	-	-	1	1	3	-	4	7
Male	56	-	-	-	-	-	1	1	1	-	2	3
Female	59	-	-	-	-	-	-	-	2	-	2	4
Black, both sexes	63	-	-	-	-	-	-	1	2	1	7	4
Male	27	-	-	-	-	-	-	1	-	-	3	2
Female	36	-	-	-	-	-	-	-	2	1	4	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Massachusetts												
All races, both sexes	195	1	-	-	-	1	-	2	2	5	7	7
Male	78	1	-	-	-	1	-	-	-	2	3	5
Female	117	-	-	-	-	-	-	2	2	3	4	2
White, both sexes	182	1	-	-	-	1	-	1	1	4	6	7
Male	74	1	-	-	-	1	-	-	-	2	2	5
Female	108	-	-	-	-	-	-	1	1	2	4	2
Black, both sexes	12	-	-	-	-	-	-	1	1	1	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	277	-	-	-	-	-	1	1	5	4	4	6
Male	125	-	-	-	-	-	1	1	2	1	2	-
Female	152	-	-	-	-	-	-	-	3	3	2	6
White, both sexes	222	-	-	-	-	-	-	-	2	1	2	5
Male	98	-	-	-	-	-	-	-	1	1	1	-
Female	124	-	-	-	-	-	-	-	1	-	1	5
Black, both sexes	50	-	-	-	-	-	1	1	3	2	2	1
Male	25	-	-	-	-	-	1	1	1	-	1	-
Female	25	-	-	-	-	-	-	-	2	2	1	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Minnesota												
All races, both sexes	108	-	-	-	-	-	-	1	1	-	4	7
Male	43	-	-	-	-	-	-	-	-	-	-	3
Female	65	-	-	-	-	-	-	1	1	-	4	4
White, both sexes	106	-	-	-	-	-	-	1	1	-	4	6
Male	43	-	-	-	-	-	-	-	-	-	-	3
Female	63	-	-	-	-	-	-	1	1	-	4	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	147	-	-	-	-	-	5	4	6	3	3	7
Male	53	-	-	-	-	-	1	1	2	1	1	2
Female	94	-	-	-	-	-	4	3	4	2	2	5
White, both sexes	92	-	-	-	-	-	1	-	2	1	2	2
Male	32	-	-	-	-	-	-	-	1	-	-	-
Female	60	-	-	-	-	-	1	-	1	1	2	2
Black, both sexes	55	-	-	-	-	-	4	4	4	2	1	5
Male	21	-	-	-	-	-	1	1	1	1	1	2
Female	34	-	-	-	-	-	3	3	3	1	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Missouri												
All races, both sexes	247	-	-	1	1	-	-	1	-	7	8	12
Male	107	-	-	-	-	-	-	1	-	4	4	5
Female	140	-	-	1	1	-	-	-	-	3	4	7
White, both sexes	213	-	-	1	1	-	-	1	-	6	6	7
Male	90	-	-	-	-	-	-	1	-	4	3	3
Female	123	-	-	1	1	-	-	-	-	2	3	4
Black, both sexes	33	-	-	-	-	-	-	-	-	1	2	5
Male	16	-	-	-	-	-	-	-	-	-	1	2
Female	17	-	-	-	-	-	-	-	-	1	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	26	-	-	-	-	1	-	1	-	1	1	1
Male	11	-	-	-	-	-	-	1	-	-	1	-
Female	15	-	-	-	-	1	-	-	-	1	-	1
White, both sexes	24	-	-	-	-	1	-	1	-	1	1	1
Male	11	-	-	-	-	-	-	1	-	-	1	-
Female	13	-	-	-	-	1	-	-	-	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Nebraska												
All races, both sexes	52	-	-	-	-	-	-	1	1	1	2	1
Male	15	-	-	-	-	-	-	-	-	1	1	-
Female	37	-	-	-	-	-	-	1	1	-	1	1
White, both sexes	47	-	-	-	-	-	-	1	-	-	2	1
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	34	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	78	-	-	-	-	-	-	-	-	2	2	4
Male	37	-	-	-	-	-	-	-	-	1	1	3
Female	41	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	66	-	-	-	-	-	-	-	-	2	-	4
Male	30	-	-	-	-	-	-	-	-	1	-	3
Female	36	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	-	2	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
New Hampshire												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	2	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	2	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	235	-	-	-	1	1	3	3	2	6	12	4
Male	81	-	-	-	1	1	1	-	-	4	4	2
Female	154	-	-	-	-	-	2	3	2	2	8	2
White, both sexes	180	-	-	-	-	-	3	1	-	4	6	2
Male	63	-	-	-	-	-	1	-	-	3	2	1
Female	117	-	-	-	-	-	2	1	-	1	4	1
Black, both sexes	53	-	-	-	1	1	-	2	2	2	5	2
Male	17	-	-	-	1	1	-	-	-	1	1	1
Female	36	-	-	-	-	-	-	2	2	1	4	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
New Mexico												
All races, both sexes	39	-	-	-	-	-	-	2	-	1	2	3
Male	13	-	-	-	-	-	-	-	-	1	-	1
Female	26	-	-	-	-	-	-	2	-	-	2	2
White, both sexes	35	-	-	-	-	-	-	1	-	1	2	2
Male	12	-	-	-	-	-	-	-	-	1	-	1
Female	23	-	-	-	-	-	-	1	-	-	2	1
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	329	-	-	-	-	-	1	1	2	12	13	14
Male	150	-	-	-	-	-	1	1	1	5	9	7
Female	179	-	-	-	-	-	-	-	1	7	4	7
White, both sexes	262	-	-	-	-	-	-	1	-	8	11	8
Male	118	-	-	-	-	-	-	1	-	4	7	5
Female	144	-	-	-	-	-	-	-	-	4	4	3
Black, both sexes	66	-	-	-	-	-	1	-	2	4	2	6
Male	32	-	-	-	-	-	1	-	1	1	2	2
Female	34	-	-	-	-	-	-	-	1	3	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
North Carolina												
All races, both sexes	328	1	-	-	-	4	2	1	4	14	12	11
Male	131	-	-	-	-	-	2	1	1	9	5	6
Female	197	1	-	-	-	4	-	-	3	5	7	5
White, both sexes	233	1	-	-	-	4	-	1	2	8	4	6
Male	92	-	-	-	-	-	-	1	1	4	2	3
Female	141	1	-	-	-	4	-	-	1	4	2	3
Black, both sexes	91	-	-	-	-	-	1	-	2	6	8	5
Male	38	-	-	-	-	-	1	-	-	5	3	3
Female	53	-	-	-	-	-	-	-	2	1	5	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Ohio												
All races, both sexes	343	1	-	-	-	1	1	7	9	11	9	23
Male	133	1	-	-	-	-	1	2	2	1	4	6
Female	210	-	-	-	-	1	-	5	7	10	5	17
White, both sexes	299	1	-	-	-	1	1	4	6	9	8	17
Male	122	1	-	-	-	-	1	1	2	1	4	3
Female	177	-	-	-	-	1	-	3	4	8	4	14
Black, both sexes	42	-	-	-	-	-	-	3	3	2	1	6
Male	10	-	-	-	-	-	-	1	-	-	-	3
Female	32	-	-	-	-	-	-	2	3	2	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	116	-	-	-	-	3	2	1	-	1	3	2
Male	48	-	-	-	-	-	-	-	-	1	2	1
Female	68	-	-	-	-	3	2	1	-	-	1	1
White, both sexes	102	-	-	-	-	3	2	1	-	1	2	2
Male	40	-	-	-	-	-	-	-	-	1	1	1
Female	62	-	-	-	-	3	2	1	-	-	1	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Oregon												
All races, both sexes	85	-	-	-	1	-	1	2	1	2	2	5
Male	36	-	-	-	-	-	1	1	1	1	1	1
Female	49	-	-	-	1	-	-	1	-	1	1	4
White, both sexes	81	-	-	-	1	-	1	2	1	1	2	5
Male	36	-	-	-	-	-	1	1	1	1	1	1
Female	45	-	-	-	1	-	-	1	-	-	1	4
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Pennsylvania												
All races, both sexes	477	-	-	-	-	4	5	3	6	6	17	21
Male	209	-	-	-	-	2	2	3	4	3	5	11
Female	268	-	-	-	-	2	3	-	2	3	12	10
White, both sexes	405	-	-	-	-	3	4	3	4	4	12	17
Male	184	-	-	-	-	2	2	3	3	1	5	9
Female	221	-	-	-	-	1	2	-	1	3	7	8
Black, both sexes	69	-	-	-	-	1	1	-	2	1	5	4
Male	23	-	-	-	-	-	-	-	1	1	-	2
Female	46	-	-	-	-	1	1	-	1	-	5	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Rhode Island												
All races, both sexes	18	-	-	-	-	-	-	-	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	1	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	148	-	-	-	-	-	-	1	3	7	7	7
Male	63	-	-	-	-	-	-	-	3	4	6	3
Female	85	-	-	-	-	-	-	1	-	3	1	4
White, both sexes	98	-	-	-	-	-	-	1	1	2	4	5
Male	39	-	-	-	-	-	-	-	1	1	3	1
Female	59	-	-	-	-	-	-	1	-	1	1	4
Black, both sexes	50	-	-	-	-	-	-	-	2	5	3	2
Male	24	-	-	-	-	-	-	-	2	3	3	2
Female	26	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
South Dakota												
All races, both sexes	30	-	-	-	-	-	-	-	2	1	1	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	2	1	1	1
White, both sexes	29	-	-	-	-	-	-	-	2	1	1	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	2	1	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	258	-	-	1	-	-	2	1	4	6	10	18
Male	101	-	-	-	-	-	1	1	1	4	4	10
Female	157	-	-	1	-	-	1	-	3	2	6	8
White, both sexes	220	-	-	-	-	-	1	1	2	5	9	16
Male	88	-	-	-	-	-	-	1	-	4	4	9
Female	132	-	-	-	-	-	1	-	2	1	5	7
Black, both sexes	38	-	-	1	-	-	1	-	2	1	1	2
Male	13	-	-	-	-	-	1	-	1	-	-	1
Female	25	-	-	1	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Texas												
All races, both sexes	580	-	-	-	1	4	3	8	8	12	14	24
Male	256	-	-	-	1	3	1	4	2	7	3	9
Female	324	-	-	-	-	1	2	4	6	5	11	15
White, both sexes	448	-	-	-	1	2	1	4	3	5	8	20
Male	186	-	-	-	1	1	-	2	-	3	1	9
Female	262	-	-	-	-	1	1	2	3	2	7	11
Black, both sexes	125	-	-	-	-	2	2	4	5	7	6	4
Male	65	-	-	-	-	2	1	2	2	4	2	-
Female	60	-	-	-	-	-	1	2	3	3	4	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	61	-	-	-	-	-	1	1	2	2	-	4
Male	34	-	-	-	-	-	1	-	1	1	-	3
Female	27	-	-	-	-	-	-	1	1	1	-	1
White, both sexes	60	-	-	-	-	-	1	1	2	2	-	3
Male	33	-	-	-	-	-	1	-	1	1	-	2
Female	27	-	-	-	-	-	-	1	1	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Vermont												
All races, both sexes	16	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	244	-	-	-	-	-	-	2	5	6	8	12
Male	114	-	-	-	-	-	-	-	1	1	2	5
Female	130	-	-	-	-	-	-	2	4	5	6	7
White, both sexes	178	-	-	-	-	-	-	-	1	6	5	5
Male	76	-	-	-	-	-	-	-	-	1	-	3
Female	102	-	-	-	-	-	-	-	1	5	5	2
Black, both sexes	64	-	-	-	-	-	-	2	4	-	3	7
Male	37	-	-	-	-	-	-	-	1	-	2	2
Female	27	-	-	-	-	-	-	2	3	-	1	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Washington												
All races, both sexes	150	-	-	-	-	-	-	1	3	2	4	7
Male	71	-	-	-	-	-	-	-	1	2	2	3
Female	79	-	-	-	-	-	-	1	2	-	2	4
White, both sexes	134	-	-	-	-	-	-	1	2	2	4	6
Male	62	-	-	-	-	-	-	-	1	2	2	2
Female	72	-	-	-	-	-	-	1	1	-	2	4
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	1	-	-	-
West Virginia												
All races, both sexes	58	-	-	-	-	-	-	-	2	1	-	4
Male	25	-	-	-	-	-	-	-	2	1	-	2
Female	33	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	57	-	-	-	-	-	-	-	2	1	-	4
Male	25	-	-	-	-	-	-	-	2	1	-	2
Female	32	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Pulmonary embolism (I26)												
Wisconsin												
All races, both sexes	181	-	-	-	-	1	1	1	-	5	6	5
Male	85	-	-	-	-	-	1	-	-	3	4	3
Female	96	-	-	-	-	1	-	1	-	2	2	2
White, both sexes	170	-	-	-	-	1	-	-	-	4	6	4
Male	80	-	-	-	-	-	-	-	-	3	4	3
Female	90	-	-	-	-	1	-	-	-	1	2	1
Black, both sexes	9	-	-	-	-	-	1	1	-	1	-	1
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	1	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	26	-	-	-	-	-	-	-	2	2	1	2
Male	5	-	-	-	-	-	-	-	1	2	1	-
Female	21	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	25	-	-	-	-	-	-	-	2	2	1	2
Male	5	-	-	-	-	-	-	-	1	2	1	-
Female	20	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
United States												
All races, both sexes	499	573	606	712	834	1,031	1,169	978	510	161	22	-
Male	268	295	299	353	383	465	447	319	130	33	3	-
Female	231	278	307	359	451	566	722	659	380	128	19	-
White, both sexes	382	452	485	575	699	867	1,009	879	463	139	19	-
Male	212	239	239	288	323	394	391	284	115	27	3	-
Female	170	213	246	287	376	473	618	595	348	112	16	-
Black, both sexes	112	108	110	127	127	154	145	84	43	22	3	-
Male	51	52	54	61	57	68	50	28	13	6	-	-
Female	61	56	56	66	70	86	95	56	30	16	3	-
American Indian 1/, both sexes	2	8	2	5	4	1	3	2	3	-	-	-
Male	2	3	2	-	2	-	2	1	1	-	-	-
Female	-	5	-	5	2	1	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	9	5	4	9	12	13	1	-	-	-
Male	3	1	4	4	1	3	4	6	1	-	-	-
Female	-	4	5	1	3	6	8	7	-	-	-	-
Alabama												
All races, both sexes	15	19	25	21	16	41	29	16	7	8	-	-
Male	5	8	10	7	6	16	14	4	-	-	-	-
Female	10	11	15	14	10	25	15	12	7	8	-	-
White, both sexes	10	13	17	13	13	27	21	14	5	7	-	-
Male	4	5	7	4	5	12	9	4	-	-	-	-
Female	6	8	10	9	8	15	12	10	5	7	-	-
Black, both sexes	5	6	8	8	3	14	8	2	2	1	-	-
Male	1	3	3	3	1	4	5	-	-	-	-	-
Female	4	3	5	5	2	10	3	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Alaska												
All races, both sexes	1	-	1	-	1	1	1	2	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	2	-	-	-	-
White, both sexes	-	-	1	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	10	8	7	13	17	15	15	15	8	1	1	-
Male	7	6	5	6	4	7	9	6	-	-	-	-
Female	3	2	2	7	13	8	6	9	8	1	1	-
White, both sexes	9	7	7	11	17	14	15	15	8	1	1	-
Male	7	5	5	5	4	7	9	6	-	-	-	-
Female	2	2	2	6	13	7	6	9	8	1	1	-
Black, both sexes	1	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Arkansas												
All races, both sexes	12	5	4	9	9	19	27	16	13	2	1	-
Male	7	5	3	6	3	9	9	6	5	1	-	-
Female	5	-	1	3	6	10	18	10	8	1	1	-
White, both sexes	10	5	3	8	8	17	25	12	11	2	-	-
Male	5	5	2	5	3	9	8	4	4	1	-	-
Female	5	-	1	3	5	8	17	8	7	1	-	-
Black, both sexes	2	-	1	1	1	2	2	4	2	-	1	-
Male	2	-	1	1	-	-	1	2	1	-	-	-
Female	-	-	-	-	1	2	1	2	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	23	32	28	28	30	37	45	49	24	5	1	-
Male	15	15	15	13	15	14	14	15	7	-	-	-
Female	8	17	13	15	15	23	31	34	17	5	1	-
White, both sexes	20	29	24	22	22	27	41	42	22	5	1	-
Male	13	13	14	11	12	12	13	12	7	-	-	-
Female	7	16	10	11	10	15	28	30	15	5	1	-
Black, both sexes	1	3	2	5	5	5	1	2	2	-	-	-
Male	-	2	1	1	2	-	-	1	-	-	-	-
Female	1	1	1	4	3	5	1	1	2	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	1	3	5	3	5	-	-	-	-
Male	1	-	-	1	1	2	1	2	-	-	-	-
Female	-	-	2	-	2	3	2	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Colorado												
All races, both sexes	5	12	11	9	10	14	21	19	8	2	-	-
Male	2	7	5	7	6	6	11	11	2	-	-	-
Female	3	5	6	2	4	8	10	8	6	2	-	-
White, both sexes	4	10	10	8	10	13	18	18	8	2	-	-
Male	2	5	5	6	6	6	9	10	2	-	-	-
Female	2	5	5	2	4	7	9	8	6	2	-	-
Black, both sexes	1	1	1	1	-	1	2	1	-	-	-	-
Male	-	1	-	1	-	-	1	1	-	-	-	-
Female	1	-	1	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	4	8	-	6	9	11	12	20	8	2	3	-
Male	3	4	-	3	4	5	2	7	-	1	-	-
Female	1	4	-	3	5	6	10	13	8	1	3	-
White, both sexes	4	5	-	5	7	11	11	20	8	2	3	-
Male	3	2	-	2	4	5	2	7	-	1	-	-
Female	1	3	-	3	3	6	9	13	8	1	3	-
Black, both sexes	-	3	-	1	2	-	1	-	-	-	-	-
Male	-	2	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	2	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Delaware												
All races, both sexes	2	2	1	2	3	8	3	2	2	-	-	-
Male	1	1	1	1	1	6	2	-	1	-	-	-
Female	1	1	-	1	2	2	1	2	1	-	-	-
White, both sexes	2	2	-	2	2	7	3	2	1	-	-	-
Male	1	1	-	1	1	5	2	-	1	-	-	-
Female	1	1	-	1	1	2	1	2	-	-	-	-
Black, both sexes	-	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	1	2	4	3	1	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	4	2	1	1	-	-
White, both sexes	-	-	-	-	-	1	-	2	-	1	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	1	1	4	1	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	4	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Florida												
All races, both sexes	18	28	33	42	52	57	76	54	30	11	3	-
Male	7	12	16	17	24	33	30	21	5	5	-	-
Female	11	16	17	25	28	24	46	33	25	6	3	-
White, both sexes	16	23	25	34	45	50	67	48	28	9	2	-
Male	7	12	12	16	19	29	26	17	5	3	-	-
Female	9	11	13	18	26	21	41	31	23	6	2	-
Black, both sexes	2	4	8	8	7	7	9	6	2	2	1	-
Male	-	-	4	1	5	4	4	4	-	2	-	-
Female	2	4	4	7	2	3	5	2	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	19	15	15	28	22	31	29	22	19	5	1	-
Male	10	9	9	10	10	14	10	4	4	1	-	-
Female	9	6	6	18	12	17	19	18	15	4	1	-
White, both sexes	11	11	11	20	16	24	21	17	15	3	1	-
Male	8	7	7	7	8	10	6	2	4	-	-	-
Female	3	4	4	13	8	14	15	15	11	3	1	-
Black, both sexes	8	4	4	8	6	7	7	5	4	2	-	-
Male	2	2	2	3	2	4	3	2	-	1	-	-
Female	6	2	2	5	4	3	4	3	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Hawaii												
All races, both sexes	1	-	1	1	1	2	4	2	1	-	-	-
Male	-	-	-	1	-	-	3	2	1	-	-	-
Female	1	-	1	-	1	2	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	2	3	1	-	-	-	-
Male	-	-	-	1	-	-	2	1	-	-	-	-
Female	-	-	1	-	1	2	1	-	-	-	-	-
Idaho												
All races, both sexes	3	2	8	1	4	3	8	12	2	1	-	-
Male	-	-	4	1	3	-	4	6	1	-	-	-
Female	3	2	4	-	1	3	4	6	1	1	-	-
White, both sexes	3	2	8	1	4	3	8	11	2	1	-	-
Male	-	-	4	1	3	-	4	6	1	-	-	-
Female	3	2	4	-	1	3	4	5	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Illinois												
All races, both sexes	25	30	32	37	36	50	63	49	31	9	2	-
Male	16	18	15	20	20	20	18	12	11	1	-	-
Female	9	12	17	17	16	30	45	37	20	8	2	-
White, both sexes	17	18	25	29	30	43	50	46	28	9	2	-
Male	11	12	14	15	19	16	15	12	10	1	-	-
Female	6	6	11	14	11	27	35	34	18	8	2	-
Black, both sexes	8	11	5	8	6	6	12	3	2	-	-	-
Male	5	6	1	5	1	4	3	-	-	-	-	-
Female	3	5	4	3	5	2	9	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	2	-	-	1	1	-	-	-	-	-
Indiana												
All races, both sexes	16	16	19	20	28	28	36	25	10	2	-	-
Male	10	11	8	8	12	12	16	11	4	-	-	-
Female	6	5	11	12	16	16	20	14	6	2	-	-
White, both sexes	15	15	16	18	23	25	31	25	9	2	-	-
Male	9	10	6	7	10	12	14	11	3	-	-	-
Female	6	5	10	11	13	13	17	14	6	2	-	-
Black, both sexes	1	1	3	2	5	3	5	-	1	-	-	-
Male	1	1	2	1	2	-	2	-	1	-	-	-
Female	-	-	1	1	3	3	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Iowa												
All races, both sexes	8	7	16	13	19	12	9	17	7	-	-	-
Male	7	5	7	10	7	4	2	4	1	-	-	-
Female	1	2	9	3	12	8	7	13	6	-	-	-
White, both sexes	7	6	15	13	18	12	9	16	7	-	-	-
Male	6	4	7	10	6	4	2	4	1	-	-	-
Female	1	2	8	3	12	8	7	12	6	-	-	-
Black, both sexes	1	1	1	-	1	-	-	1	-	-	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	5	4	7	9	8	16	20	5	5	-	-
Male	1	3	3	2	5	2	8	5	1	3	-	-
Female	2	2	1	5	4	6	8	15	4	2	-	-
White, both sexes	3	4	4	6	8	8	16	19	5	5	-	-
Male	1	2	3	1	4	2	8	5	1	3	-	-
Female	2	2	1	5	4	6	8	14	4	2	-	-
Black, both sexes	-	1	-	1	1	-	-	1	-	-	-	-
Male	-	1	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Kentucky												
All races, both sexes	9	14	9	14	19	23	20	21	9	1	-	-
Male	5	8	4	9	6	10	6	9	2	-	-	-
Female	4	6	5	5	13	13	14	12	7	1	-	-
White, both sexes	8	14	8	12	16	23	18	21	9	1	-	-
Male	4	8	4	9	6	10	6	9	2	-	-	-
Female	4	6	4	3	10	13	12	12	7	1	-	-
Black, both sexes	1	-	1	2	3	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	2	3	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	18	15	11	16	18	29	21	17	8	1	-	-
Male	12	7	5	7	9	12	6	4	2	-	-	-
Female	6	8	6	9	9	17	15	13	6	1	-	-
White, both sexes	13	9	5	12	12	16	17	14	8	-	-	-
Male	8	5	2	6	5	7	5	4	2	-	-	-
Female	5	4	3	6	7	9	12	10	6	-	-	-
Black, both sexes	5	6	6	4	6	13	4	3	-	1	-	-
Male	4	2	3	1	4	5	1	-	-	-	-	-
Female	1	4	3	3	2	8	3	3	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Maine												
All races, both sexes	2	1	2	3	1	4	5	4	2	-	-	-
Male	1	-	1	2	1	1	1	2	-	-	-	-
Female	1	1	1	1	-	3	4	2	2	-	-	-
White, both sexes	2	1	2	3	1	4	5	4	2	-	-	-
Male	1	-	1	2	1	1	1	2	-	-	-	-
Female	1	1	1	1	-	3	4	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	15	18	12	13	18	18	27	20	7	2	-	-
Male	9	8	6	8	6	12	13	6	1	1	-	-
Female	6	10	6	5	12	6	14	14	6	1	-	-
White, both sexes	7	13	8	7	11	13	18	15	7	-	-	-
Male	4	6	4	5	5	10	8	5	1	-	-	-
Female	3	7	4	2	6	3	10	10	6	-	-	-
Black, both sexes	7	5	4	6	6	5	9	4	-	2	-	-
Male	4	2	2	3	1	2	5	1	-	1	-	-
Female	3	3	2	3	5	3	4	3	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Massachusetts												
All races, both sexes	12	7	10	18	11	23	35	30	19	5	-	-
Male	7	4	6	9	5	11	10	9	4	1	-	-
Female	5	3	4	9	6	12	25	21	15	4	-	-
White, both sexes	10	6	9	17	11	22	34	29	18	5	-	-
Male	6	4	5	9	5	10	10	9	4	1	-	-
Female	4	2	4	8	6	12	24	20	14	4	-	-
Black, both sexes	2	1	1	1	-	1	-	1	1	-	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	1	1	-	1	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Michigan												
All races, both sexes	20	22	23	21	29	44	33	40	15	9	-	-
Male	9	10	13	8	15	22	16	17	6	2	-	-
Female	11	12	10	13	14	22	17	23	9	7	-	-
White, both sexes	17	17	21	15	24	37	28	33	12	8	-	-
Male	7	8	11	6	13	17	14	14	3	2	-	-
Female	10	9	10	9	11	20	14	19	9	6	-	-
Black, both sexes	3	4	2	5	5	7	4	6	3	1	-	-
Male	2	1	2	2	2	5	2	2	3	-	-	-
Female	1	3	-	3	3	2	2	4	-	1	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Minnesota												
All races, both sexes	7	10	4	8	13	5	17	14	11	6	-	-
Male	5	3	2	4	6	3	5	8	4	-	-	-
Female	2	7	2	4	7	2	12	6	7	6	-	-
White, both sexes	7	10	4	7	13	5	17	14	11	6	-	-
Male	5	3	2	4	6	3	5	8	4	-	-	-
Female	2	7	2	3	7	2	12	6	7	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	5	11	8	16	13	18	20	11	14	3	-	-
Male	1	2	3	7	7	10	7	2	6	-	-	-
Female	4	9	5	9	6	8	13	9	8	3	-	-
White, both sexes	2	5	6	12	9	14	16	9	9	2	-	-
Male	-	1	2	5	4	8	5	2	4	-	-	-
Female	2	4	4	7	5	6	11	7	5	2	-	-
Black, both sexes	3	6	2	4	4	4	4	2	5	1	-	-
Male	1	1	1	2	3	2	2	-	2	-	-	-
Female	2	5	1	2	1	2	2	2	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Missouri												
All races, both sexes	23	16	13	24	29	31	29	35	12	4	1	-
Male	14	9	3	13	12	10	14	13	4	1	-	-
Female	9	7	10	11	17	21	15	22	8	3	1	-
White, both sexes	21	13	11	23	23	26	24	33	12	4	1	-
Male	14	7	2	12	9	7	11	12	4	1	-	-
Female	7	6	9	11	14	19	13	21	8	3	1	-
Black, both sexes	2	3	2	1	6	5	4	2	-	-	-	-
Male	-	2	1	1	3	3	2	1	-	-	-	-
Female	2	1	1	-	3	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	4	-	4	3	1	5	1	1	-	-	-
Male	2	2	-	-	1	1	2	-	1	-	-	-
Female	-	2	-	4	2	-	3	1	-	-	-	-
White, both sexes	2	3	-	4	3	1	4	1	1	-	-	-
Male	2	2	-	-	1	1	2	-	1	-	-	-
Female	-	1	-	4	2	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Nebraska												
All races, both sexes	3	3	2	5	9	6	7	5	6	-	-	-
Male	-	-	1	2	3	2	3	-	2	-	-	-
Female	3	3	1	3	6	4	4	5	4	-	-	-
White, both sexes	3	1	2	5	9	5	7	5	6	-	-	-
Male	-	-	1	2	3	1	3	-	2	-	-	-
Female	3	1	1	3	6	4	4	5	4	-	-	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	6	1	13	9	7	6	12	9	5	2	-	-
Male	2	1	8	5	6	2	5	3	-	-	-	-
Female	4	-	5	4	1	4	7	6	5	2	-	-
White, both sexes	6	1	11	6	4	5	11	9	5	2	-	-
Male	2	1	8	2	4	2	4	3	-	-	-	-
Female	4	-	3	4	-	3	7	6	5	2	-	-
Black, both sexes	-	-	2	2	3	1	1	-	-	-	-	-
Male	-	-	-	2	2	-	1	-	-	-	-	-
Female	-	-	2	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
New Hampshire												
All races, both sexes	1	2	2	3	3	2	8	1	4	1	-	-
Male	1	-	2	2	1	-	4	-	1	1	-	-
Female	-	2	-	1	2	2	4	1	3	-	-	-
White, both sexes	1	2	2	3	3	2	8	1	4	1	-	-
Male	1	-	2	2	1	-	4	-	1	1	-	-
Female	-	2	-	1	2	2	4	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	19	12	18	22	21	19	38	32	18	3	1	-
Male	7	4	9	12	9	7	6	7	6	-	1	-
Female	12	8	9	10	12	12	32	25	12	3	-	-
White, both sexes	12	9	13	17	17	18	29	29	17	2	1	-
Male	5	3	6	9	6	7	6	7	6	-	1	-
Female	7	6	7	8	11	11	23	22	11	2	-	-
Black, both sexes	7	3	5	5	4	1	8	3	1	1	-	-
Male	2	1	3	3	3	-	-	-	-	-	-	-
Female	5	2	2	2	1	1	8	3	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
New Mexico												
All races, both sexes	2	2	2	5	2	3	8	3	4	-	-	-
Male	-	-	1	4	1	-	3	-	2	-	-	-
Female	2	2	1	1	1	3	5	3	2	-	-	-
White, both sexes	2	1	2	5	2	3	8	3	3	-	-	-
Male	-	-	1	4	1	-	3	-	1	-	-	-
Female	2	1	1	1	1	3	5	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	18	25	28	23	31	39	47	49	18	7	1	-
Male	9	11	14	14	13	19	21	17	5	3	-	-
Female	9	14	14	9	18	20	26	32	13	4	1	-
White, both sexes	12	19	22	17	26	31	40	43	16	7	1	-
Male	6	8	11	9	11	15	19	15	4	3	-	-
Female	6	11	11	8	15	16	21	28	12	4	1	-
Black, both sexes	6	6	6	6	5	8	7	5	2	-	-	-
Male	3	3	3	5	2	4	2	2	1	-	-	-
Female	3	3	3	1	3	4	5	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
North Carolina												
All races, both sexes	15	14	32	29	35	40	41	38	23	12	-	-
Male	6	9	18	17	15	15	12	13	1	1	-	-
Female	9	5	14	12	20	25	29	25	22	11	-	-
White, both sexes	7	8	24	17	28	29	34	34	18	8	-	-
Male	2	7	13	10	13	13	10	12	1	-	-	-
Female	5	1	11	7	15	16	24	22	17	8	-	-
Black, both sexes	8	5	8	11	7	11	7	4	4	4	-	-
Male	4	2	5	7	2	2	2	1	-	1	-	-
Female	4	3	3	4	5	9	5	3	4	3	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	1	3	1	6	3	5	2	1	1	-
Male	-	-	1	2	1	2	2	-	-	-	-	-
Female	-	1	-	1	-	4	1	5	2	1	1	-
White, both sexes	-	1	1	3	1	6	3	5	2	1	1	-
Male	-	-	1	2	1	2	2	-	-	-	-	-
Female	-	1	-	1	-	4	1	5	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Ohio												
All races, both sexes	17	28	30	25	28	46	54	38	13	2	-	-
Male	6	15	14	8	15	20	24	11	3	-	-	-
Female	11	13	16	17	13	26	30	27	10	2	-	-
White, both sexes	15	26	26	22	26	41	48	35	12	1	-	-
Male	6	14	13	7	14	20	22	10	3	-	-	-
Female	9	12	13	15	12	21	26	25	9	1	-	-
Black, both sexes	2	1	4	3	2	5	6	2	1	1	-	-
Male	-	1	1	1	1	-	2	-	-	-	-	-
Female	2	-	3	2	1	5	4	2	1	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	5	12	12	9	14	13	17	16	5	1	-	-
Male	1	6	5	6	6	6	6	6	2	-	-	-
Female	4	6	7	3	8	7	11	10	3	1	-	-
White, both sexes	4	10	9	9	10	12	15	16	5	1	-	-
Male	1	6	2	6	3	5	6	6	2	-	-	-
Female	3	4	7	3	7	7	9	10	3	1	-	-
Black, both sexes	1	1	1	-	2	-	2	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	1	1	-	-	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	1	2	-	2	-	-	-	-	-	-	-
Male	-	-	2	-	2	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Oregon												
All races, both sexes	2	7	7	4	13	12	11	8	4	3	-	-
Male	1	6	6	1	4	6	4	2	-	-	-	-
Female	1	1	1	3	9	6	7	6	4	3	-	-
White, both sexes	1	7	7	4	13	12	11	6	4	3	-	-
Male	1	6	6	1	4	6	4	2	-	-	-	-
Female	-	1	1	3	9	6	7	4	4	3	-	-
Black, both sexes	1	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	29	34	28	49	53	58	68	46	37	12	1	-
Male	15	18	12	23	26	29	29	12	10	4	1	-
Female	14	16	16	26	27	29	39	34	27	8	-	-
White, both sexes	21	30	22	42	39	53	59	44	36	11	1	-
Male	11	14	10	22	22	27	27	12	9	4	1	-
Female	10	16	12	20	17	26	32	32	27	7	-	-
Black, both sexes	7	4	6	6	14	5	9	2	1	1	-	-
Male	3	4	2	1	4	2	2	-	1	-	-	-
Female	4	-	4	5	10	3	7	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Rhode Island												
All races, both sexes	1	3	1	-	-	3	2	5	1	-	-	-
Male	1	1	-	-	-	2	-	3	1	-	-	-
Female	-	2	1	-	-	1	2	2	-	-	-	-
White, both sexes	-	3	1	-	-	3	2	5	1	-	-	-
Male	-	1	-	-	-	2	-	3	1	-	-	-
Female	-	2	1	-	-	1	2	2	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	12	11	17	17	8	24	15	11	7	-	1	-
Male	4	5	6	6	5	11	8	1	1	-	-	-
Female	8	6	11	11	3	13	7	10	6	-	1	-
White, both sexes	7	5	13	12	7	16	9	8	7	-	1	-
Male	3	2	5	5	4	8	5	-	1	-	-	-
Female	4	3	8	7	3	8	4	8	6	-	1	-
Black, both sexes	5	6	4	5	1	8	6	3	-	-	-	-
Male	1	3	1	1	1	3	3	1	-	-	-	-
Female	4	3	3	4	-	5	3	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
South Dakota												
All races, both sexes	4	2	2	2	-	5	5	2	1	2	-	-
Male	3	1	1	2	-	2	4	1	-	-	-	-
Female	1	1	1	-	-	3	1	1	1	2	-	-
White, both sexes	4	2	2	2	-	5	5	2	-	2	-	-
Male	3	1	1	2	-	2	4	1	-	-	-	-
Female	1	1	1	-	-	3	1	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	13	18	16	21	34	36	30	26	17	5	-	-
Male	9	7	6	9	17	14	10	5	2	1	-	-
Female	4	11	10	12	17	22	20	21	15	4	-	-
White, both sexes	10	15	13	20	31	30	26	22	15	4	-	-
Male	9	5	6	8	15	11	10	3	2	1	-	-
Female	1	10	7	12	16	19	16	19	13	3	-	-
Black, both sexes	3	3	3	1	3	6	4	4	2	1	-	-
Male	-	2	-	1	2	3	-	2	-	-	-	-
Female	3	1	3	-	1	3	4	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Texas												
All races, both sexes	26	44	39	61	69	78	78	61	35	13	2	-
Male	16	22	20	35	34	41	27	19	9	3	-	-
Female	10	22	19	26	35	37	51	42	26	10	2	-
White, both sexes	20	34	23	44	58	62	69	52	31	10	1	-
Male	11	19	8	25	28	30	24	15	7	2	-	-
Female	9	15	15	19	30	32	45	37	24	8	1	-
Black, both sexes	6	9	14	15	10	16	9	8	4	3	1	-
Male	5	2	10	9	6	11	3	3	2	1	-	-
Female	1	7	4	6	4	5	6	5	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	2	-	-	-	1	-	-	-	-
Male	-	1	2	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	5	1	4	4	6	6	10	9	5	-	1	-
Male	2	-	3	3	3	1	8	5	2	-	1	-
Female	3	1	1	1	3	5	2	4	3	-	-	-
White, both sexes	5	1	4	4	6	6	10	9	5	-	1	-
Male	2	-	3	3	3	1	8	5	2	-	1	-
Female	3	1	1	1	3	5	2	4	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Vermont												
All races, both sexes	-	-	1	3	2	1	4	3	-	1	-	-
Male	-	-	-	2	1	1	3	1	-	-	-	-
Female	-	-	1	1	1	-	1	2	-	1	-	-
White, both sexes	-	-	1	3	2	1	4	3	-	1	-	-
Male	-	-	-	2	1	1	3	1	-	-	-	-
Female	-	-	1	1	1	-	1	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	19	15	22	20	27	34	38	29	4	2	1	-
Male	11	9	13	10	13	16	15	14	2	2	-	-
Female	8	6	9	10	14	18	23	15	2	-	1	-
White, both sexes	10	9	17	17	24	25	33	20	3	2	1	-
Male	5	5	10	8	11	10	12	8	1	2	-	-
Female	5	4	7	9	13	15	21	12	2	-	1	-
Black, both sexes	9	6	5	3	3	9	5	7	1	-	-	-
Male	6	4	3	2	2	6	3	5	1	-	-	-
Female	3	2	2	1	1	3	2	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Washington												
All races, both sexes	12	10	13	8	14	12	29	20	11	4	-	-
Male	8	8	6	5	7	6	12	7	4	-	-	-
Female	4	2	7	3	7	6	17	13	7	4	-	-
White, both sexes	10	7	12	7	12	12	26	20	10	3	-	-
Male	7	6	5	5	5	6	11	7	3	-	-	-
Female	3	1	7	2	7	6	15	13	7	3	-	-
Black, both sexes	1	2	1	1	2	-	1	-	1	1	-	-
Male	-	2	1	-	2	-	1	-	1	-	-	-
Female	1	-	-	1	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
West Virginia												
All races, both sexes	3	4	4	4	12	11	8	2	3	-	-	-
Male	2	2	4	1	3	5	2	-	1	-	-	-
Female	1	2	-	3	9	6	6	2	2	-	-	-
White, both sexes	3	4	4	3	12	11	8	2	3	-	-	-
Male	2	2	4	1	3	5	2	-	1	-	-	-
Female	1	2	-	2	9	6	6	2	2	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Pulmonary embolism (I26)												
Wisconsin												
All races, both sexes	8	15	12	11	20	32	23	24	12	5	-	-
Male	7	13	4	5	11	16	7	7	3	1	-	-
Female	1	2	8	6	9	16	16	17	9	4	-	-
White, both sexes	7	14	11	10	19	31	22	24	12	5	-	-
Male	7	12	3	4	11	15	7	7	3	1	-	-
Female	-	2	8	6	8	16	15	17	9	4	-	-
Black, both sexes	1	-	-	1	1	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	1	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	2	3	1	4	4	3	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	2	2	1	4	4	3	-	1	-	-	-
White, both sexes	1	2	3	1	3	4	3	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	2	2	1	3	4	3	-	1	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
United States												
All races, both sexes	5,221	94	20	6	4	19	25	41	63	84	140	230
Male	1,814	52	11	2	2	7	10	9	21	26	43	80
Female	3,407	42	9	4	2	12	15	32	42	58	97	150
White, both sexes	4,307	63	16	3	3	16	19	27	37	55	99	159
Male	1,526	36	9	1	1	5	7	5	10	18	31	61
Female	2,781	27	7	2	2	11	12	22	27	37	68	98
Black, both sexes	807	28	4	2	1	2	5	11	24	26	36	63
Male	256	16	2	1	1	1	3	4	11	7	10	17
Female	551	12	2	1	-	1	2	7	13	19	26	46
American Indian 1/, both sexes	31	1	-	-	-	-	-	2	-	-	2	2
Male	13	-	-	-	-	-	-	-	-	-	2	-
Female	18	1	-	-	-	-	-	2	-	-	-	2
Asian or Pacific Islander 2/, both sexes	76	2	-	1	-	1	1	1	2	3	3	6
Male	19	-	-	-	-	1	-	-	-	1	-	2
Female	57	2	-	1	-	-	1	1	2	2	3	4
Alabama												
All races, both sexes	102	1	1	1	-	-	-	1	1	-	3	5
Male	40	-	-	1	-	-	-	-	-	-	-	3
Female	62	1	1	-	-	-	-	1	1	-	3	2
White, both sexes	67	1	1	1	-	-	-	1	1	-	1	3
Male	26	-	-	1	-	-	-	-	-	-	-	1
Female	41	1	1	-	-	-	-	1	1	-	1	2
Black, both sexes	35	-	-	-	-	-	-	-	-	-	2	2
Male	14	-	-	-	-	-	-	-	-	-	-	2
Female	21	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Alaska												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Arizona												
All races, both sexes	85	1	-	-	-	-	-	1	-	-	1	6
Male	36	1	-	-	-	-	-	-	-	-	-	3
Female	49	-	-	-	-	-	-	1	-	-	1	3
White, both sexes	80	1	-	-	-	-	-	-	-	-	1	6
Male	34	1	-	-	-	-	-	-	-	-	-	3
Female	46	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Arkansas												
All races, both sexes	67	1	-	-	-	-	1	-	-	3	2	4
Male	26	-	-	-	-	-	1	-	-	1	1	1
Female	41	1	-	-	-	-	-	-	-	2	1	3
White, both sexes	56	1	-	-	-	-	-	-	-	2	2	3
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	35	1	-	-	-	-	-	-	-	2	1	2
Black, both sexes	11	-	-	-	-	-	1	-	-	1	-	1
Male	5	-	-	-	-	-	1	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	421	7	2	2	-	1	1	8	12	13	21	21
Male	129	4	-	-	-	-	-	2	1	2	7	4
Female	292	3	2	2	-	1	1	6	11	11	14	17
White, both sexes	333	5	2	1	-	1	-	6	8	7	16	15
Male	105	3	-	-	-	-	-	1	1	2	6	2
Female	228	2	2	1	-	1	-	5	7	5	10	13
Black, both sexes	59	2	-	-	-	-	-	2	3	5	4	5
Male	17	1	-	-	-	-	-	1	-	-	1	1
Female	42	1	-	-	-	-	-	1	3	5	3	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	-	-	1	-	-	1	-	1	1	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	1	-	-	1	-	1	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Colorado												
All races, both sexes	108	2	-	-	-	-	-	-	2	3	1	3
Male	44	-	-	-	-	-	-	-	2	2	1	1
Female	64	2	-	-	-	-	-	-	-	1	-	2
White, both sexes	102	2	-	-	-	-	-	-	-	3	1	2
Male	41	-	-	-	-	-	-	-	-	2	1	1
Female	61	2	-	-	-	-	-	-	-	1	-	1
Black, both sexes	4	-	-	-	-	-	-	-	2	-	-	1
Male	2	-	-	-	-	-	-	-	2	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	56	-	-	1	1	-	-	-	-	1	4	1
Male	31	-	-	1	1	-	-	-	-	1	2	1
Female	25	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	49	-	-	-	-	-	-	-	-	1	4	1
Male	27	-	-	-	-	-	-	-	-	1	2	1
Female	22	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	7	-	-	1	1	-	-	-	-	-	-	-
Male	4	-	-	1	1	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Delaware												
All races, both sexes	24	2	-	-	-	-	-	-	-	-	-	1
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	17	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	2	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	11	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Florida												
All races, both sexes	303	6	1	-	-	2	-	1	3	5	11	11
Male	121	4	-	-	-	1	-	1	-	2	3	3
Female	182	2	1	-	-	1	-	-	3	3	8	8
White, both sexes	257	3	-	-	-	1	-	-	2	5	8	4
Male	109	2	-	-	-	-	-	-	-	2	2	3
Female	148	1	-	-	-	1	-	-	2	3	6	1
Black, both sexes	43	3	1	-	-	-	-	1	1	-	3	6
Male	10	2	-	-	-	-	-	1	-	-	1	-
Female	33	1	1	-	-	-	-	-	1	-	2	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Georgia												
All races, both sexes	175	5	-	-	-	1	1	1	3	6	4	7
Male	68	3	-	-	-	-	1	-	1	3	1	2
Female	107	2	-	-	-	1	-	1	2	3	3	5
White, both sexes	119	2	-	-	-	1	-	-	1	3	3	2
Male	50	-	-	-	-	-	-	-	1	2	1	1
Female	69	2	-	-	-	1	-	-	-	1	2	1
Black, both sexes	50	3	-	-	-	-	1	-	2	2	1	5
Male	17	3	-	-	-	-	1	-	-	1	-	1
Female	33	-	-	-	-	-	-	-	2	1	1	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	1	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Hawaii												
All races, both sexes	13	1	-	-	-	-	-	-	-	1	1	2
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	9	1	-	-	-	-	-	-	-	-	1	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	1	-	-	-	-	-	-	-	1	1	2
Male	2	-	-	-	-	-	-	-	-	1	-	1
Female	8	1	-	-	-	-	-	-	-	-	1	1
Idaho												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Illinois												
All races, both sexes	227	6	1	-	-	-	1	1	3	3	4	13
Male	72	5	1	-	-	-	-	1	-	1	-	3
Female	155	1	-	-	-	-	1	-	3	2	4	10
White, both sexes	178	4	-	-	-	-	1	1	2	1	2	10
Male	59	3	-	-	-	-	-	1	-	-	-	3
Female	119	1	-	-	-	-	1	-	2	1	2	7
Black, both sexes	44	2	1	-	-	-	-	-	1	2	2	3
Male	12	2	1	-	-	-	-	-	-	1	-	-
Female	32	-	-	-	-	-	-	-	1	1	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	139	5	1	-	-	2	-	-	1	-	2	7
Male	46	2	-	-	-	1	-	-	-	-	1	2
Female	93	3	1	-	-	1	-	-	1	-	1	5
White, both sexes	130	4	1	-	-	2	-	-	1	-	2	7
Male	43	1	-	-	-	1	-	-	-	-	1	2
Female	87	3	1	-	-	1	-	-	1	-	1	5
Black, both sexes	9	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Iowa												
All races, both sexes	68	-	-	-	-	-	-	-	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	66	-	-	-	-	-	-	-	-	-	1	1
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	48	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	58	1	-	-	-	-	-	-	-	1	4	2
Male	15	-	-	-	-	-	-	-	-	1	2	1
Female	43	1	-	-	-	-	-	-	-	-	2	1
White, both sexes	55	1	-	-	-	-	-	-	-	1	3	2
Male	14	-	-	-	-	-	-	-	-	1	2	1
Female	41	1	-	-	-	-	-	-	-	-	1	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Kentucky												
All races, both sexes	87	1	-	1	-	-	1	2	-	2	3	5
Male	25	-	-	-	-	-	1	-	-	-	3	-
Female	62	1	-	1	-	-	-	2	-	2	-	5
White, both sexes	79	1	-	1	-	-	1	1	-	2	1	3
Male	22	-	-	-	-	-	1	-	-	-	1	-
Female	57	1	-	1	-	-	-	1	-	2	-	3
Black, both sexes	8	-	-	-	-	-	-	1	-	-	2	2
Male	3	-	-	-	-	-	-	-	-	-	2	-
Female	5	-	-	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	83	1	1	-	-	-	3	1	2	1	2	4
Male	25	1	1	-	-	-	2	-	1	-	-	1
Female	58	-	-	-	-	-	1	1	1	1	2	3
White, both sexes	55	1	1	-	-	-	2	-	1	1	2	1
Male	17	1	1	-	-	-	1	-	1	-	-	-
Female	38	-	-	-	-	-	1	-	-	1	2	1
Black, both sexes	27	-	-	-	-	-	1	1	1	-	-	3
Male	8	-	-	-	-	-	1	-	-	-	-	1
Female	19	-	-	-	-	-	-	1	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Maine												
All races, both sexes	24	-	-	-	-	-	-	1	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	24	-	-	-	-	-	-	1	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	81	4	-	-	-	-	-	-	1	2	1	4
Male	38	3	-	-	-	-	-	-	1	1	-	2
Female	43	1	-	-	-	-	-	-	-	1	1	2
White, both sexes	49	2	-	-	-	-	-	-	-	1	-	1
Male	23	2	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	31	2	-	-	-	-	-	-	1	1	1	2
Male	15	1	-	-	-	-	-	-	1	1	-	1
Female	16	1	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Massachusetts												
All races, both sexes	114	2	-	-	-	-	-	1	1	1	2	2
Male	42	1	-	-	-	-	-	-	1	-	2	1
Female	72	1	-	-	-	-	-	1	-	1	-	1
White, both sexes	109	2	-	-	-	-	-	1	1	1	1	2
Male	40	1	-	-	-	-	-	-	1	-	1	1
Female	69	1	-	-	-	-	-	1	-	1	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	187	2	2	-	-	1	-	2	2	1	5	10
Male	58	-	2	-	-	-	-	-	1	-	1	1
Female	129	2	-	-	-	1	-	2	1	1	4	9
White, both sexes	146	1	1	-	-	1	-	1	1	-	2	8
Male	48	-	1	-	-	-	-	-	-	-	-	1
Female	98	1	-	-	-	1	-	1	1	-	2	7
Black, both sexes	38	1	1	-	-	-	-	1	1	1	2	2
Male	9	-	1	-	-	-	-	-	1	-	1	-
Female	29	1	-	-	-	-	-	1	-	1	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Minnesota												
All races, both sexes	96	-	-	-	-	-	1	-	-	4	-	4
Male	30	-	-	-	-	-	-	-	-	2	-	2
Female	66	-	-	-	-	-	1	-	-	2	-	2
White, both sexes	93	-	-	-	-	-	1	-	-	3	-	4
Male	29	-	-	-	-	-	-	-	-	2	-	2
Female	64	-	-	-	-	-	1	-	-	1	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	58	1	1	-	-	-	1	-	-	1	3	2
Male	17	-	1	-	-	-	1	-	-	-	-	1
Female	41	1	-	-	-	-	-	-	-	1	3	1
White, both sexes	39	-	1	-	-	-	1	-	-	1	2	1
Male	9	-	1	-	-	-	1	-	-	-	-	1
Female	30	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	19	1	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	11	1	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Missouri												
All races, both sexes	107	4	4	-	-	1	-	-	4	2	2	2
Male	39	2	3	-	-	-	-	-	-	1	-	1
Female	68	2	1	-	-	1	-	-	4	1	2	1
White, both sexes	91	3	4	-	-	1	-	-	3	2	1	2
Male	33	2	3	-	-	-	-	-	-	1	-	1
Female	58	1	1	-	-	1	-	-	3	1	1	1
Black, both sexes	16	1	-	-	-	-	-	-	1	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	1	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	25	1	-	-	-	-	-	-	-	-	-	-
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	1	-	-	-	-	-	-	-	-	-	-
Male	6	1	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Nebraska												
All races, both sexes	44	-	-	-	-	-	1	-	-	-	-	2
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	1	-	-	-	-	2
White, both sexes	42	-	-	-	-	-	1	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	37	-	-	-	-	-	-	2	-	-	-	2
Male	19	-	-	-	-	-	-	2	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	33	-	-	-	-	-	-	2	-	-	-	2
Male	19	-	-	-	-	-	-	2	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
New Hampshire												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	140	4	1	-	-	1	-	-	1	1	3	6
Male	56	3	1	-	-	-	-	-	1	-	1	2
Female	84	1	-	-	-	1	-	-	-	1	2	4
White, both sexes	118	3	1	-	-	-	-	-	1	1	1	6
Male	51	3	1	-	-	-	-	-	1	-	-	2
Female	67	-	-	-	-	-	-	-	-	1	1	4
Black, both sexes	20	-	-	-	-	1	-	-	-	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	15	-	-	-	-	1	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
New Mexico												
All races, both sexes	35	-	-	-	1	-	-	1	1	1	3	-
Male	15	-	-	-	1	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	1	1	1	2	-
White, both sexes	32	-	-	-	1	-	-	1	1	1	2	-
Male	14	-	-	-	1	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	1	1	1	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	254	6	1	-	-	3	1	4	7	4	5	9
Male	88	4	-	-	-	3	1	1	2	1	1	3
Female	166	2	1	-	-	-	-	3	5	3	4	6
White, both sexes	213	5	-	-	-	2	1	2	4	3	3	7
Male	73	3	-	-	-	2	1	-	1	1	1	2
Female	140	2	-	-	-	-	-	2	3	2	2	5
Black, both sexes	38	1	1	-	-	1	-	2	3	1	2	2
Male	14	1	-	-	-	1	-	1	1	-	-	1
Female	24	-	1	-	-	-	-	1	2	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
North Carolina												
All races, both sexes	204	4	-	-	-	1	1	2	-	4	5	14
Male	62	2	-	-	-	-	-	-	-	1	-	6
Female	142	2	-	-	-	1	1	2	-	3	5	8
White, both sexes	127	-	-	-	-	1	-	1	-	1	3	5
Male	35	-	-	-	-	-	-	-	-	-	-	3
Female	92	-	-	-	-	1	-	1	-	1	3	2
Black, both sexes	72	4	-	-	-	-	1	-	-	3	2	9
Male	24	2	-	-	-	-	-	-	-	1	-	3
Female	48	2	-	-	-	-	1	-	-	2	2	6
American Indian 1/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Ohio												
All races, both sexes	232	1	-	-	-	-	2	3	3	5	4	12
Male	84	1	-	-	-	-	-	1	1	2	3	5
Female	148	-	-	-	-	-	2	2	2	3	1	7
White, both sexes	192	1	-	-	-	-	1	2	1	4	3	9
Male	71	1	-	-	-	-	-	-	-	1	2	3
Female	121	-	-	-	-	-	1	2	1	3	1	6
Black, both sexes	38	-	-	-	-	-	1	1	2	1	1	3
Male	13	-	-	-	-	-	-	1	1	1	1	2
Female	25	-	-	-	-	-	1	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	57	1	2	-	-	-	1	-	1	1	1	5
Male	21	-	1	-	-	-	-	-	-	-	1	2
Female	36	1	1	-	-	-	1	-	1	1	-	3
White, both sexes	45	-	2	-	-	-	1	-	1	-	1	4
Male	17	-	1	-	-	-	-	-	-	-	1	2
Female	28	-	1	-	-	-	1	-	1	-	-	2
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	4	1	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Oregon												
All races, both sexes	76	-	-	-	-	1	-	-	-	1	2	4
Male	34	-	-	-	-	-	-	-	-	-	1	2
Female	42	-	-	-	-	1	-	-	-	1	1	2
White, both sexes	72	-	-	-	-	1	-	-	-	1	2	4
Male	33	-	-	-	-	-	-	-	-	-	1	2
Female	39	-	-	-	-	1	-	-	-	1	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	283	4	-	-	1	1	1	1	-	4	5	10
Male	85	2	-	-	-	-	1	-	-	-	-	5
Female	198	2	-	-	1	1	-	1	-	4	5	5
White, both sexes	245	3	-	-	1	1	1	1	-	3	3	8
Male	72	1	-	-	-	-	1	-	-	-	-	4
Female	173	2	-	-	1	1	-	1	-	3	3	4
Black, both sexes	38	1	-	-	-	-	-	-	-	1	2	2
Male	13	1	-	-	-	-	-	-	-	-	-	1
Female	25	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Rhode Island												
All races, both sexes	35	2	-	-	-	-	-	1	-	2	2	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	29	2	-	-	-	-	-	1	-	2	2	-
White, both sexes	31	1	-	-	-	-	-	-	-	1	2	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	25	1	-	-	-	-	-	-	-	1	2	-
Black, both sexes	4	1	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	96	2	-	-	-	-	-	-	1	-	3	8
Male	25	-	-	-	-	-	-	-	1	-	-	2
Female	71	2	-	-	-	-	-	-	-	-	3	6
White, both sexes	62	1	-	-	-	-	-	-	-	-	2	3
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	46	1	-	-	-	-	-	-	-	-	2	3
Black, both sexes	34	1	-	-	-	-	-	-	1	-	1	5
Male	9	-	-	-	-	-	-	-	1	-	-	2
Female	25	1	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
South Dakota												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	97	3	-	-	-	1	-	-	-	3	3	5
Male	39	2	-	-	-	-	-	-	-	1	1	4
Female	58	1	-	-	-	1	-	-	-	2	2	1
White, both sexes	84	2	-	-	-	1	-	-	-	-	3	5
Male	36	2	-	-	-	-	-	-	-	-	1	4
Female	48	-	-	-	-	1	-	-	-	-	2	1
Black, both sexes	13	1	-	-	-	-	-	-	-	3	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	10	1	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Texas												
All races, both sexes	320	7	-	-	1	-	4	4	7	2	10	18
Male	117	4	-	-	-	-	-	1	4	-	4	5
Female	203	3	-	-	1	-	4	3	3	2	6	13
White, both sexes	261	6	-	-	1	-	4	4	3	2	9	13
Male	98	3	-	-	-	-	-	1	2	-	3	5
Female	163	3	-	-	1	-	4	3	1	2	6	8
Black, both sexes	55	1	-	-	-	-	-	-	3	-	1	5
Male	17	1	-	-	-	-	-	-	2	-	1	-
Female	38	-	-	-	-	-	-	-	1	-	-	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Utah												
All races, both sexes	61	-	-	-	-	1	1	1	2	1	3	2
Male	22	-	-	-	-	1	1	-	1	1	1	1
Female	39	-	-	-	-	-	-	1	1	-	2	1
White, both sexes	60	-	-	-	-	1	1	1	2	1	3	2
Male	22	-	-	-	-	1	1	-	1	1	1	1
Female	38	-	-	-	-	-	-	1	1	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Vermont												
All races, both sexes	20	-	-	-	-	-	1	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	20	-	-	-	-	-	1	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	139	3	-	1	-	-	-	1	1	3	4	3
Male	36	3	-	-	-	-	-	-	1	1	1	2
Female	103	-	-	1	-	-	-	1	-	2	3	1
White, both sexes	97	3	-	-	-	-	-	-	-	1	-	2
Male	27	3	-	-	-	-	-	-	-	1	-	1
Female	70	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	41	-	-	1	-	-	-	1	1	2	4	1
Male	9	-	-	-	-	-	-	-	1	-	1	1
Female	32	-	-	1	-	-	-	1	-	2	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Washington												
All races, both sexes	112	-	1	-	-	1	-	1	2	2	5	5
Male	38	-	-	-	-	-	-	-	-	1	2	2
Female	74	-	1	-	-	1	-	1	2	1	3	3
White, both sexes	103	-	1	-	-	1	-	1	2	2	5	4
Male	36	-	-	-	-	-	-	-	-	1	2	2
Female	67	-	1	-	-	1	-	1	2	1	3	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	45	-	1	-	-	-	1	-	-	-	-	2
Male	18	-	1	-	-	-	-	-	-	-	-	2
Female	27	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	44	-	1	-	-	-	1	-	-	-	-	2
Male	17	-	1	-	-	-	-	-	-	-	-	2
Female	27	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Wisconsin												
All races, both sexes	115	2	-	-	-	1	1	-	1	-	3	3
Male	46	2	-	-	-	1	1	-	1	-	1	1
Female	69	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	107	2	-	-	-	1	1	-	1	-	2	1
Male	43	2	-	-	-	1	1	-	1	-	-	-
Female	64	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	17	1	-	-	-	-	-	-	-	-	-	-
Male	8	1	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	1	-	-	-	-	-	-	-	-	-	-
Male	8	1	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
United States												
All races, both sexes	243	330	377	448	599	620	775	640	345	103	15	-
Male	96	129	139	193	227	218	268	179	80	19	3	-
Female	147	201	238	255	372	402	507	461	265	84	12	-
White, both sexes	174	254	278	366	507	547	695	570	318	88	13	-
Male	70	109	107	163	200	199	246	156	73	16	3	-
Female	104	145	171	203	307	348	449	414	245	72	10	-
Black, both sexes	62	70	94	76	74	65	68	56	25	13	2	-
Male	22	19	30	29	20	18	19	18	6	2	-	-
Female	40	51	64	47	54	47	49	38	19	11	2	-
American Indian 1/, both sexes	3	2	2	2	6	-	5	3	1	-	-	-
Male	2	-	1	1	3	-	2	2	-	-	-	-
Female	1	2	1	1	3	-	3	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	4	3	4	12	8	7	11	1	2	-	-
Male	2	1	1	-	4	1	1	3	1	1	-	-
Female	2	3	2	4	8	7	6	8	-	1	-	-
Alabama												
All races, both sexes	3	10	11	8	14	11	11	12	6	3	-	-
Male	1	4	5	4	5	7	4	3	3	-	-	-
Female	2	6	6	4	9	4	7	9	3	3	-	-
White, both sexes	3	3	5	7	11	7	8	9	4	1	-	-
Male	1	1	2	3	4	5	3	3	2	-	-	-
Female	2	2	3	4	7	2	5	6	2	1	-	-
Black, both sexes	-	7	6	1	3	4	3	3	2	2	-	-
Male	-	3	3	1	1	2	1	-	1	-	-	-
Female	-	4	3	-	2	2	2	3	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Alaska												
All races, both sexes	-	-	1	2	-	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	2	-	-	2	-	-	-	-	-
White, both sexes	-	-	1	1	-	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	9	7	4	5	14	16	9	4	6	2	-	-
Male	3	4	2	3	2	10	2	1	4	1	-	-
Female	6	3	2	2	12	6	7	3	2	1	-	-
White, both sexes	7	7	4	4	14	16	9	3	6	2	-	-
Male	3	4	2	2	2	10	2	-	4	1	-	-
Female	4	3	2	2	12	6	7	3	2	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Arkansas												
All races, both sexes	1	7	4	10	7	2	11	6	7	1	-	-
Male	1	1	4	5	2	-	7	1	1	-	-	-
Female	-	6	-	5	5	2	4	5	6	1	-	-
White, both sexes	-	6	4	8	6	2	11	4	6	1	-	-
Male	-	1	4	4	2	-	7	1	-	-	-	-
Female	-	5	-	4	4	2	4	3	6	1	-	-
Black, both sexes	1	1	-	2	1	-	-	2	1	-	-	-
Male	1	-	-	1	-	-	-	-	1	-	-	-
Female	-	1	-	1	1	-	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	22	22	31	27	39	44	58	51	29	9	1	-
Male	11	10	10	10	14	15	20	8	7	3	1	-
Female	11	12	21	17	25	29	38	43	22	6	-	-
White, both sexes	18	18	25	22	34	33	47	42	26	6	1	-
Male	11	9	10	8	14	10	14	6	6	1	1	-
Female	7	9	15	14	20	23	33	36	20	5	-	-
Black, both sexes	2	2	5	4	3	8	6	4	2	2	-	-
Male	-	1	-	2	-	4	4	1	-	1	-	-
Female	2	1	5	2	3	4	2	3	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	1	2	3	4	5	1	1	-	-
Male	-	-	-	-	-	1	1	1	1	1	-	-
Female	2	2	1	1	2	2	3	4	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Colorado												
All races, both sexes	6	7	5	8	9	6	21	22	10	2	1	-
Male	1	3	3	7	6	3	8	5	2	-	-	-
Female	5	4	2	1	3	3	13	17	8	2	1	-
White, both sexes	5	6	5	8	8	6	21	22	10	2	1	-
Male	1	3	3	7	5	3	8	5	2	-	-	-
Female	4	3	2	1	3	3	13	17	8	2	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2	3	2	4	5	8	13	4	6	1	-	-
Male	1	-	1	2	3	7	7	2	2	-	-	-
Female	1	3	1	2	2	1	6	2	4	1	-	-
White, both sexes	1	3	2	2	4	7	13	4	6	1	-	-
Male	1	-	1	1	2	7	7	2	2	-	-	-
Female	-	3	1	1	2	-	6	2	4	1	-	-
Black, both sexes	1	-	-	2	1	1	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Delaware												
All races, both sexes	2	1	-	4	1	5	4	3	1	-	-	-
Male	2	1	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	3	1	3	4	3	1	-	-	-
White, both sexes	1	1	-	3	-	5	4	3	1	-	-	-
Male	1	1	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	2	-	3	4	3	1	-	-	-
Black, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	1	-	-	1	1	2	1	3	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	1	1	-	-	1	-	1	1	2	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	1	1	2	1	3	-	-	-
Male	-	-	-	-	-	1	1	-	1	-	-	-
Female	1	1	-	-	1	-	1	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Florida												
All races, both sexes	16	17	23	25	37	35	42	40	22	4	2	-
Male	8	4	9	12	12	15	14	20	8	3	2	-
Female	8	13	14	13	25	20	28	20	14	1	-	-
White, both sexes	8	11	17	25	30	35	40	40	22	4	2	-
Male	4	4	8	12	10	15	14	20	8	3	2	-
Female	4	7	9	13	20	20	26	20	14	1	-	-
Black, both sexes	7	6	6	-	7	-	2	-	-	-	-	-
Male	3	-	1	-	2	-	-	-	-	-	-	-
Female	4	6	5	-	5	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	9	11	10	18	22	24	27	17	4	4	1	-
Male	5	5	6	7	12	8	9	4	-	1	-	-
Female	4	6	4	11	10	16	18	13	4	3	1	-
White, both sexes	5	5	7	12	16	21	20	15	3	2	1	-
Male	3	3	4	4	11	8	8	4	-	-	-	-
Female	2	2	3	8	5	13	12	11	3	2	1	-
Black, both sexes	4	6	3	6	5	2	6	1	1	2	-	-
Male	2	2	2	3	-	-	1	-	-	1	-	-
Female	2	4	1	3	5	2	5	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Hawaii												
All races, both sexes	-	-	-	-	1	3	1	2	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	2	-	2	-	1	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	2	-	-	-	-
Idaho												
All races, both sexes	1	1	-	2	2	6	4	5	1	2	-	-
Male	-	-	-	1	1	2	1	1	-	-	-	-
Female	1	1	-	1	1	4	3	4	1	2	-	-
White, both sexes	1	1	-	2	2	6	4	5	1	2	-	-
Male	-	-	-	1	1	2	1	1	-	-	-	-
Female	1	1	-	1	1	4	3	4	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Illinois												
All races, both sexes	12	15	21	16	25	28	30	31	11	6	-	-
Male	1	5	9	10	6	8	8	11	3	-	-	-
Female	11	10	12	6	19	20	22	20	8	6	-	-
White, both sexes	9	12	14	13	20	20	29	24	11	5	-	-
Male	1	5	6	9	5	5	8	10	3	-	-	-
Female	8	7	8	4	15	15	21	14	8	5	-	-
Black, both sexes	3	3	5	2	4	8	1	7	-	-	-	-
Male	-	-	2	1	1	3	-	1	-	-	-	-
Female	3	3	3	1	3	5	1	6	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	-	-	-	1	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	1	-	-
Indiana												
All races, both sexes	5	9	13	15	16	20	22	12	9	-	-	-
Male	2	6	5	8	3	8	4	3	1	-	-	-
Female	3	3	8	7	13	12	18	9	8	-	-	-
White, both sexes	5	9	8	15	14	20	22	12	8	-	-	-
Male	2	6	3	8	3	8	4	3	1	-	-	-
Female	3	3	5	7	11	12	18	9	7	-	-	-
Black, both sexes	-	-	5	-	2	-	-	-	1	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	3	-	2	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Iowa												
All races, both sexes	1	5	3	6	7	8	12	15	8	-	1	-
Male	1	3	-	4	2	4	4	1	-	-	-	-
Female	-	2	3	2	5	4	8	14	8	-	1	-
White, both sexes	1	5	2	6	6	8	12	15	8	-	1	-
Male	1	3	-	4	1	4	4	1	-	-	-	-
Female	-	2	2	2	5	4	8	14	8	-	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	3	7	-	4	8	7	9	7	3	-	-
Male	1	-	1	-	2	2	1	1	2	1	-	-
Female	1	3	6	-	2	6	6	8	5	2	-	-
White, both sexes	2	3	6	-	3	8	7	9	7	3	-	-
Male	1	-	1	-	1	2	1	1	2	1	-	-
Female	1	3	5	-	2	6	6	8	5	2	-	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Kentucky												
All races, both sexes	3	2	7	8	18	9	16	4	5	-	-	-
Male	-	-	2	3	7	3	5	-	1	-	-	-
Female	3	2	5	5	11	6	11	4	4	-	-	-
White, both sexes	2	2	7	7	17	9	16	4	5	-	-	-
Male	-	-	2	2	7	3	5	-	1	-	-	-
Female	2	2	5	5	10	6	11	4	4	-	-	-
Black, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	5	13	7	7	5	8	9	11	3	-	-	-
Male	1	7	1	2	2	1	2	2	1	-	-	-
Female	4	6	6	5	3	7	7	9	2	-	-	-
White, both sexes	4	6	2	3	5	7	8	9	2	-	-	-
Male	1	5	-	1	2	1	2	1	-	-	-	-
Female	3	1	2	2	3	6	6	8	2	-	-	-
Black, both sexes	1	7	4	4	-	1	1	2	1	-	-	-
Male	-	2	1	1	-	-	-	1	1	-	-	-
Female	1	5	3	3	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Maine												
All races, both sexes	1	1	1	3	4	2	4	3	2	-	-	-
Male	1	1	-	2	1	-	-	2	1	-	-	-
Female	-	-	1	1	3	2	4	1	1	-	-	-
White, both sexes	1	1	1	3	4	2	4	3	2	-	-	-
Male	1	1	-	2	1	-	-	2	1	-	-	-
Female	-	-	1	1	3	2	4	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	7	4	4	11	10	6	15	8	3	1	-	-
Male	4	2	2	3	6	5	5	3	1	-	-	-
Female	3	2	2	8	4	1	10	5	2	1	-	-
White, both sexes	4	3	1	4	7	4	14	6	2	-	-	-
Male	2	2	-	-	6	3	5	1	1	-	-	-
Female	2	1	1	4	1	1	9	5	1	-	-	-
Black, both sexes	3	1	3	7	3	2	1	2	1	1	-	-
Male	2	-	2	3	-	2	-	2	-	-	-	-
Female	1	1	1	4	3	-	1	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Massachusetts												
All races, both sexes	5	6	10	6	14	18	21	15	6	4	-	-
Male	5	3	4	3	5	4	9	3	1	-	-	-
Female	-	3	6	3	9	14	12	12	5	4	-	-
White, both sexes	4	6	9	6	13	18	21	15	6	3	-	-
Male	4	3	4	3	5	4	9	3	1	-	-	-
Female	-	3	5	3	8	14	12	12	5	3	-	-
Black, both sexes	1	-	1	-	1	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	8	16	17	20	20	19	18	24	17	2	1	-
Male	3	6	7	9	3	4	7	10	4	-	-	-
Female	5	10	10	11	17	15	11	14	13	2	1	-
White, both sexes	6	12	13	16	17	11	17	22	15	1	1	-
Male	2	6	5	7	2	4	7	9	4	-	-	-
Female	4	6	8	9	15	7	10	13	11	1	1	-
Black, both sexes	2	4	4	4	1	8	1	2	2	1	-	-
Male	1	-	2	2	-	-	-	1	-	-	-	-
Female	1	4	2	2	1	8	1	1	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Minnesota												
All races, both sexes	2	9	4	7	7	7	25	17	8	1	-	-
Male	-	2	1	4	4	2	8	4	1	-	-	-
Female	2	7	3	3	3	5	17	13	7	1	-	-
White, both sexes	2	9	4	7	6	7	24	17	8	1	-	-
Male	-	2	1	4	3	2	8	4	1	-	-	-
Female	2	7	3	3	3	5	16	13	7	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	9	2	4	8	9	8	7	1	-	-	-
Male	1	2	1	1	4	1	2	1	1	-	-	-
Female	-	7	1	3	4	8	6	6	-	-	-	-
White, both sexes	-	4	1	2	4	9	7	5	1	-	-	-
Male	-	1	-	-	1	1	2	-	1	-	-	-
Female	-	3	1	2	3	8	5	5	-	-	-	-
Black, both sexes	1	5	1	2	4	-	1	2	-	-	-	-
Male	1	1	1	1	3	-	-	1	-	-	-	-
Female	-	4	-	1	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Missouri												
All races, both sexes	4	4	6	14	12	11	14	13	9	1	-	-
Male	2	2	2	6	6	3	6	2	3	-	-	-
Female	2	2	4	8	6	8	8	11	6	1	-	-
White, both sexes	3	4	5	12	10	9	10	12	9	1	-	-
Male	1	2	2	5	4	3	5	1	3	-	-	-
Female	2	2	3	7	6	6	5	11	6	1	-	-
Black, both sexes	1	-	1	2	2	2	4	1	-	-	-	-
Male	1	-	-	1	2	-	1	1	-	-	-	-
Female	-	-	1	1	-	2	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	4	1	2	5	-	5	3	3	-	1	-
Male	-	1	-	-	1	-	4	-	-	-	-	-
Female	-	3	1	2	4	-	1	3	3	-	1	-
White, both sexes	-	4	1	1	4	-	5	3	3	-	1	-
Male	-	1	-	-	-	-	4	-	-	-	-	-
Female	-	3	1	1	4	-	1	3	3	-	1	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Nebraska												
All races, both sexes	1	-	-	4	2	2	12	7	5	7	1	-
Male	-	-	-	1	1	-	6	1	1	3	-	-
Female	1	-	-	3	1	2	6	6	4	4	1	-
White, both sexes	1	-	-	3	2	2	12	7	5	7	1	-
Male	-	-	-	1	1	-	6	1	1	3	-	-
Female	1	-	-	2	1	2	6	6	4	4	1	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	1	2	4	11	5	3	2	1	-	-	-
Male	2	-	1	-	10	2	1	-	-	-	-	-
Female	2	1	1	4	1	3	2	2	1	-	-	-
White, both sexes	3	1	2	3	11	4	3	1	1	-	-	-
Male	2	-	1	-	10	2	1	-	-	-	-	-
Female	1	1	1	3	1	2	2	1	1	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
New Hampshire												
All races, both sexes	1	-	1	3	1	7	3	4	4	1	-	-
Male	-	-	-	1	1	3	1	-	-	-	-	-
Female	1	-	1	2	-	4	2	4	4	1	-	-
White, both sexes	1	-	1	3	1	7	3	4	4	1	-	-
Male	-	-	-	1	1	3	1	-	-	-	-	-
Female	1	-	1	2	-	4	2	4	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4	10	13	9	15	20	27	17	7	1	-	-
Male	2	3	2	5	6	6	14	8	1	1	-	-
Female	2	7	11	4	9	14	13	9	6	-	-	-
White, both sexes	2	8	9	7	13	18	26	15	6	1	-	-
Male	2	3	1	4	5	6	14	7	1	1	-	-
Female	-	5	8	3	8	12	12	8	5	-	-	-
Black, both sexes	2	2	4	2	2	1	1	2	1	-	-	-
Male	-	-	1	1	1	-	-	1	-	-	-	-
Female	2	2	3	1	1	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
New Mexico												
All races, both sexes	1	6	-	4	6	3	2	2	2	2	-	-
Male	1	4	-	-	4	2	-	2	-	-	-	-
Female	-	2	-	4	2	1	2	-	2	2	-	-
White, both sexes	1	6	-	4	5	3	2	2	1	2	-	-
Male	1	4	-	-	4	2	-	2	-	-	-	-
Female	-	2	-	4	1	1	2	-	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	12	14	15	20	22	34	48	26	16	7	-	-
Male	4	8	7	7	9	12	14	6	4	1	-	-
Female	8	6	8	13	13	22	34	20	12	6	-	-
White, both sexes	9	10	11	15	20	29	44	25	16	7	-	-
Male	3	5	5	6	9	11	13	5	4	1	-	-
Female	6	5	6	9	11	18	31	20	12	6	-	-
Black, both sexes	3	3	4	4	2	5	4	-	-	-	-	-
Male	1	3	2	1	-	1	1	-	-	-	-	-
Female	2	-	2	3	2	4	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
North Carolina												
All races, both sexes	9	11	21	24	22	24	28	21	8	4	1	-
Male	3	3	10	7	7	6	10	5	2	-	-	-
Female	6	8	11	17	15	18	18	16	6	4	1	-
White, both sexes	3	8	11	15	15	23	18	16	4	3	-	-
Male	-	3	4	4	5	6	7	3	-	-	-	-
Female	3	5	7	11	10	17	11	13	4	3	-	-
Black, both sexes	6	3	9	9	5	1	10	4	4	1	1	-
Male	3	-	5	3	1	-	3	1	2	-	-	-
Female	3	3	4	6	4	1	7	3	2	1	1	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	2	-	4	2	1	3	-	1	1	-	-
Male	-	-	-	1	2	-	-	-	-	-	-	-
Female	-	2	-	3	-	1	3	-	1	1	-	-
White, both sexes	-	2	-	4	2	1	3	-	1	1	-	-
Male	-	-	-	1	2	-	-	-	-	-	-	-
Female	-	2	-	3	-	1	3	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Ohio												
All races, both sexes	8	15	19	18	27	25	35	30	17	7	1	-
Male	3	8	5	12	9	12	11	9	1	1	-	-
Female	5	7	14	6	18	13	24	21	16	6	1	-
White, both sexes	6	12	14	16	22	22	29	28	14	7	1	-
Male	3	7	4	10	9	11	10	8	1	1	-	-
Female	3	5	10	6	13	11	19	20	13	6	1	-
Black, both sexes	2	3	5	2	4	3	5	2	3	-	-	-
Male	-	1	1	2	-	1	1	1	-	-	-	-
Female	2	2	4	-	4	2	4	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	4	3	8	5	6	6	6	5	2	-	-	-
Male	2	1	3	-	3	1	3	2	2	-	-	-
Female	2	2	5	5	3	5	3	3	-	-	-	-
White, both sexes	1	3	6	5	6	6	4	3	2	-	-	-
Male	1	1	2	-	3	1	2	1	2	-	-	-
Female	-	2	4	5	3	5	2	2	-	-	-	-
Black, both sexes	2	-	2	-	-	-	1	2	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	2	-	1	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Oregon												
All races, both sexes	2	4	6	6	7	6	15	16	3	3	-	-
Male	1	3	3	2	5	3	8	4	2	-	-	-
Female	1	1	3	4	2	3	7	12	1	3	-	-
White, both sexes	2	3	5	4	7	6	15	16	3	3	-	-
Male	1	2	3	2	5	3	8	4	2	-	-	-
Female	1	1	2	2	2	3	7	12	1	3	-	-
Black, both sexes	-	-	1	2	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	9	7	20	24	40	34	40	55	20	6	1	-
Male	5	2	5	12	12	5	17	13	6	-	-	-
Female	4	5	15	12	28	29	23	42	14	6	1	-
White, both sexes	6	4	16	18	34	32	36	51	20	6	1	-
Male	3	1	5	10	10	5	14	12	6	-	-	-
Female	3	3	11	8	24	27	22	39	14	6	1	-
Black, both sexes	3	3	4	6	6	2	4	4	-	-	-	-
Male	2	1	-	2	2	-	3	1	-	-	-	-
Female	1	2	4	4	4	2	1	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Rhode Island												
All races, both sexes	1	3	1	2	6	5	4	5	1	-	-	-
Male	1	1	-	-	-	3	-	1	-	-	-	-
Female	-	2	1	2	6	2	4	4	1	-	-	-
White, both sexes	1	3	1	2	6	5	3	5	1	-	-	-
Male	1	1	-	-	-	3	-	1	-	-	-	-
Female	-	2	1	2	6	2	3	4	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	5	12	9	3	12	9	13	9	7	2	1	-
Male	2	3	2	1	3	2	3	5	1	-	-	-
Female	3	9	7	2	9	7	10	4	6	2	1	-
White, both sexes	3	7	2	2	9	6	11	7	7	2	-	-
Male	2	2	-	1	2	1	3	4	1	-	-	-
Female	1	5	2	1	7	5	8	3	6	2	-	-
Black, both sexes	2	5	7	1	3	3	2	2	-	-	1	-
Male	-	1	2	-	1	1	-	1	-	-	-	-
Female	2	4	5	1	2	2	2	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
South Dakota												
All races, both sexes	1	-	1	-	-	4	4	2	-	1	-	-
Male	1	-	-	-	-	1	2	2	-	-	-	-
Female	-	-	1	-	-	3	2	-	-	1	-	-
White, both sexes	-	-	1	-	-	4	4	2	-	1	-	-
Male	-	-	-	-	-	1	2	2	-	-	-	-
Female	-	-	1	-	-	3	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	11	10	9	12	9	10	4	11	-	-	-
Male	2	8	2	5	5	4	3	1	1	-	-	-
Female	4	3	8	4	7	5	7	3	10	-	-	-
White, both sexes	5	11	8	8	11	8	8	4	10	-	-	-
Male	1	8	2	5	5	3	3	1	1	-	-	-
Female	4	3	6	3	6	5	5	3	9	-	-	-
Black, both sexes	1	-	2	1	1	1	2	-	1	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	2	1	1	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Texas												
All races, both sexes	19	20	26	43	41	36	39	29	9	4	1	-
Male	4	7	13	20	15	11	15	12	2	-	-	-
Female	15	13	13	23	26	25	24	17	7	4	1	-
White, both sexes	14	16	20	39	36	29	31	24	8	1	1	-
Male	1	5	12	19	13	10	13	9	2	-	-	-
Female	13	11	8	20	23	19	18	15	6	1	1	-
Black, both sexes	4	4	6	4	3	7	8	5	1	3	-	-
Male	2	2	1	1	1	1	2	3	-	-	-	-
Female	2	2	5	3	2	6	6	2	1	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	2	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Utah												
All races, both sexes	3	4	4	4	6	7	5	9	7	1	-	-
Male	1	1	1	1	2	5	2	2	-	1	-	-
Female	2	3	3	3	4	2	3	7	7	-	-	-
White, both sexes	3	4	4	4	5	7	5	9	7	1	-	-
Male	1	1	1	1	2	5	2	2	-	1	-	-
Female	2	3	3	3	3	2	3	7	7	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Vermont												
All races, both sexes	1	1	1	-	-	6	3	3	2	-	-	-
Male	-	1	-	-	-	2	2	2	-	-	-	-
Female	1	-	1	-	-	4	1	1	2	-	-	-
White, both sexes	1	1	1	-	-	6	3	3	2	-	-	-
Male	-	1	-	-	-	2	2	2	-	-	-	-
Female	1	-	1	-	-	4	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	9	6	10	12	18	22	17	20	8	1	-	-
Male	1	1	-	3	8	7	4	3	1	-	-	-
Female	8	5	10	9	10	15	13	17	7	1	-	-
White, both sexes	6	4	7	8	14	17	14	13	7	1	-	-
Male	1	-	-	2	6	6	3	3	1	-	-	-
Female	5	4	7	6	8	11	11	10	6	1	-	-
Black, both sexes	3	2	3	4	4	5	2	7	1	-	-	-
Male	-	1	-	1	2	1	1	-	-	-	-	-
Female	3	1	3	3	2	4	1	7	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Washington												
All races, both sexes	7	5	6	3	14	18	14	12	14	2	-	-
Male	3	3	5	2	7	4	2	4	2	1	-	-
Female	4	2	1	1	7	14	12	8	12	1	-	-
White, both sexes	6	3	6	3	12	18	12	11	14	2	-	-
Male	3	2	5	2	7	4	2	3	2	1	-	-
Female	3	1	1	1	5	14	10	8	12	1	-	-
Black, both sexes	1	2	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
West Virginia												
All races, both sexes	4	3	3	4	6	5	7	3	6	-	-	-
Male	-	-	2	2	2	2	1	2	4	-	-	-
Female	4	3	1	2	4	3	6	1	2	-	-	-
White, both sexes	4	3	3	4	6	5	7	2	6	-	-	-
Male	-	-	2	2	2	2	1	1	4	-	-	-
Female	4	3	1	2	4	3	6	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)												
Wisconsin												
All races, both sexes	4	5	7	11	12	15	20	18	6	5	1	-
Male	3	-	3	4	6	6	8	6	2	1	-	-
Female	1	5	4	7	6	9	12	12	4	4	1	-
White, both sexes	4	4	6	10	11	15	20	17	6	5	1	-
Male	3	-	2	4	6	6	8	6	2	1	-	-
Female	1	4	4	6	5	9	12	11	4	4	1	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	4	3	5	2	1	1	-	-
Male	-	-	-	-	1	3	1	1	-	1	-	-
Female	-	-	-	-	3	-	4	1	1	-	-	-
White, both sexes	-	-	-	-	4	3	5	2	1	1	-	-
Male	-	-	-	-	1	3	1	1	-	1	-	-
Female	-	-	-	-	3	-	4	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
United States												
All races, both sexes	157,081	244	123	88	127	310	483	639	888	1,351	2,221	3,411
Male	69,737	133	66	42	70	228	336	455	609	885	1,491	2,255
Female	87,344	111	57	46	57	82	147	184	279	466	730	1,156
White, both sexes	136,389	152	71	50	81	202	320	422	545	873	1,546	2,362
Male	59,758	79	44	26	48	147	228	315	380	594	1,065	1,612
Female	76,631	73	27	24	33	55	92	107	165	279	481	750
Black, both sexes	17,892	73	40	31	38	98	140	199	306	431	617	959
Male	8,586	44	16	13	20	75	94	128	205	264	384	584
Female	9,306	29	24	18	18	23	46	71	101	167	233	375
American Indian 1/, both sexes	643	4	1	-	2	4	8	5	12	17	21	38
Male	332	1	1	-	-	3	4	4	10	9	15	24
Female	311	3	-	-	2	1	4	1	2	8	6	14
Asian or Pacific Islander 2/, both sexes	2,157	15	11	7	6	6	15	13	25	30	37	52
Male	1,061	9	5	3	2	3	10	8	14	18	27	35
Female	1,096	6	6	4	4	3	5	5	11	12	10	17
Alabama												
All races, both sexes	6,008	11	2	-	2	7	18	28	45	64	112	169
Male	2,746	5	-	-	1	6	13	19	31	43	76	119
Female	3,262	6	2	-	1	1	5	9	14	21	36	50
White, both sexes	4,596	5	1	-	-	3	7	19	21	40	78	107
Male	2,082	3	-	-	-	2	6	14	14	27	54	76
Female	2,514	2	1	-	-	1	1	5	7	13	24	31
Black, both sexes	1,399	6	1	-	2	4	11	9	24	22	34	61
Male	656	2	-	-	1	4	7	5	17	16	22	42
Female	743	4	1	-	1	-	4	4	7	6	12	19
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Alaska												
All races, both sexes	179	1	-	-	1	1	4	2	1	2	6	4
Male	99	-	-	-	1	1	3	1	-	2	3	2
Female	80	1	-	-	-	-	1	1	1	-	3	2
White, both sexes	125	-	-	-	1	-	2	1	-	2	4	2
Male	74	-	-	-	1	-	2	1	-	2	2	1
Female	51	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	39	1	-	-	-	-	1	1	1	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	25	1	-	-	-	-	1	1	1	-	1	1
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	1	1	-	-	-	-	1
Male	6	-	-	-	-	1	1	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1,968	7	1	1	2	6	5	8	15	23	48	39
Male	955	1	1	1	1	4	3	6	9	16	32	26
Female	1,013	6	-	-	1	2	2	2	6	7	16	13
White, both sexes	1,846	4	1	1	2	6	3	5	10	17	43	29
Male	887	-	1	1	1	4	2	4	6	12	29	20
Female	959	4	-	-	1	2	1	1	4	5	14	9
Black, both sexes	41	-	-	-	-	-	1	2	4	3	4	5
Male	20	-	-	-	-	-	1	1	2	1	2	2
Female	21	-	-	-	-	-	-	1	2	2	2	3
American Indian 1/, both sexes	63	1	-	-	-	-	1	1	1	3	1	5
Male	42	1	-	-	-	-	-	1	1	3	1	4
Female	21	-	-	-	-	-	1	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	18	2	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	12	2	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Arkansas												
All races, both sexes	1,761	3	1	3	-	5	6	7	5	17	26	29
Male	787	1	-	1	-	5	5	5	4	11	18	17
Female	974	2	1	2	-	-	1	2	1	6	8	12
White, both sexes	1,517	2	1	3	-	4	2	2	5	8	14	19
Male	669	1	-	1	-	4	2	2	4	4	9	11
Female	848	1	1	2	-	-	-	-	1	4	5	8
Black, both sexes	241	1	-	-	-	1	4	5	-	9	12	10
Male	117	-	-	-	-	1	3	3	-	7	9	6
Female	124	1	-	-	-	-	1	2	-	2	3	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	11,927	24	13	7	15	27	48	72	81	135	200	303
Male	5,534	15	10	2	9	23	34	57	61	100	152	214
Female	6,393	9	3	5	6	4	14	15	20	35	48	89
White, both sexes	10,409	20	8	3	10	18	35	50	60	91	154	228
Male	4,789	11	7	-	5	16	25	45	47	71	121	167
Female	5,620	9	1	3	5	2	10	5	13	20	33	61
Black, both sexes	868	1	2	-	2	5	8	17	15	31	36	62
Male	448	1	-	-	2	4	5	10	10	20	23	36
Female	420	-	2	-	-	1	3	7	5	11	13	26
American Indian 1/, both sexes	52	-	-	-	-	1	-	-	2	2	2	2
Male	25	-	-	-	-	1	-	-	2	2	1	2
Female	27	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	598	3	3	4	3	3	5	5	4	11	8	11
Male	272	3	3	2	2	2	4	2	2	7	7	9
Female	326	-	-	2	1	1	1	3	2	4	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Colorado												
All races, both sexes	1,692	3	-	2	1	7	2	2	11	25	30	37
Male	720	1	-	1	-	5	1	1	8	18	18	24
Female	972	2	-	1	1	2	1	1	3	7	12	13
White, both sexes	1,630	2	-	2	-	7	2	2	11	21	27	32
Male	688	1	-	1	-	5	1	1	8	16	16	21
Female	942	1	-	1	-	2	1	1	3	5	11	11
Black, both sexes	42	-	-	-	-	-	-	-	-	3	2	5
Male	20	-	-	-	-	-	-	-	-	1	1	3
Female	22	-	-	-	-	-	-	-	-	2	1	2
American Indian 1/, both sexes	7	-	-	-	1	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	1	1	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	1	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	2,256	5	2	-	1	3	6	2	12	22	27	50
Male	1,008	1	1	-	-	2	4	2	9	18	17	41
Female	1,248	4	1	-	1	1	2	-	3	4	10	9
White, both sexes	2,105	5	2	-	-	2	6	1	8	21	24	44
Male	932	1	1	-	-	2	4	1	6	17	15	35
Female	1,173	4	1	-	-	-	2	-	2	4	9	9
Black, both sexes	139	-	-	-	1	1	-	1	4	1	3	5
Male	73	-	-	-	-	-	-	1	3	1	2	5
Female	66	-	-	-	1	1	-	-	1	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Delaware												
All races, both sexes	425	-	1	-	-	1	-	2	3	1	9	8
Male	188	-	-	-	-	1	-	1	3	1	8	7
Female	237	-	1	-	-	-	-	1	-	-	1	1
White, both sexes	347	-	-	-	-	-	-	2	1	1	5	4
Male	151	-	-	-	-	-	-	1	1	1	5	4
Female	196	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	69	-	1	-	-	1	-	-	2	-	4	3
Male	33	-	-	-	-	1	-	-	2	-	3	2
Female	36	-	1	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	178	-	-	-	1	1	1	-	2	2	4	10
Male	70	-	-	-	1	1	-	-	2	1	2	3
Female	108	-	-	-	-	-	1	-	-	1	2	7
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	144	-	-	-	1	1	1	-	2	1	4	10
Male	56	-	-	-	1	1	-	-	2	1	2	3
Female	88	-	-	-	-	-	1	-	-	-	2	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Florida												
All races, both sexes	8,244	12	6	2	3	17	28	26	44	68	103	176
Male	3,919	7	4	-	2	15	19	18	27	42	70	112
Female	4,325	5	2	2	1	2	9	8	17	26	33	64
White, both sexes	7,353	8	2	1	2	10	18	13	23	43	72	119
Male	3,491	4	1	-	2	9	10	8	13	31	56	77
Female	3,862	4	1	1	-	1	8	5	10	12	16	42
Black, both sexes	842	3	1	1	1	7	9	12	20	24	30	55
Male	400	2	1	-	-	6	8	9	14	11	13	34
Female	442	1	-	1	1	1	1	3	6	13	17	21
American Indian 1/, both sexes	6	-	1	-	-	-	-	1	-	-	1	-
Male	4	-	1	-	-	-	-	1	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	43	1	2	-	-	-	1	-	1	1	-	2
Male	24	1	1	-	-	-	1	-	-	-	-	1
Female	19	-	1	-	-	-	-	-	1	1	-	1
Georgia												
All races, both sexes	6,415	10	8	3	8	16	17	32	42	59	106	173
Male	2,879	8	5	1	3	11	12	21	29	32	66	108
Female	3,536	2	3	2	5	5	5	11	13	27	40	65
White, both sexes	4,748	4	1	2	5	4	3	17	21	19	51	94
Male	2,091	3	1	1	2	3	3	13	15	10	33	65
Female	2,657	1	-	1	3	1	-	4	6	9	18	29
Black, both sexes	1,615	5	6	1	3	12	12	15	20	39	50	78
Male	758	4	4	-	1	8	7	8	13	22	31	42
Female	857	1	2	1	2	4	5	7	7	17	19	36
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	38	1	1	-	-	-	2	-	1	-	5	1
Male	24	1	-	-	-	-	2	-	1	-	2	1
Female	14	-	1	-	-	-	-	-	-	-	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Hawaii												
All races, both sexes	980	1	-	1	1	-	1	3	6	7	21	23
Male	511	1	-	-	1	-	1	1	5	3	16	12
Female	469	-	-	1	-	-	-	2	1	4	5	11
White, both sexes	249	-	-	-	-	-	-	1	1	-	5	4
Male	121	-	-	-	-	-	-	1	1	-	4	1
Female	128	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	9	-	-	-	1	-	-	-	-	-	1	1
Male	7	-	-	-	1	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	719	1	-	1	-	-	1	2	5	7	15	18
Male	382	1	-	-	-	-	1	-	4	3	11	10
Female	337	-	-	1	-	-	-	2	1	4	4	8
Idaho												
All races, both sexes	655	-	1	-	1	1	1	2	3	5	5	8
Male	276	-	1	-	-	1	-	2	2	1	4	8
Female	379	-	-	-	1	-	1	-	1	4	1	-
White, both sexes	649	-	1	-	1	1	1	2	3	5	5	8
Male	273	-	1	-	-	1	-	2	2	1	4	8
Female	376	-	-	-	1	-	1	-	1	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Illinois												
All races, both sexes	7,073	9	5	2	8	24	28	41	43	69	114	155
Male	3,051	4	1	-	4	12	16	33	29	50	82	105
Female	4,022	5	4	2	4	12	12	8	14	19	32	50
White, both sexes	6,098	5	2	1	4	16	18	25	22	48	81	88
Male	2,559	1	1	-	2	7	11	22	16	38	58	66
Female	3,539	4	1	1	2	9	7	3	6	10	23	22
Black, both sexes	910	4	3	1	3	8	9	16	20	20	32	66
Male	456	3	-	-	2	5	5	11	12	11	23	38
Female	454	1	3	1	1	3	4	5	8	9	9	28
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	62	-	-	-	1	-	1	-	1	-	1	1
Male	35	-	-	-	-	-	-	-	1	-	1	1
Female	27	-	-	-	1	-	1	-	-	-	-	-
Indiana												
All races, both sexes	4,317	5	3	3	3	12	14	13	27	38	52	97
Male	1,843	3	1	2	1	8	7	9	20	22	35	55
Female	2,474	2	2	1	2	4	7	4	7	16	17	42
White, both sexes	4,009	4	1	1	2	10	14	12	22	34	38	70
Male	1,705	3	1	1	-	7	7	8	15	20	26	38
Female	2,304	1	-	-	2	3	7	4	7	14	12	32
Black, both sexes	294	1	1	2	1	2	-	1	5	4	14	26
Male	133	-	-	1	1	1	-	1	5	2	9	17
Female	161	1	1	1	-	1	-	-	-	2	5	9
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	12	-	1	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Iowa												
All races, both sexes	1,177	-	2	1	1	2	3	5	5	14	11	28
Male	488	-	2	1	1	1	2	4	1	9	6	15
Female	689	-	-	-	-	1	1	1	4	5	5	13
White, both sexes	1,154	-	2	1	1	2	3	4	4	12	9	25
Male	478	-	2	1	1	1	2	3	1	8	4	14
Female	676	-	-	-	-	1	1	1	3	4	5	11
Black, both sexes	21	-	-	-	-	-	-	1	-	2	2	2
Male	10	-	-	-	-	-	-	1	-	1	2	1
Female	11	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Kansas												
All races, both sexes	1,823	2	-	2	-	4	4	7	6	13	15	34
Male	759	2	-	1	-	3	3	5	5	8	12	21
Female	1,064	-	-	1	-	1	1	2	1	5	3	13
White, both sexes	1,708	1	-	2	-	2	4	7	5	9	14	23
Male	699	1	-	1	-	1	3	5	5	6	11	13
Female	1,009	-	-	1	-	1	1	2	-	3	3	10
Black, both sexes	91	1	-	-	-	2	-	-	1	2	1	9
Male	47	1	-	-	-	2	-	-	-	1	1	8
Female	44	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	1	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	2
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Kentucky												
All races, both sexes	3,142	11	2	-	3	2	7	6	17	23	39	54
Male	1,362	7	1	-	2	2	4	3	12	13	26	37
Female	1,780	4	1	-	1	-	3	3	5	10	13	17
White, both sexes	2,926	4	2	-	3	2	6	5	14	19	35	43
Male	1,259	3	1	-	2	2	4	3	9	9	22	31
Female	1,667	1	1	-	1	-	2	2	5	10	13	12
Black, both sexes	206	5	-	-	-	-	1	1	3	4	4	11
Male	96	3	-	-	-	-	-	-	3	4	4	6
Female	110	2	-	-	-	-	1	1	-	-	-	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	2	-	-	-	-	-	-	-	-	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3,225	6	2	4	3	8	12	17	13	32	46	72
Male	1,429	2	2	2	2	6	9	13	7	17	31	40
Female	1,796	4	-	2	1	2	3	4	6	15	15	32
White, both sexes	2,339	4	1	1	2	4	7	5	8	12	25	36
Male	1,038	1	1	-	2	2	5	5	6	7	18	23
Female	1,301	3	-	1	-	2	2	-	2	5	7	13
Black, both sexes	875	2	1	3	1	4	5	12	5	19	21	36
Male	388	1	1	2	-	4	4	8	1	9	13	17
Female	487	1	-	1	1	-	1	4	4	10	8	19
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Maine												
All races, both sexes	854	1	2	-	-	2	3	3	3	6	9	15
Male	378	-	-	-	-	1	2	2	2	5	4	12
Female	476	1	2	-	-	1	1	1	1	1	5	3
White, both sexes	849	1	1	-	-	2	3	3	3	6	8	15
Male	375	-	-	-	-	1	2	2	2	5	3	12
Female	474	1	1	-	-	1	1	1	1	1	5	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,286	5	6	1	2	9	14	18	26	24	53	75
Male	1,026	3	2	-	1	8	9	16	16	14	36	50
Female	1,260	2	4	1	1	1	5	2	10	10	17	25
White, both sexes	1,706	1	1	-	2	5	9	9	7	10	27	41
Male	749	1	1	-	1	4	6	8	4	6	20	28
Female	957	-	-	-	1	1	3	1	3	4	7	13
Black, both sexes	546	3	4	1	-	4	4	9	18	13	26	32
Male	259	1	-	-	-	4	3	8	12	7	16	20
Female	287	2	4	1	-	-	1	1	6	6	10	12
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	33	1	1	-	-	-	1	-	1	-	-	2
Male	17	1	1	-	-	-	-	-	-	-	-	2
Female	16	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Massachusetts												
All races, both sexes	4,309	3	3	1	2	2	10	12	7	19	40	61
Male	1,803	2	1	1	2	2	8	11	3	14	27	45
Female	2,506	1	2	-	-	-	2	1	4	5	13	16
White, both sexes	4,103	2	2	1	1	2	8	11	5	14	35	55
Male	1,704	2	1	1	1	2	6	10	1	11	23	41
Female	2,399	-	1	-	-	-	2	1	4	3	12	14
Black, both sexes	170	1	1	-	1	-	2	1	1	4	4	5
Male	78	-	-	-	1	-	2	1	1	2	3	3
Female	92	1	1	-	-	-	-	-	-	2	1	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	1	1	1	1
Male	21	-	-	-	-	-	-	-	1	1	1	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	5,648	8	2	9	5	13	17	22	33	55	83	130
Male	2,440	5	-	3	4	11	11	14	26	37	59	90
Female	3,208	3	2	6	1	2	6	8	7	18	24	40
White, both sexes	4,972	6	1	4	3	10	12	16	18	30	60	83
Male	2,106	4	-	3	3	8	8	12	16	20	47	63
Female	2,866	2	1	1	-	2	4	4	2	10	13	20
Black, both sexes	635	2	1	5	2	3	5	6	13	25	22	44
Male	317	1	-	-	1	3	3	2	8	17	11	25
Female	318	1	1	5	1	-	2	4	5	8	11	19
American Indian 1/, both sexes	18	-	-	-	-	-	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	1	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	1	-	1	2
Male	7	-	-	-	-	-	-	-	1	-	1	1
Female	16	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Minnesota												
All races, both sexes	2,638	1	1	1	1	3	5	8	5	6	24	35
Male	1,171	-	1	-	-	2	5	6	5	4	12	24
Female	1,467	1	-	1	1	1	-	2	-	2	12	11
White, both sexes	2,568	1	1	1	1	3	5	6	5	4	19	29
Male	1,129	-	1	-	-	2	5	4	5	3	9	19
Female	1,439	1	-	1	1	1	-	2	-	1	10	10
Black, both sexes	39	-	-	-	-	-	-	2	-	1	4	5
Male	26	-	-	-	-	-	-	2	-	1	2	4
Female	13	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	19	-	-	-	-	-	-	-	-	1	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	1
Female	8	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2,885	1	1	2	1	4	8	7	16	26	31	70
Male	1,217	1	-	2	-	3	2	2	11	14	18	41
Female	1,668	-	1	-	1	1	6	5	5	12	13	29
White, both sexes	1,933	-	-	-	-	-	-	3	5	7	11	32
Male	784	-	-	-	-	-	-	-	2	2	7	17
Female	1,149	-	-	-	-	-	-	3	3	5	4	15
Black, both sexes	943	1	1	2	1	4	8	4	11	19	20	38
Male	428	1	-	2	-	3	2	2	9	12	11	24
Female	515	-	1	-	1	1	6	2	2	7	9	14
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Missouri												
All races, both sexes	3,774	6	3	2	3	9	8	16	22	33	41	75
Male	1,644	4	1	1	2	6	7	13	16	20	26	46
Female	2,130	2	2	1	1	3	1	3	6	13	15	29
White, both sexes	3,445	6	2	2	3	4	7	10	18	25	29	58
Male	1,491	4	1	1	2	1	6	9	13	14	19	37
Female	1,954	2	1	1	1	3	1	1	5	11	10	21
Black, both sexes	311	-	-	-	-	5	1	6	4	8	12	17
Male	147	-	-	-	-	5	1	4	3	6	7	9
Female	164	-	-	-	-	-	-	2	1	2	5	8
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	1	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	646	1	2	-	2	-	-	-	5	3	6	11
Male	294	-	2	-	2	-	-	-	1	2	4	9
Female	352	1	-	-	-	-	-	-	4	1	2	2
White, both sexes	620	1	2	-	2	-	-	-	5	2	3	8
Male	278	-	2	-	2	-	-	-	1	2	1	6
Female	342	1	-	-	-	-	-	-	4	-	2	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	24	-	-	-	-	-	-	-	-	1	3	3
Male	14	-	-	-	-	-	-	-	-	-	3	3
Female	10	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Nebraska												
All races, both sexes	1,325	2	-	-	-	1	4	-	4	2	6	18
Male	560	1	-	-	-	1	1	-	2	1	4	12
Female	765	1	-	-	-	-	3	-	2	1	2	6
White, both sexes	1,283	1	-	-	-	1	2	-	2	2	6	17
Male	538	-	-	-	-	1	-	-	2	1	4	12
Female	745	1	-	-	-	-	2	-	-	1	2	5
Black, both sexes	32	1	-	-	-	-	1	-	2	-	-	1
Male	15	1	-	-	-	-	1	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	2	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1,793	3	-	-	1	4	8	16	7	22	51	58
Male	949	1	-	-	1	3	6	13	6	18	30	39
Female	844	2	-	-	-	1	2	3	1	4	21	19
White, both sexes	1,559	2	-	-	1	3	6	9	5	14	37	44
Male	819	1	-	-	1	3	4	8	4	12	20	31
Female	740	1	-	-	-	-	2	1	1	2	17	13
Black, both sexes	155	1	-	-	-	-	1	6	1	6	10	13
Male	84	-	-	-	-	-	1	4	1	4	7	7
Female	71	1	-	-	-	-	-	2	-	2	3	6
American Indian 1/, both sexes	20	-	-	-	-	-	1	-	1	-	1	-
Male	13	-	-	-	-	-	1	-	1	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	59	-	-	-	-	1	-	1	-	2	3	1
Male	33	-	-	-	-	-	-	1	-	2	2	1
Female	26	-	-	-	-	1	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
New Hampshire												
All races, both sexes	585	-	-	-	-	2	1	4	2	7	6	8
Male	248	-	-	-	-	-	1	4	2	5	4	6
Female	337	-	-	-	-	2	-	-	-	2	2	2
White, both sexes	582	-	-	-	-	2	1	4	2	7	6	8
Male	246	-	-	-	-	-	1	4	2	5	4	6
Female	336	-	-	-	-	2	-	-	-	2	2	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4,571	4	2	-	3	12	15	21	27	34	57	113
Male	2,056	2	1	-	2	7	14	17	22	24	30	78
Female	2,515	2	1	-	1	5	1	4	5	10	27	35
White, both sexes	3,985	4	1	-	1	11	12	14	12	19	36	73
Male	1,760	2	-	-	1	7	11	11	12	14	19	54
Female	2,225	2	1	-	-	4	1	3	-	5	17	19
Black, both sexes	522	-	1	-	2	1	3	6	13	14	21	36
Male	260	-	1	-	1	-	3	5	8	10	11	22
Female	262	-	-	-	1	1	-	1	5	4	10	14
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	59	-	-	-	-	-	-	1	2	1	-	3
Male	32	-	-	-	-	-	-	1	2	-	-	1
Female	27	-	-	-	-	-	-	-	-	1	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
New Mexico												
All races, both sexes	804	3	-	-	-	-	-	3	2	5	12	21
Male	369	1	-	-	-	-	-	1	2	4	7	13
Female	435	2	-	-	-	-	-	2	-	1	5	8
White, both sexes	741	1	-	-	-	-	-	3	1	4	9	19
Male	341	1	-	-	-	-	-	1	1	3	5	12
Female	400	-	-	-	-	-	-	2	-	1	4	7
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	50	2	-	-	-	-	-	-	1	1	3	1
Male	25	-	-	-	-	-	-	-	1	1	2	1
Female	25	2	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	7,422	15	7	4	4	8	19	31	33	48	68	134
Male	3,206	8	3	3	2	6	14	23	19	29	42	93
Female	4,216	7	4	1	2	2	5	8	14	19	26	41
White, both sexes	6,673	7	5	2	2	3	12	19	22	30	43	104
Male	2,881	5	2	2	1	3	8	14	13	17	26	73
Female	3,792	2	3	-	1	-	4	5	9	13	17	31
Black, both sexes	675	7	1	1	1	4	7	11	10	18	25	30
Male	295	3	1	1	1	3	6	8	6	12	16	20
Female	380	4	-	-	-	1	1	3	4	6	9	10
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	70	1	1	1	1	1	-	1	1	-	-	-
Male	28	-	-	-	-	-	-	1	-	-	-	-
Female	42	1	1	1	1	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
North Carolina												
All races, both sexes	4,553	6	9	8	3	8	17	15	33	38	63	95
Male	2,005	3	6	6	2	3	15	7	21	21	44	72
Female	2,548	3	3	2	1	5	2	8	12	17	19	23
White, both sexes	3,555	2	5	2	1	6	9	7	13	19	34	47
Male	1,528	-	3	2	1	3	9	3	10	11	23	37
Female	2,027	2	2	-	-	3	-	4	3	8	11	10
Black, both sexes	947	4	4	6	2	2	8	7	18	19	27	46
Male	454	3	3	4	1	-	6	3	10	10	20	33
Female	493	1	1	2	1	2	2	4	8	9	7	13
American Indian 1/, both sexes	35	-	-	-	-	-	-	-	-	-	2	2
Male	17	-	-	-	-	-	-	-	-	-	1	2
Female	18	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	1	2	-	-	-
Male	6	-	-	-	-	-	-	1	1	-	-	-
Female	10	-	-	-	-	-	-	-	1	-	-	-
North Dakota												
All races, both sexes	278	-	-	-	-	1	2	1	2	1	6	5
Male	110	-	-	-	-	1	1	1	1	1	5	2
Female	168	-	-	-	-	-	1	-	1	-	1	3
White, both sexes	272	-	-	-	-	1	2	1	2	1	3	4
Male	107	-	-	-	-	1	1	1	1	1	3	2
Female	165	-	-	-	-	-	1	-	1	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	2	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Ohio												
All races, both sexes	6,163	14	11	5	7	8	16	21	34	63	87	121
Male	2,708	7	9	2	6	7	11	13	25	42	54	76
Female	3,455	7	2	3	1	1	5	8	9	21	33	45
White, both sexes	5,538	6	9	4	5	6	12	16	24	43	64	100
Male	2,400	4	8	2	5	5	9	11	17	32	43	63
Female	3,138	2	1	2	-	1	3	5	7	11	21	37
Black, both sexes	615	8	2	1	2	2	4	5	10	20	23	21
Male	304	3	1	-	1	2	2	2	8	10	11	13
Female	311	5	1	1	1	-	2	3	2	10	12	8
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2,193	4	2	1	1	7	8	7	17	15	32	51
Male	900	3	1	1	-	5	7	5	10	9	22	31
Female	1,293	1	1	-	1	2	1	2	7	6	10	20
White, both sexes	1,945	2	2	1	1	4	6	4	13	10	24	32
Male	783	1	1	1	-	3	5	2	8	7	17	20
Female	1,162	1	1	-	1	1	1	2	5	3	7	12
Black, both sexes	140	2	-	-	-	2	-	1	2	3	7	11
Male	66	2	-	-	-	1	-	1	1	2	4	8
Female	74	-	-	-	-	1	-	-	1	1	3	3
American Indian 1/, both sexes	105	-	-	-	-	1	2	2	2	2	1	8
Male	51	-	-	-	-	1	2	2	1	-	1	3
Female	54	-	-	-	-	-	-	-	1	2	-	5
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Oregon												
All races, both sexes	2,099	1	-	1	3	3	6	3	9	15	16	22
Male	878	-	-	-	-	3	5	2	5	11	10	14
Female	1,221	1	-	1	3	-	1	1	4	4	6	8
White, both sexes	2,044	1	-	1	1	2	5	2	8	13	15	19
Male	851	-	-	-	-	2	4	1	5	10	9	11
Female	1,193	1	-	1	1	-	1	1	3	3	6	8
Black, both sexes	25	-	-	-	1	1	1	1	1	1	1	1
Male	10	-	-	-	-	1	1	1	-	1	1	1
Female	15	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	1	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	1	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	1	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	10,265	12	2	5	3	11	19	23	32	58	118	190
Male	4,420	6	1	3	3	11	13	18	19	39	76	121
Female	5,845	6	1	2	-	-	6	5	13	19	42	69
White, both sexes	9,412	11	-	5	3	7	13	15	24	39	83	150
Male	4,013	5	-	3	3	7	9	12	15	28	53	98
Female	5,399	6	-	2	-	-	4	3	9	11	30	52
Black, both sexes	821	-	2	-	-	4	6	7	7	19	34	39
Male	392	-	1	-	-	4	4	5	3	11	22	23
Female	429	-	1	-	-	-	2	2	4	8	12	16
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	1	-	-	-	-	-	1	-	-	1	1
Male	13	1	-	-	-	-	-	1	-	-	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Rhode Island												
All races, both sexes	378	1	2	-	3	-	1	2	4	5	9	7
Male	163	1	1	-	2	-	-	2	3	3	7	3
Female	215	-	1	-	1	-	1	-	1	2	2	4
White, both sexes	362	1	2	-	3	-	1	2	3	5	7	7
Male	151	1	1	-	2	-	-	2	2	3	6	3
Female	211	-	1	-	1	-	1	-	1	2	1	4
Black, both sexes	15	-	-	-	-	-	-	-	1	-	2	-
Male	11	-	-	-	-	-	-	-	1	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	3,014	7	2	2	5	8	9	18	30	36	64	82
Male	1,429	3	1	1	3	5	5	12	21	19	38	54
Female	1,585	4	1	1	2	3	4	6	9	17	26	28
White, both sexes	2,139	5	-	1	2	3	5	10	10	20	30	49
Male	981	2	-	-	1	3	3	6	7	9	20	34
Female	1,158	3	-	1	1	-	2	4	3	11	10	15
Black, both sexes	863	2	2	1	3	5	4	8	20	16	34	33
Male	441	1	1	1	2	2	2	6	14	10	18	20
Female	422	1	1	-	1	3	2	2	6	6	16	13
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
South Dakota												
All races, both sexes	256	1	-	-	1	-	-	1	2	1	3	5
Male	114	-	-	-	-	-	-	1	1	1	-	4
Female	142	1	-	-	1	-	-	-	1	-	3	1
White, both sexes	235	1	-	-	-	-	-	1	2	1	3	2
Male	102	-	-	-	-	-	-	1	1	1	-	2
Female	133	1	-	-	-	-	-	-	1	-	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	20	-	-	-	1	-	-	-	-	-	-	3
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	9	-	-	-	1	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2,685	5	2	1	1	9	10	26	24	35	56	90
Male	1,233	4	-	-	-	5	7	16	19	21	41	55
Female	1,452	1	2	1	1	4	3	10	5	14	15	35
White, both sexes	2,220	4	2	-	1	5	7	14	15	28	39	62
Male	987	3	-	-	-	3	5	8	12	18	27	35
Female	1,233	1	2	-	1	2	2	6	3	10	12	27
Black, both sexes	454	1	-	1	-	4	3	12	8	7	17	28
Male	243	1	-	-	-	2	2	8	7	3	14	20
Female	211	-	-	1	-	2	1	4	1	4	3	8
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Texas												
All races, both sexes	9,204	20	10	5	11	28	38	49	73	96	160	226
Male	4,119	14	6	2	7	22	29	32	51	72	111	145
Female	5,085	6	4	3	4	6	9	17	22	24	49	81
White, both sexes	8,034	14	7	4	6	19	24	40	48	64	117	159
Male	3,550	8	5	1	5	14	18	28	34	48	85	104
Female	4,484	6	2	3	1	5	6	12	14	16	32	55
Black, both sexes	1,093	6	3	-	5	9	13	9	24	30	42	64
Male	535	6	1	-	2	8	11	4	16	22	26	40
Female	558	-	2	-	3	1	2	5	8	8	16	24
American Indian 1/, both sexes	7	-	-	-	-	-	1	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	70	-	-	1	-	-	-	-	-	2	1	3
Male	31	-	-	1	-	-	-	-	-	2	-	1
Female	39	-	-	-	-	-	-	-	-	-	1	2
Utah												
All races, both sexes	1,126	-	1	-	-	-	2	2	4	9	10	11
Male	438	-	-	-	-	-	2	-	2	6	6	8
Female	688	-	1	-	-	-	-	2	2	3	4	3
White, both sexes	1,102	-	1	-	-	-	2	2	4	9	8	10
Male	427	-	-	-	-	-	2	-	2	6	4	7
Female	675	-	1	-	-	-	-	2	2	3	4	3
Black, both sexes	12	-	-	-	-	-	-	-	-	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	2	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Vermont												
All races, both sexes	263	-	-	-	-	1	1	-	1	1	2	2
Male	115	-	-	-	-	1	1	-	1	1	1	2
Female	148	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	263	-	-	-	-	1	1	-	1	1	2	2
Male	115	-	-	-	-	1	1	-	1	1	1	2
Female	148	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4,438	7	3	7	4	6	19	19	25	30	59	90
Male	1,980	4	1	5	1	3	13	12	16	20	42	61
Female	2,458	3	2	2	3	3	6	7	9	10	17	29
White, both sexes	3,541	3	1	3	3	6	11	13	11	13	40	51
Male	1,571	1	-	3	1	3	8	8	7	8	30	35
Female	1,970	2	1	-	2	3	3	5	4	5	10	16
Black, both sexes	842	4	2	4	1	-	7	6	11	17	19	38
Male	381	3	1	2	-	-	4	4	7	12	12	25
Female	461	1	1	2	1	-	3	2	4	5	7	13
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	52	-	-	-	-	-	1	-	2	-	-	1
Male	26	-	-	-	-	-	1	-	1	-	-	1
Female	26	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Washington												
All races, both sexes	2,207	1	-	2	3	4	10	7	13	19	29	45
Male	995	-	-	-	1	3	6	4	8	12	18	26
Female	1,212	1	-	2	2	1	4	3	5	7	11	19
White, both sexes	2,075	-	-	1	2	2	6	6	11	15	23	40
Male	933	-	-	-	-	2	5	3	7	8	15	25
Female	1,142	-	-	1	2	-	1	3	4	7	8	15
Black, both sexes	60	-	-	1	1	-	1	-	1	3	4	4
Male	32	-	-	-	1	-	-	-	1	3	2	-
Female	28	-	-	1	-	-	1	-	-	-	2	4
American Indian 1/, both sexes	19	-	-	-	-	2	1	-	-	-	1	1
Male	10	-	-	-	-	1	1	-	-	-	-	1
Female	9	-	-	-	-	1	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	53	1	-	-	-	-	2	1	1	1	1	-
Male	20	-	-	-	-	-	-	1	-	1	1	-
Female	33	1	-	-	-	-	2	-	1	-	-	-
West Virginia												
All races, both sexes	1,390	-	1	-	1	-	2	4	3	4	16	20
Male	594	-	-	-	-	-	1	2	3	3	14	15
Female	796	-	1	-	1	-	1	2	-	1	2	5
White, both sexes	1,341	-	1	-	1	-	2	4	1	4	16	18
Male	574	-	-	-	-	-	1	2	1	3	14	14
Female	767	-	1	-	1	-	1	2	-	1	2	4
Black, both sexes	46	-	-	-	-	-	-	-	2	-	-	2
Male	19	-	-	-	-	-	-	-	2	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other non-pulmonary forms of heart disease (I30-I51)												
Wisconsin												
All races, both sexes	3,314	2	-	-	2	3	6	5	8	14	28	50
Male	1,429	2	-	-	-	2	4	5	6	6	23	38
Female	1,885	-	-	-	2	1	2	-	2	8	5	12
White, both sexes	3,184	1	-	-	2	3	6	5	5	10	23	43
Male	1,364	1	-	-	-	2	4	5	3	5	19	34
Female	1,820	-	-	-	2	1	2	-	2	5	4	9
Black, both sexes	102	1	-	-	-	-	-	-	3	4	5	5
Male	54	1	-	-	-	-	-	-	3	1	4	3
Female	48	-	-	-	-	-	-	-	-	3	1	2
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	270	-	-	-	1	-	-	-	1	2	2	6
Male	122	-	-	-	-	-	-	-	-	-	2	4
Female	148	-	-	-	1	-	-	-	1	2	-	2
White, both sexes	268	-	-	-	1	-	-	-	1	2	2	6
Male	121	-	-	-	-	-	-	-	-	-	2	4
Female	147	-	-	-	1	-	-	-	1	2	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
United States												
All races, both sexes	4,355	5,491	6,484	7,776	11,394	17,960	26,311	30,382	24,470	10,279	2,288	6
Male	2,892	3,571	3,982	4,555	6,276	8,991	11,565	11,370	7,330	2,247	384	4
Female	1,463	1,920	2,502	3,221	5,118	8,969	14,746	19,012	17,140	8,032	1,904	2
White, both sexes	3,031	4,048	4,991	6,167	9,510	15,662	23,811	28,167	22,804	9,520	2,049	5
Male	2,085	2,666	3,128	3,654	5,269	7,935	10,578	10,603	6,889	2,075	334	4
Female	946	1,382	1,863	2,513	4,241	7,727	13,233	17,564	15,915	7,445	1,715	1
Black, both sexes	1,210	1,302	1,308	1,399	1,646	1,986	2,108	1,829	1,347	614	210	1
Male	745	808	747	770	872	913	806	608	330	121	39	-
Female	465	494	561	629	774	1,073	1,302	1,221	1,017	493	171	1
American Indian 1/, both sexes	40	40	46	59	62	74	68	61	48	28	5	-
Male	21	28	25	34	41	30	30	23	20	7	2	-
Female	19	12	21	25	21	44	38	38	28	21	3	-
Asian or Pacific Islander 2/, both sexes	74	101	139	151	176	238	324	325	271	117	24	-
Male	41	69	82	97	94	113	151	136	91	44	9	-
Female	33	32	57	54	82	125	173	189	180	73	15	-
Alabama												
All races, both sexes	245	306	381	394	527	720	958	917	767	269	66	-
Male	143	195	237	220	294	363	390	319	199	60	13	-
Female	102	111	144	174	233	357	568	598	568	209	53	-
White, both sexes	160	207	268	282	400	537	782	781	638	217	43	-
Male	100	138	172	160	226	281	325	263	166	48	7	-
Female	60	69	96	122	174	256	457	518	472	169	36	-
Black, both sexes	85	98	113	110	125	180	175	136	129	51	23	-
Male	43	56	65	59	68	79	64	56	33	11	6	-
Female	42	42	48	51	57	101	111	80	96	40	17	-
American Indian 1/, both sexes	-	1	-	1	1	2	-	-	-	-	-	-
Male	-	1	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	1	-	-	1	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Alaska												
All races, both sexes	10	3	11	12	17	26	23	25	21	8	1	-
Male	10	2	8	10	13	10	11	12	6	3	1	-
Female	-	1	3	2	4	16	12	13	15	5	-	-
White, both sexes	7	1	8	7	10	17	19	20	17	6	1	-
Male	7	1	5	7	9	8	10	9	5	3	1	-
Female	-	-	3	-	1	9	9	11	12	3	-	-
Black, both sexes	-	1	1	1	-	-	-	1	1	-	-	-
Male	-	1	1	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	3	-	1	4	5	9	3	3	3	2	-	-
Male	3	-	1	2	3	2	1	1	1	-	-	-
Female	-	-	-	2	2	7	2	2	2	2	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	2	-	1	1	-	-	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	-	-
Arizona												
All races, both sexes	50	61	89	113	144	223	336	379	290	115	13	-
Male	39	35	61	65	86	104	162	172	99	31	1	-
Female	11	26	28	48	58	119	174	207	191	84	12	-
White, both sexes	40	52	84	102	136	214	325	365	288	106	13	-
Male	32	31	59	59	80	101	154	164	97	29	1	-
Female	8	21	25	43	56	113	171	201	191	77	12	-
Black, both sexes	3	3	-	2	1	1	4	5	-	3	-	-
Male	3	1	-	-	1	-	3	3	-	-	-	-
Female	-	2	-	2	-	1	1	2	-	3	-	-
American Indian 1/, both sexes	7	5	5	8	5	5	5	5	1	4	-	-
Male	4	3	2	5	5	2	5	3	1	1	-	-
Female	3	2	3	3	-	3	-	2	-	3	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	2	3	2	4	1	2	-	-
Male	-	-	-	1	-	1	-	2	1	1	-	-
Female	-	1	-	-	2	2	2	2	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Arkansas												
All races, both sexes	49	65	87	108	133	187	286	349	274	103	18	-
Male	35	43	54	65	77	94	125	117	84	21	5	-
Female	14	22	33	43	56	93	161	232	190	82	13	-
White, both sexes	32	48	69	86	111	167	266	319	248	97	14	-
Male	21	31	46	54	67	85	117	108	78	20	4	-
Female	11	17	23	32	44	82	149	211	170	77	10	-
Black, both sexes	17	17	18	22	20	20	20	29	26	6	4	-
Male	14	12	8	11	9	9	8	9	6	1	1	-
Female	3	5	10	11	11	11	12	20	20	5	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
California												
All races, both sexes	341	401	428	530	740	1,201	2,023	2,359	1,963	827	189	-
Male	220	258	282	302	422	611	933	919	672	209	29	-
Female	121	143	146	228	318	590	1,090	1,440	1,291	618	160	-
White, both sexes	248	295	341	426	607	1,042	1,804	2,192	1,831	768	178	-
Male	165	190	225	244	357	543	836	866	629	192	27	-
Female	83	105	116	182	250	499	968	1,326	1,202	576	151	-
Black, both sexes	69	74	52	70	84	86	100	69	54	27	4	-
Male	42	45	35	38	45	42	53	18	15	4	-	-
Female	27	29	17	32	39	44	47	51	39	23	4	-
American Indian 1/, both sexes	2	5	2	6	2	9	8	6	1	2	-	-
Male	-	3	1	4	1	2	3	3	-	-	-	-
Female	2	2	1	2	1	7	5	3	1	2	-	-
Asian or Pacific Islander 2/, both sexes	22	27	33	28	47	64	111	92	77	30	7	-
Male	13	20	21	16	19	24	41	32	28	13	2	-
Female	9	7	12	12	28	40	70	60	49	17	5	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Colorado												
All races, both sexes	42	38	64	72	108	174	271	364	268	142	29	-
Male	30	26	27	42	50	77	128	141	85	29	8	-
Female	12	12	37	30	58	97	143	223	183	113	21	-
White, both sexes	38	37	62	68	100	164	266	357	265	139	28	-
Male	26	25	26	39	44	72	126	140	84	29	7	-
Female	12	12	36	29	56	92	140	217	181	110	21	-
Black, both sexes	4	-	1	3	5	5	3	6	2	2	1	-
Male	4	-	-	2	4	2	1	-	1	-	1	-
Female	-	-	1	1	1	3	2	6	1	2	-	-
American Indian 1/, both sexes	-	1	-	-	2	-	-	-	-	1	-	-
Male	-	1	-	-	2	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	5	2	1	1	-	-	-
Male	-	-	1	1	-	3	1	1	-	-	-	-
Female	-	-	-	-	1	2	1	-	1	-	-	-
Connecticut												
All races, both sexes	57	61	88	103	135	211	368	475	409	179	40	-
Male	44	40	55	58	86	113	181	186	120	27	3	-
Female	13	21	33	45	49	98	187	289	289	152	37	-
White, both sexes	47	50	71	89	114	198	353	460	400	173	37	-
Male	37	33	47	53	71	106	174	182	118	26	3	-
Female	10	17	24	36	43	92	179	278	282	147	34	-
Black, both sexes	10	10	15	14	21	11	14	12	8	5	3	-
Male	7	7	8	5	15	6	7	4	2	-	-	-
Female	3	3	7	9	6	5	7	8	6	5	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	-	-	2	1	2	1	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	1	2	-	-	1	1	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Delaware												
All races, both sexes	7	18	14	37	30	44	76	85	54	30	5	-
Male	4	12	9	21	16	18	37	30	15	4	1	-
Female	3	6	5	16	14	26	39	55	39	26	4	-
White, both sexes	4	12	10	26	21	34	66	76	53	27	5	-
Male	2	10	7	18	9	13	34	27	15	3	1	-
Female	2	2	3	8	12	21	32	49	38	24	4	-
Black, both sexes	3	6	4	10	8	9	9	6	-	3	-	-
Male	2	2	2	2	6	5	3	2	-	1	-	-
Female	1	4	2	8	2	4	6	4	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	2	1	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
Dist. of Columbia												
All races, both sexes	12	5	13	10	11	13	25	36	17	11	4	-
Male	6	5	9	7	9	6	6	10	1	1	-	-
Female	6	-	4	3	2	7	19	26	16	10	4	-
White, both sexes	-	-	1	-	1	4	7	9	3	4	3	-
Male	-	-	1	-	1	2	2	5	1	1	-	-
Female	-	-	-	-	-	2	5	4	2	3	3	-
Black, both sexes	12	4	12	10	10	9	18	27	14	7	1	-
Male	6	4	8	7	8	4	4	5	-	-	-	-
Female	6	-	4	3	2	5	14	22	14	7	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Florida												
All races, both sexes	221	246	297	386	568	969	1,384	1,712	1,344	536	95	1
Male	148	162	179	245	337	489	656	728	487	152	20	-
Female	73	84	118	141	231	480	728	984	857	384	75	1
White, both sexes	158	187	234	317	480	866	1,273	1,641	1,284	515	87	-
Male	114	128	145	206	289	441	611	713	470	145	18	-
Female	44	59	89	111	191	425	662	928	814	370	69	-
Black, both sexes	58	58	61	67	83	97	104	66	55	21	8	1
Male	32	33	33	38	44	43	41	15	14	7	2	-
Female	26	25	28	29	39	54	63	51	41	14	6	1
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	1	2	5	6	7	5	4	-	-	-
Male	2	1	-	1	4	5	4	-	3	-	-	-
Female	2	-	1	1	1	1	3	5	1	-	-	-
Georgia												
All races, both sexes	272	319	393	432	593	806	991	1,018	758	280	77	2
Male	174	197	246	263	320	388	384	360	176	58	15	2
Female	98	122	147	169	273	418	607	658	582	222	62	-
White, both sexes	160	200	245	273	429	651	821	842	640	211	53	2
Male	100	127	151	165	228	322	332	312	151	42	10	2
Female	60	73	94	108	201	329	489	530	489	169	43	-
Black, both sexes	112	116	144	153	157	150	166	174	111	67	24	-
Male	74	68	92	94	89	62	49	47	23	15	5	-
Female	38	48	52	59	68	88	117	127	88	52	19	-
American Indian 1/, both sexes	-	1	1	1	4	1	-	-	4	1	-	-
Male	-	1	1	-	1	1	-	-	2	-	-	-
Female	-	-	-	1	3	-	-	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	-	2	3	5	3	4	4	2	3	1	-	-
Male	-	1	2	4	2	3	3	1	-	1	-	-
Female	-	1	1	1	1	1	1	1	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Hawaii												
All races, both sexes	32	47	58	68	69	103	137	170	148	73	11	-
Male	19	39	39	45	39	55	80	74	54	24	3	-
Female	13	8	19	23	30	48	57	96	94	49	8	-
White, both sexes	10	10	13	18	15	28	33	49	41	18	3	-
Male	8	8	11	13	5	13	16	18	16	5	1	-
Female	2	2	2	5	10	15	17	31	25	13	2	-
Black, both sexes	1	1	-	-	1	1	1	1	-	-	-	-
Male	1	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	3	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	36	45	50	53	74	103	117	107	55	8	-
Male	10	30	28	32	33	41	64	55	38	19	2	-
Female	11	6	17	18	20	33	39	62	69	36	6	-
Idaho												
All races, both sexes	10	15	18	25	35	70	113	147	128	56	11	-
Male	8	11	13	13	20	33	43	51	51	11	3	-
Female	2	4	5	12	15	37	70	96	77	45	8	-
White, both sexes	10	15	17	25	35	68	111	146	128	56	11	-
Male	8	11	13	13	20	32	42	50	51	11	3	-
Female	2	4	4	12	15	36	69	96	77	45	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	2	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Illinois												
All races, both sexes	187	213	278	340	470	797	1,145	1,367	1,137	520	121	-
Male	130	126	170	200	271	396	521	482	311	90	18	-
Female	57	87	108	140	199	401	624	885	826	430	103	-
White, both sexes	120	149	211	252	385	682	1,031	1,282	1,074	491	111	-
Male	84	88	135	149	221	349	470	445	297	84	15	-
Female	36	61	76	103	164	333	561	837	777	407	96	-
Black, both sexes	65	63	59	81	79	111	107	73	56	25	9	-
Male	44	38	31	46	44	45	49	30	12	5	2	-
Female	21	25	28	35	35	66	58	43	44	20	7	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	7	7	6	4	6	12	7	4	1	-
Male	2	-	4	5	6	2	2	7	2	1	1	-
Female	-	1	3	2	-	2	4	5	5	3	-	-
Indiana												
All races, both sexes	100	152	196	205	349	514	706	835	657	270	66	-
Male	62	95	115	115	176	247	315	295	198	53	9	-
Female	38	57	81	90	173	267	391	540	459	217	57	-
White, both sexes	84	135	176	181	320	489	668	801	627	260	60	-
Male	52	85	102	104	163	241	308	281	187	49	7	-
Female	32	50	74	77	157	248	360	520	440	211	53	-
Black, both sexes	16	17	20	22	26	25	36	33	29	8	5	-
Male	10	10	13	10	13	6	6	14	10	3	1	-
Female	6	7	7	12	13	19	30	19	19	5	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	3	-	1	1	1	2	1	-
Male	-	-	-	1	-	-	1	-	1	1	1	-
Female	-	-	-	1	3	-	-	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Iowa												
All races, both sexes	25	25	37	41	61	118	229	225	213	101	30	-
Male	17	14	20	22	34	62	100	84	64	21	8	-
Female	8	11	17	19	27	56	129	141	149	80	22	-
White, both sexes	25	23	34	40	59	113	228	225	213	101	30	-
Male	17	14	19	22	33	60	99	84	64	21	8	-
Female	8	9	15	18	26	53	129	141	149	80	22	-
Black, both sexes	-	2	3	1	2	5	1	-	-	-	-	-
Male	-	-	1	-	1	2	1	-	-	-	-	-
Female	-	2	2	1	1	3	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	36	44	54	69	113	205	289	371	349	165	41	-
Male	24	21	33	42	55	98	135	138	103	43	7	-
Female	12	23	21	27	58	107	154	233	246	122	34	-
White, both sexes	31	42	45	66	103	184	274	355	344	157	40	-
Male	22	20	26	40	50	86	129	134	100	39	7	-
Female	9	22	19	26	53	98	145	221	244	118	33	-
Black, both sexes	4	2	9	1	9	14	12	13	4	7	-	-
Male	2	1	7	-	4	7	4	4	2	3	-	-
Female	2	1	2	1	5	7	8	9	2	4	-	-
American Indian 1/, both sexes	1	-	-	2	1	5	2	3	1	1	1	-
Male	-	-	-	2	1	4	2	-	1	1	-	-
Female	1	-	-	-	-	1	-	3	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Kentucky												
All races, both sexes	99	119	143	181	279	382	534	585	446	178	32	-
Male	72	82	101	95	160	191	208	189	123	31	3	-
Female	27	37	42	86	119	191	326	396	323	147	29	-
White, both sexes	85	105	135	167	252	354	505	559	431	170	30	-
Male	62	70	97	89	149	171	202	180	120	30	3	-
Female	23	35	38	78	103	183	303	379	311	140	27	-
Black, both sexes	14	13	7	13	26	26	29	25	14	8	2	-
Male	10	11	3	5	11	18	6	8	3	1	-	-
Female	4	2	4	8	15	8	23	17	11	7	2	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	2	-	1	1	-	-	-
Male	-	-	1	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Louisiana												
All races, both sexes	107	132	155	197	257	404	496	600	434	184	44	-
Male	69	73	90	120	133	200	221	218	126	38	10	-
Female	38	59	65	77	124	204	275	382	308	146	34	-
White, both sexes	57	64	98	129	167	299	397	489	356	144	34	-
Male	40	39	62	79	82	164	180	189	99	27	7	-
Female	17	25	36	50	85	135	217	300	257	117	27	-
Black, both sexes	49	67	57	66	89	105	97	110	77	39	10	-
Male	29	34	28	40	51	36	41	28	27	11	3	-
Female	20	33	29	26	38	69	56	82	50	28	7	-
American Indian 1/, both sexes	1	-	-	2	1	-	1	-	1	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	-	1	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Maine												
All races, both sexes	26	38	37	44	60	95	135	158	133	73	11	-
Male	17	29	21	31	41	45	55	59	40	10	2	-
Female	9	9	16	13	19	50	80	99	93	63	9	-
White, both sexes	26	38	37	44	59	94	135	157	133	73	11	-
Male	17	29	21	31	40	45	55	58	40	10	2	-
Female	9	9	16	13	19	49	80	99	93	63	9	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	70	85	103	109	167	263	395	398	313	117	33	-
Male	48	55	58	68	80	121	158	157	87	33	6	-
Female	22	30	45	41	87	142	237	241	226	84	27	-
White, both sexes	34	51	65	74	117	201	317	343	261	105	26	-
Male	26	34	34	47	52	94	132	138	76	31	6	-
Female	8	17	31	27	65	107	185	205	185	74	20	-
Black, both sexes	34	32	35	33	49	62	69	50	50	11	7	-
Male	21	21	22	19	27	27	20	19	11	1	-	-
Female	13	11	13	14	22	35	49	31	39	10	7	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	3	2	1	-	9	5	2	1	-	-
Male	1	-	2	2	1	-	6	-	-	1	-	-
Female	1	2	1	-	-	-	3	5	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Massachusetts												
All races, both sexes	71	110	112	176	279	460	722	937	816	369	97	-
Male	45	74	73	109	156	240	322	360	215	74	19	-
Female	26	36	39	67	123	220	400	577	601	295	78	-
White, both sexes	65	94	100	161	262	434	687	911	796	364	93	-
Male	42	64	66	101	151	225	305	347	213	72	19	-
Female	23	30	34	60	111	209	382	564	583	292	74	-
Black, both sexes	6	14	11	13	16	19	26	23	15	5	2	-
Male	3	9	6	7	5	10	12	10	1	2	-	-
Female	3	5	5	6	11	9	14	13	14	3	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	3	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	3	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	1	1	7	8	3	2	-	2	-
Male	-	1	1	1	-	5	5	3	1	-	-	-
Female	-	1	-	-	1	2	3	-	1	-	2	-
Michigan												
All races, both sexes	155	184	177	232	351	655	988	1,143	920	383	83	-
Male	98	113	119	129	183	312	423	433	277	85	8	-
Female	57	71	58	103	168	343	565	710	643	298	75	-
White, both sexes	102	129	134	189	300	570	908	1,077	879	363	78	-
Male	69	78	89	110	154	271	391	409	263	80	8	-
Female	33	51	45	79	146	299	517	668	616	283	70	-
Black, both sexes	52	50	41	39	49	79	75	61	38	18	5	-
Male	29	32	30	17	28	39	29	24	13	5	-	-
Female	23	18	11	22	21	40	46	37	25	13	5	-
American Indian 1/, both sexes	-	5	1	1	1	3	1	2	1	1	-	-
Male	-	3	-	1	1	1	1	-	1	-	-	-
Female	-	2	1	-	-	2	-	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	3	1	3	4	3	2	1	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	1	-	1	2	1	2	2	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Minnesota												
All races, both sexes	44	65	73	92	151	270	398	598	564	256	37	-
Male	29	37	45	70	91	163	195	229	190	58	5	-
Female	15	28	28	22	60	107	203	369	374	198	32	-
White, both sexes	35	59	66	90	146	265	393	589	560	254	36	-
Male	23	33	40	69	87	160	193	225	189	58	4	-
Female	12	26	26	21	59	105	200	364	371	196	32	-
Black, both sexes	4	4	4	1	2	3	4	2	2	1	-	-
Male	3	2	3	-	2	2	2	2	1	-	-	-
Female	1	2	1	1	-	1	2	-	1	1	-	-
American Indian 1/, both sexes	4	1	3	1	3	1	-	2	1	-	-	-
Male	3	1	2	1	2	-	-	-	-	-	-	-
Female	1	-	1	-	1	1	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	1	1	5	1	1	1	-
Male	-	1	-	-	-	1	-	2	-	-	1	-
Female	1	-	-	-	-	-	1	3	1	1	-	-
Mississippi												
All races, both sexes	79	134	159	166	223	386	449	483	411	179	49	-
Male	49	91	94	91	116	193	174	161	114	32	8	-
Female	30	43	65	75	107	193	275	322	297	147	41	-
White, both sexes	34	67	92	93	145	268	335	379	309	130	23	-
Male	22	46	59	52	77	139	133	125	82	19	2	-
Female	12	21	33	41	68	129	202	254	227	111	21	-
Black, both sexes	44	67	64	72	77	117	114	103	101	49	26	-
Male	27	45	34	38	38	53	41	36	31	13	6	-
Female	17	22	30	34	39	64	73	67	70	36	20	-
American Indian 1/, both sexes	-	-	2	-	1	1	-	-	1	-	-	-
Male	-	-	1	-	1	1	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Missouri												
All races, both sexes	92	137	171	175	276	445	652	692	597	267	52	-
Male	59	95	101	100	147	236	276	245	178	54	11	-
Female	33	42	70	75	129	209	376	447	419	213	41	-
White, both sexes	72	111	145	162	241	400	616	661	567	258	48	-
Male	48	79	87	94	135	215	262	236	167	52	9	-
Female	24	32	58	68	106	185	354	425	400	206	39	-
Black, both sexes	19	24	24	12	33	43	34	29	28	8	4	-
Male	11	16	13	5	11	21	13	9	9	2	2	-
Female	8	8	11	7	22	22	21	20	19	6	2	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	1	1	2	2	2	1	1	-	-
Male	-	-	1	1	-	-	1	-	1	-	-	-
Female	1	2	1	-	1	2	1	2	-	1	-	-
Montana												
All races, both sexes	22	17	23	32	36	63	108	123	131	50	11	-
Male	15	11	16	21	22	32	59	45	40	10	3	-
Female	7	6	7	11	14	31	49	78	91	40	8	-
White, both sexes	19	15	21	30	36	60	104	123	129	49	11	-
Male	14	10	14	19	22	31	57	45	39	10	3	-
Female	5	5	7	11	14	29	47	78	90	39	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	2	2	2	-	3	3	-	1	1	-	-
Male	1	1	2	2	-	1	1	-	-	-	-	-
Female	2	1	-	-	-	2	2	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Nebraska												
All races, both sexes	24	30	40	51	108	127	221	271	244	128	44	-
Male	21	23	26	26	63	74	97	98	74	27	9	-
Female	3	7	14	25	45	53	124	173	170	101	35	-
White, both sexes	22	29	37	49	102	122	217	262	241	128	43	-
Male	19	22	23	26	61	70	96	92	73	27	9	-
Female	3	7	14	23	41	52	121	170	168	101	34	-
Black, both sexes	1	-	3	1	6	3	4	6	3	-	-	-
Male	1	-	3	-	2	2	1	3	1	-	-	-
Female	-	-	-	1	4	1	3	3	2	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	1	-	-	1	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	2	-	2	-	-	-	-
Male	-	1	-	-	-	2	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	77	102	133	153	175	214	278	236	190	54	11	-
Male	45	74	75	104	106	121	131	95	69	10	2	-
Female	32	28	58	49	69	93	147	141	121	44	9	-
White, both sexes	56	80	110	129	147	192	265	223	176	50	10	-
Male	36	59	61	87	88	112	126	90	64	10	2	-
Female	20	21	49	42	59	80	139	133	112	40	8	-
Black, both sexes	17	13	15	12	19	15	7	9	6	3	1	-
Male	7	8	9	10	11	7	2	4	2	-	-	-
Female	10	5	6	2	8	8	5	5	4	3	1	-
American Indian 1/, both sexes	1	2	1	4	2	-	1	2	3	1	-	-
Male	-	2	1	2	2	-	1	1	1	-	-	-
Female	1	-	-	2	-	-	-	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes	3	7	7	8	7	7	5	2	5	-	-	-
Male	2	5	4	5	5	2	2	-	2	-	-	-
Female	1	2	3	3	2	5	3	2	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
New Hampshire												
All races, both sexes	14	14	11	19	43	57	101	120	105	52	19	-
Male	8	13	8	9	25	25	45	47	36	10	-	-
Female	6	1	3	10	18	32	56	73	69	42	19	-
White, both sexes	14	14	11	18	43	56	100	120	105	52	19	-
Male	8	13	8	9	25	24	44	47	36	10	-	-
Female	6	1	3	9	18	32	56	73	69	42	19	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	117	124	153	181	322	548	787	954	723	316	57	1
Male	79	81	96	122	173	287	350	363	217	78	12	1
Female	38	43	57	59	149	261	437	591	506	238	45	-
White, both sexes	77	97	108	139	268	479	719	900	673	289	52	1
Male	53	62	69	90	144	251	326	344	207	71	11	1
Female	24	35	39	49	124	228	393	556	466	218	41	-
Black, both sexes	38	26	39	35	47	63	63	47	39	24	4	-
Male	24	18	22	26	24	33	22	17	8	5	-	-
Female	14	8	17	9	23	30	41	30	31	19	4	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	1	-	1	-	-
Male	-	-	-	1	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	6	6	6	6	5	6	11	2	1	-
Male	2	1	5	5	4	3	2	2	2	1	1	-
Female	-	-	1	1	2	3	3	4	9	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
New Mexico												
All races, both sexes	17	23	23	36	61	101	130	158	126	70	13	-
Male	13	11	9	21	34	54	62	72	40	21	4	-
Female	4	12	14	15	27	47	68	86	86	49	9	-
White, both sexes	13	20	17	33	53	96	121	154	120	65	12	-
Male	11	10	8	18	30	52	57	71	38	20	3	-
Female	2	10	9	15	23	44	64	83	82	45	9	-
Black, both sexes	1	1	2	1	2	-	1	1	1	1	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	1	1	2	-	1	-	-	1	1	1	-	-
American Indian 1/, both sexes	3	2	4	2	5	5	8	3	5	4	1	-
Male	2	1	1	2	3	2	4	1	2	1	1	-
Female	1	1	3	-	2	3	4	2	3	3	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
New York												
All races, both sexes	146	220	252	287	513	867	1,240	1,562	1,281	548	134	1
Male	92	140	154	164	280	452	551	609	383	119	20	-
Female	54	80	98	123	233	415	689	953	898	429	114	1
White, both sexes	104	176	190	241	435	782	1,154	1,478	1,221	523	119	1
Male	72	118	118	135	243	425	520	580	372	117	17	-
Female	32	58	72	106	192	357	634	898	849	406	102	1
Black, both sexes	40	40	58	38	74	78	80	66	50	24	12	-
Male	20	20	34	22	35	26	29	21	7	2	2	-
Female	20	20	24	16	39	52	51	45	43	22	10	-
American Indian 1/, both sexes	-	-	1	1	-	-	-	1	-	1	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	4	3	7	4	7	6	17	10	-	3	-
Male	-	2	2	6	2	1	2	7	4	-	1	-
Female	2	2	1	1	2	6	4	10	6	-	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
North Carolina												
All races, both sexes	144	184	209	258	390	535	807	816	600	262	53	-
Male	95	121	125	148	219	261	346	263	167	46	14	-
Female	49	63	84	110	171	274	461	553	433	216	39	-
White, both sexes	80	113	128	186	297	422	673	707	531	223	50	-
Male	60	69	80	109	170	201	301	237	145	40	14	-
Female	20	44	48	77	127	221	372	470	386	183	36	-
Black, both sexes	63	67	76	69	87	105	125	105	67	37	3	-
Male	34	50	45	37	47	56	41	25	21	5	-	-
Female	29	17	31	32	40	49	84	80	46	32	3	-
American Indian 1/, both sexes	-	4	4	1	3	6	8	2	2	1	-	-
Male	-	2	-	1	2	4	3	1	1	-	-	-
Female	-	2	4	-	1	2	5	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	2	3	2	1	2	-	1	-	-
Male	1	-	-	1	-	-	1	-	-	1	-	-
Female	-	-	1	1	3	2	-	2	-	-	-	-
North Dakota												
All races, both sexes	7	5	7	1	15	22	36	69	60	32	6	-
Male	6	5	5	-	6	7	19	22	20	7	1	-
Female	1	-	2	1	9	15	17	47	40	25	5	-
White, both sexes	7	5	7	1	15	22	36	68	59	32	6	-
Male	6	5	5	-	6	7	19	22	19	7	1	-
Female	1	-	2	1	9	15	17	46	40	25	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Ohio												
All races, both sexes	161	215	242	302	457	745	1,080	1,244	903	356	71	-
Male	111	153	136	173	238	366	482	447	261	74	15	-
Female	50	62	106	129	219	379	598	797	642	282	56	-
White, both sexes	126	165	194	236	403	667	1,018	1,173	868	340	59	-
Male	86	116	111	138	212	330	456	416	255	70	11	-
Female	40	49	83	98	191	337	562	757	613	270	48	-
Black, both sexes	34	50	47	63	52	75	62	71	35	16	12	-
Male	24	37	25	33	26	35	26	31	6	4	4	-
Female	10	13	22	30	26	40	36	40	29	12	8	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	3	2	2	-	-	-	-	-	-
Male	1	-	-	2	-	1	-	-	-	-	-	-
Female	-	-	1	1	2	1	-	-	-	-	-	-
Oklahoma												
All races, both sexes	53	65	89	115	168	270	381	382	348	141	36	-
Male	29	39	55	66	77	119	160	126	88	37	10	-
Female	24	26	34	49	91	151	221	256	260	104	26	-
White, both sexes	40	50	75	92	145	248	345	353	334	130	34	-
Male	24	29	46	51	65	115	149	111	85	33	10	-
Female	16	21	29	41	80	133	196	242	249	97	24	-
Black, both sexes	9	11	9	12	9	10	22	15	7	6	2	-
Male	4	7	6	9	5	2	6	7	-	1	-	-
Female	5	4	3	3	4	8	16	8	7	5	2	-
American Indian 1/, both sexes	4	3	5	11	14	12	13	13	7	5	-	-
Male	1	3	3	6	7	2	5	8	3	3	-	-
Female	3	-	2	5	7	10	8	5	4	2	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Oregon												
All races, both sexes	39	50	58	68	106	206	352	465	446	185	45	-
Male	28	33	37	29	55	109	158	187	147	38	7	-
Female	11	17	21	39	51	97	194	278	299	147	38	-
White, both sexes	35	46	53	67	105	204	348	456	436	183	44	-
Male	26	30	34	29	55	108	155	183	145	38	6	-
Female	9	16	19	38	50	96	193	273	291	145	38	-
Black, both sexes	2	2	2	-	1	-	2	4	4	-	-	-
Male	1	1	-	-	-	-	1	1	-	-	-	-
Female	1	1	2	-	1	-	1	3	4	-	-	-
American Indian 1/, both sexes	2	1	2	1	-	1	1	2	1	-	1	-
Male	1	1	2	-	-	1	1	-	1	-	1	-
Female	1	-	-	1	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	1	1	3	5	2	-	-
Male	-	1	1	-	-	-	1	3	1	-	-	-
Female	-	-	-	-	-	1	-	-	4	2	-	-
Pennsylvania												
All races, both sexes	250	300	373	450	704	1,190	1,897	2,167	1,652	682	127	-
Male	180	195	243	263	395	595	810	828	443	139	19	-
Female	70	105	130	187	309	595	1,087	1,339	1,209	543	108	-
White, both sexes	186	228	313	386	636	1,093	1,793	2,073	1,582	653	119	-
Male	136	147	205	231	361	541	767	802	433	138	19	-
Female	50	81	108	155	275	552	1,026	1,271	1,149	515	100	-
Black, both sexes	62	68	59	63	64	93	100	92	67	27	8	-
Male	42	46	38	32	32	51	42	25	10	1	-	-
Female	20	22	21	31	32	42	58	67	57	26	8	-
American Indian 1/, both sexes	-	2	1	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	-	1	4	4	3	2	3	2	-	-
Male	2	1	-	-	2	3	1	1	-	-	-	-
Female	-	1	-	1	2	1	2	1	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Rhode Island												
All races, both sexes	13	12	13	11	20	49	50	81	63	24	8	-
Male	9	8	7	9	11	24	25	28	15	3	2	-
Female	4	4	6	2	9	25	25	53	48	21	6	-
White, both sexes	11	11	11	8	20	47	49	79	63	24	8	-
Male	7	8	6	7	11	22	24	26	15	3	2	-
Female	4	3	5	1	9	25	25	53	48	21	6	-
Black, both sexes	2	1	2	3	-	2	-	2	-	-	-	-
Male	2	-	1	2	-	2	-	2	-	-	-	-
Female	-	1	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	134	161	149	192	268	385	474	510	319	130	28	1
Male	94	107	88	114	169	196	203	174	95	22	4	1
Female	40	54	61	78	99	189	271	336	224	108	24	-
White, both sexes	69	92	90	119	181	290	362	416	266	99	19	1
Male	48	62	55	66	115	146	160	138	83	19	3	1
Female	21	30	35	53	66	144	202	278	183	80	16	-
Black, both sexes	64	69	58	72	85	95	111	91	52	30	8	-
Male	45	45	32	47	52	50	43	34	12	3	1	-
Female	19	24	26	25	33	45	68	57	40	27	7	-
American Indian 1/, both sexes	1	-	1	-	1	-	-	1	1	-	1	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	1	2	-	1	-	-
Male	-	-	-	1	1	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
South Dakota												
All races, both sexes	10	3	9	12	18	22	36	54	46	24	8	-
Male	8	2	6	9	10	13	20	28	9	2	-	-
Female	2	1	3	3	8	9	16	26	37	22	8	-
White, both sexes	7	2	8	9	17	19	33	52	46	24	8	-
Male	5	1	5	8	9	11	19	28	9	2	-	-
Female	2	1	3	1	8	8	14	24	37	22	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	1	3	1	3	3	2	-	-	-	-
Male	3	-	1	1	1	2	1	-	-	-	-	-
Female	-	-	-	2	-	1	2	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	95	155	140	185	228	266	444	438	327	113	35	-
Male	63	104	90	108	111	123	180	161	94	27	4	-
Female	32	51	50	77	117	143	264	277	233	86	31	-
White, both sexes	56	105	106	144	187	227	393	396	296	101	32	-
Male	37	74	69	93	86	103	154	146	86	25	3	-
Female	19	31	37	51	101	124	239	250	210	76	29	-
Black, both sexes	39	50	34	40	40	35	49	41	30	12	3	-
Male	26	30	21	15	24	18	26	15	8	2	1	-
Female	13	20	13	25	16	17	23	26	22	10	2	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	3	2	1	1	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	1	-	2	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Texas												
All races, both sexes	324	342	430	480	725	1,051	1,534	1,594	1,329	567	112	-
Male	225	210	248	272	404	505	655	563	394	139	13	-
Female	99	132	182	208	321	546	879	1,031	935	428	99	-
White, both sexes	219	261	342	391	629	939	1,402	1,485	1,237	528	99	-
Male	158	166	209	226	360	450	597	528	366	129	11	-
Female	61	95	133	165	269	489	805	957	871	399	88	-
Black, both sexes	101	80	79	82	92	103	120	101	84	33	13	-
Male	63	44	35	42	41	52	55	31	26	8	2	-
Female	38	36	44	40	51	51	65	70	58	25	11	-
American Indian 1/, both sexes	1	-	-	1	-	-	1	-	2	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	9	6	4	9	11	8	6	6	-	-
Male	3	-	4	4	3	3	3	4	1	2	-	-
Female	-	1	5	2	1	6	8	4	5	4	-	-
Utah												
All races, both sexes	17	22	29	39	65	131	202	283	207	84	8	-
Male	10	13	18	21	22	64	77	95	71	22	1	-
Female	7	9	11	18	43	67	125	188	136	62	7	-
White, both sexes	16	21	27	39	61	128	200	281	204	81	8	-
Male	9	13	16	21	21	63	77	94	69	22	1	-
Female	7	8	11	18	40	65	123	187	135	59	7	-
Black, both sexes	1	1	1	-	1	1	2	1	-	2	-	-
Male	1	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	1	-	2	1	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	1	-	1	2	-	-	-
Male	-	-	1	-	1	-	-	1	1	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Vermont												
All races, both sexes	3	2	10	5	15	26	40	66	55	26	7	-
Male	1	2	7	5	11	14	14	29	17	7	1	-
Female	2	-	3	-	4	12	26	37	38	19	6	-
White, both sexes	3	2	10	5	15	26	40	66	55	26	7	-
Male	1	2	7	5	11	14	14	29	17	7	1	-
Female	2	-	3	-	4	12	26	37	38	19	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	119	211	206	256	335	538	747	811	637	252	57	-
Male	75	153	122	139	178	265	317	305	186	53	9	-
Female	44	58	84	117	157	273	430	506	451	199	48	-
White, both sexes	73	144	148	177	250	425	624	708	560	224	53	-
Male	46	111	94	95	136	217	272	267	172	48	9	-
Female	27	33	54	82	114	208	352	441	388	176	44	-
Black, both sexes	42	65	53	73	79	107	116	95	72	27	4	-
Male	27	41	26	40	38	44	42	36	12	5	-	-
Female	15	24	27	33	41	63	74	59	60	22	4	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	4	6	6	6	7	7	5	1	-	-
Male	2	1	1	4	4	4	3	2	2	-	-	-
Female	2	1	3	2	2	2	4	5	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Washington												
All races, both sexes	51	56	69	93	141	233	387	485	358	155	46	-
Male	32	35	46	55	79	118	174	214	119	40	5	-
Female	19	21	23	38	62	115	213	271	239	115	41	-
White, both sexes	44	49	64	82	129	222	363	470	347	153	46	-
Male	29	31	44	50	70	111	165	207	117	39	5	-
Female	15	18	20	32	59	111	198	263	230	114	41	-
Black, both sexes	5	4	2	7	5	4	11	6	1	-	-	-
Male	3	2	-	4	3	4	6	3	-	-	-	-
Female	2	2	2	3	2	-	5	3	1	-	-	-
American Indian 1/, both sexes	1	2	1	1	3	1	3	-	2	-	-	-
Male	-	1	1	-	3	1	1	-	-	-	-	-
Female	1	1	-	1	-	-	2	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	3	4	6	10	9	8	2	-	-
Male	-	1	1	1	3	2	2	4	2	1	-	-
Female	1	-	1	2	1	4	8	5	6	1	-	-
West Virginia												
All races, both sexes	23	52	70	82	123	183	272	254	194	61	25	-
Male	13	33	41	47	60	102	109	91	43	12	5	-
Female	10	19	29	35	63	81	163	163	151	49	20	-
White, both sexes	21	52	67	75	120	179	267	242	186	60	25	-
Male	13	33	39	43	59	99	107	88	41	12	5	-
Female	8	19	28	32	61	80	160	154	145	48	20	-
Black, both sexes	2	-	2	7	3	4	5	10	8	1	-	-
Male	-	-	1	4	1	3	2	3	2	-	-	-
Female	2	-	1	3	2	1	3	7	6	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other non-pulmonary forms of heart disease (I30-I51)												
Wisconsin												
All races, both sexes	48	90	99	130	191	363	532	764	658	258	63	-
Male	38	66	58	70	104	187	260	298	208	47	7	-
Female	10	24	41	60	87	176	272	466	450	211	56	-
White, both sexes	41	77	83	123	180	347	519	750	647	258	61	-
Male	34	54	51	68	97	180	256	291	206	47	7	-
Female	7	23	32	55	83	167	263	459	441	211	54	-
Black, both sexes	6	11	12	5	8	14	9	11	6	-	2	-
Male	4	10	5	2	5	7	3	5	1	-	-	-
Female	2	1	7	3	3	7	6	6	5	-	2	-
American Indian 1/, both sexes	1	2	3	2	3	1	1	3	3	-	-	-
Male	-	2	2	-	2	-	-	2	1	-	-	-
Female	1	-	1	2	1	1	1	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	3	-	2	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	2	-	2	-	-	-
Wyoming												
All races, both sexes	8	13	11	21	21	27	46	47	37	18	9	-
Male	3	9	7	12	12	13	22	13	19	5	1	-
Female	5	4	4	9	9	14	24	34	18	13	8	-
White, both sexes	8	13	10	21	21	27	46	47	36	18	9	-
Male	3	9	6	12	12	13	22	13	19	5	1	-
Female	5	4	4	9	9	14	24	34	17	13	8	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
United States												
All races, both sexes	1,209	4	-	1	2	1	7	12	16	36	42	80
Male	695	3	-	-	-	1	4	8	12	24	27	49
Female	514	1	-	1	2	-	3	4	4	12	15	31
White, both sexes	949	3	-	-	1	1	5	9	11	19	27	54
Male	553	2	-	-	-	1	3	6	8	13	18	32
Female	396	1	-	-	1	-	2	3	3	6	9	22
Black, both sexes	229	1	-	1	1	-	2	2	4	14	14	22
Male	124	1	-	-	-	-	1	2	3	10	9	13
Female	105	-	-	1	1	-	1	-	1	4	5	9
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	1	1	4
Male	6	-	-	-	-	-	-	-	-	-	-	4
Female	6	-	-	-	-	-	-	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	1	1	2	-	-
Male	12	-	-	-	-	-	-	-	1	1	-	-
Female	7	-	-	-	-	-	-	1	-	1	-	-
Alabama												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	2	1
Male	14	-	-	-	-	-	-	-	-	-	2	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	2	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	24	-	-	-	-	-	-	1	1	1	1	3
Male	18	-	-	-	-	-	-	1	1	1	1	2
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	1	1	1	1	2
Male	17	-	-	-	-	-	-	1	1	1	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Arkansas												
All races, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	120	-	-	-	-	1	-	5	2	7	4	8
Male	79	-	-	-	-	1	-	4	2	6	3	5
Female	41	-	-	-	-	-	-	1	-	1	1	3
White, both sexes	103	-	-	-	-	1	-	4	1	4	4	7
Male	69	-	-	-	-	1	-	4	1	4	3	5
Female	34	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	10	-	-	-	-	-	-	-	-	2	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	1	1	1	-	-
Male	4	-	-	-	-	-	-	-	1	1	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Colorado												
All races, both sexes	11	-	-	-	-	-	1	-	1	1	-	-
Male	6	-	-	-	-	-	1	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	11	-	-	-	-	-	1	-	1	1	-	-
Male	6	-	-	-	-	-	1	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	18	-	-	-	-	-	-	-	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Delaware												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Florida												
All races, both sexes	107	-	-	-	-	-	-	1	1	2	5	9
Male	65	-	-	-	-	-	-	-	1	1	5	6
Female	42	-	-	-	-	-	-	1	-	1	-	3
White, both sexes	85	-	-	-	-	-	-	1	1	-	2	6
Male	51	-	-	-	-	-	-	-	1	-	2	4
Female	34	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	22	-	-	-	-	-	-	-	-	2	3	3
Male	14	-	-	-	-	-	-	-	-	1	3	2
Female	8	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	34	-	-	-	-	-	-	-	1	-	2	2
Male	17	-	-	-	-	-	-	-	1	-	2	1
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	22	-	-	-	-	-	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	12	-	-	-	-	-	-	-	-	-	2	1
Male	7	-	-	-	-	-	-	-	-	-	2	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Hawaii												
All races, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Idaho												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Illinois												
All races, both sexes	42	-	-	-	-	-	-	1	-	1	-	4
Male	18	-	-	-	-	-	-	1	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	28	-	-	-	-	-	-	1	-	-	-	2
Male	13	-	-	-	-	-	-	1	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	1	-	2
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	21	1	-	-	-	-	-	-	-	-	-	1
Male	13	1	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	1	-	-	-	-	-	-	-	-	-	-
Male	11	1	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Iowa												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	9	-	-	-	-	-	-	1	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	1	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Kentucky												
All races, both sexes	18	1	-	-	-	-	1	-	-	-	-	1
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	16	1	-	-	-	-	1	-	-	-	-	1
Male	7	1	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	18	1	-	-	-	-	-	1	-	-	-	1
Male	11	1	-	-	-	-	-	1	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	1	-	-	-	-	-	1	-	-	-	1
Male	6	1	-	-	-	-	-	1	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Maine												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	2	1
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	12	-	-	-	-	-	-	-	-	-	2	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Massachusetts												
All races, both sexes	17	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	1	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	1	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	50	-	-	-	-	-	1	-	1	-	2	6
Male	32	-	-	-	-	-	1	-	1	-	-	4
Female	18	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	35	-	-	-	-	-	-	-	1	-	-	4
Male	24	-	-	-	-	-	-	-	1	-	-	2
Female	11	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	14	-	-	-	-	-	1	-	-	-	2	1
Male	7	-	-	-	-	-	1	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Minnesota												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	18	-	-	-	-	-	-	-	2	1	1	3
Male	14	-	-	-	-	-	-	-	2	1	1	3
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	2	1	1	1
Male	8	-	-	-	-	-	-	-	2	1	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Missouri												
All races, both sexes	21	-	-	-	-	-	-	-	-	3	-	-
Male	12	-	-	-	-	-	-	-	-	3	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	2	-	-
Male	10	-	-	-	-	-	-	-	-	2	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Nebraska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	-	-	-	-	-	-	1	-	-	-	-
Male	7	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
New Hampshire												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	29	-	-	-	-	-	-	-	-	3	1	3
Male	17	-	-	-	-	-	-	-	-	3	-	1
Female	12	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	19	-	-	-	-	-	-	-	-	1	-	1
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	2	1	2
Male	4	-	-	-	-	-	-	-	-	2	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
New Mexico												
All races, both sexes	6	-	-	-	-	-	-	-	-	1	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	110	-	-	-	1	-	1	-	-	2	2	2
Male	59	-	-	-	-	-	-	-	-	-	1	-
Female	51	-	-	-	1	-	1	-	-	2	1	2
White, both sexes	93	-	-	-	1	-	1	-	-	1	1	2
Male	53	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	1	-	1	-	-	1	1	2
Black, both sexes	16	-	-	-	-	-	-	-	-	1	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
North Carolina												
All races, both sexes	27	-	-	-	-	-	-	1	-	1	1	2
Male	14	-	-	-	-	-	-	-	-	1	-	2
Female	13	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	18	-	-	-	-	-	-	1	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	1	1	1
Male	4	-	-	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	1	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	1	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Ohio												
All races, both sexes	54	-	-	-	-	-	-	-	1	-	2	5
Male	29	-	-	-	-	-	-	-	-	-	1	4
Female	25	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	41	-	-	-	-	-	-	-	1	-	2	4
Male	19	-	-	-	-	-	-	-	-	-	1	3
Female	22	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	18	-	-	-	-	-	-	-	1	1	1	2
Male	12	-	-	-	-	-	-	-	1	-	1	2
Female	6	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	15	-	-	-	-	-	-	-	1	1	1	1
Male	11	-	-	-	-	-	-	-	1	-	1	1
Female	4	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Oregon												
All races, both sexes	13	-	-	-	-	-	-	-	-	1	-	1
Male	11	-	-	-	-	-	-	-	-	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	1	-	1
Male	9	-	-	-	-	-	-	-	-	1	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	63	-	-	-	-	-	2	-	-	2	-	5
Male	33	-	-	-	-	-	1	-	-	1	-	2
Female	30	-	-	-	-	-	1	-	-	1	-	3
White, both sexes	56	-	-	-	-	-	1	-	-	2	-	4
Male	32	-	-	-	-	-	1	-	-	1	-	2
Female	24	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	7	-	-	-	-	-	1	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Rhode Island												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	16	-	-	-	-	-	-	-	1	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
South Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	3
Male	22	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	2
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Texas												
All races, both sexes	79	1	-	-	1	-	1	-	1	2	6	4
Male	38	-	-	-	-	-	1	-	-	2	2	2
Female	41	1	-	-	1	-	-	-	1	-	4	2
White, both sexes	62	1	-	-	-	-	1	-	1	-	5	3
Male	30	-	-	-	-	-	1	-	-	-	2	2
Female	32	1	-	-	-	-	-	-	1	-	3	1
Black, both sexes	17	-	-	-	1	-	-	-	-	2	1	1
Male	8	-	-	-	-	-	-	-	-	2	-	-
Female	9	-	-	-	1	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Vermont												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	30	-	-	1	-	-	-	-	-	-	-	4
Male	15	-	-	-	-	-	-	-	-	-	-	3
Female	15	-	-	1	-	-	-	-	-	-	-	1
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	1	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	1	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Washington												
All races, both sexes	30	-	-	-	-	-	-	-	-	-	2	2
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	2
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Acute and subacute endocarditis (I33)												
Wisconsin												
All races, both sexes	17	-	-	-	-	-	-	-	-	1	1	1
Male	11	-	-	-	-	-	-	-	-	1	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	1	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
United States												
All races, both sexes	93	102	96	112	122	177	151	104	44	6	1	-
Male	64	67	58	56	68	97	76	53	24	3	1	-
Female	29	35	38	56	54	80	75	51	20	3	-	-
White, both sexes	58	62	73	88	109	154	132	94	42	6	1	-
Male	42	42	42	48	60	90	69	49	24	3	1	-
Female	16	20	31	40	49	64	63	45	18	3	-	-
Black, both sexes	33	37	21	22	9	20	16	8	2	-	-	-
Male	21	23	14	8	5	6	5	3	-	-	-	-
Female	12	14	7	14	4	14	11	5	2	-	-	-
American Indian 1/, both sexes	1	2	-	2	1	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	1	1	-	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	-	3	3	3	2	-	-	-	-
Male	1	1	2	-	2	1	2	1	-	-	-	-
Female	-	-	-	-	1	2	1	1	-	-	-	-
Alabama												
All races, both sexes	2	2	2	4	-	8	2	-	1	-	-	-
Male	2	2	2	1	-	3	1	-	-	-	-	-
Female	-	-	-	3	-	5	1	-	1	-	-	-
White, both sexes	-	1	1	2	-	6	2	-	1	-	-	-
Male	-	1	1	-	-	3	1	-	-	-	-	-
Female	-	-	-	2	-	3	1	-	1	-	-	-
Black, both sexes	2	1	1	2	-	2	-	-	-	-	-	-
Male	2	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	2	1	2	1	3	4	1	1	-	-	-
Male	2	1	1	1	-	2	3	1	1	-	-	-
Female	-	1	-	1	1	1	1	-	-	-	-	-
White, both sexes	2	1	1	2	1	3	4	1	1	-	-	-
Male	2	1	1	1	-	2	3	1	1	-	-	-
Female	-	-	-	1	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Arkansas												
All races, both sexes	1	1	1	1	1	-	-	-	-	-	-	-
Male	1	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
White, both sexes	1	1	1	1	1	-	-	-	-	-	-	-
Male	1	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7	10	7	14	10	20	7	11	6	1	-	-
Male	4	7	4	6	8	13	6	5	4	1	-	-
Female	3	3	3	8	2	7	1	6	2	-	-	-
White, both sexes	6	8	7	12	9	18	5	10	6	1	-	-
Male	4	5	4	5	7	11	5	5	4	1	-	-
Female	2	3	3	7	2	7	-	5	2	-	-	-
Black, both sexes	1	2	-	1	1	1	-	1	-	-	-	-
Male	-	2	-	1	1	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Colorado												
All races, both sexes	-	1	1	-	2	1	2	1	-	-	-	-
Male	-	1	-	-	1	-	1	1	-	-	-	-
Female	-	-	1	-	1	1	1	-	-	-	-	-
White, both sexes	-	1	1	-	2	1	2	1	-	-	-	-
Male	-	1	-	-	1	-	1	1	-	-	-	-
Female	-	-	1	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	1	2	1	2	4	3	1	2	-	-	-
Male	1	-	-	-	1	3	2	1	2	-	-	-
Female	-	1	2	1	1	1	1	-	-	-	-	-
White, both sexes	-	-	2	1	2	4	2	1	2	-	-	-
Male	-	-	-	-	1	3	1	1	2	-	-	-
Female	-	-	2	1	1	1	1	-	-	-	-	-
Black, both sexes	1	1	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Delaware												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	1	1	-	-	1	1	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	1	-	-	1	1	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Florida												
All races, both sexes	10	7	12	10	9	15	12	9	4	1	-	-
Male	8	5	7	6	5	8	6	4	3	-	-	-
Female	2	2	5	4	4	7	6	5	1	1	-	-
White, both sexes	6	4	10	7	9	14	11	9	4	1	-	-
Male	5	4	5	4	5	8	6	4	3	-	-	-
Female	1	-	5	3	4	6	5	5	1	1	-	-
Black, both sexes	4	3	2	3	-	1	1	-	-	-	-	-
Male	3	1	2	2	-	-	-	-	-	-	-	-
Female	1	2	-	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	5	3	3	2	2	3	5	4	2	-	-	-
Male	2	2	2	2	1	3	-	1	-	-	-	-
Female	3	1	1	-	1	-	5	3	2	-	-	-
White, both sexes	-	1	3	2	1	3	5	3	2	-	-	-
Male	-	1	2	2	-	3	-	1	-	-	-	-
Female	-	-	1	-	1	-	5	2	2	-	-	-
Black, both sexes	5	2	-	-	1	-	-	1	-	-	-	-
Male	2	1	-	-	1	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Hawaii												
All races, both sexes	1	2	-	-	2	-	1	-	-	-	-	-
Male	1	2	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	2	-	1	-	-	-	-	-
Male	1	1	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Illinois												
All races, both sexes	3	3	4	2	4	9	5	5	1	-	-	-
Male	1	2	2	-	3	1	2	4	1	-	-	-
Female	2	1	2	2	1	8	3	1	-	-	-	-
White, both sexes	2	1	2	2	4	5	5	3	1	-	-	-
Male	-	-	1	-	3	1	2	3	1	-	-	-
Female	2	1	1	2	1	4	3	-	-	-	-	-
Black, both sexes	1	2	2	-	-	4	-	2	-	-	-	-
Male	1	2	1	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	4	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	1	2	2	3	5	3	2	-	-	-	-
Male	1	1	1	2	2	3	-	1	-	-	-	-
Female	-	-	1	-	1	2	3	1	-	-	-	-
White, both sexes	1	-	2	2	3	5	3	2	-	-	-	-
Male	1	-	1	2	2	3	-	1	-	-	-	-
Female	-	-	1	-	1	2	3	1	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Iowa												
All races, both sexes	-	-	-	1	2	2	2	1	-	-	-	-
Male	-	-	-	1	-	1	2	1	-	-	-	-
Female	-	-	-	-	2	1	-	-	-	-	-	-
White, both sexes	-	-	-	1	2	2	2	1	-	-	-	-
Male	-	-	-	1	-	1	2	1	-	-	-	-
Female	-	-	-	-	2	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	1	1	1	1	1	-	1	-	-	-	-
Male	1	-	-	1	-	1	-	1	-	-	-	-
Female	-	1	1	-	1	-	-	-	-	-	-	-
White, both sexes	1	1	1	1	1	1	-	1	-	-	-	-
Male	1	-	-	1	-	1	-	1	-	-	-	-
Female	-	1	1	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Kentucky												
All races, both sexes	2	1	1	3	1	4	2	-	1	-	-	-
Male	2	-	1	1	-	2	1	-	1	-	-	-
Female	-	1	-	2	1	2	1	-	-	-	-	-
White, both sexes	1	1	1	3	1	3	2	-	1	-	-	-
Male	1	-	1	1	-	1	1	-	1	-	-	-
Female	-	1	-	2	1	2	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	2	1	2	4	3	1	1	-	-	-	-
Male	1	1	1	1	1	1	1	1	-	-	-	-
Female	-	1	-	1	3	2	-	-	-	-	-	-
White, both sexes	1	-	1	-	3	2	1	1	-	-	-	-
Male	1	-	1	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	3	1	-	-	-	-	-	-
Black, both sexes	-	2	-	2	1	1	-	-	-	-	-	-
Male	-	1	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Maine												
All races, both sexes	-	2	1	-	1	-	-	1	-	-	1	-
Male	-	1	-	-	1	-	-	1	-	-	1	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	2	1	-	1	-	-	1	-	-	1	-
Male	-	1	-	-	1	-	-	1	-	-	1	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	2	-	1	-	4	5	2	-	-	-	-
Male	3	1	-	-	-	3	2	1	-	-	-	-
Female	3	1	-	1	-	1	3	1	-	-	-	-
White, both sexes	2	-	-	-	-	2	4	2	-	-	-	-
Male	1	-	-	-	-	2	2	1	-	-	-	-
Female	1	-	-	-	-	-	2	1	-	-	-	-
Black, both sexes	4	2	-	1	-	2	1	-	-	-	-	-
Male	2	1	-	-	-	1	-	-	-	-	-	-
Female	2	1	-	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Massachusetts												
All races, both sexes	-	-	1	1	-	2	3	7	1	-	-	-
Male	-	-	1	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	2	6	1	-	-	-
White, both sexes	-	-	1	1	-	2	3	7	1	-	-	-
Male	-	-	1	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	2	6	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	7	1	6	4	2	6	8	5	1	-	-	-
Male	4	-	5	3	2	5	3	3	1	-	-	-
Female	3	1	1	1	-	1	5	2	-	-	-	-
White, both sexes	4	-	5	2	2	4	7	5	1	-	-	-
Male	3	-	4	2	2	4	2	3	1	-	-	-
Female	1	-	1	-	-	-	5	2	-	-	-	-
Black, both sexes	3	1	1	2	-	2	1	-	-	-	-	-
Male	1	-	1	1	-	1	1	-	-	-	-	-
Female	2	1	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Minnesota												
All races, both sexes	-	2	-	-	2	1	1	1	2	-	-	-
Male	-	1	-	-	2	1	-	-	1	-	-	-
Female	-	1	-	-	-	-	1	1	1	-	-	-
White, both sexes	-	2	-	-	2	1	1	-	2	-	-	-
Male	-	1	-	-	2	1	-	-	1	-	-	-
Female	-	1	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Mississippi												
All races, both sexes	2	1	1	-	2	3	2	-	-	-	-	-
Male	2	1	1	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	3	1	-	-	-	-	-
White, both sexes	1	-	-	-	2	2	1	-	-	-	-	-
Male	1	-	-	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Black, both sexes	1	1	1	-	-	1	1	-	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Missouri												
All races, both sexes	2	1	1	5	1	4	3	1	-	-	-	-
Male	1	1	-	1	1	3	1	1	-	-	-	-
Female	1	-	1	4	-	1	2	-	-	-	-	-
White, both sexes	2	-	1	5	1	4	2	1	-	-	-	-
Male	1	-	-	1	1	3	1	1	-	-	-	-
Female	1	-	1	4	-	1	1	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	1	-	-	-	1	-	1	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	1	1	-	-	-	1	-	1	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Nebraska												
All races, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	2	1	1	1	2	-	-	-	-	-
Male	-	1	2	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	1	1	1	1	1	2	-	-	-	-	-
Male	-	1	1	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
New Hampshire												
All races, both sexes	1	1	-	1	2	4	-	-	-	-	-	-
Male	-	1	-	1	2	2	-	-	-	-	-	-
Female	1	-	-	-	-	2	-	-	-	-	-	-
White, both sexes	1	1	-	1	2	4	-	-	-	-	-	-
Male	-	1	-	1	2	2	-	-	-	-	-	-
Female	1	-	-	-	-	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	2	2	1	5	2	5	3	1	-	-	-
Male	-	1	2	-	3	1	4	1	1	-	-	-
Female	1	1	-	1	2	1	1	2	-	-	-	-
White, both sexes	-	1	-	1	4	2	5	3	1	-	-	-
Male	-	1	-	-	3	1	4	1	1	-	-	-
Female	-	-	-	1	1	1	1	2	-	-	-	-
Black, both sexes	1	1	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
New Mexico												
All races, both sexes	-	1	-	-	1	-	-	-	1	-	-	-
Male	-	1	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	8	12	11	17	12	18	14	5	2	-	-
Male	3	6	9	5	8	8	9	7	2	1	-	-
Female	-	2	3	6	9	4	9	7	3	1	-	-
White, both sexes	3	6	8	8	15	12	16	12	5	2	-	-
Male	3	5	7	4	8	8	9	6	2	1	-	-
Female	-	1	1	4	7	4	7	6	3	1	-	-
Black, both sexes	-	2	4	3	2	-	2	1	-	-	-	-
Male	-	1	2	1	-	-	-	-	-	-	-	-
Female	-	1	2	2	2	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
North Carolina												
All races, both sexes	4	1	1	4	3	4	3	-	1	1	-	-
Male	2	-	1	1	2	3	1	-	-	1	-	-
Female	2	1	-	3	1	1	2	-	1	-	-	-
White, both sexes	3	-	-	2	3	4	2	-	1	1	-	-
Male	1	-	-	1	2	3	1	-	-	1	-	-
Female	2	-	-	1	1	1	1	-	1	-	-	-
Black, both sexes	1	1	1	2	-	-	1	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	2	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Ohio												
All races, both sexes	2	8	6	4	3	11	3	8	-	1	-	-
Male	2	7	4	1	-	5	1	4	-	-	-	-
Female	-	1	2	3	3	6	2	4	-	1	-	-
White, both sexes	1	3	4	3	2	10	3	7	-	1	-	-
Male	1	2	2	1	-	5	1	3	-	-	-	-
Female	-	1	2	2	2	5	2	4	-	1	-	-
Black, both sexes	1	5	2	1	-	1	-	1	-	-	-	-
Male	1	5	2	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	-	2	4	-	2	1	1	-	-	-	-
Male	2	-	-	2	-	2	1	1	-	-	-	-
Female	1	-	2	2	-	-	-	-	-	-	-	-
White, both sexes	2	-	2	3	-	2	1	1	-	-	-	-
Male	2	-	-	2	-	2	1	1	-	-	-	-
Female	-	-	2	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Oregon												
All races, both sexes	2	2	-	-	-	4	-	2	1	-	-	-
Male	2	2	-	-	-	3	-	2	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
White, both sexes	1	1	-	-	-	3	-	2	1	-	-	-
Male	1	1	-	-	-	3	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	-	5	6	5	9	10	11	6	2	-	-	-
Male	-	3	2	4	4	6	6	3	1	-	-	-
Female	-	2	4	1	5	4	5	3	1	-	-	-
White, both sexes	-	4	4	4	9	10	10	6	2	-	-	-
Male	-	3	2	3	4	6	6	3	1	-	-	-
Female	-	1	2	1	5	4	4	3	1	-	-	-
Black, both sexes	-	1	2	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	2	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Rhode Island												
All races, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	1	2	6	-	3	1	1	-	-	-	-
Male	-	1	2	3	-	-	-	-	-	-	-	-
Female	-	-	-	3	-	3	1	1	-	-	-	-
White, both sexes	-	1	2	6	-	3	1	1	-	-	-	-
Male	-	1	2	3	-	-	-	-	-	-	-	-
Female	-	-	-	3	-	3	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	2	1	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	1	-	-	-	-	-	-	2	1	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	4	2	8	8	3	5	2	1	-	-	-
Male	2	3	2	4	3	2	2	2	1	-	-	-
Female	-	1	-	4	5	1	3	-	-	-	-	-
White, both sexes	2	2	1	6	8	2	4	2	1	-	-	-
Male	2	2	1	4	3	1	1	2	1	-	-	-
Female	-	-	-	2	5	1	3	-	-	-	-	-
Black, both sexes	-	2	1	2	-	1	1	-	-	-	-	-
Male	-	1	1	-	-	1	1	-	-	-	-	-
Female	-	1	-	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Texas												
All races, both sexes	10	9	7	5	9	7	9	4	3	-	-	-
Male	7	4	3	3	5	3	4	-	2	-	-	-
Female	3	5	4	2	4	4	5	4	1	-	-	-
White, both sexes	5	8	6	4	7	6	8	4	3	-	-	-
Male	4	4	3	3	4	2	3	-	2	-	-	-
Female	1	4	3	1	3	4	5	4	1	-	-	-
Black, both sexes	5	1	1	1	2	1	1	-	-	-	-	-
Male	3	-	-	-	1	1	1	-	-	-	-	-
Female	2	1	1	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Vermont												
All races, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	5	1	1	3	3	6	2	3	-	-	-
Male	1	2	-	-	3	-	4	2	-	-	-	-
Female	-	3	1	1	-	3	2	-	3	-	-	-
White, both sexes	1	3	1	-	2	2	3	2	1	-	-	-
Male	1	1	-	-	2	-	3	2	-	-	-	-
Female	-	2	1	-	-	2	-	-	1	-	-	-
Black, both sexes	-	2	-	1	1	1	3	-	2	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	1	-	1	-	1	2	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Washington												
All races, both sexes	1	3	2	3	6	2	7	1	1	-	-	-
Male	-	2	-	2	2	1	5	1	1	-	-	-
Female	1	1	2	1	4	1	2	-	-	-	-	-
White, both sexes	1	2	2	3	6	1	7	1	1	-	-	-
Male	-	1	-	2	2	1	5	1	1	-	-	-
Female	1	1	2	1	4	-	2	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	-	1	2	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	1	1	1	-	-	-	-	-
White, both sexes	1	-	-	-	1	2	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Acute and subacute endocarditis (I33)												
Wisconsin												
All races, both sexes	2	2	1	-	-	2	5	1	1	-	-	-
Male	2	1	-	-	-	1	3	-	1	-	-	-
Female	-	1	1	-	-	1	2	1	-	-	-	-
White, both sexes	2	1	-	-	-	2	5	-	1	-	-	-
Male	2	-	-	-	-	1	3	-	1	-	-	-
Female	-	1	-	-	-	1	2	-	-	-	-	-
Black, both sexes	-	1	1	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
United States												
All races, both sexes	864	21	17	15	15	18	16	28	25	34	50	69
Male	439	9	10	4	5	10	11	17	14	14	32	40
Female	425	12	7	11	10	8	5	11	11	20	18	29
White, both sexes	661	8	4	6	9	13	13	19	13	23	38	50
Male	344	3	4	2	4	8	10	10	8	12	21	32
Female	317	5	-	4	5	5	3	9	5	11	17	18
Black, both sexes	164	11	11	6	3	4	1	6	9	11	6	18
Male	73	5	5	1	-	2	1	5	5	2	5	8
Female	91	6	6	5	3	2	-	1	4	9	1	10
American Indian 1/, both sexes	3	-	-	-	-	-	-	1	1	-	-	-
Male	3	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	36	2	2	3	3	1	2	2	2	-	6	1
Male	19	1	1	1	1	-	-	1	-	-	6	-
Female	17	1	1	2	2	1	2	1	2	-	-	1
Alabama												
All races, both sexes	19	1	-	-	-	-	1	-	1	-	-	1
Male	9	-	-	-	-	-	1	-	-	-	-	1
Female	10	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	12	-	-	-	-	-	1	-	-	-	-	1
Male	7	-	-	-	-	-	1	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	1	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	1	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	16	-	-	-	-	1	-	1	1	1	1	-
Male	5	-	-	-	-	-	-	1	-	-	1	-
Female	11	-	-	-	-	1	-	-	1	1	-	-
White, both sexes	14	-	-	-	-	1	-	-	1	1	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	11	-	-	-	-	1	-	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Arkansas												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	73	1	1	2	3	2	1	3	2	4	7	7
Male	41	-	1	1	1	1	-	3	1	3	5	5
Female	32	1	-	1	2	1	1	-	1	1	2	2
White, both sexes	57	1	1	1	1	1	1	2	1	4	4	6
Male	30	-	1	-	-	1	-	2	-	3	3	4
Female	27	1	-	1	1	-	1	-	1	1	1	2
Black, both sexes	6	-	-	-	-	-	-	-	1	-	1	1
Male	4	-	-	-	-	-	-	-	1	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	1	2	1	-	1	-	-	2	-
Male	7	-	-	1	1	-	-	1	-	-	2	-
Female	3	-	-	-	1	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Colorado												
All races, both sexes	9	1	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	3	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	1	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	3	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	16	-	-	-	-	-	-	-	-	2	1	-
Male	10	-	-	-	-	-	-	-	-	1	1	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	2	1	-
Male	9	-	-	-	-	-	-	-	-	1	1	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Delaware												
All races, both sexes	3	-	1	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Florida												
All races, both sexes	52	2	2	1	-	3	-	1	-	3	-	7
Male	25	1	1	-	-	2	-	-	-	1	-	5
Female	27	1	1	1	-	1	-	1	-	2	-	2
White, both sexes	42	1	1	1	-	3	-	-	-	2	-	6
Male	24	1	1	-	-	2	-	-	-	1	-	5
Female	18	-	-	1	-	1	-	-	-	1	-	1
Black, both sexes	9	1	-	-	-	-	-	1	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	1	-	-	-	-	-	1	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	32	-	1	-	-	-	-	4	-	-	4	3
Male	14	-	-	-	-	-	-	3	-	-	3	1
Female	18	-	1	-	-	-	-	1	-	-	1	2
White, both sexes	19	-	-	-	-	-	-	3	-	-	2	-
Male	9	-	-	-	-	-	-	2	-	-	1	-
Female	10	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	12	-	1	-	-	-	-	1	-	-	1	3
Male	4	-	-	-	-	-	-	1	-	-	1	1
Female	8	-	1	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Hawaii												
All races, both sexes	8	1	-	1	-	-	-	1	-	-	1	-
Male	4	1	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	1	-	-	-	1	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	1	-	1	-	-	-	1	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	1	-	-	-	1	-	-	-	-
Idaho												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Illinois												
All races, both sexes	35	-	-	-	-	-	-	1	2	2	3	2
Male	16	-	-	-	-	-	-	1	1	1	1	1
Female	19	-	-	-	-	-	-	-	1	1	2	1
White, both sexes	26	-	-	-	-	-	-	-	1	1	2	-
Male	10	-	-	-	-	-	-	-	1	1	-	-
Female	16	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	8	-	-	-	-	-	-	1	1	1	-	2
Male	5	-	-	-	-	-	-	1	-	-	-	1
Female	3	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	25	1	-	-	1	1	1	1	1	1	1	2
Male	14	-	-	-	-	-	1	-	1	-	-	2
Female	11	1	-	-	1	1	-	1	-	1	1	-
White, both sexes	22	-	-	-	1	-	1	1	1	1	1	1
Male	13	-	-	-	-	-	1	-	1	-	-	1
Female	9	-	-	-	1	-	-	1	-	1	1	-
Black, both sexes	3	1	-	-	-	1	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Iowa												
All races, both sexes	11	-	1	-	-	-	1	-	-	1	1	2
Male	7	-	1	-	-	-	1	-	-	-	1	2
Female	4	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	9	-	1	-	-	-	1	-	-	1	-	2
Male	6	-	1	-	-	-	1	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	11	-	-	1	-	-	1	-	-	-	-	-
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	5	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	1	-	-	1	-	-	-	-	-
Male	6	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Kentucky												
All races, both sexes	19	-	-	-	1	-	-	-	-	-	2	-
Male	6	-	-	-	-	-	-	-	-	-	2	-
Female	13	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	1	-	-	-	-	-	2	-
Male	6	-	-	-	-	-	-	-	-	-	2	-
Female	12	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	13	-	-	1	-	1	-	-	-	-	-	3
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	1	-	1	-	-	-	-	-	3
White, both sexes	9	-	-	-	-	1	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	1	-	-	-	-	-	2
Black, both sexes	4	-	-	1	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	1	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Maine												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	16	-	2	-	-	-	1	-	-	-	-	1
Male	7	-	1	-	-	-	1	-	-	-	-	-
Female	9	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	-	1	-	-	-	-	-
Male	3	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	1	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	1	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Massachusetts												
All races, both sexes	13	-	1	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	1	-	-	-	-	-	-	1	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	1	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	33	-	1	2	2	1	-	-	-	2	2	-
Male	19	-	-	-	1	1	-	-	-	-	1	-
Female	14	-	1	2	1	-	-	-	-	2	1	-
White, both sexes	25	-	-	1	1	1	-	-	-	1	2	-
Male	16	-	-	-	1	1	-	-	-	-	1	-
Female	9	-	-	1	-	-	-	-	-	1	1	-
Black, both sexes	8	-	1	1	1	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	1	1	1	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Minnesota												
All races, both sexes	7	-	-	-	1	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	1	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	6	-	-	-	-	-	-	1	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	1	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Missouri												
All races, both sexes	20	1	-	-	-	-	1	1	1	1	2	4
Male	12	1	-	-	-	-	1	1	-	1	2	1
Female	8	-	-	-	-	-	-	-	1	-	-	3
White, both sexes	16	1	-	-	-	-	1	-	1	1	2	2
Male	9	1	-	-	-	-	1	-	-	1	2	-
Female	7	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	2
Male	2	-	-	-	-	-	-	1	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	1	-	-	1	-	-	-	1	-	-	-
Male	3	-	-	-	1	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	6	1	-	-	1	-	-	-	1	-	-	-
Male	3	-	-	-	1	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Nebraska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	-	-	-	-	-	2	-	-	-	1	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	8	-	-	-	-	-	2	-	-	-	1	-
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
New Hampshire												
All races, both sexes	4	-	-	-	-	-	-	1	-	-	1	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	4	-	-	-	-	-	-	1	-	-	1	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	29	-	-	-	-	-	-	2	2	2	-	4
Male	14	-	-	-	-	-	-	1	2	1	-	2
Female	15	-	-	-	-	-	-	1	-	1	-	2
White, both sexes	23	-	-	-	-	-	-	2	1	2	-	2
Male	11	-	-	-	-	-	-	1	1	1	-	1
Female	12	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	1
Male	2	-	-	-	-	-	-	-	1	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
New Mexico												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	58	2	2	2	1	1	1	1	4	2	2	3
Male	30	1	2	1	-	1	1	-	3	2	1	2
Female	28	1	-	1	1	-	-	1	1	-	1	1
White, both sexes	40	-	1	-	-	1	-	1	2	-	2	3
Male	20	-	1	-	-	1	-	-	2	-	1	2
Female	20	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	14	2	1	1	-	-	1	-	1	2	-	-
Male	9	1	1	1	-	-	1	-	1	2	-	-
Female	5	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	1	1	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	1	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
North Carolina												
All races, both sexes	28	2	2	1	-	-	1	-	1	-	-	2
Male	15	1	2	1	-	-	1	-	-	-	-	2
Female	13	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	17	1	-	1	-	-	1	-	-	-	-	2
Male	9	-	-	1	-	-	1	-	-	-	-	2
Female	8	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	1	2	-	-	-	-	-	1	-	-	-
Male	6	1	2	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Ohio												
All races, both sexes	26	3	1	1	-	-	-	1	1	1	4	3
Male	13	1	1	-	-	-	-	-	1	1	3	-
Female	13	2	-	1	-	-	-	1	-	-	1	3
White, both sexes	18	-	-	-	-	-	-	1	-	1	3	3
Male	9	-	-	-	-	-	-	-	-	1	2	-
Female	9	-	-	-	-	-	-	1	-	-	1	3
Black, both sexes	8	3	1	1	-	-	-	-	1	-	1	-
Male	4	1	1	-	-	-	-	-	1	-	1	-
Female	4	2	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	10	-	-	-	-	1	1	-	-	-	-	2
Male	5	-	-	-	-	-	1	-	-	-	-	1
Female	5	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	1	1	-	-	-	-	2
Male	5	-	-	-	-	-	1	-	-	-	-	1
Female	5	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Oregon												
All races, both sexes	7	-	-	-	1	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	47	-	-	1	2	-	1	1	-	4	3	1
Male	26	-	-	1	2	-	1	1	-	-	2	-
Female	21	-	-	-	-	-	-	-	-	4	1	1
White, both sexes	38	-	-	1	2	-	1	1	-	1	2	1
Male	23	-	-	1	2	-	1	1	-	-	1	-
Female	15	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	3	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Rhode Island												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	16	1	1	-	-	1	-	2	1	-	-	-
Male	10	-	1	-	-	-	-	1	1	-	-	-
Female	6	1	-	-	-	1	-	1	-	-	-	-
White, both sexes	11	1	-	-	-	-	-	2	1	-	-	-
Male	7	-	-	-	-	-	-	1	1	-	-	-
Female	4	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	5	-	1	-	-	1	-	-	-	-	-	-
Male	3	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
South Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	29	1	-	1	-	1	1	1	1	1	1	4
Male	17	1	-	-	-	-	-	1	1	1	1	3
Female	12	-	-	1	-	1	1	-	-	-	-	1
White, both sexes	21	-	-	-	-	1	1	1	1	1	1	2
Male	12	-	-	-	-	-	-	1	1	1	1	1
Female	9	-	-	-	-	1	1	-	-	-	-	1
Black, both sexes	8	1	-	1	-	-	-	-	-	-	-	2
Male	5	1	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Texas												
All races, both sexes	63	-	1	-	-	5	-	2	4	2	2	3
Male	35	-	-	-	-	5	-	1	2	1	1	2
Female	28	-	1	-	-	-	-	1	2	1	1	1
White, both sexes	50	-	-	-	-	3	-	2	2	1	2	3
Male	28	-	-	-	-	3	-	1	1	1	1	2
Female	22	-	-	-	-	-	-	1	1	-	1	1
Black, both sexes	10	-	1	-	-	2	-	-	1	1	-	-
Male	5	-	-	-	-	2	-	-	-	-	-	-
Female	5	-	1	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	3	-	-	-	-	-	-	1	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	1	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	22	2	-	-	1	-	-	2	1	-	1	2
Male	15	2	-	-	-	-	-	2	1	-	-	2
Female	7	-	-	-	1	-	-	-	-	-	1	-
White, both sexes	16	1	-	-	-	-	-	-	-	-	1	2
Male	11	1	-	-	-	-	-	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	6	1	-	-	1	-	-	2	1	-	-	-
Male	4	1	-	-	-	-	-	2	1	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Washington												
All races, both sexes	18	1	-	1	-	-	2	-	1	1	3	3
Male	6	-	-	-	-	-	-	-	-	1	1	1
Female	12	1	-	1	-	-	2	-	1	-	2	2
White, both sexes	10	-	-	-	-	-	-	-	-	1	2	2
Male	5	-	-	-	-	-	-	-	-	1	-	1
Female	5	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	3	-	-	1	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	1	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	1	-	-	-	-	2	-	1	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	4	1	-	-	-	-	2	-	1	-	-	-
West Virginia												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Wisconsin												
All races, both sexes	13	-	-	-	-	-	-	-	-	3	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	3	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
United States												
All races, both sexes	63	53	57	66	66	82	73	57	28	9	2	-
Male	37	26	31	34	37	45	33	21	5	4	-	-
Female	26	27	26	32	29	37	40	36	23	5	2	-
White, both sexes	52	39	43	58	56	69	62	52	23	9	2	-
Male	31	21	21	30	33	37	29	20	4	4	-	-
Female	21	18	22	28	23	32	33	32	19	5	2	-
Black, both sexes	10	13	12	5	9	11	10	4	4	-	-	-
Male	5	4	8	2	3	7	4	-	1	-	-	-
Female	5	9	4	3	6	4	6	4	3	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	3	1	2	1	1	1	-	-	-
Male	-	1	2	2	1	1	-	1	-	-	-	-
Female	-	-	-	1	-	1	1	-	1	-	-	-
Alabama												
All races, both sexes	-	-	-	-	7	2	4	1	1	-	-	-
Male	-	-	-	-	5	1	1	-	-	-	-	-
Female	-	-	-	-	2	1	3	1	1	-	-	-
White, both sexes	-	-	-	-	6	1	2	1	-	-	-	-
Male	-	-	-	-	4	1	-	-	-	-	-	-
Female	-	-	-	-	2	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	1	1	2	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1	2	1	3	-	3	-	1	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	3	-	2	-	1	-	-	-	-
White, both sexes	-	2	1	3	-	3	-	1	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	3	-	2	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Arkansas												
All races, both sexes	-	1	-	2	-	1	-	2	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	1	-	2	-	-	-	-
White, both sexes	-	-	-	2	-	1	-	2	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	2	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	6	4	6	4	8	3	3	2	1	2	1	-
Male	2	3	3	2	5	1	1	2	-	1	-	-
Female	4	1	3	2	3	2	2	-	1	1	1	-
White, both sexes	5	4	6	1	6	3	3	2	1	2	1	-
Male	2	3	3	-	3	1	1	2	-	1	-	-
Female	3	1	3	1	3	2	2	-	1	1	1	-
Black, both sexes	1	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Colorado												
All races, both sexes	2	1	-	-	-	1	-	2	-	-	-	-
Male	2	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	2	1	-	-	-	1	-	2	-	-	-	-
Male	2	-	-	-	-	1	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	-	1	4	3	-	2	1	1	-	-	-
Male	-	-	1	2	2	-	2	1	-	-	-	-
Female	1	-	-	2	1	-	-	-	1	-	-	-
White, both sexes	1	-	-	3	3	-	2	1	1	-	-	-
Male	-	-	-	2	2	-	2	1	-	-	-	-
Female	1	-	-	1	1	-	-	-	1	-	-	-
Black, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Delaware												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Florida												
All races, both sexes	4	3	2	2	3	7	4	4	2	2	-	-
Male	3	-	1	1	1	4	3	2	-	-	-	-
Female	1	3	1	1	2	3	1	2	2	2	-	-
White, both sexes	3	2	-	2	3	6	4	4	2	2	-	-
Male	3	-	-	1	1	4	3	2	-	-	-	-
Female	-	2	-	1	2	2	1	2	2	2	-	-
Black, both sexes	1	1	2	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	1	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	6	1	-	2	6	1	-	2	2	-	-	-
Male	3	-	-	-	3	1	-	-	-	-	-	-
Female	3	1	-	2	3	-	-	2	2	-	-	-
White, both sexes	5	-	-	1	5	1	-	1	1	-	-	-
Male	2	-	-	-	3	1	-	-	-	-	-	-
Female	3	-	-	1	2	-	-	1	1	-	-	-
Black, both sexes	1	1	-	1	1	-	-	1	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	1	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Hawaii												
All races, both sexes	-	1	-	-	-	-	1	1	-	1	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Illinois												
All races, both sexes	2	1	2	2	2	5	3	6	2	-	-	-
Male	1	1	2	1	-	2	2	2	-	-	-	-
Female	1	-	-	1	2	3	1	4	2	-	-	-
White, both sexes	2	1	1	2	2	5	1	6	2	-	-	-
Male	1	1	1	1	-	2	-	2	-	-	-	-
Female	1	-	-	1	2	3	1	4	2	-	-	-
Black, both sexes	-	-	1	-	-	-	2	-	-	-	-	-
Male	-	-	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	4	2	3	1	1	4	-	-	-	-	-
Male	-	3	1	3	1	1	1	-	-	-	-	-
Female	-	1	1	-	-	-	3	-	-	-	-	-
White, both sexes	-	4	2	3	1	1	4	-	-	-	-	-
Male	-	3	1	3	1	1	1	-	-	-	-	-
Female	-	1	1	-	-	-	3	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Iowa												
All races, both sexes	-	1	-	2	-	1	1	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	2	-	1	1	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	-	-	1	1	4	-	-	1	-	-	-
Male	2	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	1	-	-	-
White, both sexes	2	-	-	1	-	4	-	-	1	-	-	-
Male	2	-	-	1	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	1	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Kentucky												
All races, both sexes	1	-	1	5	2	2	4	-	1	-	-	-
Male	-	-	-	2	1	-	1	-	-	-	-	-
Female	1	-	1	3	1	2	3	-	1	-	-	-
White, both sexes	-	-	1	5	2	2	4	-	1	-	-	-
Male	-	-	-	2	1	-	1	-	-	-	-	-
Female	-	-	1	3	1	2	3	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	-	3	1	1	1	1	-	-	-	-	-
Male	1	-	2	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	1	-	-	-	-	-
White, both sexes	-	-	2	1	1	1	1	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	1	-	-	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Maine												
All races, both sexes	-	-	2	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	2	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	1	2	-	2	2	2	-	-	-	-	-
Male	2	-	-	-	1	1	1	-	-	-	-	-
Female	1	1	2	-	1	1	1	-	-	-	-	-
White, both sexes	2	-	2	-	1	1	2	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	2	-	1	1	1	-	-	-	-	-
Black, both sexes	1	1	-	-	1	1	-	-	-	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Massachusetts												
All races, both sexes	-	3	-	2	-	1	2	2	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	3	-	2	-	1	-	2	-	-	-	-
White, both sexes	-	3	-	2	-	-	2	2	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	3	-	2	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Michigan												
All races, both sexes	2	2	3	2	3	4	3	2	1	1	-	-
Male	2	2	3	1	1	4	-	1	1	1	-	-
Female	-	-	-	1	2	-	3	1	-	-	-	-
White, both sexes	2	1	2	2	3	4	2	2	-	1	-	-
Male	2	1	2	1	1	4	-	1	-	1	-	-
Female	-	-	-	1	2	-	2	1	-	-	-	-
Black, both sexes	-	1	1	-	-	-	1	-	1	-	-	-
Male	-	1	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Minnesota												
All races, both sexes	1	-	-	1	1	-	1	1	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	1	-	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Missouri												
All races, both sexes	-	-	1	2	1	-	2	-	1	2	-	-
Male	-	-	1	-	1	-	1	-	1	1	-	-
Female	-	-	-	2	-	-	1	-	-	1	-	-
White, both sexes	-	-	-	2	1	-	2	-	1	2	-	-
Male	-	-	-	-	1	-	1	-	1	1	-	-
Female	-	-	-	2	-	-	1	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	2	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	2	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	2	1	1	1	1	-	-	-	-	-
Male	-	-	-	1	1	1	1	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	2	1	1	-	1	-	-	-	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
New Hampshire												
All races, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	-	3	-	4	5	1	4	1	-	-	-
Male	1	-	2	-	3	1	-	1	-	-	-	-
Female	-	-	1	-	1	4	1	3	1	-	-	-
White, both sexes	1	-	2	-	4	5	-	3	1	-	-	-
Male	1	-	1	-	3	1	-	1	-	-	-	-
Female	-	-	1	-	1	4	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
New Mexico												
All races, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3	5	5	7	2	3	3	4	5	-	-	-
Male	2	3	1	4	-	2	1	1	2	-	-	-
Female	1	2	4	3	2	1	2	3	3	-	-	-
White, both sexes	2	4	4	5	1	3	2	4	5	-	-	-
Male	1	3	1	2	-	2	1	1	2	-	-	-
Female	1	1	3	3	1	1	1	3	3	-	-	-
Black, both sexes	1	1	1	1	1	-	1	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	1	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
North Carolina												
All races, both sexes	3	2	3	1	2	6	1	1	-	-	-	-
Male	2	-	2	-	1	3	-	-	-	-	-	-
Female	1	2	1	1	1	3	1	1	-	-	-	-
White, both sexes	2	1	2	1	1	3	1	1	-	-	-	-
Male	2	-	1	-	1	1	-	-	-	-	-	-
Female	-	1	1	1	-	2	1	1	-	-	-	-
Black, both sexes	1	1	1	-	1	3	-	-	-	-	-	-
Male	-	-	1	-	-	2	-	-	-	-	-	-
Female	1	1	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Ohio												
All races, both sexes	3	3	2	-	1	-	2	-	-	-	-	-
Male	2	2	-	-	-	-	2	-	-	-	-	-
Female	1	1	2	-	1	-	-	-	-	-	-	-
White, both sexes	3	2	2	-	1	-	2	-	-	-	-	-
Male	2	2	-	-	-	-	2	-	-	-	-	-
Female	1	-	2	-	1	-	-	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	1	-	1	1	2	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
White, both sexes	1	-	-	1	-	1	1	2	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Oregon												
All races, both sexes	-	1	-	-	1	2	1	1	-	-	-	-
Male	-	1	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	-	1	-	-	1	2	1	1	-	-	-	-
Male	-	1	-	-	-	2	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	2	3	1	4	6	7	5	2	1	-	-
Male	1	1	3	1	3	4	3	2	-	1	-	-
Female	2	1	-	-	1	2	4	3	2	-	-	-
White, both sexes	3	1	2	1	3	5	7	5	1	1	-	-
Male	1	-	2	1	3	4	3	2	-	1	-	-
Female	2	1	-	-	-	1	4	3	1	-	-	-
Black, both sexes	-	1	1	-	1	1	-	-	1	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Rhode Island												
All races, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	-	2	2	-	2	3	1	-	-	-	-
Male	-	-	2	2	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
White, both sexes	-	-	1	2	-	1	3	-	-	-	-	-
Male	-	-	1	2	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
South Dakota												
All races, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	3	3	2	-	2	4	1	-	-	-	-
Male	1	1	2	-	-	2	2	1	-	-	-	-
Female	1	2	1	2	-	-	2	-	-	-	-	-
White, both sexes	2	1	2	2	-	1	4	1	-	-	-	-
Male	1	-	2	-	-	1	2	1	-	-	-	-
Female	1	1	-	2	-	-	2	-	-	-	-	-
Black, both sexes	-	2	1	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Texas												
All races, both sexes	8	6	2	6	4	9	3	3	3	-	-	-
Male	4	3	-	2	3	8	2	1	-	-	-	-
Female	4	3	2	4	1	1	1	2	3	-	-	-
White, both sexes	7	5	1	5	4	8	2	3	2	-	-	-
Male	3	2	-	2	3	7	1	1	-	-	-	-
Female	4	3	1	3	1	1	1	2	2	-	-	-
Black, both sexes	1	1	1	1	-	-	1	-	-	-	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Utah												
All races, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Vermont												
All races, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	2	2	1	1	2	1	1	-	1	-
Male	1	1	2	1	-	-	1	1	1	-	-	-
Female	-	-	-	1	1	1	1	-	-	-	1	-
White, both sexes	1	1	2	2	-	1	2	1	1	-	1	-
Male	1	1	2	1	-	-	1	1	1	-	-	-
Female	-	-	-	1	-	1	1	-	-	-	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Washington												
All races, both sexes	-	1	-	-	2	-	3	-	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	1	-	-	1	-	1	-	-	-	-	-
White, both sexes	-	1	-	-	2	-	2	-	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	-	-	1	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	1	-	-	-	-
White, both sexes	-	-	1	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of pericardium and acute myocarditis (I30-I31,I40)												
Wisconsin												
All races, both sexes	2	1	-	1	-	-	1	3	1	-	-	-
Male	2	-	-	1	-	-	-	2	-	-	-	-
Female	-	1	-	-	-	-	1	1	1	-	-	-
White, both sexes	2	1	-	1	-	-	1	3	1	-	-	-
Male	2	-	-	1	-	-	-	2	-	-	-	-
Female	-	1	-	-	-	-	1	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
United States												
All races, both sexes	20,891	6	8	2	10	13	38	50	78	106	176	308
Male	8,008	3	8	2	5	10	22	36	43	67	108	186
Female	12,883	3	-	-	5	3	16	14	35	39	68	122
White, both sexes	19,317	4	6	1	8	11	31	36	58	72	133	231
Male	7,379	2	6	1	4	9	18	26	34	50	92	144
Female	11,938	2	-	-	4	2	13	10	24	22	41	87
Black, both sexes	1,251	2	1	1	2	1	5	13	17	32	38	67
Male	518	1	1	1	1	1	2	9	8	16	12	39
Female	733	1	-	-	1	-	3	4	9	16	26	28
American Indian 1/, both sexes	65	-	1	-	-	-	-	-	1	1	2	4
Male	28	-	1	-	-	-	-	-	-	-	1	2
Female	37	-	-	-	-	-	-	-	1	1	1	2
Asian or Pacific Islander 2/, both sexes	258	-	-	-	-	1	2	1	2	1	3	6
Male	83	-	-	-	-	-	2	1	1	1	3	1
Female	175	-	-	-	-	1	-	-	1	-	-	5
Alabama												
All races, both sexes	303	-	-	-	-	-	-	1	2	4	2	7
Male	103	-	-	-	-	-	-	-	2	1	-	5
Female	200	-	-	-	-	-	-	1	-	3	2	2
White, both sexes	254	-	-	-	-	-	-	-	2	3	2	4
Male	86	-	-	-	-	-	-	-	2	1	-	2
Female	168	-	-	-	-	-	-	-	-	2	2	2
Black, both sexes	49	-	-	-	-	-	-	1	-	1	-	3
Male	17	-	-	-	-	-	-	-	-	-	-	3
Female	32	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Alaska												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	312	-	-	-	-	-	-	1	2	-	5	3
Male	139	-	-	-	-	-	-	1	1	-	3	2
Female	173	-	-	-	-	-	-	-	1	-	2	1
White, both sexes	300	-	-	-	-	-	-	1	1	-	5	2
Male	130	-	-	-	-	-	-	1	-	-	3	1
Female	170	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Arkansas												
All races, both sexes	162	-	-	-	-	-	-	-	-	2	1	3
Male	62	-	-	-	-	-	-	-	-	1	1	2
Female	100	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	147	-	-	-	-	-	-	-	-	1	1	3
Male	54	-	-	-	-	-	-	-	-	-	1	2
Female	93	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	15	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2,119	1	2	-	1	2	4	3	7	13	18	28
Male	848	-	2	-	1	2	3	2	6	8	13	18
Female	1,271	1	-	-	-	-	1	1	1	5	5	10
White, both sexes	1,930	1	2	-	-	2	2	3	6	11	16	23
Male	777	-	2	-	-	2	1	2	5	7	12	15
Female	1,153	1	-	-	-	-	1	1	1	4	4	8
Black, both sexes	72	-	-	-	1	-	1	-	1	1	2	3
Male	37	-	-	-	1	-	1	-	1	-	1	3
Female	35	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	111	-	-	-	-	-	1	-	-	1	-	2
Male	32	-	-	-	-	-	1	-	-	1	-	-
Female	79	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Colorado												
All races, both sexes	287	-	-	-	-	-	-	1	2	4	2	1
Male	121	-	-	-	-	-	-	1	2	3	2	-
Female	166	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	278	-	-	-	-	-	-	1	2	3	1	1
Male	117	-	-	-	-	-	-	1	2	3	1	-
Female	161	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	289	-	-	-	-	1	-	-	1	-	1	4
Male	103	-	-	-	-	1	-	-	-	-	1	3
Female	186	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	280	-	-	-	-	1	-	-	1	-	1	4
Male	99	-	-	-	-	1	-	-	-	-	1	3
Female	181	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Delaware												
All races, both sexes	73	-	-	-	-	-	-	1	-	-	1	-
Male	22	-	-	-	-	-	-	-	-	-	1	-
Female	51	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	1	-	-	1	-
Male	18	-	-	-	-	-	-	-	-	-	1	-
Female	47	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	23	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Florida												
All races, both sexes	1,398	-	1	-	-	-	2	4	5	4	7	21
Male	567	-	1	-	-	-	1	3	2	2	5	11
Female	831	-	-	-	-	-	1	1	3	2	2	10
White, both sexes	1,316	-	-	-	-	-	2	2	3	3	5	17
Male	536	-	-	-	-	-	1	1	1	2	4	9
Female	780	-	-	-	-	-	1	1	2	1	1	8
Black, both sexes	81	-	-	-	-	-	-	2	2	1	2	4
Male	30	-	-	-	-	-	-	2	1	-	1	2
Female	51	-	-	-	-	-	-	-	1	1	1	2
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	417	-	-	-	-	-	1	1	5	3	2	8
Male	173	-	-	-	-	-	1	1	3	2	-	5
Female	244	-	-	-	-	-	-	-	2	1	2	3
White, both sexes	332	-	-	-	-	-	1	1	3	1	-	4
Male	139	-	-	-	-	-	1	1	2	1	-	3
Female	193	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	85	-	-	-	-	-	-	-	2	2	2	4
Male	34	-	-	-	-	-	-	-	1	1	-	2
Female	51	-	-	-	-	-	-	-	1	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Hawaii												
All races, both sexes	97	-	-	-	-	-	-	-	-	-	2	2
Male	29	-	-	-	-	-	-	-	-	-	1	-
Female	68	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	32	-	-	-	-	-	-	-	-	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	65	-	-	-	-	-	-	-	-	-	1	2
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	46	-	-	-	-	-	-	-	-	-	-	2
Idaho												
All races, both sexes	136	-	-	-	-	-	-	-	-	1	-	-
Male	59	-	-	-	-	-	-	-	-	1	-	-
Female	77	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	133	-	-	-	-	-	-	-	-	1	-	-
Male	57	-	-	-	-	-	-	-	-	1	-	-
Female	76	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Illinois												
All races, both sexes	772	-	-	-	3	1	4	2	5	8	4	11
Male	296	-	-	-	-	-	1	1	2	5	2	7
Female	476	-	-	-	3	1	3	1	3	3	2	4
White, both sexes	698	-	-	-	2	1	3	2	4	6	3	6
Male	258	-	-	-	-	-	1	1	2	5	2	4
Female	440	-	-	-	2	1	2	1	2	1	1	2
Black, both sexes	72	-	-	-	1	-	1	-	1	2	1	5
Male	36	-	-	-	-	-	-	-	-	-	-	3
Female	36	-	-	-	1	-	1	-	1	2	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	509	-	-	-	-	-	2	-	2	2	3	16
Male	177	-	-	-	-	-	-	-	-	1	1	8
Female	332	-	-	-	-	-	2	-	2	1	2	8
White, both sexes	487	-	-	-	-	-	2	-	2	1	3	14
Male	170	-	-	-	-	-	-	-	-	-	1	7
Female	317	-	-	-	-	-	2	-	2	1	2	7
Black, both sexes	19	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	1	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Iowa												
All races, both sexes	281	-	1	-	-	-	-	1	-	-	-	5
Male	90	-	1	-	-	-	-	-	-	-	-	1
Female	191	-	-	-	-	-	-	1	-	-	-	4
White, both sexes	276	-	1	-	-	-	-	1	-	-	-	4
Male	89	-	1	-	-	-	-	-	-	-	-	1
Female	187	-	-	-	-	-	-	1	-	-	-	3
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	234	-	-	-	-	-	-	-	-	-	-	3
Male	80	-	-	-	-	-	-	-	-	-	-	2
Female	154	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	225	-	-	-	-	-	-	-	-	-	-	1
Male	73	-	-	-	-	-	-	-	-	-	-	-
Female	152	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Kentucky												
All races, both sexes	227	1	-	-	-	1	-	-	1	2	4	3
Male	83	1	-	-	-	1	-	-	-	1	-	2
Female	144	-	-	-	-	-	-	-	1	1	4	1
White, both sexes	217	-	-	-	-	1	-	-	1	2	4	3
Male	78	-	-	-	-	1	-	-	-	1	-	2
Female	139	-	-	-	-	-	-	-	1	1	4	1
Black, both sexes	9	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	249	-	1	-	1	1	1	1	1	2	2	3
Male	108	-	1	-	1	1	1	-	-	1	2	1
Female	141	-	-	-	-	-	-	1	1	1	-	2
White, both sexes	197	-	-	-	1	-	1	-	-	1	2	2
Male	89	-	-	-	1	-	1	-	-	-	2	1
Female	108	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	52	-	1	-	-	1	-	1	1	1	-	1
Male	19	-	1	-	-	1	-	-	-	1	-	-
Female	33	-	-	-	-	-	-	1	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Maine												
All races, both sexes	133	-	-	-	-	-	-	-	-	1	1	1
Male	56	-	-	-	-	-	-	-	-	1	1	-
Female	77	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	133	-	-	-	-	-	-	-	-	1	1	1
Male	56	-	-	-	-	-	-	-	-	1	1	-
Female	77	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	316	1	-	-	1	-	1	2	2	3	5	4
Male	116	1	-	-	-	-	-	2	1	1	3	3
Female	200	-	-	-	1	-	1	-	1	2	2	1
White, both sexes	261	1	-	-	1	-	1	2	2	1	2	2
Male	95	1	-	-	-	-	-	2	1	1	2	2
Female	166	-	-	-	1	-	1	-	1	-	-	-
Black, both sexes	53	-	-	-	-	-	-	-	-	2	3	2
Male	20	-	-	-	-	-	-	-	-	-	1	1
Female	33	-	-	-	-	-	-	-	-	2	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Massachusetts												
All races, both sexes	502	-	-	1	-	1	3	-	-	2	3	4
Male	176	-	-	1	-	1	2	-	-	1	1	3
Female	326	-	-	-	-	-	1	-	-	1	2	1
White, both sexes	482	-	-	1	-	1	3	-	-	-	2	4
Male	168	-	-	1	-	1	2	-	-	-	-	3
Female	314	-	-	-	-	-	1	-	-	-	2	1
Black, both sexes	18	-	-	-	-	-	-	-	-	2	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	699	-	-	-	-	2	-	1	4	4	14	15
Male	263	-	-	-	-	2	-	-	3	2	8	13
Female	436	-	-	-	-	-	-	1	1	2	6	2
White, both sexes	644	-	-	-	-	2	-	1	4	2	10	11
Male	234	-	-	-	-	2	-	-	3	-	7	10
Female	410	-	-	-	-	-	-	1	1	2	3	1
Black, both sexes	51	-	-	-	-	-	-	-	-	2	3	4
Male	27	-	-	-	-	-	-	-	-	2	-	3
Female	24	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Minnesota												
All races, both sexes	484	-	-	-	-	-	1	-	1	1	1	8
Male	180	-	-	-	-	-	1	-	1	1	1	3
Female	304	-	-	-	-	-	-	-	-	-	-	5
White, both sexes	475	-	-	-	-	-	1	-	1	1	-	8
Male	176	-	-	-	-	-	1	-	1	1	-	3
Female	299	-	-	-	-	-	-	-	-	-	-	5
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	135	-	-	-	-	-	-	1	1	4	-	5
Male	58	-	-	-	-	-	-	1	1	1	-	3
Female	77	-	-	-	-	-	-	-	-	3	-	2
White, both sexes	101	-	-	-	-	-	-	-	1	-	-	3
Male	46	-	-	-	-	-	-	-	1	-	-	2
Female	55	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	32	-	-	-	-	-	-	1	-	4	-	2
Male	12	-	-	-	-	-	-	1	-	1	-	1
Female	20	-	-	-	-	-	-	-	-	3	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Missouri												
All races, both sexes	447	-	-	-	1	-	-	1	2	4	2	5
Male	171	-	-	-	1	-	-	1	1	1	1	5
Female	276	-	-	-	-	-	-	-	1	3	1	-
White, both sexes	419	-	-	-	1	-	-	1	2	4	2	2
Male	160	-	-	-	1	-	-	1	1	1	1	2
Female	259	-	-	-	-	-	-	-	1	3	1	-
Black, both sexes	26	-	-	-	-	-	-	-	-	-	-	3
Male	11	-	-	-	-	-	-	-	-	-	-	3
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	97	-	-	-	-	-	-	-	1	-	1	-
Male	35	-	-	-	-	-	-	-	-	-	1	-
Female	62	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	96	-	-	-	-	-	-	-	1	-	-	-
Male	34	-	-	-	-	-	-	-	-	-	-	-
Female	62	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Nebraska												
All races, both sexes	176	-	-	-	-	-	-	-	-	-	-	1
Male	63	-	-	-	-	-	-	-	-	-	-	-
Female	113	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	171	-	-	-	-	-	-	-	-	-	-	1
Male	60	-	-	-	-	-	-	-	-	-	-	-
Female	111	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	137	1	-	-	-	1	1	1	-	1	3	5
Male	53	-	-	-	-	-	-	1	-	1	1	2
Female	84	1	-	-	-	1	1	-	-	-	2	3
White, both sexes	123	1	-	-	-	-	1	-	-	1	2	4
Male	49	-	-	-	-	-	-	-	-	1	1	1
Female	74	1	-	-	-	-	1	-	-	-	1	3
Black, both sexes	9	-	-	-	-	-	-	1	-	-	1	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	1	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
New Hampshire												
All races, both sexes	110	-	-	-	-	-	-	-	-	-	-	1
Male	40	-	-	-	-	-	-	-	-	-	-	1
Female	70	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	110	-	-	-	-	-	-	-	-	-	-	1
Male	40	-	-	-	-	-	-	-	-	-	-	1
Female	70	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	752	-	-	-	-	1	1	2	3	4	9	14
Male	264	-	-	-	-	-	-	1	2	3	4	8
Female	488	-	-	-	-	1	1	1	1	1	5	6
White, both sexes	699	-	-	-	-	1	1	1	2	4	5	8
Male	244	-	-	-	-	-	-	-	2	3	3	6
Female	455	-	-	-	-	1	1	1	-	1	2	2
Black, both sexes	48	-	-	-	-	-	-	-	1	-	4	6
Male	17	-	-	-	-	-	-	-	-	-	1	2
Female	31	-	-	-	-	-	-	-	1	-	3	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
New Mexico												
All races, both sexes	81	-	-	-	-	-	-	1	-	-	-	1
Male	39	-	-	-	-	-	-	1	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	78	-	-	-	-	-	-	1	-	-	-	1
Male	38	-	-	-	-	-	-	1	-	-	-	1
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,187	1	-	-	-	1	2	5	3	4	10	14
Male	435	-	-	-	-	1	1	4	-	4	7	9
Female	752	1	-	-	-	-	1	1	3	-	3	5
White, both sexes	1,124	-	-	-	-	1	2	4	3	2	9	10
Male	407	-	-	-	-	1	1	3	-	2	6	7
Female	717	-	-	-	-	-	1	1	3	-	3	3
Black, both sexes	54	1	-	-	-	-	-	1	-	2	1	4
Male	23	-	-	-	-	-	-	1	-	2	1	2
Female	31	1	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
North Carolina												
All races, both sexes	608	-	-	1	-	-	-	1	2	1	2	9
Male	238	-	-	1	-	-	-	1	1	-	2	6
Female	370	-	-	-	-	-	-	-	1	1	-	3
White, both sexes	520	-	-	-	-	-	-	1	-	-	1	6
Male	213	-	-	-	-	-	-	1	-	-	1	5
Female	307	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	80	-	-	1	-	-	-	-	1	1	1	3
Male	24	-	-	1	-	-	-	-	1	-	1	1
Female	56	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	1	-	-	-
North Dakota												
All races, both sexes	52	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	52	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Ohio												
All races, both sexes	912	-	2	-	2	-	5	3	4	6	16	12
Male	363	-	2	-	2	-	4	2	3	5	11	8
Female	549	-	-	-	-	-	1	1	1	1	5	4
White, both sexes	851	-	2	-	2	-	4	2	4	5	13	7
Male	336	-	2	-	2	-	4	2	3	5	11	5
Female	515	-	-	-	-	-	-	-	1	-	2	2
Black, both sexes	58	-	-	-	-	-	1	1	-	1	3	5
Male	26	-	-	-	-	-	-	-	-	-	-	3
Female	32	-	-	-	-	-	1	1	-	1	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	221	-	1	-	-	-	-	-	1	2	3	5
Male	80	-	1	-	-	-	-	-	-	1	3	5
Female	141	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	204	-	1	-	-	-	-	-	-	1	3	3
Male	71	-	1	-	-	-	-	-	-	1	3	3
Female	133	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	1	1	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Oregon												
All races, both sexes	494	-	-	-	-	-	-	1	2	1	1	2
Male	197	-	-	-	-	-	-	1	1	1	1	1
Female	297	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	485	-	-	-	-	-	-	-	2	1	1	2
Male	194	-	-	-	-	-	-	-	1	1	1	1
Female	291	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,508	-	-	-	-	-	1	2	3	5	12	16
Male	539	-	-	-	-	-	-	2	1	4	10	9
Female	969	-	-	-	-	-	1	-	2	1	2	7
White, both sexes	1,437	-	-	-	-	-	1	2	1	5	9	13
Male	510	-	-	-	-	-	-	2	1	4	8	8
Female	927	-	-	-	-	-	1	-	-	1	1	5
Black, both sexes	67	-	-	-	-	-	-	-	2	-	3	2
Male	29	-	-	-	-	-	-	-	-	-	2	1
Female	38	-	-	-	-	-	-	-	2	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Rhode Island												
All races, both sexes	80	-	-	-	-	-	-	-	-	1	1	1
Male	31	-	-	-	-	-	-	-	-	-	1	-
Female	49	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	78	-	-	-	-	-	-	-	-	1	1	1
Male	29	-	-	-	-	-	-	-	-	-	1	-
Female	49	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	315	-	-	-	-	1	-	-	2	1	5	11
Male	113	-	-	-	-	1	-	-	1	1	-	6
Female	202	-	-	-	-	-	-	-	1	-	5	5
White, both sexes	246	-	-	-	-	1	-	-	-	-	1	8
Male	91	-	-	-	-	1	-	-	-	-	-	6
Female	155	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	66	-	-	-	-	-	-	-	2	1	4	3
Male	20	-	-	-	-	-	-	-	1	1	-	-
Female	46	-	-	-	-	-	-	-	1	-	4	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
South Dakota												
All races, both sexes	66	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	352	1	-	-	-	-	2	2	1	1	2	6
Male	131	1	-	-	-	-	2	2	-	1	1	2
Female	221	-	-	-	-	-	-	-	1	-	1	4
White, both sexes	330	1	-	-	-	-	2	-	1	1	2	6
Male	123	1	-	-	-	-	2	-	-	1	1	2
Female	207	-	-	-	-	-	-	-	1	-	1	4
Black, both sexes	22	-	-	-	-	-	-	2	-	-	-	-
Male	8	-	-	-	-	-	-	2	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Texas												
All races, both sexes	1,070	-	-	-	-	-	1	5	6	10	17	21
Male	431	-	-	-	-	-	1	4	3	8	11	15
Female	639	-	-	-	-	-	-	1	3	2	6	6
White, both sexes	991	-	-	-	-	-	-	5	4	6	12	18
Male	397	-	-	-	-	-	-	4	2	5	9	12
Female	594	-	-	-	-	-	-	1	2	1	3	6
Black, both sexes	75	-	-	-	-	-	1	-	2	4	5	3
Male	34	-	-	-	-	-	1	-	1	3	2	3
Female	41	-	-	-	-	-	-	-	1	1	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	132	-	-	-	-	-	1	-	1	-	1	1
Male	43	-	-	-	-	-	1	-	1	-	1	1
Female	89	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	131	-	-	-	-	-	1	-	1	-	1	1
Male	43	-	-	-	-	-	1	-	1	-	1	1
Female	88	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Vermont												
All races, both sexes	64	-	-	-	-	-	-	-	-	-	1	1
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	64	-	-	-	-	-	-	-	-	-	1	1
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	431	-	-	-	-	-	4	4	4	2	8	7
Male	167	-	-	-	-	-	2	3	3	1	6	2
Female	264	-	-	-	-	-	2	1	1	1	2	5
White, both sexes	379	-	-	-	-	-	2	2	2	2	8	3
Male	146	-	-	-	-	-	1	2	1	1	6	1
Female	233	-	-	-	-	-	1	-	1	1	2	2
Black, both sexes	45	-	-	-	-	-	1	2	1	-	-	4
Male	17	-	-	-	-	-	-	1	1	-	-	1
Female	28	-	-	-	-	-	1	1	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	1	-	1	-	-	-
Male	4	-	-	-	-	-	1	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Washington												
All races, both sexes	689	-	-	-	-	-	1	1	2	2	1	12
Male	287	-	-	-	-	-	1	-	2	2	-	7
Female	402	-	-	-	-	-	-	1	-	-	1	5
White, both sexes	666	-	-	-	-	-	1	1	2	1	-	12
Male	276	-	-	-	-	-	1	-	2	1	-	7
Female	390	-	-	-	-	-	-	1	-	-	-	5
Black, both sexes	8	-	-	-	-	-	-	-	-	1	1	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	137	-	-	-	-	-	-	1	-	-	-	2
Male	57	-	-	-	-	-	-	1	-	-	-	1
Female	80	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	135	-	-	-	-	-	-	1	-	-	-	2
Male	56	-	-	-	-	-	-	1	-	-	-	1
Female	79	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of endocardium (I34-I38)												
Wisconsin												
All races, both sexes	544	-	-	-	1	-	-	-	-	-	2	4
Male	213	-	-	-	-	-	-	-	-	-	1	4
Female	331	-	-	-	1	-	-	-	-	-	1	-
White, both sexes	528	-	-	-	1	-	-	-	-	-	2	3
Male	205	-	-	-	-	-	-	-	-	-	1	3
Female	323	-	-	-	1	-	-	-	-	-	1	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
United States												
All races, both sexes	402	502	617	813	1,234	2,311	3,783	4,863	3,794	1,539	236	2
Male	240	309	348	397	576	1,053	1,505	1,690	1,055	302	43	-
Female	162	193	269	416	658	1,258	2,278	3,173	2,739	1,237	193	2
White, both sexes	287	396	497	709	1,082	2,149	3,575	4,674	3,666	1,472	218	1
Male	178	241	290	359	515	988	1,433	1,630	1,032	288	39	-
Female	109	155	207	350	567	1,161	2,142	3,044	2,634	1,184	179	1
Black, both sexes	106	96	102	87	123	132	149	131	86	48	11	1
Male	60	61	52	33	52	55	51	40	12	9	2	-
Female	46	35	50	54	71	77	98	91	74	39	9	1
American Indian 1/, both sexes	2	5	5	5	4	6	7	12	5	5	-	-
Male	-	3	2	2	3	-	6	5	1	2	-	-
Female	2	2	3	3	1	6	1	7	4	3	-	-
Asian or Pacific Islander 2/, both sexes	7	5	13	12	25	24	52	46	37	14	7	-
Male	2	4	4	3	6	10	15	15	10	3	2	-
Female	5	1	9	9	19	14	37	31	27	11	5	-
Alabama												
All races, both sexes	12	9	7	20	24	43	55	51	42	19	5	-
Male	6	3	1	8	11	16	25	13	8	4	-	-
Female	6	6	6	12	13	27	30	38	34	15	5	-
White, both sexes	5	6	5	17	20	36	49	47	37	18	3	-
Male	2	2	1	7	9	15	23	11	8	3	-	-
Female	3	4	4	10	11	21	26	36	29	15	3	-
Black, both sexes	7	3	2	3	4	7	6	4	5	1	2	-
Male	4	1	-	1	2	1	2	2	-	1	-	-
Female	3	2	2	2	2	6	4	2	5	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Alaska												
All races, both sexes	2	-	-	1	5	5	5	7	7	1	-	-
Male	2	-	-	-	3	2	2	2	2	-	-	-
Female	-	-	-	1	2	3	3	5	5	1	-	-
White, both sexes	2	-	-	-	3	3	4	4	6	-	-	-
Male	2	-	-	-	2	2	2	1	2	-	-	-
Female	-	-	-	-	1	1	2	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	2	1	2	1	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	2	1	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Arizona												
All races, both sexes	10	9	12	23	26	38	46	74	50	12	1	-
Male	7	6	9	10	14	9	20	37	16	3	1	-
Female	3	3	3	13	12	29	26	37	34	9	-	-
White, both sexes	8	8	12	22	26	37	43	72	50	12	1	-
Male	6	5	9	9	14	9	18	35	16	3	1	-
Female	2	3	3	13	12	28	25	37	34	9	-	-
Black, both sexes	1	-	-	-	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	1	-	-	1	1	-	-	-	-
Male	-	1	-	1	-	-	1	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Arkansas												
All races, both sexes	4	7	4	9	11	19	22	45	23	12	-	-
Male	3	5	2	5	4	8	7	14	7	3	-	-
Female	1	2	2	4	7	11	15	31	16	9	-	-
White, both sexes	2	6	3	9	7	17	21	43	22	12	-	-
Male	1	4	1	5	3	6	7	14	7	3	-	-
Female	1	2	2	4	4	11	14	29	15	9	-	-
Black, both sexes	2	1	1	-	4	2	1	2	1	-	-	-
Male	2	1	1	-	1	2	-	-	-	-	-	-
Female	-	-	-	-	3	-	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	41	39	53	62	106	214	398	499	429	161	38	-
Male	26	30	36	28	55	104	172	182	116	39	5	-
Female	15	9	17	34	51	110	226	317	313	122	33	-
White, both sexes	32	33	46	53	90	193	358	472	404	150	33	-
Male	20	24	32	27	51	95	156	175	111	36	4	-
Female	12	9	14	26	39	98	202	297	293	114	29	-
Black, both sexes	5	3	3	6	5	7	15	4	9	5	1	-
Male	4	3	2	1	1	5	9	2	1	2	-	-
Female	1	-	1	5	4	2	6	2	8	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	3	-	1	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	2	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	4	3	4	3	11	12	25	20	16	5	4	-
Male	2	3	2	-	3	4	7	3	4	1	1	-
Female	2	-	2	3	8	8	18	17	12	4	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Colorado												
All races, both sexes	6	4	9	13	14	34	48	74	49	24	2	-
Male	4	1	2	7	5	14	19	37	16	7	1	-
Female	2	3	7	6	9	20	29	37	33	17	1	-
White, both sexes	6	4	9	13	12	30	47	74	49	24	2	-
Male	4	1	2	7	4	12	19	37	16	7	1	-
Female	2	3	7	6	8	18	28	37	33	17	1	-
Black, both sexes	-	-	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	1	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Connecticut												
All races, both sexes	7	3	9	7	14	16	44	78	72	25	7	-
Male	7	3	4	7	9	7	19	22	15	4	1	-
Female	-	-	5	-	5	9	25	56	57	21	6	-
White, both sexes	6	2	6	6	14	16	43	76	72	25	7	-
Male	6	2	3	6	9	7	19	22	15	4	1	-
Female	-	-	3	-	5	9	24	54	57	21	6	-
Black, both sexes	1	1	2	1	-	-	1	2	-	-	-	-
Male	1	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Delaware												
All races, both sexes	-	1	2	6	7	11	11	14	14	4	1	-
Male	-	-	2	2	3	3	3	6	2	-	-	-
Female	-	1	-	4	4	8	8	8	12	4	1	-
White, both sexes	-	1	1	5	6	9	10	12	14	4	1	-
Male	-	-	1	2	2	2	3	5	2	-	-	-
Female	-	1	-	3	4	7	7	7	12	4	1	-
Black, both sexes	-	-	1	1	1	2	1	2	-	-	-	-
Male	-	-	1	-	1	1	-	1	-	-	-	-
Female	-	-	-	1	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	1	3	7	2	1	4	1	-
Male	2	-	-	-	1	1	1	-	1	1	-	-
Female	-	-	-	-	-	2	6	2	-	3	1	-
White, both sexes	-	-	-	-	-	3	3	1	1	3	1	-
Male	-	-	-	-	-	1	1	-	1	1	-	-
Female	-	-	-	-	-	2	2	1	-	2	1	-
Black, both sexes	2	-	-	-	1	-	4	1	-	1	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	4	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Florida												
All races, both sexes	25	35	42	50	71	169	271	326	257	100	7	1
Male	13	24	21	24	34	79	108	124	86	28	1	-
Female	12	11	21	26	37	90	163	202	171	72	6	1
White, both sexes	20	26	33	42	65	163	262	321	249	97	6	-
Male	12	19	18	20	33	76	105	124	84	27	-	-
Female	8	7	15	22	32	87	157	197	165	70	6	-
Black, both sexes	5	9	9	8	6	6	9	5	8	3	1	1
Male	1	5	3	4	1	3	3	-	2	1	1	-
Female	4	4	6	4	5	3	6	5	6	2	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	15	21	17	24	38	62	70	66	63	20	1	-
Male	9	10	11	13	24	29	29	23	9	4	-	-
Female	6	11	6	11	14	33	41	43	54	16	1	-
White, both sexes	9	11	10	19	27	55	60	57	57	17	-	-
Male	5	5	7	9	18	28	27	19	9	4	-	-
Female	4	6	3	10	9	27	33	38	48	13	-	-
Black, both sexes	6	10	7	5	11	7	10	9	6	3	1	-
Male	4	5	4	4	6	1	2	4	-	-	-	-
Female	2	5	3	1	5	6	8	5	6	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Hawaii												
All races, both sexes	1	2	4	3	7	5	17	19	24	10	1	-
Male	-	1	1	1	2	1	6	5	8	3	-	-
Female	1	1	3	2	5	4	11	14	16	7	1	-
White, both sexes	-	2	1	-	4	1	5	7	9	2	-	-
Male	-	1	-	-	2	-	2	1	3	1	-	-
Female	-	1	1	-	2	1	3	6	6	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	3	3	4	12	12	15	8	1	-
Male	-	-	1	1	-	1	4	4	5	2	-	-
Female	1	-	2	2	3	3	8	8	10	6	1	-
Idaho												
All races, both sexes	5	2	2	2	6	14	23	36	31	14	-	-
Male	3	2	-	1	4	7	7	16	12	6	-	-
Female	2	-	2	1	2	7	16	20	19	8	-	-
White, both sexes	5	2	1	2	6	14	22	35	31	14	-	-
Male	3	2	-	1	4	7	6	15	12	6	-	-
Female	2	-	1	1	2	7	16	20	19	8	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Illinois												
All races, both sexes	22	15	26	29	35	75	131	183	131	65	22	-
Male	15	8	15	16	19	37	65	56	33	9	5	-
Female	7	7	11	13	16	38	66	127	98	56	17	-
White, both sexes	12	10	19	26	32	67	121	177	124	63	20	-
Male	9	5	11	14	17	31	60	53	31	9	3	-
Female	3	5	8	12	15	36	61	124	93	54	17	-
Black, both sexes	10	5	7	3	2	8	10	6	7	2	1	-
Male	6	3	4	2	1	6	5	3	2	-	1	-
Female	4	2	3	1	1	2	5	3	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	1	-
Male	-	-	-	-	1	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	4	13	23	23	33	63	83	104	82	50	6	-
Male	2	8	10	7	13	32	29	35	24	7	-	-
Female	2	5	13	16	20	31	54	69	58	43	6	-
White, both sexes	3	13	21	21	28	62	81	104	78	48	6	-
Male	2	8	9	7	11	32	29	35	22	7	-	-
Female	1	5	12	14	17	30	52	69	56	41	6	-
Black, both sexes	1	-	2	2	4	1	2	-	4	1	-	-
Male	-	-	1	-	2	-	-	-	2	-	-	-
Female	1	-	1	2	2	1	2	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Iowa												
All races, both sexes	1	5	6	10	15	23	57	62	64	25	6	-
Male	1	2	2	5	6	12	20	17	16	5	2	-
Female	-	3	4	5	9	11	37	45	48	20	4	-
White, both sexes	1	5	4	9	15	22	57	62	64	25	6	-
Male	1	2	1	5	6	12	20	17	16	5	2	-
Female	-	3	3	4	9	10	37	45	48	20	4	-
Black, both sexes	-	-	2	1	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	8	4	6	9	14	27	34	55	44	26	4	-
Male	4	1	2	2	5	14	15	17	12	5	1	-
Female	4	3	4	7	9	13	19	38	32	21	3	-
White, both sexes	6	4	6	9	14	26	32	55	44	24	4	-
Male	4	1	2	2	5	13	13	17	12	3	1	-
Female	2	3	4	7	9	13	19	38	32	21	3	-
Black, both sexes	1	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	2	-	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Kentucky												
All races, both sexes	9	6	7	10	16	24	31	52	41	18	1	-
Male	7	5	3	4	8	12	9	12	18	-	-	-
Female	2	1	4	6	8	12	22	40	23	18	1	-
White, both sexes	8	4	6	10	14	23	31	52	40	17	1	-
Male	6	4	3	4	7	11	9	12	18	-	-	-
Female	2	-	3	6	7	12	22	40	22	17	1	-
Black, both sexes	1	2	1	-	2	-	-	-	1	1	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	-	1	1	-	1	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	8	11	12	12	24	33	46	46	27	14	3	-
Male	5	4	4	8	11	15	21	19	6	5	2	-
Female	3	7	8	4	13	18	25	27	21	9	1	-
White, both sexes	3	4	9	10	13	31	42	39	24	12	3	-
Male	2	1	3	7	8	15	19	16	6	5	2	-
Female	1	3	6	3	5	16	23	23	18	7	1	-
Black, both sexes	5	7	3	2	11	2	4	7	3	2	-	-
Male	3	3	1	1	3	-	2	3	-	-	-	-
Female	2	4	2	1	8	2	2	4	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Maine												
All races, both sexes	3	6	2	3	10	19	30	31	16	10	-	-
Male	3	4	1	2	8	9	11	9	6	1	-	-
Female	-	2	1	1	2	10	19	22	10	9	-	-
White, both sexes	3	6	2	3	10	19	30	31	16	10	-	-
Male	3	4	1	2	8	9	11	9	6	1	-	-
Female	-	2	1	1	2	10	19	22	10	9	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	13	10	12	14	24	35	55	65	47	19	3	-
Male	9	7	7	8	7	13	21	21	10	1	1	-
Female	4	3	5	6	17	22	34	44	37	18	2	-
White, both sexes	5	7	10	8	17	29	50	59	44	17	3	-
Male	4	4	6	6	3	11	19	21	10	1	1	-
Female	1	3	4	2	14	18	31	38	34	16	2	-
Black, both sexes	8	3	2	5	7	6	5	5	3	2	-	-
Male	5	3	1	1	4	2	2	-	-	-	-	-
Female	3	-	1	4	3	4	3	5	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Massachusetts												
All races, both sexes	3	9	11	18	26	45	79	131	108	46	12	-
Male	1	6	9	13	13	19	25	49	22	7	3	-
Female	2	3	2	5	13	26	54	82	86	39	9	-
White, both sexes	2	9	9	17	24	43	76	128	106	46	11	-
Male	1	6	8	12	13	18	23	48	22	7	3	-
Female	1	3	1	5	11	25	53	80	84	39	8	-
Black, both sexes	1	-	2	1	2	2	3	3	2	-	-	-
Male	-	-	1	1	-	1	2	1	-	-	-	-
Female	1	-	1	-	2	1	1	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Michigan												
All races, both sexes	20	22	24	31	43	85	122	162	108	35	7	-
Male	11	9	16	17	18	33	49	52	24	5	1	-
Female	9	13	8	14	25	52	73	110	84	30	6	-
White, both sexes	14	18	21	25	38	75	118	158	108	32	7	-
Male	7	5	14	15	15	30	46	51	24	4	1	-
Female	7	13	7	10	23	45	72	107	84	28	6	-
Black, both sexes	6	4	3	6	5	9	3	4	-	2	-	-
Male	4	4	2	2	3	3	2	1	-	1	-	-
Female	2	-	1	4	2	6	1	3	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Minnesota												
All races, both sexes	6	7	9	5	18	45	89	125	118	45	5	-
Male	1	2	5	3	7	22	45	43	33	12	-	-
Female	5	5	4	2	11	23	44	82	85	33	5	-
White, both sexes	5	7	7	5	18	44	88	124	116	45	5	-
Male	-	2	4	3	7	22	45	43	32	12	-	-
Female	5	5	3	2	11	22	43	81	84	33	5	-
Black, both sexes	1	-	2	-	-	1	1	-	1	-	-	-
Male	1	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Mississippi												
All races, both sexes	8	7	4	7	17	17	20	24	14	4	2	-
Male	3	5	1	3	8	12	5	8	7	-	-	-
Female	5	2	3	4	9	5	15	16	7	4	2	-
White, both sexes	4	2	4	4	13	14	19	19	13	4	1	-
Male	3	1	1	3	5	10	5	8	7	-	-	-
Female	1	1	3	1	8	4	14	11	6	4	1	-
Black, both sexes	3	5	-	3	4	3	1	4	1	-	1	-
Male	-	4	-	-	3	2	-	-	-	-	-	-
Female	3	1	-	3	1	1	1	4	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Missouri												
All races, both sexes	4	14	17	15	17	55	85	110	82	33	-	-
Male	3	9	10	10	9	26	32	36	22	4	-	-
Female	1	5	7	5	8	29	53	74	60	29	-	-
White, both sexes	1	12	14	14	15	51	80	107	80	33	-	-
Male	1	8	10	10	9	22	32	35	22	4	-	-
Female	-	4	4	4	6	29	48	72	58	29	-	-
Black, both sexes	2	2	3	1	2	4	4	3	2	-	-	-
Male	2	1	-	-	-	4	-	1	-	-	-	-
Female	-	1	3	1	2	-	4	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	2	-	-	1	4	7	20	27	22	10	2	-
Male	-	-	-	1	1	5	7	10	8	1	1	-
Female	2	-	-	-	3	2	13	17	14	9	1	-
White, both sexes	2	-	-	1	4	7	20	27	22	10	2	-
Male	-	-	-	1	1	5	7	10	8	1	1	-
Female	2	-	-	-	3	2	13	17	14	9	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Nebraska												
All races, both sexes	3	3	3	8	15	13	32	36	35	22	5	-
Male	3	3	2	4	3	8	16	13	8	3	-	-
Female	-	-	1	4	12	5	16	23	27	19	5	-
White, both sexes	3	3	3	7	15	12	32	33	35	22	5	-
Male	3	3	2	4	3	7	16	11	8	3	-	-
Female	-	-	1	3	12	5	16	22	27	19	5	-
Black, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	3	8	12	10	9	25	26	18	6	3	-
Male	1	3	2	10	4	2	11	9	6	-	-	-
Female	3	-	6	2	6	7	14	17	12	6	3	-
White, both sexes	2	3	7	10	9	7	25	26	16	6	3	-
Male	1	3	2	9	3	2	11	9	6	-	-	-
Female	1	-	5	1	6	5	14	17	10	6	3	-
Black, both sexes	2	-	-	1	1	2	-	-	1	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	2	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
New Hampshire												
All races, both sexes	1	6	2	4	6	9	25	27	16	11	2	-
Male	1	6	1	2	2	3	9	9	5	1	-	-
Female	-	-	1	2	4	6	16	18	11	10	2	-
White, both sexes	1	6	2	4	6	9	25	27	16	11	2	-
Male	1	6	1	2	2	3	9	9	5	1	-	-
Female	-	-	1	2	4	6	16	18	11	10	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	11	22	23	25	41	89	130	193	125	51	8	-
Male	5	10	13	15	14	37	48	65	26	10	3	-
Female	6	12	10	10	27	52	82	128	99	41	5	-
White, both sexes	6	20	15	23	36	84	125	190	120	50	8	-
Male	3	9	9	14	12	36	47	64	24	9	3	-
Female	3	11	6	9	24	48	78	126	96	41	5	-
Black, both sexes	5	2	7	2	5	5	4	2	5	-	-	-
Male	2	1	4	1	2	1	1	-	2	-	-	-
Female	3	1	3	1	3	4	3	2	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
New Mexico												
All races, both sexes	4	4	3	4	2	10	13	19	17	3	-	-
Male	3	2	1	2	2	7	4	10	5	1	-	-
Female	1	2	2	2	-	3	9	9	12	2	-	-
White, both sexes	4	4	1	4	2	10	13	19	16	3	-	-
Male	3	2	-	2	2	7	4	10	5	1	-	-
Female	1	2	1	2	-	3	9	9	11	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	2	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	14	28	36	47	61	141	223	270	217	102	7	1
Male	8	18	21	24	33	70	72	89	56	17	1	-
Female	6	10	15	23	28	71	151	181	161	85	6	1
White, both sexes	12	23	28	40	54	133	214	264	215	102	7	1
Male	8	16	17	22	30	68	68	85	55	17	1	-
Female	4	7	11	18	24	65	146	179	160	85	6	1
Black, both sexes	2	3	8	5	7	8	7	4	1	-	-	-
Male	-	1	4	1	3	2	3	3	-	-	-	-
Female	2	2	4	4	4	6	4	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	-	-	2	2	1	-	-	-
Male	-	1	-	-	-	-	1	1	1	-	-	-
Female	-	1	-	1	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
North Carolina												
All races, both sexes	10	21	28	25	46	66	122	131	98	43	2	-
Male	4	12	13	11	24	29	53	41	31	8	1	-
Female	6	9	15	14	22	37	69	90	67	35	1	-
White, both sexes	5	12	19	18	41	59	107	116	95	38	2	-
Male	3	7	10	10	23	25	49	39	31	8	1	-
Female	2	5	9	8	18	34	58	77	64	30	1	-
Black, both sexes	5	8	9	7	2	7	14	13	3	5	-	-
Male	1	5	3	1	1	4	3	2	-	-	-	-
Female	4	3	6	6	1	3	11	11	3	5	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	3	-	-	1	-	-	-	-
North Dakota												
All races, both sexes	1	1	2	1	2	3	6	15	13	7	-	-
Male	1	1	2	-	1	-	2	4	4	1	-	-
Female	-	-	-	1	1	3	4	11	9	6	-	-
White, both sexes	1	1	2	1	2	3	6	15	13	7	-	-
Male	1	1	2	-	1	-	2	4	4	1	-	-
Female	-	-	-	1	1	3	4	11	9	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Ohio												
All races, both sexes	13	22	25	33	60	89	166	218	163	64	9	-
Male	9	15	20	19	20	36	71	71	50	14	1	-
Female	4	7	5	14	40	53	95	147	113	50	8	-
White, both sexes	12	16	18	27	52	84	162	211	159	63	8	-
Male	8	10	15	16	19	33	68	69	49	14	1	-
Female	4	6	3	11	33	51	94	142	110	49	7	-
Black, both sexes	1	6	7	4	7	5	4	7	4	1	1	-
Male	1	5	5	2	1	3	3	2	1	-	-	-
Female	-	1	2	2	6	2	1	5	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	5	7	10	9	17	30	30	43	40	15	3	-
Male	2	3	5	1	11	11	10	10	12	4	1	-
Female	3	4	5	8	6	19	20	33	28	11	2	-
White, both sexes	4	5	10	9	15	29	29	40	38	14	3	-
Male	2	1	5	1	10	11	9	8	11	4	1	-
Female	2	4	5	8	5	18	20	32	27	10	2	-
Black, both sexes	1	1	-	-	1	1	1	2	-	1	-	-
Male	-	1	-	-	1	-	1	1	-	-	-	-
Female	1	-	-	-	-	1	-	1	-	1	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	1	2	-	-	-
Male	-	1	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Oregon												
All races, both sexes	5	7	6	13	14	48	86	131	128	46	3	-
Male	2	4	4	2	4	29	42	57	38	10	-	-
Female	3	3	2	11	10	19	44	74	90	36	3	-
White, both sexes	5	7	6	12	14	48	86	127	125	46	3	-
Male	2	4	4	2	4	29	42	55	38	10	-	-
Female	3	3	2	10	10	19	44	72	87	36	3	-
Black, both sexes	-	-	-	-	-	-	-	2	2	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Pennsylvania												
All races, both sexes	24	29	32	50	85	153	310	364	303	108	11	-
Male	16	20	21	23	44	67	116	120	72	11	3	-
Female	8	9	11	27	41	86	194	244	231	97	8	-
White, both sexes	17	21	28	47	80	144	301	355	302	100	11	-
Male	11	15	18	22	42	63	111	119	72	11	3	-
Female	6	6	10	25	38	81	190	236	230	89	8	-
Black, both sexes	7	8	3	3	4	9	8	9	1	8	-	-
Male	5	5	3	1	2	4	5	1	-	-	-	-
Female	2	3	-	2	2	5	3	8	1	8	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Rhode Island												
All races, both sexes	1	3	1	2	7	3	13	15	20	11	1	-
Male	1	2	-	2	4	-	8	6	6	1	-	-
Female	-	1	1	-	3	3	5	9	14	10	1	-
White, both sexes	-	3	1	2	7	3	13	14	20	11	1	-
Male	-	2	-	2	4	-	8	5	6	1	-	-
Female	-	1	1	-	3	3	5	9	14	10	1	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	6	10	8	19	28	41	61	69	41	9	3	-
Male	5	6	5	9	11	14	19	20	11	3	1	-
Female	1	4	3	10	17	27	42	49	30	6	2	-
White, both sexes	4	7	3	14	22	34	50	57	34	8	3	-
Male	3	5	2	5	9	12	18	16	10	3	1	-
Female	1	2	1	9	13	22	32	41	24	5	2	-
Black, both sexes	2	3	5	5	5	7	10	11	7	1	-	-
Male	2	1	3	4	1	2	1	3	1	-	-	-
Female	-	2	2	1	4	5	9	8	6	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
South Dakota												
All races, both sexes	1	-	2	2	2	7	9	17	16	9	1	-
Male	-	-	1	2	1	4	5	7	5	-	-	-
Female	1	-	1	-	1	3	4	10	11	9	1	-
White, both sexes	1	-	1	2	2	7	9	17	16	9	1	-
Male	-	-	-	2	1	4	5	7	5	-	-	-
Female	1	-	1	-	1	3	4	10	11	9	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	4	12	18	22	15	38	84	74	52	15	3	-
Male	3	7	11	12	2	17	28	24	16	1	1	-
Female	1	5	7	10	13	21	56	50	36	14	2	-
White, both sexes	2	10	15	20	14	35	82	71	51	14	3	-
Male	2	5	10	12	2	16	28	23	16	1	1	-
Female	-	5	5	8	12	19	54	48	35	13	2	-
Black, both sexes	2	2	3	2	1	3	2	3	1	1	-	-
Male	1	2	1	-	-	1	-	1	-	-	-	-
Female	1	-	2	2	1	2	2	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Texas												
All races, both sexes	30	26	38	62	89	153	177	219	142	64	10	-
Male	15	14	22	25	43	73	70	67	42	17	1	-
Female	15	12	16	37	46	80	107	152	100	47	9	-
White, both sexes	23	23	33	56	80	142	170	211	139	60	9	-
Male	11	13	19	23	39	69	68	65	42	15	1	-
Female	12	10	14	33	41	73	102	146	97	45	8	-
Black, both sexes	7	3	3	6	9	10	7	7	3	4	1	-
Male	4	1	3	2	4	4	2	2	-	2	-	-
Female	3	2	-	4	5	6	5	5	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	2	-	-	1	-	1	-	-	-	-
Utah												
All races, both sexes	4	2	1	8	6	23	26	29	20	8	1	-
Male	3	1	1	2	2	8	8	7	6	1	-	-
Female	1	1	-	6	4	15	18	22	14	7	1	-
White, both sexes	4	2	1	8	6	22	26	29	20	8	1	-
Male	3	1	1	2	2	8	8	7	6	1	-	-
Female	1	1	-	6	4	14	18	22	14	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Vermont												
All races, both sexes	-	-	3	1	4	3	10	13	16	10	2	-
Male	-	-	3	1	2	3	2	9	4	5	-	-
Female	-	-	-	-	2	-	8	4	12	5	2	-
White, both sexes	-	-	3	1	4	3	10	13	16	10	2	-
Male	-	-	3	1	2	3	2	9	4	5	-	-
Female	-	-	-	-	2	-	8	4	12	5	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5	11	16	15	37	48	74	96	69	27	4	-
Male	2	8	9	5	19	21	25	31	22	8	-	-
Female	3	3	7	10	18	27	49	65	47	19	4	-
White, both sexes	4	8	13	12	32	44	62	93	65	24	3	-
Male	1	5	8	3	17	18	22	31	22	7	-	-
Female	3	3	5	9	15	26	40	62	43	17	3	-
Black, both sexes	1	3	3	3	4	3	10	2	4	3	1	-
Male	1	3	1	2	2	2	2	-	-	1	-	-
Female	-	-	2	1	2	1	8	2	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	2	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Washington												
All races, both sexes	8	11	13	19	26	65	130	195	137	53	13	-
Male	5	7	8	8	12	29	55	86	49	12	4	-
Female	3	4	5	11	14	36	75	109	88	41	9	-
White, both sexes	8	9	11	18	22	65	125	191	134	53	13	-
Male	5	6	7	8	9	29	53	83	49	12	4	-
Female	3	3	4	10	13	36	72	108	85	41	9	-
Black, both sexes	-	-	1	-	2	-	-	3	-	-	-	-
Male	-	-	-	-	1	-	-	2	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	2	-	-	1	-	1	-	1	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	-	4	1	2	-	-	-
Male	-	-	1	-	1	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	3	-	2	-	-	-
West Virginia												
All races, both sexes	3	5	7	7	7	14	30	34	22	5	-	-
Male	2	5	3	4	2	9	12	12	5	1	-	-
Female	1	-	4	3	5	5	18	22	17	4	-	-
White, both sexes	3	5	7	7	6	14	30	33	22	5	-	-
Male	2	5	3	4	1	9	12	12	5	1	-	-
Female	1	-	4	3	5	5	18	21	17	4	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of endocardium (I34-I38)												
Wisconsin												
All races, both sexes	3	6	12	17	25	64	92	154	115	42	7	-
Male	2	5	6	8	13	31	44	55	35	8	1	-
Female	1	1	6	9	12	33	48	99	80	34	6	-
White, both sexes	1	4	11	16	23	62	91	152	114	42	6	-
Male	1	4	5	8	11	31	44	53	35	8	1	-
Female	-	-	6	8	12	31	47	99	79	34	5	-
Black, both sexes	2	2	1	1	1	2	1	1	1	-	1	-
Male	1	1	1	-	1	-	-	1	-	-	-	-
Female	1	1	-	1	-	2	1	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	2	-	1	3	6	10	7	5	2	1	-
Male	-	2	-	1	1	4	3	3	2	1	-	-
Female	1	-	-	-	2	2	7	4	3	1	1	-
White, both sexes	1	2	-	1	3	6	10	7	5	2	1	-
Male	-	2	-	1	1	4	3	3	2	1	-	-
Female	1	-	-	-	2	2	7	4	3	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
United States												
All races, both sexes	2,605	-	1	1	4	6	9	17	19	29	41	63
Male	896	-	1	1	1	3	5	13	9	14	20	22
Female	1,709	-	-	-	3	3	4	4	10	15	21	41
White, both sexes	2,371	-	1	1	4	5	7	10	16	25	35	49
Male	823	-	1	1	1	3	4	7	7	12	18	19
Female	1,548	-	-	-	3	2	3	3	9	13	17	30
Black, both sexes	186	-	-	-	-	-	1	7	3	4	6	12
Male	56	-	-	-	-	-	-	6	2	2	2	3
Female	130	-	-	-	-	-	1	1	1	2	4	9
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	45	-	-	-	-	1	1	-	-	-	-	2
Male	16	-	-	-	-	-	1	-	-	-	-	-
Female	29	-	-	-	-	1	-	-	-	-	-	2
Alabama												
All races, both sexes	48	-	-	-	-	-	-	-	2	1	2	1
Male	16	-	-	-	-	-	-	-	2	-	-	1
Female	32	-	-	-	-	-	-	-	-	1	2	-
White, both sexes	42	-	-	-	-	-	-	-	2	1	2	1
Male	14	-	-	-	-	-	-	-	2	-	-	1
Female	28	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Alaska												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Arkansas												
All races, both sexes	19	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	18	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	236	-	-	-	-	-	-	-	-	4	6	5
Male	86	-	-	-	-	-	-	-	-	1	3	2
Female	150	-	-	-	-	-	-	-	-	3	3	3
White, both sexes	211	-	-	-	-	-	-	-	-	4	5	4
Male	75	-	-	-	-	-	-	-	-	1	2	2
Female	136	-	-	-	-	-	-	-	-	3	3	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Colorado												
All races, both sexes	34	-	-	-	-	-	-	-	-	3	-	-
Male	16	-	-	-	-	-	-	-	-	3	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	-	-	-	3	-	-
Male	15	-	-	-	-	-	-	-	-	3	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	33	-	-	-	-	1	-	-	-	-	-	-
Male	14	-	-	-	-	1	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	1	-	-	-	-	-	-
Male	13	-	-	-	-	1	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Delaware												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Florida												
All races, both sexes	168	-	-	-	-	-	1	2	3	1	3	6
Male	55	-	-	-	-	-	1	2	1	-	2	1
Female	113	-	-	-	-	-	-	-	2	1	1	5
White, both sexes	146	-	-	-	-	-	1	1	2	1	3	4
Male	50	-	-	-	-	-	1	1	-	-	2	-
Female	96	-	-	-	-	-	-	-	2	1	1	4
Black, both sexes	22	-	-	-	-	-	-	1	1	-	-	2
Male	5	-	-	-	-	-	-	1	1	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	46	-	-	-	-	-	-	-	2	-	-	1
Male	16	-	-	-	-	-	-	-	2	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	37	-	-	-	-	-	-	-	1	-	-	-
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Hawaii												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Illinois												
All races, both sexes	83	-	-	-	2	1	-	1	2	3	1	2
Male	24	-	-	-	-	-	-	1	-	1	-	1
Female	59	-	-	-	2	1	-	-	2	2	1	1
White, both sexes	67	-	-	-	2	1	-	1	2	2	1	2
Male	18	-	-	-	-	-	-	1	-	1	-	1
Female	49	-	-	-	2	1	-	-	2	1	1	1
Black, both sexes	16	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	60	-	-	-	-	-	-	-	-	1	-	5
Male	17	-	-	-	-	-	-	-	-	-	-	2
Female	43	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	58	-	-	-	-	-	-	-	-	1	-	5
Male	17	-	-	-	-	-	-	-	-	-	-	2
Female	41	-	-	-	-	-	-	-	-	1	-	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Iowa												
All races, both sexes	49	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	37	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	47	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Kentucky												
All races, both sexes	31	-	-	-	-	-	-	-	-	1	2	-
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	22	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	29	-	-	-	-	-	-	-	-	1	2	-
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	21	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	22	-	-	-	-	-	-	1	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Maine												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	40	-	-	-	-	-	-	-	1	1	-	-
Male	15	-	-	-	-	-	-	-	1	1	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	1	1	-	-
Male	14	-	-	-	-	-	-	-	1	1	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Massachusetts												
All races, both sexes	58	-	-	1	-	1	1	-	-	-	-	-
Male	19	-	-	1	-	1	-	-	-	-	-	-
Female	39	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	57	-	-	1	-	1	1	-	-	-	-	-
Male	19	-	-	1	-	1	-	-	-	-	-	-
Female	38	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	97	-	-	-	-	1	-	-	-	1	6	3
Male	38	-	-	-	-	1	-	-	-	-	2	1
Female	59	-	-	-	-	-	-	-	-	1	4	2
White, both sexes	90	-	-	-	-	1	-	-	-	1	5	1
Male	35	-	-	-	-	1	-	-	-	-	2	-
Female	55	-	-	-	-	-	-	-	-	1	3	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Minnesota												
All races, both sexes	68	-	-	-	-	-	-	-	-	1	-	4
Male	21	-	-	-	-	-	-	-	-	1	-	1
Female	47	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	67	-	-	-	-	-	-	-	-	1	-	4
Male	20	-	-	-	-	-	-	-	-	1	-	1
Female	47	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	20	-	-	-	-	-	-	1	-	-	-	1
Male	8	-	-	-	-	-	-	1	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Missouri												
All races, both sexes	69	-	-	-	-	-	-	-	1	3	1	-
Male	21	-	-	-	-	-	-	-	1	-	-	-
Female	48	-	-	-	-	-	-	-	-	3	1	-
White, both sexes	62	-	-	-	-	-	-	-	1	3	1	-
Male	18	-	-	-	-	-	-	-	1	-	-	-
Female	44	-	-	-	-	-	-	-	-	3	1	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Nebraska												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	14	-	-	-	-	1	-	1	-	1	-	1
Male	7	-	-	-	-	-	-	1	-	1	-	-
Female	7	-	-	-	-	1	-	-	-	-	-	1
White, both sexes	11	-	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
New Hampshire												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	85	-	-	-	-	1	-	1	-	-	1	3
Male	28	-	-	-	-	-	-	-	-	-	-	1
Female	57	-	-	-	-	1	-	1	-	-	1	2
White, both sexes	80	-	-	-	-	1	-	1	-	-	1	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	53	-	-	-	-	1	-	1	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	3
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
New Mexico												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	142	-	-	-	-	-	-	2	2	1	2	4
Male	57	-	-	-	-	-	-	2	-	1	2	2
Female	85	-	-	-	-	-	-	-	2	-	-	2
White, both sexes	133	-	-	-	-	-	-	2	2	-	1	3
Male	52	-	-	-	-	-	-	2	-	-	1	2
Female	81	-	-	-	-	-	-	-	2	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	1	1	1
Male	3	-	-	-	-	-	-	-	-	1	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
North Carolina												
All races, both sexes	75	-	-	-	-	-	-	1	-	-	1	1
Male	23	-	-	-	-	-	-	1	-	-	1	-
Female	52	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	61	-	-	-	-	-	-	1	-	-	1	-
Male	21	-	-	-	-	-	-	1	-	-	1	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Ohio												
All races, both sexes	117	-	1	-	1	-	2	1	1	3	5	2
Male	51	-	1	-	1	-	2	1	1	2	2	1
Female	66	-	-	-	-	-	-	-	-	1	3	1
White, both sexes	105	-	1	-	1	-	2	1	1	2	3	2
Male	49	-	1	-	1	-	2	1	1	2	2	1
Female	56	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	12	-	-	-	-	-	-	-	-	1	2	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Oregon												
All races, both sexes	63	-	-	-	-	-	-	1	1	-	-	-
Male	25	-	-	-	-	-	-	1	-	-	-	-
Female	38	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	62	-	-	-	-	-	-	-	1	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	38	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	198	-	-	-	-	-	1	-	-	-	4	3
Male	59	-	-	-	-	-	-	-	-	-	3	1
Female	139	-	-	-	-	-	1	-	-	-	1	2
White, both sexes	187	-	-	-	-	-	1	-	-	-	4	2
Male	58	-	-	-	-	-	-	-	-	-	3	1
Female	129	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Rhode Island												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	52	-	-	-	-	-	-	-	1	-	1	4
Male	17	-	-	-	-	-	-	-	-	-	-	2
Female	35	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	39	-	-	-	-	-	-	-	-	-	-	3
Male	13	-	-	-	-	-	-	-	-	-	-	2
Female	26	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	12	-	-	-	-	-	-	-	1	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
South Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	51	-	-	-	-	-	-	1	-	1	1	2
Male	13	-	-	-	-	-	-	1	-	1	1	-
Female	38	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	48	-	-	-	-	-	-	-	-	1	1	2
Male	12	-	-	-	-	-	-	-	-	1	1	-
Female	36	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Texas												
All races, both sexes	145	-	-	-	-	-	-	1	1	1	1	2
Male	43	-	-	-	-	-	-	-	-	1	1	-
Female	102	-	-	-	-	-	-	1	1	-	-	2
White, both sexes	135	-	-	-	-	-	-	1	1	-	1	2
Male	40	-	-	-	-	-	-	-	-	-	1	-
Female	95	-	-	-	-	-	-	1	1	-	-	2
Black, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	14	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	1	-	-	-	-	-
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Vermont												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	70	-	-	-	-	-	3	2	2	1	2	2
Male	27	-	-	-	-	-	1	2	1	-	2	1
Female	43	-	-	-	-	-	2	-	1	1	-	1
White, both sexes	61	-	-	-	-	-	1	1	2	1	2	2
Male	23	-	-	-	-	-	-	1	1	-	2	1
Female	38	-	-	-	-	-	1	-	1	1	-	1
Black, both sexes	8	-	-	-	-	-	1	1	-	-	-	-
Male	3	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Washington												
All races, both sexes	92	-	-	-	-	-	-	1	-	-	-	4
Male	40	-	-	-	-	-	-	-	-	-	-	2
Female	52	-	-	-	-	-	-	1	-	-	-	2
White, both sexes	89	-	-	-	-	-	-	1	-	-	-	4
Male	39	-	-	-	-	-	-	-	-	-	-	2
Female	50	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic mitral valve disorders (I34)												
Wisconsin												
All races, both sexes	55	-	-	-	1	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	1	-	-	-	-	-	1	-
White, both sexes	55	-	-	-	1	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	42	-	-	-	1	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
United States												
All races, both sexes	79	86	101	129	183	317	480	538	358	128	15	1
Male	41	43	46	44	68	124	159	165	89	25	3	-
Female	38	43	55	85	115	193	321	373	269	103	12	1
White, both sexes	55	68	78	114	161	297	451	515	342	122	14	1
Male	28	34	39	41	64	115	153	160	89	24	3	-
Female	27	34	39	73	97	182	298	355	253	98	11	1
Black, both sexes	22	14	19	15	15	15	23	15	11	4	-	-
Male	11	5	6	3	2	7	5	2	-	-	-	-
Female	11	9	13	12	13	8	18	13	11	4	-	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	3	-	7	4	6	7	5	2	1	-
Male	2	4	1	-	2	2	1	2	-	1	-	-
Female	-	-	2	-	5	2	5	5	5	1	1	-
Alabama												
All races, both sexes	2	1	2	5	2	8	11	5	3	3	-	-
Male	1	-	1	3	1	2	4	1	-	-	-	-
Female	1	1	1	2	1	6	7	4	3	3	-	-
White, both sexes	-	1	2	5	2	7	9	5	2	3	-	-
Male	-	-	1	3	1	1	4	1	-	-	-	-
Female	-	1	1	2	1	6	5	4	2	3	-	-
Black, both sexes	2	-	-	-	-	1	2	-	1	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Alaska												
All races, both sexes	1	-	-	-	1	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	1	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	-	1	1	5	5	2	9	2	-	-	-
Male	2	-	-	1	2	-	-	7	-	-	-	-
Female	-	-	1	-	3	5	2	2	2	-	-	-
White, both sexes	2	-	1	1	5	5	2	8	2	-	-	-
Male	2	-	-	1	2	-	-	6	-	-	-	-
Female	-	-	1	-	3	5	2	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Arkansas												
All races, both sexes	-	2	-	2	1	2	2	5	3	-	-	-
Male	-	1	-	-	-	-	1	2	1	-	-	-
Female	-	1	-	2	1	2	1	3	2	-	-	-
White, both sexes	-	2	-	2	1	2	2	4	3	-	-	-
Male	-	1	-	-	-	-	1	2	1	-	-	-
Female	-	1	-	2	1	2	1	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	8	10	8	11	15	24	45	50	36	11	3	-
Male	4	7	3	3	8	11	20	14	5	5	-	-
Female	4	3	5	8	7	13	25	36	31	6	3	-
White, both sexes	5	7	7	11	13	21	41	47	33	11	2	-
Male	2	4	3	3	7	9	18	14	5	5	-	-
Female	3	3	4	8	6	12	23	33	28	6	2	-
Black, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	1	-	2	2	3	3	3	-	1	-
Male	2	3	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	-	1	1	2	3	3	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Colorado												
All races, both sexes	2	-	3	4	-	4	3	9	5	1	-	-
Male	-	-	1	2	-	4	-	4	1	1	-	-
Female	2	-	2	2	-	-	3	5	4	-	-	-
White, both sexes	2	-	3	4	-	3	3	9	5	1	-	-
Male	-	-	1	2	-	3	-	4	1	1	-	-
Female	2	-	2	2	-	-	3	5	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	2	2	-	1	1	3	11	7	3	1	-
Male	1	2	2	-	-	1	1	3	2	1	-	-
Female	-	-	-	-	1	-	2	8	5	2	1	-
White, both sexes	1	1	2	-	1	1	3	10	7	3	1	-
Male	1	1	2	-	-	1	1	3	2	1	-	-
Female	-	-	-	-	1	-	2	7	5	2	1	-
Black, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Delaware												
All races, both sexes	-	-	-	-	2	2	2	2	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	2	1	2	1	-	-	-	-
White, both sexes	-	-	-	-	2	2	2	2	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	-	2	1	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	1	3	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Florida												
All races, both sexes	6	5	8	10	14	23	24	25	29	7	1	-
Male	2	2	2	4	5	8	8	7	10	-	-	-
Female	4	3	6	6	9	15	16	18	19	7	1	-
White, both sexes	6	4	3	8	13	21	23	24	26	5	1	-
Male	2	2	1	4	5	7	8	7	10	-	-	-
Female	4	2	2	4	8	14	15	17	16	5	1	-
Black, both sexes	-	1	5	2	1	2	1	1	3	2	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	1	4	2	1	1	1	1	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	5	1	3	4	3	8	7	6	3	-	-
Male	1	2	-	1	3	1	1	3	-	2	-	-
Female	2	3	1	2	1	2	7	4	6	1	-	-
White, both sexes	1	3	1	2	3	3	8	7	5	3	-	-
Male	-	1	-	-	3	1	1	3	-	2	-	-
Female	1	2	1	2	-	2	7	4	5	1	-	-
Black, both sexes	2	2	-	1	1	-	-	-	1	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	1	1	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Hawaii												
All races, both sexes	-	-	1	-	4	1	1	3	4	2	-	-
Male	-	-	1	-	-	-	1	1	1	1	-	-
Female	-	-	-	-	4	1	-	2	3	1	-	-
White, both sexes	-	-	-	-	2	-	1	-	2	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	2	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	1	-	3	2	2	-	-
Male	-	-	1	-	-	-	-	1	-	1	-	-
Female	-	-	-	-	2	1	-	2	2	1	-	-
Idaho												
All races, both sexes	1	-	-	1	-	-	3	3	3	1	-	-
Male	1	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	1	-	-	3	2	2	1	-	-
White, both sexes	1	-	-	1	-	-	3	3	3	1	-	-
Male	1	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	1	-	-	3	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Illinois												
All races, both sexes	3	1	5	4	3	7	14	18	12	3	1	-
Male	2	-	4	2	3	1	4	3	1	1	-	-
Female	1	1	1	2	-	6	10	15	11	2	1	-
White, both sexes	-	-	2	3	2	6	11	17	11	3	1	-
Male	-	-	2	2	2	1	3	3	1	1	-	-
Female	-	-	-	1	-	5	8	14	10	2	1	-
Black, both sexes	3	1	3	1	1	1	3	1	1	-	-	-
Male	2	-	2	-	1	-	1	-	-	-	-	-
Female	1	1	1	1	-	1	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	-	3	5	5	4	8	13	5	6	5	-	-
Male	-	2	2	1	1	2	5	1	1	-	-	-
Female	-	1	3	4	3	6	8	4	5	5	-	-
White, both sexes	-	3	4	4	4	8	13	5	6	5	-	-
Male	-	2	2	1	1	2	5	1	1	-	-	-
Female	-	1	2	3	3	6	8	4	5	5	-	-
Black, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Iowa												
All races, both sexes	-	1	2	4	6	5	8	8	10	4	-	-
Male	-	-	1	2	-	4	3	-	2	-	-	-
Female	-	1	1	2	6	1	5	8	8	4	-	-
White, both sexes	-	1	1	3	6	5	8	8	10	4	-	-
Male	-	-	1	2	-	4	3	-	2	-	-	-
Female	-	1	-	1	6	1	5	8	8	4	-	-
Black, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	1	1	1	1	5	1	4	2	4	-	-
Male	-	-	-	1	-	2	-	1	-	-	-	-
Female	1	1	1	-	1	3	1	3	2	4	-	-
White, both sexes	-	1	1	1	1	5	1	4	2	4	-	-
Male	-	-	-	1	-	2	-	1	-	-	-	-
Female	-	1	1	-	1	3	1	3	2	4	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Kentucky												
All races, both sexes	2	1	-	2	2	2	6	8	3	2	-	-
Male	2	1	-	1	1	-	1	1	1	-	-	-
Female	-	-	-	1	1	2	5	7	2	2	-	-
White, both sexes	1	1	-	2	2	2	6	8	2	2	-	-
Male	1	1	-	1	1	-	1	1	1	-	-	-
Female	-	-	-	1	1	2	5	7	1	2	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3	1	2	1	3	5	4	1	1	-	-	-
Male	2	-	1	-	1	2	-	1	-	-	-	-
Female	1	1	1	1	2	3	4	-	1	-	-	-
White, both sexes	1	-	1	-	2	3	4	1	1	-	-	-
Male	1	-	-	-	1	2	-	1	-	-	-	-
Female	-	-	1	-	1	1	4	-	1	-	-	-
Black, both sexes	2	1	1	1	1	2	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	1	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Maine												
All races, both sexes	-	1	1	1	2	1	3	5	5	1	-	-
Male	-	1	-	-	2	-	-	2	2	-	-	-
Female	-	-	1	1	-	1	3	3	3	1	-	-
White, both sexes	-	1	1	1	2	1	3	5	5	1	-	-
Male	-	1	-	-	2	-	-	2	2	-	-	-
Female	-	-	1	1	-	1	3	3	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	1	3	1	-	4	12	7	6	1	-	-
Male	1	1	1	-	-	1	5	3	1	-	-	-
Female	2	-	2	1	-	3	7	4	5	1	-	-
White, both sexes	2	1	3	1	-	4	10	5	6	1	-	-
Male	1	1	1	-	-	1	4	3	1	-	-	-
Female	1	-	2	1	-	3	6	2	5	1	-	-
Black, both sexes	1	-	-	-	-	-	2	2	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Massachusetts												
All races, both sexes	-	-	1	2	6	6	14	9	12	4	1	-
Male	-	-	1	1	2	4	1	4	4	-	-	-
Female	-	-	-	1	4	2	13	5	8	4	1	-
White, both sexes	-	-	1	2	5	6	14	9	12	4	1	-
Male	-	-	1	1	2	4	1	4	4	-	-	-
Female	-	-	-	1	3	2	13	5	8	4	1	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	4	3	5	3	9	14	16	21	9	2	-	-
Male	3	-	4	2	6	4	6	6	2	1	-	-
Female	1	3	1	1	3	10	10	15	7	1	-	-
White, both sexes	3	3	5	3	7	14	16	20	9	2	-	-
Male	2	-	4	2	5	4	6	6	2	1	-	-
Female	1	3	1	1	2	10	10	14	7	1	-	-
Black, both sexes	1	-	-	-	2	-	-	1	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Minnesota												
All races, both sexes	1	3	1	-	4	7	15	16	14	1	1	-
Male	1	-	-	-	1	4	7	3	3	-	-	-
Female	-	3	1	-	3	3	8	13	11	1	1	-
White, both sexes	-	3	1	-	4	7	15	16	14	1	1	-
Male	-	-	-	-	1	4	7	3	3	-	-	-
Female	-	3	1	-	3	3	8	13	11	1	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1	-	2	2	4	1	1	3	2	2	-	-
Male	-	-	-	1	2	1	-	1	1	-	-	-
Female	1	-	2	1	2	-	1	2	1	2	-	-
White, both sexes	-	-	2	1	4	1	1	2	2	2	-	-
Male	-	-	-	1	2	1	-	1	1	-	-	-
Female	-	-	2	-	2	-	1	1	1	2	-	-
Black, both sexes	1	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Missouri												
All races, both sexes	2	1	4	2	1	10	15	14	12	3	-	-
Male	2	1	1	1	-	4	2	6	2	1	-	-
Female	-	-	3	1	1	6	13	8	10	2	-	-
White, both sexes	-	1	2	2	1	9	14	14	11	3	-	-
Male	-	1	1	1	-	3	2	6	2	1	-	-
Female	-	-	1	1	1	6	12	8	9	2	-	-
Black, both sexes	2	-	2	-	-	1	1	-	1	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	2	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	1	1	4	2	1	1	-
Male	-	-	-	-	-	1	-	-	2	-	1	-
Female	1	-	-	-	-	-	1	4	-	1	-	-
White, both sexes	1	-	-	-	-	1	1	4	2	1	1	-
Male	-	-	-	-	-	1	-	-	2	-	1	-
Female	1	-	-	-	-	-	1	4	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Nebraska												
All races, both sexes	-	1	1	-	-	2	-	3	2	3	-	-
Male	-	1	-	-	-	1	-	2	-	-	-	-
Female	-	-	1	-	-	1	-	1	2	3	-	-
White, both sexes	-	1	1	-	-	2	-	2	2	3	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	1	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	-	-	2	2	4	-	-	1	-	-
Male	-	1	-	-	1	1	2	-	-	-	-	-
Female	-	-	-	-	1	1	2	-	-	1	-	-
White, both sexes	-	1	-	-	2	1	4	-	-	1	-	-
Male	-	1	-	-	1	1	2	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
New Hampshire												
All races, both sexes	1	-	-	-	-	1	2	5	2	1	-	-
Male	1	-	-	-	-	-	1	2	1	-	-	-
Female	-	-	-	-	-	1	1	3	1	1	-	-
White, both sexes	1	-	-	-	-	1	2	5	2	1	-	-
Male	1	-	-	-	-	-	1	2	1	-	-	-
Female	-	-	-	-	-	1	1	3	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	5	4	7	5	10	14	14	17	2	-	-
Male	-	2	3	1	1	4	8	6	2	-	-	-
Female	1	3	1	6	4	6	6	8	15	2	-	-
White, both sexes	1	5	3	7	5	10	14	14	16	2	-	-
Male	-	2	3	1	1	4	8	6	2	-	-	-
Female	1	3	-	6	4	6	6	8	14	2	-	-
Black, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
New Mexico												
All races, both sexes	-	1	2	-	-	1	1	3	1	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	2	-	-	-	1	2	1	-	-	-
White, both sexes	-	1	1	-	-	1	1	3	1	-	-	-
Male	-	1	-	-	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	-	1	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2	5	2	5	6	22	29	29	20	10	-	1
Male	2	4	2	1	3	12	9	13	3	1	-	-
Female	-	1	-	4	3	10	20	16	17	9	-	1
White, both sexes	2	3	2	5	5	22	27	28	20	10	-	1
Male	2	3	2	1	3	12	8	12	3	1	-	-
Female	-	-	-	4	2	10	19	16	17	9	-	1
Black, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
North Carolina												
All races, both sexes	3	6	4	5	11	9	13	8	9	4	-	-
Male	1	3	2	-	4	3	5	1	1	1	-	-
Female	2	3	2	5	7	6	8	7	8	3	-	-
White, both sexes	2	3	3	2	9	9	11	8	9	3	-	-
Male	1	2	1	-	4	3	5	1	1	1	-	-
Female	1	1	2	2	5	6	6	7	8	2	-	-
Black, both sexes	1	3	1	3	1	-	2	-	-	1	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	1	2	-	3	1	-	2	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	1	2	3	3	2	-	-
Male	-	-	-	-	-	-	1	2	2	-	-	-
Female	-	-	-	-	-	1	1	1	1	2	-	-
White, both sexes	-	-	-	-	-	1	2	3	3	2	-	-
Male	-	-	-	-	-	-	1	2	2	-	-	-
Female	-	-	-	-	-	1	1	1	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Ohio												
All races, both sexes	4	1	2	5	9	11	17	30	17	5	-	-
Male	2	1	1	4	-	5	5	10	9	3	-	-
Female	2	-	1	1	9	6	12	20	8	2	-	-
White, both sexes	4	-	1	4	5	11	16	29	17	5	-	-
Male	2	-	1	3	-	5	5	10	9	3	-	-
Female	2	-	-	1	5	6	11	19	8	2	-	-
Black, both sexes	-	1	1	1	4	-	1	1	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	4	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	-	2	3	2	8	4	2	2	-	-	-
Male	1	-	2	1	1	1	1	-	1	-	-	-
Female	1	-	-	2	1	7	3	2	1	-	-	-
White, both sexes	2	-	2	3	2	8	4	2	2	-	-	-
Male	1	-	2	1	1	1	1	-	1	-	-	-
Female	1	-	-	2	1	7	3	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Oregon												
All races, both sexes	3	1	2	2	1	8	10	19	7	8	-	-
Male	1	-	1	-	-	6	4	8	2	2	-	-
Female	2	1	1	2	1	2	6	11	5	6	-	-
White, both sexes	3	1	2	2	1	8	10	19	7	8	-	-
Male	1	-	1	-	-	6	4	8	2	2	-	-
Female	2	1	1	2	1	2	6	11	5	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	4	6	6	11	14	22	46	52	23	5	1	-
Male	1	3	3	2	7	8	15	11	4	1	-	-
Female	3	3	3	9	7	14	31	41	19	4	1	-
White, both sexes	3	5	6	11	13	21	43	50	23	4	1	-
Male	1	3	3	2	7	8	14	11	4	1	-	-
Female	2	2	3	9	6	13	29	39	19	3	1	-
Black, both sexes	1	1	-	-	-	1	2	2	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	1	-	-	-	1	1	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Rhode Island												
All races, both sexes	-	2	-	-	3	-	4	3	1	-	-	-
Male	-	1	-	-	3	-	-	1	1	-	-	-
Female	-	1	-	-	-	-	4	2	-	-	-	-
White, both sexes	-	2	-	-	3	-	4	3	1	-	-	-
Male	-	1	-	-	3	-	-	1	1	-	-	-
Female	-	1	-	-	-	-	4	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1	2	-	3	3	10	14	10	3	-	-	-
Male	1	1	-	2	1	2	6	1	1	-	-	-
Female	-	1	-	1	2	8	8	9	2	-	-	-
White, both sexes	-	-	-	2	2	9	12	9	2	-	-	-
Male	-	-	-	1	-	2	6	1	1	-	-	-
Female	-	-	-	1	2	7	6	8	1	-	-	-
Black, both sexes	1	2	-	1	-	1	2	1	1	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
South Dakota												
All races, both sexes	-	-	1	-	-	2	-	3	3	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	1	-	-	2	-	2	2	1	-	-
White, both sexes	-	-	1	-	-	2	-	3	3	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	1	-	-	2	-	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1	3	6	2	2	3	11	8	6	3	1	-
Male	1	-	2	-	1	1	2	1	2	-	-	-
Female	-	3	4	2	1	2	9	7	4	3	1	-
White, both sexes	1	3	6	1	2	3	10	8	6	3	1	-
Male	1	-	2	-	1	1	2	1	2	-	-	-
Female	-	3	4	1	1	2	8	7	4	3	1	-
Black, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Texas												
All races, both sexes	7	6	5	9	20	18	30	25	13	6	-	-
Male	3	2	2	2	7	6	11	5	2	1	-	-
Female	4	4	3	7	13	12	19	20	11	5	-	-
White, both sexes	6	6	4	8	18	16	28	25	13	6	-	-
Male	3	2	2	2	7	4	11	5	2	1	-	-
Female	3	4	2	6	11	12	17	20	11	5	-	-
Black, both sexes	1	-	-	1	2	2	2	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	1	-	-	1	2	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	3	1	3	2	3	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	3	1	2	2	2	-	-	-	-
White, both sexes	-	-	-	3	1	2	2	3	1	-	-	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	-	3	1	1	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Vermont												
All races, both sexes	-	-	-	1	-	-	3	3	1	1	1	-
Male	-	-	-	1	-	-	1	2	1	1	-	-
Female	-	-	-	-	-	-	2	1	-	-	1	-
White, both sexes	-	-	-	1	-	-	3	3	1	1	1	-
Male	-	-	-	1	-	-	1	2	1	1	-	-
Female	-	-	-	-	-	-	2	1	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1	1	3	-	4	10	12	14	9	4	-	-
Male	-	-	2	-	-	4	5	5	3	1	-	-
Female	1	1	1	-	4	6	7	9	6	3	-	-
White, both sexes	1	1	1	-	4	8	12	13	8	4	-	-
Male	-	-	1	-	-	3	5	5	3	1	-	-
Female	1	1	-	-	4	5	7	8	5	3	-	-
Black, both sexes	-	-	2	-	-	2	-	1	1	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Washington												
All races, both sexes	1	-	1	5	3	14	18	24	13	6	2	-
Male	-	-	-	3	1	7	7	13	5	1	1	-
Female	1	-	1	2	2	7	11	11	8	5	1	-
White, both sexes	1	-	-	5	3	14	17	23	13	6	2	-
Male	-	-	-	3	1	7	7	12	5	1	1	-
Female	1	-	-	2	2	7	10	11	8	5	1	-
Black, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
West Virginia												
All races, both sexes	-	1	1	1	-	4	2	6	2	-	-	-
Male	-	1	1	1	-	1	-	2	-	-	-	-
Female	-	-	-	-	-	3	2	4	2	-	-	-
White, both sexes	-	1	1	1	-	4	2	6	2	-	-	-
Male	-	1	1	1	-	1	-	2	-	-	-	-
Female	-	-	-	-	-	3	2	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic mitral valve disorders (I34)												
Wisconsin												
All races, both sexes	1	2	1	2	2	5	11	19	7	2	1	-
Male	1	2	-	-	-	1	5	1	2	-	1	-
Female	-	-	1	2	2	4	6	18	5	2	-	-
White, both sexes	1	2	1	2	2	5	11	19	7	2	1	-
Male	1	2	-	-	-	1	5	1	2	-	1	-
Female	-	-	1	2	2	4	6	18	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
United States												
All races, both sexes	13,137	2	2	-	4	4	10	12	27	26	53	95
Male	5,095	1	2	-	3	4	7	11	19	19	41	65
Female	8,042	1	-	-	1	-	3	1	8	7	12	30
White, both sexes	12,397	1	1	-	3	3	9	10	20	18	44	72
Male	4,793	-	1	-	2	3	6	9	16	16	33	51
Female	7,604	1	-	-	1	-	3	1	4	2	11	21
Black, both sexes	572	1	-	-	1	1	1	1	6	8	6	18
Male	242	1	-	-	1	1	1	1	3	3	5	13
Female	330	-	-	-	-	-	-	-	3	5	1	5
American Indian 1/, both sexes	35	-	1	-	-	-	-	-	1	-	1	2
Male	20	-	1	-	-	-	-	-	-	-	1	1
Female	15	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	133	-	-	-	-	-	-	1	-	-	2	3
Male	40	-	-	-	-	-	-	1	-	-	2	-
Female	93	-	-	-	-	-	-	-	-	-	-	3
Alabama												
All races, both sexes	162	-	-	-	-	-	-	-	-	1	-	3
Male	62	-	-	-	-	-	-	-	-	1	-	2
Female	100	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	140	-	-	-	-	-	-	-	-	1	-	1
Male	52	-	-	-	-	-	-	-	-	1	-	-
Female	88	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Alaska												
All races, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	179	-	-	-	-	-	-	1	1	-	1	-
Male	76	-	-	-	-	-	-	1	1	-	1	-
Female	103	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	173	-	-	-	-	-	-	1	-	-	1	-
Male	72	-	-	-	-	-	-	1	-	-	1	-
Female	101	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Arkansas												
All races, both sexes	94	-	-	-	-	-	-	-	-	-	1	1
Male	31	-	-	-	-	-	-	-	-	-	1	1
Female	63	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	89	-	-	-	-	-	-	-	-	-	1	1
Male	30	-	-	-	-	-	-	-	-	-	1	1
Female	59	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,515	-	-	-	1	1	1	2	5	3	6	11
Male	602	-	-	-	1	1	1	2	4	3	6	5
Female	913	-	-	-	-	-	-	-	1	-	-	6
White, both sexes	1,406	-	-	-	-	1	1	2	5	3	6	9
Male	561	-	-	-	-	1	1	2	4	3	6	4
Female	845	-	-	-	-	-	-	-	1	-	-	5
Black, both sexes	47	-	-	-	1	-	-	-	-	-	-	1
Male	24	-	-	-	1	-	-	-	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	60	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Colorado												
All races, both sexes	169	-	-	-	-	-	-	-	-	1	1	-
Male	79	-	-	-	-	-	-	-	-	-	1	-
Female	90	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	165	-	-	-	-	-	-	-	-	-	-	-
Male	77	-	-	-	-	-	-	-	-	-	-	-
Female	88	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	1	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	196	-	-	-	-	-	-	-	-	-	-	2
Male	57	-	-	-	-	-	-	-	-	-	-	2
Female	139	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	193	-	-	-	-	-	-	-	-	-	-	2
Male	57	-	-	-	-	-	-	-	-	-	-	2
Female	136	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Delaware												
All races, both sexes	45	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Florida												
All races, both sexes	850	-	1	-	-	-	1	-	1	2	1	7
Male	368	-	1	-	-	-	-	-	-	1	1	5
Female	482	-	-	-	-	-	1	-	1	1	-	2
White, both sexes	822	-	-	-	-	-	1	-	-	1	1	6
Male	355	-	-	-	-	-	-	-	-	1	1	4
Female	467	-	-	-	-	-	1	-	-	-	-	2
Black, both sexes	27	-	-	-	-	-	-	-	1	1	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	246	-	-	-	-	-	1	1	2	1	-	3
Male	106	-	-	-	-	-	1	1	1	1	-	2
Female	140	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	207	-	-	-	-	-	1	1	1	1	-	2
Male	89	-	-	-	-	-	1	1	1	1	-	1
Female	118	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	39	-	-	-	-	-	-	-	1	-	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Hawaii												
All races, both sexes	60	-	-	-	-	-	-	-	-	-	2	1
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	41	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	37	-	-	-	-	-	-	-	-	-	1	1
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	91	-	-	-	-	-	-	-	-	1	-	-
Male	44	-	-	-	-	-	-	-	-	1	-	-
Female	47	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	88	-	-	-	-	-	-	-	-	1	-	-
Male	42	-	-	-	-	-	-	-	-	1	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Illinois												
All races, both sexes	504	-	-	-	-	-	-	-	-	2	2	-
Male	190	-	-	-	-	-	-	-	-	2	1	-
Female	314	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	475	-	-	-	-	-	-	-	-	2	1	-
Male	172	-	-	-	-	-	-	-	-	2	1	-
Female	303	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	27	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	300	-	-	-	-	-	1	-	-	-	1	4
Male	121	-	-	-	-	-	-	-	-	-	-	3
Female	179	-	-	-	-	-	1	-	-	-	1	1
White, both sexes	285	-	-	-	-	-	1	-	-	-	1	3
Male	115	-	-	-	-	-	-	-	-	-	-	2
Female	170	-	-	-	-	-	1	-	-	-	1	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Iowa												
All races, both sexes	205	-	-	-	-	-	-	1	-	-	-	4
Male	66	-	-	-	-	-	-	-	-	-	-	1
Female	139	-	-	-	-	-	-	1	-	-	-	3
White, both sexes	202	-	-	-	-	-	-	1	-	-	-	3
Male	65	-	-	-	-	-	-	-	-	-	-	1
Female	137	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	153	-	-	-	-	-	-	-	-	-	-	2
Male	57	-	-	-	-	-	-	-	-	-	-	1
Female	96	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	147	-	-	-	-	-	-	-	-	-	-	1
Male	51	-	-	-	-	-	-	-	-	-	-	-
Female	96	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Kentucky												
All races, both sexes	132	1	-	-	-	1	-	-	1	-	1	-
Male	50	1	-	-	-	1	-	-	-	-	-	-
Female	82	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	127	-	-	-	-	1	-	-	1	-	1	-
Male	47	-	-	-	-	1	-	-	-	-	-	-
Female	80	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	149	-	-	-	1	1	-	-	-	1	1	1
Male	70	-	-	-	1	1	-	-	-	1	1	1
Female	79	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	129	-	-	-	1	-	-	-	-	-	1	1
Male	62	-	-	-	1	-	-	-	-	-	1	1
Female	67	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	1	-	-	-	1	-	-
Male	8	-	-	-	-	1	-	-	-	1	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Maine												
All races, both sexes	86	-	-	-	-	-	-	-	-	-	-	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	86	-	-	-	-	-	-	-	-	-	-	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	200	-	-	-	1	-	1	-	-	-	1	1
Male	67	-	-	-	-	-	-	-	-	-	1	1
Female	133	-	-	-	1	-	1	-	-	-	-	-
White, both sexes	174	-	-	-	1	-	1	-	-	-	1	-
Male	57	-	-	-	-	-	-	-	-	-	1	-
Female	117	-	-	-	1	-	1	-	-	-	-	-
Black, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Massachusetts												
All races, both sexes	347	-	-	-	-	-	1	-	-	-	2	-
Male	113	-	-	-	-	-	1	-	-	-	1	-
Female	234	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	334	-	-	-	-	-	1	-	-	-	1	-
Male	106	-	-	-	-	-	1	-	-	-	-	-
Female	228	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	406	-	-	-	-	1	-	-	1	-	4	4
Male	143	-	-	-	-	1	-	-	1	-	4	4
Female	263	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	387	-	-	-	-	1	-	-	1	-	4	4
Male	134	-	-	-	-	1	-	-	1	-	4	4
Female	253	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Minnesota												
All races, both sexes	323	-	-	-	-	-	-	-	1	-	1	1
Male	123	-	-	-	-	-	-	-	1	-	1	-
Female	200	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	317	-	-	-	-	-	-	-	1	-	-	1
Male	120	-	-	-	-	-	-	-	1	-	-	-
Female	197	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	86	-	-	-	-	-	-	-	1	3	-	2
Male	38	-	-	-	-	-	-	-	1	1	-	2
Female	48	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	66	-	-	-	-	-	-	-	1	-	-	1
Male	32	-	-	-	-	-	-	-	1	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	18	-	-	-	-	-	-	-	-	3	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	1
Female	12	-	-	-	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Missouri												
All races, both sexes	268	-	-	-	-	-	-	1	-	-	1	1
Male	105	-	-	-	-	-	-	1	-	-	1	1
Female	163	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	262	-	-	-	-	-	-	1	-	-	1	-
Male	103	-	-	-	-	-	-	1	-	-	1	-
Female	159	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	63	-	-	-	-	-	-	-	1	-	1	-
Male	23	-	-	-	-	-	-	-	-	-	1	-
Female	40	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	62	-	-	-	-	-	-	-	1	-	-	-
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Nebraska												
All races, both sexes	102	-	-	-	-	-	-	-	-	-	-	1
Male	42	-	-	-	-	-	-	-	-	-	-	-
Female	60	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	100	-	-	-	-	-	-	-	-	-	-	1
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	59	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	86	1	-	-	-	-	-	-	-	-	-	-
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	55	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	82	1	-	-	-	-	-	-	-	-	-	-
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	52	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
New Hampshire												
All races, both sexes	70	-	-	-	-	-	-	-	-	-	-	1
Male	26	-	-	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	70	-	-	-	-	-	-	-	-	-	-	1
Male	26	-	-	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	456	-	-	-	-	-	-	1	2	1	-	4
Male	165	-	-	-	-	-	-	1	2	1	-	3
Female	291	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	438	-	-	-	-	-	-	-	2	1	-	4
Male	155	-	-	-	-	-	-	-	2	1	-	3
Female	283	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
New Mexico												
All races, both sexes	47	-	-	-	-	-	-	1	-	-	-	1
Male	24	-	-	-	-	-	-	1	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	45	-	-	-	-	-	-	1	-	-	-	1
Male	23	-	-	-	-	-	-	1	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	691	-	-	-	-	-	-	1	1	1	5	4
Male	244	-	-	-	-	-	-	1	-	1	3	2
Female	447	-	-	-	-	-	-	-	1	-	2	2
White, both sexes	663	-	-	-	-	-	-	-	1	1	5	3
Male	234	-	-	-	-	-	-	-	-	1	3	1
Female	429	-	-	-	-	-	-	-	1	-	2	2
Black, both sexes	24	-	-	-	-	-	-	1	-	-	-	1
Male	8	-	-	-	-	-	-	1	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
North Carolina												
All races, both sexes	380	-	-	-	-	-	-	-	-	-	1	3
Male	156	-	-	-	-	-	-	-	-	-	1	3
Female	224	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	336	-	-	-	-	-	-	-	-	-	-	3
Male	142	-	-	-	-	-	-	-	-	-	-	3
Female	194	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	40	-	-	-	-	-	-	-	-	-	1	-
Male	13	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	29	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Ohio												
All races, both sexes	556	-	1	-	1	-	2	1	1	-	5	6
Male	227	-	1	-	1	-	2	1	1	-	4	4
Female	329	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	531	-	1	-	1	-	2	1	1	-	5	3
Male	214	-	1	-	1	-	2	1	1	-	4	3
Female	317	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	23	-	-	-	-	-	-	-	-	-	-	3
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	124	-	-	-	-	-	-	-	1	-	-	2
Male	47	-	-	-	-	-	-	-	-	-	-	2
Female	77	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	115	-	-	-	-	-	-	-	-	-	-	-
Male	41	-	-	-	-	-	-	-	-	-	-	-
Female	74	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Oregon												
All races, both sexes	341	-	-	-	-	-	-	-	1	-	-	-
Male	137	-	-	-	-	-	-	-	1	-	-	-
Female	204	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	333	-	-	-	-	-	-	-	1	-	-	-
Male	135	-	-	-	-	-	-	-	1	-	-	-
Female	198	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	980	-	-	-	-	-	-	1	2	1	3	3
Male	350	-	-	-	-	-	-	1	1	1	3	2
Female	630	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	947	-	-	-	-	-	-	1	1	1	2	2
Male	338	-	-	-	-	-	-	1	1	1	2	2
Female	609	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	32	-	-	-	-	-	-	-	1	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Rhode Island												
All races, both sexes	52	-	-	-	-	-	-	-	-	1	1	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	33	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	52	-	-	-	-	-	-	-	-	1	1	-
Male	19	-	-	-	-	-	-	-	-	-	1	-
Female	33	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	189	-	-	-	-	-	-	-	1	-	1	4
Male	68	-	-	-	-	-	-	-	1	-	-	3
Female	121	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	158	-	-	-	-	-	-	-	-	-	1	3
Male	58	-	-	-	-	-	-	-	-	-	-	3
Female	100	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	29	-	-	-	-	-	-	-	1	-	-	1
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
South Dakota												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	219	-	-	-	-	-	-	-	-	-	-	1
Male	88	-	-	-	-	-	-	-	-	-	-	1
Female	131	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	208	-	-	-	-	-	-	-	-	-	-	1
Male	85	-	-	-	-	-	-	-	-	-	-	1
Female	123	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Texas												
All races, both sexes	652	-	-	-	-	-	1	-	1	7	7	9
Male	266	-	-	-	-	-	1	-	1	5	5	7
Female	386	-	-	-	-	-	-	-	-	2	2	2
White, both sexes	611	-	-	-	-	-	-	-	-	5	6	9
Male	251	-	-	-	-	-	-	-	-	4	4	7
Female	360	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	38	-	-	-	-	-	1	-	1	2	1	-
Male	15	-	-	-	-	-	1	-	1	1	1	-
Female	23	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	75	-	-	-	-	-	-	-	1	-	-	-
Male	25	-	-	-	-	-	-	-	1	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	75	-	-	-	-	-	-	-	1	-	-	-
Male	25	-	-	-	-	-	-	-	1	-	-	-
Female	50	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Vermont												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	40	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	249	-	-	-	-	-	1	1	-	-	2	1
Male	97	-	-	-	-	-	1	1	-	-	1	-
Female	152	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	226	-	-	-	-	-	1	1	-	-	2	-
Male	90	-	-	-	-	-	1	1	-	-	1	-
Female	136	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Washington												
All races, both sexes	445	-	-	-	-	-	-	-	2	-	-	5
Male	183	-	-	-	-	-	-	-	2	-	-	4
Female	262	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	434	-	-	-	-	-	-	-	2	-	-	5
Male	177	-	-	-	-	-	-	-	2	-	-	4
Female	257	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	73	-	-	-	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	72	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Nonrheumatic aortic valve disorders (I35)												
Wisconsin												
All races, both sexes	354	-	-	-	-	-	-	-	-	-	1	1
Male	145	-	-	-	-	-	-	-	-	-	1	1
Female	209	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	349	-	-	-	-	-	-	-	-	-	1	1
Male	143	-	-	-	-	-	-	-	-	-	1	1
Female	206	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
United States												
All races, both sexes	135	214	280	376	667	1,316	2,428	3,368	2,776	1,166	175	1
Male	92	141	178	216	340	648	1,023	1,216	801	234	34	-
Female	43	73	102	160	327	668	1,405	2,152	1,975	932	141	1
White, both sexes	104	185	242	331	595	1,244	2,310	3,240	2,687	1,114	164	-
Male	76	119	154	197	308	616	979	1,173	781	221	32	-
Female	28	66	88	134	287	628	1,331	2,067	1,906	893	132	-
Black, both sexes	28	26	34	35	57	62	86	93	60	39	8	1
Male	16	20	23	17	25	29	32	30	11	9	1	-
Female	12	6	11	18	32	33	54	63	49	30	7	1
American Indian 1/, both sexes	1	2	2	3	3	-	5	7	4	3	-	-
Male	-	2	1	1	3	-	4	3	1	2	-	-
Female	1	-	1	2	-	-	1	4	3	1	-	-
Asian or Pacific Islander 2/, both sexes	2	1	2	7	12	10	27	28	25	10	3	-
Male	-	-	-	1	4	3	8	10	8	2	1	-
Female	2	1	2	6	8	7	19	18	17	8	2	-
Alabama												
All races, both sexes	5	6	1	4	12	20	34	36	28	10	2	-
Male	2	2	-	3	7	9	17	9	6	4	-	-
Female	3	4	1	1	5	11	17	27	22	6	2	-
White, both sexes	3	5	1	2	9	17	31	34	26	9	1	-
Male	1	2	-	2	6	9	15	7	6	3	-	-
Female	2	3	1	-	3	8	16	27	20	6	1	-
Black, both sexes	2	1	-	2	3	3	3	2	2	1	1	-
Male	1	-	-	1	1	-	2	2	-	1	-	-
Female	1	1	-	1	2	3	1	-	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Alaska												
All races, both sexes	1	-	-	1	2	2	4	6	6	-	-	-
Male	1	-	-	-	2	2	1	2	1	-	-	-
Female	-	-	-	1	-	-	3	4	5	-	-	-
White, both sexes	1	-	-	-	1	2	3	3	5	-	-	-
Male	1	-	-	-	1	2	1	1	1	-	-	-
Female	-	-	-	-	-	-	2	2	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	2	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	5	5	11	8	19	32	52	35	7	-	-
Male	1	4	5	5	3	5	15	23	12	-	-	-
Female	1	1	-	6	5	14	17	29	23	7	-	-
White, both sexes	1	4	5	11	8	19	30	51	35	7	-	-
Male	1	3	5	5	3	5	14	22	12	-	-	-
Female	-	1	-	6	5	14	16	29	23	7	-	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Arkansas												
All races, both sexes	1	2	1	3	7	9	12	33	16	8	-	-
Male	1	1	-	3	3	4	3	9	4	1	-	-
Female	-	1	1	-	4	5	9	24	12	7	-	-
White, both sexes	-	2	1	3	5	9	11	33	15	8	-	-
Male	-	1	-	3	3	4	3	9	4	1	-	-
Female	-	1	1	-	2	5	8	24	11	7	-	-
Black, both sexes	1	-	-	-	2	-	1	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	18	21	34	36	63	149	289	379	336	131	29	-
Male	15	17	23	18	31	73	131	140	97	30	4	-
Female	3	4	11	18	32	76	158	239	239	101	25	-
White, both sexes	15	18	31	30	54	138	263	360	319	123	28	-
Male	12	14	22	17	29	67	121	134	93	27	4	-
Female	3	4	9	13	25	71	142	226	226	96	24	-
Black, both sexes	3	3	2	4	3	5	10	4	6	5	-	-
Male	3	3	1	1	-	4	5	2	1	2	-	-
Female	-	-	1	3	3	1	5	2	5	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	6	6	16	13	11	3	1	-
Male	-	-	-	-	2	2	5	2	3	1	-	-
Female	-	-	1	2	4	4	11	11	8	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Colorado												
All races, both sexes	3	2	3	4	9	16	30	48	33	17	2	-
Male	3	1	1	3	4	8	12	28	14	3	1	-
Female	-	1	2	1	5	8	18	20	19	14	1	-
White, both sexes	3	2	3	4	7	16	30	48	33	17	2	-
Male	3	1	1	3	3	8	12	28	14	3	1	-
Female	-	1	2	1	4	8	18	20	19	14	1	-
Black, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	3	1	10	11	32	54	58	21	4	-
Male	-	-	-	1	6	5	13	15	12	2	1	-
Female	-	-	3	-	4	6	19	39	46	19	3	-
White, both sexes	-	-	2	1	10	11	31	53	58	21	4	-
Male	-	-	-	1	6	5	13	15	12	2	1	-
Female	-	-	2	-	4	6	18	38	46	19	3	-
Black, both sexes	-	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Delaware												
All races, both sexes	-	-	2	2	3	7	6	10	10	4	1	-
Male	-	-	2	1	1	2	2	5	-	-	-	-
Female	-	-	-	1	2	5	4	5	10	4	1	-
White, both sexes	-	-	1	2	2	5	5	8	10	4	1	-
Male	-	-	1	1	-	1	2	4	-	-	-	-
Female	-	-	-	1	2	4	3	4	10	4	1	-
Black, both sexes	-	-	1	-	1	2	1	2	-	-	-	-
Male	-	-	1	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	1	2	2	2	1	4	1	-
Male	-	-	-	-	1	1	-	-	1	1	-	-
Female	-	-	-	-	-	1	2	2	-	3	1	-
White, both sexes	-	-	-	-	-	2	1	1	1	3	1	-
Male	-	-	-	-	-	1	-	-	1	1	-	-
Female	-	-	-	-	-	1	1	1	-	2	1	-
Black, both sexes	-	-	-	-	1	-	1	1	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Florida												
All races, both sexes	8	10	24	29	35	93	167	209	181	76	4	1
Male	6	8	14	15	20	50	75	91	57	24	-	-
Female	2	2	10	14	15	43	92	118	124	52	4	1
White, both sexes	7	9	23	27	33	90	162	207	176	75	4	-
Male	5	7	13	13	20	48	74	91	55	23	-	-
Female	2	2	10	14	13	42	88	116	121	52	4	-
Black, both sexes	1	1	1	2	2	3	5	2	5	1	-	1
Male	1	1	1	2	-	2	1	-	2	1	-	-
Female	-	-	-	-	2	1	4	2	3	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	3	9	6	12	22	36	40	51	45	13	1	-
Male	3	4	6	8	15	19	19	18	6	2	-	-
Female	-	5	-	4	7	17	21	33	39	11	1	-
White, both sexes	3	5	5	10	16	33	34	43	41	11	-	-
Male	3	1	5	6	11	19	17	14	6	2	-	-
Female	-	4	-	4	5	14	17	29	35	9	-	-
Black, both sexes	-	4	1	2	6	3	6	8	4	2	1	-
Male	-	3	1	2	4	-	2	4	-	-	-	-
Female	-	1	-	-	2	3	4	4	4	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Hawaii												
All races, both sexes	1	2	-	2	3	3	10	12	15	8	1	-
Male	-	1	-	-	2	-	3	4	6	2	-	-
Female	1	1	-	2	1	3	7	8	9	6	1	-
White, both sexes	-	2	-	-	2	1	4	5	6	2	-	-
Male	-	1	-	-	2	-	1	1	2	1	-	-
Female	-	1	-	-	-	1	3	4	4	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	1	2	6	7	9	6	1	-
Male	-	-	-	-	-	-	2	3	4	1	-	-
Female	1	-	-	2	1	2	4	4	5	5	1	-
Idaho												
All races, both sexes	2	1	2	1	5	9	16	23	21	10	-	-
Male	1	1	-	1	3	5	7	12	8	5	-	-
Female	1	-	2	-	2	4	9	11	13	5	-	-
White, both sexes	2	1	1	1	5	9	15	22	21	10	-	-
Male	1	1	-	1	3	5	6	11	8	5	-	-
Female	1	-	1	-	2	4	9	11	13	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Illinois												
All races, both sexes	5	4	12	15	22	48	86	132	104	54	18	-
Male	4	1	7	8	12	30	47	38	28	7	5	-
Female	1	3	5	7	10	18	39	94	76	47	13	-
White, both sexes	4	4	10	15	20	44	80	128	99	52	16	-
Male	3	1	5	8	11	26	43	36	26	7	3	-
Female	1	3	5	7	9	18	37	92	73	45	13	-
Black, both sexes	1	-	2	-	1	4	6	4	5	2	1	-
Male	1	-	2	-	-	4	4	2	2	-	1	-
Female	-	-	-	-	1	-	2	2	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	1	-
Male	-	-	-	-	1	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	4	8	8	18	34	53	68	53	41	5	-
Male	1	4	5	4	12	21	18	26	20	7	-	-
Female	1	-	3	4	6	13	35	42	33	34	5	-
White, both sexes	1	4	7	7	15	33	51	68	50	39	5	-
Male	1	4	4	4	10	21	18	26	18	7	-	-
Female	-	-	3	3	5	12	33	42	32	32	5	-
Black, both sexes	1	-	1	1	3	1	2	-	3	1	-	-
Male	-	-	1	-	2	-	-	-	2	-	-	-
Female	1	-	-	1	1	1	2	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Iowa												
All races, both sexes	-	4	2	6	6	17	43	47	50	19	6	-
Male	-	2	1	3	4	7	15	15	12	4	2	-
Female	-	2	1	3	2	10	28	32	38	15	4	-
White, both sexes	-	4	1	6	6	16	43	47	50	19	6	-
Male	-	2	-	3	4	7	15	15	12	4	2	-
Female	-	2	1	3	2	9	28	32	38	15	4	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	3	4	2	9	15	21	39	34	20	3	-
Male	1	1	2	-	3	8	11	14	10	5	1	-
Female	-	2	2	2	6	7	10	25	24	15	2	-
White, both sexes	1	3	4	2	9	14	19	39	34	18	3	-
Male	1	1	2	-	3	7	9	14	10	3	1	-
Female	-	2	2	2	6	7	10	25	24	15	2	-
Black, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	1	-	-
Male	-	-	-	-	-	-	2	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Kentucky												
All races, both sexes	2	3	4	3	3	15	18	35	33	12	-	-
Male	2	3	1	1	2	7	6	10	16	-	-	-
Female	-	-	3	2	1	8	12	25	17	12	-	-
White, both sexes	2	2	3	3	2	15	18	35	33	11	-	-
Male	2	2	1	1	1	7	6	10	16	-	-	-
Female	-	-	2	2	1	8	12	25	17	11	-	-
Black, both sexes	-	1	1	-	1	-	-	-	-	1	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	4	3	5	15	20	30	32	20	10	3	-
Male	1	3	2	4	8	8	13	14	6	4	2	-
Female	1	1	1	1	7	12	17	18	14	6	1	-
White, both sexes	2	2	3	5	10	20	27	27	17	10	3	-
Male	1	1	2	4	7	8	12	12	6	4	2	-
Female	1	1	1	1	3	12	15	15	11	6	1	-
Black, both sexes	-	2	-	-	5	-	3	5	3	-	-	-
Male	-	2	-	-	1	-	1	2	-	-	-	-
Female	-	-	-	-	4	-	2	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Maine												
All races, both sexes	-	4	1	2	5	13	22	22	10	7	-	-
Male	-	2	1	2	3	7	9	7	4	1	-	-
Female	-	2	-	-	2	6	13	15	6	6	-	-
White, both sexes	-	4	1	2	5	13	22	22	10	7	-	-
Male	-	2	1	2	3	7	9	7	4	1	-	-
Female	-	2	-	-	2	6	13	15	6	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5	5	7	4	19	21	32	46	39	16	2	-
Male	4	4	5	2	4	9	10	16	9	1	1	-
Female	1	1	2	2	15	12	22	30	30	15	1	-
White, both sexes	1	4	5	1	15	19	30	44	36	14	2	-
Male	1	3	4	1	2	9	9	16	9	1	1	-
Female	-	1	1	-	13	10	21	28	27	13	1	-
Black, both sexes	4	1	2	3	4	2	2	2	3	2	-	-
Male	3	1	1	1	2	-	1	-	-	-	-	-
Female	1	-	1	2	2	2	1	2	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Massachusetts												
All races, both sexes	1	6	4	9	10	24	50	104	88	39	9	-
Male	-	4	3	7	5	7	19	41	17	6	2	-
Female	1	2	1	2	5	17	31	63	71	33	7	-
White, both sexes	-	6	3	8	10	23	47	101	86	39	9	-
Male	-	4	2	6	5	6	17	40	17	6	2	-
Female	-	2	1	2	5	17	30	61	69	33	7	-
Black, both sexes	1	-	1	1	-	1	3	3	2	-	-	-
Male	-	-	1	1	-	1	2	1	-	-	-	-
Female	1	-	-	-	-	-	1	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	10	9	15	22	44	74	109	80	29	3	-
Male	-	5	6	8	10	15	30	36	20	3	-	-
Female	1	5	3	7	12	29	44	73	60	26	3	-
White, both sexes	-	7	9	12	20	40	73	107	80	26	3	-
Male	-	2	6	7	9	14	29	35	20	2	-	-
Female	-	5	3	5	11	26	44	72	60	24	3	-
Black, both sexes	1	3	-	3	2	4	1	2	-	2	-	-
Male	-	3	-	1	1	1	1	1	-	1	-	-
Female	1	-	-	2	1	3	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Minnesota												
All races, both sexes	1	3	6	3	9	30	57	81	89	37	4	-
Male	-	1	4	2	5	12	28	32	27	10	-	-
Female	1	2	2	1	4	18	29	49	62	27	4	-
White, both sexes	1	3	5	3	9	29	56	81	87	37	4	-
Male	-	1	3	2	5	12	28	32	26	10	-	-
Female	1	2	2	1	4	17	28	49	61	27	4	-
Black, both sexes	-	-	1	-	-	1	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Mississippi												
All races, both sexes	5	1	1	4	9	11	15	19	11	2	2	-
Male	3	1	-	1	5	9	5	5	5	-	-	-
Female	2	-	1	3	4	2	10	14	6	2	2	-
White, both sexes	3	-	1	2	6	10	14	15	10	2	1	-
Male	3	-	-	1	3	8	5	5	5	-	-	-
Female	-	-	1	1	3	2	9	10	5	2	1	-
Black, both sexes	1	1	-	2	3	1	1	3	1	-	1	-
Male	-	1	-	-	2	1	-	-	-	-	-	-
Female	1	-	-	2	1	-	1	3	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Missouri												
All races, both sexes	1	5	9	10	10	28	53	72	55	22	-	-
Male	1	5	7	7	6	13	24	21	15	3	-	-
Female	-	-	2	3	4	15	29	51	40	19	-	-
White, both sexes	1	5	9	10	10	27	52	70	54	22	-	-
Male	1	5	7	7	6	12	24	21	15	3	-	-
Female	-	-	2	3	4	15	28	49	39	19	-	-
Black, both sexes	-	-	-	-	-	1	1	2	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	4	4	14	14	16	7	1	-
Male	-	-	-	-	1	4	6	7	4	-	-	-
Female	1	-	-	-	3	-	8	7	12	7	1	-
White, both sexes	1	-	-	-	4	4	14	14	16	7	1	-
Male	-	-	-	-	1	4	6	7	4	-	-	-
Female	1	-	-	-	3	-	8	7	12	7	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Nebraska												
All races, both sexes	3	2	1	4	10	5	15	23	23	12	3	-
Male	3	2	1	3	3	4	8	9	7	2	-	-
Female	-	-	-	1	7	1	7	14	16	10	3	-
White, both sexes	3	2	1	4	10	4	15	22	23	12	3	-
Male	3	2	1	3	3	3	8	9	7	2	-	-
Female	-	-	-	1	7	1	7	13	16	10	3	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	1	5	7	6	3	16	22	16	5	3	-
Male	-	1	2	6	2	-	7	7	6	-	-	-
Female	1	-	3	1	4	3	9	15	10	5	3	-
White, both sexes	-	1	5	5	6	3	16	22	15	5	3	-
Male	-	1	2	5	2	-	7	7	6	-	-	-
Female	-	-	3	-	4	3	9	15	9	5	3	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
New Hampshire												
All races, both sexes	-	6	1	1	5	5	16	17	8	9	1	-
Male	-	6	-	1	2	2	6	5	2	1	-	-
Female	-	-	1	-	3	3	10	12	6	8	1	-
White, both sexes	-	6	1	1	5	5	16	17	8	9	1	-
Male	-	6	-	1	2	2	6	5	2	1	-	-
Female	-	-	1	-	3	3	10	12	6	8	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	5	7	8	4	18	46	84	140	88	42	6	-
Male	4	3	6	4	6	21	33	48	20	10	3	-
Female	1	4	2	-	12	25	51	92	68	32	3	-
White, both sexes	4	7	6	3	15	44	81	138	86	41	6	-
Male	3	3	4	3	5	21	32	47	19	9	3	-
Female	1	4	2	-	10	23	49	91	67	32	3	-
Black, both sexes	1	-	2	1	3	2	3	1	2	-	-	-
Male	1	-	2	1	1	-	1	-	1	-	-	-
Female	-	-	-	-	2	2	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
New Mexico												
All races, both sexes	1	1	1	3	2	6	7	13	9	2	-	-
Male	1	-	1	1	2	4	2	7	3	1	-	-
Female	-	1	-	2	-	2	5	6	6	1	-	-
White, both sexes	1	1	-	3	2	6	7	13	8	2	-	-
Male	1	-	-	1	2	4	2	7	3	1	-	-
Female	-	1	-	2	-	2	5	6	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	9	21	19	34	68	129	171	149	68	5	-
Male	5	5	11	14	17	34	46	51	40	13	1	-
Female	1	4	10	5	17	34	83	120	109	55	4	-
White, both sexes	5	7	17	17	32	62	124	169	147	68	5	-
Male	5	5	9	13	16	33	45	50	39	13	1	-
Female	-	2	8	4	16	29	79	119	108	55	4	-
Black, both sexes	1	1	4	-	2	6	5	2	1	-	-	-
Male	-	-	2	-	1	1	1	1	-	-	-	-
Female	1	1	2	-	1	5	4	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
North Carolina												
All races, both sexes	4	5	13	13	25	41	83	94	65	31	2	-
Male	1	3	7	9	18	20	38	28	21	6	1	-
Female	3	2	6	4	7	21	45	66	44	25	1	-
White, both sexes	1	2	8	11	23	37	76	82	64	27	2	-
Male	-	1	5	8	17	17	36	27	21	6	1	-
Female	1	1	3	3	6	20	40	55	43	21	1	-
Black, both sexes	3	3	5	2	1	4	6	10	1	4	-	-
Male	1	2	2	1	1	3	1	1	-	-	-	-
Female	2	1	3	1	-	1	5	9	1	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
North Dakota												
All races, both sexes	1	-	2	1	2	2	2	8	8	3	-	-
Male	1	-	2	-	1	-	-	2	1	1	-	-
Female	-	-	-	1	1	2	2	6	7	2	-	-
White, both sexes	1	-	2	1	2	2	2	8	8	3	-	-
Male	1	-	2	-	1	-	-	2	1	1	-	-
Female	-	-	-	1	1	2	2	6	7	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Ohio												
All races, both sexes	3	10	14	18	25	52	105	145	115	45	7	-
Male	1	6	13	12	11	25	51	48	37	8	1	-
Female	2	4	1	6	14	27	54	97	78	37	6	-
White, both sexes	3	10	11	16	24	50	102	140	111	44	6	-
Male	1	6	11	10	11	23	48	46	36	8	1	-
Female	2	4	-	6	13	27	54	94	75	36	5	-
Black, both sexes	-	-	3	1	-	2	3	5	4	1	1	-
Male	-	-	2	1	-	2	3	2	1	-	-	-
Female	-	-	1	-	-	-	-	3	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	5	3	3	9	9	20	32	26	10	3	-
Male	1	1	1	-	7	5	7	9	11	2	1	-
Female	-	4	2	3	2	4	13	23	15	8	2	-
White, both sexes	1	5	3	3	9	8	19	30	25	9	3	-
Male	1	1	1	-	7	5	6	7	10	2	1	-
Female	-	4	2	3	2	3	13	23	15	7	2	-
Black, both sexes	-	-	-	-	-	1	1	1	-	1	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Oregon												
All races, both sexes	2	1	4	7	8	27	59	93	104	32	3	-
Male	1	-	3	2	4	14	30	40	34	8	-	-
Female	1	1	1	5	4	13	29	53	70	24	3	-
White, both sexes	2	1	4	6	8	27	59	89	101	32	3	-
Male	1	-	3	2	4	14	30	38	34	8	-	-
Female	1	1	1	4	4	13	29	51	67	24	3	-
Black, both sexes	-	-	-	-	-	-	-	2	2	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Pennsylvania												
All races, both sexes	8	16	7	21	47	80	213	254	225	90	9	-
Male	7	12	6	12	23	38	83	92	56	10	3	-
Female	1	4	1	9	24	42	130	162	169	80	6	-
White, both sexes	7	14	7	18	44	76	209	247	225	84	9	-
Male	6	10	6	11	22	35	81	91	56	10	3	-
Female	1	4	1	7	22	41	128	156	169	74	6	-
Black, both sexes	1	2	-	3	3	4	4	7	-	6	-	-
Male	1	2	-	1	1	3	2	1	-	-	-	-
Female	-	-	-	2	2	1	2	6	-	6	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Rhode Island												
All races, both sexes	-	1	1	1	2	2	7	10	17	8	1	-
Male	-	1	-	1	1	-	7	3	5	-	-	-
Female	-	-	1	-	1	2	-	7	12	8	1	-
White, both sexes	-	1	1	1	2	2	7	10	17	8	1	-
Male	-	1	-	1	1	-	7	3	5	-	-	-
Female	-	-	1	-	1	2	-	7	12	8	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	5	3	2	9	19	22	32	51	30	8	2	-
Male	4	2	1	3	8	9	7	18	8	3	1	-
Female	1	1	1	6	11	13	25	33	22	5	1	-
White, both sexes	4	3	1	9	18	17	27	41	25	7	2	-
Male	3	2	-	3	8	7	7	14	7	3	1	-
Female	1	1	1	6	10	10	20	27	18	4	1	-
Black, both sexes	1	-	1	-	1	5	4	9	5	1	-	-
Male	1	-	1	-	-	2	-	3	1	-	-	-
Female	-	-	-	-	1	3	4	6	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
South Dakota												
All races, both sexes	1	-	-	1	1	2	4	11	8	6	1	-
Male	-	-	-	1	1	2	2	5	3	-	-	-
Female	1	-	-	-	-	-	2	6	5	6	1	-
White, both sexes	1	-	-	1	1	2	4	11	8	6	1	-
Male	-	-	-	1	1	2	2	5	3	-	-	-
Female	1	-	-	-	-	-	2	6	5	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	4	5	12	9	19	57	58	40	9	2	-
Male	2	3	5	9	-	10	22	21	13	1	1	-
Female	1	1	-	3	9	9	35	37	27	8	1	-
White, both sexes	1	4	4	11	8	18	56	55	39	9	2	-
Male	1	3	4	9	-	10	22	20	13	1	1	-
Female	-	1	-	2	8	8	34	35	26	8	1	-
Black, both sexes	2	-	1	1	1	1	1	3	1	-	-	-
Male	1	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	1	1	1	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Texas												
All races, both sexes	7	14	18	32	44	95	106	147	106	50	8	-
Male	3	8	10	19	24	47	42	48	31	14	1	-
Female	4	6	8	13	20	48	64	99	75	36	7	-
White, both sexes	6	12	16	28	41	91	102	139	103	46	7	-
Male	3	8	9	17	23	46	40	46	31	12	1	-
Female	3	4	7	11	18	45	62	93	72	34	6	-
Black, both sexes	1	2	1	4	3	3	4	7	3	4	1	-
Male	-	-	1	2	1	1	2	2	-	2	-	-
Female	1	2	-	2	2	2	2	5	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	1	-	-	-	-
Utah												
All races, both sexes	3	1	1	2	2	11	15	20	14	5	-	-
Male	2	-	1	2	-	5	6	4	4	-	-	-
Female	1	1	-	-	2	6	9	16	10	5	-	-
White, both sexes	3	1	1	2	2	11	15	20	14	5	-	-
Male	2	-	1	2	-	5	6	4	4	-	-	-
Female	1	1	-	-	2	6	9	16	10	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Vermont												
All races, both sexes	-	-	2	-	1	3	6	10	12	6	-	-
Male	-	-	2	-	1	3	1	7	3	3	-	-
Female	-	-	-	-	-	-	5	3	9	3	-	-
White, both sexes	-	-	2	-	1	3	6	10	12	6	-	-
Male	-	-	2	-	1	3	1	7	3	3	-	-
Female	-	-	-	-	-	-	5	3	9	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	5	5	7	23	26	42	64	48	19	3	-
Male	1	4	3	1	13	14	15	20	16	7	-	-
Female	1	1	2	6	10	12	27	44	32	12	3	-
White, both sexes	1	4	4	5	20	25	37	63	45	16	2	-
Male	-	3	3	-	12	13	14	20	16	6	-	-
Female	1	1	1	5	8	12	23	43	29	10	2	-
Black, both sexes	1	1	1	2	2	1	5	1	3	3	1	-
Male	1	1	-	1	1	1	1	-	-	1	-	-
Female	-	-	1	1	1	-	4	1	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Washington												
All races, both sexes	4	6	6	9	16	40	85	123	98	43	8	-
Male	3	6	5	4	7	20	32	54	33	11	2	-
Female	1	-	1	5	9	20	53	69	65	32	6	-
White, both sexes	4	5	6	9	13	40	82	121	96	43	8	-
Male	3	5	5	4	4	20	31	53	33	11	2	-
Female	1	-	1	5	9	20	51	68	63	32	6	-
Black, both sexes	-	-	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	1	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	3	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-
West Virginia												
All races, both sexes	3	2	2	1	6	7	17	18	14	3	-	-
Male	2	2	-	-	1	5	8	6	4	-	-	-
Female	1	-	2	1	5	2	9	12	10	3	-	-
White, both sexes	3	2	2	1	5	7	17	18	14	3	-	-
Male	2	2	-	-	-	5	8	6	4	-	-	-
Female	1	-	2	1	5	2	9	12	10	3	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Nonrheumatic aortic valve disorders (I35)												
Wisconsin												
All races, both sexes	-	-	7	9	11	41	60	102	81	35	6	-
Male	-	-	3	5	9	23	29	43	24	7	-	-
Female	-	-	4	4	2	18	31	59	57	28	6	-
White, both sexes	-	-	6	9	10	41	59	102	80	35	5	-
Male	-	-	2	5	8	23	29	43	24	7	-	-
Female	-	-	4	4	2	18	30	59	56	28	5	-
Black, both sexes	-	-	1	-	-	-	1	-	1	-	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	1	-	-	1	4	6	7	5	1	1	-
Male	-	1	-	-	1	3	2	3	2	1	-	-
Female	1	-	-	-	-	1	4	4	3	-	1	-
White, both sexes	1	1	-	-	1	4	6	7	5	1	1	-
Male	-	1	-	-	1	3	2	3	2	1	-	-
Female	1	-	-	-	-	1	4	4	3	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
United States												
All races, both sexes	5,149	4	5	1	2	3	19	21	32	51	82	150
Male	2,017	2	5	1	1	3	10	12	15	34	47	99
Female	3,132	2	-	-	1	-	9	9	17	17	35	51
White, both sexes	4,549	3	4	-	1	3	15	16	22	29	54	110
Male	1,763	2	4	-	1	3	8	10	11	22	41	74
Female	2,786	1	-	-	-	-	7	6	11	7	13	36
Black, both sexes	493	1	1	1	1	-	3	5	8	20	26	37
Male	220	-	1	1	-	-	1	2	3	11	5	23
Female	273	1	-	-	1	-	2	3	5	9	21	14
American Indian 1/, both sexes	27	-	-	-	-	-	-	-	-	1	1	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	1	1	1
Asian or Pacific Islander 2/, both sexes	80	-	-	-	-	-	1	-	2	1	1	1
Male	27	-	-	-	-	-	1	-	1	1	1	1
Female	53	-	-	-	-	-	-	-	1	-	-	-
Alabama												
All races, both sexes	93	-	-	-	-	-	-	1	-	2	-	3
Male	25	-	-	-	-	-	-	-	-	-	-	2
Female	68	-	-	-	-	-	-	1	-	2	-	1
White, both sexes	72	-	-	-	-	-	-	-	-	1	-	2
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	21	-	-	-	-	-	-	1	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Alaska												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	105	-	-	-	-	-	-	-	1	-	4	2
Male	51	-	-	-	-	-	-	-	-	-	2	2
Female	54	-	-	-	-	-	-	-	1	-	2	-
White, both sexes	100	-	-	-	-	-	-	-	1	-	4	1
Male	47	-	-	-	-	-	-	-	-	-	2	1
Female	53	-	-	-	-	-	-	-	1	-	2	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Arkansas												
All races, both sexes	49	-	-	-	-	-	-	-	-	1	-	1
Male	26	-	-	-	-	-	-	-	-	1	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	368	1	2	-	-	1	3	1	2	6	6	12
Male	160	-	2	-	-	1	2	-	2	4	4	11
Female	208	1	-	-	-	-	1	1	-	2	2	1
White, both sexes	313	1	2	-	-	1	1	1	1	4	5	10
Male	141	-	2	-	-	1	-	-	1	3	4	9
Female	172	1	-	-	-	-	1	1	-	1	1	1
Black, both sexes	21	-	-	-	-	-	1	-	1	1	1	2
Male	10	-	-	-	-	-	1	-	1	-	-	2
Female	11	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	1	-	-	1	-	-
Male	9	-	-	-	-	-	1	-	-	1	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Colorado												
All races, both sexes	84	-	-	-	-	-	-	1	2	-	1	1
Male	26	-	-	-	-	-	-	1	2	-	1	-
Female	58	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	80	-	-	-	-	-	-	1	2	-	1	1
Male	25	-	-	-	-	-	-	1	2	-	1	-
Female	55	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	60	-	-	-	-	-	-	-	1	-	1	2
Male	32	-	-	-	-	-	-	-	-	-	1	1
Female	28	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	56	-	-	-	-	-	-	-	1	-	1	2
Male	29	-	-	-	-	-	-	-	-	-	1	1
Female	27	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Delaware												
All races, both sexes	19	-	-	-	-	-	-	1	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	1	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Florida												
All races, both sexes	380	-	-	-	-	-	-	2	1	1	3	8
Male	144	-	-	-	-	-	-	1	1	1	2	5
Female	236	-	-	-	-	-	-	1	-	-	1	3
White, both sexes	348	-	-	-	-	-	-	1	1	1	1	7
Male	131	-	-	-	-	-	-	-	1	1	1	5
Female	217	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	32	-	-	-	-	-	-	1	-	-	2	1
Male	13	-	-	-	-	-	-	1	-	-	1	-
Female	19	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	125	-	-	-	-	-	-	-	1	2	2	4
Male	51	-	-	-	-	-	-	-	-	1	-	3
Female	74	-	-	-	-	-	-	-	1	1	2	1
White, both sexes	88	-	-	-	-	-	-	-	1	-	-	2
Male	38	-	-	-	-	-	-	-	-	-	-	2
Female	50	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	37	-	-	-	-	-	-	-	-	2	2	2
Male	13	-	-	-	-	-	-	-	-	1	-	1
Female	24	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Hawaii												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Illinois												
All races, both sexes	185	-	-	-	1	-	4	1	3	3	1	9
Male	82	-	-	-	-	-	1	-	2	2	1	6
Female	103	-	-	-	1	-	3	1	1	1	-	3
White, both sexes	156	-	-	-	-	-	3	1	2	2	1	4
Male	68	-	-	-	-	-	1	-	2	2	1	3
Female	88	-	-	-	-	-	2	1	-	-	-	1
Black, both sexes	29	-	-	-	1	-	1	-	1	1	-	5
Male	14	-	-	-	-	-	-	-	-	-	-	3
Female	15	-	-	-	1	-	1	-	1	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	149	-	-	-	-	-	1	-	2	1	2	7
Male	39	-	-	-	-	-	-	-	-	1	1	3
Female	110	-	-	-	-	-	1	-	2	-	1	4
White, both sexes	144	-	-	-	-	-	1	-	2	-	2	6
Male	38	-	-	-	-	-	-	-	-	-	1	3
Female	106	-	-	-	-	-	1	-	2	-	1	3
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Iowa												
All races, both sexes	27	-	1	-	-	-	-	-	-	-	-	-
Male	12	-	1	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	27	-	1	-	-	-	-	-	-	-	-	-
Male	12	-	1	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	60	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	58	-	-	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	40	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Kentucky												
All races, both sexes	64	-	-	-	-	-	-	-	-	1	1	3
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	40	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	61	-	-	-	-	-	-	-	-	1	1	3
Male	23	-	-	-	-	-	-	-	-	-	-	2
Female	38	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	78	-	1	-	-	-	1	-	1	1	1	2
Male	31	-	1	-	-	-	1	-	-	-	1	-
Female	47	-	-	-	-	-	-	-	1	1	-	2
White, both sexes	55	-	-	-	-	-	1	-	-	1	1	1
Male	22	-	-	-	-	-	1	-	-	-	1	-
Female	33	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	23	-	1	-	-	-	-	-	1	-	-	1
Male	9	-	1	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Maine												
All races, both sexes	27	-	-	-	-	-	-	-	-	1	1	1
Male	13	-	-	-	-	-	-	-	-	1	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	27	-	-	-	-	-	-	-	-	1	1	1
Male	13	-	-	-	-	-	-	-	-	1	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	76	1	-	-	-	-	-	2	1	2	4	3
Male	34	1	-	-	-	-	-	2	-	-	2	2
Female	42	-	-	-	-	-	-	-	1	2	2	1
White, both sexes	52	1	-	-	-	-	-	2	1	-	1	2
Male	24	1	-	-	-	-	-	2	-	-	1	2
Female	28	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	22	-	-	-	-	-	-	-	-	2	3	1
Male	9	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	2	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Massachusetts												
All races, both sexes	97	-	-	-	-	-	1	-	-	2	1	4
Male	44	-	-	-	-	-	1	-	-	1	-	3
Female	53	-	-	-	-	-	-	-	-	1	1	1
White, both sexes	91	-	-	-	-	-	1	-	-	-	1	4
Male	43	-	-	-	-	-	1	-	-	-	-	3
Female	48	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	5	-	-	-	-	-	-	-	-	2	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	196	-	-	-	-	-	-	1	3	3	4	8
Male	82	-	-	-	-	-	-	-	2	2	2	8
Female	114	-	-	-	-	-	-	1	1	1	2	-
White, both sexes	167	-	-	-	-	-	-	1	3	1	1	6
Male	65	-	-	-	-	-	-	-	2	-	1	6
Female	102	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	26	-	-	-	-	-	-	-	-	2	2	2
Male	15	-	-	-	-	-	-	-	-	2	-	2
Female	11	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Minnesota												
All races, both sexes	93	-	-	-	-	-	1	-	-	-	-	3
Male	36	-	-	-	-	-	1	-	-	-	-	2
Female	57	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	91	-	-	-	-	-	1	-	-	-	-	3
Male	36	-	-	-	-	-	1	-	-	-	-	2
Female	55	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	29	-	-	-	-	-	-	-	-	1	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Missouri												
All races, both sexes	110	-	-	-	1	-	-	-	1	1	-	4
Male	45	-	-	-	1	-	-	-	-	1	-	4
Female	65	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	95	-	-	-	1	-	-	-	1	1	-	2
Male	39	-	-	-	1	-	-	-	-	1	-	2
Female	56	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	2
Male	6	-	-	-	-	-	-	-	-	-	-	2
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Nebraska												
All races, both sexes	62	-	-	-	-	-	-	-	-	-	-	-
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	45	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	60	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	37	-	-	-	-	-	1	-	-	-	3	4
Male	15	-	-	-	-	-	-	-	-	-	1	2
Female	22	-	-	-	-	-	1	-	-	-	2	2
White, both sexes	30	-	-	-	-	-	1	-	-	-	2	3
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	17	-	-	-	-	-	1	-	-	-	1	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
New Hampshire												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	211	-	-	-	-	-	1	-	1	3	8	7
Male	71	-	-	-	-	-	-	-	-	2	4	4
Female	140	-	-	-	-	-	1	-	1	1	4	3
White, both sexes	181	-	-	-	-	-	1	-	-	3	4	4
Male	62	-	-	-	-	-	-	-	-	2	3	3
Female	119	-	-	-	-	-	1	-	-	1	1	1
Black, both sexes	28	-	-	-	-	-	-	-	1	-	4	3
Male	9	-	-	-	-	-	-	-	-	-	1	1
Female	19	-	-	-	-	-	-	-	1	-	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
New Mexico												
All races, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	354	1	-	-	-	1	2	2	-	2	3	6
Male	134	-	-	-	-	1	1	1	-	2	2	5
Female	220	1	-	-	-	-	1	1	-	-	1	1
White, both sexes	328	-	-	-	-	1	2	2	-	1	3	4
Male	121	-	-	-	-	1	1	1	-	1	2	4
Female	207	-	-	-	-	-	1	1	-	-	1	-
Black, both sexes	24	1	-	-	-	-	-	-	-	1	-	2
Male	12	-	-	-	-	-	-	-	-	1	-	1
Female	12	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
North Carolina												
All races, both sexes	153	-	-	1	-	-	-	-	2	1	-	5
Male	59	-	-	1	-	-	-	-	1	-	-	3
Female	94	-	-	-	-	-	-	-	1	1	-	2
White, both sexes	123	-	-	-	-	-	-	-	-	-	-	3
Male	50	-	-	-	-	-	-	-	-	-	-	2
Female	73	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	27	-	-	1	-	-	-	-	1	1	-	2
Male	9	-	-	1	-	-	-	-	1	-	-	1
Female	18	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
North Dakota												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Ohio												
All races, both sexes	239	-	-	-	-	-	1	1	2	3	6	4
Male	85	-	-	-	-	-	-	-	1	3	5	3
Female	154	-	-	-	-	-	1	1	1	-	1	1
White, both sexes	215	-	-	-	-	-	-	-	2	3	5	2
Male	73	-	-	-	-	-	-	-	1	3	5	1
Female	142	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	23	-	-	-	-	-	1	1	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	-	2
Female	11	-	-	-	-	-	1	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	70	-	1	-	-	-	-	-	-	2	2	2
Male	23	-	1	-	-	-	-	-	-	1	2	2
Female	47	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	62	-	1	-	-	-	-	-	-	1	2	2
Male	20	-	1	-	-	-	-	-	-	1	2	2
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Oregon												
All races, both sexes	90	-	-	-	-	-	-	-	-	1	1	2
Male	35	-	-	-	-	-	-	-	-	1	1	1
Female	55	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	90	-	-	-	-	-	-	-	-	1	1	2
Male	35	-	-	-	-	-	-	-	-	1	1	1
Female	55	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	330	-	-	-	-	-	-	1	1	4	5	10
Male	130	-	-	-	-	-	-	1	-	3	4	6
Female	200	-	-	-	-	-	-	-	1	1	1	4
White, both sexes	303	-	-	-	-	-	-	1	-	4	3	9
Male	114	-	-	-	-	-	-	1	-	3	3	5
Female	189	-	-	-	-	-	-	-	-	1	-	4
Black, both sexes	26	-	-	-	-	-	-	-	1	-	2	1
Male	16	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Rhode Island												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	74	-	-	-	-	1	-	-	-	1	3	3
Male	28	-	-	-	-	1	-	-	-	1	-	1
Female	46	-	-	-	-	-	-	-	-	-	3	2
White, both sexes	49	-	-	-	-	1	-	-	-	-	-	2
Male	20	-	-	-	-	1	-	-	-	-	-	1
Female	29	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	25	-	-	-	-	-	-	-	-	1	3	1
Male	8	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	3	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
South Dakota												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	82	1	-	-	-	-	2	1	1	-	1	3
Male	30	1	-	-	-	-	2	1	-	-	-	1
Female	52	-	-	-	-	-	-	-	1	-	1	2
White, both sexes	74	1	-	-	-	-	2	-	1	-	1	3
Male	26	1	-	-	-	-	2	-	-	-	-	1
Female	48	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	8	-	-	-	-	-	-	1	-	-	-	-
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Texas												
All races, both sexes	273	-	-	-	-	-	-	4	4	2	9	10
Male	122	-	-	-	-	-	-	4	2	2	5	8
Female	151	-	-	-	-	-	-	-	2	-	4	2
White, both sexes	245	-	-	-	-	-	-	4	3	1	5	7
Male	106	-	-	-	-	-	-	4	2	1	4	5
Female	139	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	28	-	-	-	-	-	-	-	1	1	4	3
Male	16	-	-	-	-	-	-	-	-	1	1	3
Female	12	-	-	-	-	-	-	-	1	-	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	43	-	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	43	-	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Vermont												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	112	-	-	-	-	-	-	1	2	1	4	4
Male	43	-	-	-	-	-	-	-	2	1	3	1
Female	69	-	-	-	-	-	-	1	-	-	1	3
White, both sexes	92	-	-	-	-	-	-	-	-	1	4	1
Male	33	-	-	-	-	-	-	-	-	1	3	-
Female	59	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	15	-	-	-	-	-	-	1	1	-	-	3
Male	7	-	-	-	-	-	-	-	1	-	-	1
Female	8	-	-	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Washington												
All races, both sexes	152	-	-	-	-	-	1	-	-	2	1	3
Male	64	-	-	-	-	-	1	-	-	2	-	1
Female	88	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	143	-	-	-	-	-	1	-	-	1	-	3
Male	60	-	-	-	-	-	1	-	-	1	-	1
Female	83	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	3	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	47	-	-	-	-	-	-	1	-	-	-	2
Male	23	-	-	-	-	-	-	1	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	46	-	-	-	-	-	-	1	-	-	-	2
Male	23	-	-	-	-	-	-	1	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other diseases of endocardium (I36-I38)												
Wisconsin												
All races, both sexes	135	-	-	-	-	-	-	-	-	-	-	3
Male	55	-	-	-	-	-	-	-	-	-	-	3
Female	80	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	124	-	-	-	-	-	-	-	-	-	-	2
Male	49	-	-	-	-	-	-	-	-	-	-	2
Female	75	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
United States												
All races, both sexes	188	202	236	308	384	678	875	957	660	245	46	-
Male	107	125	124	137	168	281	323	309	165	43	6	-
Female	81	77	112	171	216	397	552	648	495	202	40	-
White, both sexes	128	143	177	264	326	608	814	919	637	236	40	-
Male	74	88	97	121	143	257	301	297	162	43	4	-
Female	54	55	80	143	183	351	513	622	475	193	36	-
Black, both sexes	56	56	49	37	51	55	40	23	15	5	3	-
Male	33	36	23	13	25	19	14	8	1	-	1	-
Female	23	20	26	24	26	36	26	15	14	5	2	-
American Indian 1/, both sexes	1	3	2	2	1	5	2	4	1	2	-	-
Male	-	1	1	1	-	-	2	1	-	-	-	-
Female	1	2	1	1	1	5	-	3	1	2	-	-
Asian or Pacific Islander 2/, both sexes	3	-	8	5	6	10	19	11	7	2	3	-
Male	-	-	3	2	-	5	6	3	2	-	1	-
Female	3	-	5	3	6	5	13	8	5	2	2	-
Alabama												
All races, both sexes	5	2	4	11	10	15	10	10	11	6	3	-
Male	3	1	-	2	3	5	4	3	2	-	-	-
Female	2	1	4	9	7	10	6	7	9	6	3	-
White, both sexes	2	-	2	10	9	12	9	8	9	6	2	-
Male	1	-	-	2	2	5	4	3	2	-	-	-
Female	1	-	2	8	7	7	5	5	7	6	2	-
Black, both sexes	3	2	2	1	1	3	1	2	2	-	1	-
Male	2	1	-	-	1	-	-	-	-	-	-	-
Female	1	1	2	1	-	3	1	2	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Alaska												
All races, both sexes	-	-	-	-	2	3	1	1	-	1	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	3	-	1	-	1	-	-
White, both sexes	-	-	-	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Arizona												
All races, both sexes	6	4	6	11	13	14	12	13	13	5	1	-
Male	4	2	4	4	9	4	5	7	4	3	1	-
Female	2	2	2	7	4	10	7	6	9	2	-	-
White, both sexes	5	4	6	10	13	13	11	13	13	5	1	-
Male	3	2	4	3	9	4	4	7	4	3	1	-
Female	2	2	2	7	4	9	7	6	9	2	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Arkansas												
All races, both sexes	3	3	3	4	3	8	8	7	4	4	-	-
Male	2	3	2	2	1	4	3	3	2	2	-	-
Female	1	-	1	2	2	4	5	4	2	2	-	-
White, both sexes	2	2	2	4	1	6	8	6	4	4	-	-
Male	1	2	1	2	-	2	3	3	2	2	-	-
Female	1	-	1	2	1	4	5	3	2	2	-	-
Black, both sexes	1	1	1	-	2	2	-	1	-	-	-	-
Male	1	1	1	-	1	2	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	15	8	11	15	28	41	64	70	57	19	6	-
Male	7	6	10	7	16	20	21	28	14	4	1	-
Female	8	2	1	8	12	21	43	42	43	15	5	-
White, both sexes	12	8	8	12	23	34	54	65	52	16	3	-
Male	6	6	7	7	15	19	17	27	13	4	-	-
Female	6	2	1	5	8	15	37	38	39	12	3	-
Black, both sexes	1	-	1	2	2	1	4	-	3	-	1	-
Male	1	-	1	-	1	-	3	-	-	-	-	-
Female	-	-	-	2	1	1	1	-	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	-	2	1	3	4	6	4	2	2	2	-
Male	-	-	2	-	-	1	1	1	1	-	1	-
Female	2	-	-	1	3	3	5	3	1	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Colorado												
All races, both sexes	1	2	3	5	5	14	15	17	11	6	-	-
Male	1	-	-	2	1	2	7	5	1	3	-	-
Female	-	2	3	3	4	12	8	12	10	3	-	-
White, both sexes	1	2	3	5	5	11	14	17	11	6	-	-
Male	1	-	-	2	1	1	7	5	1	3	-	-
Female	-	2	3	3	4	10	7	12	10	3	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Connecticut												
All races, both sexes	6	1	4	6	3	4	9	13	7	1	2	-
Male	6	1	2	6	3	1	5	4	1	1	-	-
Female	-	-	2	-	-	3	4	9	6	-	2	-
White, both sexes	5	1	2	5	3	4	9	13	7	1	2	-
Male	5	1	1	5	3	1	5	4	1	1	-	-
Female	-	-	1	-	-	3	4	9	6	-	2	-
Black, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	1	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Delaware												
All races, both sexes	-	1	-	4	2	2	3	2	3	-	-	-
Male	-	-	-	1	2	-	1	-	1	-	-	-
Female	-	1	-	3	-	2	2	2	2	-	-	-
White, both sexes	-	1	-	3	2	2	3	2	3	-	-	-
Male	-	-	-	1	2	-	1	-	1	-	-	-
Female	-	1	-	2	-	2	2	2	2	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	2	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Florida												
All races, both sexes	11	20	10	11	22	53	80	92	47	17	2	-
Male	5	14	5	5	9	21	25	26	19	4	1	-
Female	6	6	5	6	13	32	55	66	28	13	1	-
White, both sexes	7	13	7	7	19	52	77	90	47	17	1	-
Male	5	10	4	3	8	21	23	26	19	4	-	-
Female	2	3	3	4	11	31	54	64	28	13	1	-
Black, both sexes	4	7	3	4	3	1	3	2	-	-	1	-
Male	-	4	1	2	1	-	2	-	-	-	1	-
Female	4	3	2	2	2	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	9	7	10	9	12	23	22	8	12	4	-	-
Male	5	4	5	4	6	9	9	2	3	-	-	-
Female	4	3	5	5	6	14	13	6	9	4	-	-
White, both sexes	5	3	4	7	8	19	18	7	11	3	-	-
Male	2	3	2	3	4	8	9	2	3	-	-	-
Female	3	-	2	4	4	11	9	5	8	3	-	-
Black, both sexes	4	4	6	2	4	4	4	1	1	1	-	-
Male	3	1	3	1	2	1	-	-	-	-	-	-
Female	1	3	3	1	2	3	4	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Hawaii												
All races, both sexes	-	-	3	1	-	1	6	4	5	-	-	-
Male	-	-	-	1	-	1	2	-	1	-	-	-
Female	-	-	3	-	-	-	4	4	4	-	-	-
White, both sexes	-	-	1	-	-	-	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	1	6	2	4	-	-	-
Male	-	-	-	1	-	1	2	-	1	-	-	-
Female	-	-	2	-	-	-	4	2	3	-	-	-
Idaho												
All races, both sexes	2	1	-	-	1	5	4	10	7	3	-	-
Male	1	1	-	-	1	2	-	3	3	1	-	-
Female	1	-	-	-	-	3	4	7	4	2	-	-
White, both sexes	2	1	-	-	1	5	4	10	7	3	-	-
Male	1	1	-	-	1	2	-	3	3	1	-	-
Female	1	-	-	-	-	3	4	7	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Illinois												
All races, both sexes	14	10	9	10	10	20	31	33	15	8	3	-
Male	9	7	4	6	4	6	14	15	4	1	-	-
Female	5	3	5	4	6	14	17	18	11	7	3	-
White, both sexes	8	6	7	8	10	17	30	32	14	8	3	-
Male	6	4	4	4	4	4	14	14	4	1	-	-
Female	2	2	3	4	6	13	16	18	10	7	3	-
Black, both sexes	6	4	2	2	-	3	1	1	1	-	-	-
Male	3	3	-	2	-	2	-	1	-	-	-	-
Female	3	1	2	-	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	6	10	10	11	21	17	31	23	4	1	-
Male	1	2	3	2	-	9	6	8	3	-	-	-
Female	1	4	7	8	11	12	11	23	20	4	1	-
White, both sexes	2	6	10	10	9	21	17	31	22	4	1	-
Male	1	2	3	2	-	9	6	8	3	-	-	-
Female	1	4	7	8	9	12	11	23	19	4	1	-
Black, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Iowa												
All races, both sexes	1	-	2	-	3	1	6	7	4	2	-	-
Male	1	-	-	-	2	1	2	2	2	1	-	-
Female	-	-	2	-	1	-	4	5	2	1	-	-
White, both sexes	1	-	2	-	3	1	6	7	4	2	-	-
Male	1	-	-	-	2	1	2	2	2	1	-	-
Female	-	-	2	-	1	-	4	5	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	6	-	1	6	4	7	12	12	8	2	1	-
Male	3	-	-	1	2	4	4	2	2	-	-	-
Female	3	-	1	5	2	3	8	10	6	2	1	-
White, both sexes	5	-	1	6	4	7	12	12	8	2	1	-
Male	3	-	-	1	2	4	4	2	2	-	-	-
Female	2	-	1	5	2	3	8	10	6	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Kentucky												
All races, both sexes	5	2	3	5	11	7	7	9	5	4	1	-
Male	3	1	2	2	5	5	2	1	1	-	-	-
Female	2	1	1	3	6	2	5	8	4	4	1	-
White, both sexes	5	1	3	5	10	6	7	9	5	4	1	-
Male	3	1	2	2	5	4	2	1	1	-	-	-
Female	2	-	1	3	5	2	5	8	4	4	1	-
Black, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3	6	7	6	6	8	12	13	6	4	-	-
Male	2	1	1	4	2	5	8	4	-	1	-	-
Female	1	5	6	2	4	3	4	9	6	3	-	-
White, both sexes	-	2	5	5	1	8	11	11	6	2	-	-
Male	-	-	1	3	-	5	7	3	-	1	-	-
Female	-	2	4	2	1	3	4	8	6	1	-	-
Black, both sexes	3	4	2	1	5	-	1	2	-	2	-	-
Male	2	1	-	1	2	-	1	1	-	-	-	-
Female	1	3	2	-	3	-	-	1	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Maine												
All races, both sexes	3	1	-	-	3	5	5	4	1	2	-	-
Male	3	1	-	-	3	2	2	-	-	-	-	-
Female	-	-	-	-	-	3	3	4	1	2	-	-
White, both sexes	3	1	-	-	3	5	5	4	1	2	-	-
Male	3	1	-	-	3	2	2	-	-	-	-	-
Female	-	-	-	-	-	3	3	4	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5	4	2	9	5	10	11	12	2	2	1	-
Male	4	2	1	6	3	3	6	2	-	-	-	-
Female	1	2	1	3	2	7	5	10	2	2	1	-
White, both sexes	2	2	2	6	2	6	10	10	2	2	1	-
Male	2	-	1	5	1	1	6	2	-	-	-	-
Female	-	2	1	1	1	5	4	8	2	2	1	-
Black, both sexes	3	2	-	2	3	4	1	1	-	-	-	-
Male	2	2	-	-	2	2	-	-	-	-	-	-
Female	1	-	-	2	1	2	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Massachusetts												
All races, both sexes	2	3	6	7	10	15	15	18	8	3	2	-
Male	1	2	5	5	6	8	5	4	1	1	1	-
Female	1	1	1	2	4	7	10	14	7	2	1	-
White, both sexes	2	3	5	7	9	14	15	18	8	3	1	-
Male	1	2	5	5	6	8	5	4	1	1	1	-
Female	1	1	-	2	3	6	10	14	7	2	-	-
Black, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Michigan												
All races, both sexes	15	9	10	13	12	27	32	32	19	4	4	-
Male	8	4	6	7	2	14	13	10	2	1	1	-
Female	7	5	4	6	10	13	19	22	17	3	3	-
White, both sexes	11	8	7	10	11	21	29	31	19	4	4	-
Male	5	3	4	6	1	12	11	10	2	1	1	-
Female	6	5	3	4	10	9	18	21	17	3	3	-
Black, both sexes	4	1	3	3	1	5	2	1	-	-	-	-
Male	3	1	2	1	1	2	1	-	-	-	-	-
Female	1	-	1	2	-	3	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Minnesota												
All races, both sexes	4	1	2	2	5	8	17	28	15	7	-	-
Male	-	1	1	1	1	6	10	8	3	2	-	-
Female	4	-	1	1	4	2	7	20	12	5	-	-
White, both sexes	4	1	1	2	5	8	17	27	15	7	-	-
Male	-	1	1	1	1	6	10	8	3	2	-	-
Female	4	-	-	1	4	2	7	19	12	5	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	6	1	1	4	5	4	2	1	-	-	-
Male	-	4	1	1	1	2	-	2	1	-	-	-
Female	2	2	-	-	3	3	4	-	-	-	-	-
White, both sexes	1	2	1	1	3	3	4	2	1	-	-	-
Male	-	1	1	1	-	1	-	2	1	-	-	-
Female	1	1	-	-	3	2	4	-	-	-	-	-
Black, both sexes	1	4	-	-	1	2	-	-	-	-	-	-
Male	-	3	-	-	1	1	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Missouri												
All races, both sexes	1	8	4	3	6	17	17	24	15	8	-	-
Male	-	3	2	2	3	9	6	9	5	-	-	-
Female	1	5	2	1	3	8	11	15	10	8	-	-
White, both sexes	-	6	3	2	4	15	14	23	15	8	-	-
Male	-	2	2	2	3	7	6	8	5	-	-	-
Female	-	4	1	-	1	8	8	15	10	8	-	-
Black, both sexes	-	2	1	1	2	2	2	1	-	-	-	-
Male	-	1	-	-	-	2	-	1	-	-	-	-
Female	-	1	1	1	2	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	1	-	2	5	9	4	2	-	-
Male	-	-	-	1	-	-	1	3	2	1	-	-
Female	-	-	-	-	-	2	4	6	2	1	-	-
White, both sexes	-	-	-	1	-	2	5	9	4	2	-	-
Male	-	-	-	1	-	-	1	3	2	1	-	-
Female	-	-	-	-	-	2	4	6	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Nebraska												
All races, both sexes	-	-	1	4	5	6	17	10	10	7	2	-
Male	-	-	1	1	-	3	8	2	1	1	-	-
Female	-	-	-	3	5	3	9	8	9	6	2	-
White, both sexes	-	-	1	3	5	6	17	9	10	7	2	-
Male	-	-	1	1	-	3	8	1	1	1	-	-
Female	-	-	-	2	5	3	9	8	9	6	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	1	3	5	2	4	5	4	2	-	-	-
Male	1	1	-	4	1	1	2	2	-	-	-	-
Female	2	-	3	1	1	3	3	2	2	-	-	-
White, both sexes	2	1	2	5	1	3	5	4	1	-	-	-
Male	1	1	-	4	-	1	2	2	-	-	-	-
Female	1	-	2	1	1	2	3	2	1	-	-	-
Black, both sexes	1	-	-	-	1	1	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
New Hampshire												
All races, both sexes	-	-	1	3	1	3	7	5	6	1	1	-
Male	-	-	1	1	-	1	2	2	2	-	-	-
Female	-	-	-	2	1	2	5	3	4	1	1	-
White, both sexes	-	-	1	3	1	3	7	5	6	1	1	-
Male	-	-	1	1	-	1	2	2	2	-	-	-
Female	-	-	-	2	1	2	5	3	4	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	5	10	11	14	18	33	32	39	20	7	2	-
Male	1	5	4	10	7	12	7	11	4	-	-	-
Female	4	5	7	4	11	21	25	28	16	7	2	-
White, both sexes	1	8	6	13	16	30	30	38	18	7	2	-
Male	-	4	2	10	6	11	7	11	3	-	-	-
Female	1	4	4	3	10	19	23	27	15	7	2	-
Black, both sexes	4	2	4	1	2	3	1	1	2	-	-	-
Male	1	1	2	-	1	1	-	-	1	-	-	-
Female	3	1	2	1	1	2	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
New Mexico												
All races, both sexes	3	2	-	1	-	3	5	3	7	1	-	-
Male	2	1	-	1	-	2	2	2	2	-	-	-
Female	1	1	-	-	-	1	3	1	5	1	-	-
White, both sexes	3	2	-	1	-	3	5	3	7	1	-	-
Male	2	1	-	1	-	2	2	2	2	-	-	-
Female	1	1	-	-	-	1	3	1	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	14	13	23	21	51	65	70	48	24	2	-
Male	1	9	8	9	13	24	17	25	13	3	-	-
Female	5	5	5	14	8	27	48	45	35	21	2	-
White, both sexes	5	13	9	18	17	49	63	67	48	24	2	-
Male	1	8	6	8	11	23	15	23	13	3	-	-
Female	4	5	3	10	6	26	48	44	35	21	2	-
Black, both sexes	1	1	4	5	4	2	1	2	-	-	-	-
Male	-	1	2	1	2	1	1	2	-	-	-	-
Female	1	-	2	4	2	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
North Carolina												
All races, both sexes	3	10	11	7	10	16	26	29	24	8	-	-
Male	2	6	4	2	2	6	10	12	9	1	-	-
Female	1	4	7	5	8	10	16	17	15	7	-	-
White, both sexes	2	7	8	5	9	13	20	26	22	8	-	-
Male	2	4	4	2	2	5	8	11	9	1	-	-
Female	-	3	4	3	7	8	12	15	13	7	-	-
Black, both sexes	1	2	3	2	-	3	6	3	2	-	-	-
Male	-	2	-	-	-	1	2	1	-	-	-	-
Female	1	-	3	2	-	2	4	2	2	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	-	-	-	-	2	4	2	2	-	-
Male	-	1	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	4	1	2	-	-
White, both sexes	-	1	-	-	-	-	2	4	2	2	-	-
Male	-	1	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	4	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Ohio												
All races, both sexes	6	11	9	10	26	26	44	43	31	14	2	-
Male	6	8	6	3	9	6	15	13	4	3	-	-
Female	-	3	3	7	17	20	29	30	27	11	2	-
White, both sexes	5	6	6	7	23	23	44	42	31	14	2	-
Male	5	4	3	3	8	5	15	13	4	3	-	-
Female	-	2	3	4	15	18	29	29	27	11	2	-
Black, both sexes	1	5	3	2	3	3	-	1	-	-	-	-
Male	1	4	3	-	1	1	-	-	-	-	-	-
Female	-	1	-	2	2	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	2	5	3	6	13	6	9	12	5	-	-
Male	-	2	2	-	3	5	2	1	-	2	-	-
Female	2	-	3	3	3	8	4	8	12	3	-	-
White, both sexes	1	-	5	3	4	13	6	8	11	5	-	-
Male	-	-	2	-	2	5	2	1	-	2	-	-
Female	1	-	3	3	2	8	4	7	11	3	-	-
Black, both sexes	1	1	-	-	1	-	-	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Oregon												
All races, both sexes	-	5	-	4	5	13	17	19	17	6	-	-
Male	-	4	-	-	-	9	8	9	2	-	-	-
Female	-	1	-	4	5	4	9	10	15	6	-	-
White, both sexes	-	5	-	4	5	13	17	19	17	6	-	-
Male	-	4	-	-	-	9	8	9	2	-	-	-
Female	-	1	-	4	5	4	9	10	15	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	12	7	19	18	24	51	51	58	55	13	1	-
Male	8	5	12	9	14	21	18	17	12	-	-	-
Female	4	2	7	9	10	30	33	41	43	13	1	-
White, both sexes	7	2	15	18	23	47	49	58	54	12	1	-
Male	4	2	9	9	13	20	16	17	12	-	-	-
Female	3	-	6	9	10	27	33	41	42	12	1	-
Black, both sexes	5	5	3	-	1	4	2	-	1	1	-	-
Male	4	3	3	-	1	1	2	-	-	-	-	-
Female	1	2	-	-	-	3	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Rhode Island												
All races, both sexes	1	-	-	1	2	1	2	2	2	3	-	-
Male	1	-	-	1	-	-	1	2	-	1	-	-
Female	-	-	-	-	2	1	1	-	2	2	-	-
White, both sexes	-	-	-	1	2	1	2	1	2	3	-	-
Male	-	-	-	1	-	-	1	1	-	1	-	-
Female	-	-	-	-	2	1	1	-	2	2	-	-
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	5	6	7	6	9	15	8	8	1	1	-
Male	-	3	4	4	2	3	6	1	2	-	-	-
Female	-	2	2	3	4	6	9	7	6	1	1	-
White, both sexes	-	4	2	3	2	8	11	7	7	1	1	-
Male	-	3	2	1	1	3	5	1	2	-	-	-
Female	-	1	-	2	1	5	6	6	5	1	1	-
Black, both sexes	-	1	4	4	4	1	4	1	1	-	-	-
Male	-	-	2	3	1	-	1	-	-	-	-	-
Female	-	1	2	1	3	1	3	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
South Dakota												
All races, both sexes	-	-	1	1	1	3	5	3	5	2	-	-
Male	-	-	1	1	-	2	3	1	1	-	-	-
Female	-	-	-	-	1	1	2	2	4	2	-	-
White, both sexes	-	-	-	1	1	3	5	3	5	2	-	-
Male	-	-	-	1	-	2	3	1	1	-	-	-
Female	-	-	-	-	1	1	2	2	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	5	7	8	4	16	16	8	6	3	-	-
Male	-	4	4	3	1	6	4	2	1	-	-	-
Female	-	1	3	5	3	10	12	6	5	3	-	-
White, both sexes	-	3	5	8	4	14	16	8	6	2	-	-
Male	-	2	4	3	1	5	4	2	1	-	-	-
Female	-	1	1	5	3	9	12	6	5	2	-	-
Black, both sexes	-	2	2	-	-	2	-	-	-	1	-	-
Male	-	2	-	-	-	1	-	-	-	-	-	-
Female	-	-	2	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Texas												
All races, both sexes	16	6	15	21	25	40	41	47	23	8	2	-
Male	9	4	10	4	12	20	17	14	9	2	-	-
Female	7	2	5	17	13	20	24	33	14	6	2	-
White, both sexes	11	5	13	20	21	35	40	47	23	8	2	-
Male	5	3	8	4	9	19	17	14	9	2	-	-
Female	6	2	5	16	12	16	23	33	14	6	2	-
Black, both sexes	5	1	2	1	4	5	1	-	-	-	-	-
Male	4	1	2	-	3	1	-	-	-	-	-	-
Female	1	-	-	1	1	4	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	1	-	3	3	9	9	6	5	3	1	-
Male	1	1	-	-	2	2	2	2	1	1	-	-
Female	-	-	-	3	1	7	7	4	4	2	1	-
White, both sexes	1	1	-	3	3	9	9	6	5	3	1	-
Male	1	1	-	-	2	2	2	2	1	1	-	-
Female	-	-	-	3	1	7	7	4	4	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Vermont												
All races, both sexes	-	-	1	-	3	-	1	-	3	3	1	-
Male	-	-	1	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	2	-	1	-	3	2	1	-
White, both sexes	-	-	1	-	3	-	1	-	3	3	1	-
Male	-	-	1	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	2	-	1	-	3	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	5	8	8	10	12	20	18	12	4	1	-
Male	1	4	4	4	6	3	5	6	3	-	-	-
Female	1	1	4	4	4	9	15	12	9	4	1	-
White, both sexes	2	3	8	7	8	11	13	17	12	4	1	-
Male	1	2	4	3	5	2	3	6	3	-	-	-
Female	1	1	4	4	3	9	10	11	9	4	1	-
Black, both sexes	-	2	-	1	2	-	5	-	-	-	-	-
Male	-	2	-	1	1	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	4	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Washington												
All races, both sexes	3	5	6	5	7	11	27	48	26	4	3	-
Male	2	1	3	1	4	2	16	19	11	-	1	-
Female	1	4	3	4	3	9	11	29	15	4	2	-
White, both sexes	3	4	5	4	6	11	26	47	25	4	3	-
Male	2	1	2	1	4	2	15	18	11	-	1	-
Female	1	3	3	3	2	9	11	29	14	4	2	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	1	1	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
West Virginia												
All races, both sexes	-	2	4	5	1	3	11	10	6	2	-	-
Male	-	2	2	3	1	3	4	4	1	1	-	-
Female	-	-	2	2	-	-	7	6	5	1	-	-
White, both sexes	-	2	4	5	1	3	11	9	6	2	-	-
Male	-	2	2	3	1	3	4	4	1	1	-	-
Female	-	-	2	2	-	-	7	5	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other diseases of endocardium (I36-I38)												
Wisconsin												
All races, both sexes	2	4	4	6	12	18	21	33	27	5	-	-
Male	1	3	3	3	4	7	10	11	9	1	-	-
Female	1	1	1	3	8	11	11	22	18	4	-	-
White, both sexes	-	2	4	5	11	16	21	31	27	5	-	-
Male	-	2	3	3	3	7	10	9	9	1	-	-
Female	-	-	1	2	8	9	11	22	18	4	-	-
Black, both sexes	2	2	-	1	1	2	-	1	-	-	-	-
Male	1	1	-	-	1	-	-	1	-	-	-	-
Female	1	1	-	1	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	1	-	1	1	1	3	-	-	1	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	2	-	-	1	-	-
White, both sexes	-	1	-	1	1	1	3	-	-	1	-	-
Male	-	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	2	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
United States												
All races, both sexes	25,505	82	36	27	45	118	197	258	364	541	836	1,215
Male	14,963	38	18	13	28	90	142	194	268	370	628	868
Female	10,542	44	18	14	17	28	55	64	96	171	208	347
White, both sexes	20,536	57	24	18	30	73	115	157	211	322	555	786
Male	12,014	26	13	9	18	58	85	126	158	231	434	582
Female	8,522	31	11	9	12	15	30	31	53	91	121	204
Black, both sexes	4,408	19	8	8	14	42	72	92	142	196	257	399
Male	2,630	11	3	3	10	30	50	62	102	126	177	265
Female	1,778	8	5	5	4	12	22	30	40	70	80	134
American Indian 1/, both sexes	139	1	-	-	-	1	3	3	3	5	11	13
Male	81	-	-	-	-	1	1	2	3	3	8	9
Female	58	1	-	-	-	-	2	1	-	2	3	4
Asian or Pacific Islander 2/, both sexes	422	5	4	1	1	2	7	6	8	18	13	17
Male	238	1	2	1	-	1	6	4	5	10	9	12
Female	184	4	2	-	1	1	1	2	3	8	4	5
Alabama												
All races, both sexes	462	-	1	-	-	1	4	5	15	10	10	15
Male	261	-	-	-	-	1	3	4	9	7	7	10
Female	201	-	1	-	-	-	1	1	6	3	3	5
White, both sexes	306	-	-	-	-	1	-	3	5	2	3	6
Male	177	-	-	-	-	1	-	2	3	2	2	4
Female	129	-	-	-	-	-	-	1	2	-	1	2
Black, both sexes	154	-	1	-	-	-	4	2	10	6	7	9
Male	84	-	-	-	-	-	3	2	6	5	5	6
Female	70	-	1	-	-	-	1	-	4	1	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Alaska												
All races, both sexes	37	-	-	-	-	-	4	2	-	-	2	2
Male	22	-	-	-	-	-	3	1	-	-	1	1
Female	15	-	-	-	-	-	1	1	-	-	1	1
White, both sexes	22	-	-	-	-	-	2	1	-	-	2	-
Male	14	-	-	-	-	-	2	1	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	1	1	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	1	1	-	-	-	1
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	1	-	-	-	-	1
Male	2	-	-	-	-	-	1	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	373	4	-	1	1	3	1	3	4	11	18	16
Male	238	-	-	1	1	2	1	1	3	8	14	12
Female	135	4	-	-	-	1	-	2	1	3	4	4
White, both sexes	334	3	-	1	1	3	1	1	2	8	15	14
Male	217	-	-	1	1	2	1	-	2	6	11	11
Female	117	3	-	-	-	1	-	1	-	2	4	3
Black, both sexes	18	-	-	-	-	-	-	2	2	2	2	1
Male	10	-	-	-	-	-	-	1	1	1	2	-
Female	8	-	-	-	-	-	-	1	1	1	-	1
American Indian 1/, both sexes	16	-	-	-	-	-	-	-	-	1	1	1
Male	10	-	-	-	-	-	-	-	-	1	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Arkansas												
All races, both sexes	306	2	1	2	-	2	4	4	2	9	11	9
Male	155	-	-	-	-	2	3	3	1	6	9	4
Female	151	2	1	2	-	-	1	1	1	3	2	5
White, both sexes	244	1	1	2	-	2	1	-	2	2	5	5
Male	121	-	-	-	-	2	1	-	1	1	5	1
Female	123	1	1	2	-	-	-	-	1	1	-	4
Black, both sexes	62	1	-	-	-	-	3	4	-	7	6	4
Male	34	-	-	-	-	-	2	3	-	5	4	3
Female	28	1	-	-	-	-	1	1	-	2	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2,938	9	8	2	8	14	25	49	43	81	115	176
Male	1,819	4	5	-	4	11	18	37	32	60	94	129
Female	1,119	5	3	2	4	3	7	12	11	21	21	47
White, both sexes	2,377	9	4	2	7	7	18	32	32	48	92	127
Male	1,494	4	3	-	3	5	14	28	25	37	79	98
Female	883	5	1	2	4	2	4	4	7	11	13	29
Black, both sexes	393	-	2	-	1	5	5	14	9	23	20	42
Male	234	-	-	-	1	4	2	8	6	17	14	24
Female	159	-	2	-	-	1	3	6	3	6	6	18
American Indian 1/, both sexes	17	-	-	-	-	1	-	-	-	1	1	1
Male	10	-	-	-	-	1	-	-	-	1	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	151	-	2	-	-	1	2	3	2	9	2	6
Male	81	-	2	-	-	1	2	1	1	5	1	6
Female	70	-	-	-	-	-	-	2	1	4	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Colorado												
All races, both sexes	259	2	-	1	-	4	-	-	4	10	11	21
Male	153	1	-	-	-	4	-	-	1	8	7	15
Female	106	1	-	1	-	-	-	-	3	2	4	6
White, both sexes	248	1	-	1	-	4	-	-	4	9	9	18
Male	146	1	-	-	-	4	-	-	1	7	6	13
Female	102	-	-	1	-	-	-	-	3	2	3	5
Black, both sexes	9	-	-	-	-	-	-	-	-	1	1	3
Male	6	-	-	-	-	-	-	-	-	1	-	2
Female	3	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	334	2	1	-	1	1	1	2	5	6	9	20
Male	204	-	-	-	-	1	-	2	5	4	8	17
Female	130	2	1	-	1	-	1	-	-	2	1	3
White, both sexes	310	2	1	-	-	1	1	1	3	6	8	18
Male	188	-	-	-	-	1	-	1	3	4	7	15
Female	122	2	1	-	-	-	1	-	-	2	1	3
Black, both sexes	23	-	-	-	1	-	-	1	2	-	1	2
Male	16	-	-	-	-	-	-	1	2	-	1	2
Female	7	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Delaware												
All races, both sexes	85	-	-	-	-	-	-	-	2	-	4	4
Male	53	-	-	-	-	-	-	-	2	-	3	3
Female	32	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	65	-	-	-	-	-	-	-	1	-	1	1
Male	40	-	-	-	-	-	-	-	1	-	1	1
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	19	-	-	-	-	-	-	-	1	-	3	3
Male	12	-	-	-	-	-	-	-	1	-	2	2
Female	7	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	45	-	-	-	1	1	-	-	-	1	3	5
Male	23	-	-	-	1	1	-	-	-	-	1	2
Female	22	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	40	-	-	-	1	1	-	-	-	-	3	5
Male	21	-	-	-	1	1	-	-	-	-	1	2
Female	19	-	-	-	-	-	-	-	-	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Florida												
All races, both sexes	2,385	7	1	1	2	8	17	12	30	35	57	79
Male	1,399	5	-	-	1	8	12	9	20	23	44	55
Female	986	2	1	1	1	-	5	3	10	12	13	24
White, both sexes	2,017	5	1	-	1	4	9	4	15	21	42	44
Male	1,193	3	-	-	1	4	5	3	9	16	37	32
Female	824	2	1	-	-	-	4	1	6	5	5	12
Black, both sexes	348	1	-	1	1	4	7	7	14	14	14	34
Male	195	1	-	-	-	4	6	5	11	7	6	23
Female	153	-	-	1	1	-	1	2	3	7	8	11
American Indian 1/, both sexes	3	-	-	-	-	-	-	1	-	-	1	-
Male	3	-	-	-	-	-	-	1	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	1	-	-	-	-	1	-	1	-	-	1
Male	8	1	-	-	-	-	1	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	1	-	-	1
Georgia												
All races, both sexes	818	2	2	1	4	5	6	6	12	20	13	45
Male	468	2	1	1	-	3	5	4	8	11	7	29
Female	350	-	1	-	4	2	1	2	4	9	6	16
White, both sexes	535	1	-	1	3	1	-	3	4	6	3	21
Male	300	1	-	1	-	1	-	2	2	4	3	14
Female	235	-	-	-	3	-	-	1	2	2	-	7
Black, both sexes	275	1	1	-	1	4	4	3	8	14	10	23
Male	163	1	1	-	-	2	3	2	6	7	4	14
Female	112	-	-	-	1	2	1	1	2	7	6	9
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	1	-	-	-	2	-	-	-	-	1
Male	5	-	-	-	-	-	2	-	-	-	-	1
Female	3	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Hawaii												
All races, both sexes	156	-	-	-	-	-	-	-	3	3	10	6
Male	95	-	-	-	-	-	-	-	3	2	8	3
Female	61	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	34	-	-	-	-	-	-	-	1	-	1	-
Male	23	-	-	-	-	-	-	-	1	-	1	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	119	-	-	-	-	-	-	-	2	3	9	6
Male	72	-	-	-	-	-	-	-	2	2	7	3
Female	47	-	-	-	-	-	-	-	-	1	2	3
Idaho												
All races, both sexes	79	-	1	-	-	1	1	-	-	3	1	4
Male	37	-	1	-	-	1	-	-	-	-	-	4
Female	42	-	-	-	-	-	1	-	-	3	1	-
White, both sexes	78	-	1	-	-	1	1	-	-	3	1	4
Male	37	-	1	-	-	1	-	-	-	-	-	4
Female	41	-	-	-	-	-	1	-	-	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Illinois												
All races, both sexes	1,134	2	-	-	2	8	10	18	16	35	56	64
Male	659	1	-	-	1	5	5	16	11	26	45	49
Female	475	1	-	-	1	3	5	2	5	9	11	15
White, both sexes	851	1	-	-	-	4	4	8	4	22	39	32
Male	474	-	-	-	-	3	2	8	4	16	31	26
Female	377	1	-	-	-	1	2	-	-	6	8	6
Black, both sexes	269	1	-	-	1	4	5	10	11	12	17	32
Male	177	1	-	-	1	2	3	8	6	9	14	23
Female	92	-	-	-	-	2	2	2	5	3	3	9
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	1	-	1	-	1	-	-	-
Male	7	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	1	-	1	-	-	-	-	-
Indiana												
All races, both sexes	610	2	1	1	1	4	7	8	11	17	23	28
Male	358	1	-	-	1	2	5	6	10	13	18	18
Female	252	1	1	1	-	2	2	2	1	4	5	10
White, both sexes	534	2	-	-	-	4	7	7	7	15	16	18
Male	313	1	-	-	-	2	5	5	6	12	13	11
Female	221	1	-	-	-	2	2	2	1	3	3	7
Black, both sexes	72	-	-	1	1	-	-	1	4	2	7	10
Male	43	-	-	-	1	-	-	1	4	1	5	7
Female	29	-	-	1	-	-	-	-	-	1	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	1	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Iowa												
All races, both sexes	212	-	-	-	-	1	-	1	3	5	6	9
Male	113	-	-	-	-	1	-	1	1	4	4	5
Female	99	-	-	-	-	-	-	-	2	1	2	4
White, both sexes	205	-	-	-	-	1	-	1	3	4	5	7
Male	110	-	-	-	-	1	-	1	1	3	3	4
Female	95	-	-	-	-	-	-	-	2	1	2	3
Black, both sexes	7	-	-	-	-	-	-	-	-	1	1	2
Male	3	-	-	-	-	-	-	-	-	1	1	1
Female	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	181	-	-	1	-	3	-	4	3	3	5	8
Male	98	-	-	1	-	2	-	3	2	2	4	6
Female	83	-	-	-	-	1	-	1	1	1	1	2
White, both sexes	160	-	-	1	-	2	-	4	2	2	5	7
Male	86	-	-	1	-	1	-	3	2	2	4	5
Female	74	-	-	-	-	1	-	1	-	-	1	2
Black, both sexes	19	-	-	-	-	1	-	-	1	1	-	1
Male	10	-	-	-	-	1	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Kentucky												
All races, both sexes	338	2	2	-	2	1	2	2	3	3	9	16
Male	183	-	1	-	2	1	2	-	3	3	6	11
Female	155	2	1	-	-	-	-	2	-	-	3	5
White, both sexes	304	1	2	-	2	1	2	1	1	2	6	12
Male	160	-	1	-	2	1	2	-	1	2	3	8
Female	144	1	1	-	-	-	-	1	-	-	3	4
Black, both sexes	33	-	-	-	-	-	-	1	2	1	3	4
Male	23	-	-	-	-	-	-	-	2	1	3	3
Female	10	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	476	-	-	1	-	1	5	10	4	19	11	21
Male	270	-	-	1	-	1	4	8	4	11	9	16
Female	206	-	-	-	-	-	1	2	-	8	2	5
White, both sexes	304	-	-	-	-	1	1	3	4	6	3	10
Male	170	-	-	-	-	1	1	3	4	4	3	9
Female	134	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	170	-	-	1	-	-	4	7	-	12	8	11
Male	99	-	-	1	-	-	3	5	-	6	6	7
Female	71	-	-	-	-	-	1	2	-	6	2	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Maine												
All races, both sexes	112	-	1	-	-	1	1	-	1	1	-	5
Male	63	-	-	-	-	1	-	-	-	-	-	5
Female	49	-	1	-	-	-	1	-	1	1	-	-
White, both sexes	112	-	1	-	-	1	1	-	1	1	-	5
Male	63	-	-	-	-	1	-	-	-	-	-	5
Female	49	-	1	-	-	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	461	2	2	-	-	3	2	4	13	6	12	22
Male	266	1	1	-	-	3	1	4	10	3	9	16
Female	195	1	1	-	-	-	1	-	3	3	3	6
White, both sexes	307	-	1	-	-	1	2	3	3	4	5	11
Male	179	-	1	-	-	1	1	3	2	2	4	10
Female	128	-	-	-	-	-	1	-	1	2	1	1
Black, both sexes	144	2	1	-	-	2	-	1	10	2	7	11
Male	84	1	-	-	-	2	-	1	8	1	5	6
Female	60	1	1	-	-	-	-	-	2	1	2	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Massachusetts												
All races, both sexes	412	1	-	-	-	-	4	5	1	3	9	21
Male	263	1	-	-	-	-	4	5	-	3	7	15
Female	149	-	-	-	-	-	-	-	1	-	2	6
White, both sexes	382	1	-	-	-	-	2	5	1	3	8	17
Male	244	1	-	-	-	-	2	5	-	3	6	13
Female	138	-	-	-	-	-	-	-	1	-	2	4
Black, both sexes	27	-	-	-	-	-	2	-	-	-	1	4
Male	18	-	-	-	-	-	2	-	-	-	1	2
Female	9	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,029	3	-	4	3	3	8	16	22	31	41	80
Male	621	2	-	1	3	3	5	11	19	22	32	54
Female	408	1	-	3	-	-	3	5	3	9	9	26
White, both sexes	817	2	-	1	2	2	5	12	11	16	29	52
Male	490	1	-	1	2	2	4	10	11	13	26	39
Female	327	1	-	-	-	-	1	2	-	3	3	13
Black, both sexes	207	1	-	3	1	1	3	4	10	15	12	27
Male	129	1	-	-	1	1	1	1	7	9	6	15
Female	78	-	-	3	-	-	2	3	3	6	6	12
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Minnesota												
All races, both sexes	324	-	-	-	-	1	2	3	2	2	13	11
Male	202	-	-	-	-	1	2	3	2	1	7	9
Female	122	-	-	-	-	-	-	-	-	1	6	2
White, both sexes	304	-	-	-	-	1	2	2	2	1	10	8
Male	186	-	-	-	-	1	2	2	2	-	6	7
Female	118	-	-	-	-	-	-	-	-	1	4	1
Black, both sexes	14	-	-	-	-	-	-	1	-	1	3	2
Male	10	-	-	-	-	-	-	1	-	1	1	1
Female	4	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	276	-	-	1	-	2	2	-	5	2	5	9
Male	152	-	-	1	-	1	-	-	4	1	4	8
Female	124	-	-	-	-	1	2	-	1	1	1	1
White, both sexes	154	-	-	-	-	-	-	-	1	-	2	2
Male	91	-	-	-	-	-	-	-	-	-	2	2
Female	63	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	122	-	-	1	-	2	2	-	4	2	3	7
Male	61	-	-	1	-	1	-	-	4	1	2	6
Female	61	-	-	-	-	1	2	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Missouri												
All races, both sexes	555	2	-	1	1	3	3	4	10	10	18	21
Male	306	1	-	-	1	2	3	2	8	6	11	13
Female	249	1	-	1	-	1	-	2	2	4	7	8
White, both sexes	472	2	-	1	1	1	3	2	8	7	9	17
Male	260	1	-	-	1	-	3	2	6	4	6	10
Female	212	1	-	1	-	1	-	-	2	3	3	7
Black, both sexes	81	-	-	-	-	2	-	2	2	3	9	4
Male	45	-	-	-	-	2	-	-	2	2	5	3
Female	36	-	-	-	-	-	-	2	-	1	4	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	66	-	1	-	1	-	-	-	1	1	4	4
Male	35	-	1	-	1	-	-	-	-	1	2	3
Female	31	-	-	-	-	-	-	-	1	-	2	1
White, both sexes	62	-	1	-	1	-	-	-	1	1	3	2
Male	31	-	1	-	1	-	-	-	-	1	1	1
Female	31	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	2
Male	4	-	-	-	-	-	-	-	-	-	1	2
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Nebraska												
All races, both sexes	117	-	-	-	-	1	2	-	1	-	2	3
Male	73	-	-	-	-	1	1	-	-	-	2	2
Female	44	-	-	-	-	-	1	-	1	-	-	1
White, both sexes	106	-	-	-	-	1	-	-	-	-	2	3
Male	67	-	-	-	-	1	-	-	-	-	2	2
Female	39	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	1	-	1	-	-	-
Male	4	-	-	-	-	-	1	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	267	1	-	-	-	-	4	10	2	10	17	16
Male	170	-	-	-	-	-	4	8	2	7	14	12
Female	97	1	-	-	-	-	-	2	-	3	3	4
White, both sexes	226	-	-	-	-	-	3	8	2	8	12	13
Male	145	-	-	-	-	-	3	7	2	6	10	10
Female	81	-	-	-	-	-	-	1	-	2	2	3
Black, both sexes	30	1	-	-	-	-	-	2	-	2	3	3
Male	16	-	-	-	-	-	-	1	-	1	2	2
Female	14	1	-	-	-	-	-	1	-	1	1	1
American Indian 1/, both sexes	4	-	-	-	-	-	1	-	-	-	1	-
Male	4	-	-	-	-	-	1	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
New Hampshire												
All races, both sexes	93	-	-	-	-	1	-	3	-	3	2	4
Male	55	-	-	-	-	-	-	3	-	3	1	4
Female	38	-	-	-	-	1	-	-	-	-	1	-
White, both sexes	91	-	-	-	-	1	-	3	-	3	2	4
Male	53	-	-	-	-	-	-	3	-	3	1	4
Female	38	-	-	-	-	1	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	845	-	-	-	1	4	6	8	10	18	25	47
Male	537	-	-	-	1	2	6	7	9	13	17	34
Female	308	-	-	-	-	2	-	1	1	5	8	13
White, both sexes	694	-	-	-	1	3	4	5	6	9	17	32
Male	436	-	-	-	1	2	4	5	6	8	10	25
Female	258	-	-	-	-	1	-	-	-	1	7	7
Black, both sexes	140	-	-	-	-	1	2	3	3	8	8	15
Male	92	-	-	-	-	-	2	2	2	5	7	9
Female	48	-	-	-	-	1	-	1	1	3	1	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	1	1	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
New Mexico												
All races, both sexes	106	1	-	-	-	-	-	1	-	2	9	6
Male	58	-	-	-	-	-	-	-	-	2	6	3
Female	48	1	-	-	-	-	-	1	-	-	3	3
White, both sexes	91	-	-	-	-	-	-	1	-	2	6	6
Male	50	-	-	-	-	-	-	-	-	2	4	3
Female	41	-	-	-	-	-	-	1	-	-	2	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	1	-	-	-	-	-	-	-	-	3	-
Male	8	-	-	-	-	-	-	-	-	-	2	-
Female	5	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,111	9	2	1	1	4	4	7	10	13	28	46
Male	650	5	1	1	1	2	3	6	5	11	19	36
Female	461	4	1	-	-	2	1	1	5	2	9	10
White, both sexes	944	6	2	1	-	1	2	3	8	6	13	36
Male	549	4	1	1	-	1	1	3	5	5	8	26
Female	395	2	1	-	-	-	1	-	3	1	5	10
Black, both sexes	152	2	-	-	1	2	2	3	2	7	15	10
Male	92	1	-	-	1	1	2	2	-	6	11	10
Female	60	1	-	-	-	1	-	1	2	1	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	1	-	-	-	1	-	1	-	-	-	-
Male	9	-	-	-	-	-	-	1	-	-	-	-
Female	6	1	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
North Carolina												
All races, both sexes	866	1	1	2	2	2	11	6	14	19	31	39
Male	492	-	-	2	1	1	10	3	7	9	24	30
Female	374	1	1	-	1	1	1	3	7	10	7	9
White, both sexes	602	1	1	1	1	2	7	3	7	11	18	17
Male	338	-	-	1	1	1	7	1	4	5	13	13
Female	264	1	1	-	-	1	-	2	3	6	5	4
Black, both sexes	254	-	-	1	1	-	4	2	7	8	13	21
Male	149	-	-	1	-	-	3	1	3	4	11	16
Female	105	-	-	-	1	-	1	1	4	4	2	5
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	45	-	-	-	-	-	-	1	1	-	3	1
Male	24	-	-	-	-	-	-	1	-	-	2	-
Female	21	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	41	-	-	-	-	-	-	1	1	-	-	-
Male	22	-	-	-	-	-	-	1	-	-	-	-
Female	19	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	2	1
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Ohio												
All races, both sexes	1,084	7	4	2	3	5	7	10	16	30	25	37
Male	635	3	3	1	3	5	4	6	13	21	16	27
Female	449	4	1	1	-	-	3	4	3	9	9	10
White, both sexes	907	5	4	2	2	5	5	8	10	19	13	28
Male	524	3	3	1	2	5	3	4	8	14	8	21
Female	383	2	1	1	-	-	2	4	2	5	5	7
Black, both sexes	177	2	-	-	1	-	2	2	6	11	12	9
Male	111	-	-	-	1	-	1	2	5	7	8	6
Female	66	2	-	-	-	-	1	-	1	4	4	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	271	-	-	-	-	3	2	5	6	6	9	20
Male	159	-	-	-	-	2	2	4	4	4	5	15
Female	112	-	-	-	-	1	-	1	2	2	4	5
White, both sexes	211	-	-	-	-	1	2	3	4	3	6	13
Male	128	-	-	-	-	1	2	2	2	2	3	11
Female	83	-	-	-	-	-	-	1	2	1	3	2
Black, both sexes	39	-	-	-	-	2	-	1	1	3	3	6
Male	22	-	-	-	-	1	-	1	1	2	2	4
Female	17	-	-	-	-	1	-	-	-	1	1	2
American Indian 1/, both sexes	20	-	-	-	-	-	-	1	1	-	-	1
Male	9	-	-	-	-	-	-	1	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Oregon												
All races, both sexes	254	-	-	-	1	3	3	-	3	8	6	9
Male	150	-	-	-	-	3	3	-	3	6	3	5
Female	104	-	-	-	1	-	-	-	-	2	3	4
White, both sexes	235	-	-	-	1	2	2	-	3	6	6	8
Male	139	-	-	-	-	2	2	-	3	5	3	4
Female	96	-	-	-	1	-	-	-	-	1	3	4
Black, both sexes	10	-	-	-	-	1	1	-	-	1	-	-
Male	5	-	-	-	-	1	1	-	-	1	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,563	6	1	1	-	5	12	8	12	18	54	64
Male	881	2	1	-	-	5	9	7	7	13	42	48
Female	682	4	-	1	-	-	3	1	5	5	12	16
White, both sexes	1,361	6	-	1	-	3	8	4	9	12	33	47
Male	750	2	-	-	-	3	5	3	6	8	26	35
Female	611	4	-	1	-	-	3	1	3	4	7	12
Black, both sexes	198	-	1	-	-	2	4	4	3	6	21	17
Male	129	-	1	-	-	2	4	4	1	5	16	13
Female	69	-	-	-	-	-	-	-	2	1	5	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Rhode Island												
All races, both sexes	50	1	-	-	2	-	1	-	3	1	2	-
Male	30	1	-	-	1	-	-	-	2	-	2	-
Female	20	-	-	-	1	-	1	-	1	1	-	-
White, both sexes	47	1	-	-	2	-	1	-	2	1	2	-
Male	28	1	-	-	1	-	-	-	1	-	2	-
Female	19	-	-	-	1	-	1	-	1	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	485	3	1	1	2	3	1	6	7	8	14	22
Male	286	2	-	-	1	2	-	5	5	6	8	17
Female	199	1	1	1	1	1	1	1	2	2	6	5
White, both sexes	295	1	-	1	1	1	-	3	2	4	6	9
Male	171	1	-	-	-	1	-	2	1	3	4	8
Female	124	-	-	1	1	-	-	1	1	1	2	1
Black, both sexes	189	2	1	-	1	2	1	3	5	4	8	13
Male	114	1	-	-	1	1	-	3	4	3	4	9
Female	75	1	1	-	-	1	1	-	1	1	4	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
South Dakota												
All races, both sexes	41	-	-	-	-	-	-	-	2	1	-	1
Male	17	-	-	-	-	-	-	-	1	1	-	1
Female	24	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	37	-	-	-	-	-	-	-	2	1	-	-
Male	15	-	-	-	-	-	-	-	1	1	-	-
Female	22	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	504	-	-	-	-	5	3	8	8	8	17	26
Male	282	-	-	-	-	3	2	5	7	3	15	15
Female	222	-	-	-	-	2	1	3	1	5	2	11
White, both sexes	327	-	-	-	-	1	-	1	2	3	8	13
Male	179	-	-	-	-	1	-	1	2	1	7	7
Female	148	-	-	-	-	-	-	-	-	2	1	6
Black, both sexes	175	-	-	-	-	4	3	7	6	5	9	13
Male	102	-	-	-	-	2	2	4	5	2	8	8
Female	73	-	-	-	-	2	1	3	1	3	1	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Texas												
All races, both sexes	1,499	8	4	2	3	7	18	18	29	36	63	80
Male	891	4	3	2	3	5	12	13	24	29	49	55
Female	608	4	1	-	-	2	6	5	5	7	14	25
White, both sexes	1,211	5	3	1	2	5	9	13	20	21	49	49
Male	714	1	2	1	2	3	5	10	17	17	39	35
Female	497	4	1	-	-	2	4	3	3	4	10	14
Black, both sexes	274	3	1	-	1	2	9	5	9	14	13	30
Male	168	3	1	-	1	2	7	3	7	11	10	19
Female	106	-	-	-	-	-	2	2	2	3	3	11
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	1	-	-	-	-	-	1	1	1
Male	8	-	-	1	-	-	-	-	-	1	-	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
Utah												
All races, both sexes	72	-	-	-	-	-	1	-	1	4	1	2
Male	34	-	-	-	-	-	1	-	-	2	-	2
Female	38	-	-	-	-	-	-	-	1	2	1	-
White, both sexes	71	-	-	-	-	-	1	-	1	4	1	2
Male	34	-	-	-	-	-	1	-	-	2	-	2
Female	37	-	-	-	-	-	-	-	1	2	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Vermont												
All races, both sexes	38	-	-	-	-	-	-	-	1	-	1	1
Male	23	-	-	-	-	-	-	-	1	-	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	1	-	1	1
Male	23	-	-	-	-	-	-	-	1	-	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	630	2	-	1	2	2	7	4	10	12	15	28
Male	363	1	-	1	-	-	5	2	8	8	11	19
Female	267	1	-	-	2	2	2	2	2	4	4	9
White, both sexes	442	-	-	1	2	2	4	4	5	5	7	11
Male	253	-	-	1	-	-	3	2	5	4	5	5
Female	189	-	-	-	2	2	1	2	-	1	2	6
Black, both sexes	178	2	-	-	-	-	3	-	3	7	8	17
Male	103	1	-	-	-	-	2	-	2	4	6	14
Female	75	1	-	-	-	-	1	-	1	3	2	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Washington												
All races, both sexes	412	-	-	-	1	1	5	4	5	11	19	16
Male	256	-	-	-	1	1	3	3	5	5	15	12
Female	156	-	-	-	-	-	2	1	-	6	4	4
White, both sexes	378	-	-	-	-	1	4	3	4	10	17	14
Male	238	-	-	-	-	1	3	2	4	4	14	11
Female	140	-	-	-	-	-	1	1	-	6	3	3
Black, both sexes	23	-	-	-	1	-	1	-	1	-	2	1
Male	11	-	-	-	1	-	-	-	1	-	1	-
Female	12	-	-	-	-	-	1	-	-	-	1	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	1	-	1	-	-
Male	4	-	-	-	-	-	-	1	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	145	-	1	-	-	-	-	1	2	1	5	4
Male	83	-	-	-	-	-	-	1	2	-	5	3
Female	62	-	1	-	-	-	-	-	-	1	-	1
White, both sexes	140	-	1	-	-	-	-	1	1	1	5	4
Male	79	-	-	-	-	-	-	1	1	-	5	3
Female	61	-	1	-	-	-	-	-	-	1	-	1
Black, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cardiomyopathy (I42)												
Wisconsin												
All races, both sexes	528	1	-	-	-	1	1	-	6	5	12	19
Male	289	1	-	-	-	1	1	-	5	2	10	13
Female	239	-	-	-	-	-	-	-	1	3	2	6
White, both sexes	484	1	-	-	-	1	1	-	3	4	9	15
Male	262	1	-	-	-	1	1	-	2	2	8	11
Female	222	-	-	-	-	-	-	-	1	2	1	4
Black, both sexes	39	-	-	-	-	-	-	-	3	1	3	3
Male	26	-	-	-	-	-	-	-	3	-	2	2
Female	13	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	7	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	7	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
United States												
All races, both sexes	1,337	1,531	1,613	1,760	2,252	3,127	3,708	3,471	2,129	731	125	2
Male	967	1,058	1,048	1,164	1,413	1,836	2,098	1,648	844	200	28	2
Female	370	473	565	596	839	1,291	1,610	1,823	1,285	531	97	-
White, both sexes	881	1,079	1,192	1,333	1,839	2,689	3,283	3,155	1,951	676	108	2
Male	653	762	786	889	1,151	1,613	1,890	1,528	793	183	24	2
Female	228	317	406	444	688	1,076	1,393	1,627	1,158	493	84	-
Black, both sexes	416	403	368	378	356	400	367	262	142	50	17	-
Male	293	265	228	240	222	206	180	97	41	15	4	-
Female	123	138	140	138	134	194	187	165	101	35	13	-
American Indian 1/, both sexes	12	9	16	18	13	6	9	10	5	1	-	-
Male	7	4	10	12	9	3	5	4	-	-	-	-
Female	5	5	6	6	4	3	4	6	5	1	-	-
Asian or Pacific Islander 2/, both sexes	28	40	37	31	44	32	49	44	31	4	-	-
Male	14	27	24	23	31	14	23	19	10	2	-	-
Female	14	13	13	8	13	18	26	25	21	2	-	-
Alabama												
All races, both sexes	22	32	41	30	53	62	59	64	31	6	1	-
Male	13	22	27	20	36	34	28	29	9	2	-	-
Female	9	10	14	10	17	28	31	35	22	4	1	-
White, both sexes	13	14	30	18	38	42	48	53	25	5	-	-
Male	9	11	21	14	27	24	23	24	9	1	-	-
Female	4	3	9	4	11	18	25	29	16	4	-	-
Black, both sexes	9	18	11	12	15	20	11	11	6	1	1	-
Male	4	11	6	6	9	10	5	5	-	1	-	-
Female	5	7	5	6	6	10	6	6	6	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Alaska												
All races, both sexes	2	1	3	3	3	2	5	4	2	2	-	-
Male	2	-	1	3	2	1	4	2	1	-	-	-
Female	-	1	2	-	1	1	1	2	1	2	-	-
White, both sexes	1	-	3	1	-	-	5	4	2	1	-	-
Male	1	-	1	1	-	-	4	2	1	-	-	-
Female	-	-	2	-	-	-	1	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	2	3	2	-	-	-	1	-	-
Male	1	-	-	2	2	1	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	16	15	27	27	34	42	63	44	28	14	1	-
Male	14	11	19	17	20	26	39	28	17	4	-	-
Female	2	4	8	10	14	16	24	16	11	10	1	-
White, both sexes	12	13	23	22	33	41	60	39	28	13	1	-
Male	11	10	17	15	20	26	37	25	17	4	-	-
Female	1	3	6	7	13	15	23	14	11	9	1	-
Black, both sexes	2	-	-	1	-	-	3	2	-	1	-	-
Male	2	-	-	-	-	-	2	1	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	1	-	-
American Indian 1/, both sexes	2	2	4	4	-	-	-	1	-	-	-	-
Male	1	1	2	2	-	-	-	1	-	-	-	-
Female	1	1	2	2	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Arkansas												
All races, both sexes	13	8	23	24	22	31	47	48	31	10	3	-
Male	11	6	13	16	17	15	22	15	7	2	3	-
Female	2	2	10	8	5	16	25	33	24	8	-	-
White, both sexes	8	5	16	19	19	25	45	43	30	10	3	-
Male	6	4	11	13	15	14	22	13	7	2	3	-
Female	2	1	5	6	4	11	23	30	23	8	-	-
Black, both sexes	5	3	7	5	3	6	2	5	1	-	-	-
Male	5	2	2	3	2	1	-	2	-	-	-	-
Female	-	1	5	2	1	5	2	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	184	211	200	194	218	288	404	373	259	63	14	-
Male	127	140	139	128	143	173	239	193	119	20	4	-
Female	57	71	61	66	75	115	165	180	140	43	10	-
White, both sexes	129	145	157	148	172	247	352	339	239	58	13	-
Male	91	102	107	96	118	154	213	183	111	19	4	-
Female	38	43	50	52	54	93	139	156	128	39	9	-
Black, both sexes	43	48	28	33	28	30	30	18	9	4	1	-
Male	30	26	22	22	17	16	18	3	4	-	-	-
Female	13	22	6	11	11	14	12	15	5	4	1	-
American Indian 1/, both sexes	1	3	1	2	2	1	2	1	-	-	-	-
Male	-	2	1	2	1	-	-	1	-	-	-	-
Female	1	1	-	-	1	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	15	14	11	16	10	20	15	11	1	-	-
Male	6	10	9	8	7	3	8	6	4	1	-	-
Female	5	5	5	3	9	7	12	9	7	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Colorado												
All races, both sexes	12	20	15	11	21	25	38	35	14	13	2	-
Male	8	18	7	6	14	11	24	20	4	4	1	-
Female	4	2	8	5	7	14	14	15	10	9	1	-
White, both sexes	11	20	14	11	19	25	38	35	14	13	2	-
Male	7	18	7	6	12	11	24	20	4	4	1	-
Female	4	2	7	5	7	14	14	15	10	9	1	-
Black, both sexes	1	-	1	-	2	-	-	-	-	-	-	-
Male	1	-	-	-	2	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	9	19	19	26	17	31	56	61	28	18	2	-
Male	8	14	11	15	13	17	36	36	11	6	-	-
Female	1	5	8	11	4	14	20	25	17	12	2	-
White, both sexes	9	17	16	24	14	28	54	59	28	18	2	-
Male	8	13	10	14	11	15	35	34	11	6	-	-
Female	1	4	6	10	3	13	19	25	17	12	2	-
Black, both sexes	-	2	2	2	3	3	2	2	-	-	-	-
Male	-	1	1	1	2	2	1	2	-	-	-	-
Female	-	1	1	1	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Delaware												
All races, both sexes	3	5	5	9	5	8	15	11	9	5	-	-
Male	2	4	3	6	3	4	11	7	3	2	-	-
Female	1	1	2	3	2	4	4	4	6	3	-	-
White, both sexes	1	3	5	7	3	7	13	10	9	4	-	-
Male	-	3	3	5	2	4	10	6	3	1	-	-
Female	1	-	2	2	1	3	3	4	6	3	-	-
Black, both sexes	2	2	-	2	2	1	2	-	-	1	-	-
Male	2	1	-	1	1	-	1	-	-	1	-	-
Female	-	1	-	1	1	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	4	9	3	1	2	3	6	3	1	-	-
Male	1	4	5	2	1	2	2	1	-	-	-	-
Female	1	-	4	1	-	-	1	5	3	1	-	-
White, both sexes	-	-	-	-	-	-	2	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	2	3	9	3	1	2	1	6	3	-	-	-
Male	1	3	5	2	1	2	1	1	-	-	-	-
Female	1	-	4	1	-	-	-	5	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Florida												
All races, both sexes	109	104	119	148	204	297	404	367	266	102	16	-
Male	78	73	83	101	136	175	255	174	106	36	5	-
Female	31	31	36	47	68	122	149	193	160	66	11	-
White, both sexes	72	78	94	120	173	258	364	346	254	98	14	-
Male	56	55	66	86	117	153	239	172	101	34	4	-
Female	16	23	28	34	56	105	125	174	153	64	10	-
Black, both sexes	34	26	24	28	31	34	36	20	12	4	2	-
Male	21	18	16	15	19	18	15	2	5	2	1	-
Female	13	8	8	13	12	16	21	18	7	2	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	5	4	1	-	-	-	-
Male	1	-	-	-	-	4	1	-	-	-	-	-
Female	2	-	-	-	-	1	3	1	-	-	-	-
Georgia												
All races, both sexes	44	67	74	61	100	101	99	84	47	20	4	1
Male	32	49	46	38	60	53	59	36	17	5	1	1
Female	12	18	28	23	40	48	40	48	30	15	3	-
White, both sexes	18	41	41	30	72	78	81	68	42	17	3	1
Male	12	30	25	20	39	43	50	33	14	4	1	1
Female	6	11	16	10	33	35	31	35	28	13	2	-
Black, both sexes	26	25	33	31	27	23	17	16	4	3	1	-
Male	20	19	21	18	20	10	8	3	3	1	-	-
Female	6	6	12	13	7	13	9	13	1	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Hawaii												
All races, both sexes	11	17	15	8	15	15	13	22	15	3	-	-
Male	7	13	12	4	11	8	9	9	5	1	-	-
Female	4	4	3	4	4	7	4	13	10	2	-	-
White, both sexes	2	3	4	3	2	5	2	7	3	1	-	-
Male	2	2	4	1	-	3	2	4	2	1	-	-
Female	-	1	-	2	2	2	-	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	14	11	5	13	10	10	13	12	2	-	-
Male	5	11	8	3	11	5	7	5	3	-	-	-
Female	4	3	3	2	2	5	3	8	9	2	-	-
Idaho												
All races, both sexes	1	5	5	3	7	13	13	9	7	5	-	-
Male	1	3	3	1	3	8	5	3	4	-	-	-
Female	-	2	2	2	4	5	8	6	3	5	-	-
White, both sexes	1	5	5	3	7	13	12	9	7	5	-	-
Male	1	3	3	1	3	8	5	3	4	-	-	-
Female	-	2	2	2	4	5	7	6	3	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Illinois												
All races, both sexes	71	56	58	79	84	148	167	141	87	27	5	-
Male	56	36	37	59	48	82	94	53	30	4	1	-
Female	15	20	21	20	36	66	73	88	57	23	4	-
White, both sexes	37	40	43	49	55	120	150	129	83	26	5	-
Male	27	24	30	36	32	71	81	49	29	4	1	-
Female	10	16	13	13	23	49	69	80	54	22	4	-
Black, both sexes	33	16	14	27	28	27	15	11	4	1	-	-
Male	28	12	7	20	15	11	12	4	1	-	-	-
Female	5	4	7	7	13	16	3	7	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	3	1	1	2	1	-	-	-	-
Male	1	-	-	3	1	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	1	1	-	-	-	-
Indiana												
All races, both sexes	26	38	41	48	61	78	75	83	35	22	-	-
Male	19	22	24	27	36	55	43	36	18	4	-	-
Female	7	16	17	21	25	23	32	47	17	18	-	-
White, both sexes	18	31	35	41	54	73	72	81	31	22	-	-
Male	13	18	20	23	32	53	43	36	16	4	-	-
Female	5	13	15	18	22	20	29	45	15	18	-	-
Black, both sexes	8	7	6	6	6	5	3	2	3	-	-	-
Male	6	4	4	3	4	2	-	-	1	-	-	-
Female	2	3	2	3	2	3	3	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Iowa												
All races, both sexes	9	12	8	14	17	29	36	39	17	5	1	-
Male	7	7	5	6	12	15	19	17	8	1	-	-
Female	2	5	3	8	5	14	17	22	9	4	1	-
White, both sexes	9	11	8	14	16	28	36	39	17	5	1	-
Male	7	7	5	6	12	15	19	17	8	1	-	-
Female	2	4	3	8	4	13	17	22	9	4	1	-
Black, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	5	9	10	13	19	25	20	32	16	5	-	-
Male	2	4	8	11	10	10	11	16	5	1	-	-
Female	3	5	2	2	9	15	9	16	11	4	-	-
White, both sexes	3	7	7	12	19	22	18	28	16	5	-	-
Male	1	3	5	10	10	9	9	15	5	1	-	-
Female	2	4	2	2	9	13	9	13	11	4	-	-
Black, both sexes	2	2	3	-	-	2	2	4	-	-	-	-
Male	1	1	3	-	-	-	2	1	-	-	-	-
Female	1	1	-	-	-	2	-	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Kentucky												
All races, both sexes	13	22	18	34	27	47	54	42	30	7	2	-
Male	12	13	15	20	18	27	24	17	6	2	-	-
Female	1	9	3	14	9	20	30	25	24	5	2	-
White, both sexes	10	17	17	29	25	46	52	40	29	7	2	-
Male	9	8	14	16	18	26	24	17	6	2	-	-
Female	1	9	3	13	7	20	28	23	23	5	2	-
Black, both sexes	3	5	1	5	2	1	2	2	1	-	-	-
Male	3	5	1	4	-	1	-	-	-	-	-	-
Female	-	-	-	1	2	-	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	27	31	26	34	48	57	59	60	43	13	6	-
Male	21	16	15	28	23	35	30	26	20	1	1	-
Female	6	15	11	6	25	22	29	34	23	12	5	-
White, both sexes	16	18	13	24	30	35	45	49	35	8	3	-
Male	12	10	7	19	13	24	23	20	17	-	-	-
Female	4	8	6	5	17	11	22	29	18	8	3	-
Black, both sexes	10	13	13	10	18	22	14	11	8	5	3	-
Male	9	6	8	9	10	11	7	6	3	1	1	-
Female	1	7	5	1	8	11	7	5	5	4	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Maine												
All races, both sexes	7	7	7	10	8	14	19	18	9	2	1	-
Male	4	6	5	6	7	7	9	8	4	-	1	-
Female	3	1	2	4	1	7	10	10	5	2	-	-
White, both sexes	7	7	7	10	8	14	19	18	9	2	1	-
Male	4	6	5	6	7	7	9	8	4	-	1	-
Female	3	1	2	4	1	7	10	10	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	18	27	28	38	39	48	74	66	40	12	5	-
Male	13	19	15	27	24	28	38	32	15	7	-	-
Female	5	8	13	11	15	20	36	34	25	5	5	-
White, both sexes	8	12	18	21	26	43	55	52	29	9	4	-
Male	8	10	9	15	15	26	29	25	13	5	-	-
Female	-	2	9	6	11	17	26	27	16	4	4	-
Black, both sexes	9	14	9	17	13	5	15	13	10	2	1	-
Male	5	9	6	12	9	2	7	7	2	1	-	-
Female	4	5	3	5	4	3	8	6	8	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	-	-	-	4	1	1	1	-	-
Male	-	-	-	-	-	-	2	-	-	1	-	-
Female	1	1	1	-	-	-	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Massachusetts												
All races, both sexes	12	17	18	20	38	63	70	61	53	14	2	-
Male	10	15	11	15	25	42	48	30	27	5	-	-
Female	2	2	7	5	13	21	22	31	26	9	2	-
White, both sexes	12	14	15	16	38	61	66	57	51	13	2	-
Male	10	12	9	12	25	40	47	28	27	4	-	-
Female	2	2	6	4	13	21	19	29	24	9	2	-
Black, both sexes	-	3	2	4	-	2	3	4	1	1	-	-
Male	-	3	1	3	-	2	1	2	-	1	-	-
Female	-	-	1	1	-	-	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Michigan												
All races, both sexes	72	72	58	60	78	115	136	134	66	25	2	-
Male	51	54	43	39	50	66	69	62	29	6	-	-
Female	21	18	15	21	28	49	67	72	37	19	2	-
White, both sexes	47	47	41	50	67	99	123	122	64	23	2	-
Male	34	35	30	34	43	55	63	55	28	4	-	-
Female	13	12	11	16	24	44	60	67	36	19	2	-
Black, both sexes	25	25	16	8	11	16	13	12	2	2	-	-
Male	17	19	13	4	7	11	6	7	1	2	-	-
Female	8	6	3	4	4	5	7	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Minnesota												
All races, both sexes	14	22	20	15	32	40	45	45	40	15	2	-
Male	11	13	12	11	20	27	30	29	19	4	1	-
Female	3	9	8	4	12	13	15	16	21	11	1	-
White, both sexes	11	19	19	15	32	39	43	43	40	15	2	-
Male	8	11	11	11	20	26	28	27	19	4	1	-
Female	3	8	8	4	12	13	15	16	21	11	1	-
Black, both sexes	2	2	-	-	-	1	2	-	-	-	-	-
Male	2	1	-	-	-	1	2	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	2	-	-	-	-
Male	-	1	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	12	18	19	20	32	46	39	33	18	10	3	-
Male	9	12	12	11	18	27	22	16	5	-	1	-
Female	3	6	7	9	14	19	17	17	13	10	2	-
White, both sexes	4	7	10	10	20	30	27	24	12	4	1	-
Male	2	6	8	5	13	21	16	14	2	-	-	-
Female	2	1	2	5	7	9	11	10	10	4	1	-
Black, both sexes	8	11	9	10	12	16	12	9	6	6	2	-
Male	7	6	4	6	5	6	6	2	3	-	1	-
Female	1	5	5	4	7	10	6	7	3	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Missouri												
All races, both sexes	21	29	39	32	60	72	77	77	48	23	4	-
Male	16	23	27	21	30	47	42	29	17	7	-	-
Female	5	6	12	11	30	25	35	48	31	16	4	-
White, both sexes	17	18	30	27	49	62	71	75	46	22	4	-
Male	13	14	20	19	27	42	39	29	17	7	-	-
Female	4	4	10	8	22	20	32	46	29	15	4	-
Black, both sexes	4	10	9	4	11	10	6	2	2	1	-	-
Male	3	9	7	1	3	5	3	-	-	-	-	-
Female	1	1	2	3	8	5	3	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	3	5	4	7	9	10	4	4	2	-	-
Male	4	1	1	3	4	5	5	3	1	-	-	-
Female	2	2	4	1	3	4	5	1	3	2	-	-
White, both sexes	6	3	5	3	7	9	10	4	4	2	-	-
Male	4	1	1	2	4	5	5	3	1	-	-	-
Female	2	2	4	1	3	4	5	1	3	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Nebraska												
All races, both sexes	2	6	7	9	16	14	21	19	6	7	1	-
Male	2	5	4	6	10	12	13	9	4	2	-	-
Female	-	1	3	3	6	2	8	10	2	5	1	-
White, both sexes	2	5	6	9	13	14	20	17	6	7	1	-
Male	2	4	3	6	9	12	12	8	4	2	-	-
Female	-	1	3	3	4	2	8	9	2	5	1	-
Black, both sexes	-	-	1	-	3	-	1	1	-	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	2	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	15	12	22	22	25	30	36	25	19	1	-	-
Male	8	8	11	16	19	20	16	14	11	-	-	-
Female	7	4	11	6	6	10	20	11	8	1	-	-
White, both sexes	10	8	17	21	19	29	33	24	18	1	-	-
Male	6	5	9	15	14	20	14	14	10	-	-	-
Female	4	3	8	6	5	9	19	10	8	1	-	-
Black, both sexes	5	4	3	1	3	1	1	-	1	-	-	-
Male	2	3	-	1	2	-	1	-	1	-	-	-
Female	3	1	3	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	-	2	1	-	-	-	-
Male	-	-	1	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
New Hampshire												
All races, both sexes	5	5	2	6	10	8	18	15	6	3	2	-
Male	4	4	1	4	5	5	9	10	-	2	-	-
Female	1	1	1	2	5	3	9	5	6	1	2	-
White, both sexes	5	5	2	6	10	7	17	15	6	3	2	-
Male	4	4	1	4	5	4	8	10	-	2	-	-
Female	1	1	1	2	5	3	9	5	6	1	2	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	49	38	40	50	77	120	123	116	86	22	5	-
Male	36	31	28	37	49	73	73	67	44	8	2	-
Female	13	7	12	13	28	47	50	49	42	14	3	-
White, both sexes	33	27	29	36	62	103	112	109	82	20	4	-
Male	23	22	22	24	39	65	66	62	44	6	2	-
Female	10	5	7	12	23	38	46	47	38	14	2	-
Black, both sexes	15	10	11	11	13	15	11	7	4	2	1	-
Male	12	8	6	10	8	7	7	5	-	2	-	-
Female	3	2	5	1	5	8	4	2	4	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	3	2	2	-	-	-	-	-	-
Male	1	1	-	3	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
New Mexico												
All races, both sexes	5	6	9	4	9	17	9	17	8	3	-	-
Male	5	1	4	3	5	12	6	8	3	-	-	-
Female	-	5	5	1	4	5	3	9	5	3	-	-
White, both sexes	3	4	7	3	8	16	8	16	8	3	-	-
Male	3	1	4	2	5	11	5	7	3	-	-	-
Female	-	3	3	1	3	5	3	9	5	3	-	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	1	1	1	1	1	1	-	-	-	-
Male	2	-	-	1	-	1	1	1	-	-	-	-
Female	-	1	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	47	68	61	58	110	157	168	175	105	31	6	-
Male	29	48	40	41	68	103	98	83	41	8	1	-
Female	18	20	21	17	42	54	70	92	64	23	5	-
White, both sexes	28	55	46	46	93	147	153	167	97	29	5	-
Male	18	40	30	35	54	98	93	80	38	8	-	-
Female	10	15	16	11	39	49	60	87	59	21	5	-
Black, both sexes	18	11	14	11	15	10	14	6	6	2	1	-
Male	11	7	9	5	12	5	5	2	1	-	1	-
Female	7	4	5	6	3	5	9	4	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	1	2	-	1	2	2	-	-	-
Male	-	1	1	1	2	-	-	1	2	-	-	-
Female	1	1	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
North Carolina												
All races, both sexes	49	52	67	74	82	97	117	118	57	20	5	-
Male	34	34	48	48	53	52	64	42	26	2	2	-
Female	15	18	19	26	29	45	53	76	31	18	3	-
White, both sexes	24	32	37	49	61	73	91	100	45	17	4	-
Male	19	20	29	32	41	40	50	38	19	2	2	-
Female	5	12	8	17	20	33	41	62	26	15	2	-
Black, both sexes	25	19	28	24	19	23	25	18	12	3	1	-
Male	15	14	19	15	11	12	13	4	7	-	-	-
Female	10	5	9	9	8	11	12	14	5	3	1	-
American Indian 1/, both sexes	-	1	1	1	2	-	1	-	-	-	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	-	1	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	2	1	-	2	5	7	10	8	2	-	-
Male	2	2	1	-	1	3	4	5	3	-	-	-
Female	-	-	-	-	1	2	3	5	5	2	-	-
White, both sexes	2	2	1	-	2	5	7	10	8	2	-	-
Male	2	2	1	-	1	3	4	5	3	-	-	-
Female	-	-	-	-	1	2	3	5	5	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Ohio												
All races, both sexes	59	59	66	67	94	150	164	161	82	34	2	-
Male	42	42	37	46	68	89	97	70	32	10	-	-
Female	17	17	29	21	26	61	67	91	50	24	2	-
White, both sexes	45	44	53	49	82	125	149	148	77	32	2	-
Male	33	30	29	33	58	75	92	62	31	9	-	-
Female	12	14	24	16	24	50	57	86	46	23	2	-
Black, both sexes	14	15	13	18	12	25	15	13	5	2	-	-
Male	9	12	8	13	10	14	5	8	1	1	-	-
Female	5	3	5	5	2	11	10	5	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	13	13	19	23	22	36	37	34	16	6	1	-
Male	7	10	12	16	13	22	17	16	6	4	-	-
Female	6	3	7	7	9	14	20	18	10	2	1	-
White, both sexes	10	12	11	18	17	35	28	27	14	6	1	-
Male	5	9	8	15	8	22	13	15	6	4	-	-
Female	5	3	3	3	9	13	15	12	8	2	1	-
Black, both sexes	2	-	7	3	2	-	6	3	-	-	-	-
Male	1	-	4	1	2	-	2	1	-	-	-	-
Female	1	-	3	2	-	-	4	2	-	-	-	-
American Indian 1/, both sexes	1	1	1	2	3	1	3	3	2	-	-	-
Male	1	1	-	-	3	-	2	-	-	-	-	-
Female	-	-	1	2	-	1	1	3	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Oregon												
All races, both sexes	13	18	20	21	27	28	35	23	27	5	4	-
Male	9	15	14	10	15	20	17	12	14	1	-	-
Female	4	3	6	11	12	8	18	11	13	4	4	-
White, both sexes	11	15	15	21	26	28	33	22	27	5	4	-
Male	9	13	11	10	15	20	15	12	14	1	-	-
Female	2	2	4	11	11	8	18	10	13	4	4	-
Black, both sexes	1	2	2	-	1	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	1	1	2	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	2	-	-	-	1	1	-	-	-	-
Male	-	-	2	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	70	81	86	102	118	218	257	248	141	54	7	-
Male	54	58	64	63	71	122	134	122	42	14	3	-
Female	16	23	22	39	47	96	123	126	99	40	4	-
White, both sexes	52	60	69	90	107	201	240	234	128	51	6	-
Male	39	41	52	57	64	112	127	115	39	13	3	-
Female	13	19	17	33	43	89	113	119	89	38	3	-
Black, both sexes	18	20	17	12	10	17	17	13	12	3	1	-
Male	15	17	12	6	6	10	7	6	3	1	-	-
Female	3	3	5	6	4	7	10	7	9	2	1	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	1	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Rhode Island												
All races, both sexes	2	1	3	3	3	9	6	9	3	1	-	-
Male	1	1	1	2	3	7	2	6	1	-	-	-
Female	1	-	2	1	-	2	4	3	2	1	-	-
White, both sexes	2	1	3	1	3	9	6	9	3	1	-	-
Male	1	1	1	1	3	7	2	6	1	-	-	-
Female	1	-	2	-	-	2	4	3	2	1	-	-
Black, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	31	40	34	41	60	55	63	61	21	8	2	1
Male	23	27	20	28	42	33	34	26	5	1	-	1
Female	8	13	14	13	18	22	29	35	16	7	2	-
White, both sexes	12	20	21	22	44	38	38	44	18	7	2	1
Male	9	15	13	13	32	23	23	16	5	1	-	1
Female	3	5	8	9	12	15	15	28	13	6	2	-
Black, both sexes	19	20	13	19	16	17	25	16	3	1	-	-
Male	14	12	7	15	10	10	11	9	-	-	-	-
Female	5	8	6	4	6	7	14	7	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
South Dakota												
All races, both sexes	2	1	1	5	3	5	4	10	3	2	1	-
Male	2	1	-	3	-	-	1	7	-	-	-	-
Female	-	-	1	2	3	5	3	3	3	2	1	-
White, both sexes	2	1	1	3	3	5	4	9	3	2	1	-
Male	2	1	-	2	-	-	1	7	-	-	-	-
Female	-	-	1	1	3	5	3	2	3	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	29	34	36	51	60	43	82	54	27	10	3	-
Male	17	21	23	32	39	29	38	19	11	3	-	-
Female	12	13	13	19	21	14	44	35	16	7	3	-
White, both sexes	13	18	19	39	39	32	62	45	23	6	3	-
Male	7	12	14	28	22	21	28	16	10	2	-	-
Female	6	6	5	11	17	11	34	29	13	4	3	-
Black, both sexes	16	16	17	12	20	11	19	9	4	4	-	-
Male	10	9	9	4	16	8	10	3	1	1	-	-
Female	6	7	8	8	4	3	9	6	3	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Texas												
All races, both sexes	113	98	102	126	126	168	201	155	112	28	2	-
Male	91	64	62	83	73	81	108	81	39	10	-	-
Female	22	34	40	43	53	87	93	74	73	18	2	-
White, both sexes	75	75	80	90	110	146	181	150	98	28	1	-
Male	64	53	48	61	66	70	95	79	36	10	-	-
Female	11	22	32	29	44	76	86	71	62	18	1	-
Black, both sexes	37	23	20	31	16	22	19	5	13	-	1	-
Male	26	11	12	20	7	11	12	2	3	-	-	-
Female	11	12	8	11	9	11	7	3	10	-	1	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	4	-	-	1	-	-	-	-	-
Male	-	-	2	2	-	-	1	-	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	2	6	6	4	8	9	11	9	8	-	-	-
Male	1	5	3	4	2	5	5	1	3	-	-	-
Female	1	1	3	-	6	4	6	8	5	-	-	-
White, both sexes	2	6	6	4	8	9	10	9	8	-	-	-
Male	1	5	3	4	2	5	5	1	3	-	-	-
Female	1	1	3	-	6	4	5	8	5	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Vermont												
All races, both sexes	2	-	5	1	4	4	7	8	3	-	1	-
Male	1	-	3	1	4	2	5	4	-	-	-	-
Female	1	-	2	-	-	2	2	4	3	-	1	-
White, both sexes	2	-	5	1	4	4	7	8	3	-	1	-
Male	1	-	3	1	4	2	5	4	-	-	-	-
Female	1	-	2	-	-	2	2	4	3	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	37	55	40	49	39	91	91	83	48	13	1	-
Male	26	39	27	31	26	55	48	38	16	2	-	-
Female	11	16	13	18	13	36	43	45	32	11	1	-
White, both sexes	26	34	23	31	29	67	73	68	38	12	-	-
Male	20	27	15	18	16	43	41	31	15	2	-	-
Female	6	7	8	13	13	24	32	37	23	10	-	-
Black, both sexes	10	21	15	17	8	24	17	14	10	1	1	-
Male	6	12	10	12	8	12	6	7	1	-	-	-
Female	4	9	5	5	-	12	11	7	9	1	1	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	2	-	1	1	-	-	-	-
Male	-	-	1	1	2	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Washington												
All races, both sexes	26	21	27	30	44	51	50	58	28	14	1	-
Male	18	14	19	20	25	33	34	32	12	4	-	-
Female	8	7	8	10	19	18	16	26	16	10	1	-
White, both sexes	20	19	25	25	40	49	49	56	27	14	1	-
Male	15	14	18	17	21	32	34	32	12	4	-	-
Female	5	5	7	8	19	17	15	24	15	10	1	-
Black, both sexes	5	2	1	5	2	-	-	2	-	-	-	-
Male	3	-	-	3	2	-	-	-	-	-	-	-
Female	2	2	1	2	-	-	-	2	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	-	1	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	1	-	-	-
West Virginia												
All races, both sexes	3	12	9	14	15	26	21	17	12	1	1	-
Male	3	9	7	9	7	12	12	10	3	-	-	-
Female	-	3	2	5	8	14	9	7	9	1	1	-
White, both sexes	3	12	9	12	15	26	20	16	12	1	1	-
Male	3	9	7	7	7	12	12	9	3	-	-	-
Female	-	3	2	5	8	14	8	7	9	1	1	-
Black, both sexes	-	-	-	2	-	-	1	1	-	-	-	-
Male	-	-	-	2	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cardiomyopathy (I42)												
Wisconsin												
All races, both sexes	19	30	39	30	46	76	79	81	57	22	4	-
Male	13	20	19	19	29	46	45	38	21	5	1	-
Female	6	10	20	11	17	30	34	43	36	17	3	-
White, both sexes	16	27	30	28	44	69	78	77	55	22	4	-
Male	11	17	15	17	28	41	44	36	21	5	1	-
Female	5	10	15	11	16	28	34	41	34	17	3	-
Black, both sexes	3	3	8	2	2	7	1	3	-	-	-	-
Male	2	3	4	2	1	5	1	1	-	-	-	-
Female	1	-	4	-	1	2	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Wyoming												
All races, both sexes	1	2	1	2	2	2	1	2	-	-	-	-
Male	-	1	1	1	2	-	1	1	-	-	-	-
Female	1	1	-	1	-	2	-	1	-	-	-	-
White, both sexes	1	2	1	2	2	2	1	2	-	-	-	-
Male	-	1	1	1	2	-	1	1	-	-	-	-
Female	1	1	-	1	-	2	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
United States												
All races, both sexes	38,503	40	24	18	30	88	109	141	173	320	546	858
Male	16,959	23	9	11	20	63	75	90	112	195	315	544
Female	21,544	17	15	7	10	25	34	51	61	125	231	314
White, both sexes	32,955	24	17	14	19	63	76	104	108	227	410	633
Male	14,331	12	6	8	14	41	53	72	69	140	243	421
Female	18,624	12	11	6	5	22	23	32	39	87	167	212
Black, both sexes	4,690	12	6	3	9	23	28	35	57	83	127	206
Male	2,167	7	3	3	6	20	19	17	36	48	69	109
Female	2,523	5	3	-	3	3	9	18	21	35	58	97
American Indian 1/, both sexes	129	-	-	-	2	1	3	-	5	4	2	5
Male	70	-	-	-	-	1	2	-	4	1	2	3
Female	59	-	-	-	2	-	1	-	1	3	-	2
Asian or Pacific Islander 2/, both sexes	729	4	1	1	-	1	2	2	3	6	7	14
Male	391	4	-	-	-	1	1	1	3	6	1	11
Female	338	-	1	1	-	-	1	1	-	-	6	3
Alabama												
All races, both sexes	2,655	9	1	-	-	2	11	20	18	40	72	106
Male	1,286	4	-	-	-	2	8	14	13	28	48	77
Female	1,369	5	1	-	-	-	3	6	5	12	24	29
White, both sexes	2,043	4	1	-	-	1	6	16	12	30	60	76
Male	984	2	-	-	-	1	5	12	8	21	41	57
Female	1,059	2	1	-	-	-	1	4	4	9	19	19
Black, both sexes	608	5	-	-	-	1	5	4	6	10	12	30
Male	299	2	-	-	-	1	3	2	5	7	7	20
Female	309	3	-	-	-	-	2	2	1	3	5	10
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Alaska												
All races, both sexes	24	-	-	-	1	1	-	-	1	-	-	1
Male	16	-	-	-	1	1	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	20	-	-	-	1	-	-	-	-	-	-	1
Male	13	-	-	-	1	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	337	1	-	-	-	1	2	1	1	2	4	4
Male	154	-	-	-	-	1	1	1	-	1	-	2
Female	183	1	-	-	-	-	1	-	1	1	4	2
White, both sexes	321	1	-	-	-	1	2	1	1	2	4	2
Male	145	-	-	-	-	1	1	1	-	1	-	1
Female	176	1	-	-	-	-	1	-	1	1	4	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Arkansas												
All races, both sexes	380	-	-	-	-	1	1	3	-	4	7	7
Male	161	-	-	-	-	1	1	2	-	3	3	4
Female	219	-	-	-	-	-	-	1	-	1	4	3
White, both sexes	314	-	-	-	-	1	1	2	-	4	4	3
Male	130	-	-	-	-	1	1	2	-	3	-	2
Female	184	-	-	-	-	-	-	-	-	1	4	1
Black, both sexes	65	-	-	-	-	-	-	1	-	-	3	4
Male	31	-	-	-	-	-	-	-	-	-	3	2
Female	34	-	-	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,424	3	-	1	1	3	10	5	6	7	11	12
Male	560	2	-	-	1	3	7	4	6	5	7	9
Female	864	1	-	1	-	-	3	1	-	2	4	3
White, both sexes	1,285	1	-	-	1	3	8	3	5	6	9	11
Male	496	-	-	-	1	3	6	3	5	5	6	8
Female	789	1	-	-	-	-	2	-	-	1	3	3
Black, both sexes	49	-	-	-	-	-	1	2	-	1	1	1
Male	20	-	-	-	-	-	1	1	-	-	-	1
Female	29	-	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	83	2	-	1	-	-	1	-	-	-	1	-
Male	42	2	-	-	-	-	-	-	-	-	1	-
Female	41	-	-	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Colorado												
All races, both sexes	313	-	-	-	1	2	1	1	-	5	7	5
Male	115	-	-	-	-	-	-	-	-	4	3	1
Female	198	-	-	-	1	2	1	1	-	1	4	4
White, both sexes	305	-	-	-	-	2	1	1	-	5	7	5
Male	113	-	-	-	-	-	-	-	-	4	3	1
Female	192	-	-	-	-	2	1	1	-	1	4	4
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	786	-	-	-	-	-	1	-	3	8	7	15
Male	360	-	-	-	-	-	-	-	2	7	3	12
Female	426	-	-	-	-	-	1	-	1	1	4	3
White, both sexes	739	-	-	-	-	-	1	-	3	7	5	14
Male	339	-	-	-	-	-	-	-	2	6	2	11
Female	400	-	-	-	-	-	1	-	1	1	3	3
Black, both sexes	43	-	-	-	-	-	-	-	-	1	2	-
Male	20	-	-	-	-	-	-	-	-	1	1	-
Female	23	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Delaware												
All races, both sexes	91	-	-	-	-	1	-	1	-	1	-	1
Male	36	-	-	-	-	1	-	1	-	1	-	1
Female	55	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	72	-	-	-	-	-	-	1	-	1	-	1
Male	28	-	-	-	-	-	-	1	-	1	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	18	-	-	-	-	1	-	-	-	-	-	-
Male	8	-	-	-	-	1	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	37	-	-	-	-	-	-	-	1	-	-	2
Male	14	-	-	-	-	-	-	-	1	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	33	-	-	-	-	-	-	-	1	-	-	2
Male	13	-	-	-	-	-	-	-	1	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Florida												
All races, both sexes	1,538	2	1	-	1	5	5	2	2	11	15	30
Male	652	1	1	-	1	5	3	2	1	7	4	16
Female	886	1	-	-	-	-	2	-	1	4	11	14
White, both sexes	1,370	1	-	-	1	3	4	1	1	8	10	23
Male	584	-	-	-	1	3	2	1	1	6	3	12
Female	786	1	-	-	-	-	2	-	-	2	7	11
Black, both sexes	159	1	1	-	-	2	1	1	1	3	5	6
Male	63	1	1	-	-	2	1	1	-	1	1	3
Female	96	-	-	-	-	-	-	-	1	2	4	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,223	5	2	1	1	5	3	8	13	15	41	61
Male	939	4	2	-	1	4	2	5	8	5	27	37
Female	1,284	1	-	1	-	1	1	3	5	10	14	24
White, both sexes	1,642	1	-	1	-	-	1	4	8	5	20	36
Male	688	1	-	-	-	-	1	3	6	2	12	26
Female	954	-	-	1	-	-	-	1	2	3	8	10
Black, both sexes	558	3	2	-	1	5	2	4	5	9	18	25
Male	240	2	2	-	1	4	1	2	2	3	15	11
Female	318	1	-	-	-	1	1	2	3	6	3	14
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	16	1	-	-	-	-	-	-	-	-	3	-
Male	8	1	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Hawaii												
All races, both sexes	499	-	-	-	-	-	1	1	1	1	2	6
Male	260	-	-	-	-	-	1	-	1	1	-	5
Female	239	-	-	-	-	-	-	1	-	-	2	1
White, both sexes	107	-	-	-	-	-	-	-	-	-	-	1
Male	48	-	-	-	-	-	-	-	-	-	-	1
Female	59	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	391	-	-	-	-	-	1	1	1	1	2	5
Male	211	-	-	-	-	-	1	-	1	1	-	4
Female	180	-	-	-	-	-	-	1	-	-	2	1
Idaho												
All races, both sexes	154	-	-	-	-	-	-	2	-	1	2	2
Male	57	-	-	-	-	-	-	2	-	-	2	2
Female	97	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	154	-	-	-	-	-	-	2	-	1	2	2
Male	57	-	-	-	-	-	-	2	-	-	2	2
Female	97	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Illinois												
All races, both sexes	1,495	3	1	1	3	13	9	10	13	18	32	41
Male	660	2	-	-	3	7	6	7	11	15	20	21
Female	835	1	1	1	-	6	3	3	2	3	12	20
White, both sexes	1,290	2	1	1	2	9	6	8	9	15	22	25
Male	559	1	-	-	2	4	4	7	7	13	14	15
Female	731	1	1	1	-	5	2	1	2	2	8	10
Black, both sexes	190	1	-	-	1	4	3	2	4	3	10	16
Male	91	1	-	-	1	3	2	-	4	2	6	6
Female	99	-	-	-	-	1	1	2	-	1	4	10
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,017	-	1	1	1	1	1	1	7	8	16	34
Male	459	-	-	1	-	-	-	-	4	2	10	19
Female	558	-	1	-	1	1	1	1	3	6	6	15
White, both sexes	938	-	-	-	1	1	1	1	6	8	12	28
Male	421	-	-	-	-	-	-	-	3	2	8	15
Female	517	-	-	-	1	1	1	1	3	6	4	13
Black, both sexes	76	-	1	1	-	-	-	-	1	-	4	6
Male	36	-	-	1	-	-	-	-	1	-	2	4
Female	40	-	1	-	-	-	-	-	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Iowa												
All races, both sexes	303	-	-	1	1	1	1	1	-	5	3	6
Male	140	-	-	1	1	-	1	1	-	3	-	3
Female	163	-	-	-	-	1	-	-	-	2	3	3
White, both sexes	302	-	-	1	1	1	1	1	-	5	3	6
Male	140	-	-	1	1	-	1	1	-	3	-	3
Female	162	-	-	-	-	1	-	-	-	2	3	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	376	1	-	-	-	-	1	-	-	8	4	10
Male	179	1	-	-	-	-	-	-	-	5	4	5
Female	197	-	-	-	-	-	1	-	-	3	-	5
White, both sexes	351	1	-	-	-	-	1	-	-	6	3	8
Male	161	1	-	-	-	-	-	-	-	4	3	3
Female	190	-	-	-	-	-	1	-	-	2	-	5
Black, both sexes	18	-	-	-	-	-	-	-	-	-	1	2
Male	14	-	-	-	-	-	-	-	-	-	1	2
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Kentucky												
All races, both sexes	892	2	-	-	-	-	1	2	5	8	12	18
Male	391	2	-	-	-	-	1	2	4	4	9	11
Female	501	-	-	-	-	-	-	-	1	4	3	7
White, both sexes	817	1	-	-	-	-	1	2	4	6	11	13
Male	358	1	-	-	-	-	1	2	3	2	8	9
Female	459	-	-	-	-	-	-	-	1	4	3	4
Black, both sexes	72	-	-	-	-	-	-	-	1	2	1	5
Male	31	-	-	-	-	-	-	-	1	2	1	2
Female	41	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	669	-	-	1	1	3	3	-	1	3	13	20
Male	298	-	-	-	1	2	2	-	-	1	7	13
Female	371	-	-	1	-	1	1	-	1	2	6	7
White, both sexes	468	-	-	1	1	1	3	-	-	1	10	13
Male	219	-	-	-	1	-	2	-	-	-	7	8
Female	249	-	-	1	-	1	1	-	-	1	3	5
Black, both sexes	200	-	-	-	-	2	-	-	1	2	3	7
Male	79	-	-	-	-	2	-	-	-	1	-	5
Female	121	-	-	-	-	-	-	-	1	1	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Maine												
All races, both sexes	216	-	-	-	-	1	-	1	2	-	3	2
Male	103	-	-	-	-	-	-	1	2	-	2	1
Female	113	-	-	-	-	1	-	-	-	-	1	1
White, both sexes	213	-	-	-	-	1	-	1	2	-	2	2
Male	100	-	-	-	-	-	-	1	2	-	1	1
Female	113	-	-	-	-	1	-	-	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	573	-	-	-	-	4	6	6	4	7	10	15
Male	241	-	-	-	-	4	4	5	2	6	5	9
Female	332	-	-	-	-	-	2	1	2	1	5	6
White, both sexes	426	-	-	-	-	2	4	2	-	3	7	10
Male	177	-	-	-	-	2	3	2	-	2	5	5
Female	249	-	-	-	-	-	1	-	-	1	2	5
Black, both sexes	135	-	-	-	-	2	2	4	4	3	3	3
Male	54	-	-	-	-	2	1	3	2	3	-	2
Female	81	-	-	-	-	-	1	1	2	-	3	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Massachusetts												
All races, both sexes	1,281	1	2	-	1	1	2	4	5	7	21	25
Male	584	1	1	-	1	1	1	3	3	4	14	17
Female	697	-	1	-	-	-	1	1	2	3	7	8
White, both sexes	1,207	1	2	-	-	1	2	4	3	5	20	23
Male	543	1	1	-	-	1	1	3	1	3	13	15
Female	664	-	1	-	-	-	1	1	2	2	7	8
Black, both sexes	54	-	-	-	1	-	-	-	1	1	1	1
Male	29	-	-	-	1	-	-	-	1	-	1	1
Female	25	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	1	1	-	1
Male	12	-	-	-	-	-	-	-	1	1	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,038	2	1	-	-	5	3	3	3	9	11	13
Male	404	1	-	-	-	3	2	2	1	5	8	9
Female	634	1	1	-	-	2	1	1	2	4	3	4
White, both sexes	921	1	1	-	-	4	3	2	1	7	8	9
Male	356	1	-	-	-	2	2	1	-	4	5	7
Female	565	-	1	-	-	2	1	1	1	3	3	2
Black, both sexes	108	1	-	-	-	1	-	1	2	2	3	3
Male	43	-	-	-	-	1	-	1	1	1	3	1
Female	65	1	-	-	-	-	-	-	1	1	-	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Minnesota												
All races, both sexes	543	1	-	-	-	-	2	1	-	2	5	6
Male	258	-	-	-	-	-	2	-	-	1	1	6
Female	285	1	-	-	-	-	-	1	-	1	4	-
White, both sexes	527	1	-	-	-	-	2	1	-	1	4	4
Male	249	-	-	-	-	-	2	-	-	1	-	4
Female	278	1	-	-	-	-	-	1	-	-	4	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	744	-	-	1	1	1	3	1	3	9	16	26
Male	322	-	-	1	-	1	1	-	1	5	7	13
Female	422	-	-	-	1	-	2	1	2	4	9	13
White, both sexes	454	-	-	-	-	-	-	1	1	3	6	12
Male	182	-	-	-	-	-	-	-	1	1	4	5
Female	272	-	-	-	-	-	-	1	-	2	2	7
Black, both sexes	285	-	-	1	1	1	3	-	2	6	10	14
Male	136	-	-	1	-	1	1	-	-	4	3	8
Female	149	-	-	-	1	-	2	-	2	2	7	6
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Missouri												
All races, both sexes	1,000	2	2	-	-	3	1	5	3	9	11	29
Male	462	1	1	-	-	2	1	5	2	4	6	17
Female	538	1	1	-	-	1	-	-	1	5	5	12
White, both sexes	914	2	2	-	-	2	-	4	1	7	10	24
Male	428	1	1	-	-	1	-	4	1	3	6	16
Female	486	1	1	-	-	1	-	-	-	4	4	8
Black, both sexes	82	-	-	-	-	1	1	1	2	2	1	5
Male	34	-	-	-	-	1	1	1	1	1	-	1
Female	48	-	-	-	-	-	-	-	1	1	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	150	-	1	-	-	-	-	-	2	-	-	4
Male	69	-	1	-	-	-	-	-	1	-	-	4
Female	81	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	145	-	1	-	-	-	-	-	2	-	-	3
Male	67	-	1	-	-	-	-	-	1	-	-	3
Female	78	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Nebraska												
All races, both sexes	392	-	-	-	-	-	1	-	1	1	2	4
Male	172	-	-	-	-	-	-	-	1	-	1	3
Female	220	-	-	-	-	-	1	-	-	1	1	1
White, both sexes	382	-	-	-	-	-	1	-	1	1	2	3
Male	165	-	-	-	-	-	-	-	1	-	1	3
Female	217	-	-	-	-	-	1	-	-	1	1	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	718	1	-	-	1	3	1	4	5	11	23	34
Male	416	1	-	-	1	3	1	3	4	10	11	23
Female	302	-	-	-	-	-	-	1	1	1	12	11
White, both sexes	598	1	-	-	1	3	-	1	3	5	18	25
Male	343	1	-	-	1	3	-	1	2	5	8	19
Female	255	-	-	-	-	-	-	-	1	-	10	6
Black, both sexes	80	-	-	-	-	-	1	2	1	4	4	9
Male	48	-	-	-	-	-	1	1	1	3	3	4
Female	32	-	-	-	-	-	-	1	-	1	1	5
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	1	-	2	1	-
Male	20	-	-	-	-	-	-	1	-	2	-	-
Female	12	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
New Hampshire												
All races, both sexes	106	-	-	-	-	-	-	-	1	2	-	-
Male	44	-	-	-	-	-	-	-	1	-	-	-
Female	62	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	105	-	-	-	-	-	-	-	1	2	-	-
Male	44	-	-	-	-	-	-	-	1	-	-	-
Female	61	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,085	1	1	-	1	6	3	2	7	5	6	22
Male	484	-	-	-	1	4	3	1	5	3	2	17
Female	601	1	1	-	-	2	-	1	2	2	4	5
White, both sexes	950	1	1	-	-	6	2	2	2	2	5	21
Male	407	-	-	-	-	4	2	1	2	1	2	16
Female	543	1	1	-	-	2	-	1	-	1	3	5
Black, both sexes	117	-	-	-	1	-	1	-	4	3	1	1
Male	67	-	-	-	1	-	1	-	2	2	-	1
Female	50	-	-	-	-	-	-	-	2	1	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	1	-	-	-
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
New Mexico												
All races, both sexes	159	-	-	-	-	-	-	1	2	-	1	4
Male	82	-	-	-	-	-	-	-	2	-	-	2
Female	77	-	-	-	-	-	-	1	-	-	1	2
White, both sexes	153	-	-	-	-	-	-	1	1	-	1	4
Male	78	-	-	-	-	-	-	-	1	-	-	2
Female	75	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,715	-	2	1	1	2	7	8	8	15	7	26
Male	722	-	-	1	1	2	5	5	6	5	4	14
Female	993	-	2	-	-	-	2	3	2	10	3	12
White, both sexes	1,552	-	1	1	1	-	4	5	6	12	7	18
Male	659	-	-	1	1	-	3	3	4	4	4	11
Female	893	-	1	-	-	-	1	2	2	8	3	7
Black, both sexes	143	-	-	-	-	2	3	3	2	3	-	8
Male	57	-	-	-	-	2	2	2	2	1	-	3
Female	86	-	-	-	-	-	1	1	-	2	-	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	1	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
North Carolina												
All races, both sexes	1,199	1	-	-	-	4	-	1	4	9	12	21
Male	525	1	-	-	-	2	-	1	4	4	6	17
Female	674	-	-	-	-	2	-	-	-	5	6	4
White, both sexes	950	-	-	-	-	4	-	-	1	4	4	11
Male	400	-	-	-	-	2	-	-	1	2	2	9
Female	550	-	-	-	-	2	-	-	-	2	2	2
Black, both sexes	237	1	-	-	-	-	-	1	3	5	8	10
Male	118	1	-	-	-	-	-	1	3	2	4	8
Female	119	-	-	-	-	-	-	-	-	3	4	2
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	72	-	-	-	-	1	-	-	-	-	2	3
Male	28	-	-	-	-	1	-	-	-	-	2	2
Female	44	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	72	-	-	-	-	1	-	-	-	-	2	3
Male	28	-	-	-	-	1	-	-	-	-	2	2
Female	44	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Ohio												
All races, both sexes	1,573	-	1	1	1	-	1	2	7	10	20	34
Male	641	-	1	1	1	-	1	2	5	7	10	21
Female	932	-	-	-	-	-	-	-	2	3	10	13
White, both sexes	1,422	-	1	1	1	-	1	2	5	7	16	30
Male	575	-	1	1	1	-	1	2	4	5	9	20
Female	847	-	-	-	-	-	-	-	1	2	7	10
Black, both sexes	148	-	-	-	-	-	-	-	2	3	4	4
Male	64	-	-	-	-	-	-	-	1	2	1	1
Female	84	-	-	-	-	-	-	-	1	1	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	417	-	1	1	-	3	3	1	4	2	11	5
Male	166	-	-	1	-	3	3	-	-	1	7	3
Female	251	-	1	-	-	-	-	1	4	1	4	2
White, both sexes	369	-	1	1	-	2	2	1	3	2	9	4
Male	150	-	-	1	-	2	2	-	-	1	6	3
Female	219	-	1	-	-	-	-	1	3	1	3	1
Black, both sexes	28	-	-	-	-	-	-	-	1	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	20	-	-	-	-	1	1	-	-	-	-	1
Male	8	-	-	-	-	1	1	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Oregon												
All races, both sexes	449	-	-	1	-	-	2	2	1	4	5	5
Male	170	-	-	-	-	-	1	1	-	3	4	4
Female	279	-	-	1	-	-	1	1	1	1	1	1
White, both sexes	443	-	-	1	-	-	2	2	-	4	4	4
Male	167	-	-	-	-	-	1	1	-	3	3	3
Female	276	-	-	1	-	-	1	1	-	1	1	1
Black, both sexes	3	-	-	-	-	-	-	-	1	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3,047	1	-	3	-	1	2	8	11	21	33	63
Male	1,343	1	-	2	-	1	2	5	7	15	15	36
Female	1,704	-	-	1	-	-	-	3	4	6	18	27
White, both sexes	2,761	1	-	3	-	1	2	6	9	14	26	53
Male	1,215	1	-	2	-	1	2	5	5	11	13	31
Female	1,546	-	-	1	-	-	-	1	4	3	13	22
Black, both sexes	275	-	-	-	-	-	-	2	1	7	7	10
Male	121	-	-	-	-	-	-	-	1	4	2	5
Female	154	-	-	-	-	-	-	2	-	3	5	5
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Rhode Island												
All races, both sexes	85	-	1	-	-	-	-	1	-	-	-	-
Male	36	-	-	-	-	-	-	1	-	-	-	-
Female	49	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	82	-	1	-	-	-	-	1	-	-	-	-
Male	34	-	-	-	-	-	-	1	-	-	-	-
Female	48	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	780	1	-	-	3	1	4	4	5	8	16	15
Male	363	-	-	-	2	-	3	1	3	2	10	10
Female	417	1	-	-	1	1	1	3	2	6	6	5
White, both sexes	552	1	-	-	1	-	1	1	2	5	8	11
Male	242	-	-	-	1	-	1	-	1	-	4	7
Female	310	1	-	-	-	-	-	1	1	5	4	4
Black, both sexes	223	-	-	-	2	1	3	3	3	3	8	4
Male	117	-	-	-	1	-	2	1	2	2	6	3
Female	106	-	-	-	1	1	1	2	1	1	2	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
South Dakota												
All races, both sexes	51	-	-	-	1	-	-	1	-	-	2	-
Male	28	-	-	-	-	-	-	1	-	-	-	-
Female	23	-	-	-	1	-	-	-	-	-	2	-
White, both sexes	43	-	-	-	-	-	-	1	-	-	2	-
Male	23	-	-	-	-	-	-	1	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	1	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	823	1	-	-	-	-	2	7	5	13	23	22
Male	387	1	-	-	-	-	1	2	4	8	15	16
Female	436	-	-	-	-	-	1	5	1	5	8	6
White, both sexes	703	1	-	-	-	-	2	6	4	11	19	18
Male	321	1	-	-	-	-	1	2	3	7	12	13
Female	382	-	-	-	-	-	1	4	1	4	7	5
Black, both sexes	116	-	-	-	-	-	-	1	1	2	4	4
Male	65	-	-	-	-	-	-	-	1	1	3	3
Female	51	-	-	-	-	-	-	1	-	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Texas												
All races, both sexes	1,784	1	3	-	4	7	7	7	6	12	25	37
Male	750	-	2	-	3	5	5	4	2	8	14	20
Female	1,034	1	1	-	1	2	2	3	4	4	11	17
White, both sexes	1,540	1	2	-	2	7	5	6	3	8	19	23
Male	637	-	2	-	2	5	4	4	-	6	11	15
Female	903	1	-	-	-	2	1	2	3	2	8	8
Black, both sexes	222	-	1	-	2	-	1	1	3	3	6	12
Male	103	-	-	-	1	-	1	-	2	1	3	5
Female	119	-	1	-	1	-	-	1	1	2	3	7
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	1	-	2
Male	10	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	2
Utah												
All races, both sexes	183	-	1	-	-	-	-	1	-	2	2	1
Male	68	-	-	-	-	-	-	-	-	2	-	-
Female	115	-	1	-	-	-	-	1	-	-	2	1
White, both sexes	182	-	1	-	-	-	-	1	-	2	2	1
Male	67	-	-	-	-	-	-	-	-	2	-	-
Female	115	-	1	-	-	-	-	1	-	-	2	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Vermont												
All races, both sexes	60	-	-	-	-	1	-	-	-	-	-	-
Male	27	-	-	-	-	1	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	60	-	-	-	-	1	-	-	-	-	-	-
Male	27	-	-	-	-	1	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,460	1	2	3	1	3	4	7	7	5	21	31
Male	656	-	-	3	1	2	3	3	2	3	14	21
Female	804	1	2	-	-	1	1	4	5	2	7	10
White, both sexes	1,166	1	1	2	1	3	3	5	4	2	18	20
Male	527	-	-	2	1	2	2	2	1	1	14	15
Female	639	1	1	-	-	1	1	3	3	1	4	5
Black, both sexes	275	-	1	1	-	-	1	2	3	3	3	10
Male	119	-	-	1	-	-	1	1	1	2	-	5
Female	156	-	1	-	-	-	-	1	2	1	3	5
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Washington												
All races, both sexes	387	-	-	1	2	-	2	1	2	4	1	3
Male	155	-	-	-	-	-	2	-	1	3	1	1
Female	232	-	-	1	2	-	-	1	1	1	-	2
White, both sexes	365	-	-	1	2	-	1	1	2	2	1	3
Male	138	-	-	-	-	-	1	-	1	1	1	1
Female	227	-	-	1	2	-	-	1	1	1	-	2
Black, both sexes	7	-	-	-	-	-	-	-	-	2	-	-
Male	7	-	-	-	-	-	-	-	-	2	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	487	-	-	-	1	-	1	1	1	3	4	7
Male	215	-	-	-	-	-	1	-	1	3	4	4
Female	272	-	-	-	1	-	-	1	-	-	-	3
White, both sexes	464	-	-	-	1	-	1	1	-	3	4	6
Male	206	-	-	-	-	-	1	-	-	3	4	4
Female	258	-	-	-	1	-	-	1	-	-	-	2
Black, both sexes	21	-	-	-	-	-	-	-	1	-	-	1
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Wisconsin												
All races, both sexes	649	-	-	-	1	2	1	3	2	3	5	18
Male	274	-	-	-	-	1	-	3	1	1	5	14
Female	375	-	-	-	1	1	1	-	1	2	-	4
White, both sexes	628	-	-	-	1	2	1	3	2	3	4	18
Male	265	-	-	-	-	1	-	3	1	1	4	14
Female	363	-	-	-	1	1	1	-	1	2	-	4
Black, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	64	-	-	-	-	-	-	-	-	2	-	2
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	-	-	2	-	1
White, both sexes	64	-	-	-	-	-	-	-	-	2	-	2
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	-	-	2	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
United States												
All races, both sexes	1,226	1,644	1,964	2,219	3,165	4,521	6,281	7,015	5,391	2,216	514	-
Male	798	1,099	1,242	1,321	1,782	2,154	2,551	2,466	1,512	491	86	-
Female	428	545	722	898	1,383	2,367	3,730	4,549	3,879	1,725	428	-
White, both sexes	898	1,248	1,543	1,747	2,588	3,867	5,631	6,373	4,923	2,000	442	-
Male	622	854	1,002	1,050	1,468	1,863	2,292	2,221	1,374	439	67	-
Female	276	394	541	697	1,120	2,004	3,339	4,152	3,549	1,561	375	-
Black, both sexes	293	351	363	392	501	533	531	530	377	170	60	-
Male	156	210	206	218	269	227	194	194	103	39	14	-
Female	137	141	157	174	232	306	337	336	274	131	46	-
American Indian 1/, both sexes	11	9	8	13	11	21	11	9	9	3	2	-
Male	5	9	6	7	9	12	3	1	5	-	-	-
Female	6	-	2	6	2	9	8	8	4	3	2	-
Asian or Pacific Islander 2/, both sexes	24	36	50	67	65	100	108	103	82	43	10	-
Male	15	26	28	46	36	52	62	50	30	13	5	-
Female	9	10	22	21	29	48	46	53	52	30	5	-
Alabama												
All races, both sexes	128	168	208	211	237	295	380	346	288	96	19	-
Male	85	102	144	117	131	151	146	125	64	24	3	-
Female	43	66	64	94	106	144	234	221	224	72	16	-
White, both sexes	92	119	147	160	182	212	308	288	242	79	8	-
Male	67	75	101	90	101	110	124	98	50	21	-	-
Female	25	44	46	70	81	102	184	190	192	58	8	-
Black, both sexes	36	49	61	49	55	81	72	58	46	17	11	-
Male	18	27	43	26	30	39	22	27	14	3	3	-
Female	18	22	18	23	25	42	50	31	32	14	8	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Alaska												
All races, both sexes	1	-	1	-	3	4	3	3	4	1	-	-
Male	1	-	1	-	3	2	1	3	1	1	-	-
Female	-	-	-	-	-	2	2	-	3	-	-	-
White, both sexes	1	-	-	-	2	4	3	3	4	1	-	-
Male	1	-	-	-	2	2	1	3	1	1	-	-
Female	-	-	-	-	-	2	2	-	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	4	12	23	20	28	44	59	64	48	17	2	-
Male	4	7	14	13	18	23	22	30	12	5	-	-
Female	-	5	9	7	10	21	37	34	36	12	2	-
White, both sexes	4	10	23	17	27	42	57	61	48	16	2	-
Male	4	5	14	11	18	22	20	30	12	4	-	-
Female	-	5	9	6	9	20	37	31	36	12	2	-
Black, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	3	-	2	2	2	-	-	-	-
Male	-	1	-	2	-	1	2	-	-	-	-	-
Female	-	-	-	1	-	1	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Arkansas												
All races, both sexes	14	20	33	25	30	38	51	71	54	16	5	-
Male	8	13	21	13	18	17	19	24	11	3	-	-
Female	6	7	12	12	12	21	32	47	43	13	5	-
White, both sexes	9	12	27	20	24	34	48	58	48	16	3	-
Male	6	7	17	12	14	15	16	21	10	3	-	-
Female	3	5	10	8	10	19	32	37	38	13	3	-
Black, both sexes	5	8	6	5	6	4	3	12	6	-	2	-
Male	2	6	4	1	4	2	3	3	1	-	-	-
Female	3	2	2	4	2	2	-	9	5	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
California												
All races, both sexes	24	18	40	51	96	163	258	334	247	119	15	-
Male	14	11	23	21	52	72	100	114	83	24	2	-
Female	10	7	17	30	44	91	158	220	164	95	13	-
White, both sexes	19	15	30	42	81	144	233	315	237	107	15	-
Male	11	9	19	17	43	67	87	106	79	19	2	-
Female	8	6	11	25	38	77	146	209	158	88	13	-
Black, both sexes	2	-	6	4	8	4	6	6	4	3	-	-
Male	1	-	2	2	4	1	4	1	1	1	-	-
Female	1	-	4	2	4	3	2	5	3	2	-	-
American Indian 1/, both sexes	-	-	-	1	-	3	1	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	2	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	3	4	4	7	12	18	13	6	8	-	-
Male	2	2	2	2	5	3	9	7	3	4	-	-
Female	1	1	2	2	2	9	9	6	3	4	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Colorado												
All races, both sexes	7	4	9	19	23	39	43	62	59	23	3	-
Male	6	2	3	13	9	18	22	18	13	3	-	-
Female	1	2	6	6	14	21	21	44	46	20	3	-
White, both sexes	7	3	9	17	22	38	43	60	59	23	3	-
Male	6	1	3	12	9	18	22	18	13	3	-	-
Female	1	2	6	5	13	20	21	42	46	20	3	-
Black, both sexes	-	-	-	2	-	1	-	2	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	26	28	35	48	62	75	126	157	130	56	9	-
Male	21	19	28	25	43	40	60	59	34	7	-	-
Female	5	9	7	23	19	35	66	98	96	49	9	-
White, both sexes	23	22	32	42	52	72	123	154	126	54	9	-
Male	19	15	26	23	36	40	60	58	34	7	-	-
Female	4	7	6	19	16	32	63	96	92	47	9	-
Black, both sexes	3	5	3	6	10	3	2	3	3	2	-	-
Male	2	4	2	2	7	-	-	1	-	-	-	-
Female	1	1	1	4	3	3	2	2	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Delaware												
All races, both sexes	1	6	4	8	5	11	12	22	10	7	1	-
Male	1	3	2	4	3	2	3	8	4	2	-	-
Female	-	3	2	4	2	9	9	14	6	5	1	-
White, both sexes	1	3	2	5	3	8	10	20	10	6	1	-
Male	1	2	1	3	1	1	3	7	4	2	-	-
Female	-	1	1	2	2	7	7	13	6	4	1	-
Black, both sexes	-	3	2	3	2	2	2	2	-	1	-	-
Male	-	1	1	1	2	1	-	1	-	-	-	-
Female	-	2	1	2	-	1	2	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	-	2	1	7	2	6	9	3	1	-	-
Male	1	-	2	1	5	1	2	1	-	-	-	-
Female	2	-	-	-	2	1	4	8	3	1	-	-
White, both sexes	-	-	1	-	-	-	1	2	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	3	-	1	1	7	2	5	7	3	1	-	-
Male	1	-	1	1	5	1	2	1	-	-	-	-
Female	2	-	-	-	2	1	3	6	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Florida												
All races, both sexes	43	53	49	66	130	175	264	329	248	88	19	-
Male	28	35	25	42	70	75	97	127	90	18	4	-
Female	15	18	24	24	60	100	167	202	158	70	15	-
White, both sexes	36	44	38	52	104	153	239	317	236	81	18	-
Male	26	31	21	33	56	67	90	122	88	17	4	-
Female	10	13	17	19	48	86	149	195	148	64	14	-
Black, both sexes	6	9	10	13	23	21	25	11	12	7	1	-
Male	2	4	4	8	12	7	7	5	2	1	-	-
Female	4	5	6	5	11	14	18	6	10	6	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	3	1	-	1	-	-	-	-
Male	-	-	-	1	2	1	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
Georgia												
All races, both sexes	93	103	138	144	204	300	348	354	249	102	33	-
Male	58	59	75	94	106	133	120	116	59	19	5	-
Female	35	44	63	50	98	167	228	238	190	83	28	-
White, both sexes	62	62	97	100	144	242	276	289	197	75	22	-
Male	38	36	55	65	76	107	98	100	46	13	3	-
Female	24	26	42	35	68	135	178	189	151	62	19	-
Black, both sexes	31	41	38	41	57	55	71	65	49	25	11	-
Male	20	23	18	26	29	24	21	16	13	5	2	-
Female	11	18	20	15	28	31	50	49	36	20	9	-
American Indian 1/, both sexes	-	-	1	-	2	1	-	-	1	1	-	-
Male	-	-	1	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	3	1	2	1	-	2	1	-	-
Male	-	-	1	3	-	1	1	-	-	1	-	-
Female	-	-	1	-	1	1	-	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Hawaii												
All races, both sexes	12	20	29	44	38	65	75	85	77	36	6	-
Male	5	17	18	30	19	38	46	39	31	8	1	-
Female	7	3	11	14	19	27	29	46	46	28	5	-
White, both sexes	3	2	5	11	8	11	15	21	20	8	2	-
Male	2	2	4	8	2	6	8	6	7	2	-	-
Female	1	-	1	3	6	5	7	15	13	6	2	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	18	24	33	29	54	60	64	57	28	4	-
Male	3	15	14	22	16	32	38	33	24	6	1	-
Female	6	3	10	11	13	22	22	31	33	22	3	-
Idaho												
All races, both sexes	3	8	5	8	7	13	23	34	30	12	4	-
Male	3	6	5	4	3	3	4	11	9	3	-	-
Female	-	2	-	4	4	10	19	23	21	9	4	-
White, both sexes	3	8	5	8	7	13	23	34	30	12	4	-
Male	3	6	5	4	3	3	4	11	9	3	-	-
Female	-	2	-	4	4	10	19	23	21	9	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Illinois												
All races, both sexes	37	62	88	71	116	175	237	272	193	82	18	-
Male	26	36	59	43	76	81	103	81	46	12	5	-
Female	11	26	29	28	40	94	134	191	147	70	13	-
White, both sexes	28	44	68	59	96	151	213	255	182	77	17	-
Male	22	25	47	37	61	75	89	74	46	12	4	-
Female	6	19	21	22	35	76	124	181	136	65	13	-
Black, both sexes	9	17	16	11	18	24	22	14	10	4	1	-
Male	4	11	9	5	13	6	13	4	-	-	1	-
Female	5	6	7	6	5	18	9	10	10	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	4	1	2	-	2	3	1	1	-	-
Male	-	-	3	1	2	-	1	3	-	-	-	-
Female	-	1	1	-	-	-	1	-	1	1	-	-
Indiana												
All races, both sexes	39	47	60	53	90	132	169	169	125	48	14	-
Male	25	31	40	33	51	57	78	61	32	12	3	-
Female	14	16	20	20	39	75	91	108	93	36	11	-
White, both sexes	35	44	54	48	84	123	163	156	120	43	10	-
Male	23	29	36	32	48	55	76	53	30	10	1	-
Female	12	15	18	16	36	68	87	103	90	33	9	-
Black, both sexes	4	3	6	5	5	9	5	13	5	5	3	-
Male	2	2	4	1	3	2	1	8	2	2	1	-
Female	2	1	2	4	2	7	4	5	3	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	1	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Iowa												
All races, both sexes	7	5	14	9	12	30	64	59	47	29	8	-
Male	4	4	8	4	7	17	28	29	16	9	4	-
Female	3	1	6	5	5	13	36	30	31	20	4	-
White, both sexes	7	5	13	9	12	30	64	59	47	29	8	-
Male	4	4	8	4	7	17	28	29	16	9	4	-
Female	3	1	5	5	5	13	36	30	31	20	4	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	8	14	17	17	26	52	50	70	68	23	7	-
Male	6	7	14	12	17	26	26	25	22	8	1	-
Female	2	7	3	5	9	26	24	45	46	15	6	-
White, both sexes	8	14	15	17	22	47	48	68	66	20	7	-
Male	6	7	13	12	14	22	24	25	20	6	1	-
Female	2	7	2	5	8	25	24	43	46	14	6	-
Black, both sexes	-	-	2	-	4	2	2	1	1	3	-	-
Male	-	-	1	-	3	2	2	-	1	2	-	-
Female	-	-	1	-	1	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Kentucky												
All races, both sexes	32	39	56	53	84	104	158	155	106	50	7	-
Male	19	28	38	27	50	47	66	43	26	13	1	-
Female	13	11	18	26	34	57	92	112	80	37	6	-
White, both sexes	28	38	53	48	75	93	145	146	100	47	6	-
Male	17	27	36	26	45	39	65	39	25	12	1	-
Female	11	11	17	22	30	54	80	107	75	35	5	-
Black, both sexes	4	1	2	5	8	11	13	9	6	3	1	-
Male	2	1	1	1	5	8	1	4	1	1	-	-
Female	2	-	1	4	3	3	12	5	5	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	32	44	39	52	54	80	86	119	87	25	6	-
Male	20	25	23	31	32	32	40	36	24	6	3	-
Female	12	19	16	21	22	48	46	83	63	19	3	-
White, both sexes	20	20	26	31	37	59	72	86	67	15	5	-
Male	16	15	17	17	21	27	36	31	17	2	2	-
Female	4	5	9	14	16	32	36	55	50	13	3	-
Black, both sexes	12	23	13	21	17	21	14	33	20	10	1	-
Male	4	10	6	14	11	5	4	5	7	4	1	-
Female	8	13	7	7	6	16	10	28	13	6	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Maine												
All races, both sexes	7	12	7	16	21	23	29	32	36	20	4	-
Male	5	11	5	13	14	12	11	12	11	3	-	-
Female	2	1	2	3	7	11	18	20	25	17	4	-
White, both sexes	7	12	7	16	20	23	29	31	36	20	4	-
Male	5	11	5	13	13	12	11	11	11	3	-	-
Female	2	1	2	3	7	11	18	20	25	17	4	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	12	22	30	27	39	67	107	102	80	32	3	-
Male	8	14	18	16	16	24	37	46	18	8	1	-
Female	4	8	12	11	23	43	70	56	62	24	2	-
White, both sexes	5	13	14	20	27	52	87	84	64	30	2	-
Male	4	8	7	12	11	20	34	37	16	8	1	-
Female	1	5	7	8	16	32	53	47	48	22	1	-
Black, both sexes	6	9	14	6	11	15	17	17	16	2	1	-
Male	3	6	9	3	4	4	1	9	2	-	-	-
Female	3	3	5	3	7	11	16	8	14	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	1	1	-	3	1	-	-	-	-
Male	1	-	2	1	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Massachusetts												
All races, both sexes	42	58	60	74	116	132	197	239	191	81	22	-
Male	24	41	39	45	66	69	91	95	50	17	1	-
Female	18	17	21	29	50	63	106	144	141	64	21	-
White, both sexes	38	49	56	69	109	122	183	233	188	78	21	-
Male	21	37	37	44	64	61	83	90	50	16	1	-
Female	17	12	19	25	45	61	100	143	138	62	20	-
Black, both sexes	4	7	4	5	6	6	8	5	1	3	-	-
Male	3	3	2	1	2	4	5	4	-	1	-	-
Female	1	4	2	4	4	2	3	1	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	1	4	5	1	1	-	1	-
Male	-	1	-	-	-	4	3	1	-	-	-	-
Female	-	1	-	-	1	-	2	-	1	-	1	-
Michigan												
All races, both sexes	23	30	27	44	65	113	183	216	189	82	16	-
Male	10	20	14	28	31	51	71	77	50	21	-	-
Female	13	10	13	16	34	62	112	139	139	61	16	-
White, both sexes	15	25	21	36	53	97	168	202	177	77	14	-
Male	8	17	12	24	24	42	69	73	45	20	-	-
Female	7	8	9	12	29	55	99	129	132	57	14	-
Black, both sexes	8	4	6	6	11	14	13	14	12	5	2	-
Male	2	2	2	3	6	8	2	4	5	1	-	-
Female	6	2	4	3	5	6	11	10	7	4	2	-
American Indian 1/, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	1	2	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Minnesota												
All races, both sexes	11	14	23	33	38	58	71	123	102	47	6	-
Male	7	10	13	27	28	36	33	43	34	15	2	-
Female	4	4	10	6	10	22	38	80	68	32	4	-
White, both sexes	7	14	22	33	37	56	71	121	102	46	5	-
Male	6	10	12	27	27	35	33	42	34	15	1	-
Female	1	4	10	6	10	21	38	79	68	31	4	-
Black, both sexes	1	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	1	1	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	1	-	1	1	-
Male	-	-	-	-	-	1	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	1	-	1	-	-
Mississippi												
All races, both sexes	23	54	48	49	61	96	116	93	91	44	8	-
Male	13	39	29	31	31	40	44	26	25	13	2	-
Female	10	15	19	18	30	56	72	67	66	31	6	-
White, both sexes	10	26	27	24	36	63	87	64	67	24	3	-
Male	7	18	17	16	18	28	32	12	17	5	1	-
Female	3	8	10	8	18	35	55	52	50	19	2	-
Black, both sexes	13	28	19	24	25	32	29	29	23	20	5	-
Male	6	21	11	14	13	11	12	14	7	8	1	-
Female	7	7	8	10	12	21	17	15	16	12	4	-
American Indian 1/, both sexes	-	-	1	-	-	1	-	-	1	-	-	-
Male	-	-	1	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Missouri												
All races, both sexes	37	58	58	56	90	106	162	146	141	64	17	-
Male	19	41	37	34	51	50	68	59	43	17	4	-
Female	18	17	21	22	39	56	94	87	98	47	13	-
White, both sexes	30	52	52	51	79	98	154	136	133	63	14	-
Male	17	38	34	31	46	48	65	57	40	17	2	-
Female	13	14	18	20	33	50	89	79	93	46	12	-
Black, both sexes	7	6	6	5	11	6	8	9	8	-	3	-
Male	2	3	3	3	5	2	3	2	3	-	2	-
Female	5	3	3	2	6	4	5	7	5	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	1	-	1	-	-
Montana												
All races, both sexes	7	4	9	8	8	15	19	27	32	14	-	-
Male	5	1	8	5	6	10	10	8	6	4	-	-
Female	2	3	1	3	2	5	9	19	26	10	-	-
White, both sexes	5	4	9	8	8	14	19	27	32	13	-	-
Male	5	1	8	5	6	9	10	8	6	4	-	-
Female	-	3	1	3	2	5	9	19	26	9	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Nebraska												
All races, both sexes	7	12	17	15	33	37	66	76	70	40	10	-
Male	6	8	11	7	27	17	31	26	21	9	4	-
Female	1	4	6	8	6	20	35	50	49	31	6	-
White, both sexes	6	12	15	14	32	36	66	75	69	40	9	-
Male	5	8	9	7	26	16	31	25	20	9	4	-
Female	1	4	6	7	6	20	35	50	49	31	5	-
Black, both sexes	1	-	2	1	1	1	-	1	1	-	-	-
Male	1	-	2	-	1	1	-	1	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	48	60	77	79	81	82	73	73	47	11	4	-
Male	33	47	51	54	54	45	30	27	16	1	1	-
Female	15	13	26	25	27	37	43	46	31	10	3	-
White, both sexes	37	47	64	66	65	72	68	65	42	11	4	-
Male	26	38	42	44	44	40	29	23	15	1	1	-
Female	11	9	22	22	21	32	39	42	27	10	3	-
Black, both sexes	8	6	10	6	10	7	3	7	2	-	-	-
Male	5	3	7	6	6	3	1	4	-	-	-	-
Female	3	3	3	-	4	4	2	3	2	-	-	-
American Indian 1/, both sexes	1	2	-	2	1	-	-	-	1	-	-	-
Male	-	2	-	1	1	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	3	5	5	3	2	1	2	-	-	-
Male	2	4	2	3	3	2	-	-	1	-	-	-
Female	-	1	1	2	2	1	2	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
New Hampshire												
All races, both sexes	2	1	2	5	8	9	16	29	19	6	6	-
Male	1	1	2	1	4	1	10	11	10	2	-	-
Female	1	-	-	4	4	8	6	18	9	4	6	-
White, both sexes	2	1	2	4	8	9	16	29	19	6	6	-
Male	1	1	2	1	4	1	10	11	10	2	-	-
Female	1	-	-	3	4	8	6	18	9	4	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	32	36	39	57	77	139	189	219	157	67	19	-
Male	20	22	27	38	43	70	83	80	44	16	5	-
Female	12	14	12	19	34	69	106	139	113	51	14	-
White, both sexes	21	31	27	41	58	123	177	204	146	64	16	-
Male	13	19	17	27	32	60	77	75	40	15	4	-
Female	8	12	10	14	26	63	100	129	106	49	12	-
Black, both sexes	10	5	10	14	17	15	11	11	8	3	2	-
Male	6	3	8	10	10	10	5	4	3	1	-	-
Female	4	2	2	4	7	5	6	7	5	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	2	2	1	1	3	3	-	1	-
Male	1	-	2	1	1	-	1	1	1	-	1	-
Female	-	-	-	1	1	1	-	2	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
New Mexico												
All races, both sexes	2	6	5	7	20	16	29	34	19	11	2	-
Male	2	2	1	4	14	10	13	19	7	5	1	-
Female	-	4	4	3	6	6	16	15	12	6	1	-
White, both sexes	2	6	4	7	18	16	28	34	18	11	2	-
Male	2	2	1	4	13	10	12	19	6	5	1	-
Female	-	4	3	3	5	6	16	15	12	6	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
New York												
All races, both sexes	35	48	66	74	137	204	268	348	300	122	36	-
Male	23	30	45	44	66	110	121	111	93	31	5	-
Female	12	18	21	30	71	94	147	237	207	91	31	-
White, both sexes	28	41	54	65	115	179	250	332	287	115	31	-
Male	22	27	37	35	61	102	112	108	91	30	3	-
Female	6	14	17	30	54	77	138	224	196	85	28	-
Black, both sexes	6	7	12	6	20	21	16	14	10	7	3	-
Male	1	3	8	6	5	7	8	3	2	1	1	-
Female	5	4	4	-	15	14	8	11	8	6	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	3	2	4	2	2	3	-	2	-
Male	-	-	-	3	-	1	1	-	-	-	1	-
Female	1	-	-	-	2	3	1	2	3	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
North Carolina												
All races, both sexes	45	62	56	65	122	162	215	183	144	72	21	-
Male	32	49	34	38	71	74	79	56	37	15	5	-
Female	13	13	22	27	51	88	136	127	107	57	16	-
White, both sexes	28	42	38	52	92	130	187	157	122	58	20	-
Male	21	31	24	31	54	55	72	47	32	12	5	-
Female	7	11	14	21	38	75	115	110	90	46	15	-
Black, both sexes	16	18	17	13	29	29	25	25	22	14	1	-
Male	10	16	10	7	16	17	7	8	5	3	-	-
Female	6	2	7	6	13	12	18	17	17	11	1	-
American Indian 1/, both sexes	-	2	1	-	1	2	3	1	-	-	-	-
Male	-	2	-	-	1	2	-	1	-	-	-	-
Female	-	-	1	-	-	-	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	1	3	-	7	5	11	16	16	5	-	-
Male	2	1	1	-	1	-	7	6	5	-	-	-
Female	-	-	2	-	6	5	4	10	11	5	-	-
White, both sexes	2	1	3	-	7	5	11	16	16	5	-	-
Male	2	1	1	-	1	-	7	6	5	-	-	-
Female	-	-	2	-	6	5	4	10	11	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Ohio												
All races, both sexes	48	63	65	94	114	186	292	296	232	87	19	-
Male	30	44	35	53	58	88	116	91	57	16	4	-
Female	18	19	30	41	56	98	176	205	175	71	15	-
White, both sexes	36	54	47	74	100	166	279	275	227	83	17	-
Male	22	37	27	41	52	81	110	85	57	15	4	-
Female	14	17	20	33	48	85	169	190	170	68	13	-
Black, both sexes	12	9	17	19	14	19	13	21	5	4	2	-
Male	8	7	8	11	6	6	6	6	-	1	-	-
Female	4	2	9	8	8	13	7	15	5	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	16	14	14	35	39	53	62	67	58	20	8	-
Male	9	8	12	20	15	21	18	22	15	6	2	-
Female	7	6	2	15	24	32	44	45	43	14	6	-
White, both sexes	12	12	13	26	36	46	55	64	55	18	7	-
Male	8	8	11	14	14	21	17	21	14	5	2	-
Female	4	4	2	12	22	25	38	43	41	13	5	-
Black, both sexes	3	2	-	4	1	3	6	2	1	2	1	-
Male	1	-	-	3	-	-	1	1	-	1	-	-
Female	2	2	-	1	1	3	5	1	1	1	1	-
American Indian 1/, both sexes	1	-	1	5	2	4	1	1	2	-	-	-
Male	-	-	1	3	1	-	-	-	1	-	-	-
Female	1	-	-	2	1	4	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Oregon												
All races, both sexes	14	11	14	14	17	45	92	91	88	40	3	-
Male	10	7	10	7	11	12	38	34	21	7	-	-
Female	4	4	4	7	6	33	54	57	67	33	3	-
White, both sexes	14	11	14	14	17	45	91	91	86	40	3	-
Male	10	7	10	7	11	12	37	34	21	7	-	-
Female	4	4	4	7	6	33	54	57	65	33	3	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Pennsylvania												
All races, both sexes	95	114	150	174	246	376	555	606	394	160	34	-
Male	66	78	88	102	140	188	219	229	109	36	4	-
Female	29	36	62	72	106	188	336	377	285	124	30	-
White, both sexes	70	91	133	143	216	340	523	573	372	154	31	-
Male	51	63	80	87	123	168	207	220	105	36	4	-
Female	19	28	53	56	93	172	316	353	267	118	27	-
Black, both sexes	23	21	17	30	28	35	31	32	22	6	3	-
Male	13	13	8	15	16	19	12	9	4	-	-	-
Female	10	8	9	15	12	16	19	23	18	6	3	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	1	2	1	-	1	-	-	-	-
Male	2	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Rhode Island												
All races, both sexes	3	3	4	4	5	10	9	20	16	5	4	-
Male	3	2	3	3	2	5	5	6	3	1	2	-
Female	-	1	1	1	3	5	4	14	13	4	2	-
White, both sexes	3	3	3	4	5	9	8	20	16	5	4	-
Male	3	2	3	3	2	4	4	6	3	1	2	-
Female	-	1	-	1	3	5	4	14	13	4	2	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	29	36	35	49	70	107	138	136	79	37	7	-
Male	21	23	22	28	50	55	50	50	28	5	-	-
Female	8	13	13	21	20	52	88	86	51	32	7	-
White, both sexes	14	20	20	24	45	80	108	115	63	29	4	-
Male	9	15	14	12	32	43	40	38	20	5	-	-
Female	5	5	6	12	13	37	68	77	43	24	4	-
Black, both sexes	14	16	14	24	24	27	30	21	16	8	2	-
Male	11	8	7	15	17	12	10	12	8	-	-	-
Female	3	8	7	9	7	15	20	9	8	8	2	-
American Indian 1/, both sexes	1	-	1	-	1	-	-	-	-	-	1	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
South Dakota												
All races, both sexes	5	1	3	3	3	5	12	5	7	3	-	-
Male	5	1	2	3	1	4	8	3	-	-	-	-
Female	-	-	1	-	2	1	4	2	7	3	-	-
White, both sexes	2	-	3	3	3	3	11	5	7	3	-	-
Male	2	-	2	3	1	3	8	3	-	-	-	-
Female	-	-	1	-	2	-	3	2	7	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	2	1	-	-	-	-	-
Male	3	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	41	63	52	57	73	83	105	146	92	27	11	-
Male	28	49	36	34	34	36	37	51	27	7	1	-
Female	13	14	16	23	39	47	68	95	65	20	10	-
White, both sexes	28	47	46	40	62	75	95	133	80	26	10	-
Male	20	37	31	26	27	33	31	46	24	7	-	-
Female	8	10	15	14	35	42	64	87	56	19	10	-
Black, both sexes	13	16	6	17	11	6	9	12	12	1	1	-
Male	8	12	5	8	7	2	6	5	3	-	1	-
Female	5	4	1	9	4	4	3	7	9	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Texas												
All races, both sexes	50	65	85	89	153	194	322	330	252	112	23	-
Male	34	35	51	49	96	92	128	110	62	26	4	-
Female	16	30	34	40	57	102	194	220	190	86	19	-
White, both sexes	34	52	65	77	132	173	285	295	225	106	20	-
Male	25	31	43	42	84	84	108	95	49	24	3	-
Female	9	21	22	35	48	89	177	200	176	82	17	-
Black, both sexes	14	12	18	11	21	18	33	32	25	6	3	-
Male	7	4	8	6	12	7	18	12	13	2	1	-
Female	7	8	10	5	9	11	15	20	12	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	2	1	-	3	4	3	2	-	-	-
Male	2	-	-	1	-	1	2	3	-	-	-	-
Female	-	1	2	-	-	2	2	-	2	-	-	-
Utah												
All races, both sexes	3	1	9	5	12	26	27	44	33	13	3	-
Male	1	-	7	3	2	14	9	14	13	3	-	-
Female	2	1	2	2	10	12	18	30	20	10	3	-
White, both sexes	3	1	8	5	12	26	27	44	33	13	3	-
Male	1	-	6	3	2	14	9	14	13	3	-	-
Female	2	1	2	2	10	12	18	30	20	10	3	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Vermont												
All races, both sexes	1	1	-	1	5	8	5	17	15	4	2	-
Male	-	1	-	1	3	3	2	7	7	1	1	-
Female	1	-	-	-	2	5	3	10	8	3	1	-
White, both sexes	1	1	-	1	5	8	5	17	15	4	2	-
Male	-	1	-	1	3	3	2	7	7	1	1	-
Female	1	-	-	-	2	5	3	10	8	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	47	88	79	95	127	176	234	261	188	65	15	-
Male	28	69	45	53	61	81	102	98	54	10	3	-
Female	19	19	34	42	66	95	132	163	134	55	12	-
White, both sexes	28	64	58	62	89	139	205	227	163	56	15	-
Male	16	54	36	34	47	68	88	83	49	9	3	-
Female	12	10	22	28	42	71	117	144	114	47	12	-
Black, both sexes	18	23	19	30	36	34	29	30	24	8	-	-
Male	11	14	9	16	12	12	14	14	5	1	-	-
Female	7	9	10	14	24	22	15	16	19	7	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	3	2	3	-	3	1	1	-	-
Male	1	1	-	3	2	1	-	1	-	-	-	-
Female	-	-	2	-	-	2	-	2	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Washington												
All races, both sexes	5	12	15	19	19	41	70	90	60	30	10	-
Male	3	7	11	11	12	21	23	28	21	10	-	-
Female	2	5	4	8	7	20	47	62	39	20	10	-
White, both sexes	5	10	15	17	17	38	65	87	59	29	10	-
Male	3	5	11	10	10	18	20	27	20	9	-	-
Female	2	5	4	7	7	20	45	60	39	20	10	-
Black, both sexes	-	1	-	-	-	2	2	-	-	-	-	-
Male	-	1	-	-	-	2	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	1	1	3	3	1	1	-	-
Male	-	1	-	1	1	1	1	1	1	1	-	-
Female	-	-	-	1	-	-	2	2	-	-	-	-
West Virginia												
All races, both sexes	9	18	35	37	42	65	98	85	59	12	9	-
Male	5	10	22	23	22	38	34	29	16	1	2	-
Female	4	8	13	14	20	27	64	56	43	11	7	-
White, both sexes	8	18	33	33	41	64	95	78	57	12	9	-
Male	5	10	20	21	22	37	33	27	16	1	2	-
Female	3	8	13	12	19	27	62	51	41	11	7	-
Black, both sexes	1	-	1	4	1	1	3	6	2	-	-	-
Male	-	-	1	2	-	1	1	2	-	-	-	-
Female	1	-	-	2	1	-	2	4	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Conduction disorders and cardiac dysrhythmias (I44-I49)												
Wisconsin												
All races, both sexes	10	23	24	26	55	79	89	139	105	50	14	-
Male	9	21	19	15	31	39	39	39	29	8	-	-
Female	1	2	5	11	24	40	50	100	76	42	14	-
White, both sexes	10	21	22	24	52	78	86	135	102	50	14	-
Male	9	19	18	15	29	39	39	37	28	8	-	-
Female	1	2	4	9	23	39	47	98	74	42	14	-
Black, both sexes	-	1	1	1	3	1	3	3	1	-	-	-
Male	-	1	-	-	2	-	-	2	-	-	-	-
Female	-	-	1	1	1	1	3	1	1	-	-	-
American Indian 1/, both sexes	-	1	1	1	-	-	-	1	1	-	-	-
Male	-	1	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Wyoming												
All races, both sexes	1	2	3	8	10	6	8	12	6	2	2	-
Male	1	2	1	4	8	3	1	3	3	-	-	-
Female	-	-	2	4	2	3	7	9	3	2	2	-
White, both sexes	1	2	3	8	10	6	8	12	6	2	2	-
Male	1	2	1	4	8	3	1	3	3	-	-	-
Female	-	-	2	4	2	3	7	9	3	2	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
United States												
All races, both sexes	58,933	27	6	3	8	12	30	33	61	96	220	384
Male	23,026	14	4	3	2	9	18	20	41	60	147	230
Female	35,907	13	2	-	6	3	12	13	20	36	73	154
White, both sexes	52,570	15	3	1	3	8	17	15	32	51	127	230
Male	20,414	8	3	1	-	5	13	10	22	32	79	140
Female	32,156	7	-	-	3	3	4	5	10	19	48	90
Black, both sexes	5,609	8	3	1	5	4	12	18	28	43	88	143
Male	2,275	4	1	1	2	4	5	10	19	27	63	83
Female	3,334	4	2	-	3	-	7	8	9	16	25	60
American Indian 1/, both sexes	228	2	-	-	-	-	1	-	-	1	2	6
Male	106	1	-	-	-	-	-	-	-	1	2	3
Female	122	1	-	-	-	-	1	-	-	-	-	3
Asian or Pacific Islander 2/, both sexes	526	2	-	1	-	-	-	-	1	1	3	5
Male	231	1	-	1	-	-	-	-	-	-	3	4
Female	295	1	-	-	-	-	-	-	1	1	-	1
Alabama												
All races, both sexes	2,224	-	-	-	1	1	2	2	8	7	20	28
Male	902	-	-	-	-	-	1	1	6	6	15	16
Female	1,322	-	-	-	1	1	1	1	2	1	5	12
White, both sexes	1,724	-	-	-	-	1	-	-	2	4	8	13
Male	689	-	-	-	-	-	-	-	1	3	7	7
Female	1,035	-	-	-	-	1	-	-	1	1	1	6
Black, both sexes	495	-	-	-	1	-	2	2	6	3	12	15
Male	210	-	-	-	-	-	1	1	5	3	8	9
Female	285	-	-	-	1	-	1	1	1	-	4	6
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Alaska												
All races, both sexes	61	-	-	-	-	-	-	-	-	-	-	-
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	46	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	736	2	-	-	-	-	1	-	-	2	4	6
Male	301	1	-	-	-	-	-	-	-	-	2	4
Female	435	1	-	-	-	-	1	-	-	2	2	2
White, both sexes	705	-	-	-	-	-	-	-	-	1	4	4
Male	287	-	-	-	-	-	-	-	-	-	2	3
Female	418	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	15	1	-	-	-	-	1	-	-	-	-	-
Male	8	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	1	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Arkansas												
All races, both sexes	720	-	-	-	-	1	1	-	1	1	5	2
Male	313	-	-	-	-	1	1	-	1	1	4	2
Female	407	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	652	-	-	-	-	-	-	-	1	-	4	2
Male	284	-	-	-	-	-	-	-	1	-	3	2
Female	368	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	67	-	-	-	-	1	1	-	-	1	1	-
Male	28	-	-	-	-	1	1	-	-	1	1	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4,336	1	-	1	-	-	2	1	3	6	18	22
Male	1,755	1	-	1	-	-	2	1	2	4	10	13
Female	2,581	-	-	-	-	-	-	-	1	2	8	9
White, both sexes	3,877	1	-	-	-	-	2	1	3	3	12	16
Male	1,563	1	-	-	-	-	2	1	2	2	5	11
Female	2,314	-	-	-	-	-	-	-	1	1	7	5
Black, both sexes	261	-	-	-	-	-	-	-	-	3	4	6
Male	112	-	-	-	-	-	-	-	-	2	3	2
Female	149	-	-	-	-	-	-	-	-	1	1	4
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	183	-	-	1	-	-	-	-	-	-	1	-
Male	73	-	-	1	-	-	-	-	-	-	1	-
Female	110	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Colorado												
All races, both sexes	699	-	-	-	-	-	-	-	1	1	2	2
Male	259	-	-	-	-	-	-	-	1	1	2	2
Female	440	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	676	-	-	-	-	-	-	-	1	-	2	1
Male	247	-	-	-	-	-	-	-	1	-	2	1
Female	429	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	698	1	1	-	-	-	-	-	-	2	1	4
Male	264	-	1	-	-	-	-	-	-	2	-	4
Female	434	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	645	1	1	-	-	-	-	-	-	2	1	2
Male	238	-	1	-	-	-	-	-	-	2	-	2
Female	407	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	47	-	-	-	-	-	-	-	-	-	-	2
Male	24	-	-	-	-	-	-	-	-	-	-	2
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Delaware												
All races, both sexes	139	-	-	-	-	-	-	-	1	-	-	2
Male	57	-	-	-	-	-	-	-	1	-	-	2
Female	82	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	115	-	-	-	-	-	-	-	-	-	-	1
Male	49	-	-	-	-	-	-	-	-	-	-	1
Female	66	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	52	-	-	-	-	-	-	-	-	-	1	-
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	41	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Florida												
All races, both sexes	1,899	-	-	-	-	-	1	1	2	3	2	9
Male	790	-	-	-	-	-	-	-	2	1	1	4
Female	1,109	-	-	-	-	-	1	1	-	2	1	5
White, both sexes	1,748	-	-	-	-	-	1	1	-	1	1	6
Male	725	-	-	-	-	-	-	-	-	-	-	2
Female	1,023	-	-	-	-	-	1	1	-	1	1	4
Black, both sexes	141	-	-	-	-	-	-	-	2	2	1	3
Male	60	-	-	-	-	-	-	-	2	1	1	2
Female	81	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	2,076	1	-	-	-	3	3	3	-	9	21	25
Male	809	-	-	-	-	2	1	1	-	6	10	14
Female	1,267	1	-	-	-	1	2	2	-	3	11	11
White, both sexes	1,604	1	-	-	-	2	-	1	-	2	11	14
Male	606	-	-	-	-	1	-	1	-	1	5	9
Female	998	1	-	-	-	1	-	-	-	1	6	5
Black, both sexes	463	-	-	-	-	1	3	2	-	7	10	11
Male	197	-	-	-	-	1	1	-	-	5	5	5
Female	266	-	-	-	-	-	2	2	-	2	5	6
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Hawaii												
All races, both sexes	161	-	-	-	1	-	-	-	-	1	3	2
Male	89	-	-	-	1	-	-	-	-	-	3	-
Female	72	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	56	-	-	-	-	-	-	-	-	-	1	1
Male	29	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	1	-	-	-	-	-	-	-
Male	4	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	100	-	-	-	-	-	-	-	-	1	2	1
Male	55	-	-	-	-	-	-	-	-	-	2	-
Female	45	-	-	-	-	-	-	-	-	1	-	1
Idaho												
All races, both sexes	252	-	-	-	1	-	-	-	1	-	-	-
Male	103	-	-	-	-	-	-	-	1	-	-	-
Female	149	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	250	-	-	-	1	-	-	-	1	-	-	-
Male	102	-	-	-	-	-	-	-	1	-	-	-
Female	148	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Illinois												
All races, both sexes	3,128	-	2	-	-	-	3	3	1	2	12	21
Male	1,190	-	1	-	-	-	2	3	1	1	9	15
Female	1,938	-	1	-	-	-	1	-	-	1	3	6
White, both sexes	2,814	-	1	-	-	-	3	1	1	1	8	13
Male	1,062	-	1	-	-	-	2	1	1	1	6	11
Female	1,752	-	-	-	-	-	1	-	-	-	2	2
Black, both sexes	287	-	1	-	-	-	-	2	-	1	4	7
Male	117	-	-	-	-	-	-	2	-	-	3	3
Female	170	-	1	-	-	-	-	-	-	1	1	4
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,963	1	-	-	-	2	1	2	2	3	3	10
Male	734	1	-	-	-	2	1	2	1	2	3	4
Female	1,229	-	-	-	-	-	-	-	1	1	-	6
White, both sexes	1,847	1	-	-	-	2	1	2	2	2	2	6
Male	694	1	-	-	-	2	1	2	1	2	2	3
Female	1,153	-	-	-	-	-	-	-	1	-	-	3
Black, both sexes	112	-	-	-	-	-	-	-	-	1	1	4
Male	39	-	-	-	-	-	-	-	-	-	1	1
Female	73	-	-	-	-	-	-	-	-	1	-	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Iowa												
All races, both sexes	302	-	-	-	-	-	-	-	-	-	-	1
Male	108	-	-	-	-	-	-	-	-	-	-	1
Female	194	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	297	-	-	-	-	-	-	-	-	-	-	1
Male	104	-	-	-	-	-	-	-	-	-	-	1
Female	193	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	912	-	-	-	-	-	1	-	3	1	3	2
Male	339	-	-	-	-	-	1	-	3	1	3	1
Female	573	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	860	-	-	-	-	-	1	-	3	-	3	-
Male	321	-	-	-	-	-	1	-	3	-	3	-
Female	539	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	42	-	-	-	-	-	-	-	-	1	-	1
Male	14	-	-	-	-	-	-	-	-	1	-	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Kentucky												
All races, both sexes	1,398	3	-	-	-	-	1	-	4	3	4	7
Male	537	1	-	-	-	-	-	-	2	-	2	5
Female	861	2	-	-	-	-	1	-	2	3	2	2
White, both sexes	1,314	-	-	-	-	-	-	-	4	3	4	5
Male	504	-	-	-	-	-	-	-	2	-	2	4
Female	810	-	-	-	-	-	-	-	2	3	2	1
Black, both sexes	80	3	-	-	-	-	1	-	-	-	-	2
Male	30	1	-	-	-	-	-	-	-	-	-	1
Female	50	2	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,512	1	-	-	1	-	1	2	3	3	11	13
Male	606	-	-	-	-	-	1	1	1	2	6	5
Female	906	1	-	-	1	-	-	1	2	1	5	8
White, both sexes	1,133	1	-	-	-	-	1	-	1	2	4	2
Male	450	-	-	-	-	-	1	-	-	2	1	1
Female	683	1	-	-	-	-	-	-	1	-	3	1
Black, both sexes	371	-	-	-	1	-	-	2	2	1	7	11
Male	154	-	-	-	-	-	-	1	1	-	5	4
Female	217	-	-	-	1	-	-	1	1	1	2	7
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Maine												
All races, both sexes	271	1	-	-	-	-	1	-	-	1	1	-
Male	92	-	-	-	-	-	1	-	-	1	-	-
Female	179	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	270	1	-	-	-	-	1	-	-	1	1	-
Male	92	-	-	-	-	-	1	-	-	1	-	-
Female	178	1	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	738	-	-	-	-	-	-	-	1	1	5	7
Male	299	-	-	-	-	-	-	-	-	-	3	5
Female	439	-	-	-	-	-	-	-	1	1	2	2
White, both sexes	590	-	-	-	-	-	-	-	-	-	1	3
Male	235	-	-	-	-	-	-	-	-	-	-	1
Female	355	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	143	-	-	-	-	-	-	-	1	1	4	4
Male	62	-	-	-	-	-	-	-	-	-	3	4
Female	81	-	-	-	-	-	-	-	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Massachusetts												
All races, both sexes	1,908	1	-	-	-	-	-	-	-	1	4	5
Male	682	-	-	-	-	-	-	-	-	1	3	5
Female	1,226	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	1,840	-	-	-	-	-	-	-	-	1	3	5
Male	659	-	-	-	-	-	-	-	-	1	2	5
Female	1,181	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	59	1	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	1	-
Female	43	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2,484	1	-	-	-	1	-	-	-	5	5	9
Male	948	1	-	-	-	1	-	-	-	4	3	5
Female	1,536	-	-	-	-	-	-	-	-	1	2	4
White, both sexes	2,258	1	-	-	-	-	-	-	-	2	3	1
Male	850	1	-	-	-	-	-	-	-	1	1	-
Female	1,408	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	207	-	-	-	-	1	-	-	-	3	2	8
Male	93	-	-	-	-	1	-	-	-	3	2	5
Female	114	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Minnesota												
All races, both sexes	1,144	-	-	-	-	1	-	-	-	-	1	3
Male	461	-	-	-	-	1	-	-	-	-	1	1
Female	683	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	1,123	-	-	-	-	1	-	-	-	-	1	2
Male	450	-	-	-	-	1	-	-	-	-	1	-
Female	673	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,423	-	-	-	-	1	3	3	3	6	3	15
Male	527	-	-	-	-	1	1	1	2	3	3	8
Female	896	-	-	-	-	-	2	2	1	3	-	7
White, both sexes	1,029	-	-	-	-	-	-	-	1	2	-	7
Male	365	-	-	-	-	-	-	-	-	-	-	3
Female	664	-	-	-	-	-	-	-	1	2	-	4
Black, both sexes	392	-	-	-	-	1	3	3	2	4	3	8
Male	161	-	-	-	-	1	1	1	2	3	3	5
Female	231	-	-	-	-	-	2	2	-	1	-	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Missouri												
All races, both sexes	1,489	1	-	1	1	-	-	2	-	3	2	8
Male	545	1	-	1	-	-	-	1	-	2	2	4
Female	944	-	-	-	1	-	-	1	-	1	-	4
White, both sexes	1,392	1	-	1	1	-	-	1	-	2	2	5
Male	503	1	-	1	-	-	-	-	-	1	2	3
Female	889	-	-	-	1	-	-	1	-	1	-	2
Black, both sexes	93	-	-	-	-	-	-	1	-	1	-	3
Male	39	-	-	-	-	-	-	1	-	1	-	1
Female	54	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	286	-	-	-	-	-	-	-	-	1	-	-
Male	128	-	-	-	-	-	-	-	-	1	-	-
Female	158	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	274	-	-	-	-	-	-	-	-	1	-	-
Male	122	-	-	-	-	-	-	-	-	1	-	-
Female	152	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Nebraska												
All races, both sexes	559	-	-	-	-	-	-	-	-	-	1	9
Male	209	-	-	-	-	-	-	-	-	-	-	6
Female	350	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	546	-	-	-	-	-	-	-	-	-	1	9
Male	204	-	-	-	-	-	-	-	-	-	-	6
Female	342	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	561	-	-	-	-	-	-	-	-	-	4	3
Male	260	-	-	-	-	-	-	-	-	-	2	2
Female	301	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	517	-	-	-	-	-	-	-	-	-	2	2
Male	241	-	-	-	-	-	-	-	-	-	-	1
Female	276	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	27	-	-	-	-	-	-	-	-	-	2	1
Male	12	-	-	-	-	-	-	-	-	-	2	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
New Hampshire												
All races, both sexes	188	-	-	-	-	-	-	-	1	-	1	1
Male	66	-	-	-	-	-	-	-	1	-	1	-
Female	122	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	188	-	-	-	-	-	-	-	1	-	1	1
Male	66	-	-	-	-	-	-	-	1	-	1	-
Female	122	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,589	1	-	-	1	-	1	1	2	1	5	15
Male	616	1	-	-	-	-	1	1	1	-	2	11
Female	973	-	-	-	1	-	-	-	1	1	3	4
White, both sexes	1,401	1	-	-	-	-	1	1	-	1	-	3
Male	547	1	-	-	-	-	1	1	-	-	-	2
Female	854	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	168	-	-	-	1	-	-	-	2	-	5	10
Male	61	-	-	-	-	-	-	-	1	-	2	7
Female	107	-	-	-	1	-	-	-	1	-	3	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
New Mexico												
All races, both sexes	385	2	-	-	-	-	-	-	-	-	-	2
Male	150	1	-	-	-	-	-	-	-	-	-	-
Female	235	1	-	-	-	-	-	-	-	-	-	2
White, both sexes	355	1	-	-	-	-	-	-	-	-	-	1
Male	141	1	-	-	-	-	-	-	-	-	-	-
Female	214	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	20	1	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	14	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2,825	1	-	-	-	-	1	4	-	2	4	14
Male	1,111	-	-	-	-	-	1	3	-	-	2	9
Female	1,714	1	-	-	-	-	-	1	-	2	2	5
White, both sexes	2,559	-	-	-	-	-	1	2	-	2	1	11
Male	1,016	-	-	-	-	-	1	1	-	-	-	7
Female	1,543	-	-	-	-	-	-	1	-	2	1	4
Black, both sexes	246	1	-	-	-	-	-	2	-	-	3	3
Male	87	-	-	-	-	-	-	2	-	-	2	2
Female	159	1	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
North Carolina												
All races, both sexes	1,493	-	2	-	1	-	1	2	3	2	7	10
Male	562	-	2	-	1	-	1	-	1	1	6	6
Female	931	-	-	-	-	-	-	2	2	1	1	4
White, both sexes	1,218	-	1	-	-	-	1	1	-	1	5	5
Male	454	-	1	-	-	-	1	-	-	1	4	3
Female	764	-	-	-	-	-	-	1	-	-	1	2
Black, both sexes	264	-	1	-	1	-	-	1	3	1	2	5
Male	103	-	1	-	1	-	-	-	1	-	2	3
Female	161	-	-	-	-	-	-	1	2	1	-	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	85	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	83	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	57	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Ohio												
All races, both sexes	2,075	-	1	-	-	-	1	1	-	1	3	5
Male	803	-	-	-	-	-	1	1	-	1	1	4
Female	1,272	-	1	-	-	-	-	-	-	-	2	1
White, both sexes	1,908	-	-	-	-	-	-	1	-	1	2	4
Male	741	-	-	-	-	-	-	1	-	1	1	3
Female	1,167	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	164	-	1	-	-	-	1	-	-	-	1	1
Male	61	-	-	-	-	-	1	-	-	-	-	1
Female	103	-	1	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,139	3	-	-	-	-	-	-	2	1	4	11
Male	419	2	-	-	-	-	-	-	2	1	2	3
Female	720	1	-	-	-	-	-	-	-	-	2	8
White, both sexes	1,036	1	-	-	-	-	-	-	2	1	2	5
Male	370	-	-	-	-	-	-	-	2	1	1	-
Female	666	1	-	-	-	-	-	-	-	-	1	5
Black, both sexes	57	2	-	-	-	-	-	-	-	-	1	4
Male	26	2	-	-	-	-	-	-	-	-	-	3
Female	31	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	44	-	-	-	-	-	-	-	-	-	1	2
Male	23	-	-	-	-	-	-	-	-	-	1	-
Female	21	-	-	-	-	-	-	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Oregon												
All races, both sexes	796	1	-	-	-	-	1	-	2	-	1	3
Male	308	-	-	-	-	-	1	-	1	-	1	1
Female	488	1	-	-	-	-	-	-	1	-	-	2
White, both sexes	781	1	-	-	-	-	1	-	2	-	1	2
Male	300	-	-	-	-	-	1	-	1	-	1	-
Female	481	1	-	-	-	-	-	-	1	-	-	2
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3,555	1	-	-	-	1	1	1	2	4	6	19
Male	1,361	1	-	-	-	1	-	-	2	4	3	14
Female	2,194	-	-	-	-	-	1	1	-	-	3	5
White, both sexes	3,318	-	-	-	-	1	-	1	2	3	4	13
Male	1,266	-	-	-	-	1	-	-	2	3	2	12
Female	2,052	-	-	-	-	-	-	1	-	-	2	1
Black, both sexes	229	-	-	-	-	-	1	-	-	1	2	6
Male	91	-	-	-	-	-	-	-	-	1	1	2
Female	138	-	-	-	-	-	1	-	-	-	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Rhode Island												
All races, both sexes	113	-	-	-	-	-	-	-	-	-	2	-
Male	39	-	-	-	-	-	-	-	-	-	2	-
Female	74	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	110	-	-	-	-	-	-	-	-	-	2	-
Male	36	-	-	-	-	-	-	-	-	-	2	-
Female	74	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1,055	-	-	1	-	-	1	-	3	5	7	8
Male	450	-	-	1	-	-	-	-	2	1	5	3
Female	605	-	-	-	-	-	1	-	1	4	2	5
White, both sexes	784	-	-	-	-	-	1	-	1	2	1	4
Male	333	-	-	-	-	-	-	-	-	1	-	1
Female	451	-	-	-	-	-	1	-	1	1	1	3
Black, both sexes	269	-	-	1	-	-	-	-	2	3	6	4
Male	117	-	-	1	-	-	-	-	2	-	5	2
Female	152	-	-	-	-	-	-	-	-	3	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
South Dakota												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	49	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	708	1	-	-	1	1	-	2	1	4	4	14
Male	281	1	-	-	-	-	-	2	-	2	3	9
Female	427	-	-	-	1	1	-	-	1	2	1	5
White, both sexes	609	1	-	-	1	1	-	-	-	4	2	10
Male	240	1	-	-	-	-	-	-	-	2	2	5
Female	369	-	-	-	1	1	-	-	-	2	-	5
Black, both sexes	95	-	-	-	-	-	-	2	-	-	2	4
Male	40	-	-	-	-	-	-	2	-	-	1	4
Female	55	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Texas												
All races, both sexes	3,822	1	-	-	-	-	1	2	9	7	24	40
Male	1,545	1	-	-	-	-	1	1	6	5	18	22
Female	2,277	-	-	-	-	-	-	1	3	2	6	18
White, both sexes	3,415	1	-	-	-	-	1	1	4	4	15	31
Male	1,382	1	-	-	-	-	1	1	3	3	11	16
Female	2,033	-	-	-	-	-	-	-	1	1	4	15
Black, both sexes	380	-	-	-	-	-	-	1	5	3	9	9
Male	153	-	-	-	-	-	-	-	3	2	7	6
Female	227	-	-	-	-	-	-	1	2	1	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	555	-	-	-	-	-	-	-	-	-	1	4
Male	214	-	-	-	-	-	-	-	-	-	1	3
Female	341	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	538	-	-	-	-	-	-	-	-	-	-	3
Male	207	-	-	-	-	-	-	-	-	-	-	2
Female	331	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Vermont												
All races, both sexes	83	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	57	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	83	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	57	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,610	2	-	-	-	-	-	-	2	7	8	10
Male	628	1	-	-	-	-	-	-	1	6	7	8
Female	982	1	-	-	-	-	-	-	1	1	1	2
White, both sexes	1,337	1	-	-	-	-	-	-	-	2	5	8
Male	529	-	-	-	-	-	-	-	-	2	4	6
Female	808	1	-	-	-	-	-	-	-	-	1	2
Black, both sexes	256	1	-	-	-	-	-	-	2	5	3	2
Male	93	1	-	-	-	-	-	-	1	4	3	2
Female	163	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Washington												
All races, both sexes	558	-	-	-	-	-	-	1	-	-	1	4
Male	220	-	-	-	-	-	-	1	-	-	-	-
Female	338	-	-	-	-	-	-	-	-	-	1	4
White, both sexes	526	-	-	-	-	-	-	1	-	-	1	2
Male	209	-	-	-	-	-	-	1	-	-	-	-
Female	317	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	557	-	-	-	-	-	-	-	-	-	4	6
Male	216	-	-	-	-	-	-	-	-	-	4	6
Female	341	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	541	-	-	-	-	-	-	-	-	-	4	5
Male	210	-	-	-	-	-	-	-	-	-	4	5
Female	331	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	16	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Heart failure (I50)												
Wisconsin												
All races, both sexes	1,438	-	-	-	-	-	1	-	-	-	1	1
Male	557	-	-	-	-	-	-	-	-	-	-	1
Female	881	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	1,401	-	-	-	-	-	1	-	-	-	1	-
Male	543	-	-	-	-	-	-	-	-	-	-	-
Female	858	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	119	-	-	-	-	-	-	-	-	-	1	1
Male	54	-	-	-	-	-	-	-	-	-	1	1
Female	65	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	117	-	-	-	-	-	-	-	-	-	1	1
Male	53	-	-	-	-	-	-	-	-	-	1	1
Female	64	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
United States												
All races, both sexes	703	1,078	1,532	2,176	3,712	6,580	10,751	13,180	11,799	5,264	1,278	-
Male	423	641	868	1,163	1,906	3,184	4,620	4,843	3,485	1,138	207	-
Female	280	437	664	1,013	1,806	3,396	6,131	8,337	8,314	4,126	1,071	-
White, both sexes	456	776	1,165	1,723	3,121	5,739	9,732	12,271	11,029	4,898	1,158	-
Male	281	465	670	929	1,617	2,802	4,245	4,563	3,283	1,059	187	-
Female	175	311	495	794	1,504	2,937	5,487	7,708	7,746	3,839	971	-
Black, both sexes	229	282	336	408	538	741	890	776	642	303	111	-
Male	133	164	180	204	261	340	319	231	155	52	17	-
Female	96	118	156	204	277	401	571	545	487	251	94	-
American Indian 1/, both sexes	9	10	13	19	25	35	37	24	26	16	2	-
Male	5	7	6	13	14	13	14	9	12	5	1	-
Female	4	3	7	6	11	22	23	15	14	11	1	-
Asian or Pacific Islander 2/, both sexes	9	10	18	26	28	65	92	109	102	47	7	-
Male	4	5	12	17	14	29	42	40	35	22	2	-
Female	5	5	6	9	14	36	50	69	67	25	5	-
Alabama												
All races, both sexes	64	72	91	100	173	274	405	421	376	139	40	-
Male	29	48	46	53	94	136	166	138	109	29	9	-
Female	35	24	45	47	79	138	239	283	267	110	31	-
White, both sexes	38	48	58	63	128	214	331	363	311	111	31	-
Male	17	35	32	32	71	110	136	117	92	23	6	-
Female	21	13	26	31	57	104	195	246	219	88	25	-
Black, both sexes	26	23	33	37	43	59	74	58	65	27	9	-
Male	12	12	14	21	23	25	30	21	17	5	3	-
Female	14	11	19	16	20	34	44	37	48	22	6	-
American Indian 1/, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Alaska												
All races, both sexes	3	1	4	4	4	14	9	9	8	4	1	-
Male	3	1	3	3	3	5	4	3	2	2	1	-
Female	-	-	1	1	1	9	5	6	6	2	-	-
White, both sexes	2	-	4	3	3	10	7	7	5	4	1	-
Male	2	-	3	3	3	4	3	1	1	2	1	-
Female	-	-	1	-	-	6	4	6	4	2	-	-
Black, both sexes	-	1	-	-	-	-	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	1	1	4	1	1	2	-	-	-
Male	1	-	-	-	-	1	1	1	1	-	-	-
Female	-	-	-	1	1	3	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Arizona												
All races, both sexes	4	10	12	27	43	79	151	176	146	64	9	-
Male	1	3	6	16	27	35	72	69	47	18	-	-
Female	3	7	6	11	16	44	79	107	99	46	9	-
White, both sexes	3	9	12	25	41	75	148	173	144	57	9	-
Male	1	3	6	15	25	33	70	67	45	17	-	-
Female	2	6	6	10	16	42	78	106	99	40	9	-
Black, both sexes	-	1	-	1	-	-	-	1	-	2	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	1	-	-	-	2	3	2	-	1	4	-	-
Male	-	-	-	-	2	1	2	-	1	1	-	-
Female	1	-	-	-	-	2	-	-	-	3	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	1	2	1	1	-	-
Male	-	-	-	1	-	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Arkansas												
All races, both sexes	12	15	18	34	52	80	141	148	147	55	7	-
Male	9	9	13	19	29	44	61	55	51	12	1	-
Female	3	6	5	15	23	36	80	93	96	43	6	-
White, both sexes	8	14	17	26	45	73	133	142	130	51	6	-
Male	5	9	13	14	26	41	59	53	46	11	1	-
Female	3	5	4	12	19	32	74	89	84	40	5	-
Black, both sexes	4	1	1	8	6	7	8	6	17	4	1	-
Male	4	-	-	5	2	3	2	2	5	1	-	-
Female	-	1	1	3	4	4	6	4	12	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	42	68	92	177	254	430	817	978	891	424	109	-
Male	28	40	57	96	138	206	362	367	304	106	17	-
Female	14	28	35	81	116	224	455	611	587	318	92	-
White, both sexes	31	47	72	146	211	370	733	903	825	397	104	-
Male	22	25	45	81	119	182	326	344	280	99	16	-
Female	9	22	27	65	92	188	407	559	545	298	88	-
Black, both sexes	8	17	12	23	35	34	43	35	26	13	2	-
Male	5	12	7	9	17	13	22	10	9	1	-	-
Female	3	5	5	14	18	21	21	25	17	12	2	-
American Indian 1/, both sexes	1	-	1	2	-	2	5	2	1	-	-	-
Male	-	-	-	2	-	1	3	-	-	-	-	-
Female	1	-	1	-	-	1	2	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	7	6	8	24	36	38	39	14	3	-
Male	1	3	5	4	2	10	11	13	15	6	1	-
Female	1	1	2	2	6	14	25	25	24	8	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Colorado												
All races, both sexes	9	7	20	18	38	63	127	175	133	82	21	-
Male	6	3	9	9	18	26	54	59	48	15	6	-
Female	3	4	11	9	20	37	73	116	85	67	15	-
White, both sexes	8	7	19	18	36	59	124	171	131	79	20	-
Male	5	3	8	9	16	24	53	58	47	15	5	-
Female	3	4	11	9	20	35	71	113	84	64	15	-
Black, both sexes	1	-	-	-	1	3	2	3	2	2	1	-
Male	1	-	-	-	1	2	-	-	1	-	1	-
Female	-	-	-	-	-	1	2	3	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	1	1	-	-	-	-
Male	-	-	1	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	5	5	16	13	30	70	124	160	170	75	21	-
Male	5	1	8	5	15	37	57	61	56	10	2	-
Female	-	4	8	8	15	33	67	99	114	65	19	-
White, both sexes	2	4	10	10	24	63	118	152	165	71	19	-
Male	2	-	5	5	10	32	54	60	54	9	2	-
Female	-	4	5	5	14	31	64	92	111	62	17	-
Black, both sexes	3	1	6	3	6	5	6	5	5	3	2	-
Male	3	1	3	-	5	4	3	1	2	-	-	-
Female	-	-	3	3	1	1	3	4	3	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	2	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Delaware												
All races, both sexes	2	3	2	12	12	12	28	33	18	11	3	-
Male	1	3	1	7	6	7	13	9	6	-	1	-
Female	1	-	1	5	6	5	15	24	12	11	2	-
White, both sexes	2	3	1	7	8	9	25	29	17	10	3	-
Male	1	3	1	6	3	5	13	9	6	-	1	-
Female	1	-	-	1	5	4	12	20	11	10	2	-
Black, both sexes	-	-	1	4	3	3	2	2	-	1	-	-
Male	-	-	-	-	2	2	-	-	-	-	-	-
Female	-	-	1	4	1	1	2	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	1	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	6	2	4	8	15	9	3	2	-
Male	1	-	-	4	2	1	1	7	-	-	-	-
Female	1	-	-	2	-	3	7	8	9	3	2	-
White, both sexes	-	-	-	-	1	1	1	5	2	-	1	-
Male	-	-	-	-	1	1	-	4	-	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	1	-
Black, both sexes	2	-	-	6	1	3	7	10	7	3	1	-
Male	1	-	-	4	1	-	1	3	-	-	-	-
Female	1	-	-	2	-	3	6	7	7	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Florida												
All races, both sexes	12	24	42	65	93	202	283	498	434	188	40	-
Male	5	14	22	41	51	99	121	216	147	59	7	-
Female	7	10	20	24	42	103	162	282	287	129	33	-
White, both sexes	8	21	31	55	76	185	260	472	412	182	36	-
Male	3	12	17	36	42	91	108	211	140	56	7	-
Female	5	9	14	19	34	94	152	261	272	126	29	-
Black, both sexes	4	3	11	10	16	17	21	23	18	6	4	-
Male	2	2	5	5	8	8	11	5	5	3	-	-
Female	2	1	6	5	8	9	10	18	13	3	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	2	3	3	-	-	-
Male	-	-	-	-	1	-	2	-	2	-	-	-
Female	-	-	-	-	-	-	-	3	1	-	-	-
Georgia												
All races, both sexes	45	56	89	125	164	248	376	415	339	117	37	-
Male	23	32	57	69	77	123	138	144	78	26	8	-
Female	22	24	32	56	87	125	238	271	261	91	29	-
White, both sexes	21	34	47	73	120	199	316	349	298	88	28	-
Male	10	20	25	38	54	101	120	127	70	18	6	-
Female	11	14	22	35	66	98	196	222	228	70	22	-
Black, both sexes	24	21	42	51	43	48	59	65	38	29	9	-
Male	13	11	32	31	23	21	17	16	6	8	2	-
Female	11	10	10	20	20	27	42	49	32	21	7	-
American Indian 1/, both sexes	-	1	-	1	1	-	-	-	3	-	-	-
Male	-	1	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	1	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Hawaii												
All races, both sexes	3	4	4	8	6	17	26	36	27	19	4	-
Male	3	4	4	7	4	8	17	17	9	10	2	-
Female	-	-	-	1	2	9	9	19	18	9	2	-
White, both sexes	1	2	2	2	1	10	8	13	9	5	1	-
Male	1	2	2	2	1	4	3	7	4	1	1	-
Female	-	-	-	-	-	6	5	6	5	4	-	-
Black, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	6	5	6	18	22	18	14	3	-
Male	1	1	2	5	3	3	14	9	5	9	1	-
Female	-	-	-	1	2	3	4	13	13	5	2	-
Idaho												
All races, both sexes	-	-	5	8	10	29	48	64	57	23	6	-
Male	-	-	4	5	5	14	25	20	25	1	3	-
Female	-	-	1	3	5	15	23	44	32	22	3	-
White, both sexes	-	-	5	8	10	27	48	64	57	23	6	-
Male	-	-	4	5	5	13	25	20	25	1	3	-
Female	-	-	1	3	5	14	23	44	32	22	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Illinois												
All races, both sexes	30	55	64	133	201	334	533	681	662	321	70	-
Male	16	31	35	69	112	166	225	254	185	58	7	-
Female	14	24	29	64	89	168	308	427	477	263	63	-
White, both sexes	20	39	49	98	174	293	480	641	628	300	64	-
Male	12	23	28	52	97	148	210	236	174	52	7	-
Female	8	16	21	46	77	145	270	405	454	248	57	-
Black, both sexes	10	16	12	33	25	39	51	34	28	18	6	-
Male	4	8	6	17	13	17	15	15	9	5	-	-
Female	6	8	6	16	12	22	36	19	19	13	6	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	2	2	1	6	6	3	-	-
Male	-	-	1	-	2	1	-	3	2	1	-	-
Female	-	-	1	2	-	1	1	3	4	2	-	-
Indiana												
All races, both sexes	22	43	58	68	143	219	344	458	400	144	40	-
Male	12	27	33	39	62	87	147	158	118	29	6	-
Female	10	16	25	29	81	132	197	300	282	115	34	-
White, both sexes	19	37	53	58	133	210	318	439	383	141	38	-
Male	10	24	29	33	59	85	142	152	113	27	6	-
Female	9	13	24	25	74	125	176	287	270	114	32	-
Black, both sexes	3	6	5	9	10	9	25	18	17	2	2	-
Male	2	3	4	6	3	2	5	6	5	1	-	-
Female	1	3	1	3	7	7	20	12	12	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Iowa												
All races, both sexes	5	2	7	3	13	28	57	59	76	39	12	-
Male	3	1	4	2	8	15	28	17	21	6	2	-
Female	2	1	3	1	5	13	29	42	55	33	10	-
White, both sexes	5	2	7	3	12	25	56	59	76	39	12	-
Male	3	1	4	2	7	13	27	17	21	6	2	-
Female	2	1	3	1	5	12	29	42	55	33	10	-
Black, both sexes	-	-	-	-	1	3	1	-	-	-	-	-
Male	-	-	-	-	1	2	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	6	13	13	25	51	80	172	198	210	107	27	-
Male	4	8	7	12	22	35	76	74	59	28	5	-
Female	2	5	6	13	29	45	96	124	151	79	22	-
White, both sexes	5	13	10	23	46	68	163	188	207	104	26	-
Male	3	8	5	11	20	29	76	71	58	28	5	-
Female	2	5	5	12	26	39	87	117	149	76	21	-
Black, both sexes	1	-	3	1	4	9	8	8	3	3	-	-
Male	1	-	2	-	1	4	-	3	1	-	-	-
Female	-	-	1	1	3	5	8	5	2	3	-	-
American Indian 1/, both sexes	-	-	-	1	1	3	-	2	-	-	1	-
Male	-	-	-	1	1	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Kentucky												
All races, both sexes	19	33	43	56	124	170	259	307	246	97	22	-
Male	12	23	33	26	70	85	93	103	65	15	2	-
Female	7	10	10	30	54	85	166	204	181	82	20	-
White, both sexes	16	29	40	52	110	156	245	295	241	93	21	-
Male	10	20	32	25	65	76	88	100	63	15	2	-
Female	6	9	8	27	45	80	157	195	178	78	19	-
Black, both sexes	3	3	3	3	14	13	14	12	4	4	1	-
Male	2	2	1	-	5	8	5	3	2	-	-	-
Female	1	1	2	3	9	5	9	9	2	4	1	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Louisiana												
All races, both sexes	22	32	62	82	110	196	261	326	246	116	24	-
Male	11	20	36	42	57	102	111	118	67	24	2	-
Female	11	12	26	40	53	94	150	208	179	92	22	-
White, both sexes	9	14	38	51	70	146	205	273	202	95	19	-
Male	3	10	24	27	31	86	90	104	51	18	1	-
Female	6	4	14	24	39	60	115	169	151	77	18	-
Black, both sexes	13	18	24	29	39	50	54	52	43	20	5	-
Male	8	10	12	14	26	16	21	13	16	6	1	-
Female	5	8	12	15	13	34	33	39	27	14	4	-
American Indian 1/, both sexes	-	-	-	2	1	-	1	-	1	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Maine												
All races, both sexes	2	4	6	6	10	32	42	64	60	37	4	-
Male	-	2	2	4	5	13	19	22	17	6	-	-
Female	2	2	4	2	5	19	23	42	43	31	4	-
White, both sexes	2	4	6	6	10	31	42	64	60	37	4	-
Male	-	2	2	4	5	13	19	22	17	6	-	-
Female	2	2	4	2	5	18	23	42	43	31	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	11	18	22	23	53	95	143	151	140	49	19	-
Male	9	12	13	12	24	45	57	55	44	16	4	-
Female	2	6	9	11	29	50	86	96	96	33	15	-
White, both sexes	6	16	14	20	39	64	112	136	121	44	14	-
Male	5	11	7	10	17	29	45	53	37	16	4	-
Female	1	5	7	10	22	35	67	83	84	28	10	-
Black, both sexes	5	2	8	3	14	31	29	13	18	5	5	-
Male	4	1	6	2	7	16	10	2	7	-	-	-
Female	1	1	2	1	7	15	19	11	11	5	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	2	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Massachusetts												
All races, both sexes	10	15	15	57	86	198	340	468	431	217	60	-
Male	7	6	6	33	41	102	139	174	106	44	15	-
Female	3	9	9	24	45	96	201	294	325	173	45	-
White, both sexes	9	13	13	53	79	187	328	456	419	216	58	-
Male	7	5	5	30	39	98	135	169	104	44	15	-
Female	2	8	8	23	40	89	193	287	315	172	43	-
Black, both sexes	1	2	2	3	7	9	10	10	10	1	2	-
Male	-	1	1	2	2	3	2	3	1	-	-	-
Female	1	1	1	1	5	6	8	7	9	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	2	2	1	-	-	-
Male	-	-	-	1	-	1	2	2	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Michigan												
All races, both sexes	23	38	50	68	135	298	490	571	510	224	56	-
Male	15	16	32	28	66	133	214	219	155	49	7	-
Female	8	22	18	40	69	165	276	352	355	175	49	-
White, both sexes	16	23	38	55	116	262	449	536	488	214	54	-
Male	11	9	22	23	57	117	195	207	151	48	7	-
Female	5	14	16	32	59	145	254	329	337	166	47	-
Black, both sexes	6	12	12	13	18	33	39	30	19	9	2	-
Male	4	6	10	5	9	15	17	12	3	1	-	-
Female	2	6	2	8	9	18	22	18	16	8	2	-
American Indian 1/, both sexes	-	3	-	-	-	1	1	2	1	-	-	-
Male	-	1	-	-	-	-	1	-	1	-	-	-
Female	-	2	-	-	-	1	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	2	1	3	2	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	1	1	-	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Minnesota												
All races, both sexes	8	9	17	30	52	112	178	287	283	140	23	-
Male	7	5	13	22	30	69	77	111	96	26	2	-
Female	1	4	4	8	22	43	101	176	187	114	21	-
White, both sexes	7	6	15	28	48	111	177	284	281	139	23	-
Male	6	3	12	21	27	68	77	110	96	26	2	-
Female	1	3	3	7	21	43	100	174	185	113	21	-
Black, both sexes	-	2	-	1	2	1	-	1	1	1	-	-
Male	-	1	-	-	2	1	-	1	-	-	-	-
Female	-	1	-	1	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	1	1	2	1	2	-	-	1	1	-	-	-
Male	1	1	1	1	1	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Mississippi												
All races, both sexes	27	37	69	67	88	181	230	298	250	109	33	-
Male	16	22	36	31	43	90	86	96	68	15	5	-
Female	11	15	33	36	45	91	144	202	182	94	28	-
White, both sexes	12	22	38	42	60	131	172	248	189	88	17	-
Male	6	14	22	19	30	64	67	80	49	10	1	-
Female	6	8	16	23	30	67	105	168	140	78	16	-
Black, both sexes	15	15	30	25	27	50	58	50	61	21	16	-
Male	10	8	14	12	12	26	19	16	19	5	4	-
Female	5	7	16	13	15	24	39	34	42	16	12	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Missouri												
All races, both sexes	16	23	43	52	81	182	286	318	302	138	30	-
Male	10	13	19	25	38	94	114	104	86	24	7	-
Female	6	10	24	27	43	88	172	214	216	114	23	-
White, both sexes	12	18	37	50	73	162	273	306	287	132	29	-
Male	7	11	16	24	36	85	107	100	79	23	7	-
Female	5	7	21	26	37	77	166	206	208	109	22	-
Black, both sexes	4	4	6	2	8	20	12	12	13	6	1	-
Male	3	2	3	1	2	9	6	4	5	1	-	-
Female	1	2	3	1	6	11	6	8	8	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	3	9	7	14	15	29	52	56	70	21	9	-
Male	2	8	5	9	10	10	32	20	24	5	2	-
Female	1	1	2	5	5	19	20	36	46	16	7	-
White, both sexes	2	7	6	13	15	27	49	56	68	21	9	-
Male	1	7	4	8	10	10	31	20	23	5	2	-
Female	1	-	2	5	5	17	18	36	45	16	7	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	2	1	1	-	2	2	-	1	-	-	-
Male	1	1	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	2	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Nebraska												
All races, both sexes	10	6	8	15	38	55	91	124	120	57	25	-
Male	8	5	5	6	19	31	33	43	36	13	4	-
Female	2	1	3	9	19	24	58	81	84	44	21	-
White, both sexes	9	6	8	15	36	52	88	122	118	57	25	-
Male	7	5	5	6	19	29	33	41	36	13	4	-
Female	2	1	3	9	17	23	55	81	82	44	21	-
Black, both sexes	-	-	-	-	2	2	3	1	2	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	2	1	3	-	2	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	19	18	33	48	78	122	99	92	32	4	-
Male	3	12	7	18	24	44	64	41	34	8	1	-
Female	6	7	11	15	24	34	58	58	58	24	3	-
White, both sexes	6	15	15	25	45	72	120	96	86	30	3	-
Male	3	10	5	13	23	42	63	41	31	8	1	-
Female	3	5	10	12	22	30	57	55	55	22	2	-
Black, both sexes	2	2	2	4	3	3	1	2	2	2	1	-
Male	-	1	2	2	1	2	-	-	1	-	-	-
Female	2	1	-	2	2	1	1	2	1	2	1	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	1	2	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	1	2	-	3	1	-	2	-	-	-
Male	-	1	-	2	-	-	1	-	1	-	-	-
Female	1	1	1	-	-	3	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
New Hampshire												
All races, both sexes	1	-	4	1	7	15	34	36	51	28	8	-
Male	-	-	3	-	6	8	14	11	18	4	-	-
Female	1	-	1	1	1	7	20	25	33	24	8	-
White, both sexes	1	-	4	1	7	15	34	36	51	28	8	-
Male	-	-	3	-	6	8	14	11	18	4	-	-
Female	1	-	1	1	1	7	20	25	33	24	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	13	15	39	39	98	165	307	382	321	161	22	-
Male	10	9	20	26	48	94	127	132	93	38	2	-
Female	3	6	19	13	50	71	180	250	228	123	20	-
White, both sexes	8	10	31	31	85	137	273	357	297	144	21	-
Male	7	6	18	20	42	79	117	125	90	36	2	-
Female	1	4	13	11	43	58	156	232	207	108	19	-
Black, both sexes	5	5	8	7	10	25	32	23	18	16	1	-
Male	3	3	2	5	4	13	9	7	3	2	-	-
Female	2	2	6	2	6	12	23	16	15	14	1	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	3	2	2	6	1	-	-
Male	-	-	-	-	1	2	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	2	6	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
New Mexico												
All races, both sexes	3	3	6	18	22	52	69	80	70	49	9	-
Male	1	2	3	10	8	21	36	32	19	15	2	-
Female	2	1	3	8	14	31	33	48	51	34	7	-
White, both sexes	2	3	5	16	19	48	62	77	67	45	9	-
Male	1	2	3	8	6	20	33	32	19	14	2	-
Female	1	1	2	8	13	28	29	45	48	31	7	-
Black, both sexes	1	-	1	1	2	-	1	1	1	1	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	1	-	1	-	1	-	-	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	1	1	4	6	2	2	3	-	-
Male	-	-	-	1	1	1	2	-	-	1	-	-
Female	-	-	-	-	-	3	4	2	2	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	26	44	57	72	157	307	521	674	594	271	76	-
Male	14	22	29	33	88	141	236	286	178	58	11	-
Female	12	22	28	39	69	166	285	388	416	213	65	-
White, both sexes	15	32	39	60	133	271	482	627	560	256	67	-
Male	7	15	20	25	76	131	223	269	173	57	11	-
Female	8	17	19	35	57	140	259	358	387	199	56	-
Black, both sexes	11	12	16	11	24	34	38	37	32	14	8	-
Male	7	7	8	7	12	10	13	12	4	1	-	-
Female	4	5	8	4	12	24	25	25	28	13	8	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	2	1	9	2	-	1	-
Male	-	-	1	1	-	-	-	4	1	-	-	-
Female	-	-	-	-	-	2	1	5	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
North Carolina												
All races, both sexes	22	26	39	68	116	166	304	324	264	111	25	-
Male	13	11	21	36	58	83	133	103	61	19	6	-
Female	9	15	18	32	58	83	171	221	203	92	19	-
White, both sexes	10	14	25	49	86	130	247	283	238	98	24	-
Male	8	4	13	24	44	65	114	96	54	16	6	-
Female	2	10	12	25	42	65	133	187	184	82	18	-
Black, both sexes	12	12	13	17	30	35	53	40	25	12	1	-
Male	5	7	8	11	14	18	17	7	6	2	-	-
Female	7	5	5	6	16	17	36	33	19	10	1	-
American Indian 1/, both sexes	-	-	1	-	-	1	3	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	1	-	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	-	1	1	-	1	-	-
Male	-	-	-	1	-	-	1	-	-	1	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	3	3	9	26	21	17	6	-
Male	-	-	-	-	2	2	3	7	7	5	1	-
Female	-	-	-	-	1	1	6	19	14	12	5	-
White, both sexes	-	-	-	-	3	3	9	25	20	17	6	-
Male	-	-	-	-	2	2	3	7	6	5	1	-
Female	-	-	-	-	1	1	6	18	14	12	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Ohio												
All races, both sexes	16	31	53	81	153	257	401	492	388	153	38	-
Male	11	20	26	38	70	121	169	189	108	33	10	-
Female	5	11	27	43	83	136	232	303	280	120	28	-
White, both sexes	13	22	48	64	138	234	372	466	369	144	30	-
Male	9	16	25	34	64	111	158	177	104	31	6	-
Female	4	6	23	30	74	123	214	289	265	113	24	-
Black, both sexes	2	9	5	17	15	21	29	26	19	9	8	-
Male	1	4	1	4	6	10	11	12	4	2	4	-
Female	1	5	4	13	9	11	18	14	15	7	4	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Oklahoma												
All races, both sexes	12	27	44	40	80	139	223	214	222	93	24	-
Male	7	16	26	24	33	56	99	67	53	21	7	-
Female	5	11	18	16	47	83	124	147	169	72	17	-
White, both sexes	9	18	39	32	68	128	206	201	216	85	23	-
Male	6	10	22	16	28	53	96	58	52	18	7	-
Female	3	8	17	16	40	75	110	143	164	67	16	-
Black, both sexes	2	8	2	5	5	5	7	7	5	3	1	-
Male	1	6	2	5	2	1	-	4	-	-	-	-
Female	1	2	-	-	3	4	7	3	5	3	1	-
American Indian 1/, both sexes	1	-	3	3	7	6	9	6	1	5	-	-
Male	-	-	2	3	3	2	3	5	1	3	-	-
Female	1	-	1	-	4	4	6	1	-	2	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Oregon												
All races, both sexes	3	9	13	17	44	70	119	202	189	90	32	-
Male	3	4	7	8	23	38	50	77	68	20	6	-
Female	-	5	6	9	21	32	69	125	121	70	26	-
White, both sexes	2	9	13	17	44	69	118	198	184	89	31	-
Male	2	4	7	8	23	37	50	75	66	20	5	-
Female	-	5	6	9	21	32	68	123	118	69	26	-
Black, both sexes	-	-	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	1	-	1	-
Male	1	-	-	-	-	1	-	-	1	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	3	1	-	-
Male	-	-	-	-	-	-	-	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	2	1	-	-
Pennsylvania												
All races, both sexes	39	51	72	90	203	384	673	870	736	333	69	-
Male	28	26	46	52	105	184	297	321	198	72	7	-
Female	11	25	26	38	98	200	376	549	538	261	62	-
White, both sexes	27	36	56	76	183	352	634	836	706	323	65	-
Male	21	17	33	45	97	164	281	314	195	72	7	-
Female	6	19	23	31	86	188	353	522	511	251	58	-
Black, both sexes	12	15	16	14	20	30	37	34	28	9	4	-
Male	7	9	13	7	8	18	15	7	3	-	-	-
Female	5	6	3	7	12	12	22	27	25	9	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	2	-	2	1	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Rhode Island												
All races, both sexes	1	1	1	2	4	24	19	30	20	7	2	-
Male	1	-	-	2	2	10	8	9	4	1	-	-
Female	-	1	1	-	2	14	11	21	16	6	2	-
White, both sexes	-	1	1	1	4	23	19	30	20	7	2	-
Male	-	-	-	1	2	9	8	9	4	1	-	-
Female	-	1	1	-	2	14	11	21	16	6	2	-
Black, both sexes	1	-	-	1	-	1	-	-	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	31	38	41	52	89	151	176	215	154	68	15	-
Male	20	26	23	26	52	74	87	72	45	10	3	-
Female	11	12	18	26	37	77	89	143	109	58	12	-
White, both sexes	21	19	23	35	59	114	140	176	129	50	9	-
Male	15	12	14	18	34	53	69	63	43	8	2	-
Female	6	7	9	17	25	61	71	113	86	42	7	-
Black, both sexes	10	19	18	17	30	37	36	38	24	18	6	-
Male	5	14	9	8	18	21	18	9	2	2	1	-
Female	5	5	9	9	12	16	18	29	22	16	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
South Dakota												
All races, both sexes	1	-	-	-	3	3	4	15	12	9	6	-
Male	-	-	-	-	2	3	3	7	2	1	-	-
Female	1	-	-	-	1	-	1	8	10	8	6	-
White, both sexes	1	-	-	-	2	2	2	15	12	9	6	-
Male	-	-	-	-	1	2	2	7	2	1	-	-
Female	1	-	-	-	1	-	-	8	10	8	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	11	22	17	36	47	76	135	139	132	52	13	-
Male	7	14	10	22	20	27	61	55	32	14	2	-
Female	4	8	7	14	27	49	74	84	100	38	11	-
White, both sexes	7	15	14	30	40	62	118	125	121	47	11	-
Male	3	11	8	20	19	22	52	51	29	13	2	-
Female	4	4	6	10	21	40	66	74	92	34	9	-
Black, both sexes	4	7	3	6	7	12	17	14	10	5	2	-
Male	4	3	2	2	1	4	9	4	3	1	-	-
Female	-	4	1	4	6	8	8	10	7	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Texas												
All races, both sexes	62	98	146	159	268	449	695	767	714	318	62	-
Male	38	64	80	90	146	213	288	269	221	75	7	-
Female	24	34	66	69	122	236	407	498	493	243	55	-
White, both sexes	39	70	115	135	230	399	642	710	672	291	55	-
Male	25	46	69	79	128	186	272	253	212	70	6	-
Female	14	24	46	56	102	213	370	457	460	221	49	-
Black, both sexes	23	28	29	23	35	46	49	53	39	21	7	-
Male	13	18	10	10	15	26	16	15	8	3	1	-
Female	10	10	19	13	20	20	33	38	31	18	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	3	4	3	4	3	6	-	-
Male	-	-	1	1	3	1	-	1	1	2	-	-
Female	-	-	1	-	-	3	3	3	2	4	-	-
Utah												
All races, both sexes	6	9	10	13	30	50	99	143	131	55	4	-
Male	4	6	7	7	11	24	41	52	41	16	1	-
Female	2	3	3	6	19	26	58	91	90	39	3	-
White, both sexes	5	8	9	13	26	49	98	142	128	53	4	-
Male	3	6	6	7	10	24	41	52	39	16	1	-
Female	2	2	3	6	16	25	57	90	89	37	3	-
Black, both sexes	1	1	-	-	1	-	1	1	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	1	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	1	-	-	2	-	-	-
Male	-	-	1	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Vermont												
All races, both sexes	-	1	2	2	1	8	12	26	18	12	1	-
Male	-	1	1	2	1	4	3	8	5	1	-	-
Female	-	-	1	-	-	4	9	18	13	11	1	-
White, both sexes	-	1	2	2	1	8	12	26	18	12	1	-
Male	-	1	1	2	1	4	3	8	5	1	-	-
Female	-	-	1	-	-	4	9	18	13	11	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	14	38	47	71	109	191	311	327	305	136	32	-
Male	8	24	26	32	58	91	125	118	87	31	5	-
Female	6	14	21	39	51	100	186	209	218	105	27	-
White, both sexes	5	25	37	52	81	153	258	283	273	124	30	-
Male	3	17	23	26	45	76	108	104	81	29	5	-
Female	2	8	14	26	36	77	150	179	192	95	25	-
Black, both sexes	7	12	10	17	27	36	49	42	29	12	2	-
Male	4	7	3	6	13	13	16	13	5	2	-	-
Female	3	5	7	11	14	23	33	29	24	10	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	2	1	2	4	2	3	-	-	-
Male	1	-	-	-	-	2	1	1	1	-	-	-
Female	1	1	-	2	1	-	3	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Washington												
All races, both sexes	7	7	9	15	35	54	111	124	120	50	20	-
Male	3	5	7	9	21	25	48	55	34	11	1	-
Female	4	2	2	6	14	29	63	69	86	39	19	-
White, both sexes	6	7	9	12	33	50	100	120	116	49	20	-
Male	3	5	7	8	21	22	44	53	33	11	1	-
Female	3	2	2	4	12	28	56	67	83	38	19	-
Black, both sexes	-	-	-	2	1	2	8	1	-	-	-	-
Male	-	-	-	1	-	2	4	1	-	-	-	-
Female	-	-	-	1	1	-	4	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	1	2	3	3	1	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	1	-	-	-	1	1	2	2	2	1	-	-
West Virginia												
All races, both sexes	4	13	16	19	54	70	111	108	96	42	14	-
Male	2	6	7	8	26	39	47	39	19	10	3	-
Female	2	7	9	11	28	31	64	69	77	32	11	-
White, both sexes	3	13	15	19	53	67	110	106	90	42	14	-
Male	2	6	7	8	26	37	46	39	17	10	3	-
Female	1	7	8	11	27	30	64	67	73	32	11	-
Black, both sexes	1	-	1	-	1	3	1	2	6	-	-	-
Male	-	-	-	-	-	2	1	-	2	-	-	-
Female	1	-	1	-	1	1	-	2	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Heart failure (I50)												
Wisconsin												
All races, both sexes	5	20	16	47	56	128	252	367	365	143	36	-
Male	4	14	8	23	26	60	122	154	116	25	4	-
Female	1	6	8	24	30	68	130	213	249	118	32	-
White, both sexes	3	16	13	45	52	124	244	364	360	143	35	-
Male	3	10	7	23	24	60	119	153	115	25	4	-
Female	-	6	6	22	28	64	125	211	245	118	31	-
Black, both sexes	1	3	1	1	2	2	4	3	4	-	1	-
Male	1	3	-	-	1	-	2	1	1	-	-	-
Female	-	-	1	1	1	2	2	2	3	-	1	-
American Indian 1/, both sexes	1	1	1	1	2	1	1	-	1	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	1	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	3	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	1	-	-	1	2	-	-	-	-	-
Wyoming												
All races, both sexes	-	4	3	5	4	9	23	26	23	14	6	-
Male	-	2	2	3	-	4	17	6	13	4	1	-
Female	-	2	1	2	4	5	6	20	10	10	5	-
White, both sexes	-	4	2	5	4	9	23	26	22	14	6	-
Male	-	2	1	3	-	4	17	6	13	4	1	-
Female	-	2	1	2	4	5	6	20	9	10	5	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
United States												
All races, both sexes	54,881	7	4	2	3	7	22	24	46	75	184	304
Male	21,244	4	3	2	-	6	13	16	31	47	121	185
Female	33,637	3	1	-	3	1	9	8	15	28	63	119
White, both sexes	49,258	5	3	1	2	4	13	8	21	43	101	186
Male	18,968	2	3	1	-	3	10	6	14	27	62	112
Female	30,290	3	-	-	2	1	3	2	7	16	39	74
Black, both sexes	4,940	2	1	-	1	3	8	16	25	30	78	110
Male	1,971	2	-	-	-	3	3	10	17	19	54	67
Female	2,969	-	1	-	1	-	5	6	8	11	24	43
American Indian 1/, both sexes	209	-	-	-	-	-	1	-	-	1	2	5
Male	97	-	-	-	-	-	-	-	-	1	2	3
Female	112	-	-	-	-	-	1	-	-	-	-	2
Asian or Pacific Islander 2/, both sexes	474	-	-	1	-	-	-	-	-	1	3	3
Male	208	-	-	1	-	-	-	-	-	-	3	3
Female	266	-	-	-	-	-	-	-	-	1	-	-
Alabama												
All races, both sexes	1,899	-	-	-	-	1	2	2	6	7	16	17
Male	751	-	-	-	-	-	1	1	5	6	12	9
Female	1,148	-	-	-	-	1	1	1	1	1	4	8
White, both sexes	1,525	-	-	-	-	1	-	-	1	4	6	12
Male	592	-	-	-	-	-	-	-	1	3	5	6
Female	933	-	-	-	-	1	-	-	-	1	1	6
Black, both sexes	370	-	-	-	-	-	2	2	5	3	10	5
Male	156	-	-	-	-	-	1	1	4	3	7	3
Female	214	-	-	-	-	-	1	1	1	-	3	2
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Alaska												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	-	-
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	-	-	-	-
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	702	-	-	-	-	-	1	-	-	1	3	6
Male	285	-	-	-	-	-	-	-	-	-	1	4
Female	417	-	-	-	-	-	1	-	-	1	2	2
White, both sexes	675	-	-	-	-	-	-	-	-	1	3	4
Male	273	-	-	-	-	-	-	-	-	-	1	3
Female	402	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	14	-	-	-	-	-	1	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Arkansas												
All races, both sexes	688	-	-	-	-	1	1	-	1	1	5	2
Male	294	-	-	-	-	1	1	-	1	1	4	2
Female	394	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	621	-	-	-	-	-	-	-	1	-	4	2
Male	265	-	-	-	-	-	-	-	1	-	3	2
Female	356	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	66	-	-	-	-	1	1	-	-	1	1	-
Male	28	-	-	-	-	1	1	-	-	1	1	-
Female	38	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	4,272	1	-	1	-	-	2	-	3	6	17	21
Male	1,732	1	-	1	-	-	2	-	2	4	9	13
Female	2,540	-	-	-	-	-	-	-	1	2	8	8
White, both sexes	3,815	1	-	-	-	-	2	-	3	3	12	15
Male	1,542	1	-	-	-	-	2	-	2	2	5	11
Female	2,273	-	-	-	-	-	-	-	1	1	7	4
Black, both sexes	259	-	-	-	-	-	-	-	-	3	3	6
Male	110	-	-	-	-	-	-	-	-	2	2	2
Female	149	-	-	-	-	-	-	-	-	1	1	4
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	183	-	-	1	-	-	-	-	-	-	1	-
Male	73	-	-	1	-	-	-	-	-	-	1	-
Female	110	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Colorado												
All races, both sexes	644	-	-	-	-	-	-	-	-	1	1	1
Male	234	-	-	-	-	-	-	-	-	1	1	1
Female	410	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	624	-	-	-	-	-	-	-	-	-	1	-
Male	223	-	-	-	-	-	-	-	-	-	1	-
Female	401	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	627	-	1	-	-	-	-	-	-	1	1	4
Male	231	-	1	-	-	-	-	-	-	1	-	4
Female	396	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	580	-	1	-	-	-	-	-	-	1	1	2
Male	209	-	1	-	-	-	-	-	-	1	-	2
Female	371	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	41	-	-	-	-	-	-	-	-	-	-	2
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	21	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Delaware												
All races, both sexes	125	-	-	-	-	-	-	-	1	-	-	2
Male	52	-	-	-	-	-	-	-	1	-	-	2
Female	73	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	103	-	-	-	-	-	-	-	-	-	-	1
Male	45	-	-	-	-	-	-	-	-	-	-	1
Female	58	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	16	-	-	-	-	-	-	-	1	-	-	-
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	1	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	40	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Florida												
All races, both sexes	1,774	-	-	-	-	-	-	1	2	3	2	7
Male	737	-	-	-	-	-	-	-	2	1	1	3
Female	1,037	-	-	-	-	-	-	1	-	2	1	4
White, both sexes	1,637	-	-	-	-	-	-	1	-	1	1	4
Male	678	-	-	-	-	-	-	-	-	-	-	1
Female	959	-	-	-	-	-	-	1	-	1	1	3
Black, both sexes	129	-	-	-	-	-	-	-	2	2	1	3
Male	55	-	-	-	-	-	-	-	2	1	1	2
Female	74	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	1,810	1	-	-	-	1	1	1	-	7	19	21
Male	696	-	-	-	-	1	-	-	-	6	9	11
Female	1,114	1	-	-	-	-	1	1	-	1	10	10
White, both sexes	1,416	1	-	-	-	-	-	-	-	2	10	11
Male	531	-	-	-	-	-	-	-	-	1	5	6
Female	885	1	-	-	-	-	-	-	-	1	5	5
Black, both sexes	386	-	-	-	-	1	1	1	-	5	9	10
Male	159	-	-	-	-	1	-	-	-	5	4	5
Female	227	-	-	-	-	-	1	1	-	-	5	5
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Hawaii												
All races, both sexes	129	-	-	-	-	-	-	-	-	1	3	1
Male	71	-	-	-	-	-	-	-	-	-	3	-
Female	58	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	46	-	-	-	-	-	-	-	-	-	1	1
Male	22	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	79	-	-	-	-	-	-	-	-	1	2	-
Male	45	-	-	-	-	-	-	-	-	-	2	-
Female	34	-	-	-	-	-	-	-	-	1	-	-
Idaho												
All races, both sexes	244	-	-	-	1	-	-	-	1	-	-	-
Male	99	-	-	-	-	-	-	-	1	-	-	-
Female	145	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	242	-	-	-	1	-	-	-	1	-	-	-
Male	98	-	-	-	-	-	-	-	1	-	-	-
Female	144	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Illinois												
All races, both sexes	2,947	-	2	-	-	-	2	3	-	2	10	18
Male	1,123	-	1	-	-	-	1	3	-	1	8	14
Female	1,824	-	1	-	-	-	1	-	-	1	2	4
White, both sexes	2,664	-	1	-	-	-	2	1	-	1	6	13
Male	1,006	-	1	-	-	-	1	1	-	1	5	11
Female	1,658	-	-	-	-	-	1	-	-	-	1	2
Black, both sexes	257	-	1	-	-	-	-	2	-	1	4	4
Male	106	-	-	-	-	-	-	2	-	-	3	2
Female	151	-	1	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,863	-	-	-	-	1	1	2	2	3	3	7
Male	687	-	-	-	-	1	1	2	1	2	3	4
Female	1,176	-	-	-	-	-	-	-	1	1	-	3
White, both sexes	1,753	-	-	-	-	1	1	2	2	2	2	5
Male	649	-	-	-	-	1	1	2	1	2	2	3
Female	1,104	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	106	-	-	-	-	-	-	-	-	1	1	2
Male	37	-	-	-	-	-	-	-	-	-	1	1
Female	69	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Iowa												
All races, both sexes	282	-	-	-	-	-	-	-	-	-	-	1
Male	101	-	-	-	-	-	-	-	-	-	-	1
Female	181	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	277	-	-	-	-	-	-	-	-	-	-	1
Male	97	-	-	-	-	-	-	-	-	-	-	1
Female	180	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	884	-	-	-	-	-	-	-	3	1	3	2
Male	329	-	-	-	-	-	-	-	3	1	3	1
Female	555	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	832	-	-	-	-	-	-	-	3	-	3	-
Male	311	-	-	-	-	-	-	-	3	-	3	-
Female	521	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	42	-	-	-	-	-	-	-	-	1	-	1
Male	14	-	-	-	-	-	-	-	-	1	-	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Kentucky												
All races, both sexes	1,329	1	-	-	-	-	-	-	4	3	4	7
Male	516	1	-	-	-	-	-	-	2	-	2	5
Female	813	-	-	-	-	-	-	-	2	3	2	2
White, both sexes	1,250	-	-	-	-	-	-	-	4	3	4	5
Male	484	-	-	-	-	-	-	-	2	-	2	4
Female	766	-	-	-	-	-	-	-	2	3	2	1
Black, both sexes	76	1	-	-	-	-	-	-	-	-	-	2
Male	30	1	-	-	-	-	-	-	-	-	-	1
Female	46	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,441	1	-	-	-	-	1	2	2	2	8	12
Male	573	-	-	-	-	-	1	1	1	1	4	4
Female	868	1	-	-	-	-	-	1	1	1	4	8
White, both sexes	1,076	1	-	-	-	-	1	-	-	1	2	1
Male	423	-	-	-	-	-	1	-	-	1	-	-
Female	653	1	-	-	-	-	-	-	-	-	2	1
Black, both sexes	358	-	-	-	-	-	-	2	2	1	6	11
Male	148	-	-	-	-	-	-	1	1	-	4	4
Female	210	-	-	-	-	-	-	1	1	1	2	7
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Maine												
All races, both sexes	251	-	-	-	-	-	-	-	-	-	1	-
Male	85	-	-	-	-	-	-	-	-	-	-	-
Female	166	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	250	-	-	-	-	-	-	-	-	-	1	-
Male	85	-	-	-	-	-	-	-	-	-	-	-
Female	165	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	706	-	-	-	-	-	-	-	1	1	5	6
Male	283	-	-	-	-	-	-	-	-	-	3	4
Female	423	-	-	-	-	-	-	-	1	1	2	2
White, both sexes	568	-	-	-	-	-	-	-	-	-	1	3
Male	226	-	-	-	-	-	-	-	-	-	-	1
Female	342	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	133	-	-	-	-	-	-	-	1	1	4	3
Male	55	-	-	-	-	-	-	-	-	-	3	3
Female	78	-	-	-	-	-	-	-	1	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Massachusetts												
All races, both sexes	1,787	-	-	-	-	-	-	-	-	1	3	4
Male	631	-	-	-	-	-	-	-	-	1	2	4
Female	1,156	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1,726	-	-	-	-	-	-	-	-	1	2	4
Male	611	-	-	-	-	-	-	-	-	1	1	4
Female	1,115	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	53	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	39	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2,390	-	-	-	-	-	-	-	-	3	3	9
Male	903	-	-	-	-	-	-	-	-	2	2	5
Female	1,487	-	-	-	-	-	-	-	-	1	1	4
White, both sexes	2,179	-	-	-	-	-	-	-	-	2	1	1
Male	814	-	-	-	-	-	-	-	-	1	-	-
Female	1,365	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes	192	-	-	-	-	-	-	-	-	1	2	8
Male	84	-	-	-	-	-	-	-	-	1	2	5
Female	108	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Minnesota												
All races, both sexes	1,085	-	-	-	-	1	-	-	-	-	1	3
Male	431	-	-	-	-	1	-	-	-	-	1	1
Female	654	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	1,066	-	-	-	-	1	-	-	-	-	1	2
Male	420	-	-	-	-	1	-	-	-	-	1	-
Female	646	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,296	-	-	-	-	1	3	3	3	5	3	10
Male	477	-	-	-	-	1	1	1	2	2	3	6
Female	819	-	-	-	-	-	2	2	1	3	-	4
White, both sexes	952	-	-	-	-	-	-	-	1	2	-	5
Male	335	-	-	-	-	-	-	-	-	-	-	2
Female	617	-	-	-	-	-	-	-	1	2	-	3
Black, both sexes	342	-	-	-	-	1	3	3	2	3	3	5
Male	141	-	-	-	-	1	1	1	2	2	3	4
Female	201	-	-	-	-	-	2	2	-	1	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Missouri												
All races, both sexes	1,409	-	-	1	-	-	-	1	-	2	2	6
Male	516	-	-	1	-	-	-	1	-	1	2	4
Female	893	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	1,324	-	-	1	-	-	-	-	-	2	2	4
Male	478	-	-	1	-	-	-	-	-	1	2	3
Female	846	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	82	-	-	-	-	-	-	1	-	-	-	2
Male	35	-	-	-	-	-	-	1	-	-	-	1
Female	47	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	266	-	-	-	-	-	-	-	-	1	-	-
Male	120	-	-	-	-	-	-	-	-	1	-	-
Female	146	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	256	-	-	-	-	-	-	-	-	1	-	-
Male	115	-	-	-	-	-	-	-	-	1	-	-
Female	141	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Nebraska												
All races, both sexes	467	-	-	-	-	-	-	-	-	-	-	6
Male	161	-	-	-	-	-	-	-	-	-	-	3
Female	306	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	455	-	-	-	-	-	-	-	-	-	-	6
Male	157	-	-	-	-	-	-	-	-	-	-	3
Female	298	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	527	-	-	-	-	-	-	-	-	-	4	2
Male	250	-	-	-	-	-	-	-	-	-	2	1
Female	277	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	487	-	-	-	-	-	-	-	-	-	2	1
Male	232	-	-	-	-	-	-	-	-	-	-	-
Female	255	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	24	-	-	-	-	-	-	-	-	-	2	1
Male	11	-	-	-	-	-	-	-	-	-	2	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
New Hampshire												
All races, both sexes	174	-	-	-	-	-	-	-	-	-	1	-
Male	60	-	-	-	-	-	-	-	-	-	1	-
Female	114	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	174	-	-	-	-	-	-	-	-	-	1	-
Male	60	-	-	-	-	-	-	-	-	-	1	-
Female	114	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,474	-	-	-	1	-	1	1	2	1	4	12
Male	570	-	-	-	-	-	1	1	1	-	1	8
Female	904	-	-	-	1	-	-	-	1	1	3	4
White, both sexes	1,304	-	-	-	-	-	1	1	-	1	-	3
Male	510	-	-	-	-	-	1	1	-	-	-	2
Female	794	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	151	-	-	-	1	-	-	-	2	-	4	8
Male	53	-	-	-	-	-	-	-	1	-	1	5
Female	98	-	-	-	1	-	-	-	1	-	3	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
New Mexico												
All races, both sexes	372	1	-	-	-	-	-	-	-	-	-	2
Male	146	1	-	-	-	-	-	-	-	-	-	-
Female	226	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	345	1	-	-	-	-	-	-	-	-	-	1
Male	138	1	-	-	-	-	-	-	-	-	-	-
Female	207	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	2,630	-	-	-	-	-	1	3	-	-	4	11
Male	1,027	-	-	-	-	-	1	3	-	-	2	8
Female	1,603	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	2,387	-	-	-	-	-	1	1	-	-	1	9
Male	939	-	-	-	-	-	1	1	-	-	-	6
Female	1,448	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	224	-	-	-	-	-	-	2	-	-	3	2
Male	80	-	-	-	-	-	-	2	-	-	2	2
Female	144	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
North Carolina												
All races, both sexes	1,386	-	1	-	-	-	1	1	3	2	5	8
Male	519	-	1	-	-	-	1	-	1	1	4	6
Female	867	-	-	-	-	-	-	1	2	1	1	2
White, both sexes	1,144	-	1	-	-	-	1	1	-	1	4	3
Male	423	-	1	-	-	-	1	-	-	1	3	3
Female	721	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	233	-	-	-	-	-	-	-	3	1	1	5
Male	92	-	-	-	-	-	-	-	1	-	1	3
Female	141	-	-	-	-	-	-	-	2	1	-	2
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	82	-	-	-	-	-	-	-	-	-	-	-
Male	25	-	-	-	-	-	-	-	-	-	-	-
Female	57	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	80	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	56	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Ohio												
All races, both sexes	1,968	-	-	-	-	-	-	-	-	1	2	4
Male	753	-	-	-	-	-	-	-	-	1	-	3
Female	1,215	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	1,817	-	-	-	-	-	-	-	-	1	1	4
Male	696	-	-	-	-	-	-	-	-	1	-	3
Female	1,121	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	149	-	-	-	-	-	-	-	-	-	1	-
Male	57	-	-	-	-	-	-	-	-	-	-	-
Female	92	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,049	2	-	-	-	-	-	-	-	1	3	9
Male	387	1	-	-	-	-	-	-	-	1	2	3
Female	662	1	-	-	-	-	-	-	-	-	1	6
White, both sexes	957	1	-	-	-	-	-	-	-	1	1	5
Male	345	-	-	-	-	-	-	-	-	1	1	-
Female	612	1	-	-	-	-	-	-	-	-	-	5
Black, both sexes	50	1	-	-	-	-	-	-	-	-	1	3
Male	22	1	-	-	-	-	-	-	-	-	-	3
Female	28	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	40	-	-	-	-	-	-	-	-	-	1	1
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Oregon												
All races, both sexes	780	-	-	-	-	-	1	-	1	-	1	3
Male	299	-	-	-	-	-	1	-	-	-	1	1
Female	481	-	-	-	-	-	-	-	1	-	-	2
White, both sexes	766	-	-	-	-	-	1	-	1	-	1	2
Male	291	-	-	-	-	-	1	-	-	-	1	-
Female	475	-	-	-	-	-	-	-	1	-	-	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3,277	-	-	-	-	1	1	-	2	3	5	14
Male	1,230	-	-	-	-	1	-	-	2	3	3	10
Female	2,047	-	-	-	-	-	1	-	-	-	2	4
White, both sexes	3,073	-	-	-	-	1	-	-	2	2	3	9
Male	1,156	-	-	-	-	1	-	-	2	2	2	8
Female	1,917	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	197	-	-	-	-	-	1	-	-	1	2	5
Male	71	-	-	-	-	-	-	-	-	1	1	2
Female	126	-	-	-	-	-	1	-	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Rhode Island												
All races, both sexes	107	-	-	-	-	-	-	-	-	-	-	-
Male	35	-	-	-	-	-	-	-	-	-	-	-
Female	72	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	104	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	72	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	884	-	-	-	-	-	1	-	2	3	6	5
Male	363	-	-	-	-	-	-	-	1	-	4	2
Female	521	-	-	-	-	-	1	-	1	3	2	3
White, both sexes	666	-	-	-	-	-	1	-	1	1	1	3
Male	273	-	-	-	-	-	-	-	-	-	-	1
Female	393	-	-	-	-	-	1	-	1	1	1	2
Black, both sexes	216	-	-	-	-	-	-	-	1	2	5	2
Male	90	-	-	-	-	-	-	-	1	-	4	1
Female	126	-	-	-	-	-	-	-	-	2	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
South Dakota												
All races, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	45	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	656	-	-	-	1	-	-	2	-	4	3	11
Male	250	-	-	-	-	-	-	2	-	2	2	7
Female	406	-	-	-	1	-	-	-	-	2	1	4
White, both sexes	571	-	-	-	1	-	-	-	-	4	1	8
Male	217	-	-	-	-	-	-	-	-	2	1	4
Female	354	-	-	-	1	-	-	-	-	2	-	4
Black, both sexes	83	-	-	-	-	-	-	2	-	-	2	3
Male	32	-	-	-	-	-	-	2	-	-	1	3
Female	51	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Texas												
All races, both sexes	3,527	-	-	-	-	-	1	2	6	5	18	29
Male	1,405	-	-	-	-	-	1	1	4	4	14	15
Female	2,122	-	-	-	-	-	-	1	2	1	4	14
White, both sexes	3,156	-	-	-	-	-	1	1	1	3	11	23
Male	1,264	-	-	-	-	-	1	1	1	3	9	11
Female	1,892	-	-	-	-	-	-	-	-	-	2	12
Black, both sexes	345	-	-	-	-	-	-	1	5	2	7	6
Male	131	-	-	-	-	-	-	-	3	1	5	4
Female	214	-	-	-	-	-	-	1	2	1	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	531	-	-	-	-	-	-	-	-	-	1	4
Male	201	-	-	-	-	-	-	-	-	-	1	3
Female	330	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	516	-	-	-	-	-	-	-	-	-	-	3
Male	196	-	-	-	-	-	-	-	-	-	-	2
Female	320	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Vermont												
All races, both sexes	76	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	76	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	52	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,378	-	-	-	-	-	-	-	1	3	7	7
Male	530	-	-	-	-	-	-	-	1	3	6	5
Female	848	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	1,170	-	-	-	-	-	-	-	-	2	4	5
Male	457	-	-	-	-	-	-	-	-	2	3	3
Female	713	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	196	-	-	-	-	-	-	-	1	1	3	2
Male	69	-	-	-	-	-	-	-	1	1	3	2
Female	127	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Washington												
All races, both sexes	516	-	-	-	-	-	-	-	-	-	1	2
Male	201	-	-	-	-	-	-	-	-	-	-	-
Female	315	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	490	-	-	-	-	-	-	-	-	-	1	-
Male	192	-	-	-	-	-	-	-	-	-	-	-
Female	298	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	499	-	-	-	-	-	-	-	-	-	4	5
Male	192	-	-	-	-	-	-	-	-	-	4	5
Female	307	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	484	-	-	-	-	-	-	-	-	-	4	4
Male	187	-	-	-	-	-	-	-	-	-	4	4
Female	297	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Congestive heart failure (I50.0)												
Wisconsin												
All races, both sexes	1,404	-	-	-	-	-	1	-	-	-	1	1
Male	545	-	-	-	-	-	-	-	-	-	-	1
Female	859	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	1,371	-	-	-	-	-	1	-	-	-	1	-
Male	531	-	-	-	-	-	-	-	-	-	-	-
Female	840	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	113	-	-	-	-	-	-	-	-	-	-	1
Male	51	-	-	-	-	-	-	-	-	-	-	1
Female	62	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	111	-	-	-	-	-	-	-	-	-	-	1
Male	50	-	-	-	-	-	-	-	-	-	-	1
Female	61	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
United States												
All races, both sexes	572	914	1,345	1,956	3,379	6,073	10,100	12,437	11,226	4,992	1,209	-
Male	340	540	746	1,034	1,718	2,945	4,335	4,566	3,321	1,077	194	-
Female	232	374	599	922	1,661	3,128	5,765	7,871	7,905	3,915	1,015	-
White, both sexes	375	665	1,035	1,554	2,851	5,321	9,194	11,609	10,512	4,659	1,096	-
Male	231	396	581	827	1,460	2,598	4,005	4,312	3,135	1,008	175	-
Female	144	269	454	727	1,391	2,723	5,189	7,297	7,377	3,651	921	-
Black, both sexes	184	233	281	362	479	659	791	703	595	274	105	-
Male	104	133	148	181	231	308	281	208	142	44	16	-
Female	80	100	133	181	248	351	510	495	453	230	89	-
American Indian 1/, both sexes	7	9	13	19	23	33	31	24	24	15	2	-
Male	3	6	6	13	13	13	12	9	11	4	1	-
Female	4	3	7	6	10	20	19	15	13	11	1	-
Asian or Pacific Islander 2/, both sexes	6	7	16	21	26	60	84	101	95	44	6	-
Male	2	5	11	13	14	26	37	37	33	21	2	-
Female	4	2	5	8	12	34	47	64	62	23	4	-
Alabama												
All races, both sexes	41	54	71	77	138	232	357	369	348	124	37	-
Male	19	32	35	42	71	118	146	118	101	26	9	-
Female	22	22	36	35	67	114	211	251	247	98	28	-
White, both sexes	26	38	46	51	104	184	302	329	292	100	29	-
Male	13	25	25	25	53	95	123	106	85	21	6	-
Female	13	13	21	26	51	89	179	223	207	79	23	-
Black, both sexes	15	15	25	26	33	47	55	40	56	23	8	-
Male	6	6	10	17	18	22	23	12	16	4	3	-
Female	9	9	15	9	15	25	32	28	40	19	5	-
American Indian 1/, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Alaska												
All races, both sexes	3	-	4	4	3	12	7	8	8	4	1	-
Male	3	-	3	3	3	5	4	3	2	2	1	-
Female	-	-	1	1	-	7	3	5	6	2	-	-
White, both sexes	2	-	4	3	3	9	5	6	5	4	1	-
Male	2	-	3	3	3	4	3	1	1	2	1	-
Female	-	-	1	-	-	5	2	5	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	3	1	1	2	-	-	-
Male	1	-	-	-	-	1	1	1	1	-	-	-
Female	-	-	-	1	-	2	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Arizona												
All races, both sexes	4	8	12	24	40	74	149	169	141	62	8	-
Male	1	3	6	13	26	33	71	64	46	17	-	-
Female	3	5	6	11	14	41	78	105	95	45	8	-
White, both sexes	3	7	12	22	38	71	146	166	139	55	8	-
Male	1	3	6	12	24	32	69	62	44	16	-	-
Female	2	4	6	10	14	39	77	104	95	39	8	-
Black, both sexes	-	1	-	1	-	-	-	1	-	2	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	1	-	-	-	2	3	2	-	1	4	-	-
Male	-	-	-	-	2	1	2	-	1	1	-	-
Female	1	-	-	-	-	2	-	-	-	3	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	2	1	1	-	-
Male	-	-	-	1	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Arkansas												
All races, both sexes	12	13	16	31	48	76	133	142	144	55	7	-
Male	9	8	11	17	26	42	57	51	50	12	1	-
Female	3	5	5	14	22	34	76	91	94	43	6	-
White, both sexes	8	12	15	23	41	69	125	137	127	51	6	-
Male	5	8	11	12	23	39	55	49	45	11	1	-
Female	3	4	4	11	18	30	70	88	82	40	5	-
Black, both sexes	4	1	1	8	6	7	8	5	17	4	1	-
Male	4	-	-	5	2	3	2	2	5	1	-	-
Female	-	1	1	3	4	4	6	3	12	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	40	68	90	172	250	423	805	967	881	417	108	-
Male	27	40	56	91	137	202	357	366	302	105	17	-
Female	13	28	34	81	113	221	448	601	579	312	91	-
White, both sexes	30	47	70	141	207	363	721	892	815	390	103	-
Male	22	25	44	76	118	178	321	343	278	98	16	-
Female	8	22	26	65	89	185	400	549	537	292	87	-
Black, both sexes	7	17	12	23	35	34	43	35	26	13	2	-
Male	4	12	7	9	17	13	22	10	9	1	-	-
Female	3	5	5	14	18	21	21	25	17	12	2	-
American Indian 1/, both sexes	1	-	1	2	-	2	5	2	1	-	-	-
Male	-	-	-	2	-	1	3	-	-	-	-	-
Female	1	-	1	-	-	1	2	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	7	6	8	24	36	38	39	14	3	-
Male	1	3	5	4	2	10	11	13	15	6	1	-
Female	1	1	2	2	6	14	25	25	24	8	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Colorado												
All races, both sexes	7	7	18	17	36	55	113	165	123	79	21	-
Male	5	3	9	9	17	23	48	54	43	14	6	-
Female	2	4	9	8	19	32	65	111	80	65	15	-
White, both sexes	6	7	17	17	34	52	112	161	121	76	20	-
Male	4	3	8	9	15	21	48	53	42	14	5	-
Female	2	4	9	8	19	31	64	108	79	62	15	-
Black, both sexes	1	-	-	-	1	2	1	3	2	2	1	-
Male	1	-	-	-	1	2	-	-	1	-	1	-
Female	-	-	-	-	-	-	1	3	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes	4	3	11	11	23	61	117	140	158	72	20	-
Male	4	-	5	4	9	31	56	55	50	10	1	-
Female	-	3	6	7	14	30	61	85	108	62	19	-
White, both sexes	1	3	7	9	18	55	111	132	153	68	18	-
Male	1	-	3	4	5	27	53	54	48	9	1	-
Female	-	3	4	5	13	28	58	78	105	59	17	-
Black, both sexes	3	-	4	2	5	4	6	5	5	3	2	-
Male	3	-	2	-	4	3	3	1	2	-	-	-
Female	-	-	2	2	1	1	3	4	3	3	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	2	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Delaware												
All races, both sexes	2	3	1	9	12	10	24	31	17	10	3	-
Male	1	3	-	5	6	6	12	9	6	-	1	-
Female	1	-	1	4	6	4	12	22	11	10	2	-
White, both sexes	2	3	-	6	8	7	21	27	16	9	3	-
Male	1	3	-	5	3	4	12	9	6	-	1	-
Female	1	-	-	1	5	3	9	18	10	9	2	-
Black, both sexes	-	-	1	3	3	3	2	2	-	1	-	-
Male	-	-	-	-	2	2	-	-	-	-	-	-
Female	-	-	1	3	1	1	2	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	6	2	4	8	14	8	3	2	-
Male	1	-	-	4	2	1	1	6	-	-	-	-
Female	1	-	-	2	-	3	7	8	8	3	2	-
White, both sexes	-	-	-	-	1	1	1	4	2	-	1	-
Male	-	-	-	-	1	1	-	3	-	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	1	-
Black, both sexes	2	-	-	6	1	3	7	10	6	3	1	-
Male	1	-	-	4	1	-	1	3	-	-	-	-
Female	1	-	-	2	-	3	6	7	6	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Florida												
All races, both sexes	9	20	38	61	85	187	265	464	413	178	39	-
Male	4	12	19	39	45	89	114	205	139	57	7	-
Female	5	8	19	22	40	98	151	259	274	121	32	-
White, both sexes	5	17	29	52	68	172	247	440	393	172	35	-
Male	2	10	15	34	36	82	104	201	132	54	7	-
Female	3	7	14	18	32	90	143	239	261	118	28	-
Black, both sexes	4	3	9	9	16	15	17	22	16	6	4	-
Male	2	2	4	5	8	7	9	4	5	3	-	-
Female	2	1	5	4	8	8	8	18	11	3	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	2	3	-	-	-
Male	-	-	-	-	1	-	1	-	2	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
Georgia												
All races, both sexes	41	48	73	109	138	210	333	368	301	103	35	-
Male	22	27	45	56	62	107	121	128	74	20	7	-
Female	19	21	28	53	76	103	212	240	227	83	28	-
White, both sexes	19	32	39	63	102	170	283	310	267	81	26	-
Male	10	19	20	29	44	91	106	113	66	16	5	-
Female	9	13	19	34	58	79	177	197	201	65	21	-
Black, both sexes	22	15	34	45	35	39	49	57	32	22	9	-
Male	12	7	25	27	18	15	14	14	6	4	2	-
Female	10	8	9	18	17	24	35	43	26	18	7	-
American Indian 1/, both sexes	-	1	-	1	1	-	-	-	2	-	-	-
Male	-	1	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Hawaii												
All races, both sexes	3	4	3	6	6	14	20	27	21	17	3	-
Male	3	4	3	5	4	7	12	11	8	9	2	-
Female	-	-	-	1	2	7	8	16	13	8	1	-
White, both sexes	1	2	1	1	1	9	6	10	7	5	1	-
Male	1	2	1	1	1	4	1	4	4	1	1	-
Female	-	-	-	-	-	5	5	6	3	4	-	-
Black, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	5	5	4	14	16	14	12	2	-
Male	1	1	2	4	3	2	11	6	4	8	1	-
Female	-	-	-	1	2	2	3	10	10	4	1	-
Idaho												
All races, both sexes	-	-	4	8	10	26	46	63	56	23	6	-
Male	-	-	3	5	5	13	23	20	25	1	3	-
Female	-	-	1	3	5	13	23	43	31	22	3	-
White, both sexes	-	-	4	8	10	24	46	63	56	23	6	-
Male	-	-	3	5	5	12	23	20	25	1	3	-
Female	-	-	1	3	5	12	23	43	31	22	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Illinois												
All races, both sexes	24	51	62	122	185	310	501	654	633	304	64	-
Male	12	30	33	62	103	157	213	243	180	55	7	-
Female	12	21	29	60	82	153	288	411	453	249	57	-
White, both sexes	16	37	49	90	161	271	453	617	602	286	58	-
Male	9	22	28	47	89	139	199	226	170	50	7	-
Female	7	15	21	43	72	132	254	391	432	236	51	-
Black, both sexes	8	14	10	31	22	37	46	31	25	15	6	-
Male	3	8	4	15	12	17	14	14	8	4	-	-
Female	5	6	6	16	10	20	32	17	17	11	6	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	2	2	1	6	6	3	-	-
Male	-	-	1	-	2	1	-	3	2	1	-	-
Female	-	-	1	1	-	1	1	3	4	2	-	-
Indiana												
All races, both sexes	21	37	54	62	134	206	325	443	382	140	40	-
Male	11	24	30	35	56	80	140	152	110	29	6	-
Female	10	13	24	27	78	126	185	291	272	111	34	-
White, both sexes	18	32	50	52	124	197	300	424	366	137	38	-
Male	9	21	27	29	53	78	135	146	106	27	6	-
Female	9	11	23	23	71	119	165	278	260	110	32	-
Black, both sexes	3	5	4	9	10	9	24	18	16	2	2	-
Male	2	3	3	6	3	2	5	6	4	1	-	-
Female	1	2	1	3	7	7	19	12	12	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Iowa												
All races, both sexes	4	2	6	3	12	24	52	56	72	39	11	-
Male	2	1	3	2	8	15	25	16	20	6	2	-
Female	2	1	3	1	4	9	27	40	52	33	9	-
White, both sexes	4	2	6	3	11	21	51	56	72	39	11	-
Male	2	1	3	2	7	13	24	16	20	6	2	-
Female	2	1	3	1	4	8	27	40	52	33	9	-
Black, both sexes	-	-	-	-	1	3	1	-	-	-	-	-
Male	-	-	-	-	1	2	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	6	12	13	21	48	75	169	197	205	103	26	-
Male	4	7	7	10	22	32	74	74	59	27	5	-
Female	2	5	6	11	26	43	95	123	146	76	21	-
White, both sexes	5	12	10	19	43	63	160	187	202	100	25	-
Male	3	7	5	9	20	26	74	71	58	27	5	-
Female	2	5	5	10	23	37	86	116	144	73	20	-
Black, both sexes	1	-	3	1	4	9	8	8	3	3	-	-
Male	1	-	2	-	1	4	-	3	1	-	-	-
Female	-	-	1	1	3	5	8	5	2	3	-	-
American Indian 1/, both sexes	-	-	-	1	1	3	-	2	-	-	1	-
Male	-	-	-	1	1	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Kentucky												
All races, both sexes	16	30	41	49	121	161	250	299	230	92	21	-
Male	10	22	31	24	67	83	90	100	62	15	2	-
Female	6	8	10	25	54	78	160	199	168	77	19	-
White, both sexes	13	27	38	45	107	147	236	288	225	88	20	-
Male	8	20	30	23	62	74	85	97	60	15	2	-
Female	5	7	8	22	45	73	151	191	165	73	18	-
Black, both sexes	3	3	3	3	14	13	14	11	4	4	1	-
Male	2	2	1	-	5	8	5	3	2	-	-	-
Female	1	1	2	3	9	5	9	8	2	4	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Louisiana												
All races, both sexes	19	30	59	79	104	188	245	316	241	110	22	-
Male	9	19	33	40	55	98	102	115	65	23	2	-
Female	10	11	26	39	49	90	143	201	176	87	20	-
White, both sexes	8	12	36	48	66	140	193	264	197	89	17	-
Male	2	9	22	25	31	82	82	101	49	17	1	-
Female	6	3	14	23	35	58	111	163	148	72	16	-
Black, both sexes	11	18	23	29	37	48	51	51	43	20	5	-
Male	7	10	11	14	24	16	20	13	16	6	1	-
Female	4	8	12	15	13	32	31	38	27	14	4	-
American Indian 1/, both sexes	-	-	-	2	1	-	-	-	1	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Maine												
All races, both sexes	2	4	5	5	9	29	42	57	57	36	4	-
Male	-	2	2	3	5	12	19	20	17	5	-	-
Female	2	2	3	2	4	17	23	37	40	31	4	-
White, both sexes	2	4	5	5	9	28	42	57	57	36	4	-
Male	-	2	2	3	5	12	19	20	17	5	-	-
Female	2	2	3	2	4	16	23	37	40	31	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	9	16	19	22	49	88	138	146	138	49	19	-
Male	7	11	10	11	23	42	56	53	43	16	4	-
Female	2	5	9	11	26	46	82	93	95	33	15	-
White, both sexes	6	14	13	19	36	61	107	131	119	44	14	-
Male	5	10	6	9	16	28	44	51	36	16	4	-
Female	1	4	7	10	20	33	63	80	83	28	10	-
Black, both sexes	3	2	6	3	13	27	29	13	18	5	5	-
Male	2	1	4	2	7	14	10	2	7	-	-	-
Female	1	1	2	1	6	13	19	11	11	5	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	2	1	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Massachusetts												
All races, both sexes	8	13	14	52	80	187	318	442	405	204	56	-
Male	5	5	5	30	37	95	128	165	99	41	14	-
Female	3	8	9	22	43	92	190	277	306	163	42	-
White, both sexes	7	11	12	50	73	176	307	433	393	203	54	-
Male	5	4	4	29	35	91	124	161	97	41	14	-
Female	2	7	8	21	38	85	183	272	296	162	40	-
Black, both sexes	1	2	2	2	7	9	9	7	10	1	2	-
Male	-	1	1	1	2	3	2	2	1	-	-	-
Female	1	1	1	1	5	6	7	5	9	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	2	2	1	-	-	-
Male	-	-	-	-	-	1	2	2	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Michigan												
All races, both sexes	22	36	48	65	124	282	470	557	497	221	53	-
Male	14	16	30	26	61	126	208	210	149	48	6	-
Female	8	20	18	39	63	156	262	347	348	173	47	-
White, both sexes	15	21	37	52	107	249	433	524	475	211	51	-
Male	10	9	21	21	53	111	191	199	145	47	6	-
Female	5	12	16	31	54	138	242	325	330	164	45	-
Black, both sexes	6	12	11	13	16	30	35	28	19	9	2	-
Male	4	6	9	5	8	14	15	11	3	1	-	-
Female	2	6	2	8	8	16	20	17	16	8	2	-
American Indian 1/, both sexes	-	3	-	-	-	1	1	2	1	-	-	-
Male	-	1	-	-	-	-	1	-	1	-	-	-
Female	-	2	-	-	-	1	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	2	1	3	2	1	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	1	-	-	-	1	1	-	3	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Minnesota												
All races, both sexes	7	8	17	27	47	109	172	268	272	132	21	-
Male	6	5	13	20	25	66	73	102	91	25	2	-
Female	1	3	4	7	22	43	99	166	181	107	19	-
White, both sexes	6	6	15	26	43	108	171	265	270	131	21	-
Male	5	3	12	19	22	65	73	101	91	25	2	-
Female	1	3	3	7	21	43	98	164	179	106	19	-
Black, both sexes	-	1	-	-	2	1	-	1	1	1	-	-
Male	-	1	-	-	2	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	1	1	2	1	2	-	-	1	1	-	-	-
Male	1	1	1	1	1	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Mississippi												
All races, both sexes	20	32	58	61	78	170	213	275	233	99	29	-
Male	10	18	31	29	39	86	78	89	63	14	4	-
Female	10	14	27	32	39	84	135	186	170	85	25	-
White, both sexes	9	17	34	40	55	124	165	228	177	80	15	-
Male	4	10	20	19	27	61	63	73	46	9	1	-
Female	5	7	14	21	28	63	102	155	131	71	14	-
Black, both sexes	11	15	23	21	22	46	48	47	56	19	14	-
Male	6	8	11	10	11	25	15	16	17	5	3	-
Female	5	7	12	11	11	21	33	31	39	14	11	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Missouri												
All races, both sexes	13	18	39	46	78	172	277	301	286	137	30	-
Male	9	10	17	23	36	91	112	100	79	23	7	-
Female	4	8	22	23	42	81	165	201	207	114	23	-
White, both sexes	11	15	33	44	71	154	264	289	274	131	29	-
Male	7	8	14	22	34	82	105	96	74	22	7	-
Female	4	7	19	22	37	72	159	193	200	109	22	-
Black, both sexes	2	3	6	2	7	18	12	12	10	6	1	-
Male	2	2	3	1	2	9	6	4	3	1	-	-
Female	-	1	3	1	5	9	6	8	7	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	2	7	6	13	12	27	47	53	68	21	9	-
Male	1	6	4	9	8	10	30	20	24	5	2	-
Female	1	1	2	4	4	17	17	33	44	16	7	-
White, both sexes	2	5	5	12	12	25	45	53	66	21	9	-
Male	1	5	3	8	8	10	29	20	23	5	2	-
Female	1	-	2	4	4	15	16	33	43	16	7	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	1	1	-	2	1	-	1	-	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Nebraska												
All races, both sexes	5	2	6	8	29	41	79	108	110	52	21	-
Male	4	2	4	2	12	23	25	38	33	12	3	-
Female	1	-	2	6	17	18	54	70	77	40	18	-
White, both sexes	5	2	6	8	27	38	76	106	108	52	21	-
Male	4	2	4	2	12	21	25	36	33	12	3	-
Female	1	-	2	6	15	17	51	70	75	40	18	-
Black, both sexes	-	-	-	-	2	2	3	1	2	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	2	1	3	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	16	16	30	46	71	120	94	86	29	4	-
Male	3	10	6	18	23	43	63	40	33	7	1	-
Female	6	6	10	12	23	28	57	54	53	22	3	-
White, both sexes	6	14	13	23	43	65	118	91	81	27	3	-
Male	3	9	4	13	22	41	62	40	30	7	1	-
Female	3	5	9	10	21	24	56	51	51	20	2	-
Black, both sexes	2	1	2	3	3	3	1	2	1	2	1	-
Male	-	-	2	2	1	2	-	-	1	-	-	-
Female	2	1	-	1	2	1	1	2	-	2	1	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	1	2	-	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	2	-	3	1	-	2	-	-	-
Male	-	1	-	2	-	-	1	-	1	-	-	-
Female	1	-	1	-	-	3	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
New Hampshire												
All races, both sexes	1	-	4	1	7	13	31	35	45	28	8	-
Male	-	-	3	-	6	7	12	11	16	4	-	-
Female	1	-	1	1	1	6	19	24	29	24	8	-
White, both sexes	1	-	4	1	7	13	31	35	45	28	8	-
Male	-	-	3	-	6	7	12	11	16	4	-	-
Female	1	-	1	1	1	6	19	24	29	24	8	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	11	13	29	37	91	155	291	355	300	150	20	-
Male	8	8	16	25	44	90	121	122	87	35	2	-
Female	3	5	13	12	47	65	170	233	213	115	18	-
White, both sexes	8	9	22	29	79	130	260	330	278	134	19	-
Male	7	6	14	19	38	75	112	115	85	33	2	-
Female	1	3	8	10	41	55	148	215	193	101	17	-
Black, both sexes	3	4	7	7	9	22	29	23	16	15	1	-
Male	1	2	2	5	4	13	8	7	2	2	-	-
Female	2	2	5	2	5	9	21	16	14	13	1	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	3	2	2	6	1	-	-
Male	-	-	-	-	1	2	1	-	-	-	-	-
Female	-	-	-	-	1	1	1	2	6	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
New Mexico												
All races, both sexes	3	3	6	16	21	50	67	77	68	49	9	-
Male	1	2	3	9	8	20	35	31	19	15	2	-
Female	2	1	3	7	13	30	32	46	49	34	7	-
White, both sexes	2	3	5	14	18	46	62	74	65	45	9	-
Male	1	2	3	7	6	19	33	31	19	14	2	-
Female	1	1	2	7	12	27	29	43	46	31	7	-
Black, both sexes	1	-	1	1	2	-	1	1	1	1	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	1	-	1	-	1	-	-	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	1	1	4	4	2	2	3	-	-
Male	-	-	-	1	1	1	1	-	-	1	-	-
Female	-	-	-	-	-	3	3	2	2	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	21	37	46	62	144	282	484	634	572	257	72	-
Male	12	18	24	29	81	125	215	271	175	54	9	-
Female	9	19	22	33	63	157	269	363	397	203	63	-
White, both sexes	10	26	33	50	122	248	450	588	540	244	64	-
Male	5	12	16	21	71	115	205	254	170	53	9	-
Female	5	14	17	29	51	133	245	334	370	191	55	-
Black, both sexes	11	11	11	11	22	33	33	36	30	12	7	-
Male	7	6	7	7	10	10	10	12	4	1	-	-
Female	4	5	4	4	12	23	23	24	26	11	7	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	1	1	9	2	-	1	-
Male	-	-	1	1	-	-	-	4	1	-	-	-
Female	-	-	-	-	-	1	1	5	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
North Carolina												
All races, both sexes	20	25	33	59	105	152	278	310	254	104	25	-
Male	12	11	17	30	52	76	125	100	58	18	6	-
Female	8	14	16	29	53	76	153	210	196	86	19	-
White, both sexes	9	13	20	42	79	123	228	271	230	94	24	-
Male	7	4	10	19	40	61	107	93	51	16	6	-
Female	2	9	10	23	39	62	121	178	179	78	18	-
Black, both sexes	11	12	12	16	26	28	47	38	23	9	1	-
Male	5	7	7	11	12	15	16	7	6	1	-	-
Female	6	5	5	5	14	13	31	31	17	8	1	-
American Indian 1/, both sexes	-	-	1	-	-	1	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	1	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	1	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	1	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	3	3	7	25	21	17	6	-
Male	-	-	-	-	2	2	2	6	7	5	1	-
Female	-	-	-	-	1	1	5	19	14	12	5	-
White, both sexes	-	-	-	-	3	3	7	24	20	17	6	-
Male	-	-	-	-	2	2	2	6	6	5	1	-
Female	-	-	-	-	1	1	5	18	14	12	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Ohio												
All races, both sexes	14	27	47	74	147	240	380	471	377	147	37	-
Male	9	18	23	34	66	111	160	181	106	31	10	-
Female	5	9	24	40	81	129	220	290	271	116	27	-
White, both sexes	12	19	42	60	133	220	352	447	359	138	29	-
Male	8	14	22	31	60	102	149	169	102	29	6	-
Female	4	5	20	29	73	118	203	278	257	109	23	-
Black, both sexes	2	8	5	14	14	18	28	24	18	9	8	-
Male	1	4	1	3	6	9	11	12	4	2	4	-
Female	1	4	4	11	8	9	17	12	14	7	4	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Oklahoma												
All races, both sexes	9	21	41	35	76	130	213	200	205	84	20	-
Male	5	13	24	20	31	53	97	64	49	18	6	-
Female	4	8	17	15	45	77	116	136	156	66	14	-
White, both sexes	7	14	36	28	65	119	198	187	199	77	19	-
Male	4	9	20	13	27	50	95	55	48	16	6	-
Female	3	5	16	15	38	69	103	132	151	61	13	-
Black, both sexes	1	6	2	4	5	5	6	7	5	3	1	-
Male	1	4	2	4	2	1	-	4	-	-	-	-
Female	-	2	-	-	3	4	6	3	5	3	1	-
American Indian 1/, both sexes	1	-	3	3	6	6	8	6	1	4	-	-
Male	-	-	2	3	2	2	2	5	1	2	-	-
Female	1	-	1	-	4	4	6	1	-	2	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Oregon												
All races, both sexes	3	9	13	17	41	68	117	198	186	90	32	-
Male	3	4	7	8	20	36	49	75	68	20	6	-
Female	-	5	6	9	21	32	68	123	118	70	26	-
White, both sexes	2	9	13	17	41	67	116	195	181	89	31	-
Male	2	4	7	8	20	35	49	73	66	20	5	-
Female	-	5	6	9	21	32	67	122	115	69	26	-
Black, both sexes	-	-	-	-	-	-	1	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	1	-	1	-
Male	1	-	-	-	-	1	-	-	1	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	2	3	1	-	-
Male	-	-	-	-	-	-	-	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	2	1	-	-
Pennsylvania												
All races, both sexes	29	38	58	82	183	357	628	816	687	309	64	-
Male	21	18	34	50	95	172	274	294	181	68	4	-
Female	8	20	24	32	88	185	354	522	506	241	60	-
White, both sexes	21	26	46	69	165	332	593	784	659	300	61	-
Male	18	11	25	44	88	156	261	288	178	68	4	-
Female	3	15	21	25	77	176	332	496	481	232	57	-
Black, both sexes	8	12	12	13	18	23	33	32	26	8	3	-
Male	3	7	9	6	7	14	12	6	3	-	-	-
Female	5	5	3	7	11	9	21	26	23	8	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	2	-	2	1	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Rhode Island												
All races, both sexes	1	1	1	2	4	23	17	30	19	7	2	-
Male	1	-	-	2	2	10	7	9	3	1	-	-
Female	-	1	1	-	2	13	10	21	16	6	2	-
White, both sexes	-	1	1	1	4	22	17	30	19	7	2	-
Male	-	-	-	1	2	9	7	9	3	1	-	-
Female	-	1	1	-	2	13	10	21	16	6	2	-
Black, both sexes	1	-	-	1	-	1	-	-	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	17	23	31	39	73	124	161	179	144	62	14	-
Male	12	15	16	18	42	60	78	61	41	10	3	-
Female	5	8	15	21	31	64	83	118	103	52	11	-
White, both sexes	12	11	15	27	50	93	130	148	120	44	9	-
Male	9	6	8	13	28	41	63	54	40	8	2	-
Female	3	5	7	14	22	52	67	94	80	36	7	-
Black, both sexes	5	12	16	12	23	31	31	30	23	18	5	-
Male	3	9	8	5	14	19	15	7	1	2	1	-
Female	2	3	8	7	9	12	16	23	22	16	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
South Dakota												
All races, both sexes	-	-	-	-	3	3	4	13	12	8	6	-
Male	-	-	-	-	2	3	3	6	2	1	-	-
Female	-	-	-	-	1	-	1	7	10	7	6	-
White, both sexes	-	-	-	-	2	2	2	13	12	8	6	-
Male	-	-	-	-	1	2	2	6	2	1	-	-
Female	-	-	-	-	1	-	-	7	10	7	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	9	18	16	34	43	72	127	130	125	48	13	-
Male	5	10	9	21	17	24	57	51	29	12	2	-
Female	4	8	7	13	26	48	70	79	96	36	11	-
White, both sexes	6	12	13	28	36	61	113	118	116	43	11	-
Male	2	8	7	19	16	22	50	47	26	11	2	-
Female	4	4	6	9	20	39	63	71	90	32	9	-
Black, both sexes	3	6	3	6	7	9	14	12	9	5	2	-
Male	3	2	2	2	1	1	7	4	3	1	-	-
Female	-	4	1	4	6	8	7	8	6	4	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Texas												
All races, both sexes	52	84	125	144	243	414	646	730	676	296	56	-
Male	29	56	63	79	134	197	268	254	212	67	7	-
Female	23	28	62	65	109	217	378	476	464	229	49	-
White, both sexes	32	62	99	122	207	366	595	677	635	272	49	-
Male	19	41	54	70	118	171	253	240	204	62	6	-
Female	13	21	45	52	89	195	342	437	431	210	43	-
Black, both sexes	20	22	24	21	33	44	47	49	38	19	7	-
Male	10	15	8	8	13	25	15	13	7	3	1	-
Female	10	7	16	13	20	19	32	36	31	16	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	3	4	3	4	3	5	-	-
Male	-	-	1	1	3	1	-	1	1	2	-	-
Female	-	-	1	-	-	3	3	3	2	3	-	-
Utah												
All races, both sexes	5	9	9	13	29	46	95	138	125	53	4	-
Male	3	6	6	7	10	21	40	48	39	16	1	-
Female	2	3	3	6	19	25	55	90	86	37	3	-
White, both sexes	4	8	9	13	25	45	94	137	123	51	4	-
Male	2	6	6	7	9	21	40	48	38	16	1	-
Female	2	2	3	6	16	24	54	89	85	35	3	-
Black, both sexes	1	1	-	-	1	-	1	1	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	-	2	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	1	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Vermont												
All races, both sexes	-	1	2	2	1	7	12	24	14	12	1	-
Male	-	1	1	2	1	4	3	7	4	1	-	-
Female	-	-	1	-	-	3	9	17	10	11	1	-
White, both sexes	-	1	2	2	1	7	12	24	14	12	1	-
Male	-	1	1	2	1	4	3	7	4	1	-	-
Female	-	-	1	-	-	3	9	17	10	11	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	8	26	40	59	79	165	276	288	275	117	27	-
Male	6	18	23	25	43	78	110	103	76	28	5	-
Female	2	8	17	34	36	87	166	185	199	89	22	-
White, both sexes	3	19	33	44	59	135	231	254	247	109	25	-
Male	2	14	20	21	33	66	96	92	73	27	5	-
Female	1	5	13	23	26	69	135	162	174	82	20	-
Black, both sexes	5	7	7	13	20	29	41	32	25	8	2	-
Male	4	4	3	4	10	11	13	10	2	1	-	-
Female	1	3	4	9	10	18	28	22	23	7	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	1	4	2	3	-	-	-
Male	-	-	-	-	-	1	1	1	1	-	-	-
Female	-	-	-	2	-	-	3	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Washington												
All races, both sexes	6	5	8	15	34	50	101	117	112	46	19	-
Male	2	3	6	9	20	23	43	52	32	10	1	-
Female	4	2	2	6	14	27	58	65	80	36	18	-
White, both sexes	5	5	8	12	32	46	93	114	110	45	19	-
Male	2	3	6	8	20	20	40	50	32	10	1	-
Female	3	2	2	4	12	26	53	64	78	35	18	-
Black, both sexes	-	-	-	2	1	2	6	1	-	-	-	-
Male	-	-	-	1	-	2	3	1	-	-	-	-
Female	-	-	-	1	1	-	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	1	1	2	1	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	1	1	1	1	1	1	-	-
West Virginia												
All races, both sexes	3	10	14	17	46	63	105	92	90	38	12	-
Male	1	5	7	6	24	34	44	33	17	10	2	-
Female	2	5	7	11	22	29	61	59	73	28	10	-
White, both sexes	2	10	13	17	45	61	104	90	84	38	12	-
Male	1	5	7	6	24	33	43	33	15	10	2	-
Female	1	5	6	11	21	28	61	57	69	28	10	-
Black, both sexes	1	-	1	-	1	2	1	2	6	-	-	-
Male	-	-	-	-	-	1	1	-	2	-	-	-
Female	1	-	1	-	1	1	-	2	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Congestive heart failure (I50.0)												
Wisconsin												
All races, both sexes	5	18	15	43	55	123	244	358	363	141	36	-
Male	4	12	8	20	25	59	118	154	115	25	4	-
Female	1	6	7	23	30	64	126	204	248	116	32	-
White, both sexes	3	14	13	41	51	120	238	355	358	141	35	-
Male	3	8	7	20	23	59	115	153	114	25	4	-
Female	-	6	6	21	28	61	123	202	244	116	31	-
Black, both sexes	1	3	1	1	2	2	3	3	4	-	1	-
Male	1	3	-	-	1	-	2	1	1	-	-	-
Female	-	-	1	1	1	2	1	2	3	-	1	-
American Indian 1/, both sexes	1	1	1	1	2	-	1	-	1	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	1	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Wyoming												
All races, both sexes	-	4	3	5	4	9	21	24	22	14	6	-
Male	-	2	2	3	-	4	16	6	12	4	1	-
Female	-	2	1	2	4	5	5	18	10	10	5	-
White, both sexes	-	4	2	5	4	9	21	24	21	14	6	-
Male	-	2	1	3	-	4	16	6	12	4	1	-
Female	-	2	1	2	4	5	5	18	9	10	5	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
United States												
All races, both sexes	4,052	20	2	1	5	5	8	9	15	21	36	80
Male	1,782	10	1	1	2	3	5	4	10	13	26	45
Female	2,270	10	1	-	3	2	3	5	5	8	10	35
White, both sexes	3,312	10	-	-	1	4	4	7	11	8	26	44
Male	1,446	6	-	-	-	2	3	4	8	5	17	28
Female	1,866	4	-	-	1	2	1	3	3	3	9	16
Black, both sexes	669	6	2	1	4	1	4	2	3	13	10	33
Male	304	2	1	1	2	1	2	-	2	8	9	16
Female	365	4	1	-	2	-	2	2	1	5	1	17
American Indian 1/, both sexes	19	2	-	-	-	-	-	-	-	-	-	1
Male	9	1	-	-	-	-	-	-	-	-	-	-
Female	10	1	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	52	2	-	-	-	-	-	-	1	-	-	2
Male	23	1	-	-	-	-	-	-	-	-	-	1
Female	29	1	-	-	-	-	-	-	1	-	-	1
Alabama												
All races, both sexes	325	-	-	-	1	-	-	-	2	-	4	11
Male	151	-	-	-	-	-	-	-	1	-	3	7
Female	174	-	-	-	1	-	-	-	1	-	1	4
White, both sexes	199	-	-	-	-	-	-	-	1	-	2	1
Male	97	-	-	-	-	-	-	-	-	-	2	1
Female	102	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	125	-	-	-	1	-	-	-	1	-	2	10
Male	54	-	-	-	-	-	-	-	1	-	1	6
Female	71	-	-	-	1	-	-	-	-	-	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Alaska												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	34	2	-	-	-	-	-	-	-	1	1	-
Male	16	1	-	-	-	-	-	-	-	-	1	-
Female	18	1	-	-	-	-	-	-	-	1	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Arkansas												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	64	-	-	-	-	-	-	1	-	-	1	1
Male	23	-	-	-	-	-	-	1	-	-	1	-
Female	41	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	62	-	-	-	-	-	-	1	-	-	-	1
Male	21	-	-	-	-	-	-	1	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Colorado												
All races, both sexes	55	-	-	-	-	-	-	-	1	-	1	1
Male	25	-	-	-	-	-	-	-	1	-	1	1
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	-	1	-	1	1
Male	24	-	-	-	-	-	-	-	1	-	1	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	71	1	-	-	-	-	-	-	-	1	-	-
Male	33	-	-	-	-	-	-	-	-	1	-	-
Female	38	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	65	1	-	-	-	-	-	-	-	1	-	-
Male	29	-	-	-	-	-	-	-	-	1	-	-
Female	36	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Delaware												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Florida												
All races, both sexes	125	-	-	-	-	-	1	-	-	-	-	2
Male	53	-	-	-	-	-	-	-	-	-	-	1
Female	72	-	-	-	-	-	1	-	-	-	-	1
White, both sexes	111	-	-	-	-	-	1	-	-	-	-	2
Male	47	-	-	-	-	-	-	-	-	-	-	1
Female	64	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	266	-	-	-	-	2	2	2	-	2	2	4
Male	113	-	-	-	-	1	1	1	-	-	1	3
Female	153	-	-	-	-	1	1	1	-	2	1	1
White, both sexes	188	-	-	-	-	2	-	1	-	-	1	3
Male	75	-	-	-	-	1	-	1	-	-	-	3
Female	113	-	-	-	-	1	-	-	-	-	1	-
Black, both sexes	77	-	-	-	-	-	2	1	-	2	1	1
Male	38	-	-	-	-	-	1	-	-	-	1	-
Female	39	-	-	-	-	-	1	1	-	2	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Hawaii												
All races, both sexes	32	-	-	-	1	-	-	-	-	-	-	1
Male	18	-	-	-	1	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Illinois												
All races, both sexes	181	-	-	-	-	-	1	-	1	-	2	3
Male	67	-	-	-	-	-	1	-	1	-	1	1
Female	114	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	150	-	-	-	-	-	1	-	1	-	2	-
Male	56	-	-	-	-	-	1	-	1	-	1	-
Female	94	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	30	-	-	-	-	-	-	-	-	-	-	3
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	100	1	-	-	-	1	-	-	-	-	-	3
Male	47	1	-	-	-	1	-	-	-	-	-	-
Female	53	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	94	1	-	-	-	1	-	-	-	-	-	1
Male	45	1	-	-	-	1	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Iowa												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	28	-	-	-	-	-	1	-	-	-	-	-
Male	10	-	-	-	-	-	1	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	1	-	-	-	-	-
Male	10	-	-	-	-	-	1	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Kentucky												
All races, both sexes	69	2	-	-	-	-	1	-	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	48	2	-	-	-	-	1	-	-	-	-	-
White, both sexes	64	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	2	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	2	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	71	-	-	-	1	-	-	-	1	1	3	1
Male	33	-	-	-	-	-	-	-	-	1	2	1
Female	38	-	-	-	1	-	-	-	1	-	1	-
White, both sexes	57	-	-	-	-	-	-	-	1	1	2	1
Male	27	-	-	-	-	-	-	-	-	1	1	1
Female	30	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	13	-	-	-	1	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Maine												
All races, both sexes	20	1	-	-	-	-	1	-	-	1	-	-
Male	7	-	-	-	-	-	1	-	-	1	-	-
Female	13	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	20	1	-	-	-	-	1	-	-	1	-	-
Male	7	-	-	-	-	-	1	-	-	1	-	-
Female	13	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Massachusetts												
All races, both sexes	121	1	-	-	-	-	-	-	-	-	1	1
Male	51	-	-	-	-	-	-	-	-	-	1	1
Female	70	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	114	-	-	-	-	-	-	-	-	-	1	1
Male	48	-	-	-	-	-	-	-	-	-	1	1
Female	66	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	94	1	-	-	-	1	-	-	-	2	2	-
Male	45	1	-	-	-	1	-	-	-	2	1	-
Female	49	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	79	1	-	-	-	-	-	-	-	-	2	-
Male	36	1	-	-	-	-	-	-	-	-	1	-
Female	43	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	15	-	-	-	-	1	-	-	-	2	-	-
Male	9	-	-	-	-	1	-	-	-	2	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Minnesota												
All races, both sexes	59	-	-	-	-	-	-	-	-	-	-	-
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	57	-	-	-	-	-	-	-	-	-	-	-
Male	30	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	127	-	-	-	-	-	-	-	-	1	-	5
Male	50	-	-	-	-	-	-	-	-	1	-	2
Female	77	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	77	-	-	-	-	-	-	-	-	-	-	2
Male	30	-	-	-	-	-	-	-	-	-	-	1
Female	47	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	50	-	-	-	-	-	-	-	-	1	-	3
Male	20	-	-	-	-	-	-	-	-	1	-	1
Female	30	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Missouri												
All races, both sexes	80	1	-	-	1	-	-	1	-	1	-	2
Male	29	1	-	-	-	-	-	-	-	1	-	-
Female	51	-	-	-	1	-	-	1	-	-	-	2
White, both sexes	68	1	-	-	1	-	-	1	-	-	-	1
Male	25	1	-	-	-	-	-	-	-	-	-	-
Female	43	-	-	-	1	-	-	1	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Nebraska												
All races, both sexes	92	-	-	-	-	-	-	-	-	-	1	3
Male	48	-	-	-	-	-	-	-	-	-	-	3
Female	44	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	91	-	-	-	-	-	-	-	-	-	1	3
Male	47	-	-	-	-	-	-	-	-	-	-	3
Female	44	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
New Hampshire												
All races, both sexes	14	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	14	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	115	1	-	-	-	-	-	-	-	-	1	3
Male	46	1	-	-	-	-	-	-	-	-	1	3
Female	69	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	97	1	-	-	-	-	-	-	-	-	-	-
Male	37	1	-	-	-	-	-	-	-	-	-	-
Female	60	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	1	2
Male	8	-	-	-	-	-	-	-	-	-	1	2
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
New Mexico												
All races, both sexes	13	1	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	195	1	-	-	-	-	-	1	-	2	-	3
Male	84	-	-	-	-	-	-	-	-	-	-	1
Female	111	1	-	-	-	-	-	1	-	2	-	2
White, both sexes	172	-	-	-	-	-	-	1	-	2	-	2
Male	77	-	-	-	-	-	-	-	-	-	-	1
Female	95	-	-	-	-	-	-	1	-	2	-	1
Black, both sexes	22	1	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
North Carolina												
All races, both sexes	107	-	1	-	1	-	-	1	-	-	2	2
Male	43	-	1	-	1	-	-	-	-	-	2	-
Female	64	-	-	-	-	-	-	1	-	-	-	2
White, both sexes	74	-	-	-	-	-	-	-	-	-	1	2
Male	31	-	-	-	-	-	-	-	-	-	1	-
Female	43	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	31	-	1	-	1	-	-	1	-	-	1	-
Male	11	-	1	-	1	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Ohio												
All races, both sexes	107	-	1	-	-	-	1	1	-	-	1	1
Male	50	-	-	-	-	-	1	1	-	-	1	1
Female	57	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	91	-	-	-	-	-	-	1	-	-	1	-
Male	45	-	-	-	-	-	-	1	-	-	1	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	1	-	-	-	1	-	-	-	-	1
Male	4	-	-	-	-	-	1	-	-	-	-	1
Female	11	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	90	1	-	-	-	-	-	-	2	-	1	2
Male	32	1	-	-	-	-	-	-	2	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	79	-	-	-	-	-	-	-	2	-	1	-
Male	25	-	-	-	-	-	-	-	2	-	-	-
Female	54	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	7	1	-	-	-	-	-	-	-	-	-	1
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Oregon												
All races, both sexes	16	1	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	7	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	15	1	-	-	-	-	-	-	1	-	-	-
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	278	1	-	-	-	-	-	1	-	1	1	5
Male	131	1	-	-	-	-	-	-	-	1	-	4
Female	147	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	245	-	-	-	-	-	-	1	-	1	1	4
Male	110	-	-	-	-	-	-	-	-	1	-	4
Female	135	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	32	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Rhode Island												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	2	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	2	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	171	-	-	1	-	-	-	-	1	2	1	3
Male	87	-	-	1	-	-	-	-	1	1	1	1
Female	84	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	118	-	-	-	-	-	-	-	-	1	-	1
Male	60	-	-	-	-	-	-	-	-	1	-	-
Female	58	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	53	-	-	1	-	-	-	-	1	1	1	2
Male	27	-	-	1	-	-	-	-	1	-	1	1
Female	26	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
South Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	52	1	-	-	-	1	-	-	1	-	1	3
Male	31	1	-	-	-	-	-	-	-	-	1	2
Female	21	-	-	-	-	1	-	-	1	-	-	1
White, both sexes	38	1	-	-	-	1	-	-	-	-	1	2
Male	23	1	-	-	-	-	-	-	-	-	1	1
Female	15	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Texas												
All races, both sexes	295	1	-	-	-	-	-	-	3	2	6	11
Male	140	1	-	-	-	-	-	-	2	1	4	7
Female	155	-	-	-	-	-	-	-	1	1	2	4
White, both sexes	259	1	-	-	-	-	-	-	3	1	4	8
Male	118	1	-	-	-	-	-	-	2	-	2	5
Female	141	-	-	-	-	-	-	-	1	1	2	3
Black, both sexes	35	-	-	-	-	-	-	-	-	1	2	3
Male	22	-	-	-	-	-	-	-	-	1	2	2
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Vermont												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	232	2	-	-	-	-	-	-	1	4	1	3
Male	98	1	-	-	-	-	-	-	-	3	1	3
Female	134	1	-	-	-	-	-	-	1	1	-	-
White, both sexes	167	1	-	-	-	-	-	-	-	-	1	3
Male	72	-	-	-	-	-	-	-	-	-	1	3
Female	95	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	60	1	-	-	-	-	-	-	1	4	-	-
Male	24	1	-	-	-	-	-	-	-	3	-	-
Female	36	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Washington												
All races, both sexes	42	-	-	-	-	-	-	1	-	-	-	2
Male	19	-	-	-	-	-	-	1	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	36	-	-	-	-	-	-	1	-	-	-	2
Male	17	-	-	-	-	-	-	1	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	58	-	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	57	-	-	-	-	-	-	-	-	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other and unspecified heart failure (I50.1-I50.9)												
Wisconsin												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
United States												
All races, both sexes	131	164	187	220	333	507	651	743	573	272	69	-
Male	83	101	122	129	188	239	285	277	164	61	13	-
Female	48	63	65	91	145	268	366	466	409	211	56	-
White, both sexes	81	111	130	169	270	418	538	662	517	239	62	-
Male	50	69	89	102	157	204	240	251	148	51	12	-
Female	31	42	41	67	113	214	298	411	369	188	50	-
Black, both sexes	45	49	55	46	59	82	99	73	47	29	6	-
Male	29	31	32	23	30	32	38	23	13	8	1	-
Female	16	18	23	23	29	50	61	50	34	21	5	-
American Indian 1/, both sexes	2	1	-	-	2	2	6	-	2	1	-	-
Male	2	1	-	-	1	-	2	-	1	1	-	-
Female	-	-	-	-	1	2	4	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	3	2	5	2	5	8	8	7	3	1	-
Male	2	-	1	4	-	3	5	3	2	1	-	-
Female	1	3	1	1	2	2	3	5	5	2	1	-
Alabama												
All races, both sexes	23	18	20	23	35	42	48	52	28	15	3	-
Male	10	16	11	11	23	18	20	20	8	3	-	-
Female	13	2	9	12	12	24	28	32	20	12	3	-
White, both sexes	12	10	12	12	24	30	29	34	19	11	2	-
Male	4	10	7	7	18	15	13	11	7	2	-	-
Female	8	-	5	5	6	15	16	23	12	9	2	-
Black, both sexes	11	8	8	11	10	12	19	18	9	4	1	-
Male	6	6	4	4	5	3	7	9	1	1	-	-
Female	5	2	4	7	5	9	12	9	8	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Alaska												
All races, both sexes	-	1	-	-	1	2	2	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	2	2	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	1	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	2	-	3	3	5	2	7	5	2	1	-
Male	-	-	-	3	1	2	1	5	1	1	-	-
Female	-	2	-	-	2	3	1	2	4	1	1	-
White, both sexes	-	2	-	3	3	4	2	7	5	2	1	-
Male	-	-	-	3	1	1	1	5	1	1	-	-
Female	-	2	-	-	2	3	1	2	4	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Arkansas												
All races, both sexes	-	2	2	3	4	4	8	6	3	-	-	-
Male	-	1	2	2	3	2	4	4	1	-	-	-
Female	-	1	-	1	1	2	4	2	2	-	-	-
White, both sexes	-	2	2	3	4	4	8	5	3	-	-	-
Male	-	1	2	2	3	2	4	4	1	-	-	-
Female	-	1	-	1	1	2	4	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2	-	2	5	4	7	12	11	10	7	1	-
Male	1	-	1	5	1	4	5	1	2	1	-	-
Female	1	-	1	-	3	3	7	10	8	6	1	-
White, both sexes	1	-	2	5	4	7	12	11	10	7	1	-
Male	-	-	1	5	1	4	5	1	2	1	-	-
Female	1	-	1	-	3	3	7	10	8	6	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Colorado												
All races, both sexes	2	-	2	1	2	8	14	10	10	3	-	-
Male	1	-	-	-	1	3	6	5	5	1	-	-
Female	1	-	2	1	1	5	8	5	5	2	-	-
White, both sexes	2	-	2	1	2	7	12	10	10	3	-	-
Male	1	-	-	-	1	3	5	5	5	1	-	-
Female	1	-	2	1	1	4	7	5	5	2	-	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	2	5	2	7	9	7	20	12	3	1	-
Male	1	1	3	1	6	6	1	6	6	-	1	-
Female	-	1	2	1	1	3	6	14	6	3	-	-
White, both sexes	1	1	3	1	6	8	7	20	12	3	1	-
Male	1	-	2	1	5	5	1	6	6	-	1	-
Female	-	1	1	-	1	3	6	14	6	3	-	-
Black, both sexes	-	1	2	1	1	1	-	-	-	-	-	-
Male	-	1	1	-	1	1	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Delaware												
All races, both sexes	-	-	1	3	-	2	4	2	1	1	-	-
Male	-	-	1	2	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	1	3	2	1	1	-	-
White, both sexes	-	-	1	1	-	2	4	2	1	1	-	-
Male	-	-	1	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	3	2	1	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Florida												
All races, both sexes	3	4	4	4	8	15	18	34	21	10	1	-
Male	1	2	3	2	6	10	7	11	8	2	-	-
Female	2	2	1	2	2	5	11	23	13	8	1	-
White, both sexes	3	4	2	3	8	13	13	32	19	10	1	-
Male	1	2	2	2	6	9	4	10	8	2	-	-
Female	2	2	-	1	2	4	9	22	11	8	1	-
Black, both sexes	-	-	2	1	-	2	4	1	2	-	-	-
Male	-	-	1	-	-	1	2	1	-	-	-	-
Female	-	-	1	1	-	1	2	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes	4	8	16	16	26	38	43	47	38	14	2	-
Male	1	5	12	13	15	16	17	16	4	6	1	-
Female	3	3	4	3	11	22	26	31	34	8	1	-
White, both sexes	2	2	8	10	18	29	33	39	31	7	2	-
Male	-	1	5	9	10	10	14	14	4	2	1	-
Female	2	1	3	1	8	19	19	25	27	5	1	-
Black, both sexes	2	6	8	6	8	9	10	8	6	7	-	-
Male	1	4	7	4	5	6	3	2	-	4	-	-
Female	1	2	1	2	3	3	7	6	6	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Hawaii												
All races, both sexes	-	-	1	2	-	3	6	9	6	2	1	-
Male	-	-	1	2	-	1	5	6	1	1	-	-
Female	-	-	-	-	-	2	1	3	5	1	1	-
White, both sexes	-	-	1	1	-	1	2	3	2	-	-	-
Male	-	-	1	1	-	-	2	3	-	-	-	-
Female	-	-	-	-	-	1	-	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	4	6	4	2	1	-
Male	-	-	-	1	-	1	3	3	1	1	-	-
Female	-	-	-	-	-	1	1	3	3	1	1	-
Idaho												
All races, both sexes	-	-	1	-	-	3	2	1	1	-	-	-
Male	-	-	1	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	2	-	1	1	-	-	-
White, both sexes	-	-	1	-	-	3	2	1	1	-	-	-
Male	-	-	1	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	2	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Illinois												
All races, both sexes	6	4	2	11	16	24	32	27	29	17	6	-
Male	4	1	2	7	9	9	12	11	5	3	-	-
Female	2	3	-	4	7	15	20	16	24	14	6	-
White, both sexes	4	2	-	8	13	22	27	24	26	14	6	-
Male	3	1	-	5	8	9	11	10	4	2	-	-
Female	1	1	-	3	5	13	16	14	22	12	6	-
Black, both sexes	2	2	2	2	3	2	5	3	3	3	-	-
Male	1	-	2	2	1	-	1	1	1	1	-	-
Female	1	2	-	-	2	2	4	2	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	6	4	6	9	13	19	15	18	4	-	-
Male	1	3	3	4	6	7	7	6	8	-	-	-
Female	-	3	1	2	3	6	12	9	10	4	-	-
White, both sexes	1	5	3	6	9	13	18	15	17	4	-	-
Male	1	3	2	4	6	7	7	6	7	-	-	-
Female	-	2	1	2	3	6	11	9	10	4	-	-
Black, both sexes	-	1	1	-	-	-	1	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Iowa												
All races, both sexes	1	-	1	-	1	4	5	3	4	-	1	-
Male	1	-	1	-	-	-	3	1	1	-	-	-
Female	-	-	-	-	1	4	2	2	3	-	1	-
White, both sexes	1	-	1	-	1	4	5	3	4	-	1	-
Male	1	-	1	-	-	-	3	1	1	-	-	-
Female	-	-	-	-	1	4	2	2	3	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	-	1	-	4	3	5	3	1	5	4	1	-
Male	-	1	-	2	-	3	2	-	-	1	-	-
Female	-	-	-	2	3	2	1	1	5	3	1	-
White, both sexes	-	1	-	4	3	5	3	1	5	4	1	-
Male	-	1	-	2	-	3	2	-	-	1	-	-
Female	-	-	-	2	3	2	1	1	5	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Kentucky												
All races, both sexes	3	3	2	7	3	9	9	8	16	5	1	-
Male	2	1	2	2	3	2	3	3	3	-	-	-
Female	1	2	-	5	-	7	6	5	13	5	1	-
White, both sexes	3	2	2	7	3	9	9	7	16	5	1	-
Male	2	-	2	2	3	2	3	3	3	-	-	-
Female	1	2	-	5	-	7	6	4	13	5	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3	2	3	3	6	8	16	10	5	6	2	-
Male	2	1	3	2	2	4	9	3	2	1	-	-
Female	1	1	-	1	4	4	7	7	3	5	2	-
White, both sexes	1	2	2	3	4	6	12	9	5	6	2	-
Male	1	1	2	2	-	4	8	3	2	1	-	-
Female	-	1	-	1	4	2	4	6	3	5	2	-
Black, both sexes	2	-	1	-	2	2	3	1	-	-	-	-
Male	1	-	1	-	2	-	1	-	-	-	-	-
Female	1	-	-	-	-	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Maine												
All races, both sexes	-	-	1	1	1	3	-	7	3	1	-	-
Male	-	-	-	1	-	1	-	2	-	1	-	-
Female	-	-	1	-	1	2	-	5	3	-	-	-
White, both sexes	-	-	1	1	1	3	-	7	3	1	-	-
Male	-	-	-	1	-	1	-	2	-	1	-	-
Female	-	-	1	-	1	2	-	5	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2	2	3	1	4	7	5	5	2	-	-	-
Male	2	1	3	1	1	3	1	2	1	-	-	-
Female	-	1	-	-	3	4	4	3	1	-	-	-
White, both sexes	-	2	1	1	3	3	5	5	2	-	-	-
Male	-	1	1	1	1	1	1	2	1	-	-	-
Female	-	1	-	-	2	2	4	3	1	-	-	-
Black, both sexes	2	-	2	-	1	4	-	-	-	-	-	-
Male	2	-	2	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Massachusetts												
All races, both sexes	2	2	1	5	6	11	22	26	26	13	4	-
Male	2	1	1	3	4	7	11	9	7	3	1	-
Female	-	1	-	2	2	4	11	17	19	10	3	-
White, both sexes	2	2	1	3	6	11	21	23	26	13	4	-
Male	2	1	1	1	4	7	11	8	7	3	1	-
Female	-	1	-	2	2	4	10	15	19	10	3	-
Black, both sexes	-	-	-	1	-	-	1	3	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	2	2	3	11	16	20	14	13	3	3	-
Male	1	-	2	2	5	7	6	9	6	1	1	-
Female	-	2	-	1	6	9	14	5	7	2	2	-
White, both sexes	1	2	1	3	9	13	16	12	13	3	3	-
Male	1	-	1	2	4	6	4	8	6	1	1	-
Female	-	2	-	1	5	7	12	4	7	2	2	-
Black, both sexes	-	-	1	-	2	3	4	2	-	-	-	-
Male	-	-	1	-	1	1	2	1	-	-	-	-
Female	-	-	-	-	1	2	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Minnesota												
All races, both sexes	1	1	-	3	5	3	6	19	11	8	2	-
Male	1	-	-	2	5	3	4	9	5	1	-	-
Female	-	1	-	1	-	-	2	10	6	7	2	-
White, both sexes	1	-	-	2	5	3	6	19	11	8	2	-
Male	1	-	-	2	5	3	4	9	5	1	-	-
Female	-	-	-	-	-	-	2	10	6	7	2	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	7	5	11	6	10	11	17	23	17	10	4	-
Male	6	4	5	2	4	4	8	7	5	1	1	-
Female	1	1	6	4	6	7	9	16	12	9	3	-
White, both sexes	3	5	4	2	5	7	7	20	12	8	2	-
Male	2	4	2	-	3	3	4	7	3	1	-	-
Female	1	1	2	2	2	4	3	13	9	7	2	-
Black, both sexes	4	-	7	4	5	4	10	3	5	2	2	-
Male	4	-	3	2	1	1	4	-	2	-	1	-
Female	-	-	4	2	4	3	6	3	3	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Missouri												
All races, both sexes	3	5	4	6	3	10	9	17	16	1	-	-
Male	1	3	2	2	2	3	2	4	7	1	-	-
Female	2	2	2	4	1	7	7	13	9	-	-	-
White, both sexes	1	3	4	6	2	8	9	17	13	1	-	-
Male	-	3	2	2	2	3	2	4	5	1	-	-
Female	1	-	2	4	-	5	7	13	8	-	-	-
Black, both sexes	2	1	-	-	1	2	-	-	3	-	-	-
Male	1	-	-	-	-	-	-	-	2	-	-	-
Female	1	1	-	-	1	2	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	2	1	1	3	2	5	3	2	-	-	-
Male	1	2	1	-	2	-	2	-	-	-	-	-
Female	-	-	-	1	1	2	3	3	2	-	-	-
White, both sexes	-	2	1	1	3	2	4	3	2	-	-	-
Male	-	2	1	-	2	-	2	-	-	-	-	-
Female	-	-	-	1	1	2	2	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Nebraska												
All races, both sexes	5	4	2	7	9	14	12	16	10	5	4	-
Male	4	3	1	4	7	8	8	5	3	1	1	-
Female	1	1	1	3	2	6	4	11	7	4	3	-
White, both sexes	4	4	2	7	9	14	12	16	10	5	4	-
Male	3	3	1	4	7	8	8	5	3	1	1	-
Female	1	1	1	3	2	6	4	11	7	4	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	3	2	3	2	7	2	5	6	3	-	-
Male	-	2	1	-	1	1	1	1	1	1	-	-
Female	-	1	1	3	1	6	1	4	5	2	-	-
White, both sexes	-	1	2	2	2	7	2	5	5	3	-	-
Male	-	1	1	-	1	1	1	1	1	1	-	-
Female	-	-	1	2	1	6	1	4	4	2	-	-
Black, both sexes	-	1	-	1	-	-	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
New Hampshire												
All races, both sexes	-	-	-	-	-	2	3	1	6	-	-	-
Male	-	-	-	-	-	1	2	-	2	-	-	-
Female	-	-	-	-	-	1	1	1	4	-	-	-
White, both sexes	-	-	-	-	-	2	3	1	6	-	-	-
Male	-	-	-	-	-	1	2	-	2	-	-	-
Female	-	-	-	-	-	1	1	1	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	2	10	2	7	10	16	27	21	11	2	-
Male	2	1	4	1	4	4	6	10	6	3	-	-
Female	-	1	6	1	3	6	10	17	15	8	2	-
White, both sexes	-	1	9	2	6	7	13	27	19	10	2	-
Male	-	-	4	1	4	4	5	10	5	3	-	-
Female	-	1	5	1	2	3	8	17	14	7	2	-
Black, both sexes	2	1	1	-	1	3	3	-	2	1	-	-
Male	2	1	-	-	-	-	1	-	1	-	-	-
Female	-	-	1	-	1	3	2	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
New Mexico												
All races, both sexes	-	-	-	2	1	2	2	3	2	-	-	-
Male	-	-	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	1	1	1	1	2	2	-	-	-
White, both sexes	-	-	-	2	1	2	-	3	2	-	-	-
Male	-	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	1	1	1	-	2	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	5	7	11	10	13	25	37	40	22	14	4	-
Male	2	4	5	4	7	16	21	15	3	4	2	-
Female	3	3	6	6	6	9	16	25	19	10	2	-
White, both sexes	5	6	6	10	11	23	32	39	20	12	3	-
Male	2	3	4	4	5	16	18	15	3	4	2	-
Female	3	3	2	6	6	7	14	24	17	8	1	-
Black, both sexes	-	1	5	-	2	1	5	1	2	2	1	-
Male	-	1	1	-	2	-	3	-	-	-	-	-
Female	-	-	4	-	-	1	2	1	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
North Carolina												
All races, both sexes	2	1	6	9	11	14	26	14	10	7	-	-
Male	1	-	4	6	6	7	8	3	3	1	-	-
Female	1	1	2	3	5	7	18	11	7	6	-	-
White, both sexes	1	1	5	7	7	7	19	12	8	4	-	-
Male	1	-	3	5	4	4	7	3	3	-	-	-
Female	-	1	2	2	3	3	12	9	5	4	-	-
Black, both sexes	1	-	1	1	4	7	6	2	2	3	-	-
Male	-	-	1	-	2	3	1	-	-	1	-	-
Female	1	-	-	1	2	4	5	2	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Ohio												
All races, both sexes	2	4	6	7	6	17	21	21	11	6	1	-
Male	2	2	3	4	4	10	9	8	2	2	-	-
Female	-	2	3	3	2	7	12	13	9	4	1	-
White, both sexes	1	3	6	4	5	14	20	19	10	6	1	-
Male	1	2	3	3	4	9	9	8	2	2	-	-
Female	-	1	3	1	1	5	11	11	8	4	1	-
Black, both sexes	-	1	-	3	1	3	1	2	1	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	1	-	2	1	2	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	3	6	3	5	4	9	10	14	17	9	4	-
Male	2	3	2	4	2	3	2	3	4	3	1	-
Female	1	3	1	1	2	6	8	11	13	6	3	-
White, both sexes	2	4	3	4	3	9	8	14	17	8	4	-
Male	2	1	2	3	1	3	1	3	4	2	1	-
Female	-	3	1	1	2	6	7	11	13	6	3	-
Black, both sexes	1	2	-	1	-	-	1	-	-	-	-	-
Male	-	2	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	1	-	-
Male	-	-	-	-	1	-	1	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Oregon												
All races, both sexes	-	-	-	-	3	2	2	4	3	-	-	-
Male	-	-	-	-	3	2	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	2	3	-	-	-
White, both sexes	-	-	-	-	3	2	2	3	3	-	-	-
Male	-	-	-	-	3	2	1	2	-	-	-	-
Female	-	-	-	-	-	-	1	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	10	13	14	8	20	27	45	54	49	24	5	-
Male	7	8	12	2	10	12	23	27	17	4	3	-
Female	3	5	2	6	10	15	22	27	32	20	2	-
White, both sexes	6	10	10	7	18	20	41	52	47	23	4	-
Male	3	6	8	1	9	8	20	26	17	4	3	-
Female	3	4	2	6	9	12	21	26	30	19	1	-
Black, both sexes	4	3	4	1	2	7	4	2	2	1	1	-
Male	4	2	4	1	1	4	3	1	-	-	-	-
Female	-	1	-	-	1	3	1	1	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Rhode Island												
All races, both sexes	-	-	-	-	-	1	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	2	-	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	14	15	10	13	16	27	15	36	10	6	1	-
Male	8	11	7	8	10	14	9	11	4	-	-	-
Female	6	4	3	5	6	13	6	25	6	6	1	-
White, both sexes	9	8	8	8	9	21	10	28	9	6	-	-
Male	6	6	6	5	6	12	6	9	3	-	-	-
Female	3	2	2	3	3	9	4	19	6	6	-	-
Black, both sexes	5	7	2	5	7	6	5	8	1	-	1	-
Male	2	5	1	3	4	2	3	2	1	-	-	-
Female	3	2	1	2	3	4	2	6	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
South Dakota												
All races, both sexes	1	-	-	-	-	-	-	2	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	1	-	-	-	-	-	-	2	-	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	2	4	1	2	4	4	8	9	7	4	-	-
Male	2	4	1	1	3	3	4	4	3	2	-	-
Female	-	-	-	1	1	1	4	5	4	2	-	-
White, both sexes	1	3	1	2	4	1	5	7	5	4	-	-
Male	1	3	1	1	3	-	2	4	3	2	-	-
Female	-	-	-	1	1	1	3	3	2	2	-	-
Black, both sexes	1	1	-	-	-	3	3	2	1	-	-	-
Male	1	1	-	-	-	3	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Texas												
All races, both sexes	10	14	21	15	25	35	49	37	38	22	6	-
Male	9	8	17	11	12	16	20	15	9	8	-	-
Female	1	6	4	4	13	19	29	22	29	14	6	-
White, both sexes	7	8	16	13	23	33	47	33	37	19	6	-
Male	6	5	15	9	10	15	19	13	8	8	-	-
Female	1	3	1	4	13	18	28	20	29	11	6	-
Black, both sexes	3	6	5	2	2	2	2	4	1	2	-	-
Male	3	3	2	2	2	1	1	2	1	-	-	-
Female	-	3	3	-	-	1	1	2	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Utah												
All races, both sexes	1	-	1	-	1	4	4	5	6	2	-	-
Male	1	-	1	-	1	3	1	4	2	-	-	-
Female	-	-	-	-	-	1	3	1	4	2	-	-
White, both sexes	1	-	-	-	1	4	4	5	5	2	-	-
Male	1	-	-	-	1	3	1	4	1	-	-	-
Female	-	-	-	-	-	1	3	1	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Vermont												
All races, both sexes	-	-	-	-	-	1	-	2	4	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	1	-	1	3	-	-	-
White, both sexes	-	-	-	-	-	1	-	2	4	-	-	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	1	-	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	6	12	7	12	30	26	35	39	30	19	5	-
Male	2	6	3	7	15	13	15	15	11	3	-	-
Female	4	6	4	5	15	13	20	24	19	16	5	-
White, both sexes	2	6	4	8	22	18	27	29	26	15	5	-
Male	1	3	3	5	12	10	12	12	8	2	-	-
Female	1	3	1	3	10	8	15	17	18	13	5	-
Black, both sexes	2	5	3	4	7	7	8	10	4	4	-	-
Male	-	3	-	2	3	2	3	3	3	1	-	-
Female	2	2	3	2	4	5	5	7	1	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	1	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	1	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Washington												
All races, both sexes	1	2	1	-	1	4	10	7	8	4	1	-
Male	1	2	1	-	1	2	5	3	2	1	-	-
Female	-	-	-	-	-	2	5	4	6	3	1	-
White, both sexes	1	2	1	-	1	4	7	6	6	4	1	-
Male	1	2	1	-	1	2	4	3	1	1	-	-
Female	-	-	-	-	-	2	3	3	5	3	1	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
West Virginia												
All races, both sexes	1	3	2	2	8	7	6	16	6	4	2	-
Male	1	1	-	2	2	5	3	6	2	-	1	-
Female	-	2	2	-	6	2	3	10	4	4	1	-
White, both sexes	1	3	2	2	8	6	6	16	6	4	2	-
Male	1	1	-	2	2	4	3	6	2	-	1	-
Female	-	2	2	-	6	2	3	10	4	4	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other and unspecified heart failure (I50.1-I50.9)												
Wisconsin												
All races, both sexes	-	2	1	4	1	5	8	9	2	2	-	-
Male	-	2	-	3	1	1	4	-	1	-	-	-
Female	-	-	1	1	-	4	4	9	1	2	-	-
White, both sexes	-	2	-	4	1	4	6	9	2	2	-	-
Male	-	2	-	3	1	1	4	-	1	-	-	-
Female	-	-	-	1	-	3	2	9	1	2	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	-	2	2	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
White, both sexes	-	-	-	-	-	-	2	2	1	-	-	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
United States												
All races, both sexes	696	29	26	17	6	27	26	30	37	52	60	75
Male	354	18	13	8	3	18	14	21	20	31	30	37
Female	342	11	13	9	3	9	12	9	17	21	30	38
White, both sexes	515	16	13	7	4	15	17	20	27	40	39	56
Male	264	9	10	4	3	9	9	15	17	25	18	28
Female	251	7	3	3	1	6	8	5	10	15	21	28
Black, both sexes	159	12	10	10	1	10	8	9	7	12	17	17
Male	78	8	2	4	-	7	5	5	2	6	9	9
Female	81	4	8	6	1	3	3	4	5	6	8	8
American Indian 1/, both sexes	4	-	-	-	-	1	-	-	-	-	2	1
Male	2	-	-	-	-	1	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	18	1	3	-	1	1	1	1	3	-	2	1
Male	10	1	1	-	-	1	-	1	1	-	2	-
Female	8	-	2	-	1	-	1	-	2	-	-	1
Alabama												
All races, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	6	-	1	-	-	-	1	-	1	-	1	-
Male	5	-	1	-	-	-	1	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	5	-	1	-	-	-	-	-	1	-	1	-
Male	4	-	1	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Arkansas												
All races, both sexes	7	1	-	1	-	1	-	-	-	-	-	-
Male	3	1	-	1	-	1	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	1	-	1	-	1	-	-	-	-	-	-
Male	3	1	-	1	-	1	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	38	5	2	-	1	2	-	3	1	6	2	9
Male	25	4	2	-	1	2	-	3	1	3	1	3
Female	13	1	-	-	-	-	-	-	-	3	1	6
White, both sexes	29	3	1	-	1	1	-	3	1	5	1	7
Male	18	2	1	-	1	1	-	3	1	3	-	2
Female	11	1	-	-	-	-	-	-	-	2	1	5
Black, both sexes	4	1	-	-	-	-	-	-	-	1	-	2
Male	2	1	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	1	1	-	-	1	-	-	-	-	1	-
Male	5	1	1	-	-	1	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Colorado												
All races, both sexes	14	-	-	1	-	1	-	-	1	-	3	2
Male	8	-	-	1	-	1	-	-	1	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	14	-	-	1	-	1	-	-	1	-	3	2
Male	8	-	-	1	-	1	-	-	1	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	14	1	-	-	-	-	2	-	1	1	3	2
Male	7	-	-	-	-	-	2	-	-	1	1	2
Female	7	1	-	-	-	-	-	-	1	-	2	-
White, both sexes	12	1	-	-	-	-	2	-	-	1	3	2
Male	7	-	-	-	-	-	2	-	-	1	1	2
Female	5	1	-	-	-	-	-	-	-	-	2	-
Black, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Delaware												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Florida												
All races, both sexes	48	-	-	-	-	-	1	2	1	4	4	8
Male	29	-	-	-	-	-	1	1	-	3	2	5
Female	19	-	-	-	-	-	-	1	1	1	2	3
White, both sexes	39	-	-	-	-	-	-	2	-	4	2	7
Male	23	-	-	-	-	-	-	1	-	3	1	4
Female	16	-	-	-	-	-	-	1	-	1	1	3
Black, both sexes	7	-	-	-	-	-	1	-	1	-	2	1
Male	4	-	-	-	-	-	1	-	-	-	1	1
Female	3	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	32	1	3	1	-	2	1	-	1	1	4	2
Male	13	1	2	-	-	1	-	-	1	1	2	1
Female	19	-	1	1	-	1	1	-	-	-	2	1
White, both sexes	16	-	1	-	-	-	-	-	1	-	2	-
Male	6	-	1	-	-	-	-	-	1	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	16	1	2	1	-	2	1	-	-	1	2	2
Male	7	1	1	-	-	1	-	-	-	1	1	1
Female	9	-	1	1	-	1	1	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Hawaii												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Illinois												
All races, both sexes	32	2	2	1	-	2	-	3	3	1	1	-
Male	15	1	-	-	-	-	-	2	1	1	1	-
Female	17	1	2	1	-	2	-	1	2	-	-	-
White, both sexes	22	-	-	-	-	2	-	3	2	1	1	-
Male	13	-	-	-	-	-	-	2	1	1	1	-
Female	9	-	-	-	-	2	-	1	1	-	-	-
Black, both sexes	10	2	2	1	-	-	-	-	1	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	8	1	2	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	18	-	-	-	-	2	2	1	1	2	2	-
Male	6	-	-	-	-	2	-	1	1	-	-	-
Female	12	-	-	-	-	-	2	-	-	2	2	-
White, both sexes	16	-	-	-	-	2	2	1	1	2	2	-
Male	6	-	-	-	-	2	-	1	1	-	-	-
Female	10	-	-	-	-	-	2	-	-	2	2	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Iowa												
All races, both sexes	7	-	-	-	-	-	1	-	1	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	6	-	-	-	-	-	1	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Kansas												
All races, both sexes	9	1	-	-	-	-	-	-	-	1	-	-
Male	5	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	8	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Kentucky												
All races, both sexes	13	2	-	-	-	-	-	-	2	2	-	-
Male	7	2	-	-	-	-	-	-	2	-	-	-
Female	6	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	11	1	-	-	-	-	-	-	2	2	-	-
Male	5	1	-	-	-	-	-	-	2	-	-	-
Female	6	-	-	-	-	-	-	-	-	2	-	-
Black, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	14	1	-	1	-	-	1	-	2	1	2	1
Male	7	-	-	1	-	-	-	-	1	-	2	-
Female	7	1	-	-	-	-	1	-	1	1	-	1
White, both sexes	9	1	-	-	-	-	1	-	1	1	2	-
Male	5	-	-	-	-	-	-	-	1	-	2	-
Female	4	1	-	-	-	-	1	-	-	1	-	-
Black, both sexes	5	-	-	1	-	-	-	-	1	-	-	1
Male	2	-	-	1	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Maine												
All races, both sexes	3	-	1	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	38	1	2	1	1	2	2	2	2	4	6	10
Male	19	-	-	-	1	1	1	2	-	2	4	6
Female	19	1	2	1	-	1	1	-	2	2	2	4
White, both sexes	18	-	-	-	1	2	-	-	1	2	2	5
Male	10	-	-	-	1	1	-	-	-	1	2	3
Female	8	-	-	-	-	1	-	-	1	1	-	2
Black, both sexes	18	1	2	1	-	-	1	2	-	2	4	5
Male	9	-	-	-	-	-	1	2	-	1	2	3
Female	9	1	2	1	-	-	-	-	-	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Massachusetts												
All races, both sexes	5	-	-	-	-	-	-	-	1	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	1	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	22	1	-	2	-	1	2	1	1	-	2	-
Male	15	-	-	2	-	1	1	-	1	-	1	-
Female	7	1	-	-	-	-	1	1	-	-	1	-
White, both sexes	18	1	-	2	-	1	1	-	-	-	2	-
Male	13	-	-	2	-	1	-	-	-	-	1	-
Female	5	1	-	-	-	-	1	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	1	1	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Minnesota												
All races, both sexes	11	-	1	1	-	-	-	2	-	-	1	2
Male	4	-	1	-	-	-	-	1	-	-	-	1
Female	7	-	-	1	-	-	-	1	-	-	1	1
White, both sexes	10	-	1	1	-	-	-	1	-	-	1	2
Male	3	-	1	-	-	-	-	-	-	-	-	1
Female	7	-	-	1	-	-	-	1	-	-	1	1
Black, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	7	-	1	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	1	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	1	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Missouri												
All races, both sexes	14	-	1	-	-	2	1	-	2	-	1	1
Male	7	-	-	-	-	1	-	-	2	-	1	1
Female	7	-	1	-	-	1	1	-	-	-	-	-
White, both sexes	10	-	-	-	-	1	1	-	2	-	-	1
Male	5	-	-	-	-	-	-	-	2	-	-	1
Female	5	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes	3	-	-	-	-	1	-	-	-	-	1	-
Male	2	-	-	-	-	1	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Nebraska												
All races, both sexes	3	1	-	-	-	-	-	-	1	1	-	-
Male	2	1	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	-	-	-	-	-	1	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
New Hampshire												
All races, both sexes	2	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	2	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	17	1	1	-	-	-	-	1	-	-	4	2
Male	7	1	1	-	-	-	-	1	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	4	1
White, both sexes	12	1	-	-	-	-	-	-	-	-	3	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	3	-
Black, both sexes	4	-	1	-	-	-	-	1	-	-	1	1
Male	3	-	1	-	-	-	-	1	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
New Mexico												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	15	-	1	-	-	-	1	-	1	1	2	1
Male	6	-	-	-	-	-	1	-	-	-	1	-
Female	9	-	1	-	-	-	-	-	1	1	1	1
White, both sexes	11	-	1	-	-	-	-	-	1	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	1	-	-	-	-	-	1	-	-	1
Black, both sexes	4	-	-	-	-	-	1	-	-	1	1	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
North Carolina												
All races, both sexes	25	1	3	4	-	1	-	1	1	3	3	1
Male	14	-	2	2	-	-	-	-	1	3	2	1
Female	11	1	1	2	-	1	-	1	-	-	1	-
White, both sexes	10	-	2	-	-	-	-	-	1	1	1	1
Male	7	-	2	-	-	-	-	-	1	1	1	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	1	1	4	-	1	-	1	-	2	-	-
Male	6	-	-	2	-	-	-	-	-	2	-	-
Female	7	1	1	2	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	-	-	-	1	-	1	-	-	-
Male	2	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	1	-	1	-	-	-
Male	2	-	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Ohio												
All races, both sexes	66	3	2	-	1	2	-	2	1	8	2	8
Male	30	2	2	-	-	1	-	1	1	3	-	4
Female	36	1	-	-	1	1	-	1	-	5	2	4
White, both sexes	51	1	2	-	-	1	-	1	-	4	2	7
Male	22	1	2	-	-	-	-	1	-	2	-	3
Female	29	-	-	-	-	1	-	-	-	2	2	4
Black, both sexes	15	2	-	-	1	1	-	1	1	4	-	1
Male	8	1	-	-	-	1	-	-	1	1	-	1
Female	7	1	-	-	1	-	-	1	-	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	10	-	-	-	1	-	-	-	1	1	-	3
Male	4	-	-	-	-	-	-	-	1	1	-	-
Female	6	-	-	-	1	-	-	-	-	-	-	3
White, both sexes	8	-	-	-	1	-	-	-	1	1	-	2
Male	3	-	-	-	-	-	-	-	1	1	-	-
Female	5	-	-	-	1	-	-	-	-	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Oregon												
All races, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	24	1	1	-	1	2	-	2	3	-	1	3
Male	11	-	-	-	1	2	-	2	1	-	-	-
Female	13	1	1	-	-	-	-	-	2	-	1	3
White, both sexes	18	1	-	-	1	-	-	-	3	-	1	3
Male	7	-	-	-	1	-	-	-	1	-	-	-
Female	11	1	-	-	-	-	-	-	2	-	1	3
Black, both sexes	5	-	1	-	-	2	-	1	-	-	-	-
Male	3	-	-	-	-	2	-	1	-	-	-	-
Female	2	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Rhode Island												
All races, both sexes	7	-	1	-	-	-	-	1	-	3	-	2
Male	6	-	1	-	-	-	-	1	-	3	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	7	-	1	-	-	-	-	1	-	3	-	2
Male	6	-	1	-	-	-	-	1	-	3	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	20	1	-	-	-	-	1	2	-	1	-	1
Male	9	-	-	-	-	-	-	1	-	1	-	1
Female	11	1	-	-	-	-	1	1	-	-	-	-
White, both sexes	15	1	-	-	-	-	1	2	-	1	-	1
Male	7	-	-	-	-	-	-	1	-	1	-	1
Female	8	1	-	-	-	-	1	1	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
South Dakota												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	17	1	1	-	-	1	-	1	1	1	1	3
Male	6	-	-	-	-	1	-	1	-	1	-	-
Female	11	1	1	-	-	-	-	-	1	-	1	3
White, both sexes	15	1	1	-	-	1	-	1	1	1	1	2
Male	6	-	-	-	-	1	-	1	-	1	-	-
Female	9	1	1	-	-	-	-	-	1	-	1	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Texas												
All races, both sexes	53	4	2	2	-	4	3	5	2	4	-	3
Male	28	4	1	-	-	2	2	3	2	2	-	2
Female	25	-	1	2	-	2	1	2	-	2	-	1
White, both sexes	41	3	2	2	-	2	3	4	2	4	-	2
Male	22	3	1	-	-	1	2	3	2	2	-	2
Female	19	-	1	2	-	1	1	1	-	2	-	-
Black, both sexes	12	1	-	-	-	2	-	1	-	-	-	1
Male	6	1	-	-	-	1	-	-	-	-	-	-
Female	6	-	-	-	-	1	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	10	-	-	-	-	-	-	-	2	1	3	1
Male	5	-	-	-	-	-	-	-	1	1	2	-
Female	5	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	9	-	-	-	-	-	-	-	2	1	2	1
Male	4	-	-	-	-	-	-	-	1	1	1	-
Female	5	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Vermont												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	18	-	-	2	-	-	3	1	1	2	5	1
Male	11	-	-	1	-	-	2	1	1	1	3	1
Female	7	-	-	1	-	-	1	-	-	1	2	-
White, both sexes	8	-	-	-	-	-	2	1	-	1	1	1
Male	5	-	-	-	-	-	2	1	-	-	1	1
Female	3	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	10	-	-	2	-	-	1	-	1	1	4	-
Male	6	-	-	1	-	-	-	-	1	1	2	-
Female	4	-	-	1	-	-	1	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Washington												
All races, both sexes	9	-	-	-	-	1	-	-	1	-	1	1
Male	5	-	-	-	-	1	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	1	-	1	-
White, both sexes	8	-	-	-	-	-	-	-	1	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Wisconsin												
All races, both sexes	10	-	-	-	-	-	2	-	-	-	1	2
Male	8	-	-	-	-	-	2	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	10	-	-	-	-	-	2	-	-	-	1	2
Male	8	-	-	-	-	-	2	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
United States												
All races, both sexes	49	31	31	18	23	25	36	42	29	17	10	-
Male	27	13	18	13	16	13	11	19	8	3	-	-
Female	22	18	13	5	7	12	25	23	21	14	10	-
White, both sexes	38	25	24	16	17	19	32	39	27	16	8	-
Male	19	11	14	12	12	11	10	17	8	3	-	-
Female	19	14	10	4	5	8	22	22	19	13	8	-
Black, both sexes	9	6	7	2	5	6	4	3	2	-	2	-
Male	6	2	4	1	3	2	1	2	-	-	-	-
Female	3	4	3	1	2	4	3	1	2	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	1	-	-	-	-	1	-	-
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Alabama												
All races, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Alaska												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Arkansas												
All races, both sexes	1	-	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	1	-	1	-	-
White, both sexes	1	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3	-	-	-	1	1	-	2	-	-	-	-
Male	2	-	-	-	-	1	-	2	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	1	1	-	2	-	-	-	-
Male	1	-	-	-	-	1	-	2	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Colorado												
All races, both sexes	1	-	1	1	1	-	-	-	1	-	1	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	1	-	1	-
White, both sexes	1	-	1	1	1	-	-	-	1	-	1	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	-	-	-	-	1	1	1	-	-	1	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	1	-
White, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Delaware												
All races, both sexes	1	-	-	1	-	-	2	-	-	1	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	1	-	-
White, both sexes	-	-	-	1	-	-	2	-	-	1	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Florida												
All races, both sexes	5	4	2	1	3	1	4	3	4	-	1	-
Male	4	2	1	1	3	-	2	2	2	-	-	-
Female	1	2	1	-	-	1	2	1	2	-	1	-
White, both sexes	4	4	2	1	2	1	4	1	4	-	1	-
Male	3	2	1	1	2	-	2	1	2	-	-	-
Female	1	2	1	-	-	1	2	-	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	4	1	-	1	-	-	3	3	3	1	-	-
Male	1	-	-	-	-	-	2	1	-	-	-	-
Female	3	1	-	1	-	-	1	2	3	1	-	-
White, both sexes	2	-	-	1	-	-	3	3	2	1	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	2	-	-	1	-	-	1	2	2	1	-	-
Black, both sexes	2	1	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Hawaii												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	-	-	-	-	-	-	1	-	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	1	-
White, both sexes	-	-	-	-	-	-	1	-	1	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Illinois												
All races, both sexes	1	3	3	-	2	2	2	1	3	-	-	-
Male	1	1	2	-	2	1	-	1	1	-	-	-
Female	-	2	1	-	-	1	2	-	2	-	-	-
White, both sexes	1	2	3	-	2	-	2	1	2	-	-	-
Male	1	1	2	-	2	-	-	1	1	-	-	-
Female	-	1	1	-	-	-	2	-	1	-	-	-
Black, both sexes	-	1	-	-	-	2	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	1	2	-	-	-	1	1	2	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	1	1	1	-	-	-	1	1	1	-	-	-
White, both sexes	1	1	1	-	-	-	-	1	2	-	-	-
Male	-	-	1	-	-	-	-	-	1	-	-	-
Female	1	1	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Iowa												
All races, both sexes	1	-	1	-	-	-	-	1	-	-	1	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	-	1	-	-	-	-	1	-	-	1	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	1	1	-	1	-	1	-	-	-	1	-
Male	2	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	1	-
White, both sexes	2	1	1	-	1	-	1	-	-	-	1	-
Male	2	-	1	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Kentucky												
All races, both sexes	1	-	1	-	-	1	-	2	1	1	-	-
Male	1	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	1	1	1	-	-
White, both sexes	-	-	1	-	-	1	-	2	1	1	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	1	1	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1	1	-	-	1	-	2	-	-	-	-	-
Male	1	-	-	-	1	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	1	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Maine												
All races, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1	1	1	1	-	-	-	-	-	1	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	1	-	-
White, both sexes	1	1	1	1	-	-	-	-	-	1	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Massachusetts												
All races, both sexes	-	1	1	-	1	-	-	-	-	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1	1	-	1	1	1	-	5	1	1	-	-
Male	-	1	-	1	1	1	-	4	1	-	-	-
Female	1	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	-	1	-	1	1	1	-	5	1	1	-	-
Male	-	1	-	1	1	1	-	4	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Minnesota												
All races, both sexes	1	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	1	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	1	1	1	-	-	-	-	-	-	1	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	1	-
White, both sexes	-	1	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Missouri												
All races, both sexes	-	-	-	-	2	2	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	2	1	-	1	-	-	-	-
White, both sexes	-	-	-	-	2	1	-	2	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	2	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Nebraska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
New Hampshire												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	2	1	1	-	1	1	1	1	-	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	1	1	-	-	-	-
White, both sexes	2	1	1	-	1	1	1	1	-	-	-	-
Male	1	-	1	-	1	-	-	-	-	-	-	-
Female	1	1	-	-	-	1	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
New Mexico												
All races, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1	-	1	1	-	-	-	1	1	1	2	-
Male	1	-	1	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	2	-
White, both sexes	1	-	-	1	-	-	-	1	1	1	2	-
Male	1	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	2	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
North Carolina												
All races, both sexes	1	-	-	-	1	2	2	-	1	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	1	-	-	-
White, both sexes	1	-	-	-	-	-	2	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-
Black, both sexes	-	-	-	-	1	2	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Ohio												
All races, both sexes	5	5	1	4	-	5	3	8	3	3	-	-
Male	4	4	-	2	-	-	2	4	-	-	-	-
Female	1	1	1	2	-	5	1	4	3	3	-	-
White, both sexes	4	5	1	3	-	4	3	7	3	3	-	-
Male	3	4	-	1	-	-	2	3	-	-	-	-
Female	1	1	1	2	-	4	1	4	3	3	-	-
Black, both sexes	1	-	-	1	-	1	-	1	-	-	-	-
Male	1	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1	-	-	-	1	-	-	1	-	1	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
White, both sexes	-	-	-	-	1	-	-	1	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Oregon												
All races, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1	1	1	3	1	2	-	-	1	-	-	-
Male	-	-	1	3	-	1	-	-	-	-	-	-
Female	1	1	-	-	1	1	-	-	1	-	-	-
White, both sexes	1	1	1	3	-	2	-	-	1	-	-	-
Male	-	-	1	3	-	1	-	-	-	-	-	-
Female	1	1	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Rhode Island												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	3	4	-	2	-	-	3	-	2	-	-
Male	-	1	1	-	2	-	-	1	-	1	-	-
Female	-	2	3	-	-	-	-	2	-	1	-	-
White, both sexes	-	1	3	-	1	-	-	3	-	1	-	-
Male	-	-	1	-	1	-	-	1	-	1	-	-
Female	-	1	2	-	-	-	-	2	-	-	-	-
Black, both sexes	-	2	1	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
South Dakota												
All races, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	-	2	-	-	-	-	1	2	2	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	1	-	-	-	-	-	1	2	-	-	-
White, both sexes	-	1	-	-	-	-	1	2	2	-	-	-
Male	-	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Texas												
All races, both sexes	7	2	4	-	3	1	3	1	1	1	1	-
Male	4	1	2	-	1	-	1	-	-	1	-	-
Female	3	1	2	-	2	1	2	1	1	-	1	-
White, both sexes	6	1	2	-	1	1	2	1	1	1	1	-
Male	3	-	1	-	-	-	1	-	-	1	-	-
Female	3	1	1	-	1	1	1	1	1	-	1	-
Black, both sexes	1	1	2	-	2	-	1	-	-	-	-	-
Male	1	1	1	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Vermont												
All races, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Washington												
All races, both sexes	-	-	1	1	-	1	1	1	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	1	1	-	1	1	1	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)												
Wisconsin												
All races, both sexes	1	-	1	-	-	1	1	-	1	-	-	-
Male	1	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	1	-	-	1	1	-	1	-	-	-
Male	1	-	-	-	-	1	1	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	1	-	-	-	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
White, both sexes	1	1	-	-	-	-	2	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
United States												
All races, both sexes	10,480	35	6	5	11	33	60	87	134	166	291	422
Male	5,293	25	4	1	7	27	50	69	99	124	204	301
Female	5,187	10	2	4	4	6	10	18	35	42	87	121
White, both sexes	8,886	25	4	3	7	18	46	62	85	119	217	322
Male	4,459	17	2	1	5	16	37	50	64	91	160	233
Female	4,427	8	2	2	2	2	9	12	21	28	57	89
Black, both sexes	1,382	8	1	1	3	14	12	24	42	40	70	87
Male	721	7	1	-	1	11	11	18	30	29	40	58
Female	661	1	-	1	2	3	1	6	12	11	30	29
American Indian 1/, both sexes	63	1	-	-	-	1	1	1	2	5	1	5
Male	36	-	-	-	-	-	1	1	2	4	1	3
Female	27	1	-	-	-	1	-	-	-	1	-	2
Asian or Pacific Islander 2/, both sexes	149	1	1	1	1	-	1	-	5	2	3	8
Male	77	1	1	-	1	-	1	-	3	-	3	7
Female	72	-	-	1	-	-	-	-	2	2	-	1
Alabama												
All races, both sexes	319	1	-	-	1	2	-	-	1	3	6	11
Male	170	1	-	-	1	2	-	-	1	1	4	9
Female	149	-	-	-	-	-	-	-	-	2	2	2
White, both sexes	241	1	-	-	-	-	-	-	-	1	3	7
Male	131	1	-	-	-	-	-	-	-	-	2	5
Female	110	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	76	-	-	-	1	2	-	-	1	2	3	3
Male	37	-	-	-	1	2	-	-	1	1	2	3
Female	39	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Alaska												
All races, both sexes	19	1	-	-	-	-	-	-	-	1	1	1
Male	15	-	-	-	-	-	-	-	-	1	1	-
Female	4	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	13	-	-	-	-	-	-	-	-	1	1	1
Male	12	-	-	-	-	-	-	-	-	1	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	164	-	-	-	1	1	-	1	5	6	14	7
Male	95	-	-	-	-	1	-	1	4	6	10	4
Female	69	-	-	-	1	-	-	-	1	-	4	3
White, both sexes	145	-	-	-	1	1	-	1	3	4	12	5
Male	84	-	-	-	-	1	-	1	3	4	10	3
Female	61	-	-	-	1	-	-	-	-	-	2	2
Black, both sexes	8	-	-	-	-	-	-	-	1	-	2	2
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	1	-	2	1
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	1	2	-	-
Male	9	-	-	-	-	-	-	-	1	2	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Arkansas												
All races, both sexes	174	-	-	-	-	-	-	-	1	1	2	8
Male	88	-	-	-	-	-	-	-	1	-	1	5
Female	86	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	143	-	-	-	-	-	-	-	1	1	-	6
Male	72	-	-	-	-	-	-	-	1	-	-	4
Female	71	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	30	-	-	-	-	-	-	-	-	-	2	2
Male	16	-	-	-	-	-	-	-	-	-	1	1
Female	14	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	879	4	-	1	1	3	6	3	17	11	25	41
Male	407	4	-	-	1	3	4	3	11	11	19	32
Female	472	-	-	1	-	-	2	-	6	-	6	9
White, both sexes	751	4	-	-	-	3	4	2	11	10	16	31
Male	342	4	-	-	-	3	2	2	8	10	13	24
Female	409	-	-	-	-	-	2	-	3	-	3	7
Black, both sexes	73	-	-	-	-	-	1	1	4	-	8	6
Male	33	-	-	-	-	-	1	1	2	-	5	4
Female	40	-	-	-	-	-	-	-	2	-	3	2
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	1	1	-	1
Male	4	-	-	-	-	-	-	-	1	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	-	-	1	1	-	1	-	1	-	1	3
Male	28	-	-	-	1	-	1	-	-	-	1	3
Female	21	-	-	1	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Colorado												
All races, both sexes	100	-	-	-	-	-	-	-	2	4	4	5
Male	52	-	-	-	-	-	-	-	2	2	2	3
Female	48	-	-	-	-	-	-	-	-	2	2	2
White, both sexes	89	-	-	-	-	-	-	-	2	3	4	4
Male	45	-	-	-	-	-	-	-	2	2	2	3
Female	44	-	-	-	-	-	-	-	-	1	2	1
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	101	1	-	-	-	1	2	-	2	2	5	5
Male	49	1	-	-	-	-	2	-	2	2	3	3
Female	52	-	-	-	-	1	-	-	-	-	2	2
White, both sexes	90	1	-	-	-	-	2	-	1	2	5	4
Male	43	1	-	-	-	-	2	-	1	2	3	2
Female	47	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	11	-	-	-	-	1	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	1	-	-	1
Female	5	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Delaware												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	1	1
Male	10	-	-	-	-	-	-	-	-	-	1	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Florida												
All races, both sexes	817	1	1	-	-	1	2	3	3	6	13	13
Male	392	-	1	-	-	-	2	3	1	4	9	10
Female	425	1	-	-	-	1	-	-	2	2	4	3
White, both sexes	736	1	-	-	-	-	2	2	3	4	10	10
Male	355	-	-	-	-	-	2	2	1	3	9	9
Female	381	1	-	-	-	-	-	-	2	1	1	1
Black, both sexes	75	-	-	-	-	1	-	1	-	1	3	3
Male	33	-	-	-	-	-	-	1	-	1	-	1
Female	42	-	-	-	-	1	-	-	-	-	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	1	-	-	-	-	-	-	1	-	-
Male	4	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Georgia												
All races, both sexes	783	1	-	-	3	1	3	10	10	11	19	27
Male	446	1	-	-	2	1	3	7	8	7	15	20
Female	337	-	-	-	1	-	-	3	2	4	4	7
White, both sexes	578	1	-	-	2	1	1	5	4	5	13	18
Male	333	1	-	-	2	1	1	4	3	2	11	13
Female	245	-	-	-	-	-	-	1	1	3	2	5
Black, both sexes	194	-	-	-	1	-	2	5	5	6	5	9
Male	106	-	-	-	-	-	2	3	4	5	3	7
Female	88	-	-	-	1	-	-	2	1	1	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	1	-	1	-
Male	7	-	-	-	-	-	-	-	1	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Hawaii												
All races, both sexes	51	-	-	-	-	-	-	1	2	1	2	7
Male	27	-	-	-	-	-	-	1	1	-	2	4
Female	24	-	-	-	-	-	-	-	1	1	-	3
White, both sexes	17	-	-	-	-	-	-	1	-	-	1	2
Male	9	-	-	-	-	-	-	1	-	-	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	32	-	-	-	-	-	-	-	2	1	1	4
Male	17	-	-	-	-	-	-	-	1	-	1	3
Female	15	-	-	-	-	-	-	-	1	1	-	1
Idaho												
All races, both sexes	27	-	-	-	-	-	-	-	2	-	2	2
Male	17	-	-	-	-	-	-	-	1	-	2	2
Female	10	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	27	-	-	-	-	-	-	-	2	-	2	2
Male	17	-	-	-	-	-	-	-	1	-	2	2
Female	10	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Illinois												
All races, both sexes	435	2	-	-	-	-	2	3	3	2	6	12
Male	197	-	-	-	-	-	2	2	2	1	4	11
Female	238	2	-	-	-	-	-	1	1	1	2	1
White, both sexes	369	2	-	-	-	-	2	2	1	2	6	10
Male	170	-	-	-	-	-	2	2	-	1	4	9
Female	199	2	-	-	-	-	-	-	1	1	2	1
Black, both sexes	60	-	-	-	-	-	-	1	2	-	-	2
Male	23	-	-	-	-	-	-	-	2	-	-	2
Female	37	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	154	-	1	1	-	2	-	-	3	5	4	6
Male	82	-	1	1	-	2	-	-	3	4	3	3
Female	72	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	146	-	1	1	-	1	-	-	3	5	2	3
Male	77	-	1	1	-	1	-	-	3	4	2	1
Female	69	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	8	-	-	-	-	1	-	-	-	-	2	3
Male	5	-	-	-	-	1	-	-	-	-	1	2
Female	3	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Iowa												
All races, both sexes	52	-	-	-	-	-	-	2	1	3	1	3
Male	22	-	-	-	-	-	-	2	-	2	1	2
Female	30	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	50	-	-	-	-	-	-	1	1	2	1	3
Male	21	-	-	-	-	-	-	1	-	2	1	2
Female	29	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	2	-	-	-	-	-	-	1	-	1	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	91	-	-	-	-	1	1	2	-	-	3	10
Male	47	-	-	-	-	1	1	2	-	-	1	6
Female	44	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	85	-	-	-	-	-	1	2	-	-	3	6
Male	43	-	-	-	-	-	1	2	-	-	1	4
Female	42	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	5	-	-	-	-	1	-	-	-	-	-	3
Male	4	-	-	-	-	1	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Kentucky												
All races, both sexes	237	-	-	-	-	-	2	2	2	5	8	9
Male	146	-	-	-	-	-	1	1	1	5	7	8
Female	91	-	-	-	-	-	1	1	1	-	1	1
White, both sexes	229	-	-	-	-	-	2	2	2	4	8	9
Male	141	-	-	-	-	-	1	1	1	4	7	8
Female	88	-	-	-	-	-	1	1	1	-	1	1
Black, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	274	3	1	-	-	2	1	3	2	4	7	10
Male	125	1	1	-	-	2	1	3	1	2	5	4
Female	149	2	-	-	-	-	-	-	1	2	2	6
White, both sexes	210	2	1	-	-	1	-	2	2	1	4	7
Male	98	1	1	-	-	1	-	2	1	1	3	4
Female	112	1	-	-	-	-	-	-	1	-	1	3
Black, both sexes	64	1	-	-	-	1	1	1	-	3	3	3
Male	27	-	-	-	-	1	1	1	-	1	2	-
Female	37	1	-	-	-	-	-	-	-	2	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Maine												
All races, both sexes	109	-	-	-	-	-	1	2	-	3	2	7
Male	56	-	-	-	-	-	1	1	-	3	-	6
Female	53	-	-	-	-	-	-	1	-	-	2	1
White, both sexes	109	-	-	-	-	-	1	2	-	3	2	7
Male	56	-	-	-	-	-	1	1	-	3	-	6
Female	53	-	-	-	-	-	-	1	-	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	121	1	-	-	-	-	2	4	4	3	13	15
Male	67	1	-	-	-	-	2	3	3	2	11	11
Female	54	-	-	-	-	-	-	1	1	1	2	4
White, both sexes	83	-	-	-	-	-	1	2	1	-	8	10
Male	43	-	-	-	-	-	1	1	1	-	6	7
Female	40	-	-	-	-	-	-	1	-	-	2	3
Black, both sexes	36	-	-	-	-	-	1	2	3	3	5	5
Male	23	-	-	-	-	-	1	2	2	2	5	4
Female	13	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Massachusetts												
All races, both sexes	171	-	-	-	1	-	1	3	-	3	3	4
Male	84	-	-	-	1	-	1	3	-	3	2	3
Female	87	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	160	-	-	-	1	-	1	2	-	2	2	4
Male	77	-	-	-	1	-	1	2	-	2	2	3
Female	83	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	10	-	-	-	-	-	-	1	-	1	1	-
Male	7	-	-	-	-	-	-	1	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	293	1	-	1	-	-	3	1	2	4	6	7
Male	138	1	-	-	-	-	2	1	1	4	6	5
Female	155	-	-	1	-	-	1	-	1	-	-	2
White, both sexes	254	1	-	-	-	-	3	1	1	2	6	6
Male	123	1	-	-	-	-	2	1	1	2	6	5
Female	131	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	37	-	-	1	-	-	-	-	1	2	-	1
Male	14	-	-	-	-	-	-	-	-	2	-	-
Female	23	-	-	1	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Minnesota												
All races, both sexes	114	-	-	-	-	1	-	2	2	1	1	4
Male	57	-	-	-	-	-	-	2	2	1	-	4
Female	57	-	-	-	-	1	-	-	-	-	1	-
White, both sexes	113	-	-	-	-	1	-	2	2	1	1	4
Male	56	-	-	-	-	-	-	2	2	1	-	4
Female	57	-	-	-	-	1	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	276	1	-	-	-	-	-	1	2	4	5	9
Male	137	1	-	-	-	-	-	-	1	3	2	5
Female	139	-	-	-	-	-	-	1	1	1	3	4
White, both sexes	181	-	-	-	-	-	-	1	1	2	3	6
Male	90	-	-	-	-	-	-	-	-	1	1	3
Female	91	-	-	-	-	-	-	1	1	1	2	3
Black, both sexes	95	1	-	-	-	-	-	-	1	2	2	3
Male	47	1	-	-	-	-	-	-	1	2	1	2
Female	48	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Missouri												
All races, both sexes	228	-	-	-	-	1	2	3	4	3	5	7
Male	129	-	-	-	-	1	2	3	3	3	3	5
Female	99	-	-	-	-	-	-	-	1	-	2	2
White, both sexes	204	-	-	-	-	-	2	2	4	2	4	7
Male	116	-	-	-	-	-	2	2	3	2	2	5
Female	88	-	-	-	-	-	-	-	1	-	2	2
Black, both sexes	20	-	-	-	-	1	-	1	-	1	1	-
Male	12	-	-	-	-	1	-	1	-	1	1	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	1	3
Male	21	-	-	-	-	-	-	-	-	-	1	2
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	32	-	-	-	-	-	-	-	-	-	-	3
Male	18	-	-	-	-	-	-	-	-	-	-	2
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Nebraska												
All races, both sexes	75	1	-	-	-	-	1	-	1	-	1	-
Male	39	-	-	-	-	-	-	-	1	-	1	-
Female	36	1	-	-	-	-	1	-	-	-	-	-
White, both sexes	74	1	-	-	-	-	1	-	1	-	1	-
Male	39	-	-	-	-	-	-	-	1	-	1	-
Female	35	1	-	-	-	-	1	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	89	-	-	-	-	-	-	-	-	-	2	-
Male	37	-	-	-	-	-	-	-	-	-	1	-
Female	52	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	78	-	-	-	-	-	-	-	-	-	2	-
Male	32	-	-	-	-	-	-	-	-	-	1	-
Female	46	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
New Hampshire												
All races, both sexes	73	-	-	-	-	1	1	-	-	1	2	1
Male	34	-	-	-	-	-	1	-	-	1	2	1
Female	39	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	73	-	-	-	-	1	1	-	-	1	2	1
Male	34	-	-	-	-	-	1	-	-	1	2	1
Female	39	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	225	1	-	-	-	1	4	5	3	1	7	6
Male	117	-	-	-	-	1	4	5	3	1	5	4
Female	108	1	-	-	-	-	-	-	-	-	2	2
White, both sexes	187	1	-	-	-	1	4	3	1	-	6	6
Male	99	-	-	-	-	1	4	3	1	-	4	4
Female	88	1	-	-	-	-	-	-	-	-	2	2
Black, both sexes	32	-	-	-	-	-	-	2	2	1	1	-
Male	14	-	-	-	-	-	-	2	2	1	1	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
New Mexico												
All races, both sexes	60	-	-	-	-	-	-	-	-	2	1	4
Male	32	-	-	-	-	-	-	-	-	2	-	4
Female	28	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	53	-	-	-	-	-	-	-	-	1	1	4
Male	28	-	-	-	-	-	-	-	-	1	-	4
Female	25	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	401	2	-	-	-	-	2	6	7	9	13	28
Male	193	2	-	-	-	-	2	5	5	7	7	23
Female	208	-	-	-	-	-	-	1	2	2	6	5
White, both sexes	350	1	-	-	-	-	2	4	2	7	9	23
Male	173	1	-	-	-	-	2	4	2	6	6	20
Female	177	-	-	-	-	-	-	-	-	1	3	3
Black, both sexes	46	1	-	-	-	-	-	2	5	2	4	5
Male	20	1	-	-	-	-	-	1	3	1	1	3
Female	26	-	-	-	-	-	-	1	2	1	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
North Carolina												
All races, both sexes	307	1	1	-	-	1	4	3	8	3	7	11
Male	145	1	-	-	-	-	3	2	7	3	4	8
Female	162	-	1	-	-	1	1	1	1	-	3	3
White, both sexes	220	-	1	-	-	-	-	1	4	2	5	4
Male	97	-	-	-	-	-	-	1	4	2	2	3
Female	123	-	1	-	-	-	-	-	-	-	3	1
Black, both sexes	79	1	-	-	-	1	4	2	3	1	2	6
Male	44	1	-	-	-	-	3	1	2	1	2	4
Female	35	-	-	-	-	1	1	1	1	-	-	2
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	15	-	-	-	-	-	1	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	15	-	-	-	-	-	1	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Ohio												
All races, both sexes	373	1	-	1	-	1	2	2	4	7	15	17
Male	194	1	-	-	-	1	1	1	2	4	12	8
Female	179	-	-	1	-	-	1	1	2	3	3	9
White, both sexes	340	-	-	1	-	-	2	1	4	6	13	17
Male	174	-	-	-	-	-	1	1	2	4	11	8
Female	166	-	-	1	-	-	1	-	2	2	2	9
Black, both sexes	33	1	-	-	-	1	-	1	-	1	2	-
Male	20	1	-	-	-	1	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	107	1	-	-	-	-	2	1	2	2	4	3
Male	55	1	-	-	-	-	1	1	2	1	4	2
Female	52	-	-	-	-	-	1	-	-	1	-	1
White, both sexes	92	1	-	-	-	-	1	-	2	1	3	2
Male	45	1	-	-	-	-	-	-	2	1	3	1
Female	47	-	-	-	-	-	1	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	1	1	-	1	-	1
Male	6	-	-	-	-	-	1	1	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Oregon												
All races, both sexes	83	-	-	-	-	-	-	-	1	1	3	2
Male	38	-	-	-	-	-	-	-	-	-	1	2
Female	45	-	-	-	-	-	-	-	1	1	2	-
White, both sexes	82	-	-	-	-	-	-	-	1	1	3	2
Male	38	-	-	-	-	-	-	-	-	-	1	2
Female	44	-	-	-	-	-	-	-	1	1	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	458	3	-	-	-	2	-	1	1	4	9	19
Male	226	2	-	-	-	2	-	1	1	2	4	12
Female	232	1	-	-	-	-	-	-	-	2	5	7
White, both sexes	423	3	-	-	-	2	-	1	-	2	8	16
Male	210	2	-	-	-	2	-	1	-	1	3	10
Female	213	1	-	-	-	-	-	-	-	1	5	6
Black, both sexes	32	-	-	-	-	-	-	-	1	2	1	3
Male	16	-	-	-	-	-	-	-	1	1	1	2
Female	16	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Rhode Island												
All races, both sexes	36	-	-	-	1	-	-	-	1	-	4	2
Male	16	-	-	-	1	-	-	-	1	-	2	1
Female	20	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	32	-	-	-	1	-	-	-	1	-	2	2
Male	14	-	-	-	1	-	-	-	1	-	1	1
Female	18	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	327	1	-	-	-	2	2	4	11	12	22	25
Male	192	1	-	-	-	2	2	4	9	8	15	17
Female	135	-	-	-	-	-	-	-	2	4	7	8
White, both sexes	221	1	-	-	-	1	2	2	4	7	14	16
Male	124	1	-	-	-	1	2	2	4	4	12	11
Female	97	-	-	-	-	-	-	-	-	3	2	5
Black, both sexes	106	-	-	-	-	1	-	2	7	5	8	9
Male	68	-	-	-	-	1	-	2	5	4	3	6
Female	38	-	-	-	-	-	-	-	2	1	5	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
South Dakota												
All races, both sexes	36	1	-	-	-	-	-	-	-	-	-	3
Male	21	-	-	-	-	-	-	-	-	-	-	2
Female	15	1	-	-	-	-	-	-	-	-	-	1
White, both sexes	33	1	-	-	-	-	-	-	-	-	-	2
Male	21	-	-	-	-	-	-	-	-	-	-	2
Female	12	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	214	-	1	-	-	1	2	5	7	7	8	12
Male	107	-	-	-	-	1	2	3	7	5	6	9
Female	107	-	1	-	-	-	-	2	-	2	2	3
White, both sexes	185	-	1	-	-	1	2	5	6	7	6	9
Male	89	-	-	-	-	1	2	3	6	5	4	7
Female	96	-	1	-	-	-	-	2	-	2	2	2
Black, both sexes	28	-	-	-	-	-	-	-	1	-	2	3
Male	18	-	-	-	-	-	-	-	1	-	2	2
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Texas												
All races, both sexes	834	5	-	1	3	5	7	10	16	23	23	38
Male	401	5	-	-	1	5	7	6	12	17	16	27
Female	433	-	-	1	2	-	-	4	4	6	7	11
White, both sexes	724	3	-	1	2	2	5	9	12	20	15	30
Male	340	3	-	-	1	2	5	5	9	14	12	20
Female	384	-	-	1	1	-	-	4	3	6	3	10
Black, both sexes	103	2	-	-	1	3	2	1	4	3	8	8
Male	58	2	-	-	-	3	2	1	3	3	4	7
Female	45	-	-	-	1	-	-	-	1	-	4	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	166	-	-	-	-	-	-	-	-	2	1	-
Male	69	-	-	-	-	-	-	-	-	1	1	-
Female	97	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	163	-	-	-	-	-	-	-	-	2	1	-
Male	67	-	-	-	-	-	-	-	-	1	1	-
Female	96	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Vermont												
All races, both sexes	13	-	-	-	-	-	1	-	-	1	-	-
Male	7	-	-	-	-	-	1	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	1	-	-	1	-	-
Male	7	-	-	-	-	-	1	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	237	-	1	-	-	1	1	1	-	2	1	7
Male	125	-	1	-	-	1	1	1	-	1	1	5
Female	112	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	176	-	-	-	-	1	-	1	-	1	-	4
Male	89	-	-	-	-	1	-	1	-	-	-	3
Female	87	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	59	-	1	-	-	-	1	-	-	1	1	3
Male	35	-	1	-	-	-	1	-	-	1	1	2
Female	24	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Washington												
All races, both sexes	104	-	-	-	-	2	-	-	2	1	1	4
Male	51	-	-	-	-	1	-	-	-	1	1	3
Female	53	-	-	-	-	1	-	-	2	-	-	1
White, both sexes	95	-	-	-	-	1	-	-	2	1	-	4
Male	49	-	-	-	-	1	-	-	-	1	-	3
Female	46	-	-	-	-	-	-	-	2	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	50	-	-	-	-	-	1	1	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	31	-	-	-	-	-	1	1	-	-	1	-
White, both sexes	47	-	-	-	-	-	1	1	-	-	1	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	28	-	-	-	-	-	1	1	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Wisconsin												
All races, both sexes	115	1	-	-	-	-	1	2	-	2	6	4
Male	71	1	-	-	-	-	1	2	-	2	5	3
Female	44	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	110	-	-	-	-	-	1	2	-	2	6	3
Male	67	-	-	-	-	-	1	2	-	2	5	3
Female	43	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	1	-	-	-	-	-	-	-	-	-	1
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
United States												
All races, both sexes	482	550	574	612	820	1,137	1,528	1,650	1,256	497	122	2
Male	336	358	369	407	478	609	671	630	397	106	19	2
Female	146	192	205	205	342	528	857	1,020	859	391	103	-
White, both sexes	361	423	454	493	698	976	1,364	1,509	1,143	443	112	2
Male	259	270	303	337	413	531	610	575	371	96	16	2
Female	102	153	151	156	285	445	754	934	772	347	96	-
Black, both sexes	114	114	99	105	105	143	141	115	92	43	9	-
Male	71	79	55	64	57	70	52	41	18	6	2	-
Female	43	35	44	41	48	73	89	74	74	37	7	-
American Indian 1/, both sexes	4	5	4	2	8	6	4	6	3	3	1	-
Male	3	4	1	-	5	2	2	4	2	-	1	-
Female	1	1	3	2	3	4	2	2	1	3	-	-
Asian or Pacific Islander 2/, both sexes	3	8	17	12	9	12	19	20	18	8	-	-
Male	3	5	10	6	3	6	7	10	6	4	-	-
Female	-	3	7	6	6	6	12	10	12	4	-	-
Alabama												
All races, both sexes	17	23	32	29	32	36	53	34	28	9	1	-
Male	8	18	17	21	17	22	23	14	9	1	1	-
Female	9	5	15	8	15	14	30	20	19	8	-	-
White, both sexes	12	19	27	22	25	26	42	29	22	4	1	-
Male	5	14	16	17	14	18	18	13	7	-	1	-
Female	7	5	11	5	11	8	24	16	15	4	-	-
Black, both sexes	5	4	5	7	7	10	10	5	6	5	-	-
Male	3	4	1	4	3	4	4	1	2	1	-	-
Female	2	-	4	3	4	6	6	4	4	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Alaska												
All races, both sexes	2	1	2	4	2	1	1	2	-	-	-	-
Male	2	1	2	4	2	-	-	2	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	1	1	1	3	2	-	-	2	-	-	-	-
Male	1	1	1	3	2	-	-	2	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	13	11	12	11	12	13	13	19	17	8	-	-
Male	10	6	11	8	7	7	6	7	6	1	-	-
Female	3	5	1	3	5	6	7	12	11	7	-	-
White, both sexes	11	9	11	11	8	12	13	18	17	8	-	-
Male	8	6	11	8	3	7	6	6	6	1	-	-
Female	3	3	-	3	5	5	7	12	11	7	-	-
Black, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	1	-	3	-	-	1	-	-	-	-
Male	2	-	-	-	3	-	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Arkansas												
All races, both sexes	4	13	8	12	17	18	25	34	19	9	3	-
Male	3	10	4	10	9	10	16	9	8	1	1	-
Female	1	3	4	2	8	8	9	25	11	8	2	-
White, both sexes	3	10	5	9	15	17	19	30	18	7	2	-
Male	2	7	3	8	9	9	13	7	8	1	-	-
Female	1	3	2	1	6	8	6	23	10	6	2	-
Black, both sexes	1	3	3	3	1	1	6	4	1	2	1	-
Male	1	3	1	2	-	1	3	2	-	-	1	-
Female	-	-	2	1	1	-	3	2	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
California												
All races, both sexes	34	51	30	28	47	82	136	160	130	57	12	-
Male	17	27	20	21	21	41	53	54	46	18	1	-
Female	17	24	10	7	26	41	83	106	84	39	11	-
White, both sexes	24	43	23	24	37	66	120	149	119	53	12	-
Male	14	22	15	18	16	32	48	49	44	17	1	-
Female	10	21	8	6	21	34	72	100	75	36	11	-
Black, both sexes	9	4	3	2	6	10	6	5	6	2	-	-
Male	2	2	2	2	4	6	-	2	-	-	-	-
Female	7	2	1	-	2	4	6	3	6	2	-	-
American Indian 1/, both sexes	-	2	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	4	2	4	5	10	6	5	2	-	-
Male	1	2	3	1	1	3	5	3	2	1	-	-
Female	-	-	1	1	3	2	5	3	3	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Colorado												
All races, both sexes	5	1	9	10	9	11	13	15	12	-	-	-
Male	4	1	6	6	2	7	8	5	4	-	-	-
Female	1	-	3	4	7	4	5	10	8	-	-	-
White, both sexes	3	1	9	8	8	10	12	14	11	-	-	-
Male	2	1	6	4	1	6	7	5	4	-	-	-
Female	1	-	3	4	7	4	5	9	7	-	-	-
Black, both sexes	2	-	-	1	-	-	1	1	-	-	-	-
Male	2	-	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	1	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Connecticut												
All races, both sexes	8	5	6	4	7	14	12	16	6	5	-	-
Male	2	3	3	4	3	8	5	6	2	-	-	-
Female	6	2	3	-	4	6	7	10	4	5	-	-
White, both sexes	6	5	5	3	5	14	10	16	6	5	-	-
Male	2	3	3	3	2	8	3	6	2	-	-	-
Female	4	2	2	-	3	6	7	10	4	5	-	-
Black, both sexes	2	-	1	1	2	-	2	-	-	-	-	-
Male	-	-	-	1	1	-	2	-	-	-	-	-
Female	2	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Delaware												
All races, both sexes	-	2	1	1	-	2	8	5	3	2	-	-
Male	-	1	1	1	-	2	6	-	-	-	-	-
Female	-	1	-	-	-	-	2	5	3	2	-	-
White, both sexes	-	1	1	1	-	1	6	5	3	2	-	-
Male	-	1	1	1	-	1	4	-	-	-	-	-
Female	-	-	-	-	-	-	2	5	3	2	-	-
Black, both sexes	-	1	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	1	-	-	1	-	3	1	2	1	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	2	1	2	1	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	1	-	-	1	-	2	1	2	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	2	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Florida												
All races, both sexes	13	16	29	44	55	103	142	176	129	55	12	-
Male	9	9	19	29	37	49	64	79	53	11	3	-
Female	4	7	10	15	18	54	78	97	76	44	9	-
White, both sexes	9	8	26	38	48	86	129	171	123	54	12	-
Male	6	5	17	25	33	42	58	77	52	11	3	-
Female	3	3	9	13	15	44	71	94	71	43	9	-
Black, both sexes	4	7	3	5	7	17	12	5	5	1	-	-
Male	3	3	2	4	4	7	5	2	-	-	-	-
Female	1	4	1	1	3	10	7	3	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	60	67	72	73	79	91	90	90	53	20	2	1
Male	46	45	55	47	49	46	36	39	13	4	1	1
Female	14	22	17	26	30	45	54	51	40	16	1	-
White, both sexes	43	51	47	47	60	73	80	72	41	13	-	1
Male	33	35	37	31	38	39	35	31	12	3	-	1
Female	10	16	10	16	22	34	45	41	29	10	-	-
Black, both sexes	17	15	24	24	17	17	9	17	12	7	2	-
Male	13	9	17	15	10	6	1	8	1	1	1	-
Female	4	6	7	9	7	11	8	9	11	6	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	1	1	1	1	-	-	-	-
Male	-	1	1	1	1	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Hawaii												
All races, both sexes	4	1	6	5	1	1	4	7	5	4	-	-
Male	3	1	4	3	1	-	1	3	1	2	-	-
Female	1	-	2	2	-	1	3	4	4	2	-	-
White, both sexes	4	-	1	2	-	1	3	1	-	1	-	-
Male	3	-	1	2	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	2	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	5	3	1	-	1	5	5	3	-	-
Male	-	1	3	1	1	-	-	3	1	2	-	-
Female	-	-	2	2	-	-	1	2	4	1	-	-
Idaho												
All races, both sexes	1	-	-	4	4	1	4	3	2	2	-	-
Male	1	-	-	2	4	1	1	1	1	1	-	-
Female	-	-	-	2	-	-	3	2	1	1	-	-
White, both sexes	1	-	-	4	4	1	4	3	2	2	-	-
Male	1	-	-	2	4	1	1	1	1	1	-	-
Female	-	-	-	2	-	-	3	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Illinois												
All races, both sexes	21	18	33	24	26	49	67	78	58	25	6	-
Male	14	11	18	12	11	26	30	31	15	7	-	-
Female	7	7	15	12	15	23	37	47	43	18	6	-
White, both sexes	18	12	26	16	20	41	59	70	52	25	5	-
Male	12	9	15	9	9	21	28	27	15	7	-	-
Female	6	3	11	7	11	20	31	43	37	18	5	-
Black, both sexes	2	6	7	7	6	7	7	6	6	-	1	-
Male	1	2	3	2	2	4	2	3	-	-	-	-
Female	1	4	4	5	4	3	5	3	6	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	1	1	2	-	-	-	-
Male	1	-	-	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Indiana												
All races, both sexes	7	5	8	8	18	16	27	18	13	6	6	-
Male	3	3	5	4	11	12	17	4	5	1	-	-
Female	4	2	3	4	7	4	10	14	8	5	6	-
White, both sexes	7	5	8	8	17	15	27	18	13	6	6	-
Male	3	3	5	4	10	12	17	4	5	1	-	-
Female	4	2	3	4	7	3	10	14	8	5	6	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Iowa												
All races, both sexes	2	-	1	2	2	5	12	4	9	3	2	-
Male	2	-	-	2	1	2	3	2	3	-	-	-
Female	-	-	1	-	1	3	9	2	6	3	2	-
White, both sexes	2	-	1	2	2	5	12	4	9	3	2	-
Male	2	-	-	2	1	2	3	2	3	-	-	-
Female	-	-	1	-	1	3	9	2	6	3	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	4	2	6	3	-	16	12	15	10	4	2	-
Male	3	1	1	3	-	10	7	5	5	1	-	-
Female	1	1	5	-	-	6	5	10	5	3	2	-
White, both sexes	4	2	5	3	-	16	12	15	10	4	2	-
Male	3	1	-	3	-	10	7	5	5	1	-	-
Female	1	1	5	-	-	6	5	10	5	3	2	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Kentucky												
All races, both sexes	22	18	16	20	25	30	26	27	20	5	-	-
Male	19	13	11	15	13	17	14	13	7	1	-	-
Female	3	5	5	5	12	13	12	14	13	4	-	-
White, both sexes	22	16	16	20	25	30	26	24	18	5	-	-
Male	19	11	11	15	13	17	14	11	7	1	-	-
Female	3	5	5	5	12	13	12	13	11	4	-	-
Black, both sexes	-	2	-	-	-	-	-	2	2	-	-	-
Male	-	2	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	15	11	12	14	15	34	40	48	31	16	5	-
Male	9	7	9	9	8	15	17	18	9	2	2	-
Female	6	4	3	5	7	19	23	30	22	14	3	-
White, both sexes	7	7	9	12	12	25	31	41	28	14	4	-
Male	5	3	9	8	8	11	11	17	8	2	2	-
Female	2	4	-	4	4	14	20	24	20	12	2	-
Black, both sexes	8	4	3	2	3	9	9	7	3	2	1	-
Male	4	4	-	1	-	4	6	1	1	-	-	-
Female	4	-	3	1	3	5	3	6	2	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Maine												
All races, both sexes	7	6	12	8	10	7	15	12	12	4	1	-
Male	5	4	7	5	6	4	5	7	2	-	-	-
Female	2	2	5	3	4	3	10	5	10	4	1	-
White, both sexes	7	6	12	8	10	7	15	12	12	4	1	-
Male	5	4	7	5	6	4	5	7	2	-	-	-
Female	2	2	5	3	4	3	10	5	10	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	4	8	5	10	12	9	12	6	4	3	-
Male	4	2	4	4	8	7	2	2	-	1	-	-
Female	2	2	4	1	2	5	7	10	6	3	3	-
White, both sexes	5	2	6	4	7	10	7	10	3	4	3	-
Male	3	1	4	3	6	6	2	1	-	1	-	-
Female	2	1	2	1	1	4	5	9	3	3	3	-
Black, both sexes	1	1	2	1	3	2	2	2	3	-	-	-
Male	1	1	-	1	2	1	-	1	-	-	-	-
Female	-	-	2	-	1	1	2	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Massachusetts												
All races, both sexes	4	7	6	4	12	19	31	29	32	11	1	-
Male	3	5	6	2	10	7	16	11	10	1	-	-
Female	1	2	-	2	2	12	15	18	22	10	1	-
White, both sexes	4	5	6	3	11	19	29	28	31	11	1	-
Male	3	3	6	2	9	7	14	11	10	1	-	-
Female	1	2	-	1	2	12	15	17	21	10	1	-
Black, both sexes	-	2	-	-	1	-	2	1	1	-	-	-
Male	-	2	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	7	18	9	22	24	33	46	48	44	15	2	-
Male	5	11	6	12	14	19	17	15	16	3	-	-
Female	2	7	3	10	10	14	29	33	28	12	2	-
White, both sexes	4	14	6	18	20	28	41	47	40	15	1	-
Male	4	10	5	10	11	18	16	15	13	3	-	-
Female	-	4	1	8	9	10	25	32	27	12	1	-
Black, both sexes	3	3	2	4	4	5	5	1	4	-	1	-
Male	1	-	1	2	3	1	1	-	3	-	-	-
Female	2	3	1	2	1	4	4	1	1	-	1	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Minnesota												
All races, both sexes	3	11	4	8	8	13	13	15	18	9	1	-
Male	3	6	2	6	3	7	10	3	7	1	-	-
Female	-	5	2	2	5	6	3	12	11	8	1	-
White, both sexes	3	11	3	8	8	13	13	15	18	9	1	-
Male	3	6	1	6	3	7	10	3	7	1	-	-
Female	-	5	2	2	5	6	3	12	11	8	1	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	7	16	16	22	23	42	41	35	38	12	2	-
Male	6	12	13	14	14	23	15	15	9	4	-	-
Female	1	4	3	8	9	19	26	20	29	8	2	-
White, both sexes	3	9	11	12	14	28	28	24	28	10	1	-
Male	3	7	9	8	9	16	11	11	7	4	-	-
Female	-	2	2	4	5	12	17	13	21	6	1	-
Black, both sexes	4	7	5	10	9	14	13	11	10	2	1	-
Male	3	5	4	6	5	7	4	4	2	-	-	-
Female	1	2	1	4	4	7	9	7	8	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Missouri												
All races, both sexes	12	12	12	13	24	24	37	38	23	7	1	-
Male	10	8	7	9	17	15	18	15	9	1	-	-
Female	2	4	5	4	7	9	19	23	14	6	1	-
White, both sexes	10	11	11	13	21	22	34	34	20	6	1	-
Male	9	8	7	9	15	14	17	13	8	-	-	-
Female	1	3	4	4	6	8	17	21	12	6	1	-
Black, both sexes	2	1	-	-	1	2	3	3	3	1	-	-
Male	1	-	-	-	1	1	1	2	1	1	-	-
Female	1	1	-	-	-	1	2	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	1	-	-	-	-
Montana												
All races, both sexes	3	-	2	3	2	2	7	7	3	2	-	-
Male	3	-	2	2	1	1	5	3	1	-	-	-
Female	-	-	-	1	1	1	2	4	2	2	-	-
White, both sexes	3	-	1	3	2	2	6	7	3	2	-	-
Male	3	-	1	2	1	1	4	3	1	-	-	-
Female	-	-	-	1	1	1	2	4	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Nebraska												
All races, both sexes	2	3	5	4	6	7	11	16	12	2	3	-
Male	2	2	4	3	4	5	4	7	5	-	1	-
Female	-	1	1	1	2	2	7	9	7	2	2	-
White, both sexes	2	3	5	4	6	7	11	15	12	2	3	-
Male	2	2	4	3	4	5	4	7	5	-	1	-
Female	-	1	1	1	2	2	7	8	7	2	2	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	7	3	5	9	13	19	13	13	4	-	-
Male	-	3	2	4	3	9	8	4	2	1	-	-
Female	1	4	1	1	6	4	11	9	11	3	-	-
White, both sexes	1	6	3	5	7	11	16	12	13	2	-	-
Male	-	2	2	4	2	8	7	3	2	1	-	-
Female	1	4	1	1	5	3	9	9	11	1	-	-
Black, both sexes	-	1	-	-	2	1	2	-	-	1	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	1	-	1	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
New Hampshire												
All races, both sexes	3	-	1	2	10	12	8	13	13	4	1	-
Male	2	-	1	1	6	6	3	6	3	1	-	-
Female	1	-	-	1	4	6	5	7	10	3	1	-
White, both sexes	3	-	1	2	10	12	8	13	13	4	1	-
Male	2	-	1	1	6	6	3	6	3	1	-	-
Female	1	-	-	1	4	6	5	7	10	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	8	10	6	9	19	27	31	36	32	15	3	1
Male	6	8	3	6	12	11	15	17	9	6	-	1
Female	2	2	3	3	7	16	16	19	23	9	3	-
White, both sexes	6	7	3	7	18	24	26	33	26	11	3	1
Male	5	5	1	5	12	9	15	16	8	5	-	1
Female	1	2	2	2	6	15	11	17	18	6	3	-
Black, both sexes	2	3	2	1	1	3	4	3	4	3	-	-
Male	1	3	1	-	-	2	-	1	-	-	-	-
Female	1	-	1	1	1	1	4	2	4	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	1	-	2	1	-	-
Male	-	-	1	1	-	-	-	-	1	1	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
New Mexico												
All races, both sexes	3	3	-	2	6	6	10	8	10	3	2	-
Male	2	3	-	2	3	4	3	3	5	-	1	-
Female	1	-	-	-	3	2	7	5	5	3	1	-
White, both sexes	2	2	-	2	5	6	10	8	9	2	1	-
Male	2	2	-	2	3	4	3	3	4	-	-	-
Female	-	-	-	-	2	2	7	5	5	2	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	1	-	-	-	1	1	1	-
Male	-	1	-	-	-	-	-	-	1	-	1	-
Female	1	-	-	-	1	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	17	19	14	17	29	43	39	76	54	19	7	-
Male	12	13	8	12	17	18	14	32	10	4	2	-
Female	5	6	6	5	12	25	25	44	44	15	5	-
White, both sexes	15	15	11	16	24	37	37	71	51	18	7	-
Male	12	12	6	11	14	16	14	31	10	4	2	-
Female	3	3	5	5	10	21	23	40	41	14	5	-
Black, both sexes	2	4	2	1	5	5	2	4	1	1	-	-
Male	-	1	2	1	3	2	-	1	-	-	-	-
Female	2	3	-	-	2	3	2	3	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
North Carolina												
All races, both sexes	10	20	15	21	18	32	43	59	35	15	-	-
Male	7	15	6	14	9	16	16	21	12	1	-	-
Female	3	5	9	7	9	16	27	38	23	14	-	-
White, both sexes	7	12	7	15	13	23	36	50	29	11	-	-
Male	5	7	3	11	5	12	15	17	9	1	-	-
Female	2	5	4	4	8	11	21	33	20	10	-	-
Black, both sexes	3	8	7	6	5	6	7	9	5	3	-	-
Male	2	8	3	3	4	2	1	4	3	-	-	-
Female	1	-	4	3	1	4	6	5	2	3	-	-
American Indian 1/, both sexes	-	-	1	-	-	3	-	-	1	1	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	1	-	1	3	3	2	2	1	-	-
Male	1	-	1	-	1	2	3	-	1	1	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-
White, both sexes	1	-	1	-	1	3	3	2	2	1	-	-
Male	1	-	1	-	1	2	3	-	1	1	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Ohio												
All races, both sexes	15	24	24	19	32	47	49	61	35	14	3	-
Male	11	19	14	14	22	27	24	18	14	1	-	-
Female	4	5	10	5	10	20	25	43	21	13	3	-
White, both sexes	12	19	21	16	28	44	48	59	33	14	2	-
Male	8	15	13	12	19	25	23	17	14	1	-	-
Female	4	4	8	4	9	19	25	42	19	13	2	-
Black, both sexes	3	5	3	3	4	3	1	2	2	-	1	-
Male	3	4	1	2	3	2	1	1	-	-	-	-
Female	-	1	2	1	1	1	-	1	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	4	-	3	9	9	27	20	12	6	-	-
Male	-	2	-	2	4	6	15	10	2	2	-	-
Female	2	2	-	1	5	3	12	10	10	4	-	-
White, both sexes	2	3	-	3	8	7	25	17	11	6	-	-
Male	-	1	-	2	4	5	13	8	2	2	-	-
Female	2	2	-	1	4	2	12	9	9	4	-	-
Black, both sexes	-	-	-	-	-	1	2	1	1	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	1	-	2	-	-	-	-
Male	-	1	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Oregon												
All races, both sexes	2	2	5	3	3	9	18	14	13	4	3	-
Male	2	-	2	2	2	5	11	4	6	-	1	-
Female	-	2	3	1	1	4	7	10	7	4	2	-
White, both sexes	2	2	5	3	3	9	18	14	13	3	3	-
Male	2	-	2	2	2	5	11	4	6	-	1	-
Female	-	2	3	1	1	4	7	10	7	3	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
Pennsylvania												
All races, both sexes	18	17	23	25	38	41	84	68	73	26	6	-
Male	15	9	18	15	28	23	35	31	21	5	2	-
Female	3	8	5	10	10	18	49	37	52	21	4	-
White, both sexes	16	14	20	22	38	39	78	64	70	24	6	-
Male	13	8	17	13	28	23	32	29	21	5	2	-
Female	3	6	3	9	10	16	46	35	49	19	4	-
Black, both sexes	2	2	3	3	-	1	6	4	3	1	-	-
Male	2	1	1	2	-	-	3	2	-	-	-	-
Female	-	1	2	1	-	1	3	2	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Rhode Island												
All races, both sexes	3	3	4	-	1	3	3	6	4	-	1	-
Male	1	2	3	-	-	2	2	-	1	-	-	-
Female	2	1	1	-	1	1	1	6	3	-	1	-
White, both sexes	3	2	3	-	1	3	3	6	4	-	1	-
Male	1	2	2	-	-	2	2	-	1	-	-	-
Female	2	-	1	-	1	1	1	6	3	-	1	-
Black, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	37	33	23	23	19	26	32	24	24	6	1	-
Male	25	23	13	18	12	19	11	5	6	2	-	-
Female	12	10	10	5	7	7	21	19	18	4	1	-
White, both sexes	18	24	17	16	10	20	22	20	22	4	1	-
Male	12	14	8	13	7	15	8	4	5	1	-	-
Female	6	10	9	3	3	5	14	16	17	3	1	-
Black, both sexes	19	9	6	7	9	6	10	4	2	2	-	-
Male	13	9	5	5	5	4	3	1	1	1	-	-
Female	6	-	1	2	4	2	7	3	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
South Dakota												
All races, both sexes	-	1	3	2	6	2	7	5	5	1	-	-
Male	-	-	3	1	5	2	3	3	1	1	-	-
Female	-	1	-	1	1	-	4	2	4	-	-	-
White, both sexes	-	1	3	1	6	2	7	4	5	1	-	-
Male	-	-	3	1	5	2	3	3	1	1	-	-
Female	-	1	-	-	1	-	4	1	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	6	15	12	9	25	21	28	20	21	9	5	-
Male	5	8	6	4	13	10	11	8	7	2	-	-
Female	1	7	6	5	12	11	17	12	14	7	5	-
White, both sexes	2	11	9	7	24	20	27	17	18	8	5	-
Male	2	6	3	3	13	9	11	6	6	2	-	-
Female	-	5	6	4	11	11	16	11	12	6	5	-
Black, both sexes	4	4	3	1	1	1	1	3	3	1	-	-
Male	3	2	3	1	-	1	-	2	1	-	-	-
Female	1	2	-	-	1	-	1	1	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Texas												
All races, both sexes	44	38	46	33	73	70	124	115	102	44	14	-
Male	32	25	28	20	37	35	54	35	28	10	1	-
Female	12	13	18	13	36	35	70	80	74	34	13	-
White, both sexes	30	27	40	24	65	64	112	111	97	42	13	-
Male	23	17	26	16	36	32	49	35	25	9	1	-
Female	7	10	14	8	29	32	63	76	72	33	12	-
Black, both sexes	13	11	5	9	7	6	9	4	4	2	1	-
Male	8	8	1	4	1	3	5	-	2	1	-	-
Female	5	3	4	5	6	3	4	4	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	1	-	3	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	3	-	-	-	-	-
Utah												
All races, both sexes	1	4	3	8	9	22	36	58	15	7	-	-
Male	1	1	-	4	5	12	14	21	8	1	-	-
Female	-	3	3	4	4	10	22	37	7	6	-	-
White, both sexes	1	4	3	8	9	21	36	57	15	6	-	-
Male	1	1	-	4	5	11	14	20	8	1	-	-
Female	-	3	3	4	4	10	22	37	7	5	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Vermont												
All races, both sexes	-	-	-	-	1	-	5	1	3	-	1	-
Male	-	-	-	-	1	-	2	1	1	-	-	-
Female	-	-	-	-	-	-	3	-	2	-	1	-
White, both sexes	-	-	-	-	1	-	5	1	3	-	1	-
Male	-	-	-	-	1	-	2	1	1	-	-	-
Female	-	-	-	-	-	-	3	-	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	12	13	21	23	19	28	28	41	23	11	4	-
Male	8	10	13	17	11	17	12	17	6	2	1	-
Female	4	3	8	6	8	11	16	24	17	9	3	-
White, both sexes	7	9	14	18	17	19	20	34	19	8	4	-
Male	4	6	10	13	9	12	9	15	4	1	1	-
Female	3	3	4	5	8	7	11	19	15	7	3	-
Black, both sexes	5	4	6	5	2	9	8	7	3	3	-	-
Male	4	4	3	4	2	5	3	2	1	1	-	-
Female	1	-	3	1	-	4	5	5	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Washington												
All races, both sexes	4	1	2	6	9	19	15	16	12	8	2	-
Male	3	-	-	4	6	8	7	12	2	3	-	-
Female	1	1	2	2	3	11	8	4	10	5	2	-
White, both sexes	4	1	1	6	9	18	14	14	10	8	2	-
Male	3	-	-	4	6	8	7	11	2	3	-	-
Female	1	1	1	2	3	10	7	3	8	5	2	-
Black, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	2	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	1	-	-	1	-	1	1	-	-	-
West Virginia												
All races, both sexes	2	4	2	4	4	5	9	9	5	1	1	-
Male	1	3	2	2	3	3	3	1	-	-	-	-
Female	1	1	-	2	1	2	6	8	5	1	1	-
White, both sexes	2	4	2	3	4	5	9	8	5	-	1	-
Male	1	3	2	2	3	3	3	1	-	-	-	-
Female	1	1	-	1	1	2	6	7	5	-	1	-
Black, both sexes	-	-	-	1	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and ill-defined forms of heart disease (I51.0-I51.3,I51.6-I51.9)												
Wisconsin												
All races, both sexes	6	8	6	9	9	13	13	19	13	1	2	-
Male	5	5	6	4	5	9	6	10	5	1	1	-
Female	1	3	-	5	4	4	7	9	8	-	1	-
White, both sexes	6	7	6	9	9	11	13	19	13	1	2	-
Male	5	4	6	4	5	7	6	10	5	1	1	-
Female	1	3	-	5	4	4	7	9	8	-	1	-
Black, both sexes	-	1	-	-	-	2	-	-	-	-	-	-
Male	-	1	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	2	2	4	1	3	2	-	3	-	-	-
Male	1	2	2	3	-	1	-	-	1	-	-	-
Female	2	-	-	1	1	2	2	-	2	-	-	-
White, both sexes	3	2	2	4	1	3	2	-	3	-	-	-
Male	1	2	2	3	-	1	-	-	1	-	-	-
Female	2	-	-	1	1	2	2	-	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
United States												
All races, both sexes	143,579	126	62	52	43	76	120	192	354	693	1,567	2,629
Male	56,586	73	41	27	28	34	60	109	192	355	793	1,345
Female	86,993	53	21	25	15	42	60	83	162	338	774	1,284
White, both sexes	121,868	79	45	35	29	51	85	118	238	461	989	1,674
Male	47,194	46	28	17	18	21	44	68	129	241	513	880
Female	74,674	33	17	18	11	30	41	50	109	220	476	794
Black, both sexes	17,541	40	14	14	10	19	32	64	99	188	501	847
Male	7,519	23	10	9	6	8	16	36	53	93	239	413
Female	10,022	17	4	5	4	11	16	28	46	95	262	434
American Indian 1/, both sexes	627	2	-	-	-	1	1	1	3	13	19	21
Male	249	1	-	-	-	1	-	1	1	3	13	8
Female	378	1	-	-	-	-	1	-	2	10	6	13
Asian or Pacific Islander 2/, both sexes	3,543	5	3	3	4	5	2	9	14	31	58	87
Male	1,624	3	3	1	4	4	-	4	9	18	28	44
Female	1,919	2	-	2	-	1	2	5	5	13	30	43
Alabama												
All races, both sexes	2,952	4	-	-	1	1	3	7	13	18	44	77
Male	1,133	2	-	-	-	1	2	5	9	6	24	38
Female	1,819	2	-	-	1	-	1	2	4	12	20	39
White, both sexes	2,135	-	-	-	1	-	3	5	5	14	28	30
Male	800	-	-	-	-	-	2	4	4	4	16	12
Female	1,335	-	-	-	1	-	1	1	1	10	12	18
Black, both sexes	810	4	-	-	-	1	-	2	7	4	16	47
Male	328	2	-	-	-	1	-	1	4	2	8	26
Female	482	2	-	-	-	-	-	1	3	2	8	21
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Alaska												
All races, both sexes	178	-	-	-	-	-	1	-	-	2	5	2
Male	82	-	-	-	-	-	-	-	-	-	2	-
Female	96	-	-	-	-	-	1	-	-	2	3	2
White, both sexes	112	-	-	-	-	-	1	-	-	1	1	1
Male	54	-	-	-	-	-	-	-	-	-	1	-
Female	58	-	-	-	-	-	1	-	-	1	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	51	-	-	-	-	-	-	-	-	1	4	1
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	30	-	-	-	-	-	-	-	-	1	3	1
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2,364	2	1	2	-	2	1	5	4	13	28	38
Male	967	1	1	-	-	2	-	2	1	6	15	15
Female	1,397	1	-	2	-	-	1	3	3	7	13	23
White, both sexes	2,220	2	1	2	-	2	1	4	3	10	25	29
Male	912	1	1	-	-	2	-	2	-	4	13	14
Female	1,308	1	-	2	-	-	1	2	3	6	12	15
Black, both sexes	68	-	-	-	-	-	-	-	1	2	2	6
Male	25	-	-	-	-	-	-	-	1	2	1	1
Female	43	-	-	-	-	-	-	-	-	-	1	5
American Indian 1/, both sexes	48	-	-	-	-	-	-	-	-	1	1	2
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	1	-	2
Asian or Pacific Islander 2/, both sexes	28	-	-	-	-	-	-	1	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	1	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Arkansas												
All races, both sexes	1,847	1	-	-	1	-	1	2	5	10	23	36
Male	724	1	-	-	-	-	-	1	1	3	13	25
Female	1,123	-	-	-	1	-	1	1	4	7	10	11
White, both sexes	1,530	1	-	-	1	-	1	-	2	7	13	26
Male	593	1	-	-	-	-	-	-	-	2	8	20
Female	937	-	-	-	1	-	1	-	2	5	5	6
Black, both sexes	306	-	-	-	-	-	-	2	3	3	10	10
Male	129	-	-	-	-	-	-	1	1	1	5	5
Female	177	-	-	-	-	-	-	1	2	2	5	5
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	15,585	9	7	4	2	11	12	23	31	89	180	295
Male	6,286	6	6	2	2	5	8	15	13	46	102	158
Female	9,299	3	1	2	-	6	4	8	18	43	78	137
White, both sexes	12,748	8	4	4	1	9	11	16	22	55	121	217
Male	5,058	5	3	2	1	4	8	11	10	32	72	119
Female	7,690	3	1	2	-	5	3	5	12	23	49	98
Black, both sexes	1,176	-	1	-	1	1	-	4	4	21	37	46
Male	480	-	1	-	1	-	-	1	1	7	17	22
Female	696	-	-	-	-	1	-	3	3	14	20	24
American Indian 1/, both sexes	80	-	-	-	-	1	-	-	1	1	4	3
Male	31	-	-	-	-	1	-	-	-	-	3	-
Female	49	-	-	-	-	-	-	-	1	1	1	3
Asian or Pacific Islander 2/, both sexes	1,581	1	2	-	-	-	1	3	4	12	18	29
Male	717	1	2	-	-	-	-	3	2	7	10	17
Female	864	-	-	-	-	-	1	-	2	5	8	12

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Colorado												
All races, both sexes	1,599	1	2	2	-	-	3	4	5	9	19	25
Male	658	-	2	1	-	-	1	1	2	6	7	15
Female	941	1	-	1	-	-	2	3	3	3	12	10
White, both sexes	1,516	1	2	2	-	-	3	4	5	6	18	21
Male	621	-	2	1	-	-	1	1	2	4	7	12
Female	895	1	-	1	-	-	2	3	3	2	11	9
Black, both sexes	59	-	-	-	-	-	-	-	-	-	1	3
Male	27	-	-	-	-	-	-	-	-	-	-	2
Female	32	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	3	-	-
Male	9	-	-	-	-	-	-	-	-	2	-	-
Female	12	-	-	-	-	-	-	-	-	1	-	-
Connecticut												
All races, both sexes	1,528	2	-	3	1	1	2	2	5	6	10	18
Male	614	1	-	2	-	1	1	1	3	3	4	10
Female	914	1	-	1	1	-	1	1	2	3	6	8
White, both sexes	1,403	1	-	3	-	1	1	1	3	5	7	13
Male	557	1	-	2	-	1	1	-	2	2	3	7
Female	846	-	-	1	-	-	-	1	1	3	4	6
Black, both sexes	114	1	-	-	1	-	1	-	2	1	3	5
Male	51	-	-	-	-	-	-	-	1	1	1	3
Female	63	1	-	-	1	-	1	-	1	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	1	-	-	-	-
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Delaware												
All races, both sexes	384	-	1	-	-	-	-	1	-	2	1	4
Male	166	-	-	-	-	-	-	1	-	1	-	2
Female	218	-	1	-	-	-	-	-	-	1	1	2
White, both sexes	311	-	1	-	-	-	-	1	-	1	1	2
Male	136	-	-	-	-	-	-	1	-	1	-	1
Female	175	-	1	-	-	-	-	-	-	-	1	1
Black, both sexes	69	-	-	-	-	-	-	-	-	1	-	2
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	42	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	231	-	-	-	-	-	-	-	1	5	4	11
Male	95	-	-	-	-	-	-	-	-	3	2	8
Female	136	-	-	-	-	-	-	-	1	2	2	3
White, both sexes	53	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	175	-	-	-	-	-	-	-	1	5	4	11
Male	70	-	-	-	-	-	-	-	-	3	2	8
Female	105	-	-	-	-	-	-	-	1	2	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Florida												
All races, both sexes	9,361	10	4	3	2	11	10	8	25	42	92	158
Male	4,002	4	3	2	1	2	5	5	16	22	50	85
Female	5,359	6	1	1	1	9	5	3	9	20	42	73
White, both sexes	8,020	8	2	1	2	6	6	3	15	27	58	106
Male	3,411	4	2	1	1	1	2	-	8	17	31	61
Female	4,609	4	-	-	1	5	4	3	7	10	27	45
Black, both sexes	1,254	2	2	2	-	3	4	5	10	14	31	50
Male	557	-	1	1	-	-	3	5	8	5	17	23
Female	697	2	1	1	-	3	1	-	2	9	14	27
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	76	-	-	-	-	2	-	-	-	1	3	2
Male	29	-	-	-	-	1	-	-	-	-	2	1
Female	47	-	-	-	-	1	-	-	-	1	1	1
Georgia												
All races, both sexes	3,854	7	4	3	2	3	3	8	12	29	69	111
Male	1,526	6	3	1	1	1	2	4	6	14	35	53
Female	2,328	1	1	2	1	2	1	4	6	15	34	58
White, both sexes	2,686	4	2	2	1	2	1	4	4	18	40	55
Male	1,021	3	1	1	1	-	1	1	1	10	21	27
Female	1,665	1	1	1	-	2	-	3	3	8	19	28
Black, both sexes	1,129	3	2	1	1	1	2	4	8	9	29	55
Male	490	3	2	-	-	1	1	3	5	4	14	25
Female	639	-	-	1	1	-	1	1	3	5	15	30
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	-	-	-	1	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	18	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Hawaii												
All races, both sexes	688	1	1	1	1	-	-	-	-	2	10	17
Male	321	-	1	-	1	-	-	-	-	2	3	9
Female	367	1	-	1	-	-	-	-	-	-	7	8
White, both sexes	135	-	-	-	-	-	-	-	-	-	1	4
Male	56	-	-	-	-	-	-	-	-	-	-	3
Female	79	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	553	1	1	1	1	-	-	-	-	2	9	13
Male	265	-	1	-	1	-	-	-	-	2	3	6
Female	288	1	-	1	-	-	-	-	-	-	6	7
Idaho												
All races, both sexes	718	-	1	-	-	-	-	-	1	2	5	5
Male	293	-	1	-	-	-	-	-	1	1	3	2
Female	425	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	710	-	1	-	-	-	-	-	1	2	5	5
Male	289	-	1	-	-	-	-	-	1	1	3	2
Female	421	-	-	-	-	-	-	-	-	1	2	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Illinois												
All races, both sexes	6,252	11	4	4	3	2	5	3	12	26	57	95
Male	2,413	7	1	1	3	1	4	1	5	18	31	46
Female	3,839	4	3	3	-	1	1	2	7	8	26	49
White, both sexes	5,223	8	4	3	1	1	3	3	7	12	38	49
Male	1,963	4	1	-	1	-	2	1	3	7	21	23
Female	3,260	4	3	3	-	1	1	2	4	5	17	26
Black, both sexes	930	3	-	1	2	1	2	-	5	13	18	45
Male	405	3	-	1	2	1	2	-	2	11	9	22
Female	525	-	-	-	-	-	-	-	3	2	9	23
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	94	-	-	-	-	-	-	-	-	1	1	1
Male	43	-	-	-	-	-	-	-	-	-	1	1
Female	51	-	-	-	-	-	-	-	-	1	-	-
Indiana												
All races, both sexes	3,296	4	3	-	1	-	3	5	7	16	40	56
Male	1,266	-	2	-	1	-	2	3	3	6	15	16
Female	2,030	4	1	-	-	-	1	2	4	10	25	40
White, both sexes	2,988	3	3	-	1	-	2	4	4	10	30	43
Male	1,128	-	2	-	1	-	1	2	1	4	10	12
Female	1,860	3	1	-	-	-	1	2	3	6	20	31
Black, both sexes	300	1	-	-	-	-	1	1	3	6	9	13
Male	133	-	-	-	-	-	1	1	2	2	4	4
Female	167	1	-	-	-	-	-	-	1	4	5	9
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Iowa												
All races, both sexes	1,902	1	-	1	-	2	3	2	5	3	13	23
Male	733	1	-	-	-	-	-	1	3	1	3	11
Female	1,169	-	-	1	-	2	3	1	2	2	10	12
White, both sexes	1,878	1	-	1	-	2	2	2	5	3	13	23
Male	720	1	-	-	-	-	-	1	3	1	3	11
Female	1,158	-	-	1	-	2	2	1	2	2	10	12
Black, both sexes	18	-	-	-	-	-	1	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1,571	3	-	-	-	-	1	2	3	3	10	27
Male	607	3	-	-	-	-	-	-	2	3	5	12
Female	964	-	-	-	-	-	1	2	1	-	5	15
White, both sexes	1,466	2	-	-	-	-	1	1	3	3	8	21
Male	559	2	-	-	-	-	-	-	2	3	3	8
Female	907	-	-	-	-	-	1	1	1	-	5	13
Black, both sexes	82	1	-	-	-	-	-	1	-	-	-	5
Male	42	1	-	-	-	-	-	-	-	-	-	4
Female	40	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	2	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Kentucky												
All races, both sexes	2,168	3	-	-	3	-	2	2	3	8	24	40
Male	821	3	-	-	3	-	1	1	2	5	8	18
Female	1,347	-	-	-	-	-	1	1	1	3	16	22
White, both sexes	1,998	2	-	-	2	-	2	2	3	7	18	28
Male	754	2	-	-	2	-	1	1	2	5	6	12
Female	1,244	-	-	-	-	-	1	1	1	2	12	16
Black, both sexes	166	1	-	-	1	-	-	-	-	1	6	12
Male	65	1	-	-	1	-	-	-	-	-	2	6
Female	101	-	-	-	-	-	-	-	-	1	4	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2,469	4	3	1	2	3	3	7	7	8	41	76
Male	962	3	1	-	-	1	-	4	5	4	18	40
Female	1,507	1	2	1	2	2	3	3	2	4	23	36
White, both sexes	1,591	2	3	-	1	1	3	2	6	4	11	18
Male	601	1	1	-	-	-	-	2	4	1	6	11
Female	990	1	2	-	1	1	3	-	2	3	5	7
Black, both sexes	862	2	-	1	1	2	-	5	1	4	30	57
Male	352	2	-	-	-	1	-	2	1	3	12	29
Female	510	-	-	1	1	1	-	3	-	1	18	28
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Maine												
All races, both sexes	693	-	-	-	-	-	-	-	1	2	2	10
Male	262	-	-	-	-	-	-	-	1	1	1	4
Female	431	-	-	-	-	-	-	-	-	1	1	6
White, both sexes	692	-	-	-	-	-	-	-	1	2	2	10
Male	262	-	-	-	-	-	-	-	1	1	1	4
Female	430	-	-	-	-	-	-	-	-	1	1	6
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	2,476	4	2	-	-	1	2	2	2	10	37	48
Male	977	1	-	-	-	1	1	2	1	5	17	25
Female	1,499	3	2	-	-	-	1	-	1	5	20	23
White, both sexes	1,809	3	1	-	-	-	1	-	1	3	9	15
Male	686	1	-	-	-	-	-	-	-	2	5	10
Female	1,123	2	1	-	-	-	1	-	1	1	4	5
Black, both sexes	616	-	1	-	-	1	1	2	1	7	25	32
Male	269	-	-	-	-	1	1	2	1	3	11	14
Female	347	-	1	-	-	-	-	-	-	4	14	18
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	1	-	-	-	-	-	-	-	-	3	1
Male	21	-	-	-	-	-	-	-	-	-	1	1
Female	28	1	-	-	-	-	-	-	-	-	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Massachusetts												
All races, both sexes	2,977	1	1	1	1	-	2	3	10	8	21	35
Male	1,096	1	1	-	1	-	1	2	8	3	12	20
Female	1,881	-	-	1	-	-	1	1	2	5	9	15
White, both sexes	2,814	1	1	1	-	-	2	2	8	7	17	28
Male	1,023	1	1	-	-	-	1	2	6	2	10	17
Female	1,791	-	-	1	-	-	1	-	2	5	7	11
Black, both sexes	119	-	-	-	1	-	-	1	1	1	3	6
Male	51	-	-	-	1	-	-	-	1	1	1	2
Female	68	-	-	-	-	-	-	1	-	-	2	4
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	40	-	-	-	-	-	-	-	1	-	-	1
Male	19	-	-	-	-	-	-	-	1	-	-	1
Female	21	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	5,057	2	1	1	1	2	2	7	10	38	43	69
Male	1,975	1	1	1	-	1	-	5	5	18	21	32
Female	3,082	1	-	-	1	1	2	2	5	20	22	37
White, both sexes	4,323	1	1	1	1	-	-	5	5	27	33	43
Male	1,662	1	1	1	-	-	-	3	3	16	18	20
Female	2,661	-	-	-	1	-	-	2	2	11	15	23
Black, both sexes	684	1	-	-	-	2	2	1	4	11	9	25
Male	294	-	-	-	-	1	-	1	2	2	3	11
Female	390	1	-	-	-	1	2	-	2	9	6	14
American Indian 1/, both sexes	25	-	-	-	-	-	-	1	-	-	-	1
Male	9	-	-	-	-	-	-	1	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	1	-	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Minnesota												
All races, both sexes	2,379	1	-	-	2	-	2	4	6	8	29	27
Male	904	-	-	-	2	-	-	2	3	5	14	13
Female	1,475	1	-	-	-	-	2	2	3	3	15	14
White, both sexes	2,282	1	-	-	1	-	2	1	3	7	20	20
Male	853	-	-	-	1	-	-	1	1	5	10	9
Female	1,429	1	-	-	-	-	2	-	2	2	10	11
Black, both sexes	39	-	-	-	-	-	-	1	2	-	5	6
Male	21	-	-	-	-	-	-	1	1	-	2	3
Female	18	-	-	-	-	-	-	-	1	-	3	3
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	1	1	-
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	44	-	-	-	1	-	-	2	1	-	3	1
Male	22	-	-	-	1	-	-	-	1	-	1	1
Female	22	-	-	-	-	-	-	2	-	-	2	-
Mississippi												
All races, both sexes	1,622	3	-	1	1	1	1	5	3	12	35	49
Male	724	3	-	1	1	-	-	3	3	6	23	17
Female	898	-	-	-	-	1	1	2	-	6	12	32
White, both sexes	1,054	1	-	-	1	-	-	3	2	2	18	20
Male	443	1	-	-	1	-	-	2	2	1	10	7
Female	611	-	-	-	-	-	-	1	-	1	8	13
Black, both sexes	556	2	-	1	-	1	1	2	1	10	17	29
Male	277	2	-	1	-	-	-	1	1	5	13	10
Female	279	-	-	-	-	1	1	1	-	5	4	19
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Missouri												
All races, both sexes	3,347	2	1	2	-	2	-	9	3	13	32	64
Male	1,256	-	1	1	-	1	-	5	1	7	16	28
Female	2,091	2	-	1	-	1	-	4	2	6	16	36
White, both sexes	2,988	1	-	1	-	2	-	7	2	8	21	44
Male	1,108	-	-	-	-	1	-	3	1	4	10	21
Female	1,880	1	-	1	-	1	-	4	1	4	11	23
Black, both sexes	341	1	1	1	-	-	-	2	1	4	11	19
Male	143	-	1	1	-	-	-	2	-	2	6	7
Female	198	1	-	-	-	-	-	-	1	2	5	12
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	1	-	1
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Montana												
All races, both sexes	522	-	-	-	-	1	-	1	2	3	2	6
Male	208	-	-	-	-	-	-	1	2	-	2	5
Female	314	-	-	-	-	1	-	-	-	3	-	1
White, both sexes	501	-	-	-	-	1	-	1	2	1	2	5
Male	203	-	-	-	-	-	-	1	2	-	2	4
Female	298	-	-	-	-	1	-	-	-	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	2	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Nebraska												
All races, both sexes	986	-	-	-	-	-	1	2	4	1	3	11
Male	408	-	-	-	-	-	1	2	3	1	-	6
Female	578	-	-	-	-	-	-	-	1	-	3	5
White, both sexes	944	-	-	-	-	-	1	1	3	1	3	8
Male	386	-	-	-	-	-	1	1	2	1	-	4
Female	558	-	-	-	-	-	-	-	1	-	3	4
Black, both sexes	27	-	-	-	-	-	-	1	1	-	-	3
Male	20	-	-	-	-	-	-	1	1	-	-	2
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	945	-	2	1	-	2	-	3	7	9	13	31
Male	439	-	1	1	-	1	-	3	4	5	9	14
Female	506	-	1	-	-	1	-	-	3	4	4	17
White, both sexes	795	-	2	1	-	1	-	2	4	7	8	18
Male	362	-	1	1	-	-	-	2	2	3	5	7
Female	433	-	1	-	-	1	-	-	2	4	3	11
Black, both sexes	84	-	-	-	-	-	-	1	1	2	5	10
Male	44	-	-	-	-	-	-	1	1	2	4	6
Female	40	-	-	-	-	-	-	-	-	-	1	4
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	63	-	-	-	-	1	-	-	2	-	-	3
Male	32	-	-	-	-	1	-	-	1	-	-	1
Female	31	-	-	-	-	-	-	-	1	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
New Hampshire												
All races, both sexes	497	-	-	-	-	1	-	-	-	2	5	5
Male	167	-	-	-	-	1	-	-	-	1	3	2
Female	330	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	492	-	-	-	-	1	-	-	-	2	5	4
Male	166	-	-	-	-	1	-	-	-	1	3	2
Female	326	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
New Jersey												
All races, both sexes	3,614	2	4	-	2	-	3	4	8	15	33	63
Male	1,423	1	3	-	1	-	1	2	4	9	23	38
Female	2,191	1	1	-	1	-	2	2	4	6	10	25
White, both sexes	3,005	2	3	-	1	-	2	1	4	8	21	38
Male	1,177	1	2	-	-	-	-	1	2	4	16	25
Female	1,828	1	1	-	1	-	2	-	2	4	5	13
Black, both sexes	523	-	1	-	-	-	1	3	4	7	12	24
Male	205	-	1	-	-	-	1	1	2	5	7	13
Female	318	-	-	-	-	-	-	2	2	2	5	11
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	83	-	-	-	1	-	-	-	-	-	-	1
Male	41	-	-	-	1	-	-	-	-	-	-	-
Female	42	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
New Mexico												
All races, both sexes	730	2	-	1	-	-	-	-	4	2	4	22
Male	281	1	-	-	-	-	-	-	-	-	3	12
Female	449	1	-	1	-	-	-	-	4	2	1	10
White, both sexes	679	1	-	1	-	-	-	-	3	2	3	20
Male	258	-	-	-	-	-	-	-	-	-	2	11
Female	421	1	-	1	-	-	-	-	3	2	1	9
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	34	1	-	-	-	-	-	-	1	-	1	1
Male	15	1	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6,622	5	2	3	3	3	9	9	18	27	87	102
Male	2,571	4	2	2	2	1	6	5	10	12	36	56
Female	4,051	1	-	1	1	2	3	4	8	15	51	46
White, both sexes	5,627	2	-	1	3	3	6	3	9	20	49	64
Male	2,150	1	-	1	2	1	4	2	7	10	22	33
Female	3,477	1	-	-	1	2	2	1	2	10	27	31
Black, both sexes	811	3	2	1	-	-	3	5	8	6	36	32
Male	339	3	2	-	-	-	2	3	2	2	14	19
Female	472	-	-	1	-	-	1	2	6	4	22	13
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	173	-	-	1	-	-	-	1	1	1	2	6
Male	77	-	-	1	-	-	-	-	1	-	-	4
Female	96	-	-	-	-	-	-	1	-	1	2	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
North Carolina												
All races, both sexes	4,861	4	4	1	-	2	4	11	16	34	69	92
Male	1,907	1	3	-	-	2	1	9	9	15	34	53
Female	2,954	3	1	1	-	-	3	2	7	19	35	39
White, both sexes	3,669	-	2	1	-	2	2	4	11	19	32	49
Male	1,412	-	2	-	-	2	1	4	6	7	15	28
Female	2,257	-	-	1	-	-	1	-	5	12	17	21
Black, both sexes	1,121	4	2	-	-	-	1	7	4	14	34	42
Male	463	1	1	-	-	-	-	5	2	7	18	24
Female	658	3	1	-	-	-	1	2	2	7	16	18
American Indian 1/, both sexes	41	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	30	-	-	-	-	-	1	-	1	1	2	1
Male	17	-	-	-	-	-	-	-	1	1	1	1
Female	13	-	-	-	-	-	1	-	-	-	1	-
North Dakota												
All races, both sexes	368	-	-	-	-	-	-	-	1	-	-	-
Male	147	-	-	-	-	-	-	-	1	-	-	-
Female	221	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	360	-	-	-	-	-	-	-	1	-	-	-
Male	142	-	-	-	-	-	-	-	1	-	-	-
Female	218	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Ohio												
All races, both sexes	6,279	7	-	3	-	2	6	6	18	26	50	126
Male	2,379	5	-	2	-	2	3	3	7	10	24	64
Female	3,900	2	-	1	-	-	3	3	11	16	26	62
White, both sexes	5,604	6	-	1	-	2	3	3	12	21	33	95
Male	2,099	4	-	1	-	2	2	2	4	8	17	50
Female	3,505	2	-	-	-	-	1	1	8	13	16	45
Black, both sexes	643	1	-	1	-	-	3	3	6	5	17	30
Male	267	1	-	1	-	-	1	1	3	2	7	14
Female	376	-	-	-	-	-	2	2	3	3	10	16
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	30	-	-	1	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	1	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2,235	2	2	1	1	-	3	1	5	13	18	40
Male	851	1	1	-	1	-	-	-	3	6	9	16
Female	1,384	1	1	1	-	-	3	1	2	7	9	24
White, both sexes	1,981	1	1	1	1	-	1	-	4	8	12	27
Male	746	1	-	-	1	-	-	-	2	3	4	11
Female	1,235	-	1	1	-	-	1	-	2	5	8	16
Black, both sexes	136	1	1	-	-	-	1	1	1	3	4	7
Male	57	-	1	-	-	-	-	-	1	2	3	3
Female	79	1	-	-	-	-	1	1	-	1	1	4
American Indian 1/, both sexes	97	-	-	-	-	-	1	-	-	2	1	4
Male	36	-	-	-	-	-	-	-	-	1	1	1
Female	61	-	-	-	-	-	1	-	-	1	-	3
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	1	2
Male	12	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Oregon												
All races, both sexes	2,289	1	-	1	-	1	2	1	5	7	17	36
Male	883	-	-	1	-	-	2	-	4	4	8	19
Female	1,406	1	-	-	-	1	-	1	1	3	9	17
White, both sexes	2,191	1	-	1	-	1	2	1	4	7	15	31
Male	837	-	-	1	-	-	2	-	3	4	7	17
Female	1,354	1	-	-	-	1	-	1	1	3	8	14
Black, both sexes	34	-	-	-	-	-	-	-	1	-	1	1
Male	18	-	-	-	-	-	-	-	1	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	44	-	-	-	-	-	-	-	-	-	-	3
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	2
Pennsylvania												
All races, both sexes	7,650	4	4	3	1	4	7	11	15	16	58	99
Male	2,883	3	3	2	-	1	4	5	8	11	26	45
Female	4,767	1	1	1	1	3	3	6	7	5	32	54
White, both sexes	6,873	2	3	-	1	3	5	10	13	12	40	61
Male	2,575	2	3	-	-	1	4	5	8	9	17	31
Female	4,298	-	-	-	1	2	1	5	5	3	23	30
Black, both sexes	716	2	1	3	-	1	2	1	2	4	16	35
Male	278	1	-	2	-	-	-	-	-	2	8	13
Female	438	1	1	1	-	1	2	1	2	2	8	22
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	57	-	-	-	-	-	-	-	-	-	2	3
Male	29	-	-	-	-	-	-	-	-	-	1	1
Female	28	-	-	-	-	-	-	-	-	-	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Rhode Island												
All races, both sexes	533	-	-	-	-	-	-	-	2	4	4	14
Male	212	-	-	-	-	-	-	-	-	1	2	5
Female	321	-	-	-	-	-	-	-	2	3	2	9
White, both sexes	509	-	-	-	-	-	-	-	1	4	3	9
Male	199	-	-	-	-	-	-	-	-	1	2	3
Female	310	-	-	-	-	-	-	-	1	3	1	6
Black, both sexes	17	-	-	-	-	-	-	-	1	-	-	3
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	7	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	1	1
South Carolina												
All races, both sexes	2,458	2	1	-	-	2	-	5	10	10	36	59
Male	1,008	1	-	-	-	2	-	3	5	6	17	35
Female	1,450	1	1	-	-	-	-	2	5	4	19	24
White, both sexes	1,703	-	1	-	-	1	-	3	8	4	14	22
Male	666	-	-	-	-	1	-	2	5	2	8	13
Female	1,037	-	1	-	-	-	-	1	3	2	6	9
Black, both sexes	748	2	-	-	-	1	-	2	2	6	22	37
Male	340	1	-	-	-	1	-	1	-	4	9	22
Female	408	1	-	-	-	-	-	1	2	2	13	15
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
South Dakota												
All races, both sexes	511	-	-	-	-	1	-	-	1	1	3	5
Male	203	-	-	-	-	-	-	-	1	1	1	2
Female	308	-	-	-	-	1	-	-	-	-	2	3
White, both sexes	495	-	-	-	-	1	-	-	1	1	2	4
Male	197	-	-	-	-	-	-	-	1	1	1	1
Female	298	-	-	-	-	1	-	-	-	-	1	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3,659	4	2	2	1	2	2	6	9	15	56	88
Male	1,412	2	1	2	1	1	-	4	7	5	26	49
Female	2,247	2	1	-	-	1	2	2	2	10	30	39
White, both sexes	3,097	2	2	1	-	1	2	4	7	10	31	56
Male	1,159	1	1	1	-	1	-	3	5	3	16	33
Female	1,938	1	1	-	-	-	2	1	2	7	15	23
Black, both sexes	547	2	-	1	1	1	-	2	2	5	25	31
Male	247	1	-	1	1	-	-	1	2	2	10	16
Female	300	1	-	-	-	1	-	1	-	3	15	15
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Texas												
All races, both sexes	9,366	10	4	3	8	5	12	9	33	58	143	226
Male	3,744	4	2	2	4	1	9	3	16	37	79	131
Female	5,622	6	2	1	4	4	3	6	17	21	64	95
White, both sexes	8,004	6	4	3	7	5	9	5	24	44	96	157
Male	3,154	2	2	2	4	1	7	3	11	29	51	99
Female	4,850	4	2	1	3	4	2	2	13	15	45	58
Black, both sexes	1,218	3	-	-	1	-	3	4	6	10	40	64
Male	521	1	-	-	-	-	2	-	3	5	24	30
Female	697	2	-	-	1	-	1	4	3	5	16	34
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	135	1	-	-	-	-	-	-	3	3	7	4
Male	63	1	-	-	-	-	-	-	2	2	4	1
Female	72	-	-	-	-	-	-	-	1	1	3	3
Utah												
All races, both sexes	794	-	1	2	-	1	1	1	1	8	2	13
Male	321	-	1	1	-	1	-	-	-	6	2	5
Female	473	-	-	1	-	-	1	1	1	2	-	8
White, both sexes	766	-	1	2	-	-	1	1	1	8	2	11
Male	304	-	1	1	-	-	-	-	-	6	2	3
Female	462	-	-	1	-	-	1	1	1	2	-	8
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	1	-	-	-	-	-	1
Male	10	-	-	-	-	1	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Vermont												
All races, both sexes	260	1	-	-	-	-	-	-	-	-	2	3
Male	111	1	-	-	-	-	-	-	-	-	1	1
Female	149	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	260	1	-	-	-	-	-	-	-	-	2	3
Male	111	1	-	-	-	-	-	-	-	-	1	1
Female	149	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	3,675	-	-	1	1	3	3	3	7	27	39	72
Male	1,387	-	-	1	1	-	3	1	5	17	14	32
Female	2,288	-	-	-	-	3	-	2	2	10	25	40
White, both sexes	2,779	-	-	-	1	1	1	1	3	15	16	36
Male	994	-	-	-	1	-	1	-	2	9	7	17
Female	1,785	-	-	-	-	1	-	1	1	6	9	19
Black, both sexes	826	-	-	1	-	2	2	1	4	9	22	34
Male	368	-	-	1	-	-	2	1	3	6	7	15
Female	458	-	-	-	-	2	-	-	1	3	15	19
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	66	-	-	-	-	-	-	1	-	3	1	2
Male	24	-	-	-	-	-	-	-	-	2	-	-
Female	42	-	-	-	-	-	-	1	-	1	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Washington												
All races, both sexes	2,895	4	-	-	1	2	-	2	6	15	14	38
Male	1,133	3	-	-	1	2	-	1	5	6	6	23
Female	1,762	1	-	-	-	-	-	1	1	9	8	15
White, both sexes	2,693	2	-	-	-	1	-	2	6	11	12	32
Male	1,040	2	-	-	-	1	-	1	5	5	5	19
Female	1,653	-	-	-	-	-	-	1	1	6	7	13
Black, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	23	-	-	-	-	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	30	1	-	-	-	-	-	-	-	2	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	17	1	-	-	-	-	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	122	1	-	-	1	1	-	-	-	2	1	4
Male	57	1	-	-	1	1	-	-	-	1	-	3
Female	65	-	-	-	-	-	-	-	-	1	1	1
West Virginia												
All races, both sexes	1,151	-	-	-	-	-	2	2	3	5	10	19
Male	415	-	-	-	-	-	1	-	2	1	5	9
Female	736	-	-	-	-	-	1	2	1	4	5	10
White, both sexes	1,113	-	-	-	-	-	2	2	3	5	10	17
Male	403	-	-	-	-	-	1	-	2	1	5	8
Female	710	-	-	-	-	-	1	2	1	4	5	9
Black, both sexes	38	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebrovascular diseases (I60-I69)												
Wisconsin												
All races, both sexes	2,960	2	-	1	1	2	3	4	5	10	24	37
Male	1,149	2	-	1	1	2	1	3	2	5	19	18
Female	1,811	-	-	-	-	-	2	1	3	5	5	19
White, both sexes	2,830	2	-	1	1	1	2	4	4	10	21	30
Male	1,081	2	-	1	1	1	1	3	2	5	17	14
Female	1,749	-	-	-	-	-	1	1	2	5	4	16
Black, both sexes	98	-	-	-	-	1	1	-	1	-	1	7
Male	51	-	-	-	-	1	-	-	-	-	-	4
Female	47	-	-	-	-	-	1	-	1	-	1	3
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	-	-	2	-
Male	11	-	-	-	-	-	-	-	-	-	2	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	221	-	-	-	-	-	-	-	-	1	1	6
Male	81	-	-	-	-	-	-	-	-	1	-	3
Female	140	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	213	-	-	-	-	-	-	-	-	-	1	6
Male	79	-	-	-	-	-	-	-	-	-	-	3
Female	134	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
United States												
All races, both sexes	3,752	4,480	5,548	7,473	11,366	18,800	28,059	28,909	20,446	7,406	1,422	4
Male	2,101	2,535	3,101	3,996	5,692	8,598	11,090	9,662	5,237	1,350	163	4
Female	1,651	1,945	2,447	3,477	5,674	10,202	16,969	19,247	15,209	6,056	1,259	-
White, both sexes	2,409	3,101	4,039	5,695	9,134	15,990	24,842	26,229	18,645	6,750	1,230	-
Male	1,359	1,742	2,244	3,038	4,607	7,419	9,845	8,794	4,781	1,220	140	-
Female	1,050	1,359	1,795	2,657	4,527	8,571	14,997	17,435	13,864	5,530	1,090	-
Black, both sexes	1,157	1,181	1,249	1,434	1,801	2,229	2,517	2,054	1,408	513	166	4
Male	637	683	734	778	875	911	933	632	321	86	19	4
Female	520	498	515	656	926	1,318	1,584	1,422	1,087	427	147	-
American Indian 1/, both sexes	43	41	48	64	78	81	82	65	45	13	6	-
Male	19	25	18	29	36	36	24	23	9	2	-	-
Female	24	16	30	35	42	45	58	42	36	11	6	-
Asian or Pacific Islander 2/, both sexes	143	157	212	280	353	500	618	561	348	130	20	-
Male	86	85	105	151	174	232	288	213	126	42	4	-
Female	57	72	107	129	179	268	330	348	222	88	16	-
Alabama												
All races, both sexes	101	142	161	196	273	406	543	490	318	126	28	-
Male	55	72	102	93	135	165	188	153	63	18	2	-
Female	46	70	59	103	138	241	355	337	255	108	26	-
White, both sexes	49	74	88	129	196	313	420	406	266	94	14	-
Male	27	37	60	59	96	134	147	128	54	16	-	-
Female	22	37	28	70	100	179	273	278	212	78	14	-
Black, both sexes	52	68	72	66	76	93	122	82	52	32	14	-
Male	28	35	41	33	39	31	40	24	9	2	2	-
Female	24	33	31	33	37	62	82	58	43	30	12	-
American Indian 1/, both sexes	-	-	1	-	1	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	2	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Alaska												
All races, both sexes	10	6	13	9	14	29	31	33	11	7	5	-
Male	5	5	8	4	9	13	13	16	4	1	2	-
Female	5	1	5	5	5	16	18	17	7	6	3	-
White, both sexes	4	5	5	5	6	19	20	24	9	6	5	-
Male	2	4	4	2	3	9	10	13	3	1	2	-
Female	2	1	1	3	3	10	10	11	6	5	3	-
Black, both sexes	1	1	2	-	-	-	2	1	-	-	-	-
Male	1	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	2	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	5	-	6	4	6	8	5	8	2	1	-	-
Male	2	-	4	2	6	3	-	2	1	-	-	-
Female	3	-	2	2	-	5	5	6	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	2	4	-	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	2	1	2	-	-	-	-	-
Arizona												
All races, both sexes	61	55	89	129	213	319	458	505	317	103	19	-
Male	43	32	52	57	115	148	189	173	91	22	2	-
Female	18	23	37	72	98	171	269	332	226	81	17	-
White, both sexes	48	43	79	116	199	301	441	492	303	101	18	-
Male	36	25	48	49	109	143	183	171	87	22	2	-
Female	12	18	31	67	90	158	258	321	216	79	16	-
Black, both sexes	7	6	4	7	6	7	7	4	7	1	1	-
Male	4	3	2	5	2	2	1	-	1	-	-	-
Female	3	3	2	2	4	5	6	4	6	1	1	-
American Indian 1/, both sexes	6	5	4	5	7	3	3	6	4	1	-	-
Male	3	4	2	3	3	1	1	2	1	-	-	-
Female	3	1	2	2	4	2	2	4	3	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	1	8	7	3	3	-	-	-
Male	-	-	-	-	1	2	4	-	2	-	-	-
Female	-	1	2	1	-	6	3	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Arkansas												
All races, both sexes	56	77	80	108	138	247	344	358	244	93	23	-
Male	33	39	41	60	77	115	127	114	58	12	4	-
Female	23	38	39	48	61	132	217	244	186	81	19	-
White, both sexes	35	58	57	85	113	210	299	320	211	77	14	-
Male	23	29	29	48	63	98	113	101	47	9	2	-
Female	12	29	28	37	50	112	186	219	164	68	12	-
Black, both sexes	20	18	23	22	25	35	43	36	32	15	9	-
Male	10	9	12	12	14	17	13	13	11	3	2	-
Female	10	9	11	10	11	18	30	23	21	12	7	-
American Indian 1/, both sexes	1	-	-	1	-	1	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	-	1	-	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	2	1	-	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
California												
All races, both sexes	412	468	583	806	1,186	1,950	3,072	3,211	2,262	820	151	1
Male	229	253	311	403	581	865	1,279	1,174	643	166	18	1
Female	183	215	272	403	605	1,085	1,793	2,037	1,619	654	133	-
White, both sexes	272	331	419	575	879	1,575	2,597	2,788	1,991	725	128	-
Male	157	177	230	285	437	699	1,069	1,025	555	143	14	-
Female	115	154	189	290	442	876	1,528	1,763	1,436	582	114	-
Black, both sexes	74	66	75	118	128	133	179	152	89	35	11	1
Male	37	36	37	58	59	57	71	44	24	4	2	1
Female	37	30	38	60	69	76	108	108	65	31	9	-
American Indian 1/, both sexes	5	8	3	9	13	8	7	8	6	2	1	-
Male	1	6	1	3	4	4	2	5	1	-	-	-
Female	4	2	2	6	9	4	5	3	5	2	1	-
Asian or Pacific Islander 2/, both sexes	61	63	86	104	166	234	289	263	176	58	11	-
Male	34	34	43	57	81	105	137	100	63	19	2	-
Female	27	29	43	47	85	129	152	163	113	39	9	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Colorado												
All races, both sexes	45	50	50	82	112	213	292	353	240	77	15	-
Male	25	29	26	48	58	105	124	132	60	15	1	-
Female	20	21	24	34	54	108	168	221	180	62	14	-
White, both sexes	38	44	41	74	104	202	280	345	234	77	15	-
Male	20	28	22	43	56	99	120	128	59	15	1	-
Female	18	16	19	31	48	103	160	217	175	62	14	-
Black, both sexes	7	5	4	5	4	9	8	7	6	-	-	-
Male	5	1	2	3	2	4	4	3	1	-	-	-
Female	2	4	2	2	2	5	4	4	5	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	4	3	3	2	4	1	-	-	-	-
Male	-	-	2	2	-	2	-	1	-	-	-	-
Female	-	1	2	1	3	-	4	-	-	-	-	-
Connecticut												
All races, both sexes	32	36	43	58	126	171	308	320	268	102	14	-
Male	18	23	24	39	66	83	139	97	82	16	1	-
Female	14	13	19	19	60	88	169	223	186	86	13	-
White, both sexes	26	30	34	45	110	155	292	302	263	99	12	-
Male	13	19	18	32	58	73	133	93	82	16	1	-
Female	13	11	16	13	52	82	159	209	181	83	11	-
Black, both sexes	5	6	7	10	16	16	14	17	5	3	1	-
Male	4	4	5	4	8	10	6	4	-	-	-	-
Female	1	2	2	6	8	6	8	13	5	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	3	-	-	2	1	-	-	1	-
Male	1	-	1	3	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	2	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Delaware												
All races, both sexes	7	14	14	18	42	46	78	85	49	18	4	-
Male	3	9	7	7	22	25	30	38	17	4	-	-
Female	4	5	7	11	20	21	48	47	32	14	4	-
White, both sexes	4	8	8	12	31	40	69	72	42	16	3	-
Male	2	5	4	4	17	24	27	33	14	3	-	-
Female	2	3	4	8	14	16	42	39	28	13	3	-
Black, both sexes	3	6	6	6	8	5	9	13	7	2	1	-
Male	1	4	3	3	3	-	3	5	3	1	-	-
Female	2	2	3	3	5	5	6	8	4	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	1	-	-	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	15	11	17	18	17	22	38	37	22	11	2	-
Male	8	3	15	7	10	8	12	11	7	1	-	-
Female	7	8	2	11	7	14	26	26	15	10	2	-
White, both sexes	3	1	2	1	3	8	11	11	7	6	-	-
Male	2	-	2	-	2	4	7	4	3	-	-	-
Female	1	1	-	1	1	4	4	7	4	6	-	-
Black, both sexes	12	10	15	16	14	14	26	25	15	5	2	-
Male	6	3	13	7	8	4	5	6	4	1	-	-
Female	6	7	2	9	6	10	21	19	11	4	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Florida												
All races, both sexes	237	251	378	514	751	1,256	1,915	1,836	1,297	468	92	1
Male	133	143	230	288	398	585	806	696	405	109	13	1
Female	104	108	148	226	353	671	1,109	1,140	892	359	79	-
White, both sexes	155	171	289	369	617	1,086	1,724	1,677	1,188	432	78	-
Male	86	98	173	206	336	516	729	651	375	103	10	-
Female	69	73	116	163	281	570	995	1,026	813	329	68	-
Black, both sexes	79	75	81	136	126	158	178	144	104	35	14	1
Male	45	41	53	78	60	65	72	41	29	6	3	1
Female	34	34	28	58	66	93	106	103	75	29	11	-
American Indian 1/, both sexes	1	1	1	1	1	1	3	1	1	-	-	-
Male	1	1	-	-	1	-	1	1	-	-	-	-
Female	-	-	1	1	-	1	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	7	8	7	11	10	14	4	1	-	-
Male	1	3	4	4	1	4	4	3	1	-	-	-
Female	1	1	3	4	6	7	6	11	3	1	-	-
Georgia												
All races, both sexes	150	175	202	254	378	507	653	678	452	121	33	-
Male	100	105	113	146	170	209	230	212	88	24	3	-
Female	50	70	89	108	208	298	423	466	364	97	30	-
White, both sexes	53	77	109	162	269	365	492	564	353	86	23	-
Male	32	42	59	93	125	155	183	176	71	15	3	-
Female	21	35	50	69	144	210	309	388	282	71	20	-
Black, both sexes	94	92	90	92	104	134	153	113	97	35	10	-
Male	66	60	53	53	43	50	45	36	17	9	-	-
Female	28	32	37	39	61	84	108	77	80	26	10	-
American Indian 1/, both sexes	-	1	1	-	1	2	2	-	1	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	3	5	2	-	4	6	6	1	1	-	-	-
Male	2	2	1	-	1	3	2	-	-	-	-	-
Female	1	3	1	-	3	3	4	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Hawaii												
All races, both sexes	20	23	30	33	50	86	122	134	100	54	3	-
Male	15	11	18	19	31	46	59	49	38	19	-	-
Female	5	12	12	14	19	40	63	85	62	35	3	-
White, both sexes	-	3	10	4	11	18	26	29	19	10	-	-
Male	-	2	6	3	5	10	12	10	5	-	-	-
Female	-	1	4	1	6	8	14	19	14	10	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	20	20	29	39	68	96	105	81	44	3	-
Male	15	9	12	16	26	36	47	39	33	19	-	-
Female	5	11	8	13	13	32	49	66	48	25	3	-
Idaho												
All races, both sexes	10	16	26	42	51	79	144	165	129	40	2	-
Male	5	10	15	27	22	39	65	66	30	6	-	-
Female	5	6	11	15	29	40	79	99	99	34	2	-
White, both sexes	8	16	26	40	51	77	143	164	129	40	2	-
Male	5	10	15	27	22	37	64	65	30	6	-	-
Female	3	6	11	13	29	40	79	99	99	34	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Illinois												
All races, both sexes	129	169	216	303	446	778	1,336	1,316	937	346	54	-
Male	76	91	124	163	209	380	535	419	233	62	3	-
Female	53	78	92	140	237	398	801	897	704	284	51	-
White, both sexes	72	106	150	215	350	641	1,174	1,177	854	313	42	-
Male	43	51	81	109	172	324	484	370	207	56	3	-
Female	29	55	69	106	178	317	690	807	647	257	39	-
Black, both sexes	53	57	58	78	89	122	143	120	78	31	11	-
Male	30	36	39	48	33	53	45	39	24	5	-	-
Female	23	21	19	30	56	69	98	81	54	26	11	-
American Indian 1/, both sexes	-	-	1	1	1	-	1	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	6	7	9	6	15	18	18	5	2	1	-
Male	3	4	4	5	4	3	6	9	2	1	-	-
Female	1	2	3	4	2	12	12	9	3	1	1	-
Indiana												
All races, both sexes	82	86	144	181	278	458	629	686	427	154	36	-
Male	44	56	79	102	125	212	238	224	108	24	6	-
Female	38	30	65	79	153	246	391	462	319	130	30	-
White, both sexes	56	66	117	151	251	421	580	658	411	147	30	-
Male	32	45	63	85	109	193	222	212	105	23	6	-
Female	24	21	54	66	142	228	358	446	306	124	24	-
Black, both sexes	26	20	25	28	27	35	48	28	16	7	6	-
Male	12	11	15	16	16	17	16	12	3	1	-	-
Female	14	9	10	12	11	18	32	16	13	6	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	-	2	1	-	-	-	-	-
Male	-	-	1	1	-	2	-	-	-	-	-	-
Female	-	-	1	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Iowa												
All races, both sexes	30	24	50	76	112	201	398	436	353	140	29	-
Male	20	16	29	45	64	98	163	158	95	23	2	-
Female	10	8	21	31	48	103	235	278	258	117	27	-
White, both sexes	25	24	48	74	112	198	397	429	350	140	29	-
Male	18	16	27	44	64	96	162	154	94	23	2	-
Female	7	8	21	30	48	102	235	275	256	117	27	-
Black, both sexes	4	-	2	-	-	2	-	6	3	-	-	-
Male	1	-	2	-	-	1	-	3	1	-	-	-
Female	3	-	-	-	-	1	-	3	2	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	1	1	-	-	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	29	46	48	58	114	193	304	356	243	105	26	-
Male	17	29	27	28	54	101	121	120	66	15	4	-
Female	12	17	21	30	60	92	183	236	177	90	22	-
White, both sexes	22	39	41	46	103	180	291	345	232	102	26	-
Male	11	24	25	24	51	93	117	116	62	14	4	-
Female	11	15	16	22	52	87	174	229	170	88	22	-
Black, both sexes	7	7	6	10	3	12	10	9	8	3	-	-
Male	6	5	2	3	1	8	4	4	3	1	-	-
Female	1	2	4	7	2	4	6	5	5	2	-	-
American Indian 1/, both sexes	-	-	-	1	4	1	2	1	3	-	-	-
Male	-	-	-	1	1	-	-	-	1	-	-	-
Female	-	-	-	-	3	1	2	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	4	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	1	3	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Kentucky												
All races, both sexes	61	68	99	129	211	287	404	423	300	82	19	-
Male	36	39	55	66	109	135	150	118	61	10	1	-
Female	25	29	44	63	102	152	254	305	239	72	18	-
White, both sexes	45	56	84	119	197	268	382	401	285	79	18	-
Male	26	34	46	61	102	126	144	115	58	10	1	-
Female	19	22	38	58	95	142	238	286	227	69	17	-
Black, both sexes	16	9	15	10	14	19	22	21	15	3	1	-
Male	10	3	9	5	7	9	6	3	3	-	-	-
Female	6	6	6	5	7	10	16	18	12	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	-	-	-	-	-	1	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Louisiana												
All races, both sexes	100	130	145	165	233	341	448	382	276	77	16	1
Male	50	70	84	72	115	146	148	123	66	9	2	1
Female	50	60	61	93	118	195	300	259	210	68	14	-
White, both sexes	40	48	70	90	140	233	328	306	214	59	12	-
Male	19	23	36	36	71	107	115	106	55	5	2	-
Female	21	25	34	54	69	126	213	200	159	54	10	-
Black, both sexes	58	80	73	73	93	105	118	76	60	18	4	1
Male	30	46	46	34	44	37	32	17	11	4	-	1
Female	28	34	27	39	49	68	86	59	49	14	4	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	2	2	-	3	2	-	1	-	-	-
Male	-	1	2	2	-	2	1	-	-	-	-	-
Female	1	1	-	-	-	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Maine												
All races, both sexes	11	17	26	24	66	84	137	146	116	44	7	-
Male	8	12	9	12	38	42	49	48	31	5	1	-
Female	3	5	17	12	28	42	88	98	85	39	6	-
White, both sexes	11	17	26	24	66	84	136	146	116	44	7	-
Male	8	12	9	12	38	42	49	48	31	5	1	-
Female	3	5	17	12	28	42	87	98	85	39	6	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	65	87	125	119	168	333	485	465	367	133	21	-
Male	33	52	74	62	87	143	211	147	89	24	2	-
Female	32	35	51	57	81	190	274	318	278	109	19	-
White, both sexes	32	42	54	63	106	258	395	385	313	114	14	-
Male	16	29	33	34	55	112	170	116	79	22	2	-
Female	16	13	21	29	51	146	225	269	234	92	12	-
Black, both sexes	32	39	67	55	60	71	76	72	50	18	6	-
Male	17	21	39	27	31	28	35	27	9	2	-	-
Female	15	18	28	28	29	43	41	45	41	16	6	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	6	4	-	2	4	14	7	4	1	1	-
Male	-	2	2	-	1	3	6	4	1	-	-	-
Female	1	4	2	-	1	1	8	3	3	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Massachusetts												
All races, both sexes	50	72	65	126	175	368	601	681	512	213	31	-
Male	30	39	36	72	91	162	220	228	126	39	4	-
Female	20	33	29	54	84	206	381	453	386	174	27	-
White, both sexes	41	63	56	108	160	342	580	662	499	207	29	-
Male	26	33	29	63	83	152	213	220	122	38	4	-
Female	15	30	27	45	77	190	367	442	377	169	25	-
Black, both sexes	7	6	8	12	11	17	16	14	11	3	1	-
Male	3	3	6	7	6	7	3	8	2	-	-	-
Female	4	3	2	5	5	10	13	6	9	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	1	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	2	3	1	6	4	8	4	5	2	3	-	-
Male	1	3	1	2	2	2	3	-	2	1	-	-
Female	1	-	-	4	2	6	1	5	-	2	-	-
Michigan												
All races, both sexes	138	151	192	250	408	705	935	1,057	727	271	47	-
Male	79	86	117	120	195	338	359	346	204	41	5	-
Female	59	65	75	130	213	367	576	711	523	230	42	-
White, both sexes	86	92	139	197	330	591	842	965	667	255	42	-
Male	52	54	85	87	157	293	324	312	192	38	5	-
Female	34	38	54	110	173	298	518	653	475	217	37	-
Black, both sexes	50	55	49	45	71	109	88	84	58	15	5	-
Male	27	31	30	27	34	42	35	33	12	3	-	-
Female	23	24	19	18	37	67	53	51	46	12	5	-
American Indian 1/, both sexes	1	3	2	3	4	3	3	3	-	1	-	-
Male	-	1	1	2	2	1	-	-	-	-	-	-
Female	1	2	1	1	2	2	3	3	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	5	3	2	2	5	2	-	-	-
Male	-	-	1	4	2	2	-	1	-	-	-	-
Female	1	1	1	1	1	-	2	4	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Minnesota												
All races, both sexes	54	49	58	96	160	265	441	549	448	153	27	-
Male	28	37	33	56	76	133	174	200	101	26	1	-
Female	26	12	25	40	84	132	267	349	347	127	26	-
White, both sexes	47	45	51	86	149	256	432	538	443	153	27	-
Male	23	33	30	50	70	127	171	196	99	26	1	-
Female	24	12	21	36	79	129	261	342	344	127	26	-
Black, both sexes	3	1	3	3	3	4	3	5	-	-	-	-
Male	3	1	2	2	2	2	-	2	-	-	-	-
Female	-	-	1	1	1	2	3	3	-	-	-	-
American Indian 1/, both sexes	-	1	1	5	2	1	-	1	1	-	-	-
Male	-	1	-	3	2	1	-	-	-	-	-	-
Female	-	-	1	2	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	4	2	3	2	6	4	6	5	4	-	-	-
Male	2	2	1	1	2	3	3	2	2	-	-	-
Female	2	-	2	1	4	1	3	3	2	-	-	-
Mississippi												
All races, both sexes	79	65	88	126	146	213	285	266	168	63	12	-
Male	39	40	63	71	95	106	117	85	37	12	2	-
Female	40	25	25	55	51	107	168	181	131	51	10	-
White, both sexes	35	32	42	79	82	150	207	208	122	42	8	-
Male	19	21	29	42	53	76	86	62	26	4	1	-
Female	16	11	13	37	29	74	121	146	96	38	7	-
Black, both sexes	44	32	45	45	62	62	75	56	46	21	4	-
Male	20	19	33	28	42	30	30	22	11	8	1	-
Female	24	13	12	17	20	32	45	34	35	13	3	-
American Indian 1/, both sexes	-	1	1	2	-	-	1	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	2	2	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	2	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Missouri												
All races, both sexes	85	94	142	151	274	426	659	666	507	181	34	-
Male	44	51	75	89	122	198	267	203	105	39	3	-
Female	41	43	67	62	152	228	392	463	402	142	31	-
White, both sexes	58	68	112	124	231	392	615	623	481	167	31	-
Male	24	39	57	76	106	185	251	193	99	36	2	-
Female	34	29	55	48	125	207	364	430	382	131	29	-
Black, both sexes	27	25	30	26	38	30	43	43	23	13	3	-
Male	20	12	18	13	14	11	16	10	6	3	1	-
Female	7	13	12	13	24	19	27	33	17	10	2	-
American Indian 1/, both sexes	-	-	-	1	2	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	2	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	3	3	1	-	2	1	-	-
Male	-	-	-	-	2	2	-	-	-	-	-	-
Female	-	1	-	-	1	1	1	-	2	1	-	-
Montana												
All races, both sexes	6	11	10	26	41	56	101	126	102	23	5	-
Male	2	6	5	16	18	34	39	47	26	5	-	-
Female	4	5	5	10	23	22	62	79	76	18	5	-
White, both sexes	4	10	9	24	40	54	95	124	101	23	5	-
Male	2	6	5	15	18	33	38	46	26	5	-	-
Female	2	4	4	9	22	21	57	78	75	18	5	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	1	2	1	2	6	1	1	-	-	-
Male	-	-	-	1	-	1	1	-	-	-	-	-
Female	2	1	1	1	1	1	5	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Nebraska												
All races, both sexes	20	17	24	40	72	125	205	221	171	59	10	-
Male	12	9	12	25	41	60	101	76	43	13	3	-
Female	8	8	12	15	31	65	104	145	128	46	7	-
White, both sexes	17	12	21	36	65	120	204	215	168	59	10	-
Male	10	6	11	22	37	58	100	76	41	13	3	-
Female	7	6	10	14	28	62	104	139	127	46	7	-
Black, both sexes	2	4	1	2	4	2	1	3	3	-	-	-
Male	2	3	1	2	3	2	1	-	2	-	-	-
Female	-	1	-	-	1	-	-	3	1	-	-	-
American Indian 1/, both sexes	1	1	1	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	1	-	1	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	2	2	-	2	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	1	1	2	-	2	-	-	-	-
Nevada												
All races, both sexes	25	38	54	76	108	142	162	160	83	26	3	-
Male	14	26	23	48	62	75	75	51	24	3	-	-
Female	11	12	31	28	46	67	87	109	59	23	3	-
White, both sexes	16	29	42	66	84	122	145	147	73	25	3	-
Male	9	21	19	41	49	67	66	46	20	3	-	-
Female	7	8	23	25	35	55	79	101	53	22	3	-
Black, both sexes	7	4	6	4	14	7	9	8	6	-	-	-
Male	3	2	1	3	8	3	6	2	2	-	-	-
Female	4	2	5	1	6	4	3	6	4	-	-	-
American Indian 1/, both sexes	-	2	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	6	6	10	12	8	5	4	1	-	-
Male	2	2	3	4	5	5	3	3	2	-	-	-
Female	-	1	3	2	5	7	5	2	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
New Hampshire												
All races, both sexes	11	11	14	21	30	69	109	105	82	24	8	-
Male	4	5	11	13	14	32	40	28	10	3	-	-
Female	7	6	3	8	16	37	69	77	72	21	8	-
White, both sexes	10	11	13	21	30	69	108	105	81	24	8	-
Male	4	5	10	13	14	32	40	28	10	3	-	-
Female	6	6	3	8	16	37	68	77	71	21	8	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
New Jersey												
All races, both sexes	103	113	130	195	275	469	703	715	546	190	41	-
Male	47	66	82	106	124	219	273	250	144	26	4	-
Female	56	47	48	89	151	250	430	465	402	164	37	-
White, both sexes	65	73	86	139	214	389	610	651	487	174	37	-
Male	29	43	55	75	99	189	244	233	131	24	4	-
Female	36	30	31	64	115	200	366	418	356	150	33	-
Black, both sexes	34	36	35	42	53	71	75	53	54	14	4	-
Male	14	19	23	20	23	26	23	14	12	1	-	-
Female	20	17	12	22	30	45	52	39	42	13	4	-
American Indian 1/, both sexes	-	-	1	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	4	8	13	8	9	17	11	5	2	-	-
Male	4	4	4	11	2	4	6	3	1	1	-	-
Female	-	-	4	2	6	5	11	8	4	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
New Mexico												
All races, both sexes	19	19	33	37	63	96	140	144	92	39	13	-
Male	9	9	20	19	32	34	59	54	23	5	1	-
Female	10	10	13	18	31	62	81	90	69	34	12	-
White, both sexes	16	16	28	36	61	87	136	135	86	36	12	-
Male	7	8	17	19	31	30	57	49	21	5	1	-
Female	9	8	11	17	30	57	79	86	65	31	11	-
Black, both sexes	1	1	-	1	-	6	3	-	-	1	-	-
Male	1	-	-	-	-	2	1	-	-	-	-	-
Female	-	1	-	1	-	4	2	-	-	1	-	-
American Indian 1/, both sexes	2	2	5	-	2	2	-	8	6	2	1	-
Male	1	1	3	-	1	1	-	4	2	-	-	-
Female	1	1	2	-	1	1	-	4	4	2	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	181	226	238	339	518	839	1,248	1,357	986	353	69	-
Male	108	131	135	162	260	404	494	437	239	57	8	-
Female	73	95	103	177	258	435	754	920	747	296	61	-
White, both sexes	125	163	180	251	406	708	1,104	1,227	909	327	67	-
Male	76	92	102	113	219	347	432	397	224	57	8	-
Female	49	71	78	138	187	361	672	830	685	270	59	-
Black, both sexes	45	52	46	68	97	105	116	101	62	21	2	-
Male	26	34	24	38	37	44	48	31	10	-	-	-
Female	19	18	22	30	60	61	68	70	52	21	2	-
American Indian 1/, both sexes	1	1	1	-	3	1	1	2	1	-	-	-
Male	1	1	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	3	-	1	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	10	10	11	20	12	25	27	27	14	5	-	-
Male	5	4	8	11	4	12	14	8	5	-	-	-
Female	5	6	3	9	8	13	13	19	9	5	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
North Carolina												
All races, both sexes	160	175	205	278	399	630	931	973	607	223	43	-
Male	89	105	111	162	193	270	353	297	158	37	5	-
Female	71	70	94	116	206	360	578	676	449	186	38	-
White, both sexes	78	91	131	184	293	482	740	801	522	194	31	-
Male	47	52	74	111	143	216	282	242	145	32	3	-
Female	31	39	57	73	150	266	458	559	377	162	28	-
Black, both sexes	80	77	67	84	100	139	179	161	85	29	12	-
Male	41	49	36	45	49	50	67	48	13	5	2	-
Female	39	28	31	39	51	89	112	113	72	24	10	-
American Indian 1/, both sexes	-	3	5	7	3	6	9	7	-	-	-	-
Male	-	1	-	5	-	2	2	5	-	-	-	-
Female	-	2	5	2	3	4	7	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	4	2	3	3	3	3	4	-	-	-	-
Male	1	3	1	1	1	2	2	2	-	-	-	-
Female	1	1	1	2	2	1	1	2	-	-	-	-
North Dakota												
All races, both sexes	6	6	6	14	24	52	68	78	82	28	3	-
Male	3	4	4	9	11	25	34	29	21	6	-	-
Female	3	2	2	5	13	27	34	49	61	22	3	-
White, both sexes	6	4	6	14	23	50	67	77	82	27	3	-
Male	3	3	4	9	10	24	33	29	21	5	-	-
Female	3	1	2	5	13	26	34	48	61	22	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	2	-	-	1	1	1	1	-	1	-	-
Male	-	1	-	-	1	1	1	-	-	1	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Ohio												
All races, both sexes	149	168	235	316	474	849	1,313	1,248	907	308	68	-
Male	86	96	121	176	238	378	511	400	200	51	2	-
Female	63	72	114	140	236	471	802	848	707	257	66	-
White, both sexes	117	134	196	262	391	758	1,206	1,167	851	285	61	-
Male	68	78	101	146	199	344	459	376	185	51	2	-
Female	49	56	95	116	192	414	747	791	666	234	59	-
Black, both sexes	31	34	38	50	77	87	98	78	54	23	7	-
Male	17	18	20	28	37	32	47	24	14	-	-	-
Female	14	16	18	22	40	55	51	54	40	23	7	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	4	6	4	9	3	1	-	-	-
Male	1	-	-	2	2	2	5	-	1	-	-	-
Female	-	-	1	2	4	2	4	3	-	-	-	-
Oklahoma												
All races, both sexes	57	87	100	131	186	324	417	418	289	115	25	-
Male	35	45	56	66	86	140	158	134	71	21	2	-
Female	22	42	44	65	100	184	259	284	218	94	23	-
White, both sexes	37	71	89	114	155	290	383	393	263	109	21	-
Male	21	36	52	57	73	126	142	127	68	20	2	-
Female	16	35	37	57	82	164	241	266	195	89	19	-
Black, both sexes	12	13	4	5	13	14	16	19	15	3	3	-
Male	6	7	1	2	6	7	10	6	2	-	-	-
Female	6	6	3	3	7	7	6	13	13	3	3	-
American Indian 1/, both sexes	7	3	6	9	13	17	15	5	10	3	1	-
Male	7	2	2	5	6	5	4	-	1	1	-	-
Female	-	1	4	4	7	12	11	5	9	2	1	-
Asian or Pacific Islander 2/, both sexes	1	-	1	3	5	3	3	1	1	-	-	-
Male	1	-	1	2	1	2	2	1	-	-	-	-
Female	-	-	-	1	4	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Oregon												
All races, both sexes	44	61	71	80	148	287	490	511	369	139	18	-
Male	24	34	42	44	64	124	199	177	102	30	5	-
Female	20	27	29	36	84	163	291	334	267	109	13	-
White, both sexes	39	51	65	72	137	272	475	499	362	139	17	-
Male	24	27	39	42	56	117	191	173	99	30	5	-
Female	15	24	26	30	81	155	284	326	263	109	12	-
Black, both sexes	3	5	1	1	2	4	9	3	3	-	-	-
Male	-	4	1	1	2	1	5	1	1	-	-	-
Female	3	1	-	-	-	3	4	2	2	-	-	-
American Indian 1/, both sexes	1	4	-	2	3	3	1	3	1	-	-	-
Male	-	2	-	-	2	2	-	-	1	-	-	-
Female	1	2	-	2	1	1	1	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	5	5	6	8	5	6	3	-	1	-
Male	-	1	2	1	4	4	3	3	1	-	-	-
Female	1	-	3	4	2	4	2	3	2	-	1	-
Pennsylvania												
All races, both sexes	154	204	225	340	580	1,007	1,597	1,690	1,129	423	79	-
Male	75	120	111	178	294	452	627	551	282	75	10	-
Female	79	84	114	162	286	555	970	1,139	847	348	69	-
White, both sexes	107	158	170	283	500	902	1,475	1,601	1,061	393	73	-
Male	53	93	80	151	250	409	587	525	267	70	10	-
Female	54	65	90	132	250	493	888	1,076	794	323	63	-
Black, both sexes	42	44	50	54	70	103	111	82	63	24	6	-
Male	20	25	28	26	35	41	36	25	12	4	-	-
Female	22	19	22	28	35	62	75	57	51	20	6	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	1	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	5	2	5	3	9	2	10	6	5	5	-	-
Male	2	2	3	1	8	2	4	1	3	1	-	-
Female	3	-	2	2	1	-	6	5	2	4	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Rhode Island												
All races, both sexes	8	17	12	27	22	71	104	127	90	31	-	-
Male	6	9	9	16	14	35	41	40	26	8	-	-
Female	2	8	3	11	8	36	63	87	64	23	-	-
White, both sexes	7	17	12	26	20	69	100	122	89	30	-	-
Male	5	9	9	15	14	33	38	37	25	8	-	-
Female	2	8	3	11	6	36	62	85	64	22	-	-
Black, both sexes	1	-	-	-	2	2	3	3	1	1	-	-
Male	1	-	-	-	-	2	2	2	1	-	-	-
Female	-	-	-	-	2	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	2	-	-	-	-
Male	-	-	-	1	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
South Carolina												
All races, both sexes	91	101	111	145	246	350	487	392	292	94	24	-
Male	53	66	64	84	128	150	171	135	67	17	4	-
Female	38	35	47	61	118	200	316	257	225	77	20	-
White, both sexes	28	45	59	88	162	259	372	311	231	77	18	-
Male	15	27	36	48	84	117	127	110	54	13	4	-
Female	13	18	23	40	78	142	245	201	177	64	14	-
Black, both sexes	63	56	51	56	84	87	115	81	60	17	6	-
Male	38	39	27	35	44	33	44	25	13	4	-	-
Female	25	17	24	21	40	54	71	56	47	13	6	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	4	-	-	1	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	4	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
South Dakota												
All races, both sexes	9	9	23	13	38	52	110	113	84	42	7	-
Male	6	6	15	7	21	25	44	40	24	8	2	-
Female	3	3	8	6	17	27	66	73	60	34	5	-
White, both sexes	9	8	23	11	37	45	110	112	83	41	7	-
Male	6	5	15	7	21	21	44	40	24	8	2	-
Female	3	3	8	4	16	24	66	72	59	33	5	-
Black, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	1	-	2	1	6	-	-	1	-	-	-
Male	-	1	-	-	-	4	-	-	-	-	-	-
Female	-	-	-	2	1	2	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Tennessee												
All races, both sexes	98	134	143	220	306	506	695	690	451	191	38	-
Male	53	65	78	127	148	242	242	217	105	31	6	-
Female	45	69	65	93	158	264	453	473	346	160	32	-
White, both sexes	71	101	104	174	249	425	634	615	407	169	32	-
Male	36	50	55	102	116	202	217	193	92	28	4	-
Female	35	51	49	72	133	223	417	422	315	141	28	-
Black, both sexes	27	33	38	45	54	76	60	74	42	22	6	-
Male	17	15	22	24	32	38	25	24	11	3	2	-
Female	10	18	16	21	22	38	35	50	31	19	4	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	3	4	-	1	2	-	-	-
Male	-	-	1	1	-	2	-	-	2	-	-	-
Female	-	-	-	-	3	2	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Texas												
All races, both sexes	288	359	416	522	789	1,264	1,786	1,686	1,235	437	73	-
Male	172	194	226	272	408	563	711	515	303	87	5	-
Female	116	165	190	250	381	701	1,075	1,171	932	350	68	-
White, both sexes	215	274	325	419	651	1,067	1,578	1,528	1,124	398	65	-
Male	133	146	177	222	338	474	627	467	279	77	3	-
Female	82	128	148	197	313	593	951	1,061	845	321	62	-
Black, both sexes	66	78	81	85	131	180	186	132	104	36	8	-
Male	34	45	46	43	64	80	73	36	23	10	2	-
Female	32	33	35	42	67	100	113	96	81	26	6	-
American Indian 1/, both sexes	-	-	-	1	-	1	3	1	1	-	-	-
Male	-	-	-	-	-	1	2	1	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	7	7	10	17	7	16	19	25	6	3	-	-
Male	5	3	3	7	6	8	9	11	1	-	-	-
Female	2	4	7	10	1	8	10	14	5	3	-	-
Utah												
All races, both sexes	16	16	31	46	58	107	161	184	110	31	4	-
Male	6	9	11	22	33	43	62	69	43	7	-	-
Female	10	7	20	24	25	64	99	115	67	24	4	-
White, both sexes	16	14	29	44	57	102	156	179	108	30	4	-
Male	6	7	11	21	32	41	59	66	42	6	-	-
Female	10	7	18	23	25	61	97	113	66	24	4	-
Black, both sexes	-	-	-	1	-	-	2	-	-	1	-	-
Male	-	-	-	-	-	-	1	-	-	1	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	2	-	1	1	2	-	1	-	-	-
Male	-	-	-	-	1	1	1	-	1	-	-	-
Female	-	-	2	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	-	4	1	5	1	-	-	-
Male	-	2	-	1	-	1	1	3	-	-	-	-
Female	-	-	-	-	-	3	-	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Vermont												
All races, both sexes	3	4	7	6	21	23	49	71	49	18	3	-
Male	1	2	3	3	10	17	23	27	18	4	-	-
Female	2	2	4	3	11	6	26	44	31	14	3	-
White, both sexes	3	4	7	6	21	23	49	71	49	18	3	-
Male	1	2	3	3	10	17	23	27	18	4	-	-
Female	2	2	4	3	11	6	26	44	31	14	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	88	136	158	202	313	522	672	727	491	173	36	1
Male	43	72	80	108	153	231	244	227	118	30	6	1
Female	45	64	78	94	160	291	428	500	373	143	30	-
White, both sexes	45	91	97	137	199	387	536	611	422	150	30	-
Male	19	45	49	71	90	177	187	192	98	25	4	-
Female	26	46	48	66	109	210	349	419	324	125	26	-
Black, both sexes	41	42	56	58	105	124	124	109	63	22	6	1
Male	23	26	30	34	58	49	51	34	20	5	2	1
Female	18	16	26	24	47	75	73	75	43	17	4	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	5	7	9	10	12	6	5	1	-	-
Male	1	-	1	3	5	5	6	1	-	-	-	-
Female	1	2	4	4	4	5	6	5	5	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Washington												
All races, both sexes	69	67	89	144	211	408	561	628	433	161	42	-
Male	41	39	46	73	123	178	227	212	110	30	7	-
Female	28	28	43	71	88	230	334	416	323	131	35	-
White, both sexes	55	62	75	126	187	379	523	611	412	159	38	-
Male	33	35	39	67	108	164	211	204	106	30	5	-
Female	22	27	36	59	79	215	312	407	306	129	33	-
Black, both sexes	4	3	3	2	4	12	11	3	7	-	-	-
Male	2	3	3	1	3	6	5	-	-	-	-	-
Female	2	-	-	1	1	6	6	3	7	-	-	-
American Indian 1/, both sexes	2	-	3	3	4	4	5	2	-	-	2	-
Male	1	-	2	1	3	2	2	-	-	-	-	-
Female	1	-	1	2	1	2	3	2	-	-	2	-
Asian or Pacific Islander 2/, both sexes	8	2	8	13	16	13	22	12	14	2	2	-
Male	5	1	2	4	9	6	9	8	4	-	2	-
Female	3	1	6	9	7	7	13	4	10	2	-	-
West Virginia												
All races, both sexes	26	28	42	54	99	170	251	208	160	65	7	-
Male	17	13	24	34	43	72	86	54	35	17	2	-
Female	9	15	18	20	56	98	165	154	125	48	5	-
White, both sexes	25	26	39	53	94	165	245	200	158	62	7	-
Male	16	12	22	33	41	70	84	54	35	17	2	-
Female	9	14	17	20	53	95	161	146	123	45	5	-
Black, both sexes	1	2	3	1	5	5	6	8	2	3	-	-
Male	1	1	2	1	2	2	2	-	-	-	-	-
Female	-	1	1	-	3	3	4	8	2	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebrovascular diseases (I60-I69)												
Wisconsin												
All races, both sexes	49	88	82	121	186	365	583	666	486	194	51	-
Male	18	51	45	65	93	177	239	234	130	35	8	-
Female	31	37	37	56	93	188	344	432	356	159	43	-
White, both sexes	32	74	72	107	175	351	563	654	484	193	49	-
Male	13	43	39	56	88	167	228	228	129	35	8	-
Female	19	31	33	51	87	184	335	426	355	158	41	-
Black, both sexes	15	13	8	12	8	12	10	5	2	-	2	-
Male	5	8	6	9	4	8	3	2	1	-	-	-
Female	10	5	2	3	4	4	7	3	1	-	2	-
American Indian 1/, both sexes	2	-	1	2	1	1	5	-	-	-	-	-
Male	-	-	-	-	1	1	4	-	-	-	-	-
Female	2	-	1	2	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	2	1	5	7	-	1	-	-
Male	-	-	-	-	-	1	4	4	-	-	-	-
Female	-	1	1	-	2	-	1	3	-	1	-	-
Wyoming												
All races, both sexes	3	3	5	9	17	34	56	37	30	17	2	-
Male	1	1	-	5	7	18	24	11	7	2	1	-
Female	2	2	5	4	10	16	32	26	23	15	1	-
White, both sexes	1	3	4	9	17	32	55	36	30	17	2	-
Male	1	1	-	5	7	17	24	11	7	2	1	-
Female	-	2	4	4	10	15	31	25	23	15	1	-
Black, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
United States												
All races, both sexes	6,095	4	6	7	10	27	54	58	137	239	476	649
Male	2,163	2	4	4	6	11	27	37	63	97	196	228
Female	3,932	2	2	3	4	16	27	21	74	142	280	421
White, both sexes	4,886	2	4	5	5	20	37	37	96	170	326	471
Male	1,745	1	2	2	4	8	21	26	42	71	148	173
Female	3,141	1	2	3	1	12	16	11	54	99	178	298
Black, both sexes	899	1	2	2	5	4	16	16	33	50	120	143
Male	303	-	2	2	2	1	6	10	16	17	37	40
Female	596	1	-	-	3	3	10	6	17	33	83	103
American Indian 1/, both sexes	50	-	-	-	-	1	-	-	1	4	6	9
Male	13	-	-	-	-	1	-	-	-	-	2	2
Female	37	-	-	-	-	-	-	-	1	4	4	7
Asian or Pacific Islander 2/, both sexes	260	1	-	-	-	2	1	5	7	15	24	26
Male	102	1	-	-	-	1	-	1	5	9	9	13
Female	158	-	-	-	-	1	1	4	2	6	15	13
Alabama												
All races, both sexes	121	-	-	-	-	-	1	-	2	6	12	14
Male	38	-	-	-	-	-	1	-	1	1	5	3
Female	83	-	-	-	-	-	-	-	1	5	7	11
White, both sexes	91	-	-	-	-	-	1	-	1	4	9	6
Male	28	-	-	-	-	-	1	-	-	-	5	1
Female	63	-	-	-	-	-	-	-	1	4	4	5
Black, both sexes	30	-	-	-	-	-	-	-	1	2	3	8
Male	10	-	-	-	-	-	-	-	1	1	-	2
Female	20	-	-	-	-	-	-	-	-	1	3	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Alaska												
All races, both sexes	17	-	-	-	-	-	1	-	-	2	2	2
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	1	-	-	2	2	2
White, both sexes	9	-	-	-	-	-	1	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	1	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	1	2	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	2	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	101	-	-	-	-	1	1	2	1	5	7	9
Male	29	-	-	-	-	1	-	1	-	3	3	1
Female	72	-	-	-	-	-	1	1	1	2	4	8
White, both sexes	86	-	-	-	-	1	1	1	1	3	6	6
Male	27	-	-	-	-	1	-	1	-	2	3	1
Female	59	-	-	-	-	-	1	-	1	1	3	5
Black, both sexes	6	-	-	-	-	-	-	-	-	1	1	1
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	1	-	2
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	2
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Arkansas												
All races, both sexes	54	-	-	-	-	-	-	-	3	1	2	5
Male	19	-	-	-	-	-	-	-	1	-	-	4
Female	35	-	-	-	-	-	-	-	2	1	2	1
White, both sexes	46	-	-	-	-	-	-	-	1	1	-	5
Male	17	-	-	-	-	-	-	-	-	-	-	4
Female	29	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	8	-	-	-	-	-	-	-	2	-	2	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	1	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	623	1	2	1	-	5	7	8	9	32	48	76
Male	227	1	1	1	-	3	5	6	2	18	23	26
Female	396	-	1	-	-	2	2	2	7	14	25	50
White, both sexes	475	1	2	1	-	4	6	6	8	19	33	59
Male	176	1	1	1	-	2	5	5	2	12	18	19
Female	299	-	1	-	-	2	1	1	6	7	15	40
Black, both sexes	43	-	-	-	-	-	-	1	-	5	7	8
Male	13	-	-	-	-	-	-	-	-	1	1	3
Female	30	-	-	-	-	-	-	1	-	4	6	5
American Indian 1/, both sexes	6	-	-	-	-	1	-	-	-	-	1	1
Male	3	-	-	-	-	1	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	99	-	-	-	-	-	1	1	1	8	7	8
Male	35	-	-	-	-	-	-	1	-	5	3	4
Female	64	-	-	-	-	-	1	-	1	3	4	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Colorado												
All races, both sexes	68	-	-	-	-	-	1	-	1	2	4	6
Male	24	-	-	-	-	-	-	-	-	2	1	2
Female	44	-	-	-	-	-	1	-	1	-	3	4
White, both sexes	61	-	-	-	-	-	1	-	1	1	4	6
Male	21	-	-	-	-	-	-	-	-	1	1	2
Female	40	-	-	-	-	-	1	-	1	-	3	4
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	72	-	-	-	1	-	2	-	3	2	3	5
Male	26	-	-	-	-	-	1	-	1	1	1	3
Female	46	-	-	-	1	-	1	-	2	1	2	2
White, both sexes	63	-	-	-	-	-	1	-	2	1	2	4
Male	22	-	-	-	-	-	1	-	1	-	-	2
Female	41	-	-	-	-	-	-	-	1	1	2	2
Black, both sexes	8	-	-	-	1	-	1	-	1	1	1	1
Male	4	-	-	-	-	-	-	-	-	1	1	1
Female	4	-	-	-	1	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Delaware												
All races, both sexes	20	-	-	-	-	-	-	1	-	-	1	1
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	17	-	-	-	-	-	-	1	-	-	1	-
Male	6	-	-	-	-	-	-	1	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	22	-	-	-	-	-	-	-	-	1	4	1
Male	6	-	-	-	-	-	-	-	-	1	2	1
Female	16	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	1	4	1
Male	5	-	-	-	-	-	-	-	-	1	2	1
Female	10	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Florida												
All races, both sexes	422	-	-	1	-	6	5	2	7	10	30	39
Male	148	-	-	1	-	1	1	1	3	3	12	16
Female	274	-	-	-	-	5	4	1	4	7	18	23
White, both sexes	357	-	-	1	-	3	4	1	6	8	22	32
Male	129	-	-	1	-	1	1	-	2	2	9	13
Female	228	-	-	-	-	2	3	1	4	6	13	19
Black, both sexes	58	-	-	-	-	2	1	1	1	2	6	6
Male	16	-	-	-	-	-	-	1	1	1	1	2
Female	42	-	-	-	-	2	1	-	-	1	5	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	1	-	-	-	-	2	1
Male	3	-	-	-	-	-	-	-	-	-	2	1
Female	3	-	-	-	-	1	-	-	-	-	-	-
Georgia												
All races, both sexes	166	-	-	-	-	-	3	3	3	9	19	18
Male	55	-	-	-	-	-	2	2	-	1	8	4
Female	111	-	-	-	-	-	1	1	3	8	11	14
White, both sexes	107	-	-	-	-	-	1	1	2	7	12	8
Male	32	-	-	-	-	-	1	1	-	1	6	2
Female	75	-	-	-	-	-	-	-	2	6	6	6
Black, both sexes	56	-	-	-	-	-	2	2	1	2	7	9
Male	20	-	-	-	-	-	1	1	-	-	2	1
Female	36	-	-	-	-	-	1	1	1	2	5	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Hawaii												
All races, both sexes	41	-	-	-	-	-	-	-	-	-	4	7
Male	15	-	-	-	-	-	-	-	-	-	1	4
Female	26	-	-	-	-	-	-	-	-	-	3	3
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	40	-	-	-	-	-	-	-	-	-	4	7
Male	15	-	-	-	-	-	-	-	-	-	1	4
Female	25	-	-	-	-	-	-	-	-	-	3	3
Idaho												
All races, both sexes	24	-	-	-	-	-	-	-	-	2	1	3
Male	9	-	-	-	-	-	-	-	-	1	-	1
Female	15	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	23	-	-	-	-	-	-	-	-	2	1	3
Male	9	-	-	-	-	-	-	-	-	1	-	1
Female	14	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Illinois												
All races, both sexes	220	1	-	-	-	1	2	-	5	6	17	25
Male	81	-	-	-	-	-	1	-	2	5	9	12
Female	139	1	-	-	-	1	1	-	3	1	8	13
White, both sexes	170	1	-	-	-	1	1	-	3	5	15	18
Male	58	-	-	-	-	-	-	-	1	4	7	7
Female	112	1	-	-	-	1	1	-	2	1	8	11
Black, both sexes	44	-	-	-	-	-	1	-	2	1	2	7
Male	21	-	-	-	-	-	1	-	1	1	2	5
Female	23	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	135	-	-	-	-	-	2	2	5	5	9	18
Male	42	-	-	-	-	-	1	1	1	2	3	3
Female	93	-	-	-	-	-	1	1	4	3	6	15
White, both sexes	114	-	-	-	-	-	2	2	3	3	7	15
Male	39	-	-	-	-	-	1	1	-	1	2	3
Female	75	-	-	-	-	-	1	1	3	2	5	12
Black, both sexes	21	-	-	-	-	-	-	-	2	2	2	3
Male	3	-	-	-	-	-	-	-	1	1	1	-
Female	18	-	-	-	-	-	-	-	1	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Iowa												
All races, both sexes	58	-	-	-	-	1	1	1	1	2	4	9
Male	20	-	-	-	-	-	-	-	-	1	1	2
Female	38	-	-	-	-	1	1	1	1	1	3	7
White, both sexes	55	-	-	-	-	1	1	1	1	2	4	9
Male	19	-	-	-	-	-	-	-	-	1	1	2
Female	36	-	-	-	-	1	1	1	1	1	3	7
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	60	-	-	-	-	-	-	-	2	-	2	8
Male	19	-	-	-	-	-	-	-	2	-	1	3
Female	41	-	-	-	-	-	-	-	-	-	1	5
White, both sexes	55	-	-	-	-	-	-	-	2	-	2	7
Male	17	-	-	-	-	-	-	-	2	-	1	2
Female	38	-	-	-	-	-	-	-	-	-	1	5
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Kentucky												
All races, both sexes	92	-	-	-	2	-	2	1	-	2	10	10
Male	27	-	-	-	2	-	1	1	-	-	3	3
Female	65	-	-	-	-	-	1	-	-	2	7	7
White, both sexes	83	-	-	-	1	-	2	1	-	2	9	8
Male	25	-	-	-	1	-	1	1	-	-	3	3
Female	58	-	-	-	-	-	1	-	-	2	6	5
Black, both sexes	9	-	-	-	1	-	-	-	-	-	1	2
Male	2	-	-	-	1	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	105	-	-	-	2	1	1	-	4	3	9	16
Male	34	-	-	-	-	1	-	-	3	1	4	7
Female	71	-	-	-	2	-	1	-	1	2	5	9
White, both sexes	62	-	-	-	1	-	1	-	3	2	4	5
Male	21	-	-	-	-	-	-	-	2	-	2	4
Female	41	-	-	-	1	-	1	-	1	2	2	1
Black, both sexes	42	-	-	-	1	1	-	-	1	1	5	11
Male	13	-	-	-	-	1	-	-	1	1	2	3
Female	29	-	-	-	1	-	-	-	-	-	3	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Maine												
All races, both sexes	19	-	-	-	-	-	-	-	-	2	-	3
Male	7	-	-	-	-	-	-	-	-	1	-	2
Female	12	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	19	-	-	-	-	-	-	-	-	2	-	3
Male	7	-	-	-	-	-	-	-	-	1	-	2
Female	12	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	107	-	-	-	-	-	1	-	2	2	11	11
Male	36	-	-	-	-	-	1	-	1	-	2	4
Female	71	-	-	-	-	-	-	-	1	2	9	7
White, both sexes	64	-	-	-	-	-	-	-	1	-	2	4
Male	19	-	-	-	-	-	-	-	-	-	-	2
Female	45	-	-	-	-	-	-	-	1	-	2	2
Black, both sexes	37	-	-	-	-	-	1	-	1	2	8	7
Male	15	-	-	-	-	-	1	-	1	-	2	2
Female	22	-	-	-	-	-	-	-	-	2	6	5
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Massachusetts												
All races, both sexes	132	-	-	-	-	-	-	2	4	2	15	13
Male	51	-	-	-	-	-	-	2	2	1	7	5
Female	81	-	-	-	-	-	-	-	2	1	8	8
White, both sexes	121	-	-	-	-	-	-	2	4	1	12	11
Male	44	-	-	-	-	-	-	2	2	-	6	4
Female	77	-	-	-	-	-	-	-	2	1	6	7
Black, both sexes	9	-	-	-	-	-	-	-	-	1	3	2
Male	5	-	-	-	-	-	-	-	-	1	1	1
Female	4	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	206	-	-	-	-	-	2	4	3	13	14	21
Male	78	-	-	-	-	-	-	3	1	5	7	7
Female	128	-	-	-	-	-	2	1	2	8	7	14
White, both sexes	169	-	-	-	-	-	-	3	2	11	12	17
Male	66	-	-	-	-	-	-	2	-	5	7	6
Female	103	-	-	-	-	-	-	1	2	6	5	11
Black, both sexes	34	-	-	-	-	-	2	1	1	2	2	3
Male	10	-	-	-	-	-	-	1	1	-	-	-
Female	24	-	-	-	-	-	2	-	-	2	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Minnesota												
All races, both sexes	114	-	-	-	-	-	1	2	4	3	16	7
Male	44	-	-	-	-	-	-	1	2	2	6	1
Female	70	-	-	-	-	-	1	1	2	1	10	6
White, both sexes	96	-	-	-	-	-	1	1	1	2	10	5
Male	37	-	-	-	-	-	-	1	-	2	4	1
Female	59	-	-	-	-	-	1	-	1	-	6	4
Black, both sexes	10	-	-	-	-	-	-	-	2	-	5	2
Male	4	-	-	-	-	-	-	-	1	-	2	-
Female	6	-	-	-	-	-	-	-	1	-	3	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	1	1	-	1	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	1	-	-	1	-
Mississippi												
All races, both sexes	56	-	-	-	1	-	-	1	1	1	7	6
Male	23	-	-	-	1	-	-	1	1	-	4	2
Female	33	-	-	-	-	-	-	-	-	1	3	4
White, both sexes	39	-	-	-	1	-	-	1	-	-	6	5
Male	17	-	-	-	1	-	-	1	-	-	4	2
Female	22	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	17	-	-	-	-	-	-	-	1	1	1	1
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	11	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Missouri												
All races, both sexes	148	-	-	-	-	-	-	3	-	7	10	16
Male	47	-	-	-	-	-	-	3	-	3	3	5
Female	101	-	-	-	-	-	-	-	-	4	7	11
White, both sexes	126	-	-	-	-	-	-	2	-	6	8	12
Male	42	-	-	-	-	-	-	2	-	3	3	4
Female	84	-	-	-	-	-	-	-	-	3	5	8
Black, both sexes	19	-	-	-	-	-	-	1	-	1	2	3
Male	5	-	-	-	-	-	-	1	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	1	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Montana												
All races, both sexes	18	-	-	-	-	-	-	1	1	1	2	1
Male	6	-	-	-	-	-	-	1	1	-	2	-
Female	12	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	16	-	-	-	-	-	-	1	1	-	2	1
Male	6	-	-	-	-	-	-	1	1	-	2	-
Female	10	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Nebraska												
All races, both sexes	30	-	-	-	-	-	1	-	1	-	1	3
Male	13	-	-	-	-	-	1	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	27	-	-	-	-	-	1	-	-	-	1	3
Male	12	-	-	-	-	-	1	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	50	-	-	-	-	1	-	1	4	4	2	6
Male	22	-	-	-	-	-	-	1	2	2	2	3
Female	28	-	-	-	-	1	-	-	2	2	-	3
White, both sexes	39	-	-	-	-	1	-	-	2	3	2	4
Male	15	-	-	-	-	-	-	-	1	1	2	2
Female	24	-	-	-	-	1	-	-	1	2	-	2
Black, both sexes	7	-	-	-	-	-	-	1	-	1	-	1
Male	5	-	-	-	-	-	-	1	-	1	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	2	-	-	1
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
New Hampshire												
All races, both sexes	29	-	-	-	-	1	-	-	-	2	1	4
Male	12	-	-	-	-	1	-	-	-	1	1	1
Female	17	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	28	-	-	-	-	1	-	-	-	2	1	3
Male	12	-	-	-	-	1	-	-	-	1	1	1
Female	16	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
New Jersey												
All races, both sexes	177	-	1	-	-	-	2	2	5	4	12	11
Male	71	-	-	-	-	-	1	1	1	1	7	4
Female	106	-	1	-	-	-	1	1	4	3	5	7
White, both sexes	136	-	1	-	-	-	1	1	3	2	7	8
Male	58	-	-	-	-	-	-	1	1	-	5	4
Female	78	-	1	-	-	-	1	-	2	2	2	4
Black, both sexes	34	-	-	-	-	-	1	1	2	2	5	3
Male	9	-	-	-	-	-	1	-	-	1	2	-
Female	25	-	-	-	-	-	-	1	2	1	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
New Mexico												
All races, both sexes	35	-	-	1	-	-	-	-	1	1	1	7
Male	9	-	-	-	-	-	-	-	-	-	1	4
Female	26	-	-	1	-	-	-	-	1	1	-	3
White, both sexes	33	-	-	1	-	-	-	-	-	1	1	7
Male	9	-	-	-	-	-	-	-	-	-	1	4
Female	24	-	-	1	-	-	-	-	-	1	-	3
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	365	-	2	-	-	1	4	3	12	14	36	29
Male	127	-	2	-	-	-	3	-	6	6	13	11
Female	238	-	-	-	-	1	1	3	6	8	23	18
White, both sexes	283	-	-	-	-	1	3	1	6	10	21	20
Male	102	-	-	-	-	-	2	-	4	6	9	7
Female	181	-	-	-	-	1	1	1	2	4	12	13
Black, both sexes	63	-	2	-	-	-	1	1	5	3	14	6
Male	19	-	2	-	-	-	1	-	1	-	4	2
Female	44	-	-	-	-	-	-	1	4	3	10	4
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	1	1	1	1	3
Male	6	-	-	-	-	-	-	-	1	-	-	2
Female	12	-	-	-	-	-	-	1	-	1	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
North Carolina												
All races, both sexes	223	-	-	-	-	-	1	4	9	15	25	20
Male	75	-	-	-	-	-	-	3	7	5	10	8
Female	148	-	-	-	-	-	1	1	2	10	15	12
White, both sexes	158	-	-	-	-	-	-	-	6	9	12	16
Male	53	-	-	-	-	-	-	-	5	4	6	6
Female	105	-	-	-	-	-	-	-	1	5	6	10
Black, both sexes	59	-	-	-	-	-	1	4	2	5	11	4
Male	19	-	-	-	-	-	-	3	1	-	4	2
Female	40	-	-	-	-	-	1	1	1	5	7	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	1	1	1	-
Male	3	-	-	-	-	-	-	-	1	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
North Dakota												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Ohio												
All races, both sexes	253	-	-	-	-	-	-	2	12	14	12	34
Male	86	-	-	-	-	-	-	2	4	4	4	14
Female	167	-	-	-	-	-	-	-	8	10	8	20
White, both sexes	217	-	-	-	-	-	-	1	9	12	9	28
Male	72	-	-	-	-	-	-	1	2	4	3	11
Female	145	-	-	-	-	-	-	-	7	8	6	17
Black, both sexes	34	-	-	-	-	-	-	1	3	2	3	5
Male	14	-	-	-	-	-	-	1	2	-	1	3
Female	20	-	-	-	-	-	-	-	1	2	2	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	87	1	-	-	-	-	1	-	2	5	6	13
Male	32	-	-	-	-	-	-	-	1	2	2	4
Female	55	1	-	-	-	-	1	-	1	3	4	9
White, both sexes	68	-	-	-	-	-	-	-	2	4	4	9
Male	23	-	-	-	-	-	-	-	1	1	1	2
Female	45	-	-	-	-	-	-	-	1	3	3	7
Black, both sexes	12	1	-	-	-	-	1	-	-	1	2	2
Male	5	-	-	-	-	-	-	-	-	1	1	1
Female	7	1	-	-	-	-	1	-	-	-	1	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Oregon												
All races, both sexes	83	-	-	-	-	1	1	-	-	3	5	10
Male	28	-	-	-	-	-	1	-	-	1	2	4
Female	55	-	-	-	-	1	-	-	-	2	3	6
White, both sexes	77	-	-	-	-	1	1	-	-	3	5	10
Male	26	-	-	-	-	-	1	-	-	1	2	4
Female	51	-	-	-	-	1	-	-	-	2	3	6
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	298	-	-	-	-	1	2	2	7	7	19	31
Male	113	-	-	-	-	1	1	-	5	5	6	9
Female	185	-	-	-	-	-	1	2	2	2	13	22
White, both sexes	251	-	-	-	-	1	1	2	6	5	14	21
Male	98	-	-	-	-	1	1	-	5	4	4	7
Female	153	-	-	-	-	-	-	2	1	1	10	14
Black, both sexes	41	-	-	-	-	-	1	-	1	2	4	9
Male	13	-	-	-	-	-	-	-	-	1	1	2
Female	28	-	-	-	-	-	1	-	1	1	3	7
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	1
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Rhode Island												
All races, both sexes	25	-	-	-	-	-	-	-	-	2	3	6
Male	7	-	-	-	-	-	-	-	-	-	2	-
Female	18	-	-	-	-	-	-	-	-	2	1	6
White, both sexes	21	-	-	-	-	-	-	-	-	2	2	4
Male	6	-	-	-	-	-	-	-	-	-	2	-
Female	15	-	-	-	-	-	-	-	-	2	-	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
South Carolina												
All races, both sexes	104	-	-	-	-	-	-	2	3	5	5	10
Male	40	-	-	-	-	-	-	1	1	2	2	2
Female	64	-	-	-	-	-	-	1	2	3	3	8
White, both sexes	72	-	-	-	-	-	-	2	2	3	3	3
Male	27	-	-	-	-	-	-	1	1	1	2	1
Female	45	-	-	-	-	-	-	1	1	2	1	2
Black, both sexes	32	-	-	-	-	-	-	-	1	2	2	7
Male	13	-	-	-	-	-	-	-	-	1	-	1
Female	19	-	-	-	-	-	-	-	1	1	2	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
South Dakota												
All races, both sexes	19	-	-	-	-	1	-	-	-	1	1	1
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	12	-	-	-	-	1	-	-	-	-	1	1
White, both sexes	18	-	-	-	-	1	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	1	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	152	-	-	1	1	1	-	2	3	4	15	15
Male	56	-	-	1	1	1	-	1	2	1	6	3
Female	96	-	-	-	-	-	-	1	1	3	9	12
White, both sexes	121	-	-	-	-	1	-	1	3	2	8	11
Male	44	-	-	-	-	1	-	-	2	1	4	3
Female	77	-	-	-	-	-	-	1	1	1	4	8
Black, both sexes	30	-	-	1	1	-	-	1	-	2	7	4
Male	11	-	-	1	1	-	-	1	-	-	2	-
Female	19	-	-	-	-	-	-	-	-	2	5	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Texas												
All races, both sexes	406	-	1	1	3	1	5	2	13	15	35	52
Male	156	-	1	-	2	-	3	1	6	5	15	21
Female	250	-	-	1	1	1	2	1	7	10	20	31
White, both sexes	334	-	1	1	2	1	4	1	11	10	26	35
Male	132	-	1	-	2	-	3	1	5	3	11	19
Female	202	-	-	1	-	1	1	-	6	7	15	16
Black, both sexes	58	-	-	-	1	-	1	1	1	4	6	16
Male	18	-	-	-	-	-	-	-	-	1	3	2
Female	40	-	-	-	1	-	1	1	1	3	3	14
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	1	1	3	1
Male	6	-	-	-	-	-	-	-	1	1	1	-
Female	8	-	-	-	-	-	-	-	-	-	2	1
Utah												
All races, both sexes	42	-	-	1	-	1	-	-	1	2	1	2
Male	16	-	-	-	-	1	-	-	-	1	1	1
Female	26	-	-	1	-	-	-	-	1	1	-	1
White, both sexes	40	-	-	1	-	-	-	-	1	2	1	2
Male	14	-	-	-	-	-	-	-	-	1	1	1
Female	26	-	-	1	-	-	-	-	1	1	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Vermont												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	1	3
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	13	-	-	-	-	-	-	-	-	-	1	3
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	146	-	-	1	-	2	1	1	1	9	8	15
Male	43	-	-	1	-	-	1	-	1	4	2	7
Female	103	-	-	-	-	2	-	1	-	5	6	8
White, both sexes	109	-	-	-	-	1	-	-	-	6	4	11
Male	29	-	-	-	-	-	-	-	-	2	-	5
Female	80	-	-	-	-	1	-	-	-	4	4	6
Black, both sexes	31	-	-	1	-	1	1	-	1	1	3	4
Male	12	-	-	1	-	-	1	-	1	1	2	2
Female	19	-	-	-	-	1	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	1	-	2	1	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	1	-	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Washington												
All races, both sexes	133	1	-	-	-	1	-	1	1	4	8	10
Male	46	1	-	-	-	1	-	1	1	1	2	5
Female	87	-	-	-	-	-	-	-	-	3	6	5
White, both sexes	118	-	-	-	-	1	-	1	1	3	7	8
Male	39	-	-	-	-	1	-	1	1	1	1	4
Female	79	-	-	-	-	-	-	-	-	2	6	4
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	1	-	-	-	-	-	-	-	1	-	1
Male	4	1	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	1	-	-
West Virginia												
All races, both sexes	58	-	-	-	-	-	1	1	1	2	4	5
Male	19	-	-	-	-	-	1	-	1	1	1	1
Female	39	-	-	-	-	-	-	1	-	1	3	4
White, both sexes	56	-	-	-	-	-	1	1	1	2	4	4
Male	18	-	-	-	-	-	1	-	1	1	1	1
Female	38	-	-	-	-	-	-	1	-	1	3	3
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Subarachnoid hemorrhage (I60)												
Wisconsin												
All races, both sexes	124	-	-	-	-	-	2	2	-	5	12	11
Male	54	-	-	-	-	-	1	2	-	2	9	4
Female	70	-	-	-	-	-	1	-	-	3	3	7
White, both sexes	113	-	-	-	-	-	1	2	-	5	10	9
Male	52	-	-	-	-	-	1	2	-	2	8	3
Female	61	-	-	-	-	-	-	-	-	3	2	6
Black, both sexes	8	-	-	-	-	-	1	-	-	-	1	2
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	1	-	-	-	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
United States												
All races, both sexes	661	608	510	469	475	535	549	376	178	62	5	-
Male	261	244	191	160	157	161	150	107	46	10	1	-
Female	400	364	319	309	318	374	399	269	132	52	4	-
White, both sexes	491	501	414	400	390	471	480	342	168	52	4	-
Male	196	207	152	131	129	144	134	100	45	8	1	-
Female	295	294	262	269	261	327	346	242	123	44	3	-
Black, both sexes	131	84	67	43	56	44	44	23	8	6	1	-
Male	46	30	29	17	22	13	7	4	-	2	-	-
Female	85	54	38	26	34	31	37	19	8	4	1	-
American Indian 1/, both sexes	10	4	5	3	5	-	2	-	-	-	-	-
Male	2	1	1	2	2	-	-	-	-	-	-	-
Female	8	3	4	1	3	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	19	24	23	24	20	23	11	2	4	-	-
Male	17	6	9	10	4	4	9	3	1	-	-	-
Female	12	13	15	13	20	16	14	8	1	4	-	-
Alabama												
All races, both sexes	13	13	7	13	11	9	9	8	3	-	-	-
Male	6	4	5	3	3	4	-	2	-	-	-	-
Female	7	9	2	10	8	5	9	6	3	-	-	-
White, both sexes	10	9	6	11	8	9	7	7	3	-	-	-
Male	5	2	4	1	3	4	-	2	-	-	-	-
Female	5	7	2	10	5	5	7	5	3	-	-	-
Black, both sexes	3	4	1	2	3	-	2	1	-	-	-	-
Male	1	2	1	2	-	-	-	-	-	-	-	-
Female	2	2	-	-	3	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Alaska												
All races, both sexes	1	1	1	2	2	1	2	-	-	-	-	-
Male	-	-	1	1	1	-	1	-	-	-	-	-
Female	1	1	-	1	1	1	1	-	-	-	-	-
White, both sexes	-	1	-	1	1	1	2	-	-	-	-	-
Male	-	-	-	1	-	-	1	-	-	-	-	-
Female	-	1	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	1	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	9	6	14	13	10	12	4	6	1	-	-	-
Male	3	2	4	3	2	2	2	2	-	-	-	-
Female	6	4	10	10	8	10	2	4	1	-	-	-
White, both sexes	5	6	12	12	10	11	4	6	1	-	-	-
Male	3	2	4	2	2	2	2	2	-	-	-	-
Female	2	4	8	10	8	9	2	4	1	-	-	-
Black, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Arkansas												
All races, both sexes	5	10	7	7	2	6	1	3	1	1	-	-
Male	2	4	3	1	1	1	-	1	-	1	-	-
Female	3	6	4	6	1	5	1	2	1	-	-	-
White, both sexes	5	9	5	6	2	6	1	3	1	1	-	-
Male	2	4	3	-	1	1	-	1	-	1	-	-
Female	3	5	2	6	1	5	1	2	1	-	-	-
Black, both sexes	-	1	2	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	72	58	55	47	43	62	47	31	15	4	-	-
Male	29	24	21	16	11	15	12	8	4	1	-	-
Female	43	34	34	31	32	47	35	23	11	3	-	-
White, both sexes	47	50	39	38	29	55	38	22	15	3	-	-
Male	20	21	14	13	8	13	10	6	4	1	-	-
Female	27	29	25	25	21	42	28	16	11	2	-	-
Black, both sexes	8	4	4	1	3	-	1	1	-	-	-	-
Male	1	2	3	-	2	-	-	-	-	-	-	-
Female	7	2	1	1	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	4	11	7	11	7	8	8	-	1	-	-
Male	8	1	4	2	1	2	2	2	-	-	-	-
Female	8	3	7	5	10	5	6	6	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Colorado												
All races, both sexes	7	9	7	8	5	4	6	6	2	-	-	-
Male	3	5	2	2	-	-	3	4	-	-	-	-
Female	4	4	5	6	5	4	3	2	2	-	-	-
White, both sexes	6	9	5	8	4	4	5	5	2	-	-	-
Male	3	5	1	2	-	-	3	3	-	-	-	-
Female	3	4	4	6	4	4	2	2	2	-	-	-
Black, both sexes	1	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	9	5	9	2	12	5	3	4	5	-	2	-
Male	1	3	4	2	3	2	1	1	2	-	-	-
Female	8	2	5	-	9	3	2	3	3	-	2	-
White, both sexes	9	5	7	2	11	5	3	4	5	-	2	-
Male	1	3	3	2	3	2	1	1	2	-	-	-
Female	8	2	4	-	8	3	2	3	3	-	2	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Delaware												
All races, both sexes	2	1	-	3	4	3	2	2	-	-	-	-
Male	-	-	-	-	2	1	1	1	-	-	-	-
Female	2	1	-	3	2	2	1	1	-	-	-	-
White, both sexes	2	-	-	3	4	3	2	1	-	-	-	-
Male	-	-	-	-	2	1	1	1	-	-	-	-
Female	2	-	-	3	2	2	1	-	-	-	-	-
Black, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	3	-	1	2	1	3	2	-	2	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	3	-	1	2	1	3	1	-	2	-	-
White, both sexes	-	1	-	1	1	1	-	1	-	2	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	1	-	1	1	1	-	-	-	2	-	-
Black, both sexes	2	2	-	-	1	-	3	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	2	-	-	1	-	3	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Florida												
All races, both sexes	49	43	38	45	33	34	35	33	9	3	-	-
Male	19	17	15	13	14	10	11	9	2	-	-	-
Female	30	26	23	32	19	24	24	24	7	3	-	-
White, both sexes	38	36	34	38	30	29	34	29	9	3	-	-
Male	14	14	13	13	14	10	11	9	2	-	-	-
Female	24	22	21	25	16	19	23	20	7	3	-	-
Black, both sexes	11	7	3	7	2	4	1	4	-	-	-	-
Male	5	3	2	-	-	-	-	-	-	-	-	-
Female	6	4	1	7	2	4	1	4	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Georgia												
All races, both sexes	16	16	18	10	12	8	14	9	7	1	-	-
Male	10	7	7	3	3	1	2	1	3	1	-	-
Female	6	9	11	7	9	7	12	8	4	-	-	-
White, both sexes	9	8	10	9	10	5	9	9	7	-	-	-
Male	4	3	2	3	2	1	2	1	3	-	-	-
Female	5	5	8	6	8	4	7	8	4	-	-	-
Black, both sexes	6	7	8	1	2	3	5	-	-	1	-	-
Male	5	3	5	-	1	-	-	-	-	1	-	-
Female	1	4	3	1	1	3	5	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Hawaii												
All races, both sexes	4	2	-	3	2	6	8	2	2	1	-	-
Male	3	-	-	1	-	2	3	-	1	-	-	-
Female	1	2	-	2	2	4	5	2	1	1	-	-
White, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	-	3	2	6	8	2	2	1	-	-
Male	3	-	-	1	-	2	3	-	1	-	-	-
Female	1	1	-	2	2	4	5	2	1	1	-	-
Idaho												
All races, both sexes	2	1	4	3	1	2	-	2	1	2	-	-
Male	-	1	2	2	-	-	-	1	1	-	-	-
Female	2	-	2	1	1	2	-	1	-	2	-	-
White, both sexes	1	1	4	3	1	2	-	2	1	2	-	-
Male	-	1	2	2	-	-	-	1	1	-	-	-
Female	1	-	2	1	1	2	-	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Illinois												
All races, both sexes	22	16	18	21	13	25	31	11	4	2	-	-
Male	6	4	6	5	6	13	7	4	1	-	-	-
Female	16	12	12	16	7	12	24	7	3	2	-	-
White, both sexes	11	11	14	17	11	19	28	10	3	2	-	-
Male	3	2	5	3	6	10	6	3	1	-	-	-
Female	8	9	9	14	5	9	22	7	2	2	-	-
Black, both sexes	11	4	2	3	2	4	3	1	1	-	-	-
Male	3	2	-	1	-	3	1	1	-	-	-	-
Female	8	2	2	2	2	1	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	-	2	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	2	-	-	-	-	-	-
Indiana												
All races, both sexes	13	12	11	8	10	9	18	8	5	-	-	-
Male	5	4	4	4	3	1	6	2	2	-	-	-
Female	8	8	7	4	7	8	12	6	3	-	-	-
White, both sexes	12	7	10	8	7	9	17	7	5	-	-	-
Male	5	4	4	4	3	1	6	2	2	-	-	-
Female	7	3	6	4	4	8	11	5	3	-	-	-
Black, both sexes	1	5	1	-	3	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	5	1	-	3	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Iowa												
All races, both sexes	10	2	6	5	3	4	8	1	-	-	-	-
Male	5	1	3	1	2	2	2	-	-	-	-	-
Female	5	1	3	4	1	2	6	1	-	-	-	-
White, both sexes	7	2	6	5	3	4	8	1	-	-	-	-
Male	4	1	3	1	2	2	2	-	-	-	-	-
Female	3	1	3	4	1	2	6	1	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	9	7	6	7	2	7	6	4	-	-	-	-
Male	1	2	2	3	-	3	1	1	-	-	-	-
Female	8	5	4	4	2	4	5	3	-	-	-	-
White, both sexes	9	6	5	5	2	7	6	4	-	-	-	-
Male	1	1	2	3	-	3	1	1	-	-	-	-
Female	8	5	3	2	2	4	5	3	-	-	-	-
Black, both sexes	-	1	1	2	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	2	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Kentucky												
All races, both sexes	8	9	8	10	10	4	9	3	3	1	-	-
Male	1	3	4	2	1	2	3	-	1	-	-	-
Female	7	6	4	8	9	2	6	3	2	1	-	-
White, both sexes	6	8	7	10	10	4	9	3	2	1	-	-
Male	1	3	3	2	1	2	3	-	1	-	-	-
Female	5	5	4	8	9	2	6	3	1	1	-	-
Black, both sexes	2	1	1	-	-	-	-	-	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	11	10	9	9	13	7	6	4	-	-	-	-
Male	2	3	3	2	4	2	1	1	-	-	-	-
Female	9	7	6	7	9	5	5	3	-	-	-	-
White, both sexes	3	7	7	8	7	5	6	3	-	-	-	-
Male	-	3	3	2	1	2	1	1	-	-	-	-
Female	3	4	4	6	6	3	5	2	-	-	-	-
Black, both sexes	8	2	2	1	6	2	-	1	-	-	-	-
Male	2	-	-	-	3	-	-	-	-	-	-	-
Female	6	2	2	1	3	2	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Maine												
All races, both sexes	3	3	2	1	-	1	1	2	-	1	-	-
Male	2	1	-	-	-	-	-	1	-	-	-	-
Female	1	2	2	1	-	1	1	1	-	1	-	-
White, both sexes	3	3	2	1	-	1	1	2	-	1	-	-
Male	2	1	-	-	-	-	-	1	-	-	-	-
Female	1	2	2	1	-	1	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	11	10	12	7	2	10	14	7	3	4	-	-
Male	4	5	7	2	1	3	3	1	1	1	-	-
Female	7	5	5	5	1	7	11	6	2	3	-	-
White, both sexes	9	4	5	6	1	9	11	6	3	3	-	-
Male	3	2	3	2	-	2	2	1	1	1	-	-
Female	6	2	2	4	1	7	9	5	2	2	-	-
Black, both sexes	2	4	5	1	1	1	2	1	-	1	-	-
Male	1	2	3	-	1	1	1	-	-	-	-	-
Female	1	2	2	1	-	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	2	-	-	-	1	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	1	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Massachusetts												
All races, both sexes	8	14	8	9	5	13	19	15	3	2	-	-
Male	5	5	4	4	2	4	5	5	-	-	-	-
Female	3	9	4	5	3	9	14	10	3	2	-	-
White, both sexes	6	14	7	8	5	13	18	15	3	2	-	-
Male	4	5	3	3	2	4	4	5	-	-	-	-
Female	2	9	4	5	3	9	14	10	3	2	-	-
Black, both sexes	1	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	25	23	16	15	17	16	18	13	6	-	-	-
Male	14	10	7	3	4	7	4	3	3	-	-	-
Female	11	13	9	12	13	9	14	10	3	-	-	-
White, both sexes	21	20	11	14	13	11	16	13	5	-	-	-
Male	12	10	5	2	3	4	4	3	3	-	-	-
Female	9	10	6	12	10	7	12	10	2	-	-	-
Black, both sexes	4	3	4	-	4	5	2	-	1	-	-	-
Male	2	-	2	-	1	3	-	-	-	-	-	-
Female	2	3	2	-	3	2	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Minnesota												
All races, both sexes	16	15	4	4	7	13	11	7	2	2	-	-
Male	6	9	1	2	2	5	2	4	-	1	-	-
Female	10	6	3	2	5	8	9	3	2	1	-	-
White, both sexes	16	13	4	4	6	12	10	7	2	2	-	-
Male	6	7	1	2	2	4	2	4	-	1	-	-
Female	10	6	3	2	4	8	8	3	2	1	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	-	-	-
Mississippi												
All races, both sexes	11	4	6	4	1	3	4	2	3	1	-	-
Male	3	2	4	-	-	1	2	-	2	-	-	-
Female	8	2	2	4	1	2	2	2	1	1	-	-
White, both sexes	6	3	3	3	1	3	2	1	3	1	-	-
Male	2	2	1	-	-	1	1	-	2	-	-	-
Female	4	1	2	3	1	2	1	1	1	1	-	-
Black, both sexes	5	1	3	1	-	-	2	1	-	-	-	-
Male	1	-	3	-	-	-	1	-	-	-	-	-
Female	4	1	-	1	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Missouri												
All races, both sexes	21	13	16	9	11	17	12	9	3	1	-	-
Male	8	3	5	5	3	5	3	1	-	-	-	-
Female	13	10	11	4	8	12	9	8	3	1	-	-
White, both sexes	16	9	15	8	11	16	11	8	3	1	-	-
Male	6	2	5	5	3	5	3	1	-	-	-	-
Female	10	7	10	3	8	11	8	7	3	1	-	-
Black, both sexes	5	3	1	1	-	-	1	1	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	3	2	1	1	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	1	1	1	2	2	1	-	2	1	1	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	1	1	1	1	2	1	-	1	1	1	-	-
White, both sexes	1	1	-	2	2	1	-	2	1	1	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	1	1	-	1	2	1	-	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Nebraska												
All races, both sexes	3	3	4	3	2	1	4	3	1	-	-	-
Male	2	1	1	2	1	1	3	-	-	-	-	-
Female	1	2	3	1	1	-	1	3	1	-	-	-
White, both sexes	3	2	4	2	2	1	4	3	1	-	-	-
Male	2	1	1	2	1	1	3	-	-	-	-	-
Female	1	1	3	-	1	-	1	3	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	4	2	6	5	7	4	4	-	-	-	-	-
Male	1	1	1	2	4	2	1	-	-	-	-	-
Female	3	1	5	3	3	2	3	-	-	-	-	-
White, both sexes	3	1	5	5	5	4	4	-	-	-	-	-
Male	1	1	-	2	2	2	1	-	-	-	-	-
Female	2	-	5	3	3	2	3	-	-	-	-	-
Black, both sexes	1	1	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
New Hampshire												
All races, both sexes	5	4	1	1	1	1	4	3	1	-	-	-
Male	1	1	-	-	-	1	3	1	1	-	-	-
Female	4	3	1	1	1	-	1	2	-	-	-	-
White, both sexes	5	4	1	1	1	1	4	3	1	-	-	-
Male	1	1	-	-	-	1	3	1	1	-	-	-
Female	4	3	1	1	1	-	1	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	23	23	6	13	12	15	25	15	6	2	-	-
Male	7	13	2	6	4	4	11	8	1	-	-	-
Female	16	10	4	7	8	11	14	7	5	2	-	-
White, both sexes	17	17	4	9	10	12	23	14	5	2	-	-
Male	6	10	1	4	4	4	10	7	1	-	-	-
Female	11	7	3	5	6	8	13	7	4	2	-	-
Black, both sexes	6	5	-	3	-	3	2	-	1	-	-	-
Male	1	2	-	1	-	-	1	-	-	-	-	-
Female	5	3	-	2	-	3	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	2	1	2	-	-	1	-	-	-	-
Male	-	1	1	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	2	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
New Mexico												
All races, both sexes	4	3	2	1	4	5	3	1	1	-	-	-
Male	1	1	-	-	-	1	1	-	-	-	-	-
Female	3	2	2	1	4	4	2	1	1	-	-	-
White, both sexes	4	3	2	1	4	4	3	1	1	-	-	-
Male	1	1	-	-	-	1	1	-	-	-	-	-
Female	3	2	2	1	4	3	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	40	37	28	23	27	29	33	27	14	6	-	-
Male	19	11	10	5	12	7	9	8	4	1	-	-
Female	21	26	18	18	15	22	24	19	10	5	-	-
White, both sexes	30	31	25	20	19	25	28	26	13	4	-	-
Male	15	10	9	5	9	6	7	8	4	1	-	-
Female	15	21	16	15	10	19	21	18	9	3	-	-
Black, both sexes	8	4	2	1	6	3	4	1	1	1	-	-
Male	3	1	-	-	3	1	1	-	-	-	-	-
Female	5	3	2	1	3	2	3	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	1	2	1	1	1	-	-	1	-	-
Male	1	-	1	-	-	-	1	-	-	-	-	-
Female	1	2	-	2	1	1	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
North Carolina												
All races, both sexes	21	11	19	15	18	20	18	19	6	1	1	-
Male	6	7	6	6	1	8	3	4	1	-	-	-
Female	15	4	13	9	17	12	15	15	5	1	1	-
White, both sexes	8	8	17	11	16	18	15	16	5	1	-	-
Male	1	5	6	4	1	7	3	4	1	-	-	-
Female	7	3	11	7	15	11	12	12	4	1	-	-
Black, both sexes	12	3	1	4	2	2	3	3	1	-	1	-
Male	4	2	-	2	-	1	-	-	-	-	-	-
Female	8	1	1	2	2	1	3	3	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	3	1	1	1	1	3	-	3	-	1	-	-
Male	2	1	-	-	1	1	-	-	-	-	-	-
Female	1	-	1	1	-	2	-	3	-	1	-	-
White, both sexes	3	1	1	1	1	3	-	3	-	1	-	-
Male	2	1	-	-	1	1	-	-	-	-	-	-
Female	1	-	1	1	-	2	-	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Ohio												
All races, both sexes	17	18	24	22	22	25	17	15	16	3	-	-
Male	3	5	13	9	10	5	6	3	3	1	-	-
Female	14	13	11	13	12	20	11	12	13	2	-	-
White, both sexes	16	15	22	18	18	23	15	14	16	1	-	-
Male	2	4	13	6	9	5	5	3	3	1	-	-
Female	14	11	9	12	9	18	10	11	13	-	-	-
Black, both sexes	1	3	2	4	4	2	1	1	-	2	-	-
Male	1	1	-	3	1	-	1	-	-	-	-	-
Female	-	2	2	1	3	2	-	1	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	10	12	9	5	8	2	6	3	4	-	-	-
Male	5	4	2	3	3	1	3	1	1	-	-	-
Female	5	8	7	2	5	1	3	2	3	-	-	-
White, both sexes	8	10	9	4	4	2	5	3	4	-	-	-
Male	4	4	2	2	1	1	2	1	1	-	-	-
Female	4	6	7	2	3	1	3	2	3	-	-	-
Black, both sexes	2	1	-	-	2	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	1	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	2	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Oregon												
All races, both sexes	6	9	9	1	7	9	10	4	6	1	1	-
Male	3	4	4	-	4	1	1	-	2	-	1	-
Female	3	5	5	1	3	8	9	4	4	1	-	-
White, both sexes	6	8	7	1	5	9	9	4	6	1	1	-
Male	3	4	3	-	3	1	1	-	2	-	1	-
Female	3	4	4	1	2	8	8	4	4	1	-	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	34	24	16	20	33	27	31	28	11	5	-	-
Male	11	14	7	9	13	10	10	9	2	1	-	-
Female	23	10	9	11	20	17	21	19	9	4	-	-
White, both sexes	25	21	11	18	31	25	30	26	10	4	-	-
Male	8	12	5	8	13	9	10	8	2	1	-	-
Female	17	9	6	10	18	16	20	18	8	3	-	-
Black, both sexes	7	2	5	2	2	2	1	2	1	-	-	-
Male	3	1	2	1	-	1	-	1	-	-	-	-
Female	4	1	3	1	2	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	-	-	-	-	-	-	1	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Rhode Island												
All races, both sexes	2	2	-	2	-	3	3	1	-	1	-	-
Male	1	1	-	1	-	2	-	-	-	-	-	-
Female	1	1	-	1	-	1	3	1	-	1	-	-
White, both sexes	2	2	-	1	-	3	3	1	-	1	-	-
Male	1	1	-	-	-	2	-	-	-	-	-	-
Female	1	1	-	1	-	1	3	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	15	11	10	8	11	8	9	6	-	1	-	-
Male	7	7	2	3	6	4	-	3	-	-	-	-
Female	8	4	8	5	5	4	9	3	-	1	-	-
White, both sexes	9	9	6	5	9	8	7	5	-	1	-	-
Male	4	5	1	1	4	4	-	2	-	-	-	-
Female	5	4	5	4	5	4	7	3	-	1	-	-
Black, both sexes	6	2	4	3	2	-	2	1	-	-	-	-
Male	3	2	1	2	2	-	-	1	-	-	-	-
Female	3	-	3	1	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
South Dakota												
All races, both sexes	3	-	5	-	4	2	-	-	-	1	-	-
Male	3	-	2	-	-	1	-	-	-	-	-	-
Female	-	-	3	-	4	1	-	-	-	1	-	-
White, both sexes	3	-	5	-	4	2	-	-	-	1	-	-
Male	3	-	2	-	-	1	-	-	-	-	-	-
Female	-	-	3	-	4	1	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	15	18	9	15	12	14	14	11	2	-	-	-
Male	7	9	3	8	4	4	2	3	-	-	-	-
Female	8	9	6	7	8	10	12	8	2	-	-	-
White, both sexes	11	16	6	13	11	12	13	11	2	-	-	-
Male	5	8	1	6	4	4	2	3	-	-	-	-
Female	6	8	5	7	7	8	11	8	2	-	-	-
Black, both sexes	4	2	3	1	1	2	1	-	-	-	-	-
Male	2	1	2	1	-	-	-	-	-	-	-	-
Female	2	1	1	-	1	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Texas												
All races, both sexes	43	53	37	29	32	32	27	10	14	1	-	-
Male	21	20	12	13	14	11	6	2	2	1	-	-
Female	22	33	25	16	18	21	21	8	12	-	-	-
White, both sexes	36	45	33	26	26	26	25	10	14	1	-	-
Male	17	16	12	12	11	9	5	2	2	1	-	-
Female	19	29	21	14	15	17	20	8	12	-	-	-
Black, both sexes	6	6	4	1	4	5	2	-	-	-	-	-
Male	3	4	-	-	2	2	1	-	-	-	-	-
Female	3	2	4	1	2	3	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	-	2	2	1	-	-	-	-	-	-
Male	1	-	-	1	1	-	-	-	-	-	-	-
Female	-	2	-	1	1	1	-	-	-	-	-	-
Utah												
All races, both sexes	7	3	5	3	5	5	3	-	2	1	-	-
Male	3	1	1	1	1	1	1	-	2	1	-	-
Female	4	2	4	2	4	4	2	-	-	-	-	-
White, both sexes	7	3	5	3	5	5	3	-	2	-	-	-
Male	3	1	1	1	1	1	1	-	2	-	-	-
Female	4	2	4	2	4	4	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Vermont												
All races, both sexes	1	1	1	-	1	2	-	2	-	1	-	-
Male	-	1	-	-	-	1	-	2	-	-	-	-
Female	1	-	1	-	1	1	-	-	-	1	-	-
White, both sexes	1	1	1	-	1	2	-	2	-	1	-	-
Male	-	1	-	-	-	1	-	2	-	-	-	-
Female	1	-	1	-	1	1	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	9	19	14	6	11	19	12	9	6	3	-	-
Male	3	6	5	1	5	3	2	1	1	-	-	-
Female	6	13	9	5	6	16	10	8	5	3	-	-
White, both sexes	6	15	12	6	8	15	8	9	5	3	-	-
Male	2	6	5	1	2	3	1	1	1	-	-	-
Female	4	9	7	5	6	12	7	8	4	3	-	-
Black, both sexes	3	3	2	-	3	4	3	-	1	-	-	-
Male	1	-	-	-	3	-	-	-	-	-	-	-
Female	2	3	2	-	-	4	3	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Washington												
All races, both sexes	17	18	8	17	10	9	12	9	3	3	1	-
Male	8	7	2	5	3	1	3	3	2	-	-	-
Female	9	11	6	12	7	8	9	6	1	3	1	-
White, both sexes	14	18	7	13	9	9	11	9	3	3	1	-
Male	6	7	2	3	3	1	3	3	2	-	-	-
Female	8	11	5	10	6	8	8	6	1	3	1	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	1	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	3	1	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	2	1	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	6	7	5	4	6	6	7	-	2	1	-	-
Male	4	1	2	2	2	1	2	-	-	-	-	-
Female	2	6	3	2	4	5	5	-	2	1	-	-
White, both sexes	6	7	5	4	5	6	7	-	2	1	-	-
Male	4	1	2	2	1	1	2	-	-	-	-	-
Female	2	6	3	2	4	5	5	-	2	1	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Subarachnoid hemorrhage (I60)												
Wisconsin												
All races, both sexes	13	20	7	7	5	10	16	10	4	-	-	-
Male	4	9	2	3	1	4	8	4	1	-	-	-
Female	9	11	5	4	4	6	8	6	3	-	-	-
White, both sexes	10	18	7	7	5	10	15	10	4	-	-	-
Male	4	9	2	3	1	4	8	4	1	-	-	-
Female	6	9	5	4	4	6	7	6	3	-	-	-
Black, both sexes	2	2	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	2	1	-	1	1	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	1	-	1	1	-	1	-	1	-	-
White, both sexes	-	2	1	-	1	1	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	2	1	-	1	1	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
United States												
All races, both sexes	27,785	45	12	16	16	19	42	64	123	252	623	1,090
Male	13,077	28	11	9	10	9	22	33	81	151	351	617
Female	14,708	17	1	7	6	10	20	31	42	101	272	473
White, both sexes	22,670	31	8	10	11	13	30	39	74	158	363	636
Male	10,515	20	7	5	6	6	16	20	52	94	206	374
Female	12,155	11	1	5	5	7	14	19	22	64	157	262
Black, both sexes	3,960	13	3	6	3	6	10	22	42	78	223	403
Male	2,009	7	3	4	2	3	6	11	25	47	122	216
Female	1,951	6	-	2	1	3	4	11	17	31	101	187
American Indian 1/, both sexes	140	1	-	-	-	-	1	1	-	5	9	6
Male	76	1	-	-	-	-	-	1	-	3	8	2
Female	64	-	-	-	-	-	1	-	-	2	1	4
Asian or Pacific Islander 2/, both sexes	1,015	-	1	-	2	-	1	2	7	11	28	45
Male	477	-	1	-	2	-	-	1	4	7	15	25
Female	538	-	-	-	-	-	1	1	3	4	13	20
Alabama												
All races, both sexes	634	1	-	-	-	1	1	1	7	7	13	36
Male	296	-	-	-	-	1	-	1	4	4	9	18
Female	338	1	-	-	-	-	1	-	3	3	4	18
White, both sexes	444	-	-	-	-	-	1	1	2	6	8	12
Male	206	-	-	-	-	-	-	1	2	3	5	6
Female	238	-	-	-	-	-	1	-	-	3	3	6
Black, both sexes	188	1	-	-	-	1	-	-	5	1	5	24
Male	88	-	-	-	-	1	-	-	2	1	4	12
Female	100	1	-	-	-	-	-	-	3	-	1	12
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Alaska												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	2	-
Male	20	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	402	-	-	1	-	-	-	3	-	4	11	16
Male	192	-	-	-	-	-	-	1	-	1	6	8
Female	210	-	-	1	-	-	-	2	-	3	5	8
White, both sexes	377	-	-	1	-	-	-	3	-	4	11	14
Male	181	-	-	-	-	-	-	1	-	1	6	8
Female	196	-	-	1	-	-	-	2	-	3	5	6
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	2
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Arkansas												
All races, both sexes	307	-	-	-	-	-	-	-	1	5	10	17
Male	137	-	-	-	-	-	-	-	-	1	6	13
Female	170	-	-	-	-	-	-	-	1	4	4	4
White, both sexes	255	-	-	-	-	-	-	-	-	3	7	10
Male	112	-	-	-	-	-	-	-	-	1	4	9
Female	143	-	-	-	-	-	-	-	-	2	3	1
Black, both sexes	50	-	-	-	-	-	-	-	1	2	3	7
Male	24	-	-	-	-	-	-	-	-	-	2	4
Female	26	-	-	-	-	-	-	-	1	2	1	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	2,947	2	3	1	1	2	4	13	12	37	79	126
Male	1,397	-	3	1	1	-	2	7	8	23	51	76
Female	1,550	2	-	-	-	2	2	6	4	14	28	50
White, both sexes	2,238	2	1	1	-	2	4	9	7	24	51	86
Male	1,056	-	1	1	-	-	2	5	6	17	33	56
Female	1,182	2	-	-	-	2	2	4	1	7	18	30
Black, both sexes	261	-	1	-	1	-	-	3	2	9	18	21
Male	125	-	1	-	1	-	-	1	-	4	11	10
Female	136	-	-	-	-	-	-	2	2	5	7	11
American Indian 1/, both sexes	20	-	-	-	-	-	-	-	-	1	1	2
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	10	-	-	-	-	-	-	-	-	1	-	2
Asian or Pacific Islander 2/, both sexes	428	-	1	-	-	-	-	1	3	3	9	17
Male	206	-	1	-	-	-	-	1	2	2	6	10
Female	222	-	-	-	-	-	-	-	1	1	3	7

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Colorado												
All races, both sexes	298	1	-	-	-	-	2	2	2	4	9	8
Male	158	-	-	-	-	-	1	-	2	2	6	5
Female	140	1	-	-	-	-	1	2	-	2	3	3
White, both sexes	280	1	-	-	-	-	2	2	2	2	8	5
Male	148	-	-	-	-	-	1	-	2	1	6	3
Female	132	1	-	-	-	-	1	2	-	1	2	2
Black, both sexes	11	-	-	-	-	-	-	-	-	-	1	2
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	2	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	1	-	-
Connecticut												
All races, both sexes	306	-	-	-	-	1	-	1	2	1	4	6
Male	138	-	-	-	-	1	-	-	2	1	3	3
Female	168	-	-	-	-	-	-	1	-	-	1	3
White, both sexes	274	-	-	-	-	1	-	1	1	1	3	4
Male	122	-	-	-	-	1	-	-	1	1	3	2
Female	152	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	28	-	-	-	-	-	-	-	1	-	1	2
Male	13	-	-	-	-	-	-	-	1	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Delaware												
All races, both sexes	85	-	-	-	-	-	-	-	-	1	-	3
Male	40	-	-	-	-	-	-	-	-	-	-	2
Female	45	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	69	-	-	-	-	-	-	-	-	-	-	2
Male	32	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	16	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	63	-	-	-	-	-	-	-	-	2	-	8
Male	31	-	-	-	-	-	-	-	-	1	-	6
Female	32	-	-	-	-	-	-	-	-	1	-	2
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	50	-	-	-	-	-	-	-	-	2	-	8
Male	23	-	-	-	-	-	-	-	-	1	-	6
Female	27	-	-	-	-	-	-	-	-	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Florida												
All races, both sexes	1,874	6	1	-	2	2	4	5	11	18	35	62
Male	944	3	1	-	1	-	3	3	9	11	21	38
Female	930	3	-	-	1	2	1	2	2	7	14	24
White, both sexes	1,572	5	1	-	2	1	2	2	5	10	17	40
Male	788	3	1	-	1	-	1	-	4	7	11	25
Female	784	2	-	-	1	1	1	2	1	3	6	15
Black, both sexes	273	1	-	-	-	1	2	3	6	8	17	22
Male	143	-	-	-	-	-	2	3	5	4	10	13
Female	130	1	-	-	-	1	-	-	1	4	7	9
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	1	-
Georgia												
All races, both sexes	794	5	1	1	1	1	-	2	7	6	29	56
Male	371	4	1	-	-	-	-	-	5	4	17	34
Female	423	1	-	1	1	1	-	2	2	2	12	22
White, both sexes	515	4	1	1	-	1	-	2	2	2	18	26
Male	227	3	1	-	-	-	-	-	1	2	12	16
Female	288	1	-	1	-	1	-	2	1	-	6	10
Black, both sexes	265	1	-	-	1	-	-	-	5	4	11	30
Male	139	1	-	-	-	-	-	-	4	2	5	18
Female	126	-	-	-	1	-	-	-	1	2	6	12
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Hawaii												
All races, both sexes	154	-	-	-	-	-	-	-	-	1	2	5
Male	81	-	-	-	-	-	-	-	-	1	-	1
Female	73	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	128	-	-	-	-	-	-	-	-	1	2	4
Male	68	-	-	-	-	-	-	-	-	1	-	1
Female	60	-	-	-	-	-	-	-	-	-	2	3
Idaho												
All races, both sexes	101	-	-	-	-	-	-	-	-	-	2	1
Male	47	-	-	-	-	-	-	-	-	-	2	1
Female	54	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	100	-	-	-	-	-	-	-	-	-	2	1
Male	46	-	-	-	-	-	-	-	-	-	2	1
Female	54	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Illinois												
All races, both sexes	1,144	4	-	1	2	1	2	-	4	10	22	36
Male	531	3	-	-	2	1	2	-	2	5	11	19
Female	613	1	-	1	-	-	-	-	2	5	11	17
White, both sexes	918	3	-	1	1	-	2	-	3	2	12	18
Male	415	2	-	-	1	-	2	-	1	-	7	10
Female	503	1	-	1	-	-	-	-	2	2	5	8
Black, both sexes	201	1	-	-	1	1	-	-	1	7	9	17
Male	102	1	-	-	1	1	-	-	1	5	3	8
Female	99	-	-	-	-	-	-	-	-	2	6	9
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	25	-	-	-	-	-	-	-	-	1	1	1
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	11	-	-	-	-	-	-	-	-	1	-	-
Indiana												
All races, both sexes	604	-	-	-	1	-	1	-	2	7	17	16
Male	267	-	-	-	1	-	1	-	2	3	7	5
Female	337	-	-	-	-	-	-	-	-	4	10	11
White, both sexes	534	-	-	-	1	-	-	-	1	4	14	12
Male	233	-	-	-	1	-	-	-	1	2	5	4
Female	301	-	-	-	-	-	-	-	-	2	9	8
Black, both sexes	67	-	-	-	-	-	1	-	1	3	2	4
Male	32	-	-	-	-	-	1	-	1	1	1	1
Female	35	-	-	-	-	-	-	-	-	2	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Iowa												
All races, both sexes	280	1	-	1	-	-	2	-	2	1	5	7
Male	128	1	-	-	-	-	-	-	1	-	1	3
Female	152	-	-	1	-	-	2	-	1	1	4	4
White, both sexes	276	1	-	1	-	-	1	-	2	1	5	7
Male	126	1	-	-	-	-	-	-	1	-	1	3
Female	150	-	-	1	-	-	1	-	1	1	4	4
Black, both sexes	3	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	277	3	-	-	-	-	1	-	-	2	7	14
Male	132	3	-	-	-	-	-	-	-	2	3	7
Female	145	-	-	-	-	-	1	-	-	-	4	7
White, both sexes	247	2	-	-	-	-	1	-	-	2	5	9
Male	114	2	-	-	-	-	-	-	-	2	1	4
Female	133	-	-	-	-	-	1	-	-	-	4	5
Black, both sexes	23	1	-	-	-	-	-	-	-	-	-	4
Male	16	1	-	-	-	-	-	-	-	-	-	3
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	2	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Kentucky												
All races, both sexes	444	1	-	-	-	-	-	-	2	2	7	20
Male	195	1	-	-	-	-	-	-	1	2	2	8
Female	249	-	-	-	-	-	-	-	1	-	5	12
White, both sexes	402	1	-	-	-	-	-	-	2	2	5	14
Male	173	1	-	-	-	-	-	-	1	2	2	5
Female	229	-	-	-	-	-	-	-	1	-	3	9
Black, both sexes	39	-	-	-	-	-	-	-	-	-	2	6
Male	20	-	-	-	-	-	-	-	-	-	-	3
Female	19	-	-	-	-	-	-	-	-	-	2	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	490	3	-	-	-	-	1	4	1	3	17	26
Male	232	2	-	-	-	-	-	1	1	2	11	13
Female	258	1	-	-	-	-	1	3	-	1	6	13
White, both sexes	305	2	-	-	-	-	1	-	1	-	4	7
Male	129	1	-	-	-	-	-	-	1	-	3	3
Female	176	1	-	-	-	-	1	-	-	-	1	4
Black, both sexes	182	1	-	-	-	-	-	4	-	3	13	19
Male	101	1	-	-	-	-	-	1	-	2	8	10
Female	81	-	-	-	-	-	-	3	-	1	5	9
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Maine												
All races, both sexes	123	-	-	-	-	-	-	-	-	-	2	4
Male	58	-	-	-	-	-	-	-	-	-	1	1
Female	65	-	-	-	-	-	-	-	-	-	1	3
White, both sexes	123	-	-	-	-	-	-	-	-	-	2	4
Male	58	-	-	-	-	-	-	-	-	-	1	1
Female	65	-	-	-	-	-	-	-	-	-	1	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	498	-	1	-	-	1	-	1	-	5	15	26
Male	236	-	-	-	-	1	-	1	-	3	9	15
Female	262	-	1	-	-	-	-	-	-	2	6	11
White, both sexes	324	-	1	-	-	-	-	-	-	2	1	7
Male	153	-	-	-	-	-	-	-	-	1	1	4
Female	171	-	1	-	-	-	-	-	-	1	-	3
Black, both sexes	164	-	-	-	-	1	-	1	-	3	12	19
Male	79	-	-	-	-	1	-	1	-	2	7	11
Female	85	-	-	-	-	-	-	-	-	1	5	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	2	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	6	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Massachusetts												
All races, both sexes	585	-	-	-	-	-	1	1	5	1	4	14
Male	277	-	-	-	-	-	1	-	5	-	4	10
Female	308	-	-	-	-	-	-	1	-	1	-	4
White, both sexes	537	-	-	-	-	-	1	-	3	1	4	10
Male	256	-	-	-	-	-	1	-	3	-	4	8
Female	281	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	33	-	-	-	-	-	-	1	1	-	-	3
Male	15	-	-	-	-	-	-	-	1	-	-	1
Female	18	-	-	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	1	-	-	1
Male	6	-	-	-	-	-	-	-	1	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	970	-	-	-	-	-	-	2	4	14	15	19
Male	453	-	-	-	-	-	-	2	1	7	5	8
Female	517	-	-	-	-	-	-	-	3	7	10	11
White, both sexes	789	-	-	-	-	-	-	1	1	8	10	9
Male	368	-	-	-	-	-	-	1	1	5	4	5
Female	421	-	-	-	-	-	-	-	-	3	6	4
Black, both sexes	167	-	-	-	-	-	-	-	2	6	4	10
Male	81	-	-	-	-	-	-	-	-	2	1	3
Female	86	-	-	-	-	-	-	-	2	4	3	7
American Indian 1/, both sexes	4	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	1	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Minnesota												
All races, both sexes	466	-	-	-	-	-	-	1	1	2	11	12
Male	208	-	-	-	-	-	-	-	1	1	8	6
Female	258	-	-	-	-	-	-	1	-	1	3	6
White, both sexes	437	-	-	-	-	-	-	-	1	2	8	9
Male	193	-	-	-	-	-	-	-	1	1	6	4
Female	244	-	-	-	-	-	-	-	-	1	2	5
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	2
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	1	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	10	-	-	-	-	-	-	1	-	-	1	-
Mississippi												
All races, both sexes	310	-	-	1	-	1	-	2	-	7	15	23
Male	164	-	-	1	-	-	-	1	-	5	11	7
Female	146	-	-	-	-	1	-	1	-	2	4	16
White, both sexes	180	-	-	-	-	-	-	2	-	2	6	9
Male	95	-	-	-	-	-	-	1	-	1	4	4
Female	85	-	-	-	-	-	-	1	-	1	2	5
Black, both sexes	126	-	-	1	-	1	-	-	-	5	9	14
Male	67	-	-	1	-	-	-	-	-	4	7	3
Female	59	-	-	-	-	1	-	-	-	1	2	11
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Missouri												
All races, both sexes	601	-	1	1	-	2	-	5	2	3	14	24
Male	271	-	1	1	-	1	-	1	-	3	9	13
Female	330	-	-	-	-	1	-	4	2	-	5	11
White, both sexes	503	-	-	-	-	2	-	5	1	1	7	12
Male	216	-	-	-	-	1	-	1	-	1	4	7
Female	287	-	-	-	-	1	-	4	1	-	3	5
Black, both sexes	93	-	1	1	-	-	-	-	1	1	7	12
Male	52	-	1	1	-	-	-	-	-	1	5	6
Female	41	-	-	-	-	-	-	-	1	-	2	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	75	-	-	-	-	-	-	-	1	2	-	2
Male	36	-	-	-	-	-	-	-	1	-	-	2
Female	39	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	68	-	-	-	-	-	-	-	1	1	-	2
Male	34	-	-	-	-	-	-	-	1	-	-	2
Female	34	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Nebraska												
All races, both sexes	178	-	-	-	-	-	-	-	3	-	2	4
Male	85	-	-	-	-	-	-	-	2	-	-	3
Female	93	-	-	-	-	-	-	-	1	-	2	1
White, both sexes	170	-	-	-	-	-	-	-	3	-	2	2
Male	82	-	-	-	-	-	-	-	2	-	-	2
Female	88	-	-	-	-	-	-	-	1	-	2	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	218	-	-	-	-	-	-	2	-	3	7	15
Male	117	-	-	-	-	-	-	2	-	2	5	6
Female	101	-	-	-	-	-	-	-	-	1	2	9
White, both sexes	166	-	-	-	-	-	-	2	-	2	3	7
Male	92	-	-	-	-	-	-	2	-	1	2	2
Female	74	-	-	-	-	-	-	-	-	1	1	5
Black, both sexes	23	-	-	-	-	-	-	-	-	1	4	6
Male	12	-	-	-	-	-	-	-	-	1	3	3
Female	11	-	-	-	-	-	-	-	-	-	1	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	-	2
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	14	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
New Hampshire												
All races, both sexes	91	-	-	-	-	-	-	-	-	-	3	1
Male	40	-	-	-	-	-	-	-	-	-	1	1
Female	51	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	88	-	-	-	-	-	-	-	-	-	3	1
Male	39	-	-	-	-	-	-	-	-	-	1	1
Female	49	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	863	1	-	-	1	-	-	2	2	7	12	37
Male	397	-	-	-	1	-	-	1	2	6	10	23
Female	466	1	-	-	-	-	-	1	-	1	2	14
White, both sexes	702	1	-	-	-	-	-	-	1	4	8	21
Male	317	-	-	-	-	-	-	-	1	3	7	14
Female	385	1	-	-	-	-	-	-	-	1	1	7
Black, both sexes	132	-	-	-	-	-	-	2	1	3	4	15
Male	64	-	-	-	-	-	-	1	1	3	3	9
Female	68	-	-	-	-	-	-	1	-	-	1	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	1	-	-	-	-	-	-	1
Male	16	-	-	-	1	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
New Mexico												
All races, both sexes	140	1	-	-	-	-	-	-	-	-	3	13
Male	63	1	-	-	-	-	-	-	-	-	2	7
Female	77	-	-	-	-	-	-	-	-	-	1	6
White, both sexes	123	-	-	-	-	-	-	-	-	-	2	11
Male	52	-	-	-	-	-	-	-	-	-	1	6
Female	71	-	-	-	-	-	-	-	-	-	1	5
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	12	1	-	-	-	-	-	-	-	-	1	1
Male	7	1	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,505	3	-	2	1	1	1	1	4	7	23	39
Male	706	2	-	1	1	-	-	1	3	4	12	23
Female	799	1	-	1	-	1	1	-	1	3	11	16
White, both sexes	1,247	2	-	1	1	1	1	1	3	6	12	21
Male	577	1	-	1	1	-	-	1	3	4	7	12
Female	670	1	-	-	-	1	1	-	-	2	5	9
Black, both sexes	204	1	-	1	-	-	-	-	1	1	10	15
Male	106	1	-	-	-	-	-	-	-	-	5	9
Female	98	-	-	1	-	-	-	-	1	1	5	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	53	-	-	-	-	-	-	-	-	-	1	3
Male	22	-	-	-	-	-	-	-	-	-	-	2
Female	31	-	-	-	-	-	-	-	-	-	1	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
North Carolina												
All races, both sexes	981	2	1	-	-	2	2	4	5	9	27	42
Male	444	1	1	-	-	2	-	4	1	4	14	26
Female	537	1	-	-	-	-	2	-	4	5	13	16
White, both sexes	722	-	1	-	-	2	1	2	4	6	11	17
Male	326	-	1	-	-	2	-	2	1	1	4	12
Female	396	-	-	-	-	-	1	-	3	5	7	5
Black, both sexes	241	2	-	-	-	-	-	2	1	3	15	25
Male	111	1	-	-	-	-	-	2	-	3	9	14
Female	130	1	-	-	-	-	-	-	1	-	6	11
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	1	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	1	-	-	-	-	-
North Dakota												
All races, both sexes	66	-	-	-	-	-	-	-	1	-	-	-
Male	31	-	-	-	-	-	-	-	1	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	63	-	-	-	-	-	-	-	1	-	-	-
Male	29	-	-	-	-	-	-	-	1	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Ohio												
All races, both sexes	1,185	2	-	1	-	-	6	1	3	8	22	49
Male	520	2	-	1	-	-	3	1	1	4	10	25
Female	665	-	-	-	-	-	3	-	2	4	12	24
White, both sexes	1,040	2	-	-	-	-	3	1	1	7	14	35
Male	459	2	-	-	-	-	2	1	-	3	7	19
Female	581	-	-	-	-	-	1	-	1	4	7	16
Black, both sexes	135	-	-	1	-	-	3	-	2	1	8	14
Male	57	-	-	1	-	-	1	-	1	1	3	6
Female	78	-	-	-	-	-	2	-	1	-	5	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	370	-	1	1	1	-	2	1	2	1	8	16
Male	170	-	1	-	1	-	-	-	2	1	6	8
Female	200	-	-	1	-	-	2	1	-	-	2	8
White, both sexes	320	-	-	1	1	-	1	-	1	-	4	10
Male	144	-	-	-	1	-	-	-	1	-	2	6
Female	176	-	-	1	-	-	1	-	-	-	2	4
Black, both sexes	27	-	1	-	-	-	-	1	1	-	2	3
Male	16	-	1	-	-	-	-	-	1	-	2	1
Female	11	-	-	-	-	-	-	1	-	-	-	2
American Indian 1/, both sexes	18	-	-	-	-	-	1	-	-	1	1	1
Male	7	-	-	-	-	-	-	-	-	1	1	-
Female	11	-	-	-	-	-	1	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	2
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Oregon												
All races, both sexes	361	-	-	1	-	-	1	-	3	4	5	11
Male	182	-	-	1	-	-	1	-	3	3	3	7
Female	179	-	-	-	-	-	-	-	-	1	2	4
White, both sexes	329	-	-	1	-	-	1	-	2	4	4	9
Male	165	-	-	1	-	-	1	-	2	3	2	6
Female	164	-	-	-	-	-	-	-	-	1	2	3
Black, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	2
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	1,536	2	2	2	1	-	3	4	4	5	23	36
Male	705	1	2	1	-	-	2	2	2	4	11	16
Female	831	1	-	1	1	-	1	2	2	1	12	20
White, both sexes	1,349	1	2	-	1	-	2	3	3	5	14	19
Male	621	1	2	-	-	-	2	2	2	4	6	10
Female	728	-	-	-	1	-	-	1	1	1	8	9
Black, both sexes	167	1	-	2	-	-	1	1	1	-	8	16
Male	70	-	-	1	-	-	-	-	-	-	5	5
Female	97	1	-	1	-	-	1	1	1	-	3	11
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Rhode Island												
All races, both sexes	114	-	-	-	-	-	-	-	2	2	-	5
Male	54	-	-	-	-	-	-	-	-	1	-	3
Female	60	-	-	-	-	-	-	-	2	1	-	2
White, both sexes	106	-	-	-	-	-	-	-	1	2	-	2
Male	50	-	-	-	-	-	-	-	-	1	-	1
Female	56	-	-	-	-	-	-	-	1	1	-	1
Black, both sexes	6	-	-	-	-	-	-	-	1	-	-	2
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	2	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
South Carolina												
All races, both sexes	521	1	-	-	-	1	-	2	4	2	22	24
Male	260	-	-	-	-	1	-	1	3	2	10	18
Female	261	1	-	-	-	-	-	1	1	-	12	6
White, both sexes	326	-	-	-	-	1	-	-	3	-	9	10
Male	149	-	-	-	-	1	-	-	3	-	5	7
Female	177	-	-	-	-	-	-	-	-	-	4	3
Black, both sexes	189	1	-	-	-	-	-	2	1	2	13	14
Male	109	-	-	-	-	-	-	1	-	2	5	11
Female	80	1	-	-	-	-	-	1	1	-	8	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
South Dakota												
All races, both sexes	72	-	-	-	-	-	-	-	-	-	1	1
Male	38	-	-	-	-	-	-	-	-	-	-	1
Female	34	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	70	-	-	-	-	-	-	-	-	-	1	-
Male	37	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	691	-	-	-	-	1	-	2	4	8	20	37
Male	340	-	-	-	-	-	-	2	4	2	11	23
Female	351	-	-	-	-	1	-	-	-	6	9	14
White, both sexes	579	-	-	-	-	-	-	2	2	6	8	19
Male	280	-	-	-	-	-	-	2	2	1	3	12
Female	299	-	-	-	-	-	-	-	-	5	5	7
Black, both sexes	110	-	-	-	-	1	-	-	2	2	12	17
Male	59	-	-	-	-	-	-	-	2	1	8	11
Female	51	-	-	-	-	1	-	-	-	1	4	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Texas												
All races, both sexes	1,807	4	1	1	3	2	4	1	11	27	64	93
Male	870	2	1	1	-	1	4	-	6	19	36	61
Female	937	2	-	-	3	1	-	1	5	8	28	32
White, both sexes	1,546	2	1	1	3	2	3	-	7	21	43	68
Male	733	1	1	1	-	1	3	-	3	15	24	45
Female	813	1	-	-	3	1	-	-	4	6	19	23
Black, both sexes	224	2	-	-	-	-	1	1	2	3	18	23
Male	121	1	-	-	-	-	1	-	2	2	10	15
Female	103	1	-	-	-	-	-	1	-	1	8	8
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	2	2	3	2
Male	14	-	-	-	-	-	-	-	1	1	2	1
Female	21	-	-	-	-	-	-	-	1	1	1	1
Utah												
All races, both sexes	127	-	-	-	-	-	1	-	-	2	1	4
Male	62	-	-	-	-	-	-	-	-	2	1	2
Female	65	-	-	-	-	-	1	-	-	-	-	2
White, both sexes	119	-	-	-	-	-	1	-	-	2	1	3
Male	55	-	-	-	-	-	-	-	-	2	1	1
Female	64	-	-	-	-	-	1	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Vermont												
All races, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	41	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	751	-	-	-	1	-	2	1	4	14	22	40
Male	345	-	-	-	1	-	2	1	3	10	9	20
Female	406	-	-	-	-	-	-	-	1	4	13	20
White, both sexes	544	-	-	-	1	-	1	-	2	7	7	20
Male	241	-	-	-	1	-	1	-	1	5	4	9
Female	303	-	-	-	-	-	-	-	1	2	3	11
Black, both sexes	185	-	-	-	-	-	1	1	2	7	15	19
Male	97	-	-	-	-	-	1	1	2	5	5	11
Female	88	-	-	-	-	-	-	-	-	2	10	8
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Washington												
All races, both sexes	547	2	-	-	1	-	-	-	2	2	3	14
Male	252	2	-	-	1	-	-	-	1	1	1	9
Female	295	-	-	-	-	-	-	-	1	1	2	5
White, both sexes	486	2	-	-	-	-	-	-	2	1	2	12
Male	224	2	-	-	-	-	-	-	1	-	1	7
Female	262	-	-	-	-	-	-	-	1	1	1	5
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	47	-	-	-	1	-	-	-	-	1	1	2
Male	21	-	-	-	1	-	-	-	-	1	-	2
Female	26	-	-	-	-	-	-	-	-	-	1	-
West Virginia												
All races, both sexes	217	-	-	-	-	-	1	-	2	2	3	7
Male	99	-	-	-	-	-	-	-	1	-	2	5
Female	118	-	-	-	-	-	1	-	1	2	1	2
White, both sexes	211	-	-	-	-	-	1	-	2	2	3	6
Male	95	-	-	-	-	-	-	-	1	-	2	4
Female	116	-	-	-	-	-	1	-	1	2	1	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Intracerebral and other intracranial hemorrhage (I61-I62)												
Wisconsin												
All races, both sexes	499	-	-	1	-	-	-	-	1	3	4	13
Male	223	-	-	1	-	-	-	-	1	3	3	7
Female	276	-	-	-	-	-	-	-	-	-	1	6
White, both sexes	460	-	-	1	-	-	-	-	1	3	3	11
Male	199	-	-	1	-	-	-	-	1	3	2	6
Female	261	-	-	-	-	-	-	-	-	-	1	5
Black, both sexes	29	-	-	-	-	-	-	-	-	-	-	2
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	36	-	-	-	-	-	-	-	-	1	1	2
Male	18	-	-	-	-	-	-	-	-	1	-	1
Female	18	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	33	-	-	-	-	-	-	-	-	-	1	2
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
United States												
All races, both sexes	1,565	1,693	1,752	2,081	2,831	4,153	5,025	3,930	1,908	487	56	2
Male	925	1,026	992	1,142	1,465	1,971	2,088	1,457	547	130	10	2
Female	640	667	760	939	1,366	2,182	2,937	2,473	1,361	357	46	-
White, both sexes	928	1,098	1,276	1,629	2,371	3,644	4,528	3,584	1,752	443	44	-
Male	550	659	720	905	1,236	1,768	1,885	1,351	509	119	7	-
Female	378	439	556	724	1,135	1,876	2,643	2,233	1,243	324	37	-
Black, both sexes	551	494	370	350	331	338	333	233	110	28	11	2
Male	327	302	223	184	167	128	130	66	26	5	3	2
Female	224	192	147	166	164	210	203	167	84	23	8	-
American Indian 1/, both sexes	10	15	14	11	18	21	15	6	5	2	-	-
Male	4	13	7	6	8	10	6	4	3	-	-	-
Female	6	2	7	5	10	11	9	2	2	2	-	-
Asian or Pacific Islander 2/, both sexes	76	86	92	91	111	150	149	107	41	14	1	-
Male	44	52	42	47	54	65	67	36	9	6	-	-
Female	32	34	50	44	57	85	82	71	32	8	1	-
Alabama												
All races, both sexes	41	52	47	56	72	82	107	63	32	10	5	-
Male	21	30	30	27	37	36	40	26	8	2	2	-
Female	20	22	17	29	35	46	67	37	24	8	3	-
White, both sexes	15	29	22	42	58	68	90	57	26	7	-	-
Male	8	16	18	21	30	32	31	24	7	2	-	-
Female	7	13	4	21	28	36	59	33	19	5	-	-
Black, both sexes	26	23	25	13	14	14	17	5	6	3	5	-
Male	13	14	12	5	7	4	9	1	1	-	2	-
Female	13	9	13	8	7	10	8	4	5	3	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Alaska												
All races, both sexes	3	4	4	1	5	8	5	2	2	-	-	-
Male	2	4	4	-	2	4	1	1	1	-	-	-
Female	1	-	-	1	3	4	4	1	1	-	-	-
White, both sexes	2	3	3	1	2	6	3	2	2	-	-	-
Male	1	3	3	-	1	4	1	1	1	-	-	-
Female	1	-	-	1	1	2	2	1	1	-	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	2	1	-	-	-	-	-
Male	-	-	1	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	1	-	-	-	-	-
Arizona												
All races, both sexes	26	19	27	23	51	69	55	62	33	2	-	-
Male	21	11	16	16	31	31	22	21	6	1	-	-
Female	5	8	11	7	20	38	33	41	27	1	-	-
White, both sexes	22	14	26	23	47	66	54	60	31	1	-	-
Male	17	8	15	16	29	31	21	21	6	1	-	-
Female	5	6	11	7	18	35	33	39	25	-	-	-
Black, both sexes	2	-	-	-	1	1	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	1	-	-	-	-
American Indian 1/, both sexes	2	4	1	-	3	1	-	-	1	1	-	-
Male	2	3	1	-	2	-	-	-	-	-	-	-
Female	-	1	-	-	1	1	-	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	1	1	1	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	1	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Arkansas												
All races, both sexes	17	23	22	20	40	40	51	38	17	6	-	-
Male	8	13	10	8	19	17	21	16	5	-	-	-
Female	9	10	12	12	21	23	30	22	12	6	-	-
White, both sexes	8	17	16	17	37	37	47	34	16	6	-	-
Male	4	9	6	8	17	16	20	13	5	-	-	-
Female	4	8	10	9	20	21	27	21	11	6	-	-
Black, both sexes	8	5	6	3	3	3	4	4	1	-	-	-
Male	4	3	4	-	2	1	1	3	-	-	-	-
Female	4	2	2	3	1	2	3	1	1	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	182	194	203	227	302	388	524	402	192	46	6	1
Male	106	108	113	111	152	168	227	172	53	13	1	1
Female	76	86	90	116	150	220	297	230	139	33	5	-
White, both sexes	116	124	151	158	224	294	437	338	167	37	5	-
Male	65	68	86	81	111	128	187	152	46	10	1	-
Female	51	56	65	77	113	166	250	186	121	27	4	-
Black, both sexes	36	28	19	27	27	19	22	20	6	1	-	1
Male	23	15	9	11	15	6	8	6	3	-	-	1
Female	13	13	10	16	12	13	14	14	3	1	-	-
American Indian 1/, both sexes	2	3	1	3	3	1	1	2	-	-	-	-
Male	1	3	1	1	1	1	-	1	-	-	-	-
Female	1	-	-	2	2	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	28	39	32	39	48	74	64	42	19	8	1	-
Male	17	22	17	18	25	33	32	13	4	3	-	-
Female	11	17	15	21	23	41	32	29	15	5	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Colorado												
All races, both sexes	18	13	18	21	27	63	53	42	11	4	-	-
Male	10	8	12	15	16	30	27	20	3	1	-	-
Female	8	5	6	6	11	33	26	22	8	3	-	-
White, both sexes	16	10	15	21	26	61	53	41	11	4	-	-
Male	8	7	10	15	15	30	27	19	3	1	-	-
Female	8	3	5	6	11	31	26	22	8	3	-	-
Black, both sexes	2	2	1	-	1	2	-	-	-	-	-	-
Male	2	1	1	-	1	-	-	-	-	-	-	-
Female	-	1	-	-	-	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	13	9	7	22	44	42	64	55	28	7	-	-
Male	11	5	4	14	21	19	32	12	9	1	-	-
Female	2	4	3	8	23	23	32	43	19	6	-	-
White, both sexes	9	7	5	17	39	41	58	52	28	7	-	-
Male	7	3	3	11	19	19	30	12	9	1	-	-
Female	2	4	2	6	20	22	28	40	19	6	-	-
Black, both sexes	3	2	2	3	5	1	5	3	-	-	-	-
Male	3	2	1	1	2	-	2	-	-	-	-	-
Female	-	-	1	2	3	1	3	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	-	-	1	-	-	-	-	-
Male	1	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Delaware												
All races, both sexes	3	4	5	6	16	11	13	16	3	4	-	-
Male	1	3	3	2	8	6	4	11	-	-	-	-
Female	2	1	2	4	8	5	9	5	3	4	-	-
White, both sexes	-	1	4	5	14	10	11	15	3	4	-	-
Male	-	-	3	2	6	6	4	10	-	-	-	-
Female	-	1	1	3	8	4	7	5	3	4	-	-
Black, both sexes	3	3	1	1	2	1	2	1	-	-	-	-
Male	1	3	-	-	2	-	-	1	-	-	-	-
Female	2	-	1	1	-	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	7	7	7	5	3	7	6	11	-	-	-	-
Male	4	3	7	2	2	2	2	2	-	-	-	-
Female	3	4	-	3	1	5	4	9	-	-	-	-
White, both sexes	2	-	1	-	1	3	2	4	-	-	-	-
Male	2	-	1	-	1	2	1	1	-	-	-	-
Female	-	-	-	-	-	1	1	3	-	-	-	-
Black, both sexes	5	7	6	5	2	4	4	7	-	-	-	-
Male	2	3	6	2	1	-	1	1	-	-	-	-
Female	3	4	-	3	1	4	3	6	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Florida												
All races, both sexes	96	105	110	140	198	264	382	287	112	31	3	-
Male	61	70	63	79	112	132	164	120	44	9	-	-
Female	35	35	47	61	86	132	218	167	68	22	3	-
White, both sexes	62	61	86	107	170	244	356	263	104	31	3	-
Male	39	40	50	63	99	126	154	114	41	9	-	-
Female	23	21	36	44	71	118	202	149	63	22	3	-
Black, both sexes	32	40	19	30	26	19	21	19	7	-	-	-
Male	21	26	11	15	13	6	7	4	3	-	-	-
Female	11	14	8	15	13	13	14	15	4	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	5	3	2	1	4	5	-	-	-	-
Male	1	3	2	1	-	-	2	2	-	-	-	-
Female	1	-	3	2	2	1	2	3	-	-	-	-
Georgia												
All races, both sexes	59	57	62	68	90	101	108	88	44	7	1	-
Male	42	31	34	41	42	44	42	21	8	1	-	-
Female	17	26	28	27	48	57	66	67	36	6	1	-
White, both sexes	14	23	27	46	70	73	92	76	32	5	-	-
Male	8	9	15	28	35	34	38	18	7	-	-	-
Female	6	14	12	18	35	39	54	58	25	5	-	-
Black, both sexes	44	30	33	22	19	24	14	12	12	2	1	-
Male	33	21	19	13	6	8	4	3	1	1	-	-
Female	11	9	14	9	13	16	10	9	11	1	1	-
American Indian 1/, both sexes	-	-	1	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	1	-	-	3	2	-	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	3	1	-	-	2	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Hawaii												
All races, both sexes	12	11	14	9	15	21	28	25	7	4	-	-
Male	8	6	10	7	9	12	13	9	2	3	-	-
Female	4	5	4	2	6	9	15	16	5	1	-	-
White, both sexes	-	-	4	2	3	4	3	6	3	-	-	-
Male	-	-	3	2	2	4	-	1	1	-	-	-
Female	-	-	1	-	1	-	3	5	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	11	10	7	12	17	25	19	4	4	-	-
Male	8	6	7	5	7	8	13	8	1	3	-	-
Female	4	5	3	2	5	9	12	11	3	1	-	-
Idaho												
All races, both sexes	2	6	6	8	13	12	14	21	13	3	-	-
Male	1	4	3	4	6	6	8	8	3	1	-	-
Female	1	2	3	4	7	6	6	13	10	2	-	-
White, both sexes	2	6	6	8	13	11	14	21	13	3	-	-
Male	1	4	3	4	6	5	8	8	3	1	-	-
Female	1	2	3	4	7	6	6	13	10	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Illinois												
All races, both sexes	54	71	61	85	116	166	222	164	100	22	1	-
Male	33	41	35	45	64	93	94	50	26	5	-	-
Female	21	30	26	40	52	73	128	114	74	17	1	-
White, both sexes	31	41	44	55	98	147	201	147	90	21	1	-
Male	19	20	22	28	58	82	89	47	22	5	-	-
Female	12	21	22	27	40	65	112	100	68	16	1	-
Black, both sexes	22	26	15	27	15	16	19	15	8	1	-	-
Male	13	18	12	16	4	9	5	2	3	-	-	-
Female	9	8	3	11	11	7	14	13	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	4	2	3	3	3	2	2	2	-	-	-
Male	1	3	1	1	2	2	-	1	1	-	-	-
Female	-	1	1	2	1	1	2	1	1	-	-	-
Indiana												
All races, both sexes	33	34	41	54	56	113	99	77	36	16	1	-
Male	21	22	22	29	24	58	37	22	9	4	-	-
Female	12	12	19	25	32	55	62	55	27	12	1	-
White, both sexes	25	26	34	45	52	105	89	75	35	16	-	-
Male	17	18	17	26	21	52	35	21	9	4	-	-
Female	8	8	17	19	31	53	54	54	26	12	-	-
Black, both sexes	8	8	7	8	4	7	10	2	1	-	1	-
Male	4	4	5	3	3	5	2	1	-	-	-	-
Female	4	4	2	5	1	2	8	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Iowa												
All races, both sexes	11	9	17	27	20	40	65	43	24	5	-	-
Male	8	6	8	17	12	23	24	16	7	1	-	-
Female	3	3	9	10	8	17	41	27	17	4	-	-
White, both sexes	9	9	17	26	20	40	65	43	24	5	-	-
Male	7	6	8	16	12	23	24	16	7	1	-	-
Female	2	3	9	10	8	17	41	27	17	4	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	14	9	13	14	32	46	54	40	21	6	1	-
Male	12	7	7	7	18	19	24	15	6	2	-	-
Female	2	2	6	7	14	27	30	25	15	4	1	-
White, both sexes	7	7	11	12	29	45	49	40	21	6	1	-
Male	6	5	7	6	17	18	23	15	6	2	-	-
Female	1	2	4	6	12	27	26	25	15	4	1	-
Black, both sexes	7	2	2	2	1	1	3	-	-	-	-	-
Male	6	2	-	1	1	1	1	-	-	-	-	-
Female	1	-	2	1	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Kentucky												
All races, both sexes	28	29	29	33	48	83	82	56	20	4	-	-
Male	17	18	11	15	25	41	30	20	4	-	-	-
Female	11	11	18	18	23	42	52	36	16	4	-	-
White, both sexes	19	22	25	30	45	81	81	53	18	4	-	-
Male	11	15	9	12	23	39	29	20	4	-	-	-
Female	8	7	16	18	22	42	52	33	14	4	-	-
Black, both sexes	9	4	4	3	3	2	1	3	2	-	-	-
Male	6	1	2	3	2	2	1	-	-	-	-	-
Female	3	3	2	-	1	-	-	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	-	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	45	42	34	50	54	60	60	49	28	10	2	1
Male	26	24	20	24	29	23	27	15	9	3	1	1
Female	19	18	14	26	25	37	33	34	19	7	1	-
White, both sexes	13	13	20	32	38	48	51	44	24	5	2	-
Male	6	7	10	13	19	18	24	13	9	1	1	-
Female	7	6	10	19	19	30	27	31	15	4	1	-
Black, both sexes	31	29	13	18	16	11	9	5	4	5	-	1
Male	20	17	9	11	10	4	3	2	-	2	-	1
Female	11	12	4	7	6	7	6	3	4	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Maine												
All races, both sexes	3	8	8	7	15	18	23	20	13	1	1	-
Male	2	7	3	4	9	8	8	10	5	-	-	-
Female	1	1	5	3	6	10	15	10	8	1	1	-
White, both sexes	3	8	8	7	15	18	23	20	13	1	1	-
Male	2	7	3	4	9	8	8	10	5	-	-	-
Female	1	1	5	3	6	10	15	10	8	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	34	41	37	31	46	56	90	64	40	10	-	-
Male	17	26	22	16	23	17	43	26	12	5	-	-
Female	17	15	15	15	23	39	47	38	28	5	-	-
White, both sexes	13	21	16	15	29	44	77	54	35	9	-	-
Male	9	16	9	7	16	15	37	22	11	5	-	-
Female	4	5	7	8	13	29	40	32	24	4	-	-
Black, both sexes	21	19	20	16	16	12	11	8	4	1	-	-
Male	8	10	13	9	7	2	4	3	1	-	-	-
Female	13	9	7	7	9	10	7	5	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	1	-	2	2	1	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	-	1	1	-	1	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Massachusetts												
All races, both sexes	19	25	22	44	52	96	123	94	66	15	3	-
Male	9	18	13	23	29	45	51	37	24	8	-	-
Female	10	7	9	21	23	51	72	57	42	7	3	-
White, both sexes	16	20	17	37	46	91	120	91	63	15	2	-
Male	9	14	9	19	27	44	50	37	23	8	-	-
Female	7	6	8	18	19	47	70	54	40	7	2	-
Black, both sexes	2	4	4	5	4	3	2	1	2	-	1	-
Male	-	3	3	3	2	1	1	-	-	-	-	-
Female	2	1	1	2	2	2	1	1	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	1	2	2	2	1	2	1	-	-	-
Male	-	1	1	1	-	-	-	-	1	-	-	-
Female	1	-	-	1	2	2	1	2	-	-	-	-
Michigan												
All races, both sexes	62	56	58	68	101	170	173	139	69	16	4	-
Male	32	37	39	37	47	80	62	61	33	2	-	-
Female	30	19	19	31	54	90	111	78	36	14	4	-
White, both sexes	34	29	42	55	87	143	157	128	65	16	4	-
Male	19	19	29	27	42	68	56	57	33	2	-	-
Female	15	10	13	28	45	75	101	71	32	14	4	-
Black, both sexes	27	25	16	10	13	26	14	10	4	-	-	-
Male	13	17	10	8	5	12	6	4	-	-	-	-
Female	14	8	6	2	8	14	8	6	4	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	3	-	-	2	1	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	1	1	-	1	-	-	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Minnesota												
All races, both sexes	19	17	20	31	60	73	74	87	46	11	1	-
Male	11	13	9	20	27	40	30	29	10	2	1	-
Female	8	4	11	11	33	33	44	58	36	9	-	-
White, both sexes	15	16	15	26	57	72	73	86	45	11	1	-
Male	8	12	8	16	26	39	30	29	10	2	1	-
Female	7	4	7	10	31	33	43	57	35	9	-	-
Black, both sexes	2	1	2	3	-	-	-	-	-	-	-	-
Male	2	1	1	2	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	2	-	-	-	-	-	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	2	-	3	1	1	1	1	-	-	-
Male	1	-	-	-	1	1	-	-	-	-	-	-
Female	1	-	2	-	2	-	1	1	1	-	-	-
Mississippi												
All races, both sexes	30	23	28	29	25	39	42	29	14	1	1	-
Male	17	11	22	17	20	20	19	10	3	-	-	-
Female	13	12	6	12	5	19	23	19	11	1	1	-
White, both sexes	10	9	12	17	13	32	31	23	12	1	1	-
Male	7	5	10	9	10	17	16	8	3	-	-	-
Female	3	4	2	8	3	15	15	15	9	1	1	-
Black, both sexes	20	14	15	12	11	7	9	6	2	-	-	-
Male	10	6	11	8	10	3	2	2	-	-	-	-
Female	10	8	4	4	1	4	7	4	2	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Missouri												
All races, both sexes	30	41	42	38	65	80	111	90	39	12	1	-
Male	16	24	26	20	28	35	48	35	7	3	-	-
Female	14	17	16	18	37	45	63	55	32	9	1	-
White, both sexes	17	26	31	34	54	77	103	83	39	10	1	-
Male	5	16	18	19	25	33	42	34	7	3	-	-
Female	12	10	13	15	29	44	61	49	32	7	1	-
Black, both sexes	13	15	11	4	9	2	8	7	-	1	-	-
Male	11	8	8	1	2	1	6	1	-	-	-	-
Female	2	7	3	3	7	1	2	6	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	-	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	1	-	-
Montana												
All races, both sexes	4	3	2	5	6	13	15	13	8	-	1	-
Male	1	2	1	4	4	8	5	6	2	-	-	-
Female	3	1	1	1	2	5	10	7	6	-	1	-
White, both sexes	2	3	2	5	6	11	13	13	8	-	1	-
Male	1	2	1	4	4	7	4	6	2	-	-	-
Female	1	1	1	1	2	4	9	7	6	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	2	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	2	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Nebraska												
All races, both sexes	7	8	8	15	12	28	43	26	17	4	1	-
Male	5	5	3	8	8	11	21	10	6	2	1	-
Female	2	3	5	7	4	17	22	16	11	2	-	-
White, both sexes	7	6	5	15	12	27	43	26	17	4	1	-
Male	5	4	2	8	8	11	21	10	6	2	1	-
Female	2	2	3	7	4	16	22	16	11	2	-	-
Black, both sexes	-	2	1	-	-	-	-	-	-	-	-	-
Male	-	1	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Nevada												
All races, both sexes	18	20	19	23	26	33	25	20	7	-	-	-
Male	11	11	10	14	16	20	11	7	2	-	-	-
Female	7	9	9	9	10	13	14	13	5	-	-	-
White, both sexes	10	15	14	20	19	28	23	17	6	-	-	-
Male	6	9	9	12	11	19	11	6	2	-	-	-
Female	4	6	5	8	8	9	12	11	4	-	-	-
Black, both sexes	6	1	1	-	2	-	-	1	1	-	-	-
Male	3	-	-	-	2	-	-	-	-	-	-	-
Female	3	1	1	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	2	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	4	3	5	4	2	2	-	-	-	-
Male	2	1	1	2	3	1	-	1	-	-	-	-
Female	-	1	3	1	2	3	2	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
New Hampshire												
All races, both sexes	5	5	6	4	9	23	16	13	5	1	-	-
Male	2	3	5	3	6	11	5	-	2	1	-	-
Female	3	2	1	1	3	12	11	13	3	-	-	-
White, both sexes	4	5	5	4	9	23	15	13	5	1	-	-
Male	2	3	4	3	6	11	5	-	2	1	-	-
Female	2	2	1	1	3	12	10	13	3	-	-	-
Black, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
New Jersey												
All races, both sexes	45	44	50	64	83	115	157	142	84	16	1	-
Male	21	24	30	38	36	50	71	55	26	3	-	-
Female	24	20	20	26	47	65	86	87	58	13	1	-
White, both sexes	28	23	35	51	69	98	138	135	75	14	1	-
Male	12	11	22	28	32	43	64	54	23	3	-	-
Female	16	12	13	23	37	55	74	81	52	11	1	-
Black, both sexes	15	18	12	6	11	16	14	5	8	2	-	-
Male	7	10	7	3	3	7	6	1	3	-	-	-
Female	8	8	5	3	8	9	8	4	5	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	3	3	7	3	1	5	2	1	-	-	-
Male	2	3	1	7	1	-	1	-	-	-	-	-
Female	-	-	2	-	2	1	4	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
New Mexico												
All races, both sexes	8	10	10	13	16	15	27	13	6	5	-	-
Male	3	5	7	5	7	8	10	5	2	1	-	-
Female	5	5	3	8	9	7	17	8	4	4	-	-
White, both sexes	6	9	8	13	15	13	25	12	5	4	-	-
Male	2	4	6	5	7	6	9	4	1	1	-	-
Female	4	5	2	8	8	7	16	8	4	3	-	-
Black, both sexes	1	-	-	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	1	2	-	1	1	-	1	1	1	-	-
Male	-	1	1	-	-	1	-	1	1	-	-	-
Female	1	-	1	-	1	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	71	98	85	98	143	228	306	235	126	30	3	-
Male	44	66	48	49	74	116	137	87	33	4	1	-
Female	27	32	37	49	69	112	169	148	93	26	2	-
White, both sexes	47	71	62	73	119	198	274	212	111	28	3	-
Male	30	48	34	38	63	101	120	79	29	4	1	-
Female	17	23	28	35	56	97	154	133	82	24	2	-
Black, both sexes	19	25	19	23	19	22	21	15	10	2	-	-
Male	13	16	12	10	9	10	13	6	2	-	-	-
Female	6	9	7	13	10	12	8	9	8	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	2	4	2	5	8	11	7	5	-	-	-
Male	1	2	2	1	2	5	4	1	2	-	-	-
Female	4	-	2	1	3	3	7	6	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
North Carolina												
All races, both sexes	76	73	54	80	85	140	182	123	59	14	1	-
Male	45	46	27	38	45	61	74	35	15	5	-	-
Female	31	27	27	42	40	79	108	88	44	9	1	-
White, both sexes	39	34	34	53	65	120	160	106	53	14	-	-
Male	23	21	19	30	37	58	66	31	13	5	-	-
Female	16	13	15	23	28	62	94	75	40	9	-	-
Black, both sexes	36	36	19	24	18	18	19	16	6	-	1	-
Male	22	23	7	6	8	3	7	4	2	-	-	-
Female	14	13	12	18	10	15	12	12	4	-	1	-
American Indian 1/, both sexes	-	-	-	2	2	1	2	-	-	-	-	-
Male	-	-	-	2	-	-	1	-	-	-	-	-
Female	-	-	-	-	2	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	1	1	-	1	1	1	-	-	-	-
Male	-	2	1	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	-	1	1	1	-	-	-	-
North Dakota												
All races, both sexes	1	2	2	4	8	12	14	13	9	-	-	-
Male	-	2	2	4	1	5	6	6	4	-	-	-
Female	1	-	-	-	7	7	8	7	5	-	-	-
White, both sexes	1	1	2	4	8	10	14	13	9	-	-	-
Male	-	1	2	4	1	4	6	6	4	-	-	-
Female	1	-	-	-	7	6	8	7	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Ohio												
All races, both sexes	56	65	70	79	106	201	245	160	85	24	2	-
Male	32	39	35	43	54	97	85	57	23	8	-	-
Female	24	26	35	36	52	104	160	103	62	16	2	-
White, both sexes	43	49	62	69	91	184	227	145	83	22	2	-
Male	27	30	31	38	46	92	77	53	23	8	-	-
Female	16	19	31	31	45	92	150	92	60	14	2	-
Black, both sexes	13	16	8	7	12	16	15	15	2	2	-	-
Male	5	9	4	4	8	4	6	4	-	-	-	-
Female	8	7	4	3	4	12	9	11	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	3	1	3	-	-	-	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	-	-	-	2	3	-	1	-	-	-	-	-
Oklahoma												
All races, both sexes	12	30	26	26	40	58	52	59	24	9	1	-
Male	7	22	14	12	18	25	22	25	5	1	-	-
Female	5	8	12	14	22	33	30	34	19	8	1	-
White, both sexes	8	25	20	25	36	51	48	56	23	9	1	-
Male	4	19	11	11	17	23	19	24	5	1	-	-
Female	4	6	9	14	19	28	29	32	18	8	1	-
Black, both sexes	3	4	2	-	1	3	3	2	1	-	-	-
Male	2	2	1	-	1	1	3	1	-	-	-	-
Female	1	2	1	-	-	2	-	1	1	-	-	-
American Indian 1/, both sexes	1	1	4	1	2	3	1	1	-	-	-	-
Male	1	1	2	1	-	-	-	-	-	-	-	-
Female	-	-	2	-	2	3	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Oregon												
All races, both sexes	17	24	23	21	31	57	69	55	34	5	-	-
Male	13	14	13	10	19	32	29	17	14	3	-	-
Female	4	10	10	11	12	25	40	38	20	2	-	-
White, both sexes	15	20	21	18	27	52	66	52	32	5	-	-
Male	13	10	12	10	16	29	28	16	13	3	-	-
Female	2	10	9	8	11	23	38	36	19	2	-	-
Black, both sexes	1	2	-	-	-	-	-	-	-	-	-	-
Male	-	2	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	1	1	1	-	1	-	-	-
Male	-	1	-	-	1	1	-	-	1	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	2	2	3	4	2	3	1	-	-	-
Male	-	1	1	-	2	2	1	1	-	-	-	-
Female	1	-	1	2	1	2	1	2	1	-	-	-
Pennsylvania												
All races, both sexes	58	71	92	104	155	247	314	267	108	35	3	-
Male	27	42	45	65	78	136	135	99	28	9	-	-
Female	31	29	47	39	77	111	179	168	80	26	3	-
White, both sexes	36	51	71	91	138	226	295	249	106	33	3	-
Male	16	31	34	55	69	126	131	94	28	8	-	-
Female	20	20	37	36	69	100	164	155	78	25	3	-
Black, both sexes	20	19	16	12	13	21	16	16	2	2	-	-
Male	9	10	8	9	6	10	2	4	-	1	-	-
Female	11	9	8	3	7	11	14	12	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	5	1	3	-	3	2	-	-	-	-
Male	2	1	3	1	2	-	2	1	-	-	-	-
Female	-	-	2	-	1	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Rhode Island												
All races, both sexes	3	3	3	15	9	15	19	26	7	5	-	-
Male	2	-	3	10	5	10	8	8	1	3	-	-
Female	1	3	-	5	4	5	11	18	6	2	-	-
White, both sexes	2	3	3	15	8	15	18	25	7	5	-	-
Male	1	-	3	10	5	10	7	8	1	3	-	-
Female	1	3	-	5	3	5	11	17	6	2	-	-
Black, both sexes	1	-	-	-	1	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
South Carolina												
All races, both sexes	36	36	41	39	54	77	86	56	34	6	-	-
Male	24	28	26	25	31	30	30	20	10	1	-	-
Female	12	8	15	14	23	47	56	36	24	5	-	-
White, both sexes	5	12	21	24	38	57	66	46	28	6	-	-
Male	2	9	13	15	21	25	25	16	6	1	-	-
Female	3	3	8	9	17	32	41	30	22	5	-	-
Black, both sexes	31	24	19	14	16	16	20	10	6	-	-	-
Male	22	19	12	9	10	5	5	4	4	-	-	-
Female	9	5	7	5	6	11	15	6	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	4	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	4	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
South Dakota												
All races, both sexes	3	3	7	3	9	9	17	13	5	1	-	-
Male	1	2	5	2	5	4	8	8	1	1	-	-
Female	2	1	2	1	4	5	9	5	4	-	-	-
White, both sexes	3	3	7	3	9	8	17	13	5	1	-	-
Male	1	2	5	2	5	4	8	8	1	1	-	-
Female	2	1	2	1	4	4	9	5	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Tennessee												
All races, both sexes	38	53	47	54	70	110	118	81	35	12	1	-
Male	27	21	31	37	34	55	48	32	11	2	-	-
Female	11	32	16	17	36	55	70	49	24	10	1	-
White, both sexes	30	42	37	45	60	97	112	75	34	9	1	-
Male	20	19	24	33	27	50	43	31	11	2	-	-
Female	10	23	13	12	33	47	69	44	23	7	1	-
Black, both sexes	8	11	10	9	10	12	6	6	1	3	-	-
Male	7	2	7	4	7	4	5	1	-	-	-	-
Female	1	9	3	5	3	8	1	5	1	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Texas												
All races, both sexes	137	119	125	139	166	249	306	224	103	21	7	-
Male	85	72	69	73	83	108	138	70	35	5	1	-
Female	52	47	56	66	83	141	168	154	68	16	6	-
White, both sexes	96	94	101	117	146	229	282	211	95	19	5	-
Male	64	57	57	61	73	99	125	65	34	4	-	-
Female	32	37	44	56	73	130	157	146	61	15	5	-
Black, both sexes	36	25	21	19	18	16	21	7	7	2	2	-
Male	18	15	12	12	8	8	12	2	1	1	1	-
Female	18	10	9	7	10	8	9	5	6	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	3	3	2	4	3	5	1	-	-	-
Male	3	-	-	-	2	1	1	2	-	-	-	-
Female	2	-	3	3	-	3	2	3	1	-	-	-
Utah												
All races, both sexes	4	3	11	12	15	23	23	21	7	-	-	-
Male	2	2	7	6	8	8	9	13	2	-	-	-
Female	2	1	4	6	7	15	14	8	5	-	-	-
White, both sexes	4	1	11	11	15	22	23	19	6	-	-	-
Male	2	-	7	5	8	8	9	11	1	-	-	-
Female	2	1	4	6	7	14	14	8	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	1	-	1	-	2	-	-	-	-
Male	-	2	-	1	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Vermont												
All races, both sexes	1	1	4	3	10	3	4	9	5	1	-	-
Male	1	-	1	2	5	2	-	3	2	-	-	-
Female	-	1	3	1	5	1	4	6	3	1	-	-
White, both sexes	1	1	4	3	10	3	4	9	5	1	-	-
Male	1	-	1	2	5	2	-	3	2	-	-	-
Female	-	1	3	1	5	1	4	6	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	43	43	48	67	75	117	123	93	44	13	1	-
Male	21	28	25	39	38	58	44	33	10	3	-	-
Female	22	15	23	28	37	59	79	60	34	10	1	-
White, both sexes	18	30	31	48	52	93	102	83	36	12	1	-
Male	8	18	17	29	25	48	36	28	8	3	-	-
Female	10	12	14	19	27	45	66	55	28	9	1	-
Black, both sexes	23	13	13	18	18	20	18	9	7	1	-	-
Male	12	10	8	10	10	9	6	5	2	-	-	-
Female	11	3	5	8	8	11	12	4	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	4	1	5	4	3	1	1	-	-	-
Male	1	-	-	-	3	1	2	-	-	-	-	-
Female	1	-	4	1	2	3	1	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Washington												
All races, both sexes	31	23	31	41	62	101	100	90	33	9	2	-
Male	20	14	16	19	41	48	36	32	7	2	2	-
Female	11	9	15	22	21	53	64	58	26	7	-	-
White, both sexes	23	20	23	34	59	89	92	87	30	8	2	-
Male	15	12	13	18	38	42	33	31	7	2	2	-
Female	8	8	10	16	21	47	59	56	23	6	-	-
Black, both sexes	2	1	1	1	-	2	1	1	-	-	-	-
Male	2	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	2	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	2	7	4	3	8	6	2	3	1	-	-
Male	3	1	2	1	3	4	2	1	-	-	-	-
Female	3	1	5	3	-	4	4	1	3	1	-	-
West Virginia												
All races, both sexes	12	8	16	12	26	27	46	25	19	11	-	-
Male	9	5	9	10	13	11	18	7	3	6	-	-
Female	3	3	7	2	13	16	28	18	16	5	-	-
White, both sexes	11	6	15	12	26	27	46	24	19	11	-	-
Male	8	4	8	10	13	11	18	7	3	6	-	-
Female	3	2	7	2	13	16	28	17	16	5	-	-
Black, both sexes	1	2	1	-	-	-	-	1	-	-	-	-
Male	1	1	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intracerebral and other intracranial hemorrhage (I61-I62)												
Wisconsin												
All races, both sexes	17	39	29	40	44	97	83	85	31	11	1	-
Male	5	23	14	24	21	43	33	36	6	3	-	-
Female	12	16	15	16	23	54	50	49	25	8	1	-
White, both sexes	9	32	23	35	42	96	78	83	31	11	1	-
Male	2	18	10	19	20	42	30	36	6	3	-	-
Female	7	14	13	16	22	54	48	47	25	8	1	-
Black, both sexes	8	7	5	5	-	-	2	-	-	-	-	-
Male	3	5	4	5	-	-	1	-	-	-	-	-
Female	5	2	1	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	2	-	-	-	-
Wyoming												
All races, both sexes	1	-	1	3	5	7	7	4	3	1	-	-
Male	-	-	-	2	3	5	5	1	-	-	-	-
Female	1	-	1	1	2	2	2	3	3	1	-	-
White, both sexes	-	-	1	3	5	6	7	4	3	1	-	-
Male	-	-	-	2	3	4	5	1	-	-	-	-
Female	-	-	1	1	2	2	2	3	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
United States												
All races, both sexes	7,563	14	5	1	2	10	4	20	29	48	90	132
Male	3,087	4	2	1	1	4	2	9	15	30	49	86
Female	4,476	10	3	-	1	6	2	11	14	18	41	46
White, both sexes	6,468	7	4	1	2	8	4	14	23	34	66	83
Male	2,606	2	2	1	1	2	2	6	11	24	37	57
Female	3,862	5	2	-	1	6	2	8	12	10	29	26
Black, both sexes	894	4	1	-	-	-	-	5	5	13	23	46
Male	393	1	-	-	-	-	-	2	4	6	11	27
Female	501	3	1	-	-	-	-	3	1	7	12	19
American Indian 1/, both sexes	42	-	-	-	-	-	-	-	1	-	1	2
Male	20	-	-	-	-	-	-	-	-	-	1	2
Female	22	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	159	3	-	-	-	2	-	1	-	1	-	1
Male	68	1	-	-	-	2	-	1	-	-	-	-
Female	91	2	-	-	-	-	-	-	-	1	-	1
Alabama												
All races, both sexes	129	-	-	-	-	-	-	2	1	-	2	1
Male	51	-	-	-	-	-	-	1	1	-	1	1
Female	78	-	-	-	-	-	-	1	-	-	1	-
White, both sexes	96	-	-	-	-	-	-	2	-	-	1	-
Male	38	-	-	-	-	-	-	1	-	-	-	-
Female	58	-	-	-	-	-	-	1	-	-	1	-
Black, both sexes	33	-	-	-	-	-	-	-	1	-	1	1
Male	13	-	-	-	-	-	-	-	1	-	1	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Alaska												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	136	1	-	-	-	-	-	-	2	1	-	2
Male	65	1	-	-	-	-	-	-	1	-	-	-
Female	71	-	-	-	-	-	-	-	1	1	-	2
White, both sexes	121	1	-	-	-	-	-	-	1	1	-	-
Male	61	1	-	-	-	-	-	-	-	-	-	-
Female	60	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	11	-	-	-	-	-	-	-	1	-	-	1
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Arkansas												
All races, both sexes	77	-	-	-	1	-	-	-	-	-	2	2
Male	33	-	-	-	-	-	-	-	-	-	1	2
Female	44	-	-	-	1	-	-	-	-	-	1	-
White, both sexes	63	-	-	-	1	-	-	-	-	-	1	1
Male	28	-	-	-	-	-	-	-	-	-	1	1
Female	35	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	644	1	-	-	-	1	-	1	3	4	3	15
Male	267	1	-	-	-	-	-	1	1	2	1	12
Female	377	-	-	-	-	1	-	-	2	2	2	3
White, both sexes	541	1	-	-	-	1	-	1	1	2	3	14
Male	224	1	-	-	-	-	-	1	-	2	1	11
Female	317	-	-	-	-	1	-	-	1	-	2	3
Black, both sexes	42	-	-	-	-	-	-	-	1	1	-	1
Male	15	-	-	-	-	-	-	-	1	-	-	1
Female	27	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	59	-	-	-	-	-	-	-	-	1	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Colorado												
All races, both sexes	50	-	1	-	-	-	-	1	1	-	1	1
Male	20	-	1	-	-	-	-	-	-	-	-	1
Female	30	-	-	-	-	-	-	1	1	-	1	-
White, both sexes	48	-	1	-	-	-	-	1	1	-	1	1
Male	19	-	1	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	-	-	1	1	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	104	-	-	-	-	-	-	1	-	-	1	1
Male	48	-	-	-	-	-	-	1	-	-	-	1
Female	56	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	94	-	-	-	-	-	-	-	-	-	1	1
Male	43	-	-	-	-	-	-	-	-	-	-	1
Female	51	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Delaware												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Florida												
All races, both sexes	529	-	-	-	-	2	-	-	2	6	7	13
Male	257	-	-	-	-	1	-	-	1	5	3	8
Female	272	-	-	-	-	1	-	-	1	1	4	5
White, both sexes	448	-	-	-	-	1	-	-	2	5	5	8
Male	219	-	-	-	-	-	-	-	1	5	2	6
Female	229	-	-	-	-	1	-	-	1	-	3	2
Black, both sexes	75	-	-	-	-	-	-	-	-	1	2	5
Male	35	-	-	-	-	-	-	-	-	-	1	2
Female	40	-	-	-	-	-	-	-	-	1	1	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	148	-	-	1	-	-	-	2	-	2	2	7
Male	62	-	-	1	-	-	-	1	-	2	2	5
Female	86	-	-	-	-	-	-	1	-	-	-	2
White, both sexes	106	-	-	1	-	-	-	1	-	2	1	4
Male	40	-	-	1	-	-	-	-	-	2	1	4
Female	66	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	39	-	-	-	-	-	-	1	-	-	1	3
Male	21	-	-	-	-	-	-	1	-	-	1	1
Female	18	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Hawaii												
All races, both sexes	39	1	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	27	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	34	1	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	25	1	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Illinois												
All races, both sexes	351	1	-	-	-	-	-	-	-	1	5	4
Male	133	1	-	-	-	-	-	-	-	1	2	2
Female	218	-	-	-	-	-	-	-	-	-	3	2
White, both sexes	290	-	-	-	-	-	-	-	-	-	2	2
Male	108	-	-	-	-	-	-	-	-	-	1	1
Female	182	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	57	1	-	-	-	-	-	-	-	1	3	2
Male	23	1	-	-	-	-	-	-	-	1	1	1
Female	34	-	-	-	-	-	-	-	-	-	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	152	-	1	-	-	-	-	-	-	1	3	4
Male	58	-	1	-	-	-	-	-	-	1	1	3
Female	94	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	134	-	1	-	-	-	-	-	-	1	2	3
Male	51	-	1	-	-	-	-	-	-	1	1	2
Female	83	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	18	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Iowa												
All races, both sexes	122	-	-	-	-	-	-	-	1	-	3	1
Male	49	-	-	-	-	-	-	-	1	-	1	1
Female	73	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	121	-	-	-	-	-	-	-	1	-	3	1
Male	49	-	-	-	-	-	-	-	1	-	1	1
Female	72	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	68	-	-	-	-	-	-	-	1	-	-	2
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	43	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	64	-	-	-	-	-	-	-	1	-	-	2
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	41	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Kentucky												
All races, both sexes	102	-	-	-	-	-	-	1	-	2	4	1
Male	41	-	-	-	-	-	-	-	-	2	-	1
Female	61	-	-	-	-	-	-	1	-	-	4	-
White, both sexes	96	-	-	-	-	-	-	1	-	2	3	1
Male	39	-	-	-	-	-	-	-	-	2	-	1
Female	57	-	-	-	-	-	-	1	-	-	3	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	131	-	1	-	-	-	-	-	2	2	1	5
Male	50	-	-	-	-	-	-	-	1	1	-	4
Female	81	-	1	-	-	-	-	-	1	1	1	1
White, both sexes	86	-	1	-	-	-	-	-	2	2	-	1
Male	29	-	-	-	-	-	-	-	1	1	-	1
Female	57	-	1	-	-	-	-	-	1	1	-	-
Black, both sexes	45	-	-	-	-	-	-	-	-	-	1	4
Male	21	-	-	-	-	-	-	-	-	-	-	3
Female	24	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Maine												
All races, both sexes	27	-	-	-	-	-	-	-	1	-	-	1
Male	13	-	-	-	-	-	-	-	1	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	27	-	-	-	-	-	-	-	1	-	-	1
Male	13	-	-	-	-	-	-	-	1	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	217	3	-	-	-	-	-	1	-	1	2	1
Male	84	-	-	-	-	-	-	1	-	-	2	-
Female	133	3	-	-	-	-	-	-	-	1	-	1
White, both sexes	167	2	-	-	-	-	-	-	-	-	2	-
Male	63	-	-	-	-	-	-	-	-	-	2	-
Female	104	2	-	-	-	-	-	-	-	-	-	-
Black, both sexes	44	-	-	-	-	-	-	1	-	1	-	1
Male	21	-	-	-	-	-	-	1	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	6	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Massachusetts												
All races, both sexes	143	-	-	-	-	-	-	-	-	-	1	2
Male	53	-	-	-	-	-	-	-	-	-	1	1
Female	90	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	138	-	-	-	-	-	-	-	-	-	-	2
Male	50	-	-	-	-	-	-	-	-	-	-	1
Female	88	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	361	1	-	-	-	-	-	-	1	5	5	9
Male	139	-	-	-	-	-	-	-	1	1	3	4
Female	222	1	-	-	-	-	-	-	-	4	2	5
White, both sexes	315	-	-	-	-	-	-	-	1	2	5	5
Male	118	-	-	-	-	-	-	-	1	1	3	1
Female	197	-	-	-	-	-	-	-	-	1	2	4
Black, both sexes	43	1	-	-	-	-	-	-	-	3	-	4
Male	19	-	-	-	-	-	-	-	-	-	-	3
Female	24	1	-	-	-	-	-	-	-	3	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Minnesota												
All races, both sexes	137	-	-	-	-	-	1	-	1	1	2	3
Male	54	-	-	-	-	-	-	-	-	1	-	2
Female	83	-	-	-	-	-	1	-	1	-	2	1
White, both sexes	134	-	-	-	-	-	1	-	1	1	2	3
Male	52	-	-	-	-	-	-	-	-	1	-	2
Female	82	-	-	-	-	-	1	-	1	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	78	-	-	-	-	-	-	-	-	-	2	4
Male	36	-	-	-	-	-	-	-	-	-	2	2
Female	42	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	51	-	-	-	-	-	-	-	-	-	1	1
Male	23	-	-	-	-	-	-	-	-	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	26	-	-	-	-	-	-	-	-	-	1	3
Male	13	-	-	-	-	-	-	-	-	-	1	2
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Missouri												
All races, both sexes	197	-	-	-	-	-	-	-	-	-	1	4
Male	81	-	-	-	-	-	-	-	-	-	1	1
Female	116	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	170	-	-	-	-	-	-	-	-	-	1	1
Male	74	-	-	-	-	-	-	-	-	-	1	1
Female	96	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	26	-	-	-	-	-	-	-	-	-	-	3
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	16	-	-	-	-	1	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	1	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Nebraska												
All races, both sexes	37	-	-	-	-	-	-	1	-	-	-	-
Male	18	-	-	-	-	-	-	1	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	-	-	-	-	-	-	1	-	-	-	-
Male	17	-	-	-	-	-	-	1	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	33	-	-	-	-	-	-	-	1	-	-	1
Male	16	-	-	-	-	-	-	-	1	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	1	-	-	1
Male	14	-	-	-	-	-	-	-	1	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
New Hampshire												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	19	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	165	-	-	-	-	-	-	-	-	-	1	1
Male	61	-	-	-	-	-	-	-	-	-	1	1
Female	104	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	141	-	-	-	-	-	-	-	-	-	-	1
Male	52	-	-	-	-	-	-	-	-	-	-	1
Female	89	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
New Mexico												
All races, both sexes	32	1	-	-	-	-	-	-	1	1	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	20	1	-	-	-	-	-	-	1	1	-	-
White, both sexes	30	1	-	-	-	-	-	-	1	1	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	20	1	-	-	-	-	-	-	1	1	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	361	-	-	-	-	-	1	-	-	1	3	6
Male	152	-	-	-	-	-	1	-	-	1	2	4
Female	209	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	321	-	-	-	-	-	1	-	-	-	3	3
Male	133	-	-	-	-	-	1	-	-	-	2	1
Female	188	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	30	-	-	-	-	-	-	-	-	1	-	3
Male	17	-	-	-	-	-	-	-	-	1	-	3
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
North Carolina												
All races, both sexes	257	1	-	-	-	-	1	1	1	3	3	6
Male	116	-	-	-	-	-	1	1	-	2	2	4
Female	141	1	-	-	-	-	-	-	1	1	1	2
White, both sexes	202	-	-	-	-	-	1	1	1	2	2	4
Male	90	-	-	-	-	-	1	1	-	1	2	2
Female	112	-	-	-	-	-	-	-	1	1	-	2
Black, both sexes	52	1	-	-	-	-	-	-	-	1	1	2
Male	24	-	-	-	-	-	-	-	-	1	-	2
Female	28	1	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Ohio												
All races, both sexes	392	-	-	-	-	1	-	2	1	-	5	8
Male	143	-	-	-	-	1	-	-	1	-	4	6
Female	249	-	-	-	-	-	-	2	-	-	1	2
White, both sexes	358	-	-	-	-	1	-	-	1	-	4	4
Male	125	-	-	-	-	1	-	-	1	-	3	4
Female	233	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	31	-	-	-	-	-	-	2	-	-	1	4
Male	16	-	-	-	-	-	-	-	-	-	1	2
Female	15	-	-	-	-	-	-	2	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	106	-	-	-	-	-	-	-	-	4	1	-
Male	43	-	-	-	-	-	-	-	-	2	-	-
Female	63	-	-	-	-	-	-	-	-	2	1	-
White, both sexes	90	-	-	-	-	-	-	-	-	3	1	-
Male	34	-	-	-	-	-	-	-	-	2	-	-
Female	56	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	9	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Oregon												
All races, both sexes	106	-	-	-	-	-	-	-	1	-	3	-
Male	36	-	-	-	-	-	-	-	1	-	-	-
Female	70	-	-	-	-	-	-	-	-	-	3	-
White, both sexes	102	-	-	-	-	-	-	-	1	-	2	-
Male	33	-	-	-	-	-	-	-	1	-	-	-
Female	69	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	511	-	1	-	-	2	-	2	2	2	4	5
Male	182	-	-	-	-	-	-	1	-	1	2	4
Female	329	-	1	-	-	2	-	1	2	1	2	1
White, both sexes	461	-	-	-	-	2	-	2	2	1	3	4
Male	161	-	-	-	-	-	-	1	-	-	1	3
Female	300	-	-	-	-	2	-	1	2	1	2	1
Black, both sexes	45	-	1	-	-	-	-	-	-	1	1	1
Male	18	-	-	-	-	-	-	-	-	1	1	1
Female	27	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Rhode Island												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	163	-	-	-	-	-	-	-	-	1	2	2
Male	54	-	-	-	-	-	-	-	-	1	1	1
Female	109	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	121	-	-	-	-	-	-	-	-	-	1	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	85	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	42	-	-	-	-	-	-	-	-	1	1	2
Male	18	-	-	-	-	-	-	-	-	1	1	1
Female	24	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
South Dakota												
All races, both sexes	28	-	-	-	-	-	-	-	-	-	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	-	-	-	1	1
Male	12	-	-	-	-	-	-	-	-	-	1	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	207	1	-	-	-	-	1	-	-	-	6	1
Male	88	-	-	-	-	-	-	-	-	-	3	1
Female	119	1	-	-	-	-	1	-	-	-	3	-
White, both sexes	176	-	-	-	-	-	1	-	-	-	5	-
Male	74	-	-	-	-	-	-	-	-	-	3	-
Female	102	-	-	-	-	-	1	-	-	-	2	-
Black, both sexes	31	1	-	-	-	-	-	-	-	-	1	1
Male	14	-	-	-	-	-	-	-	-	-	-	1
Female	17	1	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Texas												
All races, both sexes	456	3	1	-	1	1	-	3	5	4	5	10
Male	199	1	-	-	1	-	-	1	3	4	5	6
Female	257	2	1	-	-	1	-	2	2	-	-	4
White, both sexes	388	2	1	-	1	1	-	2	3	3	3	6
Male	163	-	-	-	1	-	-	1	2	3	3	4
Female	225	2	1	-	-	1	-	1	1	-	-	2
Black, both sexes	60	-	-	-	-	-	-	1	2	1	2	3
Male	29	-	-	-	-	-	-	-	1	1	2	1
Female	31	-	-	-	-	-	-	1	1	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	1	-	-	-	-	-	-	-	-	-	-
Male	6	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	25	-	-	-	-	-	-	1	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	25	-	-	-	-	-	-	1	-	1	-	-
Male	11	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Vermont												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	204	-	-	-	-	-	-	-	-	-	3	-
Male	79	-	-	-	-	-	-	-	-	-	1	-
Female	125	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	152	-	-	-	-	-	-	-	-	-	1	-
Male	59	-	-	-	-	-	-	-	-	-	1	-
Female	93	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	49	-	-	-	-	-	-	-	-	-	2	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Washington												
All races, both sexes	181	-	-	-	-	1	-	1	1	4	-	3
Male	82	-	-	-	-	1	-	-	1	2	-	3
Female	99	-	-	-	-	-	-	1	-	2	-	-
White, both sexes	170	-	-	-	-	-	-	1	1	4	-	2
Male	76	-	-	-	-	-	-	-	1	2	-	2
Female	94	-	-	-	-	-	-	1	-	2	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	65	-	-	-	-	-	-	-	-	-	2	2
Male	21	-	-	-	-	-	-	-	-	-	2	-
Female	44	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	63	-	-	-	-	-	-	-	-	-	2	2
Male	21	-	-	-	-	-	-	-	-	-	2	-
Female	42	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Cerebral infarction (I63)												
Wisconsin												
All races, both sexes	138	-	-	-	-	1	-	-	-	1	2	3
Male	64	-	-	-	-	1	-	-	-	-	2	2
Female	74	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	126	-	-	-	-	1	-	-	-	1	2	3
Male	58	-	-	-	-	1	-	-	-	-	2	2
Female	68	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
United States												
All races, both sexes	198	270	358	455	635	966	1,429	1,441	1,054	342	60	-
Male	122	167	222	255	308	400	578	478	286	64	4	-
Female	76	103	136	200	327	566	851	963	768	278	56	-
White, both sexes	144	194	267	351	532	808	1,294	1,312	951	315	54	-
Male	89	123	163	201	256	350	536	426	255	58	4	-
Female	55	71	104	150	276	458	758	886	696	257	50	-
Black, both sexes	46	63	79	82	82	129	107	101	82	22	4	-
Male	28	36	51	45	41	41	32	40	23	5	-	-
Female	18	27	28	37	41	88	75	61	59	17	4	-
American Indian 1/, both sexes	3	7	2	7	7	3	2	3	4	-	-	-
Male	2	4	2	3	2	-	2	2	-	-	-	-
Female	1	3	-	4	5	3	-	1	4	-	-	-
Asian or Pacific Islander 2/, both sexes	5	6	10	15	14	26	26	25	17	5	2	-
Male	3	4	6	6	9	9	8	10	8	1	-	-
Female	2	2	4	9	5	17	18	15	9	4	2	-
Alabama												
All races, both sexes	4	12	6	13	12	16	20	21	13	6	-	-
Male	3	6	4	6	7	5	5	10	1	-	-	-
Female	1	6	2	7	5	11	15	11	12	6	-	-
White, both sexes	4	5	4	7	10	15	16	18	10	4	-	-
Male	3	4	3	4	5	5	5	8	-	-	-	-
Female	1	1	1	3	5	10	11	10	10	4	-	-
Black, both sexes	-	7	2	6	2	1	4	3	3	2	-	-
Male	-	2	1	2	2	-	-	2	1	-	-	-
Female	-	5	1	4	-	1	4	1	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Alaska												
All races, both sexes	1	-	-	1	-	1	3	1	-	-	-	-
Male	1	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	3	1	-	-	-	-
White, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	1	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	5	7	8	10	21	21	19	18	16	4	1	-
Male	4	5	8	6	9	9	7	7	7	-	1	-
Female	1	2	-	4	12	12	12	11	9	4	-	-
White, both sexes	4	5	8	9	19	18	17	17	16	4	1	-
Male	3	4	8	6	9	8	7	7	7	-	1	-
Female	1	1	-	3	10	10	10	10	9	4	-	-
Black, both sexes	1	2	-	1	1	1	2	1	-	-	-	-
Male	1	1	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	1	1	-	2	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Arkansas												
All races, both sexes	4	2	2	4	4	16	11	9	14	3	3	-
Male	2	-	1	4	3	6	6	2	5	-	1	-
Female	2	2	1	-	1	10	5	7	9	3	2	-
White, both sexes	4	1	1	3	3	12	10	8	13	3	2	-
Male	2	-	1	3	2	5	5	2	5	-	1	-
Female	2	1	-	-	1	7	5	6	8	3	1	-
Black, both sexes	-	1	1	1	1	4	1	1	1	-	1	-
Male	-	-	-	1	1	1	1	-	-	-	-	-
Female	-	1	1	-	-	3	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	19	33	31	36	46	83	125	131	81	29	2	-
Male	10	22	15	18	25	28	49	52	22	8	-	-
Female	9	11	16	18	21	55	76	79	59	21	2	-
White, both sexes	16	28	23	28	37	68	107	114	67	28	2	-
Male	9	19	12	16	18	23	41	46	16	8	-	-
Female	7	9	11	12	19	45	66	68	51	20	2	-
Black, both sexes	1	3	2	4	4	6	7	4	8	-	-	-
Male	-	1	1	-	2	1	5	-	3	-	-	-
Female	1	2	1	4	2	5	2	4	5	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	2	2	6	4	5	9	11	13	5	1	-	-
Male	1	2	2	2	5	4	3	6	3	-	-	-
Female	1	-	4	2	-	5	8	7	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Colorado												
All races, both sexes	2	3	2	2	2	6	10	8	9	1	-	-
Male	1	1	2	2	1	3	6	1	1	-	-	-
Female	1	2	-	-	1	3	4	7	8	1	-	-
White, both sexes	2	2	2	2	2	5	10	8	9	1	-	-
Male	1	1	2	2	1	2	6	1	1	-	-	-
Female	1	1	-	-	1	3	4	7	8	1	-	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	4	5	4	6	14	20	21	22	4	1	-
Male	-	4	4	2	2	7	11	6	8	2	-	-
Female	-	-	1	2	4	7	9	15	14	2	1	-
White, both sexes	-	4	3	4	4	13	20	18	22	3	1	-
Male	-	4	2	2	1	7	11	5	8	2	-	-
Female	-	-	1	2	3	6	9	13	14	1	1	-
Black, both sexes	-	-	1	-	2	1	-	3	-	1	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Delaware												
All races, both sexes	-	1	2	-	2	4	-	3	-	1	-	-
Male	-	1	-	-	1	2	-	1	-	1	-	-
Female	-	-	2	-	1	2	-	2	-	-	-	-
White, both sexes	-	-	1	-	1	4	-	3	-	1	-	-
Male	-	-	-	-	1	2	-	1	-	1	-	-
Female	-	-	1	-	-	2	-	2	-	-	-	-
Black, both sexes	-	1	1	-	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	-	-	1	3	2	1	1	-	-	-
Male	-	-	-	-	1	1	1	1	-	-	-	-
Female	-	-	-	-	-	2	1	-	1	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	1	3	1	1	1	-	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	2	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Florida												
All races, both sexes	16	8	30	34	67	69	86	89	74	22	4	-
Male	11	6	22	20	41	37	35	37	27	3	-	-
Female	5	2	8	14	26	32	51	52	47	19	4	-
White, both sexes	12	7	22	25	57	59	80	74	69	19	3	-
Male	8	5	15	15	34	36	34	28	27	3	-	-
Female	4	2	7	10	23	23	46	46	42	16	3	-
Black, both sexes	4	1	8	8	8	9	6	14	5	3	1	-
Male	3	1	7	5	6	1	1	8	-	-	-	-
Female	1	-	1	3	2	8	5	6	5	3	1	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Georgia												
All races, both sexes	8	9	7	11	11	18	30	23	12	2	3	-
Male	5	8	3	7	6	2	9	9	2	-	-	-
Female	3	1	4	4	5	16	21	14	10	2	3	-
White, both sexes	4	3	5	9	9	15	22	19	7	1	3	-
Male	3	2	2	5	5	1	7	7	-	-	-	-
Female	1	1	3	4	4	14	15	12	7	1	3	-
Black, both sexes	4	5	2	2	2	3	6	4	5	1	-	-
Male	2	5	1	2	1	1	2	2	2	-	-	-
Female	2	-	1	-	1	2	4	2	3	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Hawaii												
All races, both sexes	-	1	-	3	4	5	4	8	11	1	1	-
Male	-	1	-	-	2	1	1	2	5	-	-	-
Female	-	-	-	3	2	4	3	6	6	1	1	-
White, both sexes	-	1	-	-	-	-	1	2	1	-	-	-
Male	-	1	-	-	-	-	-	1	1	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	3	4	5	3	6	10	1	1	-
Male	-	-	-	-	2	1	1	1	4	-	-	-
Female	-	-	-	3	2	4	2	5	6	1	1	-
Idaho												
All races, both sexes	-	1	3	3	4	3	4	6	7	4	-	-
Male	-	1	1	2	2	2	3	2	1	1	-	-
Female	-	-	2	1	2	1	1	4	6	3	-	-
White, both sexes	-	1	3	3	4	3	4	6	7	4	-	-
Male	-	1	1	2	2	2	3	2	1	1	-	-
Female	-	-	2	1	2	1	1	4	6	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Illinois												
All races, both sexes	9	9	17	17	26	49	74	71	49	17	2	-
Male	5	5	9	11	10	19	28	25	13	1	1	-
Female	4	4	8	6	16	30	46	46	36	16	1	-
White, both sexes	4	5	13	13	21	43	65	63	42	16	1	-
Male	3	2	6	8	9	17	27	22	10	1	1	-
Female	1	3	7	5	12	26	38	41	32	15	-	-
Black, both sexes	5	4	4	4	5	6	7	6	7	1	1	-
Male	2	3	3	3	1	2	-	2	3	-	-	-
Female	3	1	1	1	4	4	7	4	4	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Indiana												
All races, both sexes	6	8	5	10	17	23	25	30	12	5	2	-
Male	3	5	2	7	6	8	6	9	5	1	-	-
Female	3	3	3	3	11	15	19	21	7	4	2	-
White, both sexes	4	7	4	8	16	21	23	25	12	5	2	-
Male	3	4	1	5	6	8	6	7	5	1	-	-
Female	1	3	3	3	10	13	17	18	7	4	2	-
Black, both sexes	2	1	1	2	1	2	2	5	-	-	-	-
Male	-	1	1	2	-	-	-	2	-	-	-	-
Female	2	-	-	-	1	2	2	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Iowa												
All races, both sexes	3	4	4	3	10	14	18	21	26	12	2	-
Male	2	3	3	1	5	6	8	5	10	3	-	-
Female	1	1	1	2	5	8	10	16	16	9	2	-
White, both sexes	3	4	4	2	10	14	18	21	26	12	2	-
Male	2	3	3	1	5	6	8	5	10	3	-	-
Female	1	1	1	1	5	8	10	16	16	9	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	4	3	3	8	6	15	16	5	4	-	-
Male	1	1	2	2	2	-	6	8	2	-	-	-
Female	-	3	1	1	6	6	9	8	3	4	-	-
White, both sexes	1	4	3	3	8	5	15	14	4	4	-	-
Male	1	1	2	2	2	-	6	6	2	-	-	-
Female	-	3	1	1	6	5	9	8	2	4	-	-
Black, both sexes	-	-	-	-	-	1	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	2	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Kentucky												
All races, both sexes	3	1	8	5	12	10	22	18	12	2	1	-
Male	2	1	8	3	5	5	6	6	2	-	-	-
Female	1	-	-	2	7	5	16	12	10	2	1	-
White, both sexes	3	1	6	5	12	9	21	17	12	2	1	-
Male	2	1	6	3	5	5	6	6	2	-	-	-
Female	1	-	-	2	7	4	15	11	10	2	1	-
Black, both sexes	-	-	2	-	-	1	1	1	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	6	11	7	13	17	18	23	18	3	-	-
Male	2	2	7	5	9	6	2	8	2	1	-	-
Female	2	4	4	2	4	11	16	15	16	2	-	-
White, both sexes	3	2	4	4	7	10	14	18	15	3	-	-
Male	1	-	2	4	4	6	-	6	2	1	-	-
Female	2	2	2	-	3	4	14	12	13	2	-	-
Black, both sexes	1	4	7	3	6	7	4	5	3	-	-	-
Male	1	2	5	1	5	-	2	2	-	-	-	-
Female	-	2	2	2	1	7	2	3	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Maine												
All races, both sexes	1	1	2	-	2	2	6	5	4	2	-	-
Male	-	1	1	-	2	2	1	3	2	-	-	-
Female	1	-	1	-	-	-	5	2	2	2	-	-
White, both sexes	1	1	2	-	2	2	6	5	4	2	-	-
Male	-	1	1	-	2	2	1	3	2	-	-	-
Female	1	-	1	-	-	-	5	2	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	4	13	12	7	32	39	42	41	12	4	-
Male	2	4	8	5	5	15	17	13	9	3	-	-
Female	1	-	5	7	2	17	22	29	32	9	4	-
White, both sexes	2	3	4	6	6	23	33	36	35	12	3	-
Male	2	3	2	2	4	11	17	10	7	3	-	-
Female	-	-	2	4	2	12	16	26	28	9	3	-
Black, both sexes	-	1	9	6	1	9	3	6	6	-	-	-
Male	-	1	6	3	1	4	-	3	2	-	-	-
Female	-	-	3	3	-	5	3	3	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	3	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	3	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Massachusetts												
All races, both sexes	1	2	3	6	7	15	37	24	32	11	2	-
Male	-	2	2	2	3	9	16	4	9	3	1	-
Female	1	-	1	4	4	6	21	20	23	8	1	-
White, both sexes	1	1	2	6	7	15	36	23	32	11	2	-
Male	-	1	1	2	3	9	16	4	9	3	1	-
Female	1	-	1	4	4	6	20	19	23	8	1	-
Black, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Michigan												
All races, both sexes	5	10	26	17	28	45	62	77	53	15	2	-
Male	4	5	18	9	13	16	27	24	13	1	-	-
Female	1	5	8	8	15	29	35	53	40	14	2	-
White, both sexes	3	5	21	12	25	38	57	75	50	14	2	-
Male	2	3	14	6	10	13	26	24	13	1	-	-
Female	1	2	7	6	15	25	31	51	37	13	2	-
Black, both sexes	2	4	4	5	3	6	5	2	3	1	-	-
Male	2	2	3	3	3	2	1	-	-	-	-	-
Female	-	2	1	2	-	4	4	2	3	1	-	-
American Indian 1/, both sexes	-	1	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Minnesota												
All races, both sexes	3	1	4	9	11	19	25	24	22	7	4	-
Male	2	1	3	7	4	11	10	7	5	1	-	-
Female	1	-	1	2	7	8	15	17	17	6	4	-
White, both sexes	2	1	3	9	11	19	25	24	21	7	4	-
Male	1	1	2	7	4	11	10	7	5	1	-	-
Female	1	-	1	2	7	8	15	17	16	6	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	6	5	4	6	5	22	8	9	3	-	-
Male	2	2	3	3	2	3	12	2	2	1	-	-
Female	2	4	2	1	4	2	10	6	7	2	-	-
White, both sexes	4	3	2	2	5	4	15	8	5	1	-	-
Male	2	2	1	1	2	2	9	2	1	-	-	-
Female	2	1	1	1	3	2	6	6	4	1	-	-
Black, both sexes	-	2	3	2	1	1	7	-	4	2	-	-
Male	-	-	2	2	-	1	3	-	1	1	-	-
Female	-	2	1	-	1	-	4	-	3	1	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Missouri												
All races, both sexes	8	9	14	16	16	25	41	32	25	4	2	-
Male	5	6	6	9	8	11	19	9	4	2	-	-
Female	3	3	8	7	8	14	22	23	21	2	2	-
White, both sexes	6	7	11	14	14	24	35	28	24	3	2	-
Male	3	5	5	9	8	10	18	9	3	2	-	-
Female	3	2	6	5	6	14	17	19	21	1	2	-
Black, both sexes	2	2	3	2	2	-	6	4	1	1	-	-
Male	2	1	1	-	-	-	1	-	1	-	-	-
Female	-	1	2	2	2	-	5	4	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	1	-	1	2	1	4	4	2	-	-	-
Male	-	1	-	1	-	-	1	3	-	-	-	-
Female	-	-	-	-	2	1	3	1	2	-	-	-
White, both sexes	-	1	-	1	2	1	4	4	2	-	-	-
Male	-	1	-	1	-	-	1	3	-	-	-	-
Female	-	-	-	-	2	1	3	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Nebraska												
All races, both sexes	1	1	4	1	4	7	6	8	3	1	-	-
Male	-	-	3	1	2	2	4	2	2	1	-	-
Female	1	1	1	-	2	5	2	6	1	-	-	-
White, both sexes	-	1	4	1	3	5	6	7	3	1	-	-
Male	-	-	3	1	2	1	4	2	2	1	-	-
Female	-	1	1	-	1	4	2	5	1	-	-	-
Black, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	-	1	4	2	4	10	4	4	2	-	-	-
Male	-	1	2	2	2	5	2	-	-	-	-	-
Female	-	-	2	-	2	5	2	4	2	-	-	-
White, both sexes	-	1	3	2	4	9	3	4	2	-	-	-
Male	-	1	1	2	2	5	1	-	-	-	-	-
Female	-	-	2	-	2	4	2	4	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	1	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
New Hampshire												
All races, both sexes	1	-	2	1	-	1	5	6	2	-	-	-
Male	1	-	2	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	5	5	2	-	-	-
White, both sexes	1	-	2	1	-	1	5	6	2	-	-	-
Male	1	-	2	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	5	5	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	2	3	10	9	21	37	43	24	10	1	-
Male	3	-	2	8	5	6	13	14	7	1	-	-
Female	-	2	1	2	4	15	24	29	17	9	1	-
White, both sexes	1	1	3	4	8	15	35	40	23	9	1	-
Male	1	-	2	4	5	5	13	13	7	1	-	-
Female	-	1	1	-	3	10	22	27	16	8	1	-
Black, both sexes	2	1	-	5	1	4	2	3	1	-	-	-
Male	2	-	-	3	-	-	-	1	-	-	-	-
Female	-	1	-	2	1	4	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	-	-	-	1	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
New Mexico												
All races, both sexes	1	1	1	2	5	2	5	3	5	3	1	-
Male	-	1	1	2	3	1	2	1	-	1	-	-
Female	1	-	-	-	2	1	3	2	5	2	1	-
White, both sexes	1	1	1	2	5	1	5	2	5	3	1	-
Male	-	1	1	2	3	-	2	-	-	1	-	-
Female	1	-	-	-	2	1	3	2	5	2	1	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	11	17	12	15	29	33	64	78	71	16	4	-
Male	8	11	6	8	19	16	29	23	21	3	-	-
Female	3	6	6	7	10	17	35	55	50	13	4	-
White, both sexes	9	13	8	11	20	29	64	75	65	16	4	-
Male	6	8	4	6	15	14	29	23	21	3	-	-
Female	3	5	4	5	5	15	35	52	44	13	4	-
Black, both sexes	1	3	4	2	6	3	-	3	4	-	-	-
Male	1	3	2	1	4	2	-	-	-	-	-	-
Female	-	-	2	1	2	1	-	3	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	2	1	1	-	-	1	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	1	1	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
North Carolina												
All races, both sexes	14	11	12	17	26	37	39	49	21	15	-	-
Male	8	6	7	12	9	18	19	19	6	2	-	-
Female	6	5	5	5	17	19	20	30	15	13	-	-
White, both sexes	6	6	10	12	21	27	33	43	18	15	-	-
Male	4	4	6	8	7	13	18	16	5	2	-	-
Female	2	2	4	4	14	14	15	27	13	13	-	-
Black, both sexes	8	3	2	4	5	10	6	6	3	-	-	-
Male	4	1	1	3	2	5	1	3	1	-	-	-
Female	4	2	1	1	3	5	5	3	2	-	-	-
American Indian 1/, both sexes	-	2	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	2	-	-	2	1	4	1	4	3	1	-	-
Male	1	-	-	2	1	2	-	1	2	-	-	-
Female	1	-	-	-	-	2	1	3	1	1	-	-
White, both sexes	2	-	-	2	1	4	1	4	3	1	-	-
Male	1	-	-	2	1	2	-	1	2	-	-	-
Female	1	-	-	-	-	2	1	3	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Ohio												
All races, both sexes	9	11	17	25	25	41	89	83	53	17	5	-
Male	5	9	7	15	11	12	36	22	13	1	-	-
Female	4	2	10	10	14	29	53	61	40	16	5	-
White, both sexes	7	9	13	20	23	36	84	83	51	17	5	-
Male	3	7	5	12	10	10	33	22	13	1	-	-
Female	4	2	8	8	13	26	51	61	38	16	5	-
Black, both sexes	2	2	4	5	1	4	4	-	2	-	-	-
Male	2	2	2	3	-	2	2	-	-	-	-	-
Female	-	-	2	2	1	2	2	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Oklahoma												
All races, both sexes	6	3	6	5	6	11	15	25	19	4	1	-
Male	4	2	5	2	2	6	6	8	5	1	-	-
Female	2	1	1	3	4	5	9	17	14	3	1	-
White, both sexes	3	2	5	2	5	10	14	24	17	3	1	-
Male	2	1	4	1	1	6	5	7	4	1	-	-
Female	1	1	1	1	4	4	9	17	13	2	1	-
Black, both sexes	2	-	-	2	-	-	-	1	2	1	-	-
Male	1	-	-	1	-	-	-	1	1	-	-	-
Female	1	-	-	1	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	1	1	-	1	1	1	1	-	-	-	-	-
Male	1	1	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Oregon												
All races, both sexes	1	4	4	5	6	11	22	26	16	7	-	-
Male	1	3	2	3	2	4	4	8	6	2	-	-
Female	-	1	2	2	4	7	18	18	10	5	-	-
White, both sexes	1	3	4	5	5	10	22	26	16	7	-	-
Male	1	2	2	3	1	3	4	8	6	2	-	-
Female	-	1	2	2	4	7	18	18	10	5	-	-
Black, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	9	17	10	26	31	57	117	103	93	26	4	-
Male	5	9	6	12	11	19	51	32	24	5	-	-
Female	4	8	4	14	20	38	66	71	69	21	4	-
White, both sexes	8	16	8	22	23	50	109	98	85	24	4	-
Male	5	8	4	11	8	17	49	30	20	4	-	-
Female	3	8	4	11	15	33	60	68	65	20	4	-
Black, both sexes	1	1	2	3	7	7	7	5	7	1	-	-
Male	-	1	2	1	2	2	2	2	3	-	-	-
Female	1	-	-	2	5	5	5	3	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	1	-	1	1	-	-
Male	-	-	-	-	1	-	-	-	1	1	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Rhode Island												
All races, both sexes	-	2	1	1	1	2	4	6	1	-	-	-
Male	-	2	1	1	-	1	1	2	-	-	-	-
Female	-	-	-	-	1	1	3	4	1	-	-	-
White, both sexes	-	2	1	1	1	2	4	6	1	-	-	-
Male	-	2	1	1	-	1	1	2	-	-	-	-
Female	-	-	-	-	1	1	3	4	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	4	3	8	10	16	31	39	17	20	9	1	-
Male	2	1	5	2	6	11	15	4	3	2	-	-
Female	2	2	3	8	10	20	24	13	17	7	1	-
White, both sexes	1	1	5	10	12	21	34	12	16	7	1	-
Male	1	-	3	2	5	9	11	1	3	1	-	-
Female	-	1	2	8	7	12	23	11	13	6	1	-
Black, both sexes	3	2	3	-	4	10	5	5	4	2	-	-
Male	1	1	2	-	1	2	4	3	-	1	-	-
Female	2	1	1	-	3	8	1	2	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
South Dakota												
All races, both sexes	-	3	1	3	2	5	6	2	2	2	-	-
Male	-	2	1	1	-	3	2	-	2	-	-	-
Female	-	1	-	2	2	2	4	2	-	2	-	-
White, both sexes	-	2	1	2	1	5	6	2	2	2	-	-
Male	-	1	1	1	-	3	2	-	2	-	-	-
Female	-	1	-	1	1	2	4	2	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	1	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	11	7	15	19	28	27	41	31	13	3	-
Male	2	8	3	7	10	13	10	21	5	5	-	-
Female	1	3	4	8	9	15	17	20	26	8	3	-
White, both sexes	2	7	5	11	17	26	21	39	28	12	2	-
Male	1	4	3	6	9	13	7	19	4	5	-	-
Female	1	3	2	5	8	13	14	20	24	7	2	-
Black, both sexes	1	4	2	4	2	2	6	2	3	1	1	-
Male	1	4	-	1	1	-	3	2	1	-	-	-
Female	-	-	2	3	1	2	3	-	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Texas												
All races, both sexes	12	20	22	29	53	59	96	76	39	15	2	-
Male	7	9	17	13	22	28	44	23	12	3	-	-
Female	5	11	5	16	31	31	52	53	27	12	2	-
White, both sexes	11	15	18	24	42	47	86	69	38	14	2	-
Male	6	7	13	9	18	21	40	21	12	2	-	-
Female	5	8	5	15	24	26	46	48	26	12	2	-
Black, both sexes	1	3	4	3	11	12	9	6	1	1	-	-
Male	1	1	4	2	4	7	3	1	-	1	-	-
Female	-	2	-	1	7	5	6	5	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	2	-	-	1	1	-	-	-	-
Male	-	1	-	2	-	-	1	1	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	1	1	5	3	1	3	5	3	1	-	-
Male	-	1	-	1	3	-	1	3	1	-	-	-
Female	-	-	1	4	-	1	2	2	2	1	-	-
White, both sexes	-	1	1	5	3	1	3	5	3	1	-	-
Male	-	1	-	1	3	-	1	3	1	-	-	-
Female	-	-	1	4	-	1	2	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Vermont												
All races, both sexes	-	-	-	-	1	-	1	2	3	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	3	1	-	-
White, both sexes	-	-	-	-	1	-	1	2	3	1	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	1	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4	6	15	24	16	29	28	39	31	8	1	-
Male	2	1	9	15	10	11	8	9	11	2	-	-
Female	2	5	6	9	6	18	20	30	20	6	1	-
White, both sexes	2	4	8	18	15	19	23	34	24	3	1	-
Male	1	1	6	10	9	9	7	8	7	-	-	-
Female	1	3	2	8	6	10	16	26	17	3	1	-
Black, both sexes	2	2	7	6	1	9	4	5	7	4	-	-
Male	1	-	3	5	1	1	1	1	4	2	-	-
Female	1	2	4	1	-	8	3	4	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Washington												
All races, both sexes	4	5	9	15	14	25	38	33	19	9	-	-
Male	3	5	7	9	6	9	20	11	4	1	-	-
Female	1	-	2	6	8	16	18	22	15	8	-	-
White, both sexes	4	4	8	13	13	22	38	32	19	9	-	-
Male	3	4	6	8	6	9	20	10	4	1	-	-
Female	1	-	2	5	7	13	18	22	15	8	-	-
Black, both sexes	-	1	-	-	1	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	1	3	4	2	12	17	13	6	2	-	-
Male	1	-	1	3	1	5	5	2	1	-	-	-
Female	-	1	2	1	1	7	12	11	5	2	-	-
White, both sexes	1	1	3	4	2	11	17	12	6	2	-	-
Male	1	-	1	3	1	5	5	2	1	-	-	-
Female	-	1	2	1	1	6	12	10	5	2	-	-
Black, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Cerebral infarction (I63)												
Wisconsin												
All races, both sexes	2	2	4	7	16	15	21	39	16	8	1	-
Male	2	1	3	2	8	12	11	14	4	2	-	-
Female	-	1	1	5	8	3	10	25	12	6	1	-
White, both sexes	1	1	4	4	14	14	20	37	15	8	1	-
Male	1	1	3	1	7	11	10	13	4	2	-	-
Female	-	-	1	3	7	3	10	24	11	6	1	-
Black, both sexes	1	1	-	2	1	1	-	1	1	-	-	-
Male	1	-	-	1	1	1	-	1	-	-	-	-
Female	-	1	-	1	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Wyoming												
All races, both sexes	-	1	1	-	1	2	3	2	1	-	-	-
Male	-	1	-	-	1	1	3	1	-	-	-	-
Female	-	-	1	-	-	1	-	1	1	-	-	-
White, both sexes	-	1	1	-	1	2	3	2	1	-	-	-
Male	-	1	-	-	1	1	3	1	-	-	-	-
Female	-	-	1	-	-	1	-	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
United States												
All races, both sexes	74,416	4	4	8	3	5	7	30	40	109	292	571
Male	27,963	1	2	3	1	2	2	17	20	55	163	324
Female	46,453	3	2	5	2	3	5	13	20	54	129	247
White, both sexes	63,946	4	3	7	3	4	6	16	24	63	180	357
Male	23,570	1	1	3	1	1	1	9	11	33	99	211
Female	40,376	3	2	4	2	3	5	7	13	30	81	146
Black, both sexes	8,637	-	1	1	-	1	1	14	15	38	106	198
Male	3,573	-	1	-	-	1	1	8	8	20	59	106
Female	5,064	-	-	1	-	-	-	6	7	18	47	92
American Indian 1/, both sexes	298	-	-	-	-	-	-	-	1	4	1	4
Male	101	-	-	-	-	-	-	-	1	-	1	2
Female	197	-	-	-	-	-	-	-	-	4	-	2
Asian or Pacific Islander 2/, both sexes	1,535	-	-	-	-	-	-	-	-	4	5	12
Male	719	-	-	-	-	-	-	-	-	2	4	5
Female	816	-	-	-	-	-	-	-	-	2	1	7
Alabama												
All races, both sexes	1,651	-	-	-	1	-	1	4	2	4	12	22
Male	598	-	-	-	-	-	1	3	2	1	7	14
Female	1,053	-	-	-	1	-	-	1	-	3	5	8
White, both sexes	1,198	-	-	-	1	-	1	2	1	3	7	8
Male	419	-	-	-	-	-	1	2	1	1	5	3
Female	779	-	-	-	1	-	-	-	-	2	2	5
Black, both sexes	449	-	-	-	-	-	-	2	-	1	5	14
Male	177	-	-	-	-	-	-	1	-	-	2	11
Female	272	-	-	-	-	-	-	1	-	1	3	3
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Alaska												
All races, both sexes	86	-	-	-	-	-	-	-	-	-	1	-
Male	44	-	-	-	-	-	-	-	-	-	1	-
Female	42	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	54	-	-	-	-	-	-	-	-	-	1	-
Male	28	-	-	-	-	-	-	-	-	-	1	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	1,326	1	-	-	-	-	-	-	1	2	6	9
Male	524	-	-	-	-	-	-	-	-	1	4	5
Female	802	1	-	-	-	-	-	-	1	1	2	4
White, both sexes	1,252	1	-	-	-	-	-	-	1	1	5	7
Male	495	-	-	-	-	-	-	-	-	-	3	4
Female	757	1	-	-	-	-	-	-	1	1	2	3
Black, both sexes	37	-	-	-	-	-	-	-	-	1	1	2
Male	15	-	-	-	-	-	-	-	-	1	1	1
Female	22	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Arkansas												
All races, both sexes	1,087	-	-	-	-	-	1	2	1	4	9	8
Male	417	-	-	-	-	-	-	1	-	2	6	6
Female	670	-	-	-	-	-	1	1	1	2	3	2
White, both sexes	916	-	-	-	-	-	1	-	1	3	5	7
Male	340	-	-	-	-	-	-	-	-	1	3	6
Female	576	-	-	-	-	-	1	-	1	2	2	1
Black, both sexes	165	-	-	-	-	-	-	2	-	1	4	1
Male	76	-	-	-	-	-	-	1	-	1	3	-
Female	89	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	7,391	-	-	1	-	-	-	-	4	12	38	48
Male	2,923	-	-	-	-	-	-	-	1	3	22	31
Female	4,468	-	-	1	-	-	-	-	3	9	16	17
White, both sexes	6,198	-	-	1	-	-	-	-	3	8	27	38
Male	2,413	-	-	-	-	-	-	-	1	1	16	24
Female	3,785	-	-	1	-	-	-	-	2	7	11	14
Black, both sexes	501	-	-	-	-	-	-	-	1	4	8	8
Male	195	-	-	-	-	-	-	-	-	2	4	5
Female	306	-	-	-	-	-	-	-	1	2	4	3
American Indian 1/, both sexes	38	-	-	-	-	-	-	-	-	-	1	-
Male	14	-	-	-	-	-	-	-	-	-	1	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	654	-	-	-	-	-	-	-	-	-	2	2
Male	301	-	-	-	-	-	-	-	-	-	1	2
Female	353	-	-	-	-	-	-	-	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Colorado												
All races, both sexes	833	-	1	-	-	-	-	-	-	2	2	8
Male	332	-	1	-	-	-	-	-	-	2	-	5
Female	501	-	-	-	-	-	-	-	-	-	2	3
White, both sexes	791	-	1	-	-	-	-	-	-	2	2	7
Male	314	-	1	-	-	-	-	-	-	2	-	4
Female	477	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	776	-	-	-	-	-	-	-	-	1	2	5
Male	302	-	-	-	-	-	-	-	-	-	-	2
Female	474	-	-	-	-	-	-	-	-	1	2	3
White, both sexes	721	-	-	-	-	-	-	-	-	1	1	3
Male	278	-	-	-	-	-	-	-	-	-	-	1
Female	443	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	52	-	-	-	-	-	-	-	-	-	1	2
Male	23	-	-	-	-	-	-	-	-	-	-	1
Female	29	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Delaware												
All races, both sexes	200	-	-	-	-	-	-	-	-	1	-	-
Male	94	-	-	-	-	-	-	-	-	1	-	-
Female	106	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	161	-	-	-	-	-	-	-	-	1	-	-
Male	75	-	-	-	-	-	-	-	-	1	-	-
Female	86	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	112	-	-	-	-	-	-	-	1	2	-	2
Male	44	-	-	-	-	-	-	-	-	1	-	1
Female	68	-	-	-	-	-	-	-	1	1	-	1
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	86	-	-	-	-	-	-	-	1	2	-	2
Male	33	-	-	-	-	-	-	-	-	1	-	1
Female	53	-	-	-	-	-	-	-	1	1	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Florida												
All races, both sexes	4,626	-	-	-	-	-	1	-	4	6	16	32
Male	1,908	-	-	-	-	-	1	-	3	1	12	17
Female	2,718	-	-	-	-	-	-	-	1	5	4	15
White, both sexes	3,974	-	-	-	-	-	-	-	2	2	12	17
Male	1,633	-	-	-	-	-	-	-	1	1	8	12
Female	2,341	-	-	-	-	-	-	-	1	1	4	5
Black, both sexes	620	-	-	-	-	-	1	-	2	3	4	14
Male	266	-	-	-	-	-	1	-	2	-	4	5
Female	354	-	-	-	-	-	-	-	-	3	-	9
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	27	-	-	-	-	-	-	-	-	1	-	1
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	1	-	1
Georgia												
All races, both sexes	2,149	-	1	1	-	-	-	1	1	8	14	24
Male	829	-	-	-	-	-	-	1	1	4	7	8
Female	1,320	-	1	1	-	-	-	-	-	4	7	16
White, both sexes	1,544	-	1	-	-	-	-	-	-	3	6	13
Male	579	-	-	-	-	-	-	-	-	2	1	4
Female	965	-	1	-	-	-	-	-	-	1	5	9
Black, both sexes	589	-	-	1	-	-	-	1	1	3	8	11
Male	245	-	-	-	-	-	-	1	1	2	6	4
Female	344	-	-	1	-	-	-	-	-	1	2	7
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Hawaii												
All races, both sexes	369	-	-	-	-	-	-	-	-	1	3	5
Male	178	-	-	-	-	-	-	-	-	1	2	4
Female	191	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	87	-	-	-	-	-	-	-	-	-	1	3
Male	38	-	-	-	-	-	-	-	-	-	-	3
Female	49	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	282	-	-	-	-	-	-	-	-	1	2	2
Male	140	-	-	-	-	-	-	-	-	1	2	1
Female	142	-	-	-	-	-	-	-	-	-	-	1
Idaho												
All races, both sexes	390	-	-	-	-	-	-	-	1	-	2	1
Male	163	-	-	-	-	-	-	-	1	-	1	-
Female	227	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	384	-	-	-	-	-	-	-	1	-	2	1
Male	160	-	-	-	-	-	-	-	1	-	1	-
Female	224	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Illinois												
All races, both sexes	3,326	1	1	1	-	-	-	3	2	7	8	20
Male	1,207	-	-	-	-	-	-	1	-	6	6	10
Female	2,119	1	1	1	-	-	-	2	2	1	2	10
White, both sexes	2,837	1	1	1	-	-	-	3	-	3	4	7
Male	1,009	-	-	-	-	-	-	1	-	2	3	4
Female	1,828	1	1	1	-	-	-	2	-	1	1	3
Black, both sexes	436	-	-	-	-	-	-	-	2	4	4	13
Male	177	-	-	-	-	-	-	-	-	4	3	6
Female	259	-	-	-	-	-	-	-	2	-	1	7
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	49	-	-	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1,864	-	-	-	-	-	-	2	-	1	6	14
Male	692	-	-	-	-	-	-	1	-	-	2	5
Female	1,172	-	-	-	-	-	-	1	-	1	4	9
White, both sexes	1,718	-	-	-	-	-	-	2	-	-	3	11
Male	624	-	-	-	-	-	-	1	-	-	1	3
Female	1,094	-	-	-	-	-	-	1	-	-	2	8
Black, both sexes	141	-	-	-	-	-	-	-	-	1	3	3
Male	65	-	-	-	-	-	-	-	-	-	1	2
Female	76	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Iowa												
All races, both sexes	1,110	-	-	-	-	1	-	-	-	-	1	4
Male	408	-	-	-	-	-	-	-	-	-	-	4
Female	702	-	-	-	-	1	-	-	-	-	1	-
White, both sexes	1,099	-	-	-	-	1	-	-	-	-	1	4
Male	400	-	-	-	-	-	-	-	-	-	-	4
Female	699	-	-	-	-	1	-	-	-	-	1	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	893	-	-	-	-	-	-	2	-	1	1	3
Male	332	-	-	-	-	-	-	-	-	1	1	1
Female	561	-	-	-	-	-	-	2	-	-	-	2
White, both sexes	839	-	-	-	-	-	-	1	-	1	1	3
Male	307	-	-	-	-	-	-	-	-	1	1	1
Female	532	-	-	-	-	-	-	1	-	-	-	2
Black, both sexes	41	-	-	-	-	-	-	1	-	-	-	-
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Kentucky												
All races, both sexes	1,242	-	-	-	-	-	-	-	1	1	3	8
Male	447	-	-	-	-	-	-	-	1	-	3	5
Female	795	-	-	-	-	-	-	-	-	1	-	3
White, both sexes	1,154	-	-	-	-	-	-	-	1	-	1	5
Male	415	-	-	-	-	-	-	-	1	-	1	3
Female	739	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	87	-	-	-	-	-	-	-	-	1	2	3
Male	32	-	-	-	-	-	-	-	-	-	2	2
Female	55	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	1,333	-	-	-	-	1	-	1	-	-	12	27
Male	512	-	-	-	-	-	-	1	-	-	3	16
Female	821	-	-	-	-	1	-	-	-	-	9	11
White, both sexes	863	-	-	-	-	1	-	1	-	-	2	5
Male	331	-	-	-	-	-	-	1	-	-	1	3
Female	532	-	-	-	-	1	-	-	-	-	1	2
Black, both sexes	460	-	-	-	-	-	-	-	-	-	10	21
Male	175	-	-	-	-	-	-	-	-	-	2	13
Female	285	-	-	-	-	-	-	-	-	-	8	8
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Maine												
All races, both sexes	407	-	-	-	-	-	-	-	-	-	-	2
Male	131	-	-	-	-	-	-	-	-	-	-	1
Female	276	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	406	-	-	-	-	-	-	-	-	-	-	2
Male	131	-	-	-	-	-	-	-	-	-	-	1
Female	275	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	1,223	-	-	-	-	-	-	-	-	2	7	9
Male	469	-	-	-	-	-	-	-	-	2	3	6
Female	754	-	-	-	-	-	-	-	-	-	4	3
White, both sexes	929	-	-	-	-	-	-	-	-	1	4	4
Male	337	-	-	-	-	-	-	-	-	1	2	4
Female	592	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	270	-	-	-	-	-	-	-	-	1	3	4
Male	120	-	-	-	-	-	-	-	-	1	1	1
Female	150	-	-	-	-	-	-	-	-	-	2	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	11	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Massachusetts												
All races, both sexes	1,493	-	-	-	-	-	1	-	-	3	-	5
Male	479	-	-	-	-	-	-	-	-	-	-	3
Female	1,014	-	-	-	-	-	1	-	-	3	-	2
White, both sexes	1,414	-	-	-	-	-	1	-	-	3	-	4
Male	447	-	-	-	-	-	-	-	-	-	-	3
Female	967	-	-	-	-	-	1	-	-	3	-	1
Black, both sexes	58	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	2,277	-	-	-	-	-	-	-	1	3	7	14
Male	873	-	-	-	-	-	-	-	1	3	5	10
Female	1,404	-	-	-	-	-	-	-	-	-	2	4
White, both sexes	1,985	-	-	-	-	-	-	-	-	3	5	8
Male	751	-	-	-	-	-	-	-	-	3	3	6
Female	1,234	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	274	-	-	-	-	-	-	-	1	-	2	6
Male	115	-	-	-	-	-	-	-	1	-	2	4
Female	159	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Minnesota												
All races, both sexes	1,158	-	-	-	-	-	-	1	-	-	-	2
Male	405	-	-	-	-	-	-	1	-	-	-	1
Female	753	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	1,133	-	-	-	-	-	-	-	-	-	-	1
Male	390	-	-	-	-	-	-	-	-	-	-	-
Female	743	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	8	-	-	-	-	-	-	1	-	-	-	1
Male	5	-	-	-	-	-	-	1	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	1,006	-	-	-	-	-	-	2	2	1	10	11
Male	419	-	-	-	-	-	-	1	2	-	6	3
Female	587	-	-	-	-	-	-	1	-	1	4	8
White, both sexes	667	-	-	-	-	-	-	-	2	-	4	3
Male	255	-	-	-	-	-	-	-	2	-	1	-
Female	412	-	-	-	-	-	-	-	-	-	3	3
Black, both sexes	332	-	-	-	-	-	-	2	-	1	6	8
Male	162	-	-	-	-	-	-	1	-	-	5	3
Female	170	-	-	-	-	-	-	1	-	1	1	5
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Missouri												
All races, both sexes	1,755	-	-	1	-	-	-	-	-	3	7	15
Male	623	-	-	-	-	-	-	-	-	1	3	9
Female	1,132	-	-	1	-	-	-	-	-	2	4	6
White, both sexes	1,610	-	-	1	-	-	-	-	-	1	5	15
Male	567	-	-	-	-	-	-	-	-	-	2	9
Female	1,043	-	-	1	-	-	-	-	-	1	3	6
Black, both sexes	137	-	-	-	-	-	-	-	-	2	2	-
Male	55	-	-	-	-	-	-	-	-	1	1	-
Female	82	-	-	-	-	-	-	-	-	1	1	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	288	-	-	-	-	-	-	-	-	-	-	2
Male	110	-	-	-	-	-	-	-	-	-	-	2
Female	178	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	280	-	-	-	-	-	-	-	-	-	-	1
Male	108	-	-	-	-	-	-	-	-	-	-	1
Female	172	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Nebraska												
All races, both sexes	547	-	-	-	-	-	-	1	-	1	-	2
Male	215	-	-	-	-	-	-	1	-	1	-	2
Female	332	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	528	-	-	-	-	-	-	-	-	1	-	1
Male	203	-	-	-	-	-	-	-	-	1	-	1
Female	325	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	-	1	-	-	-	1
Male	12	-	-	-	-	-	-	1	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	542	-	-	1	-	-	-	-	2	1	3	6
Male	239	-	-	1	-	-	-	-	1	1	2	3
Female	303	-	-	-	-	-	-	-	1	-	1	3
White, both sexes	466	-	-	1	-	-	-	-	1	1	2	3
Male	198	-	-	1	-	-	-	-	-	1	1	1
Female	268	-	-	-	-	-	-	-	1	-	1	2
Black, both sexes	52	-	-	-	-	-	-	-	1	-	1	3
Male	27	-	-	-	-	-	-	-	1	-	1	2
Female	25	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
New Hampshire												
All races, both sexes	242	-	-	-	-	-	-	-	-	-	-	-
Male	75	-	-	-	-	-	-	-	-	-	-	-
Female	167	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	242	-	-	-	-	-	-	-	-	-	-	-
Male	75	-	-	-	-	-	-	-	-	-	-	-
Female	167	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1,955	-	-	-	1	-	-	-	1	4	7	11
Male	719	-	-	-	-	-	-	-	1	2	4	9
Female	1,236	-	-	-	1	-	-	-	-	2	3	2
White, both sexes	1,633	-	-	-	1	-	-	-	-	2	6	5
Male	601	-	-	-	-	-	-	-	-	1	4	5
Female	1,032	-	-	-	1	-	-	-	-	1	2	-
Black, both sexes	285	-	-	-	-	-	-	-	1	2	1	6
Male	103	-	-	-	-	-	-	-	1	1	-	4
Female	182	-	-	-	-	-	-	-	-	1	1	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
New Mexico												
All races, both sexes	351	-	-	-	-	-	-	-	1	-	-	1
Male	120	-	-	-	-	-	-	-	-	-	-	1
Female	231	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	328	-	-	-	-	-	-	-	1	-	-	1
Male	112	-	-	-	-	-	-	-	-	-	-	1
Female	216	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	3,448	-	-	-	-	-	-	3	2	1	19	21
Male	1,245	-	-	-	-	-	-	3	1	-	7	13
Female	2,203	-	-	-	-	-	-	-	1	1	12	8
White, both sexes	2,955	-	-	-	-	-	-	1	-	1	10	16
Male	1,044	-	-	-	-	-	-	1	-	-	3	11
Female	1,911	-	-	-	-	-	-	-	-	1	7	5
Black, both sexes	404	-	-	-	-	-	-	2	2	-	9	5
Male	153	-	-	-	-	-	-	2	1	-	4	2
Female	251	-	-	-	-	-	-	-	1	-	5	3
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	85	-	-	-	-	-	-	-	-	-	-	-
Male	46	-	-	-	-	-	-	-	-	-	-	-
Female	39	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
North Carolina												
All races, both sexes	2,518	-	1	1	-	-	-	2	1	4	10	19
Male	954	-	1	-	-	-	-	1	1	2	5	12
Female	1,564	-	-	1	-	-	-	1	-	2	5	7
White, both sexes	1,927	-	-	1	-	-	-	1	-	1	5	10
Male	709	-	-	-	-	-	-	1	-	-	2	7
Female	1,218	-	-	1	-	-	-	-	-	1	3	3
Black, both sexes	559	-	1	-	-	-	-	1	1	3	5	8
Male	229	-	1	-	-	-	-	-	1	2	3	4
Female	330	-	-	-	-	-	-	1	-	1	2	4
American Indian 1/, both sexes	21	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	207	-	-	-	-	-	-	-	-	-	-	-
Male	74	-	-	-	-	-	-	-	-	-	-	-
Female	133	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	204	-	-	-	-	-	-	-	-	-	-	-
Male	72	-	-	-	-	-	-	-	-	-	-	-
Female	132	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Ohio												
All races, both sexes	2,909	1	-	1	-	1	-	-	-	4	9	28
Male	1,052	-	-	1	-	1	-	-	-	2	5	14
Female	1,857	1	-	-	-	-	-	-	-	2	4	14
White, both sexes	2,621	1	-	1	-	1	-	-	-	2	5	23
Male	947	-	-	1	-	1	-	-	-	1	4	11
Female	1,674	1	-	-	-	-	-	-	-	1	1	12
Black, both sexes	279	-	-	-	-	-	-	-	-	2	4	5
Male	102	-	-	-	-	-	-	-	-	1	1	3
Female	177	-	-	-	-	-	-	-	-	1	3	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	1,316	-	-	-	-	-	-	-	1	3	3	10
Male	490	-	-	-	-	-	-	-	-	1	1	4
Female	826	-	-	-	-	-	-	-	1	2	2	6
White, both sexes	1,178	-	-	-	-	-	-	-	1	1	3	7
Male	439	-	-	-	-	-	-	-	-	-	1	3
Female	739	-	-	-	-	-	-	-	1	1	2	4
Black, both sexes	74	-	-	-	-	-	-	-	-	1	-	2
Male	27	-	-	-	-	-	-	-	-	1	-	1
Female	47	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	52	-	-	-	-	-	-	-	-	1	-	1
Male	17	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Oregon												
All races, both sexes	1,229	-	-	-	-	-	-	-	1	-	3	12
Male	446	-	-	-	-	-	-	-	-	-	2	6
Female	783	-	-	-	-	-	-	-	1	-	1	6
White, both sexes	1,192	-	-	-	-	-	-	-	1	-	3	10
Male	429	-	-	-	-	-	-	-	-	-	2	5
Female	763	-	-	-	-	-	-	-	1	-	1	5
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3,999	-	-	-	-	-	1	2	2	1	10	22
Male	1,415	-	-	-	-	-	-	1	1	-	6	14
Female	2,584	-	-	-	-	-	1	1	1	1	4	8
White, both sexes	3,639	-	-	-	-	-	1	2	2	-	7	13
Male	1,273	-	-	-	-	-	-	1	1	-	5	9
Female	2,366	-	-	-	-	-	1	1	1	-	2	4
Black, both sexes	337	-	-	-	-	-	-	-	-	1	3	8
Male	134	-	-	-	-	-	-	-	-	-	1	5
Female	203	-	-	-	-	-	-	-	-	1	2	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Rhode Island												
All races, both sexes	299	-	-	-	-	-	-	-	-	-	-	2
Male	116	-	-	-	-	-	-	-	-	-	-	1
Female	183	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	290	-	-	-	-	-	-	-	-	-	-	2
Male	110	-	-	-	-	-	-	-	-	-	-	1
Female	180	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	1,279	-	-	-	-	1	-	-	-	2	7	19
Male	500	-	-	-	-	1	-	-	-	1	4	13
Female	779	-	-	-	-	-	-	-	-	1	3	6
White, both sexes	912	-	-	-	-	-	-	-	-	1	1	8
Male	348	-	-	-	-	-	-	-	-	1	1	5
Female	564	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	366	-	-	-	-	1	-	-	-	1	6	11
Male	152	-	-	-	-	1	-	-	-	-	3	8
Female	214	-	-	-	-	-	-	-	-	1	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
South Dakota												
All races, both sexes	290	-	-	-	-	-	-	-	-	-	-	2
Male	112	-	-	-	-	-	-	-	-	-	-	-
Female	178	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	281	-	-	-	-	-	-	-	-	-	-	2
Male	108	-	-	-	-	-	-	-	-	-	-	-
Female	173	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	1,980	-	-	1	-	-	-	-	2	3	11	27
Male	721	-	-	1	-	-	-	-	1	2	4	19
Female	1,259	-	-	-	-	-	-	-	1	1	7	8
White, both sexes	1,672	-	-	1	-	-	-	-	2	2	8	20
Male	584	-	-	1	-	-	-	-	1	1	4	15
Female	1,088	-	-	-	-	-	-	-	1	1	4	5
Black, both sexes	297	-	-	-	-	-	-	-	-	1	3	7
Male	134	-	-	-	-	-	-	-	-	1	-	4
Female	163	-	-	-	-	-	-	-	-	-	3	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Texas												
All races, both sexes	4,648	-	-	-	1	1	1	1	2	9	33	50
Male	1,767	-	-	-	1	-	-	-	-	7	22	29
Female	2,881	-	-	-	-	1	1	1	2	2	11	21
White, both sexes	4,005	-	-	-	1	1	1	-	1	8	22	36
Male	1,508	-	-	-	1	-	-	-	-	6	12	23
Female	2,497	-	-	-	-	1	1	-	1	2	10	13
Black, both sexes	582	-	-	-	-	-	-	1	1	1	10	13
Male	236	-	-	-	-	-	-	-	-	1	9	6
Female	346	-	-	-	-	-	-	1	1	-	1	7
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	57	-	-	-	-	-	-	-	-	-	1	1
Male	22	-	-	-	-	-	-	-	-	-	1	-
Female	35	-	-	-	-	-	-	-	-	-	-	1
Utah												
All races, both sexes	417	-	-	-	-	-	-	-	-	2	-	6
Male	157	-	-	-	-	-	-	-	-	2	-	2
Female	260	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	402	-	-	-	-	-	-	-	-	2	-	5
Male	150	-	-	-	-	-	-	-	-	2	-	1
Female	252	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Vermont												
All races, both sexes	142	-	-	-	-	-	-	-	-	-	-	-
Male	63	-	-	-	-	-	-	-	-	-	-	-
Female	79	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	142	-	-	-	-	-	-	-	-	-	-	-
Male	63	-	-	-	-	-	-	-	-	-	-	-
Female	79	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	1,995	-	-	-	-	-	-	-	1	3	4	13
Male	702	-	-	-	-	-	-	-	1	2	1	3
Female	1,293	-	-	-	-	-	-	-	-	1	3	10
White, both sexes	1,508	-	-	-	-	-	-	-	1	1	3	2
Male	488	-	-	-	-	-	-	-	1	1	1	1
Female	1,020	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	454	-	-	-	-	-	-	-	-	1	1	10
Male	201	-	-	-	-	-	-	-	-	-	-	2
Female	253	-	-	-	-	-	-	-	-	1	1	8
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	29	-	-	-	-	-	-	-	-	1	-	1
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Washington												
All races, both sexes	1,433	-	-	-	-	-	-	-	1	5	2	6
Male	515	-	-	-	-	-	-	-	1	2	2	3
Female	918	-	-	-	-	-	-	-	-	3	-	3
White, both sexes	1,353	-	-	-	-	-	-	-	1	3	2	5
Male	479	-	-	-	-	-	-	-	1	2	2	3
Female	874	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	24	-	-	-	-	-	-	-	-	-	-	-
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	2	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	2	-	-
Asian or Pacific Islander 2/, both sexes	45	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	1
West Virginia												
All races, both sexes	612	-	-	-	-	-	-	1	-	1	-	3
Male	209	-	-	-	-	-	-	-	-	-	-	2
Female	403	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	592	-	-	-	-	-	-	1	-	1	-	3
Male	203	-	-	-	-	-	-	-	-	-	-	2
Female	389	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Stroke, not specified as hemorrhage or infarction (I64)												
Wisconsin												
All races, both sexes	1,559	1	-	-	-	-	1	2	2	-	4	9
Male	556	1	-	-	-	-	-	1	-	-	4	4
Female	1,003	-	-	-	-	-	1	1	2	-	-	5
White, both sexes	1,516	1	-	-	-	-	1	2	1	-	4	6
Male	535	1	-	-	-	-	-	1	-	-	4	2
Female	981	-	-	-	-	-	1	1	1	-	-	4
Black, both sexes	31	-	-	-	-	-	-	-	1	-	-	3
Male	14	-	-	-	-	-	-	-	-	-	-	2
Female	17	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	128	-	-	-	-	-	-	-	-	-	-	2
Male	38	-	-	-	-	-	-	-	-	-	-	1
Female	90	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	123	-	-	-	-	-	-	-	-	-	-	2
Male	38	-	-	-	-	-	-	-	-	-	-	1
Female	85	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
United States												
All races, both sexes	1,044	1,485	2,251	3,427	5,562	9,810	15,331	16,798	12,271	4,507	856	1
Male	644	882	1,329	1,862	2,823	4,521	5,933	5,471	3,061	750	96	1
Female	400	603	922	1,565	2,739	5,289	9,398	11,327	9,210	3,757	760	-
White, both sexes	659	1,006	1,594	2,555	4,370	8,312	13,520	15,234	11,184	4,099	746	-
Male	425	597	951	1,383	2,236	3,859	5,230	4,959	2,798	677	84	-
Female	234	409	643	1,172	2,134	4,453	8,290	10,275	8,386	3,422	662	-
Black, both sexes	345	434	573	722	1,000	1,238	1,446	1,224	852	332	95	1
Male	192	261	334	401	487	531	537	379	185	52	9	1
Female	153	173	239	321	513	707	909	845	667	280	86	-
American Indian 1/, both sexes	17	11	19	34	36	42	48	39	28	9	5	-
Male	9	6	6	13	15	19	12	11	5	1	-	-
Female	8	5	13	21	21	23	36	28	23	8	5	-
Asian or Pacific Islander 2/, both sexes	23	34	65	116	156	218	317	301	207	67	10	-
Male	18	18	38	65	85	112	154	122	73	20	3	-
Female	5	16	27	51	71	106	163	179	134	47	7	-
Alabama												
All races, both sexes	34	53	83	88	138	239	332	327	207	86	18	-
Male	20	29	55	41	68	90	117	95	43	12	-	-
Female	14	24	28	47	70	149	215	232	164	74	18	-
White, both sexes	15	23	45	55	91	177	254	264	172	68	11	-
Male	8	13	30	25	45	69	91	75	39	11	-	-
Female	7	10	15	30	46	108	163	189	133	57	11	-
Black, both sexes	19	30	38	33	46	62	77	62	35	18	7	-
Male	12	16	25	16	23	21	25	20	4	1	-	-
Female	7	14	13	17	23	41	52	42	31	17	7	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Alaska												
All races, both sexes	5	1	6	3	5	13	14	22	7	6	3	-
Male	2	1	3	1	4	7	8	13	3	-	1	-
Female	3	-	3	2	1	6	6	9	4	6	2	-
White, both sexes	2	1	1	2	2	6	8	18	5	5	3	-
Male	1	1	1	1	1	3	5	11	2	-	1	-
Female	1	-	-	1	1	3	3	7	3	5	2	-
Black, both sexes	-	-	2	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	3	1	3	5	2	4	2	1	-	-
Male	1	-	2	-	3	3	-	2	1	-	-	-
Female	2	-	1	1	-	2	2	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	3	-	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Arizona												
All races, both sexes	18	15	37	67	107	164	279	338	195	71	16	-
Male	13	9	22	28	63	84	111	115	54	14	1	-
Female	5	6	15	39	44	80	168	223	141	57	15	-
White, both sexes	15	12	30	57	99	155	266	330	188	70	15	-
Male	11	7	19	22	59	81	107	115	52	14	1	-
Female	4	5	11	35	40	74	159	215	136	56	14	-
Black, both sexes	2	2	3	5	4	5	5	2	3	1	1	-
Male	1	1	2	4	2	1	1	-	-	-	-	-
Female	1	1	1	1	2	4	4	2	3	1	1	-
American Indian 1/, both sexes	1	1	3	4	3	2	3	4	2	-	-	-
Male	1	1	1	2	1	1	1	-	-	-	-	-
Female	-	-	2	2	2	1	2	4	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	2	5	2	2	-	-	-
Male	-	-	-	-	1	1	2	-	2	-	-	-
Female	-	-	1	1	-	1	3	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Arkansas												
All races, both sexes	25	39	34	60	65	143	212	240	167	62	15	-
Male	19	20	20	38	38	72	77	68	42	7	1	-
Female	6	19	14	22	27	71	135	172	125	55	14	-
White, both sexes	15	28	24	47	55	121	188	217	146	49	9	-
Male	13	14	14	31	33	58	68	61	32	5	1	-
Female	2	14	10	16	22	63	120	156	114	44	8	-
Black, both sexes	10	11	10	13	10	21	23	21	20	12	6	-
Male	6	6	6	7	5	14	8	7	10	2	-	-
Female	4	5	4	6	5	7	15	14	10	10	6	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	1	1	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
California												
All races, both sexes	95	118	201	333	542	971	1,563	1,696	1,259	432	78	-
Male	64	71	114	171	279	449	653	615	360	78	12	-
Female	31	47	87	162	263	522	910	1,081	899	354	66	-
White, both sexes	68	91	141	240	398	805	1,330	1,480	1,117	381	70	-
Male	50	53	85	117	213	375	552	532	318	67	9	-
Female	18	38	56	123	185	430	778	948	799	314	61	-
Black, both sexes	17	12	34	53	64	66	85	81	41	24	3	-
Male	7	8	16	30	27	29	30	25	9	2	1	-
Female	10	4	18	23	37	37	55	56	32	22	2	-
American Indian 1/, both sexes	1	4	1	4	7	6	5	4	3	1	1	-
Male	-	3	-	1	1	3	2	2	1	-	-	-
Female	1	1	1	3	6	3	3	2	2	1	1	-
Asian or Pacific Islander 2/, both sexes	9	11	25	36	73	94	143	131	98	26	4	-
Male	7	7	13	23	38	42	69	56	32	9	2	-
Female	2	4	12	13	35	52	74	75	66	17	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Colorado												
All races, both sexes	14	15	17	42	58	108	157	205	148	44	12	-
Male	10	8	8	24	31	53	63	78	41	8	-	-
Female	4	7	9	18	27	55	94	127	107	36	12	-
White, both sexes	10	14	13	35	53	103	150	200	145	44	12	-
Male	7	8	7	20	31	50	60	76	40	8	-	-
Female	3	6	6	15	22	53	90	124	105	36	12	-
Black, both sexes	4	1	2	4	2	4	4	5	3	-	-	-
Male	3	-	1	2	-	2	3	2	1	-	-	-
Female	1	1	1	2	2	2	1	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	3	2	1	3	-	-	-	-	-
Male	-	-	-	2	-	1	-	-	-	-	-	-
Female	-	-	2	1	2	-	3	-	-	-	-	-
Connecticut												
All races, both sexes	10	13	18	24	47	89	159	191	145	63	9	-
Male	6	9	12	17	31	40	71	63	42	8	1	-
Female	4	4	6	7	16	49	88	128	103	55	8	-
White, both sexes	8	10	15	18	41	78	154	182	141	62	7	-
Male	5	8	10	14	27	32	69	61	42	8	1	-
Female	3	2	5	4	14	46	85	121	99	54	6	-
Black, both sexes	2	3	3	5	6	11	5	8	4	1	1	-
Male	1	1	2	2	4	8	2	2	-	-	-	-
Female	1	2	1	3	2	3	3	6	4	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	1	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Delaware												
All races, both sexes	2	7	5	8	15	17	45	57	33	8	2	-
Male	2	5	4	4	10	11	19	24	11	3	-	-
Female	-	2	1	4	5	6	26	33	22	5	2	-
White, both sexes	2	7	2	3	9	12	42	48	27	6	2	-
Male	2	5	1	1	7	10	17	20	9	2	-	-
Female	-	2	1	2	2	2	25	28	18	4	2	-
Black, both sexes	-	-	3	5	4	4	3	9	6	2	-	-
Male	-	-	3	3	1	-	2	4	2	1	-	-
Female	-	-	-	2	3	4	1	5	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	5	1	7	9	10	10	19	17	19	9	1	-
Male	2	-	6	3	7	5	7	5	6	1	-	-
Female	3	1	1	6	3	5	12	12	13	8	1	-
White, both sexes	1	-	-	-	1	3	4	3	7	4	-	-
Male	-	-	-	-	1	2	3	1	3	-	-	-
Female	1	-	-	-	-	1	1	2	4	4	-	-
Black, both sexes	4	1	7	8	9	7	14	13	12	5	1	-
Male	2	-	6	3	6	3	4	3	3	1	-	-
Female	2	1	1	5	3	4	10	10	9	4	1	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Florida												
All races, both sexes	66	70	144	226	336	662	990	1,006	757	259	51	-
Male	36	40	97	129	173	315	414	382	229	53	6	-
Female	30	30	47	97	163	347	576	624	528	206	45	-
White, both sexes	36	48	105	155	270	565	871	926	688	235	42	-
Male	21	30	72	84	144	270	365	362	210	49	4	-
Female	15	18	33	71	126	295	506	564	478	186	38	-
Black, both sexes	29	22	38	68	65	90	112	74	65	24	9	-
Male	14	10	24	44	29	42	48	19	18	4	2	-
Female	15	12	14	24	36	48	64	55	47	20	7	-
American Indian 1/, both sexes	1	-	-	-	-	1	2	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	3	1	6	5	5	4	-	-	-
Male	-	-	1	1	-	3	1	-	1	-	-	-
Female	-	-	-	2	1	3	4	5	3	-	-	-
Georgia												
All races, both sexes	61	83	91	125	209	310	382	448	289	83	18	-
Male	40	54	52	72	95	136	140	148	52	17	2	-
Female	21	29	39	53	114	174	242	300	237	66	16	-
White, both sexes	25	38	55	73	148	223	283	375	228	59	14	-
Male	16	25	31	41	69	100	108	125	42	13	2	-
Female	9	13	24	32	79	123	175	250	186	46	12	-
Black, both sexes	36	45	35	52	57	85	95	72	59	24	4	-
Male	24	29	20	31	25	35	30	23	10	4	-	-
Female	12	16	15	21	32	50	65	49	49	20	4	-
American Indian 1/, both sexes	-	-	-	-	-	1	2	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	4	1	2	1	1	-	-	-
Male	-	-	1	-	1	1	2	-	-	-	-	-
Female	-	-	-	-	3	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Hawaii												
All races, both sexes	4	9	12	16	25	45	72	78	64	33	2	-
Male	4	4	8	10	17	26	38	31	24	9	-	-
Female	-	5	4	6	8	19	34	47	40	24	2	-
White, both sexes	-	1	4	1	7	13	19	18	13	7	-	-
Male	-	1	3	1	3	6	10	8	3	-	-	-
Female	-	-	1	-	4	7	9	10	10	7	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	8	8	15	18	32	53	60	51	26	2	-
Male	4	3	5	9	14	20	28	23	21	9	-	-
Female	-	5	3	6	4	12	25	37	30	17	2	-
Idaho												
All races, both sexes	6	3	9	19	25	43	90	97	71	21	2	-
Male	4	-	7	13	10	24	42	40	17	4	-	-
Female	2	3	2	6	15	19	48	57	54	17	2	-
White, both sexes	5	3	9	17	25	42	89	96	71	21	2	-
Male	4	-	7	13	10	23	41	39	17	4	-	-
Female	1	3	2	4	15	19	48	57	54	17	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	2	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Illinois												
All races, both sexes	38	53	91	138	214	381	750	799	564	219	36	-
Male	26	31	58	79	94	179	287	255	133	40	2	-
Female	12	22	33	59	120	202	463	544	431	179	34	-
White, both sexes	25	37	62	99	169	311	649	715	522	198	30	-
Male	17	19	38	53	76	153	256	223	125	37	2	-
Female	8	18	24	46	93	158	393	492	397	161	28	-
Black, both sexes	11	15	25	34	41	61	87	73	40	21	5	-
Male	8	11	18	23	16	25	26	26	8	3	-	-
Female	3	4	7	11	25	36	61	47	32	18	5	-
American Indian 1/, both sexes	-	-	1	1	1	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	3	4	3	9	13	11	2	-	1	-
Male	1	1	2	2	2	1	5	6	-	-	-	-
Female	1	-	1	2	1	8	8	5	2	-	1	-
Indiana												
All races, both sexes	21	25	76	90	148	238	380	447	290	101	25	-
Male	10	18	45	50	67	110	141	152	69	16	6	-
Female	11	7	31	40	81	128	239	295	221	85	19	-
White, both sexes	11	21	60	76	134	220	351	433	279	95	22	-
Male	5	14	36	41	58	101	130	146	67	15	6	-
Female	6	7	24	35	76	119	221	287	212	80	16	-
Black, both sexes	10	4	14	13	14	17	28	14	11	6	3	-
Male	5	4	8	8	9	8	11	6	2	1	-	-
Female	5	-	6	5	5	9	17	8	9	5	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	1	1	-	-	-	-	-
Male	-	-	1	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Iowa												
All races, both sexes	5	6	17	36	66	110	238	281	231	93	21	-
Male	5	4	10	22	36	53	100	99	58	15	2	-
Female	-	2	7	14	30	57	138	182	173	78	19	-
White, both sexes	5	6	15	36	66	108	237	277	229	93	21	-
Male	5	4	8	22	36	52	99	96	57	15	2	-
Female	-	2	7	14	30	56	138	181	172	78	19	-
Black, both sexes	-	-	2	-	-	1	-	3	2	-	-	-
Male	-	-	2	-	-	-	-	2	1	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	4	21	23	27	59	108	185	219	163	61	16	-
Male	2	17	14	16	28	62	69	70	39	9	3	-
Female	2	4	9	11	31	46	116	149	124	52	13	-
White, both sexes	4	17	20	20	51	99	179	213	156	58	16	-
Male	2	15	12	13	26	55	66	68	36	8	3	-
Female	2	2	8	7	25	44	113	145	120	50	13	-
Black, both sexes	-	4	3	5	2	8	5	5	5	3	-	-
Male	-	2	2	2	-	7	3	2	2	1	-	-
Female	-	2	1	3	2	1	2	3	3	2	-	-
American Indian 1/, both sexes	-	-	-	1	3	1	-	1	2	-	-	-
Male	-	-	-	1	1	-	-	-	1	-	-	-
Female	-	-	-	-	2	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	3	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	2	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Kentucky												
All races, both sexes	19	26	49	68	107	149	236	281	212	68	14	-
Male	15	16	29	41	52	68	91	72	43	10	1	-
Female	4	10	20	27	55	81	145	209	169	58	13	-
White, both sexes	14	23	41	64	97	137	220	268	204	66	13	-
Male	11	14	25	39	48	64	87	70	41	10	1	-
Female	3	9	16	25	49	73	133	198	163	56	12	-
Black, both sexes	5	3	8	4	10	12	16	12	8	2	1	-
Male	4	2	4	2	4	4	4	2	2	-	-	-
Female	1	1	4	2	6	8	12	10	6	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Louisiana												
All races, both sexes	33	62	77	80	116	203	270	225	176	42	8	-
Male	19	36	44	35	57	92	87	73	44	4	1	-
Female	14	26	33	45	59	111	183	152	132	38	7	-
White, both sexes	17	19	33	38	70	134	193	179	131	34	6	-
Male	11	9	18	16	40	64	67	64	34	2	1	-
Female	6	10	15	22	30	70	126	115	97	32	5	-
Black, both sexes	15	42	43	41	46	67	76	46	43	8	2	-
Male	7	26	25	18	17	27	19	9	10	2	-	-
Female	8	16	18	23	29	40	57	37	33	6	2	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	1	-	2	1	-	1	-	-	-
Male	-	1	1	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Maine												
All races, both sexes	3	3	10	10	37	47	83	95	76	35	6	-
Male	3	2	5	4	18	22	28	26	16	5	1	-
Female	-	1	5	6	19	25	55	69	60	30	5	-
White, both sexes	3	3	10	10	37	47	82	95	76	35	6	-
Male	3	2	5	4	18	22	28	26	16	5	1	-
Female	-	1	5	6	19	25	54	69	60	30	5	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	12	27	54	55	87	175	258	253	193	78	13	-
Male	7	16	33	32	43	84	116	67	48	11	1	-
Female	5	11	21	23	44	91	142	186	145	67	12	-
White, both sexes	6	12	25	27	51	138	212	211	167	64	7	-
Male	1	8	17	18	23	64	88	54	46	10	1	-
Female	5	4	8	9	28	74	124	157	121	54	6	-
Black, both sexes	6	13	28	27	35	33	39	38	24	13	6	-
Male	6	7	15	13	19	17	25	12	2	1	-	-
Female	-	6	13	14	16	16	14	26	22	12	6	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	-	1	4	7	3	2	1	-	-
Male	-	1	1	-	1	3	3	1	-	-	-	-
Female	-	1	-	-	-	1	4	2	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Massachusetts												
All races, both sexes	16	24	20	47	82	182	302	389	284	123	15	-
Male	12	12	8	31	43	79	99	122	53	17	-	-
Female	4	12	12	16	39	103	203	267	231	106	15	-
White, both sexes	13	22	18	39	76	163	289	380	275	117	14	-
Male	10	11	7	28	38	71	95	117	51	16	-	-
Female	3	11	11	11	38	92	194	263	224	101	14	-
Black, both sexes	3	1	2	4	4	14	11	7	8	3	-	-
Male	2	-	1	2	3	6	2	5	1	-	-	-
Female	1	1	1	2	1	8	9	2	7	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	1	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	1	-	4	2	5	1	2	1	3	-	-
Male	-	1	-	1	2	2	1	-	1	1	-	-
Female	-	-	-	3	-	3	-	2	-	2	-	-
Michigan												
All races, both sexes	37	44	58	101	175	319	441	524	379	149	25	-
Male	22	25	36	49	86	161	182	157	105	27	4	-
Female	15	19	22	52	89	158	259	367	274	122	21	-
White, both sexes	24	28	43	84	140	271	401	471	345	141	21	-
Male	16	15	26	38	72	143	167	136	96	26	4	-
Female	8	13	17	46	68	128	234	335	249	115	17	-
Black, both sexes	12	16	14	13	31	47	39	49	33	7	4	-
Male	6	10	9	8	12	17	15	21	9	1	-	-
Female	6	6	5	5	19	30	24	28	24	6	4	-
American Indian 1/, both sexes	1	-	-	3	2	-	1	2	-	1	-	-
Male	-	-	-	2	1	-	-	-	-	-	-	-
Female	1	-	-	1	1	-	1	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	2	1	-	2	1	-	-	-
Male	-	-	1	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Minnesota												
All races, both sexes	9	15	19	41	58	115	244	295	263	82	14	-
Male	6	13	13	20	27	54	95	112	54	9	-	-
Female	3	2	6	21	31	61	149	183	209	73	14	-
White, both sexes	8	15	19	37	56	110	240	290	261	82	14	-
Male	6	13	13	19	25	51	92	110	52	9	-	-
Female	2	2	6	18	31	59	148	180	209	73	14	-
Black, both sexes	-	-	-	-	1	3	1	1	-	-	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	1	2	3	3	2	-	-	-
Male	-	-	-	1	1	2	3	1	2	-	-	-
Female	1	-	-	1	-	-	-	2	-	-	-	-
Mississippi												
All races, both sexes	28	28	42	75	102	150	187	197	116	47	8	-
Male	13	22	28	40	67	74	71	62	23	6	1	-
Female	15	6	14	35	35	76	116	135	93	41	7	-
White, both sexes	10	13	24	47	57	100	133	150	86	33	5	-
Male	4	9	16	24	37	50	50	43	16	3	-	-
Female	6	4	8	23	20	50	83	107	70	30	5	-
Black, both sexes	18	15	18	26	44	49	53	45	30	14	3	-
Male	9	13	12	15	30	24	21	18	7	3	1	-
Female	9	2	6	11	14	25	32	27	23	11	2	-
American Indian 1/, both sexes	-	-	-	2	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Missouri												
All races, both sexes	19	29	61	65	134	221	364	380	316	121	19	-
Male	11	17	35	39	55	109	137	111	72	24	-	-
Female	8	12	26	26	79	112	227	269	244	97	19	-
White, both sexes	14	26	47	49	111	207	345	356	302	113	18	-
Male	8	16	27	30	45	105	131	104	68	22	-	-
Female	6	10	20	19	66	102	214	252	234	91	18	-
Black, both sexes	5	3	14	15	20	13	18	24	12	8	1	-
Male	3	1	8	9	9	4	6	7	4	2	-	-
Female	2	2	6	6	11	9	12	17	8	6	1	-
American Indian 1/, both sexes	-	-	-	1	2	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	2	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	2	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	2	-	-	-
Montana												
All races, both sexes	1	4	5	12	20	23	58	83	62	15	3	-
Male	1	2	3	7	10	15	25	27	16	2	-	-
Female	-	2	2	5	10	8	33	56	46	13	3	-
White, both sexes	1	3	5	11	20	23	55	81	62	15	3	-
Male	1	2	3	7	10	15	25	26	16	2	-	-
Female	-	1	2	4	10	8	30	55	46	13	3	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	3	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	3	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Nebraska												
All races, both sexes	6	4	6	13	40	63	119	131	114	43	4	-
Male	4	3	3	9	22	33	56	46	26	8	1	-
Female	2	1	3	4	18	30	63	85	88	35	3	-
White, both sexes	4	2	6	11	37	62	119	126	112	43	4	-
Male	2	1	3	7	20	32	56	46	25	8	1	-
Female	2	1	3	4	17	30	63	80	87	35	3	-
Black, both sexes	2	2	-	2	2	1	-	2	2	-	-	-
Male	2	2	-	2	2	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	2	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	2	-	-	-	-
Nevada												
All races, both sexes	3	13	20	44	63	79	108	114	63	20	2	-
Male	2	12	7	29	34	38	52	34	20	3	-	-
Female	1	1	13	15	29	41	56	80	43	17	2	-
White, both sexes	3	10	15	37	49	66	97	104	55	20	2	-
Male	2	9	6	24	28	31	45	30	16	3	-	-
Female	1	1	9	13	21	35	52	74	39	17	2	-
Black, both sexes	-	2	4	4	11	7	7	7	5	-	-	-
Male	-	2	-	3	5	3	6	2	2	-	-	-
Female	-	-	4	1	6	4	1	5	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	3	3	6	4	3	3	-	-	-
Male	-	1	1	2	1	4	1	2	2	-	-	-
Female	-	-	-	1	2	2	3	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
New Hampshire												
All races, both sexes	-	-	3	13	14	30	56	57	47	18	4	-
Male	-	-	2	8	4	13	22	19	6	1	-	-
Female	-	-	1	5	10	17	34	38	41	17	4	-
White, both sexes	-	-	3	13	14	30	56	57	47	18	4	-
Male	-	-	2	8	4	13	22	19	6	1	-	-
Female	-	-	1	5	10	17	34	38	41	17	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	24	38	58	93	146	254	399	419	346	131	23	-
Male	11	25	39	48	68	122	153	137	83	16	1	-
Female	13	13	19	45	78	132	246	282	263	115	22	-
White, both sexes	15	26	34	63	106	210	342	380	302	120	21	-
Male	8	18	24	34	49	106	137	126	74	14	1	-
Female	7	8	10	29	57	104	205	254	228	106	20	-
Black, both sexes	8	12	21	27	37	38	46	34	40	10	2	-
Male	2	7	14	12	18	13	13	9	8	1	-	-
Female	6	5	7	15	19	25	33	25	32	9	2	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	2	3	3	6	10	5	4	1	-	-
Male	1	-	1	2	1	3	3	2	1	1	-	-
Female	-	-	1	1	2	3	7	3	3	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
New Mexico												
All races, both sexes	6	3	14	14	27	54	64	86	51	23	7	-
Male	5	1	9	8	14	14	26	29	10	2	1	-
Female	1	2	5	6	13	40	38	57	41	21	6	-
White, both sexes	5	1	12	14	26	51	63	80	46	21	7	-
Male	4	1	7	8	13	14	26	26	9	2	1	-
Female	1	-	5	6	13	37	37	54	37	19	6	-
Black, both sexes	-	1	-	-	-	2	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	2	1	-	-	1	-	-
American Indian 1/, both sexes	1	1	2	-	1	1	-	5	5	1	-	-
Male	1	-	2	-	1	-	-	2	1	-	-	-
Female	-	1	-	-	-	1	-	3	4	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	48	62	91	178	259	441	655	800	602	222	44	-
Male	32	35	57	87	124	212	239	254	141	33	7	-
Female	16	27	34	91	135	229	416	546	461	189	37	-
White, both sexes	32	41	69	126	200	366	569	722	558	202	42	-
Male	21	23	42	54	102	180	203	232	132	33	7	-
Female	11	18	27	72	98	186	366	490	426	169	35	-
Black, both sexes	14	16	16	39	54	61	72	60	36	16	2	-
Male	9	10	9	24	20	24	27	15	6	-	-	-
Female	5	6	7	15	34	37	45	45	30	16	2	-
American Indian 1/, both sexes	-	-	1	-	-	1	1	1	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	5	13	5	13	13	17	8	4	-	-
Male	2	2	5	9	2	7	9	7	3	-	-	-
Female	-	3	-	4	3	6	4	10	5	4	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
North Carolina												
All races, both sexes	41	62	93	134	203	316	514	571	372	145	29	-
Male	24	37	54	84	103	133	195	175	97	26	4	-
Female	17	25	39	50	100	183	319	396	275	119	25	-
White, both sexes	21	29	56	89	144	233	401	473	317	123	23	-
Male	15	16	35	56	70	102	150	141	91	21	2	-
Female	6	13	21	33	74	131	251	332	226	102	21	-
Black, both sexes	20	31	34	39	57	79	106	91	55	22	6	-
Male	9	20	19	25	32	29	42	29	6	5	2	-
Female	11	11	15	14	25	50	64	62	49	17	4	-
American Indian 1/, both sexes	-	1	3	4	1	3	5	4	-	-	-	-
Male	-	-	-	2	-	1	1	3	-	-	-	-
Female	-	1	3	2	1	2	4	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	2	1	1	2	3	-	-	-	-
Male	-	1	-	1	1	1	2	2	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
North Dakota												
All races, both sexes	-	2	3	6	12	26	39	45	53	19	2	-
Male	-	-	2	2	7	12	20	18	10	3	-	-
Female	-	2	1	4	5	14	19	27	43	16	2	-
White, both sexes	-	1	3	6	11	26	38	45	53	19	2	-
Male	-	-	2	2	6	12	19	18	10	3	-	-
Female	-	1	1	4	5	14	19	27	43	16	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Ohio												
All races, both sexes	47	42	80	117	212	377	629	656	500	166	39	-
Male	34	22	41	66	112	173	247	203	108	22	1	-
Female	13	20	39	51	100	204	382	453	392	144	38	-
White, both sexes	36	34	60	97	168	337	581	617	470	153	35	-
Male	28	20	31	57	93	160	228	189	100	22	1	-
Female	8	14	29	40	75	177	353	428	370	131	34	-
Black, both sexes	11	8	19	20	42	39	45	38	29	13	4	-
Male	6	2	10	9	18	13	18	14	7	-	-	-
Female	5	6	9	11	24	26	27	24	22	13	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	1	3	1	1	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	1	-	1	1	2	1	-	-	-	-
Oklahoma												
All races, both sexes	23	36	52	84	114	198	272	250	178	73	19	-
Male	16	14	30	46	55	81	98	81	47	15	1	-
Female	7	22	22	38	59	117	174	169	131	58	18	-
White, both sexes	13	30	48	75	93	181	250	233	160	68	15	-
Male	9	10	30	41	46	74	89	76	45	14	1	-
Female	4	20	18	34	47	107	161	157	115	54	14	-
Black, both sexes	5	6	2	2	10	7	11	13	10	2	3	-
Male	2	4	-	1	4	3	6	4	1	-	-	-
Female	3	2	2	1	6	4	5	9	9	2	3	-
American Indian 1/, both sexes	4	-	2	5	7	8	9	3	8	3	1	-
Male	4	-	-	2	4	3	2	-	1	1	-	-
Female	-	-	2	3	3	5	7	3	7	2	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	4	2	2	1	-	-	-	-
Male	1	-	-	2	1	1	1	1	-	-	-	-
Female	-	-	-	-	3	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Oregon												
All races, both sexes	14	18	24	36	79	156	276	307	198	94	11	-
Male	5	12	16	19	30	65	105	112	55	17	2	-
Female	9	6	8	17	49	91	171	195	143	77	9	-
White, both sexes	11	15	23	33	77	150	269	302	194	94	10	-
Male	5	10	15	18	28	63	101	110	53	17	2	-
Female	6	5	8	15	49	87	168	192	141	77	8	-
Black, both sexes	2	2	-	1	1	2	4	2	3	-	-	-
Male	-	1	-	1	1	-	2	1	1	-	-	-
Female	2	1	-	-	-	2	2	1	2	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	1	-	1	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	2	1	3	3	2	1	-	1	-
Male	-	-	1	-	1	2	2	1	1	-	-	-
Female	-	-	-	2	-	1	1	1	-	-	1	-
Pennsylvania												
All races, both sexes	40	77	86	156	281	536	857	976	652	255	45	-
Male	28	46	46	75	149	227	318	299	161	36	8	-
Female	12	31	40	81	132	309	539	677	491	219	37	-
White, both sexes	29	58	64	125	239	487	788	928	616	237	43	-
Male	21	36	34	63	123	206	293	285	154	34	8	-
Female	8	22	30	62	116	281	495	643	462	203	35	-
Black, both sexes	10	19	22	30	38	47	65	44	33	15	2	-
Male	7	10	12	12	22	19	24	14	6	2	-	-
Female	3	9	10	18	16	28	41	30	27	13	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	4	2	3	4	3	2	-	-
Male	-	-	-	-	4	2	1	-	1	-	-	-
Female	1	-	-	1	-	-	2	4	2	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Rhode Island												
All races, both sexes	2	8	6	7	10	43	60	75	67	19	-	-
Male	2	6	4	4	8	18	24	24	21	4	-	-
Female	-	2	2	3	2	25	36	51	46	15	-	-
White, both sexes	2	8	6	7	9	42	57	72	67	18	-	-
Male	2	6	4	4	8	17	22	21	21	4	-	-
Female	-	2	2	3	1	25	35	51	46	14	-	-
Black, both sexes	-	-	-	-	1	1	2	2	-	1	-	-
Male	-	-	-	-	-	1	1	2	-	-	-	-
Female	-	-	-	-	1	-	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	31	43	42	67	137	179	254	236	188	55	18	-
Male	17	26	28	39	69	80	87	78	43	11	3	-
Female	14	17	14	28	68	99	167	158	145	44	15	-
White, both sexes	10	19	23	42	84	136	194	188	148	45	13	-
Male	6	11	18	26	44	61	63	65	36	8	3	-
Female	4	8	5	16	40	75	131	123	112	37	10	-
Black, both sexes	21	24	19	25	53	43	60	48	39	10	5	-
Male	11	15	10	13	25	19	24	13	7	3	-	-
Female	10	9	9	12	28	24	36	35	32	7	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
South Dakota												
All races, both sexes	3	2	8	5	15	30	66	69	59	27	4	-
Male	2	1	6	3	10	16	28	26	14	5	1	-
Female	1	1	2	2	5	14	38	43	45	22	3	-
White, both sexes	3	2	8	4	15	25	66	68	58	26	4	-
Male	2	1	6	3	10	12	28	26	14	5	1	-
Female	1	1	2	1	5	13	38	42	44	21	3	-
Black, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	5	-	-	1	-	-	-
Male	-	-	-	-	-	4	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	33	46	71	103	163	279	408	415	281	116	21	-
Male	14	25	36	58	79	135	131	123	70	20	3	-
Female	19	21	35	45	84	144	277	292	211	96	18	-
White, both sexes	21	32	51	74	128	228	371	364	248	105	17	-
Male	9	18	24	41	60	105	116	109	61	17	2	-
Female	12	14	27	33	68	123	255	255	187	88	15	-
Black, both sexes	12	14	20	29	32	47	36	50	31	11	4	-
Male	5	7	12	17	19	29	15	14	7	3	1	-
Female	7	7	8	12	13	18	21	36	24	8	3	-
American Indian 1/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	3	3	-	1	2	-	-	-
Male	-	-	-	-	-	1	-	-	2	-	-	-
Female	-	-	-	-	3	2	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Texas												
All races, both sexes	72	128	174	243	378	663	966	927	700	266	33	-
Male	43	73	96	131	219	292	367	271	166	48	2	-
Female	29	55	78	112	159	371	599	656	534	218	31	-
White, both sexes	54	96	133	193	304	558	854	834	638	240	31	-
Male	32	56	76	108	175	247	333	245	151	41	2	-
Female	22	40	57	85	129	311	521	589	487	199	29	-
Black, both sexes	17	31	35	41	71	98	101	79	58	23	2	-
Male	10	17	17	20	41	41	31	21	15	7	-	-
Female	7	14	18	21	30	57	70	58	43	16	2	-
American Indian 1/, both sexes	-	-	-	1	-	1	1	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	6	8	3	6	10	14	3	3	-	-
Male	1	-	3	3	3	3	3	5	-	-	-	-
Female	-	1	3	5	-	3	7	9	3	3	-	-
Utah												
All races, both sexes	2	3	8	21	22	64	88	113	67	19	2	-
Male	-	2	1	11	13	27	33	36	26	4	-	-
Female	2	1	7	10	9	37	55	77	41	15	2	-
White, both sexes	2	3	7	20	21	61	84	110	66	19	2	-
Male	-	2	1	11	12	25	31	35	26	4	-	-
Female	2	1	6	9	9	36	53	75	40	15	2	-
Black, both sexes	-	-	-	1	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	1	2	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	3	1	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Vermont												
All races, both sexes	-	2	2	2	9	14	32	37	31	10	3	-
Male	-	1	2	1	5	12	15	13	12	2	-	-
Female	-	1	-	1	4	2	17	24	19	8	3	-
White, both sexes	-	2	2	2	9	14	32	37	31	10	3	-
Male	-	1	2	1	5	12	15	13	12	2	-	-
Female	-	1	-	1	4	2	17	24	19	8	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	27	56	68	90	165	295	381	447	310	105	29	1
Male	15	30	38	46	77	126	139	131	66	20	6	1
Female	12	26	30	44	88	169	242	316	244	85	23	-
White, both sexes	16	34	37	56	95	216	300	362	266	96	23	-
Male	7	15	19	27	42	91	101	106	54	18	4	-
Female	9	19	18	29	53	125	199	256	212	78	19	-
Black, both sexes	11	20	30	30	68	73	75	80	39	9	6	1
Male	8	14	18	17	34	32	35	24	12	2	2	1
Female	3	6	12	13	34	41	40	56	27	7	4	-
American Indian 1/, both sexes	-	1	-	-	-	1	-	1	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	4	2	5	6	4	4	-	-	-
Male	-	-	1	2	1	3	3	1	-	-	-	-
Female	-	1	-	2	1	2	3	3	4	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Washington												
All races, both sexes	11	14	30	51	77	194	287	356	273	98	28	-
Male	6	9	14	28	44	86	117	117	67	17	2	-
Female	5	5	16	23	33	108	170	239	206	81	26	-
White, both sexes	9	13	27	48	65	186	267	346	258	98	25	-
Male	5	8	12	27	38	79	107	113	64	17	1	-
Female	4	5	15	21	27	107	160	233	194	81	24	-
Black, both sexes	1	1	2	-	1	5	7	1	6	-	-	-
Male	-	1	2	-	1	5	4	-	-	-	-	-
Female	1	-	-	-	-	-	3	1	6	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	1	2	2	-	-	2	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	1	-	2	2	-	-	2	-
Asian or Pacific Islander 2/, both sexes	1	-	1	3	9	2	11	7	9	-	1	-
Male	1	-	-	1	4	1	6	4	3	-	1	-
Female	-	-	1	2	5	1	5	3	6	-	-	-
West Virginia												
All races, both sexes	5	11	14	27	58	92	134	131	95	35	5	-
Male	3	7	8	13	25	42	44	32	25	7	1	-
Female	2	4	6	14	33	50	90	99	70	28	4	-
White, both sexes	5	11	12	26	55	89	131	126	94	33	5	-
Male	3	7	7	12	24	40	43	32	25	7	1	-
Female	2	4	5	14	31	49	88	94	69	26	4	-
Black, both sexes	-	-	2	1	3	3	3	5	1	2	-	-
Male	-	-	1	1	1	2	1	-	-	-	-	-
Female	-	-	1	-	2	1	2	5	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Stroke, not specified as hemorrhage or infarction (I64)												
Wisconsin												
All races, both sexes	14	21	31	53	85	174	320	367	315	127	33	-
Male	6	14	21	28	41	82	118	126	85	20	5	-
Female	8	7	10	25	44	92	202	241	230	107	28	-
White, both sexes	10	18	29	49	81	170	311	362	314	126	31	-
Male	5	12	20	26	39	80	114	122	84	20	5	-
Female	5	6	9	23	42	90	197	240	230	106	26	-
Black, both sexes	3	2	1	3	4	4	6	1	1	-	2	-
Male	1	2	1	2	2	2	1	-	1	-	-	-
Female	2	-	-	1	2	2	5	1	-	-	2	-
American Indian 1/, both sexes	1	-	1	1	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	1	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	2	4	-	1	-	-
Male	-	-	-	-	-	-	2	4	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	1	-	-
Wyoming												
All races, both sexes	2	-	1	5	6	18	37	23	23	10	1	-
Male	1	-	-	3	2	8	11	5	5	1	1	-
Female	1	-	1	2	4	10	26	18	18	9	-	-
White, both sexes	1	-	-	5	6	17	36	22	23	10	1	-
Male	1	-	-	3	2	8	11	5	5	1	1	-
Female	-	-	-	2	4	9	25	17	18	9	-	-
Black, both sexes	-	-	-	-	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
United States												
All races, both sexes	27,720	59	35	20	12	15	13	20	25	45	86	187
Male	10,296	38	22	10	10	8	7	13	13	22	34	90
Female	17,424	21	13	10	2	7	6	7	12	23	52	97
White, both sexes	23,898	35	26	12	8	6	8	12	21	36	54	127
Male	8,758	22	16	6	6	4	4	7	13	19	23	65
Female	15,140	13	10	6	2	2	4	5	8	17	31	62
Black, both sexes	3,151	22	7	5	2	8	5	7	4	9	29	57
Male	1,241	15	4	3	2	3	3	5	-	3	10	24
Female	1,910	7	3	2	-	5	2	2	4	6	19	33
American Indian 1/, both sexes	97	1	-	-	-	-	-	-	-	-	2	-
Male	39	-	-	-	-	-	-	-	-	-	1	-
Female	58	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	574	1	2	3	2	1	-	1	-	-	1	3
Male	258	1	2	1	2	1	-	1	-	-	-	1
Female	316	-	-	2	-	-	-	-	-	-	1	2
Alabama												
All races, both sexes	417	3	-	-	-	-	-	-	1	1	5	4
Male	150	2	-	-	-	-	-	-	1	-	2	2
Female	267	1	-	-	-	-	-	-	-	1	3	2
White, both sexes	306	-	-	-	-	-	-	-	1	1	3	4
Male	109	-	-	-	-	-	-	-	1	-	1	2
Female	197	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	110	3	-	-	-	-	-	-	-	-	2	-
Male	40	2	-	-	-	-	-	-	-	-	1	-
Female	70	1	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Alaska												
All races, both sexes	32	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	399	-	1	1	-	1	-	-	-	1	4	2
Male	157	-	1	-	-	1	-	-	-	1	2	1
Female	242	-	-	1	-	-	-	-	-	-	2	1
White, both sexes	384	-	1	1	-	1	-	-	-	1	3	2
Male	148	-	1	-	-	1	-	-	-	1	1	1
Female	236	-	-	1	-	-	-	-	-	-	2	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Arkansas												
All races, both sexes	322	1	-	-	-	-	-	-	-	-	-	4
Male	118	1	-	-	-	-	-	-	-	-	-	-
Female	204	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	250	1	-	-	-	-	-	-	-	-	-	3
Male	96	1	-	-	-	-	-	-	-	-	-	-
Female	154	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	69	-	-	-	-	-	-	-	-	-	-	1
Male	22	-	-	-	-	-	-	-	-	-	-	-
Female	47	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3,980	5	2	1	1	3	1	1	3	4	12	30
Male	1,472	4	2	-	1	2	1	1	1	-	5	13
Female	2,508	1	-	1	-	1	-	-	2	4	7	17
White, both sexes	3,296	4	1	1	1	2	1	-	3	2	7	20
Male	1,189	3	1	-	1	2	1	-	1	-	4	9
Female	2,107	1	-	1	-	-	-	-	2	2	3	11
Black, both sexes	329	-	-	-	-	1	-	-	-	2	4	8
Male	132	-	-	-	-	-	-	-	-	-	1	3
Female	197	-	-	-	-	1	-	-	-	2	3	5
American Indian 1/, both sexes	14	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	341	1	1	-	-	-	-	1	-	-	-	2
Male	147	1	1	-	-	-	-	1	-	-	-	1
Female	194	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Colorado												
All races, both sexes	350	-	-	2	-	-	-	1	1	1	3	2
Male	124	-	-	1	-	-	-	1	-	-	-	2
Female	226	-	-	1	-	-	-	-	1	1	3	-
White, both sexes	336	-	-	2	-	-	-	1	1	1	3	2
Male	119	-	-	1	-	-	-	1	-	-	-	2
Female	217	-	-	1	-	-	-	-	1	1	3	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	270	2	-	3	-	-	-	-	-	2	-	1
Male	100	1	-	2	-	-	-	-	-	1	-	1
Female	170	1	-	1	-	-	-	-	-	1	-	-
White, both sexes	251	1	-	3	-	-	-	-	-	2	-	1
Male	92	1	-	2	-	-	-	-	-	1	-	1
Female	159	-	-	1	-	-	-	-	-	1	-	-
Black, both sexes	18	1	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	10	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Delaware												
All races, both sexes	66	-	1	-	-	-	-	-	-	-	-	-
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	46	-	1	-	-	-	-	-	-	-	-	-
White, both sexes	54	-	1	-	-	-	-	-	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	36	-	1	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Florida												
All races, both sexes	1,910	4	3	2	-	1	-	1	1	2	4	12
Male	745	1	2	1	-	-	-	1	-	2	2	6
Female	1,165	3	1	1	-	1	-	-	1	-	2	6
White, both sexes	1,669	3	1	-	-	1	-	-	-	2	2	9
Male	642	1	1	-	-	-	-	-	-	2	1	5
Female	1,027	2	-	-	-	1	-	-	-	-	1	4
Black, both sexes	228	1	2	2	-	-	-	1	1	-	2	3
Male	97	-	1	1	-	-	-	1	-	-	1	1
Female	131	1	1	1	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	597	2	2	-	1	2	-	-	1	4	5	6
Male	209	2	2	-	1	1	-	-	-	3	1	2
Female	388	-	-	-	-	1	-	-	1	1	4	4
White, both sexes	414	-	-	-	1	1	-	-	-	4	3	4
Male	143	-	-	-	1	-	-	-	-	3	1	1
Female	271	-	-	-	-	1	-	-	-	1	2	3
Black, both sexes	180	2	2	-	-	1	-	-	1	-	2	2
Male	65	2	2	-	-	1	-	-	-	-	-	1
Female	115	-	-	-	-	-	-	-	1	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Hawaii												
All races, both sexes	85	-	1	1	1	-	-	-	-	-	1	-
Male	35	-	1	-	1	-	-	-	-	-	-	-
Female	50	-	-	1	-	-	-	-	-	-	1	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	69	-	1	1	1	-	-	-	-	-	1	-
Male	33	-	1	-	1	-	-	-	-	-	-	-
Female	36	-	-	1	-	-	-	-	-	-	1	-
Idaho												
All races, both sexes	168	-	1	-	-	-	-	-	-	-	-	-
Male	59	-	1	-	-	-	-	-	-	-	-	-
Female	109	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	168	-	1	-	-	-	-	-	-	-	-	-
Male	59	-	1	-	-	-	-	-	-	-	-	-
Female	109	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Illinois												
All races, both sexes	1,211	4	3	2	1	-	1	-	1	2	5	10
Male	461	3	1	1	1	-	1	-	1	1	3	3
Female	750	1	2	1	-	-	-	-	-	1	2	7
White, both sexes	1,008	3	3	1	-	-	-	-	1	2	5	4
Male	373	2	1	-	-	-	-	-	1	1	3	1
Female	635	1	2	1	-	-	-	-	-	1	2	3
Black, both sexes	192	1	-	1	1	-	1	-	-	-	-	6
Male	82	1	-	1	1	-	1	-	-	-	-	2
Female	110	-	-	-	-	-	-	-	-	-	-	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	541	4	2	-	-	-	-	1	-	2	5	4
Male	207	-	1	-	-	-	-	1	-	-	2	-
Female	334	4	1	-	-	-	-	-	-	2	3	4
White, both sexes	488	3	2	-	-	-	-	-	-	2	4	2
Male	181	-	1	-	-	-	-	-	-	-	1	-
Female	307	3	1	-	-	-	-	-	-	2	3	2
Black, both sexes	53	1	-	-	-	-	-	1	-	-	1	2
Male	26	-	-	-	-	-	-	1	-	-	1	-
Female	27	1	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Iowa												
All races, both sexes	332	-	-	-	-	-	-	1	1	-	-	2
Male	128	-	-	-	-	-	-	1	1	-	-	1
Female	204	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	327	-	-	-	-	-	-	1	1	-	-	2
Male	126	-	-	-	-	-	-	1	1	-	-	1
Female	201	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	273	-	-	-	-	-	-	-	-	-	-	-
Male	99	-	-	-	-	-	-	-	-	-	-	-
Female	174	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	261	-	-	-	-	-	-	-	-	-	-	-
Male	98	-	-	-	-	-	-	-	-	-	-	-
Female	163	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Kentucky												
All races, both sexes	288	2	-	-	1	-	-	-	-	1	-	1
Male	111	2	-	-	1	-	-	-	-	1	-	1
Female	177	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	263	1	-	-	1	-	-	-	-	1	-	-
Male	102	1	-	-	1	-	-	-	-	1	-	-
Female	161	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	25	1	-	-	-	-	-	-	-	-	-	1
Male	9	1	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	410	1	2	1	-	1	1	2	-	-	2	2
Male	134	1	1	-	-	-	-	2	-	-	-	-
Female	276	-	1	1	-	1	1	-	-	-	2	2
White, both sexes	275	-	2	-	-	-	1	1	-	-	1	-
Male	91	-	1	-	-	-	-	1	-	-	-	-
Female	184	-	1	-	-	-	1	-	-	-	1	-
Black, both sexes	133	1	-	1	-	1	-	1	-	-	1	2
Male	42	1	-	-	-	-	-	1	-	-	-	-
Female	91	-	-	1	-	1	-	-	-	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Maine												
All races, both sexes	117	-	-	-	-	-	-	-	-	-	-	-
Male	53	-	-	-	-	-	-	-	-	-	-	-
Female	64	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	117	-	-	-	-	-	-	-	-	-	-	-
Male	53	-	-	-	-	-	-	-	-	-	-	-
Female	64	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	431	1	1	-	-	-	1	-	-	-	2	1
Male	152	1	-	-	-	-	-	-	-	-	1	-
Female	279	-	1	-	-	-	1	-	-	-	1	1
White, both sexes	325	1	-	-	-	-	1	-	-	-	-	-
Male	114	1	-	-	-	-	-	-	-	-	-	-
Female	211	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	101	-	1	-	-	-	-	-	-	-	2	1
Male	34	-	-	-	-	-	-	-	-	-	1	-
Female	67	-	1	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Massachusetts												
All races, both sexes	624	1	1	1	1	-	-	-	1	2	1	1
Male	236	1	1	-	1	-	-	-	1	2	-	1
Female	388	-	-	1	-	-	-	-	-	-	1	-
White, both sexes	604	1	1	1	-	-	-	-	1	2	1	1
Male	226	1	1	-	-	-	-	-	1	2	-	1
Female	378	-	-	1	-	-	-	-	-	-	1	-
Black, both sexes	17	-	-	-	1	-	-	-	-	-	-	-
Male	8	-	-	-	1	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,243	1	1	1	1	2	-	1	1	3	2	6
Male	432	1	1	1	-	1	-	-	1	2	1	3
Female	811	-	-	-	1	1	-	1	-	1	1	3
White, both sexes	1,065	1	1	1	1	-	-	1	1	3	1	4
Male	359	1	1	1	-	-	-	-	1	2	1	2
Female	706	-	-	-	1	-	-	1	-	1	-	2
Black, both sexes	166	-	-	-	-	2	-	-	-	-	1	2
Male	69	-	-	-	-	1	-	-	-	-	-	1
Female	97	-	-	-	-	1	-	-	-	-	1	1
American Indian 1/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Minnesota												
All races, both sexes	504	1	-	-	2	-	-	-	-	2	-	3
Male	193	-	-	-	2	-	-	-	-	1	-	3
Female	311	1	-	-	-	-	-	-	-	1	-	-
White, both sexes	482	1	-	-	1	-	-	-	-	2	-	2
Male	181	-	-	-	1	-	-	-	-	1	-	2
Female	301	1	-	-	-	-	-	-	-	1	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	1	-	-	-	-	-	-	-
Male	3	-	-	-	1	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	172	3	-	-	-	-	1	-	-	3	1	5
Male	82	3	-	-	-	-	-	-	-	1	-	3
Female	90	-	-	-	-	-	1	-	-	2	1	2
White, both sexes	117	1	-	-	-	-	-	-	-	-	1	2
Male	53	1	-	-	-	-	-	-	-	-	-	1
Female	64	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	55	2	-	-	-	-	1	-	-	3	-	3
Male	29	2	-	-	-	-	-	-	-	1	-	2
Female	26	-	-	-	-	-	1	-	-	2	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Missouri												
All races, both sexes	646	2	-	-	-	-	-	1	1	-	-	5
Male	234	-	-	-	-	-	-	1	1	-	-	-
Female	412	2	-	-	-	-	-	-	-	-	-	5
White, both sexes	579	1	-	-	-	-	-	-	1	-	-	4
Male	209	-	-	-	-	-	-	-	1	-	-	-
Female	370	1	-	-	-	-	-	-	-	-	-	4
Black, both sexes	66	1	-	-	-	-	-	1	-	-	-	1
Male	25	-	-	-	-	-	-	1	-	-	-	-
Female	41	1	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	125	-	-	-	-	-	-	-	-	-	-	1
Male	50	-	-	-	-	-	-	-	-	-	-	1
Female	75	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	121	-	-	-	-	-	-	-	-	-	-	1
Male	49	-	-	-	-	-	-	-	-	-	-	1
Female	72	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Nebraska												
All races, both sexes	194	-	-	-	-	-	-	-	-	-	-	2
Male	77	-	-	-	-	-	-	-	-	-	-	1
Female	117	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	187	-	-	-	-	-	-	-	-	-	-	2
Male	72	-	-	-	-	-	-	-	-	-	-	1
Female	115	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	102	-	2	-	-	1	-	-	-	1	1	3
Male	45	-	1	-	-	1	-	-	-	-	-	1
Female	57	-	1	-	-	-	-	-	-	1	1	2
White, both sexes	94	-	2	-	-	-	-	-	-	1	1	3
Male	43	-	1	-	-	-	-	-	-	-	-	1
Female	51	-	1	-	-	-	-	-	-	1	1	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	1	-	-	-	-	-	-
Male	2	-	-	-	-	1	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
New Hampshire												
All races, both sexes	116	-	-	-	-	-	-	-	-	-	-	-
Male	34	-	-	-	-	-	-	-	-	-	-	-
Female	82	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	115	-	-	-	-	-	-	-	-	-	-	-
Male	34	-	-	-	-	-	-	-	-	-	-	-
Female	81	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	454	1	3	-	-	-	1	-	-	-	1	3
Male	175	1	3	-	-	-	-	-	-	-	1	1
Female	279	-	-	-	-	-	1	-	-	-	-	2
White, both sexes	393	1	2	-	-	-	1	-	-	-	-	3
Male	149	1	2	-	-	-	-	-	-	-	-	1
Female	244	-	-	-	-	-	1	-	-	-	-	2
Black, both sexes	52	-	1	-	-	-	-	-	-	-	1	-
Male	22	-	1	-	-	-	-	-	-	-	1	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
New Mexico												
All races, both sexes	172	-	-	-	-	-	-	-	1	-	-	1
Male	77	-	-	-	-	-	-	-	-	-	-	-
Female	95	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	165	-	-	-	-	-	-	-	1	-	-	1
Male	75	-	-	-	-	-	-	-	-	-	-	-
Female	90	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	943	2	-	1	2	1	3	2	-	4	6	7
Male	341	2	-	1	1	1	2	1	-	1	2	5
Female	602	-	-	-	1	-	1	1	-	3	4	2
White, both sexes	821	-	-	-	2	1	1	-	-	3	3	4
Male	294	-	-	-	1	1	1	-	-	-	1	2
Female	527	-	-	-	1	-	-	-	-	3	2	2
Black, both sexes	110	2	-	-	-	-	2	2	-	1	3	3
Male	44	2	-	-	-	-	1	1	-	1	1	3
Female	66	-	-	-	-	-	1	1	-	-	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
North Carolina												
All races, both sexes	882	1	2	-	-	-	-	-	-	3	4	5
Male	318	-	1	-	-	-	-	-	-	2	3	3
Female	564	1	1	-	-	-	-	-	-	1	1	2
White, both sexes	660	-	1	-	-	-	-	-	-	1	2	2
Male	234	-	1	-	-	-	-	-	-	1	1	1
Female	426	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	210	1	1	-	-	-	-	-	-	2	2	3
Male	80	-	-	-	-	-	-	-	-	1	2	2
Female	130	1	1	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	63	-	-	-	-	-	-	-	-	-	-	-
Male	28	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	61	-	-	-	-	-	-	-	-	-	-	-
Male	27	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Ohio												
All races, both sexes	1,540	4	-	1	-	-	-	1	2	-	2	7
Male	578	3	-	-	-	-	-	-	1	-	1	5
Female	962	1	-	1	-	-	-	1	1	-	1	2
White, both sexes	1,368	3	-	-	-	-	-	1	1	-	1	5
Male	496	2	-	-	-	-	-	-	1	-	-	5
Female	872	1	-	-	-	-	-	1	-	-	1	-
Black, both sexes	164	1	-	-	-	-	-	-	1	-	1	2
Male	78	1	-	-	-	-	-	-	-	-	1	-
Female	86	-	-	-	-	-	-	-	1	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	1	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	1	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	356	1	1	-	-	-	-	-	-	-	-	1
Male	116	1	-	-	-	-	-	-	-	-	-	-
Female	240	-	1	-	-	-	-	-	-	-	-	1
White, both sexes	325	1	1	-	-	-	-	-	-	-	-	1
Male	106	1	-	-	-	-	-	-	-	-	-	-
Female	219	-	1	-	-	-	-	-	-	-	-	1
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Oregon												
All races, both sexes	510	1	-	-	-	-	-	1	-	-	1	3
Male	191	-	-	-	-	-	-	-	-	-	1	2
Female	319	1	-	-	-	-	-	1	-	-	-	1
White, both sexes	491	1	-	-	-	-	-	1	-	-	1	2
Male	184	-	-	-	-	-	-	-	-	-	1	2
Female	307	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	1
Pennsylvania												
All races, both sexes	1,306	2	1	1	-	1	1	1	-	1	2	5
Male	468	2	1	1	-	-	1	1	-	1	1	2
Female	838	-	-	-	-	1	-	-	-	-	1	3
White, both sexes	1,173	1	1	-	-	-	1	1	-	1	2	4
Male	422	1	1	-	-	-	1	1	-	1	1	2
Female	751	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	126	1	-	1	-	1	-	-	-	-	-	1
Male	43	1	-	1	-	-	-	-	-	-	-	-
Female	83	-	-	-	-	1	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Rhode Island												
All races, both sexes	77	-	-	-	-	-	-	-	-	-	1	1
Male	27	-	-	-	-	-	-	-	-	-	-	1
Female	50	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	74	-	-	-	-	-	-	-	-	-	1	1
Male	25	-	-	-	-	-	-	-	-	-	-	1
Female	49	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	391	1	1	-	-	-	-	1	3	-	-	4
Male	154	1	-	-	-	-	-	1	1	-	-	1
Female	237	-	1	-	-	-	-	-	2	-	-	3
White, both sexes	272	-	1	-	-	-	-	1	3	-	-	1
Male	106	-	-	-	-	-	-	1	1	-	-	-
Female	166	-	1	-	-	-	-	-	2	-	-	1
Black, both sexes	119	1	-	-	-	-	-	-	-	-	-	3
Male	48	1	-	-	-	-	-	-	-	-	-	1
Female	71	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
South Dakota												
All races, both sexes	102	-	-	-	-	-	-	-	1	-	-	-
Male	33	-	-	-	-	-	-	-	1	-	-	-
Female	69	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	101	-	-	-	-	-	-	-	1	-	-	-
Male	33	-	-	-	-	-	-	-	1	-	-	-
Female	68	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	629	3	2	-	-	-	1	2	-	-	4	8
Male	207	2	1	-	-	-	-	1	-	-	2	3
Female	422	1	1	-	-	-	1	1	-	-	2	5
White, both sexes	549	2	2	-	-	-	1	1	-	-	2	6
Male	177	1	1	-	-	-	-	1	-	-	2	3
Female	372	1	1	-	-	-	1	-	-	-	-	3
Black, both sexes	79	1	-	-	-	-	-	1	-	-	2	2
Male	29	1	-	-	-	-	-	-	-	-	-	-
Female	50	-	-	-	-	-	-	1	-	-	2	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Texas												
All races, both sexes	2,049	3	1	1	-	-	2	2	2	3	6	21
Male	752	1	-	1	-	-	2	1	1	2	1	14
Female	1,297	2	1	-	-	-	-	1	1	1	5	7
White, both sexes	1,731	2	1	1	-	-	1	2	2	2	2	12
Male	618	1	-	1	-	-	1	1	1	2	1	8
Female	1,113	1	1	-	-	-	-	1	1	-	1	4
Black, both sexes	294	1	-	-	-	-	1	-	-	1	4	9
Male	117	-	-	-	-	-	1	-	-	-	-	6
Female	177	1	-	-	-	-	-	-	-	1	4	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	183	-	1	1	-	-	-	-	-	1	-	1
Male	75	-	1	1	-	-	-	-	-	-	-	-
Female	108	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	180	-	1	1	-	-	-	-	-	1	-	1
Male	74	-	1	1	-	-	-	-	-	-	-	-
Female	106	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Vermont												
All races, both sexes	55	1	-	-	-	-	-	-	-	-	-	-
Male	25	1	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	55	1	-	-	-	-	-	-	-	-	-	-
Male	25	1	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	579	-	-	-	-	1	-	1	1	1	2	4
Male	218	-	-	-	-	-	-	-	-	1	1	2
Female	361	-	-	-	-	1	-	1	1	-	1	2
White, both sexes	466	-	-	-	-	-	-	1	-	1	1	3
Male	177	-	-	-	-	-	-	-	-	1	1	2
Female	289	-	-	-	-	-	-	1	-	-	-	1
Black, both sexes	107	-	-	-	-	1	-	-	1	-	1	1
Male	39	-	-	-	-	-	-	-	-	-	-	-
Female	68	-	-	-	-	1	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Washington												
All races, both sexes	601	1	-	-	-	-	-	-	1	-	1	5
Male	238	-	-	-	-	-	-	-	1	-	1	3
Female	363	1	-	-	-	-	-	-	-	-	-	2
White, both sexes	566	-	-	-	-	-	-	-	1	-	1	5
Male	222	-	-	-	-	-	-	-	1	-	1	3
Female	344	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	1	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	199	-	-	-	-	-	-	-	-	-	1	2
Male	67	-	-	-	-	-	-	-	-	-	-	1
Female	132	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	191	-	-	-	-	-	-	-	-	-	1	2
Male	66	-	-	-	-	-	-	-	-	-	-	1
Female	125	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other cerebrovascular diseases and their sequelae (I67,I69)												
Wisconsin												
All races, both sexes	640	1	-	-	1	1	-	-	2	1	2	1
Male	252	1	-	-	1	1	-	-	1	-	1	1
Female	388	-	-	-	-	-	-	-	1	1	1	-
White, both sexes	615	1	-	-	1	-	-	-	2	1	2	1
Male	237	1	-	-	1	-	-	-	1	-	1	1
Female	378	-	-	-	-	-	-	-	1	1	1	-
Black, both sexes	22	-	-	-	-	1	-	-	-	-	-	-
Male	12	-	-	-	-	1	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	38	-	-	-	-	-	-	-	-	-	-	1
Male	18	-	-	-	-	-	-	-	-	-	-	1
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
United States												
All races, both sexes	284	424	677	1,041	1,863	3,336	5,725	6,364	5,035	2,008	445	1
Male	149	216	367	577	939	1,545	2,341	2,149	1,297	396	52	1
Female	135	208	310	464	924	1,791	3,384	4,215	3,738	1,612	393	-
White, both sexes	187	302	488	760	1,471	2,755	5,020	5,757	4,590	1,841	382	-
Male	99	156	258	418	750	1,298	2,060	1,958	1,174	358	44	-
Female	88	146	230	342	721	1,457	2,960	3,799	3,416	1,483	338	-
Black, both sexes	84	106	160	237	332	480	587	473	356	125	55	1
Male	44	54	97	131	158	198	227	143	87	22	7	1
Female	40	52	63	106	174	282	360	330	269	103	48	-
American Indian 1/, both sexes	3	4	8	9	12	15	15	17	8	2	1	-
Male	2	1	2	5	9	7	4	6	1	1	-	-
Female	1	3	6	4	3	8	11	11	7	1	1	-
Asian or Pacific Islander 2/, both sexes	10	12	21	35	48	86	103	117	81	40	7	-
Male	4	5	10	23	22	42	50	42	35	15	1	-
Female	6	7	11	12	26	44	53	75	46	25	6	-
Alabama												
All races, both sexes	9	12	18	26	40	60	75	71	63	24	5	-
Male	5	3	8	16	20	30	26	20	11	4	-	-
Female	4	9	10	10	20	30	49	51	52	20	5	-
White, both sexes	5	8	11	14	29	44	53	60	55	15	3	-
Male	3	2	5	8	13	24	20	19	8	3	-	-
Female	2	6	6	6	16	20	33	41	47	12	3	-
Black, both sexes	4	4	6	12	11	16	22	11	8	9	2	-
Male	2	1	2	8	7	6	6	1	3	1	-	-
Female	2	3	4	4	4	10	16	10	5	8	2	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Alaska												
All races, both sexes	-	-	2	2	2	6	7	8	2	1	2	-
Male	-	-	-	1	2	1	3	2	-	1	1	-
Female	-	-	2	1	-	5	4	6	2	-	1	-
White, both sexes	-	-	1	1	1	5	5	4	2	1	2	-
Male	-	-	-	-	1	1	3	1	-	1	1	-
Female	-	-	1	1	-	4	2	3	2	-	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	1	1	2	3	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	1	2	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	3	8	3	16	24	53	101	81	72	26	2	-
Male	2	5	2	4	10	22	47	28	24	7	-	-
Female	1	3	1	12	14	31	54	53	48	19	2	-
White, both sexes	2	6	3	15	24	51	100	79	67	26	2	-
Male	2	4	2	3	10	21	46	26	22	7	-	-
Female	-	2	1	12	14	30	54	53	45	19	2	-
Black, both sexes	1	2	-	-	-	-	-	-	4	-	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	1	1	-	-	-	-	-	-	3	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	2	1	-	-	-
Male	-	-	-	1	-	-	-	2	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	1	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Arkansas												
All races, both sexes	5	3	15	17	27	42	69	68	45	21	5	-
Male	2	2	7	9	16	19	23	27	6	4	2	-
Female	3	1	8	8	11	23	46	41	39	17	3	-
White, both sexes	3	3	11	12	16	34	53	58	35	18	3	-
Male	2	2	5	6	10	18	20	24	5	3	-	-
Female	1	1	6	6	6	16	33	34	30	15	3	-
Black, both sexes	2	-	4	4	11	7	15	10	10	3	2	-
Male	-	-	2	3	6	1	3	3	1	1	2	-
Female	2	-	2	1	5	6	12	7	9	2	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
California												
All races, both sexes	44	65	93	163	253	446	813	951	715	309	65	-
Male	20	28	48	87	114	205	338	327	204	66	5	-
Female	24	37	45	76	139	241	475	624	511	243	60	-
White, both sexes	25	38	65	111	191	353	685	834	625	276	51	-
Male	13	16	33	58	87	160	279	289	171	57	4	-
Female	12	22	32	53	104	193	406	545	454	219	47	-
Black, both sexes	12	19	16	33	30	42	64	46	34	10	8	-
Male	6	10	8	17	13	21	28	13	9	2	1	-
Female	6	9	8	16	17	21	36	33	25	8	7	-
American Indian 1/, both sexes	1	1	-	1	3	1	1	2	2	1	-	-
Male	-	-	-	-	2	-	-	2	-	-	-	-
Female	1	1	-	1	1	1	1	-	2	1	-	-
Asian or Pacific Islander 2/, both sexes	6	7	12	18	29	50	63	69	54	22	6	-
Male	1	2	7	12	12	24	31	23	24	7	-	-
Female	5	5	5	6	17	26	32	46	30	15	6	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Colorado												
All races, both sexes	4	10	6	9	20	32	66	92	70	28	3	-
Male	1	7	2	5	10	19	25	29	15	6	1	-
Female	3	3	4	4	10	13	41	63	55	22	2	-
White, both sexes	4	9	6	8	19	29	62	91	67	28	3	-
Male	1	7	2	4	9	17	24	29	15	6	1	-
Female	3	2	4	4	10	12	38	62	52	22	2	-
Black, both sexes	-	1	-	1	1	2	3	1	3	-	-	-
Male	-	-	-	1	1	1	1	-	-	-	-	-
Female	-	1	-	-	-	1	2	1	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Connecticut												
All races, both sexes	-	5	4	6	17	21	62	49	68	28	2	-
Male	-	2	-	4	9	15	24	15	21	5	-	-
Female	-	3	4	2	8	6	38	34	47	23	2	-
White, both sexes	-	4	4	4	15	18	57	46	67	27	2	-
Male	-	1	-	3	8	13	22	14	21	5	-	-
Female	-	3	4	1	7	5	35	32	46	22	2	-
Black, both sexes	-	1	-	2	2	3	4	3	1	1	-	-
Male	-	1	-	1	1	2	2	1	-	-	-	-
Female	-	-	-	1	1	1	2	2	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Delaware												
All races, both sexes	-	1	2	1	5	11	18	7	13	5	2	-
Male	-	-	-	1	1	5	6	1	6	-	-	-
Female	-	1	2	-	4	6	12	6	7	5	2	-
White, both sexes	-	-	1	1	3	11	14	5	12	5	1	-
Male	-	-	-	1	1	5	5	1	5	-	-	-
Female	-	-	1	-	2	6	9	4	7	5	1	-
Black, both sexes	-	1	1	-	1	-	4	2	1	-	1	-
Male	-	-	-	-	-	-	1	-	1	-	-	-
Female	-	1	1	-	1	-	3	2	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	-	3	3	1	1	8	6	2	-	1	-
Male	1	-	2	2	-	-	2	2	1	-	-	-
Female	-	-	1	1	1	1	6	4	1	-	1	-
White, both sexes	-	-	1	-	-	1	4	3	-	-	-	-
Male	-	-	1	-	-	-	2	1	-	-	-	-
Female	-	-	-	-	-	1	2	2	-	-	-	-
Black, both sexes	1	-	2	3	1	-	4	3	2	-	1	-
Male	1	-	1	2	-	-	-	1	1	-	-	-
Female	-	-	1	1	1	-	4	2	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Florida												
All races, both sexes	10	25	56	69	117	227	422	421	345	153	34	1
Male	6	10	33	47	58	91	182	148	103	44	7	1
Female	4	15	23	22	59	136	240	273	242	109	27	-
White, both sexes	7	19	42	44	90	189	383	385	318	144	30	-
Male	4	9	23	31	45	74	165	138	95	42	6	-
Female	3	10	19	13	45	115	218	247	223	102	24	-
Black, both sexes	3	5	13	23	25	36	38	33	27	8	4	1
Male	2	1	9	14	12	16	16	10	8	2	1	1
Female	1	4	4	9	13	20	22	23	19	6	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	2	2	2	1	3	-	1	-	-
Male	-	-	1	2	1	1	1	-	-	-	-	-
Female	-	1	-	-	1	1	-	3	-	1	-	-
Georgia												
All races, both sexes	6	10	24	40	56	70	119	110	100	28	11	-
Male	3	5	17	23	24	26	37	33	23	5	1	-
Female	3	5	7	17	32	44	82	77	77	23	10	-
White, both sexes	1	5	12	25	32	49	86	85	79	21	6	-
Male	1	3	9	16	14	19	28	25	19	2	1	-
Female	-	2	3	9	18	30	58	60	60	19	5	-
Black, both sexes	4	5	12	15	24	19	33	25	21	7	5	-
Male	2	2	8	7	10	6	9	8	4	3	-	-
Female	2	3	4	8	14	13	24	17	17	4	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Hawaii												
All races, both sexes	-	-	4	2	4	9	10	21	16	15	-	-
Male	-	-	-	1	3	5	4	7	6	7	-	-
Female	-	-	4	1	1	4	6	14	10	8	-	-
White, both sexes	-	-	2	1	1	1	3	3	2	3	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	2	1	1	1	1	3	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	3	8	7	18	14	12	-	-
Male	-	-	-	1	3	5	2	7	6	7	-	-
Female	-	-	2	-	-	3	5	11	8	5	-	-
Idaho												
All races, both sexes	-	5	4	9	8	19	36	39	37	10	-	-
Male	-	4	2	6	4	7	12	15	8	-	-	-
Female	-	1	2	3	4	12	24	24	29	10	-	-
White, both sexes	-	5	4	9	8	19	36	39	37	10	-	-
Male	-	4	2	6	4	7	12	15	8	-	-	-
Female	-	1	2	3	4	12	24	24	29	10	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Illinois												
All races, both sexes	6	20	29	42	77	157	259	271	220	86	15	-
Male	6	10	16	23	35	76	119	85	60	16	-	-
Female	-	10	13	19	42	81	140	186	160	70	15	-
White, both sexes	1	12	17	31	51	121	231	242	197	76	10	-
Male	1	8	10	17	23	62	106	75	49	13	-	-
Female	-	4	7	14	28	59	125	167	148	63	10	-
Black, both sexes	4	8	12	10	26	35	27	25	22	8	5	-
Male	4	2	6	5	12	14	13	8	10	2	-	-
Female	-	6	6	5	14	21	14	17	12	6	5	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	1	1	4	1	2	-	-
Male	1	-	-	1	-	-	-	2	1	1	-	-
Female	-	-	-	-	-	1	1	2	-	1	-	-
Indiana												
All races, both sexes	9	7	11	19	47	75	107	124	84	32	8	-
Male	5	7	6	12	25	35	48	39	23	3	-	-
Female	4	-	5	7	22	40	59	85	61	29	8	-
White, both sexes	4	5	9	14	42	66	100	118	80	31	6	-
Male	2	5	5	9	21	31	45	36	22	3	-	-
Female	2	-	4	5	21	35	55	82	58	28	6	-
Black, both sexes	5	2	2	5	5	9	7	6	4	1	2	-
Male	3	2	1	3	4	4	3	3	1	-	-	-
Female	2	-	1	2	1	5	4	3	3	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Iowa												
All races, both sexes	1	3	6	5	13	33	69	90	72	30	6	-
Male	-	2	5	4	9	14	29	38	20	4	-	-
Female	1	1	1	1	4	19	40	52	52	26	6	-
White, both sexes	1	3	6	5	13	32	69	87	71	30	6	-
Male	-	2	5	4	9	13	29	37	20	4	-	-
Female	1	1	1	1	4	19	40	50	51	26	6	-
Black, both sexes	-	-	-	-	-	1	-	3	1	-	-	-
Male	-	-	-	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	5	3	7	13	26	44	77	54	34	9	-
Male	1	2	2	-	6	17	21	26	19	4	1	-
Female	-	3	1	7	7	9	23	51	35	30	8	-
White, both sexes	1	5	2	6	13	24	42	74	51	34	9	-
Male	1	2	2	-	6	17	21	26	18	4	1	-
Female	-	3	-	6	7	7	21	48	33	30	8	-
Black, both sexes	-	-	-	1	-	2	2	2	3	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	1	-	2	2	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Kentucky												
All races, both sexes	3	3	5	13	34	41	55	65	53	7	4	-
Male	1	1	3	5	26	19	20	20	11	-	-	-
Female	2	2	2	8	8	22	35	45	42	7	4	-
White, both sexes	3	2	5	10	33	37	51	60	49	6	4	-
Male	1	1	3	5	25	16	19	19	10	-	-	-
Female	2	1	2	5	8	21	32	41	39	6	4	-
Black, both sexes	-	1	-	3	1	4	4	5	4	1	-	-
Male	-	-	-	-	1	3	1	1	1	-	-	-
Female	-	1	-	3	-	1	3	4	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	7	10	14	19	37	54	94	81	54	22	6	-
Male	1	5	10	6	16	23	31	26	11	1	-	-
Female	6	5	4	13	21	31	63	55	43	21	6	-
White, both sexes	4	7	6	8	18	36	64	62	44	17	4	-
Male	1	4	3	1	7	17	23	22	10	1	-	-
Female	3	3	3	7	11	19	41	40	34	16	4	-
Black, both sexes	3	3	8	10	19	18	29	19	10	5	2	-
Male	-	1	7	4	9	6	8	4	1	-	-	-
Female	3	2	1	6	10	12	21	15	9	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Maine												
All races, both sexes	1	2	4	6	12	16	24	24	23	5	-	-
Male	1	1	-	4	9	10	12	8	8	-	-	-
Female	-	1	4	2	3	6	12	16	15	5	-	-
White, both sexes	1	2	4	6	12	16	24	24	23	5	-	-
Male	1	1	-	4	9	10	12	8	8	-	-	-
Female	-	1	4	2	3	6	12	16	15	5	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5	5	9	14	26	60	84	99	90	29	4	-
Male	3	1	4	7	15	24	32	40	19	4	1	-
Female	2	4	5	7	11	36	52	59	71	25	3	-
White, both sexes	2	2	4	9	19	44	62	78	73	26	4	-
Male	1	-	2	5	12	20	26	29	14	3	1	-
Female	1	2	2	4	7	24	36	49	59	23	3	-
Black, both sexes	3	2	5	5	7	16	21	19	16	3	-	-
Male	2	1	2	2	3	4	5	9	4	1	-	-
Female	1	1	3	3	4	12	16	10	12	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	1	2	1	-	-	-
Male	-	-	-	-	-	-	1	2	1	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Massachusetts												
All races, both sexes	6	7	12	20	29	62	120	159	127	62	11	-
Male	4	2	9	12	14	25	49	60	40	11	3	-
Female	2	5	3	8	15	37	71	99	87	51	8	-
White, both sexes	5	6	12	18	26	60	117	153	126	62	11	-
Male	3	2	9	11	13	24	48	57	39	11	3	-
Female	2	4	3	7	13	36	69	96	87	51	8	-
Black, both sexes	1	1	-	2	3	-	2	6	1	-	-	-
Male	1	-	-	1	1	-	-	3	1	-	-	-
Female	-	1	-	1	2	-	2	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Michigan												
All races, both sexes	9	18	34	49	87	155	241	304	220	91	16	-
Male	7	9	17	22	45	74	84	101	50	11	1	-
Female	2	9	17	27	42	81	157	203	170	80	15	-
White, both sexes	4	10	22	32	65	128	211	278	202	84	15	-
Male	3	7	11	14	30	65	71	92	47	9	1	-
Female	1	3	11	18	35	63	140	186	155	75	14	-
Black, both sexes	5	7	11	17	20	25	28	23	17	7	1	-
Male	4	2	6	8	13	8	13	8	3	2	-	-
Female	1	5	5	9	7	17	15	15	14	5	1	-
American Indian 1/, both sexes	-	1	1	-	1	2	2	1	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	1	1	-	-	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	2	1	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Minnesota												
All races, both sexes	7	1	11	11	24	45	87	136	115	51	8	-
Male	3	1	7	7	16	23	37	48	32	13	-	-
Female	4	-	4	4	8	22	50	88	83	38	8	-
White, both sexes	6	-	10	10	19	43	84	131	114	51	8	-
Male	2	-	6	6	13	22	37	46	32	13	-	-
Female	4	-	4	4	6	21	47	85	82	38	8	-
Black, both sexes	1	-	1	-	2	-	2	4	-	-	-	-
Male	1	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	2	3	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	2	1	-	-	-	-	-	-
Male	-	-	-	1	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	1	1	1	1	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	1	1	-	1	-	-	-
Mississippi												
All races, both sexes	6	4	7	14	12	16	30	30	26	11	3	-
Male	4	3	6	11	6	8	13	11	7	5	1	-
Female	2	1	1	3	6	8	17	19	19	6	2	-
White, both sexes	5	4	1	10	6	11	26	26	16	6	2	-
Male	4	3	1	8	4	6	10	9	4	1	1	-
Female	1	1	-	2	2	5	16	17	12	5	1	-
Black, both sexes	1	-	6	4	6	5	4	4	10	5	1	-
Male	-	-	5	3	2	2	3	2	3	4	-	-
Female	1	-	1	1	4	3	1	2	7	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Missouri												
All races, both sexes	7	2	9	23	48	83	131	155	124	43	12	-
Male	4	1	3	16	28	38	60	47	22	10	3	-
Female	3	1	6	7	20	45	71	108	102	33	9	-
White, both sexes	5	-	8	19	41	68	121	148	113	40	10	-
Male	2	-	2	13	25	32	57	45	21	9	2	-
Female	3	-	6	6	16	36	64	103	92	31	8	-
Black, both sexes	2	2	1	4	7	15	10	7	10	3	2	-
Male	2	1	1	3	3	6	3	2	1	1	1	-
Female	-	1	-	1	4	9	7	5	9	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	2	2	6	11	18	24	24	29	7	1	-
Male	-	1	1	3	4	11	8	10	8	3	-	-
Female	-	1	1	3	7	7	16	14	21	4	1	-
White, both sexes	-	2	2	5	10	18	23	24	28	7	1	-
Male	-	1	1	2	4	11	8	10	8	3	-	-
Female	-	1	1	3	6	7	15	14	20	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	1	-	1	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Nebraska												
All races, both sexes	3	1	2	8	14	26	33	53	36	11	5	-
Male	1	-	2	5	8	13	17	18	9	2	1	-
Female	2	1	-	3	6	13	16	35	27	9	4	-
White, both sexes	3	1	2	7	11	25	32	53	35	11	5	-
Male	1	-	2	4	6	13	16	18	8	2	1	-
Female	2	1	-	3	5	12	16	35	27	9	4	-
Black, both sexes	-	-	-	-	2	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Nevada												
All races, both sexes	-	2	5	2	8	16	21	22	11	6	1	-
Male	-	1	3	1	6	10	9	10	2	-	-	-
Female	-	1	2	1	2	6	12	12	9	6	1	-
White, both sexes	-	2	5	2	7	15	18	22	10	5	1	-
Male	-	1	3	1	6	10	8	10	2	-	-	-
Female	-	1	2	1	1	5	10	12	8	5	1	-
Black, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	1	-	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
New Hampshire												
All races, both sexes	-	2	2	2	6	14	28	26	27	5	4	-
Male	-	1	2	1	4	7	10	7	1	1	-	-
Female	-	1	-	1	2	7	18	19	26	4	4	-
White, both sexes	-	2	2	2	6	14	28	26	26	5	4	-
Male	-	1	2	1	4	7	10	7	1	1	-	-
Female	-	1	-	1	2	7	18	19	25	4	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
New Jersey												
All races, both sexes	8	6	13	15	25	64	85	96	86	31	16	-
Male	5	4	9	6	11	37	25	36	27	6	3	-
Female	3	2	4	9	14	27	60	60	59	25	13	-
White, both sexes	4	6	10	12	21	54	72	82	82	29	14	-
Male	2	4	6	5	9	31	20	33	26	6	3	-
Female	2	2	4	7	12	23	52	49	56	23	11	-
Black, both sexes	3	-	2	1	4	10	11	11	4	2	2	-
Male	2	-	2	1	2	6	3	3	1	-	-	-
Female	1	-	-	-	2	4	8	8	3	2	2	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	-	2	3	-	-	-	-
Male	1	-	1	-	-	-	2	-	-	-	-	-
Female	-	-	-	1	-	-	-	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
New Mexico												
All races, both sexes	-	2	6	7	11	20	41	41	29	8	5	-
Male	-	1	3	4	8	10	20	19	11	1	-	-
Female	-	1	3	3	3	10	21	22	18	7	5	-
White, both sexes	-	2	5	6	11	18	40	40	29	8	4	-
Male	-	1	3	4	8	9	19	19	11	1	-	-
Female	-	1	2	2	3	9	21	21	18	7	4	-
Black, both sexes	-	-	-	1	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	1	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	11	12	22	25	60	108	190	217	173	79	18	-
Male	5	8	14	13	31	53	80	65	40	16	-	-
Female	6	4	8	12	29	55	110	152	133	63	18	-
White, both sexes	7	7	16	21	48	90	169	192	162	77	18	-
Male	4	3	13	10	30	46	73	55	38	16	-	-
Female	3	4	3	11	18	44	96	137	124	61	18	-
Black, both sexes	3	4	5	3	12	16	19	22	11	2	-	-
Male	-	4	1	3	1	7	7	10	2	-	-	-
Female	3	-	4	-	11	9	12	12	9	2	-	-
American Indian 1/, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	-	2	2	3	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	2	2	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
North Carolina												
All races, both sexes	8	18	27	32	67	117	178	211	149	48	12	-
Male	6	9	17	22	35	50	62	64	39	4	1	-
Female	2	9	10	10	32	67	116	147	110	44	11	-
White, both sexes	4	14	14	19	47	84	131	163	129	41	8	-
Male	4	6	8	13	28	36	45	50	35	4	1	-
Female	-	8	6	6	19	48	86	113	94	37	7	-
Black, both sexes	4	4	11	13	18	30	45	45	20	7	4	-
Male	2	3	9	9	7	12	17	12	4	-	-	-
Female	2	1	2	4	11	18	28	33	16	7	4	-
American Indian 1/, both sexes	-	-	2	-	-	2	2	3	-	-	-	-
Male	-	-	-	-	-	1	-	2	-	-	-	-
Female	-	-	2	-	-	1	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	1	-	1	2	7	14	13	17	7	1	-
Male	-	1	-	1	1	5	8	4	5	3	-	-
Female	-	-	-	-	1	2	6	9	12	4	1	-
White, both sexes	-	1	-	1	2	7	14	12	17	6	1	-
Male	-	1	-	1	1	5	8	4	5	2	-	-
Female	-	-	-	-	1	2	6	8	12	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Ohio												
All races, both sexes	20	32	44	73	109	205	333	334	253	98	22	-
Male	12	21	25	43	51	91	137	115	53	19	1	-
Female	8	11	19	30	58	114	196	219	200	79	21	-
White, both sexes	15	27	39	58	91	178	299	308	231	92	19	-
Male	8	17	21	33	41	77	116	109	46	19	1	-
Female	7	10	18	25	50	101	183	199	185	73	18	-
Black, both sexes	4	5	5	14	18	26	33	24	21	6	3	-
Male	3	4	4	9	10	13	20	6	7	-	-	-
Female	1	1	1	5	8	13	13	18	14	6	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	1	1	2	-	-	-	-
Male	1	-	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Oklahoma												
All races, both sexes	6	6	7	11	18	55	72	81	64	29	4	-
Male	3	3	5	3	8	27	29	19	13	4	1	-
Female	3	3	2	8	10	28	43	62	51	25	3	-
White, both sexes	5	4	7	8	17	46	66	77	59	29	4	-
Male	2	2	5	2	8	22	27	19	13	4	1	-
Female	3	2	2	6	9	24	39	58	46	25	3	-
Black, both sexes	-	2	-	1	-	4	2	3	2	-	-	-
Male	-	1	-	-	-	3	1	-	-	-	-	-
Female	-	1	-	1	-	1	1	3	2	-	-	-
American Indian 1/, both sexes	1	-	-	1	1	5	4	1	2	-	-	-
Male	1	-	-	1	-	2	1	-	-	-	-	-
Female	-	-	-	-	1	3	3	1	2	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Oregon												
All races, both sexes	6	6	11	17	25	54	113	119	115	32	6	-
Male	2	1	7	12	9	22	60	40	25	8	2	-
Female	4	5	4	5	16	32	53	79	90	24	4	-
White, both sexes	6	5	10	15	23	51	109	115	114	32	6	-
Male	2	1	7	11	8	21	57	39	25	8	2	-
Female	4	4	3	4	15	30	52	76	89	24	4	-
Black, both sexes	-	-	-	-	-	1	4	1	-	-	-	-
Male	-	-	-	-	-	-	3	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	1	1	-	2	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	1	1	1	-	1	1	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	1	-	1	1	-	-	1	-	-	-
Pennsylvania												
All races, both sexes	13	15	21	34	80	140	278	316	265	102	27	-
Male	4	9	7	17	43	60	113	112	67	24	2	-
Female	9	6	14	17	37	80	165	204	198	78	25	-
White, both sexes	9	12	16	27	69	114	253	300	244	95	23	-
Male	3	6	3	14	37	51	104	108	63	23	2	-
Female	6	6	13	13	32	63	149	192	181	72	21	-
Black, both sexes	4	3	5	7	10	26	22	15	20	6	4	-
Male	1	3	4	3	5	9	8	4	3	1	-	-
Female	3	-	1	4	5	17	14	11	17	5	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	3	-	1	1	-	-
Male	-	-	-	-	1	-	1	-	1	-	-	-
Female	-	-	-	-	-	-	2	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Rhode Island												
All races, both sexes	1	2	2	2	2	8	18	19	15	6	-	-
Male	1	-	1	-	1	4	8	6	4	1	-	-
Female	-	2	1	2	1	4	10	13	11	5	-	-
White, both sexes	1	2	2	2	2	7	18	18	14	6	-	-
Male	1	-	1	-	1	3	8	6	3	1	-	-
Female	-	2	1	2	1	4	10	12	11	5	-	-
Black, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	5	8	10	21	28	55	99	77	50	23	5	-
Male	3	4	3	15	16	25	39	30	11	3	1	-
Female	2	4	7	6	12	30	60	47	39	20	4	-
White, both sexes	3	4	4	7	19	37	71	60	39	18	4	-
Male	2	2	1	4	10	18	28	26	9	3	1	-
Female	1	2	3	3	9	19	43	34	30	15	3	-
Black, both sexes	2	4	6	14	9	18	28	17	11	5	1	-
Male	1	2	2	11	6	7	11	4	2	-	-	-
Female	1	2	4	3	3	11	17	13	9	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
South Dakota												
All races, both sexes	-	1	2	2	8	6	21	29	18	11	3	-
Male	-	1	1	1	6	1	6	6	7	2	1	-
Female	-	-	1	1	2	5	15	23	11	9	2	-
White, both sexes	-	1	2	2	8	5	21	29	18	11	3	-
Male	-	1	1	1	6	1	6	6	7	2	1	-
Female	-	-	1	1	2	4	15	23	11	9	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	9	6	9	33	42	75	128	142	102	50	13	-
Male	3	2	5	17	21	35	51	38	19	4	3	-
Female	6	4	4	16	21	40	77	104	83	46	10	-
White, both sexes	7	4	5	31	33	62	117	126	95	43	12	-
Male	1	1	3	16	16	30	49	31	16	4	2	-
Female	6	3	2	15	17	32	68	95	79	39	10	-
Black, both sexes	2	2	3	2	9	13	11	16	7	7	1	-
Male	2	1	1	1	5	5	2	7	3	-	1	-
Female	-	1	2	1	4	8	9	9	4	7	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Texas												
All races, both sexes	24	39	58	82	160	261	391	449	379	134	31	-
Male	16	20	32	42	70	124	156	149	88	30	2	-
Female	8	19	26	40	90	137	235	300	291	104	29	-
White, both sexes	18	24	40	59	133	207	331	404	339	124	27	-
Male	14	10	19	32	61	98	124	134	80	29	1	-
Female	4	14	21	27	72	109	207	270	259	95	26	-
Black, both sexes	6	13	17	21	27	49	53	40	38	10	4	-
Male	2	8	13	9	9	22	26	12	7	1	1	-
Female	4	5	4	12	18	27	27	28	31	9	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	1	2	-	5	5	5	2	-	-	-
Male	-	2	-	1	-	4	4	3	1	-	-	-
Female	-	-	1	1	-	1	1	2	1	-	-	-
Utah												
All races, both sexes	3	6	6	5	13	14	44	45	31	10	2	-
Male	1	3	2	3	8	7	18	17	12	2	-	-
Female	2	3	4	2	5	7	26	28	19	8	2	-
White, both sexes	3	6	5	5	13	13	43	45	31	10	2	-
Male	1	3	2	3	8	7	17	17	12	2	-	-
Female	2	3	3	2	5	6	26	28	19	8	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Vermont												
All races, both sexes	1	-	-	1	-	4	12	21	10	5	-	-
Male	-	-	-	-	-	2	8	8	4	2	-	-
Female	1	-	-	1	-	2	4	13	6	3	-	-
White, both sexes	1	-	-	1	-	4	12	21	10	5	-	-
Male	-	-	-	-	-	2	8	8	4	2	-	-
Female	1	-	-	1	-	2	4	13	6	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5	12	13	15	46	62	128	139	100	44	5	-
Male	2	7	3	7	23	33	51	53	30	5	-	-
Female	3	5	10	8	23	29	77	86	70	39	5	-
White, both sexes	3	8	9	9	29	44	103	123	91	36	5	-
Male	1	5	2	4	12	26	42	49	28	4	-	-
Female	2	3	7	5	17	18	61	74	63	32	5	-
Black, both sexes	2	4	4	4	15	18	24	15	9	8	-	-
Male	1	2	1	2	10	7	9	4	2	1	-	-
Female	1	2	3	2	5	11	15	11	7	7	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	2	-	1	1	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	1	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Washington												
All races, both sexes	6	7	11	20	48	79	124	140	105	42	11	-
Male	4	4	7	12	29	34	51	49	30	10	3	-
Female	2	3	4	8	19	45	73	91	75	32	8	-
White, both sexes	5	7	10	18	41	73	115	137	102	41	10	-
Male	4	4	6	11	23	33	48	47	29	10	2	-
Female	1	3	4	7	18	40	67	90	73	31	8	-
Black, both sexes	1	-	-	-	2	4	3	1	1	-	-	-
Male	-	-	-	-	2	-	1	-	-	-	-	-
Female	1	-	-	-	-	4	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	1	-	2	-	1	-	-	-	-	-
Male	-	-	1	-	2	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	3	2	5	2	2	1	1	-
Male	-	-	-	1	2	1	1	2	1	-	1	-
Female	-	-	-	1	1	1	4	-	1	1	-	-
West Virginia												
All races, both sexes	2	1	4	7	7	33	47	39	38	16	2	-
Male	-	-	4	6	2	13	17	13	6	4	1	-
Female	2	1	-	1	5	20	30	26	32	12	1	-
White, both sexes	2	1	4	7	6	32	44	38	37	15	2	-
Male	-	-	4	6	2	13	16	13	6	4	1	-
Female	2	1	-	1	4	19	28	25	31	11	1	-
Black, both sexes	-	-	-	-	1	1	3	1	1	1	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	1	2	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other cerebrovascular diseases and their sequelae (I67,I69)												
Wisconsin												
All races, both sexes	3	6	11	14	36	69	143	165	120	48	16	-
Male	1	4	5	8	22	36	69	54	34	10	3	-
Female	2	2	6	6	14	33	74	111	86	38	13	-
White, both sexes	2	5	9	12	33	61	139	162	120	48	16	-
Male	1	3	4	7	21	30	66	53	34	10	3	-
Female	1	2	5	5	12	31	73	109	86	38	13	-
Black, both sexes	1	1	2	2	3	7	2	3	-	-	-	-
Male	-	1	1	1	1	5	1	1	-	-	-	-
Female	1	-	1	1	2	2	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	2	-	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	1	4	6	9	7	3	5	1	-
Male	-	-	-	-	1	4	5	4	2	1	-	-
Female	-	-	1	1	3	2	4	3	1	4	1	-
White, both sexes	-	-	1	1	4	6	9	7	3	5	1	-
Male	-	-	-	-	1	4	5	4	2	1	-	-
Female	-	-	1	1	3	2	4	3	1	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
United States												
All races, both sexes	35,458	17	4	5	12	17	37	66	96	169	329	509
Male	16,895	11	4	3	10	14	24	36	69	123	219	359
Female	18,563	6	-	2	2	3	13	30	27	46	110	150
White, both sexes	31,330	11	3	4	10	14	30	40	65	105	235	360
Male	14,918	6	3	3	8	11	18	19	48	72	153	256
Female	16,412	5	-	1	2	3	12	21	17	33	82	104
Black, both sexes	3,446	4	1	1	2	3	6	22	25	58	79	128
Male	1,594	3	1	-	2	3	5	14	15	46	53	89
Female	1,852	1	-	1	-	-	1	8	10	12	26	39
American Indian 1/, both sexes	135	1	-	-	-	-	-	1	-	1	2	6
Male	61	1	-	-	-	-	-	1	-	1	1	2
Female	74	-	-	-	-	-	-	-	-	-	1	4
Asian or Pacific Islander 2/, both sexes	547	1	-	-	-	-	1	3	6	5	13	15
Male	322	1	-	-	-	-	1	2	6	4	12	12
Female	225	-	-	-	-	-	-	1	-	1	1	3
Alabama												
All races, both sexes	646	-	-	-	-	-	-	1	2	5	8	12
Male	308	-	-	-	-	-	-	1	1	3	5	9
Female	338	-	-	-	-	-	-	-	1	2	3	3
White, both sexes	513	-	-	-	-	-	-	1	2	3	6	9
Male	243	-	-	-	-	-	-	1	1	1	4	6
Female	270	-	-	-	-	-	-	-	1	2	2	3
Black, both sexes	128	-	-	-	-	-	-	-	-	2	2	3
Male	64	-	-	-	-	-	-	-	-	2	1	3
Female	64	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Alaska												
All races, both sexes	30	-	-	-	-	-	-	-	-	1	1	-
Male	18	-	-	-	-	-	-	-	-	1	1	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	1	-
Male	15	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	661	1	1	-	-	1	-	-	1	-	2	10
Male	338	1	1	-	-	1	-	-	1	-	1	8
Female	323	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	634	-	1	-	-	1	-	-	1	-	2	10
Male	324	-	1	-	-	1	-	-	1	-	1	8
Female	310	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	1	-	-	-	-	-	-	-	-	-	-
Male	4	1	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Arkansas												
All races, both sexes	390	-	1	1	-	-	3	-	1	3	3	7
Male	184	-	1	-	-	-	2	-	1	3	2	1
Female	206	-	-	1	-	-	1	-	-	-	1	6
White, both sexes	356	-	1	-	-	-	2	-	-	2	3	7
Male	168	-	1	-	-	-	1	-	-	2	2	1
Female	188	-	-	-	-	-	1	-	-	-	1	6
Black, both sexes	33	-	-	1	-	-	1	-	1	1	-	-
Male	16	-	-	-	-	-	1	-	1	1	-	-
Female	17	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	3,451	2	-	-	1	1	7	6	8	12	41	61
Male	1,651	2	-	-	1	1	5	3	8	9	28	46
Female	1,800	-	-	-	-	-	2	3	-	3	13	15
White, both sexes	2,973	-	-	-	1	1	5	3	5	8	28	40
Male	1,409	-	-	-	1	1	3	1	5	5	18	27
Female	1,564	-	-	-	-	-	2	2	-	3	10	13
Black, both sexes	229	1	-	-	-	-	1	1	2	3	8	12
Male	105	1	-	-	-	-	1	1	2	3	5	12
Female	124	-	-	-	-	-	-	-	-	-	3	-
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	239	1	-	-	-	-	1	2	1	1	5	9
Male	133	1	-	-	-	-	1	1	1	1	5	7
Female	106	-	-	-	-	-	-	1	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Colorado												
All races, both sexes	735	-	-	1	1	-	-	2	1	2	9	5
Male	300	-	-	1	1	-	-	2	-	2	5	4
Female	435	-	-	-	-	-	-	-	1	-	4	1
White, both sexes	696	-	-	1	1	-	-	1	1	2	8	5
Male	282	-	-	1	1	-	-	1	-	2	4	4
Female	414	-	-	-	-	-	-	-	1	-	4	1
Black, both sexes	30	-	-	-	-	-	-	1	-	-	1	-
Male	14	-	-	-	-	-	-	1	-	-	1	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	432	1	-	-	-	-	-	-	1	4	-	2
Male	195	1	-	-	-	-	-	-	1	3	-	2
Female	237	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	415	-	-	-	-	-	-	-	1	2	-	2
Male	185	-	-	-	-	-	-	-	1	1	-	2
Female	230	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	15	1	-	-	-	-	-	-	-	2	-	-
Male	9	1	-	-	-	-	-	-	-	2	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Delaware												
All races, both sexes	91	-	-	-	-	-	-	1	-	-	-	4
Male	49	-	-	-	-	-	-	1	-	-	-	2
Female	42	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	79	-	-	-	-	-	-	1	-	-	-	3
Male	42	-	-	-	-	-	-	1	-	-	-	2
Female	37	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	82	-	-	-	-	-	-	-	-	2	1	1
Male	34	-	-	-	-	-	-	-	-	1	1	1
Female	48	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	72	-	-	-	-	-	-	-	-	2	1	1
Male	30	-	-	-	-	-	-	-	-	1	1	1
Female	42	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Florida												
All races, both sexes	2,510	-	1	-	1	-	1	3	6	15	18	24
Male	1,264	-	1	-	1	-	1	2	4	8	11	19
Female	1,246	-	-	-	-	-	-	1	2	7	7	5
White, both sexes	2,226	-	1	-	1	-	1	-	3	8	9	14
Male	1,141	-	1	-	1	-	1	-	3	4	7	11
Female	1,085	-	-	-	-	-	-	-	-	4	2	3
Black, both sexes	263	-	-	-	-	-	-	3	3	7	9	10
Male	113	-	-	-	-	-	-	2	1	4	4	8
Female	150	-	-	-	-	-	-	1	2	3	5	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	838	2	-	1	-	-	-	3	2	9	11	22
Male	399	1	-	-	-	-	-	1	1	8	9	15
Female	439	1	-	1	-	-	-	2	1	1	2	7
White, both sexes	630	2	-	1	-	-	-	-	-	5	4	10
Male	296	1	-	-	-	-	-	-	-	4	3	9
Female	334	1	-	1	-	-	-	-	-	1	1	1
Black, both sexes	205	-	-	-	-	-	-	3	1	4	7	12
Male	100	-	-	-	-	-	-	1	-	4	6	6
Female	105	-	-	-	-	-	-	2	1	-	1	6
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	3	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Hawaii												
All races, both sexes	132	-	-	-	1	-	-	1	-	2	1	1
Male	74	-	-	-	1	-	-	1	-	2	1	1
Female	58	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	1	-	-	-	-	-	-	1
Male	18	-	-	-	1	-	-	-	-	-	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	100	-	-	-	-	-	-	1	-	2	-	-
Male	54	-	-	-	-	-	-	1	-	2	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	140	-	-	-	-	-	-	-	1	1	-	2
Male	76	-	-	-	-	-	-	-	1	1	-	2
Female	64	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	136	-	-	-	-	-	-	-	1	1	-	2
Male	74	-	-	-	-	-	-	-	1	1	-	2
Female	62	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Illinois												
All races, both sexes	1,508	2	-	-	1	-	2	1	3	9	14	22
Male	696	1	-	-	1	-	2	-	2	8	11	17
Female	812	1	-	-	-	-	-	1	1	1	3	5
White, both sexes	1,310	2	-	-	1	-	2	1	1	4	10	13
Male	590	1	-	-	1	-	2	-	1	4	7	11
Female	720	1	-	-	-	-	-	1	-	-	3	2
Black, both sexes	181	-	-	-	-	-	-	-	2	4	4	8
Male	96	-	-	-	-	-	-	-	1	4	4	5
Female	85	-	-	-	-	-	-	-	1	-	-	3
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	1	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	1	-	-
Indiana												
All races, both sexes	902	-	-	1	-	1	1	5	3	3	14	13
Male	449	-	-	1	-	1	1	2	3	3	7	12
Female	453	-	-	-	-	-	-	3	-	-	7	1
White, both sexes	846	-	-	1	-	1	1	3	3	1	13	13
Male	417	-	-	1	-	1	1	1	3	1	6	12
Female	429	-	-	-	-	-	-	2	-	-	7	1
Black, both sexes	56	-	-	-	-	-	-	2	-	2	1	-
Male	32	-	-	-	-	-	-	1	-	2	1	-
Female	24	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Iowa												
All races, both sexes	600	-	-	-	-	-	-	1	2	1	3	4
Male	282	-	-	-	-	-	-	-	-	1	1	2
Female	318	-	-	-	-	-	-	1	2	-	2	2
White, both sexes	593	-	-	-	-	-	-	1	2	1	3	4
Male	277	-	-	-	-	-	-	-	-	1	1	2
Female	316	-	-	-	-	-	-	1	2	-	2	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	617	-	-	-	-	-	-	1	1	1	3	4
Male	262	-	-	-	-	-	-	-	-	1	2	4
Female	355	-	-	-	-	-	-	1	1	-	1	-
White, both sexes	589	-	-	-	-	-	-	1	1	-	3	4
Male	249	-	-	-	-	-	-	-	-	-	2	4
Female	340	-	-	-	-	-	-	1	1	-	1	-
Black, both sexes	22	-	-	-	-	-	-	-	-	1	-	-
Male	9	-	-	-	-	-	-	-	-	1	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Kentucky												
All races, both sexes	515	-	-	-	1	1	-	-	1	-	2	4
Male	258	-	-	-	1	1	-	-	1	-	1	4
Female	257	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	481	-	-	-	1	1	-	-	1	-	2	4
Male	246	-	-	-	1	1	-	-	1	-	1	4
Female	235	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	571	-	-	-	1	-	1	2	1	7	8	17
Male	253	-	-	-	1	-	1	2	1	4	6	6
Female	318	-	-	-	-	-	-	-	-	3	2	11
White, both sexes	402	-	-	-	-	-	-	-	1	5	5	7
Male	169	-	-	-	-	-	-	-	1	4	4	2
Female	233	-	-	-	-	-	-	-	-	1	1	5
Black, both sexes	165	-	-	-	1	-	1	2	-	2	2	10
Male	81	-	-	-	1	-	1	2	-	-	1	4
Female	84	-	-	-	-	-	-	-	-	2	1	6
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Maine												
All races, both sexes	199	-	-	-	-	-	-	-	-	-	4	2
Male	93	-	-	-	-	-	-	-	-	-	2	1
Female	106	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	199	-	-	-	-	-	-	-	-	-	4	2
Male	93	-	-	-	-	-	-	-	-	-	2	1
Female	106	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	569	-	-	-	-	1	1	1	2	6	5	9
Male	254	-	-	-	-	1	-	-	1	6	5	7
Female	315	-	-	-	-	-	1	1	1	-	-	2
White, both sexes	451	-	-	-	-	-	1	1	2	3	3	3
Male	198	-	-	-	-	-	-	-	1	3	3	2
Female	253	-	-	-	-	-	1	1	1	-	-	1
Black, both sexes	110	-	-	-	-	1	-	-	-	3	2	6
Male	50	-	-	-	-	1	-	-	-	3	2	5
Female	60	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Massachusetts												
All races, both sexes	743	-	-	-	-	-	1	-	2	2	2	3
Male	317	-	-	-	-	-	-	-	2	2	1	3
Female	426	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	709	-	-	-	-	-	1	-	2	2	2	2
Male	302	-	-	-	-	-	-	-	2	2	1	2
Female	407	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	1,385	2	-	-	1	3	1	6	3	5	12	24
Male	658	1	-	-	-	2	1	3	2	4	7	18
Female	727	1	-	-	1	1	-	3	1	1	5	6
White, both sexes	1,213	2	-	-	1	3	1	5	2	3	7	16
Male	584	1	-	-	-	2	1	3	1	2	4	13
Female	629	1	-	-	1	1	-	2	1	1	3	3
Black, both sexes	156	-	-	-	-	-	-	1	1	2	4	8
Male	65	-	-	-	-	-	-	-	1	2	2	5
Female	91	-	-	-	-	-	-	1	-	-	2	3
American Indian 1/, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Minnesota												
All races, both sexes	600	-	-	-	-	-	1	1	-	2	7	2
Male	314	-	-	-	-	-	1	1	-	2	5	1
Female	286	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	585	-	-	-	-	-	1	1	-	1	5	1
Male	305	-	-	-	-	-	1	1	-	1	4	1
Female	280	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	8	-	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Mississippi												
All races, both sexes	278	-	-	-	-	-	1	-	6	3	6	8
Male	131	-	-	-	-	-	1	-	3	2	2	4
Female	147	-	-	-	-	-	-	-	3	1	4	4
White, both sexes	190	-	-	-	-	-	1	-	2	1	4	4
Male	93	-	-	-	-	-	1	-	2	-	1	1
Female	97	-	-	-	-	-	-	-	-	1	3	3
Black, both sexes	86	-	-	-	-	-	-	-	4	2	2	4
Male	36	-	-	-	-	-	-	-	1	2	1	3
Female	50	-	-	-	-	-	-	-	3	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Missouri												
All races, both sexes	805	1	-	-	-	-	1	-	1	3	13	10
Male	383	1	-	-	-	-	1	-	1	1	9	6
Female	422	-	-	-	-	-	-	-	-	2	4	4
White, both sexes	716	-	-	-	-	-	1	-	1	2	11	9
Male	346	-	-	-	-	-	1	-	1	1	9	6
Female	370	-	-	-	-	-	-	-	-	1	2	3
Black, both sexes	85	1	-	-	-	-	-	-	-	1	2	1
Male	34	1	-	-	-	-	-	-	-	-	-	-
Female	51	-	-	-	-	-	-	-	-	1	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	115	-	-	-	-	-	-	-	-	1	-	-
Male	63	-	-	-	-	-	-	-	-	1	-	-
Female	52	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	109	-	-	-	-	-	-	-	-	1	-	-
Male	63	-	-	-	-	-	-	-	-	1	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Nebraska												
All races, both sexes	248	-	-	-	-	-	-	-	-	1	-	1
Male	122	-	-	-	-	-	-	-	-	-	-	1
Female	126	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	242	-	-	-	-	-	-	-	-	1	-	1
Male	119	-	-	-	-	-	-	-	-	-	-	1
Female	123	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	308	-	-	-	1	-	-	-	-	2	5	2
Male	129	-	-	-	1	-	-	-	-	-	4	1
Female	179	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	292	-	-	-	-	-	-	-	-	2	5	2
Male	123	-	-	-	-	-	-	-	-	-	4	1
Female	169	-	-	-	-	-	-	-	-	2	1	1
Black, both sexes	9	-	-	-	1	-	-	-	-	-	-	-
Male	3	-	-	-	1	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
New Hampshire												
All races, both sexes	168	-	-	-	-	-	-	1	-	1	2	2
Male	68	-	-	-	-	-	-	-	-	1	1	1
Female	100	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	168	-	-	-	-	-	-	1	-	1	2	2
Male	68	-	-	-	-	-	-	-	-	1	1	1
Female	100	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	996	-	-	-	-	-	-	3	1	7	10	10
Male	447	-	-	-	-	-	-	-	1	4	6	9
Female	549	-	-	-	-	-	-	3	-	3	4	1
White, both sexes	877	-	-	-	-	-	-	2	-	5	7	8
Male	394	-	-	-	-	-	-	-	-	3	4	7
Female	483	-	-	-	-	-	-	2	-	2	3	1
Black, both sexes	96	-	-	-	-	-	-	1	-	2	3	2
Male	39	-	-	-	-	-	-	-	-	1	2	2
Female	57	-	-	-	-	-	-	1	-	1	1	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	20	-	-	-	-	-	-	-	1	-	-	-
Male	13	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
New Mexico												
All races, both sexes	205	-	-	-	-	-	-	-	-	-	2	2
Male	113	-	-	-	-	-	-	-	-	-	2	2
Female	92	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	196	-	-	-	-	-	-	-	-	-	2	1
Male	106	-	-	-	-	-	-	-	-	-	2	1
Female	90	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	1,881	-	-	1	1	-	3	4	5	4	24	39
Male	920	-	-	1	-	-	2	3	4	3	19	31
Female	961	-	-	-	1	-	1	1	1	1	5	8
White, both sexes	1,636	-	-	1	1	-	3	3	5	2	17	27
Male	780	-	-	1	-	-	2	2	4	1	13	21
Female	856	-	-	-	1	-	1	1	1	1	4	6
Black, both sexes	218	-	-	-	-	-	-	1	-	2	4	10
Male	119	-	-	-	-	-	-	1	-	2	3	8
Female	99	-	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	26	-	-	-	-	-	-	-	-	-	3	2
Male	20	-	-	-	-	-	-	-	-	-	3	2
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
North Carolina												
All races, both sexes	986	1	-	-	-	1	1	1	5	-	12	19
Male	495	-	-	-	-	1	-	1	1	-	8	15
Female	491	1	-	-	-	-	1	-	4	-	4	4
White, both sexes	765	1	-	-	-	1	1	1	3	-	8	14
Male	401	-	-	-	-	1	-	1	-	-	6	12
Female	364	1	-	-	-	-	1	-	3	-	2	2
Black, both sexes	211	-	-	-	-	-	-	-	2	-	4	4
Male	91	-	-	-	-	-	-	-	1	-	2	3
Female	120	-	-	-	-	-	-	-	1	-	2	1
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
North Dakota												
All races, both sexes	98	-	-	-	-	-	-	-	1	1	-	1
Male	47	-	-	-	-	-	-	-	1	1	-	-
Female	51	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	95	-	-	-	-	-	-	-	1	-	-	-
Male	46	-	-	-	-	-	-	-	1	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Ohio												
All races, both sexes	1,687	-	-	-	-	1	2	2	7	8	8	21
Male	797	-	-	-	-	1	2	2	6	6	4	15
Female	890	-	-	-	-	-	-	-	1	2	4	6
White, both sexes	1,519	-	-	-	-	1	2	-	5	6	5	13
Male	724	-	-	-	-	1	2	-	4	4	1	9
Female	795	-	-	-	-	-	-	-	1	2	4	4
Black, both sexes	166	-	-	-	-	-	-	2	2	2	3	8
Male	73	-	-	-	-	-	-	2	2	2	3	6
Female	93	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	566	-	-	-	-	1	1	3	4	-	3	11
Male	266	-	-	-	-	1	1	2	2	-	3	6
Female	300	-	-	-	-	-	-	1	2	-	-	5
White, both sexes	512	-	-	-	-	1	-	2	4	-	2	8
Male	243	-	-	-	-	1	-	1	2	-	2	5
Female	269	-	-	-	-	-	-	1	2	-	-	3
Black, both sexes	31	-	-	-	-	-	1	-	-	-	-	2
Male	9	-	-	-	-	-	1	-	-	-	-	1
Female	22	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	22	-	-	-	-	-	-	1	-	-	1	1
Male	13	-	-	-	-	-	-	1	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Oregon												
All races, both sexes	505	-	-	-	-	-	-	-	-	-	2	4
Male	255	-	-	-	-	-	-	-	-	-	1	4
Female	250	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	494	-	-	-	-	-	-	-	-	-	2	4
Male	251	-	-	-	-	-	-	-	-	-	1	4
Female	243	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	1,860	-	-	-	-	1	1	2	5	11	11	19
Male	797	-	-	-	-	1	-	-	4	10	5	14
Female	1,063	-	-	-	-	-	1	2	1	1	6	5
White, both sexes	1,740	-	-	-	-	-	1	2	5	8	10	14
Male	743	-	-	-	-	-	-	-	4	7	4	11
Female	997	-	-	-	-	-	1	2	1	1	6	3
Black, both sexes	115	-	-	-	-	1	-	-	-	3	-	5
Male	52	-	-	-	-	1	-	-	-	3	-	3
Female	63	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Rhode Island												
All races, both sexes	156	-	-	-	-	1	-	1	-	-	2	2
Male	71	-	-	-	-	1	-	1	-	-	1	-
Female	85	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	149	-	-	-	-	1	-	-	-	-	2	2
Male	67	-	-	-	-	1	-	-	-	-	1	-
Female	82	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	5	-	-	-	-	-	-	1	-	-	-	-
Male	2	-	-	-	-	-	-	1	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	489	-	-	-	-	-	2	1	5	-	5	7
Male	238	-	-	-	-	-	2	-	4	-	4	7
Female	251	-	-	-	-	-	-	1	1	-	1	-
White, both sexes	348	-	-	-	-	-	1	-	-	-	3	5
Male	177	-	-	-	-	-	1	-	-	-	3	5
Female	171	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	137	-	-	-	-	-	1	1	5	-	2	2
Male	59	-	-	-	-	-	1	-	4	-	1	2
Female	78	-	-	-	-	-	-	1	1	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
South Dakota												
All races, both sexes	116	-	-	-	-	-	-	-	-	-	-	2
Male	64	-	-	-	-	-	-	-	-	-	-	1
Female	52	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	113	-	-	-	-	-	-	-	-	-	-	2
Male	62	-	-	-	-	-	-	-	-	-	-	1
Female	51	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	765	-	-	-	-	1	1	1	2	3	10	11
Male	374	-	-	-	-	1	1	-	2	3	7	6
Female	391	-	-	-	-	-	-	1	-	-	3	5
White, both sexes	666	-	-	-	-	-	1	-	2	2	7	9
Male	331	-	-	-	-	-	1	-	2	2	5	5
Female	335	-	-	-	-	-	-	-	-	-	2	4
Black, both sexes	96	-	-	-	-	1	-	1	-	1	3	2
Male	40	-	-	-	-	1	-	-	-	1	2	1
Female	56	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Texas												
All races, both sexes	1,942	3	1	-	-	3	3	8	6	14	25	37
Male	947	2	1	-	-	1	-	6	5	5	19	26
Female	995	1	-	-	-	2	3	2	1	9	6	11
White, both sexes	1,685	2	-	-	-	3	2	6	5	10	16	27
Male	821	2	-	-	-	1	-	4	4	4	12	19
Female	864	-	-	-	-	2	2	2	1	6	4	8
Black, both sexes	245	1	1	-	-	-	1	2	-	4	9	10
Male	120	-	1	-	-	-	-	2	-	1	7	7
Female	125	1	-	-	-	-	1	-	-	3	2	3
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	1	-	-	-
Male	6	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	167	-	-	-	-	-	-	1	-	-	2	6
Male	88	-	-	-	-	-	-	1	-	-	-	4
Female	79	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	158	-	-	-	-	-	-	1	-	-	1	5
Male	81	-	-	-	-	-	-	1	-	-	-	3
Female	77	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Vermont												
All races, both sexes	87	-	-	-	-	-	-	-	-	1	1	1
Male	47	-	-	-	-	-	-	-	-	1	1	-
Female	40	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	87	-	-	-	-	-	-	-	-	1	1	1
Male	47	-	-	-	-	-	-	-	-	1	1	-
Female	40	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	738	-	-	-	-	-	1	1	4	5	6	10
Male	347	-	-	-	-	-	-	1	3	2	5	4
Female	391	-	-	-	-	-	1	-	1	3	1	6
White, both sexes	605	-	-	-	-	-	1	1	1	3	3	6
Male	284	-	-	-	-	-	-	1	1	1	3	3
Female	321	-	-	-	-	-	1	-	-	2	-	3
Black, both sexes	121	-	-	-	-	-	-	-	2	2	3	4
Male	53	-	-	-	-	-	-	-	1	1	2	1
Female	68	-	-	-	-	-	-	-	1	1	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	1	-	-	-
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Washington												
All races, both sexes	842	1	-	-	1	-	-	1	2	6	5	16
Male	413	-	-	-	1	-	-	-	2	6	4	9
Female	429	1	-	-	-	-	-	1	-	-	1	7
White, both sexes	794	1	-	-	1	-	-	1	1	4	3	13
Male	381	-	-	-	1	-	-	-	1	4	3	7
Female	413	1	-	-	-	-	-	1	-	-	-	6
Black, both sexes	22	-	-	-	-	-	-	-	-	2	1	1
Male	17	-	-	-	-	-	-	-	-	2	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	11	-	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	1	-	-	1
Male	10	-	-	-	-	-	-	-	1	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	324	1	-	-	-	-	-	-	1	-	1	4
Male	151	1	-	-	-	-	-	-	-	-	1	3
Female	173	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	313	1	-	-	-	-	-	-	1	-	1	3
Male	149	1	-	-	-	-	-	-	-	-	1	2
Female	164	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	10	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of arteries, arterioles and capillaries (I70-I78)												
Wisconsin												
All races, both sexes	718	-	-	-	-	-	1	1	-	6	4	9
Male	363	-	-	-	-	-	-	-	-	5	1	5
Female	355	-	-	-	-	-	1	1	-	1	3	4
White, both sexes	703	-	-	-	-	-	1	1	-	5	4	8
Male	356	-	-	-	-	-	-	-	-	4	1	4
Female	347	-	-	-	-	-	1	1	-	1	3	4
Black, both sexes	10	-	-	-	-	-	-	-	-	1	-	-
Male	6	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	61	-	-	-	1	-	-	-	-	-	1	-
Male	33	-	-	-	1	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	59	-	-	-	1	-	-	-	-	-	1	-
Male	32	-	-	-	1	-	-	-	-	-	-	-
Female	27	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
United States												
All races, both sexes	769	1,103	1,603	2,225	3,267	5,027	6,527	6,395	4,775	2,032	474	-
Male	524	753	1,049	1,398	1,958	2,757	3,095	2,515	1,466	435	73	-
Female	245	350	554	827	1,309	2,270	3,432	3,880	3,309	1,597	401	-
White, both sexes	553	871	1,340	1,871	2,849	4,474	5,901	5,905	4,406	1,867	416	-
Male	382	603	897	1,188	1,749	2,482	2,825	2,352	1,374	406	63	-
Female	171	268	443	683	1,100	1,992	3,076	3,553	3,032	1,461	353	-
Black, both sexes	185	211	224	299	348	440	503	406	307	143	51	-
Male	118	137	130	174	164	215	205	122	72	19	7	-
Female	67	74	94	125	184	225	298	284	235	124	44	-
American Indian 1/, both sexes	4	10	13	16	19	23	18	11	7	3	-	-
Male	2	6	6	8	11	8	9	3	2	-	-	-
Female	2	4	7	8	8	15	9	8	5	3	-	-
Asian or Pacific Islander 2/, both sexes	27	11	26	39	51	90	105	73	55	19	7	-
Male	22	7	16	28	34	52	56	38	18	10	3	-
Female	5	4	10	11	17	38	49	35	37	9	4	-
Alabama												
All races, both sexes	22	26	39	52	83	85	111	104	73	21	2	-
Male	15	20	26	27	51	45	44	37	19	5	-	-
Female	7	6	13	25	32	40	67	67	54	16	2	-
White, both sexes	16	15	26	41	72	66	86	93	60	16	1	-
Male	10	12	17	22	46	34	35	34	16	4	-	-
Female	6	3	9	19	26	32	51	59	44	12	1	-
Black, both sexes	6	10	13	11	10	19	22	11	13	5	1	-
Male	5	8	9	5	5	11	8	3	3	1	-	-
Female	1	2	4	6	5	8	14	8	10	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Alaska												
All races, both sexes	-	-	3	1	2	5	10	3	4	-	-	-
Male	-	-	2	1	2	3	5	2	1	-	-	-
Female	-	-	1	-	-	2	5	1	3	-	-	-
White, both sexes	-	-	3	1	1	5	8	3	2	-	-	-
Male	-	-	2	1	1	3	5	2	-	-	-	-
Female	-	-	1	-	-	2	3	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Arizona												
All races, both sexes	15	17	31	39	66	106	124	115	86	42	4	-
Male	8	15	21	24	39	59	62	50	28	18	1	-
Female	7	2	10	15	27	47	62	65	58	24	3	-
White, both sexes	13	16	31	38	63	98	118	112	86	41	3	-
Male	7	14	21	23	37	54	61	49	28	18	-	-
Female	6	2	10	15	26	44	57	63	58	23	3	-
Black, both sexes	2	1	-	-	2	2	3	-	-	-	1	-
Male	1	1	-	-	1	2	-	-	-	-	1	-
Female	1	-	-	-	1	-	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	5	3	1	-	1	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	3	2	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	2	-	-	-	-
Male	-	-	-	1	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Arkansas												
All races, both sexes	7	16	28	31	36	56	62	70	44	17	4	-
Male	4	12	17	18	21	28	27	33	8	6	-	-
Female	3	4	11	13	15	28	35	37	36	11	4	-
White, both sexes	6	14	24	29	34	53	56	64	40	17	4	-
Male	4	10	14	16	19	28	25	31	8	6	-	-
Female	2	4	10	13	15	25	31	33	32	11	4	-
Black, both sexes	1	2	4	2	2	3	5	6	4	-	-	-
Male	-	2	3	2	2	-	2	2	-	-	-	-
Female	1	-	1	-	-	3	3	4	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	79	103	132	194	286	457	652	650	481	218	60	-
Male	55	71	86	123	173	261	307	255	158	49	10	-
Female	24	32	46	71	113	196	345	395	323	169	50	-
White, both sexes	61	79	111	156	242	396	562	585	440	197	53	-
Male	44	55	72	98	151	231	267	228	150	44	8	-
Female	17	24	39	58	91	165	295	357	290	153	45	-
Black, both sexes	8	17	8	18	20	23	36	34	18	14	5	-
Male	3	12	6	12	10	9	14	9	3	2	-	-
Female	5	5	2	6	10	14	22	25	15	12	5	-
American Indian 1/, both sexes	-	1	1	3	1	2	1	1	-	-	-	-
Male	-	-	-	2	-	1	1	-	-	-	-	-
Female	-	1	1	1	1	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	6	12	17	23	36	53	30	23	7	2	-
Male	8	4	8	11	12	20	25	18	5	3	2	-
Female	2	2	4	6	11	16	28	12	18	4	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Colorado												
All races, both sexes	14	25	26	25	57	103	135	126	131	63	9	-
Male	9	15	18	15	35	47	58	45	31	12	-	-
Female	5	10	8	10	22	56	77	81	100	51	9	-
White, both sexes	13	23	25	22	54	93	129	123	124	62	9	-
Male	8	13	17	13	33	42	55	45	31	12	-	-
Female	5	10	8	9	21	51	74	78	93	50	9	-
Black, both sexes	1	2	1	2	2	7	5	2	5	1	-	-
Male	1	2	1	2	1	3	2	-	-	-	-	-
Female	-	-	-	-	1	4	3	2	5	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	3	1	-	2	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	2	-	-	-
Connecticut												
All races, both sexes	2	10	12	16	37	62	73	96	74	34	8	-
Male	1	9	9	12	23	27	42	31	24	8	2	-
Female	1	1	3	4	14	35	31	65	50	26	6	-
White, both sexes	2	9	10	15	37	58	71	95	72	33	8	-
Male	1	8	8	11	23	25	41	31	23	8	2	-
Female	1	1	2	4	14	33	30	64	49	25	6	-
Black, both sexes	-	1	2	1	-	2	2	1	2	1	-	-
Male	-	1	1	1	-	1	1	-	1	-	-	-
Female	-	-	1	-	-	1	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Delaware												
All races, both sexes	-	2	7	7	10	19	16	10	11	4	-	-
Male	-	-	7	5	3	14	6	7	3	1	-	-
Female	-	2	-	2	7	5	10	3	8	3	-	-
White, both sexes	-	2	5	5	8	16	16	10	10	3	-	-
Male	-	-	5	3	2	12	6	7	3	1	-	-
Female	-	2	-	2	6	4	10	3	7	2	-	-
Black, both sexes	-	-	1	2	2	3	-	-	1	1	-	-
Male	-	-	1	2	1	2	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	8	9	6	6	8	7	10	15	3	4	-
Male	2	3	4	3	3	5	3	4	4	-	-	-
Female	-	5	5	3	3	3	4	6	11	3	4	-
White, both sexes	-	-	1	-	-	1	1	1	2	1	2	-
Male	-	-	-	-	-	1	-	1	1	-	-	-
Female	-	-	1	-	-	-	1	-	1	1	2	-
Black, both sexes	2	8	8	6	6	7	5	9	13	2	2	-
Male	2	3	4	3	3	4	2	3	3	-	-	-
Female	-	5	4	3	3	3	3	6	10	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Florida												
All races, both sexes	56	79	113	162	214	378	490	451	332	138	28	-
Male	38	46	75	101	134	208	253	196	116	40	10	-
Female	18	33	38	61	80	170	237	255	216	98	18	-
White, both sexes	35	63	100	136	184	340	444	422	309	129	27	-
Male	22	37	69	91	118	192	238	185	112	39	10	-
Female	13	26	31	45	66	148	206	237	197	90	17	-
Black, both sexes	18	14	13	23	28	32	43	28	22	9	1	-
Male	13	8	6	9	15	15	13	11	3	1	-	-
Female	5	6	7	14	13	17	30	17	19	8	1	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	1	-	2	2	6	3	1	1	-	-	-
Male	3	1	-	1	1	1	2	-	1	-	-	-
Female	-	-	-	1	1	5	1	1	-	-	-	-
Georgia												
All races, both sexes	29	51	50	63	80	101	143	130	91	41	9	-
Male	20	35	31	48	46	56	58	45	14	9	2	-
Female	9	16	19	15	34	45	85	85	77	32	7	-
White, both sexes	15	31	34	44	64	84	120	109	70	31	6	-
Male	11	23	24	36	36	47	48	37	10	5	2	-
Female	4	8	10	8	28	37	72	72	60	26	4	-
Black, both sexes	14	20	16	18	15	17	23	21	21	10	3	-
Male	9	12	7	11	9	9	10	8	4	4	-	-
Female	5	8	9	7	6	8	13	13	17	6	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	-	-	-	-	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Hawaii												
All races, both sexes	6	4	6	8	9	18	19	22	21	10	3	-
Male	5	3	3	5	7	11	12	11	6	5	-	-
Female	1	1	3	3	2	7	7	11	15	5	3	-
White, both sexes	1	2	2	4	5	5	-	3	5	1	-	-
Male	1	2	1	2	5	3	-	1	1	-	-	-
Female	-	-	1	2	-	2	-	2	4	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	2	4	4	4	13	18	19	16	9	3	-
Male	4	1	2	3	2	8	11	10	5	5	-	-
Female	1	1	2	1	2	5	7	9	11	4	3	-
Idaho												
All races, both sexes	4	3	6	11	16	20	24	28	10	12	2	-
Male	3	1	3	8	13	9	16	12	2	4	1	-
Female	1	2	3	3	3	11	8	16	8	8	1	-
White, both sexes	3	3	6	11	15	19	24	27	10	12	2	-
Male	2	1	3	8	12	9	16	12	2	4	1	-
Female	1	2	3	3	3	10	8	15	8	8	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Illinois												
All races, both sexes	30	38	61	92	134	210	279	272	229	88	21	-
Male	23	26	40	55	74	130	125	99	68	10	4	-
Female	7	12	21	37	60	80	154	173	161	78	17	-
White, both sexes	18	27	53	74	108	183	249	251	211	82	20	-
Male	13	18	37	43	65	111	110	90	63	10	3	-
Female	5	9	16	31	43	72	139	161	148	72	17	-
Black, both sexes	11	10	8	17	24	24	27	19	16	6	1	-
Male	10	7	3	11	8	16	14	8	4	-	1	-
Female	1	3	5	6	16	8	13	11	12	6	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	1	3	3	1	2	-	-	-
Male	-	1	-	1	1	3	1	-	1	-	-	-
Female	1	-	-	-	-	-	2	1	1	-	-	-
Indiana												
All races, both sexes	16	30	37	64	89	125	153	169	126	45	7	-
Male	12	23	23	38	60	71	81	62	36	12	1	-
Female	4	7	14	26	29	54	72	107	90	33	6	-
White, both sexes	11	27	35	58	83	119	149	159	120	44	5	-
Male	8	20	21	34	58	69	79	57	33	12	-	-
Female	3	7	14	24	25	50	70	102	87	32	5	-
Black, both sexes	5	3	2	6	6	6	4	10	6	1	2	-
Male	4	3	2	4	2	2	2	5	3	-	1	-
Female	1	-	-	2	4	4	2	5	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Iowa												
All races, both sexes	7	13	20	25	43	96	113	113	103	41	15	-
Male	1	9	16	17	27	54	57	48	41	7	1	-
Female	6	4	4	8	16	42	56	65	62	34	14	-
White, both sexes	6	12	20	25	41	96	112	111	103	41	15	-
Male	1	8	16	17	25	54	56	47	41	7	1	-
Female	5	4	4	8	16	42	56	64	62	34	14	-
Black, both sexes	1	1	-	-	1	-	1	2	-	-	-	-
Male	-	1	-	-	1	-	1	1	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	10	13	24	31	33	50	123	132	117	58	16	-
Male	9	8	16	21	20	34	52	44	38	11	2	-
Female	1	5	8	10	13	16	71	88	79	47	14	-
White, both sexes	8	13	23	30	31	50	116	124	112	57	16	-
Male	8	8	15	20	19	34	48	42	36	11	2	-
Female	-	5	8	10	12	16	68	82	76	46	14	-
Black, both sexes	-	-	1	-	1	-	6	8	4	1	-	-
Male	-	-	1	-	-	-	3	2	2	-	-	-
Female	-	-	-	-	1	-	3	6	2	1	-	-
American Indian 1/, both sexes	1	-	-	1	1	-	1	-	1	-	-	-
Male	-	-	-	1	1	-	1	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Kentucky												
All races, both sexes	14	16	26	39	65	94	84	88	53	19	8	-
Male	10	9	17	26	38	55	40	30	18	4	3	-
Female	4	7	9	13	27	39	44	58	35	15	5	-
White, both sexes	13	14	26	35	60	91	79	82	49	17	6	-
Male	9	9	17	23	36	53	38	29	17	4	3	-
Female	4	5	9	12	24	38	41	53	32	13	3	-
Black, both sexes	1	2	-	4	5	3	5	6	4	2	2	-
Male	1	-	-	3	2	2	2	1	1	-	-	-
Female	-	2	-	1	3	1	3	5	3	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	14	19	28	46	53	74	106	92	68	24	10	-
Male	8	14	17	23	31	29	46	34	23	5	2	-
Female	6	5	11	23	22	45	60	58	45	19	8	-
White, both sexes	5	11	19	25	39	53	82	75	51	18	6	-
Male	4	8	15	11	24	19	33	26	14	3	1	-
Female	1	3	4	14	15	34	49	49	37	15	5	-
Black, both sexes	9	8	9	21	14	20	23	16	17	6	4	-
Male	4	6	2	12	7	10	12	7	9	2	1	-
Female	5	2	7	9	7	10	11	9	8	4	3	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Maine												
All races, both sexes	1	3	7	11	15	29	41	41	33	10	2	-
Male	-	3	4	7	11	14	16	19	13	3	-	-
Female	1	-	3	4	4	15	25	22	20	7	2	-
White, both sexes	1	3	7	11	15	29	41	41	33	10	2	-
Male	-	3	4	7	11	14	16	19	13	3	-	-
Female	1	-	3	4	4	15	25	22	20	7	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	17	14	20	44	57	92	93	88	72	41	6	-
Male	8	11	15	28	26	44	43	33	18	7	1	-
Female	9	3	5	16	31	48	50	55	54	34	5	-
White, both sexes	7	8	19	29	43	74	74	78	62	39	5	-
Male	4	7	15	21	24	33	32	29	16	7	1	-
Female	3	1	4	8	19	41	42	49	46	32	4	-
Black, both sexes	9	6	1	13	14	16	17	10	9	2	1	-
Male	3	4	-	6	2	9	9	4	2	-	-	-
Female	6	2	1	7	12	7	8	6	7	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	2	-	2	2	-	1	-	-	-
Male	1	-	-	1	-	2	2	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Massachusetts												
All races, both sexes	12	13	32	41	69	105	136	151	108	52	14	-
Male	8	8	17	27	40	55	64	57	25	8	-	-
Female	4	5	15	14	29	50	72	94	83	44	14	-
White, both sexes	10	11	27	40	63	100	131	148	107	49	14	-
Male	6	7	16	27	35	53	62	57	25	7	-	-
Female	4	4	11	13	28	47	69	91	82	42	14	-
Black, both sexes	2	2	3	1	4	5	4	3	1	2	-	-
Male	2	1	1	-	3	2	1	-	-	-	-	-
Female	-	1	2	1	1	3	3	3	1	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	2	-	1	-	-	1	-	-
Male	-	-	-	-	2	-	1	-	-	1	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	29	44	68	92	110	190	281	260	174	59	21	-
Male	19	29	41	46	67	95	129	107	68	16	3	-
Female	10	15	27	46	43	95	152	153	106	43	18	-
White, both sexes	19	34	57	68	96	167	262	234	164	55	17	-
Male	14	24	36	33	60	85	122	98	67	15	3	-
Female	5	10	21	35	36	82	140	136	97	40	14	-
Black, both sexes	10	10	8	21	12	21	16	25	9	4	4	-
Male	5	5	4	10	6	9	5	9	1	1	-	-
Female	5	5	4	11	6	12	11	16	8	3	4	-
American Indian 1/, both sexes	-	-	3	3	2	-	-	1	1	-	-	-
Male	-	-	1	3	1	-	-	-	-	-	-	-
Female	-	-	2	-	1	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	3	-	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Minnesota												
All races, both sexes	9	21	15	41	46	82	123	117	90	36	7	-
Male	5	14	10	30	30	54	66	50	32	12	1	-
Female	4	7	5	11	16	28	57	67	58	24	6	-
White, both sexes	8	19	14	39	45	79	123	117	89	36	7	-
Male	4	12	9	30	29	52	66	50	32	12	1	-
Female	4	7	5	9	16	27	57	67	57	24	6	-
Black, both sexes	1	1	1	-	-	3	-	-	-	-	-	-
Male	1	1	1	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	2	1	-	-	-	1	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	2	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	9	11	23	26	35	37	45	29	24	13	2	-
Male	6	6	15	16	21	21	16	12	5	1	-	-
Female	3	5	8	10	14	16	29	17	19	12	2	-
White, both sexes	6	5	16	16	29	29	33	18	17	8	1	-
Male	3	3	11	9	19	17	12	9	4	1	-	-
Female	3	2	5	7	10	12	21	9	13	7	1	-
Black, both sexes	3	6	7	9	6	8	12	10	7	5	1	-
Male	3	3	4	6	2	4	4	2	1	-	-	-
Female	-	3	3	3	4	4	8	8	6	5	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Missouri												
All races, both sexes	14	22	32	56	73	126	139	143	113	43	15	-
Male	12	14	19	37	42	71	66	56	35	7	5	-
Female	2	8	13	19	31	55	73	87	78	36	10	-
White, both sexes	10	20	25	50	65	110	127	128	106	38	13	-
Male	9	12	15	35	39	62	62	52	32	6	4	-
Female	1	8	10	15	26	48	65	76	74	32	9	-
Black, both sexes	4	2	7	6	8	13	12	14	7	5	2	-
Male	3	2	4	2	3	7	4	3	3	1	1	-
Female	1	-	3	4	5	6	8	11	4	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	2	4	3	6	13	20	17	29	11	9	-	-
Male	1	2	2	2	6	14	13	13	8	1	-	-
Female	1	2	1	4	7	6	4	16	3	8	-	-
White, both sexes	2	4	2	6	11	19	17	28	11	8	-	-
Male	1	2	2	2	6	14	13	13	8	1	-	-
Female	1	2	-	4	5	5	4	15	3	7	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	2	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	2	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Nebraska												
All races, both sexes	2	7	4	11	25	31	47	46	49	18	6	-
Male	-	7	2	9	19	16	19	22	18	8	1	-
Female	2	-	2	2	6	15	28	24	31	10	5	-
White, both sexes	2	7	4	11	23	30	47	44	49	17	6	-
Male	-	7	2	9	17	15	19	22	18	8	1	-
Female	2	-	2	2	6	15	28	22	31	9	5	-
Black, both sexes	-	-	-	-	2	1	-	2	-	1	-	-
Male	-	-	-	-	2	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	9	11	21	14	31	52	63	54	30	11	2	-
Male	5	7	13	4	14	27	27	18	5	2	1	-
Female	4	4	8	10	17	25	36	36	25	9	1	-
White, both sexes	8	11	19	14	28	49	63	52	26	11	2	-
Male	5	7	13	4	13	24	27	18	4	2	1	-
Female	3	4	6	10	15	25	36	34	22	9	1	-
Black, both sexes	1	-	2	-	1	2	-	1	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	1	-	2	-	1	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	1	2	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	1	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
New Hampshire												
All races, both sexes	3	6	3	11	9	25	28	32	20	21	4	-
Male	2	5	3	6	3	14	11	7	7	7	-	-
Female	1	1	-	5	6	11	17	25	13	14	4	-
White, both sexes	3	6	3	11	9	25	28	32	20	21	4	-
Male	2	5	3	6	3	14	11	7	7	7	-	-
Female	1	1	-	5	6	11	17	25	13	14	4	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	21	27	37	55	101	145	192	177	133	62	15	-
Male	15	15	26	29	55	83	93	65	33	11	2	-
Female	6	12	11	26	46	62	99	112	100	51	13	-
White, both sexes	16	24	26	45	84	127	177	162	123	58	13	-
Male	11	13	21	25	46	71	88	61	31	11	2	-
Female	5	11	5	20	38	56	89	101	92	47	11	-
Black, both sexes	5	3	9	9	10	12	14	12	9	4	1	-
Male	4	2	4	3	3	10	4	3	1	-	-	-
Female	1	1	5	6	7	2	10	9	8	4	1	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	7	5	1	2	1	-	1	-
Male	-	-	1	-	6	2	1	1	1	-	-	-
Female	-	-	1	-	1	3	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
New Mexico												
All races, both sexes	4	5	10	10	17	39	38	39	21	15	3	-
Male	3	2	8	6	12	22	19	19	13	5	-	-
Female	1	3	2	4	5	17	19	20	8	10	3	-
White, both sexes	3	5	9	10	16	38	37	37	21	14	3	-
Male	2	2	7	6	11	22	18	18	13	4	-	-
Female	1	3	2	4	5	16	19	19	8	10	3	-
Black, both sexes	1	-	-	-	1	1	-	1	-	1	-	-
Male	1	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	1	-	-	-	-
Male	-	-	1	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	43	73	106	98	194	256	334	315	238	114	29	-
Male	31	50	72	59	109	145	165	117	82	23	4	-
Female	12	23	34	39	85	111	169	198	156	91	25	-
White, both sexes	26	51	82	79	166	230	298	293	224	105	23	-
Male	19	37	57	47	91	131	143	111	76	22	2	-
Female	7	14	25	32	75	99	155	182	148	83	21	-
Black, both sexes	15	22	23	16	25	23	31	18	14	9	5	-
Male	10	13	14	10	15	12	19	4	6	1	1	-
Female	5	9	9	6	10	11	12	14	8	8	4	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	3	3	3	5	4	-	-	1	-
Male	2	-	-	2	3	2	3	2	-	-	1	-
Female	-	-	-	1	-	1	2	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
North Carolina												
All races, both sexes	36	34	54	65	102	135	172	179	112	47	10	-
Male	26	22	30	43	61	87	85	72	33	9	1	-
Female	10	12	24	22	41	48	87	107	79	38	9	-
White, both sexes	24	23	40	47	80	111	129	146	92	36	8	-
Male	19	14	19	32	49	72	72	65	29	9	1	-
Female	5	9	21	15	31	39	57	81	63	27	7	-
Black, both sexes	10	11	12	17	21	23	43	32	20	10	2	-
Male	6	8	9	11	12	15	13	7	4	-	-	-
Female	4	3	3	6	9	8	30	25	16	10	2	-
American Indian 1/, both sexes	1	-	2	1	1	-	-	-	-	1	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	1	-	-	1	1	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
North Dakota												
All races, both sexes	1	1	3	5	7	11	19	16	23	9	-	-
Male	1	-	3	1	4	8	14	9	3	2	-	-
Female	-	1	-	4	3	3	5	7	20	7	-	-
White, both sexes	1	-	3	5	7	11	19	16	23	9	-	-
Male	1	-	3	1	4	8	14	9	3	2	-	-
Female	-	-	-	4	3	3	5	7	20	7	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Ohio												
All races, both sexes	43	47	76	107	152	246	327	320	210	89	21	-
Male	27	36	49	63	99	127	154	130	57	17	2	-
Female	16	11	27	44	53	119	173	190	153	72	19	-
White, both sexes	34	41	68	100	132	219	296	298	197	83	19	-
Male	22	31	46	59	90	118	141	125	53	17	1	-
Female	12	10	22	41	42	101	155	173	144	66	18	-
Black, both sexes	9	6	7	7	20	27	31	21	13	6	2	-
Male	5	5	3	4	9	9	13	5	4	-	1	-
Female	4	1	4	3	11	18	18	16	9	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
Oklahoma												
All races, both sexes	9	27	26	36	48	90	101	92	77	34	3	-
Male	8	18	14	21	27	45	56	34	21	7	-	-
Female	1	9	12	15	21	45	45	58	56	27	3	-
White, both sexes	6	22	23	34	40	78	94	87	76	32	3	-
Male	6	15	13	20	25	39	53	33	21	7	-	-
Female	-	7	10	14	15	39	41	54	55	25	3	-
Black, both sexes	2	2	-	2	5	7	2	5	1	2	-	-
Male	1	1	-	1	-	3	-	1	-	-	-	-
Female	1	1	-	1	5	4	2	4	1	2	-	-
American Indian 1/, both sexes	1	3	3	-	3	5	4	-	-	-	-	-
Male	1	2	1	-	2	3	2	-	-	-	-	-
Female	-	1	2	-	1	2	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Oregon												
All races, both sexes	4	16	25	35	36	89	96	96	70	29	3	-
Male	4	10	17	22	23	44	52	49	21	7	1	-
Female	-	6	8	13	13	45	44	47	49	22	2	-
White, both sexes	4	16	24	33	35	86	96	94	70	27	3	-
Male	4	10	16	21	22	43	52	49	21	7	1	-
Female	-	6	8	12	13	43	44	45	49	20	2	-
Black, both sexes	-	-	1	1	-	1	-	-	-	2	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	2	-	-	-	-
Male	-	-	-	1	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Pennsylvania												
All races, both sexes	22	42	68	99	164	239	358	386	296	115	21	-
Male	15	32	43	64	91	123	138	139	91	26	1	-
Female	7	10	25	35	73	116	220	247	205	89	20	-
White, both sexes	18	33	60	89	149	220	336	371	292	112	20	-
Male	13	25	39	59	85	116	127	135	91	26	1	-
Female	5	8	21	30	64	104	209	236	201	86	19	-
Black, both sexes	3	9	8	9	15	18	21	15	4	3	1	-
Male	2	7	4	4	6	7	11	4	-	-	-	-
Female	1	2	4	5	9	11	10	11	4	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	1	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Rhode Island												
All races, both sexes	4	1	5	5	13	24	30	31	27	8	2	-
Male	3	1	2	4	7	16	15	13	6	1	-	-
Female	1	-	3	1	6	8	15	18	21	7	2	-
White, both sexes	4	1	5	5	13	20	28	31	27	8	2	-
Male	3	1	2	4	7	14	14	13	6	1	-	-
Female	1	-	3	1	6	6	14	18	21	7	2	-
Black, both sexes	-	-	-	-	-	3	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	2	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	9	18	35	43	53	74	87	76	51	21	2	-
Male	5	10	23	32	27	40	44	26	9	5	-	-
Female	4	8	12	11	26	34	43	50	42	16	2	-
White, both sexes	6	11	24	33	37	57	60	59	34	18	-	-
Male	3	6	13	26	21	33	35	20	8	3	-	-
Female	3	5	11	7	16	24	25	39	26	15	-	-
Black, both sexes	3	6	11	10	15	17	26	16	17	3	2	-
Male	2	3	10	6	5	7	9	6	1	2	-	-
Female	1	3	1	4	10	10	17	10	16	1	2	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
South Dakota												
All races, both sexes	3	2	5	2	10	15	22	27	19	7	2	-
Male	1	2	3	2	5	10	16	13	10	1	-	-
Female	2	-	2	-	5	5	6	14	9	6	2	-
White, both sexes	2	1	5	2	10	14	22	27	19	7	2	-
Male	-	1	3	2	5	10	16	13	10	1	-	-
Female	2	-	2	-	5	4	6	14	9	6	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	1	-	-	-	1	-	-	-	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	20	28	47	62	85	97	142	137	77	34	7	-
Male	12	13	30	48	54	46	69	52	24	5	1	-
Female	8	15	17	14	31	51	73	85	53	29	6	-
White, both sexes	16	25	38	52	76	84	129	122	65	32	6	-
Male	10	13	26	40	48	43	65	47	18	5	1	-
Female	6	12	12	12	28	41	64	75	47	27	5	-
Black, both sexes	4	3	9	10	8	13	13	14	11	2	1	-
Male	2	-	4	8	5	3	4	4	5	-	-	-
Female	2	3	5	2	3	10	9	10	6	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Texas												
All races, both sexes	52	70	95	145	222	287	324	281	242	103	21	-
Male	36	52	62	91	155	150	146	105	69	14	2	-
Female	16	18	33	54	67	137	178	176	173	89	19	-
White, both sexes	38	57	77	122	193	249	292	260	222	87	17	-
Male	28	43	48	76	136	134	132	98	66	12	2	-
Female	10	14	29	46	57	115	160	162	156	75	15	-
Black, both sexes	14	13	17	23	27	37	27	20	19	16	4	-
Male	8	9	13	15	17	15	13	7	3	2	-	-
Female	6	4	4	8	10	22	14	13	16	14	4	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	2	1	4	1	1	-	-	-
Male	-	-	1	-	2	1	1	-	-	-	-	-
Female	-	-	-	-	-	-	3	1	1	-	-	-
Utah												
All races, both sexes	6	4	4	15	18	24	36	28	13	8	2	-
Male	5	3	4	9	13	15	18	13	2	1	-	-
Female	1	1	-	6	5	9	18	15	11	7	2	-
White, both sexes	6	4	2	12	18	24	35	27	13	8	2	-
Male	5	3	2	7	13	15	17	12	2	1	-	-
Female	1	1	-	5	5	9	18	15	11	7	2	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	2	-	-	-	1	-	-	-	-
Male	-	-	2	2	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Vermont												
All races, both sexes	1	3	3	5	8	14	19	14	12	3	2	-
Male	-	2	2	4	3	10	11	9	4	-	-	-
Female	1	1	1	1	5	4	8	5	8	3	2	-
White, both sexes	1	3	3	5	8	14	19	14	12	3	2	-
Male	-	2	2	4	3	10	11	9	4	-	-	-
Female	1	1	1	1	5	4	8	5	8	3	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	16	18	46	56	65	106	131	128	99	37	9	-
Male	11	14	34	47	29	60	57	44	31	4	1	-
Female	5	4	12	9	36	46	74	84	68	33	8	-
White, both sexes	9	14	35	45	53	85	110	117	83	31	8	-
Male	5	12	27	37	25	48	48	42	26	4	1	-
Female	4	2	8	8	28	37	62	75	57	27	7	-
Black, both sexes	6	4	10	10	12	17	19	10	15	6	1	-
Male	5	2	6	9	4	9	8	1	4	-	-	-
Female	1	2	4	1	8	8	11	9	11	6	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	-	4	2	1	1	-	-	-
Male	1	-	1	1	-	3	1	1	1	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Washington												
All races, both sexes	21	24	37	53	69	97	166	161	127	40	15	-
Male	17	17	28	30	42	47	85	71	44	7	3	-
Female	4	7	9	23	27	50	81	90	83	33	12	-
White, both sexes	20	20	35	48	65	90	162	156	122	37	15	-
Male	16	13	27	27	39	43	83	69	40	5	3	-
Female	4	7	8	21	26	47	79	87	82	32	12	-
Black, both sexes	1	3	1	3	2	3	1	2	1	1	-	-
Male	1	3	1	1	2	2	-	1	1	1	-	-
Female	-	-	-	2	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	1	1	1	2	3	-	1	-	-	-	-
Male	-	1	-	1	1	2	-	-	-	-	-	-
Female	-	-	1	-	1	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	3	2	4	2	-	-
Male	-	-	-	1	-	-	2	1	3	1	-	-
Female	-	-	-	-	-	1	1	1	1	1	-	-
West Virginia												
All races, both sexes	5	8	12	23	31	48	60	58	47	20	5	-
Male	3	7	8	13	18	31	25	26	10	3	2	-
Female	2	1	4	10	13	17	35	32	37	17	3	-
White, both sexes	3	8	12	23	30	46	60	57	44	20	4	-
Male	3	7	8	13	18	31	25	25	10	3	2	-
Female	-	1	4	10	12	15	35	32	34	17	2	-
Black, both sexes	2	-	-	-	1	1	-	1	3	-	1	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	2	-	-	-	1	1	-	-	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of arteries, arterioles and capillaries (I70-I78)												
Wisconsin												
All races, both sexes	12	20	22	38	65	126	127	152	82	42	11	-
Male	9	18	19	26	42	72	62	63	31	8	2	-
Female	3	2	3	12	23	54	65	89	51	34	9	-
White, both sexes	11	18	21	35	64	125	126	151	81	42	10	-
Male	8	16	18	25	42	72	62	63	31	8	2	-
Female	3	2	3	10	22	53	64	88	50	34	8	-
Black, both sexes	1	2	1	1	1	-	1	1	-	-	1	-
Male	1	2	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	1	1	-	-	1	-
American Indian 1/, both sexes	-	-	-	2	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	-	1	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	3	6	1	3	5	9	9	11	6	4	2	-
Male	3	4	-	2	3	5	7	7	-	1	-	-
Female	-	2	1	1	2	4	2	4	6	3	2	-
White, both sexes	3	5	1	3	5	9	8	11	6	4	2	-
Male	3	4	-	2	3	5	6	7	-	1	-	-
Female	-	1	1	1	2	4	2	4	6	3	2	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
United States												
All races, both sexes	11,841	-	1	-	-	-	2	3	3	7	24	63
Male	4,472	-	1	-	-	-	1	1	2	6	14	40
Female	7,369	-	-	-	-	-	1	2	1	1	10	23
White, both sexes	10,845	-	-	-	-	-	1	2	3	6	20	54
Male	4,093	-	-	-	-	-	-	-	2	5	12	36
Female	6,752	-	-	-	-	-	1	2	1	1	8	18
Black, both sexes	844	-	1	-	-	-	1	1	-	1	4	8
Male	312	-	1	-	-	-	1	1	-	1	2	3
Female	532	-	-	-	-	-	-	-	-	-	2	5
American Indian 1/, both sexes	51	-	-	-	-	-	-	-	-	-	-	1
Male	24	-	-	-	-	-	-	-	-	-	-	1
Female	27	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	101	-	-	-	-	-	-	-	-	-	-	-
Male	43	-	-	-	-	-	-	-	-	-	-	-
Female	58	-	-	-	-	-	-	-	-	-	-	-
Alabama												
All races, both sexes	253	-	-	-	-	-	-	-	1	1	7	4
Male	107	-	-	-	-	-	-	-	1	-	4	2
Female	146	-	-	-	-	-	-	-	-	1	3	2
White, both sexes	195	-	-	-	-	-	-	-	1	1	6	4
Male	75	-	-	-	-	-	-	-	1	-	4	2
Female	120	-	-	-	-	-	-	-	-	1	2	2
Black, both sexes	56	-	-	-	-	-	-	-	-	-	1	-
Male	31	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Alaska												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	283	-	-	-	-	-	-	-	-	-	-	-
Male	125	-	-	-	-	-	-	-	-	-	-	-
Female	158	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	277	-	-	-	-	-	-	-	-	-	-	-
Male	122	-	-	-	-	-	-	-	-	-	-	-
Female	155	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Arkansas												
All races, both sexes	94	-	-	-	-	-	-	-	-	-	-	1
Male	45	-	-	-	-	-	-	-	-	-	-	-
Female	49	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	87	-	-	-	-	-	-	-	-	-	-	1
Male	43	-	-	-	-	-	-	-	-	-	-	-
Female	44	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,416	-	-	-	-	-	1	-	-	-	1	6
Male	505	-	-	-	-	-	-	-	-	-	1	3
Female	911	-	-	-	-	-	1	-	-	-	-	3
White, both sexes	1,301	-	-	-	-	-	1	-	-	-	1	5
Male	469	-	-	-	-	-	-	-	-	-	1	2
Female	832	-	-	-	-	-	1	-	-	-	-	3
Black, both sexes	64	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	47	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	31	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Colorado												
All races, both sexes	422	-	-	-	-	-	-	-	-	1	1	1
Male	136	-	-	-	-	-	-	-	-	1	1	1
Female	286	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	402	-	-	-	-	-	-	-	-	1	1	1
Male	130	-	-	-	-	-	-	-	-	1	1	1
Female	272	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	134	-	-	-	-	-	-	-	-	-	-	-
Male	48	-	-	-	-	-	-	-	-	-	-	-
Female	86	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	132	-	-	-	-	-	-	-	-	-	-	-
Male	48	-	-	-	-	-	-	-	-	-	-	-
Female	84	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Delaware												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Florida												
All races, both sexes	855	-	-	-	-	-	-	1	-	-	-	2
Male	357	-	-	-	-	-	-	1	-	-	-	1
Female	498	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	780	-	-	-	-	-	-	-	-	-	-	2
Male	332	-	-	-	-	-	-	-	-	-	-	1
Female	448	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	69	-	-	-	-	-	-	1	-	-	-	-
Male	23	-	-	-	-	-	-	1	-	-	-	-
Female	46	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	245	-	-	-	-	-	-	-	-	-	-	2
Male	94	-	-	-	-	-	-	-	-	-	-	1
Female	151	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	195	-	-	-	-	-	-	-	-	-	-	1
Male	74	-	-	-	-	-	-	-	-	-	-	1
Female	121	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	20	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Hawaii												
All races, both sexes	20	-	-	-	-	-	-	-	-	-	1	-
Male	11	-	-	-	-	-	-	-	-	-	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Illinois												
All races, both sexes	475	-	-	-	-	-	-	-	-	-	-	1
Male	171	-	-	-	-	-	-	-	-	-	-	1
Female	304	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	441	-	-	-	-	-	-	-	-	-	-	-
Male	157	-	-	-	-	-	-	-	-	-	-	-
Female	284	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	337	-	-	-	-	-	-	-	-	-	2	6
Male	130	-	-	-	-	-	-	-	-	-	1	6
Female	207	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	317	-	-	-	-	-	-	-	-	-	2	6
Male	122	-	-	-	-	-	-	-	-	-	1	6
Female	195	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	12	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Iowa												
All races, both sexes	264	-	-	-	-	-	-	-	-	-	1	-
Male	100	-	-	-	-	-	-	-	-	-	1	-
Female	164	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	262	-	-	-	-	-	-	-	-	-	1	-
Male	98	-	-	-	-	-	-	-	-	-	1	-
Female	164	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	363	-	-	-	-	-	-	1	1	-	-	1
Male	122	-	-	-	-	-	-	-	-	-	-	1
Female	241	-	-	-	-	-	-	1	1	-	-	-
White, both sexes	350	-	-	-	-	-	-	1	1	-	-	1
Male	116	-	-	-	-	-	-	-	-	-	-	1
Female	234	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Kentucky												
All races, both sexes	155	-	-	-	-	-	-	-	-	-	-	2
Male	67	-	-	-	-	-	-	-	-	-	-	2
Female	88	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	147	-	-	-	-	-	-	-	-	-	-	2
Male	66	-	-	-	-	-	-	-	-	-	-	2
Female	81	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	186	-	-	-	-	-	-	-	-	1	-	2
Male	60	-	-	-	-	-	-	-	-	1	-	-
Female	126	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	138	-	-	-	-	-	-	-	-	1	-	1
Male	43	-	-	-	-	-	-	-	-	1	-	-
Female	95	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	46	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Maine												
All races, both sexes	44	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	44	-	-	-	-	-	-	-	-	-	-	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	25	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	148	-	-	-	-	-	-	-	-	-	1	-
Male	59	-	-	-	-	-	-	-	-	-	1	-
Female	89	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	127	-	-	-	-	-	-	-	-	-	1	-
Male	52	-	-	-	-	-	-	-	-	-	1	-
Female	75	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	20	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Massachusetts												
All races, both sexes	192	-	-	-	-	-	-	-	-	-	-	1
Male	65	-	-	-	-	-	-	-	-	-	-	1
Female	127	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	187	-	-	-	-	-	-	-	-	-	-	1
Male	63	-	-	-	-	-	-	-	-	-	-	1
Female	124	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	470	-	-	-	-	-	-	-	-	1	2	5
Male	186	-	-	-	-	-	-	-	-	1	-	3
Female	284	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	427	-	-	-	-	-	-	-	-	1	2	5
Male	170	-	-	-	-	-	-	-	-	1	-	3
Female	257	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	39	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	24	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Minnesota												
All races, both sexes	136	-	-	-	-	-	-	-	-	-	-	-
Male	59	-	-	-	-	-	-	-	-	-	-	-
Female	77	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	132	-	-	-	-	-	-	-	-	-	-	-
Male	57	-	-	-	-	-	-	-	-	-	-	-
Female	75	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	48	-	-	-	-	-	-	-	-	1	-	1
Male	14	-	-	-	-	-	-	-	-	1	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	37	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	11	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Missouri												
All races, both sexes	260	-	-	-	-	-	-	-	-	-	-	2
Male	87	-	-	-	-	-	-	-	-	-	-	-
Female	173	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	225	-	-	-	-	-	-	-	-	-	-	1
Male	73	-	-	-	-	-	-	-	-	-	-	-
Female	152	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	34	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Nebraska												
All races, both sexes	106	-	-	-	-	-	-	-	-	-	-	-
Male	45	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	103	-	-	-	-	-	-	-	-	-	-	-
Male	44	-	-	-	-	-	-	-	-	-	-	-
Female	59	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	156	-	-	-	-	-	-	-	-	-	-	-
Male	52	-	-	-	-	-	-	-	-	-	-	-
Female	104	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	147	-	-	-	-	-	-	-	-	-	-	-
Male	49	-	-	-	-	-	-	-	-	-	-	-
Female	98	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
New Hampshire												
All races, both sexes	52	-	-	-	-	-	-	1	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	1	-	-	-	-
Male	18	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	297	-	-	-	-	-	-	-	-	-	2	-
Male	102	-	-	-	-	-	-	-	-	-	-	-
Female	195	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	270	-	-	-	-	-	-	-	-	-	2	-
Male	90	-	-	-	-	-	-	-	-	-	-	-
Female	180	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	18	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
New Mexico												
All races, both sexes	62	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	59	-	-	-	-	-	-	-	-	-	-	-
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	519	-	-	-	-	-	-	-	1	-	1	3
Male	220	-	-	-	-	-	-	-	1	-	-	3
Female	299	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	467	-	-	-	-	-	-	-	1	-	1	3
Male	192	-	-	-	-	-	-	-	1	-	-	3
Female	275	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	47	-	-	-	-	-	-	-	-	-	-	-
Male	24	-	-	-	-	-	-	-	-	-	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
North Carolina												
All races, both sexes	273	-	-	-	-	-	-	-	-	-	-	1
Male	97	-	-	-	-	-	-	-	-	-	-	-
Female	176	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	220	-	-	-	-	-	-	-	-	-	-	-
Male	86	-	-	-	-	-	-	-	-	-	-	-
Female	134	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	52	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	41	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	26	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Ohio												
All races, both sexes	548	-	-	-	-	-	-	-	-	-	-	1
Male	193	-	-	-	-	-	-	-	-	-	-	-
Female	355	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	513	-	-	-	-	-	-	-	-	-	-	1
Male	184	-	-	-	-	-	-	-	-	-	-	-
Female	329	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	35	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	26	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	249	-	-	-	-	-	-	-	-	-	-	-
Male	88	-	-	-	-	-	-	-	-	-	-	-
Female	161	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	223	-	-	-	-	-	-	-	-	-	-	-
Male	79	-	-	-	-	-	-	-	-	-	-	-
Female	144	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Oregon												
All races, both sexes	192	-	-	-	-	-	-	-	-	-	-	1
Male	88	-	-	-	-	-	-	-	-	-	-	1
Female	104	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	188	-	-	-	-	-	-	-	-	-	-	1
Male	88	-	-	-	-	-	-	-	-	-	-	1
Female	100	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	562	-	-	-	-	-	-	-	-	-	1	3
Male	184	-	-	-	-	-	-	-	-	-	1	1
Female	378	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	544	-	-	-	-	-	-	-	-	-	1	2
Male	180	-	-	-	-	-	-	-	-	-	1	1
Female	364	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Rhode Island												
All races, both sexes	61	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	35	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	60	-	-	-	-	-	-	-	-	-	-	-
Male	26	-	-	-	-	-	-	-	-	-	-	-
Female	34	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	68	-	-	-	-	-	1	-	-	-	-	1
Male	25	-	-	-	-	-	1	-	-	-	-	1
Female	43	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	52	-	-	-	-	-	-	-	-	-	-	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	15	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	1	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
South Dakota												
All races, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	270	-	-	-	-	-	-	-	-	-	-	3
Male	109	-	-	-	-	-	-	-	-	-	-	1
Female	161	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	238	-	-	-	-	-	-	-	-	-	-	3
Male	97	-	-	-	-	-	-	-	-	-	-	1
Female	141	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Texas												
All races, both sexes	548	-	1	-	-	-	-	-	-	-	3	7
Male	211	-	1	-	-	-	-	-	-	-	2	6
Female	337	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	493	-	-	-	-	-	-	-	-	-	1	6
Male	184	-	-	-	-	-	-	-	-	-	1	5
Female	309	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	53	-	1	-	-	-	-	-	-	-	2	1
Male	25	-	1	-	-	-	-	-	-	-	1	1
Female	28	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	56	-	-	-	-	-	-	-	-	-	-	3
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	36	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	55	-	-	-	-	-	-	-	-	-	-	3
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	35	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Vermont												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	245	-	-	-	-	-	-	-	-	-	-	-
Male	100	-	-	-	-	-	-	-	-	-	-	-
Female	145	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	214	-	-	-	-	-	-	-	-	-	-	-
Male	88	-	-	-	-	-	-	-	-	-	-	-
Female	126	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	28	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Washington												
All races, both sexes	312	-	-	-	-	-	-	-	-	1	1	2
Male	123	-	-	-	-	-	-	-	-	1	1	2
Female	189	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	299	-	-	-	-	-	-	-	-	1	1	2
Male	115	-	-	-	-	-	-	-	-	1	1	2
Female	184	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	121	-	-	-	-	-	-	-	-	-	-	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	85	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	117	-	-	-	-	-	-	-	-	-	-	-
Male	36	-	-	-	-	-	-	-	-	-	-	-
Female	81	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Atherosclerosis (I70)												
Wisconsin												
All races, both sexes	172	-	-	-	-	-	-	-	-	1	-	1
Male	71	-	-	-	-	-	-	-	-	1	-	1
Female	101	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	169	-	-	-	-	-	-	-	-	1	-	-
Male	69	-	-	-	-	-	-	-	-	1	-	-
Female	100	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
United States												
All races, both sexes	130	167	273	397	611	1,255	2,132	2,710	2,485	1,238	340	-
Male	94	100	169	227	333	624	908	946	709	246	51	-
Female	36	67	104	170	278	631	1,224	1,764	1,776	992	289	-
White, both sexes	94	138	228	334	521	1,127	1,961	2,544	2,340	1,171	301	-
Male	71	83	142	195	283	575	835	902	673	235	44	-
Female	23	55	86	139	238	552	1,126	1,642	1,667	936	257	-
Black, both sexes	30	24	39	57	74	99	139	148	122	63	33	-
Male	18	14	24	32	39	38	57	38	29	9	5	-
Female	12	10	15	25	35	61	82	110	93	54	28	-
American Indian 1/, both sexes	2	5	5	3	9	9	7	5	5	-	-	-
Male	2	3	2	-	6	3	3	2	2	-	-	-
Female	-	2	3	3	3	6	4	3	3	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	1	3	7	20	25	13	18	4	6	-
Male	3	-	1	-	5	8	13	4	5	2	2	-
Female	1	-	-	3	2	12	12	9	13	2	4	-
Alabama												
All races, both sexes	13	11	13	15	23	25	39	46	40	13	2	-
Male	9	9	8	9	13	9	18	12	8	5	-	-
Female	4	2	5	6	10	16	21	34	32	8	2	-
White, both sexes	9	7	7	7	18	16	31	42	34	11	1	-
Male	5	6	3	4	10	6	13	11	6	4	-	-
Female	4	1	4	3	8	10	18	31	28	7	1	-
Black, both sexes	4	4	6	8	5	9	6	4	6	2	1	-
Male	4	3	5	5	3	3	4	1	2	1	-	-
Female	-	1	1	3	2	6	2	3	4	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	2	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Alaska												
All races, both sexes	-	-	1	-	-	2	1	1	1	-	-	-
Male	-	-	-	-	-	1	1	1	1	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	2	1	1	-	-	-	-
Male	-	-	-	-	-	1	1	1	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	2	7	9	20	39	50	64	58	29	3	-
Male	2	2	4	4	10	24	24	23	20	12	-	-
Female	-	-	3	5	10	15	26	41	38	17	3	-
White, both sexes	2	2	7	9	20	36	47	64	58	29	3	-
Male	2	2	4	4	10	22	23	23	20	12	-	-
Female	-	-	3	5	10	14	24	41	38	17	3	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	2	-	-	-	-	-
Male	-	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Arkansas												
All races, both sexes	-	2	-	5	5	10	16	28	18	7	2	-
Male	-	2	-	4	4	6	7	16	4	2	-	-
Female	-	-	-	1	1	4	9	12	14	5	2	-
White, both sexes	-	2	-	4	4	9	14	27	17	7	2	-
Male	-	2	-	3	3	6	7	16	4	2	-	-
Female	-	-	-	1	1	3	7	11	13	5	2	-
Black, both sexes	-	-	-	1	1	1	1	1	1	-	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	12	17	22	35	65	143	268	325	307	161	53	-
Male	7	9	11	18	36	74	104	110	90	32	10	-
Female	5	8	11	17	29	69	164	215	217	129	43	-
White, both sexes	11	15	21	28	58	135	238	301	291	149	47	-
Male	6	8	10	16	33	72	95	102	87	29	8	-
Female	5	7	11	12	25	63	143	199	204	120	39	-
Black, both sexes	-	1	1	3	3	5	13	16	8	9	4	-
Male	-	1	1	2	1	1	3	6	2	2	-	-
Female	-	-	-	1	2	4	10	10	6	7	4	-
American Indian 1/, both sexes	-	1	-	1	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	3	3	3	17	7	8	3	2	-
Male	1	-	-	-	2	1	6	2	1	1	2	-
Female	-	-	-	3	1	2	11	5	7	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Colorado												
All races, both sexes	2	8	6	10	15	50	78	90	99	53	8	-
Male	2	3	4	6	9	22	31	25	23	8	-	-
Female	-	5	2	4	6	28	47	65	76	45	8	-
White, both sexes	2	7	6	9	14	47	73	87	94	52	8	-
Male	2	2	4	5	8	21	29	25	23	8	-	-
Female	-	5	2	4	6	26	44	62	71	44	8	-
Black, both sexes	-	1	-	1	-	2	4	2	3	1	-	-
Male	-	1	-	1	-	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	3	2	3	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	1	-	2	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	2	-	-	-
Connecticut												
All races, both sexes	1	-	1	1	4	12	18	35	42	13	7	-
Male	1	-	-	1	3	5	9	10	13	4	2	-
Female	-	-	1	-	1	7	9	25	29	9	5	-
White, both sexes	1	-	1	1	4	12	17	35	41	13	7	-
Male	1	-	-	1	3	5	9	10	13	4	2	-
Female	-	-	1	-	1	7	8	25	28	9	5	-
Black, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Delaware												
All races, both sexes	-	-	1	-	-	-	4	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	4	1	-	-	-	-
White, both sexes	-	-	1	-	-	-	4	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	4	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	-	1	-	-	-	2	4	6	2	3	-
Male	-	-	-	-	-	-	-	2	1	-	-	-
Female	-	-	1	-	-	-	2	2	5	2	3	-
White, both sexes	-	-	-	-	-	-	1	1	-	1	2	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	1	2	-
Black, both sexes	-	-	1	-	-	-	1	3	6	1	1	-
Male	-	-	-	-	-	-	-	1	1	-	-	-
Female	-	-	1	-	-	-	1	2	5	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Florida												
All races, both sexes	6	14	19	24	36	85	158	207	184	97	22	-
Male	5	5	14	13	19	44	69	90	61	28	7	-
Female	1	9	5	11	17	41	89	117	123	69	15	-
White, both sexes	5	12	16	16	26	75	147	193	174	92	22	-
Male	4	5	12	11	13	41	67	85	58	28	7	-
Female	1	7	4	5	13	34	80	108	116	64	15	-
Black, both sexes	-	2	3	8	10	7	11	13	9	5	-	-
Male	-	-	2	2	6	3	2	5	2	-	-	-
Female	-	2	1	6	4	4	9	8	7	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	3	-	1	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	3	-	1	-	-	-	-
Georgia												
All races, both sexes	8	13	10	14	12	32	45	51	34	17	7	-
Male	5	8	7	10	5	15	17	17	4	3	2	-
Female	3	5	3	4	7	17	28	34	30	14	5	-
White, both sexes	4	9	6	9	10	28	37	45	28	14	4	-
Male	3	5	5	7	4	14	13	16	3	1	2	-
Female	1	4	1	2	6	14	24	29	25	13	2	-
Black, both sexes	4	4	4	5	2	4	8	6	6	3	3	-
Male	2	3	2	3	1	1	4	1	1	2	-	-
Female	2	1	2	2	1	3	4	5	5	1	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Hawaii												
All races, both sexes	1	1	1	-	-	1	4	2	5	1	3	-
Male	-	1	1	-	-	-	4	2	1	1	-	-
Female	1	-	-	-	-	1	-	-	4	-	3	-
White, both sexes	-	1	-	-	-	1	-	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	3	2	4	1	3	-
Male	-	-	1	-	-	-	3	2	1	1	-	-
Female	1	-	-	-	-	-	-	-	3	-	3	-
Idaho												
All races, both sexes	-	-	-	2	-	-	2	8	3	6	2	-
Male	-	-	-	2	-	-	1	4	-	2	1	-
Female	-	-	-	-	-	-	1	4	3	4	1	-
White, both sexes	-	-	-	2	-	-	2	8	3	6	2	-
Male	-	-	-	2	-	-	1	4	-	2	1	-
Female	-	-	-	-	-	-	1	4	3	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Illinois												
All races, both sexes	2	6	12	18	24	32	79	119	116	51	15	-
Male	2	4	6	11	13	19	37	41	30	4	3	-
Female	-	2	6	7	11	13	42	78	86	47	12	-
White, both sexes	1	6	9	15	19	29	73	115	109	50	15	-
Male	1	4	5	9	12	18	32	40	29	4	3	-
Female	-	2	4	6	7	11	41	75	80	46	12	-
Black, both sexes	1	-	3	3	4	2	6	3	7	1	-	-
Male	1	-	1	2	1	-	5	-	1	-	-	-
Female	-	-	2	1	3	2	1	3	6	1	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	2	8	11	15	18	41	59	73	68	28	6	-
Male	1	6	7	8	12	18	26	23	14	7	1	-
Female	1	2	4	7	6	23	33	50	54	21	5	-
White, both sexes	2	7	10	15	16	37	58	69	63	28	4	-
Male	1	5	6	8	10	18	25	23	12	7	-	-
Female	1	2	4	7	6	19	33	46	51	21	4	-
Black, both sexes	-	1	1	-	2	4	1	4	5	-	2	-
Male	-	1	1	-	2	-	1	-	2	-	1	-
Female	-	-	-	-	-	4	-	4	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Iowa												
All races, both sexes	1	1	5	9	7	35	46	60	62	28	9	-
Male	-	1	3	5	4	16	23	20	23	4	-	-
Female	1	-	2	4	3	19	23	40	39	24	9	-
White, both sexes	1	1	5	9	7	35	45	59	62	28	9	-
Male	-	1	3	5	4	16	22	19	23	4	-	-
Female	1	-	2	4	3	19	23	40	39	24	9	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	2	5	7	10	25	75	87	87	46	13	-
Male	3	1	3	4	7	14	27	25	26	10	1	-
Female	-	1	2	3	3	11	48	62	61	36	12	-
White, both sexes	3	2	5	7	10	25	72	82	83	45	13	-
Male	3	1	3	4	7	14	24	24	24	10	1	-
Female	-	1	2	3	3	11	48	58	59	35	12	-
Black, both sexes	-	-	-	-	-	-	3	5	3	1	-	-
Male	-	-	-	-	-	-	3	1	2	-	-	-
Female	-	-	-	-	-	-	-	4	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Kentucky												
All races, both sexes	4	3	8	3	10	21	21	36	30	12	5	-
Male	4	-	6	2	3	13	8	13	11	2	3	-
Female	-	3	2	1	7	8	13	23	19	10	2	-
White, both sexes	4	2	8	3	9	20	20	36	27	12	4	-
Male	4	-	6	2	3	13	8	13	10	2	3	-
Female	-	2	2	1	6	7	12	23	17	10	1	-
Black, both sexes	-	1	-	-	1	1	1	-	3	-	1	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	1	1	1	-	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	3	2	3	8	13	22	30	42	38	16	6	-
Male	2	1	2	3	8	8	13	12	9	-	1	-
Female	1	1	1	5	5	14	17	30	29	16	5	-
White, both sexes	1	1	3	4	7	14	26	32	32	14	2	-
Male	1	1	2	1	5	6	10	8	8	-	-	-
Female	-	-	1	3	2	8	16	24	24	14	2	-
Black, both sexes	2	1	-	4	6	7	3	10	6	2	4	-
Male	1	-	-	2	3	2	2	4	1	-	1	-
Female	1	1	-	2	3	5	1	6	5	2	3	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Maine												
All races, both sexes	-	1	-	4	-	3	7	10	14	4	1	-
Male	-	1	-	3	-	3	3	2	6	1	-	-
Female	-	-	-	1	-	-	4	8	8	3	1	-
White, both sexes	-	1	-	4	-	3	7	10	14	4	1	-
Male	-	1	-	3	-	3	3	2	6	1	-	-
Female	-	-	-	1	-	-	4	8	8	3	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	6	-	2	3	12	15	21	27	29	28	4	-
Male	3	-	2	2	6	8	11	12	10	4	-	-
Female	3	-	-	1	6	7	10	15	19	24	4	-
White, both sexes	1	-	2	3	10	13	17	24	27	26	3	-
Male	1	-	2	2	5	8	9	11	9	4	-	-
Female	-	-	-	1	5	5	8	13	18	22	3	-
Black, both sexes	4	-	-	-	2	2	4	3	2	2	1	-
Male	1	-	-	-	1	-	2	1	1	-	-	-
Female	3	-	-	-	1	2	2	2	1	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Massachusetts												
All races, both sexes	-	1	6	13	7	25	33	41	36	20	9	-
Male	-	1	1	8	4	12	14	14	6	4	-	-
Female	-	-	5	5	3	13	19	27	30	16	9	-
White, both sexes	-	1	6	13	7	22	32	41	36	19	9	-
Male	-	1	1	8	4	10	14	14	6	4	-	-
Female	-	-	5	5	3	12	18	27	30	15	9	-
Black, both sexes	-	-	-	-	-	3	1	-	-	1	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	10	3	16	20	28	51	86	117	83	35	13	-
Male	8	2	10	8	15	22	37	37	31	9	3	-
Female	2	1	6	12	13	29	49	80	52	26	10	-
White, both sexes	6	2	11	14	25	42	83	109	79	35	13	-
Male	6	2	7	6	12	19	35	36	31	9	3	-
Female	-	-	4	8	13	23	48	73	48	26	10	-
Black, both sexes	4	1	4	6	2	8	3	7	4	-	-	-
Male	2	-	3	2	2	3	2	1	-	-	-	-
Female	2	1	1	4	-	5	1	6	4	-	-	-
American Indian 1/, both sexes	-	-	1	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Minnesota												
All races, both sexes	-	2	-	5	4	17	21	33	36	16	2	-
Male	-	2	-	3	3	13	12	9	10	7	-	-
Female	-	-	-	2	1	4	9	24	26	9	2	-
White, both sexes	-	2	-	4	3	16	21	33	35	16	2	-
Male	-	2	-	3	2	12	12	9	10	7	-	-
Female	-	-	-	1	1	4	9	24	25	9	2	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	-	2	6	6	4	4	8	9	6	1	-
Male	-	-	-	3	2	3	2	3	-	-	-	-
Female	-	-	2	3	4	1	2	5	9	6	1	-
White, both sexes	-	-	2	4	5	3	2	6	8	5	1	-
Male	-	-	-	2	2	2	1	2	-	-	-	-
Female	-	-	2	2	3	1	1	4	8	5	1	-
Black, both sexes	-	-	-	2	1	1	2	2	1	1	-	-
Male	-	-	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	1	1	-	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Missouri												
All races, both sexes	-	3	7	13	11	25	38	67	58	25	11	-
Male	-	1	4	6	5	10	19	19	17	3	3	-
Female	-	2	3	7	6	15	19	48	41	22	8	-
White, both sexes	-	3	4	12	8	22	32	59	54	21	9	-
Male	-	1	2	6	4	9	16	16	15	2	2	-
Female	-	2	2	6	4	13	16	43	39	19	7	-
Black, both sexes	-	-	3	1	3	3	6	7	4	4	2	-
Male	-	-	2	-	1	1	3	2	2	1	1	-
Female	-	-	1	1	2	2	3	5	2	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	1	1	3	2	5	9	7	3	-	-
Male	-	-	-	-	2	1	2	4	5	1	-	-
Female	-	-	1	1	1	1	3	5	2	2	-	-
White, both sexes	-	-	-	1	3	2	5	9	7	3	-	-
Male	-	-	-	-	2	1	2	4	5	1	-	-
Female	-	-	-	1	1	1	3	5	2	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Nebraska												
All races, both sexes	-	1	1	2	10	9	22	17	29	13	2	-
Male	-	1	1	1	8	5	6	9	8	6	-	-
Female	-	-	-	1	2	4	16	8	21	7	2	-
White, both sexes	-	1	1	2	9	9	22	16	29	12	2	-
Male	-	1	1	1	7	5	6	9	8	6	-	-
Female	-	-	-	1	2	4	16	7	21	6	2	-
Black, both sexes	-	-	-	-	1	-	-	1	-	1	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	3	3	4	5	12	24	34	38	22	9	2	-
Male	3	2	3	-	4	11	11	13	3	1	1	-
Female	-	1	1	5	8	13	23	25	19	8	1	-
White, both sexes	3	3	3	5	10	22	34	37	19	9	2	-
Male	3	2	3	-	3	9	11	13	3	1	1	-
Female	-	1	-	5	7	13	23	24	16	8	1	-
Black, both sexes	-	-	1	-	-	2	-	1	1	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
New Hampshire												
All races, both sexes	-	-	-	1	-	6	6	10	12	13	3	-
Male	-	-	-	1	-	3	2	3	4	5	-	-
Female	-	-	-	-	-	3	4	7	8	8	3	-
White, both sexes	-	-	-	1	-	6	6	10	12	13	3	-
Male	-	-	-	1	-	3	2	3	4	5	-	-
Female	-	-	-	-	-	3	4	7	8	8	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	4	4	3	9	17	36	58	61	61	32	10	-
Male	3	2	3	6	11	17	27	19	11	3	-	-
Female	1	2	-	3	6	19	31	42	50	29	10	-
White, both sexes	2	3	3	8	16	28	54	57	58	31	8	-
Male	1	1	3	5	10	13	25	18	11	3	-	-
Female	1	2	-	3	6	15	29	39	47	28	8	-
Black, both sexes	2	1	-	1	-	3	3	3	3	1	1	-
Male	2	1	-	1	-	3	1	1	-	-	-	-
Female	-	-	-	-	-	-	2	2	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	4	1	1	-	-	1	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	3	-	1	-	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
New Mexico												
All races, both sexes	1	-	3	2	2	5	16	13	11	7	2	-
Male	1	-	3	1	1	2	10	5	7	2	-	-
Female	-	-	-	1	1	3	6	8	4	5	2	-
White, both sexes	1	-	2	2	2	5	15	13	11	6	2	-
Male	1	-	2	1	1	2	9	5	7	1	-	-
Female	-	-	-	1	1	3	6	8	4	5	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	1	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	9	9	9	15	34	49	91	108	102	68	20	-
Male	7	5	7	10	21	27	40	47	39	12	1	-
Female	2	4	2	5	13	22	51	61	63	56	19	-
White, both sexes	7	7	7	13	23	41	84	103	97	64	16	-
Male	6	4	6	8	12	23	36	45	36	12	-	-
Female	1	3	1	5	11	18	48	58	61	52	16	-
Black, both sexes	2	2	2	2	10	6	6	4	5	4	4	-
Male	1	1	1	2	8	2	3	2	3	-	1	-
Female	1	1	1	-	2	4	3	2	2	4	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	2	1	1	-	-	-	-
Male	-	-	-	-	1	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
North Carolina												
All races, both sexes	3	1	7	8	21	26	57	79	45	19	6	-
Male	1	-	3	4	12	13	20	27	14	2	1	-
Female	2	1	4	4	9	13	37	52	31	17	5	-
White, both sexes	1	1	6	6	15	24	42	64	41	15	5	-
Male	1	-	2	3	9	11	18	26	13	2	1	-
Female	-	1	4	3	6	13	24	38	28	13	4	-
Black, both sexes	2	-	1	2	5	2	15	15	4	4	1	-
Male	-	-	1	1	3	2	2	1	1	-	-	-
Female	2	-	-	1	2	-	13	14	3	4	1	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	-	2	4	5	12	3	-	-
Male	-	-	-	-	-	1	2	3	-	1	-	-
Female	-	-	-	-	-	1	2	2	12	2	-	-
White, both sexes	-	-	-	-	-	2	4	5	12	3	-	-
Male	-	-	-	-	-	1	2	3	-	1	-	-
Female	-	-	-	-	-	1	2	2	12	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Ohio												
All races, both sexes	10	5	15	9	23	60	94	153	108	55	15	-
Male	6	2	10	5	14	26	41	53	23	11	2	-
Female	4	3	5	4	9	34	53	100	85	44	13	-
White, both sexes	8	5	14	9	21	56	90	144	100	52	13	-
Male	5	2	10	5	14	24	39	52	21	11	1	-
Female	3	3	4	4	7	32	51	92	79	41	12	-
Black, both sexes	2	-	1	-	2	4	4	9	8	3	2	-
Male	1	-	-	-	-	2	2	1	2	-	1	-
Female	1	-	1	-	2	2	2	8	6	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	7	5	7	14	38	44	55	52	22	3	-
Male	2	3	3	4	6	15	24	15	13	3	-	-
Female	-	4	2	3	8	23	20	40	39	19	3	-
White, both sexes	-	4	3	7	11	31	42	51	51	20	3	-
Male	-	2	2	4	5	11	24	15	13	3	-	-
Female	-	2	1	3	6	20	18	36	38	17	3	-
Black, both sexes	1	1	-	-	2	5	1	4	1	2	-	-
Male	1	-	-	-	-	3	-	-	-	-	-	-
Female	-	1	-	-	2	2	1	4	1	2	-	-
American Indian 1/, both sexes	1	2	2	-	1	2	1	-	-	-	-	-
Male	1	1	1	-	1	1	-	-	-	-	-	-
Female	-	1	1	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Oregon												
All races, both sexes	-	4	5	10	13	24	32	46	30	24	3	-
Male	-	1	3	6	8	11	17	24	10	6	1	-
Female	-	3	2	4	5	13	15	22	20	18	2	-
White, both sexes	-	4	5	9	13	23	32	46	30	22	3	-
Male	-	1	3	6	8	11	17	24	10	6	1	-
Female	-	3	2	3	5	12	15	22	20	16	2	-
Black, both sexes	-	-	-	1	-	-	-	-	-	2	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	3	4	10	10	26	53	98	128	140	68	18	-
Male	3	3	4	5	13	25	26	42	44	16	1	-
Female	-	1	6	5	13	28	72	86	96	52	17	-
White, both sexes	3	3	9	10	23	52	93	125	140	66	17	-
Male	3	3	4	5	13	25	24	40	44	16	1	-
Female	-	-	5	5	10	27	69	85	96	50	16	-
Black, both sexes	-	1	1	-	3	1	4	3	-	2	1	-
Male	-	-	-	-	-	-	2	2	-	-	-	-
Female	-	1	1	-	3	1	2	1	-	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Rhode Island												
All races, both sexes	-	-	-	2	5	5	11	11	20	7	-	-
Male	-	-	-	1	3	2	8	6	5	1	-	-
Female	-	-	-	1	2	3	3	5	15	6	-	-
White, both sexes	-	-	-	2	5	4	11	11	20	7	-	-
Male	-	-	-	1	3	2	8	6	5	1	-	-
Female	-	-	-	1	2	2	3	5	15	6	-	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	-	1	2	5	5	9	15	11	12	6	-	-
Male	-	1	-	4	3	3	8	2	2	-	-	-
Female	-	-	2	1	2	6	7	9	10	6	-	-
White, both sexes	-	-	2	4	4	8	10	10	7	6	-	-
Male	-	-	-	3	2	3	6	2	2	-	-	-
Female	-	-	2	1	2	5	4	8	5	6	-	-
Black, both sexes	-	-	-	1	1	1	5	1	5	-	-	-
Male	-	-	-	1	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	1	3	1	5	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
South Dakota												
All races, both sexes	1	1	-	-	-	1	5	9	11	5	1	-
Male	1	1	-	-	-	-	3	4	5	1	-	-
Female	-	-	-	-	-	1	2	5	6	4	1	-
White, both sexes	-	1	-	-	-	1	5	9	11	5	1	-
Male	-	1	-	-	-	-	3	4	5	1	-	-
Female	-	-	-	-	-	1	2	5	6	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	5	8	8	10	22	27	54	62	43	23	5	-
Male	4	3	7	8	8	13	29	19	12	4	1	-
Female	1	5	1	2	14	14	25	43	31	19	4	-
White, both sexes	5	8	6	6	21	23	49	54	37	21	5	-
Male	4	3	6	5	7	13	27	17	9	4	1	-
Female	1	5	-	1	14	10	22	37	28	17	4	-
Black, both sexes	-	-	2	4	1	4	5	8	5	2	-	-
Male	-	-	1	3	1	-	2	2	2	-	-	-
Female	-	-	1	1	-	4	3	6	3	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Texas												
All races, both sexes	4	7	17	22	38	66	93	106	115	59	10	-
Male	2	7	10	13	21	36	35	36	34	7	1	-
Female	2	-	7	9	17	30	58	70	81	52	9	-
White, both sexes	3	5	13	20	33	59	84	98	110	53	8	-
Male	1	5	6	12	19	33	30	32	33	6	1	-
Female	2	-	7	8	14	26	54	66	77	47	7	-
Black, both sexes	1	2	4	2	4	6	9	8	5	6	2	-
Male	1	2	4	1	1	2	5	4	1	1	-	-
Female	-	-	-	1	3	4	4	4	4	5	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	1	1	-	5	2	6	12	11	7	6	2	-
Male	1	1	-	1	2	3	7	3	-	-	-	-
Female	-	-	-	4	-	3	5	8	7	6	2	-
White, both sexes	1	1	-	4	2	6	12	11	7	6	2	-
Male	1	1	-	1	2	3	7	3	-	-	-	-
Female	-	-	-	3	-	3	5	8	7	6	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Vermont												
All races, both sexes	1	1	2	1	-	2	1	1	5	1	1	-
Male	-	-	2	1	-	2	-	-	1	-	-	-
Female	1	1	-	-	-	-	1	1	4	1	1	-
White, both sexes	1	1	2	1	-	2	1	1	5	1	1	-
Male	-	-	2	1	-	2	-	-	1	-	-	-
Female	1	1	-	-	-	-	1	1	4	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	2	2	10	15	13	26	56	44	48	23	6	-
Male	1	2	7	12	5	13	23	18	15	4	-	-
Female	1	-	3	3	8	13	33	26	33	19	6	-
White, both sexes	1	2	9	13	11	22	48	41	40	22	5	-
Male	-	2	7	10	4	10	20	18	13	4	-	-
Female	1	-	2	3	7	12	28	23	27	18	5	-
Black, both sexes	1	-	1	2	2	1	8	3	8	1	1	-
Male	1	-	-	2	1	1	3	-	2	-	-	-
Female	-	-	1	-	1	-	5	3	6	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	3	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Washington												
All races, both sexes	2	2	7	12	11	27	63	81	67	24	12	-
Male	2	1	4	7	5	16	28	31	19	3	3	-
Female	-	1	3	5	6	11	35	50	48	21	9	-
White, both sexes	2	1	7	12	9	23	62	79	65	23	12	-
Male	2	-	4	7	3	14	28	31	17	2	3	-
Female	-	1	3	5	6	9	34	48	48	21	9	-
Black, both sexes	-	-	-	-	1	2	1	1	-	1	-	-
Male	-	-	-	-	1	1	-	-	-	1	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	1	2	-	-	-	-	-	-
Male	-	1	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	2	-	-	-
Male	-	-	-	-	-	-	-	-	2	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
West Virginia												
All races, both sexes	-	2	1	2	5	15	29	25	27	12	3	-
Male	-	2	1	1	1	6	7	9	8	-	1	-
Female	-	-	-	1	4	9	22	16	19	12	2	-
White, both sexes	-	2	1	2	5	14	29	25	25	12	2	-
Male	-	2	1	1	1	6	7	9	8	-	1	-
Female	-	-	-	1	4	8	22	16	17	12	1	-
Black, both sexes	-	-	-	-	-	-	-	-	2	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Atherosclerosis (I70)												
Wisconsin												
All races, both sexes	3	4	4	5	5	21	24	42	34	20	8	-
Male	3	4	4	3	4	13	14	11	8	4	1	-
Female	-	-	-	2	1	8	10	31	26	16	7	-
White, both sexes	3	3	4	5	5	21	24	42	34	20	7	-
Male	3	3	4	3	4	13	14	11	8	4	1	-
Female	-	-	-	2	1	8	10	31	26	16	6	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	1	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	-	-	1	3	3	2	2	1	-
Male	-	-	-	-	-	1	1	1	-	1	-	-
Female	-	-	-	-	-	-	2	2	2	1	1	-
White, both sexes	-	-	-	-	-	1	3	3	2	2	1	-
Male	-	-	-	-	-	1	1	1	-	1	-	-
Female	-	-	-	-	-	-	2	2	2	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
United States												
All races, both sexes	13,843	1	-	-	4	13	21	47	67	115	214	323
Male	8,168	1	-	-	4	10	16	28	55	94	171	256
Female	5,675	-	-	-	-	3	5	19	12	21	43	67
White, both sexes	12,156	1	-	-	3	12	19	30	46	66	147	216
Male	7,203	1	-	-	3	9	14	16	37	52	116	176
Female	4,953	-	-	-	-	3	5	14	9	14	31	40
Black, both sexes	1,304	-	-	-	1	1	2	14	15	44	54	92
Male	719	-	-	-	1	1	2	9	12	38	43	69
Female	585	-	-	-	-	-	-	5	3	6	11	23
American Indian 1/, both sexes	43	-	-	-	-	-	-	1	-	-	1	1
Male	21	-	-	-	-	-	-	1	-	-	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	340	-	-	-	-	-	-	2	6	5	12	14
Male	225	-	-	-	-	-	-	2	6	4	11	11
Female	115	-	-	-	-	-	-	-	-	1	1	3
Alabama												
All races, both sexes	208	-	-	-	-	-	-	-	-	3	1	7
Male	132	-	-	-	-	-	-	-	-	3	1	6
Female	76	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	171	-	-	-	-	-	-	-	-	1	-	5
Male	110	-	-	-	-	-	-	-	-	1	-	4
Female	61	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	35	-	-	-	-	-	-	-	-	2	1	2
Male	22	-	-	-	-	-	-	-	-	2	1	2
Female	13	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Alaska												
All races, both sexes	12	-	-	-	-	-	-	-	-	1	1	-
Male	9	-	-	-	-	-	-	-	-	1	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	229	-	-	-	-	1	-	-	1	-	-	8
Male	146	-	-	-	-	1	-	-	1	-	-	7
Female	83	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	219	-	-	-	-	1	-	-	1	-	-	8
Male	138	-	-	-	-	1	-	-	1	-	-	7
Female	81	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Arkansas												
All races, both sexes	161	-	-	-	-	-	1	-	1	3	2	4
Male	83	-	-	-	-	-	1	-	1	3	2	1
Female	78	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	149	-	-	-	-	-	1	-	-	2	2	4
Male	77	-	-	-	-	-	1	-	-	2	2	1
Female	72	-	-	-	-	-	-	-	-	-	-	3
Black, both sexes	12	-	-	-	-	-	-	-	1	1	-	-
Male	6	-	-	-	-	-	-	-	1	1	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	1,241	-	-	-	-	-	2	5	6	8	28	44
Male	764	-	-	-	-	-	2	3	6	7	23	36
Female	477	-	-	-	-	-	-	2	-	1	5	8
White, both sexes	1,000	-	-	-	-	-	1	3	4	4	19	26
Male	616	-	-	-	-	-	1	1	4	3	14	20
Female	384	-	-	-	-	-	-	2	-	1	5	6
Black, both sexes	91	-	-	-	-	-	1	1	1	3	4	9
Male	53	-	-	-	-	-	1	1	1	3	4	9
Female	38	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	144	-	-	-	-	-	-	1	1	1	5	9
Male	91	-	-	-	-	-	-	1	1	1	5	7
Female	53	-	-	-	-	-	-	-	-	-	-	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Colorado												
All races, both sexes	191	-	-	-	-	-	-	1	1	1	4	4
Male	109	-	-	-	-	-	-	1	-	1	3	3
Female	82	-	-	-	-	-	-	-	1	-	1	1
White, both sexes	178	-	-	-	-	-	-	-	1	1	3	4
Male	100	-	-	-	-	-	-	-	-	1	2	3
Female	78	-	-	-	-	-	-	-	1	-	1	1
Black, both sexes	10	-	-	-	-	-	-	1	-	-	1	-
Male	7	-	-	-	-	-	-	1	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	181	-	-	-	-	-	-	-	1	2	-	2
Male	93	-	-	-	-	-	-	-	1	2	-	2
Female	88	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	175	-	-	-	-	-	-	-	1	1	-	2
Male	90	-	-	-	-	-	-	-	1	1	-	2
Female	85	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Delaware												
All races, both sexes	45	-	-	-	-	-	-	1	-	-	-	3
Male	27	-	-	-	-	-	-	1	-	-	-	2
Female	18	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	40	-	-	-	-	-	-	1	-	-	-	3
Male	25	-	-	-	-	-	-	1	-	-	-	2
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	35	-	-	-	-	-	-	-	-	1	1	-
Male	21	-	-	-	-	-	-	-	-	1	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	34	-	-	-	-	-	-	-	-	1	1	-
Male	20	-	-	-	-	-	-	-	-	1	1	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Florida												
All races, both sexes	1,006	-	-	-	-	-	1	2	3	9	12	15
Male	604	-	-	-	-	-	1	1	3	8	10	14
Female	402	-	-	-	-	-	-	1	-	1	2	1
White, both sexes	899	-	-	-	-	-	1	-	2	5	7	8
Male	544	-	-	-	-	-	1	-	2	4	6	8
Female	355	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	95	-	-	-	-	-	-	2	1	4	5	7
Male	52	-	-	-	-	-	-	1	1	4	4	6
Female	43	-	-	-	-	-	-	1	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	331	-	-	-	-	-	-	2	1	5	10	14
Male	197	-	-	-	-	-	-	1	1	4	8	11
Female	134	-	-	-	-	-	-	1	-	1	2	3
White, both sexes	256	-	-	-	-	-	-	-	-	2	4	7
Male	149	-	-	-	-	-	-	-	-	1	3	7
Female	107	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	73	-	-	-	-	-	-	2	-	3	6	7
Male	46	-	-	-	-	-	-	1	-	3	5	4
Female	27	-	-	-	-	-	-	1	-	-	1	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Hawaii												
All races, both sexes	81	-	-	-	-	-	-	1	-	2	-	1
Male	50	-	-	-	-	-	-	1	-	2	-	1
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	18	-	-	-	-	-	-	-	-	-	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	63	-	-	-	-	-	-	1	-	2	-	-
Male	40	-	-	-	-	-	-	1	-	2	-	-
Female	23	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	78	-	-	-	-	-	-	-	-	-	-	2
Male	47	-	-	-	-	-	-	-	-	-	-	2
Female	31	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	74	-	-	-	-	-	-	-	-	-	-	2
Male	45	-	-	-	-	-	-	-	-	-	-	2
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Illinois												
All races, both sexes	595	1	-	-	-	-	2	-	2	7	10	17
Male	329	1	-	-	-	-	2	-	1	6	10	15
Female	266	-	-	-	-	-	-	-	1	1	-	2
White, both sexes	502	1	-	-	-	-	2	-	1	3	6	11
Male	274	1	-	-	-	-	2	-	1	3	6	10
Female	228	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	80	-	-	-	-	-	-	-	1	3	4	5
Male	47	-	-	-	-	-	-	-	-	3	4	4
Female	33	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	13	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	1	-	-
Indiana												
All races, both sexes	329	-	-	-	-	-	1	5	3	3	8	5
Male	204	-	-	-	-	-	1	2	3	3	4	5
Female	125	-	-	-	-	-	-	3	-	-	4	-
White, both sexes	310	-	-	-	-	-	1	3	3	1	8	5
Male	192	-	-	-	-	-	1	1	3	1	4	5
Female	118	-	-	-	-	-	-	2	-	-	4	-
Black, both sexes	19	-	-	-	-	-	-	2	-	2	-	-
Male	12	-	-	-	-	-	-	1	-	2	-	-
Female	7	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Iowa												
All races, both sexes	211	-	-	-	-	-	-	1	-	1	1	3
Male	130	-	-	-	-	-	-	-	-	1	-	2
Female	81	-	-	-	-	-	-	1	-	-	1	1
White, both sexes	206	-	-	-	-	-	-	1	-	1	1	3
Male	127	-	-	-	-	-	-	-	-	1	-	2
Female	79	-	-	-	-	-	-	1	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	149	-	-	-	-	-	-	-	-	-	2	1
Male	89	-	-	-	-	-	-	-	-	-	1	1
Female	60	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	142	-	-	-	-	-	-	-	-	-	2	1
Male	87	-	-	-	-	-	-	-	-	-	1	1
Female	55	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Kentucky												
All races, both sexes	205	-	-	-	1	1	-	-	1	-	1	1
Male	119	-	-	-	1	1	-	-	1	-	1	1
Female	86	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	194	-	-	-	1	1	-	-	1	-	1	1
Male	113	-	-	-	1	1	-	-	1	-	1	1
Female	81	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	197	-	-	-	1	-	1	1	-	5	6	9
Male	105	-	-	-	1	-	1	1	-	3	5	5
Female	92	-	-	-	-	-	-	-	-	2	1	4
White, both sexes	148	-	-	-	-	-	-	-	-	4	3	2
Male	78	-	-	-	-	-	-	-	-	3	3	2
Female	70	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	48	-	-	-	1	-	1	1	-	1	2	7
Male	26	-	-	-	1	-	1	1	-	-	1	3
Female	22	-	-	-	-	-	-	-	-	1	1	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Maine												
All races, both sexes	95	-	-	-	-	-	-	-	-	-	1	-
Male	46	-	-	-	-	-	-	-	-	-	1	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	95	-	-	-	-	-	-	-	-	-	1	-
Male	46	-	-	-	-	-	-	-	-	-	1	-
Female	49	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	248	-	-	-	-	-	-	-	1	5	4	8
Male	127	-	-	-	-	-	-	-	1	5	4	6
Female	121	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	203	-	-	-	-	-	-	-	1	3	2	2
Male	102	-	-	-	-	-	-	-	1	3	2	1
Female	101	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	40	-	-	-	-	-	-	-	-	2	2	6
Male	21	-	-	-	-	-	-	-	-	2	2	5
Female	19	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Massachusetts												
All races, both sexes	314	-	-	-	-	-	-	-	2	2	1	1
Male	163	-	-	-	-	-	-	-	2	2	-	1
Female	151	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	297	-	-	-	-	-	-	-	2	2	1	1
Male	154	-	-	-	-	-	-	-	2	2	-	1
Female	143	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	568	-	-	-	-	3	-	6	3	3	9	17
Male	328	-	-	-	-	2	-	3	2	3	6	13
Female	240	-	-	-	-	1	-	3	1	-	3	4
White, both sexes	493	-	-	-	-	3	-	5	2	1	4	10
Male	288	-	-	-	-	2	-	3	1	1	3	9
Female	205	-	-	-	-	1	-	2	1	-	1	1
Black, both sexes	66	-	-	-	-	-	-	1	1	2	4	7
Male	34	-	-	-	-	-	-	-	1	2	2	4
Female	32	-	-	-	-	-	-	1	-	-	2	3
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Minnesota												
All races, both sexes	284	-	-	-	-	-	1	1	-	1	7	1
Male	172	-	-	-	-	-	1	1	-	1	5	1
Female	112	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	277	-	-	-	-	-	1	1	-	-	5	1
Male	166	-	-	-	-	-	1	1	-	-	4	1
Female	111	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	2	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Mississippi												
All races, both sexes	138	-	-	-	-	-	1	-	3	-	3	2
Male	84	-	-	-	-	-	1	-	2	-	-	2
Female	54	-	-	-	-	-	-	-	1	-	3	-
White, both sexes	100	-	-	-	-	-	1	-	1	-	2	-
Male	64	-	-	-	-	-	1	-	1	-	-	-
Female	36	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	36	-	-	-	-	-	-	-	2	-	1	2
Male	18	-	-	-	-	-	-	-	1	-	-	2
Female	18	-	-	-	-	-	-	-	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Missouri												
All races, both sexes	305	-	-	-	-	-	1	-	1	1	11	6
Male	188	-	-	-	-	-	1	-	1	-	9	5
Female	117	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	284	-	-	-	-	-	1	-	1	-	11	6
Male	178	-	-	-	-	-	1	-	1	-	9	5
Female	106	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	19	-	-	-	-	-	-	-	-	1	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	48	-	-	-	-	-	-	-	-	1	-	-
Male	34	-	-	-	-	-	-	-	-	1	-	-
Female	14	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	45	-	-	-	-	-	-	-	-	1	-	-
Male	34	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Nebraska												
All races, both sexes	81	-	-	-	-	-	-	-	-	1	-	1
Male	45	-	-	-	-	-	-	-	-	-	-	1
Female	36	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	78	-	-	-	-	-	-	-	-	1	-	1
Male	43	-	-	-	-	-	-	-	-	-	-	1
Female	35	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	95	-	-	-	-	-	-	-	-	-	3	1
Male	51	-	-	-	-	-	-	-	-	-	3	1
Female	44	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	91	-	-	-	-	-	-	-	-	-	3	1
Male	49	-	-	-	-	-	-	-	-	-	3	1
Female	42	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
New Hampshire												
All races, both sexes	64	-	-	-	-	-	-	-	-	1	1	2
Male	31	-	-	-	-	-	-	-	-	1	1	1
Female	33	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	64	-	-	-	-	-	-	-	-	1	1	2
Male	31	-	-	-	-	-	-	-	-	1	1	1
Female	33	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	408	-	-	-	-	-	-	2	1	4	6	6
Male	239	-	-	-	-	-	-	-	1	4	5	6
Female	169	-	-	-	-	-	-	2	-	-	1	-
White, both sexes	360	-	-	-	-	-	-	2	-	3	3	5
Male	211	-	-	-	-	-	-	-	-	3	3	5
Female	149	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	38	-	-	-	-	-	-	-	-	1	3	1
Male	20	-	-	-	-	-	-	-	-	1	2	1
Female	18	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
New Mexico												
All races, both sexes	89	-	-	-	-	-	-	-	-	-	1	1
Male	55	-	-	-	-	-	-	-	-	-	1	1
Female	34	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	86	-	-	-	-	-	-	-	-	-	1	1
Male	53	-	-	-	-	-	-	-	-	-	1	1
Female	33	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	870	-	-	-	-	-	2	1	3	4	19	28
Male	499	-	-	-	-	-	2	1	2	3	17	26
Female	371	-	-	-	-	-	-	-	1	1	2	2
White, both sexes	746	-	-	-	-	-	2	1	3	2	14	17
Male	419	-	-	-	-	-	2	1	2	1	13	16
Female	327	-	-	-	-	-	-	-	1	1	1	1
Black, both sexes	107	-	-	-	-	-	-	-	-	2	3	9
Male	67	-	-	-	-	-	-	-	-	2	2	8
Female	40	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	16	-	-	-	-	-	-	-	-	-	2	2
Male	12	-	-	-	-	-	-	-	-	-	2	2
Female	4	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
North Carolina												
All races, both sexes	391	-	-	-	-	1	1	1	2	-	9	11
Male	251	-	-	-	-	1	-	1	1	-	6	8
Female	140	-	-	-	-	-	1	-	1	-	3	3
White, both sexes	325	-	-	-	-	1	1	1	1	-	6	9
Male	218	-	-	-	-	1	-	1	-	-	5	7
Female	107	-	-	-	-	-	1	-	1	-	1	2
Black, both sexes	61	-	-	-	-	-	-	-	1	-	3	1
Male	32	-	-	-	-	-	-	-	1	-	1	1
Female	29	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
North Dakota												
All races, both sexes	42	-	-	-	-	-	-	-	1	-	-	-
Male	27	-	-	-	-	-	-	-	1	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	42	-	-	-	-	-	-	-	1	-	-	-
Male	27	-	-	-	-	-	-	-	1	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Ohio												
All races, both sexes	664	-	-	-	-	1	1	2	5	6	7	15
Male	393	-	-	-	-	1	1	2	4	5	4	12
Female	271	-	-	-	-	-	-	-	1	1	3	3
White, both sexes	588	-	-	-	-	1	1	-	3	4	4	7
Male	358	-	-	-	-	1	1	-	2	3	1	6
Female	230	-	-	-	-	-	-	-	1	1	3	1
Black, both sexes	74	-	-	-	-	-	-	2	2	2	3	8
Male	35	-	-	-	-	-	-	2	2	2	3	6
Female	39	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	193	-	-	-	-	1	-	3	4	-	2	7
Male	124	-	-	-	-	1	-	2	2	-	2	4
Female	69	-	-	-	-	-	-	1	2	-	-	3
White, both sexes	178	-	-	-	-	1	-	2	4	-	1	4
Male	115	-	-	-	-	1	-	1	2	-	1	3
Female	63	-	-	-	-	-	-	1	2	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	2
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	8	-	-	-	-	-	-	1	-	-	1	1
Male	5	-	-	-	-	-	-	1	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Oregon												
All races, both sexes	178	-	-	-	-	-	-	-	-	-	-	3
Male	113	-	-	-	-	-	-	-	-	-	-	3
Female	65	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	173	-	-	-	-	-	-	-	-	-	-	3
Male	110	-	-	-	-	-	-	-	-	-	-	3
Female	63	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	777	-	-	-	-	1	-	1	4	9	5	14
Male	412	-	-	-	-	1	-	-	3	8	4	12
Female	365	-	-	-	-	-	-	1	1	1	1	2
White, both sexes	709	-	-	-	-	-	-	1	4	6	4	10
Male	374	-	-	-	-	-	-	-	3	5	3	9
Female	335	-	-	-	-	-	-	1	1	1	1	1
Black, both sexes	65	-	-	-	-	1	-	-	-	3	-	4
Male	36	-	-	-	-	1	-	-	-	3	-	3
Female	29	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Rhode Island												
All races, both sexes	61	-	-	-	-	1	-	-	-	-	2	2
Male	33	-	-	-	-	1	-	-	-	-	1	-
Female	28	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	58	-	-	-	-	1	-	-	-	-	2	2
Male	31	-	-	-	-	1	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	229	-	-	-	-	-	1	-	4	-	4	3
Male	132	-	-	-	-	-	1	-	4	-	4	3
Female	97	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	178	-	-	-	-	-	1	-	-	-	3	2
Male	106	-	-	-	-	-	1	-	-	-	3	2
Female	72	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	49	-	-	-	-	-	-	-	4	-	1	1
Male	25	-	-	-	-	-	-	-	4	-	1	1
Female	24	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
South Dakota												
All races, both sexes	51	-	-	-	-	-	-	-	-	-	-	1
Male	34	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	50	-	-	-	-	-	-	-	-	-	-	1
Male	33	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	275	-	-	-	-	-	1	1	2	3	7	8
Male	163	-	-	-	-	-	1	-	2	3	6	5
Female	112	-	-	-	-	-	-	1	-	-	1	3
White, both sexes	242	-	-	-	-	-	1	-	2	2	5	6
Male	148	-	-	-	-	-	1	-	2	2	4	4
Female	94	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	31	-	-	-	-	-	-	1	-	1	2	2
Male	13	-	-	-	-	-	-	-	-	1	2	1
Female	18	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Texas												
All races, both sexes	703	-	-	-	-	3	2	7	6	11	13	19
Male	431	-	-	-	-	1	-	5	5	4	12	14
Female	272	-	-	-	-	2	2	2	1	7	1	5
White, both sexes	606	-	-	-	-	3	2	6	5	7	7	12
Male	383	-	-	-	-	1	-	4	4	3	7	9
Female	223	-	-	-	-	2	2	2	1	4	-	3
Black, both sexes	88	-	-	-	-	-	-	1	-	4	6	7
Male	44	-	-	-	-	-	-	1	-	1	5	5
Female	44	-	-	-	-	-	-	-	-	3	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	73	-	-	-	-	-	-	1	-	-	-	2
Male	49	-	-	-	-	-	-	1	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	69	-	-	-	-	-	-	1	-	-	-	2
Male	45	-	-	-	-	-	-	1	-	-	-	1
Female	24	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Vermont												
All races, both sexes	50	-	-	-	-	-	-	-	-	1	1	-
Male	33	-	-	-	-	-	-	-	-	1	1	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	50	-	-	-	-	-	-	-	-	1	1	-
Male	33	-	-	-	-	-	-	-	-	1	1	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	283	-	-	-	-	-	1	1	3	4	4	6
Male	164	-	-	-	-	-	-	1	2	2	4	3
Female	119	-	-	-	-	-	1	-	1	2	-	3
White, both sexes	235	-	-	-	-	-	1	1	1	2	3	3
Male	136	-	-	-	-	-	-	1	1	1	3	2
Female	99	-	-	-	-	-	1	-	-	1	-	1
Black, both sexes	44	-	-	-	-	-	-	-	1	2	1	3
Male	24	-	-	-	-	-	-	-	-	1	1	1
Female	20	-	-	-	-	-	-	-	1	1	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Washington												
All races, both sexes	318	-	-	-	1	-	-	-	2	4	3	10
Male	204	-	-	-	1	-	-	-	2	4	3	6
Female	114	-	-	-	-	-	-	-	-	-	-	4
White, both sexes	292	-	-	-	1	-	-	-	1	2	2	8
Male	184	-	-	-	1	-	-	-	1	2	2	4
Female	108	-	-	-	-	-	-	-	-	-	-	4
Black, both sexes	12	-	-	-	-	-	-	-	-	2	1	1
Male	11	-	-	-	-	-	-	-	-	2	1	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	1	-	-	1
Male	7	-	-	-	-	-	-	-	1	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	111	-	-	-	-	-	-	-	-	-	1	2
Male	71	-	-	-	-	-	-	-	-	-	1	2
Female	40	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	107	-	-	-	-	-	-	-	-	-	1	1
Male	70	-	-	-	-	-	-	-	-	-	1	1
Female	37	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Aortic aneurysm and dissection (I71)												
Wisconsin												
All races, both sexes	316	-	-	-	-	-	1	1	-	3	2	6
Male	189	-	-	-	-	-	-	-	-	2	1	4
Female	127	-	-	-	-	-	1	1	-	1	1	2
White, both sexes	311	-	-	-	-	-	1	1	-	3	2	6
Male	188	-	-	-	-	-	-	-	-	2	1	4
Female	123	-	-	-	-	-	1	1	-	1	1	2
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	36	-	-	-	1	-	-	-	-	-	1	-
Male	25	-	-	-	1	-	-	-	-	-	-	-
Female	11	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	34	-	-	-	1	-	-	-	-	-	1	-
Male	24	-	-	-	1	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
United States												
All races, both sexes	435	578	878	1,185	1,685	2,340	2,631	2,003	1,020	257	26	-
Male	321	445	618	807	1,104	1,365	1,413	950	422	85	3	-
Female	114	133	260	378	581	975	1,218	1,053	598	172	23	-
White, both sexes	301	467	742	1,019	1,520	2,126	2,406	1,848	934	228	25	-
Male	229	366	537	701	1,022	1,255	1,308	884	397	77	3	-
Female	72	101	205	318	498	871	1,098	964	537	151	22	-
Black, both sexes	113	102	111	131	124	153	159	110	58	19	1	-
Male	75	72	66	79	54	76	67	39	14	2	-	-
Female	38	30	45	52	70	77	92	71	44	17	1	-
American Indian 1/, both sexes	1	1	3	8	5	8	7	3	2	2	-	-
Male	-	1	1	6	4	3	4	-	-	-	-	-
Female	1	-	2	2	1	5	3	3	2	2	-	-
Asian or Pacific Islander 2/, both sexes	20	8	22	27	36	53	59	42	26	8	-	-
Male	17	6	14	21	24	31	34	27	11	6	-	-
Female	3	2	8	6	12	22	25	15	15	2	-	-
Alabama												
All races, both sexes	5	9	13	21	41	35	36	21	12	4	-	-
Male	4	9	9	13	29	23	17	11	7	-	-	-
Female	1	-	4	8	12	12	19	10	5	4	-	-
White, both sexes	3	5	9	20	38	30	30	17	12	1	-	-
Male	3	5	7	13	28	19	14	9	7	-	-	-
Female	-	-	2	7	10	11	16	8	5	1	-	-
Black, both sexes	2	4	4	1	2	5	5	4	-	3	-	-
Male	1	4	2	-	1	4	3	2	-	-	-	-
Female	1	-	2	1	1	1	2	2	-	3	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Alaska												
All races, both sexes	-	-	1	-	2	1	5	-	1	-	-	-
Male	-	-	1	-	2	1	3	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	1	-	-	-
White, both sexes	-	-	1	-	1	1	4	-	1	-	-	-
Male	-	-	1	-	1	1	3	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	9	9	20	19	27	43	40	31	16	5	-	-
Male	6	9	16	16	17	25	23	15	7	3	-	-
Female	3	-	4	3	10	18	17	16	9	2	-	-
White, both sexes	7	8	20	18	25	40	40	30	16	5	-	-
Male	5	8	16	15	15	23	23	14	7	3	-	-
Female	2	-	4	3	10	17	17	16	9	2	-	-
Black, both sexes	2	1	-	-	1	1	-	-	-	-	-	-
Male	1	1	-	-	1	1	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	1	1	-	1	-	-	-	-
Male	-	-	-	1	1	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Arkansas												
All races, both sexes	6	6	10	20	25	28	22	23	7	3	-	-
Male	3	4	6	11	15	12	11	10	1	2	-	-
Female	3	2	4	9	10	16	11	13	6	1	-	-
White, both sexes	5	6	8	19	24	27	20	22	6	3	-	-
Male	3	4	4	10	14	12	11	10	1	2	-	-
Female	2	2	4	9	10	15	9	12	5	1	-	-
Black, both sexes	1	-	2	1	1	1	2	1	1	-	-	-
Male	-	-	2	1	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	53	55	76	105	138	196	250	173	83	17	2	-
Male	39	41	51	69	87	120	142	89	39	10	-	-
Female	14	14	25	36	51	76	108	84	44	7	2	-
White, both sexes	39	40	61	85	114	162	212	148	66	14	2	-
Male	31	31	41	54	75	103	123	72	35	8	-	-
Female	8	9	20	31	39	59	89	76	31	6	2	-
Black, both sexes	7	10	4	9	9	7	11	8	6	1	-	-
Male	2	7	3	6	5	3	5	3	-	-	-	-
Female	5	3	1	3	4	4	6	5	6	1	-	-
American Indian 1/, both sexes	-	-	1	2	-	2	1	-	-	-	-	-
Male	-	-	-	2	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	5	10	9	15	25	26	17	11	2	-	-
Male	6	3	7	7	7	13	13	14	4	2	-	-
Female	1	2	3	2	8	12	13	3	7	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Colorado												
All races, both sexes	7	16	15	12	26	37	33	17	13	4	-	-
Male	4	11	11	8	17	16	17	8	6	3	-	-
Female	3	5	4	4	9	21	16	9	7	1	-	-
White, both sexes	6	15	14	10	25	32	33	17	13	4	-	-
Male	3	10	10	7	17	13	17	8	6	3	-	-
Female	3	5	4	3	8	19	16	9	7	1	-	-
Black, both sexes	1	1	1	1	1	3	-	-	-	-	-	-
Male	1	1	1	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	1	2	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	6	5	10	23	28	36	38	17	13	-	-
Male	-	6	5	7	14	7	22	16	8	3	-	-
Female	-	-	-	3	9	21	14	22	9	10	-	-
White, both sexes	-	6	5	9	23	26	36	38	16	12	-	-
Male	-	6	5	6	14	7	22	16	7	3	-	-
Female	-	-	-	3	9	19	14	22	9	9	-	-
Black, both sexes	-	-	-	1	-	1	-	-	1	1	-	-
Male	-	-	-	1	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Delaware												
All races, both sexes	-	1	5	3	3	10	9	3	5	2	-	-
Male	-	-	5	3	1	6	4	2	2	1	-	-
Female	-	1	-	-	2	4	5	1	3	1	-	-
White, both sexes	-	1	4	2	2	9	9	3	5	1	-	-
Male	-	-	4	2	1	6	4	2	2	1	-	-
Female	-	1	-	-	1	3	5	1	3	-	-	-
Black, both sexes	-	-	-	1	1	1	-	-	-	1	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	6	5	2	4	4	3	3	4	-	-	-
Male	2	3	4	1	2	2	2	1	2	-	-	-
Female	-	3	1	1	2	2	1	2	2	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	6	5	2	4	4	2	3	4	-	-	-
Male	2	3	4	1	2	2	1	1	2	-	-	-
Female	-	3	1	1	2	2	1	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Florida												
All races, both sexes	34	41	68	92	120	180	208	142	65	12	2	-
Male	24	31	44	62	82	107	118	68	26	4	1	-
Female	10	10	24	30	38	73	90	74	39	8	1	-
White, both sexes	20	33	60	83	112	165	192	136	61	12	2	-
Male	13	26	40	57	78	101	112	65	26	4	1	-
Female	7	7	20	26	34	64	80	71	35	8	1	-
Black, both sexes	12	7	8	8	6	12	13	6	4	-	-	-
Male	9	4	4	4	3	5	4	3	-	-	-	-
Female	3	3	4	4	3	7	9	3	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	-	1	2	3	3	-	-	-	-	-
Male	2	1	-	1	1	1	2	-	-	-	-	-
Female	-	-	-	-	1	2	1	-	-	-	-	-
Georgia												
All races, both sexes	13	27	27	28	38	39	57	40	25	5	-	-
Male	11	20	19	24	22	28	24	14	7	3	-	-
Female	2	7	8	4	16	11	33	26	18	2	-	-
White, both sexes	7	16	22	21	35	33	51	36	19	3	-	-
Male	6	13	16	19	20	23	22	12	5	2	-	-
Female	1	3	6	2	15	10	29	24	14	1	-	-
Black, both sexes	6	11	5	7	2	6	6	4	6	2	-	-
Male	5	7	3	5	1	5	2	2	2	1	-	-
Female	1	4	2	2	1	1	4	2	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Hawaii												
All races, both sexes	5	2	4	6	8	11	11	15	10	5	-	-
Male	5	2	2	4	6	6	8	7	3	3	-	-
Female	-	-	2	2	2	5	3	8	7	2	-	-
White, both sexes	1	1	2	2	4	2	-	2	2	1	-	-
Male	1	1	1	1	4	1	-	-	-	-	-	-
Female	-	-	1	1	-	1	-	2	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	1	2	4	4	9	11	13	8	4	-	-
Male	4	1	1	3	2	5	8	7	3	3	-	-
Female	-	-	1	1	2	4	3	6	5	1	-	-
Idaho												
All races, both sexes	3	2	3	7	10	15	15	12	4	5	-	-
Male	3	1	2	4	8	6	11	7	1	2	-	-
Female	-	1	1	3	2	9	4	5	3	3	-	-
White, both sexes	2	2	3	7	9	14	15	11	4	5	-	-
Male	2	1	2	4	7	6	11	7	1	2	-	-
Female	-	1	1	3	2	8	4	4	3	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Illinois												
All races, both sexes	21	15	31	45	74	110	119	80	49	11	1	-
Male	16	11	24	26	38	67	59	31	20	2	-	-
Female	5	4	7	19	36	43	60	49	29	9	1	-
White, both sexes	12	10	28	37	60	99	107	72	42	10	1	-
Male	9	6	22	22	34	59	54	27	16	2	-	-
Female	3	4	6	15	26	40	53	45	26	8	1	-
Black, both sexes	8	4	3	7	13	9	10	7	5	1	-	-
Male	7	4	2	3	3	6	4	4	3	-	-	-
Female	1	-	1	4	10	3	6	3	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	1	-	1	1	2	2	1	2	-	-	-
Male	-	1	-	1	1	2	1	-	1	-	-	-
Female	1	-	-	-	-	-	1	1	1	-	-	-
Indiana												
All races, both sexes	10	15	16	27	44	52	54	51	28	7	-	-
Male	9	12	11	16	34	31	34	24	12	3	-	-
Female	1	3	5	11	10	21	20	27	16	4	-	-
White, both sexes	7	15	16	25	42	51	53	46	28	6	-	-
Male	6	12	11	15	34	30	34	20	12	3	-	-
Female	1	3	5	10	8	21	19	26	16	3	-	-
Black, both sexes	3	-	-	2	2	1	1	5	-	1	-	-
Male	3	-	-	1	-	1	-	4	-	-	-	-
Female	-	-	-	1	2	-	1	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Iowa												
All races, both sexes	5	7	12	15	29	39	43	34	15	4	2	-
Male	1	6	11	11	18	24	24	21	9	2	-	-
Female	4	1	1	4	11	15	19	13	6	2	2	-
White, both sexes	4	6	12	15	27	39	43	33	15	4	2	-
Male	1	5	11	11	16	24	24	21	9	2	-	-
Female	3	1	1	4	11	15	19	12	6	2	2	-
Black, both sexes	1	1	-	-	1	-	-	1	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	4	8	13	16	15	13	34	25	15	1	2	-
Male	3	5	9	12	10	11	19	11	7	-	-	-
Female	1	3	4	4	5	2	15	14	8	1	2	-
White, both sexes	3	8	13	15	14	13	32	23	15	1	2	-
Male	3	5	9	11	10	11	19	10	7	-	-	-
Female	-	3	4	4	4	2	13	13	8	1	2	-
Black, both sexes	-	-	-	-	1	-	2	2	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	1	-	2	1	-	-	-	-
American Indian 1/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Kentucky												
All races, both sexes	7	6	14	25	34	41	34	26	11	1	1	-
Male	4	4	10	16	24	22	18	11	4	1	-	-
Female	3	2	4	9	10	19	16	15	7	-	1	-
White, both sexes	6	5	14	23	34	40	31	23	11	1	1	-
Male	3	4	10	15	24	21	16	10	4	1	-	-
Female	3	1	4	8	10	19	15	13	7	-	1	-
Black, both sexes	1	1	-	2	-	1	3	3	-	-	-	-
Male	1	-	-	1	-	1	2	1	-	-	-	-
Female	-	1	-	1	-	-	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	7	9	16	22	18	26	38	24	12	1	1	-
Male	3	9	11	13	12	11	16	8	6	-	-	-
Female	4	-	5	9	6	15	22	16	6	1	1	-
White, both sexes	2	7	12	13	17	23	31	23	10	-	1	-
Male	2	7	10	7	11	8	12	8	5	-	-	-
Female	-	-	2	6	6	15	19	15	5	-	1	-
Black, both sexes	5	2	4	9	1	3	7	1	2	1	-	-
Male	1	2	1	6	1	3	4	-	1	-	-	-
Female	4	-	3	3	-	-	3	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Maine												
All races, both sexes	-	1	3	5	12	19	17	21	13	2	1	-
Male	-	1	1	3	9	9	7	9	5	1	-	-
Female	-	-	2	2	3	10	10	12	8	1	1	-
White, both sexes	-	1	3	5	12	19	17	21	13	2	1	-
Male	-	1	1	3	9	9	7	9	5	1	-	-
Female	-	-	2	2	3	10	10	12	8	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	8	7	11	30	28	42	40	33	23	7	1	-
Male	4	7	8	20	13	20	16	14	6	2	1	-
Female	4	-	3	10	15	22	24	19	17	5	-	-
White, both sexes	5	5	11	19	22	38	34	32	21	7	1	-
Male	3	5	8	16	13	17	11	14	5	2	1	-
Female	2	-	3	3	9	21	23	18	16	5	-	-
Black, both sexes	3	2	-	9	6	3	4	1	2	-	-	-
Male	1	2	-	3	-	2	3	-	1	-	-	-
Female	2	-	-	6	6	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	2	-	1	2	-	-	-	-	-
Male	-	-	-	1	-	1	2	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Massachusetts												
All races, both sexes	5	8	18	12	35	60	61	66	33	9	1	-
Male	4	4	11	9	20	38	33	25	12	2	-	-
Female	1	4	7	3	15	22	28	41	21	7	1	-
White, both sexes	4	7	14	12	29	59	60	65	33	7	1	-
Male	3	4	10	9	15	38	32	25	12	1	-	-
Female	1	3	4	3	14	21	28	40	21	6	1	-
Black, both sexes	1	1	2	-	4	1	1	1	-	1	-	-
Male	1	-	1	-	3	-	1	-	-	-	-	-
Female	-	1	1	-	1	1	-	1	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	2	-	-	-	-	1	-	-
Male	-	-	-	-	2	-	-	-	-	1	-	-
Female	-	-	2	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	12	27	38	51	56	88	119	81	43	10	2	-
Male	7	18	25	31	33	48	69	46	18	4	-	-
Female	5	9	13	20	23	40	50	35	25	6	2	-
White, both sexes	7	21	34	41	48	82	113	71	40	9	2	-
Male	4	15	24	24	31	46	65	39	18	3	-	-
Female	3	6	10	17	17	36	48	32	22	6	2	-
Black, both sexes	5	6	4	8	7	5	3	10	2	1	-	-
Male	3	3	1	5	2	1	2	7	-	1	-	-
Female	2	3	3	3	5	4	1	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	2	1	-	-	-	1	-	-	-
Male	-	-	-	2	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	3	-	-	-	-	-
Male	-	-	-	-	-	1	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Minnesota												
All races, both sexes	6	14	12	30	27	45	64	45	23	6	1	-
Male	4	10	8	22	19	26	39	22	11	2	-	-
Female	2	4	4	8	8	19	25	23	12	4	1	-
White, both sexes	5	13	11	30	27	44	64	45	23	6	1	-
Male	3	9	7	22	19	25	39	22	11	2	-	-
Female	2	4	4	8	8	19	25	23	12	4	1	-
Black, both sexes	1	1	1	-	-	1	-	-	-	-	-	-
Male	1	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	5	6	10	15	23	20	25	15	8	2	-	-
Male	4	4	8	12	17	14	10	7	3	-	-	-
Female	1	2	2	3	6	6	15	8	5	2	-	-
White, both sexes	3	3	7	9	21	17	21	8	6	1	-	-
Male	2	2	7	7	15	12	9	5	3	-	-	-
Female	1	1	-	2	6	5	12	3	3	1	-	-
Black, both sexes	2	3	3	5	2	3	4	6	2	1	-	-
Male	2	2	1	4	2	2	1	1	-	-	-	-
Female	-	1	2	1	-	1	3	5	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Missouri												
All races, both sexes	11	8	18	26	35	65	58	39	22	3	-	-
Male	9	5	12	20	24	43	29	22	8	-	-	-
Female	2	3	6	6	11	22	29	17	14	3	-	-
White, both sexes	7	8	16	23	34	61	57	36	20	3	-	-
Male	6	5	11	19	24	39	29	22	7	-	-	-
Female	1	3	5	4	10	22	28	14	13	3	-	-
Black, both sexes	4	-	2	3	1	2	1	3	2	-	-	-
Male	3	-	1	1	-	2	-	-	1	-	-	-
Female	1	-	1	2	1	-	1	3	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	1	1	3	3	14	9	10	2	3	-	-
Male	1	1	1	2	2	10	8	7	1	-	-	-
Female	-	-	-	1	1	4	1	3	1	3	-	-
White, both sexes	1	1	1	3	3	13	9	9	2	2	-	-
Male	1	1	1	2	2	10	8	7	1	-	-	-
Female	-	-	-	1	1	3	1	2	1	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Nebraska												
All races, both sexes	1	5	2	7	7	19	14	15	7	1	1	-
Male	-	5	-	6	6	9	7	7	4	-	-	-
Female	1	-	2	1	1	10	7	8	3	1	1	-
White, both sexes	1	5	2	7	6	18	14	14	7	1	1	-
Male	-	5	-	6	5	8	7	7	4	-	-	-
Female	1	-	2	1	1	10	7	7	3	1	1	-
Black, both sexes	-	-	-	-	1	1	-	1	-	-	-	-
Male	-	-	-	-	1	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	5	3	12	6	14	19	19	6	5	2	-	-
Male	2	2	7	3	8	11	11	1	1	1	-	-
Female	3	1	5	3	6	8	8	5	4	1	-	-
White, both sexes	5	3	11	6	14	18	19	5	4	2	-	-
Male	2	2	7	3	8	10	11	1	-	1	-	-
Female	3	1	4	3	6	8	8	4	4	1	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
New Hampshire												
All races, both sexes	1	4	3	7	4	14	15	9	-	3	-	-
Male	-	4	3	4	2	6	7	1	-	1	-	-
Female	1	-	-	3	2	8	8	8	-	2	-	-
White, both sexes	1	4	3	7	4	14	15	9	-	3	-	-
Male	-	4	3	4	2	6	7	1	-	1	-	-
Female	1	-	-	3	2	8	8	8	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	14	13	25	30	48	66	90	61	30	10	2	-
Male	11	12	19	18	30	41	47	30	12	3	-	-
Female	3	1	6	12	18	25	43	31	18	7	2	-
White, both sexes	11	12	18	25	40	62	85	54	29	9	2	-
Male	9	11	15	16	24	37	45	28	12	3	-	-
Female	2	1	3	9	16	25	40	26	17	6	2	-
Black, both sexes	3	1	5	5	4	3	5	5	1	1	-	-
Male	2	1	3	2	2	3	2	1	-	-	-	-
Female	1	-	2	3	2	-	3	4	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	4	1	-	1	-	-	-	-
Male	-	-	1	-	4	1	-	1	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
New Mexico												
All races, both sexes	2	5	4	5	8	26	15	14	6	2	-	-
Male	2	2	2	4	6	17	7	8	4	1	-	-
Female	-	3	2	1	2	9	8	6	2	1	-	-
White, both sexes	1	5	4	5	7	25	15	14	6	2	-	-
Male	1	2	2	4	5	17	7	8	4	1	-	-
Female	-	3	2	1	2	8	8	6	2	1	-	-
Black, both sexes	1	-	-	-	1	1	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	25	46	71	62	112	141	150	124	69	12	1	-
Male	18	37	49	39	70	85	79	42	23	5	1	-
Female	7	9	22	23	42	56	71	82	46	7	-	-
White, both sexes	12	32	55	50	101	129	135	116	64	12	1	-
Male	8	26	40	30	64	78	69	40	23	5	1	-
Female	4	6	15	20	37	51	66	76	41	7	-	-
Black, both sexes	11	14	15	9	9	11	13	6	5	-	-	-
Male	8	11	8	7	4	7	9	1	-	-	-	-
Female	3	3	7	2	5	4	4	5	5	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	3	2	1	2	2	-	-	-	-
Male	2	-	-	2	2	-	1	1	-	-	-	-
Female	-	-	-	1	-	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
North Carolina												
All races, both sexes	18	18	30	37	43	68	62	54	25	11	-	-
Male	16	13	17	28	31	47	39	28	11	4	-	-
Female	2	5	13	9	12	21	23	26	14	7	-	-
White, both sexes	14	13	22	27	39	59	52	48	22	10	-	-
Male	13	10	11	21	28	43	37	26	11	4	-	-
Female	1	3	11	6	11	16	15	22	11	6	-	-
Black, both sexes	3	5	8	9	4	8	10	6	3	-	-	-
Male	2	3	6	7	3	4	2	2	-	-	-	-
Female	1	2	2	2	1	4	8	4	3	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	2	2	5	6	8	9	6	2	-	-
Male	1	-	2	1	4	4	7	5	2	-	-	-
Female	-	-	-	1	1	2	1	4	4	2	-	-
White, both sexes	1	-	2	2	5	6	8	9	6	2	-	-
Male	1	-	2	1	4	4	7	5	2	-	-	-
Female	-	-	-	1	1	2	1	4	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Ohio												
All races, both sexes	20	24	39	59	84	112	143	87	43	14	2	-
Male	14	22	24	36	58	66	75	48	18	3	-	-
Female	6	2	15	23	26	46	68	39	25	11	2	-
White, both sexes	16	20	34	57	76	104	125	81	41	12	2	-
Male	11	19	22	35	55	66	69	46	18	3	-	-
Female	5	1	12	22	21	38	56	35	23	9	2	-
Black, both sexes	4	4	4	2	8	8	18	5	2	2	-	-
Male	3	3	2	1	3	-	6	2	-	-	-	-
Female	1	1	2	1	5	8	12	3	2	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
Oklahoma												
All races, both sexes	5	11	12	17	20	32	39	25	14	1	-	-
Male	5	10	9	12	13	18	25	16	5	-	-	-
Female	-	1	3	5	7	14	14	9	9	1	-	-
White, both sexes	5	10	12	17	19	29	35	24	14	1	-	-
Male	5	9	9	12	12	17	23	15	5	-	-	-
Female	-	1	3	5	7	12	12	9	9	1	-	-
Black, both sexes	-	1	-	-	-	1	1	1	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	2	2	-	-	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Oregon												
All races, both sexes	3	9	14	17	13	35	38	28	17	1	-	-
Male	3	7	11	12	9	19	23	18	7	1	-	-
Female	-	2	3	5	4	16	15	10	10	-	-	-
White, both sexes	3	9	13	16	13	33	38	27	17	1	-	-
Male	3	7	10	11	9	18	23	18	7	1	-	-
Female	-	2	3	5	4	15	15	9	10	-	-	-
Black, both sexes	-	-	1	-	-	1	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes	10	24	41	55	90	118	169	148	73	14	1	-
Male	8	19	30	37	56	63	76	61	28	6	-	-
Female	2	5	11	18	34	55	93	87	45	8	1	-
White, both sexes	7	19	35	48	81	108	157	141	73	14	1	-
Male	7	14	27	33	51	58	70	60	28	6	-	-
Female	-	5	8	15	30	50	87	81	45	8	1	-
Black, both sexes	2	5	6	6	9	10	12	7	-	-	-	-
Male	1	5	3	3	5	5	6	1	-	-	-	-
Female	1	-	3	3	4	5	6	6	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Rhode Island												
All races, both sexes	4	1	4	3	6	12	12	10	4	-	-	-
Male	3	1	2	3	3	10	4	4	1	-	-	-
Female	1	-	2	-	3	2	8	6	3	-	-	-
White, both sexes	4	1	4	3	6	11	10	10	4	-	-	-
Male	3	1	2	3	3	9	3	4	1	-	-	-
Female	1	-	2	-	3	2	7	6	3	-	-	-
Black, both sexes	-	-	-	-	-	1	1	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	7	8	22	26	27	37	40	28	16	5	1	-
Male	5	5	15	21	18	20	19	11	4	2	-	-
Female	2	3	7	5	9	17	21	17	12	3	1	-
White, both sexes	4	6	15	22	21	29	30	26	14	5	-	-
Male	3	4	9	19	16	17	16	10	4	2	-	-
Female	1	2	6	3	5	12	14	16	10	3	-	-
Black, both sexes	3	2	7	4	5	8	9	2	2	-	1	-
Male	2	1	6	2	1	3	3	1	-	-	-	-
Female	1	1	1	2	4	5	6	1	2	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
South Dakota												
All races, both sexes	2	1	4	1	9	12	8	9	3	1	-	-
Male	-	1	3	1	4	9	5	7	3	-	-	-
Female	2	-	1	-	5	3	3	2	-	1	-	-
White, both sexes	2	-	4	1	9	12	8	9	3	1	-	-
Male	-	-	3	1	4	9	5	7	3	-	-	-
Female	2	-	1	-	5	3	3	2	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	7	11	25	29	40	40	50	37	12	2	-	-
Male	5	5	15	23	30	19	25	18	6	-	-	-
Female	2	6	10	6	10	21	25	19	6	2	-	-
White, both sexes	5	10	20	26	36	35	46	35	11	2	-	-
Male	4	5	14	21	28	18	24	16	5	-	-	-
Female	1	5	6	5	8	17	22	19	6	2	-	-
Black, both sexes	2	1	5	3	3	5	4	1	1	-	-	-
Male	1	-	1	2	1	1	1	1	1	-	-	-
Female	1	1	4	1	2	4	3	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Texas												
All races, both sexes	30	35	36	72	105	124	114	69	45	12	-	-
Male	23	28	27	50	79	66	61	34	19	3	-	-
Female	7	7	9	22	26	58	53	35	26	9	-	-
White, both sexes	20	32	31	61	97	107	104	63	39	10	-	-
Male	17	26	23	44	74	60	56	34	18	3	-	-
Female	3	6	8	17	23	47	48	29	21	7	-	-
Black, both sexes	10	3	4	11	7	17	6	5	5	2	-	-
Male	6	2	3	6	4	6	4	-	1	-	-	-
Female	4	1	1	5	3	11	2	5	4	2	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	1	-	4	1	1	-	-	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	3	1	1	-	-	-
Utah												
All races, both sexes	5	3	3	9	13	14	13	7	2	1	-	-
Male	4	2	3	7	9	10	6	4	2	-	-	-
Female	1	1	-	2	4	4	7	3	-	1	-	-
White, both sexes	5	3	1	8	13	14	12	7	2	1	-	-
Male	4	2	1	6	9	10	5	4	2	-	-	-
Female	1	1	-	2	4	4	7	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	1	-	-	-	-	-	-	-	-
Male	-	-	2	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Vermont												
All races, both sexes	-	1	-	3	5	9	15	11	2	2	-	-
Male	-	1	-	2	3	7	8	8	2	-	-	-
Female	-	-	-	1	2	2	7	3	-	2	-	-
White, both sexes	-	1	-	3	5	9	15	11	2	2	-	-
Male	-	1	-	2	3	7	8	8	2	-	-	-
Female	-	-	-	1	2	2	7	3	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	10	8	25	26	30	48	42	53	18	4	-	-
Male	9	6	22	22	18	31	20	18	6	-	-	-
Female	1	2	3	4	12	17	22	35	12	4	-	-
White, both sexes	6	6	18	22	24	41	38	49	16	4	-	-
Male	5	5	16	18	15	27	19	18	5	-	-	-
Female	1	1	2	4	9	14	19	31	11	4	-	-
Black, both sexes	3	2	6	4	6	7	4	4	1	-	-	-
Male	3	1	5	4	3	4	1	-	-	-	-	-
Female	-	1	1	-	3	3	3	4	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	-	-	-	-	-	1	-	-	-
Male	1	-	1	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Washington												
All races, both sexes	14	14	20	29	40	46	55	46	30	4	-	-
Male	13	12	16	17	26	22	37	27	17	1	-	-
Female	1	2	4	12	14	24	18	19	13	3	-	-
White, both sexes	13	12	18	25	40	43	52	45	27	3	-	-
Male	12	10	15	14	26	20	35	26	15	1	-	-
Female	1	2	3	11	14	23	17	19	12	2	-	-
Black, both sexes	1	2	1	2	-	1	-	-	1	-	-	-
Male	1	2	1	1	-	1	-	-	1	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	1	-	1	-	-	-	-	-	-
Male	-	-	-	1	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	3	1	2	1	-	-
Male	-	-	-	1	-	-	2	1	1	-	-	-
Female	-	-	-	-	-	1	1	-	1	1	-	-
West Virginia												
All races, both sexes	3	5	5	12	20	17	19	16	7	4	-	-
Male	1	4	5	9	15	14	10	8	1	1	-	-
Female	2	1	-	3	5	3	9	8	6	3	-	-
White, both sexes	1	5	5	12	19	17	19	16	7	4	-	-
Male	1	4	5	9	15	14	10	8	1	1	-	-
Female	-	1	-	3	4	3	9	8	6	3	-	-
Black, both sexes	2	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Aortic aneurysm and dissection (I71)												
Wisconsin												
All races, both sexes	7	12	13	21	39	60	56	64	26	5	-	-
Male	6	10	12	15	28	34	27	35	15	-	-	-
Female	1	2	1	6	11	26	29	29	11	5	-	-
White, both sexes	6	12	13	20	38	60	56	63	25	5	-	-
Male	5	10	12	15	28	34	27	35	15	-	-	-
Female	1	2	1	5	10	26	29	28	10	5	-	-
Black, both sexes	1	-	-	-	1	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	2	5	1	3	5	4	5	5	1	2	1	-
Male	2	3	-	2	3	4	5	5	-	-	-	-
Female	-	2	1	1	2	-	-	-	1	2	1	-
White, both sexes	2	4	1	3	5	4	4	5	1	2	1	-
Male	2	3	-	2	3	4	4	5	-	-	-	-
Female	-	1	1	1	2	-	-	-	1	2	1	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
United States												
All races, both sexes	9,774	16	3	5	8	4	14	16	26	47	91	123
Male	4,255	10	3	3	6	4	7	7	12	23	34	63
Female	5,519	6	-	2	2	-	7	9	14	24	57	60
White, both sexes	8,329	10	3	4	7	2	10	8	16	33	68	90
Male	3,622	5	3	3	5	2	4	3	9	15	25	44
Female	4,707	5	-	1	2	-	6	5	7	18	43	46
Black, both sexes	1,298	4	-	1	1	2	3	7	10	13	21	28
Male	563	3	-	-	1	2	2	4	3	7	8	17
Female	735	1	-	1	-	-	1	3	7	6	13	11
American Indian 1/, both sexes	41	1	-	-	-	-	-	-	-	1	1	4
Male	16	1	-	-	-	-	-	-	-	1	-	1
Female	25	-	-	-	-	-	-	-	-	-	1	3
Asian or Pacific Islander 2/, both sexes	106	1	-	-	-	-	1	1	-	-	1	1
Male	54	1	-	-	-	-	1	-	-	-	1	1
Female	52	-	-	-	-	-	-	1	-	-	-	-
Alabama												
All races, both sexes	185	-	-	-	-	-	-	1	1	1	-	1
Male	69	-	-	-	-	-	-	1	-	-	-	1
Female	116	-	-	-	-	-	-	-	1	1	-	-
White, both sexes	147	-	-	-	-	-	-	1	1	1	-	-
Male	58	-	-	-	-	-	-	1	-	-	-	-
Female	89	-	-	-	-	-	-	-	1	1	-	-
Black, both sexes	37	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	26	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Alaska												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	10	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	149	1	1	-	-	-	-	-	-	-	2	2
Male	67	1	1	-	-	-	-	-	-	-	1	1
Female	82	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	138	-	1	-	-	-	-	-	-	-	2	2
Male	64	-	1	-	-	-	-	-	-	-	1	1
Female	74	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	6	1	-	-	-	-	-	-	-	-	-	-
Male	2	1	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Arkansas												
All races, both sexes	135	-	1	1	-	-	2	-	-	-	1	2
Male	56	-	1	-	-	-	1	-	-	-	-	-
Female	79	-	-	1	-	-	1	-	-	-	1	2
White, both sexes	120	-	1	-	-	-	1	-	-	-	1	2
Male	48	-	1	-	-	-	-	-	-	-	-	-
Female	72	-	-	-	-	-	1	-	-	-	1	2
Black, both sexes	15	-	-	1	-	-	1	-	-	-	-	-
Male	8	-	-	-	-	-	1	-	-	-	-	-
Female	7	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	794	2	-	-	1	1	4	1	2	4	12	11
Male	382	2	-	-	1	1	3	-	2	2	4	7
Female	412	-	-	-	-	-	1	1	-	2	8	4
White, both sexes	672	-	-	-	1	1	3	-	1	4	8	9
Male	324	-	-	-	1	1	2	-	1	2	3	5
Female	348	-	-	-	-	-	1	-	-	2	5	4
Black, both sexes	74	1	-	-	-	-	-	-	1	-	4	2
Male	32	1	-	-	-	-	-	-	1	-	1	2
Female	42	-	-	-	-	-	-	-	-	-	3	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	48	1	-	-	-	-	1	1	-	-	-	-
Male	26	1	-	-	-	-	1	-	-	-	-	-
Female	22	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Colorado												
All races, both sexes	122	-	-	1	1	-	-	1	-	-	4	-
Male	55	-	-	1	1	-	-	1	-	-	1	-
Female	67	-	-	-	-	-	-	-	-	-	3	-
White, both sexes	116	-	-	1	1	-	-	1	-	-	4	-
Male	52	-	-	1	1	-	-	1	-	-	1	-
Female	64	-	-	-	-	-	-	-	-	-	3	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	117	1	-	-	-	-	-	-	-	2	-	-
Male	54	1	-	-	-	-	-	-	-	1	-	-
Female	63	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	108	-	-	-	-	-	-	-	-	1	-	-
Male	47	-	-	-	-	-	-	-	-	-	-	-
Female	61	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	8	1	-	-	-	-	-	-	-	1	-	-
Male	6	1	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Delaware												
All races, both sexes	40	-	-	-	-	-	-	-	-	-	-	1
Male	21	-	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	29	-	-	-	-	-	-	-	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	25	-	-	-	-	-	-	-	-	1	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	17	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Florida												
All races, both sexes	649	-	1	-	1	-	-	-	3	6	6	7
Male	303	-	1	-	1	-	-	-	1	-	1	4
Female	346	-	-	-	-	-	-	-	2	6	5	3
White, both sexes	547	-	1	-	1	-	-	-	1	3	2	4
Male	265	-	1	-	1	-	-	-	1	-	1	2
Female	282	-	-	-	-	-	-	-	-	3	1	2
Black, both sexes	99	-	-	-	-	-	-	-	2	3	4	3
Male	38	-	-	-	-	-	-	-	-	-	-	2
Female	61	-	-	-	-	-	-	-	2	3	4	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	262	2	-	1	-	-	-	1	1	4	1	6
Male	108	1	-	-	-	-	-	-	-	4	1	3
Female	154	1	-	1	-	-	-	1	1	-	-	3
White, both sexes	179	2	-	1	-	-	-	-	-	3	-	2
Male	73	1	-	-	-	-	-	-	-	3	-	1
Female	106	1	-	1	-	-	-	-	-	-	-	1
Black, both sexes	82	-	-	-	-	-	-	1	1	1	1	4
Male	34	-	-	-	-	-	-	-	-	1	1	2
Female	48	-	-	-	-	-	-	1	1	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Hawaii												
All races, both sexes	31	-	-	-	1	-	-	-	-	-	-	-
Male	13	-	-	-	1	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	1	-	-	-	-	-	-	-
Male	7	-	-	-	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	22	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	39	-	-	-	-	-	-	-	1	1	-	-
Male	19	-	-	-	-	-	-	-	1	1	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	39	-	-	-	-	-	-	-	1	1	-	-
Male	19	-	-	-	-	-	-	-	1	1	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Illinois												
All races, both sexes	438	1	-	-	1	-	-	1	1	2	4	4
Male	196	-	-	-	1	-	-	-	1	2	1	1
Female	242	1	-	-	-	-	-	1	-	-	3	3
White, both sexes	367	1	-	-	1	-	-	1	-	1	4	2
Male	159	-	-	-	1	-	-	-	-	1	1	1
Female	208	1	-	-	-	-	-	1	-	-	3	1
Black, both sexes	70	-	-	-	-	-	-	-	1	1	-	2
Male	37	-	-	-	-	-	-	-	1	1	-	-
Female	33	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	236	-	-	1	-	1	-	-	-	-	4	2
Male	115	-	-	1	-	1	-	-	-	-	2	1
Female	121	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	219	-	-	1	-	1	-	-	-	-	3	2
Male	103	-	-	1	-	1	-	-	-	-	1	1
Female	116	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	17	-	-	-	-	-	-	-	-	-	1	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Iowa												
All races, both sexes	125	-	-	-	-	-	-	-	2	-	1	1
Male	52	-	-	-	-	-	-	-	-	-	-	-
Female	73	-	-	-	-	-	-	-	2	-	1	1
White, both sexes	125	-	-	-	-	-	-	-	2	-	1	1
Male	52	-	-	-	-	-	-	-	-	-	-	-
Female	73	-	-	-	-	-	-	-	2	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	105	-	-	-	-	-	-	-	-	1	1	2
Male	51	-	-	-	-	-	-	-	-	1	1	2
Female	54	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	97	-	-	-	-	-	-	-	-	-	1	2
Male	46	-	-	-	-	-	-	-	-	-	1	2
Female	51	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Kentucky												
All races, both sexes	155	-	-	-	-	-	-	-	-	-	1	1
Male	72	-	-	-	-	-	-	-	-	-	-	1
Female	83	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	140	-	-	-	-	-	-	-	-	-	1	1
Male	67	-	-	-	-	-	-	-	-	-	-	1
Female	73	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	188	-	-	-	-	-	-	1	1	1	2	6
Male	88	-	-	-	-	-	-	1	1	-	1	1
Female	100	-	-	-	-	-	-	-	-	1	1	5
White, both sexes	116	-	-	-	-	-	-	-	1	-	2	4
Male	48	-	-	-	-	-	-	-	1	-	1	-
Female	68	-	-	-	-	-	-	-	-	-	1	4
Black, both sexes	71	-	-	-	-	-	-	1	-	1	-	2
Male	39	-	-	-	-	-	-	1	-	-	-	1
Female	32	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Maine												
All races, both sexes	60	-	-	-	-	-	-	-	-	-	3	2
Male	28	-	-	-	-	-	-	-	-	-	1	1
Female	32	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	60	-	-	-	-	-	-	-	-	-	3	2
Male	28	-	-	-	-	-	-	-	-	-	1	1
Female	32	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	173	-	-	-	-	1	1	1	1	1	-	1
Male	68	-	-	-	-	1	-	-	-	1	-	1
Female	105	-	-	-	-	-	1	1	1	-	-	-
White, both sexes	121	-	-	-	-	-	1	1	1	-	-	1
Male	44	-	-	-	-	-	-	-	-	-	-	1
Female	77	-	-	-	-	-	1	1	1	-	-	-
Black, both sexes	50	-	-	-	-	1	-	-	-	1	-	-
Male	23	-	-	-	-	1	-	-	-	1	-	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Massachusetts												
All races, both sexes	237	-	-	-	-	-	1	-	-	-	1	1
Male	89	-	-	-	-	-	-	-	-	-	1	1
Female	148	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	225	-	-	-	-	-	1	-	-	-	1	-
Male	85	-	-	-	-	-	-	-	-	-	1	-
Female	140	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	8	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	347	2	-	-	1	-	1	-	-	1	1	2
Male	144	1	-	-	-	-	1	-	-	-	1	2
Female	203	1	-	-	1	-	-	-	-	1	-	-
White, both sexes	293	2	-	-	1	-	1	-	-	1	1	1
Male	126	1	-	-	-	-	1	-	-	-	1	1
Female	167	1	-	-	1	-	-	-	-	1	-	-
Black, both sexes	51	-	-	-	-	-	-	-	-	-	-	1
Male	16	-	-	-	-	-	-	-	-	-	-	1
Female	35	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Minnesota												
All races, both sexes	180	-	-	-	-	-	-	-	-	1	-	1
Male	83	-	-	-	-	-	-	-	-	1	-	-
Female	97	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	176	-	-	-	-	-	-	-	-	1	-	-
Male	82	-	-	-	-	-	-	-	-	1	-	-
Female	94	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	92	-	-	-	-	-	-	-	3	2	3	5
Male	33	-	-	-	-	-	-	-	1	1	2	2
Female	59	-	-	-	-	-	-	-	2	1	1	3
White, both sexes	53	-	-	-	-	-	-	-	1	1	2	3
Male	20	-	-	-	-	-	-	-	1	-	1	1
Female	33	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	39	-	-	-	-	-	-	-	2	1	1	2
Male	13	-	-	-	-	-	-	-	-	1	1	1
Female	26	-	-	-	-	-	-	-	2	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Missouri												
All races, both sexes	240	1	-	-	-	-	-	-	-	2	2	2
Male	108	1	-	-	-	-	-	-	-	1	-	1
Female	132	-	-	-	-	-	-	-	-	1	2	1
White, both sexes	207	-	-	-	-	-	-	-	-	2	-	2
Male	95	-	-	-	-	-	-	-	-	1	-	1
Female	112	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	32	1	-	-	-	-	-	-	-	-	2	-
Male	13	1	-	-	-	-	-	-	-	-	-	-
Female	19	-	-	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	36	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	-	-
Male	14	-	-	-	-	-	-	-	-	-	-	-
Female	20	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Nebraska												
All races, both sexes	61	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	61	-	-	-	-	-	-	-	-	-	-	-
Male	32	-	-	-	-	-	-	-	-	-	-	-
Female	29	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	57	-	-	-	1	-	-	-	-	2	2	1
Male	26	-	-	-	1	-	-	-	-	-	1	-
Female	31	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	54	-	-	-	-	-	-	-	-	2	2	1
Male	25	-	-	-	-	-	-	-	-	-	1	-
Female	29	-	-	-	-	-	-	-	-	2	1	1
Black, both sexes	3	-	-	-	1	-	-	-	-	-	-	-
Male	1	-	-	-	1	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
New Hampshire												
All races, both sexes	52	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	52	-	-	-	-	-	-	-	-	-	1	-
Male	19	-	-	-	-	-	-	-	-	-	-	-
Female	33	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	291	-	-	-	-	-	-	1	-	3	2	4
Male	106	-	-	-	-	-	-	-	-	-	1	3
Female	185	-	-	-	-	-	-	1	-	3	1	1
White, both sexes	247	-	-	-	-	-	-	-	-	2	2	3
Male	93	-	-	-	-	-	-	-	-	-	1	2
Female	154	-	-	-	-	-	-	-	-	2	1	1
Black, both sexes	40	-	-	-	-	-	-	1	-	1	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	30	-	-	-	-	-	-	1	-	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
New Mexico												
All races, both sexes	54	-	-	-	-	-	-	-	-	-	1	1
Male	26	-	-	-	-	-	-	-	-	-	1	1
Female	28	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	51	-	-	-	-	-	-	-	-	-	1	-
Male	24	-	-	-	-	-	-	-	-	-	1	-
Female	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	492	-	-	1	1	-	1	3	1	-	4	8
Male	201	-	-	1	-	-	-	2	1	-	2	2
Female	291	-	-	-	1	-	1	1	-	-	2	6
White, both sexes	423	-	-	1	1	-	1	2	1	-	2	7
Male	169	-	-	1	-	-	-	1	1	-	-	2
Female	254	-	-	-	1	-	1	1	-	-	2	5
Black, both sexes	64	-	-	-	-	-	-	1	-	-	1	1
Male	28	-	-	-	-	-	-	1	-	-	1	-
Female	36	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	1	-
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
North Carolina												
All races, both sexes	322	1	-	-	-	-	-	-	3	-	3	7
Male	147	-	-	-	-	-	-	-	-	-	2	7
Female	175	1	-	-	-	-	-	-	3	-	1	-
White, both sexes	220	1	-	-	-	-	-	-	2	-	2	5
Male	97	-	-	-	-	-	-	-	-	-	1	5
Female	123	1	-	-	-	-	-	-	2	-	1	-
Black, both sexes	98	-	-	-	-	-	-	-	1	-	1	2
Male	48	-	-	-	-	-	-	-	-	-	1	2
Female	50	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	30	-	-	-	-	-	-	-	-	1	-	1
Male	13	-	-	-	-	-	-	-	-	1	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	27	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	1	-	1
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Ohio												
All races, both sexes	475	-	-	-	-	-	1	-	2	2	1	5
Male	211	-	-	-	-	-	1	-	2	1	-	3
Female	264	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	418	-	-	-	-	-	1	-	2	2	1	5
Male	182	-	-	-	-	-	1	-	2	1	-	3
Female	236	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	57	-	-	-	-	-	-	-	-	-	-	-
Male	29	-	-	-	-	-	-	-	-	-	-	-
Female	28	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	124	-	-	-	-	-	1	-	-	-	1	4
Male	54	-	-	-	-	-	1	-	-	-	1	2
Female	70	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	111	-	-	-	-	-	-	-	-	-	1	4
Male	49	-	-	-	-	-	-	-	-	-	1	2
Female	62	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	8	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	1	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Oregon												
All races, both sexes	135	-	-	-	-	-	-	-	-	-	2	-
Male	54	-	-	-	-	-	-	-	-	-	1	-
Female	81	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	133	-	-	-	-	-	-	-	-	-	2	-
Male	53	-	-	-	-	-	-	-	-	-	1	-
Female	80	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	521	-	-	-	-	-	1	1	1	2	5	2
Male	201	-	-	-	-	-	-	-	1	2	-	1
Female	320	-	-	-	-	-	1	1	-	-	5	1
White, both sexes	487	-	-	-	-	-	1	1	1	2	5	2
Male	189	-	-	-	-	-	-	-	1	2	-	1
Female	298	-	-	-	-	-	1	1	-	-	5	1
Black, both sexes	33	-	-	-	-	-	-	-	-	-	-	-
Male	12	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Rhode Island												
All races, both sexes	34	-	-	-	-	-	-	1	-	-	-	-
Male	12	-	-	-	-	-	-	1	-	-	-	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	31	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	1	-	-	-	-
Male	1	-	-	-	-	-	-	1	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	192	-	-	-	-	-	-	1	1	-	1	3
Male	81	-	-	-	-	-	-	-	-	-	-	3
Female	111	-	-	-	-	-	-	1	1	-	1	-
White, both sexes	118	-	-	-	-	-	-	-	-	-	-	2
Male	52	-	-	-	-	-	-	-	-	-	-	2
Female	66	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	73	-	-	-	-	-	-	1	1	-	1	1
Male	29	-	-	-	-	-	-	-	-	-	-	1
Female	44	-	-	-	-	-	-	1	1	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
South Dakota												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	30	-	-	-	-	-	-	-	-	-	-	1
Male	15	-	-	-	-	-	-	-	-	-	-	-
Female	15	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	220	-	-	-	-	1	-	-	-	-	3	-
Male	102	-	-	-	-	1	-	-	-	-	1	-
Female	118	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	186	-	-	-	-	-	-	-	-	-	2	-
Male	86	-	-	-	-	-	-	-	-	-	1	-
Female	100	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	34	-	-	-	-	1	-	-	-	-	1	-
Male	16	-	-	-	-	1	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Texas												
All races, both sexes	691	3	-	-	-	-	1	1	-	3	9	11
Male	305	2	-	-	-	-	-	1	-	1	5	6
Female	386	1	-	-	-	-	1	-	-	2	4	5
White, both sexes	586	2	-	-	-	-	-	-	-	3	8	9
Male	254	2	-	-	-	-	-	-	-	1	4	5
Female	332	-	-	-	-	-	-	-	-	2	4	4
Black, both sexes	104	1	-	-	-	-	1	1	-	-	1	2
Male	51	-	-	-	-	-	-	1	-	-	1	1
Female	53	1	-	-	-	-	1	-	-	-	-	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	38	-	-	-	-	-	-	-	-	-	2	1
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	34	-	-	-	-	-	-	-	-	-	1	-
Male	16	-	-	-	-	-	-	-	-	-	-	-
Female	18	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Vermont												
All races, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	21	-	-	-	-	-	-	-	-	-	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	210	-	-	-	-	-	-	-	1	1	2	4
Male	83	-	-	-	-	-	-	-	1	-	1	1
Female	127	-	-	-	-	-	-	-	-	1	1	3
White, both sexes	156	-	-	-	-	-	-	-	-	1	-	3
Male	60	-	-	-	-	-	-	-	-	-	-	1
Female	96	-	-	-	-	-	-	-	-	1	-	2
Black, both sexes	49	-	-	-	-	-	-	-	1	-	2	1
Male	19	-	-	-	-	-	-	-	1	-	1	-
Female	30	-	-	-	-	-	-	-	-	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Washington												
All races, both sexes	212	1	-	-	-	-	-	1	-	1	1	4
Male	86	-	-	-	-	-	-	-	-	1	-	1
Female	126	1	-	-	-	-	-	1	-	-	1	3
White, both sexes	203	1	-	-	-	-	-	1	-	1	-	3
Male	82	-	-	-	-	-	-	-	-	1	-	1
Female	121	1	-	-	-	-	-	1	-	-	-	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	4	-	-	-	-	-	-	-	-	-	1	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	1
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	92	1	-	-	-	-	-	-	1	-	-	2
Male	44	1	-	-	-	-	-	-	-	-	-	1
Female	48	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	89	1	-	-	-	-	-	-	1	-	-	2
Male	43	1	-	-	-	-	-	-	-	-	-	1
Female	46	-	-	-	-	-	-	-	1	-	-	1
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Wisconsin												
All races, both sexes	230	-	-	-	-	-	-	-	-	2	2	2
Male	103	-	-	-	-	-	-	-	-	2	-	-
Female	127	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	223	-	-	-	-	-	-	-	-	1	2	2
Male	99	-	-	-	-	-	-	-	-	1	-	-
Female	124	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
United States												
All races, both sexes	204	358	452	643	971	1,432	1,764	1,682	1,270	537	108	-
Male	109	208	262	364	521	768	774	619	335	104	19	-
Female	95	150	190	279	450	664	990	1,063	935	433	89	-
White, both sexes	158	266	370	518	808	1,221	1,534	1,513	1,132	468	90	-
Male	82	154	218	292	444	652	682	566	304	94	16	-
Female	76	112	152	226	364	569	852	947	828	374	74	-
Black, both sexes	42	85	74	111	150	188	205	148	127	61	17	-
Male	25	51	40	63	71	101	81	45	29	8	2	-
Female	17	34	34	48	79	87	124	103	98	53	15	-
American Indian 1/, both sexes	1	4	5	5	5	6	4	3	-	1	-	-
Male	-	2	3	2	1	2	2	1	-	-	-	-
Female	1	2	2	3	4	4	2	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes	3	3	3	9	8	17	21	18	11	7	1	-
Male	2	1	1	7	5	13	9	7	2	2	1	-
Female	1	2	2	2	3	4	12	11	9	5	-	-
Alabama												
All races, both sexes	4	6	13	16	19	25	36	37	21	4	-	-
Male	2	2	9	5	9	13	9	14	4	-	-	-
Female	2	4	4	11	10	12	27	23	17	4	-	-
White, both sexes	4	3	10	14	16	20	25	34	14	4	-	-
Male	2	1	7	5	8	9	8	14	3	-	-	-
Female	2	2	3	9	8	11	17	20	11	4	-	-
Black, both sexes	-	2	3	2	3	5	11	3	7	-	-	-
Male	-	1	2	-	1	4	1	-	1	-	-	-
Female	-	1	1	2	2	1	10	3	6	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Alaska												
All races, both sexes	-	-	1	1	-	2	4	2	2	-	-	-
Male	-	-	1	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	3	1	2	-	-	-
White, both sexes	-	-	1	1	-	2	3	2	1	-	-	-
Male	-	-	1	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	1	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Arizona												
All races, both sexes	4	6	4	11	19	24	34	20	12	8	1	-
Male	-	4	1	4	12	10	15	12	1	3	1	-
Female	4	2	3	7	7	14	19	8	11	5	-	-
White, both sexes	4	6	4	11	18	22	31	18	12	7	-	-
Male	-	4	1	4	12	9	15	12	1	3	-	-
Female	4	2	3	7	6	13	16	6	11	4	-	-
Black, both sexes	-	-	-	-	1	-	2	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	2	1	1	-	1	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Arkansas												
All races, both sexes	1	8	18	6	6	18	24	19	19	7	2	-
Male	1	6	11	3	2	10	9	7	3	2	-	-
Female	-	2	7	3	4	8	15	12	16	5	2	-
White, both sexes	1	6	16	6	6	17	22	15	17	7	2	-
Male	1	4	10	3	2	10	7	5	3	2	-	-
Female	-	2	6	3	4	7	15	10	14	5	2	-
Black, both sexes	-	2	2	-	-	1	2	4	2	-	-	-
Male	-	2	1	-	-	-	2	2	-	-	-	-
Female	-	-	1	-	-	1	-	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	14	31	34	54	83	118	134	152	91	40	5	-
Male	9	21	24	36	50	67	61	56	29	7	-	-
Female	5	10	10	18	33	51	73	96	62	33	5	-
White, both sexes	11	24	29	43	70	99	112	136	83	34	4	-
Male	7	16	21	28	43	56	49	54	28	7	-	-
Female	4	8	8	15	27	43	63	82	55	27	4	-
Black, both sexes	1	6	3	6	8	11	12	10	4	4	1	-
Male	1	4	2	4	4	5	6	-	1	-	-	-
Female	-	2	1	2	4	6	6	10	3	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	1	2	5	5	8	10	6	4	2	-	-
Male	1	1	1	4	3	6	6	2	-	-	-	-
Female	1	-	1	1	2	2	4	4	4	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Colorado												
All races, both sexes	5	1	5	3	16	16	24	19	19	6	1	-
Male	3	1	3	1	9	9	10	12	2	1	-	-
Female	2	-	2	2	7	7	14	7	17	5	1	-
White, both sexes	5	1	5	3	15	14	23	19	17	6	1	-
Male	3	1	3	1	8	8	9	12	2	1	-	-
Female	2	-	2	2	7	6	14	7	15	5	1	-
Black, both sexes	-	-	-	-	1	2	1	-	2	-	-	-
Male	-	-	-	-	1	1	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	1	4	6	5	10	22	19	23	15	8	1	-
Male	-	3	4	4	6	15	11	5	3	1	-	-
Female	1	1	2	1	4	7	8	18	12	7	1	-
White, both sexes	1	3	4	5	10	20	18	22	15	8	1	-
Male	-	2	3	4	6	13	10	5	3	1	-	-
Female	1	1	1	1	4	7	8	17	12	7	1	-
Black, both sexes	-	1	2	-	-	1	1	1	-	-	-	-
Male	-	1	1	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Delaware												
All races, both sexes	-	1	1	4	7	9	3	6	6	2	-	-
Male	-	-	1	2	2	8	2	5	1	-	-	-
Female	-	1	-	2	5	1	1	1	5	2	-	-
White, both sexes	-	1	-	3	6	7	3	6	5	2	-	-
Male	-	-	-	1	1	6	2	5	1	-	-	-
Female	-	1	-	2	5	1	1	1	4	2	-	-
Black, both sexes	-	-	1	1	1	2	-	-	1	-	-	-
Male	-	-	1	1	1	2	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	-	2	3	4	2	4	2	3	5	1	1	-
Male	-	-	-	2	1	3	1	1	1	-	-	-
Female	-	2	3	2	1	1	1	2	4	1	1	-
White, both sexes	-	-	1	-	-	1	-	-	2	-	-	-
Male	-	-	-	-	-	1	-	-	1	-	-	-
Female	-	-	1	-	-	-	-	-	1	-	-	-
Black, both sexes	-	2	2	4	2	3	2	3	3	1	1	-
Male	-	-	-	2	1	2	1	1	-	-	-	-
Female	-	2	2	2	1	1	1	2	3	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Florida												
All races, both sexes	16	24	26	46	58	113	124	102	83	29	4	-
Male	9	10	17	26	33	57	66	38	29	8	2	-
Female	7	14	9	20	25	56	58	64	54	21	2	-
White, both sexes	10	18	24	37	46	100	105	93	74	25	3	-
Male	5	6	17	23	27	50	59	35	28	7	2	-
Female	5	12	7	14	19	50	46	58	46	18	1	-
Black, both sexes	6	5	2	7	12	13	19	9	9	4	1	-
Male	4	4	-	3	6	7	7	3	1	1	-	-
Female	2	1	2	4	6	6	12	6	8	3	1	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	8	11	13	21	30	30	41	39	32	19	2	-
Male	4	7	5	14	19	13	17	14	3	3	-	-
Female	4	4	8	7	11	17	24	25	29	16	2	-
White, both sexes	4	6	6	14	19	23	32	28	23	14	2	-
Male	2	5	3	10	12	10	13	9	2	2	-	-
Female	2	1	3	4	7	13	19	19	21	12	2	-
Black, both sexes	4	5	7	6	11	7	9	11	9	5	-	-
Male	2	2	2	3	7	3	4	5	1	1	-	-
Female	2	3	5	3	4	4	5	6	8	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Hawaii												
All races, both sexes	-	1	1	2	1	6	4	5	6	4	-	-
Male	-	-	-	1	1	5	-	2	2	1	-	-
Female	-	1	1	1	-	1	4	3	4	3	-	-
White, both sexes	-	-	-	2	1	2	-	1	2	-	-	-
Male	-	-	-	1	1	2	-	1	1	-	-	-
Female	-	-	-	1	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	1	-	-	4	4	4	4	4	-	-
Male	-	-	-	-	-	3	-	1	1	1	-	-
Female	-	1	1	-	-	1	4	3	3	3	-	-
Idaho												
All races, both sexes	1	1	3	2	6	5	7	8	3	1	-	-
Male	-	-	1	2	5	3	4	1	1	-	-	-
Female	1	1	2	-	1	2	3	7	2	1	-	-
White, both sexes	1	1	3	2	6	5	7	8	3	1	-	-
Male	-	-	1	2	5	3	4	1	1	-	-	-
Female	1	1	2	-	1	2	3	7	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Illinois												
All races, both sexes	7	17	18	29	36	68	81	73	64	26	5	-
Male	5	11	10	18	23	44	29	27	18	4	1	-
Female	2	6	8	11	13	24	52	46	46	22	4	-
White, both sexes	5	11	16	22	29	55	69	64	60	22	4	-
Male	3	8	10	12	19	34	24	23	18	4	-	-
Female	2	3	6	10	10	21	45	41	42	18	4	-
Black, both sexes	2	6	2	7	7	13	11	9	4	4	1	-
Male	2	3	-	6	4	10	5	4	-	-	1	-
Female	-	3	2	1	3	3	6	5	4	4	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Indiana												
All races, both sexes	4	7	10	22	27	32	40	45	30	10	1	-
Male	2	5	5	14	14	22	21	15	10	2	-	-
Female	2	2	5	8	13	10	19	30	20	8	1	-
White, both sexes	2	5	9	18	25	31	38	44	29	10	1	-
Male	1	3	4	11	14	21	20	14	9	2	-	-
Female	1	2	5	7	11	10	18	30	20	8	1	-
Black, both sexes	2	2	1	4	2	1	2	1	1	-	-	-
Male	1	2	1	3	-	1	1	1	1	-	-	-
Female	1	-	-	1	2	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Iowa												
All races, both sexes	1	5	3	1	7	22	24	19	26	9	4	-
Male	-	2	2	1	5	14	10	7	9	1	1	-
Female	1	3	1	-	2	8	14	12	17	8	3	-
White, both sexes	1	5	3	1	7	22	24	19	26	9	4	-
Male	-	2	2	1	5	14	10	7	9	1	1	-
Female	1	3	1	-	2	8	14	12	17	8	3	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	3	6	8	8	12	14	20	15	11	1	-
Male	3	2	4	5	3	9	6	8	5	1	1	-
Female	-	1	2	3	5	3	8	12	10	10	-	-
White, both sexes	2	3	5	8	7	12	12	19	14	11	1	-
Male	2	2	3	5	2	9	5	8	5	1	1	-
Female	-	1	2	3	5	3	7	11	9	10	-	-
Black, both sexes	-	-	1	-	-	-	1	1	1	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Kentucky												
All races, both sexes	3	7	4	11	21	32	29	26	12	6	2	-
Male	2	5	1	8	11	20	14	6	3	1	-	-
Female	1	2	3	3	10	12	15	20	9	5	2	-
White, both sexes	3	7	4	9	17	31	28	23	11	4	1	-
Male	2	5	1	6	9	19	14	6	3	1	-	-
Female	1	2	3	3	8	12	14	17	8	3	1	-
Black, both sexes	-	-	-	2	4	1	1	3	1	2	1	-
Male	-	-	-	2	2	1	-	-	-	-	-	-
Female	-	-	-	-	2	-	1	3	1	2	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	4	8	9	16	22	26	38	26	18	7	3	-
Male	3	4	4	7	11	10	17	14	8	5	1	-
Female	1	4	5	9	11	16	21	12	10	2	2	-
White, both sexes	2	3	4	8	15	16	25	20	9	4	3	-
Male	1	-	3	3	8	5	11	10	1	3	1	-
Female	1	3	1	5	7	11	14	10	8	1	2	-
Black, both sexes	2	5	5	8	7	10	13	5	9	3	-	-
Male	2	4	1	4	3	5	6	3	7	2	-	-
Female	-	1	4	4	4	5	7	2	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Maine												
All races, both sexes	1	1	4	2	3	7	17	10	6	4	-	-
Male	-	1	3	1	2	2	6	8	2	1	-	-
Female	1	-	1	1	1	5	11	2	4	3	-	-
White, both sexes	1	1	4	2	3	7	17	10	6	4	-	-
Male	-	1	3	1	2	2	6	8	2	1	-	-
Female	1	-	1	1	1	5	11	2	4	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	3	7	7	11	17	35	32	28	20	6	1	-
Male	1	4	5	6	7	16	16	7	2	1	-	-
Female	2	3	2	5	10	19	16	21	18	5	1	-
White, both sexes	1	3	6	7	11	23	23	22	14	6	1	-
Male	-	2	5	3	6	8	12	4	2	1	-	-
Female	1	1	1	4	5	15	11	18	12	5	1	-
Black, both sexes	2	4	1	4	6	11	9	6	5	-	-	-
Male	1	2	-	3	1	7	4	3	-	-	-	-
Female	1	2	1	1	5	4	5	3	5	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	1	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Massachusetts												
All races, both sexes	7	4	8	16	27	20	42	44	39	23	4	-
Male	4	3	5	10	16	5	17	18	7	2	-	-
Female	3	1	3	6	11	15	25	26	32	21	4	-
White, both sexes	6	3	7	15	27	19	39	42	38	23	4	-
Male	3	2	5	10	16	5	16	18	7	2	-	-
Female	3	1	2	5	11	14	23	24	31	21	4	-
Black, both sexes	1	1	1	1	-	1	2	2	1	-	-	-
Male	1	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	1	-	1	2	2	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	7	14	14	21	26	51	76	62	48	14	6	-
Male	4	9	6	7	19	25	23	24	19	3	-	-
Female	3	5	8	14	7	26	53	38	29	11	6	-
White, both sexes	6	11	12	13	23	43	66	54	45	11	2	-
Male	4	7	5	3	17	20	22	23	18	3	-	-
Female	2	4	7	10	6	23	44	31	27	8	2	-
Black, both sexes	1	3	-	7	3	8	10	8	3	3	4	-
Male	-	2	-	3	2	5	1	1	1	-	-	-
Female	1	1	-	4	1	3	9	7	2	3	4	-
American Indian 1/, both sexes	-	-	2	1	-	-	-	-	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Minnesota												
All races, both sexes	3	5	3	6	15	20	38	39	31	14	4	-
Male	1	2	2	5	8	15	15	19	11	3	1	-
Female	2	3	1	1	7	5	23	20	20	11	3	-
White, both sexes	3	4	3	5	15	19	38	39	31	14	4	-
Male	1	1	2	5	8	15	15	19	11	3	1	-
Female	2	3	1	-	7	4	23	20	20	11	3	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	4	5	11	5	6	13	16	6	7	5	1	-
Male	2	2	7	1	2	4	4	2	2	1	-	-
Female	2	3	4	4	4	9	12	4	5	4	1	-
White, both sexes	3	2	7	3	3	9	10	4	3	2	-	-
Male	1	1	4	-	2	3	2	2	1	1	-	-
Female	2	1	3	3	1	6	8	2	2	1	-	-
Black, both sexes	1	3	4	2	3	4	6	2	4	3	1	-
Male	1	1	3	1	-	1	2	-	1	-	-	-
Female	-	2	1	1	3	3	4	2	3	3	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Missouri												
All races, both sexes	3	11	7	17	27	36	43	37	33	15	4	-
Male	3	8	3	11	13	18	18	15	10	4	2	-
Female	-	3	4	6	14	18	25	22	23	11	2	-
White, both sexes	3	9	5	15	23	27	38	33	32	14	4	-
Male	3	6	2	10	11	14	17	14	10	4	2	-
Female	-	3	3	5	12	13	21	19	22	10	2	-
Black, both sexes	-	2	2	2	4	8	5	4	1	1	-	-
Male	-	2	1	1	2	4	1	1	-	-	-	-
Female	-	-	1	1	2	4	4	3	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Montana												
All races, both sexes	1	3	1	2	7	4	3	10	2	3	-	-
Male	-	1	1	-	2	3	3	2	2	-	-	-
Female	1	2	-	2	5	1	-	8	-	3	-	-
White, both sexes	1	3	1	2	5	4	3	10	2	3	-	-
Male	-	1	1	-	2	3	3	2	2	-	-	-
Female	1	2	-	2	3	1	-	8	-	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	2	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	2	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Nebraska												
All races, both sexes	1	1	1	2	8	3	11	14	13	4	3	-
Male	-	1	1	2	5	2	6	6	6	2	1	-
Female	1	-	-	-	3	1	5	8	7	2	2	-
White, both sexes	1	1	1	2	8	3	11	14	13	4	3	-
Male	-	1	1	2	5	2	6	6	6	2	1	-
Female	1	-	-	-	3	1	5	8	7	2	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	5	5	3	5	9	10	10	3	-	-	-
Male	-	3	3	1	2	5	5	4	1	-	-	-
Female	1	2	2	2	3	4	5	6	2	-	-	-
White, both sexes	-	5	5	3	4	9	10	10	3	-	-	-
Male	-	3	3	1	2	5	5	4	1	-	-	-
Female	-	2	2	2	2	4	5	6	2	-	-	-
Black, both sexes	1	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
New Hampshire												
All races, both sexes	2	2	-	3	5	5	7	13	8	5	1	-
Male	2	1	-	1	1	5	2	3	3	1	-	-
Female	-	1	-	2	4	-	5	10	5	4	1	-
White, both sexes	2	2	-	3	5	5	7	13	8	5	1	-
Male	2	1	-	1	1	5	2	3	3	1	-	-
Female	-	1	-	2	4	-	5	10	5	4	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	3	10	9	16	36	43	44	55	42	20	3	-
Male	1	1	4	5	14	25	19	16	10	5	2	-
Female	2	9	5	11	22	18	25	39	32	15	1	-
White, both sexes	3	9	5	12	28	37	38	51	36	18	3	-
Male	1	1	3	4	12	21	18	15	8	5	2	-
Female	2	8	2	8	16	16	20	36	28	13	1	-
Black, both sexes	-	1	4	3	6	6	6	4	5	2	-	-
Male	-	-	1	-	1	4	1	1	1	-	-	-
Female	-	1	3	3	5	2	5	3	4	2	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	2	-	-	-	1	-	-	-
Male	-	-	-	-	1	-	-	-	1	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
New Mexico												
All races, both sexes	1	-	3	3	7	8	7	12	4	6	1	-
Male	-	-	3	1	5	3	2	6	2	2	-	-
Female	1	-	-	2	2	5	5	6	2	4	1	-
White, both sexes	1	-	3	3	7	8	7	10	4	6	1	-
Male	-	-	3	1	5	3	2	5	2	2	-	-
Female	1	-	-	2	2	5	5	5	2	4	1	-
Black, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	9	18	26	21	48	66	93	83	67	34	8	-
Male	6	8	16	10	18	33	46	28	20	6	2	-
Female	3	10	10	11	30	33	47	55	47	28	6	-
White, both sexes	7	12	20	16	42	60	79	74	63	29	6	-
Male	5	7	11	9	15	30	38	26	17	5	1	-
Female	2	5	9	7	27	30	41	48	46	24	5	-
Black, both sexes	2	6	6	5	6	6	12	8	4	5	1	-
Male	1	1	5	1	3	3	7	1	3	1	-	-
Female	1	5	1	4	3	3	5	7	1	4	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	2	1	-	-	1	-
Male	-	-	-	-	-	-	1	1	-	-	1	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
North Carolina												
All races, both sexes	15	15	17	20	38	41	53	46	42	17	4	-
Male	9	9	10	11	18	27	26	17	8	3	-	-
Female	6	6	7	9	20	14	27	29	34	14	4	-
White, both sexes	9	9	12	14	26	28	35	34	29	11	3	-
Male	5	4	6	8	12	18	17	13	5	3	-	-
Female	4	5	6	6	14	10	18	21	24	8	3	-
Black, both sexes	5	6	3	6	12	13	18	11	13	6	1	-
Male	4	5	2	3	6	9	9	4	3	-	-	-
Female	1	1	1	3	6	4	9	7	10	6	1	-
American Indian 1/, both sexes	1	-	2	-	-	-	-	-	-	-	-	-
Male	-	-	2	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
North Dakota												
All races, both sexes	-	1	1	3	2	3	7	2	5	4	-	-
Male	-	-	1	-	-	3	5	1	1	1	-	-
Female	-	1	-	3	2	-	2	1	4	3	-	-
White, both sexes	-	-	1	3	2	3	7	2	5	4	-	-
Male	-	-	1	-	-	3	5	1	1	1	-	-
Female	-	-	-	3	2	-	2	1	4	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Ohio												
All races, both sexes	13	18	22	39	45	74	90	80	59	20	4	-
Male	7	12	15	22	27	35	38	29	16	3	-	-
Female	6	6	7	17	18	39	52	51	43	17	4	-
White, both sexes	10	16	20	34	35	59	81	73	56	19	4	-
Male	6	10	14	19	21	28	33	27	14	3	-	-
Female	4	6	6	15	14	31	48	46	42	16	4	-
Black, both sexes	3	2	2	5	10	15	9	7	3	1	-	-
Male	1	2	1	3	6	7	5	2	2	-	-	-
Female	2	-	1	2	4	8	4	5	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	9	9	12	14	20	18	12	11	11	-	-
Male	1	5	2	5	8	12	7	3	3	4	-	-
Female	1	4	7	7	6	8	11	9	8	7	-	-
White, both sexes	1	8	8	10	10	18	17	12	11	11	-	-
Male	1	4	2	4	8	11	6	3	3	4	-	-
Female	-	4	6	6	2	7	11	9	8	7	-	-
Black, both sexes	1	-	-	2	3	1	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	1	3	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	1	1	-	1	1	1	-	-	-	-	-
Male	-	1	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Oregon												
All races, both sexes	1	3	6	8	10	30	26	22	23	4	-	-
Male	1	2	3	4	6	14	12	7	4	-	-	-
Female	-	1	3	4	4	16	14	15	19	4	-	-
White, both sexes	1	3	6	8	9	30	26	21	23	4	-	-
Male	1	2	3	4	5	14	12	7	4	-	-	-
Female	-	1	3	4	4	16	14	14	19	4	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Pennsylvania												
All races, both sexes	9	14	17	34	48	68	91	110	83	33	2	-
Male	4	10	9	22	22	35	36	36	19	4	-	-
Female	5	4	8	12	26	33	55	74	64	29	2	-
White, both sexes	8	11	16	31	45	60	86	105	79	32	2	-
Male	3	8	8	21	21	33	33	35	19	4	-	-
Female	5	3	8	10	24	27	53	70	60	28	2	-
Black, both sexes	1	3	1	3	3	7	5	5	4	1	-	-
Male	1	2	1	1	1	2	3	1	-	-	-	-
Female	-	1	-	2	2	5	2	4	4	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Rhode Island												
All races, both sexes	-	-	1	-	2	7	7	10	3	1	2	-
Male	-	-	-	-	1	4	3	3	-	-	-	-
Female	-	-	1	-	1	3	4	7	3	1	2	-
White, both sexes	-	-	1	-	2	5	7	10	3	1	2	-
Male	-	-	-	-	1	3	3	3	-	-	-	-
Female	-	-	1	-	1	2	4	7	3	1	2	-
Black, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	2	9	11	12	21	28	32	37	23	10	1	-
Male	-	4	8	7	6	17	17	13	3	3	-	-
Female	2	5	3	5	15	11	15	24	20	7	1	-
White, both sexes	2	5	7	7	12	20	20	23	13	7	-	-
Male	-	2	4	4	3	13	13	8	2	1	-	-
Female	2	3	3	3	9	7	7	15	11	6	-	-
Black, both sexes	-	4	4	5	9	8	12	13	10	3	1	-
Male	-	2	4	3	3	4	4	5	1	2	-	-
Female	-	2	-	2	6	4	8	8	9	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
South Dakota												
All races, both sexes	-	-	1	1	1	2	9	9	5	1	1	-
Male	-	-	-	1	1	1	8	2	2	-	-	-
Female	-	-	1	-	-	1	1	7	3	1	1	-
White, both sexes	-	-	1	1	1	1	9	9	5	1	1	-
Male	-	-	-	1	1	1	8	2	2	-	-	-
Female	-	-	1	-	-	-	1	7	3	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	8	9	14	23	23	30	38	38	22	9	2	-
Male	3	5	8	17	16	14	15	15	6	1	-	-
Female	5	4	6	6	7	16	23	23	16	8	2	-
White, both sexes	6	7	12	20	19	26	34	33	17	9	1	-
Male	2	5	6	14	13	12	14	14	4	1	-	-
Female	4	2	6	6	6	14	20	19	13	8	1	-
Black, both sexes	2	2	2	3	4	4	4	5	5	-	1	-
Male	1	-	2	3	3	2	1	1	2	-	-	-
Female	1	2	-	-	1	2	3	4	3	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Texas												
All races, both sexes	18	28	42	51	79	97	117	106	82	32	11	-
Male	11	17	25	28	55	48	50	35	16	4	1	-
Female	7	11	17	23	24	49	67	71	66	28	10	-
White, both sexes	15	20	33	41	63	83	104	99	73	24	9	-
Male	10	12	19	20	43	41	46	32	15	3	1	-
Female	5	8	14	21	20	42	58	67	58	21	8	-
Black, both sexes	3	8	9	10	16	14	12	7	9	8	2	-
Male	1	5	6	8	12	7	4	3	1	1	-	-
Female	2	3	3	2	4	7	8	4	8	7	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	1	1	3	4	11	10	4	1	-	-
Male	-	-	1	1	2	2	5	6	-	1	-	-
Female	-	-	-	-	1	2	6	4	4	-	-	-
White, both sexes	-	-	1	-	3	4	11	9	4	1	-	-
Male	-	-	1	-	2	2	5	5	-	1	-	-
Female	-	-	-	-	1	2	6	4	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Vermont												
All races, both sexes	-	1	1	1	3	3	3	2	5	-	1	-
Male	-	1	-	1	-	1	3	1	1	-	-	-
Female	-	-	1	-	3	2	-	1	4	-	1	-
White, both sexes	-	1	1	1	3	3	3	2	5	-	1	-
Male	-	1	-	1	-	1	3	1	1	-	-	-
Female	-	-	1	-	3	2	-	1	4	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	4	8	11	15	22	32	33	31	33	10	3	-
Male	1	6	5	13	6	16	14	8	10	-	1	-
Female	3	2	6	2	16	16	19	23	23	10	2	-
White, both sexes	2	6	8	10	18	22	24	27	27	5	3	-
Male	-	5	4	9	6	11	9	6	8	-	1	-
Female	2	1	4	1	12	11	15	21	19	5	2	-
Black, both sexes	2	2	3	4	4	9	7	3	6	5	-	-
Male	1	1	1	3	-	4	4	1	2	-	-	-
Female	1	1	2	1	4	5	3	2	4	5	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	2	1	-	-	-	-
Male	-	-	-	1	-	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Washington												
All races, both sexes	5	8	10	12	18	24	48	34	30	12	3	-
Male	2	4	8	6	11	9	20	13	8	3	-	-
Female	3	4	2	6	7	15	28	21	22	9	3	-
White, both sexes	5	7	10	11	16	24	48	32	30	11	3	-
Male	2	3	8	6	10	9	20	12	8	2	-	-
Female	3	4	2	5	6	15	28	20	22	9	3	-
Black, both sexes	-	1	-	1	1	-	-	1	-	-	-	-
Male	-	1	-	-	1	-	-	1	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	1	6	9	6	16	12	17	13	4	2	-
Male	2	1	2	3	2	11	8	9	1	2	1	-
Female	-	-	4	6	4	5	4	8	12	2	1	-
White, both sexes	2	1	6	9	6	15	12	16	12	4	2	-
Male	2	1	2	3	2	11	8	8	1	2	1	-
Female	-	-	4	6	4	4	4	8	11	2	1	-
Black, both sexes	-	-	-	-	-	1	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other diseases of arteries, arterioles and capillaries (I72-I78)												
Wisconsin												
All races, both sexes	2	4	5	12	21	45	47	46	22	17	3	-
Male	-	4	3	8	10	25	21	17	8	4	1	-
Female	2	-	2	4	11	20	26	29	14	13	2	-
White, both sexes	2	3	4	10	21	44	46	46	22	17	3	-
Male	-	3	2	7	10	25	21	17	8	4	1	-
Female	2	-	2	3	11	19	25	29	14	13	2	-
Black, both sexes	-	1	1	1	-	-	1	-	-	-	-	-
Male	-	1	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	1	-	-	-	4	1	3	3	-	-	-
Male	1	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	4	-	2	3	-	-	-
White, both sexes	1	1	-	-	-	4	1	3	3	-	-	-
Male	1	1	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	4	-	2	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
United States												
All races, both sexes	4,813	26	6	4	1	11	25	50	82	124	208	298
Male	2,156	13	3	2	1	4	11	23	39	55	109	180
Female	2,657	13	3	2	-	7	14	27	43	69	99	118
White, both sexes	3,936	21	3	4	1	6	15	35	52	88	135	220
Male	1,723	11	2	2	1	3	5	14	27	42	72	131
Female	2,213	10	1	2	-	3	10	21	25	46	63	89
Black, both sexes	796	4	2	-	-	5	10	15	27	32	67	72
Male	388	2	-	-	-	1	6	9	9	11	33	46
Female	408	2	2	-	-	4	4	6	18	21	34	26
American Indian 1/, both sexes	27	1	1	-	-	-	-	-	-	1	5	4
Male	16	-	1	-	-	-	-	-	-	1	3	2
Female	11	1	-	-	-	-	-	-	-	-	2	2
Asian or Pacific Islander 2/, both sexes	54	-	-	-	-	-	-	-	3	3	1	2
Male	29	-	-	-	-	-	-	-	3	1	1	1
Female	25	-	-	-	-	-	-	-	-	2	-	1
Alabama												
All races, both sexes	94	-	-	-	-	-	-	-	-	3	1	4
Male	39	-	-	-	-	-	-	-	-	1	-	3
Female	55	-	-	-	-	-	-	-	-	2	1	1
White, both sexes	71	-	-	-	-	-	-	-	-	1	-	3
Male	27	-	-	-	-	-	-	-	-	-	-	3
Female	44	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	23	-	-	-	-	-	-	-	-	2	1	1
Male	12	-	-	-	-	-	-	-	-	1	-	-
Female	11	-	-	-	-	-	-	-	-	1	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Alaska												
All races, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	96	-	-	-	-	-	1	-	3	1	6	5
Male	50	-	-	-	-	-	-	-	2	1	1	3
Female	46	-	-	-	-	-	1	-	1	-	5	2
White, both sexes	86	-	-	-	-	-	-	-	3	1	3	4
Male	46	-	-	-	-	-	-	-	2	1	1	2
Female	40	-	-	-	-	-	-	-	1	-	2	2
Black, both sexes	8	-	-	-	-	-	1	-	-	-	3	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	1	-	-	-	3	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Arkansas												
All races, both sexes	51	-	-	-	-	1	-	1	2	1	3	2
Male	21	-	-	-	-	-	-	-	1	-	1	2
Female	30	-	-	-	-	1	-	1	1	1	2	-
White, both sexes	43	-	-	-	-	-	-	-	1	1	2	1
Male	17	-	-	-	-	-	-	-	1	-	-	1
Female	26	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	7	-	-	-	-	1	-	1	1	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	4	-	-	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	541	2	-	-	-	-	4	6	10	25	27	38
Male	243	1	-	-	-	-	-	3	6	8	17	23
Female	298	1	-	-	-	-	4	3	4	17	10	15
White, both sexes	447	2	-	-	-	-	4	4	6	19	17	29
Male	194	1	-	-	-	-	-	1	3	7	10	17
Female	253	1	-	-	-	-	4	3	3	12	7	12
Black, both sexes	74	-	-	-	-	-	-	2	2	6	10	8
Male	40	-	-	-	-	-	-	2	1	1	7	6
Female	34	-	-	-	-	-	-	-	1	5	3	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	19	-	-	-	-	-	-	-	2	-	-	1
Male	9	-	-	-	-	-	-	-	2	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Colorado												
All races, both sexes	57	-	1	-	-	-	-	1	1	1	2	3
Male	25	-	1	-	-	-	-	-	-	-	2	3
Female	32	-	-	-	-	-	-	1	1	1	-	-
White, both sexes	50	-	1	-	-	-	-	1	-	-	1	3
Male	21	-	1	-	-	-	-	-	-	-	1	3
Female	29	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	6	-	-	-	-	-	-	-	1	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	59	1	-	-	-	-	-	1	-	-	-	5
Male	20	1	-	-	-	-	-	-	-	-	-	5
Female	39	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	54	1	-	-	-	-	-	1	-	-	-	5
Male	18	1	-	-	-	-	-	-	-	-	-	5
Female	36	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Delaware												
All races, both sexes	9	-	-	-	-	-	-	-	1	1	-	-
Male	7	-	-	-	-	-	-	-	1	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	15	-	-	-	-	-	1	-	-	1	1	3
Male	9	-	-	-	-	-	1	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	13	-	-	-	-	-	1	-	-	1	1	3
Male	7	-	-	-	-	-	1	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Florida												
All races, both sexes	300	1	-	-	-	-	-	-	3	7	15	16
Male	137	1	-	-	-	-	-	-	-	4	8	8
Female	163	-	-	-	-	-	-	-	3	3	7	8
White, both sexes	250	1	-	-	-	-	-	-	2	6	11	13
Male	118	1	-	-	-	-	-	-	-	4	5	7
Female	132	-	-	-	-	-	-	-	2	2	6	6
Black, both sexes	48	-	-	-	-	-	-	-	1	1	4	3
Male	18	-	-	-	-	-	-	-	-	-	3	1
Female	30	-	-	-	-	-	-	-	1	1	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	98	-	1	-	-	-	-	3	2	2	2	10
Male	43	-	-	-	-	-	-	2	1	-	2	6
Female	55	-	1	-	-	-	-	1	1	2	-	4
White, both sexes	61	-	-	-	-	-	-	2	1	2	-	5
Male	24	-	-	-	-	-	-	1	1	-	-	2
Female	37	-	-	-	-	-	-	1	-	2	-	3
Black, both sexes	37	-	1	-	-	-	-	1	1	-	2	5
Male	19	-	-	-	-	-	-	1	-	-	2	4
Female	18	-	1	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Hawaii												
All races, both sexes	19	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	19	-	-	-	-	-	-	-	1	-	2	1
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	19	-	-	-	-	-	-	-	1	-	2	1
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	2	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Illinois												
All races, both sexes	219	2	1	1	-	-	2	4	7	9	7	11
Male	104	1	1	1	-	-	1	2	4	6	5	8
Female	115	1	-	-	-	-	1	2	3	3	2	3
White, both sexes	164	2	1	1	-	-	1	3	2	6	3	5
Male	74	1	1	1	-	-	-	1	1	4	3	3
Female	90	1	-	-	-	-	1	2	1	2	-	2
Black, both sexes	53	-	-	-	-	-	1	1	5	3	4	6
Male	29	-	-	-	-	-	1	1	3	2	2	5
Female	24	-	-	-	-	-	-	-	2	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	73	-	-	-	-	1	-	-	-	3	2	2
Male	29	-	-	-	-	-	-	-	-	1	2	2
Female	44	-	-	-	-	1	-	-	-	2	-	-
White, both sexes	66	-	-	-	-	-	-	-	-	2	2	1
Male	27	-	-	-	-	-	-	-	-	1	2	1
Female	39	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	6	-	-	-	-	1	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Iowa												
All races, both sexes	50	-	-	-	-	1	1	-	-	2	2	2
Male	22	-	-	-	-	-	-	-	-	-	1	1
Female	28	-	-	-	-	1	1	-	-	2	1	1
White, both sexes	49	-	-	-	-	1	1	-	-	2	2	1
Male	21	-	-	-	-	-	-	-	-	-	1	-
Female	28	-	-	-	-	1	1	-	-	2	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	34	-	-	-	-	-	-	1	-	-	5	3
Male	13	-	-	-	-	-	-	-	-	-	2	-
Female	21	-	-	-	-	-	-	1	-	-	3	3
White, both sexes	34	-	-	-	-	-	-	1	-	-	5	3
Male	13	-	-	-	-	-	-	-	-	-	2	-
Female	21	-	-	-	-	-	-	1	-	-	3	3
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Kentucky												
All races, both sexes	61	-	-	-	-	-	1	-	-	1	-	5
Male	29	-	-	-	-	-	-	-	-	1	-	3
Female	32	-	-	-	-	-	1	-	-	-	-	2
White, both sexes	54	-	-	-	-	-	1	-	-	1	-	4
Male	25	-	-	-	-	-	-	-	-	1	-	2
Female	29	-	-	-	-	-	1	-	-	-	-	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	57	-	-	-	-	-	-	1	-	1	1	5
Male	23	-	-	-	-	-	-	1	-	1	1	4
Female	34	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	35	-	-	-	-	-	-	-	-	-	1	2
Male	14	-	-	-	-	-	-	-	-	-	1	2
Female	21	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	22	-	-	-	-	-	-	1	-	1	-	3
Male	9	-	-	-	-	-	-	1	-	1	-	2
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Maine												
All races, both sexes	22	-	-	-	-	-	-	-	-	1	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	21	-	-	-	-	-	-	-	-	1	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	96	-	-	-	-	2	-	4	3	6	5	14
Male	30	-	-	-	-	1	-	3	1	1	-	8
Female	66	-	-	-	-	1	-	1	2	5	5	6
White, both sexes	53	-	-	-	-	1	-	1	1	2	3	6
Male	19	-	-	-	-	1	-	1	1	1	-	4
Female	34	-	-	-	-	-	-	-	-	1	3	2
Black, both sexes	43	-	-	-	-	1	-	3	2	4	2	8
Male	11	-	-	-	-	-	-	2	-	-	-	4
Female	32	-	-	-	-	1	-	1	2	4	2	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Massachusetts												
All races, both sexes	87	1	-	-	-	-	-	-	-	-	2	4
Male	46	-	-	-	-	-	-	-	-	-	2	4
Female	41	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	82	1	-	-	-	-	-	-	-	-	2	4
Male	42	-	-	-	-	-	-	-	-	-	2	4
Female	40	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	208	1	-	-	-	-	2	4	5	6	13	18
Male	101	1	-	-	-	-	-	3	1	4	5	9
Female	107	-	-	-	-	-	2	1	4	2	8	9
White, both sexes	165	1	-	-	-	-	1	3	3	5	6	14
Male	76	1	-	-	-	-	-	2	-	3	3	7
Female	89	-	-	-	-	-	1	1	3	2	3	7
Black, both sexes	40	-	-	-	-	-	1	1	2	-	7	4
Male	22	-	-	-	-	-	-	1	1	-	2	2
Female	18	-	-	-	-	-	1	-	1	-	5	2
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Minnesota												
All races, both sexes	57	2	-	-	-	-	-	-	1	-	1	6
Male	19	-	-	-	-	-	-	-	-	-	1	1
Female	38	2	-	-	-	-	-	-	1	-	-	5
White, both sexes	55	2	-	-	-	-	-	-	1	-	1	5
Male	18	-	-	-	-	-	-	-	-	-	1	1
Female	37	2	-	-	-	-	-	-	1	-	-	4
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	48	-	-	-	-	-	-	-	-	-	5	6
Male	28	-	-	-	-	-	-	-	-	-	3	4
Female	20	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	33	-	-	-	-	-	-	-	-	-	4	2
Male	20	-	-	-	-	-	-	-	-	-	2	2
Female	13	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	14	-	-	-	-	-	-	-	-	-	1	4
Male	8	-	-	-	-	-	-	-	-	-	1	2
Female	6	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Missouri												
All races, both sexes	106	-	-	1	-	1	-	3	4	2	6	2
Male	42	-	-	-	-	-	-	-	1	2	2	2
Female	64	-	-	1	-	1	-	3	3	-	4	-
White, both sexes	93	-	-	1	-	1	-	2	3	1	4	2
Male	35	-	-	-	-	-	-	-	1	1	2	2
Female	58	-	-	1	-	1	-	2	2	-	2	-
Black, both sexes	13	-	-	-	-	-	-	1	1	1	2	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	6	-	-	-	-	-	-	1	1	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	1	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	7	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	14	-	-	-	-	-	-	-	-	-	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Nebraska												
All races, both sexes	29	-	-	-	-	-	-	-	1	-	-	2
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	29	-	-	-	-	-	-	-	1	-	-	2
Male	12	-	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	27	-	-	-	-	-	-	1	1	2	1	1
Male	13	-	-	-	-	-	-	1	-	2	1	-
Female	14	-	-	-	-	-	-	-	1	-	-	1
White, both sexes	26	-	-	-	-	-	-	1	-	2	1	1
Male	13	-	-	-	-	-	-	1	-	2	1	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
New Hampshire												
All races, both sexes	28	-	-	-	-	-	-	-	1	-	1	3
Male	13	-	-	-	-	-	-	-	-	-	1	3
Female	15	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	1	-	1	3
Male	13	-	-	-	-	-	-	-	-	-	1	3
Female	15	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	169	-	-	-	1	-	1	-	2	3	7	10
Male	78	-	-	-	1	-	1	-	1	2	4	6
Female	91	-	-	-	-	-	-	-	1	1	3	4
White, both sexes	138	-	-	-	1	-	-	-	-	2	5	9
Male	62	-	-	-	1	-	-	-	-	2	3	6
Female	76	-	-	-	-	-	-	-	-	-	2	3
Black, both sexes	27	-	-	-	-	-	1	-	2	1	2	1
Male	15	-	-	-	-	-	1	-	1	-	1	-
Female	12	-	-	-	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
New Mexico												
All races, both sexes	41	1	-	-	-	-	-	1	1	1	3	3
Male	21	-	-	-	-	-	-	-	-	-	2	2
Female	20	1	-	-	-	-	-	1	1	1	1	1
White, both sexes	36	-	-	-	-	-	-	1	1	1	2	2
Male	19	-	-	-	-	-	-	-	-	-	1	2
Female	17	-	-	-	-	-	-	1	1	1	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	3	1	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	1	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	331	1	-	1	-	1	2	5	6	12	15	16
Male	159	1	-	-	-	-	1	3	4	7	6	9
Female	172	-	-	1	-	1	1	2	2	5	9	7
White, both sexes	256	1	-	1	-	-	-	2	5	9	11	11
Male	123	1	-	-	-	-	-	2	4	5	5	7
Female	133	-	-	1	-	-	-	-	1	4	6	4
Black, both sexes	71	-	-	-	-	1	2	3	1	2	4	4
Male	34	-	-	-	-	-	1	1	-	2	1	1
Female	37	-	-	-	-	1	1	2	1	-	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	1	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
North Carolina												
All races, both sexes	162	4	-	-	-	-	2	-	4	4	6	8
Male	77	3	-	-	-	-	1	-	1	1	3	5
Female	85	1	-	-	-	-	1	-	3	3	3	3
White, both sexes	110	2	-	-	-	-	1	-	3	4	1	6
Male	51	1	-	-	-	-	1	-	1	1	1	4
Female	59	1	-	-	-	-	-	-	2	3	-	2
Black, both sexes	48	2	-	-	-	-	1	-	1	-	4	2
Male	25	2	-	-	-	-	-	-	-	-	2	1
Female	23	-	-	-	-	-	1	-	1	-	2	1
American Indian 1/, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	14	-	-	-	-	-	1	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	1	-	-	-	-	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Ohio												
All races, both sexes	255	1	-	1	-	1	2	3	3	6	7	21
Male	109	1	-	1	-	1	2	1	3	1	5	13
Female	146	-	-	-	-	-	-	2	-	5	2	8
White, both sexes	217	1	-	1	-	1	1	3	3	4	4	16
Male	91	1	-	1	-	1	1	1	3	1	4	8
Female	126	-	-	-	-	-	-	2	-	3	-	8
Black, both sexes	38	-	-	-	-	-	1	-	-	2	3	5
Male	18	-	-	-	-	-	1	-	-	-	1	5
Female	20	-	-	-	-	-	-	-	-	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	57	-	-	-	-	1	-	-	1	-	7	3
Male	27	-	-	-	-	1	-	-	-	-	4	2
Female	30	-	-	-	-	-	-	-	1	-	3	1
White, both sexes	49	-	-	-	-	1	-	-	1	-	4	3
Male	22	-	-	-	-	1	-	-	-	-	1	2
Female	27	-	-	-	-	-	-	-	1	-	3	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	2	-
Male	3	-	-	-	-	-	-	-	-	-	2	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Oregon												
All races, both sexes	64	1	-	-	-	-	-	1	1	1	3	3
Male	26	-	-	-	-	-	-	1	1	-	-	1
Female	38	1	-	-	-	-	-	-	-	1	3	2
White, both sexes	64	1	-	-	-	-	-	1	1	1	3	3
Male	26	-	-	-	-	-	-	1	1	-	-	1
Female	38	1	-	-	-	-	-	-	-	1	3	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	254	2	-	-	-	-	1	3	4	6	15	14
Male	103	-	-	-	-	-	1	1	1	5	8	8
Female	151	2	-	-	-	-	-	2	3	1	7	6
White, both sexes	214	2	-	-	-	-	1	3	3	3	10	12
Male	82	-	-	-	-	-	1	1	1	2	6	7
Female	132	2	-	-	-	-	-	2	2	1	4	5
Black, both sexes	39	-	-	-	-	-	-	-	1	2	5	2
Male	20	-	-	-	-	-	-	-	-	2	2	1
Female	19	-	-	-	-	-	-	-	1	-	3	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Rhode Island												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	80	-	-	-	-	-	1	1	-	-	4	6
Male	45	-	-	-	-	-	1	1	-	-	3	4
Female	35	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	56	-	-	-	-	-	1	1	-	-	2	3
Male	31	-	-	-	-	-	1	1	-	-	2	1
Female	25	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	24	-	-	-	-	-	-	-	-	-	2	3
Male	14	-	-	-	-	-	-	-	-	-	1	3
Female	10	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
South Dakota												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	2	-
Male	8	-	-	-	-	-	-	-	-	-	2	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	132	-	-	-	-	-	2	1	2	2	6	10
Male	58	-	-	-	-	-	2	-	-	2	4	4
Female	74	-	-	-	-	-	-	1	2	-	2	6
White, both sexes	104	-	-	-	-	-	1	1	1	2	5	7
Male	49	-	-	-	-	-	1	-	-	2	4	3
Female	55	-	-	-	-	-	-	1	1	-	1	4
Black, both sexes	28	-	-	-	-	-	1	-	1	-	1	3
Male	9	-	-	-	-	-	1	-	-	-	-	1
Female	19	-	-	-	-	-	-	-	1	-	1	2
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Texas												
All races, both sexes	288	1	1	-	-	1	1	1	5	4	8	13
Male	119	1	-	-	-	-	-	1	4	1	6	9
Female	169	-	1	-	-	1	1	-	1	3	2	4
White, both sexes	244	1	1	-	-	1	1	1	2	2	6	12
Male	96	1	-	-	-	-	-	1	2	1	4	9
Female	148	-	1	-	-	1	1	-	-	1	2	3
Black, both sexes	40	-	-	-	-	-	-	-	3	2	1	1
Male	22	-	-	-	-	-	-	-	2	-	1	-
Female	18	-	-	-	-	-	-	-	1	2	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	23	2	1	-	-	-	-	-	1	-	-	-
Male	13	2	1	-	-	-	-	-	1	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	2	-	-	-	-	-	-	1	-	-	-
Male	12	2	-	-	-	-	-	-	1	-	-	-
Female	10	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Vermont												
All races, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	7	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	113	2	1	-	-	1	-	3	3	2	6	5
Male	47	-	-	-	-	1	-	-	-	-	2	3
Female	66	2	1	-	-	-	-	3	3	2	4	2
White, both sexes	80	-	-	-	-	-	-	2	3	-	3	4
Male	34	-	-	-	-	-	-	-	-	-	1	2
Female	46	-	-	-	-	-	-	2	3	-	2	2
Black, both sexes	32	2	1	-	-	1	-	1	-	2	3	1
Male	12	-	-	-	-	1	-	-	-	-	1	1
Female	20	2	1	-	-	-	-	1	-	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Washington												
All races, both sexes	99	1	-	-	-	-	-	-	1	4	2	5
Male	44	-	-	-	-	-	-	-	1	1	1	4
Female	55	1	-	-	-	-	-	-	-	3	1	1
White, both sexes	93	1	-	-	-	-	-	-	-	4	2	5
Male	41	-	-	-	-	-	-	-	-	1	1	4
Female	52	1	-	-	-	-	-	-	-	3	1	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	34	-	-	-	-	-	-	-	2	-	2	2
Male	17	-	-	-	-	-	-	-	2	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	2	2
White, both sexes	33	-	-	-	-	-	-	-	1	-	2	2
Male	16	-	-	-	-	-	-	-	1	-	-	-
Female	17	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other disorders of circulatory system (I80-I99)												
Wisconsin												
All races, both sexes	77	-	-	-	-	-	-	1	-	3	1	5
Male	29	-	-	-	-	-	-	-	-	2	1	4
Female	48	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	70	-	-	-	-	-	-	1	-	3	1	4
Male	24	-	-	-	-	-	-	-	-	2	1	3
Female	46	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	5	-	-	-	-	-	-	-	-	-	-	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
United States												
All races, both sexes	322	333	308	336	377	517	611	610	395	151	18	-
Male	202	197	184	188	191	207	239	180	104	24	-	-
Female	120	136	124	148	186	310	372	430	291	127	18	-
White, both sexes	237	256	236	274	310	438	530	560	361	139	15	-
Male	155	154	141	152	153	177	205	164	92	20	-	-
Female	82	102	95	122	157	261	325	396	269	119	15	-
Black, both sexes	80	72	65	54	59	69	72	46	31	11	3	-
Male	43	39	38	33	32	27	32	14	9	4	-	-
Female	37	33	27	21	27	42	40	32	22	7	3	-
American Indian 1/, both sexes	3	-	2	3	1	3	3	-	-	-	-	-
Male	3	-	2	1	1	2	-	-	-	-	-	-
Female	-	-	-	2	-	1	3	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	5	5	5	7	7	6	4	3	1	-	-
Male	1	4	3	2	5	1	2	2	3	-	-	-
Female	1	1	2	3	2	6	4	2	-	1	-	-
Alabama												
All races, both sexes	2	9	10	9	7	15	14	9	8	2	1	-
Male	1	7	5	5	1	9	3	2	2	-	-	-
Female	1	2	5	4	6	6	11	7	6	2	1	-
White, both sexes	2	5	4	7	7	10	13	9	7	2	1	-
Male	1	4	2	3	1	6	3	2	2	-	-	-
Female	1	1	2	4	6	4	10	7	5	2	1	-
Black, both sexes	-	4	6	2	-	5	1	-	1	-	-	-
Male	-	3	3	2	-	3	-	-	-	-	-	-
Female	-	1	3	-	-	2	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	11	4	8	10	7	10	11	9	6	4	-	-
Male	9	2	4	8	5	3	5	4	1	2	-	-
Female	2	2	4	2	2	7	6	5	5	2	-	-
White, both sexes	10	3	8	8	6	10	11	9	6	4	-	-
Male	8	2	4	6	5	3	5	4	1	2	-	-
Female	2	1	4	2	1	7	6	5	5	2	-	-
Black, both sexes	1	1	-	1	1	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Arkansas												
All races, both sexes	4	-	3	1	4	7	7	4	9	2	-	-
Male	2	-	3	1	1	2	3	3	1	1	-	-
Female	2	-	-	-	3	5	4	1	8	1	-	-
White, both sexes	4	-	3	1	4	7	6	3	8	2	-	-
Male	2	-	3	1	1	2	2	2	1	1	-	-
Female	2	-	-	-	3	5	4	1	7	1	-	-
Black, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	40	43	31	34	39	49	63	71	42	15	2	-
Male	22	26	16	18	23	21	24	27	7	1	-	-
Female	18	17	15	16	16	28	39	44	35	14	2	-
White, both sexes	29	35	23	30	31	45	52	63	42	14	2	-
Male	16	22	10	16	19	20	20	24	7	1	-	-
Female	13	13	13	14	12	25	32	39	35	13	2	-
Black, both sexes	10	8	5	3	3	3	7	7	-	-	-	-
Male	5	4	4	2	1	1	4	2	-	-	-	-
Female	5	4	1	1	2	2	3	5	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	3	1	5	1	3	1	-	1	-	-
Male	1	-	2	-	3	-	-	1	-	-	-	-
Female	-	-	1	1	2	1	3	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Colorado												
All races, both sexes	3	1	8	5	4	7	5	6	5	3	1	-
Male	2	1	5	2	4	1	2	1	1	-	-	-
Female	1	-	3	3	-	6	3	5	4	3	1	-
White, both sexes	3	1	8	3	3	7	5	6	5	2	1	-
Male	2	1	5	-	3	1	2	1	1	-	-	-
Female	1	-	3	3	-	6	3	5	4	2	1	-
Black, both sexes	-	-	-	2	1	-	-	-	-	1	-	-
Male	-	-	-	2	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3	4	4	3	1	4	11	10	9	3	-	-
Male	2	1	2	1	-	1	3	4	-	-	-	-
Female	1	3	2	2	1	3	8	6	9	3	-	-
White, both sexes	3	4	3	3	-	4	8	10	9	3	-	-
Male	2	1	1	1	-	1	2	4	-	-	-	-
Female	1	3	2	2	-	3	6	6	9	3	-	-
Black, both sexes	-	-	1	-	1	-	3	-	-	-	-	-
Male	-	-	1	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Delaware												
All races, both sexes	-	1	1	1	-	1	1	1	1	-	-	-
Male	-	1	1	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	1	1	-	1	-	1	-	-	-	-
Male	-	-	1	1	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	3	1	-	-	1	1	3	-	-	-	-	-
Male	2	1	-	-	1	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	2	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	1	-	-	1	1	2	-	-	-	-	-
Male	1	1	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Florida												
All races, both sexes	22	20	24	22	29	30	41	40	21	7	2	-
Male	14	15	13	15	13	11	17	10	5	3	-	-
Female	8	5	11	7	16	19	24	30	16	4	2	-
White, both sexes	18	16	21	16	21	27	40	33	19	5	1	-
Male	13	13	13	10	9	11	17	8	5	2	-	-
Female	5	3	8	6	12	16	23	25	14	3	1	-
Black, both sexes	3	4	3	5	8	3	1	7	2	2	1	-
Male	1	2	-	4	4	-	-	2	-	1	-	-
Female	2	2	3	1	4	3	1	5	2	1	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	9	4	6	7	9	11	11	11	5	4	1	-
Male	6	2	4	7	4	3	4	1	1	-	-	-
Female	3	2	2	-	5	8	7	10	4	4	1	-
White, both sexes	5	2	3	4	7	8	7	8	2	4	1	-
Male	3	1	2	4	3	3	3	1	-	-	-	-
Female	2	1	1	-	4	5	4	7	2	4	1	-
Black, both sexes	4	2	3	3	2	3	4	3	3	-	-	-
Male	3	1	2	3	1	-	1	-	1	-	-	-
Female	1	1	1	-	1	3	3	3	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Hawaii												
All races, both sexes	-	3	-	1	3	1	6	1	3	-	-	-
Male	-	3	-	-	1	-	3	1	3	-	-	-
Female	-	-	-	1	2	1	3	-	-	-	-	-
White, both sexes	-	-	-	-	2	1	4	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	2	1	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	3	-	1	1	-	2	1	3	-	-	-
Male	-	3	-	-	1	-	1	1	3	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Idaho												
All races, both sexes	1	2	1	-	-	1	5	2	3	-	-	-
Male	1	2	1	-	-	1	3	1	2	-	-	-
Female	-	-	-	-	-	-	2	1	1	-	-	-
White, both sexes	1	2	1	-	-	1	5	2	3	-	-	-
Male	1	2	1	-	-	1	3	1	2	-	-	-
Female	-	-	-	-	-	-	2	1	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Illinois												
All races, both sexes	13	14	14	18	19	27	15	35	12	7	1	-
Male	8	8	9	11	10	11	7	7	3	1	-	-
Female	5	6	5	7	9	16	8	28	9	6	1	-
White, both sexes	9	11	6	13	16	23	13	30	12	6	1	-
Male	8	6	4	7	9	10	6	5	3	1	-	-
Female	1	5	2	6	7	13	7	25	9	5	1	-
Black, both sexes	4	3	7	5	3	4	2	4	-	1	-	-
Male	-	2	4	4	1	1	1	2	-	-	-	-
Female	4	1	3	1	2	3	1	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Indiana												
All races, both sexes	4	4	4	10	7	8	10	8	5	5	-	-
Male	4	4	1	4	2	2	4	2	1	-	-	-
Female	-	-	3	6	5	6	6	6	4	5	-	-
White, both sexes	4	4	4	10	6	7	9	7	5	5	-	-
Male	4	4	1	4	1	2	4	2	1	-	-	-
Female	-	-	3	6	5	5	5	5	4	5	-	-
Black, both sexes	-	-	-	-	1	1	1	1	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Iowa												
All races, both sexes	2	5	4	4	2	5	4	5	8	3	-	-
Male	2	5	3	3	1	1	2	-	3	-	-	-
Female	-	-	1	1	1	4	2	5	5	3	-	-
White, both sexes	2	5	4	4	2	5	4	5	8	3	-	-
Male	2	5	3	3	1	1	2	-	3	-	-	-
Female	-	-	1	1	1	4	2	5	5	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	3	1	1	1	2	3	2	5	4	2	1	-
Male	1	-	-	-	2	2	1	1	3	1	-	-
Female	2	1	1	1	-	1	1	4	1	1	1	-
White, both sexes	3	1	1	1	2	3	2	5	4	2	1	-
Male	1	-	-	-	2	2	1	1	3	1	-	-
Female	2	1	1	1	-	1	1	4	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Kentucky												
All races, both sexes	5	4	5	4	3	12	9	5	7	-	-	-
Male	3	1	4	-	1	8	4	2	2	-	-	-
Female	2	3	1	4	2	4	5	3	5	-	-	-
White, both sexes	3	4	4	3	3	10	9	5	7	-	-	-
Male	2	1	3	-	1	7	4	2	2	-	-	-
Female	1	3	1	3	2	3	5	3	5	-	-	-
Black, both sexes	1	-	1	1	-	2	-	-	-	-	-	-
Male	-	-	1	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	8	4	3	3	4	9	6	7	3	1	1	-
Male	1	1	2	1	3	1	4	3	-	-	-	-
Female	7	3	1	2	1	8	2	4	3	1	1	-
White, both sexes	5	1	1	1	2	6	5	6	3	1	1	-
Male	1	-	1	1	1	1	3	3	-	-	-	-
Female	4	1	-	-	1	5	2	3	3	1	1	-
Black, both sexes	3	3	2	2	2	3	1	1	-	-	-	-
Male	-	1	1	-	2	-	1	-	-	-	-	-
Female	3	2	1	2	-	3	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Maine												
All races, both sexes	1	2	-	2	3	3	2	4	2	1	-	-
Male	1	-	-	-	1	2	-	1	-	-	-	-
Female	-	2	-	2	2	1	2	3	2	1	-	-
White, both sexes	1	2	-	2	3	3	2	4	2	1	-	-
Male	1	-	-	-	1	2	-	1	-	-	-	-
Female	-	2	-	2	2	1	2	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5	6	5	3	3	5	9	13	7	4	2	-
Male	3	2	3	2	-	-	1	3	2	-	-	-
Female	2	4	2	1	3	5	8	10	5	4	2	-
White, both sexes	1	4	3	1	-	2	6	12	7	3	-	-
Male	1	1	2	1	-	-	1	3	2	-	-	-
Female	-	3	1	-	-	2	5	9	5	3	-	-
Black, both sexes	4	2	2	2	3	3	3	1	-	1	2	-
Male	2	1	1	1	-	-	-	-	-	-	-	-
Female	2	1	1	1	3	3	3	1	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Massachusetts												
All races, both sexes	1	4	1	7	9	10	14	16	14	4	-	-
Male	1	3	1	5	7	5	7	5	6	-	-	-
Female	-	1	-	2	2	5	7	11	8	4	-	-
White, both sexes	1	4	1	5	9	9	13	16	13	4	-	-
Male	1	3	1	3	7	5	6	5	5	-	-	-
Female	-	1	-	2	2	4	7	11	8	4	-	-
Black, both sexes	-	-	-	2	-	1	1	-	1	-	-	-
Male	-	-	-	2	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	16	20	13	13	14	21	22	30	6	4	-	-
Male	10	13	8	7	9	8	8	12	2	1	-	-
Female	6	7	5	6	5	13	14	18	4	3	-	-
White, both sexes	10	17	13	13	11	17	17	26	5	3	-	-
Male	8	10	8	7	6	6	4	10	1	-	-	-
Female	2	7	5	6	5	11	13	16	4	3	-	-
Black, both sexes	5	3	-	-	3	3	5	4	1	1	-	-
Male	1	3	-	-	3	1	4	2	1	1	-	-
Female	4	-	-	-	-	2	1	2	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	1	-	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Minnesota												
All races, both sexes	1	4	5	5	4	4	8	9	5	2	-	-
Male	-	1	3	-	3	2	3	3	2	-	-	-
Female	1	3	2	5	1	2	5	6	3	2	-	-
White, both sexes	1	3	5	5	4	4	8	9	5	2	-	-
Male	-	-	3	-	3	2	3	3	2	-	-	-
Female	1	3	2	5	1	2	5	6	3	2	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	3	4	5	2	6	7	8	-	-	-	-
Male	2	1	4	4	1	2	4	3	-	-	-	-
Female	-	2	-	1	1	4	3	5	-	-	-	-
White, both sexes	1	3	2	5	1	3	7	5	-	-	-	-
Male	1	1	2	4	1	1	4	2	-	-	-	-
Female	-	2	-	1	-	2	3	3	-	-	-	-
Black, both sexes	1	-	2	-	1	2	-	3	-	-	-	-
Male	1	-	2	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	1	1	-	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Missouri												
All races, both sexes	6	5	4	9	13	11	14	10	11	4	-	-
Male	2	4	2	4	11	4	6	-	2	-	-	-
Female	4	1	2	5	2	7	8	10	9	4	-	-
White, both sexes	6	3	3	8	13	9	13	10	10	4	-	-
Male	2	2	1	4	11	3	5	-	1	-	-	-
Female	4	1	2	4	2	6	8	10	9	4	-	-
Black, both sexes	-	2	1	1	-	2	1	-	1	-	-	-
Male	-	2	1	-	-	1	1	-	1	-	-	-
Female	-	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	1	4	2	2	4	-	-	-
Male	1	-	-	-	1	3	1	1	1	-	-	-
Female	-	-	-	-	-	1	1	1	3	-	-	-
White, both sexes	1	-	-	-	1	3	2	2	4	-	-	-
Male	1	-	-	-	1	3	1	1	1	-	-	-
Female	-	-	-	-	-	-	1	1	3	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Nebraska												
All races, both sexes	2	-	2	1	2	2	5	4	6	2	-	-
Male	2	-	2	-	-	2	2	1	2	-	-	-
Female	-	-	-	1	2	-	3	3	4	2	-	-
White, both sexes	2	-	2	1	2	2	5	4	6	2	-	-
Male	2	-	2	-	-	2	2	1	2	-	-	-
Female	-	-	-	1	2	-	3	3	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	2	1	1	4	3	-	3	4	3	-	-	-
Male	-	-	1	2	1	-	1	2	2	-	-	-
Female	2	1	-	2	2	-	2	2	1	-	-	-
White, both sexes	2	1	1	4	3	-	3	4	3	-	-	-
Male	-	-	1	2	1	-	1	2	2	-	-	-
Female	2	1	-	2	2	-	2	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
New Hampshire												
All races, both sexes	2	2	1	-	3	4	1	6	4	-	-	-
Male	2	-	1	-	2	1	1	2	-	-	-	-
Female	-	2	-	-	1	3	-	4	4	-	-	-
White, both sexes	2	2	1	-	3	4	1	6	4	-	-	-
Male	2	-	1	-	2	1	1	2	-	-	-	-
Female	-	2	-	-	1	3	-	4	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	11	14	11	9	7	27	32	15	12	6	1	-
Male	6	10	7	4	4	11	10	4	4	3	-	-
Female	5	4	4	5	3	16	22	11	8	3	1	-
White, both sexes	10	11	10	8	6	22	24	13	11	5	1	-
Male	5	8	6	4	3	8	8	3	3	2	-	-
Female	5	3	4	4	3	14	16	10	8	3	1	-
Black, both sexes	1	2	1	1	1	3	7	2	1	1	-	-
Male	1	1	1	-	1	3	2	1	1	1	-	-
Female	-	1	-	1	-	-	5	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	2	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	2	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
New Mexico												
All races, both sexes	1	2	3	3	2	4	7	2	6	1	-	-
Male	1	2	3	2	1	1	4	-	3	-	-	-
Female	-	-	-	1	1	3	3	2	3	1	-	-
White, both sexes	1	2	2	3	2	4	6	2	6	1	-	-
Male	1	2	2	2	1	1	4	-	3	-	-	-
Female	-	-	-	1	1	3	2	2	3	1	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	22	33	17	16	40	43	44	35	14	8	-	-
Male	12	18	13	10	21	22	17	10	4	1	-	-
Female	10	15	4	6	19	21	27	25	10	7	-	-
White, both sexes	17	23	14	12	29	34	37	31	11	8	-	-
Male	8	14	12	8	12	17	15	8	4	1	-	-
Female	9	9	2	4	17	17	22	23	7	7	-	-
Black, both sexes	5	10	3	4	10	8	7	4	3	-	-	-
Male	4	4	1	2	8	5	2	2	-	-	-	-
Female	1	6	2	2	2	3	5	2	3	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
North Carolina												
All races, both sexes	13	11	9	10	9	17	20	24	16	2	3	-
Male	11	8	6	6	5	9	8	7	3	-	-	-
Female	2	3	3	4	4	8	12	17	13	2	3	-
White, both sexes	11	6	5	6	7	9	14	22	9	1	3	-
Male	10	4	3	4	3	4	5	7	2	-	-	-
Female	1	2	2	2	4	5	9	15	7	1	3	-
Black, both sexes	2	5	3	3	2	7	6	2	7	1	-	-
Male	1	4	3	2	2	4	3	-	1	-	-	-
Female	1	1	-	1	-	3	3	2	6	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	1	1	2	-	4	4	-	-	-	-
Male	1	-	1	-	2	-	3	1	-	-	-	-
Female	-	-	-	1	-	-	1	3	-	-	-	-
White, both sexes	1	-	1	-	2	-	4	4	-	-	-	-
Male	1	-	1	-	2	-	3	1	-	-	-	-
Female	-	-	-	-	-	-	1	3	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Ohio												
All races, both sexes	10	13	25	17	15	21	32	40	28	9	-	-
Male	5	4	12	8	8	7	13	16	6	2	-	-
Female	5	9	13	9	7	14	19	24	22	7	-	-
White, both sexes	8	9	17	15	13	20	28	39	26	8	-	-
Male	4	3	9	7	8	7	10	16	5	1	-	-
Female	4	6	8	8	5	13	18	23	21	7	-	-
Black, both sexes	2	4	8	2	2	1	4	1	2	1	-	-
Male	1	1	3	1	-	-	3	-	1	1	-	-
Female	1	3	5	1	2	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	4	3	3	4	7	6	3	10	4	1	-	-
Male	3	2	2	2	3	1	1	4	2	-	-	-
Female	1	1	1	2	4	5	2	6	2	1	-	-
White, both sexes	2	2	3	3	6	6	3	10	4	1	-	-
Male	2	2	2	2	2	1	1	4	2	-	-	-
Female	-	-	1	1	4	5	2	6	2	1	-	-
Black, both sexes	2	1	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Oregon												
All races, both sexes	1	3	5	11	3	4	7	8	5	7	-	-
Male	1	3	3	7	1	2	1	2	2	1	-	-
Female	-	-	2	4	2	2	6	6	3	6	-	-
White, both sexes	1	3	5	11	3	4	7	8	5	7	-	-
Male	1	3	3	7	1	2	1	2	2	1	-	-
Female	-	-	2	4	2	2	6	6	3	6	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	19	15	11	20	22	29	31	30	26	5	1	-
Male	13	5	8	9	8	8	11	7	10	-	-	-
Female	6	10	3	11	14	21	20	23	16	5	1	-
White, both sexes	11	12	10	17	18	26	27	29	24	5	1	-
Male	7	3	8	8	6	7	9	7	9	-	-	-
Female	4	9	2	9	12	19	18	22	15	5	1	-
Black, both sexes	8	3	1	3	4	3	4	1	2	-	-	-
Male	6	2	-	1	2	1	2	-	1	-	-	-
Female	2	1	1	2	2	2	2	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Rhode Island												
All races, both sexes	-	1	2	4	1	-	2	2	1	1	-	-
Male	-	1	1	3	1	-	-	1	-	-	-	-
Female	-	-	1	1	-	-	2	1	1	1	-	-
White, both sexes	-	1	2	4	1	-	2	2	1	1	-	-
Male	-	1	1	3	1	-	-	1	-	-	-	-
Female	-	-	1	1	-	-	2	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	10	6	8	5	1	9	12	7	9	1	-	-
Male	6	3	6	3	1	4	6	2	4	1	-	-
Female	4	3	2	2	-	5	6	5	5	-	-	-
White, both sexes	5	3	4	4	1	8	10	6	7	1	-	-
Male	2	3	3	2	1	4	4	2	4	1	-	-
Female	3	-	1	2	-	4	6	4	3	-	-	-
Black, both sexes	5	3	4	1	-	1	2	1	2	-	-	-
Male	4	-	3	1	-	-	2	-	-	-	-	-
Female	1	3	1	-	-	1	-	1	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
South Dakota												
All races, both sexes	-	1	1	3	-	1	2	2	-	2	-	-
Male	-	1	1	3	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	2	-	2	-	-
White, both sexes	-	1	-	3	-	1	2	2	-	2	-	-
Male	-	1	-	3	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	2	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	11	11	7	11	11	10	17	17	11	3	-	-
Male	4	10	4	8	3	6	8	1	2	-	-	-
Female	7	1	3	3	8	4	9	16	9	3	-	-
White, both sexes	7	8	5	7	9	8	16	16	9	2	-	-
Male	3	7	3	7	3	6	8	1	1	-	-	-
Female	4	1	2	-	6	2	8	15	8	2	-	-
Black, both sexes	4	3	2	4	2	2	1	1	2	1	-	-
Male	1	3	1	1	-	-	-	-	1	-	-	-
Female	3	-	1	3	2	2	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Texas												
All races, both sexes	23	22	22	19	26	36	38	36	20	10	1	-
Male	16	12	10	9	11	12	12	9	3	3	-	-
Female	7	10	12	10	15	24	26	27	17	7	1	-
White, both sexes	17	17	18	14	21	30	35	35	19	10	1	-
Male	12	9	7	7	8	9	11	9	3	3	-	-
Female	5	8	11	7	13	21	24	26	16	7	1	-
Black, both sexes	6	5	4	4	5	5	3	-	1	-	-	-
Male	4	3	3	2	3	3	1	-	-	-	-	-
Female	2	2	1	2	2	2	2	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	1	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	1	-	1	-	-	-	-
Utah												
All races, both sexes	-	3	-	1	7	4	1	2	-	1	-	-
Male	-	1	-	1	4	2	-	1	-	-	-	-
Female	-	2	-	-	3	2	1	1	-	1	-	-
White, both sexes	-	3	-	1	7	4	1	2	-	1	-	-
Male	-	1	-	1	4	2	-	1	-	-	-	-
Female	-	2	-	-	3	2	1	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Vermont												
All races, both sexes	-	1	-	2	2	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	2	-	-	-	-	-	-	-
White, both sexes	-	1	-	2	2	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	2	2	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	11	2	7	5	8	11	15	12	12	7	-	-
Male	9	1	3	4	3	7	6	4	3	1	-	-
Female	2	1	4	1	5	4	9	8	9	6	-	-
White, both sexes	6	1	3	4	6	9	11	11	11	6	-	-
Male	6	1	1	3	2	6	4	4	3	1	-	-
Female	-	-	2	1	4	3	7	7	8	5	-	-
Black, both sexes	5	1	4	1	2	1	4	1	1	1	-	-
Male	3	-	2	1	1	-	2	-	-	-	-	-
Female	2	1	2	-	1	1	2	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Washington												
All races, both sexes	5	13	4	9	12	12	8	11	9	3	-	-
Male	2	7	3	5	5	5	4	3	2	1	-	-
Female	3	6	1	4	7	7	4	8	7	2	-	-
White, both sexes	4	12	4	8	11	12	7	11	9	3	-	-
Male	1	7	3	4	5	5	4	3	2	1	-	-
Female	3	5	1	4	6	7	3	8	7	2	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	2	3	3	-	2	2	6	7	2	1	-	-
Male	2	3	2	-	1	1	3	2	1	-	-	-
Female	-	-	1	-	1	1	3	5	1	1	-	-
White, both sexes	2	3	3	-	2	2	6	7	2	1	-	-
Male	2	3	2	-	1	1	3	2	1	-	-	-
Female	-	-	1	-	1	1	3	5	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Other disorders of circulatory system (I80-I99)												
Wisconsin												
All races, both sexes	6	5	5	3	2	10	17	9	6	4	-	-
Male	6	1	-	2	1	1	7	2	1	1	-	-
Female	-	4	5	1	1	9	10	7	5	3	-	-
White, both sexes	4	4	4	3	2	9	16	9	6	4	-	-
Male	4	1	-	2	1	-	6	2	1	1	-	-
Female	-	3	4	1	1	9	10	7	5	3	-	-
Black, both sexes	2	1	1	-	-	1	1	-	-	-	-	-
Male	2	-	-	-	-	1	1	-	-	-	-	-
Female	-	1	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	1	-	-	2	1	1	-	-	-
Male	-	-	1	1	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	1	1	-	-	2	1	1	-	-	-
Male	-	-	1	1	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
United States												
All races, both sexes	3,435	1	1	3	-	10	25	44	71	89	155	208
Male	1,434	1	-	1	-	4	11	18	31	37	65	113
Female	2,001	-	1	2	-	6	14	26	40	52	90	95
White, both sexes	2,791	1	-	3	-	5	15	30	43	63	101	154
Male	1,152	1	-	1	-	3	5	10	21	30	45	83
Female	1,639	-	-	2	-	2	10	20	22	33	56	71
Black, both sexes	609	-	1	-	-	5	10	14	26	24	52	51
Male	266	-	-	-	-	1	6	8	8	6	19	29
Female	343	-	1	-	-	4	4	6	18	18	33	22
American Indian 1/, both sexes	12	-	-	-	-	-	-	-	-	-	2	3
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	7	-	-	-	-	-	-	-	-	-	1	2
Asian or Pacific Islander 2/, both sexes	23	-	-	-	-	-	-	-	2	2	-	-
Male	11	-	-	-	-	-	-	-	2	1	-	-
Female	12	-	-	-	-	-	-	-	-	1	-	-
Alabama												
All races, both sexes	58	-	-	-	-	-	-	-	-	1	1	2
Male	22	-	-	-	-	-	-	-	-	-	-	2
Female	36	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	43	-	-	-	-	-	-	-	-	1	-	2
Male	14	-	-	-	-	-	-	-	-	-	-	2
Female	29	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	15	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	82	-	-	-	-	-	1	-	3	1	6	5
Male	40	-	-	-	-	-	-	-	2	1	1	3
Female	42	-	-	-	-	-	1	-	1	-	5	2
White, both sexes	73	-	-	-	-	-	-	-	3	1	3	4
Male	37	-	-	-	-	-	-	-	2	1	1	2
Female	36	-	-	-	-	-	-	-	1	-	2	2
Black, both sexes	8	-	-	-	-	-	1	-	-	-	3	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	1	-	-	-	3	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Arkansas												
All races, both sexes	38	-	-	-	-	1	-	1	2	1	2	1
Male	13	-	-	-	-	-	-	-	1	-	-	1
Female	25	-	-	-	-	1	-	1	1	1	2	-
White, both sexes	32	-	-	-	-	-	-	-	1	1	2	-
Male	11	-	-	-	-	-	-	-	1	-	-	-
Female	21	-	-	-	-	-	-	-	-	1	2	-
Black, both sexes	6	-	-	-	-	1	-	1	1	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	1	-	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	423	-	-	-	-	-	4	5	8	20	22	30
Male	182	-	-	-	-	-	-	2	5	6	13	16
Female	241	-	-	-	-	-	4	3	3	14	9	14
White, both sexes	347	-	-	-	-	-	4	3	5	14	13	23
Male	145	-	-	-	-	-	-	-	3	5	7	11
Female	202	-	-	-	-	-	4	3	2	9	6	12
Black, both sexes	65	-	-	-	-	-	-	2	2	6	9	7
Male	33	-	-	-	-	-	-	2	1	1	6	5
Female	32	-	-	-	-	-	-	-	1	5	3	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	10	-	-	-	-	-	-	-	1	-	-	-
Male	4	-	-	-	-	-	-	-	1	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Colorado												
All races, both sexes	46	-	-	-	-	-	-	1	1	1	1	3
Male	19	-	-	-	-	-	-	-	-	-	1	3
Female	27	-	-	-	-	-	-	1	1	1	-	-
White, both sexes	41	-	-	-	-	-	-	1	-	-	1	3
Male	17	-	-	-	-	-	-	-	-	-	1	3
Female	24	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	1	1	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	40	-	-	-	-	-	-	1	-	-	-	2
Male	11	-	-	-	-	-	-	-	-	-	-	2
Female	29	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	37	-	-	-	-	-	-	1	-	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	27	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Delaware												
All races, both sexes	8	-	-	-	-	-	-	-	1	1	-	-
Male	6	-	-	-	-	-	-	-	1	1	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	1	-	-	-
Male	2	-	-	-	-	-	-	-	1	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	1	-	-
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	11	-	-	-	-	-	1	-	-	1	-	3
Male	6	-	-	-	-	-	1	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	10	-	-	-	-	-	1	-	-	1	-	3
Male	5	-	-	-	-	-	1	-	-	-	-	2
Female	5	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Florida												
All races, both sexes	220	-	-	-	-	-	-	-	3	5	10	10
Male	94	-	-	-	-	-	-	-	-	4	3	4
Female	126	-	-	-	-	-	-	-	3	1	7	6
White, both sexes	180	-	-	-	-	-	-	-	2	5	8	8
Male	80	-	-	-	-	-	-	-	-	4	2	3
Female	100	-	-	-	-	-	-	-	2	1	6	5
Black, both sexes	39	-	-	-	-	-	-	-	1	-	2	2
Male	14	-	-	-	-	-	-	-	-	-	1	1
Female	25	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	57	-	-	-	-	-	-	2	2	2	-	8
Male	22	-	-	-	-	-	-	2	1	-	-	4
Female	35	-	-	-	-	-	-	-	1	2	-	4
White, both sexes	34	-	-	-	-	-	-	1	1	2	-	4
Male	12	-	-	-	-	-	-	1	1	-	-	1
Female	22	-	-	-	-	-	-	-	-	2	-	3
Black, both sexes	23	-	-	-	-	-	-	1	1	-	-	4
Male	10	-	-	-	-	-	-	1	-	-	-	3
Female	13	-	-	-	-	-	-	-	1	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Hawaii												
All races, both sexes	7	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	15	-	-	-	-	-	-	-	1	-	2	-
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	2	-
White, both sexes	15	-	-	-	-	-	-	-	1	-	2	-
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Illinois												
All races, both sexes	150	1	-	-	-	-	2	4	6	5	6	8
Male	63	1	-	-	-	-	1	2	3	2	4	5
Female	87	-	-	-	-	-	1	2	3	3	2	3
White, both sexes	115	1	-	-	-	-	1	3	2	3	2	4
Male	47	1	-	-	-	-	-	1	1	1	2	2
Female	68	-	-	-	-	-	1	2	1	2	-	2
Black, both sexes	35	-	-	-	-	-	1	1	4	2	4	4
Male	16	-	-	-	-	-	1	1	2	1	2	3
Female	19	-	-	-	-	-	-	-	2	1	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	39	-	-	-	-	1	-	-	-	1	1	-
Male	14	-	-	-	-	-	-	-	-	1	1	-
Female	25	-	-	-	-	1	-	-	-	-	-	-
White, both sexes	36	-	-	-	-	-	-	-	-	1	1	-
Male	14	-	-	-	-	-	-	-	-	1	1	-
Female	22	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Iowa												
All races, both sexes	31	-	-	-	-	-	1	-	-	-	2	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	1	-	-	-	1	-
White, both sexes	31	-	-	-	-	-	1	-	-	-	2	-
Male	12	-	-	-	-	-	-	-	-	-	1	-
Female	19	-	-	-	-	-	1	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	24	-	-	-	-	-	-	1	-	-	3	2
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	1	-	-	2	2
White, both sexes	24	-	-	-	-	-	-	1	-	-	3	2
Male	8	-	-	-	-	-	-	-	-	-	1	-
Female	16	-	-	-	-	-	-	1	-	-	2	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Kentucky												
All races, both sexes	43	-	-	-	-	-	1	-	-	-	-	4
Male	20	-	-	-	-	-	-	-	-	-	-	2
Female	23	-	-	-	-	-	1	-	-	-	-	2
White, both sexes	38	-	-	-	-	-	1	-	-	-	-	4
Male	18	-	-	-	-	-	-	-	-	-	-	2
Female	20	-	-	-	-	-	1	-	-	-	-	2
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	43	-	-	-	-	-	-	1	-	-	1	3
Male	17	-	-	-	-	-	-	1	-	-	1	2
Female	26	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	26	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	1	2
Female	13	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	17	-	-	-	-	-	-	1	-	-	-	1
Male	4	-	-	-	-	-	-	1	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Maine												
All races, both sexes	16	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	1	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	13	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	79	-	-	-	-	2	-	2	3	4	5	11
Male	22	-	-	-	-	1	-	1	1	-	-	5
Female	57	-	-	-	-	1	-	1	2	4	5	6
White, both sexes	41	-	-	-	-	1	-	-	1	1	3	4
Male	13	-	-	-	-	1	-	-	1	-	-	2
Female	28	-	-	-	-	-	-	-	-	1	3	2
Black, both sexes	38	-	-	-	-	1	-	2	2	3	2	7
Male	9	-	-	-	-	-	-	1	-	-	-	3
Female	29	-	-	-	-	1	-	1	2	3	2	4
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Massachusetts												
All races, both sexes	51	-	-	-	-	-	-	-	-	-	-	4
Male	25	-	-	-	-	-	-	-	-	-	-	4
Female	26	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	50	-	-	-	-	-	-	-	-	-	-	4
Male	24	-	-	-	-	-	-	-	-	-	-	4
Female	26	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	165	-	-	-	-	-	2	4	5	6	12	16
Male	78	-	-	-	-	-	-	3	1	4	4	8
Female	87	-	-	-	-	-	2	1	4	2	8	8
White, both sexes	129	-	-	-	-	-	1	3	3	5	5	12
Male	57	-	-	-	-	-	-	2	-	3	2	6
Female	72	-	-	-	-	-	1	1	3	2	3	6
Black, both sexes	34	-	-	-	-	-	1	1	2	-	7	4
Male	19	-	-	-	-	-	-	1	1	-	2	2
Female	15	-	-	-	-	-	1	-	1	-	5	2
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Minnesota												
All races, both sexes	31	-	-	-	-	-	-	-	-	-	1	3
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	21	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	30	-	-	-	-	-	-	-	-	-	1	2
Male	10	-	-	-	-	-	-	-	-	-	1	-
Female	20	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	33	-	-	-	-	-	-	-	-	-	5	3
Male	19	-	-	-	-	-	-	-	-	-	3	2
Female	14	-	-	-	-	-	-	-	-	-	2	1
White, both sexes	25	-	-	-	-	-	-	-	-	-	4	2
Male	15	-	-	-	-	-	-	-	-	-	2	2
Female	10	-	-	-	-	-	-	-	-	-	2	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Missouri												
All races, both sexes	75	-	-	1	-	1	-	3	4	1	5	-
Male	27	-	-	-	-	-	-	-	1	1	1	-
Female	48	-	-	1	-	1	-	3	3	-	4	-
White, both sexes	65	-	-	1	-	1	-	2	3	1	3	-
Male	22	-	-	-	-	-	-	-	1	1	1	-
Female	43	-	-	1	-	1	-	2	2	-	2	-
Black, both sexes	10	-	-	-	-	-	-	1	1	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	1	1	-	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	1	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Nebraska												
All races, both sexes	18	-	-	-	-	-	-	-	1	-	-	2
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	2
White, both sexes	18	-	-	-	-	-	-	-	1	-	-	2
Male	9	-	-	-	-	-	-	-	1	-	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	19	-	-	-	-	-	-	1	1	2	1	-
Male	9	-	-	-	-	-	-	1	-	2	1	-
Female	10	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	18	-	-	-	-	-	-	1	-	2	1	-
Male	9	-	-	-	-	-	-	1	-	2	1	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
New Hampshire												
All races, both sexes	23	-	-	-	-	-	-	-	1	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	13	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	23	-	-	-	-	-	-	-	1	-	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	2
Female	13	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	130	-	-	-	-	-	1	-	2	3	7	6
Male	58	-	-	-	-	-	1	-	1	2	4	3
Female	72	-	-	-	-	-	-	-	1	1	3	3
White, both sexes	100	-	-	-	-	-	-	-	-	2	5	5
Male	42	-	-	-	-	-	-	-	-	2	3	3
Female	58	-	-	-	-	-	-	-	-	-	2	2
Black, both sexes	27	-	-	-	-	-	1	-	2	1	2	1
Male	15	-	-	-	-	-	1	-	1	-	1	-
Female	12	-	-	-	-	-	-	-	1	1	1	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
New Mexico												
All races, both sexes	26	-	-	-	-	-	-	1	1	1	1	1
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	16	-	-	-	-	-	-	1	1	1	1	1
White, both sexes	24	-	-	-	-	-	-	1	1	1	1	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	14	-	-	-	-	-	-	1	1	1	1	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	248	-	-	1	-	1	2	5	6	9	13	12
Male	114	-	-	-	-	-	1	3	4	5	5	7
Female	134	-	-	1	-	1	1	2	2	4	8	5
White, both sexes	183	-	-	1	-	-	-	2	5	6	9	8
Male	86	-	-	-	-	-	-	2	4	3	4	6
Female	97	-	-	1	-	-	-	-	1	3	5	2
Black, both sexes	64	-	-	-	-	1	2	3	1	2	4	4
Male	28	-	-	-	-	-	1	1	-	2	1	1
Female	36	-	-	-	-	1	1	2	1	-	3	3
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
North Carolina												
All races, both sexes	107	-	-	-	-	-	2	-	2	4	4	4
Male	47	-	-	-	-	-	1	-	-	1	2	2
Female	60	-	-	-	-	-	1	-	2	3	2	2
White, both sexes	72	-	-	-	-	-	1	-	1	4	1	3
Male	29	-	-	-	-	-	1	-	-	1	1	2
Female	43	-	-	-	-	-	-	-	1	3	-	1
Black, both sexes	34	-	-	-	-	-	1	-	1	-	3	1
Male	17	-	-	-	-	-	-	-	-	-	1	-
Female	17	-	-	-	-	-	1	-	1	-	2	1
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	9	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	8	-	-	-	-	-	1	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Ohio												
All races, both sexes	193	-	-	1	-	1	2	3	2	6	5	20
Male	82	-	-	1	-	1	2	1	2	1	3	13
Female	111	-	-	-	-	-	-	2	-	5	2	7
White, both sexes	164	-	-	1	-	1	1	3	2	4	3	15
Male	70	-	-	1	-	1	1	1	2	1	3	8
Female	94	-	-	-	-	-	-	2	-	3	-	7
Black, both sexes	29	-	-	-	-	-	1	-	-	2	2	5
Male	12	-	-	-	-	-	1	-	-	-	-	5
Female	17	-	-	-	-	-	-	-	-	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	41	-	-	-	-	1	-	-	1	-	4	3
Male	19	-	-	-	-	1	-	-	-	-	2	2
Female	22	-	-	-	-	-	-	-	1	-	2	1
White, both sexes	37	-	-	-	-	1	-	-	1	-	2	3
Male	17	-	-	-	-	1	-	-	-	-	-	2
Female	20	-	-	-	-	-	-	-	1	-	2	1
Black, both sexes	4	-	-	-	-	-	-	-	-	-	2	-
Male	2	-	-	-	-	-	-	-	-	-	2	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Oregon												
All races, both sexes	39	-	-	-	-	-	-	1	-	-	3	2
Male	16	-	-	-	-	-	-	1	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	3	1
White, both sexes	39	-	-	-	-	-	-	1	-	-	3	2
Male	16	-	-	-	-	-	-	1	-	-	-	1
Female	23	-	-	-	-	-	-	-	-	-	3	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	180	-	-	-	-	-	1	2	3	3	9	10
Male	69	-	-	-	-	-	1	-	-	3	3	4
Female	111	-	-	-	-	-	-	2	3	-	6	6
White, both sexes	151	-	-	-	-	-	1	2	2	2	6	9
Male	56	-	-	-	-	-	1	-	-	2	2	4
Female	95	-	-	-	-	-	-	2	2	-	4	5
Black, both sexes	29	-	-	-	-	-	-	-	1	1	3	1
Male	13	-	-	-	-	-	-	-	-	1	1	-
Female	16	-	-	-	-	-	-	-	1	-	2	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Rhode Island												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	6	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	54	-	-	-	-	-	1	1	-	-	4	3
Male	26	-	-	-	-	-	1	1	-	-	3	1
Female	28	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	39	-	-	-	-	-	1	1	-	-	2	2
Male	20	-	-	-	-	-	1	1	-	-	2	-
Female	19	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	15	-	-	-	-	-	-	-	-	-	2	1
Male	6	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
South Dakota												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	100	-	-	-	-	-	2	1	2	1	4	5
Male	39	-	-	-	-	-	2	-	-	1	2	2
Female	61	-	-	-	-	-	-	1	2	-	2	3
White, both sexes	78	-	-	-	-	-	1	1	1	1	3	3
Male	33	-	-	-	-	-	1	-	-	1	2	1
Female	45	-	-	-	-	-	-	1	1	-	1	2
Black, both sexes	22	-	-	-	-	-	1	-	1	-	1	2
Male	6	-	-	-	-	-	1	-	-	-	-	1
Female	16	-	-	-	-	-	-	-	1	-	1	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Texas												
All races, both sexes	193	-	-	-	-	1	1	-	4	4	3	5
Male	67	-	-	-	-	-	-	-	3	1	2	3
Female	126	-	-	-	-	1	1	-	1	3	1	2
White, both sexes	161	-	-	-	-	1	1	-	1	2	3	4
Male	52	-	-	-	-	-	-	-	1	1	2	3
Female	109	-	-	-	-	1	1	-	-	1	1	1
Black, both sexes	30	-	-	-	-	-	-	-	3	2	-	1
Male	15	-	-	-	-	-	-	-	2	-	-	-
Female	15	-	-	-	-	-	-	-	1	2	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Vermont												
All races, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	1	-
White, both sexes	6	-	-	-	-	-	-	-	-	1	1	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	76	-	1	-	-	1	-	3	3	2	4	3
Male	26	-	-	-	-	1	-	-	-	-	-	1
Female	50	-	1	-	-	-	-	3	3	2	4	2
White, both sexes	56	-	-	-	-	-	-	2	3	-	2	3
Male	19	-	-	-	-	-	-	-	-	-	-	1
Female	37	-	-	-	-	-	-	2	3	-	2	2
Black, both sexes	20	-	1	-	-	1	-	1	-	2	2	-
Male	7	-	-	-	-	1	-	-	-	-	-	-
Female	13	-	1	-	-	-	-	1	-	2	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Washington												
All races, both sexes	71	-	-	-	-	-	-	-	1	1	2	4
Male	34	-	-	-	-	-	-	-	1	1	1	3
Female	37	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	70	-	-	-	-	-	-	-	-	1	2	4
Male	33	-	-	-	-	-	-	-	-	1	1	3
Female	37	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	22	-	-	-	-	-	-	-	2	-	1	2
Male	9	-	-	-	-	-	-	-	2	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	2
White, both sexes	21	-	-	-	-	-	-	-	1	-	1	2
Male	8	-	-	-	-	-	-	-	1	-	-	-
Female	13	-	-	-	-	-	-	-	-	-	1	2
Black, both sexes	1	-	-	-	-	-	-	-	1	-	-	-
Male	1	-	-	-	-	-	-	-	1	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Wisconsin												
All races, both sexes	53	-	-	-	-	-	-	1	-	1	1	4
Male	18	-	-	-	-	-	-	-	-	-	1	3
Female	35	-	-	-	-	-	-	1	-	1	-	1
White, both sexes	48	-	-	-	-	-	-	1	-	1	1	3
Male	14	-	-	-	-	-	-	-	-	-	1	2
Female	34	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes	5	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
United States												
All races, both sexes	200	234	212	235	257	389	447	450	289	104	11	-
Male	112	126	116	127	129	156	170	133	68	16	-	-
Female	88	108	96	108	128	233	277	317	221	88	11	-
White, both sexes	141	180	164	194	207	328	383	410	265	95	9	-
Male	85	100	90	105	101	131	145	120	62	14	-	-
Female	56	80	74	89	106	197	238	290	203	81	9	-
Black, both sexes	57	51	46	39	49	55	58	37	24	8	2	-
Male	26	23	25	22	27	23	24	11	6	2	-	-
Female	31	28	21	17	22	32	34	26	18	6	2	-
American Indian 1/, both sexes	1	-	-	1	-	3	2	-	-	-	-	-
Male	1	-	-	-	-	2	-	-	-	-	-	-
Female	-	-	-	1	-	1	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	3	2	1	1	3	4	3	-	1	-	-
Male	-	3	1	-	1	-	1	2	-	-	-	-
Female	1	-	1	1	-	3	3	1	-	1	-	-
Alabama												
All races, both sexes	2	5	6	8	4	8	10	5	5	1	-	-
Male	1	3	3	5	-	5	1	2	-	-	-	-
Female	1	2	3	3	4	3	9	3	5	1	-	-
White, both sexes	2	2	2	6	4	5	9	5	4	1	-	-
Male	1	1	1	3	-	3	1	2	-	-	-	-
Female	1	1	1	3	4	2	8	3	4	1	-	-
Black, both sexes	-	3	4	2	-	3	1	-	1	-	-	-
Male	-	2	2	2	-	2	-	-	-	-	-	-
Female	-	1	2	-	-	1	1	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	9	3	8	8	5	8	9	7	6	3	-	-
Male	7	1	4	6	4	3	3	3	1	1	-	-
Female	2	2	4	2	1	5	6	4	5	2	-	-
White, both sexes	8	2	8	7	4	8	9	7	6	3	-	-
Male	6	1	4	5	4	3	3	3	1	1	-	-
Female	2	1	4	2	-	5	6	4	5	2	-	-
Black, both sexes	1	1	-	1	1	-	-	-	-	-	-	-
Male	1	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Arkansas												
All races, both sexes	4	-	2	1	3	5	5	3	6	1	-	-
Male	2	-	2	1	-	1	2	2	-	1	-	-
Female	2	-	-	-	3	4	3	1	6	-	-	-
White, both sexes	4	-	2	1	3	5	5	2	5	1	-	-
Male	2	-	2	1	-	1	2	1	-	1	-	-
Female	2	-	-	-	3	4	3	1	5	-	-	-
Black, both sexes	-	-	-	-	-	-	-	1	1	-	-	-
Male	-	-	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	28	34	23	26	28	35	55	54	38	11	2	-
Male	15	19	12	13	16	16	19	22	7	1	-	-
Female	13	15	11	13	12	19	36	32	31	10	2	-
White, both sexes	18	29	17	23	25	31	45	47	38	10	2	-
Male	10	17	8	11	15	15	16	19	7	1	-	-
Female	8	12	9	12	10	16	29	28	31	9	2	-
Black, both sexes	10	5	4	3	2	3	6	6	-	-	-	-
Male	5	2	3	2	-	1	3	2	-	-	-	-
Female	5	3	1	1	2	2	3	4	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	2	-	1	1	3	1	-	1	-	-
Male	-	-	1	-	1	-	-	1	-	-	-	-
Female	-	-	1	-	-	1	3	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Colorado												
All races, both sexes	2	-	8	4	4	6	4	4	3	3	1	-
Male	1	-	5	1	4	1	1	1	1	-	-	-
Female	1	-	3	3	-	5	3	3	2	3	1	-
White, both sexes	2	-	8	3	3	6	4	4	3	2	1	-
Male	1	-	5	-	3	1	1	1	1	-	-	-
Female	1	-	3	3	-	5	3	3	2	2	1	-
Black, both sexes	-	-	-	1	1	-	-	-	-	1	-	-
Male	-	-	-	1	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	-	3	2	2	-	4	8	9	7	2	-	-
Male	-	1	1	-	-	1	2	4	-	-	-	-
Female	-	2	1	2	-	3	6	5	7	2	-	-
White, both sexes	-	3	2	2	-	4	5	9	7	2	-	-
Male	-	1	1	-	-	1	1	4	-	-	-	-
Female	-	2	1	2	-	3	4	5	7	2	-	-
Black, both sexes	-	-	-	-	-	-	3	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Delaware												
All races, both sexes	-	1	1	-	-	1	1	1	1	-	-	-
Male	-	1	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	1	-	-	1	-	1	-	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	1	1	-	-	1	1	2	-	-	-	-	-
Male	-	1	-	-	1	-	1	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	1	-	-	1	1	1	-	-	-	-	-
Male	-	1	-	-	1	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Florida												
All races, both sexes	11	17	17	17	21	28	29	31	17	3	1	-
Male	4	12	9	11	9	10	13	9	5	1	-	-
Female	7	5	8	6	12	18	16	22	12	2	1	-
White, both sexes	7	13	14	12	14	25	28	26	15	2	1	-
Male	3	10	9	7	6	10	13	7	5	1	-	-
Female	4	3	5	5	8	15	15	19	10	1	1	-
Black, both sexes	3	4	3	5	7	3	1	5	2	1	-	-
Male	1	2	-	4	3	-	-	2	-	-	-	-
Female	2	2	3	1	4	3	1	3	2	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	4	2	3	3	5	8	7	5	3	3	-	-
Male	2	-	2	3	3	2	3	-	-	-	-	-
Female	2	2	1	-	2	6	4	5	3	3	-	-
White, both sexes	2	1	2	1	4	5	4	3	1	3	-	-
Male	1	-	1	1	2	2	2	-	-	-	-	-
Female	1	1	1	-	2	3	2	3	1	3	-	-
Black, both sexes	2	1	1	2	1	3	3	2	2	-	-	-
Male	1	-	1	2	1	-	1	-	-	-	-	-
Female	1	1	-	-	-	3	2	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Hawaii												
All races, both sexes	-	2	-	-	1	1	1	1	-	-	-	-
Male	-	2	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
White, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	1	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	2	-	-	-	-	1	1	-	-	-	-
Male	-	2	-	-	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	1	1	1	-	-	1	4	2	2	-	-	-
Male	1	1	1	-	-	1	2	1	2	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
White, both sexes	1	1	1	-	-	1	4	2	2	-	-	-
Male	1	1	1	-	-	1	2	1	2	-	-	-
Female	-	-	-	-	-	-	2	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Illinois												
All races, both sexes	8	8	7	9	17	20	7	26	10	6	-	-
Male	4	5	4	6	8	7	3	5	2	1	-	-
Female	4	3	3	3	9	13	4	21	8	5	-	-
White, both sexes	4	8	3	7	14	18	7	23	10	5	-	-
Male	4	5	2	4	7	7	3	4	2	1	-	-
Female	-	3	1	3	7	11	4	19	8	4	-	-
Black, both sexes	4	-	4	2	3	2	-	3	-	1	-	-
Male	-	-	2	2	1	-	-	1	-	-	-	-
Female	4	-	2	-	2	2	-	2	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	1	3	2	6	1	5	7	5	3	3	-	-
Male	1	3	-	2	-	2	2	1	1	-	-	-
Female	-	-	2	4	1	3	5	4	2	3	-	-
White, both sexes	1	3	2	6	1	5	6	4	3	3	-	-
Male	1	3	-	2	-	2	2	1	1	-	-	-
Female	-	-	2	4	1	3	4	3	2	3	-	-
Black, both sexes	-	-	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Iowa												
All races, both sexes	1	3	3	4	1	3	1	4	7	1	-	-
Male	1	3	2	3	-	-	-	-	2	-	-	-
Female	-	-	1	1	1	3	1	4	5	1	-	-
White, both sexes	1	3	3	4	1	3	1	4	7	1	-	-
Male	1	3	2	3	-	-	-	-	2	-	-	-
Female	-	-	1	1	1	3	1	4	5	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	1	1	-	1	1	3	2	4	3	1	1	-
Male	-	-	-	-	1	2	1	1	2	-	-	-
Female	1	1	-	1	-	1	1	3	1	1	1	-
White, both sexes	1	1	-	1	1	3	2	4	3	1	1	-
Male	-	-	-	-	1	2	1	1	2	-	-	-
Female	1	1	-	1	-	1	1	3	1	1	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Kentucky												
All races, both sexes	4	2	3	2	3	9	7	3	5	-	-	-
Male	2	-	2	-	1	7	2	2	2	-	-	-
Female	2	2	1	2	2	2	5	1	3	-	-	-
White, both sexes	2	2	3	1	3	7	7	3	5	-	-	-
Male	1	-	2	-	1	6	2	2	2	-	-	-
Female	1	2	1	1	2	1	5	1	3	-	-	-
Black, both sexes	1	-	-	1	-	2	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	1	-	-	1	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	6	4	2	3	3	7	4	7	2	-	-	-
Male	1	1	1	1	2	1	3	3	-	-	-	-
Female	5	3	1	2	1	6	1	4	2	-	-	-
White, both sexes	3	1	-	1	2	4	4	6	2	-	-	-
Male	1	-	-	1	1	1	3	3	-	-	-	-
Female	2	1	-	-	1	3	1	3	2	-	-	-
Black, both sexes	3	3	2	2	1	3	-	1	-	-	-	-
Male	-	1	1	-	1	-	-	-	-	-	-	-
Female	3	2	1	2	-	3	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Maine												
All races, both sexes	-	1	-	2	3	2	-	4	2	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	1	-	2	2	1	-	3	2	1	-	-
White, both sexes	-	1	-	2	3	2	-	4	2	1	-	-
Male	-	-	-	-	1	1	-	1	-	-	-	-
Female	-	1	-	2	2	1	-	3	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	5	3	5	3	3	2	9	12	5	3	2	-
Male	3	1	3	2	-	-	1	3	1	-	-	-
Female	2	2	2	1	3	2	8	9	4	3	2	-
White, both sexes	1	2	3	1	-	-	6	11	5	2	-	-
Male	1	-	2	1	-	-	1	3	1	-	-	-
Female	-	2	1	-	-	-	5	8	4	2	-	-
Black, both sexes	4	1	2	2	3	2	3	1	-	1	2	-
Male	2	1	1	1	-	-	-	-	-	-	-	-
Female	2	-	1	1	3	2	3	1	-	1	2	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Massachusetts												
All races, both sexes	1	1	-	4	6	5	9	10	8	3	-	-
Male	1	-	-	4	5	3	3	3	2	-	-	-
Female	-	1	-	-	1	2	6	7	6	3	-	-
White, both sexes	1	1	-	3	6	5	9	10	8	3	-	-
Male	1	-	-	3	5	3	3	3	2	-	-	-
Female	-	1	-	-	1	2	6	7	6	3	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	9	17	11	9	13	19	14	22	4	2	-	-
Male	5	12	6	5	9	8	4	7	1	1	-	-
Female	4	5	5	4	4	11	10	15	3	1	-	-
White, both sexes	6	14	11	9	10	16	11	19	3	1	-	-
Male	4	9	6	5	6	6	2	6	-	-	-	-
Female	2	5	5	4	4	10	9	13	3	1	-	-
Black, both sexes	3	3	-	-	3	2	3	3	1	1	-	-
Male	1	3	-	-	3	1	2	1	1	1	-	-
Female	2	-	-	-	-	1	1	2	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Minnesota												
All races, both sexes	1	1	5	3	2	2	6	5	2	-	-	-
Male	-	-	3	-	1	1	2	2	-	-	-	-
Female	1	1	2	3	1	1	4	3	2	-	-	-
White, both sexes	1	1	5	3	2	2	6	5	2	-	-	-
Male	-	-	3	-	1	1	2	2	-	-	-	-
Female	1	1	2	3	1	1	4	3	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	-	3	2	5	-	5	5	5	-	-	-	-
Male	-	1	2	4	-	2	3	2	-	-	-	-
Female	-	2	-	1	-	3	2	3	-	-	-	-
White, both sexes	-	3	1	5	-	2	5	3	-	-	-	-
Male	-	1	1	4	-	1	3	1	-	-	-	-
Female	-	2	-	1	-	1	2	2	-	-	-	-
Black, both sexes	-	-	1	-	-	2	-	2	-	-	-	-
Male	-	-	1	-	-	1	-	1	-	-	-	-
Female	-	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Missouri												
All races, both sexes	5	3	3	7	8	9	7	6	9	3	-	-
Male	2	2	1	3	6	3	5	-	2	-	-	-
Female	3	1	2	4	2	6	2	6	7	3	-	-
White, both sexes	5	2	2	7	8	7	6	6	8	3	-	-
Male	2	1	-	3	6	2	4	-	1	-	-	-
Female	3	1	2	4	2	5	2	6	7	3	-	-
Black, both sexes	-	1	1	-	-	2	1	-	1	-	-	-
Male	-	1	1	-	-	1	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	-	-	-	-	-	3	2	1	2	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	1	1	1	2	-	-	-
White, both sexes	-	-	-	-	-	2	2	1	2	-	-	-
Male	-	-	-	-	-	2	1	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Nebraska												
All races, both sexes	2	-	2	1	1	2	2	2	2	1	-	-
Male	2	-	2	-	-	2	1	-	1	-	-	-
Female	-	-	-	1	1	-	1	2	1	1	-	-
White, both sexes	2	-	2	1	1	2	2	2	2	1	-	-
Male	2	-	2	-	-	2	1	-	1	-	-	-
Female	-	-	-	1	1	-	1	2	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	-	1	3	2	-	2	4	1	-	-	-
Male	-	-	1	1	-	-	-	2	1	-	-	-
Female	1	-	-	2	2	-	2	2	-	-	-	-
White, both sexes	1	-	1	3	2	-	2	4	1	-	-	-
Male	-	-	1	1	-	-	-	2	1	-	-	-
Female	1	-	-	2	2	-	2	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
New Hampshire												
All races, both sexes	2	1	1	-	3	4	-	5	4	-	-	-
Male	2	-	1	-	2	1	-	2	-	-	-	-
Female	-	1	-	-	1	3	-	3	4	-	-	-
White, both sexes	2	1	1	-	3	4	-	5	4	-	-	-
Male	2	-	1	-	2	1	-	2	-	-	-	-
Female	-	1	-	-	1	3	-	3	4	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	10	11	8	3	5	21	27	10	11	4	1	-
Male	5	8	5	1	3	8	8	3	4	2	-	-
Female	5	3	3	2	2	13	19	7	7	2	1	-
White, both sexes	9	8	7	2	4	17	19	8	10	3	1	-
Male	4	6	4	1	2	5	6	2	3	1	-	-
Female	5	2	3	1	2	12	13	6	7	2	1	-
Black, both sexes	1	2	1	1	1	3	7	2	1	1	-	-
Male	1	1	1	-	1	3	2	1	1	1	-	-
Female	-	1	-	1	-	-	5	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
New Mexico												
All races, both sexes	1	1	-	3	1	4	5	1	5	-	-	-
Male	1	1	-	2	-	1	2	-	3	-	-	-
Female	-	-	-	1	1	3	3	1	2	-	-	-
White, both sexes	1	1	-	3	1	4	4	1	5	-	-	-
Male	1	1	-	2	-	1	2	-	3	-	-	-
Female	-	-	-	1	1	3	2	1	2	-	-	-
Black, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	16	26	11	11	27	32	34	26	10	6	-	-
Male	10	12	7	7	14	15	14	8	2	-	-	-
Female	6	14	4	4	13	17	20	18	8	6	-	-
White, both sexes	11	17	8	8	18	25	28	23	8	6	-	-
Male	6	9	6	6	7	11	13	7	2	-	-	-
Female	5	8	2	2	11	14	15	16	6	6	-	-
Black, both sexes	5	9	3	3	9	7	6	3	2	-	-	-
Male	4	3	1	1	7	4	1	1	-	-	-	-
Female	1	6	2	2	2	3	5	2	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
North Carolina												
All races, both sexes	7	7	5	5	6	13	15	19	12	1	1	-
Male	5	4	4	3	4	7	6	5	3	-	-	-
Female	2	3	1	2	2	6	9	14	9	1	1	-
White, both sexes	5	3	3	4	4	7	10	17	7	1	1	-
Male	4	1	2	2	2	3	3	5	2	-	-	-
Female	1	2	1	2	2	4	7	12	5	1	1	-
Black, both sexes	2	4	2	1	2	5	5	2	5	-	-	-
Male	1	3	2	1	2	3	3	-	1	-	-	-
Female	1	1	-	-	-	2	2	2	4	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	1	-	1	1	1	-	1	3	-	-	-	-
Male	1	-	1	-	1	-	1	1	-	-	-	-
Female	-	-	-	1	-	-	-	2	-	-	-	-
White, both sexes	1	-	1	-	1	-	1	3	-	-	-	-
Male	1	-	1	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	2	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Ohio												
All races, both sexes	7	8	20	14	9	15	26	29	18	7	-	-
Male	3	2	10	6	6	4	12	11	3	1	-	-
Female	4	6	10	8	3	11	14	18	15	6	-	-
White, both sexes	6	6	14	12	8	14	22	28	17	7	-	-
Male	3	2	8	5	6	4	9	11	3	1	-	-
Female	3	4	6	7	2	10	13	17	14	6	-	-
Black, both sexes	1	2	6	2	1	1	4	1	1	-	-	-
Male	-	-	2	1	-	-	3	-	-	-	-	-
Female	1	2	4	1	1	1	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	3	1	2	4	5	2	10	2	1	-	-
Male	1	2	1	1	2	1	1	4	1	-	-	-
Female	1	1	-	1	2	4	1	6	1	1	-	-
White, both sexes	1	2	1	2	4	5	2	10	2	1	-	-
Male	1	2	1	1	2	1	1	4	1	-	-	-
Female	-	-	-	1	2	4	1	6	1	1	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Oregon												
All races, both sexes	1	-	2	8	2	3	3	7	3	4	-	-
Male	1	-	1	6	-	1	1	2	1	1	-	-
Female	-	-	1	2	2	2	2	5	2	3	-	-
White, both sexes	1	-	2	8	2	3	3	7	3	4	-	-
Male	1	-	1	6	-	1	1	2	1	1	-	-
Female	-	-	1	2	2	2	2	5	2	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	10	12	5	15	17	23	25	22	18	4	1	-
Male	7	4	3	7	7	7	10	6	7	-	-	-
Female	3	8	2	8	10	16	15	16	11	4	1	-
White, both sexes	6	10	4	12	13	20	22	21	16	4	1	-
Male	4	3	3	6	5	6	8	6	6	-	-	-
Female	2	7	1	6	8	14	14	15	10	4	1	-
Black, both sexes	4	2	1	3	4	3	3	1	2	-	-	-
Male	3	1	-	1	2	1	2	-	1	-	-	-
Female	1	1	1	2	2	2	1	1	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Rhode Island												
All races, both sexes	-	-	2	4	1	-	1	-	1	1	-	-
Male	-	-	1	3	1	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	1	-	1	1	-	-
White, both sexes	-	-	2	4	1	-	1	-	1	1	-	-
Male	-	-	1	3	1	-	-	-	-	-	-	-
Female	-	-	1	1	-	-	1	-	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	4	4	5	2	1	7	8	5	8	1	-	-
Male	3	1	3	2	1	2	3	1	3	1	-	-
Female	1	3	2	-	-	5	5	4	5	-	-	-
White, both sexes	2	1	2	2	1	6	7	5	6	1	-	-
Male	2	1	1	2	1	2	2	1	3	1	-	-
Female	-	-	1	-	-	4	5	4	3	-	-	-
Black, both sexes	2	3	3	-	-	1	1	-	2	-	-	-
Male	1	-	2	-	-	-	1	-	-	-	-	-
Female	1	3	1	-	-	1	-	-	2	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
South Dakota												
All races, both sexes	-	1	-	2	-	1	2	1	-	2	-	-
Male	-	1	-	2	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	2	-	-
White, both sexes	-	1	-	2	-	1	2	1	-	2	-	-
Male	-	1	-	2	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	2	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	8	7	5	7	11	7	16	11	10	3	-	-
Male	2	7	2	4	3	4	8	-	2	-	-	-
Female	6	-	3	3	8	3	8	11	8	3	-	-
White, both sexes	6	5	3	4	9	6	15	10	8	2	-	-
Male	2	5	1	4	3	4	8	-	1	-	-	-
Female	4	-	2	-	6	2	7	10	7	2	-	-
Black, both sexes	2	2	2	3	2	1	1	1	2	1	-	-
Male	-	2	1	-	-	-	-	-	1	-	-	-
Female	2	-	1	3	2	1	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Texas												
All races, both sexes	11	13	14	13	15	25	32	27	16	8	1	-
Male	6	4	4	5	7	12	9	6	2	3	-	-
Female	5	9	10	8	8	13	23	21	14	5	1	-
White, both sexes	7	11	11	9	10	21	29	26	16	8	1	-
Male	3	4	2	4	4	9	8	6	2	3	-	-
Female	4	7	9	5	6	12	21	20	14	5	1	-
Black, both sexes	4	2	3	3	5	4	3	-	-	-	-	-
Male	3	-	2	1	3	3	1	-	-	-	-	-
Female	1	2	1	2	2	1	2	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	1	-	-	-	-
Utah												
All races, both sexes	-	3	-	-	3	1	1	1	-	-	-	-
Male	-	1	-	-	2	-	-	1	-	-	-	-
Female	-	2	-	-	1	1	1	-	-	-	-	-
White, both sexes	-	3	-	-	3	1	1	1	-	-	-	-
Male	-	1	-	-	2	-	-	1	-	-	-	-
Female	-	2	-	-	1	1	1	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Vermont												
All races, both sexes	-	1	-	1	2	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	2	-	-	-	-	-	-	-
White, both sexes	-	1	-	1	2	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	2	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	6	2	3	4	6	7	8	10	9	4	-	-
Male	5	1	1	3	1	4	5	2	2	-	-	-
Female	1	1	2	1	5	3	3	8	7	4	-	-
White, both sexes	4	1	2	3	4	7	5	9	8	3	-	-
Male	4	1	-	2	-	4	3	2	2	-	-	-
Female	-	-	2	1	4	3	2	7	6	3	-	-
Black, both sexes	2	1	1	1	2	-	3	1	1	1	-	-
Male	1	-	1	1	1	-	2	-	-	-	-	-
Female	1	1	-	-	1	-	1	1	1	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Washington												
All races, both sexes	2	10	4	7	5	12	6	9	5	3	-	-
Male	-	6	3	3	3	5	4	2	1	1	-	-
Female	2	4	1	4	2	7	2	7	4	2	-	-
White, both sexes	2	10	4	7	5	12	6	9	5	3	-	-
Male	-	6	3	3	3	5	4	2	1	1	-	-
Female	2	4	1	4	2	7	2	7	4	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	1	2	-	2	1	5	4	-	1	-	-
Male	1	1	1	-	1	-	2	1	-	-	-	-
Female	-	-	1	-	1	1	3	3	-	1	-	-
White, both sexes	1	1	2	-	2	1	5	4	-	1	-	-
Male	1	1	1	-	1	-	2	1	-	-	-	-
Female	-	-	1	-	1	1	3	3	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)												
Wisconsin												
All races, both sexes	4	4	5	2	-	6	11	7	4	3	-	-
Male	4	1	-	1	-	1	5	-	1	1	-	-
Female	-	3	5	1	-	5	6	7	3	2	-	-
White, both sexes	3	4	4	2	-	5	10	7	4	3	-	-
Male	3	1	-	1	-	-	4	-	1	1	-	-
Female	-	3	4	1	-	5	6	7	3	2	-	-
Black, both sexes	1	-	1	-	-	1	1	-	-	-	-	-
Male	1	-	-	-	-	1	1	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
United States												
All races, both sexes	1,378	25	5	1	1	1	-	6	11	35	53	90
Male	722	12	3	1	1	-	-	5	8	18	44	67
Female	656	13	2	-	-	1	-	1	3	17	9	23
White, both sexes	1,145	20	3	1	1	1	-	5	9	25	34	66
Male	571	10	2	1	1	-	-	4	6	12	27	48
Female	574	10	1	-	-	1	-	1	3	13	7	18
Black, both sexes	187	4	1	-	-	-	-	1	1	8	15	21
Male	122	2	-	-	-	-	-	1	1	5	14	17
Female	65	2	1	-	-	-	-	-	-	3	1	4
American Indian 1/, both sexes	15	1	1	-	-	-	-	-	-	1	3	1
Male	11	-	1	-	-	-	-	-	-	1	2	1
Female	4	1	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	31	-	-	-	-	-	-	-	1	1	1	2
Male	18	-	-	-	-	-	-	-	1	-	1	1
Female	13	-	-	-	-	-	-	-	-	1	-	1
Alabama												
All races, both sexes	36	-	-	-	-	-	-	-	-	2	-	2
Male	17	-	-	-	-	-	-	-	-	1	-	1
Female	19	-	-	-	-	-	-	-	-	1	-	1
White, both sexes	28	-	-	-	-	-	-	-	-	-	-	1
Male	13	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	8	-	-	-	-	-	-	-	-	2	-	1
Male	4	-	-	-	-	-	-	-	-	1	-	-
Female	4	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Alaska												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	14	-	-	-	-	-	-	-	-	-	-	-
Male	10	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	-	-	-	-	-	-	-	-	-	-	-
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Arkansas												
All races, both sexes	13	-	-	-	-	-	-	-	-	-	1	1
Male	8	-	-	-	-	-	-	-	-	-	1	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	1
Male	6	-	-	-	-	-	-	-	-	-	-	1
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	118	2	-	-	-	-	-	1	2	5	5	8
Male	61	1	-	-	-	-	-	1	1	2	4	7
Female	57	1	-	-	-	-	-	-	1	3	1	1
White, both sexes	100	2	-	-	-	-	-	1	1	5	4	6
Male	49	1	-	-	-	-	-	1	-	2	3	6
Female	51	1	-	-	-	-	-	-	1	3	1	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	1
Male	7	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	9	-	-	-	-	-	-	-	1	-	-	1
Male	5	-	-	-	-	-	-	-	1	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Colorado												
All races, both sexes	11	-	1	-	-	-	-	-	-	-	1	-
Male	6	-	1	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	1	-	-	-	-	-	-	-	-	-
Male	4	-	1	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	19	1	-	-	-	-	-	-	-	-	-	3
Male	9	1	-	-	-	-	-	-	-	-	-	3
Female	10	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	1	-	-	-	-	-	-	-	-	-	3
Male	8	1	-	-	-	-	-	-	-	-	-	3
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Delaware												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Florida												
All races, both sexes	80	1	-	-	-	-	-	-	-	2	5	6
Male	43	1	-	-	-	-	-	-	-	-	5	4
Female	37	-	-	-	-	-	-	-	-	2	-	2
White, both sexes	70	1	-	-	-	-	-	-	-	1	3	5
Male	38	1	-	-	-	-	-	-	-	-	3	4
Female	32	-	-	-	-	-	-	-	-	1	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	1	2	1
Male	4	-	-	-	-	-	-	-	-	-	2	-
Female	5	-	-	-	-	-	-	-	-	1	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	41	-	1	-	-	-	-	1	-	-	2	2
Male	21	-	-	-	-	-	-	-	-	-	2	2
Female	20	-	1	-	-	-	-	1	-	-	-	-
White, both sexes	27	-	-	-	-	-	-	1	-	-	-	1
Male	12	-	-	-	-	-	-	-	-	-	-	1
Female	15	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	14	-	1	-	-	-	-	-	-	-	2	1
Male	9	-	-	-	-	-	-	-	-	-	2	1
Female	5	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Hawaii												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	-	-
Male	7	-	-	-	-	-	-	-	-	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Idaho												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Illinois												
All races, both sexes	69	1	1	1	-	-	-	-	1	4	1	3
Male	41	-	1	1	-	-	-	-	1	4	1	3
Female	28	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	49	1	1	1	-	-	-	-	-	3	1	1
Male	27	-	1	1	-	-	-	-	-	3	1	1
Female	22	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	18	-	-	-	-	-	-	-	1	1	-	2
Male	13	-	-	-	-	-	-	-	1	1	-	2
Female	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes	34	-	-	-	-	-	-	-	-	2	1	2
Male	15	-	-	-	-	-	-	-	-	-	1	2
Female	19	-	-	-	-	-	-	-	-	2	-	-
White, both sexes	30	-	-	-	-	-	-	-	-	1	1	1
Male	13	-	-	-	-	-	-	-	-	-	1	1
Female	17	-	-	-	-	-	-	-	-	1	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Iowa												
All races, both sexes	19	-	-	-	-	1	-	-	-	2	-	2
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	9	-	-	-	-	1	-	-	-	2	-	1
White, both sexes	18	-	-	-	-	1	-	-	-	2	-	1
Male	9	-	-	-	-	-	-	-	-	-	-	-
Female	9	-	-	-	-	1	-	-	-	2	-	1
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	10	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	1	1
White, both sexes	10	-	-	-	-	-	-	-	-	-	2	1
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	5	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Kentucky												
All races, both sexes	18	-	-	-	-	-	-	-	-	1	-	1
Male	9	-	-	-	-	-	-	-	-	1	-	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	16	-	-	-	-	-	-	-	-	1	-	-
Male	7	-	-	-	-	-	-	-	-	1	-	-
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	14	-	-	-	-	-	-	-	-	1	-	2
Male	6	-	-	-	-	-	-	-	-	1	-	2
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	9	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	-	-	1	-	2
Male	5	-	-	-	-	-	-	-	-	1	-	2
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Maine												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	1
Male	1	-	-	-	-	-	-	-	-	-	-	1
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	17	-	-	-	-	-	-	2	-	2	-	3
Male	8	-	-	-	-	-	-	2	-	1	-	3
Female	9	-	-	-	-	-	-	-	-	1	-	-
White, both sexes	12	-	-	-	-	-	-	1	-	1	-	2
Male	6	-	-	-	-	-	-	1	-	1	-	2
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	5	-	-	-	-	-	-	1	-	1	-	1
Male	2	-	-	-	-	-	-	1	-	-	-	1
Female	3	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Massachusetts												
All races, both sexes	36	1	-	-	-	-	-	-	-	-	2	-
Male	21	-	-	-	-	-	-	-	-	-	2	-
Female	15	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	32	1	-	-	-	-	-	-	-	-	2	-
Male	18	-	-	-	-	-	-	-	-	-	2	-
Female	14	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	43	1	-	-	-	-	-	-	-	-	1	2
Male	23	1	-	-	-	-	-	-	-	-	1	1
Female	20	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	36	1	-	-	-	-	-	-	-	-	1	2
Male	19	1	-	-	-	-	-	-	-	-	1	1
Female	17	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Minnesota												
All races, both sexes	26	2	-	-	-	-	-	-	1	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	1
Female	17	2	-	-	-	-	-	-	1	-	-	2
White, both sexes	25	2	-	-	-	-	-	-	1	-	-	3
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	17	2	-	-	-	-	-	-	1	-	-	2
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	15	-	-	-	-	-	-	-	-	-	-	3
Male	9	-	-	-	-	-	-	-	-	-	-	2
Female	6	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	3
Male	4	-	-	-	-	-	-	-	-	-	-	2
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Missouri												
All races, both sexes	31	-	-	-	-	-	-	-	-	1	1	2
Male	15	-	-	-	-	-	-	-	-	1	1	2
Female	16	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	28	-	-	-	-	-	-	-	-	-	1	2
Male	13	-	-	-	-	-	-	-	-	-	1	2
Female	15	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	3	-	-	-	-	-	-	-	-	1	-	-
Male	2	-	-	-	-	-	-	-	-	1	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	6	-	-	-	-	-	-	-	-	-	-	-
Male	5	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Nebraska												
All races, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	11	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	8	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	8	-	-	-	-	-	-	-	-	-	-	1
Male	4	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
New Hampshire												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	1	1
Male	3	-	-	-	-	-	-	-	-	-	1	1
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	39	-	-	-	1	-	-	-	-	-	-	4
Male	20	-	-	-	1	-	-	-	-	-	-	3
Female	19	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	38	-	-	-	1	-	-	-	-	-	-	4
Male	20	-	-	-	1	-	-	-	-	-	-	3
Female	18	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
New Mexico												
All races, both sexes	15	1	-	-	-	-	-	-	-	-	2	2
Male	11	-	-	-	-	-	-	-	-	-	2	2
Female	4	1	-	-	-	-	-	-	-	-	-	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	2
Male	9	-	-	-	-	-	-	-	-	-	1	2
Female	3	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	83	1	-	-	-	-	-	-	-	3	2	4
Male	45	1	-	-	-	-	-	-	-	2	1	2
Female	38	-	-	-	-	-	-	-	-	1	1	2
White, both sexes	73	1	-	-	-	-	-	-	-	3	2	3
Male	37	1	-	-	-	-	-	-	-	2	1	1
Female	36	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes	7	-	-	-	-	-	-	-	-	-	-	-
Male	6	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	3	-	-	-	-	-	-	-	-	-	-	1
Male	2	-	-	-	-	-	-	-	-	-	-	1
Female	1	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
North Carolina												
All races, both sexes	55	4	-	-	-	-	-	-	2	-	2	4
Male	30	3	-	-	-	-	-	-	1	-	1	3
Female	25	1	-	-	-	-	-	-	1	-	1	1
White, both sexes	38	2	-	-	-	-	-	-	2	-	-	3
Male	22	1	-	-	-	-	-	-	1	-	-	2
Female	16	1	-	-	-	-	-	-	1	-	-	1
Black, both sexes	14	2	-	-	-	-	-	-	-	-	1	1
Male	8	2	-	-	-	-	-	-	-	-	1	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Ohio												
All races, both sexes	62	1	-	-	-	-	-	-	1	-	2	1
Male	27	1	-	-	-	-	-	-	1	-	2	-
Female	35	-	-	-	-	-	-	-	-	-	-	1
White, both sexes	53	1	-	-	-	-	-	-	1	-	1	1
Male	21	1	-	-	-	-	-	-	1	-	1	-
Female	32	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes	9	-	-	-	-	-	-	-	-	-	1	-
Male	6	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	16	-	-	-	-	-	-	-	-	-	3	-
Male	8	-	-	-	-	-	-	-	-	-	2	-
Female	8	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	2	-
Male	5	-	-	-	-	-	-	-	-	-	1	-
Female	7	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	2	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Oregon												
All races, both sexes	25	1	-	-	-	-	-	-	1	1	-	1
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	15	1	-	-	-	-	-	-	-	1	-	1
White, both sexes	25	1	-	-	-	-	-	-	1	1	-	1
Male	10	-	-	-	-	-	-	-	1	-	-	-
Female	15	1	-	-	-	-	-	-	-	1	-	1
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	74	2	-	-	-	-	-	1	1	3	6	4
Male	34	-	-	-	-	-	-	1	1	2	5	4
Female	40	2	-	-	-	-	-	-	-	1	1	-
White, both sexes	63	2	-	-	-	-	-	1	1	1	4	3
Male	26	-	-	-	-	-	-	1	1	-	4	3
Female	37	2	-	-	-	-	-	-	-	1	-	-
Black, both sexes	10	-	-	-	-	-	-	-	-	1	2	1
Male	7	-	-	-	-	-	-	-	-	1	1	1
Female	3	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	1	-	-
Male	1	-	-	-	-	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Rhode Island												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	4	-	-	-	-	-	-	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	26	-	-	-	-	-	-	-	-	-	-	3
Male	19	-	-	-	-	-	-	-	-	-	-	3
Female	7	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	17	-	-	-	-	-	-	-	-	-	-	1
Male	11	-	-	-	-	-	-	-	-	-	-	1
Female	6	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	9	-	-	-	-	-	-	-	-	-	-	2
Male	8	-	-	-	-	-	-	-	-	-	-	2
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
South Dakota												
All races, both sexes	4	-	-	-	-	-	-	-	-	-	1	-
Male	3	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	32	-	-	-	-	-	-	-	-	1	2	5
Male	19	-	-	-	-	-	-	-	-	1	2	2
Female	13	-	-	-	-	-	-	-	-	-	-	3
White, both sexes	26	-	-	-	-	-	-	-	-	1	2	4
Male	16	-	-	-	-	-	-	-	-	1	2	2
Female	10	-	-	-	-	-	-	-	-	-	-	2
Black, both sexes	6	-	-	-	-	-	-	-	-	-	-	1
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Texas												
All races, both sexes	95	1	1	-	-	-	-	1	1	-	5	8
Male	52	1	-	-	-	-	-	1	1	-	4	6
Female	43	-	1	-	-	-	-	-	-	-	1	2
White, both sexes	83	1	1	-	-	-	-	1	1	-	3	8
Male	44	1	-	-	-	-	-	1	1	-	2	6
Female	39	-	1	-	-	-	-	-	-	-	1	2
Black, both sexes	10	-	-	-	-	-	-	-	-	-	1	-
Male	7	-	-	-	-	-	-	-	-	-	1	-
Female	3	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	1	-
Male	1	-	-	-	-	-	-	-	-	-	1	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Utah												
All races, both sexes	14	2	1	-	-	-	-	-	1	-	-	-
Male	9	2	1	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	13	2	-	-	-	-	-	-	1	-	-	-
Male	8	2	-	-	-	-	-	-	1	-	-	-
Female	5	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	1	-	-	-	-	-	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Vermont												
All races, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	37	2	-	-	-	-	-	-	-	-	2	2
Male	21	-	-	-	-	-	-	-	-	-	2	2
Female	16	2	-	-	-	-	-	-	-	-	-	-
White, both sexes	24	-	-	-	-	-	-	-	-	-	1	1
Male	15	-	-	-	-	-	-	-	-	-	1	1
Female	9	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	12	2	-	-	-	-	-	-	-	-	1	1
Male	5	-	-	-	-	-	-	-	-	-	1	1
Female	7	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Washington												
All races, both sexes	28	1	-	-	-	-	-	-	-	3	-	1
Male	10	-	-	-	-	-	-	-	-	-	-	1
Female	18	1	-	-	-	-	-	-	-	3	-	-
White, both sexes	23	1	-	-	-	-	-	-	-	3	-	1
Male	8	-	-	-	-	-	-	-	-	-	-	1
Female	15	1	-	-	-	-	-	-	-	3	-	-
Black, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
White, both sexes	12	-	-	-	-	-	-	-	-	-	1	-
Male	8	-	-	-	-	-	-	-	-	-	-	-
Female	4	-	-	-	-	-	-	-	-	-	1	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All other and unspecified disorders of circulatory system (I83-I99)												
Wisconsin												
All races, both sexes	24	-	-	-	-	-	-	-	-	2	-	1
Male	11	-	-	-	-	-	-	-	-	2	-	1
Female	13	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	22	-	-	-	-	-	-	-	-	2	-	1
Male	10	-	-	-	-	-	-	-	-	2	-	1
Female	12	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	2	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	5	-	-	-	-	-	-	-	-	-	-	-
Male	3	-	-	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
United States												
All races, both sexes	122	99	96	101	120	128	164	160	106	47	7	-
Male	90	71	68	61	62	51	69	47	36	8	-	-
Female	32	28	28	40	58	77	95	113	70	39	7	-
White, both sexes	96	76	72	80	103	110	147	150	96	44	6	-
Male	70	54	51	47	52	46	60	44	30	6	-	-
Female	26	22	21	33	51	64	87	106	66	38	6	-
Black, both sexes	23	21	19	15	10	14	14	9	7	3	1	-
Male	17	16	13	11	5	4	8	3	3	2	-	-
Female	6	5	6	4	5	10	6	6	4	1	1	-
American Indian 1/, both sexes	2	-	2	2	1	-	1	-	-	-	-	-
Male	2	-	2	1	1	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	2	3	4	6	4	2	1	3	-	-	-
Male	1	1	2	2	4	1	1	-	3	-	-	-
Female	-	1	1	2	2	3	1	1	-	-	-	-
Alabama												
All races, both sexes	-	4	4	1	3	7	4	4	3	1	1	-
Male	-	4	2	-	1	4	2	-	2	-	-	-
Female	-	-	2	1	2	3	2	4	1	1	1	-
White, both sexes	-	3	2	1	3	5	4	4	3	1	1	-
Male	-	3	1	-	1	3	2	-	2	-	-	-
Female	-	-	1	1	2	2	2	4	1	1	1	-
Black, both sexes	-	1	2	-	-	2	-	-	-	-	-	-
Male	-	1	1	-	-	1	-	-	-	-	-	-
Female	-	-	1	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Alaska												
All races, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes	2	1	-	2	2	2	2	2	-	1	-	-
Male	2	1	-	2	1	-	2	1	-	1	-	-
Female	-	-	-	-	1	2	-	1	-	-	-	-
White, both sexes	2	1	-	1	2	2	2	2	-	1	-	-
Male	2	1	-	1	1	-	2	1	-	1	-	-
Female	-	-	-	-	1	2	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Arkansas												
All races, both sexes	-	-	1	-	1	2	2	1	3	1	-	-
Male	-	-	1	-	1	1	1	1	1	-	-	-
Female	-	-	-	-	-	1	1	-	2	1	-	-
White, both sexes	-	-	1	-	1	2	1	1	3	1	-	-
Male	-	-	1	-	1	1	-	1	1	-	-	-
Female	-	-	-	-	-	1	1	-	2	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	1	-	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes	12	9	8	8	11	14	8	17	4	4	-	-
Male	7	7	4	5	7	5	5	5	-	-	-	-
Female	5	2	4	3	4	9	3	12	4	4	-	-
White, both sexes	11	6	6	7	6	14	7	16	4	4	-	-
Male	6	5	2	5	4	5	4	5	-	-	-	-
Female	5	1	4	2	2	9	3	11	4	4	-	-
Black, both sexes	-	3	1	-	1	-	1	1	-	-	-	-
Male	-	2	1	-	1	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	1	-	1	1	4	-	-	-	-	-	-	-
Male	1	-	1	-	2	-	-	-	-	-	-	-
Female	-	-	-	1	2	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Colorado												
All races, both sexes	1	1	-	1	-	1	1	2	2	-	-	-
Male	1	1	-	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	2	-	-	-
White, both sexes	1	1	-	-	-	1	1	2	2	-	-	-
Male	1	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	1	-	2	2	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Connecticut												
All races, both sexes	3	1	2	1	1	-	3	1	2	1	-	-
Male	2	-	1	1	-	-	1	-	-	-	-	-
Female	1	1	1	-	1	-	2	1	2	1	-	-
White, both sexes	3	1	1	1	-	-	3	1	2	1	-	-
Male	2	-	-	1	-	-	1	-	-	-	-	-
Female	1	1	1	-	-	-	2	1	2	1	-	-
Black, both sexes	-	-	1	-	1	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Delaware												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Dist. of Columbia												
All races, both sexes	2	-	-	-	-	-	1	-	-	-	-	-
Male	2	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Florida												
All races, both sexes	11	3	7	5	8	2	12	9	4	4	1	-
Male	10	3	4	4	4	1	4	1	-	2	-	-
Female	1	-	3	1	4	1	8	8	4	2	1	-
White, both sexes	11	3	7	4	7	2	12	7	4	3	-	-
Male	10	3	4	3	3	1	4	1	-	1	-	-
Female	1	-	3	1	4	1	8	6	4	2	-	-
Black, both sexes	-	-	-	-	1	-	-	2	-	1	1	-
Male	-	-	-	-	1	-	-	-	-	1	-	-
Female	-	-	-	-	-	-	-	2	-	-	1	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes	5	2	3	4	4	3	4	6	2	1	1	-
Male	4	2	2	4	1	1	1	1	1	-	-	-
Female	1	-	1	-	3	2	3	5	1	1	1	-
White, both sexes	3	1	1	3	3	3	3	5	1	1	1	-
Male	2	1	1	3	1	1	1	1	-	-	-	-
Female	1	-	-	-	2	2	2	4	1	1	1	-
Black, both sexes	2	1	2	1	1	-	1	1	1	-	-	-
Male	2	1	1	1	-	-	-	-	1	-	-	-
Female	-	-	1	-	1	-	1	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Hawaii												
All races, both sexes	-	1	-	1	2	-	5	-	3	-	-	-
Male	-	1	-	-	1	-	2	-	3	-	-	-
Female	-	-	-	1	1	-	3	-	-	-	-	-
White, both sexes	-	-	-	-	1	-	4	-	-	-	-	-
Male	-	-	-	-	-	-	2	-	-	-	-	-
Female	-	-	-	-	1	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	1	-	1	-	3	-	-	-
Male	-	1	-	-	1	-	-	-	3	-	-	-
Female	-	-	-	1	-	-	1	-	-	-	-	-
Idaho												
All races, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	-	1	-	-	-	-	1	-	1	-	-	-
Male	-	1	-	-	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Illinois												
All races, both sexes	5	6	7	9	2	7	8	9	2	1	1	-
Male	4	3	5	5	2	4	4	2	1	-	-	-
Female	1	3	2	4	-	3	4	7	1	1	1	-
White, both sexes	5	3	3	6	2	5	6	7	2	1	1	-
Male	4	1	2	3	2	3	3	1	1	-	-	-
Female	1	2	1	3	-	2	3	6	1	1	1	-
Black, both sexes	-	3	3	3	-	2	2	1	-	-	-	-
Male	-	2	2	2	-	1	1	1	-	-	-	-
Female	-	1	1	1	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	1	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Indiana												
All races, both sexes	3	1	2	4	6	3	3	3	2	2	-	-
Male	3	1	1	2	2	-	2	1	-	-	-	-
Female	-	-	1	2	4	3	1	2	2	2	-	-
White, both sexes	3	1	2	4	5	2	3	3	2	2	-	-
Male	3	1	1	2	1	-	2	1	-	-	-	-
Female	-	-	1	2	4	2	1	2	2	2	-	-
Black, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Iowa												
All races, both sexes	1	2	1	-	1	2	3	1	1	2	-	-
Male	1	2	1	-	1	1	2	-	1	-	-	-
Female	-	-	-	-	-	1	1	1	-	2	-	-
White, both sexes	1	2	1	-	1	2	3	1	1	2	-	-
Male	1	2	1	-	1	1	2	-	1	-	-	-
Female	-	-	-	-	-	1	1	1	-	2	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Kansas												
All races, both sexes	2	-	1	-	1	-	-	1	1	1	-	-
Male	1	-	-	-	1	-	-	-	1	1	-	-
Female	1	-	1	-	-	-	-	1	-	-	-	-
White, both sexes	2	-	1	-	1	-	-	1	1	1	-	-
Male	1	-	-	-	1	-	-	-	1	1	-	-
Female	1	-	1	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Kentucky												
All races, both sexes	1	2	2	2	-	3	2	2	2	-	-	-
Male	1	1	2	-	-	1	2	-	-	-	-	-
Female	-	1	-	2	-	2	-	2	2	-	-	-
White, both sexes	1	2	1	2	-	3	2	2	2	-	-	-
Male	1	1	1	-	-	1	2	-	-	-	-	-
Female	-	1	-	2	-	2	-	2	2	-	-	-
Black, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana												
All races, both sexes	2	-	1	-	1	2	2	-	1	1	1	-
Male	-	-	1	-	1	-	1	-	-	-	-	-
Female	2	-	-	-	-	2	1	-	1	1	1	-
White, both sexes	2	-	1	-	-	2	1	-	1	1	1	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	2	-	-	-	-	2	1	-	1	1	1	-
Black, both sexes	-	-	-	-	1	-	1	-	-	-	-	-
Male	-	-	-	-	1	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Maine												
All races, both sexes	1	1	-	-	-	1	2	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
White, both sexes	1	1	-	-	-	1	2	-	-	-	-	-
Male	1	-	-	-	-	1	-	-	-	-	-	-
Female	-	1	-	-	-	-	2	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes	-	3	-	-	-	3	-	1	2	1	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	2	-	-	-	3	-	1	1	1	-	-
White, both sexes	-	2	-	-	-	2	-	1	2	1	-	-
Male	-	1	-	-	-	-	-	-	1	-	-	-
Female	-	1	-	-	-	2	-	1	1	1	-	-
Black, both sexes	-	1	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Massachusetts												
All races, both sexes	-	3	1	3	3	5	5	6	6	1	-	-
Male	-	3	1	1	2	2	4	2	4	-	-	-
Female	-	-	-	2	1	3	1	4	2	1	-	-
White, both sexes	-	3	1	2	3	4	4	6	5	1	-	-
Male	-	3	1	-	2	2	3	2	3	-	-	-
Female	-	-	-	2	1	2	1	4	2	1	-	-
Black, both sexes	-	-	-	1	-	1	1	-	1	-	-	-
Male	-	-	-	1	-	-	1	-	1	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Michigan												
All races, both sexes	7	3	2	4	1	2	8	8	2	2	-	-
Male	5	1	2	2	-	-	4	5	1	-	-	-
Female	2	2	-	2	1	2	4	3	1	2	-	-
White, both sexes	4	3	2	4	1	1	6	7	2	2	-	-
Male	4	1	2	2	-	-	2	4	1	-	-	-
Female	-	2	-	2	1	1	4	3	1	2	-	-
Black, both sexes	2	-	-	-	-	1	2	1	-	-	-	-
Male	-	-	-	-	-	-	2	1	-	-	-	-
Female	2	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Minnesota												
All races, both sexes	-	3	-	2	2	2	2	4	3	2	-	-
Male	-	1	-	-	2	1	1	1	2	-	-	-
Female	-	2	-	2	-	1	1	3	1	2	-	-
White, both sexes	-	2	-	2	2	2	2	4	3	2	-	-
Male	-	-	-	-	2	1	1	1	2	-	-	-
Female	-	2	-	2	-	1	1	3	1	2	-	-
Black, both sexes	-	1	-	-	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Mississippi												
All races, both sexes	2	-	2	-	2	1	2	3	-	-	-	-
Male	2	-	2	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	1	1	1	2	-	-	-	-
White, both sexes	1	-	1	-	1	1	2	2	-	-	-	-
Male	1	-	1	-	1	-	1	1	-	-	-	-
Female	-	-	-	-	-	1	1	1	-	-	-	-
Black, both sexes	1	-	1	-	1	-	-	1	-	-	-	-
Male	1	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Missouri												
All races, both sexes	1	2	1	2	5	2	7	4	2	1	-	-
Male	-	2	1	1	5	1	1	-	-	-	-	-
Female	1	-	-	1	-	1	6	4	2	1	-	-
White, both sexes	1	1	1	1	5	2	7	4	2	1	-	-
Male	-	1	1	1	5	1	1	-	-	-	-	-
Female	1	-	-	-	-	1	6	4	2	1	-	-
Black, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	1	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Montana												
All races, both sexes	1	-	-	-	1	1	-	1	2	-	-	-
Male	1	-	-	-	1	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	-	-	-	1	1	-	1	2	-	-	-
Male	1	-	-	-	1	1	-	1	1	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Nebraska												
All races, both sexes	-	-	-	-	1	-	3	2	4	1	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	1	-	2	1	3	1	-	-
White, both sexes	-	-	-	-	1	-	3	2	4	1	-	-
Male	-	-	-	-	-	-	1	1	1	-	-	-
Female	-	-	-	-	1	-	2	1	3	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Nevada												
All races, both sexes	1	1	-	1	1	-	1	-	2	-	-	-
Male	-	-	-	1	1	-	1	-	1	-	-	-
Female	1	1	-	-	-	-	-	-	1	-	-	-
White, both sexes	1	1	-	1	1	-	1	-	2	-	-	-
Male	-	-	-	1	1	-	1	-	1	-	-	-
Female	1	1	-	-	-	-	-	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
New Hampshire												
All races, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	1	-	-	-	-
Male	-	-	-	-	-	-	1	-	-	-	-	-
Female	-	1	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes	1	3	3	6	2	6	5	5	1	2	-	-
Male	1	2	2	3	1	3	2	1	-	1	-	-
Female	-	1	1	3	1	3	3	4	1	1	-	-
White, both sexes	1	3	3	6	2	5	5	5	1	2	-	-
Male	1	2	2	3	1	3	2	1	-	1	-	-
Female	-	1	1	3	1	2	3	4	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
New Mexico												
All races, both sexes	-	1	3	-	1	-	2	1	1	1	-	-
Male	-	1	3	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
White, both sexes	-	1	2	-	1	-	2	1	1	1	-	-
Male	-	1	2	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	1	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes	6	7	6	5	13	11	10	9	4	2	-	-
Male	2	6	6	3	7	7	3	2	2	1	-	-
Female	4	1	-	2	6	4	7	7	2	1	-	-
White, both sexes	6	6	6	4	11	9	9	8	3	2	-	-
Male	2	5	6	2	5	6	2	1	2	1	-	-
Female	4	1	-	2	6	3	7	7	1	1	-	-
Black, both sexes	-	1	-	1	1	1	1	1	1	-	-	-
Male	-	1	-	1	1	1	1	1	-	-	-	-
Female	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	1	1	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
North Carolina												
All races, both sexes	6	4	4	5	3	4	5	5	4	1	2	-
Male	6	4	2	3	1	2	2	2	-	-	-	-
Female	-	-	2	2	2	2	3	3	4	1	2	-
White, both sexes	6	3	2	2	3	2	4	5	2	-	2	-
Male	6	3	1	2	1	1	2	2	-	-	-	-
Female	-	-	1	-	2	1	2	3	2	-	2	-
Black, both sexes	-	1	1	2	-	2	1	-	2	1	-	-
Male	-	1	1	1	-	1	-	-	-	-	-	-
Female	-	-	-	1	-	1	1	-	2	1	-	-
American Indian 1/, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	1	-	-	-	-	-	-	-	-	-
North Dakota												
All races, both sexes	-	-	-	-	1	-	3	1	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	-	-	-	1	-	3	1	-	-	-	-
Male	-	-	-	-	1	-	2	-	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Ohio												
All races, both sexes	3	5	5	3	6	6	6	11	10	2	-	-
Male	2	2	2	2	2	3	1	5	3	1	-	-
Female	1	3	3	1	4	3	5	6	7	1	-	-
White, both sexes	2	3	3	3	5	6	6	11	9	1	-	-
Male	1	1	1	2	2	3	1	5	2	-	-	-
Female	1	2	2	1	3	3	5	6	7	1	-	-
Black, both sexes	1	2	2	-	1	-	-	-	1	1	-	-
Male	1	1	1	-	-	-	-	-	1	1	-	-
Female	-	1	1	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes	2	-	2	2	3	1	1	-	2	-	-	-
Male	2	-	1	1	1	-	-	-	1	-	-	-
Female	-	-	1	1	2	1	1	-	1	-	-	-
White, both sexes	1	-	2	1	2	1	1	-	2	-	-	-
Male	1	-	1	1	-	-	-	-	1	-	-	-
Female	-	-	1	-	2	1	1	-	1	-	-	-
Black, both sexes	1	-	-	1	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	1	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Oregon												
All races, both sexes	-	3	3	3	1	1	4	1	2	3	-	-
Male	-	3	2	1	1	1	-	-	1	-	-	-
Female	-	-	1	2	-	-	4	1	1	3	-	-
White, both sexes	-	3	3	3	1	1	4	1	2	3	-	-
Male	-	3	2	1	1	1	-	-	1	-	-	-
Female	-	-	1	2	-	-	4	1	1	3	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes	9	3	6	5	5	6	6	8	8	1	-	-
Male	6	1	5	2	1	1	1	1	3	-	-	-
Female	3	2	1	3	4	5	5	7	5	1	-	-
White, both sexes	5	2	6	5	5	6	5	8	8	1	-	-
Male	3	-	5	2	1	1	1	1	3	-	-	-
Female	2	2	1	3	4	5	4	7	5	1	-	-
Black, both sexes	4	1	-	-	-	-	1	-	-	-	-	-
Male	3	1	-	-	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Rhode Island												
All races, both sexes	-	1	-	-	-	-	1	2	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
White, both sexes	-	1	-	-	-	-	1	2	-	-	-	-
Male	-	1	-	-	-	-	-	1	-	-	-	-
Female	-	-	-	-	-	-	1	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina												
All races, both sexes	6	2	3	3	-	2	4	2	1	-	-	-
Male	3	2	3	1	-	2	3	1	1	-	-	-
Female	3	-	-	2	-	-	1	1	-	-	-	-
White, both sexes	3	2	2	2	-	2	3	1	1	-	-	-
Male	-	2	2	-	-	2	2	1	1	-	-	-
Female	3	-	-	2	-	-	1	-	-	-	-	-
Black, both sexes	3	-	1	1	-	-	1	1	-	-	-	-
Male	3	-	1	1	-	-	1	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
South Dakota												
All races, both sexes	-	-	1	1	-	-	-	1	-	-	-	-
Male	-	-	1	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	1	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	1	-	-	-	-	-	-	-	-	-
Male	-	-	1	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes	3	4	2	4	-	3	1	6	1	-	-	-
Male	2	3	2	4	-	2	-	1	-	-	-	-
Female	1	1	-	-	-	1	1	5	1	-	-	-
White, both sexes	1	3	2	3	-	2	1	6	1	-	-	-
Male	1	2	2	3	-	2	-	1	-	-	-	-
Female	-	1	-	-	-	-	1	5	1	-	-	-
Black, both sexes	2	1	-	1	-	1	-	-	-	-	-	-
Male	1	1	-	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Texas												
All races, both sexes	12	9	8	6	11	11	6	9	4	2	-	-
Male	10	8	6	4	4	-	3	3	1	-	-	-
Female	2	1	2	2	7	11	3	6	3	2	-	-
White, both sexes	10	6	7	5	11	9	6	9	3	2	-	-
Male	9	5	5	3	4	-	3	3	1	-	-	-
Female	1	1	2	2	7	9	3	6	2	2	-	-
Black, both sexes	2	3	1	1	-	1	-	-	1	-	-	-
Male	1	3	1	1	-	-	-	-	-	-	-	-
Female	1	-	-	-	-	1	-	-	1	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	1	-	-	-	-	-	-
Utah												
All races, both sexes	-	-	-	1	4	3	-	1	-	1	-	-
Male	-	-	-	1	2	2	-	-	-	-	-	-
Female	-	-	-	-	2	1	-	1	-	1	-	-
White, both sexes	-	-	-	1	4	3	-	1	-	1	-	-
Male	-	-	-	1	2	2	-	-	-	-	-	-
Female	-	-	-	-	2	1	-	1	-	1	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Vermont												
All races, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
White, both sexes	-	-	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	1	-	-	-	-	-	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes	5	-	4	1	2	4	7	2	3	3	-	-
Male	4	-	2	1	2	3	1	2	1	1	-	-
Female	1	-	2	-	-	1	6	-	2	2	-	-
White, both sexes	2	-	1	1	2	2	6	2	3	3	-	-
Male	2	-	1	1	2	2	1	2	1	1	-	-
Female	-	-	-	-	-	-	5	-	2	2	-	-
Black, both sexes	3	-	3	-	-	1	1	-	-	-	-	-
Male	2	-	1	-	-	-	-	-	-	-	-	-
Female	1	-	2	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	1	-	-	-	-	-	-
Male	-	-	-	-	-	1	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Washington												
All races, both sexes	3	3	-	2	7	-	2	2	4	-	-	-
Male	2	1	-	2	2	-	-	1	1	-	-	-
Female	1	2	-	-	5	-	2	1	3	-	-	-
White, both sexes	2	2	-	1	6	-	1	2	4	-	-	-
Male	1	1	-	1	2	-	-	1	1	-	-	-
Female	1	1	-	-	4	-	1	1	3	-	-	-
Black, both sexes	-	-	-	-	1	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes	1	-	-	-	-	-	1	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	1	-	1	-	-	-	-	-	-	-	-
Male	-	-	-	1	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
West Virginia												
All races, both sexes	1	2	1	-	-	1	1	3	2	-	-	-
Male	1	2	1	-	-	1	1	1	1	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
White, both sexes	1	2	1	-	-	1	1	3	2	-	-	-
Male	1	2	1	-	-	1	1	1	1	-	-	-
Female	-	-	-	-	-	-	-	2	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2005

[Race categories are consistent with the 1977 Office of Management and Budget standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All other and unspecified disorders of circulatory system (I83-I99)												
Wisconsin												
All races, both sexes	2	1	-	1	2	4	6	2	2	1	-	-
Male	2	-	-	1	1	-	2	2	-	-	-	-
Female	-	1	-	-	1	4	4	-	2	1	-	-
White, both sexes	1	-	-	1	2	4	6	2	2	1	-	-
Male	1	-	-	1	1	-	2	2	-	-	-	-
Female	-	-	-	-	1	4	4	-	2	1	-	-
Black, both sexes	1	1	-	-	-	-	-	-	-	-	-	-
Male	1	-	-	-	-	-	-	-	-	-	-	-
Female	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Wyoming												
All races, both sexes	-	-	-	1	-	-	2	1	1	-	-	-
Male	-	-	-	1	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes	-	-	-	1	-	-	2	1	1	-	-	-
Male	-	-	-	1	-	-	1	1	-	-	-	-
Female	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes	-	-	-	-	-	-	-	-	-	-	-	-
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	-	-	-	-	-	-	-	-	-	-	-	-