

Person with disabilities experiencing problems accessing
NCHS birth and death tables should contact
nchsed@cdc.gov, or call 301-458-4688

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
All races 3/, both sexes										
2006.....	810.4	690.7	28.4	13.9	16.6	64.4	100.2	100.9	111.9	148.7
2005.....	825.9	692.5	29.4	14.5	18.1	65.1	97.6	97.5	111.4	149.6
2004.....	816.5	685.2	29.9	14.7	18.7	66.1	94.0	96.0	107.9	151.7
2003.....	841.9	700.0	31.5	14.7	19.1	66.4	96.4	95.2	111.3	159.1
2002.....	847.3	695.0	31.2	15.2	19.5	67.8	95.2	94.7	111.6	161.3
2001.....	848.5	683.4	33.3	15.3	19.2	66.9	95.0	96.2	113.5	165.9
2000.....	854.0	736.7	32.4	15.8	20.3	67.1	93.6	91.2	111.0	159.2
1999.....	857.0	736.0	34.2	16.9	20.4	68.6	90.8	90.4	113.4	160.8
All races 3/, male										
2006.....	814.8	756.3	30.5	15.4	19.6	90.7	148.0	143.4	150.4	189.0
2005.....	827.2	762.3	33.4	15.6	21.5	91.6	143.9	138.7	148.1	189.4
2004.....	817.6	753.7	32.4	16.4	21.7	91.0	138.0	135.6	143.2	193.8
2003.....	840.3	777.4	35.1	16.2	23.1	92.3	140.3	133.5	148.9	202.7
2002.....	846.6	761.5	35.2	16.7	23.1	94.0	140.8	134.4	149.4	206.6
2001.....	846.4	749.8	37.0	16.7	22.8	93.7	141.1	135.6	151.2	213.7
2000.....	853.0	806.5	35.9	17.6	24.2	93.3	138.0	128.8	147.9	205.4
1999.....	859.2	805.0	37.9	18.6	24.4	94.7	133.1	127.0	152.0	207.6
All races 3/, female										
2006.....	806.1	622.0	26.3	12.3	13.3	36.8	49.0	56.5	72.5	107.9
2005.....	824.6	619.4	25.1	13.4	14.4	37.2	48.2	54.4	73.7	109.3
2004.....	815.4	613.4	27.3	13.0	15.4	39.9	47.3	54.5	72.0	109.3
2003.....	843.4	619.1	27.8	13.0	15.0	39.0	49.8	55.4	73.0	115.3
2002.....	848.0	625.3	27.0	13.5	15.9	40.1	47.3	53.6	73.3	116.1
2001.....	850.4	613.9	29.5	13.9	15.3	38.5	46.8	55.8	75.2	118.2
2000.....	855.0	663.4	28.7	14.0	16.1	39.3	47.1	52.8	73.6	113.2
1999.....	854.9	663.6	30.3	15.0	16.2	41.0	46.6	53.2	74.4	114.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
White, both sexes										
2006 4/.....	858.1	576.0	25.5	12.8	15.6	61.1	94.8	93.6	103.3	137.8
2005 4/.....	873.7	579.1	27.0	13.3	16.7	62.5	91.5	89.3	101.7	138.3
2004 4/.....	863.2	574.0	27.0	13.4	17.4	64.4	88.3	87.3	98.3	140.6
2003 4/.....	890.1	591.3	28.5	13.7	17.8	64.7	89.4	86.1	101.9	145.5
2002.....	895.7	586.7	28.1	14.0	18.1	66.3	88.0	85.0	101.1	146.8
2001.....	895.1	570.9	30.7	14.4	17.9	64.2	87.1	87.0	102.7	151.2
2000.....	900.2	610.6	29.2	14.8	19.2	64.5	85.1	81.0	99.0	144.0
1999.....	901.4	606.1	30.8	15.3	19.2	65.3	81.6	80.8	101.4	144.5
White, male										
2006 4/.....	852.3	632.7	27.5	14.1	18.6	84.1	139.3	132.3	138.7	176.1
2005 4/.....	864.5	640.0	30.9	14.2	19.8	86.5	133.8	126.5	135.1	176.1
2004 4/.....	854.2	631.6	29.4	15.1	20.6	87.1	129.1	123.2	130.8	180.9
2003 4/.....	877.6	658.7	31.5	15.3	21.4	88.3	129.1	120.0	137.0	187.6
2002.....	884.0	650.9	31.5	15.4	21.2	90.5	128.9	119.7	136.0	188.7
2001.....	881.9	627.6	34.2	15.3	21.3	88.3	128.2	122.4	137.4	196.7
2000.....	887.8	667.6	32.6	16.5	23.1	88.1	124.6	113.9	133.6	188.2
1999.....	892.1	667.0	33.9	16.8	22.7	88.3	118.5	113.5	136.5	188.8
White, female										
2006 4/.....	863.9	516.5	23.5	11.4	12.5	36.9	46.6	52.2	65.9	98.1
2005 4/.....	882.8	515.3	22.9	12.3	13.4	37.2	45.8	49.4	66.6	99.0
2004 4/.....	871.9	513.6	24.4	11.6	14.2	40.4	44.3	48.8	64.3	98.9
2003 4/.....	902.3	520.7	25.5	12.1	14.0	39.6	46.7	50.0	65.2	102.2
2002.....	907.0	519.4	24.5	12.6	14.8	40.4	44.4	48.1	64.8	103.9
2001.....	907.9	511.5	27.1	13.3	14.4	38.5	43.2	49.6	66.6	104.9
2000.....	912.3	550.5	25.5	13.0	15.2	39.3	43.0	46.5	63.0	99.2
1999.....	910.4	542.0	27.5	13.8	15.4	40.8	42.3	46.7	65.0	99.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
Black, both sexes										
2006 4/.....	733.0	1,303.1	43.3	19.9	22.4	84.7	142.0	163.9	192.9	252.6
2005 4/.....	749.4	1,311.2	41.8	21.1	25.2	83.6	143.1	162.0	197.3	254.1
2004 4/.....	744.3	1,284.7	44.8	21.6	25.3	81.4	138.8	167.2	191.3	254.4
2003 4/.....	763.6	1,273.8	46.8	19.7	25.7	80.0	147.3	168.6	194.1	275.2
2002.....	768.4	1,263.6	47.1	21.3	27.5	82.0	146.8	172.1	200.7	284.0
2001.....	773.5	1,304.5	47.5	20.7	25.8	86.3	151.3	168.8	200.5	286.0
2000.....	781.1	1,426.1	49.9	21.8	26.6	87.4	155.0	168.4	206.9	283.1
1999.....	788.1	1,457.3	52.6	25.2	28.0	91.9	154.4	163.5	204.9	293.5
Black, male										
2006 4/.....	786.7	1,407.1	47.1	22.9	26.7	130.8	215.7	242.0	268.0	324.7
2005 4/.....	799.2	1,437.2	46.7	22.6	30.9	127.2	220.4	239.1	270.7	323.9
2004 4/.....	792.6	1,414.2	48.6	23.1	28.6	121.0	210.1	244.7	259.9	323.6
2003 4/.....	813.7	1,410.0	53.7	21.6	31.4	120.9	224.5	249.6	262.5	344.8
2002.....	816.7	1,351.5	54.4	24.1	33.4	121.7	227.6	255.7	272.9	366.5
2001.....	823.9	1,426.4	53.2	23.8	31.6	130.4	236.3	247.0	271.0	364.1
2000.....	834.1	1,567.6	54.5	24.5	32.1	130.1	240.6	246.8	275.0	361.6
1999.....	847.4	1,592.8	59.0	29.3	35.0	137.7	239.9	238.0	278.9	372.8
Black, female										
2006 4/.....	684.0	1,194.6	39.4	16.8	17.9	37.4	66.5	89.8	124.9	188.2
2005 4/.....	703.9	1,179.7	36.7	19.6	19.3	38.8	64.5	89.1	131.0	191.5
2004 4/.....	700.3	1,149.9	40.9	20.1	21.9	40.7	67.0	94.7	129.5	192.6
2003 4/.....	717.9	1,132.2	39.7	17.8	19.8	37.8	70.6	93.6	132.5	213.1
2002.....	724.4	1,172.0	39.5	18.5	21.3	41.0	68.0	95.7	135.7	210.6
2001.....	727.7	1,176.3	41.7	17.5	19.8	40.8	69.4	97.5	137.3	216.6
2000.....	733.0	1,279.8	45.3	19.0	21.0	43.7	73.9	97.3	145.6	213.4
1999.....	734.3	1,317.4	46.1	21.0	20.7	45.2	73.2	96.1	138.6	223.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Septicemia (A40-A41)										
All races 3/, both sexes										
2006.....	11.4	6.5	0.5	0.2	0.2	0.2	0.4	0.6	0.8	1.5
2005.....	11.5	7.4	0.5	0.2	0.2	0.3	0.4	0.6	0.9	1.5
2004.....	11.4	6.6	0.5	0.2	0.1	0.2	0.4	0.6	1.0	1.5
2003.....	11.7	6.9	0.5	0.2	0.2	0.3	0.5	0.6	0.9	1.5
2002.....	11.7	7.3	0.5	0.2	0.3	0.2	0.3	0.6	0.9	1.5
2001.....	11.3	7.7	0.7	0.1	0.2	0.3	0.4	0.5	0.9	1.4
2000.....	11.1	7.2	0.6	0.2	0.1	0.2	0.3	0.5	0.9	1.4
1999.....	11.0	7.5	0.6	0.2	0.1	0.2	0.5	0.5	0.8	1.4
All races 3/, male										
2006.....	10.5	6.8	0.5	0.2	0.3	0.2	0.4	0.6	0.9	1.5
2005.....	10.5	8.1	0.6	0.2	0.2	0.3	0.5	0.7	0.8	1.4
2004.....	10.4	8.0	0.5	*	*	0.2	0.4	0.5	1.1	1.5
2003.....	10.5	7.4	0.7	*	*	0.3	0.4	0.6	1.0	1.6
2002.....	10.6	7.7	0.5	0.3	0.2	0.2	0.4	0.7	1.0	1.5
2001.....	10.2	8.4	0.8	*	0.2	0.3	0.4	0.5	0.9	1.5
2000.....	9.8	7.1	0.7	0.2	*	0.2	0.4	0.6	0.9	1.4
1999.....	9.8	9.0	0.6	0.2	*	*	0.5	0.5	0.9	1.5
All races 3/, female										
2006.....	12.3	6.2	0.6	*	*	0.3	0.4	0.6	0.8	1.6
2005.....	12.5	6.5	0.5	*	0.2	0.3	0.4	0.6	1.0	1.6
2004.....	12.3	5.2	0.6	0.2	*	0.2	0.4	0.7	1.0	1.4
2003.....	12.9	6.5	0.4	0.2	*	0.2	0.5	0.7	0.8	1.4
2002.....	12.9	7.0	0.5	*	0.3	0.3	0.3	0.6	0.9	1.5
2001.....	12.4	7.0	0.6	*	*	0.3	0.3	0.5	0.9	1.2
2000.....	12.3	7.3	0.6	*	*	*	0.2	0.4	0.8	1.4
1999.....	12.2	5.8	0.5	0.2	*	*	0.4	0.5	0.7	1.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Septicemia (A40-A41)										
White, both sexes										
2006 4/.....	11.3	4.9	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.2
2005 4/.....	11.3	5.0	0.5	0.2	0.2	0.2	0.4	0.5	0.7	1.2
2004 4/.....	11.2	5.2	0.5	0.2	*	0.2	0.3	0.5	0.8	1.3
2003 4/.....	11.5	4.6	0.4	0.2	0.2	0.2	0.4	0.5	0.7	1.2
2002.....	11.5	6.1	0.5	0.2	0.2	0.2	0.3	0.5	0.8	1.2
2001.....	11.1	5.4	0.7	0.2	0.2	0.2	0.3	0.4	0.7	1.1
2000.....	10.8	5.7	0.6	0.2	0.1	0.2	0.2	0.4	0.6	1.1
1999.....	10.6	5.5	0.5	0.2	0.1	0.2	0.4	0.4	0.7	1.0
White, male										
2006 4/.....	10.4	5.1	0.4	*	0.3	*	0.3	0.5	0.7	1.3
2005 4/.....	10.2	5.9	0.5	*	*	*	0.4	0.6	0.6	1.1
2004 4/.....	10.2	6.3	0.5	*	*	*	0.3	0.4	0.8	1.4
2003 4/.....	10.2	4.6	0.6	*	*	0.3	0.4	0.4	0.8	1.3
2002.....	10.3	6.2	0.5	0.3	0.2	*	0.4	0.6	0.8	1.2
2001.....	9.9	6.2	0.8	*	*	0.3	0.3	0.4	0.6	1.3
2000.....	9.4	5.8	0.6	*	*	*	0.3	0.4	0.7	1.2
1999.....	9.3	6.9	0.5	*	*	*	0.5	0.5	0.8	1.2
White, female										
2006 4/.....	12.2	4.6	0.5	*	*	0.2	0.3	0.5	0.7	1.2
2005 4/.....	12.4	4.1	0.4	*	*	*	0.3	0.4	0.8	1.3
2004 4/.....	12.2	4.0	0.5	*	*	*	0.3	0.6	0.8	1.2
2003 4/.....	12.8	4.5	*	*	*	*	0.4	0.6	0.7	1.1
2002.....	12.8	6.0	0.4	*	*	0.3	*	0.4	0.8	1.2
2001.....	12.3	4.6	0.6	*	*	*	0.3	0.4	0.7	0.9
2000.....	12.1	5.6	0.5	*	*	*	*	0.4	0.5	1.0
1999.....	11.9	4.0	0.5	*	*	*	*	0.4	0.6	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Septicemia (A40-A41)

Black, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	15.4	15.0	0.8	*	*	*	0.9	1.4	1.8	3.8
2005 4/.....	15.9	19.8	0.8	*	*	0.6	0.8	1.5	2.3	3.6
2004 4/.....	15.6	14.5	*	*	*	*	1.0	1.3	2.5	2.7
2003 4/.....	16.3	18.6	0.9	*	*	*	0.7	1.6	2.2	3.6
2002.....	16.3	13.5	*	*	*	*	*	1.6	2.2	3.3
2001.....	15.8	19.3	0.9	*	*	*	0.7	1.1	2.5	3.7
2000.....	16.2	16.4	1.0	*	*	*	0.8	1.5	2.5	3.7
1999.....	16.1	17.7	*	*	*	*	1.0	1.4	1.7	3.7

Black, male

2006 4/.....	14.4	15.6	*	*	*	*	*	*	2.1	3.6
2005 4/.....	14.9	20.4	*	*	*	*	*	1.6	2.1	3.6
2004 4/.....	14.6	16.3	*	*	*	*	*	*	2.8	2.5
2003 4/.....	15.3	21.1	*	*	*	*	*	1.6	2.7	3.6
2002.....	15.0	14.8	*	*	*	*	*	*	2.5	3.7
2001.....	14.9	20.1	*	*	*	*	*	*	2.5	3.7
2000.....	14.9	16.0	*	*	*	*	*	2.1	2.5	3.5
1999.....	15.3	19.8	*	*	*	*	*	*	2.1	4.0

Black, female

2006 4/.....	16.4	14.4	*	*	*	*	*	1.5	1.5	4.0
2005 4/.....	16.8	19.1	*	*	*	*	*	1.4	2.6	3.7
2004 4/.....	16.4	12.7	*	*	*	*	*	1.4	2.1	2.9
2003 4/.....	17.2	16.1	*	*	*	*	*	1.6	1.6	3.5
2002.....	17.4	12.1	*	*	*	*	*	1.9	2.0	3.1
2001.....	16.7	18.6	*	*	*	*	*	*	2.5	3.7
2000.....	17.4	16.9	*	*	*	*	*	*	2.5	3.9
1999.....	16.9	15.4	*	*	*	*	*	1.6	1.4	3.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Viral hepatitis (B15-B19)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	2.4	*	*	*	*	*	*	*	0.2	0.6
2005.....	1.9	*	*	*	*	*	*	0.1	0.2	0.6
2004.....	1.8	*	*	*	*	*	*	0.1	0.2	0.7
2003.....	1.9	*	*	*	*	*	*	0.1	0.2	0.8
2002.....	2.0	*	*	*	*	*	*	0.2	0.3	1.0
2001.....	2.0	*	*	*	*	*	*	*	0.3	0.9
2000.....	1.9	*	*	*	*	*	*	0.1	0.3	1.0
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9

All races 3/, male

2006.....	3.2	*	*	*	*	*	*	*	0.3	0.8
2005.....	2.5	*	*	*	*	*	*	*	*	0.8
2004.....	2.4	*	*	*	*	*	*	*	0.2	0.8
2003.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2002.....	2.7	*	*	*	*	*	*	0.2	0.4	1.4
2001.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.3
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.2

All races 3/, female

2006.....	1.6	*	*	*	*	*	*	*	*	0.4
2005.....	1.3	*	*	*	*	*	*	*	*	0.4
2004.....	1.3	*	*	*	*	*	*	*	*	0.5
2003.....	1.3	*	*	*	*	*	*	*	*	0.4
2002.....	1.4	*	*	*	*	*	*	*	*	0.6
2001.....	1.4	*	*	*	*	*	*	*	0.2	0.6
2000.....	1.4	*	*	*	*	*	*	*	0.2	0.6
1999.....	1.3	*	*	*	*	*	*	*	0.2	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral hepatitis (B15-B19)										
White, both sexes										
2006 4/.....	2.4	*	*	*	*	*	*	*	0.2	0.5
2005 4/.....	1.8	*	*	*	*	*	*	*	0.1	0.6
2004 4/.....	1.8	*	*	*	*	*	*	*	0.2	0.6
2003 4/.....	1.8	*	*	*	*	*	*	*	0.2	0.8
2002.....	2.0	*	*	*	*	*	*	0.1	0.3	1.0
2001.....	1.9	*	*	*	*	*	*	*	0.2	0.9
2000.....	1.8	*	*	*	*	*	*	*	0.2	1.0
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9
White, male										
2006 4/.....	3.2	*	*	*	*	*	*	*	0.3	0.7
2005 4/.....	2.5	*	*	*	*	*	*	*	*	0.9
2004 4/.....	2.4	*	*	*	*	*	*	*	*	0.8
2003 4/.....	2.4	*	*	*	*	*	*	*	0.2	1.2
2002.....	2.6	*	*	*	*	*	*	*	0.4	1.5
2001.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.3
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.3
White, female										
2006 4/.....	1.6	*	*	*	*	*	*	*	*	0.4
2005 4/.....	1.2	*	*	*	*	*	*	*	*	0.3
2004 4/.....	1.2	*	*	*	*	*	*	*	*	0.5
2003 4/.....	1.2	*	*	*	*	*	*	*	*	0.5
2002.....	1.3	*	*	*	*	*	*	*	*	0.5
2001.....	1.3	*	*	*	*	*	*	*	*	0.6
2000.....	1.3	*	*	*	*	*	*	*	*	0.6
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Human immunodeficiency virus (HIV) disease (B20-B24)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	4.0	*	*	*	*	0.2	0.8	2.0	3.9	7.5
2005.....	4.2	*	*	*	*	0.1	0.6	2.1	4.5	8.1
2004.....	4.4	*	*	*	*	0.1	0.8	2.2	5.1	9.5
2003.....	4.7	*	*	*	0.1	0.2	0.7	2.3	5.5	11.2
2002.....	4.9	*	*	*	0.1	0.2	0.7	2.6	6.4	11.9
2001.....	5.0	*	*	*	0.1	0.2	0.9	3.1	7.3	12.3
2000.....	5.1	*	*	*	0.1	0.2	0.8	3.3	8.7	12.8
1999.....	5.3	*	0.2	0.2	0.1	0.2	0.9	3.9	9.5	13.0

All races 3/, male

2006.....	5.9	*	*	*	*	0.2	0.9	2.4	4.7	10.1
2005.....	6.3	*	*	*	*	*	0.7	2.4	5.5	11.2
2004.....	6.6	*	*	*	*	*	0.9	2.6	6.4	13.3
2003.....	7.1	*	*	*	*	*	0.8	2.8	7.3	15.9
2002.....	7.4	*	*	*	*	*	0.7	3.0	8.5	17.7
2001.....	7.6	*	*	*	*	0.2	0.8	3.7	10.2	18.4
2000.....	7.9	*	*	*	*	*	0.8	3.8	11.9	19.2
1999.....	8.2	*	*	0.2	*	*	0.8	5.1	13.6	19.5

All races 3/, female

2006.....	2.2	*	*	*	*	*	0.7	1.6	3.0	4.9
2005.....	2.2	*	*	*	*	*	0.5	1.7	3.5	4.9
2004.....	2.4	*	*	*	*	*	0.7	1.8	3.7	5.6
2003.....	2.4	*	*	*	*	*	0.7	1.8	3.7	6.5
2002.....	2.5	*	*	*	*	0.2	0.6	2.3	4.2	6.2
2001.....	2.5	*	*	*	*	0.2	1.0	2.5	4.4	6.3
2000.....	2.5	*	*	*	*	*	0.7	2.9	5.5	6.4
1999.....	2.5	*	*	*	*	*	1.0	2.7	5.4	6.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Human immunodeficiency virus (HIV) disease (B20-B24)

White, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	2.1	*	*	*	*	*	0.3	0.8	1.8	4.0
2005 4/.....	2.2	*	*	*	*	*	0.3	0.8	2.1	4.2
2004 4/.....	2.4	*	*	*	*	*	0.2	0.8	2.3	5.4
2003 4/.....	2.5	*	*	*	*	*	0.3	0.9	2.9	6.1
2002.....	2.6	*	*	*	*	*	0.3	1.2	3.0	6.8
2001.....	2.7	*	*	*	*	*	0.3	1.3	3.8	7.1
2000.....	2.8	*	*	*	*	*	0.3	1.5	4.5	7.6
1999.....	3.0	*	*	*	*	*	0.4	2.0	5.4	7.9

White, male

2006 4/.....	3.5	*	*	*	*	*	0.3	1.1	2.7	6.2
2005 4/.....	3.7	*	*	*	*	*	0.4	1.1	3.0	6.8
2004 4/.....	3.9	*	*	*	*	*	0.4	1.2	3.6	8.4
2003 4/.....	4.2	*	*	*	*	*	0.3	1.4	4.5	9.7
2002.....	4.3	*	*	*	*	*	0.3	1.8	4.7	11.1
2001.....	4.5	*	*	*	*	*	0.4	1.8	5.8	11.6
2000.....	4.7	*	*	*	*	*	0.4	2.0	7.1	12.5
1999.....	5.0	*	*	*	*	*	0.4	2.8	8.6	13.0

White, female

2006 4/.....	0.7	*	*	*	*	*	0.3	0.4	0.9	1.8
2005 4/.....	0.8	*	*	*	*	*	*	0.5	1.2	1.6
2004 4/.....	0.8	*	*	*	*	*	*	0.4	1.0	2.4
2003 4/.....	0.9	*	*	*	*	*	*	0.5	1.3	2.4
2002.....	0.9	*	*	*	*	*	*	0.7	1.3	2.3
2001.....	0.9	*	*	*	*	*	0.3	0.8	1.6	2.4
2000.....	1.0	*	*	*	*	*	*	1.0	1.9	2.6
1999.....	1.0	*	*	*	*	*	0.3	1.0	2.1	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease (B20-B24)										
Black, both sexes										
2006 4/.....	17.3	*	*	*	*	0.9	3.6	9.9	17.7	31.8
2005 4/.....	18.0	*	*	*	*	0.6	2.7	9.7	20.4	34.3
2004 4/.....	18.8	*	*	*	*	0.7	3.7	10.8	23.2	37.4
2003 4/.....	19.6	*	*	*	*	0.6	3.2	11.2	23.2	46.0
2002.....	20.8	*	*	*	*	0.8	3.3	11.7	29.1	48.2
2001.....	21.1	*	*	*	*	1.1	4.4	14.5	31.5	48.8
2000.....	21.4	*	*	*	*	0.7	3.7	15.2	37.1	49.8
1999.....	21.8	*	*	0.7	*	0.7	4.1	16.6	37.8	49.9
Black, male										
2006 4/.....	23.5	*	*	*	*	*	4.1	11.3	20.0	40.2
2005 4/.....	25.1	*	*	*	*	*	2.9	11.0	23.5	44.5
2004 4/.....	25.9	*	*	*	*	*	3.8	11.9	26.5	50.1
2003 4/.....	27.7	*	*	*	*	*	3.2	12.7	28.6	61.9
2002.....	29.5	*	*	*	*	*	3.3	11.7	36.6	68.4
2001.....	30.1	*	*	*	*	*	3.5	16.8	42.3	69.2
2000.....	31.0	*	*	*	*	*	3.6	16.4	47.9	71.2
1999.....	31.9	*	*	*	*	*	3.5	21.1	50.6	69.9
Black, female										
2006 4/.....	11.7	*	*	*	*	*	3.0	8.7	15.6	24.3
2005 4/.....	11.5	*	*	*	*	*	2.5	8.5	17.6	25.1
2004 4/.....	12.4	*	*	*	*	*	3.7	9.8	20.3	25.9
2003 4/.....	12.3	*	*	*	*	*	3.1	9.8	18.3	31.9
2002.....	12.8	*	*	*	*	*	3.3	11.7	22.2	30.3
2001.....	12.9	*	*	*	*	*	5.2	12.4	21.8	30.7
2000.....	12.8	*	*	*	*	*	3.8	14.1	27.4	30.8
1999.....	12.6	*	*	*	*	*	4.7	12.5	26.3	32.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All Under 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39
ages 1/ year 2/ years years years years years years years years

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races 3/, both sexes

2006.....	2.0	4.0	0.4	0.2	0.2	0.2	0.2	0.3	0.4	0.5
2005.....	2.6	4.1	0.4	0.2	0.1	0.2	0.2	0.2	0.4	0.7
2004.....	2.6	4.1	0.4	0.1	0.1	0.2	0.3	0.3	0.4	0.7
2003.....	2.6	4.3	0.5	0.1	0.1	0.2	0.2	0.3	0.4	0.7
2002.....	2.3	5.6	0.4	0.2	0.2	0.2	0.3	0.4	0.5	0.7
2001.....	2.0	4.7	0.4	0.2	0.1	0.2	0.2	0.3	0.4	0.7
2000.....	2.0	4.4	0.5	0.1	0.2	0.2	0.3	0.3	0.4	0.7
1999.....	2.6	5.3	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.0

All races 3/, male

2006.....	2.1	4.5	0.5	*	*	0.2	0.2	0.3	0.3	0.6
2005.....	2.9	4.8	0.5	*	*	0.2	0.2	0.3	0.5	0.9
2004.....	2.8	4.8	0.4	*	*	0.2	0.4	0.2	0.6	0.8
2003.....	2.8	4.2	0.5	*	*	0.3	0.2	0.3	0.5	0.9
2002.....	2.6	6.1	0.5	*	0.2	0.2	0.3	0.4	0.5	0.9
2001.....	2.1	4.9	0.5	*	*	0.2	0.3	0.3	0.5	0.8
2000.....	2.1	4.8	0.5	*	*	0.2	0.3	0.3	0.4	0.9
1999.....	2.7	6.1	0.6	0.2	0.2	0.2	0.3	0.5	0.8	1.3

All races 3/, female

2006.....	1.9	3.6	0.3	*	0.2	0.2	0.2	0.2	0.4	0.3
2005.....	2.3	3.4	0.3	0.2	*	0.2	0.2	0.2	0.3	0.5
2004.....	2.3	3.4	0.4	*	*	0.2	*	0.4	0.3	0.6
2003.....	2.3	4.5	0.5	*	*	*	*	0.3	0.3	0.6
2002.....	2.1	5.0	0.3	0.3	*	0.2	0.2	0.3	0.4	0.6
2001.....	1.9	4.6	0.4	0.2	*	*	*	0.3	0.4	0.6
2000.....	1.9	3.9	0.4	*	*	*	0.2	0.3	0.4	0.5
1999.....	2.5	4.5	0.4	0.2	*	0.2	0.2	0.5	0.6	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

White, both sexes

2006 4/.....	2.0	2.8	0.4	0.2	0.2	0.2	0.2	0.2	0.3	0.4
2005 4/.....	2.7	3.7	0.4	0.2	0.2	0.2	0.2	0.2	0.3	0.6
2004 4/.....	2.6	3.7	0.4	*	*	0.2	0.3	0.2	0.3	0.7
2003 4/.....	2.6	3.5	0.4	0.1	*	0.2	0.2	0.2	0.4	0.6
2002.....	2.4	5.0	0.4	0.2	0.2	0.2	0.2	0.3	0.4	0.6
2001.....	2.0	3.8	0.4	0.2	0.1	0.2	0.2	0.3	0.3	0.6
2000.....	2.0	3.4	0.4	0.2	0.2	0.2	0.3	0.2	0.3	0.6
1999.....	2.5	4.5	0.5	0.2	0.2	0.2	0.2	0.4	0.5	0.8

White, male

2006 4/.....	2.1	3.3	0.5	*	*	*	0.2	0.3	0.3	0.5
2005 4/.....	3.0	4.2	0.4	*	*	*	*	0.2	0.4	0.8
2004 4/.....	2.9	4.2	0.4	*	*	*	0.3	*	0.4	0.7
2003 4/.....	2.8	3.2	0.4	*	*	0.3	*	0.3	0.4	0.8
2002.....	2.6	5.3	0.5	*	0.2	*	*	0.3	0.5	0.7
2001.....	2.1	3.9	0.4	*	*	*	*	0.3	0.4	0.7
2000.....	2.1	3.9	0.4	*	*	*	0.3	*	0.3	0.7
1999.....	2.6	5.6	0.6	*	*	*	*	0.3	0.6	1.2

White, female

2006 4/.....	2.0	2.2	*	*	*	*	*	*	*	0.2
2005 4/.....	2.4	3.1	*	*	*	*	*	*	*	0.4
2004 4/.....	2.4	3.0	0.3	*	*	*	*	0.3	*	0.6
2003 4/.....	2.4	3.9	0.5	*	*	*	*	*	0.3	0.5
2002.....	2.2	4.6	*	*	*	*	*	0.3	0.4	0.4
2001.....	1.9	3.8	*	*	*	*	*	*	0.2	0.5
2000.....	1.9	2.9	0.4	*	*	*	*	*	0.3	0.4
1999.....	2.5	3.4	*	*	*	*	*	0.4	0.5	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Black, both sexes

2006 4/.....	2.1	9.9	*	*	*	*	*	*	1.0	1.1
2005 4/.....	2.5	6.5	*	*	*	*	*	*	1.1	1.2
2004 4/.....	2.5	6.7	*	*	*	*	*	*	0.9	1.1
2003 4/.....	2.5	8.0	1.0	*	*	*	*	*	0.7	1.4
2002.....	2.5	8.7	*	*	*	*	*	*	1.0	1.8
2001.....	2.2	9.5	*	*	*	*	*	*	1.0	1.4
2000.....	2.3	9.9	*	*	*	*	*	*	1.2	1.6
1999.....	3.5	9.9	0.9	*	*	*	*	1.1	2.1	2.5

Black, male

2006 4/.....	2.3	10.1	*	*	*	*	*	*	*	*
2005 4/.....	2.8	8.0	*	*	*	*	*	*	*	1.5
2004 4/.....	2.9	7.4	*	*	*	*	*	*	*	*
2003 4/.....	2.9	8.9	*	*	*	*	*	*	*	1.6
2002.....	2.9	9.6	*	*	*	*	*	*	*	2.1
2001.....	2.4	10.8	*	*	*	*	*	*	*	1.7
2000.....	2.5	9.9	*	*	*	*	*	*	*	2.2
1999.....	3.6	10.1	*	*	*	*	*	*	2.6	2.7

Black, female

2006 4/.....	2.0	9.6	*	*	*	*	*	*	*	*
2005 4/.....	2.2	*	*	*	*	*	*	*	*	*
2004 4/.....	2.2	*	*	*	*	*	*	*	*	*
2003 4/.....	2.2	7.1	*	*	*	*	*	*	*	*
2002.....	2.1	7.9	*	*	*	*	*	*	*	1.6
2001.....	2.0	8.2	*	*	*	*	*	*	*	*
2000.....	2.1	9.9	*	*	*	*	*	*	*	*
1999.....	3.4	9.7	*	*	*	*	*	*	1.6	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
All races 3/, both sexes										
2006.....	187.0	1.8	2.3	2.3	2.2	3.2	4.6	6.7	11.5	20.8
2005.....	188.7	1.8	2.3	2.5	2.5	3.5	4.7	6.7	11.3	20.8
2004.....	188.6	1.8	2.5	2.7	2.3	3.5	4.7	6.6	11.5	21.5
2003.....	191.5	1.9	2.5	2.6	2.6	3.4	4.6	6.8	11.8	23.0
2002.....	193.2	1.8	2.6	2.7	2.5	3.5	5.0	6.6	12.5	23.5
2001.....	194.4	1.6	2.7	2.4	2.5	3.6	4.9	7.3	12.6	24.6
2000.....	196.5	2.4	2.7	2.4	2.6	3.7	5.1	7.0	12.5	24.5
1999.....	197.0	1.8	2.7	2.5	2.5	3.7	5.3	7.3	12.5	25.2
All races 3/, male										
2006.....	196.6	1.8	2.5	2.6	2.4	3.8	5.4	7.0	10.3	17.2
2005.....	198.9	2.1	2.6	2.6	2.8	4.1	5.5	7.3	10.4	17.4
2004.....	198.4	1.8	2.6	3.0	2.5	4.1	5.5	6.8	10.4	18.1
2003.....	201.3	1.7	2.8	2.9	2.7	4.2	5.1	6.8	10.9	19.3
2002.....	203.8	2.0	2.7	2.9	2.8	4.0	5.7	7.0	11.3	19.8
2001.....	205.3	1.5	2.9	2.4	2.6	4.3	5.7	7.2	11.2	21.0
2000.....	207.2	2.6	3.0	2.6	2.9	4.3	5.9	7.1	11.3	21.2
1999.....	208.9	1.9	2.9	2.6	2.7	4.3	6.1	7.6	11.6	21.6
All races 3/, female										
2006.....	177.6	1.8	2.1	2.1	2.0	2.5	3.7	6.4	12.7	24.4
2005.....	178.8	1.5	2.0	2.3	2.2	2.8	3.9	6.0	12.2	24.3
2004.....	179.1	1.9	2.4	2.3	2.2	2.9	3.8	6.3	12.6	24.9
2003.....	182.0	2.1	2.1	2.3	2.5	2.5	4.2	6.8	12.6	26.7
2002.....	183.0	1.6	2.4	2.5	2.2	3.0	4.2	6.2	13.7	27.2
2001.....	183.9	1.8	2.5	2.4	2.3	2.9	4.1	7.4	14.0	28.1
2000.....	186.2	2.3	2.5	2.2	2.2	3.0	4.2	7.0	13.7	27.8
1999.....	185.6	1.7	2.5	2.3	2.3	3.0	4.4	7.0	13.4	28.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
White, both sexes										
2006 4/.....	199.3	2.0	2.3	2.4	2.2	3.2	4.6	6.6	11.2	20.1
2005 4/.....	200.8	2.0	2.3	2.5	2.5	3.6	4.7	6.3	10.8	20.1
2004 4/.....	200.7	2.0	2.5	2.7	2.3	3.6	4.5	6.4	11.1	20.8
2003 4/.....	203.7	2.1	2.6	2.8	2.6	3.5	4.7	6.7	11.3	22.2
2002.....	205.5	1.8	2.5	2.8	2.5	3.6	5.2	6.5	12.1	22.7
2001.....	206.4	1.7	3.0	2.5	2.4	3.7	5.0	7.2	12.2	23.7
2000.....	208.6	2.5	2.8	2.5	2.6	3.6	5.1	7.0	12.1	23.3
1999.....	208.7	1.8	2.9	2.5	2.4	3.6	5.2	7.3	12.1	24.0
White, male										
2006 4/.....	208.7	2.0	2.5	2.6	2.4	3.9	5.5	7.1	10.2	16.9
2005 4/.....	210.6	2.3	2.6	2.6	2.8	4.2	5.4	7.0	10.0	17.1
2004 4/.....	209.9	1.9	2.5	3.1	2.5	4.2	5.3	6.8	10.4	18.0
2003 4/.....	213.1	1.8	2.9	3.1	2.7	4.3	5.1	7.0	10.6	19.3
2002.....	215.5	2.2	2.7	2.9	2.7	4.2	5.9	6.9	11.1	19.2
2001.....	216.4	1.5	3.1	2.4	2.4	4.4	5.8	7.2	11.2	20.5
2000.....	218.1	2.8	3.1	2.6	2.9	4.2	5.9	7.1	11.2	20.5
1999.....	219.5	2.0	3.2	2.7	2.6	4.1	5.9	7.6	11.3	21.1
White, female										
2006 4/.....	190.1	2.0	2.0	2.1	2.0	2.5	3.6	6.1	12.4	23.4
2005 4/.....	191.1	1.7	2.0	2.4	2.3	2.9	3.9	5.5	11.7	23.3
2004 4/.....	191.7	2.1	2.5	2.3	2.1	2.9	3.7	6.0	11.9	23.7
2003 4/.....	194.6	2.3	2.3	2.4	2.5	2.6	4.2	6.5	12.0	25.2
2002.....	195.8	1.4	2.3	2.6	2.4	3.0	4.4	6.1	13.2	26.3
2001.....	196.7	2.0	2.8	2.5	2.4	2.9	4.1	7.2	13.3	27.1
2000.....	199.4	2.3	2.4	2.3	2.3	3.0	4.2	7.0	13.0	26.3
1999.....	198.2	1.5	2.7	2.2	2.2	3.0	4.3	7.0	12.9	27.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
Black, both sexes										
2006 4/.....	159.5	*	2.5	2.3	2.1	3.1	4.9	8.0	15.0	28.1
2005 4/.....	161.7	*	2.4	2.6	2.6	3.0	5.1	9.2	15.6	28.1
2004 4/.....	161.9	*	2.5	2.5	2.6	3.8	5.9	8.9	15.1	28.4
2003 4/.....	164.3	*	2.2	2.3	2.8	2.9	4.9	8.5	15.7	31.0
2002.....	165.9	*	2.9	2.6	2.5	3.6	4.5	8.0	16.3	31.7
2001.....	167.1	*	1.9	2.4	2.8	3.6	5.1	9.1	16.4	32.7
2000.....	169.3	*	2.8	2.1	2.3	4.4	5.9	8.0	15.8	34.5
1999.....	171.3	*	2.0	2.4	3.0	4.5	6.6	8.6	15.8	35.8
Black, male										
2006 4/.....	172.3	*	2.6	2.5	2.4	3.7	5.4	7.5	12.8	21.9
2005 4/.....	175.4	*	2.8	2.9	3.2	3.4	6.1	9.3	14.7	21.7
2004 4/.....	176.2	*	2.8	2.5	2.4	4.1	6.8	7.9	12.2	19.7
2003 4/.....	178.3	*	2.4	2.4	3.1	3.5	5.6	7.0	13.5	21.7
2002.....	181.5	*	2.6	3.3	2.9	4.1	5.0	8.2	14.1	25.6
2001.....	184.5	*	2.4	2.5	3.4	4.3	5.9	8.5	12.4	26.8
2000.....	188.5	*	2.7	2.4	2.7	5.2	7.1	7.3	12.9	27.6
1999.....	191.0	*	2.1	2.3	3.3	5.6	7.3	8.8	14.0	27.8
Black, female										
2006 4/.....	147.7	*	2.3	2.0	1.8	2.4	4.4	8.5	16.9	33.6
2005 4/.....	149.1	*	2.1	2.3	2.0	2.7	4.1	9.0	16.3	33.8
2004 4/.....	148.9	*	2.2	2.5	2.7	3.5	5.0	9.9	17.8	36.2
2003 4/.....	151.4	*	2.1	2.1	2.5	2.3	4.3	10.0	17.7	39.2
2002.....	151.7	*	3.2	1.9	2.1	3.1	4.0	7.9	18.3	37.1
2001.....	151.3	*	*	2.2	2.2	2.9	4.3	9.7	20.0	37.9
2000.....	151.8	*	2.9	1.8	2.0	3.5	4.7	8.5	18.3	40.5
1999.....	153.3	*	1.9	2.6	2.7	3.4	5.9	8.3	17.5	43.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of stomach (C16)										
All races 3/, both sexes										
2006.....	3.8	*	*	*	*	*	*	0.2	0.4	0.8
2005.....	3.9	*	*	*	*	*	*	0.1	0.4	0.7
2004.....	4.0	*	*	*	*	*	*	0.1	0.5	0.7
2003.....	4.2	*	*	*	*	*	*	0.2	0.4	0.8
2002.....	4.2	*	*	*	*	*	0.1	0.3	0.4	0.7
2001.....	4.3	*	*	*	*	*	*	0.2	0.4	0.8
2000.....	4.5	*	*	*	*	*	*	0.2	0.4	0.7
1999.....	4.6	*	*	*	*	*	*	0.2	0.4	0.8
All races 3/, male										
2006.....	4.5	*	*	*	*	*	*	*	0.4	0.7
2005.....	4.6	*	*	*	*	*	*	*	0.4	0.8
2004.....	4.8	*	*	*	*	*	*	*	0.4	0.7
2003.....	4.9	*	*	*	*	*	*	*	0.4	0.8
2002.....	5.0	*	*	*	*	*	*	0.3	0.5	0.7
2001.....	5.1	*	*	*	*	*	*	0.2	0.5	0.9
2000.....	5.3	*	*	*	*	*	*	0.2	0.5	0.9
1999.....	5.5	*	*	*	*	*	*	0.2	0.5	0.9
All races 3/, female										
2006.....	3.1	*	*	*	*	*	*	0.2	0.4	0.9
2005.....	3.2	*	*	*	*	*	*	*	0.4	0.5
2004.....	3.3	*	*	*	*	*	*	*	0.5	0.6
2003.....	3.4	*	*	*	*	*	*	*	0.4	0.8
2002.....	3.5	*	*	*	*	*	*	*	0.4	0.7
2001.....	3.5	*	*	*	*	*	*	*	0.4	0.7
2000.....	3.7	*	*	*	*	*	*	*	0.4	0.6
1999.....	3.7	*	*	*	*	*	*	0.3	0.3	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	17.9	*	*	*	*	*	0.2	0.5	1.1	2.0
2005.....	18.0	*	*	*	*	*	0.2	0.4	1.1	1.9
2004.....	18.3	*	*	*	*	*	0.2	0.4	0.9	2.0
2003.....	19.2	*	*	*	*	*	0.2	0.5	0.9	2.1
2002.....	19.7	*	*	*	*	*	0.2	0.5	1.0	2.0
2001.....	20.0	*	*	*	*	*	0.2	0.4	1.1	2.3
2000.....	20.4	*	*	*	*	*	0.1	0.4	0.9	2.1
1999.....	20.5	*	*	*	*	*	0.2	0.4	1.0	1.9

All races 3/, male

2006.....	18.3	*	*	*	*	*	*	0.5	1.1	2.1
2005.....	18.4	*	*	*	*	*	0.2	0.5	1.1	2.1
2004.....	18.6	*	*	*	*	*	0.2	0.5	1.0	2.1
2003.....	19.6	*	*	*	*	*	*	0.6	0.9	2.2
2002.....	20.1	*	*	*	*	*	0.3	0.5	1.1	2.1
2001.....	20.2	*	*	*	*	*	*	0.4	1.1	2.6
2000.....	20.6	*	*	*	*	*	*	0.4	1.1	2.4
1999.....	20.6	*	*	*	*	*	0.2	0.6	1.0	2.0

All races 3/, female

2006.....	17.5	*	*	*	*	*	0.2	0.5	1.1	1.9
2005.....	17.6	*	*	*	*	*	*	0.4	1.0	1.7
2004.....	18.0	*	*	*	*	*	*	0.4	0.9	2.0
2003.....	18.9	*	*	*	*	*	*	0.4	0.9	2.0
2002.....	19.2	*	*	*	*	*	*	0.4	0.9	1.9
2001.....	19.8	*	*	*	*	*	*	0.5	1.0	1.9
2000.....	20.2	*	*	*	*	*	*	0.4	0.8	1.8
1999.....	20.3	*	*	*	*	*	*	0.3	0.9	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of colon, rectum and anus (C18-C21)										
White, both sexes										
2006 4/.....	18.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2005 4/.....	18.7	*	*	*	*	*	0.2	0.4	1.0	1.8
2004 4/.....	19.2	*	*	*	*	*	0.2	0.4	0.9	1.9
2003 4/.....	20.2	*	*	*	*	*	0.1	0.5	0.9	1.9
2002.....	20.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2001.....	21.0	*	*	*	*	*	0.2	0.4	1.0	2.1
2000.....	21.5	*	*	*	*	*	*	0.4	0.9	1.9
1999.....	21.6	*	*	*	*	*	0.2	0.4	0.9	1.8
White, male										
2006 4/.....	19.0	*	*	*	*	*	*	0.5	1.0	2.0
2005 4/.....	19.2	*	*	*	*	*	*	0.5	1.0	2.1
2004 4/.....	19.5	*	*	*	*	*	*	0.5	0.9	2.1
2003 4/.....	20.5	*	*	*	*	*	*	0.6	0.9	2.0
2002.....	21.1	*	*	*	*	*	0.2	0.5	1.0	2.0
2001.....	21.2	*	*	*	*	*	*	0.4	1.1	2.5
2000.....	21.6	*	*	*	*	*	*	0.4	1.0	2.2
1999.....	21.8	*	*	*	*	*	*	0.5	0.9	1.9
White, female										
2006 4/.....	18.4	*	*	*	*	*	*	0.5	0.9	1.8
2005 4/.....	18.3	*	*	*	*	*	*	0.3	1.1	1.5
2004 4/.....	18.9	*	*	*	*	*	*	0.4	0.8	1.7
2003 4/.....	19.8	*	*	*	*	*	*	0.4	0.9	1.8
2002.....	20.3	*	*	*	*	*	*	0.4	0.9	1.9
2001.....	20.7	*	*	*	*	*	*	0.4	0.9	1.8
2000.....	21.4	*	*	*	*	*	*	0.3	0.7	1.6
1999.....	21.4	*	*	*	*	*	*	0.3	0.9	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of pancreas (C25)										
All races 3/, both sexes										
2006.....	11.2	*	*	*	*	*	*	*	0.2	0.6
2005.....	11.1	*	*	*	*	*	*	0.1	0.2	0.6
2004.....	10.8	*	*	*	*	*	*	*	0.2	0.6
2003.....	10.6	*	*	*	*	*	*	0.1	0.2	0.7
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.7
2001.....	10.5	*	*	*	*	*	*	*	0.2	0.8
2000.....	10.4	*	*	*	*	*	*	*	0.2	0.6
1999.....	10.4	*	*	*	*	*	*	*	0.2	0.8
All races 3/, male										
2006.....	11.2	*	*	*	*	*	*	*	*	0.8
2005.....	11.1	*	*	*	*	*	*	*	0.2	0.8
2004.....	10.9	*	*	*	*	*	*	*	0.3	0.7
2003.....	10.5	*	*	*	*	*	*	*	0.2	0.9
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.8
2001.....	10.3	*	*	*	*	*	*	*	0.2	0.9
2000.....	10.3	*	*	*	*	*	*	*	0.2	0.8
1999.....	10.4	*	*	*	*	*	*	*	0.3	0.9
All races 3/, female										
2006.....	11.1	*	*	*	*	*	*	*	*	0.3
2005.....	11.0	*	*	*	*	*	*	*	*	0.5
2004.....	10.7	*	*	*	*	*	*	*	*	0.5
2003.....	10.6	*	*	*	*	*	*	*	*	0.5
2002.....	10.5	*	*	*	*	*	*	*	*	0.6
2001.....	10.6	*	*	*	*	*	*	*	*	0.6
2000.....	10.5	*	*	*	*	*	*	*	*	0.5
1999.....	10.5	*	*	*	*	*	*	*	0.2	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
All races 3/, both sexes										
2006.....	53.0	*	*	*	*	*	*	0.2	0.6	2.0
2005.....	53.7	*	*	*	*	*	*	0.2	0.5	2.1
2004.....	53.8	*	*	*	*	*	*	0.2	0.5	2.2
2003.....	54.4	*	*	*	*	*	*	0.2	0.6	2.4
2002.....	54.7	*	*	*	*	*	*	0.2	0.6	2.7
2001.....	54.8	*	*	*	*	*	*	0.2	0.6	2.9
2000.....	55.3	*	*	*	*	*	*	0.2	0.8	2.9
1999.....	54.5	*	*	*	*	*	*	0.2	0.7	2.9
All races 3/, male										
2006.....	60.5	*	*	*	*	*	*	0.2	0.6	2.0
2005.....	61.8	*	*	*	*	*	*	*	0.6	2.2
2004.....	62.0	*	*	*	*	*	*	0.3	0.6	2.2
2003.....	62.9	*	*	*	*	*	*	*	0.7	2.4
2002.....	63.7	*	*	*	*	*	*	*	0.7	2.6
2001.....	64.7	*	*	*	*	*	*	*	0.6	2.8
2000.....	65.5	*	*	*	*	*	*	0.3	0.7	3.3
1999.....	65.4	*	*	*	*	*	*	0.2	0.7	2.8
All races 3/, female										
2006.....	45.7	*	*	*	*	*	*	*	0.5	2.0
2005.....	45.9	*	*	*	*	*	*	0.2	0.4	2.0
2004.....	45.9	*	*	*	*	*	*	*	0.5	2.2
2003.....	46.1	*	*	*	*	*	*	0.2	0.5	2.5
2002.....	46.0	*	*	*	*	*	*	*	0.6	2.7
2001.....	45.3	*	*	*	*	*	*	*	0.6	2.9
2000.....	45.4	*	*	*	*	*	*	*	0.8	2.6
1999.....	44.1	*	*	*	*	*	*	*	0.6	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of trachea, bronchus and lung (C33-C34)										
White, both sexes										
2006 4/.....	57.3	*	*	*	*	*	*	0.2	0.5	2.0
2005 4/.....	58.1	*	*	*	*	*	*	0.2	0.4	2.1
2004 4/.....	58.1	*	*	*	*	*	*	0.2	0.5	2.2
2003 4/.....	58.7	*	*	*	*	*	*	0.2	0.6	2.4
2002.....	59.1	*	*	*	*	*	*	0.2	0.6	2.6
2001.....	59.1	*	*	*	*	*	*	0.2	0.6	2.9
2000.....	59.5	*	*	*	*	*	*	0.2	0.8	2.7
1999.....	58.5	*	*	*	*	*	*	0.2	0.7	2.7
White, male										
2006 4/.....	64.6	*	*	*	*	*	*	*	0.6	2.0
2005 4/.....	65.9	*	*	*	*	*	*	*	0.6	2.2
2004 4/.....	65.9	*	*	*	*	*	*	0.3	0.5	2.1
2003 4/.....	66.9	*	*	*	*	*	*	*	0.6	2.3
2002.....	67.7	*	*	*	*	*	*	*	0.7	2.5
2001.....	68.6	*	*	*	*	*	*	*	0.6	2.8
2000.....	69.4	*	*	*	*	*	*	0.3	0.7	3.0
1999.....	69.1	*	*	*	*	*	*	*	0.8	2.6
White, female										
2006 4/.....	50.2	*	*	*	*	*	*	*	0.5	2.0
2005 4/.....	50.4	*	*	*	*	*	*	*	0.3	2.0
2004 4/.....	50.4	*	*	*	*	*	*	*	0.4	2.3
2003 4/.....	50.7	*	*	*	*	*	*	*	0.6	2.4
2002.....	50.7	*	*	*	*	*	*	*	0.5	2.8
2001.....	49.8	*	*	*	*	*	*	*	0.6	3.0
2000.....	49.9	*	*	*	*	*	*	*	0.8	2.4
1999.....	48.2	*	*	*	*	*	*	*	0.6	2.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant melanoma of skin (C43)										
All races 3/, both sexes										
2006.....	2.8	*	*	*	*	*	0.2	0.3	0.7	0.9
2005.....	2.8	*	*	*	*	*	0.2	0.5	0.6	0.9
2004.....	2.7	*	*	*	*	*	0.1	0.3	0.6	0.9
2003.....	2.7	*	*	*	*	*	0.1	0.3	0.6	0.9
2002.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2001.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2000.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0
1999.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0
All races 3/, male										
2006.....	3.7	*	*	*	*	*	0.2	0.3	0.7	1.2
2005.....	3.6	*	*	*	*	*	*	0.6	0.8	0.9
2004.....	3.5	*	*	*	*	*	*	0.4	0.7	1.0
2003.....	3.5	*	*	*	*	*	*	0.4	0.8	1.1
2002.....	3.3	*	*	*	*	*	*	0.4	0.8	1.1
2001.....	3.4	*	*	*	*	*	*	0.4	0.6	1.2
2000.....	3.3	*	*	*	*	*	*	0.3	0.7	1.0
1999.....	3.3	*	*	*	*	*	*	0.4	0.7	1.3
All races 3/, female										
2006.....	2.0	*	*	*	*	*	*	*	0.6	0.6
2005.....	2.0	*	*	*	*	*	*	0.3	0.5	0.8
2004.....	1.9	*	*	*	*	*	*	0.3	0.5	0.8
2003.....	1.9	*	*	*	*	*	*	0.2	0.5	0.7
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.9
2001.....	1.9	*	*	*	*	*	*	0.3	0.5	0.7
2000.....	2.0	*	*	*	*	*	*	0.3	0.5	1.0
1999.....	1.9	*	*	*	*	*	*	0.2	0.6	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant melanoma of skin (C43)										
White, both sexes										
2006 4/.....	3.4	*	*	*	*	*	0.2	0.3	0.8	1.1
2005 4/.....	3.4	*	*	*	*	*	0.2	0.6	0.8	1.0
2004 4/.....	3.3	*	*	*	*	*	0.2	0.4	0.8	1.1
2003 4/.....	3.2	*	*	*	*	*	0.2	0.4	0.8	1.1
2002.....	3.1	*	*	*	*	*	0.2	0.5	0.7	1.2
2001.....	3.2	*	*	*	*	*	0.2	0.5	0.7	1.2
2000.....	3.1	*	*	*	*	*	0.2	0.4	0.8	1.2
1999.....	3.1	*	*	*	*	*	0.2	0.4	0.8	1.3
White, male										
2006 4/.....	4.5	*	*	*	*	*	0.2	0.4	0.9	1.4
2005 4/.....	4.4	*	*	*	*	*	*	0.7	0.9	1.1
2004 4/.....	4.3	*	*	*	*	*	*	0.5	0.9	1.3
2003 4/.....	4.2	*	*	*	*	*	*	0.5	0.9	1.3
2002.....	4.0	*	*	*	*	*	*	0.5	1.0	1.3
2001.....	4.1	*	*	*	*	*	*	0.5	0.7	1.4
2000.....	4.0	*	*	*	*	*	*	0.4	0.9	1.2
1999.....	4.0	*	*	*	*	*	*	0.5	0.8	1.6
White, female										
2006 4/.....	2.3	*	*	*	*	*	*	*	0.8	0.8
2005 4/.....	2.4	*	*	*	*	*	*	0.4	0.6	1.0
2004 4/.....	2.3	*	*	*	*	*	*	0.4	0.6	1.0
2003 4/.....	2.3	*	*	*	*	*	*	0.3	0.6	0.9
2002.....	2.3	*	*	*	*	*	*	0.4	0.5	1.1
2001.....	2.3	*	*	*	*	*	*	0.4	0.6	0.9
2000.....	2.3	*	*	*	*	*	*	0.3	0.6	1.2
1999.....	2.3	*	*	*	*	*	*	0.3	0.7	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Malignant neoplasm of breast (C50)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	13.8	*	*	*	*	*	*	0.4	1.5	3.8
2005.....	14.0	*	*	*	*	*	*	0.4	1.4	3.8
2004.....	14.1	*	*	*	*	*	*	0.4	1.6	3.9
2003.....	14.4	*	*	*	*	*	*	0.3	1.7	4.3
2002.....	14.5	*	*	*	*	*	*	0.3	1.7	4.1
2001.....	14.7	*	*	*	*	*	*	0.5	1.8	4.3
2000.....	15.0	*	*	*	*	*	*	0.4	1.9	4.3
1999.....	14.9	*	*	*	*	*	*	0.4	1.7	4.3

All races 3/, male

2006.....	0.3	*	*	*	*	*	*	*	*	*
2005.....	0.3	*	*	*	*	*	*	*	*	*
2004.....	0.3	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*

All races 3/, female

2006.....	26.9	*	*	*	*	*	*	0.7	3.0	7.5
2005.....	27.3	*	*	*	*	*	*	0.8	2.8	7.7
2004.....	27.5	*	*	*	*	*	*	0.8	3.2	7.8
2003.....	28.2	*	*	*	*	*	*	0.6	3.4	8.5
2002.....	28.3	*	*	*	*	*	*	0.7	3.5	8.2
2001.....	28.6	*	*	*	*	*	*	0.9	3.6	8.6
2000.....	29.2	*	*	*	*	*	*	0.8	3.8	8.5
1999.....	28.9	*	*	*	*	*	*	0.9	3.4	8.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of breast (C50)										
White, both sexes										
2006 4/.....	14.2	*	*	*	*	*	*	0.3	1.3	3.3
2005 4/.....	14.4	*	*	*	*	*	*	0.3	1.2	3.3
2004 4/.....	14.6	*	*	*	*	*	*	0.3	1.3	3.4
2003 4/.....	15.0	*	*	*	*	*	*	0.2	1.5	3.8
2002.....	15.1	*	*	*	*	*	*	0.3	1.5	3.6
2001.....	15.2	*	*	*	*	*	*	0.4	1.5	3.9
2000.....	15.7	*	*	*	*	*	*	0.4	1.6	3.9
1999.....	15.5	*	*	*	*	*	*	0.4	1.5	3.7
White, male										
2006 4/.....	0.3	*	*	*	*	*	*	*	*	*
2005 4/.....	0.3	*	*	*	*	*	*	*	*	*
2004 4/.....	0.3	*	*	*	*	*	*	*	*	*
2003 4/.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*
White, female										
2006 4/.....	27.9	*	*	*	*	*	*	0.6	2.6	6.6
2005 4/.....	28.3	*	*	*	*	*	*	0.6	2.4	6.7
2004 4/.....	28.6	*	*	*	*	*	*	0.6	2.7	6.9
2003 4/.....	29.3	*	*	*	*	*	*	0.4	3.0	7.7
2002.....	29.5	*	*	*	*	*	*	0.5	3.0	7.2
2001.....	29.8	*	*	*	*	*	*	0.8	3.0	7.8
2000.....	30.7	*	*	*	*	*	*	0.8	3.3	7.7
1999.....	30.2	*	*	*	*	*	*	0.8	3.0	7.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of breast (C50)										
Black, both sexes										
2006 4/.....	14.5	*	*	*	*	*	*	0.8	3.2	7.6
2005 4/.....	15.0	*	*	*	*	*	*	0.8	2.9	7.8
2004 4/.....	14.5	*	*	*	*	*	*	0.9	3.2	8.1
2003 4/.....	15.1	*	*	*	*	*	*	0.9	3.4	8.0
2002.....	14.9	*	*	*	*	*	*	0.8	3.7	8.1
2001.....	15.0	*	*	*	*	*	*	1.1	4.0	7.5
2000.....	14.8	*	*	*	*	*	*	0.7	3.7	7.7
1999.....	15.0	*	*	*	*	*	*	0.9	3.7	8.8
Black, male										
2006 4/.....	0.3	*	*	*	*	*	*	*	*	*
2005 4/.....	0.3	*	*	*	*	*	*	*	*	*
2004 4/.....	0.3	*	*	*	*	*	*	*	*	*
2003 4/.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*
Black, female										
2006 4/.....	27.5	*	*	*	*	*	*	1.5	6.0	14.3
2005 4/.....	28.4	*	*	*	*	*	*	1.6	5.6	14.7
2004 4/.....	27.5	*	*	*	*	*	*	1.8	6.2	15.1
2003 4/.....	28.5	*	*	*	*	*	*	1.6	6.5	15.1
2002.....	28.2	*	*	*	*	*	*	1.6	6.9	15.1
2001.....	28.3	*	*	*	*	*	*	2.2	7.6	14.2
2000.....	27.9	*	*	*	*	*	*	*	7.1	14.3
1999.....	28.2	*	*	*	*	*	*	1.8	6.9	16.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of cervix uteri (C53)										
All races 3/, both sexes										
2006.....	1.3	*	*	*	*	*	*	0.2	0.7	1.2
2005.....	1.3	*	*	*	*	*	*	0.2	0.6	1.1
2004.....	1.3	*	*	*	*	*	*	0.4	0.7	1.1
2003.....	1.3	*	*	*	*	*	*	0.3	0.7	1.2
2002.....	1.4	*	*	*	*	*	*	0.2	0.8	1.3
2001.....	1.4	*	*	*	*	*	*	0.3	0.8	1.3
2000.....	1.5	*	*	*	*	*	*	0.4	0.7	1.3
1999.....	1.5	*	*	*	*	*	*	0.3	0.8	1.3
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	2.6	*	*	*	*	*	*	0.5	1.4	2.4
2005.....	2.6	*	*	*	*	*	*	0.4	1.3	2.3
2004.....	2.6	*	*	*	*	*	*	0.7	1.4	2.2
2003.....	2.7	*	*	*	*	*	*	0.6	1.4	2.3
2002.....	2.7	*	*	*	*	*	*	0.5	1.5	2.5
2001.....	2.8	*	*	*	*	*	*	0.5	1.5	2.6
2000.....	2.9	*	*	*	*	*	*	0.8	1.5	2.6
1999.....	3.0	*	*	*	*	*	*	0.6	1.5	2.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of cervix uteri (C53)										
White, both sexes										
2006 4/.....	1.2	*	*	*	*	*	*	0.2	0.6	1.2
2005 4/.....	1.2	*	*	*	*	*	*	0.2	0.7	1.1
2004 4/.....	1.2	*	*	*	*	*	*	0.3	0.7	1.1
2003 4/.....	1.2	*	*	*	*	*	*	0.3	0.6	1.1
2002.....	1.3	*	*	*	*	*	*	0.2	0.8	1.2
2001.....	1.3	*	*	*	*	*	*	0.3	0.7	1.3
2000.....	1.4	*	*	*	*	*	*	0.4	0.7	1.2
1999.....	1.4	*	*	*	*	*	*	0.3	0.7	1.2
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	2.5	*	*	*	*	*	*	0.5	1.2	2.4
2005 4/.....	2.5	*	*	*	*	*	*	0.4	1.4	2.2
2004 4/.....	2.4	*	*	*	*	*	*	0.6	1.4	2.2
2003 4/.....	2.5	*	*	*	*	*	*	0.5	1.2	2.1
2002.....	2.5	*	*	*	*	*	*	0.5	1.5	2.4
2001.....	2.7	*	*	*	*	*	*	0.6	1.5	2.5
2000.....	2.7	*	*	*	*	*	*	0.8	1.4	2.4
1999.....	2.8	*	*	*	*	*	*	0.7	1.5	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of cervix uteri (C53)										
Black, both sexes										
2006 4/.....	2.0	*	*	*	*	*	*	*	1.2	1.6
2005 4/.....	2.0	*	*	*	*	*	*	*	0.8	1.6
2004 4/.....	2.0	*	*	*	*	*	*	0.7	0.9	1.6
2003 4/.....	2.1	*	*	*	*	*	*	*	1.2	2.0
2002.....	2.2	*	*	*	*	*	*	*	1.1	2.1
2001.....	2.1	*	*	*	*	*	*	*	1.1	1.8
2000.....	2.4	*	*	*	*	*	*	*	1.4	2.1
1999.....	2.3	*	*	*	*	*	*	*	1.0	2.1
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	3.8	*	*	*	*	*	*	*	2.2	3.0
2005 4/.....	3.8	*	*	*	*	*	*	*	1.5	3.0
2004 4/.....	3.9	*	*	*	*	*	*	1.4	1.7	3.0
2003 4/.....	4.0	*	*	*	*	*	*	*	2.2	3.8
2002.....	4.2	*	*	*	*	*	*	*	2.0	3.9
2001.....	4.0	*	*	*	*	*	*	*	2.1	3.5
2000.....	4.5	*	*	*	*	*	*	*	2.7	4.0
1999.....	4.4	*	*	*	*	*	*	*	1.9	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of ovary (C56)										
All races 3/, both sexes										
2006.....	5.0	*	*	*	*	*	*	0.1	0.3	0.5
2005.....	5.0	*	*	*	*	*	*	0.2	0.3	0.5
2004.....	5.0	*	*	*	*	*	0.1	0.2	0.3	0.6
2003.....	5.0	*	*	*	*	*	*	0.2	0.3	0.6
2002.....	5.1	*	*	*	*	*	0.1	0.2	0.3	0.7
2001.....	5.1	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	5.0	*	*	*	*	*	*	0.2	0.3	0.6
1999.....	4.9	*	*	*	*	*	*	0.1	0.2	0.7
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	9.8	*	*	*	*	*	*	0.3	0.5	1.1
2005.....	9.8	*	*	*	*	*	*	0.3	0.5	1.0
2004.....	9.9	*	*	*	*	*	0.2	0.3	0.6	1.2
2003.....	9.9	*	*	*	*	*	*	0.3	0.5	1.3
2002.....	10.0	*	*	*	*	*	0.3	0.4	0.6	1.4
2001.....	9.9	*	*	*	*	*	*	0.4	0.6	1.4
2000.....	9.8	*	*	*	*	*	*	0.4	0.6	1.3
1999.....	9.6	*	*	*	*	*	*	0.3	0.5	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of ovary (C56)										
White, both sexes										
2006 4/.....	5.5	*	*	*	*	*	*	0.1	0.3	0.6
2005 4/.....	5.5	*	*	*	*	*	*	0.2	0.3	0.5
2004 4/.....	5.5	*	*	*	*	*	*	0.1	0.3	0.6
2003 4/.....	5.6	*	*	*	*	*	*	0.2	0.3	0.6
2002.....	5.6	*	*	*	*	*	0.1	0.2	0.3	0.7
2001.....	5.6	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	5.5	*	*	*	*	*	*	0.2	0.3	0.6
1999.....	5.4	*	*	*	*	*	*	0.1	0.2	0.7
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	10.9	*	*	*	*	*	*	0.3	0.6	1.2
2005 4/.....	10.9	*	*	*	*	*	*	0.3	0.5	1.1
2004 4/.....	10.9	*	*	*	*	*	*	0.3	0.6	1.2
2003 4/.....	11.0	*	*	*	*	*	*	0.3	0.6	1.2
2002.....	11.1	*	*	*	*	*	0.3	0.4	0.6	1.4
2001.....	11.0	*	*	*	*	*	*	0.4	0.6	1.3
2000.....	10.9	*	*	*	*	*	*	0.4	0.6	1.3
1999.....	10.6	*	*	*	*	*	*	0.3	0.5	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
All races 3/, both sexes										
2006.....	4.3	*	0.6	0.9	0.6	0.5	0.5	0.8	1.1	1.6
2005.....	4.4	*	0.5	1.0	0.7	0.6	0.6	0.8	1.1	1.6
2004.....	4.4	*	0.6	0.9	0.7	0.5	0.5	0.7	1.1	1.7
2003.....	4.4	*	0.7	0.9	0.8	0.6	0.6	0.9	1.2	1.9
2002.....	4.4	*	0.7	1.0	0.6	0.5	0.5	0.7	1.2	1.7
2001.....	4.4	*	0.7	0.8	0.6	0.5	0.6	0.7	1.4	1.8
2000.....	4.5	0.5	0.7	0.8	0.7	0.5	0.6	0.8	1.2	2.0
1999.....	4.6	*	0.7	0.9	0.6	0.6	0.6	0.9	1.3	1.9
All races 3/, male										
2006.....	4.9	*	0.7	0.9	0.7	0.5	0.5	0.9	1.2	1.9
2005.....	5.0	*	0.5	0.9	0.7	0.7	0.6	1.1	1.3	2.0
2004.....	4.9	*	0.7	1.0	0.7	0.6	0.7	0.8	1.3	2.1
2003.....	5.0	*	0.7	0.9	0.8	0.6	0.6	0.9	1.4	2.3
2002.....	5.0	*	0.8	1.1	0.7	0.6	0.7	0.7	1.5	2.0
2001.....	5.0	*	0.7	0.8	0.6	0.6	0.7	0.9	1.7	2.2
2000.....	5.1	*	0.7	0.8	0.8	0.6	0.7	1.0	1.6	2.6
1999.....	5.1	*	0.7	0.9	0.6	0.7	0.7	1.1	1.7	2.3
All races 3/, female										
2006.....	3.7	*	0.6	0.8	0.6	0.4	0.5	0.7	1.0	1.2
2005.....	3.9	*	0.6	1.0	0.6	0.4	0.5	0.5	0.8	1.2
2004.....	3.9	*	0.6	0.9	0.7	0.5	0.4	0.6	0.8	1.2
2003.....	3.9	*	0.6	0.9	0.7	0.5	0.5	0.9	0.9	1.5
2002.....	3.9	*	0.7	0.9	0.5	0.4	0.3	0.7	1.0	1.4
2001.....	3.9	*	0.6	0.9	0.6	0.4	0.6	0.5	1.1	1.4
2000.....	3.9	*	0.6	0.8	0.6	0.4	0.5	0.7	0.8	1.4
1999.....	4.1	*	0.8	0.9	0.7	0.5	0.5	0.8	0.9	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
White, both sexes										
2006 4/.....	4.9	*	0.6	0.9	0.7	0.5	0.5	0.8	1.3	1.8
2005 4/.....	5.0	*	0.5	0.9	0.7	0.6	0.6	0.9	1.2	1.8
2004 4/.....	4.9	*	0.7	1.0	0.6	0.6	0.6	0.7	1.2	1.9
2003 4/.....	5.0	*	0.7	0.9	0.8	0.6	0.6	1.0	1.3	2.1
2002.....	5.0	*	0.7	1.0	0.6	0.5	0.5	0.7	1.4	1.9
2001.....	5.0	*	0.7	0.9	0.6	0.6	0.6	0.8	1.5	2.1
2000.....	5.1	*	0.7	0.8	0.7	0.5	0.7	0.9	1.3	2.2
1999.....	5.2	*	0.8	0.9	0.6	0.5	0.6	1.1	1.4	2.2
White, male										
2006 4/.....	5.5	*	0.7	1.0	0.7	0.5	0.5	1.0	1.4	2.1
2005 4/.....	5.6	*	0.4	0.9	0.8	0.8	0.7	1.2	1.5	2.2
2004 4/.....	5.5	*	0.7	1.0	0.7	0.6	0.7	0.8	1.5	2.4
2003 4/.....	5.6	*	0.8	0.9	0.8	0.7	0.7	1.0	1.6	2.5
2002.....	5.7	*	0.7	1.1	0.7	0.7	0.8	0.7	1.6	2.3
2001.....	5.7	*	0.8	0.8	0.6	0.7	0.7	1.1	1.9	2.6
2000.....	5.7	*	0.7	0.9	0.8	0.6	0.7	1.0	1.7	2.8
1999.....	5.8	*	0.8	0.9	0.6	0.6	0.7	1.2	1.8	2.6
White, female										
2006 4/.....	4.3	*	0.5	0.8	0.6	0.4	0.5	0.6	1.1	1.4
2005 4/.....	4.4	*	0.6	1.0	0.6	0.4	0.5	0.6	0.9	1.3
2004 4/.....	4.4	*	0.6	0.9	0.6	0.5	0.4	0.7	0.9	1.4
2003 4/.....	4.4	*	0.6	0.9	0.8	0.6	0.5	0.9	1.0	1.8
2002.....	4.4	*	0.6	1.0	0.6	0.4	0.3	0.7	1.2	1.6
2001.....	4.4	*	0.7	1.0	0.7	0.4	0.6	0.5	1.2	1.6
2000.....	4.4	*	0.6	0.8	0.7	0.4	0.6	0.8	0.9	1.5
1999.....	4.6	*	0.8	0.8	0.7	0.5	0.5	0.9	1.0	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	18.4	0.8	0.8	0.6	0.9	1.3	1.9	1.9	2.2	2.8
2005.....	18.6	0.6	0.8	0.8	0.9	1.3	1.8	1.8	2.2	2.7
2004.....	18.6	0.6	0.8	0.9	0.9	1.4	1.7	1.8	2.3	2.7
2003.....	19.1	0.8	0.9	0.9	1.0	1.2	1.7	1.9	2.4	3.1
2002.....	19.5	0.7	0.8	0.8	1.0	1.5	1.9	1.9	2.6	3.2
2001.....	19.8	0.7	1.0	0.8	1.0	1.5	2.0	2.3	2.7	3.5
2000.....	20.1	1.0	0.9	0.8	1.0	1.6	2.1	2.1	2.5	3.4
1999.....	20.2	0.8	0.9	0.7	1.0	1.5	2.2	2.2	2.6	3.7

All races 3/, male

2006.....	20.3	*	0.9	0.7	1.0	1.6	2.3	2.3	2.6	3.1
2005.....	20.6	*	1.0	0.9	1.0	1.6	2.2	2.1	2.4	3.0
2004.....	20.4	*	0.9	1.1	1.0	1.6	2.1	2.1	2.7	3.2
2003.....	21.0	*	1.0	1.0	1.1	1.6	2.0	2.3	3.0	3.5
2002.....	21.4	*	0.8	0.9	1.2	1.8	2.2	2.3	3.0	3.7
2001.....	21.5	*	1.0	0.9	1.1	1.8	2.3	2.6	3.1	4.2
2000.....	21.6	1.1	1.0	0.9	1.2	1.9	2.6	2.3	3.0	3.9
1999.....	21.7	*	1.0	0.8	1.2	1.7	2.7	2.4	3.1	4.4

All races 3/, female

2006.....	16.5	*	0.8	0.5	0.7	0.9	1.3	1.5	1.8	2.4
2005.....	16.6	*	0.6	0.6	0.7	1.0	1.3	1.5	2.0	2.3
2004.....	16.8	*	0.6	0.7	0.7	1.1	1.3	1.5	1.9	2.2
2003.....	17.3	1.1	0.8	0.7	0.9	0.9	1.4	1.6	1.7	2.6
2002.....	17.7	*	0.7	0.8	0.8	1.2	1.7	1.5	2.2	2.6
2001.....	18.1	*	1.0	0.7	0.8	1.2	1.7	2.0	2.2	2.7
2000.....	18.6	*	0.9	0.7	0.8	1.2	1.5	1.9	1.9	3.0
1999.....	18.7	*	0.8	0.6	0.9	1.3	1.8	1.9	2.2	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
White, both sexes										
2006 4/.....	20.1	0.8	0.9	0.6	0.9	1.3	1.8	2.0	2.1	2.6
2005 4/.....	20.3	0.6	0.7	0.8	0.9	1.4	1.8	1.6	2.1	2.7
2004 4/.....	20.2	0.6	0.8	1.0	0.9	1.4	1.7	1.7	2.2	2.6
2003 4/.....	20.9	0.9	0.9	0.9	1.0	1.3	1.7	1.9	2.3	3.0
2002.....	21.3	*	0.8	0.9	1.0	1.5	2.1	1.9	2.5	3.1
2001.....	21.5	0.7	1.1	0.8	1.0	1.6	2.0	2.1	2.6	3.4
2000.....	21.9	1.2	1.0	0.9	1.1	1.6	2.0	2.1	2.4	3.3
1999.....	21.9	0.8	1.0	0.7	1.0	1.5	2.3	2.1	2.5	3.6
White, male										
2006 4/.....	22.1	*	0.9	0.6	1.0	1.7	2.4	2.4	2.5	3.0
2005 4/.....	22.5	*	0.9	0.9	1.1	1.7	2.2	1.9	2.2	3.1
2004 4/.....	22.2	*	0.8	1.2	1.0	1.7	2.1	2.1	2.6	3.1
2003 4/.....	23.1	*	1.1	1.2	1.1	1.6	2.0	2.2	2.9	3.4
2002.....	23.4	*	0.9	0.9	1.2	1.9	2.4	2.3	2.9	3.5
2001.....	23.3	*	1.1	0.9	1.0	1.9	2.3	2.4	3.2	4.0
2000.....	23.6	*	1.0	0.9	1.2	1.9	2.5	2.3	2.9	3.8
1999.....	23.6	*	1.0	0.9	1.1	1.7	2.8	2.3	2.9	4.3
White, female										
2006 4/.....	18.0	*	0.8	0.6	0.7	0.9	1.3	1.5	1.7	2.3
2005 4/.....	18.1	*	0.5	0.7	0.7	1.0	1.4	1.3	1.9	2.3
2004 4/.....	18.3	*	0.7	0.7	0.8	1.1	1.3	1.3	1.8	2.1
2003 4/.....	18.8	1.4	0.8	0.7	0.9	0.8	1.5	1.6	1.7	2.5
2002.....	19.2	*	0.8	0.9	0.8	1.1	1.7	1.5	2.2	2.6
2001.....	19.7	*	1.1	0.7	0.9	1.2	1.6	1.9	2.0	2.8
2000.....	20.3	*	1.0	0.8	0.9	1.2	1.4	1.8	1.9	2.9
1999.....	20.4	*	0.9	0.6	0.9	1.3	1.7	1.9	2.1	2.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
Black, both sexes										
2006 4/.....	13.3	*	*	*	0.9	1.2	2.0	1.9	2.9	4.1
2005 4/.....	12.9	*	1.0	*	0.8	1.3	1.9	2.7	3.8	3.2
2004 4/.....	13.6	*	*	0.7	0.8	1.4	1.9	2.8	3.1	3.7
2003 4/.....	13.4	*	*	0.6	1.0	1.1	1.7	2.4	3.3	4.1
2002.....	13.6	*	*	*	1.0	1.4	1.5	2.3	3.4	3.8
2001.....	14.1	*	*	0.9	1.0	1.5	2.1	3.2	3.5	4.5
2000.....	13.9	*	0.8	0.6	0.7	1.9	2.6	2.8	3.1	4.4
1999.....	14.1	*	*	0.6	1.3	1.7	2.9	2.6	3.2	4.6
Black, male										
2006 4/.....	14.9	*	*	*	*	1.5	2.4	2.1	3.5	4.9
2005 4/.....	14.1	*	*	*	*	1.4	2.5	2.8	4.7	3.7
2004 4/.....	14.4	*	*	*	*	1.4	2.5	2.9	3.6	4.3
2003 4/.....	13.9	*	*	*	1.2	*	2.1	2.5	4.2	4.7
2002.....	14.6	*	*	*	1.1	1.6	1.3	2.8	4.4	5.1
2001.....	15.1	*	*	*	1.7	1.8	2.2	3.8	4.0	6.5
2000.....	14.7	*	*	*	*	2.3	3.2	2.6	4.4	4.8
1999.....	15.1	*	*	*	1.7	2.1	3.1	3.3	4.2	5.3
Black, female										
2006 4/.....	11.9	*	*	*	*	*	1.7	1.8	2.3	3.3
2005 4/.....	11.8	*	*	*	*	1.3	*	2.6	3.0	2.8
2004 4/.....	12.8	*	*	*	*	1.4	1.3	2.7	2.7	3.1
2003 4/.....	12.9	*	*	*	*	*	1.3	2.4	2.4	3.6
2002.....	12.8	*	*	*	*	1.3	1.6	2.0	2.5	2.7
2001.....	13.1	*	*	*	*	*	2.1	2.7	3.1	2.8
2000.....	13.1	*	*	*	*	1.4	2.1	3.0	1.9	4.1
1999.....	13.2	*	*	*	*	1.4	2.8	2.0	2.3	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hodgkin's disease (C81)										
All races 3/, both sexes										
2006.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3
2005.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3
2004.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.3
1999.....	0.5	*	*	*	*	0.2	0.3	0.4	0.3	0.4
All races 3/, male										
2006.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4
2005.....	0.5	*	*	*	*	*	0.2	0.3	0.3	0.3
2004.....	0.5	*	*	*	*	*	0.3	0.3	0.4	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.5
2000.....	0.5	*	*	*	*	*	0.3	0.3	0.5	0.4
1999.....	0.6	*	*	*	*	0.2	0.4	0.4	0.4	0.5
All races 3/, female										
2006.....	0.4	*	*	*	*	*	*	0.3	0.2	0.3
2005.....	0.4	*	*	*	*	*	0.3	0.2	0.3	0.2
2004.....	0.4	*	*	*	*	*	*	0.3	0.3	0.2
2003.....	0.4	*	*	*	*	*	0.2	0.4	0.4	0.3
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3
2001.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.3
2000.....	0.4	*	*	*	*	*	0.2	0.3	0.4	0.3
1999.....	0.4	*	*	*	*	*	0.3	0.4	0.3	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hodgkin's disease (C81)										
White, both sexes										
2006 4/.....	0.5	*	*	*	*	*	0.2	0.4	0.3	0.4
2005 4/.....	0.5	*	*	*	*	*	0.2	0.2	0.2	0.3
2004 4/.....	0.5	*	*	*	*	*	0.2	0.3	0.3	0.3
2003 4/.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4
1999.....	0.6	*	*	*	*	0.2	0.3	0.4	0.4	0.4
White, male										
2006 4/.....	0.6	*	*	*	*	*	0.3	0.4	0.4	0.4
2005 4/.....	0.5	*	*	*	*	*	*	0.3	*	0.3
2004 4/.....	0.5	*	*	*	*	*	0.3	0.3	0.4	0.3
2003 4/.....	0.5	*	*	*	*	*	*	0.4	0.5	0.4
2002.....	0.6	*	*	*	*	*	0.3	0.4	0.4	0.4
2001.....	0.6	*	*	*	*	*	0.3	0.4	0.5	0.5
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4
1999.....	0.6	*	*	*	*	*	0.4	0.4	0.5	0.5
White, female										
2006 4/.....	0.4	*	*	*	*	*	*	0.3	*	0.3
2005 4/.....	0.4	*	*	*	*	*	0.3	*	0.3	*
2004 4/.....	0.4	*	*	*	*	*	*	0.3	0.3	*
2003 4/.....	0.5	*	*	*	*	*	*	0.4	0.4	0.3
2002.....	0.5	*	*	*	*	*	0.4	0.3	0.5	0.3
2001.....	0.4	*	*	*	*	*	*	0.4	0.3	0.3
2000.....	0.5	*	*	*	*	*	*	0.3	0.4	0.3
1999.....	0.5	*	*	*	*	*	0.3	0.4	0.3	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Non-Hodgkin's Lymphoma (C82-C85)										
All races 3/, both sexes										
2006.....	6.9	*	*	*	0.1	0.2	0.4	0.5	0.6	0.8
2005.....	7.0	*	*	*	0.1	0.2	0.4	0.4	0.7	0.9
2004.....	7.1	*	*	0.1	*	0.3	0.4	0.6	0.7	0.9
2003.....	7.4	*	*	*	0.1	0.3	0.4	0.5	0.8	1.1
2002.....	7.6	*	*	*	0.2	0.2	0.5	0.6	0.8	1.2
2001.....	7.8	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4
2000.....	8.1	*	*	0.1	0.1	0.3	0.6	0.6	0.8	1.3
1999.....	8.2	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4
All races 3/, male										
2006.....	7.4	*	*	*	*	0.3	0.5	0.6	0.7	1.0
2005.....	7.6	*	*	*	0.2	0.3	0.5	0.5	0.8	1.1
2004.....	7.6	*	*	*	*	0.3	0.5	0.8	0.8	1.2
2003.....	8.0	*	*	*	*	0.3	0.5	0.6	1.0	1.4
2002.....	8.1	*	*	*	0.3	0.3	0.6	0.7	1.0	1.5
2001.....	8.4	*	*	*	0.2	0.4	0.5	0.8	1.0	1.8
2000.....	8.6	*	*	*	*	0.4	0.8	0.7	1.0	1.7
1999.....	8.6	*	*	*	0.2	0.3	0.6	0.7	1.1	1.8
All races 3/, female										
2006.....	6.4	*	*	*	*	*	0.3	0.4	0.5	0.7
2005.....	6.5	*	*	*	*	*	0.3	0.4	0.5	0.7
2004.....	6.6	*	*	*	*	*	0.3	0.4	0.6	0.6
2003.....	6.8	*	*	*	*	*	0.3	0.4	0.5	0.7
2002.....	7.1	*	*	*	*	*	0.4	0.4	0.6	0.8
2001.....	7.3	*	*	*	*	*	0.4	0.6	0.6	1.0
2000.....	7.6	*	*	*	*	*	0.4	0.6	0.6	0.9
1999.....	7.7	*	*	*	*	0.2	0.4	0.4	0.7	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Non-Hodgkin's Lymphoma (C82-C85)										
White, both sexes										
2006 4/.....	7.7	*	*	*	*	0.2	0.4	0.5	0.6	0.7
2005 4/.....	7.9	*	*	*	0.2	0.2	0.3	0.4	0.6	0.8
2004 4/.....	8.0	*	*	*	*	0.3	0.4	0.5	0.6	0.9
2003 4/.....	8.3	*	*	*	*	0.2	0.4	0.4	0.7	1.1
2002.....	8.5	*	*	*	0.2	0.2	0.5	0.6	0.7	1.1
2001.....	8.8	*	*	*	0.1	0.3	0.4	0.6	0.8	1.3
2000.....	9.1	*	*	*	0.1	0.3	0.6	0.6	0.8	1.2
1999.....	9.2	*	*	*	*	0.2	0.5	0.5	0.8	1.4
White, male										
2006 4/.....	8.2	*	*	*	*	0.3	0.5	0.6	0.7	0.9
2005 4/.....	8.5	*	*	*	0.3	0.4	0.5	0.4	0.7	1.1
2004 4/.....	8.5	*	*	*	*	0.4	0.5	0.8	0.7	1.2
2003 4/.....	8.9	*	*	*	*	0.3	0.5	0.6	0.9	1.4
2002.....	9.1	*	*	*	0.3	0.3	0.6	0.8	0.9	1.4
2001.....	9.3	*	*	*	*	0.4	0.5	0.7	1.0	1.6
2000.....	9.6	*	*	*	*	0.5	0.8	0.6	1.0	1.6
1999.....	9.6	*	*	*	*	0.3	0.6	0.6	1.0	1.7
White, female										
2006 4/.....	7.2	*	*	*	*	*	*	0.4	0.4	0.6
2005 4/.....	7.3	*	*	*	*	*	*	0.3	0.5	0.6
2004 4/.....	7.5	*	*	*	*	*	0.3	0.3	0.5	0.6
2003 4/.....	7.7	*	*	*	*	*	0.4	0.3	0.4	0.7
2002.....	8.0	*	*	*	*	*	0.3	0.4	0.6	0.8
2001.....	8.2	*	*	*	*	*	0.4	0.5	0.5	1.0
2000.....	8.6	*	*	*	*	*	0.4	0.5	0.6	0.8
1999.....	8.8	*	*	*	*	*	0.4	0.4	0.6	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
All races 3/, both sexes										
2006.....	7.3	0.8	0.8	0.5	0.7	1.0	1.2	1.1	1.2	1.4
2005.....	7.3	0.5	0.7	0.6	0.7	1.1	1.2	1.1	1.2	1.4
2004.....	7.3	0.6	0.8	0.8	0.7	1.0	1.1	0.9	1.2	1.4
2003.....	7.4	0.7	0.8	0.8	0.9	0.9	1.1	1.1	1.1	1.5
2002.....	7.5	0.5	0.7	0.7	0.8	1.2	1.2	1.0	1.3	1.5
2001.....	7.5	0.6	1.0	0.8	0.8	1.1	1.3	1.2	1.4	1.6
2000.....	7.6	1.0	0.9	0.7	0.8	1.2	1.2	1.2	1.2	1.6
1999.....	7.5	0.7	0.8	0.6	0.9	1.1	1.4	1.2	1.3	1.6
All races 3/, male										
2006.....	8.4	*	0.8	0.6	0.8	1.2	1.5	1.3	1.4	1.6
2005.....	8.4	*	1.0	0.8	0.8	1.2	1.5	1.3	1.2	1.4
2004.....	8.3	*	0.9	1.0	0.8	1.2	1.3	1.0	1.4	1.5
2003.....	8.4	*	0.9	0.9	1.0	1.2	1.2	1.3	1.4	1.6
2002.....	8.5	*	0.7	0.7	0.9	1.4	1.4	1.2	1.5	1.6
2001.....	8.5	*	1.0	0.8	0.9	1.3	1.5	1.4	1.6	1.8
2000.....	8.5	1.1	0.9	0.8	1.0	1.4	1.5	1.3	1.5	1.6
1999.....	8.4	*	0.8	0.7	0.9	1.3	1.7	1.3	1.4	1.8
All races 3/, female										
2006.....	6.3	*	0.8	0.5	0.6	0.7	0.9	0.8	1.0	1.3
2005.....	6.2	*	0.5	0.5	0.7	0.9	0.8	0.9	1.2	1.3
2004.....	6.3	*	0.6	0.6	0.7	0.8	0.8	0.8	0.9	1.2
2003.....	6.4	1.1	0.8	0.7	0.8	0.6	0.9	0.9	0.8	1.4
2002.....	6.5	*	0.6	0.8	0.7	1.0	1.0	0.8	1.1	1.3
2001.....	6.6	*	1.0	0.7	0.7	0.9	1.0	1.0	1.2	1.3
2000.....	6.7	*	0.9	0.6	0.6	0.9	0.9	1.0	0.9	1.6
1999.....	6.7	*	0.7	0.5	0.8	0.9	1.1	1.1	1.2	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
White, both sexes										
2006 4/.....	8.1	0.8	0.8	0.5	0.8	1.0	1.2	1.1	1.2	1.4
2005 4/.....	8.1	*	0.7	0.7	0.7	1.1	1.2	1.0	1.2	1.5
2004 4/.....	8.0	*	0.7	0.8	0.8	1.0	1.1	0.9	1.2	1.3
2003 4/.....	8.2	0.9	0.9	0.9	0.9	0.9	1.1	1.1	1.1	1.4
2002.....	8.2	*	0.7	0.8	0.8	1.2	1.3	1.0	1.3	1.5
2001.....	8.3	0.6	1.1	0.7	0.8	1.2	1.3	1.1	1.4	1.5
2000.....	8.3	1.2	1.0	0.7	0.9	1.2	1.2	1.1	1.1	1.6
1999.....	8.3	0.8	0.9	0.7	0.8	1.1	1.4	1.2	1.3	1.6
White, male										
2006 4/.....	9.3	*	0.8	0.5	0.9	1.3	1.6	1.5	1.3	1.6
2005 4/.....	9.3	*	0.9	0.8	0.8	1.3	1.5	1.2	1.3	1.5
2004 4/.....	9.1	*	0.8	1.0	0.8	1.2	1.4	1.0	1.4	1.4
2003 4/.....	9.4	*	1.0	1.0	1.0	1.2	1.2	1.3	1.4	1.4
2002.....	9.3	*	0.8	0.7	0.9	1.4	1.5	1.2	1.5	1.6
2001.....	9.3	*	1.1	0.8	0.8	1.4	1.6	1.3	1.6	1.7
2000.....	9.4	*	0.9	0.8	1.1	1.4	1.5	1.3	1.4	1.7
1999.....	9.2	*	0.9	0.8	0.9	1.2	1.8	1.3	1.4	1.8
White, female										
2006 4/.....	7.0	*	0.8	0.5	0.7	0.8	0.9	0.8	1.0	1.3
2005 4/.....	6.9	*	0.5	0.6	0.7	0.9	0.9	0.8	1.2	1.4
2004 4/.....	6.9	*	0.7	0.7	0.7	0.8	0.9	0.8	1.0	1.2
2003 4/.....	7.0	1.4	0.8	0.7	0.8	0.6	0.9	0.9	0.8	1.4
2002.....	7.1	*	0.6	0.8	0.7	1.0	1.1	0.8	1.1	1.4
2001.....	7.3	*	1.1	0.7	0.8	0.9	1.0	0.9	1.2	1.4
2000.....	7.4	*	1.0	0.7	0.7	0.9	0.9	0.9	0.9	1.6
1999.....	7.3	*	0.8	0.6	0.8	1.0	1.0	1.1	1.2	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
Black, both sexes										
2006 4/.....	4.6	*	*	*	0.7	0.9	1.2	0.9	1.3	1.5
2005 4/.....	4.4	*	1.0	*	0.7	1.0	0.9	1.5	1.5	1.1
2004 4/.....	4.8	*	*	*	0.6	1.1	0.9	1.1	1.5	1.8
2003 4/.....	4.6	*	*	*	0.8	0.7	1.0	1.0	1.4	2.0
2002.....	4.7	*	*	*	0.7	0.9	0.8	1.1	1.4	1.5
2001.....	4.7	*	*	0.8	0.9	0.9	1.2	1.4	1.4	1.8
2000.....	4.8	*	*	*	0.6	1.5	1.3	1.4	1.8	1.7
1999.....	4.7	*	*	*	1.1	1.2	1.7	1.1	1.4	1.8
Black, male										
2006 4/.....	5.3	*	*	*	*	*	1.5	*	1.6	1.7
2005 4/.....	4.9	*	*	*	*	*	1.6	1.5	*	*
2004 4/.....	5.3	*	*	*	*	*	*	*	1.8	2.1
2003 4/.....	5.0	*	*	*	*	*	1.3	*	1.8	2.5
2002.....	5.1	*	*	*	*	*	*	*	1.6	1.9
2001.....	5.2	*	*	*	1.5	*	*	1.7	1.8	2.3
2000.....	5.3	*	*	*	*	2.0	1.8	*	2.5	1.4
1999.....	5.2	*	*	*	1.5	1.5	1.7	*	1.7	1.8
Black, female										
2006 4/.....	3.9	*	*	*	*	*	*	*	*	*
2005 4/.....	4.0	*	*	*	*	*	*	1.4	1.6	*
2004 4/.....	4.4	*	*	*	*	*	*	*	*	1.5
2003 4/.....	4.2	*	*	*	*	*	*	*	*	1.7
2002.....	4.3	*	*	*	*	*	*	*	*	*
2001.....	4.3	*	*	*	*	*	*	*	*	1.4
2000.....	4.2	*	*	*	*	*	*	1.5	*	1.9
1999.....	4.3	*	*	*	*	*	1.8	*	*	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes

2006.....	21.1	0.5	0.7	0.7	0.5	1.1	1.3	1.6	1.9	2.7
2005.....	21.2	*	0.8	0.6	0.8	1.3	1.5	1.5	2.0	2.9
2004.....	21.1	0.6	0.8	0.6	0.7	1.2	1.4	1.5	2.0	3.1
2003.....	21.5	*	0.7	0.6	0.7	1.2	1.3	1.6	1.9	3.1
2002.....	21.8	0.6	0.8	0.7	0.7	1.1	1.5	1.5	2.1	3.3
2001.....	22.0	*	0.8	0.6	0.8	1.2	1.4	1.7	2.0	3.2
2000.....	22.4	0.6	0.8	0.6	0.7	1.2	1.4	1.5	2.0	3.4
1999.....	23.7	*	0.8	0.7	0.7	1.2	1.5	1.6	2.2	3.6

All races 3/, male

2006.....	21.9	*	0.7	0.8	0.5	1.3	1.8	1.9	2.4	2.9
2005.....	22.1	*	0.8	0.6	0.8	1.4	1.9	1.9	2.4	3.1
2004.....	21.9	*	0.7	0.8	0.8	1.5	1.8	1.9	2.3	3.6
2003.....	22.3	*	0.8	0.8	0.7	1.5	1.6	1.9	2.4	3.5
2002.....	22.7	*	0.8	0.8	0.7	1.3	1.9	2.0	2.5	3.7
2001.....	22.6	*	0.9	0.6	0.8	1.5	1.9	2.0	2.4	3.5
2000.....	23.0	*	1.0	0.6	0.7	1.4	1.8	1.9	2.4	3.7
1999.....	24.5	*	0.8	0.8	0.8	1.5	1.8	2.1	2.8	4.1

All races 3/, female

2006.....	20.2	*	0.6	0.6	0.5	0.9	0.8	1.3	1.4	2.5
2005.....	20.4	*	0.7	0.6	0.7	1.2	1.0	1.0	1.6	2.7
2004.....	20.3	*	0.9	0.5	0.6	1.0	0.9	1.1	1.7	2.5
2003.....	20.8	*	0.5	0.5	0.7	0.8	1.1	1.2	1.4	2.7
2002.....	21.0	*	0.8	0.6	0.8	0.9	1.0	1.0	1.7	2.8
2001.....	21.4	*	0.6	0.5	0.7	0.9	0.8	1.4	1.6	3.0
2000.....	21.9	*	0.7	0.6	0.6	1.0	1.0	1.1	1.6	3.0
1999.....	23.1	*	0.7	0.7	0.6	0.9	1.1	1.1	1.6	3.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

White, both sexes

2006 4/.....	22.6	*	0.6	0.7	0.5	1.1	1.4	1.6	2.0	2.7
2005 4/.....	22.7	*	0.8	0.6	0.7	1.4	1.5	1.4	1.9	2.9
2004 4/.....	22.6	*	0.8	0.7	0.7	1.2	1.3	1.6	2.0	3.0
2003 4/.....	23.1	*	0.6	0.7	0.7	1.3	1.4	1.6	1.8	3.1
2002.....	23.3	0.7	0.7	0.7	0.8	1.2	1.5	1.5	2.1	3.2
2001.....	23.5	*	0.8	0.6	0.8	1.2	1.5	1.7	1.9	3.1
2000.....	23.9	*	0.8	0.7	0.7	1.2	1.5	1.6	2.0	3.2
1999.....	25.3	*	0.8	0.7	0.7	1.2	1.4	1.6	2.2	3.5

White, male

2006 4/.....	23.6	*	0.7	0.8	0.5	1.3	1.9	2.0	2.5	2.9
2005 4/.....	23.6	*	0.9	0.6	0.7	1.5	2.0	1.9	2.3	3.2
2004 4/.....	23.4	*	0.7	0.8	0.8	1.5	1.8	2.0	2.4	3.5
2003 4/.....	23.8	*	0.7	0.9	0.6	1.6	1.6	2.0	2.3	3.6
2002.....	24.2	*	0.8	0.7	0.7	1.4	1.9	2.0	2.5	3.6
2001.....	24.0	*	1.0	0.6	0.8	1.5	2.0	2.0	2.3	3.4
2000.....	24.4	*	1.0	0.7	0.7	1.3	1.9	2.1	2.5	3.6
1999.....	25.9	*	0.9	0.8	0.8	1.5	1.7	2.2	2.7	4.0

White, female

2006 4/.....	21.7	*	0.5	0.6	0.6	0.9	0.8	1.1	1.5	2.4
2005 4/.....	21.9	*	0.6	0.5	0.8	1.2	1.0	0.9	1.5	2.6
2004 4/.....	21.8	*	0.8	0.5	0.6	1.0	0.9	1.1	1.5	2.4
2003 4/.....	22.5	*	0.5	0.5	0.7	0.9	1.1	1.1	1.3	2.6
2002.....	22.5	*	0.7	0.6	0.8	1.0	1.0	1.0	1.7	2.8
2001.....	22.9	*	0.7	0.5	0.7	1.0	0.8	1.4	1.5	2.8
2000.....	23.5	*	0.6	0.6	0.6	1.0	1.1	1.1	1.5	2.8
1999.....	24.6	*	0.7	0.7	0.6	1.0	1.1	1.0	1.7	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Black, both sexes

2006 4/.....	16.8	*	1.0	0.7	*	1.1	1.3	2.1	2.1	3.1
2005 4/.....	17.1	*	0.8	0.7	1.1	0.9	1.5	2.1	2.6	3.6
2004 4/.....	17.2	*	0.8	0.8	1.0	1.4	1.8	1.5	2.1	4.3
2003 4/.....	17.3	*	0.9	0.6	0.9	0.9	1.3	1.9	2.7	3.7
2002.....	18.1	*	1.1	0.8	0.7	1.1	1.6	1.9	2.6	4.5
2001.....	18.3	*	*	*	0.8	1.2	1.3	2.0	2.6	4.3
2000.....	18.3	*	1.0	*	0.7	1.5	1.3	1.3	2.0	5.2
1999.....	19.5	*	*	0.6	0.9	1.3	1.6	1.5	2.4	4.9

Black, male

2006 4/.....	17.4	*	*	*	*	1.4	1.4	2.1	2.7	3.3
2005 4/.....	18.0	*	*	*	1.4	*	1.9	2.4	3.2	3.7
2004 4/.....	18.4	*	*	*	*	1.7	2.1	1.7	2.1	4.6
2003 4/.....	18.3	*	*	*	*	1.2	1.7	1.7	3.0	3.6
2002.....	18.8	*	*	*	*	1.4	2.2	2.5	2.7	5.1
2001.....	19.4	*	*	*	*	1.5	1.9	2.5	3.0	4.4
2000.....	19.4	*	*	*	*	1.7	2.1	1.6	2.0	5.4
1999.....	20.6	*	*	*	*	1.9	2.0	1.5	3.1	5.1

Black, female

2006 4/.....	16.3	*	*	*	*	*	1.3	2.2	1.6	3.0
2005 4/.....	16.3	*	*	*	*	*	*	1.7	2.0	3.6
2004 4/.....	16.2	*	*	*	1.1	*	1.5	*	2.2	4.0
2003 4/.....	16.5	*	*	*	*	*	*	2.0	2.3	3.9
2002.....	17.4	*	*	*	*	*	*	1.4	2.6	4.0
2001.....	17.3	*	*	*	*	*	*	1.5	2.3	4.2
2000.....	17.3	*	*	*	*	1.4	*	*	1.9	5.1
1999.....	18.5	*	*	*	*	*	*	1.5	1.8	4.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races 3/, both sexes

2006.....	4.7	1.6	0.4	0.2	0.2	0.3	0.2	0.3	0.4	0.6
2005.....	4.6	1.4	0.3	0.2	0.2	0.2	0.2	0.3	0.5	0.7
2004.....	4.6	1.7	0.3	0.2	0.2	0.2	0.3	0.3	0.5	0.6
2003.....	4.7	1.5	0.3	0.2	0.2	0.2	0.3	0.4	0.5	0.7
2002.....	4.6	1.7	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.6
2001.....	4.8	1.3	0.4	0.3	0.3	0.2	0.3	0.3	0.4	0.7
2000.....	4.8	1.6	0.3	0.3	0.2	0.2	0.3	0.4	0.5	0.7
1999.....	4.7	1.5	0.4	0.3	0.2	0.3	0.2	0.3	0.6	0.6

All races 3/, male

2006.....	4.7	1.6	0.3	0.2	*	0.2	0.2	0.3	0.4	0.7
2005.....	4.7	1.0	0.4	*	*	0.3	0.2	0.4	0.5	0.7
2004.....	4.7	1.7	0.3	0.2	0.2	0.2	0.4	0.4	0.6	0.6
2003.....	4.7	1.6	0.4	0.2	0.2	0.2	0.3	0.4	0.6	0.7
2002.....	4.6	1.5	0.4	0.2	0.2	0.2	0.3	0.3	0.5	0.7
2001.....	4.8	1.3	0.4	0.3	0.2	0.2	0.3	0.3	0.4	0.7
2000.....	4.8	1.6	0.4	0.3	*	0.3	0.4	0.3	0.5	0.8
1999.....	4.8	1.5	0.4	0.3	*	0.3	0.3	0.3	0.6	0.7

All races 3/, female

2006.....	4.7	1.6	0.5	*	0.2	0.3	0.2	0.4	0.4	0.5
2005.....	4.6	1.8	0.3	0.2	*	0.2	*	0.3	0.4	0.7
2004.....	4.6	1.8	0.3	0.2	*	0.2	0.2	0.3	0.5	0.6
2003.....	4.7	1.5	0.3	*	*	*	0.2	0.3	0.5	0.6
2002.....	4.6	2.0	0.4	0.2	*	0.2	*	0.3	0.5	0.5
2001.....	4.8	1.3	0.4	0.3	0.3	0.3	0.2	0.3	0.5	0.7
2000.....	4.9	1.5	0.3	0.3	0.2	*	0.2	0.4	0.4	0.6
1999.....	4.7	1.5	0.4	0.3	*	0.3	*	0.2	0.5	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

White, both sexes

2006 4/.....	5.2	1.6	0.4	0.2	0.2	0.2	0.2	0.3	0.4	0.6
2005 4/.....	5.1	1.3	0.3	0.2	0.2	0.2	0.2	0.3	0.5	0.6
2004 4/.....	5.1	1.9	0.3	0.2	0.2	0.2	0.3	0.3	0.5	0.6
2003 4/.....	5.2	1.5	0.3	0.2	0.2	0.2	0.3	0.4	0.5	0.6
2002.....	5.1	1.9	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.6
2001.....	5.3	1.2	0.4	0.2	0.2	0.2	0.3	0.4	0.4	0.7
2000.....	5.3	1.6	0.4	0.3	0.2	0.2	0.3	0.3	0.5	0.7
1999.....	5.2	1.4	0.4	0.3	0.2	0.2	0.2	0.3	0.5	0.6

White, male

2006 4/.....	5.2	1.6	*	*	*	*	0.2	0.3	0.4	0.6
2005 4/.....	5.2	*	0.4	*	*	0.3	0.3	0.4	0.5	0.7
2004 4/.....	5.1	1.8	0.3	*	*	0.2	0.4	0.4	0.6	0.6
2003 4/.....	5.2	1.6	0.3	*	*	*	0.3	0.4	0.5	0.7
2002.....	5.1	1.7	0.3	*	0.2	*	0.2	0.3	0.5	0.7
2001.....	5.3	1.2	0.4	*	*	*	0.3	0.4	0.4	0.7
2000.....	5.2	1.6	0.4	0.3	*	*	0.4	0.3	0.5	0.7
1999.....	5.2	1.4	0.4	0.3	*	*	0.3	0.3	0.5	0.7

White, female

2006 4/.....	5.2	1.5	0.5	*	*	*	*	0.4	0.3	0.5
2005 4/.....	5.0	1.7	*	*	*	*	*	0.3	0.5	0.6
2004 4/.....	5.0	1.9	0.3	*	*	*	*	0.3	0.4	0.6
2003 4/.....	5.1	1.5	*	*	*	*	0.3	0.3	0.5	0.5
2002.....	5.1	2.1	0.4	0.3	*	*	*	*	0.4	0.5
2001.....	5.2	*	0.4	*	0.3	0.3	*	0.3	0.5	0.7
2000.....	5.4	1.6	*	0.3	*	*	*	0.3	0.4	0.6
1999.....	5.1	1.4	0.4	0.4	*	0.3	*	*	0.5	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Anemias (D50-D64)										
All races 3/, both sexes										
2006.....	1.3	*	0.2	*	*	0.2	0.3	0.4	0.3	0.4
2005.....	1.6	*	0.1	*	*	0.2	0.3	0.4	0.4	0.4
2004.....	1.6	*	0.2	0.1	0.1	0.2	0.4	0.4	0.4	0.4
2003.....	1.6	0.5	0.2	*	0.1	0.2	0.3	0.4	0.4	0.4
2002.....	1.6	*	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.5
2001.....	1.6	*	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.4
2000.....	1.6	*	0.2	0.1	0.1	0.1	0.3	0.3	0.4	0.4
1999.....	1.6	*	0.2	0.1	0.1	0.2	0.4	0.3	0.5	0.4
All races 3/, male										
2006.....	1.1	*	*	*	*	0.2	0.3	0.4	0.3	0.3
2005.....	1.2	*	*	*	*	0.2	0.3	0.4	0.4	0.4
2004.....	1.3	*	*	*	*	0.3	0.4	0.4	0.5	0.5
2003.....	1.3	*	*	*	*	*	0.3	0.4	0.4	0.3
2002.....	1.3	*	*	*	*	0.2	0.4	0.5	0.5	0.5
2001.....	1.3	*	*	*	*	0.2	0.3	0.4	0.5	0.4
2000.....	1.2	*	*	*	*	*	0.3	0.3	0.4	0.3
1999.....	1.3	*	*	*	*	0.2	0.5	0.3	0.5	0.4
All races 3/, female										
2006.....	1.6	*	*	*	*	*	0.2	0.3	0.4	0.5
2005.....	1.9	*	*	*	*	*	0.3	0.4	0.3	0.4
2004.....	1.8	*	*	*	*	*	0.4	0.5	0.3	0.3
2003.....	1.9	*	*	*	*	*	0.3	0.3	0.4	0.4
2002.....	1.9	*	*	*	*	*	0.3	0.3	0.5	0.4
2001.....	2.0	*	*	*	*	0.2	0.3	0.3	0.4	0.4
2000.....	2.0	*	*	*	*	*	0.3	0.3	0.5	0.4
1999.....	1.9	*	*	*	*	*	0.4	0.3	0.5	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Anemias (D50-D64)										
Black, both sexes										
2006 4/.....	2.3	*	*	*	*	0.6	1.3	1.9	2.1	2.2
2005 4/.....	2.6	*	*	*	*	0.9	1.5	2.3	2.4	2.2
2004 4/.....	2.7	*	*	*	*	0.9	2.0	2.2	2.4	2.4
2003 4/.....	2.7	*	*	*	*	*	1.9	1.9	2.2	2.2
2002.....	2.8	*	*	*	*	0.9	1.7	2.5	2.3	2.6
2001.....	2.7	*	*	*	*	0.9	1.5	1.9	2.5	2.1
2000.....	2.6	*	*	*	*	*	1.6	1.7	2.8	2.2
1999.....	2.8	*	0.8	*	*	0.7	2.2	2.1	3.0	2.2
Black, male										
2006 4/.....	2.1	*	*	*	*	*	1.5	2.1	2.0	1.9
2005 4/.....	2.4	*	*	*	*	*	1.7	2.4	2.7	2.7
2004 4/.....	2.6	*	*	*	*	1.2	1.9	2.3	3.0	3.0
2003 4/.....	2.7	*	*	*	*	*	1.8	2.2	2.3	2.2
2002.....	2.7	*	*	*	*	*	2.3	3.2	2.5	2.7
2001.....	2.5	*	*	*	*	*	1.6	2.5	3.3	2.2
2000.....	2.3	*	*	*	*	*	1.7	1.6	2.7	1.9
1999.....	2.7	*	*	*	*	*	2.8	2.3	3.3	2.5
Black, female										
2006 4/.....	2.6	*	*	*	*	*	*	1.8	2.1	2.4
2005 4/.....	2.8	*	*	*	*	*	1.3	2.1	2.2	1.8
2004 4/.....	2.8	*	*	*	*	*	2.1	2.2	1.9	1.9
2003 4/.....	2.8	*	*	*	*	*	2.1	1.7	2.2	2.2
2002.....	3.0	*	*	*	*	*	*	1.8	2.2	2.4
2001.....	2.8	*	*	*	*	*	1.5	*	1.8	2.1
2000.....	3.0	*	*	*	*	*	1.6	1.9	2.9	2.5
1999.....	3.0	*	*	*	*	*	1.7	1.8	2.8	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Diabetes mellitus (E10-E14)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	24.2	*	*	*	0.1	0.2	0.6	1.2	2.1	3.3
2005.....	25.3	*	*	*	0.2	0.3	0.6	1.0	2.0	3.6
2004.....	24.9	*	*	*	0.1	0.2	0.5	0.9	2.0	3.4
2003.....	25.5	*	*	*	0.1	0.2	0.5	1.2	2.1	3.3
2002.....	25.4	*	*	*	0.1	0.3	0.6	1.2	2.0	3.5
2001.....	25.1	*	*	*	0.2	0.2	0.5	1.1	1.8	3.3
2000.....	24.6	*	*	*	*	0.2	0.6	1.1	2.0	3.0
1999.....	24.5	*	*	*	0.1	0.3	0.6	1.0	1.9	3.2

All races 3/, male

2006.....	24.4	*	*	*	*	0.2	0.7	1.3	2.4	4.0
2005.....	25.0	*	*	*	*	0.4	0.7	1.1	2.5	4.3
2004.....	24.4	*	*	*	*	0.3	0.5	1.0	2.3	4.2
2003.....	24.8	*	*	*	*	0.3	0.6	1.2	2.5	3.9
2002.....	24.2	*	*	*	*	0.3	0.7	1.3	2.4	4.1
2001.....	23.5	*	*	*	0.2	0.3	0.6	1.2	2.2	3.9
2000.....	22.9	*	*	*	*	0.3	0.7	1.2	2.5	3.5
1999.....	22.8	*	*	*	*	0.2	0.6	1.0	2.2	3.8

All races 3/, female

2006.....	24.0	*	*	*	*	*	0.5	1.1	1.9	2.7
2005.....	25.7	*	*	*	0.2	0.3	0.6	0.9	1.5	2.9
2004.....	25.4	*	*	*	*	*	0.6	0.8	1.8	2.5
2003.....	26.2	*	*	*	*	*	0.5	1.1	1.7	2.8
2002.....	26.5	*	*	*	*	0.3	0.5	1.1	1.5	3.0
2001.....	26.6	*	*	*	*	0.2	0.5	1.1	1.5	2.7
2000.....	26.3	*	*	*	*	*	0.5	0.9	1.6	2.5
1999.....	26.2	*	*	*	*	0.3	0.5	1.0	1.6	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Diabetes mellitus (E10-E14)

White, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	23.6	*	*	*	*	0.2	0.5	0.9	1.7	2.8
2005 4/.....	24.9	*	*	*	0.1	0.2	0.5	0.8	1.6	3.0
2004 4/.....	24.4	*	*	*	*	0.2	0.4	0.8	1.7	2.9
2003 4/.....	25.0	*	*	*	*	0.2	0.5	0.9	1.8	2.8
2002.....	24.9	*	*	*	*	0.2	0.5	1.0	1.7	3.1
2001.....	24.6	*	*	*	*	0.2	0.4	0.8	1.6	2.8
2000.....	24.1	*	*	*	*	0.2	0.5	0.8	1.8	2.7
1999.....	23.9	*	*	*	*	0.2	0.5	0.9	1.6	2.8

White, male

2006 4/.....	24.2	*	*	*	*	*	0.5	0.9	1.8	3.3
2005 4/.....	24.9	*	*	*	*	0.3	0.6	0.9	1.9	3.5
2004 4/.....	24.3	*	*	*	*	*	0.4	0.8	2.0	3.6
2003 4/.....	24.8	*	*	*	*	*	0.6	0.8	2.2	3.4
2002.....	24.2	*	*	*	*	*	0.5	1.1	2.1	3.7
2001.....	23.5	*	*	*	*	*	0.4	0.8	1.8	3.3
2000.....	22.9	*	*	*	*	0.3	0.5	1.0	2.3	3.1
1999.....	22.7	*	*	*	*	0.3	0.5	0.9	1.8	3.4

White, female

2006 4/.....	23.0	*	*	*	*	*	0.4	0.8	1.6	2.3
2005 4/.....	24.9	*	*	*	*	*	0.4	0.7	1.2	2.5
2004 4/.....	24.5	*	*	*	*	*	0.4	0.7	1.5	2.1
2003 4/.....	25.2	*	*	*	*	*	0.4	1.0	1.4	2.3
2002.....	25.6	*	*	*	*	*	0.4	0.9	1.3	2.5
2001.....	25.7	*	*	*	*	*	0.3	0.8	1.4	2.3
2000.....	25.3	*	*	*	*	*	0.4	0.7	1.2	2.3
1999.....	25.0	*	*	*	*	*	0.4	0.9	1.4	2.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Diabetes mellitus (E10-E14)

Black, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	32.4	*	*	*	*	*	1.3	3.4	5.2	7.8
2005 4/.....	33.2	*	*	*	*	0.7	1.5	2.8	5.1	8.1
2004 4/.....	33.2	*	*	*	*	0.6	1.3	2.0	4.6	7.5
2003 4/.....	33.8	*	*	*	*	*	1.0	3.2	4.4	7.3
2002.....	33.6	*	*	*	*	*	1.4	2.8	4.3	7.3
2001.....	33.1	*	*	*	*	*	1.4	3.2	4.2	6.9
2000.....	32.8	*	*	*	*	*	1.5	2.8	4.1	6.0
1999.....	33.0	*	*	*	*	*	1.1	1.8	4.1	6.6

Black, male

2006 4/.....	30.6	*	*	*	*	*	1.5	3.6	6.7	9.8
2005 4/.....	30.7	*	*	*	*	*	1.4	2.9	6.7	10.8
2004 4/.....	30.2	*	*	*	*	*	*	2.5	5.4	9.5
2003 4/.....	29.8	*	*	*	*	*	*	4.1	5.2	8.4
2002.....	29.0	*	*	*	*	*	1.7	3.0	5.7	8.5
2001.....	28.5	*	*	*	*	*	1.5	3.9	5.6	8.7
2000.....	27.4	*	*	*	*	*	1.5	3.1	4.5	7.1
1999.....	27.7	*	*	*	*	*	*	1.6	5.0	7.7

Black, female

2006 4/.....	34.1	*	*	*	*	*	*	3.1	3.9	6.0
2005 4/.....	35.5	*	*	*	*	*	1.7	2.6	3.6	5.7
2004 4/.....	36.0	*	*	*	*	*	1.6	1.6	3.8	5.6
2003 4/.....	37.4	*	*	*	*	*	*	2.3	3.7	6.3
2002.....	37.8	*	*	*	*	*	*	2.6	3.1	6.3
2001.....	37.2	*	*	*	*	*	1.3	2.7	2.9	5.3
2000.....	37.8	*	*	*	*	*	1.5	2.5	3.8	5.0
1999.....	37.8	*	*	*	*	*	*	2.0	3.3	5.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Major cardiovascular diseases (I00-I78)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	275.1	12.3	1.3	0.7	1.1	2.2	4.3	7.2	13.2	23.4
2005.....	288.8	12.2	1.3	0.8	1.0	2.3	4.3	7.3	12.8	24.1
2004.....	293.3	13.8	1.5	0.6	1.0	2.2	4.2	7.3	12.6	24.3
2003.....	310.3	13.9	1.5	0.7	1.0	2.5	4.5	7.1	13.2	25.1
2002.....	318.6	15.9	1.4	0.6	1.1	2.4	3.9	7.1	12.6	25.3
2001.....	323.9	14.9	1.9	0.7	1.1	2.2	4.3	7.1	13.0	25.0
2000.....	332.9	16.7	1.5	0.7	1.1	2.4	4.3	6.7	12.2	24.2
1999.....	340.6	17.0	1.5	0.7	1.0	2.7	4.1	6.5	12.6	25.9

All races 3/, male

2006.....	267.7	13.4	1.4	0.8	1.2	2.6	5.4	9.5	17.6	31.5
2005.....	277.9	13.4	1.5	0.8	1.1	3.0	5.5	9.5	16.6	32.0
2004.....	281.2	14.7	1.5	0.5	1.1	2.8	5.3	9.2	16.3	32.1
2003.....	296.3	15.2	1.4	0.7	1.2	3.1	5.3	8.5	17.0	33.4
2002.....	303.3	16.6	1.5	0.6	1.3	3.1	5.0	9.0	16.1	32.9
2001.....	306.2	15.3	1.9	0.6	1.1	2.7	5.3	8.8	16.5	33.1
2000.....	315.8	17.5	1.6	0.7	1.2	2.8	5.0	8.2	15.3	31.3
1999.....	323.0	17.9	1.6	0.8	1.1	3.3	4.8	7.8	16.2	34.2

All races 3/, female

2006.....	282.4	11.2	1.3	0.6	0.9	1.7	3.1	4.8	8.8	15.3
2005.....	299.3	11.0	1.1	0.9	0.8	1.7	3.0	5.0	9.0	16.1
2004.....	304.9	12.9	1.5	0.7	1.0	1.6	3.0	5.3	8.9	16.4
2003.....	323.9	12.5	1.5	0.6	0.8	2.0	3.6	5.6	9.3	16.8
2002.....	333.3	15.1	1.3	0.7	0.9	1.7	2.8	5.1	9.0	17.7
2001.....	340.9	14.5	1.8	0.8	1.1	1.7	3.2	5.5	9.5	17.0
2000.....	349.5	15.9	1.4	0.6	1.0	2.0	3.5	5.2	9.0	17.1
1999.....	357.5	16.0	1.4	0.6	1.0	2.1	3.4	5.2	9.1	17.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Major cardiovascular diseases (I00-I78)										
White, both sexes										
2006 4/.....	292.2	9.8	1.2	0.6	1.0	1.9	3.8	5.9	10.9	19.6
2005 4/.....	307.0	9.9	1.1	0.7	0.8	2.0	3.6	5.9	10.5	20.3
2004 4/.....	311.7	11.9	1.2	0.5	0.9	1.8	3.6	5.8	10.4	21.1
2003 4/.....	330.0	11.0	1.2	0.6	0.9	2.2	3.7	5.7	11.0	21.4
2002.....	339.4	13.4	1.3	0.6	0.9	2.1	3.3	5.7	10.2	21.6
2001.....	344.8	12.2	1.5	0.6	1.0	1.9	3.4	5.9	10.9	21.4
2000.....	354.7	14.6	1.3	0.6	0.9	2.1	3.5	5.5	10.1	20.6
1999.....	362.7	14.6	1.3	0.7	0.9	2.4	3.4	5.4	10.6	22.0
White, male										
2006 4/.....	281.5	10.5	1.2	0.7	1.2	2.4	5.0	8.0	14.8	26.8
2005 4/.....	293.0	10.8	1.3	0.7	1.0	2.5	4.7	7.9	13.7	27.5
2004 4/.....	296.7	12.7	1.3	0.5	1.0	2.3	4.6	7.6	13.8	28.3
2003 4/.....	312.6	11.5	1.1	0.6	1.1	2.7	4.4	7.0	14.4	29.3
2002.....	320.9	14.3	1.5	0.5	1.1	2.7	4.3	7.5	13.3	28.5
2001.....	323.5	13.2	1.6	0.6	1.0	2.3	4.3	7.3	14.0	28.8
2000.....	334.2	15.3	1.6	0.7	1.1	2.4	4.2	6.9	13.2	27.6
1999.....	341.2	15.3	1.5	0.8	0.9	2.9	3.9	6.5	14.0	29.8
White, female										
2006 4/.....	302.7	9.1	1.2	0.5	0.7	1.4	2.6	3.8	6.7	12.2
2005 4/.....	320.8	9.0	0.9	0.7	0.7	1.5	2.5	3.7	7.1	13.0
2004 4/.....	326.5	11.0	1.1	0.6	0.8	1.3	2.5	3.9	7.0	13.7
2003 4/.....	347.0	10.4	1.4	0.5	0.7	1.7	2.8	4.3	7.4	13.2
2002.....	357.5	12.5	1.1	0.7	0.8	1.5	2.2	3.9	7.1	14.5
2001.....	365.5	11.1	1.5	0.7	0.9	1.5	2.5	4.3	7.6	13.9
2000.....	374.8	13.9	1.1	0.5	0.7	1.7	2.8	4.1	6.9	13.5
1999.....	383.6	13.8	1.2	0.6	0.9	1.9	2.9	4.2	7.2	14.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Major cardiovascular diseases (I00-I78)

Black, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	246.2	25.0	2.2	1.0	1.7	3.8	7.6	15.5	30.8	52.7
2005 4/.....	256.2	23.1	2.3	1.6	1.5	4.2	8.3	16.6	29.5	52.6
2004 4/.....	260.3	21.9	2.9	1.1	1.8	4.2	7.7	16.8	28.5	49.8
2003 4/.....	273.1	26.2	2.8	1.2	1.6	4.4	9.6	16.1	29.3	52.7
2002.....	276.1	27.6	2.0	0.8	2.0	4.0	7.8	16.6	29.6	52.9
2001.....	280.4	28.6	3.5	1.0	1.6	4.0	9.4	16.4	28.5	51.5
2000.....	285.0	27.8	2.5	1.0	1.9	4.6	9.1	15.1	27.7	51.2
1999.....	289.4	29.2	2.4	1.0	1.8	4.8	9.0	14.4	26.8	55.2

Black, male

2006 4/.....	250.3	28.2	2.6	*	1.6	4.3	8.7	20.4	39.7	69.8
2005 4/.....	255.0	26.3	2.2	1.4	1.7	5.6	10.6	20.6	38.0	69.2
2004 4/.....	257.2	23.7	2.8	*	1.8	5.6	9.8	20.3	35.2	65.4
2003 4/.....	268.6	31.2	3.3	1.4	1.7	5.6	11.6	19.5	36.3	65.7
2002.....	268.5	28.2	1.9	*	2.3	5.1	9.4	20.6	36.9	66.8
2001.....	272.4	28.1	3.6	*	1.5	5.1	11.3	20.3	35.2	65.6
2000.....	275.6	29.1	2.1	*	1.8	5.1	10.2	18.0	32.7	61.6
1999.....	282.5	30.6	2.5	*	2.1	6.3	10.9	17.7	33.0	69.6

Black, female

2006 4/.....	242.5	21.7	1.7	*	1.7	3.3	6.4	10.9	22.7	37.4
2005 4/.....	257.2	19.7	2.3	1.7	1.4	2.6	6.1	12.9	22.0	37.8
2004 4/.....	263.2	20.1	3.0	1.3	1.8	2.7	5.5	13.5	22.5	35.7
2003 4/.....	277.2	21.0	2.4	*	1.5	3.1	7.6	13.0	23.0	41.0
2002.....	283.0	26.9	2.1	*	1.6	2.8	6.3	13.0	23.0	40.6
2001.....	287.8	29.0	3.4	1.3	1.6	2.8	7.5	12.9	22.4	38.9
2000.....	293.4	26.5	2.9	*	2.0	4.2	8.1	12.4	23.3	42.0
1999.....	295.7	27.9	2.4	*	1.5	3.3	7.1	11.5	21.3	42.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Diseases of heart (I00-I09,I11,I13,I20-I51)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	211.0	8.4	1.0	0.5	0.8	1.7	3.4	5.8	10.6	18.6
2005.....	220.0	8.7	0.9	0.5	0.7	1.8	3.5	5.9	10.3	19.4
2004.....	222.2	10.3	1.2	0.4	0.8	1.8	3.2	5.7	10.0	19.4
2003.....	235.6	11.0	1.2	0.5	0.8	1.9	3.6	5.6	10.5	20.1
2002.....	241.7	12.4	1.1	0.5	0.8	2.0	3.1	5.7	10.0	20.3
2001.....	245.8	11.9	1.5	0.5	0.8	1.7	3.3	5.6	10.2	19.9
2000.....	252.6	13.0	1.2	0.5	0.8	2.0	3.3	5.2	9.5	19.1
1999.....	259.9	13.8	1.2	0.6	0.8	2.3	3.3	5.1	10.1	20.6

All races 3/, male

2006.....	214.0	8.8	1.1	0.5	0.9	2.2	4.4	7.9	14.7	25.9
2005.....	221.1	9.4	1.0	0.5	0.8	2.5	4.6	7.9	13.7	26.8
2004.....	222.8	10.9	1.1	0.3	0.8	2.3	4.2	7.4	13.6	26.5
2003.....	235.0	12.1	1.1	0.6	0.9	2.4	4.4	6.9	13.9	27.8
2002.....	240.7	12.9	1.1	0.4	0.9	2.6	4.0	7.4	13.4	27.8
2001.....	242.5	11.8	1.5	0.4	0.9	2.1	4.2	7.0	13.4	27.4
2000.....	249.8	13.3	1.4	0.6	0.9	2.4	4.0	6.5	12.4	26.0
1999.....	257.0	13.9	1.3	0.6	0.9	2.9	3.9	6.3	13.2	28.5

All races 3/, female

2006.....	208.0	7.9	0.9	0.4	0.7	1.2	2.4	3.7	6.5	11.3
2005.....	218.9	8.0	0.9	0.6	0.6	1.2	2.2	3.7	6.8	12.0
2004.....	221.6	9.7	1.2	0.5	0.7	1.2	2.2	3.9	6.4	12.3
2003.....	236.2	9.8	1.3	0.5	0.6	1.4	2.7	4.2	7.1	12.5
2002.....	242.7	11.8	1.0	0.5	0.6	1.3	2.1	3.9	6.5	12.8
2001.....	249.0	12.0	1.4	0.6	0.8	1.3	2.4	4.1	6.9	12.4
2000.....	255.3	12.5	1.0	0.4	0.7	1.6	2.6	3.9	6.5	12.2
1999.....	262.6	13.7	1.1	0.5	0.7	1.7	2.6	3.9	6.9	12.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Diseases of heart (I00-I09,I11,I13,I20-I51)

White, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	225.5	6.6	0.9	0.4	0.7	1.4	3.0	4.8	8.8	15.9
2005 4/.....	235.2	7.1	0.7	0.4	0.6	1.6	2.9	4.8	8.4	16.7
2004 4/.....	237.4	9.0	0.9	0.4	0.6	1.4	2.7	4.5	8.4	17.2
2003 4/.....	251.7	8.8	1.0	0.4	0.7	1.6	2.9	4.5	8.8	17.7
2002.....	258.5	10.6	0.9	0.4	0.7	1.7	2.5	4.6	8.1	17.7
2001.....	262.8	9.9	1.2	0.4	0.7	1.5	2.6	4.5	8.5	17.3
2000.....	270.2	11.5	1.1	0.4	0.7	1.7	2.6	4.2	7.9	16.7
1999.....	277.7	11.8	1.0	0.5	0.7	2.0	2.6	4.1	8.5	17.8

White, male

2006 4/.....	226.9	7.0	0.9	0.4	0.8	1.9	4.0	6.6	12.5	22.6
2005 4/.....	234.9	7.6	0.9	0.4	0.7	2.1	3.9	6.8	11.4	23.4
2004 4/.....	236.5	9.6	1.0	0.3	0.7	1.8	3.6	6.0	11.6	23.8
2003 4/.....	249.5	9.4	0.8	0.4	0.8	2.0	3.5	5.7	12.0	25.1
2002.....	256.0	11.3	1.0	0.4	0.8	2.2	3.3	6.1	11.2	24.6
2001.....	257.8	10.6	1.2	0.4	0.8	1.8	3.4	5.9	11.3	24.4
2000.....	265.8	11.9	1.3	0.5	0.8	2.0	3.3	5.3	10.7	23.5
1999.....	272.9	12.0	1.2	0.6	0.7	2.4	3.0	5.2	11.4	25.2

White, female

2006 4/.....	224.2	6.2	0.8	0.4	0.6	0.9	1.9	2.9	5.0	9.1
2005 4/.....	235.5	6.6	0.6	0.4	0.5	1.1	1.8	2.7	5.3	9.7
2004 4/.....	238.3	8.4	0.9	0.4	0.6	1.0	1.7	2.9	5.0	10.3
2003 4/.....	253.8	8.1	1.1	0.4	0.5	1.3	2.2	3.2	5.4	10.2
2002.....	261.0	9.9	0.8	0.5	0.5	1.1	1.6	3.0	5.0	10.7
2001.....	267.7	9.2	1.1	0.5	0.7	1.1	1.8	3.1	5.6	10.0
2000.....	274.5	11.1	0.8	0.4	0.5	1.3	1.9	2.9	5.0	9.8
1999.....	282.4	11.6	0.9	0.5	0.6	1.6	2.1	3.0	5.4	10.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of heart (I00-I09,I11,I13,I20-I51)										
Black, both sexes										
2006 4/.....	182.6	17.1	1.7	0.7	1.3	3.2	6.1	12.7	24.4	40.8
2005 4/.....	189.8	16.1	1.7	1.1	1.2	3.4	6.9	13.1	23.7	41.5
2004 4/.....	192.3	15.3	2.4	0.9	1.5	3.6	6.3	13.3	22.1	38.1
2003 4/.....	202.8	20.2	2.2	1.0	1.2	3.5	7.8	13.0	23.0	39.6
2002.....	205.6	21.3	1.6	*	1.3	3.4	6.5	13.4	23.2	40.8
2001.....	208.8	21.6	3.0	0.8	1.3	3.3	7.5	12.9	22.6	39.8
2000.....	211.8	19.8	2.0	0.9	1.4	4.0	7.4	12.8	21.5	38.2
1999.....	217.2	23.8	2.0	0.9	1.4	4.2	7.7	11.7	21.5	42.7
Black, male										
2006 4/.....	191.8	17.9	2.0	*	1.5	3.7	7.2	16.9	32.8	54.9
2005 4/.....	194.8	18.3	*	*	1.2	4.9	9.0	16.5	31.1	56.1
2004 4/.....	196.7	16.3	2.4	*	1.4	4.8	8.0	16.7	28.4	50.9
2003 4/.....	206.0	24.1	2.5	*	1.3	4.5	10.2	15.7	28.6	51.1
2002.....	206.3	22.1	*	*	1.5	4.5	8.0	17.5	30.0	54.4
2001.....	209.0	19.5	3.2	*	1.4	4.4	9.7	16.0	29.3	52.1
2000.....	211.1	19.8	1.9	*	1.4	4.8	8.7	15.7	26.5	48.6
1999.....	218.5	23.4	2.1	*	1.6	5.8	9.3	14.8	27.1	56.3
Black, female										
2006 4/.....	174.3	16.2	*	*	1.2	2.7	5.0	8.8	16.8	28.2
2005 4/.....	185.2	13.9	2.0	1.4	1.2	1.9	4.7	9.8	17.0	28.4
2004 4/.....	188.3	14.2	2.3	*	1.5	2.3	4.5	10.1	16.4	26.7
2003 4/.....	200.0	16.1	2.0	*	*	2.3	5.5	10.5	18.1	29.3
2002.....	205.0	20.6	1.8	*	*	2.4	5.0	9.7	17.0	28.6
2001.....	208.6	23.9	2.7	*	1.2	2.3	5.5	10.1	16.7	28.9
2000.....	212.6	19.8	2.0	*	1.3	3.3	6.3	10.2	16.9	29.0
1999.....	216.0	24.2	1.9	*	*	2.5	6.1	9.0	16.5	30.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
All races 3/, both sexes										
2006.....	9.9	*	*	*	*	*	0.2	0.5	0.9	2.2
2005.....	9.9	*	*	*	*	*	0.2	0.6	1.1	2.2
2004.....	9.7	*	*	*	*	*	0.1	0.5	1.0	1.9
2003.....	9.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2002.....	9.2	*	*	*	*	*	0.1	0.4	1.0	1.8
2001.....	8.7	*	*	*	*	*	*	0.3	0.9	1.6
2000.....	8.4	*	*	*	*	*	0.1	0.4	0.7	1.4
1999.....	8.1	*	*	*	*	*	0.1	0.3	0.8	1.5
All races 3/, male										
2006.....	9.3	*	*	*	*	*	0.3	0.7	1.3	3.2
2005.....	9.1	*	*	*	*	*	0.2	0.7	1.4	3.2
2004.....	8.9	*	*	*	*	*	*	0.7	1.5	2.8
2003.....	8.8	*	*	*	*	*	0.3	0.7	1.4	2.7
2002.....	8.2	*	*	*	*	*	0.2	0.7	1.4	2.6
2001.....	7.5	*	*	*	*	*	*	0.4	1.2	2.1
2000.....	7.2	*	*	*	*	*	0.2	0.4	0.9	1.9
1999.....	6.8	*	*	*	*	*	*	0.4	1.0	2.0
All races 3/, female										
2006.....	10.6	*	*	*	*	*	*	0.3	0.5	1.1
2005.....	10.7	*	*	*	*	*	*	0.4	0.7	1.2
2004.....	10.5	*	*	*	*	*	*	0.3	0.5	1.0
2003.....	10.7	*	*	*	*	*	*	0.3	0.7	1.2
2002.....	10.2	*	*	*	*	*	*	*	0.6	1.0
2001.....	9.8	*	*	*	*	*	*	0.3	0.5	1.1
2000.....	9.7	*	*	*	*	*	*	0.3	0.5	1.0
1999.....	9.4	*	*	*	*	*	*	*	0.5	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
Black, both sexes										
2006 4/.....	17.3	*	*	*	*	*	*	1.7	3.3	6.4
2005 4/.....	17.7	*	*	*	*	*	0.7	1.6	3.4	6.7
2004 4/.....	17.6	*	*	*	*	*	*	2.1	3.2	6.1
2003 4/.....	17.9	*	*	*	*	*	0.6	1.5	3.5	6.0
2002.....	17.2	*	*	*	*	*	*	1.5	3.9	5.8
2001.....	16.0	*	*	*	*	*	*	1.1	3.1	5.1
2000.....	16.1	*	*	*	*	*	*	1.4	2.8	4.9
1999.....	15.5	*	*	*	*	*	*	1.0	2.6	5.7
Black, male										
2006 4/.....	18.8	*	*	*	*	*	*	2.6	4.6	9.0
2005 4/.....	18.5	*	*	*	*	*	*	1.7	4.2	9.2
2004 4/.....	18.4	*	*	*	*	*	*	2.5	4.6	8.8
2003 4/.....	18.1	*	*	*	*	*	*	1.7	4.3	7.3
2002.....	17.5	*	*	*	*	*	*	2.4	5.4	8.5
2001.....	15.9	*	*	*	*	*	*	*	4.0	5.9
2000.....	15.6	*	*	*	*	*	*	1.6	3.5	5.8
1999.....	15.2	*	*	*	*	*	*	*	2.8	7.3
Black, female										
2006 4/.....	15.9	*	*	*	*	*	*	*	2.1	4.0
2005 4/.....	17.0	*	*	*	*	*	*	1.6	2.7	4.4
2004 4/.....	16.9	*	*	*	*	*	*	1.8	2.1	3.6
2003 4/.....	17.8	*	*	*	*	*	*	*	2.7	4.8
2002.....	16.9	*	*	*	*	*	*	*	2.6	3.4
2001.....	16.2	*	*	*	*	*	*	*	2.3	4.4
2000.....	16.5	*	*	*	*	*	*	*	2.1	4.1
1999.....	15.9	*	*	*	*	*	*	*	2.5	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Ischemic heart diseases (I20-I25)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	142.1	0.5	*	*	*	0.2	0.5	1.4	3.9	8.8
2005.....	150.4	*	*	*	*	0.1	0.6	1.4	3.6	9.1
2004.....	153.7	*	*	*	*	0.1	0.5	1.5	3.9	9.4
2003.....	165.1	0.5	*	*	*	0.2	0.6	1.4	3.9	10.1
2002.....	171.4	0.6	*	*	*	0.1	0.5	1.3	3.6	10.1
2001.....	176.3	0.6	*	*	*	0.2	0.7	1.5	4.0	9.9
2000.....	183.1	0.8	*	*	*	0.2	0.5	1.2	3.4	9.8
1999.....	189.8	0.7	*	*	*	0.3	0.5	1.3	3.7	10.4

All races 3/, male

2006.....	152.2	*	*	*	*	0.3	0.7	2.1	6.1	13.2
2005.....	159.0	*	*	*	*	0.2	0.9	2.2	5.4	13.6
2004.....	161.6	*	*	*	*	0.2	0.7	2.1	6.0	13.9
2003.....	172.2	*	*	*	*	0.2	0.8	2.0	5.9	15.2
2002.....	178.4	*	*	*	*	*	0.7	2.0	5.2	15.2
2001.....	181.7	*	*	*	*	0.2	0.9	2.2	5.9	15.0
2000.....	188.7	*	*	*	*	0.2	0.6	1.8	5.1	14.7
1999.....	195.4	*	*	*	*	0.3	0.6	1.7	5.5	15.8

All races 3/, female

2006.....	132.3	*	*	*	*	*	0.2	0.6	1.7	4.4
2005.....	142.0	*	*	*	*	*	0.2	0.6	1.9	4.6
2004.....	146.1	*	*	*	*	*	0.2	0.9	1.8	4.9
2003.....	158.1	*	*	*	*	*	0.4	0.8	1.9	4.9
2002.....	164.7	*	*	*	*	*	0.3	0.6	1.9	4.9
2001.....	171.2	*	*	*	*	*	0.4	0.7	2.1	4.9
2000.....	177.6	*	*	*	*	*	0.3	0.6	1.7	5.0
1999.....	184.5	*	*	*	*	*	0.3	0.8	1.9	5.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Ischemic heart diseases (I20-I25)

White, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	153.4	*	*	*	*	0.2	0.5	1.3	3.7	8.3
2005 4/.....	162.6	*	*	*	*	0.1	0.5	1.2	3.5	8.5
2004 4/.....	166.1	*	*	*	*	*	0.4	1.3	3.6	9.2
2003 4/.....	178.3	*	*	*	*	0.2	0.5	1.3	3.7	9.8
2002.....	185.4	*	*	*	*	0.1	0.4	1.2	3.3	9.6
2001.....	190.5	*	*	*	*	0.1	0.5	1.3	3.8	9.5
2000.....	197.9	0.8	*	*	*	0.2	0.4	1.0	3.1	9.4
1999.....	205.1	*	*	*	*	0.2	0.3	1.2	3.5	9.9

White, male

2006 4/.....	163.7	*	*	*	*	0.3	0.7	2.0	5.7	12.6
2005 4/.....	171.5	*	*	*	*	*	0.8	1.9	5.0	12.7
2004 4/.....	174.3	*	*	*	*	*	0.5	1.9	5.6	13.6
2003 4/.....	185.5	*	*	*	*	*	0.7	1.8	5.6	15.0
2002.....	192.5	*	*	*	*	*	0.6	1.9	5.0	14.6
2001.....	195.8	*	*	*	*	*	0.8	2.0	5.5	14.5
2000.....	203.6	*	*	*	*	*	0.6	1.6	4.8	14.3
1999.....	210.5	*	*	*	*	0.3	0.5	1.6	5.1	15.1

White, female

2006 4/.....	143.3	*	*	*	*	*	*	0.5	1.5	3.8
2005 4/.....	153.9	*	*	*	*	*	*	0.5	1.8	4.1
2004 4/.....	158.1	*	*	*	*	*	*	0.7	1.6	4.7
2003 4/.....	171.4	*	*	*	*	*	0.4	0.7	1.7	4.4
2002.....	178.4	*	*	*	*	*	*	0.5	1.6	4.5
2001.....	185.3	*	*	*	*	*	0.3	0.6	1.9	4.5
2000.....	192.4	*	*	*	*	*	0.3	0.5	1.4	4.5
1999.....	200.0	*	*	*	*	*	*	0.7	1.7	4.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Ischemic heart diseases (I20-I25)

Black, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	112.6	*	*	*	*	*	0.7	2.3	6.2	14.7
2005 4/.....	117.8	*	*	*	*	*	0.9	2.6	5.3	15.5
2004 4/.....	121.0	*	*	*	*	*	1.2	2.9	6.5	12.7
2003 4/.....	129.2	*	*	*	*	*	1.2	2.4	6.1	14.0
2002.....	132.9	*	*	*	*	*	0.9	2.1	5.6	15.1
2001.....	136.9	*	*	*	*	*	1.5	2.8	6.4	14.9
2000.....	140.1	*	*	*	*	*	0.7	2.4	6.0	14.6
1999.....	143.7	*	*	*	*	*	1.2	1.9	5.8	16.1

Black, male

2006 4/.....	120.2	*	*	*	*	*	*	3.2	9.3	20.7
2005 4/.....	122.9	*	*	*	*	*	1.3	4.2	7.9	23.0
2004 4/.....	125.2	*	*	*	*	*	1.8	3.4	9.6	18.4
2003 4/.....	133.4	*	*	*	*	*	1.6	3.1	9.2	19.6
2002.....	135.3	*	*	*	*	*	*	2.7	7.5	22.0
2001.....	138.9	*	*	*	*	*	1.9	3.9	9.3	21.6
2000.....	141.5	*	*	*	*	*	*	3.7	8.0	20.3
1999.....	146.0	*	*	*	*	*	*	2.9	8.4	23.3

Black, female

2006 4/.....	105.6	*	*	*	*	*	*	1.4	3.3	9.4
2005 4/.....	113.1	*	*	*	*	*	*	*	3.0	8.8
2004 4/.....	117.1	*	*	*	*	*	*	2.4	3.7	7.6
2003 4/.....	125.3	*	*	*	*	*	*	1.7	3.3	9.1
2002.....	130.8	*	*	*	*	*	*	1.5	4.0	9.0
2001.....	135.2	*	*	*	*	*	*	1.7	3.8	9.0
2000.....	138.8	*	*	*	*	*	*	*	4.3	9.7
1999.....	141.7	*	*	*	*	*	*	*	3.4	9.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
All races 3/, both sexes										
2006.....	47.2	*	*	*	*	0.1	0.2	0.6	1.5	3.6
2005.....	50.9	*	*	*	*	*	0.3	0.6	1.3	3.5
2004.....	53.4	*	*	*	*	*	0.2	0.6	1.6	3.8
2003.....	58.7	*	*	*	*	*	0.3	0.5	1.5	4.1
2002.....	62.3	*	*	*	*	*	0.2	0.5	1.5	4.0
2001.....	64.9	*	*	*	*	*	0.2	0.6	1.8	4.2
2000.....	68.5	0.5	*	*	*	*	0.2	0.6	1.3	4.0
1999.....	71.5	*	*	*	*	0.1	0.2	0.5	1.6	4.1
All races 3/, male										
2006.....	51.6	*	*	*	*	*	0.2	0.9	2.3	5.2
2005.....	54.8	*	*	*	*	*	0.4	1.0	2.0	5.1
2004.....	57.4	*	*	*	*	*	0.3	0.8	2.3	5.7
2003.....	62.6	*	*	*	*	*	0.3	0.8	2.2	6.2
2002.....	66.2	*	*	*	*	*	0.3	0.7	2.1	6.0
2001.....	68.6	*	*	*	*	*	0.3	0.7	2.5	6.2
2000.....	72.7	*	*	*	*	*	0.2	0.8	1.9	5.8
1999.....	76.1	*	*	*	*	*	0.3	0.7	2.3	6.2
All races 3/, female										
2006.....	43.0	*	*	*	*	*	*	0.3	0.8	1.9
2005.....	47.2	*	*	*	*	*	*	0.3	0.7	2.0
2004.....	49.6	*	*	*	*	*	*	0.4	0.8	1.9
2003.....	54.8	*	*	*	*	*	0.2	0.3	0.8	2.0
2002.....	58.4	*	*	*	*	*	*	0.3	0.8	2.1
2001.....	61.3	*	*	*	*	*	*	0.4	1.0	2.1
2000.....	64.6	*	*	*	*	*	*	0.3	0.7	2.2
1999.....	67.0	*	*	*	*	*	*	0.3	0.9	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
White, both sexes										
2006 4/.....	51.0	*	*	*	*	*	0.2	0.6	1.5	3.5
2005 4/.....	55.1	*	*	*	*	*	0.2	0.5	1.3	3.4
2004 4/.....	57.8	*	*	*	*	*	0.1	0.5	1.6	3.8
2003 4/.....	63.4	*	*	*	*	*	0.2	0.5	1.4	4.1
2002.....	67.3	*	*	*	*	*	0.2	0.4	1.4	4.0
2001.....	70.1	*	*	*	*	*	0.2	0.5	1.7	4.1
2000.....	74.1	*	*	*	*	*	0.2	0.5	1.2	4.0
1999.....	77.4	*	*	*	*	*	0.1	0.5	1.5	4.0
White, male										
2006 4/.....	55.8	*	*	*	*	*	0.3	0.8	2.2	5.3
2005 4/.....	59.5	*	*	*	*	*	0.4	0.8	1.9	4.9
2004 4/.....	62.3	*	*	*	*	*	*	0.7	2.4	5.6
2003 4/.....	67.7	*	*	*	*	*	0.3	0.7	2.0	6.3
2002.....	71.8	*	*	*	*	*	0.3	0.6	2.1	5.9
2001.....	74.4	*	*	*	*	*	0.3	0.6	2.5	6.1
2000.....	78.8	*	*	*	*	*	*	0.6	1.8	6.0
1999.....	82.7	*	*	*	*	*	*	0.6	2.1	6.2
White, female										
2006 4/.....	46.3	*	*	*	*	*	*	0.3	0.7	1.7
2005 4/.....	50.8	*	*	*	*	*	*	*	0.7	1.8
2004 4/.....	53.3	*	*	*	*	*	*	0.4	0.7	1.9
2003 4/.....	59.2	*	*	*	*	*	*	*	0.8	1.9
2002.....	62.8	*	*	*	*	*	*	*	0.7	1.9
2001.....	66.0	*	*	*	*	*	*	0.3	0.9	2.0
2000.....	69.6	*	*	*	*	*	*	0.3	0.6	2.0
1999.....	72.3	*	*	*	*	*	*	0.3	0.9	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
Black, both sexes										
2006 4/.....	37.4	*	*	*	*	*	*	1.0	2.4	5.2
2005 4/.....	39.8	*	*	*	*	*	*	1.4	2.0	5.4
2004 4/.....	42.0	*	*	*	*	*	*	1.2	2.2	4.9
2003 4/.....	45.7	*	*	*	*	*	*	0.9	2.3	5.6
2002.....	48.5	*	*	*	*	*	*	0.9	2.3	5.5
2001.....	50.0	*	*	*	*	*	*	1.1	2.5	5.8
2000.....	52.2	*	*	*	*	*	*	1.2	2.3	4.9
1999.....	53.3	*	*	*	*	*	*	0.8	2.4	5.5
Black, male										
2006 4/.....	39.1	*	*	*	*	*	*	1.4	3.4	6.9
2005 4/.....	40.3	*	*	*	*	*	*	2.1	2.9	7.5
2004 4/.....	42.4	*	*	*	*	*	*	*	2.7	7.7
2003 4/.....	46.6	*	*	*	*	*	*	*	3.5	7.5
2002.....	48.3	*	*	*	*	*	*	*	3.0	7.6
2001.....	49.2	*	*	*	*	*	*	*	3.4	8.0
2000.....	52.0	*	*	*	*	*	*	1.6	2.7	6.5
1999.....	53.4	*	*	*	*	*	*	*	3.4	7.7
Black, female										
2006 4/.....	35.7	*	*	*	*	*	*	*	1.4	3.7
2005 4/.....	39.2	*	*	*	*	*	*	*	*	3.6
2004 4/.....	41.6	*	*	*	*	*	*	*	1.8	2.4
2003 4/.....	44.9	*	*	*	*	*	*	*	*	3.9
2002.....	48.8	*	*	*	*	*	*	*	1.6	3.6
2001.....	50.6	*	*	*	*	*	*	*	1.7	3.8
2000.....	52.5	*	*	*	*	*	*	*	1.9	3.6
1999.....	53.2	*	*	*	*	*	*	*	1.4	3.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other forms of chronic ischemic heart disease (I20,I25)										
All races 3/, both sexes										
2006.....	93.5	*	*	*	*	*	0.3	0.8	2.3	5.1
2005.....	98.2	*	*	*	*	*	0.3	0.8	2.3	5.5
2004.....	99.5	*	*	*	*	*	0.3	0.9	2.3	5.5
2003.....	105.3	*	*	*	*	*	0.3	0.9	2.4	5.8
2002.....	108.0	*	*	*	*	*	0.3	0.8	2.0	6.0
2001.....	110.3	*	*	*	*	*	0.4	0.9	2.2	5.6
2000.....	113.3	*	*	*	*	*	0.3	0.6	2.0	5.7
1999.....	117.2	*	*	*	*	0.1	0.3	0.7	2.1	6.2
All races 3/, male										
2006.....	99.2	*	*	*	*	*	0.4	1.2	3.7	7.9
2005.....	102.9	*	*	*	*	*	0.5	1.2	3.4	8.4
2004.....	103.3	*	*	*	*	*	0.4	1.3	3.6	8.1
2003.....	108.5	*	*	*	*	*	0.4	1.2	3.7	8.8
2002.....	110.9	*	*	*	*	*	0.4	1.3	3.0	9.2
2001.....	111.9	*	*	*	*	*	0.5	1.5	3.3	8.6
2000.....	114.8	*	*	*	*	*	0.4	1.1	3.1	8.7
1999.....	118.0	*	*	*	*	0.2	0.3	1.0	3.2	9.4
All races 3/, female										
2006.....	88.0	*	*	*	*	*	*	0.3	0.9	2.3
2005.....	93.7	*	*	*	*	*	*	0.3	1.2	2.5
2004.....	95.7	*	*	*	*	*	*	0.5	0.9	3.0
2003.....	102.3	*	*	*	*	*	0.2	0.5	1.0	2.8
2002.....	105.2	*	*	*	*	*	*	0.4	1.0	2.8
2001.....	108.8	*	*	*	*	*	0.2	0.3	1.1	2.7
2000.....	111.9	*	*	*	*	*	*	0.2	1.0	2.8
1999.....	116.4	*	*	*	*	*	*	0.4	1.0	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other forms of chronic ischemic heart disease (I20,I25)										
White, both sexes										
2006 4/.....	101.0	*	*	*	*	*	0.3	0.7	2.1	4.7
2005 4/.....	106.2	*	*	*	*	*	0.3	0.7	2.2	5.0
2004 4/.....	107.5	*	*	*	*	*	0.2	0.8	2.0	5.4
2003 4/.....	113.8	*	*	*	*	*	0.3	0.8	2.2	5.6
2002.....	116.9	*	*	*	*	*	0.2	0.8	1.8	5.6
2001.....	119.1	*	*	*	*	*	0.3	0.8	2.0	5.3
2000.....	122.6	*	*	*	*	*	0.2	0.6	1.9	5.3
1999.....	126.5	*	*	*	*	*	0.2	0.7	1.9	5.8
White, male										
2006 4/.....	106.5	*	*	*	*	*	0.4	1.1	3.4	7.3
2005 4/.....	110.7	*	*	*	*	*	0.4	1.1	3.1	7.6
2004 4/.....	111.1	*	*	*	*	*	0.3	1.2	3.1	7.9
2003 4/.....	116.6	*	*	*	*	*	0.4	1.1	3.5	8.5
2002.....	119.4	*	*	*	*	*	0.4	1.3	2.8	8.6
2001.....	120.1	*	*	*	*	*	0.4	1.4	3.0	8.2
2000.....	123.5	*	*	*	*	*	0.3	0.9	2.9	8.2
1999.....	126.5	*	*	*	*	*	*	0.9	3.0	8.8
White, female										
2006 4/.....	95.6	*	*	*	*	*	*	*	0.8	2.0
2005 4/.....	101.9	*	*	*	*	*	*	0.3	1.1	2.3
2004 4/.....	103.9	*	*	*	*	*	*	0.3	0.9	2.8
2003 4/.....	111.0	*	*	*	*	*	*	0.4	0.9	2.5
2002.....	114.4	*	*	*	*	*	*	0.3	0.8	2.5
2001.....	118.1	*	*	*	*	*	*	*	0.9	2.4
2000.....	121.6	*	*	*	*	*	*	*	0.8	2.4
1999.....	126.5	*	*	*	*	*	*	0.4	0.9	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other forms of chronic ischemic heart disease (I20,I25)										
Black, both sexes										
2006 4/.....	73.9	*	*	*	*	*	*	1.3	3.6	9.0
2005 4/.....	76.6	*	*	*	*	*	*	1.1	3.3	9.8
2004 4/.....	78.0	*	*	*	*	*	*	1.6	4.1	7.7
2003 4/.....	82.3	*	*	*	*	*	*	1.5	3.7	8.2
2002.....	83.2	*	*	*	*	*	*	1.1	3.2	9.6
2001.....	85.8	*	*	*	*	*	0.8	1.7	3.8	8.9
2000.....	86.6	*	*	*	*	*	*	1.3	3.6	9.5
1999.....	89.3	*	*	*	*	*	0.7	1.1	3.4	10.1
Black, male										
2006 4/.....	79.6	*	*	*	*	*	*	1.8	5.7	13.2
2005 4/.....	81.0	*	*	*	*	*	*	2.0	4.9	15.3
2004 4/.....	81.5	*	*	*	*	*	*	1.9	6.7	10.5
2003 4/.....	85.5	*	*	*	*	*	*	2.1	5.7	11.7
2002.....	85.7	*	*	*	*	*	*	*	4.1	14.4
2001.....	88.2	*	*	*	*	*	*	2.4	5.9	13.3
2000.....	88.0	*	*	*	*	*	*	2.1	5.0	13.4
1999.....	91.1	*	*	*	*	*	*	1.7	5.0	14.9
Black, female										
2006 4/.....	68.7	*	*	*	*	*	*	*	1.8	5.2
2005 4/.....	72.6	*	*	*	*	*	*	*	1.8	4.9
2004 4/.....	74.7	*	*	*	*	*	*	*	1.7	5.2
2003 4/.....	79.4	*	*	*	*	*	*	*	2.0	5.0
2002.....	81.0	*	*	*	*	*	*	*	2.3	5.3
2001.....	83.5	*	*	*	*	*	*	*	1.9	4.9
2000.....	85.3	*	*	*	*	*	*	*	2.4	5.9
1999.....	87.6	*	*	*	*	*	*	*	1.9	5.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Atherosclerotic cardiovascular disease, so described (I25.0)										
All races 3/, both sexes										
2006.....	20.4	*	*	*	*	*	*	0.2	0.8	1.9
2005.....	21.2	*	*	*	*	*	*	0.2	0.7	1.9
2004.....	21.7	*	*	*	*	*	*	0.2	0.8	1.7
2003.....	23.1	*	*	*	*	*	*	0.3	0.8	1.9
2002.....	23.6	*	*	*	*	*	*	0.2	0.7	1.9
2001.....	23.9	*	*	*	*	*	*	0.2	0.7	1.8
2000.....	24.7	*	*	*	*	*	*	0.2	0.5	1.7
1999.....	25.8	*	*	*	*	*	*	0.2	0.6	2.0
All races 3/, male										
2006.....	23.2	*	*	*	*	*	*	0.4	1.4	3.1
2005.....	23.6	*	*	*	*	*	*	0.4	1.1	3.0
2004.....	23.9	*	*	*	*	*	*	0.3	1.2	2.5
2003.....	25.2	*	*	*	*	*	*	0.4	1.3	2.9
2002.....	25.4	*	*	*	*	*	*	0.3	1.1	3.0
2001.....	25.3	*	*	*	*	*	*	0.3	1.0	2.8
2000.....	26.0	*	*	*	*	*	*	0.3	0.9	2.8
1999.....	26.8	*	*	*	*	*	*	0.2	0.9	3.1
All races 3/, female										
2006.....	17.7	*	*	*	*	*	*	*	0.3	0.7
2005.....	18.8	*	*	*	*	*	*	*	0.3	0.8
2004.....	19.5	*	*	*	*	*	*	*	0.3	0.8
2003.....	21.1	*	*	*	*	*	*	*	0.2	0.9
2002.....	21.9	*	*	*	*	*	*	*	0.3	0.8
2001.....	22.5	*	*	*	*	*	*	*	0.4	0.8
2000.....	23.5	*	*	*	*	*	*	*	0.2	0.7
1999.....	24.8	*	*	*	*	*	*	*	0.3	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
All races 3/, both sexes										
2006.....	73.1	*	*	*	*	*	0.2	0.5	1.5	3.2
2005.....	77.0	*	*	*	*	*	0.2	0.5	1.6	3.6
2004.....	77.8	*	*	*	*	*	0.2	0.7	1.5	3.9
2003.....	82.2	*	*	*	*	*	0.3	0.6	1.6	3.9
2002.....	84.4	*	*	*	*	*	0.2	0.6	1.3	4.1
2001.....	86.4	*	*	*	*	*	0.3	0.7	1.5	3.8
2000.....	88.6	*	*	*	*	*	0.2	0.5	1.5	4.0
1999.....	91.4	*	*	*	*	0.1	0.2	0.6	1.5	4.3
All races 3/, male										
2006.....	76.1	*	*	*	*	*	0.3	0.9	2.3	4.8
2005.....	79.3	*	*	*	*	*	0.4	0.9	2.3	5.4
2004.....	79.4	*	*	*	*	*	0.3	1.0	2.3	5.6
2003.....	83.3	*	*	*	*	*	0.3	0.8	2.3	5.9
2002.....	85.5	*	*	*	*	*	0.3	0.9	1.9	6.2
2001.....	86.5	*	*	*	*	*	0.4	1.1	2.3	5.8
2000.....	88.8	*	*	*	*	*	0.2	0.8	2.2	5.9
1999.....	91.2	*	*	*	*	0.2	0.2	0.8	2.2	6.3
All races 3/, female										
2006.....	70.3	*	*	*	*	*	*	*	0.6	1.6
2005.....	74.8	*	*	*	*	*	*	0.2	0.9	1.7
2004.....	76.2	*	*	*	*	*	*	0.4	0.7	2.1
2003.....	81.1	*	*	*	*	*	*	0.4	0.8	1.9
2002.....	83.3	*	*	*	*	*	*	0.3	0.8	2.0
2001.....	86.2	*	*	*	*	*	*	0.2	0.7	1.9
2000.....	88.4	*	*	*	*	*	*	*	0.8	2.1
1999.....	91.6	*	*	*	*	*	*	0.3	0.7	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
White, both sexes										
2006 4/.....	80.3	*	*	*	*	*	0.2	0.5	1.4	2.9
2005 4/.....	84.7	*	*	*	*	*	0.2	0.5	1.6	3.3
2004 4/.....	85.4	*	*	*	*	*	0.2	0.6	1.4	3.7
2003 4/.....	90.2	*	*	*	*	*	0.2	0.5	1.5	3.7
2002.....	92.8	*	*	*	*	*	0.2	0.6	1.2	3.9
2001.....	94.8	*	*	*	*	*	0.2	0.6	1.4	3.6
2000.....	97.4	*	*	*	*	*	0.1	0.4	1.4	3.7
1999.....	100.4	*	*	*	*	*	0.1	0.5	1.4	4.0
White, male										
2006 4/.....	83.4	*	*	*	*	*	0.3	0.8	2.2	4.4
2005 4/.....	87.0	*	*	*	*	*	0.3	0.8	2.3	5.0
2004 4/.....	87.0	*	*	*	*	*	0.2	0.9	2.1	5.5
2003 4/.....	91.3	*	*	*	*	*	0.3	0.7	2.2	5.7
2002.....	93.9	*	*	*	*	*	0.3	0.9	1.8	5.9
2001.....	94.8	*	*	*	*	*	0.4	1.1	2.1	5.5
2000.....	97.5	*	*	*	*	*	*	0.7	2.1	5.6
1999.....	99.9	*	*	*	*	*	*	0.7	2.1	6.0
White, female										
2006 4/.....	77.3	*	*	*	*	*	*	*	0.6	1.3
2005 4/.....	82.4	*	*	*	*	*	*	*	0.9	1.6
2004 4/.....	83.8	*	*	*	*	*	*	0.3	0.6	2.0
2003 4/.....	89.1	*	*	*	*	*	*	0.3	0.7	1.6
2002.....	91.7	*	*	*	*	*	*	*	0.6	1.9
2001.....	94.9	*	*	*	*	*	*	*	0.6	1.7
2000.....	97.3	*	*	*	*	*	*	*	0.6	1.8
1999.....	100.8	*	*	*	*	*	*	0.3	0.6	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Other heart diseases (I26-I51)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	56.9	7.7	0.9	0.4	0.7	1.5	2.7	3.9	5.6	7.4
2005.....	57.6	8.3	0.9	0.5	0.6	1.7	2.7	3.8	5.4	7.8
2004.....	56.6	10.0	1.1	0.4	0.7	1.5	2.5	3.6	5.0	7.8
2003.....	58.5	10.4	1.1	0.5	0.7	1.7	2.7	3.6	5.4	7.8
2002.....	58.8	11.7	1.0	0.4	0.7	1.8	2.3	3.8	5.2	8.1
2001.....	58.6	11.1	1.4	0.5	0.7	1.5	2.5	3.6	5.1	8.0
2000.....	58.8	12.0	1.1	0.5	0.7	1.8	2.6	3.5	5.2	7.5
1999.....	59.4	12.9	1.2	0.5	0.7	2.0	2.6	3.5	5.3	8.2

All races 3/, male

2006.....	51.0	8.0	1.0	0.5	0.8	1.8	3.3	4.9	7.0	9.2
2005.....	51.4	8.9	0.9	0.4	0.7	2.3	3.4	4.9	6.7	9.6
2004.....	50.6	10.5	1.1	0.3	0.8	1.9	3.2	4.5	6.1	9.4
2003.....	52.3	11.4	1.0	0.5	0.8	2.1	3.2	4.1	6.5	9.5
2002.....	52.4	12.1	1.0	0.4	0.8	2.4	3.0	4.6	6.6	9.6
2001.....	51.8	10.8	1.5	0.4	0.8	1.9	3.1	4.3	6.0	9.9
2000.....	52.2	12.2	1.2	0.6	0.9	2.1	3.1	4.2	6.2	9.1
1999.....	53.1	13.1	1.3	0.5	0.8	2.4	3.0	4.0	6.4	10.3

All races 3/, female

2006.....	62.6	7.5	0.9	0.4	0.6	1.1	2.0	2.8	4.2	5.6
2005.....	63.5	7.7	0.8	0.5	0.6	1.1	1.9	2.7	4.1	6.0
2004.....	62.4	9.4	1.1	0.5	0.6	1.2	1.8	2.6	3.9	6.1
2003.....	64.5	9.3	1.2	0.4	0.5	1.3	2.1	3.1	4.3	6.1
2002.....	65.0	11.4	1.0	0.5	0.5	1.1	1.7	3.0	3.8	6.6
2001.....	65.2	11.5	1.3	0.5	0.7	1.1	1.9	2.9	4.1	6.0
2000.....	65.1	11.8	0.9	0.4	0.6	1.4	2.1	2.9	4.1	5.9
1999.....	65.5	12.7	1.0	0.4	0.7	1.5	2.2	2.9	4.2	6.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
White, both sexes										
2006 4/.....	60.9	6.0	0.8	0.4	0.6	1.2	2.3	3.2	4.4	5.9
2005 4/.....	61.4	6.9	0.7	0.4	0.5	1.5	2.3	3.2	4.1	6.4
2004 4/.....	60.4	8.7	0.9	0.3	0.6	1.3	2.2	2.8	4.0	6.5
2003 4/.....	62.2	8.2	0.9	0.4	0.6	1.5	2.2	2.8	4.4	6.4
2002.....	62.6	10.0	0.9	0.4	0.6	1.5	1.9	3.0	4.1	6.6
2001.....	62.3	9.3	1.1	0.4	0.7	1.3	2.0	2.9	4.1	6.5
2000.....	62.5	10.6	1.0	0.4	0.6	1.4	2.1	2.8	4.3	6.1
1999.....	62.9	11.0	1.0	0.4	0.6	1.8	2.1	2.7	4.4	6.7
White, male										
2006 4/.....	53.6	6.1	0.9	0.4	0.7	1.6	2.9	4.1	5.6	7.3
2005 4/.....	54.0	7.2	0.8	0.3	0.6	1.9	2.9	4.2	5.2	8.1
2004 4/.....	53.0	9.1	0.9	*	0.7	1.6	2.9	3.6	5.0	8.0
2003 4/.....	54.7	8.8	0.8	0.4	0.8	1.8	2.5	3.2	5.5	7.9
2002.....	54.9	10.5	1.0	0.4	0.7	2.1	2.5	3.6	5.2	7.9
2001.....	54.1	9.6	1.2	0.4	0.7	1.5	2.4	3.5	4.8	8.1
2000.....	54.6	10.8	1.2	0.5	0.8	1.7	2.5	3.4	5.2	7.5
1999.....	55.1	11.4	1.1	0.5	0.7	2.1	2.4	3.3	5.4	8.5
White, female										
2006 4/.....	68.1	5.9	0.8	0.3	0.5	0.9	1.6	2.2	3.2	4.5
2005 4/.....	68.7	6.5	0.6	0.4	0.5	1.0	1.5	2.1	3.0	4.6
2004 4/.....	67.6	8.3	0.8	0.4	0.5	1.0	1.5	2.0	3.1	4.9
2003 4/.....	69.6	7.6	1.0	0.3	0.4	1.1	1.8	2.4	3.2	5.0
2002.....	70.2	9.6	0.8	0.4	0.5	1.0	1.3	2.3	2.9	5.3
2001.....	70.4	8.9	1.1	0.5	0.6	1.0	1.5	2.3	3.4	4.8
2000.....	70.2	10.4	0.7	0.3	0.4	1.2	1.6	2.2	3.3	4.6
1999.....	70.5	10.7	0.8	0.4	0.6	1.4	1.9	2.2	3.4	4.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
Black, both sexes										
2006 4/.....	50.1	16.0	1.6	0.6	1.2	2.8	4.9	8.6	14.3	18.7
2005 4/.....	51.4	15.2	1.7	1.1	1.1	3.2	5.1	8.6	14.0	18.5
2004 4/.....	50.8	14.8	2.3	0.8	1.3	3.0	4.5	8.0	11.9	17.9
2003 4/.....	52.5	19.6	2.0	0.9	1.0	3.0	5.9	8.9	12.8	18.1
2002.....	52.5	20.2	1.5	*	1.2	3.1	5.1	9.5	13.0	18.9
2001.....	52.8	20.1	2.9	0.8	1.1	2.9	5.5	8.7	12.3	18.4
2000.....	52.7	19.0	1.9	0.8	1.4	3.6	6.0	8.8	11.9	17.8
1999.....	54.4	22.3	2.0	0.7	1.2	3.5	5.6	8.6	12.4	19.5
Black, male										
2006 4/.....	50.1	17.0	1.9	*	1.4	3.4	5.3	10.8	18.0	24.0
2005 4/.....	50.6	17.7	*	*	1.2	4.6	6.6	10.4	17.8	22.6
2004 4/.....	50.4	15.7	2.3	*	1.3	4.0	5.6	10.4	13.9	21.5
2003 4/.....	51.5	23.2	2.2	*	1.2	3.9	7.5	10.5	14.3	22.5
2002.....	50.8	20.9	*	*	1.3	4.0	6.2	11.9	16.4	22.8
2001.....	51.7	17.7	3.2	*	1.2	3.8	7.0	10.7	14.9	23.1
2000.....	51.5	18.9	1.9	*	1.4	4.3	6.7	10.1	14.3	21.6
1999.....	54.3	21.8	2.1	*	1.5	4.8	6.8	10.2	15.2	24.3
Black, female										
2006 4/.....	50.0	15.0	*	*	*	2.2	4.4	6.4	10.9	13.9
2005 4/.....	52.2	12.6	2.0	1.3	*	1.7	3.7	6.9	10.6	14.8
2004 4/.....	51.1	13.9	2.3	*	1.3	2.0	3.4	5.8	10.1	14.7
2003 4/.....	53.4	15.8	1.8	*	*	2.1	4.2	7.3	11.4	14.3
2002.....	54.0	19.4	1.8	*	*	2.2	4.0	7.4	9.8	15.5
2001.....	53.8	22.7	2.5	*	*	2.0	4.1	6.9	10.0	14.1
2000.....	53.9	19.2	2.0	*	1.3	2.9	5.3	7.6	9.8	14.5
1999.....	54.5	22.8	1.8	*	*	2.2	4.5	7.1	9.9	15.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
All races 3/, both sexes										
2006.....	36.0	6.9	0.7	0.3	0.6	1.3	2.4	3.5	5.1	6.6
2005.....	37.0	7.1	0.7	0.4	0.5	1.5	2.4	3.5	4.9	7.0
2004.....	36.5	8.7	0.9	0.3	0.6	1.4	2.3	3.2	4.5	6.9
2003.....	38.0	8.6	0.9	0.4	0.6	1.5	2.4	3.3	4.9	7.0
2002.....	38.5	10.5	0.8	0.3	0.6	1.6	2.1	3.5	4.6	7.3
2001.....	37.9	9.7	1.1	0.4	0.7	1.3	2.3	3.3	4.6	7.0
2000.....	38.3	10.6	0.9	0.4	0.6	1.5	2.4	3.2	4.8	6.8
1999.....	39.0	11.9	0.9	0.4	0.7	1.8	2.4	3.2	4.9	7.4
All races 3/, male										
2006.....	34.0	6.9	0.8	0.4	0.7	1.7	3.0	4.6	6.3	8.2
2005.....	34.9	7.7	0.8	0.4	0.6	2.1	3.1	4.4	6.0	8.7
2004.....	34.5	9.2	0.9	0.2	0.7	1.8	3.0	4.1	5.5	8.3
2003.....	35.8	9.2	0.9	0.4	0.7	1.8	2.9	3.7	5.8	8.6
2002.....	36.3	10.9	0.8	0.3	0.7	2.2	2.7	4.2	5.8	8.8
2001.....	35.6	9.1	1.2	0.3	0.7	1.7	2.8	3.8	5.4	8.8
2000.....	36.1	10.5	1.0	0.5	0.8	1.9	2.8	3.8	5.7	8.3
1999.....	37.1	12.1	1.0	0.4	0.8	2.2	2.7	3.7	5.9	9.3
All races 3/, female										
2006.....	38.0	6.8	0.7	0.3	0.5	1.0	1.8	2.5	3.8	4.9
2005.....	39.0	6.4	0.7	0.4	0.4	1.0	1.7	2.4	3.7	5.3
2004.....	38.5	8.2	0.8	0.4	0.6	1.0	1.6	2.4	3.5	5.5
2003.....	40.1	7.9	1.0	0.3	0.4	1.1	1.8	2.8	3.9	5.5
2002.....	40.6	10.0	0.8	0.4	0.4	1.0	1.5	2.7	3.4	5.9
2001.....	40.2	10.4	1.0	0.4	0.6	1.0	1.7	2.7	3.7	5.3
2000.....	40.4	10.7	0.7	0.3	0.5	1.2	2.0	2.6	3.8	5.3
1999.....	40.9	11.6	0.8	0.4	0.5	1.4	2.0	2.7	3.9	5.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
White, both sexes										
2006 4/.....	37.9	5.4	0.6	0.3	0.6	1.1	2.1	2.9	4.1	5.3
2005 4/.....	38.9	6.0	0.6	0.3	0.5	1.3	2.0	2.9	3.8	5.8
2004 4/.....	38.2	7.7	0.7	0.3	0.5	1.2	2.0	2.6	3.7	5.8
2003 4/.....	39.8	7.1	0.7	0.3	0.5	1.3	1.9	2.5	4.0	5.9
2002.....	40.4	9.1	0.7	0.3	0.5	1.4	1.7	2.7	3.6	6.1
2001.....	39.6	8.2	0.9	0.3	0.6	1.1	1.8	2.6	3.7	5.7
2000.....	40.0	9.6	0.8	0.3	0.5	1.3	1.9	2.5	3.9	5.5
1999.....	40.6	10.3	0.8	0.4	0.6	1.6	1.9	2.5	4.0	6.0
White, male										
2006 4/.....	35.1	5.4	0.6	0.3	0.7	1.4	2.7	3.8	5.1	6.6
2005 4/.....	36.0	6.4	0.7	0.3	0.6	1.7	2.6	3.8	4.7	7.4
2004 4/.....	35.5	8.1	0.7	*	0.6	1.4	2.6	3.3	4.6	7.0
2003 4/.....	36.9	7.4	0.7	0.3	0.7	1.6	2.2	2.9	5.0	7.2
2002.....	37.5	9.5	0.8	0.3	0.6	1.9	2.3	3.3	4.6	7.3
2001.....	36.5	8.2	0.9	0.3	0.6	1.3	2.3	3.1	4.4	7.2
2000.....	37.1	9.4	0.9	0.4	0.7	1.5	2.3	3.1	4.8	6.9
1999.....	37.9	10.7	1.0	0.4	0.6	1.9	2.2	3.0	4.9	7.7
White, female										
2006 4/.....	40.7	5.4	0.6	*	0.4	0.8	1.4	2.0	2.9	3.9
2005 4/.....	41.6	5.6	0.6	0.3	0.4	0.9	1.4	1.9	2.8	4.2
2004 4/.....	40.9	7.3	0.6	0.4	0.5	0.9	1.4	1.9	2.7	4.5
2003 4/.....	42.6	6.8	0.8	*	0.4	1.0	1.5	2.2	2.9	4.4
2002.....	43.2	8.7	0.6	0.3	0.4	0.9	1.1	2.1	2.6	4.8
2001.....	42.6	8.2	0.8	0.3	0.5	0.9	1.3	2.1	3.0	4.2
2000.....	42.8	9.7	0.7	*	0.3	1.0	1.5	1.9	3.0	4.2
1999.....	43.2	10.0	0.7	0.4	0.5	1.4	1.7	2.0	3.1	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)										
Black, both sexes										
2006 4/.....	35.1	13.7	1.4	*	1.1	2.5	4.4	7.8	12.7	16.3
2005 4/.....	36.1	12.2	1.1	0.8	0.9	2.9	4.7	7.7	12.5	16.1
2004 4/.....	36.3	12.4	1.8	*	1.2	2.7	4.1	7.3	10.7	15.7
2003 4/.....	37.5	15.0	1.6	0.7	0.8	2.6	5.2	8.0	11.3	15.9
2002.....	37.8	17.2	1.1	*	0.9	2.9	4.8	8.7	11.4	17.0
2001.....	38.0	17.2	2.3	0.6	1.1	2.6	4.9	7.9	11.2	16.3
2000.....	38.2	15.6	1.4	0.7	1.1	3.2	5.4	8.0	10.8	15.9
1999.....	39.6	20.2	1.4	0.6	1.1	3.3	5.2	7.8	11.4	17.5
Black, male										
2006 4/.....	36.8	14.1	1.6	*	1.3	3.1	4.8	10.2	15.8	20.8
2005 4/.....	37.3	14.8	*	*	1.1	4.2	6.1	9.2	15.7	19.7
2004 4/.....	37.5	13.3	1.8	*	1.1	3.7	5.1	9.4	12.2	19.3
2003 4/.....	38.6	17.8	1.8	*	*	3.4	6.7	9.6	12.2	19.9
2002.....	38.3	18.9	*	*	1.2	3.7	5.8	10.9	14.5	20.5
2001.....	38.9	15.0	2.6	*	1.1	3.4	6.3	9.3	13.5	20.3
2000.....	39.2	14.7	1.6	*	1.2	3.9	6.1	9.1	13.2	19.3
1999.....	41.5	19.8	*	*	1.5	4.6	6.2	9.2	13.8	21.8
Black, female										
2006 4/.....	33.6	13.2	*	*	*	1.9	4.0	5.6	9.9	12.3
2005 4/.....	34.9	9.6	*	*	*	1.6	3.2	6.2	9.6	12.9
2004 4/.....	35.1	11.4	1.7	*	1.3	1.7	3.1	5.3	9.3	12.4
2003 4/.....	36.5	12.1	*	*	*	1.8	3.6	6.6	10.4	12.4
2002.....	37.4	15.4	*	*	*	2.0	3.8	6.7	8.7	13.9
2001.....	37.2	19.5	2.0	*	*	1.7	3.5	6.6	9.2	12.8
2000.....	37.3	16.5	*	*	*	2.6	4.8	6.9	8.7	12.9
1999.....	37.9	20.5	*	*	*	2.0	4.2	6.5	9.2	13.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Cerebrovascular diseases (I60-I69)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	45.8	3.4	0.3	0.2	0.2	0.3	0.6	0.9	1.7	3.4
2005.....	48.4	3.1	0.4	0.3	0.2	0.4	0.6	1.0	1.8	3.3
2004.....	51.1	3.1	0.3	0.2	0.2	0.3	0.7	1.1	1.8	3.4
2003.....	54.2	2.5	0.3	0.1	0.2	0.4	0.6	1.0	1.9	3.6
2002.....	56.4	2.9	0.3	0.2	0.3	0.3	0.6	1.0	1.8	3.6
2001.....	57.4	2.7	0.4	0.2	0.2	0.3	0.7	1.1	1.9	3.8
2000.....	59.6	3.3	0.3	0.1	0.2	0.3	0.7	1.1	1.9	3.9
1999.....	60.0	2.7	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.0

All races 3/, male

2006.....	37.0	3.9	0.3	0.3	0.3	0.3	0.7	1.0	1.8	3.7
2005.....	38.8	3.5	0.5	0.3	0.3	0.3	0.6	1.1	1.9	3.4
2004.....	40.7	3.4	0.3	*	0.2	0.4	0.7	1.1	1.7	3.7
2003.....	42.9	2.8	0.3	*	0.2	0.5	0.6	1.0	2.0	3.8
2002.....	44.2	3.2	0.4	*	0.3	0.3	0.7	1.0	1.7	3.6
2001.....	45.2	3.1	0.3	0.2	0.2	0.3	0.7	1.1	2.0	3.9
2000.....	46.9	3.8	*	*	0.2	0.4	0.7	1.1	1.9	3.8
1999.....	47.1	3.3	0.3	*	*	0.3	0.6	1.0	1.9	3.9

All races 3/, female

2006.....	54.4	2.9	0.4	*	0.2	0.4	0.6	0.8	1.6	3.1
2005.....	57.8	2.6	0.3	0.3	*	0.4	0.6	0.8	1.6	3.2
2004.....	61.2	2.8	*	*	0.2	0.3	0.7	1.0	1.8	3.1
2003.....	65.1	2.2	0.3	*	*	0.4	0.7	0.9	1.8	3.3
2002.....	68.2	2.5	0.3	*	0.3	*	0.5	0.9	1.9	3.7
2001.....	69.2	2.3	0.4	*	0.2	0.3	0.6	1.0	1.9	3.6
2000.....	71.8	2.7	0.4	*	0.3	0.3	0.7	1.0	1.9	4.0
1999.....	72.3	2.1	0.3	*	0.2	0.3	0.6	1.0	1.8	4.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Cerebrovascular diseases (I60-I69)

White, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	47.9	2.8	0.3	0.2	0.2	0.4	0.6	0.8	1.4	2.6
2005 4/.....	50.7	2.5	0.4	0.2	0.2	0.3	0.5	0.7	1.5	2.7
2004 4/.....	53.7	2.5	0.2	0.2	0.2	0.3	0.6	0.9	1.4	2.9
2003 4/.....	57.0	2.0	0.2	0.1	0.2	0.4	0.5	0.8	1.6	2.7
2002.....	59.5	2.3	0.3	0.2	0.2	0.2	0.6	0.8	1.5	2.9
2001.....	60.5	1.9	0.3	0.2	0.2	0.3	0.5	0.9	1.7	3.1
2000.....	62.8	2.5	0.3	0.1	0.2	0.3	0.6	1.0	1.6	3.0
1999.....	63.3	2.3	0.3	0.1	0.2	0.3	0.6	0.9	1.5	3.2

White, male

2006 4/.....	37.7	2.9	*	*	0.3	0.4	0.7	0.9	1.6	2.8
2005 4/.....	39.7	2.8	0.4	*	*	0.2	0.5	0.8	1.6	2.8
2004 4/.....	41.8	2.8	*	*	*	0.3	0.6	1.1	1.4	3.1
2003 4/.....	44.2	2.0	*	*	*	0.4	0.6	0.8	1.7	2.8
2002.....	45.7	2.6	0.4	*	*	0.3	0.7	1.0	1.4	2.8
2001.....	46.6	2.2	*	*	*	0.3	0.7	0.9	1.8	3.1
2000.....	48.4	3.0	*	*	*	0.4	0.7	1.1	1.6	3.0
1999.....	48.7	2.8	*	*	*	0.3	0.6	0.9	1.7	3.2

White, female

2006 4/.....	57.9	2.6	0.4	*	*	0.3	0.5	0.7	1.3	2.4
2005 4/.....	61.6	2.1	*	*	*	0.4	0.5	0.7	1.4	2.7
2004 4/.....	65.3	2.2	*	*	*	0.3	0.6	0.7	1.5	2.7
2003 4/.....	69.5	2.0	*	*	*	0.3	0.5	0.8	1.5	2.5
2002.....	73.0	2.0	*	*	0.3	*	0.5	0.7	1.6	2.9
2001.....	74.0	1.6	0.4	*	*	0.3	0.4	0.9	1.6	3.0
2000.....	76.9	2.0	*	*	0.3	*	0.6	0.9	1.5	3.0
1999.....	77.6	1.9	*	*	*	*	0.6	0.9	1.4	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Cerebrovascular diseases (I60-I69)

Black, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	43.1	7.2	*	*	*	*	1.0	1.7	3.7	8.4
2005 4/.....	44.9	6.0	*	*	*	*	1.0	2.2	3.6	6.8
2004 4/.....	46.9	6.0	*	*	*	*	1.0	2.2	4.1	7.0
2003 4/.....	49.3	4.9	*	*	*	0.7	1.2	2.0	4.1	8.8
2002.....	50.0	5.5	*	*	*	*	0.8	1.9	4.2	8.2
2001.....	51.1	6.6	*	*	*	*	1.3	2.2	3.6	8.1
2000.....	52.5	7.5	*	*	*	*	1.1	1.5	4.3	9.6
1999.....	52.2	4.6	*	*	*	*	0.8	1.8	3.6	9.1

Black, male

2006 4/.....	39.3	9.5	*	*	*	*	*	1.9	3.7	9.8
2005 4/.....	40.3	6.8	*	*	*	*	*	2.6	4.1	7.1
2004 4/.....	41.5	6.8	*	*	*	*	*	1.7	4.1	8.5
2003 4/.....	43.2	5.9	*	*	*	*	*	2.6	4.1	9.7
2002.....	43.5	5.8	*	*	*	*	*	1.7	4.0	7.5
2001.....	44.6	8.1	*	*	*	*	*	2.8	3.4	8.8
2000.....	46.1	8.6	*	*	*	*	*	*	4.1	9.2
1999.....	45.9	*	*	*	*	*	*	1.9	3.7	8.9

Black, female

2006 4/.....	46.5	*	*	*	*	*	*	1.5	3.8	7.1
2005 4/.....	49.1	*	*	*	*	*	*	1.9	3.2	6.5
2004 4/.....	51.9	*	*	*	*	*	*	2.6	4.0	5.7
2003 4/.....	54.8	*	*	*	*	*	1.6	1.5	4.1	8.0
2002.....	55.8	*	*	*	*	*	*	2.2	4.3	8.9
2001.....	56.9	*	*	*	*	*	1.7	1.7	3.8	7.4
2000.....	58.3	*	*	*	*	*	*	1.6	4.6	10.0
1999.....	57.9	*	*	*	*	*	*	1.8	3.6	9.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Other diseases of circulatory system (I71-I78)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	7.5	*	*	*	*	0.1	0.2	0.3	0.4	0.8
2005.....	8.0	*	*	*	*	*	0.2	0.3	0.5	0.8
2004.....	8.1	*	*	*	*	*	0.2	0.3	0.5	0.8
2003.....	8.5	*	*	*	*	0.1	0.2	0.3	0.4	0.8
2002.....	8.6	*	*	*	*	0.2	0.2	0.3	0.5	0.7
2001.....	8.9	*	*	*	*	0.1	0.2	0.4	0.5	0.9
2000.....	9.3	*	*	*	*	*	0.2	0.3	0.5	0.7
1999.....	9.2	*	*	*	*	*	0.2	0.3	0.4	0.8

All races 3/, male

2006.....	8.0	*	*	*	*	*	0.3	0.4	0.6	1.1
2005.....	8.5	*	*	*	*	*	0.2	0.3	0.7	1.1
2004.....	8.5	*	*	*	*	*	0.3	0.5	0.6	1.1
2003.....	9.2	*	*	*	*	*	0.3	0.4	0.6	1.1
2002.....	9.4	*	*	*	*	*	0.2	0.3	0.7	0.7
2001.....	9.6	*	*	*	*	*	0.2	0.4	0.7	1.2
2000.....	10.2	*	*	*	*	*	0.2	0.4	0.6	0.8
1999.....	10.1	*	*	*	*	*	0.2	0.4	0.6	1.0

All races 3/, female

2006.....	7.0	*	*	*	*	*	*	*	0.2	0.5
2005.....	7.4	*	*	*	*	*	*	0.3	0.3	0.4
2004.....	7.6	*	*	*	*	*	*	*	0.4	0.6
2003.....	7.8	*	*	*	*	*	*	*	0.3	0.5
2002.....	7.9	*	*	*	*	*	*	0.2	0.4	0.7
2001.....	8.2	*	*	*	*	*	*	0.3	0.3	0.6
2000.....	8.3	*	*	*	*	*	*	*	0.4	0.6
1999.....	8.4	*	*	*	*	*	*	*	0.2	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	1.3	0.8	*	*	*	*	0.2	0.2	0.4	0.5
2005.....	1.6	0.6	*	*	*	*	0.1	0.2	0.4	0.6
2004.....	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.5
2003.....	1.6	0.9	*	*	*	*	0.2	0.3	0.4	0.6
2002.....	1.6	0.7	*	*	*	*	0.1	0.2	0.5	0.6
2001.....	1.6	0.5	*	*	*	*	0.1	0.2	0.4	0.4
2000.....	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.6
1999.....	1.4	0.6	*	*	*	*	*	0.2	0.3	0.4

All races 3/, male

2006.....	1.2	*	*	*	*	*	*	*	0.4	0.6
2005.....	1.5	*	*	*	*	*	*	0.2	0.4	0.5
2004.....	1.4	*	*	*	*	*	*	0.3	0.4	0.5
2003.....	1.4	1.0	*	*	*	*	*	0.2	0.4	0.5
2002.....	1.4	*	*	*	*	*	*	*	0.5	0.6
2001.....	1.4	*	*	*	*	*	*	*	0.3	0.5
2000.....	1.4	*	*	*	*	*	*	*	0.4	0.6
1999.....	1.3	*	*	*	*	*	*	*	0.3	0.5

All races 3/, female

2006.....	1.5	*	*	*	*	*	0.2	*	0.4	0.5
2005.....	1.8	*	*	*	*	*	*	0.3	0.4	0.7
2004.....	1.7	*	*	*	*	*	*	0.3	0.4	0.5
2003.....	1.9	*	*	*	*	*	0.2	0.3	0.5	0.6
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.6
2001.....	1.9	*	*	*	*	*	*	0.3	0.4	0.4
2000.....	1.8	*	*	*	*	*	*	0.4	0.4	0.5
1999.....	1.6	*	*	*	*	*	*	*	0.3	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All Under 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39
ages 1/ year 2/ years years years years years years years years

Other disorders of circulatory system (I80-I99)

White, both sexes

2006 4/.....	1.3	0.9	*	*	*	*	*	0.1	0.3	0.5
2005 4/.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.5
2004 4/.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.4
2003 4/.....	1.6	0.9	*	*	*	*	0.1	0.2	0.3	0.5
2002.....	1.6	0.6	*	*	*	*	*	0.2	0.4	0.5
2001.....	1.7	*	*	*	*	*	*	0.2	0.3	0.3
2000.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.5
1999.....	1.5	*	*	*	*	*	*	0.1	0.3	0.4

White, male

2006 4/.....	1.2	*	*	*	*	*	*	*	0.3	0.6
2005 4/.....	1.4	*	*	*	*	*	*	*	0.3	0.5
2004 4/.....	1.4	*	*	*	*	*	*	*	0.3	0.4
2003 4/.....	1.3	*	*	*	*	*	*	*	0.3	0.5
2002.....	1.3	*	*	*	*	*	*	*	0.4	0.5
2001.....	1.4	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.4	*	*	*	*	*	*	*	0.3	0.5
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.4

White, female

2006 4/.....	1.5	*	*	*	*	*	*	*	0.3	0.4
2005 4/.....	1.8	*	*	*	*	*	*	0.3	0.3	0.6
2004 4/.....	1.8	*	*	*	*	*	*	*	0.3	0.4
2003 4/.....	1.9	*	*	*	*	*	*	0.3	0.3	0.4
2002.....	1.9	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.9	*	*	*	*	*	*	*	0.3	0.3
2000.....	1.9	*	*	*	*	*	*	0.3	0.3	0.4
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Influenza and pneumonia (J10-J18)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	18.8	6.4	0.8	0.2	0.1	0.3	0.5	0.7	1.0	1.4
2005.....	21.3	6.5	0.7	0.3	0.3	0.3	0.5	0.7	1.0	1.5
2004.....	20.3	6.7	0.7	0.2	0.2	0.3	0.6	0.5	1.0	1.5
2003.....	22.4	8.0	1.0	0.4	0.3	0.4	0.7	0.8	1.1	1.8
2002.....	22.8	6.5	0.7	0.2	0.3	0.4	0.5	0.7	1.0	1.7
2001.....	21.8	7.4	0.7	0.2	0.2	0.3	0.6	0.7	1.0	1.8
2000.....	23.2	7.6	0.7	0.2	0.2	0.3	0.7	0.7	1.1	1.8
1999.....	22.8	8.4	0.8	0.2	0.2	0.4	0.6	0.7	1.0	2.0

All races 3/, male

2006.....	17.4	7.7	0.8	*	*	0.3	0.6	0.8	0.9	1.5
2005.....	19.2	7.0	0.7	0.3	0.3	0.5	0.6	0.8	1.2	1.8
2004.....	18.6	8.0	0.6	*	0.3	0.4	0.7	0.7	1.2	1.7
2003.....	20.1	9.4	1.1	0.4	0.3	0.5	0.7	0.9	1.3	1.9
2002.....	20.4	7.8	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8
2001.....	19.6	8.1	0.7	0.2	0.2	0.3	0.7	0.7	1.2	2.0
2000.....	20.8	8.4	0.7	0.3	0.2	0.3	0.7	0.8	1.2	1.9
1999.....	20.3	8.8	0.8	0.3	0.3	0.3	0.6	0.8	1.2	2.2

All races 3/, female

2006.....	20.2	5.0	0.8	0.2	*	0.4	0.5	0.6	1.0	1.3
2005.....	23.2	5.8	0.7	0.3	0.3	*	0.4	0.7	0.9	1.2
2004.....	22.0	5.4	0.9	*	0.2	0.2	0.4	0.4	0.7	1.2
2003.....	24.6	6.6	0.9	0.4	0.4	0.3	0.7	0.6	0.8	1.6
2002.....	25.1	5.1	0.6	0.2	0.3	0.3	0.4	0.7	0.8	1.6
2001.....	23.9	6.7	0.7	0.2	0.2	0.3	0.5	0.6	0.9	1.6
2000.....	25.6	6.8	0.7	0.2	*	0.3	0.6	0.6	1.0	1.7
1999.....	25.3	8.0	0.9	*	0.2	0.4	0.5	0.6	0.8	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Influenza and pneumonia (J10-J18)

White, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	20.4	5.0	0.7	0.2	0.1	0.4	0.5	0.6	0.9	1.3
2005 4/.....	23.1	4.9	0.6	0.2	0.2	0.3	0.5	0.7	1.0	1.4
2004 4/.....	22.0	5.1	0.6	*	0.2	0.3	0.5	0.5	0.9	1.3
2003 4/.....	24.4	6.7	0.9	0.3	0.3	0.4	0.7	0.7	1.0	1.6
2002.....	24.9	5.2	0.5	0.2	0.2	0.3	0.4	0.7	0.9	1.5
2001.....	23.6	5.3	0.7	0.2	0.2	0.3	0.5	0.6	0.9	1.5
2000.....	25.2	5.8	0.6	0.2	0.2	0.3	0.6	0.6	1.0	1.6
1999.....	24.8	6.2	0.6	0.2	0.2	0.3	0.5	0.5	0.8	1.7

White, male

2006 4/.....	18.6	6.3	0.7	*	*	0.3	0.5	0.7	0.9	1.3
2005 4/.....	20.5	5.9	0.6	*	0.2	0.4	0.6	0.7	1.0	1.6
2004 4/.....	19.8	6.2	0.5	*	0.2	0.4	0.7	0.6	1.1	1.4
2003 4/.....	21.4	8.0	1.1	0.3	0.3	0.4	0.7	0.9	1.2	1.8
2002.....	21.9	6.6	0.6	*	*	0.4	0.5	0.7	1.0	1.5
2001.....	20.7	6.2	0.7	*	*	0.3	0.7	0.5	1.0	1.7
2000.....	22.0	6.4	0.5	0.3	*	0.3	0.7	0.6	1.0	1.7
1999.....	21.5	6.7	0.5	0.3	0.3	0.3	0.5	0.6	1.0	1.9

White, female

2006 4/.....	22.2	3.7	0.6	*	*	0.4	0.4	0.6	0.9	1.2
2005 4/.....	25.7	4.0	0.6	0.3	0.3	*	0.4	0.7	0.9	1.1
2004 4/.....	24.2	3.9	0.7	*	*	*	0.3	0.4	0.7	1.1
2003 4/.....	27.3	5.4	0.8	0.4	0.4	0.3	0.7	0.5	0.7	1.4
2002.....	27.8	3.7	0.5	*	0.3	0.3	0.4	0.7	0.7	1.4
2001.....	26.4	4.4	0.6	*	*	0.3	0.3	0.6	0.8	1.4
2000.....	28.2	5.1	0.7	*	*	0.3	0.5	0.6	0.9	1.6
1999.....	27.9	5.6	0.7	*	*	0.4	0.4	0.5	0.6	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Influenza and pneumonia (J10-J18)

Black, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	13.4	13.2	1.3	*	*	*	0.9	1.0	1.7	2.4
2005 4/.....	14.8	14.8	1.1	*	*	*	*	1.0	1.8	2.6
2004 4/.....	14.8	14.1	1.3	*	*	*	0.8	0.9	1.5	3.0
2003 4/.....	15.4	14.1	1.1	0.6	*	*	0.8	1.6	2.1	3.3
2002.....	15.6	12.9	1.6	*	*	*	0.7	1.0	2.0	3.5
2001.....	15.5	16.3	1.0	*	*	*	1.0	1.2	2.1	3.7
2000.....	16.4	16.7	1.2	*	*	*	1.1	1.4	2.0	2.9
1999.....	16.2	19.0	1.7	*	*	*	1.3	1.7	2.2	4.1

Black, male

2006 4/.....	13.2	15.8	*	*	*	*	*	1.4	*	2.8
2005 4/.....	14.6	13.9	*	*	*	*	*	*	2.2	3.4
2004 4/.....	15.1	16.3	*	*	*	*	*	*	2.0	3.7
2003 4/.....	16.0	16.1	*	*	*	*	*	1.7	2.8	3.5
2002.....	15.4	13.9	1.7	*	*	*	*	*	2.4	4.0
2001.....	15.9	15.9	*	*	*	*	*	*	2.7	4.5
2000.....	16.7	17.9	1.7	*	*	*	*	2.1	2.5	3.0
1999.....	16.4	18.9	2.0	*	*	*	*	1.9	2.5	4.5

Black, female

2006 4/.....	13.7	10.5	*	*	*	*	*	*	2.1	2.0
2005 4/.....	14.9	15.7	*	*	*	*	*	*	1.5	1.8
2004 4/.....	14.5	11.7	1.8	*	*	*	*	*	*	2.3
2003 4/.....	14.9	12.1	*	*	*	*	*	1.6	1.4	3.1
2002.....	15.7	11.8	*	*	*	*	*	*	1.6	3.1
2001.....	15.2	16.7	*	*	*	*	*	*	1.4	3.0
2000.....	16.0	15.5	*	*	*	*	*	*	1.6	2.7
1999.....	16.1	19.1	*	*	*	*	*	1.5	1.9	3.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
All races 3/, both sexes										
2006.....	18.5	5.9	0.7	0.1	*	0.3	0.5	0.7	1.0	1.4
2005.....	20.6	6.0	0.6	0.2	0.2	0.3	0.5	0.7	1.0	1.5
2004.....	19.9	6.3	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.4
2003.....	21.8	7.3	0.7	0.2	0.3	0.3	0.6	0.8	1.0	1.7
2002.....	22.5	6.3	0.7	0.2	0.2	0.3	0.4	0.7	1.0	1.7
2001.....	21.7	7.2	0.7	0.2	0.2	0.3	0.6	0.6	1.0	1.8
2000.....	22.6	7.4	0.6	0.2	0.2	0.3	0.6	0.7	1.1	1.7
1999.....	22.2	8.1	0.8	0.2	0.2	0.3	0.5	0.7	1.0	2.0
All races 3/, male										
2006.....	17.1	7.2	0.6	*	*	0.3	0.6	0.8	0.9	1.5
2005.....	18.8	6.6	0.6	0.2	0.2	0.4	0.6	0.8	1.1	1.8
2004.....	18.3	7.5	0.5	*	0.2	0.4	0.7	0.7	1.2	1.7
2003.....	19.6	8.5	0.7	0.2	0.2	0.4	0.7	0.9	1.3	1.9
2002.....	20.2	7.6	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8
2001.....	19.5	7.8	0.7	0.2	0.2	0.3	0.7	0.6	1.2	2.0
2000.....	20.3	8.1	0.6	0.2	*	0.3	0.7	0.8	1.2	1.8
1999.....	19.8	8.5	0.7	0.2	0.2	0.3	0.6	0.8	1.1	2.2
All races 3/, female										
2006.....	19.9	4.6	0.7	*	*	0.3	0.4	0.6	1.0	1.3
2005.....	22.5	5.3	0.6	0.2	0.2	*	0.3	0.7	0.9	1.2
2004.....	21.5	5.0	0.8	*	*	0.2	0.4	0.4	0.7	1.2
2003.....	23.9	6.0	0.6	0.2	0.3	0.2	0.6	0.6	0.8	1.5
2002.....	24.8	5.0	0.6	*	0.2	0.3	0.4	0.7	0.8	1.5
2001.....	23.8	6.7	0.7	*	0.2	0.3	0.5	0.6	0.8	1.6
2000.....	24.8	6.6	0.6	*	*	0.3	0.6	0.6	0.9	1.7
1999.....	24.6	7.7	0.8	*	*	0.3	0.5	0.6	0.8	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
White, both sexes										
2006 4/.....	20.1	4.6	0.6	*	*	0.3	0.4	0.6	0.9	1.3
2005 4/.....	22.4	4.5	0.5	0.2	0.2	0.3	0.5	0.7	0.9	1.4
2004 4/.....	21.6	4.7	0.5	*	0.2	0.3	0.5	0.5	0.9	1.2
2003 4/.....	23.7	6.0	0.6	0.2	0.3	0.3	0.7	0.6	0.9	1.5
2002.....	24.6	5.0	0.5	0.2	0.2	0.3	0.4	0.7	0.9	1.4
2001.....	23.5	5.1	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.5
2000.....	24.4	5.6	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.6
1999.....	24.1	5.9	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.7
White, male										
2006 4/.....	18.3	5.7	0.5	*	*	0.3	0.5	0.7	0.8	1.3
2005 4/.....	20.0	5.3	0.4	*	*	0.4	0.6	0.7	1.0	1.6
2004 4/.....	19.5	5.8	0.4	*	*	0.4	0.7	0.6	1.1	1.4
2003 4/.....	20.9	7.1	0.7	*	0.3	0.4	0.7	0.8	1.1	1.7
2002.....	21.7	6.4	0.5	*	*	0.4	0.5	0.6	1.0	1.5
2001.....	20.6	5.8	0.6	*	*	0.3	0.7	0.5	1.0	1.6
2000.....	21.5	6.1	0.4	0.3	*	0.3	0.6	0.6	1.0	1.7
1999.....	21.0	6.7	0.4	*	0.2	0.3	0.5	0.6	0.9	1.9
White, female										
2006 4/.....	21.8	3.5	0.6	*	*	0.3	0.4	0.6	0.9	1.2
2005 4/.....	24.8	3.7	0.5	*	*	*	0.3	0.7	0.8	1.1
2004 4/.....	23.6	3.5	0.6	*	*	*	0.3	0.4	0.7	1.0
2003 4/.....	26.5	4.8	0.5	*	0.3	*	0.6	0.5	0.7	1.3
2002.....	27.4	3.6	0.5	*	*	*	0.4	0.7	0.7	1.4
2001.....	26.2	4.4	0.6	*	*	0.3	0.3	0.6	0.7	1.4
2000.....	27.3	5.0	0.6	*	*	*	0.5	0.6	0.9	1.5
1999.....	27.1	5.2	0.7	*	*	0.3	0.4	0.4	0.6	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
Black, both sexes										
2006 4/.....	13.3	12.5	1.1	*	*	*	0.9	1.0	1.6	2.4
2005 4/.....	14.6	14.0	1.0	*	*	*	*	0.9	1.8	2.5
2004 4/.....	14.6	13.5	1.2	*	*	*	0.8	0.9	1.5	3.0
2003 4/.....	15.1	13.0	0.9	*	*	*	0.7	1.6	2.0	3.3
2002.....	15.5	12.7	1.5	*	*	*	0.7	1.0	2.0	3.5
2001.....	15.5	16.3	1.0	*	*	*	1.0	1.2	2.0	3.7
2000.....	16.1	16.4	1.2	*	*	*	1.0	1.4	2.0	2.9
1999.....	16.1	18.3	1.7	*	*	*	1.3	1.7	2.2	4.0
Black, male										
2006 4/.....	13.0	15.3	*	*	*	*	*	1.4	*	2.8
2005 4/.....	14.5	13.9	*	*	*	*	*	*	2.2	3.2
2004 4/.....	14.9	15.1	*	*	*	*	*	*	1.9	3.7
2003 4/.....	15.8	15.2	*	*	*	*	*	1.6	2.8	3.5
2002.....	15.3	13.7	1.6	*	*	*	*	*	2.4	4.0
2001.....	15.8	15.9	*	*	*	*	*	*	2.7	4.5
2000.....	16.5	17.9	1.7	*	*	*	*	2.0	2.5	3.0
1999.....	16.3	17.9	1.9	*	*	*	*	1.9	2.5	4.4
Black, female										
2006 4/.....	13.6	9.6	*	*	*	*	*	*	2.1	2.0
2005 4/.....	14.8	14.2	*	*	*	*	*	*	1.5	1.8
2004 4/.....	14.4	11.7	1.7	*	*	*	*	*	*	2.3
2003 4/.....	14.6	10.8	*	*	*	*	*	1.5	1.4	3.1
2002.....	15.6	11.8	*	*	*	*	*	*	1.6	3.1
2001.....	15.1	16.7	*	*	*	*	*	*	1.4	3.0
2000.....	15.8	14.9	*	*	*	*	*	*	1.6	2.7
1999.....	15.9	18.8	*	*	*	*	*	1.5	1.9	3.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Chronic lower respiratory diseases (J40-J47)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	41.6	0.7	0.3	0.3	0.3	0.3	0.4	0.5	0.7	1.0
2005.....	44.2	0.8	0.3	0.3	0.3	0.3	0.4	0.5	0.8	1.2
2004.....	41.5	0.9	0.3	0.2	0.3	0.4	0.4	0.6	0.7	1.3
2003.....	43.5	0.8	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
2002.....	43.3	1.0	0.4	0.2	0.4	0.5	0.5	0.6	0.9	1.6
2001.....	43.2	1.0	0.3	0.2	0.3	0.4	0.5	0.5	0.9	1.6
2000.....	43.4	0.9	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
1999.....	44.5	0.9	0.4	0.2	0.4	0.5	0.5	0.7	0.9	1.3

All races 3/, male

2006.....	40.2	0.9	0.3	0.3	0.4	0.4	0.5	0.6	0.7	1.0
2005.....	42.8	*	0.4	0.3	0.3	0.4	0.5	0.6	0.7	1.2
2004.....	40.6	1.2	0.4	0.3	0.4	0.5	0.5	0.6	0.9	1.2
2003.....	42.4	1.1	0.5	0.2	0.5	0.5	0.6	0.7	0.9	1.3
2002.....	42.9	1.1	0.6	0.2	0.5	0.6	0.5	0.7	0.9	1.5
2001.....	42.7	1.1	0.4	0.2	0.3	0.4	0.6	0.6	0.9	1.5
2000.....	43.5	1.2	0.4	0.2	0.5	0.6	0.6	0.7	0.9	1.3
1999.....	45.6	*	0.4	0.3	0.6	0.6	0.7	0.7	0.8	1.2

All races 3/, female

2006.....	43.0	*	*	*	*	0.2	0.4	0.5	0.7	1.1
2005.....	45.5	*	0.3	0.2	0.2	*	0.4	0.4	0.9	1.3
2004.....	42.5	*	*	*	0.3	0.3	0.4	0.5	0.5	1.4
2003.....	44.4	*	*	*	0.3	0.3	0.4	0.5	0.7	1.5
2002.....	43.7	*	0.3	*	0.4	0.3	0.5	0.5	0.9	1.7
2001.....	43.7	*	*	*	0.2	0.3	0.4	0.5	0.9	1.7
2000.....	43.2	*	0.3	0.2	0.4	0.3	0.5	0.6	0.8	1.4
1999.....	43.4	*	0.3	0.2	0.3	0.5	0.4	0.8	1.0	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic lower respiratory diseases (J40-J47)										
White, both sexes										
2006 4/.....	47.5	*	0.2	0.2	*	0.1	0.3	0.4	0.5	0.9
2005 4/.....	50.3	0.6	0.2	0.2	0.1	0.2	0.2	0.4	0.6	1.1
2004 4/.....	47.4	0.7	0.2	*	0.2	0.2	0.3	0.4	0.5	1.1
2003 4/.....	49.5	0.7	0.2	*	0.2	0.2	0.4	0.5	0.6	1.2
2002.....	49.2	0.8	0.2	*	0.2	0.3	0.4	0.5	0.7	1.3
2001.....	49.0	0.7	0.2	*	0.2	0.3	0.3	0.4	0.6	1.2
2000.....	49.0	0.8	0.2	0.1	0.2	0.3	0.4	0.4	0.6	1.0
1999.....	50.2	*	0.3	0.2	0.3	0.4	0.4	0.5	0.6	1.1
White, male										
2006 4/.....	45.1	*	*	*	*	*	0.3	0.5	0.4	0.8
2005 4/.....	47.9	*	*	*	*	0.3	0.3	0.4	0.4	1.0
2004 4/.....	45.5	*	*	*	0.3	0.3	0.4	0.4	0.6	1.0
2003 4/.....	47.4	1.3	*	*	0.3	0.2	0.5	0.4	0.5	1.1
2002.....	47.8	*	0.3	*	*	0.4	0.4	0.5	0.8	1.1
2001.....	47.6	*	*	*	*	*	0.3	0.4	0.5	1.2
2000.....	48.3	*	*	*	0.3	0.3	0.5	0.5	0.6	0.9
1999.....	50.6	*	*	*	0.4	0.4	0.4	0.4	0.5	1.0
White, female										
2006 4/.....	49.9	*	*	*	*	*	0.3	0.3	0.6	0.9
2005 4/.....	52.8	*	*	*	*	*	*	0.4	0.7	1.2
2004 4/.....	49.3	*	*	*	*	*	0.3	0.4	0.4	1.2
2003 4/.....	51.5	*	*	*	*	*	0.3	0.5	0.7	1.3
2002.....	50.5	*	*	*	*	*	0.4	0.4	0.7	1.5
2001.....	50.3	*	*	*	*	0.3	*	0.3	0.7	1.2
2000.....	49.7	*	*	*	*	*	0.3	0.4	0.6	1.2
1999.....	49.8	*	*	*	*	0.4	0.4	0.5	0.7	1.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Chronic lower respiratory diseases (J40-J47)

Black, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	19.5	*	*	0.9	1.2	1.1	1.2	1.5	2.3	2.7
2005 4/.....	21.1	*	1.2	0.7	0.9	0.7	1.6	1.2	2.3	2.6
2004 4/.....	19.2	*	0.8	0.9	1.0	1.3	1.2	1.7	1.9	3.0
2003 4/.....	20.2	*	1.0	0.7	1.1	1.3	1.3	1.7	2.1	2.8
2002.....	20.7	*	1.2	0.6	1.6	1.4	1.2	1.6	2.2	3.8
2001.....	20.4	*	0.9	0.7	0.9	1.0	1.6	1.8	3.0	4.5
2000.....	20.8	*	0.8	0.7	1.4	1.4	1.3	1.9	2.4	3.5
1999.....	21.9	*	*	0.6	1.5	1.2	1.5	2.5	2.7	3.2

Black, male

2006 4/.....	21.9	*	*	*	1.7	1.4	1.4	1.7	2.8	2.6
2005 4/.....	23.9	*	1.6	*	1.2	1.2	1.8	1.5	2.5	2.7
2004 4/.....	22.0	*	*	1.3	1.3	1.6	1.3	2.4	2.6	3.1
2003 4/.....	23.3	*	1.6	*	1.5	1.6	1.5	2.4	3.1	2.8
2002.....	24.1	*	1.6	*	1.7	2.2	1.5	2.2	2.3	4.5
2001.....	23.6	*	*	*	*	1.4	1.8	2.1	3.4	4.3
2000.....	24.3	*	*	*	1.7	2.0	1.5	2.4	3.1	3.7
1999.....	26.2	*	*	*	2.1	1.5	2.2	2.6	2.7	3.3

Black, female

2006 4/.....	17.4	*	*	*	*	*	*	1.4	1.8	2.7
2005 4/.....	18.4	*	*	*	*	*	1.4	*	2.2	2.5
2004 4/.....	16.6	*	*	*	*	*	*	*	1.4	2.9
2003 4/.....	17.3	*	*	*	*	*	*	*	*	2.9
2002.....	17.7	*	*	*	1.4	*	*	*	2.1	3.2
2001.....	17.5	*	*	*	*	*	1.4	1.5	2.6	4.7
2000.....	17.6	*	*	*	*	*	*	1.4	1.7	3.3
1999.....	18.0	*	*	*	*	*	*	2.4	2.8	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
All races 3/, both sexes										
2006.....	1.2	*	0.2	0.2	0.3	0.3	0.4	0.4	0.5	0.7
2005.....	1.3	*	0.2	0.2	0.3	0.2	0.4	0.5	0.6	0.7
2004.....	1.3	*	0.2	0.2	0.3	0.4	0.4	0.5	0.5	0.7
2003.....	1.4	*	0.2	0.2	0.4	0.3	0.4	0.5	0.6	0.7
2002.....	1.5	*	0.3	0.2	0.4	0.4	0.4	0.5	0.6	0.9
2001.....	1.5	*	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9
2000.....	1.6	*	0.2	0.2	0.4	0.4	0.5	0.5	0.7	0.8
1999.....	1.7	*	0.2	0.2	0.4	0.5	0.5	0.6	0.7	0.9
All races 3/, male										
2006.....	0.9	*	*	0.3	0.4	0.3	0.4	0.5	0.5	0.6
2005.....	0.9	*	0.3	0.3	0.3	0.3	0.4	0.5	0.5	0.5
2004.....	1.0	*	*	0.2	0.4	0.5	0.5	0.5	0.6	0.6
2003.....	1.0	*	0.4	0.2	0.4	0.4	0.5	0.6	0.7	0.6
2002.....	1.1	*	0.4	0.2	0.5	0.6	0.5	0.7	0.6	0.8
2001.....	1.1	*	0.3	0.2	0.3	0.3	0.4	0.5	0.6	0.8
2000.....	1.2	*	0.3	0.2	0.5	0.5	0.5	0.7	0.8	0.7
1999.....	1.2	*	*	0.2	0.6	0.5	0.6	0.6	0.6	0.8
All races 3/, female										
2006.....	1.5	*	*	*	*	0.2	0.3	0.4	0.5	0.7
2005.....	1.7	*	*	*	*	*	0.3	0.4	0.7	0.8
2004.....	1.6	*	*	*	0.2	0.3	0.3	0.5	0.4	0.9
2003.....	1.8	*	*	*	0.3	0.3	0.4	0.4	0.6	0.9
2002.....	1.8	*	*	*	0.4	0.2	0.4	0.4	0.6	1.0
2001.....	1.9	*	*	*	0.2	0.3	0.4	0.4	0.8	1.0
2000.....	2.0	*	*	0.2	0.3	0.3	0.4	0.4	0.6	0.9
1999.....	2.1	*	*	*	0.3	0.4	0.4	0.6	0.8	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
White, both sexes										
2006 4/.....	1.0	*	*	*	*	*	0.2	0.3	0.3	0.5
2005 4/.....	1.1	*	*	0.1	*	0.2	0.2	0.4	0.4	0.5
2004 4/.....	1.1	*	*	*	0.2	0.2	0.3	0.3	0.3	0.6
2003 4/.....	1.2	*	*	*	0.2	0.2	0.3	0.4	0.4	0.5
2002.....	1.3	*	*	*	0.2	0.3	0.3	0.4	0.5	0.7
2001.....	1.3	*	*	*	0.2	0.2	0.2	0.3	0.4	0.5
2000.....	1.4	*	*	*	0.2	0.2	0.4	0.3	0.5	0.5
1999.....	1.5	*	*	0.1	0.2	0.4	0.3	0.4	0.4	0.7
White, male										
2006 4/.....	0.7	*	*	*	*	*	0.3	0.4	*	0.4
2005 4/.....	0.7	*	*	*	*	*	0.2	0.4	0.3	0.4
2004 4/.....	0.7	*	*	*	*	0.3	0.3	0.3	0.4	0.4
2003 4/.....	0.8	*	*	*	0.3	*	0.3	0.4	0.4	0.4
2002.....	0.9	*	*	*	*	0.3	0.3	0.4	0.5	0.5
2001.....	0.8	*	*	*	*	*	0.2	0.3	0.3	0.5
2000.....	0.9	*	*	*	0.2	0.3	0.4	0.4	0.5	0.4
1999.....	0.9	*	*	*	0.3	0.4	0.4	0.4	0.4	0.6
White, female										
2006 4/.....	1.4	*	*	*	*	*	*	*	0.4	0.5
2005 4/.....	1.5	*	*	*	*	*	*	0.4	0.5	0.6
2004 4/.....	1.5	*	*	*	*	*	*	0.4	0.3	0.7
2003 4/.....	1.6	*	*	*	*	*	*	0.4	0.5	0.7
2002.....	1.7	*	*	*	*	*	0.3	0.4	0.6	0.9
2001.....	1.7	*	*	*	*	*	*	*	0.6	0.6
2000.....	1.8	*	*	*	*	*	0.3	0.3	0.5	0.6
1999.....	2.0	*	*	*	*	0.4	*	0.4	0.5	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
Black, both sexes										
2006 4/.....	2.4	*	*	0.8	1.1	1.1	1.2	1.3	1.9	2.2
2005 4/.....	2.6	*	1.1	0.7	0.9	0.7	1.5	1.2	1.9	1.8
2004 4/.....	2.6	*	*	0.7	1.0	1.2	1.1	1.6	1.7	2.0
2003 4/.....	2.7	*	0.9	0.7	1.0	1.1	1.3	1.4	1.9	2.0
2002.....	2.9	*	0.9	*	1.5	1.3	1.2	1.6	1.6	2.9
2001.....	3.0	*	0.8	0.7	0.8	0.9	1.4	1.7	2.6	3.4
2000.....	3.2	*	*	0.7	1.4	1.3	1.2	1.8	2.0	2.7
1999.....	3.2	*	*	0.6	1.4	1.1	1.4	2.3	2.4	2.6
Black, male										
2006 4/.....	2.2	*	*	*	1.6	1.4	1.3	1.5	2.4	2.0
2005 4/.....	2.2	*	1.6	*	1.2	*	1.6	1.5	1.9	1.7
2004 4/.....	2.5	*	*	*	1.2	1.5	1.3	2.2	2.3	1.8
2003 4/.....	2.4	*	1.5	*	1.4	1.3	1.4	2.1	2.7	1.9
2002.....	2.8	*	*	*	1.7	2.1	1.5	2.2	1.7	3.4
2001.....	2.6	*	*	*	*	*	1.5	1.9	3.0	3.0
2000.....	2.8	*	*	*	1.7	1.7	*	2.3	2.6	2.7
1999.....	2.8	*	*	*	2.0	1.3	2.0	2.6	2.6	2.6
Black, female										
2006 4/.....	2.6	*	*	*	*	*	*	*	1.6	2.4
2005 4/.....	3.0	*	*	*	*	*	1.3	*	1.9	1.9
2004 4/.....	2.7	*	*	*	*	*	*	*	*	2.2
2003 4/.....	2.9	*	*	*	*	*	*	*	*	2.1
2002.....	3.0	*	*	*	1.4	*	*	*	1.4	2.3
2001.....	3.3	*	*	*	*	*	*	1.4	2.3	3.7
2000.....	3.5	*	*	*	*	*	*	*	1.4	2.6
1999.....	3.5	*	*	*	*	*	*	2.0	2.3	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races 3/, both sexes

2006.....	9.2	8.1	0.6	0.2	0.2	0.2	0.2	0.4	0.6	0.9
2005.....	9.1	7.4	0.6	0.2	0.1	0.2	0.3	0.3	0.6	0.7
2004.....	9.3	6.5	0.7	0.2	0.1	0.3	0.3	0.4	0.5	0.9
2003.....	8.8	7.0	0.5	0.1	0.2	0.3	0.4	0.4	0.6	0.9
2002.....	8.7	6.9	0.5	0.2	0.2	0.3	0.4	0.4	0.5	0.9
2001.....	9.2	7.3	0.6	0.1	0.2	0.3	0.4	0.5	0.6	1.0
2000.....	9.1	7.2	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.0
1999.....	9.0	6.6	0.6	0.2	0.2	0.3	0.3	0.5	0.8	1.0

All races 3/, male

2006.....	9.3	9.2	0.7	0.2	*	0.3	0.3	0.4	0.5	1.0
2005.....	9.0	8.9	0.7	*	*	0.3	0.3	0.4	0.5	0.7
2004.....	9.3	6.6	0.7	*	*	0.2	0.3	0.4	0.5	0.9
2003.....	8.7	8.0	0.6	*	0.2	0.3	0.4	0.5	0.6	0.9
2002.....	8.5	7.9	0.7	0.2	*	0.3	0.4	0.3	0.5	0.9
2001.....	8.9	9.0	0.6	*	0.2	0.3	0.5	0.5	0.6	0.9
2000.....	9.0	7.9	0.6	0.2	*	0.3	0.4	0.5	0.8	1.2
1999.....	8.9	7.5	0.7	0.2	*	0.4	0.3	0.5	0.8	1.1

All races 3/, female

2006.....	9.2	6.9	0.5	*	*	0.2	*	0.4	0.7	0.8
2005.....	9.2	5.8	0.5	*	*	*	0.3	0.3	0.6	0.8
2004.....	9.3	6.4	0.6	*	*	0.3	0.3	0.4	0.5	0.9
2003.....	8.9	5.9	0.5	*	*	0.3	0.3	0.3	0.6	0.9
2002.....	8.8	5.8	0.4	0.2	0.2	0.2	0.4	0.4	0.6	0.8
2001.....	9.4	5.5	0.6	*	0.2	0.3	0.4	0.6	0.6	1.0
2000.....	9.2	6.4	0.6	*	0.2	0.3	0.3	0.5	0.7	0.9
1999.....	9.1	5.8	0.5	*	0.2	0.2	0.3	0.5	0.8	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

White, both sexes

2006 4/.....	10.1	5.7	0.5	0.2	0.1	0.2	0.2	0.4	0.6	0.8
2005 4/.....	9.9	6.0	0.5	0.2	0.1	0.2	0.3	0.3	0.5	0.6
2004 4/.....	10.2	4.9	0.5	*	0.1	0.3	0.3	0.3	0.4	0.7
2003 4/.....	9.6	5.4	0.4	0.1	0.2	0.2	0.3	0.4	0.6	0.7
2002.....	9.5	4.8	0.5	0.2	0.2	0.2	0.3	0.3	0.5	0.7
2001.....	10.0	5.0	0.5	0.1	0.2	0.2	0.4	0.4	0.5	0.8
2000.....	9.9	5.6	0.5	0.2	0.2	0.2	0.3	0.5	0.7	0.8
1999.....	9.7	4.9	0.5	0.2	0.2	0.3	0.3	0.4	0.7	0.8

White, male

2006 4/.....	10.2	6.1	0.6	*	*	0.3	0.2	0.4	0.5	0.9
2005 4/.....	9.7	7.0	0.6	*	*	0.3	*	0.4	0.5	0.6
2004 4/.....	10.1	4.9	0.6	*	*	*	0.3	0.3	0.4	0.7
2003 4/.....	9.4	5.8	0.4	*	*	*	0.3	0.4	0.6	0.8
2002.....	9.2	5.4	0.6	*	*	*	0.4	0.3	0.5	0.8
2001.....	9.7	5.9	0.5	*	0.2	0.3	0.4	0.3	0.5	0.8
2000.....	9.7	6.6	0.5	*	*	*	0.4	0.4	0.7	0.9
1999.....	9.6	5.6	0.6	*	*	0.4	0.4	0.5	0.7	0.9

White, female

2006 4/.....	10.1	5.3	0.4	*	*	*	*	0.4	0.7	0.7
2005 4/.....	10.1	4.9	0.5	*	*	*	0.3	*	0.5	0.7
2004 4/.....	10.2	4.9	0.4	*	*	0.3	0.3	0.3	0.4	0.8
2003 4/.....	9.8	5.0	0.4	*	*	0.3	*	0.3	0.5	0.7
2002.....	9.7	4.2	*	*	*	*	*	0.4	0.5	0.7
2001.....	10.2	4.0	0.6	*	*	*	0.4	0.5	0.5	0.7
2000.....	10.0	4.6	0.6	*	*	0.3	*	0.5	0.6	0.7
1999.....	9.8	4.3	0.5	*	*	*	*	0.3	0.7	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Black, both sexes

2006 4/.....	6.2	19.4	0.8	*	*	*	*	*	1.1	1.5
2005 4/.....	6.3	15.4	0.9	*	*	*	*	*	1.2	1.4
2004 4/.....	6.2	14.8	1.2	*	*	*	*	*	1.1	2.2
2003 4/.....	5.9	15.0	1.1	*	*	*	0.8	0.8	1.2	1.8
2002.....	6.0	16.6	1.1	*	*	*	0.9	*	1.0	1.8
2001.....	6.7	18.0	1.0	*	*	*	0.8	1.1	1.4	2.5
2000.....	6.5	16.7	0.8	*	*	*	0.8	0.8	1.3	2.4
1999.....	6.5	15.4	1.0	*	*	*	*	1.1	1.6	2.3

Black, male

2006 4/.....	6.2	23.6	*	*	*	*	*	*	*	1.7
2005 4/.....	6.5	20.1	*	*	*	*	*	*	*	*
2004 4/.....	6.3	15.7	*	*	*	*	*	*	*	2.8
2003 4/.....	6.0	18.4	*	*	*	*	*	*	*	1.9
2002.....	6.0	19.5	*	*	*	*	*	*	*	1.9
2001.....	6.4	23.1	*	*	*	*	*	*	1.5	2.0
2000.....	6.5	16.3	*	*	*	*	*	*	*	3.0
1999.....	6.4	17.6	*	*	*	*	*	*	1.7	2.5

Black, female

2006 4/.....	6.2	15.0	*	*	*	*	*	*	*	*
2005 4/.....	6.2	10.5	*	*	*	*	*	*	*	1.7
2004 4/.....	6.1	13.9	*	*	*	*	*	*	*	1.7
2003 4/.....	5.7	11.4	*	*	*	*	*	*	*	1.7
2002.....	6.0	13.6	*	*	*	*	*	*	*	1.6
2001.....	7.0	12.6	*	*	*	*	*	*	*	2.9
2000.....	6.4	17.2	*	*	*	*	*	*	*	1.9
1999.....	6.7	13.1	*	*	*	*	*	1.7	1.6	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Chronic liver disease and cirrhosis (K70,K73-K74)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	9.2	*	*	*	*	*	0.1	0.3	1.3	3.3
2005.....	9.3	*	*	*	*	*	*	0.4	1.2	3.4
2004.....	9.2	*	*	*	*	*	*	0.3	1.3	3.7
2003.....	9.5	*	*	*	*	*	*	0.4	1.4	4.1
2002.....	9.5	*	*	*	*	*	0.1	0.4	1.4	4.2
2001.....	9.5	*	*	*	*	*	0.1	0.4	1.5	5.0
2000.....	9.4	*	*	*	*	*	0.2	0.5	1.6	4.8
1999.....	9.4	*	*	*	*	*	*	0.4	1.6	4.8

All races 3/, male

2006.....	12.1	*	*	*	*	*	*	0.4	1.7	4.3
2005.....	12.3	*	*	*	*	*	*	0.5	1.4	4.5
2004.....	12.1	*	*	*	*	*	*	0.3	1.8	4.9
2003.....	12.5	*	*	*	*	*	*	0.6	1.9	5.4
2002.....	12.3	*	*	*	*	*	*	0.5	1.8	5.5
2001.....	12.4	*	*	*	*	*	*	0.6	2.0	6.4
2000.....	12.5	*	*	*	*	*	*	0.6	2.1	6.3
1999.....	12.5	*	*	*	*	*	*	0.4	2.1	6.3

All races 3/, female

2006.....	6.4	*	*	*	*	*	*	0.2	0.8	2.4
2005.....	6.4	*	*	*	*	*	*	0.3	0.9	2.3
2004.....	6.3	*	*	*	*	*	*	*	0.7	2.5
2003.....	6.5	*	*	*	*	*	*	0.3	0.8	2.9
2002.....	6.7	*	*	*	*	*	*	0.3	1.0	3.0
2001.....	6.7	*	*	*	*	*	*	0.3	1.0	3.5
2000.....	6.5	*	*	*	*	*	*	0.3	1.1	3.2
1999.....	6.4	*	*	*	*	*	*	0.3	1.2	3.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Chronic liver disease and cirrhosis (K70,K73-K74)

White, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	10.0	*	*	*	*	*	*	0.3	1.3	3.5
2005 4/.....	10.0	*	*	*	*	*	*	0.4	1.1	3.6
2004 4/.....	9.9	*	*	*	*	*	*	0.3	1.3	3.8
2003 4/.....	10.2	*	*	*	*	*	*	0.4	1.4	4.2
2002.....	10.1	*	*	*	*	*	*	0.4	1.4	4.3
2001.....	10.1	*	*	*	*	*	*	0.3	1.5	5.1
2000.....	10.0	*	*	*	*	*	*	0.4	1.5	4.7
1999.....	9.9	*	*	*	*	*	*	0.3	1.6	4.8

White, male

2006 4/.....	13.1	*	*	*	*	*	*	0.4	1.8	4.6
2005 4/.....	13.2	*	*	*	*	*	*	0.5	1.4	4.8
2004 4/.....	13.0	*	*	*	*	*	*	0.3	1.9	5.0
2003 4/.....	13.4	*	*	*	*	*	*	0.5	2.0	5.6
2002.....	13.2	*	*	*	*	*	*	0.4	1.9	5.6
2001.....	13.1	*	*	*	*	*	*	0.5	2.1	6.5
2000.....	13.2	*	*	*	*	*	*	0.6	2.1	6.3
1999.....	13.1	*	*	*	*	*	*	0.4	2.1	6.4

White, female

2006 4/.....	6.9	*	*	*	*	*	*	*	0.8	2.3
2005 4/.....	6.9	*	*	*	*	*	*	*	0.8	2.4
2004 4/.....	6.8	*	*	*	*	*	*	*	0.7	2.6
2003 4/.....	7.0	*	*	*	*	*	*	*	0.8	2.9
2002.....	7.2	*	*	*	*	*	*	0.3	0.9	3.0
2001.....	7.1	*	*	*	*	*	*	*	1.0	3.7
2000.....	6.8	*	*	*	*	*	*	*	1.0	3.1
1999.....	6.8	*	*	*	*	*	*	*	1.1	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Alcoholic liver disease (K70)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	4.4	*	*	*	*	*	*	0.3	1.0	2.4
2005.....	4.4	*	*	*	*	*	*	0.3	0.8	2.4
2004.....	4.3	*	*	*	*	*	*	0.2	0.9	2.5
2003.....	4.3	*	*	*	*	*	*	0.3	0.9	2.9
2002.....	4.2	*	*	*	*	*	*	0.2	0.9	2.8
2001.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3
2000.....	4.3	*	*	*	*	*	*	0.3	1.1	3.3
1999.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3

All races 3/, male

2006.....	6.4	*	*	*	*	*	*	0.4	1.4	3.1
2005.....	6.5	*	*	*	*	*	*	0.4	1.0	3.2
2004.....	6.3	*	*	*	*	*	*	0.3	1.3	3.3
2003.....	6.4	*	*	*	*	*	*	0.4	1.4	3.7
2002.....	6.2	*	*	*	*	*	*	0.3	1.2	3.6
2001.....	6.4	*	*	*	*	*	*	0.3	1.4	4.4
2000.....	6.5	*	*	*	*	*	*	0.5	1.5	4.3
1999.....	6.5	*	*	*	*	*	*	0.2	1.4	4.3

All races 3/, female

2006.....	2.4	*	*	*	*	*	*	*	0.6	1.7
2005.....	2.3	*	*	*	*	*	*	*	0.6	1.6
2004.....	2.3	*	*	*	*	*	*	*	0.5	1.8
2003.....	2.2	*	*	*	*	*	*	*	0.5	2.0
2002.....	2.3	*	*	*	*	*	*	*	0.7	2.0
2001.....	2.3	*	*	*	*	*	*	*	0.7	2.3
2000.....	2.2	*	*	*	*	*	*	*	0.6	2.2
1999.....	2.1	*	*	*	*	*	*	*	0.8	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcoholic liver disease (K70)										
White, both sexes										
2006 4/.....	4.7	*	*	*	*	*	*	0.2	1.0	2.5
2005 4/.....	4.7	*	*	*	*	*	*	0.3	0.8	2.6
2004 4/.....	4.6	*	*	*	*	*	*	0.2	1.0	2.6
2003 4/.....	4.5	*	*	*	*	*	*	0.2	1.0	2.9
2002.....	4.4	*	*	*	*	*	*	0.2	0.9	2.8
2001.....	4.5	*	*	*	*	*	*	0.2	1.1	3.4
2000.....	4.5	*	*	*	*	*	*	0.3	1.1	3.2
1999.....	4.4	*	*	*	*	*	*	0.2	1.0	3.3
White, male										
2006 4/.....	7.0	*	*	*	*	*	*	0.3	1.4	3.3
2005 4/.....	6.9	*	*	*	*	*	*	0.4	1.0	3.4
2004 4/.....	6.8	*	*	*	*	*	*	0.3	1.5	3.3
2003 4/.....	6.8	*	*	*	*	*	*	0.4	1.4	3.9
2002.....	6.6	*	*	*	*	*	*	0.3	1.3	3.7
2001.....	6.7	*	*	*	*	*	*	0.3	1.5	4.5
2000.....	6.8	*	*	*	*	*	*	0.5	1.6	4.3
1999.....	6.7	*	*	*	*	*	*	0.3	1.4	4.3
White, female										
2006 4/.....	2.5	*	*	*	*	*	*	*	0.6	1.7
2005 4/.....	2.4	*	*	*	*	*	*	*	0.5	1.7
2004 4/.....	2.4	*	*	*	*	*	*	*	0.5	1.9
2003 4/.....	2.3	*	*	*	*	*	*	*	0.5	1.9
2002.....	2.3	*	*	*	*	*	*	*	0.6	2.0
2001.....	2.3	*	*	*	*	*	*	*	0.7	2.3
2000.....	2.1	*	*	*	*	*	*	*	0.5	2.2
1999.....	2.1	*	*	*	*	*	*	*	0.7	2.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other chronic liver disease and cirrhosis (K73-K74)										
All races 3/, both sexes										
2006.....	4.8	*	*	*	*	*	*	*	0.3	1.0
2005.....	4.9	*	*	*	*	*	*	*	0.4	1.0
2004.....	4.9	*	*	*	*	*	*	*	0.3	1.2
2003.....	5.2	*	*	*	*	*	*	0.2	0.4	1.3
2002.....	5.2	*	*	*	*	*	*	0.2	0.5	1.5
2001.....	5.2	*	*	*	*	*	*	0.2	0.4	1.6
2000.....	5.1	*	*	*	*	*	0.1	0.2	0.5	1.5
1999.....	5.1	*	*	*	*	*	*	0.2	0.6	1.5
All races 3/, male										
2006.....	5.7	*	*	*	*	*	*	*	0.3	1.2
2005.....	5.8	*	*	*	*	*	*	*	0.4	1.3
2004.....	5.8	*	*	*	*	*	*	*	0.5	1.7
2003.....	6.2	*	*	*	*	*	*	*	0.5	1.6
2002.....	6.1	*	*	*	*	*	*	*	0.6	1.9
2001.....	6.1	*	*	*	*	*	*	0.2	0.6	2.0
2000.....	6.0	*	*	*	*	*	*	*	0.6	2.0
1999.....	6.0	*	*	*	*	*	*	*	0.7	2.0
All races 3/, female										
2006.....	4.0	*	*	*	*	*	*	*	0.2	0.7
2005.....	4.0	*	*	*	*	*	*	*	0.4	0.7
2004.....	4.1	*	*	*	*	*	*	*	0.2	0.7
2003.....	4.3	*	*	*	*	*	*	*	0.3	0.9
2002.....	4.5	*	*	*	*	*	*	0.2	0.4	1.0
2001.....	4.4	*	*	*	*	*	*	*	0.3	1.3
2000.....	4.3	*	*	*	*	*	*	*	0.4	1.0
1999.....	4.3	*	*	*	*	*	*	*	0.4	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other chronic liver disease and cirrhosis (K73-K74)										
White, both sexes										
2006 4/.....	5.3	*	*	*	*	*	*	*	0.3	1.0
2005 4/.....	5.3	*	*	*	*	*	*	*	0.3	1.1
2004 4/.....	5.3	*	*	*	*	*	*	*	0.3	1.2
2003 4/.....	5.7	*	*	*	*	*	*	*	0.4	1.3
2002.....	5.7	*	*	*	*	*	*	0.2	0.4	1.5
2001.....	5.6	*	*	*	*	*	*	*	0.5	1.7
2000.....	5.5	*	*	*	*	*	*	*	0.5	1.5
1999.....	5.5	*	*	*	*	*	*	0.1	0.6	1.5
White, male										
2006 4/.....	6.2	*	*	*	*	*	*	*	0.4	1.3
2005 4/.....	6.2	*	*	*	*	*	*	*	0.4	1.4
2004 4/.....	6.2	*	*	*	*	*	*	*	0.5	1.7
2003 4/.....	6.6	*	*	*	*	*	*	*	0.6	1.7
2002.....	6.5	*	*	*	*	*	*	*	0.6	2.0
2001.....	6.4	*	*	*	*	*	*	*	0.6	2.0
2000.....	6.4	*	*	*	*	*	*	*	0.5	2.1
1999.....	6.4	*	*	*	*	*	*	*	0.7	2.0
White, female										
2006 4/.....	4.4	*	*	*	*	*	*	*	*	0.7
2005 4/.....	4.4	*	*	*	*	*	*	*	0.3	0.7
2004 4/.....	4.5	*	*	*	*	*	*	*	*	0.7
2003 4/.....	4.7	*	*	*	*	*	*	*	0.3	1.0
2002.....	4.9	*	*	*	*	*	*	*	0.3	1.0
2001.....	4.8	*	*	*	*	*	*	*	0.4	1.4
2000.....	4.7	*	*	*	*	*	*	*	0.4	0.9
1999.....	4.6	*	*	*	*	*	*	*	0.4	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
All races 3/, both sexes										
2006.....	15.1	3.9	*	*	*	0.1	0.3	0.4	1.0	1.3
2005.....	14.8	3.9	*	*	*	0.2	0.3	0.6	0.9	1.2
2004.....	14.5	4.3	*	*	*	0.1	0.3	0.5	0.7	1.4
2003.....	14.6	4.5	*	*	*	0.1	0.3	0.5	0.9	1.3
2002.....	14.2	4.3	*	*	*	0.1	0.3	0.6	0.8	1.3
2001.....	13.9	3.3	*	*	*	0.1	0.4	0.4	0.7	1.3
2000.....	13.2	4.3	*	*	*	0.2	0.3	0.4	0.8	1.2
1999.....	12.7	4.4	*	*	*	0.1	0.3	0.5	0.8	1.2
All races 3/, male										
2006.....	15.0	4.8	*	*	*	*	0.4	0.4	1.2	1.4
2005.....	14.6	4.2	*	*	*	*	0.3	0.5	1.0	1.3
2004.....	14.1	5.3	*	*	*	*	0.3	0.5	0.6	1.5
2003.....	14.3	5.2	*	*	*	*	0.4	0.5	0.9	1.4
2002.....	13.9	5.3	*	*	*	*	0.3	0.5	0.8	1.4
2001.....	13.5	3.7	*	*	*	*	0.3	0.5	0.6	1.4
2000.....	12.9	5.3	*	*	*	*	0.3	0.4	0.8	1.3
1999.....	12.4	4.6	*	*	*	*	0.2	0.4	0.8	1.3
All races 3/, female										
2006.....	15.3	3.0	*	*	*	*	0.3	0.4	0.8	1.1
2005.....	15.0	3.5	*	*	*	0.2	0.3	0.6	0.8	1.1
2004.....	14.8	3.2	*	*	*	*	0.2	0.5	0.8	1.3
2003.....	14.9	3.8	*	*	*	*	0.3	0.5	0.8	1.1
2002.....	14.5	3.2	*	*	*	*	0.3	0.7	0.7	1.1
2001.....	14.2	2.9	*	*	*	*	0.4	0.4	0.7	1.3
2000.....	13.6	3.2	*	*	*	*	0.3	0.4	0.8	1.1
1999.....	13.0	4.1	*	*	*	*	0.3	0.6	0.7	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)										
White, both sexes										
2006 4/.....	14.8	3.5	*	*	*	*	0.3	0.3	0.7	0.8
2005 4/.....	14.5	3.0	*	*	*	0.1	0.2	0.4	0.6	0.9
2004 4/.....	14.1	3.1	*	*	*	*	0.2	0.3	0.5	1.0
2003 4/.....	14.3	3.8	*	*	*	*	0.2	0.4	0.6	0.8
2002.....	13.9	3.6	*	*	*	*	0.2	0.4	0.6	0.8
2001.....	13.5	2.4	*	*	*	*	0.3	0.3	0.5	0.9
2000.....	12.9	2.9	*	*	*	0.1	0.3	0.3	0.5	0.8
1999.....	12.3	3.4	*	*	*	*	0.2	0.3	0.5	0.8
White, male										
2006 4/.....	14.8	4.4	*	*	*	*	0.4	0.3	0.9	0.9
2005 4/.....	14.4	3.1	*	*	*	*	0.3	0.3	0.7	1.0
2004 4/.....	13.9	3.8	*	*	*	*	0.2	0.4	0.3	1.0
2003 4/.....	14.0	4.5	*	*	*	*	0.3	0.4	0.6	0.8
2002.....	13.7	4.4	*	*	*	*	*	0.3	0.6	0.9
2001.....	13.3	2.8	*	*	*	*	*	0.3	0.4	1.1
2000.....	12.7	3.8	*	*	*	*	0.3	0.4	0.5	0.9
1999.....	12.1	4.1	*	*	*	*	*	0.3	0.5	0.9
White, female										
2006 4/.....	14.8	2.6	*	*	*	*	*	0.3	0.5	0.7
2005 4/.....	14.6	2.9	*	*	*	*	*	0.4	0.5	0.8
2004 4/.....	14.4	2.3	*	*	*	*	*	0.3	0.6	1.0
2003 4/.....	14.5	3.0	*	*	*	*	*	0.4	0.6	0.8
2002.....	14.1	2.7	*	*	*	*	*	0.5	0.5	0.8
2001.....	13.7	2.1	*	*	*	*	0.4	0.3	0.6	0.7
2000.....	13.0	2.0	*	*	*	*	0.3	0.3	0.5	0.7
1999.....	12.4	2.8	*	*	*	*	*	0.4	0.5	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Black, both sexes

2006 4/.....	21.2	6.3	*	*	*	*	0.7	1.3	2.5	4.1
2005 4/.....	20.7	8.7	*	*	*	*	*	1.6	2.7	3.5
2004 4/.....	20.3	10.1	*	*	*	*	0.8	1.3	2.4	4.1
2003 4/.....	20.6	9.1	*	*	*	*	0.7	1.4	2.9	4.4
2002.....	19.8	8.2	*	*	*	*	0.9	1.9	2.1	4.0
2001.....	19.6	8.1	*	*	*	*	0.7	1.3	2.2	4.0
2000.....	18.9	11.5	*	*	*	*	*	1.1	2.6	4.4
1999.....	18.6	9.6	*	*	*	*	0.8	1.5	2.3	4.0

Black, male

2006 4/.....	20.2	6.9	*	*	*	*	*	*	3.1	4.6
2005 4/.....	19.5	10.6	*	*	*	*	*	1.7	2.9	3.7
2004 4/.....	19.2	12.2	*	*	*	*	*	*	2.8	5.2
2003 4/.....	20.0	9.5	*	*	*	*	*	*	3.5	5.4
2002.....	19.1	10.2	*	*	*	*	*	1.6	2.2	4.4
2001.....	18.0	8.7	*	*	*	*	*	1.7	2.6	3.4
2000.....	17.7	13.4	*	*	*	*	*	*	2.8	5.1
1999.....	17.5	8.1	*	*	*	*	*	*	2.3	4.8

Black, female

2006 4/.....	22.2	*	*	*	*	*	*	1.4	2.1	3.7
2005 4/.....	21.7	6.8	*	*	*	*	*	1.6	2.6	3.4
2004 4/.....	21.3	8.0	*	*	*	*	*	1.5	2.1	3.1
2003 4/.....	21.1	8.7	*	*	*	*	*	*	2.5	3.5
2002.....	20.5	6.1	*	*	*	*	*	2.1	2.1	3.6
2001.....	21.0	7.6	*	*	*	*	*	*	1.8	4.5
2000.....	20.0	9.6	*	*	*	*	*	*	2.3	3.8
1999.....	19.5	11.1	*	*	*	*	*	1.6	2.3	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Renal failure (N17-N19)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	14.5	3.7	*	*	*	0.1	0.3	0.4	0.9	1.2
2005.....	14.5	3.7	*	*	*	0.1	0.2	0.5	0.8	1.2
2004.....	14.2	4.0	*	*	*	0.1	0.3	0.4	0.7	1.3
2003.....	14.4	4.2	*	*	*	0.1	0.3	0.5	0.8	1.2
2002.....	13.9	4.1	*	*	*	*	0.3	0.5	0.7	1.2
2001.....	13.6	3.1	*	*	*	*	0.3	0.4	0.6	1.2
2000.....	13.0	4.0	*	*	*	0.1	0.3	0.4	0.7	1.2
1999.....	12.4	4.3	*	*	*	0.1	0.2	0.4	0.7	1.1

All races 3/, male

2006.....	14.3	4.4	*	*	*	*	0.3	0.3	1.0	1.3
2005.....	14.2	4.1	*	*	*	*	0.3	0.5	0.9	1.2
2004.....	13.8	5.1	*	*	*	*	0.3	0.4	0.6	1.4
2003.....	14.1	4.8	*	*	*	*	0.3	0.5	0.9	1.3
2002.....	13.7	5.1	*	*	*	*	0.3	0.4	0.7	1.3
2001.....	13.2	3.5	*	*	*	*	0.3	0.4	0.6	1.3
2000.....	12.6	5.1	*	*	*	*	0.2	0.4	0.7	1.3
1999.....	12.1	4.5	*	*	*	*	*	0.3	0.7	1.3

All races 3/, female

2006.....	14.6	3.0	*	*	*	*	0.3	0.4	0.7	1.1
2005.....	14.7	3.2	*	*	*	*	0.2	0.5	0.7	1.1
2004.....	14.6	3.0	*	*	*	*	0.2	0.4	0.8	1.2
2003.....	14.6	3.7	*	*	*	*	0.2	0.5	0.8	1.1
2002.....	14.2	3.0	*	*	*	*	0.3	0.6	0.7	1.1
2001.....	14.0	2.7	*	*	*	*	0.4	0.4	0.7	1.2
2000.....	13.3	2.8	*	*	*	*	0.3	0.3	0.7	1.0
1999.....	12.7	4.1	*	*	*	*	0.3	0.5	0.7	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
White, both sexes										
2006 4/.....	14.1	3.3	*	*	*	*	0.3	0.3	0.7	0.8
2005 4/.....	14.2	2.8	*	*	*	*	0.2	0.4	0.5	0.8
2004 4/.....	13.9	2.9	*	*	*	*	0.2	0.3	0.4	0.9
2003 4/.....	14.0	3.5	*	*	*	*	0.2	0.4	0.5	0.8
2002.....	13.6	3.4	*	*	*	*	0.2	0.3	0.5	0.8
2001.....	13.2	2.2	*	*	*	*	0.3	0.3	0.4	0.8
2000.....	12.6	2.7	*	*	*	*	0.2	0.3	0.5	0.7
1999.....	12.0	3.3	*	*	*	*	0.2	0.3	0.5	0.7
White, male										
2006 4/.....	14.1	4.0	*	*	*	*	0.3	*	0.8	0.9
2005 4/.....	14.1	3.1	*	*	*	*	0.2	0.3	0.6	0.9
2004 4/.....	13.6	3.6	*	*	*	*	*	0.3	0.3	0.9
2003 4/.....	13.8	4.1	*	*	*	*	*	0.4	0.5	0.8
2002.....	13.4	4.2	*	*	*	*	*	*	0.5	0.9
2001.....	13.0	2.6	*	*	*	*	*	0.3	0.4	1.0
2000.....	12.4	3.6	*	*	*	*	0.3	0.3	0.5	0.8
1999.....	11.8	3.9	*	*	*	*	*	*	0.5	0.8
White, female										
2006 4/.....	14.2	2.6	*	*	*	*	*	0.3	0.5	0.7
2005 4/.....	14.2	2.6	*	*	*	*	*	0.4	0.4	0.7
2004 4/.....	14.1	2.1	*	*	*	*	*	0.3	0.5	1.0
2003 4/.....	14.2	2.8	*	*	*	*	*	0.4	0.5	0.7
2002.....	13.8	2.6	*	*	*	*	*	0.4	0.5	0.7
2001.....	13.4	1.8	*	*	*	*	0.3	0.3	0.5	0.7
2000.....	12.8	1.7	*	*	*	*	*	*	0.5	0.7
1999.....	12.2	2.8	*	*	*	*	*	0.3	0.5	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
Black, both sexes										
2006 4/.....	20.3	6.2	*	*	*	*	0.7	1.2	2.3	3.8
2005 4/.....	20.2	8.4	*	*	*	*	*	1.4	2.6	3.5
2004 4/.....	20.0	9.8	*	*	*	*	0.7	1.2	2.3	4.0
2003 4/.....	20.3	8.9	*	*	*	*	0.6	1.3	2.8	4.4
2002.....	19.5	7.7	*	*	*	*	0.8	1.7	2.0	3.9
2001.....	19.3	8.1	*	*	*	*	0.7	1.2	2.1	3.9
2000.....	18.5	11.1	*	*	*	*	*	1.0	2.5	4.1
1999.....	18.2	9.6	*	*	*	*	*	1.4	2.2	3.8
Black, male										
2006 4/.....	19.2	6.6	*	*	*	*	*	*	2.6	4.3
2005 4/.....	19.1	10.6	*	*	*	*	*	1.4	2.7	3.5
2004 4/.....	18.9	11.9	*	*	*	*	*	*	2.7	5.1
2003 4/.....	19.7	9.2	*	*	*	*	*	*	3.4	5.4
2002.....	18.8	9.6	*	*	*	*	*	*	2.0	4.4
2001.....	17.7	8.7	*	*	*	*	*	1.6	2.6	3.3
2000.....	17.3	13.4	*	*	*	*	*	*	2.7	4.9
1999.....	17.1	8.1	*	*	*	*	*	*	2.2	4.5
Black, female										
2006 4/.....	21.3	*	*	*	*	*	*	1.3	2.0	3.4
2005 4/.....	21.2	6.2	*	*	*	*	*	1.4	2.5	3.4
2004 4/.....	21.0	7.7	*	*	*	*	*	*	1.9	3.0
2003 4/.....	20.9	8.7	*	*	*	*	*	*	2.3	3.5
2002.....	20.2	*	*	*	*	*	*	1.9	2.0	3.5
2001.....	20.7	7.6	*	*	*	*	*	*	1.7	4.4
2000.....	19.6	8.6	*	*	*	*	*	*	2.3	3.4
1999.....	19.1	11.1	*	*	*	*	*	1.6	2.2	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Pregnancy, childbirth and the puerperium (000-099)

All races 3/, both sexes

2006.....	0.3	*	0.2	0.7	0.8	0.8	0.7
2005.....	0.3	*	0.2	0.7	0.8	0.8	0.6
2004.....	0.2	*	0.2	0.6	0.7	0.7	0.5
2003.....	0.2	*	0.1	0.4	0.6	0.6	0.5
2002.....	0.1	*	0.1	0.3	0.4	0.4	0.4
2001.....	0.1	*	0.2	0.4	0.5	0.5	0.3
2000.....	0.1	*	*	0.4	0.4	0.5	0.4
1999.....	0.1	*	0.2	0.3	0.5	0.4	0.4

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	0.5	*	0.3	1.4	1.7	1.6	1.3
2005.....	0.5	*	0.4	1.4	1.6	1.6	1.3
2004.....	0.5	*	0.4	1.3	1.5	1.4	0.9
2003.....	0.4	*	0.3	0.9	1.2	1.2	1.0
2002.....	0.3	*	0.3	0.6	0.9	0.9	0.7
2001.....	0.3	*	0.4	0.8	1.0	0.9	0.7
2000.....	0.3	*	*	0.8	0.9	0.9	0.8
1999.....	0.3	*	0.3	0.7	0.9	0.9	0.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Pregnancy, childbirth and the puerperium (000-099)

White, both sexes

2006 4/.....	0.2	*	0.1	0.6	0.6	0.6	0.5
2005 4/.....	0.2	*	0.1	0.5	0.5	0.6	0.5
2004 4/.....	0.2	*	0.1	0.4	0.5	0.6	0.4
2003 4/.....	0.1	*	*	0.3	0.5	0.4	0.3
2002.....	0.1	*	*	0.2	0.3	0.3	0.2
2001.....	0.1	*	0.1	0.3	0.3	0.4	0.2
2000.....	0.1	*	*	0.3	0.3	0.3	0.3
1999.....	0.1	*	*	0.2	0.3	0.3	0.3

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	0.4	*	0.3	1.2	1.2	1.2	0.9
2005 4/.....	0.4	*	0.3	1.1	1.0	1.3	1.0
2004 4/.....	0.3	*	0.3	0.8	1.0	1.2	0.7
2003 4/.....	0.3	*	*	0.6	1.0	0.7	0.7
2002.....	0.2	*	*	0.4	0.6	0.7	0.5
2001.....	0.2	*	0.3	0.6	0.7	0.8	0.4
2000.....	0.2	*	*	0.6	0.7	0.7	0.6
1999.....	0.2	*	*	0.4	0.6	0.6	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Pregnancy, childbirth and the puerperium (000-099)

Black, both sexes

2006 4/.....	0.7	*	*	1.5	2.3	1.9	1.8
2005 4/.....	0.7	*	0.6	1.5	2.3	1.9	1.5
2004 4/.....	0.7	*	*	2.0	2.2	1.7	1.1
2003 4/.....	0.5	*	*	1.0	1.2	1.8	1.4
2002.....	0.4	*	*	1.0	1.4	1.0	1.2
2001.....	0.4	*	*	1.1	1.4	1.1	1.0
2000.....	0.4	*	*	1.1	1.1	1.2	0.9
1999.....	0.4	*	*	1.1	1.4	1.2	0.9

Black, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2006 4/.....	1.3	*	*	3.1	4.5	3.6	3.5
2005 4/.....	1.3	*	1.2	3.1	4.5	3.6	2.8
2004 4/.....	1.2	*	*	4.0	4.3	3.2	2.0
2003 4/.....	1.0	*	*	2.1	2.3	3.5	2.7
2002.....	0.8	*	*	2.0	2.8	1.9	2.3
2001.....	0.8	*	*	2.1	2.6	2.1	1.9
2000.....	0.7	*	*	2.2	2.2	2.3	1.7
1999.....	0.9	*	*	2.1	2.7	2.3	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Pregnancy with abortive outcome (000-007)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Pregnancy with abortive outcome (000-007)

White, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Pregnancy with abortive outcome (000-007)

Black, both sexes

2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Black, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other complications of pregnancy, childbirth and the puerperium (010-099)

All races 3/, both sexes

2006	0.2	*	0.2	0.7	0.8	0.7	0.6
2005	0.2	*	0.2	0.7	0.7	0.8	0.6
2004	0.2	*	0.2	0.6	0.7	0.7	0.4
2003	0.2	*	0.1	0.4	0.5	0.5	0.5
2002	0.1	*	0.1	0.3	0.4	0.4	0.3
2001	0.1	*	0.2	0.3	0.5	0.4	0.3
2000	0.1	*	*	0.4	0.4	0.4	0.3
1999	0.1	*	0.1	0.3	0.4	0.4	0.4

All races 3/, male

2006
2005
2004
2003
2002
2001
2000
1999

All races 3/, female

2006	0.5	*	0.3	1.4	1.6	1.5	1.3
2005	0.5	*	0.4	1.4	1.5	1.5	1.2
2004	0.4	*	0.4	1.2	1.4	1.4	0.9
2003	0.3	*	0.2	0.8	1.1	1.1	0.9
2002	0.2	*	0.3	0.6	0.8	0.8	0.7
2001	0.3	*	0.3	0.7	0.9	0.9	0.6
2000	0.3	*	*	0.8	0.8	0.9	0.7
1999	0.3	*	0.3	0.6	0.9	0.8	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All Under 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39
ages 1/ year 2/ years years years years years years years years

Other complications of pregnancy, childbirth and the puerperium (010-099)

White, both sexes

2006 4/.....	0.2	*	0.1	0.5	0.6	0.6	0.5
2005 4/.....	0.2	*	*	0.5	0.5	0.6	0.4
2004 4/.....	0.2	*	0.1	0.4	0.5	0.6	0.4
2003 4/.....	0.1	*	*	0.3	0.4	0.3	0.3
2002.....	0.1	*	*	0.2	0.3	0.3	0.2
2001.....	0.1	*	*	0.2	0.3	0.4	0.2
2000.....	0.1	*	*	0.3	0.3	0.3	0.3
1999.....	0.1	*	*	0.2	0.3	0.3	0.3

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	0.4	*	0.3	1.1	1.1	1.1	0.9
2005 4/.....	0.4	*	*	1.1	1.0	1.3	0.9
2004 4/.....	0.3	*	0.3	0.7	1.0	1.1	0.7
2003 4/.....	0.3	*	*	0.6	0.9	0.7	0.6
2002.....	0.2	*	*	0.4	0.5	0.6	0.5
2001.....	0.2	*	*	0.5	0.6	0.7	0.3
2000.....	0.2	*	*	0.6	0.6	0.7	0.5
1999.....	0.2	*	*	0.4	0.6	0.6	0.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

Black, both sexes

2006 4/.....	0.6	*	*	1.5	2.2	1.7	1.8
2005 4/.....	0.7	*	*	1.5	2.1	1.8	1.3
2004 4/.....	0.6	*	*	1.7	2.1	1.6	0.9
2003 4/.....	0.5	*	*	0.9	1.2	1.6	1.3
2002.....	0.4	*	*	0.9	1.4	0.9	1.0
2001.....	0.4	*	*	1.0	1.3	1.0	0.9
2000.....	0.3	*	*	1.0	1.0	1.1	0.7
1999.....	0.4	*	*	1.0	1.3	1.0	0.8

Black, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2006 4/.....	1.2	*	*	3.0	4.3	3.3	3.3
2005 4/.....	1.2	*	*	3.1	4.1	3.3	2.5
2004 4/.....	1.1	*	*	3.5	4.1	3.0	1.8
2003 4/.....	0.9	*	*	1.9	2.3	3.1	2.5
2002.....	0.7	*	*	1.8	2.6	1.8	2.0
2001.....	0.7	*	*	1.9	2.5	1.8	1.7
2000.....	0.6	*	*	1.9	1.8	2.1	1.4
1999.....	0.8	*	*	1.9	2.4	2.0	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Certain conditions originating in the perinatal period (P00-P96)

All races 3/, both sexes

2006.....	4.8	346.7	0.4	*	*	*	*	*	*	*
2005.....	4.9	351.2	0.4	*	*	*	*	*	*	*
2004.....	4.8	345.3	0.4	0.1	*	*	*	*	*	*
2003.....	4.9	356.0	0.5	*	*	*	*	*	*	*
2002.....	4.9	349.7	0.4	*	*	*	*	*	*	*
2001.....	4.9	340.5	0.5	0.1	*	*	*	*	*	*
2000.....	5.0	365.7	0.5	0.1	*	*	*	*	*	*
1999.....	5.1	371.0	0.6	0.1	*	*	*	*	*	*

All races 3/, male

2006.....	5.5	379.8	0.4	*	*	*	*	*	*	*
2005.....	5.6	386.1	0.4	*	*	*	*	*	*	*
2004.....	5.6	384.2	0.5	*	*	*	*	*	*	*
2003.....	5.7	397.4	0.5	*	*	*	*	*	*	*
2002.....	5.7	385.7	0.5	*	*	*	*	*	*	*
2001.....	5.6	376.6	0.6	*	*	*	*	*	*	*
2000.....	5.8	403.9	0.7	*	*	*	*	*	*	*
1999.....	5.9	408.6	0.7	*	*	*	*	*	*	*

All races 3/, female

2006.....	4.2	312.1	0.4	*	*	*	*	*	*	*
2005.....	4.2	314.6	0.3	*	*	*	*	*	*	*
2004.....	4.1	304.6	0.3	*	*	*	*	*	*	*
2003.....	4.2	312.8	0.5	*	*	*	*	*	*	*
2002.....	4.2	312.0	0.3	*	*	*	*	*	*	*
2001.....	4.2	302.6	0.4	*	*	*	*	*	*	*
2000.....	4.3	325.6	0.4	*	*	*	*	*	*	*
1999.....	4.4	331.7	0.5	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Certain conditions originating in the perinatal period (P00-P96)

White, both sexes

2006 4/.....	3.6	270.6	0.3	*	*	*	*	*	*	*
2005 4/.....	3.7	275.7	0.3	*	*	*	*	*	*	*
2004 4/.....	3.7	272.1	0.3	*	*	*	*	*	*	*
2003 4/.....	3.8	285.1	0.4	*	*	*	*	*	*	*
2002.....	3.7	277.2	0.3	*	*	*	*	*	*	*
2001.....	3.7	268.5	0.4	*	*	*	*	*	*	*
2000.....	3.7	283.7	0.4	*	*	*	*	*	*	*
1999.....	3.8	285.9	0.5	*	*	*	*	*	*	*

White, male

2006 4/.....	4.1	298.6	0.4	*	*	*	*	*	*	*
2005 4/.....	4.2	303.8	0.3	*	*	*	*	*	*	*
2004 4/.....	4.3	305.9	0.3	*	*	*	*	*	*	*
2003 4/.....	4.4	319.1	0.4	*	*	*	*	*	*	*
2002.....	4.3	309.4	*	*	*	*	*	*	*	*
2001.....	4.2	297.1	0.5	*	*	*	*	*	*	*
2000.....	4.3	312.4	0.5	*	*	*	*	*	*	*
1999.....	4.3	315.0	0.5	*	*	*	*	*	*	*

White, female

2006 4/.....	3.1	241.3	*	*	*	*	*	*	*	*
2005 4/.....	3.2	246.3	*	*	*	*	*	*	*	*
2004 4/.....	3.1	236.8	*	*	*	*	*	*	*	*
2003 4/.....	3.2	249.5	0.3	*	*	*	*	*	*	*
2002.....	3.2	243.5	*	*	*	*	*	*	*	*
2001.....	3.1	238.6	*	*	*	*	*	*	*	*
2000.....	3.2	253.5	*	*	*	*	*	*	*	*
1999.....	3.2	255.3	0.4	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes

2006.....	3.5	140.9	3.2	0.9	0.8	1.1	1.1	1.1	1.0	1.1
2005.....	3.5	135.2	3.2	1.0	1.0	1.2	1.2	1.2	1.0	1.1
2004.....	3.6	137.9	3.6	1.0	0.9	1.2	1.1	1.1	1.0	1.0
2003.....	3.6	140.4	3.4	0.9	1.0	1.1	1.1	1.0	1.1	1.1
2002.....	3.7	139.4	3.4	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001.....	3.7	136.7	3.6	0.9	0.9	1.3	1.3	1.2	1.1	1.2
2000.....	3.8	150.9	3.2	1.0	1.0	1.1	1.1	1.1	1.3	1.2
1999.....	3.7	144.2	3.6	1.0	1.1	1.1	1.1	1.1	1.2	1.2

All races 3/, male

2006.....	3.8	146.6	2.9	0.9	0.9	1.3	1.3	1.4	1.2	1.3
2005.....	3.7	141.1	3.2	0.9	1.0	1.6	1.4	1.3	1.1	1.2
2004.....	3.8	139.0	3.9	1.1	0.9	1.4	1.3	1.2	1.1	1.2
2003.....	3.8	147.6	3.4	0.9	1.0	1.4	1.2	1.2	1.1	1.2
2002.....	3.9	142.6	3.5	1.0	1.1	1.4	1.3	1.4	1.4	1.2
2001.....	3.9	143.1	3.6	1.0	1.0	1.5	1.4	1.3	1.2	1.2
2000.....	4.0	155.4	3.3	1.0	1.0	1.3	1.4	1.2	1.5	1.4
1999.....	3.9	147.9	4.0	1.0	1.1	1.3	1.3	1.1	1.3	1.3

All races 3/, female

2006.....	3.2	134.9	3.4	1.0	0.7	0.8	0.9	0.9	0.9	0.9
2005.....	3.3	129.0	3.3	1.2	0.9	0.8	1.0	1.0	0.9	1.0
2004.....	3.4	136.8	3.2	1.0	0.8	1.1	0.9	0.9	1.0	0.8
2003.....	3.4	132.8	3.5	0.9	1.0	0.8	1.0	0.8	1.1	1.1
2002.....	3.5	136.1	3.3	1.0	1.0	1.0	1.1	0.9	1.1	1.2
2001.....	3.4	129.9	3.7	0.8	0.8	1.0	1.1	1.1	1.0	1.1
2000.....	3.5	146.2	3.1	0.9	1.0	0.9	0.9	0.9	1.1	1.1
1999.....	3.5	140.3	3.2	1.0	1.1	0.9	1.0	1.1	1.2	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All Under 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39
ages 1/ year 2/ years years years years years years years years

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, both sexes

2006 4/.....	3.4	137.2	3.0	0.9	0.8	1.0	1.0	1.1	1.1	1.1
2005 4/.....	3.4	131.0	3.1	0.9	0.9	1.1	1.2	1.1	1.0	1.1
2004 4/.....	3.5	134.5	3.2	1.0	0.9	1.1	1.0	1.1	1.1	1.0
2003 4/.....	3.5	137.6	3.2	0.9	1.0	1.1	1.0	1.1	1.1	1.2
2002.....	3.6	138.3	3.3	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001.....	3.6	135.0	3.2	0.9	0.9	1.2	1.2	1.2	1.2	1.1
2000.....	3.6	147.1	2.9	0.9	0.9	1.0	1.2	1.1	1.4	1.2
1999.....	3.6	141.3	3.2	0.9	1.0	1.1	1.1	1.1	1.3	1.2

White, male

2006 4/.....	3.7	142.3	2.9	1.0	0.9	1.2	1.2	1.4	1.2	1.3
2005 4/.....	3.6	136.9	3.1	0.8	1.0	1.5	1.4	1.2	1.1	1.2
2004 4/.....	3.6	134.3	3.5	1.1	0.9	1.2	1.1	1.3	1.1	1.1
2003 4/.....	3.7	143.2	3.1	0.8	0.9	1.3	1.1	1.2	1.1	1.2
2002.....	3.8	143.3	3.4	0.9	1.0	1.3	1.3	1.4	1.5	1.2
2001.....	3.8	143.0	3.2	1.0	1.1	1.4	1.3	1.4	1.3	1.2
2000.....	3.9	151.9	2.9	0.9	0.9	1.1	1.4	1.2	1.6	1.4
1999.....	3.8	146.5	3.5	0.8	1.0	1.3	1.3	1.2	1.3	1.4

White, female

2006 4/.....	3.1	131.8	3.0	0.8	0.6	0.8	0.9	0.8	0.9	0.9
2005 4/.....	3.2	124.8	3.0	1.0	0.8	0.7	0.9	1.0	0.9	1.1
2004 4/.....	3.4	134.7	2.9	0.8	0.8	1.0	0.9	1.0	1.1	0.8
2003 4/.....	3.4	131.8	3.2	0.9	1.0	0.8	1.0	0.9	1.1	1.1
2002.....	3.5	133.1	3.1	1.1	1.0	1.0	1.1	0.9	1.1	1.2
2001.....	3.4	126.7	3.3	0.8	0.7	0.9	1.1	1.0	1.1	1.0
2000.....	3.5	142.0	2.9	1.0	0.9	0.9	0.9	0.9	1.1	1.0
1999.....	3.5	135.7	2.9	1.0	1.0	0.8	0.9	1.1	1.2	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)										
Black, both sexes										
2006 4/.....	4.6	170.8	4.1	1.1	0.9	1.3	1.3	1.5	1.0	1.6
2005 4/.....	4.4	162.9	4.0	1.3	1.3	1.6	1.4	1.6	1.3	1.3
2004 4/.....	4.5	162.9	5.1	1.5	1.0	1.8	1.7	1.1	1.1	1.2
2003 4/.....	4.5	157.5	4.7	1.1	1.1	1.2	1.7	1.2	1.5	1.2
2002.....	4.7	155.2	4.5	1.2	1.2	1.7	1.5	1.3	1.4	1.4
2001.....	4.6	158.4	5.3	1.0	1.2	1.9	1.8	1.4	1.2	1.5
2000.....	4.9	179.2	5.0	1.3	1.4	1.8	1.3	1.4	1.4	1.6
1999.....	4.8	169.3	5.5	1.4	1.7	1.4	1.5	1.0	1.2	1.4
Black, male										
2006 4/.....	5.0	176.8	3.1	*	*	1.6	1.5	1.8	*	1.7
2005 4/.....	4.8	166.5	3.6	*	1.3	2.1	1.8	2.1	*	1.8
2004 4/.....	5.1	169.9	5.8	1.2	*	2.0	2.5	*	1.6	1.7
2003 4/.....	5.1	170.7	4.8	1.2	*	1.7	1.8	*	*	*
2002.....	5.1	151.7	4.6	1.4	1.4	2.1	1.7	1.7	1.6	*
2001.....	5.0	160.8	5.5	*	*	2.6	2.2	*	*	*
2000.....	5.4	181.4	5.5	1.6	1.4	2.5	*	1.9	1.7	1.8
1999.....	5.2	172.7	6.2	1.4	1.8	1.7	1.8	*	*	*
Black, female										
2006 4/.....	4.2	164.5	5.0	1.3	*	*	*	*	*	1.5
2005 4/.....	4.1	159.0	4.5	1.5	1.4	*	*	*	1.5	*
2004 4/.....	4.0	155.6	4.3	1.8	*	1.6	*	*	*	*
2003 4/.....	3.9	143.8	4.6	*	*	*	1.5	*	1.6	*
2002.....	4.3	158.9	4.4	*	*	1.3	1.3	*	*	1.4
2001.....	4.3	155.9	5.1	*	1.4	1.3	1.5	1.7	*	1.7
2000.....	4.4	176.9	4.6	*	1.5	*	*	*	*	1.5
1999.....	4.3	165.8	4.8	1.5	1.7	*	*	*	*	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2006.....	10.6	83.8	1.5	0.3	0.3	0.9	2.2	2.4	2.6	3.2
2005.....	10.8	87.4	1.4	0.3	0.5	1.2	2.5	2.8	3.1	4.1
2004.....	10.3	82.4	1.2	0.3	0.3	1.0	2.1	2.7	2.8	3.6
2003.....	10.8	82.9	1.6	0.4	0.5	1.5	2.6	3.2	3.4	4.8
2002.....	10.4	85.7	1.6	0.3	0.4	1.3	2.4	3.0	3.6	4.6
2001.....	11.4	86.5	1.8	0.4	0.5	1.8	3.0	3.5	4.2	6.2
2000.....	11.3	95.6	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.7
1999.....	9.6	93.2	1.5	0.3	0.4	1.1	2.0	2.3	3.1	4.1

All races 3/, male

2006.....	9.5	95.2	1.6	0.4	0.4	1.1	2.9	3.2	3.7	4.0
2005.....	10.0	101.8	1.6	0.3	0.6	1.6	3.4	3.9	4.0	5.1
2004.....	9.5	96.9	1.2	0.3	0.2	1.3	3.0	3.6	3.6	4.6
2003.....	10.2	96.5	1.8	0.3	0.6	1.9	3.4	4.4	4.5	6.4
2002.....	10.1	98.5	1.8	0.4	0.4	1.6	3.2	4.1	4.9	6.1
2001.....	11.3	98.9	2.1	0.4	0.5	2.4	4.0	4.8	5.6	8.0
2000.....	11.5	110.6	1.7	0.5	0.5	1.9	4.0	4.7	5.9	7.6
1999.....	9.4	106.6	1.7	0.3	0.4	1.5	2.8	2.9	4.2	5.3

All races 3/, female

2006.....	11.7	71.9	1.3	0.3	0.2	0.7	1.5	1.5	1.5	2.4
2005.....	11.5	72.3	1.3	0.3	0.3	0.8	1.4	1.7	2.1	3.1
2004.....	11.0	67.3	1.1	0.2	0.4	0.7	1.3	1.7	2.0	2.7
2003.....	11.4	68.6	1.5	0.5	0.4	1.0	1.7	1.9	2.3	3.3
2002.....	10.7	72.3	1.4	0.2	0.3	0.8	1.6	1.8	2.3	3.1
2001.....	11.4	73.6	1.5	0.4	0.5	1.1	1.9	2.2	2.8	4.4
2000.....	11.2	79.9	1.5	0.4	0.4	1.2	1.8	2.2	2.8	3.7
1999.....	9.7	79.2	1.3	0.3	0.3	0.8	1.2	1.6	2.1	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, both sexes

2006 4/.....	11.0	70.2	1.2	0.3	0.3	0.9	2.2	2.4	2.6	3.2
2005 4/.....	11.1	72.9	1.2	0.2	0.4	1.1	2.4	2.7	3.1	4.0
2004 4/.....	10.5	65.8	1.0	0.2	0.2	1.0	2.2	2.6	2.7	3.6
2003 4/.....	11.1	68.1	1.4	0.4	0.4	1.5	2.5	3.1	3.4	4.7
2002.....	10.5	71.0	1.3	0.2	0.3	1.2	2.3	2.8	3.5	4.4
2001.....	11.6	70.9	1.5	0.3	0.4	1.8	3.0	3.5	4.1	6.0
2000.....	11.4	77.2	1.5	0.4	0.4	1.5	2.8	3.2	4.2	5.6
1999.....	9.4	74.7	1.3	0.3	0.3	1.0	1.8	2.0	2.9	3.7

White, male

2006 4/.....	9.5	81.8	1.3	0.3	0.3	1.1	2.9	3.3	3.6	4.0
2005 4/.....	10.0	87.7	1.4	*	0.5	1.6	3.4	3.8	4.1	4.9
2004 4/.....	9.4	76.9	1.1	0.3	*	1.2	3.0	3.6	3.5	4.6
2003 4/.....	10.1	82.4	1.4	0.3	0.4	2.0	3.4	4.2	4.5	6.1
2002.....	9.8	82.5	1.4	0.3	0.4	1.6	3.2	3.9	4.7	5.8
2001.....	11.2	82.0	1.7	0.3	0.5	2.4	4.0	4.9	5.5	7.7
2000.....	11.3	89.4	1.7	0.4	0.4	1.9	3.9	4.4	5.7	7.7
1999.....	9.0	86.4	1.6	0.3	0.3	1.3	2.6	2.7	3.8	4.9

White, female

2006 4/.....	12.5	58.1	1.1	*	*	0.6	1.5	1.5	1.5	2.4
2005 4/.....	12.2	57.4	1.1	*	*	0.7	1.4	1.5	2.0	3.0
2004 4/.....	11.6	54.2	1.0	*	0.3	0.7	1.3	1.5	1.9	2.5
2003 4/.....	12.0	53.3	1.3	0.4	0.3	1.0	1.6	1.9	2.1	3.2
2002.....	11.2	59.0	1.2	*	*	0.8	1.4	1.6	2.2	3.0
2001.....	11.9	59.3	1.4	0.4	0.4	1.2	1.9	1.9	2.7	4.3
2000.....	11.5	64.3	1.3	0.3	0.4	1.1	1.6	1.9	2.6	3.5
1999.....	9.9	62.4	1.1	0.3	*	0.7	1.0	1.3	2.1	2.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, both sexes

2006 4/.....	11.1	155.8	2.8	0.6	*	1.1	2.5	2.9	3.9	4.3
2005 4/.....	12.3	169.0	2.5	0.8	0.9	1.9	3.1	4.2	4.2	6.0
2004 4/.....	11.6	171.7	1.9	*	0.6	1.3	2.3	3.9	4.2	5.2
2003 4/.....	12.3	159.3	3.1	*	1.0	1.5	3.4	4.9	5.0	7.1
2002.....	12.3	162.3	2.6	0.7	0.7	1.8	3.2	4.6	5.6	7.5
2001.....	12.6	169.5	2.8	0.6	0.6	1.8	3.8	4.6	5.3	8.2
2000.....	13.7	197.1	2.3	0.6	0.8	2.2	4.3	5.3	6.3	7.3
1999.....	12.7	192.8	2.7	*	*	1.7	3.7	4.3	5.4	7.8

Black, male

2006 4/.....	11.7	165.9	3.1	*	*	1.4	3.1	3.7	5.7	5.2
2005 4/.....	13.3	186.6	2.7	*	*	2.4	4.5	5.7	5.2	7.1
2004 4/.....	12.6	204.3	2.1	*	*	1.6	3.1	4.8	5.5	6.3
2003 4/.....	13.5	170.7	3.4	*	1.3	1.8	4.2	7.3	6.2	9.6
2002.....	14.1	183.0	3.5	*	*	2.4	3.6	5.8	7.8	10.3
2001.....	13.9	187.4	3.5	*	*	2.4	5.2	5.5	6.5	10.5
2000.....	15.5	227.1	2.3	*	*	2.0	5.6	7.1	8.2	8.3
1999.....	14.6	217.6	2.8	*	*	2.4	4.7	5.2	8.3	9.4

Black, female

2006 4/.....	10.5	145.3	2.5	*	*	*	1.9	2.1	2.3	3.5
2005 4/.....	11.4	150.7	2.2	*	*	1.3	1.7	2.8	3.3	5.0
2004 4/.....	10.8	137.7	1.7	*	*	*	1.5	3.0	2.9	4.3
2003 4/.....	11.3	147.5	2.9	*	*	*	2.7	2.7	3.9	4.9
2002.....	10.6	140.8	1.8	*	*	*	2.7	3.5	3.7	5.1
2001.....	11.4	150.6	2.0	*	*	*	2.4	3.9	4.2	6.2
2000.....	12.0	166.0	2.3	*	*	2.3	3.0	3.6	4.6	6.3
1999.....	10.9	167.1	2.7	*	*	*	2.6	3.4	2.8	6.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)										
All races 3/, both sexes										
2006.....	79.3	28.4	3.4	1.8	2.2	3.9	6.1	7.4	9.2	13.4
2005.....	73.4	29.2	3.8	1.9	2.4	4.0	5.8	7.0	8.7	13.9
2004.....	68.8	29.2	3.8	1.9	2.4	3.8	5.6	7.1	9.0	14.3
2003.....	69.3	29.7	4.0	2.1	2.4	4.1	5.8	7.4	9.3	14.5
2002.....	67.5	28.4	4.2	1.9	2.3	3.5	5.6	6.6	9.0	14.1
2001.....	65.5	28.0	4.2	1.9	2.3	3.8	5.2	6.5	8.9	14.1
2000.....	63.4	29.0	4.0	1.8	2.5	3.6	5.4	6.5	9.0	13.9
1999.....	61.5	30.2	4.2	2.1	2.4	3.9	5.3	6.3	8.9	14.0
All races 3/, male										
2006.....	65.2	32.5	3.6	1.9	2.3	4.6	7.0	8.6	10.5	15.4
2005.....	60.9	34.8	4.3	2.0	2.5	5.0	6.5	7.9	9.9	16.1
2004.....	57.2	33.8	4.0	2.0	2.6	4.4	6.4	8.2	10.1	16.6
2003.....	57.3	34.8	4.2	2.3	2.8	4.6	6.6	8.2	10.4	16.5
2002.....	55.8	33.4	4.6	2.2	2.5	4.0	6.3	7.4	10.6	16.4
2001.....	53.8	31.4	4.6	1.8	2.5	4.4	6.0	7.3	9.9	16.8
2000.....	52.3	32.3	4.3	1.9	2.6	4.2	6.5	7.5	10.4	16.5
1999.....	51.0	35.0	4.5	2.2	2.5	4.7	5.9	7.3	10.4	16.8
All races 3/, female										
2006.....	93.0	24.1	3.1	1.7	2.0	3.1	5.2	6.3	8.0	11.5
2005.....	85.6	23.2	3.3	1.7	2.2	2.9	5.0	6.1	7.5	11.7
2004.....	79.9	24.5	3.6	1.8	2.2	3.1	4.8	6.0	7.8	12.0
2003.....	81.0	24.5	3.8	1.8	2.1	3.4	4.9	6.5	8.1	12.5
2002.....	78.8	23.0	3.9	1.7	2.1	3.0	4.9	5.8	7.3	11.9
2001.....	76.7	24.5	3.8	2.1	2.0	3.1	4.4	5.6	8.0	11.5
2000.....	74.0	25.6	3.6	1.8	2.3	3.1	4.4	5.4	7.5	11.3
1999.....	71.6	25.3	3.9	2.1	2.3	3.1	4.7	5.3	7.4	11.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)										
White, both sexes										
2006 4/.....	85.2	23.8	3.1	1.7	2.1	3.6	6.0	7.1	8.5	12.3
2005 4/.....	78.5	25.1	3.5	1.7	2.1	3.9	5.6	6.5	7.9	12.6
2004 4/.....	73.3	25.4	3.5	1.8	2.4	3.5	5.2	6.4	7.9	12.9
2003 4/.....	73.9	25.5	3.7	2.0	2.3	4.0	5.5	6.7	8.3	13.0
2002.....	71.8	25.2	3.9	1.9	2.2	3.4	5.0	6.0	8.1	12.8
2001.....	69.4	23.7	3.9	2.0	2.2	3.6	4.9	5.8	7.9	12.6
2000.....	67.0	24.9	3.5	1.7	2.3	3.5	5.1	5.7	8.0	12.3
1999.....	65.0	25.2	4.0	2.0	2.3	3.8	4.8	5.9	8.1	12.1
White, male										
2006 4/.....	69.0	27.3	3.3	1.8	2.3	4.4	7.2	8.5	10.2	14.5
2005 4/.....	64.1	30.0	4.0	1.9	2.3	5.0	6.5	7.6	9.6	15.1
2004 4/.....	59.9	29.6	3.7	2.0	2.6	4.2	6.2	7.8	9.0	15.6
2003 4/.....	59.9	31.5	3.8	2.4	2.6	4.7	6.6	7.8	9.6	15.2
2002.....	58.2	30.8	4.1	2.2	2.3	4.1	5.9	6.8	9.9	15.5
2001.....	55.8	25.9	4.4	1.9	2.4	4.4	5.8	6.6	8.9	15.5
2000.....	54.1	26.8	3.9	1.9	2.6	4.1	6.2	7.0	9.6	15.0
1999.....	52.7	29.2	4.3	2.0	2.2	4.8	5.5	7.0	9.8	15.1
White, female										
2006 4/.....	101.1	20.2	2.9	1.6	1.9	2.8	4.6	5.5	6.7	10.0
2005 4/.....	92.6	19.9	3.1	1.5	1.9	2.7	4.6	5.5	6.2	10.0
2004 4/.....	86.4	21.0	3.3	1.6	2.1	2.8	4.2	4.9	6.7	10.2
2003 4/.....	87.6	19.3	3.7	1.6	2.0	3.1	4.2	5.4	6.8	10.7
2002.....	85.2	19.3	3.6	1.6	2.1	2.8	4.1	5.0	6.1	9.9
2001.....	82.7	21.4	3.4	2.0	2.0	2.8	3.9	4.8	6.8	9.7
2000.....	79.6	22.9	3.2	1.6	2.0	2.8	3.9	4.4	6.4	9.5
1999.....	77.0	20.9	3.7	1.9	2.3	2.8	4.1	4.7	6.3	9.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Black, both sexes

2006 4/.....	66.2	51.8	4.5	2.2	2.8	5.4	7.9	11.3	15.9	22.4
2005 4/.....	63.6	50.5	5.0	2.8	3.8	4.9	8.1	11.3	15.5	24.7
2004 4/.....	60.9	49.3	5.3	2.6	2.9	5.1	8.5	12.8	17.7	26.1
2003 4/.....	60.9	51.8	5.3	2.2	3.2	4.9	8.1	13.3	16.9	27.2
2002.....	59.7	45.2	5.9	2.2	2.9	4.3	9.4	12.2	16.6	25.7
2001.....	59.1	51.3	5.4	2.0	2.9	5.3	7.7	11.5	16.6	26.6
2000.....	58.1	49.6	5.9	2.3	3.3	4.8	8.3	12.8	17.1	26.9
1999.....	56.2	56.7	5.3	2.9	3.1	4.7	8.9	10.4	15.4	28.7

Black, male

2006 4/.....	59.3	59.0	4.9	2.6	3.1	6.2	7.2	11.0	14.8	23.7
2005 4/.....	56.8	62.6	5.8	2.9	4.1	5.7	7.9	11.7	15.3	25.1
2004 4/.....	56.1	55.7	5.7	2.1	3.0	5.8	8.0	12.4	19.7	26.6
2003 4/.....	55.7	55.3	6.1	1.8	3.9	4.7	7.2	12.9	17.1	28.7
2002.....	54.8	48.8	6.4	2.3	3.4	4.2	9.4	12.7	17.9	26.4
2001.....	54.6	58.4	5.1	1.5	3.3	5.6	7.7	12.3	17.5	29.7
2000.....	54.3	59.8	6.1	2.0	3.1	5.1	9.1	12.9	18.3	29.9
1999.....	52.8	64.7	5.4	3.2	3.8	4.8	9.7	10.5	16.2	31.5

Black, female

2006 4/.....	72.6	44.2	4.0	1.7	2.6	4.6	8.5	11.5	16.8	21.3
2005 4/.....	69.7	37.9	4.1	2.7	3.4	4.2	8.2	10.9	15.6	24.3
2004 4/.....	65.4	42.6	4.9	3.1	2.9	4.3	9.0	13.2	15.9	25.6
2003 4/.....	65.6	48.2	4.5	2.7	2.5	5.2	9.0	13.6	16.7	25.9
2002.....	64.1	41.5	5.4	2.2	2.5	4.3	9.5	11.8	15.5	25.0
2001.....	63.1	43.8	5.6	2.5	2.5	5.1	7.6	10.8	15.7	24.0
2000.....	61.6	39.0	5.7	2.6	3.5	4.6	7.6	12.6	16.1	24.2
1999.....	59.3	48.3	5.1	2.6	2.4	4.5	8.1	10.3	14.6	26.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	40.6	27.8	9.9	5.3	5.9	31.2	45.3	38.8	35.1	36.9
2005.....	39.7	26.4	10.3	5.5	6.4	31.4	43.4	36.0	33.8	35.3
2004.....	38.1	25.8	10.3	5.7	7.3	32.9	41.1	34.6	30.6	34.8
2003.....	37.6	23.6	10.9	5.5	7.2	33.0	41.1	32.8	30.2	35.0
2002.....	37.0	23.5	10.5	5.9	7.3	35.0	40.9	32.3	30.8	34.9
2001.....	35.7	24.2	11.2	6.4	7.4	32.8	39.5	31.4	28.5	34.1
2000.....	34.8	23.1	11.9	6.8	7.7	33.4	38.8	30.4	28.6	32.9
1999.....	35.1	22.3	12.4	7.1	8.1	33.3	37.5	30.0	29.2	32.2

All races 3/, male

2006.....	53.5	31.6	11.1	6.1	7.3	43.0	69.1	59.2	52.1	52.7
2005.....	52.3	27.5	12.4	6.3	8.1	43.0	66.2	54.8	49.7	50.6
2004.....	49.8	29.0	11.8	6.8	9.0	44.6	61.8	52.2	45.0	49.7
2003.....	49.3	26.4	13.0	6.4	9.3	44.7	61.8	49.4	45.1	50.2
2002.....	48.9	26.8	12.9	7.1	9.2	47.6	62.4	49.5	46.1	50.5
2001.....	47.2	26.2	13.3	7.6	9.5	45.3	60.7	48.0	43.1	49.5
2000.....	46.2	25.7	14.0	7.8	9.9	45.8	58.9	47.0	43.1	48.9
1999.....	46.4	25.4	14.4	8.5	10.4	44.8	57.1	46.1	43.9	47.4

All races 3/, female

2006.....	28.1	23.8	8.6	4.4	4.4	18.9	19.9	17.6	17.6	20.8
2005.....	27.5	25.2	8.0	4.7	4.7	19.2	19.2	16.3	17.5	19.8
2004.....	26.8	22.5	8.7	4.6	5.4	20.6	19.1	16.2	15.9	19.7
2003.....	26.2	20.7	8.7	4.6	4.9	20.6	19.2	15.5	15.1	19.7
2002.....	25.6	20.0	8.1	4.7	5.3	21.8	18.4	14.5	15.2	19.2
2001.....	24.5	22.1	8.9	5.0	5.3	19.5	17.2	14.3	13.8	18.7
2000.....	23.8	20.5	9.7	5.7	5.5	20.3	17.8	13.5	13.9	17.0
1999.....	24.1	19.0	10.3	5.6	5.6	21.1	17.1	13.7	14.3	17.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

White, both sexes

2006 4/.....	42.9	22.8	9.2	4.7	5.9	33.8	49.5	41.6	37.1	38.5
2005 4/.....	41.8	23.0	10.0	5.1	6.3	33.9	46.9	38.5	35.3	36.8
2004 4/.....	40.2	22.2	10.0	5.2	7.1	36.0	44.4	36.5	32.1	36.1
2003 4/.....	39.5	20.0	10.6	5.2	7.0	35.9	44.3	34.6	31.6	36.2
2002.....	38.7	18.6	9.9	5.2	7.0	37.9	43.8	33.5	32.2	35.5
2001.....	37.0	20.6	11.0	5.8	7.1	34.9	41.7	32.7	29.3	34.8
2000.....	35.9	19.1	11.1	6.2	7.6	36.0	40.7	30.9	29.1	33.4
1999.....	36.0	18.9	11.5	6.3	7.9	35.6	38.9	30.8	29.3	32.4

White, male

2006 4/.....	55.7	25.8	10.4	5.4	7.2	45.9	75.5	62.8	54.3	54.4
2005 4/.....	54.3	23.7	12.3	5.9	7.8	46.1	71.6	58.1	51.2	52.1
2004 4/.....	52.0	25.5	11.5	6.1	8.9	48.5	67.0	54.6	46.5	51.4
2003 4/.....	51.3	22.5	12.8	6.0	9.0	48.4	66.2	51.9	46.8	51.4
2002.....	50.4	22.3	11.8	6.3	8.8	51.0	66.4	50.7	47.9	50.9
2001.....	48.4	22.1	13.3	6.7	9.0	47.8	64.2	49.6	44.0	50.3
2000.....	47.0	21.7	13.2	7.2	9.6	48.8	61.7	47.2	43.7	49.5
1999.....	46.9	21.8	13.2	7.4	10.1	47.2	58.9	47.1	43.8	47.5

White, female

2006 4/.....	30.3	19.8	8.0	4.0	4.5	21.0	21.4	18.9	19.0	22.1
2005 4/.....	29.5	22.2	7.6	4.3	4.7	21.0	20.3	17.4	18.6	20.9
2004 4/.....	28.8	18.8	8.4	4.3	5.2	22.9	20.1	17.2	16.9	20.3
2003 4/.....	28.0	17.4	8.3	4.4	4.8	22.7	20.7	16.1	15.7	20.6
2002.....	27.3	14.7	7.8	4.2	5.2	23.9	19.6	15.1	15.9	19.7
2001.....	25.9	18.9	8.5	4.8	5.1	21.1	17.8	14.8	14.0	19.0
2000.....	25.1	16.4	9.0	5.3	5.5	22.3	18.3	13.8	13.8	17.0
1999.....	25.3	15.8	9.6	5.1	5.6	23.3	17.6	13.8	14.3	17.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Black, both sexes

2006 4/.....	35.2	52.7	14.0	8.1	6.3	21.8	31.1	33.4	35.0	37.9
2005 4/.....	34.9	47.4	12.5	7.5	7.6	22.8	31.8	29.4	34.9	36.1
2004 4/.....	32.8	45.8	12.5	8.5	8.5	21.6	30.1	32.2	30.6	34.9
2003 4/.....	32.4	43.5	13.4	7.3	8.1	20.7	30.6	29.8	29.9	35.5
2002.....	33.1	48.9	14.1	9.3	9.0	23.9	32.0	32.8	30.1	38.1
2001.....	33.5	43.1	13.6	9.4	9.4	24.7	33.0	30.5	30.0	35.8
2000.....	33.5	44.5	16.2	9.7	8.8	23.7	34.3	33.1	31.9	35.9
1999.....	35.2	41.6	17.7	11.3	9.6	25.0	34.6	31.5	32.9	36.6

Black, male

2006 4/.....	50.8	61.3	15.9	9.7	8.0	33.1	46.2	53.4	57.2	58.8
2005 4/.....	50.0	50.8	13.8	8.3	10.3	32.5	48.5	47.2	54.8	55.7
2004 4/.....	46.7	48.3	14.2	10.4	10.1	31.2	44.8	51.7	48.9	51.5
2003 4/.....	46.1	47.6	15.7	8.7	10.6	30.0	47.6	46.6	46.9	54.9
2002.....	47.9	51.1	18.5	11.1	12.1	34.6	49.5	53.6	47.7	58.9
2001.....	48.2	48.5	15.4	12.6	12.9	36.1	51.0	48.3	46.6	53.3
2000.....	49.0	47.7	18.8	11.9	12.1	34.5	52.9	54.1	49.6	55.7
1999.....	51.0	45.9	21.0	14.5	13.0	36.6	54.7	51.0	51.7	55.4

Black, female

2006 4/.....	20.9	43.6	12.1	6.3	4.6	10.1	15.6	14.4	14.9	19.2
2005 4/.....	21.2	43.8	11.1	6.7	4.8	12.7	14.9	12.6	16.8	18.5
2004 4/.....	20.1	43.2	10.6	6.5	6.7	11.8	15.4	14.1	14.1	20.0
2003 4/.....	19.9	39.3	11.0	5.9	5.5	11.2	13.7	14.3	14.7	18.3
2002.....	19.7	46.6	9.6	7.5	5.8	12.8	14.9	13.8	14.4	19.7
2001.....	20.1	37.5	11.9	6.2	5.8	12.9	15.6	14.2	15.1	20.2
2000.....	19.5	41.3	13.5	7.6	5.4	12.7	16.7	14.1	15.9	18.3
1999.....	20.8	37.3	14.3	8.0	6.1	13.1	15.6	13.8	16.0	19.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Transport accidents (V01-V99,Y85)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	16.2	3.4	3.8	3.1	4.1	23.9	29.8	20.9	17.4	16.4
2005.....	16.3	3.6	4.0	3.3	4.3	24.4	29.3	20.5	17.6	16.3
2004.....	16.4	3.6	4.1	3.5	5.1	26.0	28.5	20.9	16.7	16.4
2003.....	16.5	3.7	4.1	3.5	5.1	26.7	28.8	20.1	16.5	16.7
2002.....	16.8	3.1	4.1	3.5	4.9	28.4	29.9	20.5	17.1	17.1
2001.....	16.6	3.7	4.4	3.8	5.0	26.6	29.2	20.9	16.9	17.4
2000.....	16.6	4.6	4.5	4.1	5.3	27.1	29.3	20.4	16.8	17.1
1999.....	16.6	5.0	4.6	4.4	5.3	27.1	28.2	20.0	17.0	16.9

All races 3/, male

2006.....	23.1	3.3	4.0	3.5	4.7	31.2	44.6	31.8	25.9	23.7
2005.....	23.3	3.5	4.5	3.7	5.1	31.8	44.0	31.2	26.0	23.5
2004.....	23.0	3.9	4.4	3.9	6.0	33.6	42.0	31.2	24.4	24.0
2003.....	23.3	4.0	4.5	4.0	6.3	34.7	42.4	29.8	24.8	24.1
2002.....	23.7	3.5	4.8	4.0	5.9	37.0	44.4	31.0	25.4	24.7
2001.....	23.5	3.3	5.0	4.3	6.2	35.4	43.9	31.4	25.4	24.7
2000.....	23.3	4.8	5.0	4.4	6.4	35.5	43.3	30.8	24.7	24.8
1999.....	23.1	5.2	5.1	5.0	6.5	34.8	41.8	29.9	24.7	24.0

All races 3/, female

2006.....	9.4	3.6	3.6	2.8	3.3	16.2	14.0	9.5	8.6	9.1
2005.....	9.5	3.6	3.5	3.0	3.5	16.5	13.6	9.3	8.9	8.9
2004.....	9.9	3.2	3.8	3.0	4.1	18.0	14.2	10.2	8.9	8.8
2003.....	10.0	3.4	3.7	3.0	3.8	18.3	14.3	10.0	8.1	9.3
2002.....	10.1	2.8	3.4	3.1	3.9	19.3	14.6	9.7	8.7	9.6
2001.....	10.0	4.2	3.8	3.3	3.8	17.3	13.8	10.1	8.3	10.2
2000.....	10.2	4.3	4.0	3.8	4.1	18.1	14.6	9.8	8.7	9.5
1999.....	10.4	4.8	4.2	3.6	4.1	19.0	14.0	10.0	9.3	9.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Transport accidents (V01-V99,Y85)

White, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	16.7	3.1	3.5	3.0	4.2	26.0	31.9	21.4	17.6	16.8
2005 4/.....	16.9	3.1	3.9	3.3	4.3	26.4	31.1	21.2	17.9	16.6
2004 4/.....	16.9	3.3	4.0	3.4	5.2	28.6	30.3	21.4	17.0	16.7
2003 4/.....	17.0	3.6	4.0	3.5	5.2	29.2	30.4	20.7	16.8	17.0
2002.....	17.3	2.7	3.9	3.4	5.0	31.0	31.5	20.7	17.6	17.2
2001.....	17.0	3.7	4.3	3.6	5.0	28.6	30.4	21.6	17.1	17.6
2000.....	17.0	4.1	4.3	4.1	5.4	29.6	30.5	20.3	16.8	17.3
1999.....	17.0	4.6	4.1	4.1	5.5	29.4	29.2	20.4	17.2	16.8

White, male

2006 4/.....	23.7	3.2	3.6	3.2	4.8	33.7	47.7	32.4	25.8	23.9
2005 4/.....	24.0	3.3	4.3	3.7	5.1	34.2	46.7	32.0	26.0	23.5
2004 4/.....	23.7	3.3	4.2	3.8	6.3	36.7	44.8	31.5	24.3	24.3
2003 4/.....	23.9	4.0	4.5	3.9	6.5	37.6	44.4	30.6	25.1	24.2
2002.....	24.2	3.1	4.5	3.8	6.0	40.2	46.7	31.0	25.9	24.5
2001.....	23.9	3.2	4.9	4.0	6.2	37.8	45.8	32.3	25.4	24.7
2000.....	23.6	4.5	4.7	4.4	6.5	38.4	45.0	30.3	24.6	24.8
1999.....	23.4	5.2	4.4	4.7	6.7	37.3	43.1	30.2	24.6	23.7

White, female

2006 4/.....	9.8	3.1	3.4	2.7	3.5	17.9	14.8	9.6	9.1	9.5
2005 4/.....	10.0	2.9	3.5	3.0	3.6	18.2	14.2	9.7	9.5	9.4
2004 4/.....	10.3	3.2	3.8	2.9	4.1	20.0	14.7	10.5	9.3	8.9
2003 4/.....	10.3	3.1	3.4	3.1	3.9	20.2	15.3	10.2	8.2	9.6
2002.....	10.5	2.2	3.4	2.9	3.9	21.3	15.3	9.8	9.0	9.7
2001.....	10.3	4.1	3.6	3.1	3.8	18.9	14.0	10.4	8.4	10.4
2000.....	10.5	3.7	3.8	3.7	4.3	20.2	15.1	9.9	8.7	9.8
1999.....	10.7	3.9	3.8	3.5	4.3	21.0	14.4	10.1	9.6	9.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Transport accidents (V01-V99,Y85)										
Black, both sexes										
2006 4/.....	15.1	4.6	5.2	4.0	3.8	15.7	23.0	21.7	21.0	17.7
2005 4/.....	15.0	5.7	4.5	3.6	4.3	17.1	23.6	19.2	20.0	17.9
2004 4/.....	15.2	4.8	5.1	4.0	4.8	16.6	22.7	22.4	19.3	17.6
2003 4/.....	15.1	4.2	5.0	3.6	4.5	16.2	23.2	20.3	18.3	17.9
2002.....	15.4	5.3	5.0	4.6	4.6	17.9	24.6	23.0	18.0	19.2
2001.....	15.9	*	5.2	5.3	5.3	18.8	25.8	21.0	18.9	18.1
2000.....	16.0	6.5	5.6	4.5	4.8	17.3	26.6	23.5	19.4	17.9
1999.....	16.3	7.3	7.1	5.8	5.0	18.7	25.8	21.3	18.5	19.9
Black, male										
2006 4/.....	22.9	*	5.7	4.8	4.6	22.5	33.9	34.4	35.2	28.9
2005 4/.....	22.7	*	4.9	3.6	5.5	23.5	35.8	31.2	33.6	29.4
2004 4/.....	22.8	5.9	5.4	4.4	5.3	22.9	33.2	36.1	31.5	27.9
2003 4/.....	22.6	*	5.1	4.5	5.7	22.7	36.3	31.8	29.4	28.4
2002.....	23.1	*	6.4	5.0	5.6	24.5	37.1	37.5	28.9	29.8
2001.....	23.8	*	5.7	6.6	7.3	26.5	39.1	33.2	30.0	27.2
2000.....	24.1	6.7	6.2	4.9	6.5	24.2	40.3	38.2	30.7	28.6
1999.....	24.2	*	8.0	7.2	6.4	25.8	39.2	34.3	29.4	29.7
Black, female										
2006 4/.....	8.0	*	4.6	3.1	3.0	8.7	11.8	9.7	8.2	7.7
2005 4/.....	7.9	6.8	4.2	3.5	3.0	10.6	11.2	7.9	7.8	7.7
2004 4/.....	8.2	*	4.8	3.6	4.2	10.0	12.1	9.6	8.3	8.5
2003 4/.....	8.3	*	5.0	2.8	3.4	9.5	10.2	9.8	8.2	8.4
2002.....	8.3	*	3.4	4.2	3.5	11.2	12.5	9.8	8.2	9.8
2001.....	8.7	*	4.7	4.0	3.4	10.8	13.0	9.9	8.9	10.0
2000.....	8.5	*	5.0	4.1	3.0	10.3	13.7	10.2	9.2	8.4
1999.....	9.2	8.7	6.2	4.4	3.6	11.4	13.1	9.5	8.6	11.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes

2006.....	15.1	3.4	3.6	2.9	3.7	23.2	28.8	19.9	16.4	15.2
2005.....	15.3	3.6	3.8	3.2	4.0	23.6	28.2	19.6	16.5	15.1
2004.....	15.3	3.5	4.0	3.3	4.8	25.2	27.5	19.7	15.6	15.2
2003.....	15.4	3.6	3.9	3.3	4.7	25.7	27.5	19.0	15.4	15.4
2002.....	15.7	3.0	3.9	3.3	4.5	27.6	28.8	19.7	16.0	15.8
2001.....	15.4	3.6	4.1	3.6	4.6	25.7	28.0	19.6	15.6	15.9
2000.....	15.4	4.4	4.2	3.8	4.8	26.0	28.0	19.2	15.5	15.6
1999.....	15.2	4.8	4.2	3.9	4.8	25.9	26.5	18.4	15.4	15.1

All races 3/, male

2006.....	21.4	3.3	3.8	3.2	4.2	30.2	42.9	30.1	24.4	21.7
2005.....	21.7	3.5	4.2	3.5	4.7	30.7	42.3	29.6	24.2	21.8
2004.....	21.3	3.8	4.2	3.7	5.6	32.5	40.4	29.3	22.4	21.9
2003.....	21.4	3.9	4.4	3.7	5.9	33.3	40.4	28.2	23.0	21.9
2002.....	21.9	3.3	4.5	3.8	5.3	36.0	42.6	29.6	23.6	22.4
2001.....	21.5	3.2	4.5	4.0	5.7	34.0	42.1	29.5	23.1	22.3
2000.....	21.3	4.6	4.7	4.1	5.8	33.9	41.2	28.6	22.6	22.1
1999.....	20.9	5.0	4.7	4.4	5.7	33.1	39.1	27.2	22.2	21.2

All races 3/, female

2006.....	9.0	3.5	3.4	2.6	3.2	15.7	13.7	9.2	8.1	8.7
2005.....	9.1	3.6	3.4	2.9	3.2	16.2	13.2	9.1	8.6	8.4
2004.....	9.5	3.2	3.7	2.8	3.9	17.5	13.8	9.7	8.5	8.3
2003.....	9.5	3.3	3.5	2.8	3.5	17.8	13.9	9.5	7.7	8.8
2002.....	9.8	2.8	3.3	2.9	3.6	18.9	14.3	9.4	8.3	9.2
2001.....	9.5	4.0	3.6	3.0	3.5	16.9	13.4	9.5	8.0	9.6
2000.....	9.7	4.2	3.8	3.5	3.8	17.5	14.2	9.5	8.2	9.1
1999.....	9.7	4.7	3.8	3.3	3.8	18.3	13.4	9.4	8.5	9.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, both sexes

2006 4/.....	15.6	3.1	3.4	2.7	3.8	25.2	30.7	20.4	16.6	15.6
2005 4/.....	15.8	3.1	3.7	3.2	4.0	25.6	30.0	20.3	16.8	15.4
2004 4/.....	15.8	3.2	3.9	3.2	4.9	27.8	29.2	20.0	15.7	15.3
2003 4/.....	15.8	3.5	3.8	3.3	4.8	28.1	29.1	19.5	15.7	15.6
2002.....	16.2	2.6	3.7	3.2	4.6	30.2	30.4	19.8	16.4	15.8
2001.....	15.7	3.5	3.9	3.3	4.6	27.6	29.1	20.2	15.7	16.1
2000.....	15.7	3.9	4.0	3.7	5.0	28.4	29.1	19.0	15.4	15.8
1999.....	15.5	4.5	3.8	3.7	5.0	28.1	27.4	18.6	15.5	14.9

White, male

2006 4/.....	22.0	3.2	3.5	3.0	4.2	32.6	45.8	30.7	24.2	21.9
2005 4/.....	22.3	3.3	4.1	3.5	4.6	33.0	45.1	30.4	24.1	21.7
2004 4/.....	21.9	3.3	4.0	3.6	5.9	35.6	43.2	29.5	22.2	22.1
2003 4/.....	22.0	3.9	4.3	3.6	6.0	36.1	42.3	28.8	23.3	21.9
2002.....	22.4	2.9	4.2	3.6	5.5	39.0	44.8	29.5	23.9	22.2
2001.....	21.9	3.1	4.4	3.7	5.6	36.2	43.8	30.1	23.1	22.3
2000.....	21.6	4.2	4.5	4.0	5.9	36.8	42.6	27.9	22.4	22.1
1999.....	21.1	5.0	4.1	4.2	5.9	35.5	40.4	27.4	22.1	20.8

White, female

2006 4/.....	9.4	3.0	3.3	2.5	3.3	17.4	14.3	9.3	8.5	9.1
2005 4/.....	9.5	2.9	3.4	2.8	3.2	17.7	13.8	9.4	9.1	8.8
2004 4/.....	9.8	3.2	3.7	2.7	3.9	19.5	14.1	9.9	8.9	8.3
2003 4/.....	9.8	3.0	3.2	2.9	3.6	19.6	14.8	9.6	7.7	9.0
2002.....	10.1	2.2	3.2	2.7	3.7	20.9	15.0	9.5	8.6	9.3
2001.....	9.8	3.8	3.4	2.9	3.5	18.5	13.5	9.7	8.0	9.8
2000.....	10.0	3.5	3.6	3.4	4.0	19.5	14.6	9.6	8.2	9.3
1999.....	10.0	3.9	3.5	3.2	4.0	20.2	13.7	9.5	8.7	8.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2006 4/.....	14.3	4.4	5.0	3.8	3.6	15.3	22.5	20.6	20.1	16.5
2005 4/.....	14.1	5.7	4.3	3.4	4.0	16.7	22.4	18.4	19.0	17.0
2004 4/.....	14.3	4.7	4.9	3.8	4.5	15.8	21.8	21.2	18.4	16.6
2003 4/.....	14.2	4.2	4.9	3.4	4.3	15.5	22.3	19.5	17.3	16.7
2002.....	14.5	5.2	4.8	4.3	4.2	17.4	24.0	22.4	17.2	17.9
2001.....	14.8	*	5.0	4.9	4.8	18.1	25.2	20.1	17.7	16.7
2000.....	15.0	6.5	5.4	4.3	4.6	16.5	25.8	22.8	18.0	16.4
1999.....	15.1	7.1	6.4	5.3	4.6	17.8	24.8	20.1	17.0	18.2

Black, male

2006 4/.....	21.5	*	5.7	4.6	4.3	21.8	33.1	32.4	33.6	26.6
2005 4/.....	21.2	*	4.7	3.4	5.1	22.7	33.8	29.9	31.9	27.6
2004 4/.....	21.2	*	5.2	4.2	5.0	21.6	31.5	33.8	29.8	26.0
2003 4/.....	21.1	*	5.0	4.1	5.2	21.6	34.6	30.5	27.7	26.3
2002.....	21.5	*	6.3	4.8	5.0	23.7	36.1	36.2	27.4	27.6
2001.....	22.0	*	5.3	6.2	6.5	25.5	38.0	31.9	28.1	24.9
2000.....	22.5	6.7	5.8	4.5	6.2	22.8	38.8	36.7	28.5	26.0
1999.....	22.2	*	7.0	6.3	5.7	24.5	37.7	32.4	27.0	27.0

Black, female

2006 4/.....	7.7	*	4.4	3.1	2.9	8.6	11.6	9.4	7.9	7.5
2005 4/.....	7.5	6.8	3.9	3.4	3.0	10.6	10.8	7.6	7.5	7.5
2004 4/.....	7.9	*	4.7	3.5	4.0	9.8	12.0	9.4	8.1	8.2
2003 4/.....	8.0	*	4.8	2.7	3.4	9.2	10.1	9.4	8.0	8.2
2002.....	8.0	*	3.4	3.8	3.3	11.0	12.2	9.7	7.9	9.3
2001.....	8.2	*	4.6	3.6	3.1	10.5	12.9	9.5	8.4	9.5
2000.....	8.2	*	4.9	4.0	2.9	10.1	13.5	10.2	8.6	7.9
1999.....	8.7	8.7	5.8	4.2	3.5	11.0	12.6	9.0	8.1	10.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2006.....	0.4	*	*	*	0.2	0.4	0.4	0.4	0.3	0.5
2005.....	0.4	*	0.2	*	0.2	0.4	0.5	0.4	0.4	0.5
2004.....	0.4	*	*	0.1	0.2	0.5	0.4	0.5	0.4	0.4
2003.....	0.5	*	*	0.1	0.2	0.6	0.7	0.5	0.5	0.5
2002.....	0.4	*	*	0.1	0.2	0.3	0.5	0.3	0.4	0.5
2001.....	0.5	*	0.2	0.1	0.3	0.5	0.5	0.4	0.4	0.6
2000.....	0.5	*	0.2	0.2	0.3	0.6	0.6	0.5	0.6	0.7
1999.....	0.8	*	0.3	0.3	0.3	0.9	1.0	0.8	0.8	0.9

All races 3/, male

2006.....	0.6	*	*	*	0.3	0.5	0.7	0.6	0.4	0.8
2005.....	0.7	*	*	*	0.3	0.7	0.7	0.7	0.6	0.7
2004.....	0.6	*	*	*	0.2	0.7	0.7	0.8	0.7	0.7
2003.....	0.7	*	*	*	0.2	0.9	1.0	0.8	0.8	0.8
2002.....	0.6	*	*	*	0.3	0.5	0.8	0.5	0.6	0.8
2001.....	0.8	*	0.3	*	0.3	0.7	0.9	0.7	0.8	1.1
2000.....	0.8	*	*	0.2	0.4	1.0	1.0	0.8	0.9	1.1
1999.....	1.1	*	0.3	0.4	0.5	1.2	1.7	1.3	1.1	1.4

All races 3/, female

2006.....	0.2	*	*	*	*	0.2	*	*	*	*
2005.....	0.2	*	*	*	*	*	*	*	*	0.2
2004.....	0.2	*	*	*	*	0.2	*	*	*	*
2003.....	0.2	*	*	*	*	0.3	0.2	0.2	*	0.2
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	0.2	*	*	*	0.2
2000.....	0.3	*	*	*	*	0.3	0.2	*	0.3	0.2
1999.....	0.4	*	0.3	*	*	0.5	0.3	0.3	0.5	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, both sexes

2006 4/.....	0.4	*	*	*	0.2	0.4	0.5	0.3	0.4	0.5
2005 4/.....	0.4	*	0.2	*	0.2	0.5	0.4	0.4	0.4	0.5
2004 4/.....	0.4	*	*	*	0.2	0.4	0.4	0.5	0.4	0.5
2003 4/.....	0.5	*	*	*	0.2	0.7	0.7	0.6	0.5	0.5
2002.....	0.4	*	*	*	0.2	0.3	0.5	0.3	0.4	0.5
2001.....	0.5	*	0.2	*	0.2	0.5	0.6	0.5	0.4	0.5
2000.....	0.5	*	0.2	0.2	0.3	0.7	0.6	0.5	0.6	0.6
1999.....	0.8	*	0.3	0.3	0.3	0.9	1.0	0.8	0.8	0.9

White, male

2006 4/.....	0.6	*	*	*	0.3	0.6	0.8	0.5	0.5	0.8
2005 4/.....	0.6	*	*	*	0.3	0.7	0.7	0.6	0.6	0.7
2004 4/.....	0.6	*	*	*	0.2	0.7	0.6	0.7	0.7	0.7
2003 4/.....	0.7	*	*	*	0.3	1.0	1.1	0.9	0.8	0.8
2002.....	0.6	*	*	*	0.3	0.5	0.9	0.5	0.6	0.8
2001.....	0.7	*	*	*	0.3	0.7	0.9	0.8	0.7	0.9
2000.....	0.8	*	*	*	0.4	1.0	1.0	0.9	0.9	1.1
1999.....	1.1	*	*	0.4	0.5	1.2	1.7	1.3	1.1	1.4

White, female

2006 4/.....	0.2	*	*	*	*	*	*	*	*	*
2005 4/.....	0.2	*	*	*	*	*	*	*	*	0.2
2004 4/.....	0.2	*	*	*	*	*	*	*	*	*
2003 4/.....	0.2	*	*	*	*	0.4	0.3	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	0.3	*	*	0.3	0.2
1999.....	0.4	*	*	*	*	0.6	0.4	0.3	0.5	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes

2006.....	0.6	*	*	0.1	0.2	0.3	0.6	0.7	0.7	0.7
2005.....	0.6	*	*	*	0.2	0.3	0.6	0.5	0.7	0.7
2004.....	0.7	*	*	*	0.1	0.4	0.6	0.8	0.7	0.9
2003.....	0.7	*	*	0.1	0.2	0.4	0.6	0.6	0.6	0.9
2002.....	0.7	*	*	0.1	0.2	0.4	0.6	0.5	0.7	0.9
2001.....	0.7	*	0.2	0.1	0.1	0.4	0.6	0.8	0.9	0.9
2000.....	0.7	*	*	0.1	0.1	0.5	0.7	0.8	0.8	0.9
1999.....	0.7	*	*	0.2	0.2	0.4	0.6	0.8	0.8	0.9

All races 3/, male

2006.....	1.0	*	*	*	0.3	0.5	1.0	1.1	1.0	1.2
2005.....	1.0	*	*	*	*	0.4	0.9	0.9	1.1	1.1
2004.....	1.1	*	*	*	*	0.5	0.9	1.2	1.2	1.4
2003.....	1.1	*	*	*	0.2	0.6	1.0	0.8	1.0	1.4
2002.....	1.1	*	*	*	0.2	0.6	0.9	0.9	1.2	1.5
2001.....	1.2	*	*	*	0.2	0.7	1.0	1.2	1.5	1.4
2000.....	1.2	*	*	*	0.2	0.6	1.2	1.3	1.3	1.5
1999.....	1.1	*	*	0.2	0.3	0.6	0.9	1.4	1.3	1.4

All races 3/, female

2006.....	0.3	*	*	*	*	0.2	0.3	*	0.3	0.2
2005.....	0.2	*	*	*	*	0.2	*	*	0.2	0.3
2004.....	0.3	*	*	*	*	0.2	0.3	0.3	0.2	0.3
2003.....	0.2	*	*	*	*	*	0.2	0.3	0.2	0.3
2002.....	0.2	*	*	*	*	0.2	*	*	0.2	0.3
2001.....	0.3	*	*	*	*	*	0.3	0.4	0.3	0.4
2000.....	0.2	*	*	*	*	0.3	*	*	0.2	0.2
1999.....	0.3	*	*	*	*	0.2	0.3	0.3	0.3	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

White, both sexes

2006 4/.....	0.7	*	*	0.1	0.2	0.4	0.7	0.7	0.7	0.8
2005 4/.....	0.7	*	*	*	0.2	0.4	0.6	0.6	0.7	0.7
2004 4/.....	0.7	*	*	*	0.2	0.4	0.7	0.9	0.8	0.9
2003 4/.....	0.7	*	*	0.1	0.2	0.4	0.6	0.6	0.7	0.9
2002.....	0.7	*	*	*	0.2	0.5	0.6	0.6	0.8	0.9
2001.....	0.8	*	0.2	0.1	0.2	0.5	0.7	0.9	1.0	1.0
2000.....	0.7	*	*	*	0.2	0.5	0.8	0.9	0.8	0.9
1999.....	0.7	*	*	0.2	0.2	0.4	0.7	1.0	0.9	1.0

White, male

2006 4/.....	1.1	*	*	*	0.3	0.5	1.1	1.2	1.0	1.2
2005 4/.....	1.1	*	*	*	*	0.5	1.0	1.0	1.2	1.1
2004 4/.....	1.2	*	*	*	*	0.5	1.0	1.3	1.4	1.5
2003 4/.....	1.2	*	*	*	*	0.6	1.0	1.0	1.1	1.5
2002.....	1.2	*	*	*	*	0.6	1.1	1.0	1.4	1.6
2001.....	1.3	*	*	*	*	0.8	1.1	1.4	1.6	1.5
2000.....	1.3	*	*	*	*	0.6	1.4	1.5	1.4	1.6
1999.....	1.2	*	*	*	0.3	0.6	1.1	1.5	1.4	1.5

White, female

2006 4/.....	0.3	*	*	*	*	0.3	0.3	*	0.4	0.3
2005 4/.....	0.3	*	*	*	*	0.3	*	*	*	0.4
2004 4/.....	0.3	*	*	*	*	0.3	0.4	0.4	0.3	0.3
2003 4/.....	0.3	*	*	*	*	*	*	0.3	0.3	0.4
2002.....	0.2	*	*	*	*	0.3	*	*	*	0.3
2001.....	0.3	*	*	*	*	*	0.4	0.5	0.3	0.4
2000.....	0.2	*	*	*	*	0.3	*	*	*	*
1999.....	0.3	*	*	*	*	*	0.3	0.3	0.4	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006.....	24.4	24.3	6.1	2.2	1.8	7.3	15.5	17.9	17.7	20.4
2005.....	23.4	22.8	6.3	2.2	2.1	7.1	14.2	15.5	16.2	19.1
2004.....	21.8	22.2	6.1	2.3	2.2	6.9	12.6	13.7	13.8	18.3
2003.....	21.0	19.9	6.8	2.0	2.1	6.3	12.3	12.7	13.7	18.3
2002.....	20.2	20.3	6.4	2.4	2.4	6.6	11.1	11.8	13.6	17.8
2001.....	19.0	20.5	6.8	2.5	2.4	6.2	10.3	10.5	11.6	16.7
2000.....	18.2	18.6	7.4	2.7	2.5	6.3	9.5	10.0	11.8	15.8
1999.....	18.4	17.3	7.7	2.7	2.7	6.2	9.3	10.0	12.1	15.3

All races 3/, male

2006.....	30.4	28.3	7.2	2.7	2.5	11.7	24.5	27.3	26.3	29.1
2005.....	29.0	24.0	7.9	2.6	3.0	11.2	22.2	23.6	23.7	27.1
2004.....	26.8	25.1	7.4	2.9	3.1	11.0	19.8	21.1	20.5	25.7
2003.....	26.0	22.4	8.5	2.5	3.0	10.0	19.4	19.6	20.2	26.2
2002.....	25.2	23.3	8.1	3.1	3.3	10.6	18.0	18.4	20.6	25.9
2001.....	23.8	22.9	8.3	3.3	3.3	9.9	16.8	16.6	17.7	24.8
2000.....	22.9	20.9	9.0	3.4	3.5	10.2	15.5	16.2	18.4	24.2
1999.....	23.3	20.2	9.3	3.5	3.9	10.0	15.3	16.2	19.2	23.4

All races 3/, female

2006.....	18.6	20.2	5.0	1.7	1.1	2.7	5.9	8.1	8.9	11.7
2005.....	18.0	21.5	4.5	1.7	1.2	2.7	5.6	7.0	8.6	10.9
2004.....	16.9	19.3	4.8	1.6	1.3	2.6	4.9	6.0	7.0	10.9
2003.....	16.2	17.3	5.0	1.6	1.2	2.3	4.9	5.5	7.0	10.4
2002.....	15.4	17.2	4.7	1.6	1.5	2.5	3.9	4.9	6.5	9.7
2001.....	14.5	17.9	5.1	1.8	1.5	2.3	3.4	4.2	5.5	8.5
2000.....	13.6	16.2	5.7	1.9	1.4	2.2	3.2	3.7	5.2	7.4
1999.....	13.7	14.2	6.1	2.0	1.5	2.1	3.1	3.7	5.0	7.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nontransport accidents (W00-X59,Y86)										
White, both sexes										
2006 4/.....	26.2	19.7	5.7	1.8	1.7	7.8	17.6	20.2	19.5	21.7
2005 4/.....	24.9	19.8	6.1	1.8	1.9	7.5	15.8	17.3	17.4	20.2
2004 4/.....	23.3	19.0	6.0	1.8	1.9	7.4	14.1	15.2	15.1	19.4
2003 4/.....	22.5	16.4	6.6	1.7	1.8	6.8	13.9	13.9	14.8	19.2
2002.....	21.4	15.9	5.9	1.8	2.0	6.9	12.3	12.7	14.6	18.3
2001.....	20.0	16.9	6.7	2.2	2.1	6.3	11.3	11.0	12.2	17.2
2000.....	18.9	15.1	6.9	2.2	2.2	6.4	10.2	10.5	12.2	16.0
1999.....	19.0	14.3	7.3	2.2	2.4	6.2	9.7	10.4	12.1	15.6
White, male										
2006 4/.....	32.0	22.6	6.8	2.2	2.4	12.3	27.8	30.4	28.5	30.5
2005 4/.....	30.3	20.4	8.0	2.2	2.7	11.9	24.8	26.1	25.3	28.6
2004 4/.....	28.3	22.1	7.3	2.2	2.6	11.7	22.1	23.1	22.2	27.1
2003 4/.....	27.4	18.6	8.3	2.1	2.5	10.8	21.7	21.3	21.7	27.2
2002.....	26.2	19.2	7.4	2.4	2.8	10.9	19.7	19.7	22.0	26.4
2001.....	24.5	18.9	8.4	2.7	2.8	10.0	18.4	17.3	18.6	25.5
2000.....	23.4	17.3	8.4	2.8	3.1	10.3	16.7	16.9	19.1	24.6
1999.....	23.5	16.5	8.8	2.7	3.4	10.0	15.8	16.9	19.2	23.8
White, female										
2006 4/.....	20.5	16.7	4.6	1.3	1.0	3.1	6.6	9.3	9.9	12.6
2005 4/.....	19.6	19.2	4.1	1.3	1.1	2.8	6.1	7.8	9.1	11.5
2004 4/.....	18.5	15.6	4.6	1.3	1.0	2.9	5.4	6.7	7.6	11.5
2003 4/.....	17.7	14.2	4.9	1.3	0.9	2.5	5.4	5.9	7.6	11.0
2002.....	16.8	12.5	4.4	1.2	1.2	2.7	4.3	5.3	6.9	10.1
2001.....	15.6	14.8	4.8	1.7	1.3	2.3	3.8	4.4	5.6	8.6
2000.....	14.6	12.8	5.2	1.6	1.2	2.2	3.3	3.8	5.1	7.3
1999.....	14.6	11.8	5.8	1.6	1.3	2.2	3.2	3.7	4.8	7.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

Nontransport accidents (W00-X59,Y86)

Black, both sexes

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
2006 4/.....	20.1	48.1	8.9	4.1	2.5	6.0	8.1	11.7	14.0	20.2
2005 4/.....	20.0	41.6	7.9	3.9	3.3	5.6	8.2	10.2	14.8	18.2
2004 4/.....	17.6	41.0	7.4	4.5	3.7	5.0	7.5	9.9	11.3	17.2
2003 4/.....	17.2	39.3	8.4	3.7	3.6	4.5	7.4	9.5	11.6	17.7
2002.....	17.8	43.6	9.2	4.7	4.4	6.0	7.4	9.8	12.1	18.9
2001.....	17.6	40.2	8.4	4.1	4.1	5.9	7.2	9.4	11.1	17.7
2000.....	17.6	38.0	10.6	5.2	4.0	6.4	7.7	9.6	12.5	18.0
1999.....	18.9	34.4	10.6	5.6	4.6	6.3	8.8	10.2	14.4	16.7

Black, male

2006 4/.....	27.9	57.0	10.2	4.9	3.3	10.5	12.3	19.1	22.1	29.9
2005 4/.....	27.3	46.1	8.9	4.6	4.8	9.1	12.7	16.0	21.3	26.3
2004 4/.....	23.9	42.4	8.9	6.1	4.9	8.2	11.7	15.6	17.4	23.7
2003 4/.....	23.5	44.0	10.6	4.2	5.0	7.3	11.3	14.8	17.5	26.4
2002.....	24.8	45.9	12.1	6.1	6.5	10.1	12.4	16.2	18.8	29.1
2001.....	24.4	46.4	9.7	6.0	5.7	9.6	11.8	15.1	16.5	26.1
2000.....	24.9	40.9	12.6	6.9	5.6	10.3	12.6	15.9	18.9	27.1
1999.....	26.8	40.0	13.0	7.4	6.6	10.8	15.5	16.7	22.3	25.7

Black, female

2006 4/.....	12.9	38.8	7.5	3.2	1.6	1.4	3.7	4.7	6.8	11.5
2005 4/.....	13.3	37.0	6.9	3.2	1.8	2.1	3.7	4.7	9.0	10.9
2004 4/.....	11.9	39.5	5.9	2.9	2.5	1.8	3.3	4.5	5.8	11.5
2003 4/.....	11.6	34.3	6.0	3.1	2.1	1.7	3.5	4.5	6.4	9.8
2002.....	11.4	41.2	6.2	3.3	2.3	1.7	2.4	4.0	6.2	9.8
2001.....	11.5	33.7	7.1	2.2	2.5	2.1	2.7	4.3	6.2	10.2
2000.....	11.0	35.1	8.5	3.5	2.3	2.3	3.0	3.8	6.7	9.9
1999.....	11.7	28.5	8.1	3.7	2.5	1.6	2.5	4.3	7.3	8.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Falls (W00-W19)										
All races 3/, both sexes										
2006.....	7.0	0.6	0.2	*	0.1	0.4	0.8	0.7	0.9	1.1
2005.....	6.6	*	0.2	*	*	0.4	0.8	0.7	0.8	1.1
2004.....	6.4	0.6	0.3	*	0.1	0.4	0.7	0.8	0.8	1.1
2003.....	5.9	*	0.3	*	0.1	0.4	0.7	0.7	0.7	1.2
2002.....	5.6	*	0.2	*	0.1	0.4	0.8	0.7	0.8	1.2
2001.....	5.3	0.6	0.2	0.2	0.2	0.4	0.9	0.8	0.9	1.2
2000.....	4.7	*	0.2	*	0.1	0.5	0.7	0.7	0.8	1.0
1999.....	4.7	*	0.4	0.1	0.1	0.6	0.7	0.8	0.9	1.2
All races 3/, male										
2006.....	7.3	*	0.3	*	*	0.6	1.3	1.1	1.5	1.8
2005.....	7.0	*	0.2	*	*	0.6	1.2	1.2	1.3	1.8
2004.....	6.8	*	0.4	*	*	0.7	1.3	1.3	1.4	1.8
2003.....	6.2	*	0.5	*	0.2	0.7	1.1	1.3	1.1	2.0
2002.....	6.0	*	0.4	*	*	0.6	1.4	1.2	1.4	1.7
2001.....	5.8	*	0.3	0.2	0.2	0.8	1.5	1.5	1.5	2.1
2000.....	5.2	*	0.3	*	*	0.8	1.2	1.2	1.5	1.8
1999.....	5.2	*	0.5	*	0.2	0.9	1.2	1.4	1.5	2.0
All races 3/, female										
2006.....	6.6	*	*	*	*	*	*	*	0.3	0.4
2005.....	6.3	*	*	*	*	*	0.3	*	0.3	0.4
2004.....	6.0	*	*	*	*	*	*	0.2	0.2	0.5
2003.....	5.6	*	*	*	*	*	0.2	*	0.3	0.3
2002.....	5.3	*	*	*	*	*	*	*	0.3	0.6
2001.....	4.8	*	*	*	*	*	*	*	0.3	0.4
2000.....	4.3	*	*	*	*	*	0.2	*	*	0.3
1999.....	4.3	*	*	*	*	*	*	*	0.3	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Falls (W00-W19)										
White, both sexes										
2006 4/.....	7.9	*	0.2	*	*	0.4	0.9	0.7	1.0	1.2
2005 4/.....	7.5	*	*	*	*	0.4	0.9	0.8	0.8	1.1
2004 4/.....	7.3	*	0.3	*	0.1	0.5	0.8	0.8	0.9	1.2
2003 4/.....	6.7	*	0.3	*	0.2	0.5	0.8	0.8	0.8	1.2
2002.....	6.4	*	0.2	*	*	0.4	0.9	0.8	0.9	1.2
2001.....	6.0	*	0.2	0.2	0.2	0.5	1.0	0.9	0.9	1.3
2000.....	5.3	*	0.2	*	*	0.6	0.8	0.8	0.9	1.0
1999.....	5.2	*	0.4	*	0.2	0.6	0.8	0.9	0.9	1.3
White, male										
2006 4/.....	8.2	*	*	*	*	0.7	1.5	1.2	1.7	1.9
2005 4/.....	7.8	*	*	*	*	0.7	1.4	1.4	1.3	1.9
2004 4/.....	7.6	*	0.4	*	*	0.8	1.4	1.4	1.5	2.0
2003 4/.....	6.9	*	0.5	*	*	0.8	1.3	1.4	1.2	2.1
2002.....	6.6	*	*	*	*	0.7	1.6	1.4	1.5	1.8
2001.....	6.4	*	0.3	0.2	0.2	0.9	1.8	1.6	1.6	2.1
2000.....	5.7	*	*	*	*	1.0	1.4	1.4	1.5	1.7
1999.....	5.6	*	0.5	*	*	1.1	1.3	1.5	1.5	2.0
White, female										
2006 4/.....	7.7	*	*	*	*	*	*	*	0.3	0.4
2005 4/.....	7.3	*	*	*	*	*	0.3	*	*	0.4
2004 4/.....	7.0	*	*	*	*	*	*	*	0.3	0.4
2003 4/.....	6.5	*	*	*	*	*	0.3	*	0.3	0.3
2002.....	6.2	*	*	*	*	*	*	*	0.3	0.7
2001.....	5.5	*	*	*	*	*	*	*	0.3	0.4
2000.....	5.0	*	*	*	*	*	0.3	*	*	0.3
1999.....	4.9	*	*	*	*	*	*	*	0.3	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
All races 3/, both sexes										
2006.....	1.2	1.2	2.8	0.7	0.6	1.5	1.4	1.2	0.9	1.0
2005.....	1.2	1.6	3.0	0.6	0.6	1.5	1.6	1.0	0.9	1.0
2004.....	1.1	1.5	2.7	0.7	0.7	1.5	1.3	1.1	0.8	0.9
2003.....	1.1	1.4	2.9	0.6	0.7	1.4	1.4	1.0	0.8	1.0
2002.....	1.2	1.6	2.9	0.8	0.8	1.6	1.5	1.2	1.0	1.2
2001.....	1.2	1.7	3.0	0.8	0.8	1.6	1.4	1.1	0.8	1.0
2000.....	1.2	2.0	3.2	1.0	0.8	1.8	1.5	1.1	1.0	1.1
1999.....	1.3	1.8	3.2	0.9	0.9	1.8	1.5	1.1	1.1	1.1
All races 3/, male										
2006.....	1.9	1.3	3.5	1.0	0.8	2.6	2.5	2.2	1.6	1.6
2005.....	1.9	1.3	4.1	0.8	1.0	2.6	2.8	1.7	1.5	1.7
2004.....	1.8	1.8	3.2	0.9	1.0	2.6	2.3	1.9	1.4	1.6
2003.....	1.8	1.7	3.7	0.9	1.0	2.5	2.5	1.8	1.4	1.7
2002.....	1.9	1.9	3.9	1.2	1.1	2.8	2.8	2.1	1.7	2.0
2001.....	1.8	1.5	3.7	1.2	1.1	2.8	2.5	1.9	1.4	1.6
2000.....	2.0	2.2	4.0	1.3	1.2	3.3	2.5	1.9	1.7	1.9
1999.....	2.0	2.0	4.0	1.3	1.3	3.2	2.8	2.0	2.0	1.9
All races 3/, female										
2006.....	0.5	1.2	2.1	0.4	0.2	0.3	0.4	0.3	*	0.4
2005.....	0.5	1.8	2.0	0.4	0.3	0.3	0.3	0.3	0.3	0.3
2004.....	0.5	1.2	2.1	0.4	0.3	0.3	0.2	0.3	0.2	0.2
2003.....	0.5	1.2	2.0	0.4	0.3	*	0.2	*	*	0.3
2002.....	0.5	1.2	1.9	0.3	0.4	0.3	*	0.3	0.3	0.4
2001.....	0.5	1.9	2.2	0.5	0.5	0.3	0.3	0.3	0.2	0.4
2000.....	0.5	1.7	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3
1999.....	0.5	1.6	2.4	0.5	0.4	0.3	0.3	*	0.3	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
White, both sexes										
2006 4/.....	1.2	1.4	2.9	0.5	0.5	1.3	1.5	1.3	0.8	1.0
2005 4/.....	1.2	1.8	3.3	0.5	0.5	1.3	1.6	1.0	0.9	0.9
2004 4/.....	1.1	1.6	3.0	0.5	0.4	1.2	1.3	1.1	0.8	0.9
2003 4/.....	1.1	1.4	3.1	0.6	0.5	1.3	1.3	1.0	0.8	1.0
2002.....	1.1	1.6	3.0	0.6	0.5	1.4	1.5	1.1	1.0	1.1
2001.....	1.1	1.7	3.3	0.6	0.5	1.3	1.4	1.0	0.7	1.0
2000.....	1.2	1.6	3.5	0.8	0.6	1.6	1.4	1.1	1.0	1.1
1999.....	1.2	1.5	3.5	0.7	0.5	1.6	1.5	1.0	1.1	1.1
White, male										
2006 4/.....	1.8	1.4	3.6	0.8	0.7	2.1	2.4	2.2	1.5	1.6
2005 4/.....	1.8	1.3	4.5	0.6	0.6	2.2	2.8	1.8	1.4	1.5
2004 4/.....	1.7	1.8	3.5	0.7	0.6	2.1	2.2	1.8	1.3	1.5
2003 4/.....	1.8	1.5	4.0	0.8	0.7	2.3	2.4	1.7	1.3	1.7
2002.....	1.8	1.9	4.0	0.9	0.6	2.4	2.7	1.9	1.7	1.9
2001.....	1.7	1.6	4.2	0.8	0.6	2.3	2.4	1.7	1.2	1.5
2000.....	1.9	1.8	4.4	1.0	0.8	2.8	2.4	1.8	1.7	1.8
1999.....	1.9	1.7	4.3	0.9	0.7	2.8	2.5	1.7	1.9	1.8
White, female										
2006 4/.....	0.5	1.4	2.3	0.3	*	0.3	0.4	0.3	*	0.4
2005 4/.....	0.5	2.2	2.1	0.4	0.3	0.3	0.3	0.3	0.3	0.3
2004 4/.....	0.5	1.4	2.4	0.3	*	*	0.3	0.3	0.3	0.3
2003 4/.....	0.5	1.3	2.2	0.3	*	*	*	*	*	0.3
2002.....	0.4	*	2.0	0.3	0.3	0.3	*	0.3	*	0.4
2001.....	0.5	1.8	2.4	0.5	0.3	*	0.3	0.3	*	0.4
2000.....	0.5	*	2.5	0.5	0.3	0.3	0.3	0.3	*	0.4
1999.....	0.5	1.4	2.6	0.4	*	0.3	0.3	*	0.3	0.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
Black, both sexes										
2006 4/.....	1.4	*	2.6	1.4	1.0	2.4	1.4	1.1	1.2	1.0
2005 4/.....	1.4	*	2.2	1.2	1.4	2.2	1.9	1.1	1.1	1.5
2004 4/.....	1.4	*	1.9	1.3	1.8	2.4	1.2	1.3	0.9	1.1
2003 4/.....	1.3	*	2.1	1.0	1.6	1.8	1.4	1.1	1.1	0.8
2002.....	1.6	*	2.5	1.8	2.2	2.3	1.9	1.6	1.2	1.5
2001.....	1.6	*	1.9	1.7	2.3	2.9	1.5	1.2	1.1	1.2
2000.....	1.6	4.2	2.2	1.9	2.1	2.8	1.7	1.3	0.9	1.2
1999.....	1.8	3.5	2.4	2.2	2.6	2.7	1.9	1.8	1.4	1.1
Black, male										
2006 4/.....	2.4	*	3.6	2.1	1.6	4.7	2.5	2.3	2.2	2.1
2005 4/.....	2.5	*	2.7	1.6	2.6	4.2	3.3	1.9	2.1	2.7
2004 4/.....	2.3	*	2.8	1.6	2.7	4.3	2.3	2.2	1.6	2.2
2003 4/.....	2.2	*	3.0	1.4	2.4	3.4	2.6	2.1	2.2	1.7
2002.....	2.8	*	3.4	2.8	3.5	4.3	3.7	3.1	2.0	2.8
2001.....	2.7	*	2.2	2.7	3.2	5.2	3.1	2.6	2.2	2.2
2000.....	2.7	*	2.8	2.7	3.2	5.2	3.3	2.3	1.8	2.3
1999.....	3.1	*	3.2	3.4	3.9	5.0	3.8	3.5	2.7	2.2
Black, female										
2006 4/.....	0.4	*	1.5	*	*	*	*	*	*	*
2005 4/.....	0.4	*	1.7	*	*	*	*	*	*	*
2004 4/.....	0.5	*	*	*	*	*	*	*	*	*
2003 4/.....	0.4	*	*	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*	*
2001.....	0.5	*	1.6	*	1.3	*	*	*	*	*
2000.....	0.6	*	1.7	*	*	*	*	*	*	*
1999.....	0.5	*	1.6	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
All races 3/, both sexes										
2006.....	1.0	0.7	1.2	0.6	0.3	0.4	0.5	0.5	0.5	0.6
2005.....	1.1	0.8	1.3	0.7	0.4	0.3	0.5	0.6	0.5	0.7
2004.....	1.1	0.6	1.4	0.9	0.4	0.3	0.6	0.7	0.5	0.7
2003.....	1.2	0.7	1.4	0.7	0.4	0.4	0.6	0.6	0.6	0.8
2002.....	1.1	0.9	1.4	0.8	0.5	0.4	0.5	0.6	0.7	0.7
2001.....	1.2	1.2	1.5	0.8	0.4	0.4	0.7	0.6	0.6	0.9
2000.....	1.2	1.0	1.9	0.9	0.4	0.4	0.6	0.6	0.6	0.8
1999.....	1.2	1.1	2.0	0.8	0.5	0.4	0.6	0.7	0.7	0.9
All races 3/, male										
2006.....	1.2	*	1.4	0.6	0.3	0.4	0.6	0.7	0.6	0.8
2005.....	1.3	*	1.5	0.8	0.4	0.4	0.6	0.7	0.6	0.9
2004.....	1.3	*	1.5	1.1	0.4	0.4	0.6	0.9	0.7	0.9
2003.....	1.4	*	1.6	0.7	0.4	0.4	0.7	0.7	0.8	1.1
2002.....	1.4	1.0	1.6	0.9	0.6	0.6	0.7	0.8	0.9	0.9
2001.....	1.4	1.5	1.8	1.0	0.4	0.4	0.9	0.8	0.8	1.2
2000.....	1.5	1.0	2.2	1.0	0.3	0.4	0.8	0.8	0.8	1.2
1999.....	1.5	1.1	2.2	0.9	0.5	0.6	0.8	0.9	1.0	1.2
All races 3/, female										
2006.....	0.8	*	1.0	0.6	0.3	0.3	0.3	0.3	0.4	0.5
2005.....	0.9	*	1.0	0.6	0.4	0.2	0.4	0.4	0.5	0.4
2004.....	0.9	*	1.3	0.6	0.4	0.2	0.5	0.4	0.3	0.4
2003.....	1.0	1.1	1.2	0.7	0.4	0.3	0.5	0.5	0.5	0.5
2002.....	0.8	*	1.2	0.6	0.4	0.3	0.4	0.4	0.6	0.4
2001.....	0.9	1.0	1.1	0.6	0.4	0.3	0.4	0.4	0.4	0.5
2000.....	0.9	*	1.5	0.7	0.5	0.4	0.4	0.4	0.4	0.5
1999.....	0.9	1.1	1.7	0.7	0.4	0.3	0.4	0.5	0.4	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
White, both sexes										
2006 4/.....	1.0	*	0.9	0.4	0.3	0.4	0.5	0.4	0.5	0.6
2005 4/.....	1.0	0.7	1.0	0.5	0.3	0.2	0.5	0.5	0.5	0.6
2004 4/.....	1.0	*	1.1	0.6	0.3	0.3	0.5	0.6	0.5	0.7
2003 4/.....	1.1	0.7	1.2	0.5	0.2	0.3	0.6	0.6	0.7	0.8
2002.....	1.0	*	1.0	0.5	0.4	0.4	0.5	0.5	0.7	0.6
2001.....	1.0	1.1	1.2	0.7	0.4	0.3	0.7	0.5	0.6	0.8
2000.....	1.0	0.8	1.3	0.7	0.3	0.4	0.5	0.5	0.5	0.7
1999.....	1.0	0.9	1.5	0.6	0.4	0.4	0.6	0.7	0.7	0.8
White, male										
2006 4/.....	1.2	*	1.2	0.4	0.3	0.5	0.6	0.6	0.6	0.7
2005 4/.....	1.2	*	1.3	0.6	0.3	0.3	0.6	0.7	0.6	0.8
2004 4/.....	1.2	*	1.2	0.7	0.3	0.4	0.6	0.8	0.7	0.8
2003 4/.....	1.3	*	1.4	0.5	*	0.4	0.8	0.7	0.8	1.1
2002.....	1.2	*	1.1	0.6	0.5	0.5	0.6	0.7	0.9	0.8
2001.....	1.3	*	1.6	0.8	0.4	0.4	0.9	0.7	0.8	1.0
2000.....	1.3	*	1.6	0.7	0.3	0.4	0.8	0.7	0.7	1.0
1999.....	1.3	*	1.7	0.6	0.4	0.6	0.7	0.9	1.0	1.0
White, female										
2006 4/.....	0.8	*	0.7	0.5	*	0.3	0.3	0.3	0.4	0.5
2005 4/.....	0.8	*	0.7	0.4	0.3	*	0.4	0.4	0.3	0.4
2004 4/.....	0.8	*	1.0	0.5	0.3	0.3	0.5	0.4	0.4	0.5
2003 4/.....	0.9	*	1.0	0.5	0.3	0.3	0.5	0.5	0.5	0.5
2002.....	0.7	*	1.0	0.4	*	*	0.3	0.3	0.5	0.4
2001.....	0.8	*	0.8	0.5	0.3	0.3	0.4	*	0.4	0.5
2000.....	0.8	*	1.1	0.6	0.4	0.4	0.3	0.4	0.3	0.4
1999.....	0.8	*	1.3	0.6	0.3	0.3	0.4	0.4	0.3	0.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental exposure to smoke, fire and flames (X00-X09)

Black, both sexes

2006 4/.....	1.8	*	3.0	1.5	*	*	0.7	0.8	0.9	1.1
2005 4/.....	1.9	*	2.8	1.6	0.9	*	*	0.9	1.0	1.1
2004 4/.....	2.1	*	2.8	2.3	0.9	*	0.8	1.2	0.8	0.9
2003 4/.....	2.0	*	3.0	1.9	1.0	0.6	*	0.9	0.8	1.0
2002.....	2.1	*	3.5	1.9	0.9	0.7	0.9	1.2	1.1	1.0
2001.....	2.3	*	3.3	1.8	0.8	*	1.0	1.4	1.0	1.5
2000.....	2.5	*	4.8	2.1	0.8	*	1.0	1.1	1.2	1.5
1999.....	2.5	*	4.5	2.1	0.9	*	0.9	1.1	0.9	1.9

Black, male

2006 4/.....	2.1	*	3.0	1.5	*	*	*	*	*	*
2005 4/.....	2.4	*	2.8	1.7	*	*	*	*	*	*
2004 4/.....	2.6	*	2.9	3.1	*	*	*	1.6	*	*
2003 4/.....	2.4	*	3.2	1.9	*	*	*	*	*	*
2002.....	2.6	*	4.1	2.2	*	*	*	*	*	1.5
2001.....	2.9	*	3.6	2.3	*	*	*	*	*	2.5
2000.....	3.2	*	5.6	2.6	*	*	*	*	1.5	2.2
1999.....	3.0	*	5.0	2.4	*	*	*	*	*	2.8

Black, female

2006 4/.....	1.5	*	2.9	1.5	*	*	*	*	*	*
2005 4/.....	1.5	*	2.7	1.6	*	*	*	*	*	*
2004 4/.....	1.6	*	2.7	1.4	*	*	*	*	*	*
2003 4/.....	1.7	*	2.7	1.8	*	*	*	*	*	*
2002.....	1.6	*	2.8	1.7	*	*	*	*	*	*
2001.....	1.7	*	2.9	1.2	*	*	*	1.5	*	*
2000.....	2.0	*	4.1	1.6	*	*	*	*	*	*
1999.....	2.0	*	3.9	1.8	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
All races 3/, both sexes										
2006.....	9.2	*	0.2	*	0.2	3.5	10.4	13.0	13.0	15.1
2005.....	8.0	0.5	0.1	*	0.2	3.0	8.8	10.6	11.3	13.5
2004.....	7.1	*	*	*	0.2	3.1	7.7	8.9	9.3	12.8
2003.....	6.7	0.5	0.3	*	0.2	2.5	7.1	7.9	9.3	12.5
2002.....	6.1	0.6	0.2	*	0.1	2.4	5.9	6.9	8.6	11.9
2001.....	4.9	*	0.2	*	0.2	2.0	4.9	5.7	6.9	10.4
2000.....	4.5	*	0.2	*	0.1	1.7	4.3	5.1	6.8	9.5
1999.....	4.4	*	0.2	*	0.1	1.3	3.8	4.9	6.8	9.1
All races 3/, male										
2006.....	12.6	*	*	*	0.2	5.2	16.0	19.2	18.6	20.5
2005.....	10.9	*	*	*	0.2	4.5	13.4	15.5	15.7	18.0
2004.....	9.6	*	*	*	0.2	4.6	11.8	13.3	13.0	16.8
2003.....	9.2	*	0.3	*	0.2	3.8	10.9	11.7	13.1	16.8
2002.....	8.5	1.0	*	*	*	3.5	9.1	10.4	12.5	16.5
2001.....	7.1	*	0.3	*	*	3.0	7.6	8.7	9.9	14.6
2000.....	6.6	*	0.3	*	*	2.6	6.7	7.9	10.1	13.7
1999.....	6.5	*	*	*	*	1.9	5.9	7.7	10.3	13.3
All races 3/, female										
2006.....	5.9	*	*	*	*	1.6	4.4	6.5	7.3	9.6
2005.....	5.1	*	*	*	*	1.5	3.9	5.4	6.7	8.9
2004.....	4.7	*	*	*	0.2	1.5	3.4	4.4	5.4	8.7
2003.....	4.3	*	0.3	*	*	1.2	3.1	4.0	5.4	8.1
2002.....	3.7	*	*	*	*	1.2	2.6	3.3	4.7	7.2
2001.....	2.9	*	*	*	*	1.0	1.9	2.6	3.9	6.1
2000.....	2.5	*	*	*	*	0.8	1.7	2.1	3.5	5.3
1999.....	2.3	*	*	*	*	0.7	1.6	2.0	3.2	4.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
White, both sexes										
2006 4/.....	9.7	*	*	*	0.2	4.2	12.3	15.1	14.6	16.2
2005 4/.....	8.4	*	*	*	0.2	3.6	10.4	12.1	12.5	14.6
2004 4/.....	7.6	*	*	*	0.2	3.7	9.1	10.4	10.3	13.8
2003 4/.....	7.1	*	0.3	*	0.1	3.0	8.5	9.1	10.2	13.3
2002.....	6.3	*	*	*	0.2	2.8	6.9	7.8	9.5	12.5
2001.....	5.0	*	0.2	*	0.2	2.3	5.6	6.4	7.4	10.8
2000.....	4.6	*	*	*	0.1	2.0	4.9	5.5	7.2	9.8
1999.....	4.3	*	0.2	*	0.1	1.5	4.2	5.2	6.9	9.3
White, male										
2006 4/.....	13.2	*	*	*	*	6.4	19.0	22.0	20.6	21.7
2005 4/.....	11.4	*	*	*	*	5.5	15.8	17.7	17.4	19.5
2004 4/.....	10.2	*	*	*	0.3	5.6	13.8	15.3	14.5	18.2
2003 4/.....	9.7	*	*	*	*	4.6	12.9	13.4	14.4	17.7
2002.....	8.8	*	*	*	*	4.1	10.6	11.6	13.6	17.3
2001.....	7.2	*	*	*	*	3.5	8.7	9.5	10.7	15.2
2000.....	6.7	*	*	*	*	3.1	7.8	8.6	10.7	14.3
1999.....	6.4	*	*	*	*	2.1	6.5	8.3	10.6	13.7
White, female										
2006 4/.....	6.3	*	*	*	*	1.9	5.2	7.8	8.3	10.4
2005 4/.....	5.4	*	*	*	*	1.7	4.4	6.2	7.3	9.5
2004 4/.....	5.0	*	*	*	*	1.7	4.0	5.1	6.0	9.3
2003 4/.....	4.5	*	*	*	*	1.4	3.8	4.5	5.8	8.7
2002.....	3.9	*	*	*	*	1.4	3.0	3.8	5.1	7.5
2001.....	2.9	*	*	*	*	1.0	2.3	3.0	3.9	6.2
2000.....	2.5	*	*	*	*	0.9	1.8	2.3	3.6	5.2
1999.....	2.3	*	*	*	*	0.8	1.7	2.1	3.1	4.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
Black, both sexes										
2006 4/.....	8.9	*	*	*	*	0.9	3.5	6.8	9.3	14.5
2005 4/.....	7.6	*	*	*	*	0.7	3.0	5.6	8.8	11.6
2004 4/.....	6.5	*	*	*	*	0.9	2.9	4.5	6.2	11.0
2003 4/.....	6.3	*	*	*	*	*	2.3	4.0	7.3	11.8
2002.....	6.3	*	*	*	*	*	2.3	4.4	6.8	11.9
2001.....	5.8	*	*	*	*	0.9	2.3	4.0	6.3	11.1
2000.....	5.5	*	*	*	*	0.7	2.1	4.3	6.8	10.7
1999.....	5.7	*	*	*	*	*	2.8	4.2	8.0	10.3
Black, male										
2006 4/.....	12.9	*	*	*	*	1.2	5.0	11.0	14.3	20.6
2005 4/.....	10.8	*	*	*	*	*	4.2	8.6	12.0	15.5
2004 4/.....	8.9	*	*	*	*	*	4.4	6.5	8.7	13.5
2003 4/.....	8.8	*	*	*	*	*	3.5	5.7	10.2	16.7
2002.....	9.1	*	*	*	*	*	3.5	7.0	9.9	17.0
2001.....	8.4	*	*	*	*	*	3.8	6.7	8.2	15.4
2000.....	8.2	*	*	*	*	*	2.7	7.1	9.6	14.9
1999.....	8.7	*	*	*	*	*	4.7	6.7	11.5	14.9
Black, female										
2006 4/.....	5.3	*	*	*	*	*	2.1	2.8	4.8	9.0
2005 4/.....	4.8	*	*	*	*	*	1.7	2.8	5.9	8.1
2004 4/.....	4.3	*	*	*	*	*	1.4	2.5	4.0	8.8
2003 4/.....	3.9	*	*	*	*	*	*	2.4	4.8	7.4
2002.....	3.7	*	*	*	*	*	*	2.0	4.0	7.4
2001.....	3.5	*	*	*	*	*	*	1.6	4.6	7.3
2000.....	3.1	*	*	*	*	*	1.5	1.7	4.3	7.1
1999.....	3.0	*	*	*	*	*	*	2.0	4.9	6.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes

2006.....	5.8	21.5	1.6	0.6	0.6	1.2	1.9	2.1	2.2	2.5
2005.....	6.3	19.5	1.5	0.6	0.7	1.4	2.0	2.3	2.4	2.6
2004.....	5.8	19.2	1.6	0.6	0.6	1.2	1.9	1.9	2.2	2.6
2003.....	5.9	16.9	1.8	0.5	0.6	1.1	2.0	2.0	2.0	2.6
2002.....	6.0	16.8	1.6	0.6	0.8	1.3	1.8	2.0	2.1	2.6
2001.....	6.2	16.6	1.8	0.6	0.7	1.2	2.0	1.8	2.2	2.9
2000.....	6.2	15.0	1.7	0.6	0.7	1.4	2.0	2.1	2.3	3.0
1999.....	6.6	13.8	1.9	0.7	0.9	1.4	2.0	2.2	2.3	2.7

All races 3/, male

2006.....	7.0	25.2	1.7	0.8	0.8	2.0	3.2	3.5	3.5	4.2
2005.....	7.4	21.1	1.8	0.8	1.0	2.3	3.3	3.8	4.0	4.2
2004.....	6.8	21.7	2.0	0.8	1.0	2.0	3.1	3.1	3.7	4.1
2003.....	6.9	19.5	2.2	0.6	0.9	1.8	3.3	3.3	3.4	4.0
2002.....	6.9	18.8	2.0	0.7	1.1	2.1	3.1	3.2	3.6	4.3
2001.....	7.2	18.9	2.1	0.7	1.0	1.9	3.4	3.1	3.7	4.8
2000.....	7.2	16.8	2.1	0.7	1.1	2.1	3.5	3.7	3.9	5.1
1999.....	7.6	16.2	2.3	0.9	1.2	2.3	3.4	3.7	3.7	4.5

All races 3/, female

2006.....	4.7	17.6	1.5	0.4	0.3	0.4	0.6	0.7	0.8	0.8
2005.....	5.1	17.9	1.2	0.5	0.3	0.6	0.7	0.7	0.8	0.9
2004.....	4.8	16.5	1.1	0.4	0.3	0.4	0.5	0.6	0.7	1.0
2003.....	4.9	14.1	1.3	0.3	0.2	0.3	0.7	0.7	0.6	1.1
2002.....	5.0	14.6	1.2	0.4	0.4	0.5	0.5	0.7	0.6	1.0
2001.....	5.3	14.2	1.5	0.4	0.3	0.6	0.6	0.6	0.6	1.1
2000.....	5.2	13.2	1.4	0.4	0.3	0.5	0.4	0.6	0.7	0.8
1999.....	5.6	11.3	1.5	0.5	0.4	0.5	0.5	0.6	0.8	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

White, both sexes

2006 4/.....	6.2	16.9	1.5	0.6	0.6	1.2	2.2	2.3	2.3	2.6
2005 4/.....	6.5	16.6	1.5	0.6	0.7	1.5	2.1	2.5	2.5	2.6
2004 4/.....	6.2	16.0	1.5	0.6	0.6	1.4	2.0	2.1	2.3	2.6
2003 4/.....	6.3	13.6	1.7	0.5	0.6	1.2	2.1	2.1	2.2	2.6
2002.....	6.3	12.9	1.4	0.5	0.8	1.4	2.0	2.2	2.3	2.7
2001.....	6.6	13.4	1.7	0.6	0.7	1.4	2.2	1.9	2.2	3.0
2000.....	6.6	12.2	1.6	0.6	0.8	1.4	2.1	2.3	2.4	3.0
1999.....	6.9	11.2	1.8	0.7	0.9	1.6	2.2	2.2	2.3	2.9

White, male

2006 4/.....	7.4	19.6	1.6	0.8	0.9	2.0	3.7	3.8	3.8	4.4
2005 4/.....	7.7	17.8	1.9	0.8	1.0	2.4	3.4	4.1	4.2	4.4
2004 4/.....	7.2	19.0	2.0	0.7	1.0	2.2	3.4	3.4	3.8	4.2
2003 4/.....	7.3	15.7	2.2	0.6	0.9	2.0	3.6	3.6	3.6	4.1
2002.....	7.3	15.3	1.8	0.7	1.0	2.4	3.4	3.6	3.8	4.3
2001.....	7.5	15.2	2.0	0.7	1.0	2.1	3.7	3.2	3.7	5.0
2000.....	7.4	14.0	2.0	0.7	1.2	2.2	3.7	3.8	4.1	5.2
1999.....	7.8	12.9	2.1	1.0	1.3	2.5	3.7	3.9	3.8	4.8

White, female

2006 4/.....	5.1	14.1	1.4	0.3	0.3	0.4	0.5	0.6	0.7	0.8
2005 4/.....	5.4	15.3	1.1	0.5	0.3	0.5	0.6	0.7	0.8	0.8
2004 4/.....	5.2	12.8	0.9	0.4	0.3	0.4	0.5	0.7	0.8	1.0
2003 4/.....	5.3	11.4	1.2	0.3	*	0.4	0.6	0.6	0.7	1.1
2002.....	5.4	10.5	1.1	0.4	0.5	0.5	0.5	0.7	0.6	1.0
2001.....	5.8	11.5	1.4	0.5	0.3	0.6	0.6	0.6	0.6	1.0
2000.....	5.7	10.3	1.2	0.4	0.3	0.5	0.4	0.6	0.7	0.8
1999.....	6.1	9.5	1.4	0.4	0.5	0.6	0.5	0.6	0.7	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Black, both sexes

2006 4/.....	5.0	44.7	2.2	0.7	0.6	1.1	1.1	1.8	1.9	2.4
2005 4/.....	6.0	37.4	1.8	*	0.7	1.3	1.8	1.8	2.3	2.7
2004 4/.....	4.8	37.1	1.8	*	0.6	0.6	1.4	1.6	2.2	2.8
2003 4/.....	4.9	34.9	2.0	*	*	0.7	1.9	2.0	1.6	2.7
2002.....	5.2	37.5	2.4	0.7	0.9	1.0	1.2	1.5	1.9	3.0
2001.....	5.4	34.2	2.5	*	0.7	0.8	1.3	1.8	1.9	2.6
2000.....	5.4	30.6	2.3	0.7	0.7	1.3	1.8	2.0	2.4	3.1
1999.....	6.1	27.9	2.5	0.7	0.7	1.0	1.6	2.2	2.6	2.2

Black, male

2006 4/.....	6.4	53.3	2.2	*	*	1.8	1.4	2.5	3.1	4.2
2005 4/.....	7.6	41.0	1.9	*	*	2.0	2.8	2.9	3.3	4.3
2004 4/.....	6.1	38.3	1.7	*	*	*	2.2	2.8	3.8	4.6
2003 4/.....	6.4	40.5	2.8	*	*	*	2.6	3.0	2.9	4.6
2002.....	6.6	38.9	3.3	*	1.5	1.5	1.9	2.3	3.4	5.2
2001.....	6.8	40.4	2.8	*	*	*	2.0	2.8	3.6	3.7
2000.....	7.3	32.0	2.5	*	1.2	1.8	3.2	3.5	3.8	5.5
1999.....	8.0	33.2	3.4	*	1.2	1.8	3.0	3.5	4.4	3.5

Black, female

2006 4/.....	3.6	35.8	2.3	*	*	*	*	*	*	*
2005 4/.....	4.5	33.6	1.6	*	*	*	*	*	*	1.4
2004 4/.....	3.6	35.8	2.0	*	*	*	*	*	*	*
2003 4/.....	3.6	29.1	*	*	*	*	1.3	*	*	*
2002.....	4.0	36.0	*	*	*	*	*	*	*	*
2001.....	4.1	27.7	2.2	*	*	*	*	*	*	1.5
2000.....	3.8	29.1	2.0	*	*	*	*	*	*	*
1999.....	4.4	22.5	1.6	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
All races 3/, both sexes										
2006.....	11.1	*	1.0	7.3	12.5	12.4	12.3	14.7
2005.....	11.0	*	1.3	7.7	12.4	11.8	13.0	13.8
2004.....	11.0	*	1.3	8.2	12.5	12.3	13.0	14.0
2003.....	10.8	*	1.2	7.3	12.1	11.9	13.5	13.8
2002.....	11.0	*	1.2	7.4	12.4	12.8	12.5	14.3
2001 5/.....	10.8	*	1.3	7.9	12.0	12.6	13.0	14.3
2000.....	10.4	*	1.5	8.0	12.5	12.1	12.0	13.7
1999.....	10.5	*	1.2	8.0	12.3	12.3	13.1	14.4
All races 3/, male										
2006.....	17.8	*	1.4	11.5	20.8	19.7	19.7	22.7
2005.....	17.7	*	1.9	12.1	20.2	19.2	20.6	21.4
2004.....	17.7	*	1.7	12.6	20.8	20.2	20.6	21.7
2003.....	17.6	*	1.7	11.6	20.2	19.5	21.6	21.9
2002.....	17.9	*	1.8	12.2	20.8	20.8	20.2	22.7
2001 5/.....	17.6	*	1.9	12.9	20.5	21.0	21.0	22.7
2000.....	17.1	*	2.3	13.0	21.4	20.0	19.2	21.7
1999.....	17.1	*	1.9	13.0	20.8	20.3	21.1	22.7
All races 3/, female										
2006.....	4.6	*	0.6	2.8	3.6	4.7	4.7	6.6
2005.....	4.5	*	0.7	3.0	4.0	4.1	5.3	6.1
2004.....	4.6	*	1.0	3.5	3.6	4.2	5.3	6.3
2003.....	4.3	*	0.5	2.7	3.4	4.0	5.2	5.7
2002.....	4.3	*	0.6	2.4	3.5	4.5	4.7	6.0
2001 5/.....	4.1	*	0.6	2.7	3.1	4.0	4.8	5.9
2000.....	4.0	*	0.6	2.7	3.2	4.0	4.6	5.8
1999.....	4.0	*	0.5	2.7	3.4	4.1	5.1	6.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
White, both sexes										
2006 4/.....	12.4	*	1.0	7.8	13.2	13.4	13.5	16.5
2005 4/.....	12.3	*	1.3	8.3	13.1	12.5	14.3	15.4
2004 4/.....	12.3	*	1.3	8.8	13.2	13.0	14.2	15.6
2003 4/.....	12.1	*	1.1	7.9	12.5	12.6	14.7	15.3
2002.....	12.2	*	1.3	8.2	13.2	13.5	13.7	15.9
2001 5/.....	11.9	*	1.3	8.6	12.6	13.5	14.2	15.8
2000.....	11.5	*	1.5	8.5	13.1	12.7	13.0	15.1
1999.....	11.5	*	1.3	8.5	12.7	12.8	14.2	15.7
White, male										
2006 4/.....	19.8	*	1.4	12.3	21.9	21.0	21.2	25.0
2005 4/.....	19.7	*	1.9	13.3	21.3	20.3	22.2	23.7
2004 4/.....	19.6	*	1.7	13.6	22.0	21.2	22.3	24.0
2003 4/.....	19.5	*	1.7	12.7	21.1	20.3	23.3	23.9
2002.....	19.9	*	1.9	13.4	22.0	21.4	21.7	24.7
2001 5/.....	19.5	*	2.0	14.0	21.3	22.2	22.6	24.7
2000.....	18.8	*	2.4	13.7	22.3	20.8	20.7	23.6
1999.....	18.7	*	2.0	13.7	21.5	21.0	22.5	24.3
White, female										
2006 4/.....	5.2	*	0.6	3.0	3.7	5.3	5.4	7.7
2005 4/.....	5.0	*	0.6	3.1	4.3	4.2	6.0	6.8
2004 4/.....	5.1	*	0.9	3.7	3.8	4.3	5.6	7.0
2003 4/.....	4.7	*	0.5	2.9	3.3	4.3	5.7	6.5
2002.....	4.8	*	0.7	2.6	3.7	5.1	5.3	6.8
2001 5/.....	4.6	*	0.6	2.9	3.3	4.3	5.4	6.7
2000.....	4.4	*	0.7	2.9	3.3	4.1	5.1	6.5
1999.....	4.5	*	0.5	2.9	3.5	4.3	5.5	6.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
Black, both sexes										
2006 4/.....	4.9	*	1.1	4.2	8.5	8.6	8.1	8.2
2005 4/.....	5.1	*	1.1	4.4	9.1	8.8	8.5	7.2
2004 4/.....	5.2	*	1.4	4.7	9.8	10.1	8.7	7.0
2003 4/.....	5.1	*	1.2	3.8	10.5	9.5	9.1	7.5
2002.....	5.1	*	1.0	4.0	9.1	10.3	8.2	8.1
2001 5/.....	5.3	*	1.1	4.3	10.1	9.6	8.6	7.8
2000.....	5.4	*	1.2	5.5	11.0	9.9	7.6	7.3
1999.....	5.4	*	0.9	5.8	10.7	9.6	8.7	8.4
Black, male										
2006 4/.....	8.8	*	1.7	7.0	14.5	15.7	15.2	14.9
2005 4/.....	8.7	*	1.5	7.2	16.2	14.9	15.0	12.0
2004 4/.....	9.0	*	1.6	7.4	17.2	18.0	14.4	11.7
2003 4/.....	8.8	*	1.9	6.6	18.0	17.4	15.7	12.9
2002.....	9.1	*	1.6	6.9	16.1	19.5	15.2	14.5
2001 5/.....	9.2	*	1.7	7.3	19.4	17.7	15.4	13.8
2000.....	9.4	*	2.1	9.5	19.5	17.5	13.6	13.0
1999.....	9.6	*	1.3	9.9	19.5	17.9	15.7	15.6
Black, female										
2006 4/.....	1.4	*	*	1.3	2.3	1.9	1.7	2.2
2005 4/.....	1.8	*	*	1.4	2.0	3.0	2.7	2.8
2004 4/.....	1.8	*	*	1.9	2.4	2.8	3.6	2.7
2003 4/.....	1.8	*	*	*	3.1	2.2	3.1	2.6
2002.....	1.5	*	*	*	2.3	2.0	2.0	2.5
2001 5/.....	1.7	*	*	1.3	*	2.2	2.5	2.5
2000.....	1.7	*	*	1.4	3.0	3.0	2.2	2.2
1999.....	1.6	*	*	1.6	2.3	2.1	2.5	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
All races 3/, both sexes										
2006.....	5.6	*	0.3	3.3	6.0	5.4	5.2	6.2
2005.....	5.7	*	0.4	3.5	5.8	5.6	5.7	5.9
2004.....	5.7	*	0.3	3.8	6.3	5.7	5.7	5.9
2003.....	5.8	*	0.3	3.6	6.5	5.8	6.1	6.0
2002.....	5.9	*	0.4	3.6	6.7	6.4	5.6	6.6
2001.....	5.9	*	0.4	4.1	6.6	6.7	6.3	6.4
2000.....	5.9	*	0.5	4.4	7.2	6.3	6.0	6.7
1999.....	5.9	*	0.5	4.9	7.2	6.6	6.5	7.1
All races 3/, male										
2006.....	10.0	*	0.5	5.8	10.6	9.2	9.2	10.4
2005.....	10.2	*	0.7	6.1	10.3	9.7	9.7	10.1
2004.....	10.0	*	0.5	6.5	11.1	9.9	9.6	9.8
2003.....	10.4	*	0.6	6.3	11.4	10.1	10.6	10.2
2002.....	10.6	*	0.6	6.4	11.9	11.1	9.6	11.3
2001.....	10.6	*	0.6	7.1	11.8	11.7	10.8	10.8
2000.....	10.5	*	0.9	7.6	12.9	11.1	10.1	11.1
1999.....	10.6	*	0.8	8.4	12.8	11.5	10.9	11.8
All races 3/, female										
2006.....	1.4	*	*	0.6	1.2	1.4	1.2	1.9
2005.....	1.4	*	*	0.8	1.1	1.3	1.6	1.7
2004.....	1.5	*	*	1.0	1.1	1.3	1.7	1.9
2003.....	1.4	*	*	0.8	1.2	1.3	1.5	1.8
2002.....	1.4	*	*	0.7	1.2	1.6	1.7	2.0
2001.....	1.5	*	0.2	1.0	1.0	1.5	1.7	2.1
2000.....	1.5	*	0.2	1.1	1.3	1.5	1.9	2.3
1999.....	1.5	*	0.2	1.1	1.4	1.6	1.9	2.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
White, both sexes										
2006 4/.....	6.4	*	0.4	3.5	6.3	5.7	5.7	7.0
2005 4/.....	6.5	*	0.5	3.8	6.1	6.0	6.3	6.7
2004 4/.....	6.5	*	0.3	4.2	6.6	6.0	6.3	6.6
2003 4/.....	6.6	*	0.4	4.0	6.7	6.1	6.7	6.8
2002.....	6.7	*	0.4	3.9	7.0	6.8	6.2	7.4
2001.....	6.7	*	0.4	4.5	6.7	7.2	7.0	7.2
2000.....	6.6	*	0.6	4.6	7.6	6.6	6.6	7.5
1999.....	6.6	*	0.6	5.2	7.4	7.0	7.1	7.8
White, male										
2006 4/.....	11.3	*	0.6	6.3	11.0	9.6	9.9	11.6
2005 4/.....	11.5	*	0.8	6.6	10.7	10.3	10.5	11.2
2004 4/.....	11.3	*	0.5	7.1	11.7	10.2	10.4	11.0
2003 4/.....	11.7	*	0.7	6.9	11.8	10.5	11.3	11.3
2002.....	11.9	*	0.7	6.9	12.3	11.3	10.4	12.4
2001.....	11.8	*	0.7	7.6	11.9	12.3	11.7	11.9
2000.....	11.6	*	1.0	7.9	13.5	11.4	11.0	12.2
1999.....	11.7	*	0.9	8.9	13.0	11.9	11.8	12.7
White, female										
2006 4/.....	1.6	*	*	0.6	1.2	1.6	1.4	2.3
2005 4/.....	1.6	*	*	0.9	1.2	1.4	1.9	2.0
2004 4/.....	1.7	*	*	1.1	1.2	1.5	2.0	2.1
2003 4/.....	1.6	*	*	0.9	1.2	1.5	1.8	2.1
2002.....	1.6	*	*	0.8	1.3	2.0	1.9	2.3
2001.....	1.7	*	*	1.1	1.1	1.7	2.0	2.4
2000.....	1.7	*	*	1.1	1.3	1.6	2.0	2.7
1999.....	1.7	*	0.3	1.2	1.4	1.8	2.2	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
Black, both sexes										
2006 4/.....	2.6	*	*	2.4	5.3	5.0	4.1	3.4
2005 4/.....	2.6	*	*	2.4	5.3	4.6	4.2	3.5
2004 4/.....	2.6	*	*	2.2	5.9	5.4	4.2	2.9
2003 4/.....	2.6	*	*	2.1	6.5	5.5	4.8	2.8
2002.....	2.8	*	*	2.5	5.8	6.1	4.0	4.0
2001.....	2.9	*	*	2.8	7.1	5.8	4.1	3.7
2000.....	3.0	*	*	3.8	6.8	6.0	4.0	3.4
1999.....	3.1	*	*	3.9	7.6	5.7	4.2	4.2
Black, male										
2006 4/.....	4.9	*	*	4.2	9.5	9.5	7.9	6.6
2005 4/.....	4.9	*	*	4.3	9.8	8.4	8.3	6.7
2004 4/.....	4.9	*	*	3.7	10.7	10.2	7.8	5.5
2003 4/.....	5.0	*	*	3.9	11.9	11.0	9.1	5.5
2002.....	5.4	*	*	4.3	11.0	12.4	7.7	7.4
2001.....	5.5	*	*	5.0	13.9	10.9	8.0	7.1
2000.....	5.5	*	*	6.9	12.6	11.3	7.3	6.4
1999.....	5.9	*	*	7.0	14.2	11.2	8.1	8.1
Black, female										
2006 4/.....	0.4	*	*	*	*	*	*	*
2005 4/.....	0.5	*	*	*	*	*	*	*
2004 4/.....	0.5	*	*	*	*	*	*	*
2003 4/.....	0.5	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All races 3/, both sexes

2006.....	5.5	*	0.7	4.0	6.4	6.9	7.1	8.5
2005.....	5.3	*	0.9	4.2	6.5	6.2	7.3	7.8
2004.....	5.3	*	1.1	4.4	6.2	6.6	7.3	8.1
2003.....	5.0	*	0.8	3.7	5.6	6.1	7.4	7.8
2002.....	5.0	*	0.8	3.8	5.7	6.4	6.9	7.7
2001 5/.....	4.8	*	0.9	3.8	5.4	5.9	6.7	7.9
2000.....	4.5	*	0.9	3.6	5.3	5.8	6.0	7.0
1999.....	4.5	*	0.7	3.2	5.1	5.7	6.7	7.3

All races 3/, male

2006.....	7.8	*	0.9	5.7	10.2	10.5	10.5	12.3
2005.....	7.5	*	1.2	6.0	10.0	9.5	10.9	11.3
2004.....	7.6	*	1.2	6.2	9.7	10.3	11.0	11.9
2003.....	7.3	*	1.2	5.4	8.8	9.3	11.0	11.7
2002.....	7.3	*	1.2	5.8	8.9	9.7	10.6	11.4
2001 5/.....	7.1	*	1.3	5.8	8.7	9.3	10.2	11.9
2000.....	6.6	*	1.4	5.4	8.5	8.9	9.1	10.6
1999.....	6.6	*	1.1	4.6	8.0	8.8	10.2	10.9

All races 3/, female

2006.....	3.2	*	0.5	2.2	2.4	3.3	3.5	4.7
2005.....	3.1	*	0.6	2.2	2.9	2.8	3.6	4.4
2004.....	3.1	*	0.9	2.5	2.4	2.8	3.5	4.4
2003.....	2.8	*	0.4	1.9	2.2	2.7	3.7	4.0
2002.....	2.9	*	0.4	1.6	2.3	2.9	3.1	4.0
2001 5/.....	2.6	*	0.4	1.7	2.0	2.5	3.1	3.9
2000.....	2.5	*	0.4	1.7	1.9	2.6	2.7	3.4
1999.....	2.5	*	0.3	1.6	2.0	2.5	3.2	3.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
White, both sexes										
2006 4/.....	6.0	*	0.7	4.3	6.8	7.7	7.8	9.4
2005 4/.....	5.8	*	0.8	4.5	7.0	6.5	8.0	8.8
2004 4/.....	5.8	*	1.0	4.6	6.6	7.0	7.9	9.0
2003 4/.....	5.5	*	0.8	4.0	5.8	6.4	8.0	8.5
2002.....	5.5	*	0.8	4.2	6.2	6.8	7.4	8.5
2001 5/.....	5.3	*	0.9	4.2	5.9	6.3	7.2	8.6
2000.....	4.9	*	0.9	3.8	5.5	6.1	6.4	7.6
1999.....	4.9	*	0.7	3.3	5.4	5.9	7.1	7.9
White, male										
2006 4/.....	8.5	*	0.8	6.1	10.9	11.4	11.3	13.4
2005 4/.....	8.2	*	1.1	6.7	10.6	9.9	11.8	12.5
2004 4/.....	8.3	*	1.2	6.5	10.4	10.9	11.9	13.0
2003 4/.....	7.9	*	1.1	5.8	9.3	9.8	11.9	12.6
2002.....	8.0	*	1.2	6.5	9.7	10.2	11.2	12.3
2001 5/.....	7.7	*	1.3	6.4	9.4	9.8	10.8	12.8
2000.....	7.1	*	1.3	5.8	8.8	9.4	9.7	11.4
1999.....	7.0	*	1.1	4.9	8.5	9.1	10.7	11.6
White, female										
2006 4/.....	3.6	*	0.5	2.4	2.5	3.7	4.0	5.3
2005 4/.....	3.4	*	0.5	2.2	3.2	2.8	4.0	4.8
2004 4/.....	3.4	*	0.8	2.7	2.6	2.8	3.6	4.9
2003 4/.....	3.1	*	0.4	2.0	2.2	2.8	4.0	4.3
2002.....	3.2	*	0.5	1.8	2.5	3.1	3.4	4.5
2001 5/.....	2.9	*	0.4	1.8	2.2	2.6	3.4	4.3
2000.....	2.8	*	0.5	1.8	1.9	2.5	3.1	3.8
1999.....	2.8	*	0.3	1.7	2.1	2.5	3.3	4.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)										
Black, both sexes										
2006 4/.....	2.4	*	1.0	1.8	3.2	3.6	4.0	4.8
2005 4/.....	2.5	*	1.0	2.0	3.9	4.1	4.3	3.7
2004 4/.....	2.6	*	1.1	2.5	3.9	4.8	4.5	4.0
2003 4/.....	2.5	*	1.0	1.7	4.0	4.0	4.3	4.6
2002.....	2.3	*	0.7	1.6	3.2	4.2	4.2	4.1
2001 5/.....	2.3	*	0.8	1.5	3.0	3.8	4.5	4.1
2000.....	2.4	*	1.1	1.8	4.2	3.9	3.6	3.8
1999.....	2.3	*	0.7	1.9	3.1	3.9	4.6	4.2
Black, male										
2006 4/.....	4.0	*	1.6	2.8	5.0	6.1	7.4	8.3
2005 4/.....	3.8	*	1.2	2.9	6.4	6.5	6.7	5.3
2004 4/.....	4.1	*	1.2	3.7	6.5	7.8	6.5	6.2
2003 4/.....	3.8	*	1.6	2.6	6.1	6.4	6.7	7.4
2002.....	3.7	*	1.1	2.6	5.0	7.0	7.5	7.0
2001 5/.....	3.7	*	1.1	2.3	5.5	6.8	7.4	6.7
2000.....	3.9	*	1.8	2.6	6.8	6.2	6.4	6.6
1999.....	3.7	*	*	2.9	5.3	6.7	7.6	7.5
Black, female										
2006 4/.....	0.9	*	*	*	1.4	*	*	1.6
2005 4/.....	1.3	*	*	*	1.3	1.9	2.1	2.3
2004 4/.....	1.3	*	*	*	1.3	2.0	2.6	2.0
2003 4/.....	1.3	*	*	*	1.9	1.8	2.2	2.2
2002.....	1.1	*	*	*	1.5	1.6	1.4	1.6
2001 5/.....	1.1	*	*	*	*	*	1.8	1.8
2000.....	1.0	*	*	*	1.7	1.8	*	1.4
1999.....	1.0	*	*	*	*	*	1.9	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
All races 3/, both sexes										
2006.....	6.2	8.1	2.2	0.8	1.2	10.7	16.2	13.5	9.8	7.6
2005.....	6.1	7.5	2.3	0.6	1.1	9.9	16.1	13.6	10.0	7.5
2004.....	5.9	8.0	2.4	0.6	1.0	9.3	15.0	13.3	9.3	7.3
2003.....	6.1	8.5	2.4	0.6	1.0	9.5	16.5	13.4	9.4	7.6
2002.....	6.1	7.5	2.7	0.7	1.0	9.3	16.5	13.3	9.4	7.9
2001 5/.....	7.1	8.2	2.7	0.7	0.9	9.4	17.3	14.9	11.5	10.4
2000.....	6.0	9.2	2.3	0.7	1.1	9.5	16.0	11.9	9.0	7.6
1999.....	6.1	8.7	2.5	0.9	1.2	10.4	15.6	12.1	9.1	7.7
All races 3/, male										
2006.....	10.0	9.4	2.5	0.8	1.6	18.2	27.5	22.4	16.2	11.4
2005.....	9.8	8.2	2.6	0.7	1.3	16.8	27.2	22.8	16.4	11.2
2004.....	9.4	8.0	2.5	0.6	1.3	15.3	25.4	22.3	14.9	11.2
2003.....	9.7	10.0	2.5	0.7	1.3	15.9	27.6	22.4	14.8	11.4
2002.....	9.6	7.9	2.9	0.6	1.2	15.3	27.7	22.0	14.7	11.8
2001 5/.....	11.1	9.5	3.0	0.7	1.1	15.7	28.9	24.1	17.7	15.3
2000.....	9.3	10.4	2.5	0.7	1.4	15.5	26.7	19.6	14.0	10.9
1999.....	9.3	9.6	2.5	0.9	1.5	16.9	25.7	19.5	13.8	11.0
All races 3/, female										
2006.....	2.5	6.8	2.0	0.7	0.7	2.9	4.2	4.2	3.3	3.6
2005.....	2.5	6.6	2.0	0.6	0.8	2.5	4.3	4.0	3.5	3.9
2004.....	2.5	7.9	2.2	0.6	0.7	3.0	4.0	3.9	3.6	3.3
2003.....	2.6	6.9	2.3	0.5	0.6	2.6	4.8	4.0	3.9	3.9
2002.....	2.7	7.1	2.5	0.8	0.8	2.9	4.6	4.2	4.1	4.1
2001 5/.....	3.3	6.9	2.4	0.6	0.7	2.7	5.1	5.5	5.2	5.5
2000.....	2.8	7.9	2.1	0.6	0.8	3.1	4.7	4.1	4.0	4.4
1999.....	2.9	7.8	2.4	0.9	0.9	3.5	5.1	4.5	4.3	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

White, both sexes

2006 4/.....	3.7	6.9	1.5	0.6	0.8	5.4	7.9	6.8	5.2	4.7
2005 4/.....	3.7	6.1	1.7	0.4	0.6	5.3	7.9	7.2	5.6	4.7
2004 4/.....	3.6	6.5	1.6	0.5	0.7	5.1	7.9	7.0	5.2	4.6
2003 4/.....	3.7	6.9	1.7	0.4	0.6	5.0	8.4	6.9	5.3	5.0
2002.....	3.7	5.4	2.0	0.5	0.7	5.2	8.2	6.8	5.1	5.0
2001 5/.....	4.9	6.2	1.9	0.7	0.7	5.1	9.4	8.9	7.8	7.8
2000.....	3.6	6.6	1.7	0.6	0.9	5.2	7.7	6.2	5.3	4.9
1999.....	3.8	6.7	1.7	0.7	1.0	5.6	8.0	6.5	5.5	5.1

White, male

2006 4/.....	5.4	7.4	1.7	0.5	1.1	8.7	12.6	10.6	7.9	6.7
2005 4/.....	5.4	6.7	1.8	0.5	0.7	8.6	12.5	11.1	8.4	6.5
2004 4/.....	5.3	6.9	1.8	0.5	0.9	7.9	12.5	10.9	7.7	6.8
2003 4/.....	5.4	8.1	1.5	0.4	0.9	8.0	13.3	10.7	7.8	6.9
2002.....	5.4	6.2	2.1	0.4	0.7	8.2	12.9	10.5	7.3	7.0
2001 5/.....	7.2	7.3	2.2	0.6	0.8	7.9	14.5	13.5	11.3	11.6
2000.....	5.2	8.2	1.9	0.7	1.1	8.0	12.0	9.2	7.6	6.6
1999.....	5.5	7.9	1.8	0.7	1.1	8.6	12.1	9.9	7.8	7.0

White, female

2006 4/.....	1.9	6.3	1.3	0.7	0.6	1.9	2.9	2.7	2.3	2.7
2005 4/.....	1.9	5.5	1.6	0.4	0.6	1.8	2.9	3.0	2.7	2.9
2004 4/.....	1.9	6.2	1.4	0.5	0.5	2.1	3.0	2.9	2.6	2.4
2003 4/.....	2.0	5.7	1.8	0.4	0.3	1.8	3.2	2.9	2.6	3.0
2002.....	2.0	4.6	1.8	0.6	0.6	1.9	3.1	2.8	2.8	3.1
2001 5/.....	2.6	5.1	1.6	0.7	0.6	2.0	3.9	4.1	4.2	3.9
2000.....	2.1	5.0	1.5	0.5	0.7	2.1	3.2	3.0	2.9	3.2
1999.....	2.1	5.4	1.7	0.7	0.8	2.3	3.7	2.9	3.2	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
Black, both sexes										
2006 4/.....	22.8	16.0	5.8	1.6	2.9	37.5	62.4	54.0	39.7	26.9
2005 4/.....	22.2	14.3	5.3	1.4	3.0	33.5	61.2	53.0	39.2	26.5
2004 4/.....	21.1	15.6	6.3	1.4	2.4	31.6	55.9	52.8	35.6	25.2
2003 4/.....	22.0	16.1	6.1	1.7	2.7	33.2	62.4	54.5	36.3	25.4
2002.....	22.0	17.3	6.3	1.4	2.6	31.0	63.4	54.2	37.0	26.9
2001 5/.....	22.1	18.9	6.8	0.9	1.9	32.7	62.8	52.6	35.3	28.2
2000.....	21.5	22.8	5.8	1.2	2.5	33.1	64.1	48.2	33.5	25.9
1999.....	21.1	19.5	6.1	1.9	2.5	36.5	60.4	47.0	31.8	25.1
Black, male										
2006 4/.....	40.6	20.7	6.3	2.1	4.1	66.3	112.2	97.3	73.6	46.3
2005 4/.....	39.7	15.9	6.6	1.6	4.1	60.0	110.1	98.6	72.8	44.7
2004 4/.....	37.1	14.2	6.3	1.4	3.3	54.8	101.8	98.4	64.7	43.1
2003 4/.....	38.9	17.8	7.4	2.0	3.7	58.9	111.6	101.6	63.9	43.7
2002.....	38.4	16.3	7.1	1.4	3.5	53.3	115.3	99.9	65.4	45.2
2001 5/.....	38.3	21.0	8.0	1.2	2.7	58.2	116.3	96.0	62.3	42.8
2000.....	37.2	23.3	5.9	*	3.3	57.2	117.8	89.4	58.6	42.5
1999.....	36.1	19.5	5.9	2.0	3.3	62.4	110.9	84.0	54.7	40.1
Black, female										
2006 4/.....	6.6	11.1	5.4	*	1.7	7.9	11.3	12.9	9.0	9.6
2005 4/.....	6.2	12.6	4.0	1.3	1.9	6.3	11.5	9.9	8.8	10.1
2004 4/.....	6.4	17.0	6.3	1.4	1.5	7.8	9.8	10.2	9.3	9.2
2003 4/.....	6.6	14.2	4.8	1.3	1.7	6.7	13.6	10.9	11.4	9.0
2002.....	7.0	18.5	5.5	1.3	1.6	8.0	12.7	12.3	11.5	10.6
2001 5/.....	7.4	16.7	5.6	*	*	6.4	11.4	13.1	11.1	15.3
2000.....	7.2	22.2	5.7	1.4	1.7	8.4	13.2	10.9	10.9	11.1
1999.....	7.6	19.5	6.3	1.8	1.6	10.0	12.5	13.6	11.2	11.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
All races 3/, both sexes										
2006.....	4.3	*	0.3	0.3	0.8	9.1	13.7	10.8	7.7	5.2
2005.....	4.2	*	0.2	0.2	0.7	8.3	13.1	11.0	7.8	5.1
2004.....	4.0	*	0.2	0.2	0.7	7.6	12.2	10.5	7.0	4.9
2003.....	4.1	*	0.3	0.2	0.7	7.7	13.6	10.8	7.1	5.0
2002.....	4.1	*	0.3	0.3	0.7	7.7	13.6	10.6	6.9	5.2
2001.....	4.0	*	0.4	0.3	0.6	7.5	13.6	10.3	6.6	5.1
2000.....	3.8	*	0.2	0.2	0.7	7.7	12.7	9.2	6.3	4.7
1999.....	3.9	*	0.3	0.3	0.8	8.5	12.5	9.2	6.1	4.6
All races 3/, male										
2006.....	7.4	*	0.3	0.3	1.3	16.0	24.1	18.7	13.4	8.4
2005.....	7.2	*	0.3	0.3	1.0	14.7	23.1	19.3	13.5	8.1
2004.....	6.9	*	*	0.2	1.0	13.1	21.5	18.8	11.9	8.3
2003.....	7.1	*	0.2	0.3	1.0	13.7	23.8	19.1	12.0	8.0
2002.....	7.0	*	0.4	0.3	0.9	13.2	23.8	18.6	11.6	8.4
2001.....	6.8	*	0.3	0.3	0.8	13.2	24.0	17.8	11.1	8.0
2000.....	6.5	*	*	0.3	1.0	13.3	22.4	16.1	10.4	7.4
1999.....	6.5	*	0.3	0.4	1.1	14.5	21.8	15.9	10.1	7.1
All races 3/, female										
2006.....	1.3	*	*	0.3	0.4	1.8	2.6	2.5	1.9	2.0
2005.....	1.2	*	*	*	0.4	1.5	2.4	2.3	2.1	2.1
2004.....	1.1	*	*	0.2	0.3	1.8	2.2	2.0	2.0	1.4
2003.....	1.2	*	0.3	0.2	0.3	1.5	2.8	2.2	2.1	1.9
2002.....	1.3	*	*	0.3	0.5	1.8	2.9	2.4	2.1	2.0
2001.....	1.3	*	0.4	0.2	0.3	1.5	2.7	2.5	2.1	2.2
2000.....	1.3	*	*	0.2	0.3	1.7	2.7	2.2	2.1	1.9
1999.....	1.3	*	0.3	0.2	0.5	2.2	2.8	2.3	2.1	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
White, both sexes										
2006 4/.....	2.2	*	0.2	0.3	0.6	4.2	5.9	4.8	3.6	3.0
2005 4/.....	2.2	*	0.2	0.2	0.4	4.1	5.5	5.1	4.0	3.0
2004 4/.....	2.1	*	*	0.2	0.4	3.8	5.5	4.9	3.7	2.9
2003 4/.....	2.2	*	0.2	0.1	0.4	3.8	6.3	4.9	3.6	3.0
2002.....	2.2	*	0.3	0.2	0.4	4.1	6.0	4.9	3.4	3.0
2001.....	2.2	*	0.3	0.3	0.4	3.7	6.3	5.0	3.5	3.2
2000.....	2.1	*	*	0.2	0.5	3.9	5.5	4.2	3.3	2.8
1999.....	2.2	*	0.2	0.2	0.6	4.2	5.6	4.3	3.4	3.0
White, male										
2006 4/.....	3.5	*	*	*	0.8	7.2	9.9	7.9	5.9	4.4
2005 4/.....	3.5	*	*	*	0.5	7.0	9.3	8.4	6.3	4.2
2004 4/.....	3.5	*	*	*	0.6	6.3	9.3	8.3	5.8	4.7
2003 4/.....	3.5	*	*	*	0.6	6.4	10.6	8.2	5.7	4.4
2002.....	3.5	*	0.3	*	0.5	6.7	10.0	8.1	5.3	4.5
2001.....	3.5	*	*	0.3	0.6	6.2	10.5	8.2	5.4	4.9
2000.....	3.2	*	*	0.3	0.7	6.5	9.0	6.7	5.1	4.1
1999.....	3.4	*	*	0.2	0.8	6.9	9.1	7.1	5.2	4.3
White, female										
2006 4/.....	0.9	*	*	0.3	0.3	1.1	1.5	1.5	1.3	1.5
2005 4/.....	0.9	*	*	*	*	1.0	1.4	1.6	1.5	1.7
2004 4/.....	0.9	*	*	*	*	1.2	1.5	1.2	1.4	1.0
2003 4/.....	0.9	*	*	*	*	1.0	1.7	1.4	1.4	1.6
2002.....	1.0	*	*	*	0.3	1.2	1.6	1.6	1.4	1.5
2001.....	1.0	*	*	*	0.3	1.1	1.9	1.6	1.6	1.6
2000.....	1.0	*	*	*	*	1.1	1.7	1.6	1.5	1.5
1999.....	0.9	*	*	*	0.4	1.2	1.9	1.4	1.5	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/ ages 1/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
Black, both sexes										
2006 4/.....	18.0	*	0.8	*	2.4	33.5	57.1	47.4	34.4	20.6
2005 4/.....	17.2	*	*	*	2.1	30.0	55.1	46.9	33.5	19.4
2004 4/.....	16.1	*	0.8	*	1.8	27.8	50.4	46.3	29.3	18.5
2003 4/.....	16.8	*	*	0.7	2.0	29.0	55.3	47.9	30.1	18.1
2002.....	16.7	*	*	*	1.9	27.3	57.0	46.9	30.1	19.5
2001.....	15.8	*	*	*	1.3	28.7	55.7	43.6	26.9	18.4
2000.....	15.6	*	*	*	1.7	28.6	55.3	40.7	26.0	17.3
1999.....	15.4	*	*	0.7	1.8	32.0	53.4	39.3	23.7	15.9
Black, male										
2006 4/.....	33.7	*	*	*	3.7	60.8	104.4	87.9	65.8	37.7
2005 4/.....	32.5	*	*	*	3.1	54.9	101.6	89.5	64.2	35.7
2004 4/.....	30.3	*	*	*	2.9	49.8	93.9	88.9	55.2	34.6
2003 4/.....	31.6	*	*	*	2.9	53.2	101.7	92.4	56.4	33.9
2002.....	31.0	*	*	*	2.7	48.4	105.9	90.2	56.3	35.7
2001.....	29.8	*	*	*	2.1	52.8	106.5	83.2	51.3	31.5
2000.....	29.2	*	*	*	2.8	51.7	105.2	78.7	48.1	31.7
1999.....	28.4	*	*	*	2.8	56.3	101.0	74.4	44.0	27.3
Black, female										
2006 4/.....	3.6	*	*	*	*	5.5	8.6	9.0	5.9	5.4
2005 4/.....	3.1	*	*	*	*	4.4	7.6	6.6	5.8	4.7
2004 4/.....	3.0	*	*	*	*	5.1	6.6	6.4	5.9	4.0
2003 4/.....	3.3	*	*	*	1.1	4.0	9.1	6.8	6.3	4.1
2002.....	3.6	*	*	*	1.2	5.4	9.2	7.3	6.5	5.0
2001.....	3.1	*	*	*	*	3.9	7.0	7.6	5.0	6.8
2000.....	3.2	*	*	*	*	4.9	8.0	6.2	6.0	4.5
1999.....	3.7	*	*	*	*	7.0	8.2	7.5	5.5	5.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes

2006.....	1.9	8.0	2.0	0.4	0.3	1.6	2.6	2.7	2.0	2.3
2005.....	1.9	7.3	2.1	0.4	0.4	1.6	3.0	2.6	2.2	2.5
2004.....	2.0	7.8	2.1	0.4	0.3	1.7	2.9	2.7	2.2	2.4
2003.....	2.0	8.3	2.1	0.4	0.3	1.7	2.9	2.6	2.3	2.7
2002.....	2.0	7.3	2.4	0.4	0.3	1.6	2.9	2.6	2.5	2.7
2001 5/.....	3.1	8.0	2.3	0.4	0.3	1.8	3.7	4.6	4.9	5.2
2000.....	2.1	8.9	2.1	0.4	0.5	1.8	3.2	2.7	2.8	2.9
1999.....	2.2	8.5	2.1	0.6	0.4	1.9	3.1	2.9	3.0	3.0

All races 3/, male

2006.....	2.6	9.3	2.2	0.5	0.3	2.2	3.5	3.7	2.7	3.1
2005.....	2.6	7.9	2.3	0.4	0.3	2.1	4.0	3.5	3.0	3.1
2004.....	2.5	7.9	2.3	0.4	0.3	2.2	4.0	3.5	2.9	2.9
2003.....	2.6	9.7	2.3	0.4	0.4	2.3	3.8	3.3	2.8	3.4
2002.....	2.6	7.8	2.6	0.4	0.3	2.1	3.9	3.4	3.0	3.4
2001 5/.....	4.3	9.2	2.7	0.4	0.3	2.4	4.9	6.3	6.7	7.3
2000.....	2.8	10.0	2.3	0.5	0.4	2.2	4.4	3.5	3.6	3.5
1999.....	2.8	9.4	2.1	0.5	0.4	2.4	3.9	3.6	3.7	3.9

All races 3/, female

2006.....	1.3	6.6	1.8	0.4	0.3	1.1	1.6	1.7	1.3	1.6
2005.....	1.3	6.6	1.8	0.4	0.4	1.0	1.9	1.7	1.5	1.8
2004.....	1.4	7.7	2.0	0.4	0.4	1.2	1.7	1.9	1.5	1.9
2003.....	1.4	6.9	2.0	0.3	0.2	1.1	2.0	1.8	1.8	1.9
2002.....	1.4	6.8	2.2	0.5	0.3	1.1	1.8	1.8	2.0	2.1
2001 5/.....	2.0	6.7	2.0	0.4	0.4	1.2	2.4	3.0	3.2	3.2
2000.....	1.5	7.7	2.0	0.4	0.5	1.4	2.0	1.9	2.0	2.4
1999.....	1.6	7.6	2.1	0.7	0.5	1.4	2.2	2.2	2.3	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

White, both sexes

2006 4/.....	1.5	6.8	1.3	0.4	0.3	1.2	2.1	2.0	1.5	1.7
2005 4/.....	1.5	6.0	1.5	0.3	0.2	1.2	2.4	2.1	1.7	1.7
2004 4/.....	1.5	6.5	1.5	0.3	0.3	1.3	2.4	2.1	1.5	1.7
2003 4/.....	1.5	6.9	1.5	0.3	0.2	1.2	2.1	2.0	1.7	2.0
2002.....	1.5	5.3	1.7	0.3	0.2	1.1	2.2	1.9	1.7	2.0
2001 5/.....	2.6	6.0	1.6	0.4	0.3	1.4	3.0	3.9	4.3	4.6
2000.....	1.5	6.5	1.5	0.4	0.4	1.3	2.3	2.0	2.0	2.1
1999.....	1.6	6.6	1.5	0.5	0.4	1.4	2.4	2.1	2.2	2.1

White, male

2006 4/.....	1.9	7.4	1.5	0.3	0.3	1.5	2.7	2.7	2.0	2.3
2005 4/.....	1.9	6.4	1.6	0.3	*	1.6	3.2	2.7	2.2	2.3
2004 4/.....	1.9	6.8	1.7	0.3	0.2	1.6	3.2	2.6	1.9	2.1
2003 4/.....	1.9	8.0	1.4	0.3	0.3	1.6	2.6	2.4	2.2	2.5
2002.....	1.9	6.2	1.8	*	*	1.5	2.9	2.5	2.0	2.5
2001 5/.....	3.7	7.0	1.9	0.3	*	1.8	3.9	5.3	5.9	6.8
2000.....	2.0	8.1	1.7	0.5	0.4	1.6	3.0	2.5	2.5	2.5
1999.....	2.1	7.7	1.5	0.4	0.3	1.7	3.0	2.7	2.6	2.7

White, female

2006 4/.....	1.0	6.2	1.2	0.4	0.3	0.8	1.4	1.2	1.0	1.1
2005 4/.....	1.0	5.5	1.4	*	0.4	0.8	1.5	1.3	1.2	1.2
2004 4/.....	1.1	6.1	1.3	0.3	0.3	0.9	1.5	1.7	1.2	1.3
2003 4/.....	1.1	5.6	1.6	*	*	0.8	1.5	1.5	1.2	1.4
2002.....	1.1	4.5	1.6	0.4	0.3	0.7	1.5	1.2	1.4	1.5
2001 5/.....	1.6	4.9	1.4	0.4	0.3	0.9	2.0	2.5	2.6	2.3
2000.....	1.1	4.8	1.3	0.3	0.5	1.0	1.5	1.4	1.4	1.7
1999.....	1.2	5.4	1.5	0.5	0.4	1.1	1.8	1.5	1.7	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All Under 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39
ages 1/ year 2/ years years years years years years years years

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Black, both sexes

2006 4/.....	4.9	15.4	5.1	1.0	*	4.0	5.3	6.6	5.3	6.3
2005 4/.....	5.0	14.0	4.8	0.9	0.9	3.5	6.2	6.1	5.7	7.1
2004 4/.....	5.0	15.1	5.4	0.9	0.6	3.9	5.6	6.6	6.3	6.7
2003 4/.....	5.2	15.2	5.3	1.0	0.6	4.2	7.2	6.6	6.2	7.2
2002.....	5.3	16.5	5.8	1.0	0.6	3.7	6.4	7.3	7.0	7.4
2001 5/.....	6.3	18.6	6.1	*	0.6	4.0	7.1	9.0	8.4	9.8
2000.....	5.9	21.6	5.5	0.8	0.8	4.5	8.8	7.6	7.6	8.6
1999.....	5.7	19.2	5.4	1.3	0.6	4.5	7.0	7.7	8.1	9.2

Black, male

2006 4/.....	6.9	20.2	5.4	1.4	*	5.5	7.7	9.4	7.8	8.6
2005 4/.....	7.2	15.4	5.9	*	*	5.1	8.5	9.1	8.6	9.0
2004 4/.....	6.8	13.9	5.5	*	*	5.0	7.9	9.4	9.5	8.5
2003 4/.....	7.4	16.1	6.5	*	*	5.7	9.9	9.2	7.6	9.8
2002.....	7.3	15.7	6.4	*	*	4.8	9.4	9.7	9.1	9.5
2001 5/.....	8.5	20.7	7.3	*	*	5.4	9.8	12.8	11.0	11.3
2000.....	8.0	21.1	5.5	*	*	5.5	12.6	10.7	10.5	10.9
1999.....	7.7	19.2	5.4	1.2	*	6.1	9.8	9.5	10.7	12.8

Black, female

2006 4/.....	3.0	10.5	4.7	*	*	2.4	2.7	3.9	3.1	4.2
2005 4/.....	3.0	12.6	3.7	*	*	1.9	3.9	3.3	3.0	5.4
2004 4/.....	3.4	16.4	5.4	*	*	2.8	3.3	3.9	3.4	5.2
2003 4/.....	3.3	14.2	4.1	*	*	2.7	4.5	4.1	5.0	5.0
2002.....	3.4	17.3	5.1	*	*	2.6	3.5	5.0	5.0	5.6
2001 5/.....	4.3	16.4	4.8	*	*	2.5	4.5	5.5	6.1	8.4
2000.....	4.0	22.2	5.5	*	*	3.4	5.2	4.7	4.9	6.7
1999.....	3.9	19.1	5.4	1.3	*	2.9	4.3	6.1	5.7	6.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All Under 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39
ages 1/ year 2/ years years years years years years years years

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2006.....	1.7	2.2	0.3	0.1	0.2	0.8	2.0	2.0	2.1	2.6
2005.....	1.6	2.5	0.4	0.1	0.2	0.7	1.6	1.8	1.9	2.3
2004.....	1.7	2.6	0.3	0.1	0.2	0.7	1.7	1.9	2.1	2.9
2003.....	1.7	1.9	0.2	*	0.2	0.8	1.9	1.8	2.3	3.2
2002.....	1.7	2.5	0.4	0.1	0.2	0.7	1.7	1.9	2.3	3.3
2001.....	1.5	2.1	0.3	*	0.2	0.7	1.4	1.6	2.0	3.0
2000.....	1.4	1.7	0.3	*	0.2	0.6	1.3	1.6	1.8	2.6
1999.....	1.4	1.4	0.3	*	0.2	0.8	1.5	1.6	2.1	2.8

All races 3/, male

2006.....	2.2	2.7	0.3	*	0.3	1.2	2.9	2.8	2.7	3.2
2005.....	2.0	3.2	0.5	*	0.3	1.0	2.3	2.4	2.3	2.7
2004.....	2.2	3.1	0.3	*	0.2	1.1	2.4	2.6	2.9	3.8
2003.....	2.3	2.4	*	*	0.3	1.2	2.7	2.5	3.2	4.2
2002.....	2.2	2.5	0.4	*	0.2	1.0	2.4	2.7	3.1	4.5
2001.....	2.0	2.4	*	*	0.3	1.0	2.1	2.4	2.7	3.9
2000.....	1.9	1.4	0.3	*	0.4	0.9	2.0	2.3	2.6	3.7
1999.....	2.0	1.7	0.3	*	0.3	1.3	2.2	2.3	2.8	3.9

All races 3/, female

2006.....	1.2	1.7	0.3	*	*	0.5	1.0	1.1	1.4	2.0
2005.....	1.3	1.8	0.3	*	*	0.4	0.9	1.2	1.5	1.9
2004.....	1.2	2.0	0.3	*	*	0.4	0.8	1.1	1.4	1.9
2003.....	1.2	1.3	*	*	*	0.5	1.1	1.1	1.5	2.2
2002.....	1.2	2.5	0.4	*	*	0.4	1.0	1.0	1.4	2.1
2001.....	0.9	1.7	0.3	*	*	0.3	0.7	0.9	1.3	2.1
2000.....	0.8	2.0	*	*	*	0.3	0.6	0.9	1.1	1.6
1999.....	0.9	1.2	0.3	*	*	0.3	0.6	0.9	1.3	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

White, both sexes

2006 4/.....	1.7	2.0	0.2	*	0.2	1.0	2.2	2.2	2.2	2.7
2005 4/.....	1.6	2.1	0.3	*	0.2	0.7	1.7	2.0	2.0	2.4
2004 4/.....	1.7	2.1	0.2	*	0.2	0.8	1.9	2.1	2.2	3.0
2003 4/.....	1.8	1.7	0.2	*	0.2	0.9	2.1	2.0	2.6	3.4
2002.....	1.7	1.8	0.4	*	0.2	0.7	1.9	1.9	2.3	3.4
2001.....	1.5	1.4	0.3	*	0.2	0.7	1.5	1.7	2.1	3.0
2000.....	1.3	1.4	0.2	*	0.2	0.6	1.3	1.6	1.8	2.7
1999.....	1.4	1.2	0.2	*	0.2	0.8	1.5	1.6	2.1	2.8

White, male

2006 4/.....	2.2	2.5	*	*	0.3	1.4	3.2	3.0	2.9	3.2
2005 4/.....	2.0	2.8	0.4	*	0.3	1.0	2.5	2.5	2.4	2.8
2004 4/.....	2.2	2.7	*	*	*	1.1	2.7	2.9	2.9	3.9
2003 4/.....	2.3	1.9	*	*	0.3	1.3	2.9	2.7	3.3	4.4
2002.....	2.2	1.8	0.4	*	0.2	1.1	2.6	2.7	3.1	4.5
2001.....	2.0	1.7	*	*	0.3	1.1	2.2	2.5	2.9	3.8
2000.....	1.8	*	*	*	0.3	1.0	1.9	2.3	2.5	3.7
1999.....	1.9	1.4	*	*	0.3	1.3	2.3	2.3	2.8	3.7

White, female

2006 4/.....	1.3	1.4	*	*	*	0.5	1.1	1.3	1.5	2.2
2005 4/.....	1.3	1.5	*	*	*	0.4	0.9	1.4	1.6	2.0
2004 4/.....	1.3	1.5	*	*	*	0.4	0.9	1.2	1.5	2.1
2003 4/.....	1.3	1.4	*	*	*	0.5	1.2	1.2	1.7	2.3
2002.....	1.2	1.8	0.4	*	*	0.4	1.1	1.1	1.5	2.3
2001.....	1.0	*	*	*	*	0.3	0.8	0.9	1.4	2.3
2000.....	0.8	1.6	*	*	*	0.3	0.7	0.9	1.1	1.6
1999.....	0.9	*	*	*	*	*	0.6	1.0	1.4	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Black, both sexes

2006 4/.....	1.8	3.4	0.8	*	*	*	1.3	1.5	1.6	2.8
2005 4/.....	1.8	4.7	0.9	*	*	0.6	1.1	1.4	1.9	2.5
2004 4/.....	1.8	4.8	*	*	*	0.7	1.1	1.0	2.6	3.1
2003 4/.....	1.8	3.2	*	*	*	*	1.1	1.4	2.1	3.2
2002.....	2.0	5.5	*	*	*	*	1.3	2.0	2.5	4.1
2001.....	1.7	5.5	*	*	*	*	1.1	1.5	1.9	3.8
2000.....	1.8	*	*	*	*	*	1.4	1.7	2.2	3.3
1999.....	1.9	*	*	*	*	0.8	1.7	1.9	2.5	3.9

Black, male

2006 4/.....	2.7	*	*	*	*	*	2.0	2.6	2.3	4.0
2005 4/.....	2.4	*	*	*	*	*	1.6	2.3	2.3	3.1
2004 4/.....	2.6	*	*	*	*	*	1.5	1.5	4.0	4.6
2003 4/.....	2.7	*	*	*	*	*	1.9	2.5	3.4	4.4
2002.....	2.9	*	*	*	*	*	1.9	3.2	3.7	6.5
2001.....	2.7	6.0	*	*	*	*	1.9	2.5	2.7	5.9
2000.....	2.8	*	*	*	*	*	2.5	2.7	3.4	4.8
1999.....	3.0	*	*	*	*	1.3	2.7	3.4	3.7	6.3

Black, female

2006 4/.....	1.0	*	*	*	*	*	*	*	*	1.8
2005 4/.....	1.1	*	*	*	*	*	*	*	1.5	2.0
2004 4/.....	1.1	*	*	*	*	*	*	*	1.4	1.8
2003 4/.....	0.9	*	*	*	*	*	*	*	*	2.2
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.9
2001.....	0.9	*	*	*	*	*	*	*	*	2.0
2000.....	0.9	*	*	*	*	*	*	*	*	2.0
1999.....	0.9	*	*	*	*	*	*	*	1.4	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2006.....	1.6	2.2	0.3	0.1	0.2	0.7	1.8	1.8	2.0	2.5
2005.....	1.5	2.5	0.4	*	0.2	0.6	1.3	1.7	1.8	2.3
2004.....	1.6	2.6	0.3	0.1	0.2	0.6	1.5	1.8	2.0	2.8
2003.....	1.7	1.9	0.2	*	0.1	0.7	1.7	1.7	2.3	3.1
2002.....	1.6	2.5	0.4	0.1	0.2	0.5	1.5	1.7	2.2	3.2
2001.....	1.4	2.1	0.3	*	0.1	0.5	1.2	1.5	1.9	2.9
2000.....	1.3	1.7	0.3	*	0.2	0.5	1.1	1.4	1.7	2.6
1999.....	1.3	1.4	0.3	*	0.1	0.5	1.1	1.4	2.0	2.7

All races 3/, male

2006.....	2.1	2.7	0.3	*	0.2	0.9	2.6	2.6	2.6	3.0
2005.....	1.8	3.2	0.5	*	0.2	0.8	1.9	2.2	2.2	2.6
2004.....	2.1	3.1	0.3	*	0.2	0.9	2.1	2.4	2.7	3.7
2003.....	2.2	2.4	*	*	0.2	0.9	2.4	2.4	3.0	4.1
2002.....	2.1	2.5	0.3	*	0.2	0.7	2.1	2.4	3.0	4.3
2001.....	1.9	2.4	*	*	0.2	0.8	1.8	2.1	2.6	3.8
2000.....	1.8	1.4	0.3	*	0.3	0.7	1.6	2.0	2.4	3.6
1999.....	1.8	1.7	*	*	0.2	0.7	1.7	2.0	2.7	3.7

All races 3/, female

2006.....	1.2	1.7	0.3	*	*	0.4	1.0	1.1	1.3	2.0
2005.....	1.2	1.8	0.3	*	*	0.4	0.8	1.1	1.5	1.9
2004.....	1.2	2.0	0.3	*	*	0.3	0.8	1.1	1.3	1.9
2003.....	1.2	1.3	*	*	*	0.4	1.0	1.1	1.5	2.1
2002.....	1.1	2.5	0.4	*	*	0.3	0.9	1.0	1.4	2.0
2001.....	0.9	1.7	0.3	*	*	0.3	0.6	0.8	1.3	2.0
2000.....	0.8	2.0	*	*	*	0.3	0.5	0.8	1.0	1.6
1999.....	0.8	1.2	0.3	*	*	0.2	0.5	0.8	1.3	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

White, both sexes

2006 4/.....	1.7	2.0	0.2	*	0.2	0.8	2.0	2.0	2.2	2.6
2005 4/.....	1.6	2.1	0.3	*	0.1	0.6	1.5	1.9	1.9	2.4
2004 4/.....	1.7	2.1	0.2	*	0.1	0.7	1.7	2.0	2.1	2.9
2003 4/.....	1.7	1.7	0.2	*	0.1	0.8	1.9	1.9	2.5	3.3
2002.....	1.6	1.8	0.4	*	0.2	0.6	1.7	1.8	2.2	3.3
2001.....	1.4	1.4	0.3	*	0.2	0.6	1.3	1.6	2.0	2.9
2000.....	1.3	1.4	0.2	*	0.1	0.5	1.1	1.5	1.8	2.6
1999.....	1.3	1.2	0.2	*	0.1	0.5	1.2	1.5	2.0	2.6

White, male

2006 4/.....	2.1	2.5	*	*	*	1.1	2.9	2.8	2.8	3.0
2005 4/.....	1.8	2.8	0.4	*	*	0.8	2.1	2.4	2.3	2.7
2004 4/.....	2.1	2.7	*	*	*	1.0	2.4	2.8	2.8	3.8
2003 4/.....	2.2	1.9	*	*	*	1.1	2.7	2.5	3.2	4.2
2002.....	2.0	1.8	0.4	*	*	0.8	2.3	2.5	3.0	4.3
2001.....	1.9	1.7	*	*	0.3	0.9	1.9	2.2	2.7	3.6
2000.....	1.7	*	*	*	0.3	0.7	1.6	2.1	2.4	3.6
1999.....	1.7	1.4	*	*	*	0.8	1.8	2.0	2.7	3.5

White, female

2006 4/.....	1.3	1.4	*	*	*	0.5	1.1	1.2	1.5	2.1
2005 4/.....	1.3	1.5	*	*	*	0.4	0.8	1.3	1.6	2.0
2004 4/.....	1.2	1.5	*	*	*	0.3	0.9	1.2	1.4	2.0
2003 4/.....	1.3	1.4	*	*	*	0.5	1.2	1.2	1.7	2.2
2002.....	1.2	1.8	0.4	*	*	0.4	1.0	1.1	1.5	2.2
2001.....	1.0	*	*	*	*	0.3	0.8	0.9	1.4	2.1
2000.....	0.8	1.6	*	*	*	0.3	0.6	0.9	1.1	1.6
1999.....	0.8	*	*	*	*	*	0.5	0.9	1.4	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Black, both sexes

2006 4/.....	1.7	3.4	0.8	*	*	*	0.9	1.1	1.5	2.7
2005 4/.....	1.7	4.7	0.8	*	*	*	0.9	1.3	1.8	2.5
2004 4/.....	1.7	4.8	*	*	*	*	0.8	0.8	2.5	3.1
2003 4/.....	1.7	3.2	*	*	*	*	0.8	1.3	2.0	3.2
2002.....	1.9	5.5	*	*	*	*	1.1	1.8	2.4	3.9
2001.....	1.7	5.5	*	*	*	*	0.8	1.3	1.9	3.8
2000.....	1.7	*	*	*	*	*	0.9	1.4	1.9	3.2
1999.....	1.7	*	*	*	*	*	1.1	1.5	2.4	3.8

Black, male

2006 4/.....	2.5	*	*	*	*	*	1.2	1.8	2.2	3.6
2005 4/.....	2.3	*	*	*	*	*	*	1.9	2.2	3.0
2004 4/.....	2.4	*	*	*	*	*	*	*	3.7	4.5
2003 4/.....	2.6	*	*	*	*	*	1.4	2.2	3.2	4.3
2002.....	2.7	*	*	*	*	*	1.4	2.7	3.4	6.3
2001.....	2.5	6.0	*	*	*	*	1.5	2.2	2.7	5.9
2000.....	2.5	*	*	*	*	*	1.5	2.2	3.1	4.5
1999.....	2.7	*	*	*	*	*	1.8	2.7	3.5	6.1

Black, female

2006 4/.....	1.0	*	*	*	*	*	*	*	*	1.8
2005 4/.....	1.1	*	*	*	*	*	*	*	1.5	2.0
2004 4/.....	1.1	*	*	*	*	*	*	*	*	1.8
2003 4/.....	0.9	*	*	*	*	*	*	*	*	2.1
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.8
2001.....	0.8	*	*	*	*	*	*	*	*	2.0
2000.....	0.9	*	*	*	*	*	*	*	*	2.0
1999.....	0.9	*	*	*	*	*	*	*	1.4	1.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All Under 1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39
ages 1/ year 2/ years years years years years years years years

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes

2006.....	0.8	0.6	0.1	*	*	0.1	0.1	0.1	0.2	0.2
2005.....	0.9	*	*	*	*	*	*	0.2	0.2	0.3
2004.....	1.0	*	*	*	*	0.1	0.1	0.2	0.2	0.3
2003.....	1.0	*	0.2	*	*	0.1	0.1	0.1	0.2	0.4
2002.....	1.0	*	*	*	0.1	*	*	0.1	0.2	0.3
2001.....	1.1	0.6	0.2	*	*	*	0.1	0.1	0.3	0.3
2000.....	1.1	*	0.1	*	*	*	0.1	0.2	0.2	0.3
1999.....	1.0	1.4	0.2	*	*	*	*	0.1	0.2	0.3

All races 3/, male

2006.....	0.8	*	*	*	*	*	*	*	*	*
2005.....	0.8	*	*	*	*	*	*	*	*	0.2
2004.....	0.9	*	*	*	*	*	*	*	*	0.2
2003.....	0.9	*	*	*	*	*	*	*	0.2	0.3
2002.....	0.9	*	*	*	*	*	*	*	0.3	0.3
2001.....	1.0	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.0	*	*	*	*	*	*	0.2	0.2	0.3
1999.....	1.0	1.6	*	*	*	*	*	*	0.3	0.3

All races 3/, female

2006.....	0.9	*	*	*	*	*	*	0.2	*	0.2
2005.....	1.0	*	*	*	*	*	*	*	0.3	0.3
2004.....	1.0	*	*	*	*	*	*	*	0.3	0.3
2003.....	1.0	*	*	*	*	*	*	*	0.2	0.4
2002.....	1.1	*	*	*	*	*	*	*	0.2	0.3
2001.....	1.1	*	*	*	*	*	*	*	0.2	0.3
2000.....	1.1	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.1	1.2	*	*	*	*	*	0.2	*	0.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 6/

All races 3/, both sexes

2006.....	7.4	*	*	*	*	0.2	0.5	1.0	2.2	4.6
2005.....	7.3	*	*	*	*	0.2	0.5	1.0	1.7	4.7
2004.....	7.2	*	*	*	*	0.2	0.5	0.9	2.1	5.0
2003.....	7.1	*	*	*	*	0.2	0.4	0.9	2.1	5.3
2002.....	7.0	*	*	*	*	0.2	0.4	0.8	2.1	5.4
2001.....	7.1	*	*	*	*	0.2	0.3	0.9	2.3	6.0
2000.....	7.0	*	*	*	*	0.2	0.3	0.9	2.3	6.0
1999.....	7.0	*	*	*	*	0.1	0.4	0.8	2.4	6.2

All races 3/, male

2006.....	11.2	*	*	*	*	0.4	0.7	1.7	3.2	6.3
2005.....	11.1	*	*	*	*	0.3	0.9	1.5	2.5	6.6
2004.....	11.0	*	*	*	*	0.2	0.8	1.4	3.2	7.1
2003.....	10.9	*	*	*	*	0.4	0.6	1.4	3.2	7.4
2002.....	10.8	*	*	*	*	0.3	0.6	1.4	3.1	7.6
2001.....	10.8	*	*	*	*	0.3	0.6	1.3	3.3	8.5
2000.....	10.9	*	*	*	*	0.3	0.5	1.5	3.5	8.7
1999.....	10.9	*	*	*	*	0.2	0.6	1.2	3.3	8.7

All races 3/, female

2006.....	3.7	*	*	*	*	*	0.2	0.4	1.1	2.8
2005.....	3.6	*	*	*	*	*	*	0.4	1.0	2.7
2004.....	3.5	*	*	*	*	*	*	0.4	1.0	2.9
2003.....	3.4	*	*	*	*	*	*	0.3	0.9	3.2
2002.....	3.4	*	*	*	*	*	*	0.3	1.1	3.3
2001.....	3.4	*	*	*	*	*	*	0.5	1.3	3.5
2000.....	3.2	*	*	*	*	*	*	0.3	1.1	3.2
1999.....	3.2	*	*	*	*	*	*	0.3	1.5	3.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 6/

White, both sexes

2006 4/.....	7.8	*	*	*	*	0.2	0.5	1.0	2.2	4.7
2005 4/.....	7.7	*	*	*	*	0.2	0.5	1.0	1.7	4.9
2004 4/.....	7.5	*	*	*	*	0.2	0.5	0.9	2.2	5.2
2003 4/.....	7.4	*	*	*	*	0.2	0.4	0.9	2.1	5.4
2002.....	7.2	*	*	*	*	0.2	0.3	0.8	2.1	5.5
2001.....	7.2	*	*	*	*	0.2	0.3	0.9	2.3	6.0
2000.....	7.1	*	*	*	*	0.2	0.3	0.9	2.2	5.9
1999.....	7.0	*	*	*	*	*	0.4	0.8	2.4	6.0

White, male

2006 4/.....	11.8	*	*	*	*	0.4	0.8	1.6	3.4	6.5
2005 4/.....	11.7	*	*	*	*	0.3	0.9	1.5	2.5	6.9
2004 4/.....	11.5	*	*	*	*	*	0.8	1.3	3.3	7.2
2003 4/.....	11.3	*	*	*	*	0.4	0.6	1.5	3.2	7.6
2002.....	11.1	*	*	*	*	0.3	0.6	1.4	3.1	7.6
2001.....	11.0	*	*	*	*	0.3	0.5	1.3	3.3	8.4
2000.....	11.0	*	*	*	*	0.3	0.5	1.5	3.4	8.7
1999.....	10.9	*	*	*	*	*	0.7	1.3	3.2	8.5

White, female

2006 4/.....	3.9	*	*	*	*	*	*	0.3	0.9	2.7
2005 4/.....	3.7	*	*	*	*	*	*	0.4	0.9	2.7
2004 4/.....	3.6	*	*	*	*	*	*	0.4	1.0	3.1
2003 4/.....	3.5	*	*	*	*	*	*	0.3	1.0	3.2
2002.....	3.4	*	*	*	*	*	*	*	1.0	3.3
2001.....	3.4	*	*	*	*	*	*	0.4	1.2	3.6
2000.....	3.2	*	*	*	*	*	*	*	1.1	3.1
1999.....	3.1	*	*	*	*	*	*	*	1.4	3.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 6/

Black, both sexes

2006 4/.....	5.5	*	*	*	*	*	*	0.7	1.2	3.2
2005 4/.....	5.9	*	*	*	*	*	*	*	1.2	3.0
2004 4/.....	6.1	*	*	*	*	*	*	*	1.3	3.7
2003 4/.....	6.3	*	*	*	*	*	*	0.9	1.3	4.7
2002.....	6.4	*	*	*	*	*	*	*	1.9	4.4
2001.....	7.3	*	*	*	*	*	*	0.8	1.9	5.2
2000.....	7.4	*	*	*	*	*	*	0.9	2.5	5.5
1999.....	7.8	*	*	*	*	*	*	*	2.2	7.1

Black, male

2006 4/.....	8.5	*	*	*	*	*	*	*	*	4.1
2005 4/.....	9.1	*	*	*	*	*	*	*	1.5	4.1
2004 4/.....	9.7	*	*	*	*	*	*	*	2.2	5.9
2003 4/.....	10.0	*	*	*	*	*	*	*	2.2	6.5
2002.....	10.0	*	*	*	*	*	*	*	2.9	6.2
2001.....	11.6	*	*	*	*	*	*	*	2.5	8.4
2000.....	11.4	*	*	*	*	*	*	1.6	3.4	8.1
1999.....	12.2	*	*	*	*	*	*	*	2.9	10.2

Black, female

2006 4/.....	2.8	*	*	*	*	*	*	*	*	2.4
2005 4/.....	3.0	*	*	*	*	*	*	*	*	2.1
2004 4/.....	2.8	*	*	*	*	*	*	*	*	1.7
2003 4/.....	2.9	*	*	*	*	*	*	*	*	3.2
2002.....	3.2	*	*	*	*	*	*	*	*	2.9
2001.....	3.5	*	*	*	*	*	*	*	*	2.4
2000.....	3.7	*	*	*	*	*	*	*	1.7	3.2
1999.....	3.9	*	*	*	*	*	*	*	1.5	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

All races 3/, both sexes

2006.....	12.8	1.1	0.2	*	0.2	4.1	13.1	16.5	17.1	20.2
2005.....	11.3	0.9	0.2	*	0.2	3.7	11.0	13.5	15.1	18.3
2004.....	10.5	0.7	0.2	*	0.2	3.9	10.0	12.2	13.1	18.4
2003.....	9.9	0.6	0.2	*	0.2	3.3	9.4	10.9	13.3	18.3
2002.....	9.0	0.7	0.2	*	0.2	3.1	7.9	9.8	12.4	17.5
2001.....	7.6	0.5	0.2	*	0.2	2.7	6.5	8.2	10.6	15.5
2000.....	7.0	*	*	*	0.2	2.2	5.9	7.5	10.1	14.4
1999.....	6.9	0.6	0.2	*	0.1	1.8	5.3	7.4	10.3	14.2

All races 3/, male

2006.....	16.6	1.1	0.2	*	0.2	6.0	19.5	23.3	23.5	25.8
2005.....	14.5	*	0.3	*	*	5.4	16.0	18.9	20.0	23.0
2004.....	13.4	*	*	*	0.2	5.6	14.7	17.3	17.6	23.6
2003.....	12.9	*	0.3	*	*	4.5	13.9	15.4	17.7	23.8
2002.....	11.8	1.0	*	*	*	4.3	11.4	14.0	17.0	23.1
2001.....	10.2	*	0.3	*	*	3.7	9.6	11.9	14.4	20.7
2000.....	9.5	*	*	*	*	3.1	8.8	10.8	14.1	19.8
1999.....	9.4	*	*	*	*	2.4	7.7	10.8	14.6	19.6

All races 3/, female

2006.....	9.1	1.0	*	*	*	2.2	6.3	9.3	10.7	14.5
2005.....	8.2	*	*	*	0.2	2.0	5.6	7.7	10.1	13.5
2004.....	7.6	*	*	*	0.3	2.1	5.0	6.7	8.5	13.2
2003.....	7.0	*	*	*	*	1.9	4.7	6.2	8.8	12.8
2002.....	6.3	*	0.3	*	*	1.8	4.2	5.4	7.8	11.9
2001.....	5.1	*	*	*	*	1.6	3.2	4.4	6.7	10.3
2000.....	4.6	*	*	*	*	1.4	2.9	4.1	6.1	9.0
1999.....	4.4	*	*	*	*	1.2	2.8	4.0	6.0	8.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

White, both sexes

2006 4/.....	13.6	0.9	0.2	*	0.2	5.0	15.6	19.2	19.2	21.7
2005 4/.....	12.0	0.8	*	*	0.2	4.5	13.0	15.3	16.8	19.8
2004 4/.....	11.1	*	*	*	0.2	4.7	11.8	14.0	14.5	19.7
2003 4/.....	10.4	*	0.2	*	0.1	3.9	11.2	12.4	14.7	19.4
2002.....	9.4	*	*	*	0.2	3.6	9.2	10.9	13.5	18.5
2001.....	7.8	*	*	*	0.2	3.1	7.5	9.1	11.4	16.1
2000.....	7.1	*	*	*	0.2	2.6	6.7	8.2	10.7	14.8
1999.....	6.9	*	*	*	0.1	2.1	5.9	7.9	10.7	14.4

White, male

2006 4/.....	17.3	*	*	*	*	7.2	23.2	26.8	25.8	27.4
2005 4/.....	15.3	*	*	*	*	6.5	19.0	21.5	22.2	24.8
2004 4/.....	14.1	*	*	*	*	6.7	17.3	19.9	19.5	25.3
2003 4/.....	13.5	*	*	*	*	5.5	16.4	17.6	19.5	25.1
2002.....	12.2	*	*	*	*	5.0	13.2	15.5	18.5	24.2
2001.....	10.4	*	*	*	*	4.4	10.9	13.1	15.6	21.6
2000.....	9.6	*	*	*	*	3.6	10.0	11.6	14.9	20.4
1999.....	9.3	*	*	*	*	2.8	8.5	11.3	15.2	19.9

White, female

2006 4/.....	9.9	*	*	*	*	2.6	7.4	11.0	12.2	15.8
2005 4/.....	8.8	*	*	*	*	2.3	6.5	8.8	11.1	14.6
2004 4/.....	8.2	*	*	*	0.3	2.5	5.8	7.7	9.4	14.1
2003 4/.....	7.4	*	*	*	*	2.2	5.5	6.8	9.7	13.6
2002.....	6.7	*	*	*	*	2.1	4.9	6.1	8.3	12.6
2001.....	5.4	*	*	*	*	1.6	3.8	5.0	7.0	10.6
2000.....	4.7	*	*	*	*	1.5	3.2	4.6	6.3	9.1
1999.....	4.5	*	*	*	*	1.4	3.0	4.3	6.1	8.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/										
Black, both sexes										
2006 4/.....	12.1	*	*	*	*	0.9	4.3	8.2	12.2	18.8
2005 4/.....	10.5	*	*	*	*	1.0	3.4	7.5	11.5	16.1
2004 4/.....	9.4	*	*	*	*	1.0	3.5	6.2	9.6	16.3
2003 4/.....	9.2	*	*	*	*	0.7	2.9	5.8	10.4	17.9
2002.....	9.2	*	*	*	*	1.0	3.2	6.6	10.2	17.3
2001.....	8.5	*	*	*	*	0.9	3.0	5.8	9.5	16.7
2000.....	8.3	*	*	*	*	0.9	3.2	6.2	10.3	16.5
1999.....	8.6	*	*	*	*	0.7	3.7	6.8	11.2	16.5
Black, male										
2006 4/.....	17.3	*	*	*	*	1.1	5.8	12.8	18.6	25.9
2005 4/.....	14.3	*	*	*	*	*	4.8	11.1	14.8	20.1
2004 4/.....	12.8	*	*	*	*	1.4	5.0	9.0	13.2	20.5
2003 4/.....	12.7	*	*	*	*	*	4.4	7.7	13.6	24.3
2002.....	12.8	*	*	*	*	1.4	4.6	9.9	13.9	24.2
2001.....	12.2	*	*	*	*	*	5.1	9.3	12.3	22.4
2000.....	12.0	*	*	*	*	*	4.6	9.8	14.3	22.8
1999.....	12.7	*	*	*	*	*	5.6	11.0	15.7	23.9
Black, female										
2006 4/.....	7.4	*	*	*	*	*	2.8	3.9	6.5	12.4
2005 4/.....	7.0	*	*	*	*	*	2.0	4.0	8.6	12.5
2004 4/.....	6.3	*	*	*	*	*	2.0	3.6	6.4	12.5
2003 4/.....	6.1	*	*	*	*	*	1.5	4.0	7.6	12.2
2002.....	5.8	*	*	*	*	*	1.9	3.6	6.8	11.2
2001.....	5.1	*	*	*	*	*	*	2.6	7.0	11.6
2000.....	4.9	*	*	*	*	*	1.8	2.9	6.8	11.0
1999.....	4.8	*	*	*	*	*	1.9	3.1	7.3	10.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/										
All races 3/, both sexes										
2006.....	10.3	*	0.3	0.4	1.3	13.2	20.6	17.0	13.5	11.8
2005.....	10.4	*	0.4	0.3	1.3	12.5	20.0	17.2	14.2	11.6
2004.....	10.1	*	0.3	0.3	1.1	12.0	19.3	16.9	13.3	11.2
2003.....	10.4	*	0.3	0.3	1.2	12.1	21.1	17.4	13.8	11.6
2002.....	10.5	*	0.4	0.4	1.3	12.1	21.3	17.8	13.2	12.4
2001.....	10.4	*	0.5	0.4	1.2	12.4	21.2	17.7	13.5	12.2
2000.....	10.2	*	0.3	0.3	1.5	12.9	20.9	16.3	12.8	12.0
1999.....	10.3	*	0.4	0.4	1.7	14.4	21.0	16.6	13.2	12.3
All races 3/, male										
2006.....	18.1	*	0.5	0.4	2.0	23.3	36.2	29.4	23.6	19.5
2005.....	18.3	*	0.5	0.4	2.0	22.0	35.3	30.2	24.3	19.3
2004.....	17.6	*	0.4	0.3	1.8	20.7	34.2	29.8	22.5	19.0
2003.....	18.3	*	0.3	0.4	2.0	21.2	37.1	30.7	23.7	19.3
2002.....	18.4	*	0.5	0.4	1.9	21.1	37.6	31.1	22.3	20.6
2001.....	18.2	*	0.5	0.5	1.9	21.8	37.7	30.9	22.9	20.0
2000.....	17.8	*	0.4	0.4	2.4	22.4	37.0	28.5	21.4	19.5
1999.....	18.1	*	0.5	0.5	2.5	24.8	36.8	28.8	22.3	20.0
All races 3/, female										
2006.....	2.8	*	*	0.4	0.5	2.5	3.8	4.1	3.2	4.1
2005.....	2.7	*	0.3	0.3	0.6	2.4	3.6	3.7	3.8	3.9
2004.....	2.7	*	0.3	0.3	0.4	2.9	3.5	3.4	3.9	3.3
2003.....	2.7	*	0.3	0.2	0.4	2.4	4.2	3.6	3.7	3.8
2002.....	2.8	*	0.3	0.3	0.7	2.7	4.2	4.1	3.9	4.2
2001.....	2.8	*	0.4	0.3	0.5	2.6	3.8	4.1	3.9	4.6
2000.....	2.8	*	*	0.2	0.5	2.9	4.2	3.9	4.1	4.5
1999.....	2.9	*	0.4	0.3	0.8	3.4	4.4	4.1	4.1	4.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

White, both sexes

2006 4/.....	9.0	*	0.2	0.3	1.0	8.4	12.9	11.2	9.9	10.5
2005 4/.....	9.1	*	0.3	0.2	1.1	8.5	12.6	11.7	10.7	10.2
2004 4/.....	9.0	*	*	0.2	0.9	8.5	12.9	11.4	10.4	10.0
2003 4/.....	9.2	*	0.2	0.2	1.0	8.4	13.9	11.7	10.8	10.4
2002.....	9.3	*	0.3	0.3	1.1	8.7	13.9	12.3	10.2	10.9
2001.....	9.4	*	0.4	0.4	1.1	8.9	14.0	12.9	11.1	11.2
2000.....	9.1	*	0.2	0.3	1.4	9.2	13.9	11.5	10.4	10.9
1999.....	9.2	*	0.3	0.3	1.6	10.2	14.0	12.1	11.1	11.3

White, male

2006 4/.....	15.4	*	*	0.3	1.6	14.6	22.1	18.6	16.7	16.7
2005 4/.....	15.8	*	0.3	0.3	1.7	14.6	21.8	19.7	17.6	16.3
2004 4/.....	15.5	*	*	*	1.5	14.4	22.3	19.5	17.0	16.6
2003 4/.....	16.0	*	*	*	1.6	14.4	23.9	19.8	17.9	16.8
2002.....	16.1	*	0.4	0.3	1.5	14.9	23.9	20.5	16.7	17.7
2001.....	16.2	*	0.4	0.5	1.7	15.0	24.1	21.8	18.2	17.9
2000.....	15.6	*	*	0.4	2.3	15.6	23.9	19.3	16.9	17.2
1999.....	15.9	*	*	0.3	2.3	17.3	23.8	20.3	18.1	18.1

White, female

2006 4/.....	2.6	*	*	0.3	0.4	1.8	2.8	3.2	2.7	4.0
2005 4/.....	2.6	*	*	*	0.4	1.9	2.6	3.2	3.6	3.9
2004 4/.....	2.7	*	*	*	0.3	2.3	2.8	2.8	3.5	3.2
2003 4/.....	2.6	*	*	*	0.3	2.0	3.1	3.0	3.3	3.9
2002.....	2.7	*	*	0.3	0.6	2.1	3.1	3.7	3.4	4.0
2001.....	2.7	*	*	0.3	0.5	2.3	3.1	3.4	3.7	4.3
2000.....	2.7	*	*	*	0.5	2.3	3.2	3.3	3.7	4.5
1999.....	2.8	*	*	*	0.7	2.6	3.5	3.4	3.8	4.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Black, both sexes

2006 4/.....	21.3	*	1.1	0.7	2.7	37.5	64.3	54.2	39.5	24.6
2005 4/.....	20.4	*	1.0	0.8	2.4	33.6	61.8	52.6	39.4	23.8
2004 4/.....	19.3	*	1.2	0.8	2.2	30.9	58.0	53.1	34.5	22.0
2003 4/.....	20.1	*	0.9	0.8	2.5	32.2	63.8	55.5	35.8	21.5
2002.....	20.2	*	*	*	2.4	31.4	64.7	54.8	35.1	24.4
2001.....	19.3	*	0.9	*	1.9	32.9	64.4	50.4	31.5	22.6
2000.....	19.3	*	*	*	2.2	33.9	64.0	48.1	30.9	21.7
1999.....	19.4	*	1.1	0.9	2.3	38.0	63.5	46.5	29.0	20.9

Black, male

2006 4/.....	40.0	*	*	*	4.3	68.2	117.7	101.0	75.4	45.6
2005 4/.....	38.7	*	*	*	3.7	61.5	114.1	100.1	75.9	44.4
2004 4/.....	36.4	*	*	*	3.5	55.2	107.7	102.1	64.9	41.3
2003 4/.....	37.8	*	*	*	3.7	59.3	117.4	107.6	67.4	40.3
2002.....	37.8	*	*	*	3.5	56.0	120.6	106.3	65.9	44.8
2001.....	36.4	*	*	*	3.0	60.5	123.4	96.0	60.1	39.5
2000.....	36.1	*	*	*	3.7	61.5	121.3	92.9	57.0	39.9
1999.....	36.0	*	*	1.3	3.4	67.1	119.9	88.6	54.3	37.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Black, female

2006 4/.....	4.1	*	*	*	*	6.0	9.5	9.8	6.9	5.9
2005 4/.....	3.7	*	*	*	1.2	5.0	8.5	7.7	6.4	5.3
2004 4/.....	3.7	*	*	*	*	5.9	7.9	7.3	7.0	4.7
2003 4/.....	3.9	*	*	*	1.3	4.2	10.5	7.4	7.2	4.7
2002.....	4.2	*	*	*	1.3	6.0	10.2	7.7	7.4	6.3
2001.....	3.8	*	*	*	*	4.5	7.8	8.9	5.8	7.5
2000.....	4.0	*	*	*	*	5.7	9.6	7.5	7.4	5.5
1999.....	4.4	*	*	*	*	8.2	9.9	8.4	6.4	6.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

All causes

All races 3/, both sexes

2006.....	229.3	347.7	516.4	730.1	1,110.2	1,656.6	2,554.8	4,033.4	6,524.0	13,253.1
2005.....	233.4	353.1	520.8	734.6	1,136.9	1,700.0	2,657.6	4,154.0	6,712.9	13,798.6
2004.....	231.7	352.3	511.7	734.8	1,140.1	1,727.4	2,676.4	4,193.2	6,717.2	13,823.5
2003.....	241.3	359.2	517.8	755.6	1,182.7	1,793.7	2,778.3	4,363.4	6,976.2	14,593.3
2002.....	242.6	357.1	512.8	770.1	1,187.7	1,832.7	2,845.9	4,449.1	7,103.8	14,828.3
2001.....	240.5	354.8	512.4	771.8	1,210.7	1,869.7	2,878.3	4,494.0	7,151.9	15,112.8
2000.....	239.1	352.5	509.0	792.6	1,241.0	1,906.3	2,929.4	4,553.8	7,335.0	15,524.4
1999.....	236.3	346.6	501.4	801.6	1,254.0	1,956.7	2,995.3	4,608.0	7,378.5	15,554.6

All races 3/, male

2006.....	285.9	435.3	659.7	920.0	1,373.6	2,040.2	3,117.5	4,944.6	7,942.7	14,309.1
2005.....	292.7	443.7	666.0	925.7	1,410.0	2,084.2	3,267.0	5,103.4	8,147.4	14,889.4
2004.....	289.5	444.4	657.0	917.3	1,411.0	2,125.1	3,279.2	5,173.5	8,179.0	15,031.1
2003.....	304.1	457.9	660.8	942.3	1,462.2	2,216.6	3,428.7	5,400.3	8,513.7	15,794.0
2002.....	306.4	454.7	653.5	961.8	1,476.0	2,265.9	3,533.0	5,522.7	8,652.3	16,254.5
2001.....	304.8	455.4	646.8	958.4	1,496.8	2,315.6	3,587.4	5,603.2	8,781.9	16,744.8
2000.....	305.9	456.3	642.2	989.8	1,535.9	2,362.0	3,676.1	5,693.2	9,095.4	17,501.4
1999.....	304.0	448.0	632.4	1,003.8	1,560.8	2,444.1	3,785.1	5,806.4	9,216.8	17,597.2

All races 3/, female

2006.....	173.1	262.2	378.9	551.0	869.7	1,321.4	2,097.0	3,366.6	5,637.2	12,759.0
2005.....	174.7	264.6	381.6	554.3	887.9	1,364.7	2,164.0	3,464.7	5,822.0	13,297.7
2004.....	174.5	262.5	372.7	562.7	893.6	1,381.2	2,191.0	3,488.7	5,816.6	13,280.3
2003.....	179.2	263.2	380.9	579.7	928.8	1,427.1	2,256.6	3,625.3	6,036.0	14,062.5
2002.....	179.7	262.4	378.2	590.0	926.0	1,458.7	2,296.7	3,690.9	6,162.2	14,209.6
2001.....	177.2	257.2	383.7	596.9	951.3	1,485.9	2,314.8	3,717.6	6,175.1	14,429.9
2000.....	173.4	252.0	381.3	608.1	973.7	1,515.7	2,341.2	3,760.2	6,296.6	14,719.2
1999.....	169.8	248.5	375.9	612.6	976.4	1,539.6	2,378.2	3,774.8	6,302.8	14,731.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All causes

White, both sexes

2006 4/.....	214.0	323.7	479.5	686.5	1,064.0	1,607.4	2,519.6	4,020.7	6,550.9	13,480.4
2005 4/.....	217.8	326.6	480.7	689.7	1,085.7	1,649.5	2,618.1	4,128.9	6,731.9	14,021.1
2004 4/.....	215.0	325.0	471.5	688.5	1,090.5	1,675.5	2,634.1	4,160.5	6,736.0	14,010.1
2003 4/.....	222.6	329.4	474.5	710.6	1,127.8	1,739.2	2,728.6	4,329.5	6,992.1	14,792.3
2002.....	224.0	327.4	471.3	723.5	1,133.0	1,777.6	2,796.3	4,407.8	7,111.3	15,015.9
2001.....	218.8	322.8	471.4	723.2	1,156.1	1,805.8	2,823.2	4,446.8	7,149.4	15,287.5
2000.....	215.9	317.9	467.4	743.9	1,184.9	1,844.5	2,866.8	4,496.6	7,333.2	15,707.0
1999.....	210.8	310.5	460.0	747.2	1,193.5	1,895.5	2,923.9	4,542.8	7,370.9	15,718.0

White, male

2006 4/.....	269.8	408.3	612.8	861.2	1,309.5	1,973.1	3,060.0	4,914.2	7,982.5	14,576.8
2005 4/.....	276.4	413.4	616.6	864.0	1,341.8	2,016.7	3,208.8	5,065.8	8,169.0	15,156.5
2004 4/.....	272.8	412.3	607.8	855.8	1,343.7	2,055.9	3,219.3	5,123.2	8,214.4	15,250.7
2003 4/.....	284.0	423.5	606.9	884.2	1,390.6	2,146.2	3,359.5	5,349.4	8,522.6	16,037.9
2002.....	287.2	420.4	601.2	902.0	1,402.7	2,196.1	3,469.7	5,469.3	8,660.8	16,473.2
2001.....	281.4	417.1	595.5	896.0	1,427.9	2,237.6	3,513.9	5,543.5	8,771.1	16,935.4
2000.....	279.4	416.1	588.3	925.1	1,465.1	2,285.1	3,593.1	5,620.3	9,092.5	17,716.4
1999.....	274.0	402.1	579.3	934.6	1,484.0	2,367.8	3,698.3	5,727.2	9,211.2	17,752.9

White, female

2006 4/.....	157.3	239.3	348.7	518.0	835.1	1,281.6	2,071.0	3,353.8	5,646.4	12,965.7
2005 4/.....	158.6	239.9	347.5	521.9	847.6	1,322.9	2,129.8	3,435.6	5,831.3	13,498.3
2004 4/.....	156.6	238.0	338.1	527.6	855.6	1,338.1	2,153.2	3,456.3	5,817.7	13,450.9
2003 4/.....	160.7	235.8	344.9	543.9	884.5	1,379.6	2,212.3	3,592.1	6,048.6	14,240.6
2002.....	160.5	235.1	344.3	552.6	883.7	1,409.1	2,247.4	3,648.0	6,161.6	14,382.8
2001.....	155.9	229.4	350.1	558.4	905.3	1,426.7	2,263.9	3,669.9	6,169.3	14,597.6
2000.....	152.2	220.7	349.3	571.5	926.5	1,459.3	2,284.0	3,706.2	6,286.6	14,890.7
1999.....	147.4	219.9	343.6	569.0	925.9	1,483.5	2,308.5	3,711.2	6,284.5	14,900.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
Black, both sexes										
2006 4/.....	373.1	576.3	892.9	1,232.7	1,750.2	2,409.9	3,371.9	4,902.4	7,275.3	12,489.7
2005 4/.....	379.3	602.7	914.9	1,254.6	1,812.2	2,466.3	3,514.3	5,161.7	7,547.7	13,083.1
2004 4/.....	386.6	608.3	910.1	1,271.1	1,813.5	2,509.7	3,582.0	5,266.0	7,493.0	13,338.5
2003 4/.....	410.4	632.9	940.3	1,295.1	1,889.9	2,595.8	3,754.9	5,421.2	7,745.3	13,976.7
2002.....	412.9	634.1	925.9	1,321.6	1,880.7	2,632.0	3,824.9	5,571.3	7,962.2	14,260.8
2001.....	428.4	644.1	928.5	1,328.9	1,891.7	2,738.8	3,904.2	5,660.2	8,028.9	14,550.8
2000.....	434.5	666.6	938.7	1,347.0	1,929.0	2,769.0	4,019.7	5,782.6	8,161.0	14,714.3
1999.....	448.6	668.3	929.9	1,399.3	1,969.2	2,813.7	4,148.0	5,878.3	8,169.4	14,757.9
Black, male										
2006 4/.....	457.3	715.6	1,171.2	1,625.1	2,302.6	3,132.8	4,373.3	6,329.2	9,021.0	13,206.0
2005 4/.....	463.8	755.9	1,184.8	1,667.1	2,380.7	3,174.7	4,509.2	6,579.2	9,376.2	13,809.8
2004 4/.....	467.1	766.8	1,186.6	1,668.7	2,374.4	3,242.7	4,587.3	6,747.0	9,229.2	14,452.5
2003 4/.....	507.1	800.8	1,227.5	1,688.7	2,454.5	3,334.8	4,843.9	6,978.2	9,827.8	14,903.4
2002.....	502.9	801.4	1,206.6	1,719.3	2,464.7	3,357.1	4,918.1	7,088.6	9,968.4	15,635.5
2001.....	527.8	830.2	1,201.4	1,718.2	2,424.6	3,497.3	5,053.2	7,255.7	10,256.3	16,439.9
2000.....	549.1	853.0	1,227.4	1,776.2	2,462.1	3,513.8	5,203.9	7,410.0	10,405.7	16,791.0
1999.....	572.7	885.6	1,220.1	1,822.6	2,549.7	3,599.1	5,336.3	7,538.1	10,433.8	17,304.5
Black, female										
2006 4/.....	298.9	455.0	657.4	909.1	1,319.8	1,876.0	2,683.8	4,048.5	6,352.9	12,196.7
2005 4/.....	304.8	469.0	686.6	914.1	1,369.2	1,940.8	2,833.5	4,316.0	6,581.9	12,789.9
2004 4/.....	315.7	470.1	676.4	943.8	1,376.3	1,965.5	2,899.8	4,388.5	6,578.2	12,896.9
2003 4/.....	325.1	486.7	697.2	971.3	1,449.4	2,047.3	3,020.5	4,495.8	6,654.1	13,616.7
2002.....	333.6	489.0	688.0	995.2	1,423.7	2,093.9	3,091.2	4,664.6	6,917.8	13,734.2
2001.....	340.6	482.8	696.2	1,009.5	1,472.5	2,178.4	3,138.8	4,707.8	6,894.3	13,849.9
2000.....	333.3	505.0	692.6	995.2	1,508.3	2,219.3	3,233.3	4,801.6	7,030.5	13,941.3
1999.....	339.3	480.9	683.9	1,053.5	1,514.0	2,231.6	3,359.6	4,878.6	7,047.2	13,805.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Certain other intestinal infections (A04,A07-A09)

All races 3/, both sexes

2006.....	0.1	0.2	0.4	0.9	1.5	3.3	6.7	13.5	26.2	50.5
2005.....	0.1	0.1	0.4	0.5	1.2	2.8	6.5	11.6	21.5	45.8
2004.....	0.1	0.2	0.3	0.6	1.3	2.2	4.6	9.3	17.5	34.3
2003.....	*	0.1	0.2	0.4	0.9	1.7	3.3	7.0	12.3	24.8
2002.....	*	0.1	0.2	0.4	0.7	1.4	2.6	5.0	9.7	21.0
2001.....	*	*	0.2	0.2	0.5	0.8	1.8	3.2	6.4	12.8
2000.....	*	*	*	0.2	0.3	0.9	1.5	2.7	5.3	11.8
1999.....	*	*	*	*	0.3	0.7	1.2	2.3	4.5	10.3

All races 3/, male

2006.....	*	0.2	0.4	0.9	1.8	3.7	6.3	13.9	25.3	50.0
2005.....	*	*	0.4	0.5	1.4	2.9	6.3	12.4	22.1	46.1
2004.....	*	*	0.3	0.6	1.3	2.2	4.3	9.3	17.8	32.0
2003.....	*	*	0.2	0.5	0.8	1.5	3.1	6.9	12.9	21.4
2002.....	*	*	*	0.4	0.8	1.5	3.1	5.1	10.1	20.4
2001.....	*	*	*	*	0.4	0.7	2.0	3.5	6.2	12.1
2000.....	*	*	*	0.3	*	1.0	1.6	2.9	5.8	11.4
1999.....	*	*	*	*	*	0.6	1.2	2.7	4.9	9.5

All races 3/, female

2006.....	*	*	0.4	0.8	1.3	3.0	6.9	13.3	26.8	50.8
2005.....	*	*	0.4	0.6	1.0	2.6	6.6	11.0	21.1	45.6
2004.....	*	*	0.3	0.6	1.2	2.2	4.9	9.3	17.3	35.4
2003.....	*	*	0.2	0.4	0.9	1.8	3.5	7.1	12.0	26.3
2002.....	*	*	*	0.4	0.6	1.3	2.3	5.0	9.5	21.3
2001.....	*	*	0.2	*	0.5	0.8	1.8	3.0	6.5	13.1
2000.....	*	*	*	*	*	0.8	1.5	2.6	4.9	11.9
1999.....	*	*	*	*	*	0.8	1.3	2.0	4.3	10.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Certain other intestinal infections (A04,A07-A09)

White, both sexes

2006 4/.....	*	0.2	0.3	0.8	1.5	3.5	6.9	14.4	27.9	53.6
2005 4/.....	*	0.2	0.3	0.5	1.2	2.9	6.7	12.2	22.6	48.6
2004 4/.....	*	0.1	0.3	0.6	1.3	2.2	4.9	9.8	18.4	36.6
2003 4/.....	*	*	0.2	0.5	0.9	1.8	3.5	7.2	13.2	26.1
2002.....	*	0.1	0.2	0.4	0.6	1.5	2.7	5.3	10.3	22.0
2001.....	*	*	0.2	0.2	0.4	0.7	2.0	3.3	6.6	13.5
2000.....	*	*	*	0.3	0.3	0.9	1.6	2.9	5.6	12.2
1999.....	*	*	*	*	0.3	0.7	1.3	2.3	4.7	10.9

White, male

2006 4/.....	*	0.2	0.3	0.9	1.7	3.8	6.5	14.5	26.6	53.1
2005 4/.....	*	*	0.3	0.4	1.5	3.2	6.5	13.0	22.7	49.3
2004 4/.....	*	*	0.3	0.5	1.4	2.2	4.5	9.7	18.8	33.9
2003 4/.....	*	*	*	0.4	0.8	1.6	3.2	7.2	13.7	22.7
2002.....	*	*	*	0.4	0.7	1.6	3.0	5.3	10.7	21.4
2001.....	*	*	*	*	*	0.7	2.0	3.6	6.2	12.7
2000.....	*	*	*	*	*	1.0	1.6	3.0	6.0	11.9
1999.....	*	*	*	*	*	0.7	1.3	2.8	5.2	9.9

White, female

2006 4/.....	*	*	0.4	0.8	1.3	3.1	7.3	14.3	28.8	53.9
2005 4/.....	*	*	0.4	0.5	1.0	2.7	6.9	11.7	22.6	48.3
2004 4/.....	*	*	0.3	0.6	1.2	2.3	5.2	9.9	18.1	37.8
2003 4/.....	*	*	*	0.5	0.9	1.9	3.8	7.2	12.8	27.6
2002.....	*	*	*	0.4	0.6	1.4	2.4	5.3	10.1	22.3
2001.....	*	*	0.3	*	0.5	0.8	1.9	3.2	6.8	13.9
2000.....	*	*	*	*	*	0.8	1.6	2.7	5.3	12.3
1999.....	*	*	*	*	*	0.8	1.3	2.1	4.4	11.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Tuberculosis (A16-A19)

All races 3/, both sexes

2006.....	0.1	0.2	0.2	0.3	0.4	0.6	0.7	0.9	1.7	2.1
2005.....	*	0.2	0.2	0.2	0.4	0.5	0.7	1.3	1.3	2.2
2004.....	0.1	0.1	0.2	0.2	0.3	0.6	1.0	1.1	1.5	2.4
2003.....	0.1	0.2	0.3	0.4	0.3	0.5	0.8	1.4	2.0	2.3
2002.....	0.1	0.2	0.2	0.3	0.5	0.9	0.9	1.5	2.3	2.6
2001.....	0.1	0.2	0.3	0.4	0.5	0.6	1.0	1.4	2.0	2.8
2000.....	0.1	0.2	0.3	0.2	0.5	0.7	1.0	1.8	2.4	2.7
1999.....	0.1	0.3	0.3	0.5	0.6	0.7	1.2	1.8	2.4	4.2

All races 3/, male

2006.....	0.2	0.3	0.4	0.4	0.7	0.8	0.9	1.4	2.5	3.1
2005.....	*	0.2	0.3	0.3	0.6	0.7	1.1	1.7	1.7	3.3
2004.....	0.2	0.2	0.3	0.3	0.6	0.8	1.4	1.2	2.2	3.8
2003.....	*	0.4	0.4	0.6	0.4	0.6	1.2	2.0	2.7	3.1
2002.....	0.2	0.2	0.4	0.5	0.7	1.2	1.2	1.8	3.2	4.0
2001.....	*	0.3	0.4	0.7	0.7	0.9	1.5	1.7	2.4	4.4
2000.....	*	0.4	0.5	0.4	0.7	1.0	1.1	2.3	3.8	4.6
1999.....	0.2	0.4	0.5	0.8	0.9	1.1	1.7	2.4	3.7	7.1

All races 3/, female

2006.....	*	*	*	*	*	*	0.5	0.6	1.3	1.6
2005.....	*	*	*	*	*	0.4	0.4	0.9	1.0	1.7
2004.....	*	*	*	*	*	*	0.6	1.0	1.1	1.7
2003.....	*	*	*	*	*	0.4	0.5	1.0	1.7	1.9
2002.....	*	*	*	*	0.4	0.5	0.6	1.3	1.7	2.1
2001.....	*	*	*	*	*	0.4	0.7	1.1	1.8	2.2
2000.....	*	*	*	*	*	0.5	0.9	1.4	1.6	2.0
1999.....	*	*	*	*	*	0.4	0.7	1.5	1.7	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Tuberculosis (A16-A19)

White, both sexes

2006 4/.....	*	0.1	0.1	0.2	0.3	0.4	0.4	0.8	1.2	1.6
2005 4/.....	*	0.1	0.1	*	0.2	0.4	0.4	0.8	0.9	1.7
2004 4/.....	*	*	0.1	*	0.2	0.3	0.7	0.8	1.2	1.7
2003 4/.....	*	0.1	0.1	0.3	0.2	0.3	0.7	1.0	1.7	1.8
2002.....	*	*	0.1	0.2	0.4	0.5	0.6	1.2	1.8	2.1
2001.....	*	0.1	0.1	0.3	0.3	0.5	0.7	1.1	1.4	2.2
2000.....	*	*	0.2	*	0.3	0.5	0.6	1.2	1.7	2.2
1999.....	*	0.2	0.2	0.3	0.4	0.6	0.8	1.3	2.0	3.2

White, male

2006 4/.....	*	*	0.2	*	0.4	0.5	*	1.0	1.7	2.1
2005 4/.....	*	*	*	*	0.4	0.5	*	1.0	1.0	2.4
2004 4/.....	*	*	*	*	*	*	1.0	1.0	1.6	2.8
2003 4/.....	*	*	*	0.4	*	*	1.0	1.3	2.2	2.4
2002.....	*	*	*	*	0.5	0.8	1.0	1.3	2.5	3.2
2001.....	*	*	*	0.5	0.5	0.6	0.8	1.4	1.8	3.1
2000.....	*	*	0.3	*	*	0.7	0.8	1.5	2.3	3.8
1999.....	*	0.3	0.4	0.5	0.6	0.9	1.1	1.5	3.0	5.4

White, female

2006 4/.....	*	*	*	*	*	*	*	0.6	0.9	1.4
2005 4/.....	*	*	*	*	*	*	*	0.6	0.8	1.4
2004 4/.....	*	*	*	*	*	*	*	0.7	0.9	1.3
2003 4/.....	*	*	*	*	*	*	*	0.9	1.3	1.6
2002.....	*	*	*	*	*	*	*	1.1	1.3	1.6
2001.....	*	*	*	*	*	*	0.5	0.9	1.2	1.8
2000.....	*	*	*	*	*	*	0.5	1.0	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	1.1	1.4	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Respiratory tuberculosis (A16)

All races 3/, both sexes

2006.....	*	0.1	0.2	0.2	0.4	0.4	0.5	0.7	1.4	1.6
2005.....	*	0.1	0.2	0.2	0.3	0.3	0.6	1.0	1.0	1.9
2004.....	0.1	0.1	0.2	0.2	0.3	0.4	0.8	0.8	1.1	2.2
2003.....	0.1	0.2	0.2	0.3	0.2	0.4	0.6	1.2	1.6	1.8
2002.....	0.1	0.1	0.2	0.2	0.4	0.6	0.7	1.2	2.0	2.3
2001.....	*	0.1	0.2	0.4	0.4	0.5	0.8	1.1	1.6	2.4
2000.....	0.1	0.1	0.2	0.2	0.3	0.5	0.8	1.4	2.1	2.3
1999.....	0.1	0.2	0.2	0.4	0.5	0.6	0.9	1.5	2.1	3.7

All races 3/, male

2006.....	*	0.2	0.3	0.3	0.6	0.6	0.7	1.2	2.0	2.4
2005.....	*	*	0.3	0.2	0.4	0.5	0.9	1.2	1.4	2.7
2004.....	*	*	0.3	*	0.4	0.6	1.2	0.9	1.7	3.7
2003.....	*	0.3	0.3	0.5	*	0.5	1.0	1.7	2.2	2.5
2002.....	0.2	*	0.3	0.4	0.5	1.0	0.9	1.3	2.8	3.5
2001.....	*	0.2	0.3	0.6	0.7	0.7	1.2	1.5	2.0	3.8
2000.....	*	0.3	0.4	0.3	0.6	0.9	0.9	1.9	3.2	4.1
1999.....	*	0.3	0.4	0.6	0.9	0.9	1.4	2.0	3.3	6.4

All races 3/, female

2006.....	*	*	*	*	*	*	*	*	1.0	1.2
2005.....	*	*	*	*	*	*	*	0.8	0.8	1.5
2004.....	*	*	*	*	*	*	0.5	0.7	0.8	1.5
2003.....	*	*	*	*	*	*	*	0.8	1.3	1.5
2002.....	*	*	*	*	*	*	0.5	1.1	1.5	1.8
2001.....	*	*	*	*	*	*	0.5	0.8	1.4	1.8
2000.....	*	*	*	*	*	*	0.6	1.1	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	1.2	1.4	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Respiratory tuberculosis (A16)

White, both sexes

2006 4/.....	*	*	*	*	0.2	0.3	0.3	0.6	1.0	1.2
2005 4/.....	*	*	*	*	0.2	0.3	0.3	0.6	0.7	1.3
2004 4/.....	*	*	*	*	*	0.3	0.5	0.6	0.9	1.6
2003 4/.....	*	*	*	0.2	*	*	0.5	0.9	1.4	1.4
2002.....	*	*	*	*	0.3	0.4	0.5	0.9	1.5	1.8
2001.....	*	*	*	0.2	0.3	0.4	0.5	0.9	1.2	1.8
2000.....	*	*	0.2	*	*	0.3	0.5	1.0	1.5	1.9
1999.....	*	0.1	0.1	0.2	0.4	0.5	0.6	1.1	1.7	3.0

White, male

2006 4/.....	*	*	*	*	0.4	*	*	0.9	1.5	1.5
2005 4/.....	*	*	*	*	*	*	*	*	*	1.9
2004 4/.....	*	*	*	*	*	*	0.8	0.8	1.2	2.6
2003 4/.....	*	*	*	0.3	*	*	0.9	1.1	1.8	1.8
2002.....	*	*	*	*	*	0.7	0.8	0.9	2.0	2.9
2001.....	*	*	*	0.4	0.5	*	0.7	1.2	1.4	2.6
2000.....	*	*	0.3	*	*	0.5	0.7	1.3	1.9	3.4
1999.....	*	*	*	0.4	0.5	0.7	1.0	1.3	2.6	5.2

White, female

2006 4/.....	*	*	*	*	*	*	*	*	0.7	1.0
2005 4/.....	*	*	*	*	*	*	*	*	0.7	1.1
2004 4/.....	*	*	*	*	*	*	*	*	0.7	1.1
2003 4/.....	*	*	*	*	*	*	*	0.8	1.1	1.2
2002.....	*	*	*	*	*	*	*	0.9	1.1	1.3
2001.....	*	*	*	*	*	*	*	0.6	1.0	1.4
2000.....	*	*	*	*	*	*	*	0.7	1.2	1.2
1999.....	*	*	*	*	*	*	*	0.9	1.1	2.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Septicemia (A40-A41)

All races 3/, both sexes

2006.....	2.4	3.9	6.6	10.3	16.2	25.3	40.3	66.0	103.7	177.3
2005.....	2.3	4.0	6.6	10.5	16.1	25.7	40.8	63.6	104.8	187.3
2004.....	2.3	4.1	6.9	10.3	16.2	24.9	41.2	65.1	103.6	186.7
2003.....	2.6	4.2	6.5	10.3	16.7	25.2	40.9	68.4	107.8	202.5
2002.....	2.3	4.2	6.2	10.0	16.0	27.0	43.1	70.0	109.4	203.0
2001.....	2.3	4.0	6.2	9.8	15.5	25.3	40.9	67.3	103.9	205.9
2000.....	2.5	3.9	6.1	9.4	15.2	24.2	38.4	64.0	105.0	215.7
1999.....	2.2	3.7	5.6	9.0	14.4	24.7	38.3	63.8	102.9	220.7

All races 3/, male

2006.....	2.6	4.2	7.7	11.7	17.4	27.6	43.9	75.3	117.7	194.8
2005.....	2.4	4.4	7.1	11.5	17.8	28.6	45.9	71.7	119.8	201.9
2004.....	2.4	4.5	7.6	11.4	17.1	26.9	46.6	74.5	120.9	203.9
2003.....	2.5	4.6	7.1	11.3	17.7	28.7	45.1	76.7	122.5	213.8
2002.....	2.5	4.7	6.8	10.6	17.8	29.1	48.2	80.6	127.0	213.9
2001.....	2.5	4.3	6.7	10.6	16.6	27.8	46.7	78.6	122.3	224.3
2000.....	2.6	4.4	7.0	10.1	16.7	26.9	43.2	73.9	121.7	223.0
1999.....	2.6	4.2	6.2	9.7	15.7	27.7	43.7	74.8	120.6	233.5

All races 3/, female

2006.....	2.2	3.7	5.6	8.9	15.1	23.2	37.4	59.2	94.9	169.2
2005.....	2.3	3.6	6.1	9.5	14.5	23.2	36.7	57.8	95.4	180.6
2004.....	2.3	3.7	6.4	9.3	15.5	23.2	36.8	58.4	92.9	179.0
2003.....	2.7	3.8	6.0	9.4	15.7	22.2	37.5	62.5	98.8	197.5
2002.....	2.2	3.8	5.6	9.5	14.5	25.2	39.0	62.5	98.7	198.3
2001.....	2.1	3.8	5.7	9.1	14.4	23.1	36.2	59.5	92.9	198.3
2000.....	2.3	3.4	5.2	8.7	13.8	21.8	34.6	57.1	95.2	212.7
1999.....	1.9	3.3	5.1	8.4	13.1	22.1	34.1	56.2	92.5	215.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Septicemia (A40-A41)

White, both sexes

2006 4/.....	2.0	3.2	5.4	8.6	14.1	22.5	36.2	61.9	98.3	171.0
2005 4/.....	1.9	3.3	5.4	9.0	14.0	22.9	37.5	58.9	98.3	177.3
2004 4/.....	1.8	3.2	5.7	8.8	13.8	21.9	37.6	60.7	97.5	178.1
2003 4/.....	2.1	3.4	5.2	8.8	14.3	21.7	36.9	63.7	102.5	191.1
2002.....	1.9	3.6	5.0	8.4	13.5	23.6	39.2	64.4	103.1	191.9
2001.....	1.8	3.2	4.9	8.1	13.2	22.2	36.8	62.1	98.5	195.3
2000.....	2.0	3.0	4.9	7.8	12.6	20.7	34.5	58.0	97.4	203.0
1999.....	1.7	2.8	4.5	7.2	12.1	21.3	33.6	57.6	95.5	205.9

White, male

2006 4/.....	2.2	3.5	6.3	9.7	14.8	24.8	40.0	70.8	112.6	190.7
2005 4/.....	2.0	3.6	5.7	9.7	15.8	25.2	41.8	66.7	113.0	190.7
2004 4/.....	1.8	3.5	6.4	9.7	14.4	23.7	42.5	69.8	113.4	196.5
2003 4/.....	2.0	3.8	5.7	9.3	15.3	24.6	40.8	71.3	116.7	203.3
2002.....	2.0	4.0	5.4	8.9	15.3	25.2	44.0	73.4	119.0	202.9
2001.....	2.1	3.3	5.3	8.5	14.4	24.4	41.6	72.5	116.0	211.8
2000.....	2.2	3.5	5.5	8.4	14.0	23.1	39.4	66.7	114.0	209.8
1999.....	2.0	3.1	4.9	7.6	12.8	23.7	38.2	68.3	111.6	217.4

White, female

2006 4/.....	1.8	2.8	4.6	7.5	13.4	20.4	33.2	55.2	89.3	161.8
2005 4/.....	1.8	2.9	5.1	8.3	12.4	20.8	33.9	53.1	89.2	171.0
2004 4/.....	1.8	2.9	5.1	8.0	13.3	20.2	33.5	54.1	87.7	169.9
2003 4/.....	2.2	3.0	4.7	8.2	13.4	19.2	33.7	58.2	93.7	185.8
2002.....	1.9	3.1	4.5	7.8	11.8	22.3	35.2	58.0	93.4	187.1
2001.....	1.6	3.0	4.5	7.7	12.2	20.4	32.9	54.8	87.9	188.3
2000.....	1.9	2.5	4.3	7.2	11.3	18.7	30.6	51.9	87.6	200.2
1999.....	1.4	2.4	4.1	6.9	11.5	19.2	29.9	50.1	86.1	201.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Septicemia (A40-A41)

Black, both sexes

2006 4/.....	5.7	10.0	16.8	26.0	38.4	55.7	85.8	124.8	186.2	287.5
2005 4/.....	5.5	10.0	16.6	25.5	38.4	55.8	81.2	125.4	203.7	336.8
2004 4/.....	6.2	10.3	17.3	25.5	39.9	57.9	85.1	128.0	196.1	320.3
2003 4/.....	6.3	10.5	17.5	25.3	41.5	61.0	88.0	133.4	193.4	372.0
2002.....	5.5	9.5	17.0	27.2	40.9	62.0	90.6	146.5	208.8	367.0
2001.....	5.6	10.6	16.9	27.1	38.4	57.4	89.6	139.1	193.1	364.8
2000.....	5.8	11.0	16.9	24.3	40.2	59.7	86.6	146.4	216.3	392.6
1999.....	6.2	11.0	15.5	25.4	36.3	57.9	94.6	142.8	209.5	412.3

Black, male

2006 4/.....	6.4	10.1	20.4	31.8	45.4	61.6	94.4	153.0	209.5	296.7
2005 4/.....	5.5	10.9	19.5	30.5	43.4	68.2	97.1	143.5	238.8	374.6
2004 4/.....	6.7	12.0	19.0	30.4	44.5	65.2	102.3	153.8	244.6	341.7
2003 4/.....	6.4	11.8	19.7	31.8	46.1	74.9	100.2	160.2	235.5	404.5
2002.....	6.1	10.3	20.0	29.0	45.1	71.6	105.5	183.7	259.8	393.5
2001.....	5.7	11.6	19.1	33.3	41.5	66.3	111.1	171.0	236.8	424.4
2000.....	5.8	12.4	21.3	27.8	45.1	68.9	97.0	182.9	250.4	427.8
1999.....	7.7	12.8	18.1	29.7	46.5	71.4	115.0	166.2	266.8	453.5

Black, female

2006 4/.....	5.2	9.9	13.7	21.2	32.9	51.4	79.9	107.9	174.0	283.7
2005 4/.....	5.5	9.2	14.1	21.3	34.4	46.5	70.3	114.7	185.1	321.5
2004 4/.....	5.7	8.9	15.8	21.5	36.4	52.5	73.4	112.7	170.5	311.9
2003 4/.....	6.3	9.4	15.6	20.0	37.9	50.8	79.8	117.5	171.4	359.3
2002.....	5.0	8.9	14.4	25.7	37.6	54.9	80.7	124.3	182.2	356.9
2001.....	5.4	9.7	15.0	22.1	36.0	50.9	75.3	120.1	170.9	342.7
2000.....	5.8	9.7	13.2	21.5	36.3	52.8	79.7	124.5	199.1	379.5
1999.....	5.0	9.4	13.3	21.9	28.3	47.9	81.0	128.7	181.1	396.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Viral hepatitis (B15-B19)

All races 3/, both sexes

2006.....	1.8	4.8	8.9	8.0	5.0	4.5	4.5	5.0	4.0	3.1
2005.....	1.7	4.2	6.8	5.3	3.5	3.2	3.8	3.6	3.8	2.4
2004.....	2.0	4.3	6.5	4.9	3.5	3.3	3.6	3.6	3.6	2.7
2003.....	2.1	4.8	6.3	4.5	3.1	3.4	4.3	4.3	4.1	2.4
2002.....	2.3	5.6	6.1	4.3	3.7	3.9	4.5	4.7	4.4	3.9
2001.....	2.6	5.3	5.2	4.1	3.8	4.3	4.9	5.1	4.6	3.8
2000.....	2.5	5.2	4.8	3.7	3.8	4.3	4.9	5.4	5.2	4.0
1999.....	2.3	4.4	3.8	3.3	3.3	4.5	5.1	5.3	5.4	4.7

All races 3/, male

2006.....	2.5	6.8	13.4	11.9	7.0	5.2	4.7	5.1	3.8	3.0
2005.....	2.3	6.3	10.2	8.2	4.7	3.5	4.0	3.3	4.2	2.6
2004.....	2.8	6.4	9.9	7.3	4.5	4.1	3.7	3.8	3.3	3.0
2003.....	2.9	7.1	9.7	6.6	3.9	3.7	4.8	4.7	5.0	2.8
2002.....	3.2	8.5	9.5	6.2	4.6	4.3	4.7	5.7	4.8	5.0
2001.....	3.7	8.1	8.2	5.9	4.5	4.7	4.9	5.1	5.5	4.4
2000.....	3.5	8.0	7.4	5.2	4.7	4.6	5.0	6.5	6.2	5.7
1999.....	3.5	6.7	5.9	4.7	3.8	4.9	5.4	6.1	7.3	6.9

All races 3/, female

2006.....	1.1	2.8	4.6	4.2	3.3	3.8	4.3	5.0	4.1	3.1
2005.....	1.1	2.2	3.6	2.7	2.3	2.9	3.7	3.8	3.6	2.3
2004.....	1.3	2.3	3.3	2.5	2.6	2.6	3.5	3.4	3.7	2.5
2003.....	1.2	2.6	3.1	2.5	2.4	3.1	3.9	4.0	3.5	2.3
2002.....	1.4	2.8	2.8	2.5	2.9	3.4	4.4	4.1	4.2	3.3
2001.....	1.5	2.5	2.4	2.5	3.3	3.9	4.9	5.0	4.1	3.5
2000.....	1.5	2.5	2.4	2.2	3.1	4.1	4.9	4.6	4.7	3.3
1999.....	1.2	2.2	1.7	2.1	2.8	4.1	4.9	4.7	4.2	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Viral hepatitis (B15-B19)

White, both sexes

2006 4/.....	1.8	4.9	8.8	7.2	4.5	3.8	4.1	4.5	3.5	2.8
2005 4/.....	1.7	4.3	6.7	4.9	3.0	2.7	3.2	3.3	3.6	2.2
2004 4/.....	2.1	4.3	6.2	4.2	3.1	3.0	3.2	3.3	3.1	2.5
2003 4/.....	2.1	4.9	6.0	3.8	2.7	2.9	3.8	4.0	3.8	2.3
2002.....	2.2	5.5	5.8	3.8	3.2	3.4	4.1	4.5	4.3	3.7
2001.....	2.6	5.2	4.7	3.7	3.5	3.8	4.4	4.9	4.4	3.6
2000.....	2.4	4.9	4.3	3.0	3.4	3.9	4.6	5.3	5.0	3.9
1999.....	2.2	4.2	3.3	2.9	2.9	4.0	4.6	4.9	5.3	4.5

White, male

2006 4/.....	2.5	7.0	13.3	10.8	6.2	4.4	4.3	4.6	3.6	3.1
2005 4/.....	2.4	6.4	10.0	7.6	4.0	3.1	3.4	3.3	3.9	2.4
2004 4/.....	2.9	6.4	9.4	6.4	4.0	3.7	3.4	3.6	2.8	2.7
2003 4/.....	2.9	7.2	9.2	5.7	3.4	3.3	4.3	4.2	4.8	2.8
2002.....	3.1	8.3	9.1	5.5	3.9	4.0	4.3	5.5	4.8	4.9
2001.....	3.7	8.1	7.6	5.4	4.1	4.4	4.6	5.2	5.6	4.5
2000.....	3.4	7.6	6.6	4.4	4.2	4.0	4.6	6.4	5.9	5.7
1999.....	3.3	6.1	5.2	4.1	3.6	4.6	5.1	5.8	7.7	6.5

White, female

2006 4/.....	1.1	2.8	4.5	3.8	2.9	3.2	3.9	4.3	3.5	2.7
2005 4/.....	1.1	2.2	3.4	2.4	2.0	2.4	3.0	3.2	3.4	2.2
2004 4/.....	1.3	2.3	3.0	2.2	2.4	2.4	3.0	3.1	3.3	2.4
2003 4/.....	1.2	2.6	2.9	2.0	2.0	2.6	3.5	3.8	3.2	2.1
2002.....	1.3	2.6	2.7	2.2	2.6	2.8	3.9	3.8	4.0	3.2
2001.....	1.4	2.4	2.0	2.1	2.9	3.3	4.2	4.6	3.7	3.2
2000.....	1.4	2.3	2.0	1.8	2.7	3.7	4.5	4.5	4.4	3.2
1999.....	1.1	2.2	1.5	1.7	2.3	3.5	4.3	4.3	3.9	3.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Viral hepatitis (B15-B19)

Black, both sexes

2006 4/.....	1.8	4.8	11.0	14.5	9.0	8.9	5.0	7.0	6.9	*
2005 4/.....	1.7	4.7	9.7	9.2	7.2	5.8	6.4	5.5	4.9	*
2004 4/.....	2.0	5.2	10.3	10.2	6.5	4.2	5.1	3.9	6.7	*
2003 4/.....	2.7	5.2	10.1	9.4	6.3	5.1	5.7	6.5	5.3	*
2002.....	2.8	7.2	9.0	7.7	6.3	6.1	6.2	6.0	*	*
2001.....	3.3	6.8	9.4	7.3	6.0	5.1	6.2	6.1	*	*
2000.....	3.1	7.3	8.8	6.9	4.6	5.2	6.3	5.7	7.1	*
1999.....	3.5	7.0	6.7	5.7	4.2	6.6	7.2	6.8	*	6.4

Black, male

2006 4/.....	2.3	6.3	16.2	23.5	13.8	11.2	6.5	9.4	*	*
2005 4/.....	2.2	6.7	14.5	14.3	10.5	6.7	7.9	*	*	*
2004 4/.....	2.2	8.3	16.0	16.6	9.9	6.8	6.5	*	*	*
2003 4/.....	3.6	7.6	16.2	14.9	7.5	5.4	6.9	*	*	*
2002.....	3.8	10.5	14.6	12.4	8.1	8.0	*	*	*	*
2001.....	4.7	10.1	13.9	10.2	7.3	*	*	*	*	*
2000.....	4.2	11.0	13.9	10.4	5.2	6.8	7.4	*	*	*
1999.....	5.1	11.9	11.2	7.6	4.5	5.5	*	*	*	*

Black, female

2006 4/.....	1.4	3.4	6.6	7.1	5.2	7.1	*	5.6	7.6	*
2005 4/.....	*	3.0	5.7	5.0	4.7	5.1	5.4	7.5	*	*
2004 4/.....	1.8	2.5	5.5	4.9	3.9	*	*	*	*	*
2003 4/.....	1.9	3.2	5.0	4.8	5.3	4.9	4.9	5.7	*	*
2002.....	1.9	4.4	4.2	3.9	4.8	4.6	6.3	5.8	*	*
2001.....	2.0	3.9	5.5	4.9	5.0	5.4	6.7	7.2	*	*
2000.....	2.1	4.0	4.4	4.0	4.1	4.0	5.5	*	*	*
1999.....	2.0	2.8	2.8	4.2	4.0	7.5	7.7	6.6	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Human immunodeficiency virus (HIV) disease (B20-B24)

All races 3/, both sexes

2006.....	10.8	10.9	9.3	6.5	4.1	3.1	1.8	1.0	0.6	*
2005.....	11.7	11.3	9.9	6.3	4.0	2.8	1.8	1.0	0.5	*
2004.....	12.3	11.5	9.6	6.3	4.2	3.1	1.5	1.1	0.5	*
2003.....	12.8	12.1	9.5	6.4	4.1	2.8	1.9	1.0	0.4	*
2002.....	13.4	12.6	9.6	5.9	4.0	2.9	1.4	1.0	0.5	*
2001.....	13.7	12.1	8.7	5.8	4.5	2.7	1.4	0.9	0.4	*
2000.....	13.5	12.7	9.0	5.9	4.1	2.6	1.7	0.9	*	*
1999.....	14.6	12.4	8.7	5.9	3.5	2.7	1.5	0.7	*	*

All races 3/, male

2006.....	15.6	16.1	14.4	10.3	6.9	5.0	3.1	1.9	1.1	*
2005.....	17.2	17.2	15.4	10.5	6.6	4.8	3.1	1.8	*	*
2004.....	17.9	17.5	14.9	10.5	7.0	5.1	2.6	1.7	*	*
2003.....	19.1	18.8	15.3	10.7	6.9	4.6	3.3	1.8	1.0	*
2002.....	19.8	19.4	15.7	9.8	6.8	5.0	2.6	1.7	*	*
2001.....	20.5	19.0	14.4	9.8	7.1	4.2	2.7	1.8	*	*
2000.....	20.4	20.3	15.0	10.1	7.0	4.5	2.9	1.8	*	*
1999.....	22.6	20.0	14.5	10.2	5.9	4.8	2.7	1.3	*	*

All races 3/, female

2006.....	5.9	5.7	4.3	2.8	1.5	1.4	0.8	*	*	*
2005.....	6.2	5.6	4.6	2.4	1.6	1.1	0.7	*	*	*
2004.....	6.7	5.7	4.5	2.3	1.5	1.4	0.6	0.6	*	*
2003.....	6.6	5.6	3.9	2.5	1.6	1.3	0.7	*	*	*
2002.....	7.2	5.9	3.6	2.3	1.5	1.0	0.5	*	*	*
2001.....	7.0	5.4	3.3	2.0	2.0	1.4	0.4	*	*	*
2000.....	6.7	5.3	3.4	2.0	1.5	1.0	0.7	*	*	*
1999.....	6.9	5.0	3.1	1.9	1.2	1.0	0.6	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Human immunodeficiency virus (HIV) disease (B20-B24)

White, both sexes

2006 4/.....	6.0	5.6	4.8	3.1	2.2	1.6	0.9	0.5	0.4	*
2005 4/.....	6.7	5.9	5.1	3.2	1.9	1.4	0.9	0.6	*	*
2004 4/.....	6.8	6.1	4.8	3.3	2.0	1.5	0.8	0.6	*	*
2003 4/.....	7.4	6.7	4.9	3.5	2.1	1.3	1.0	0.6	*	*
2002.....	7.3	6.8	4.6	3.3	2.2	1.3	0.6	0.5	*	*
2001.....	7.5	6.4	4.5	3.3	2.0	1.2	0.8	0.5	*	*
2000.....	7.5	6.9	4.7	3.4	2.1	1.3	0.7	0.5	*	*
1999.....	8.1	6.5	4.9	3.1	1.9	1.4	0.7	0.4	*	*

White, male

2006 4/.....	9.8	9.7	8.0	5.4	4.0	2.8	1.6	1.0	*	*
2005 4/.....	11.0	10.0	8.4	5.6	3.4	2.5	1.7	1.1	*	*
2004 4/.....	11.2	10.1	8.2	5.9	3.7	2.7	1.5	1.2	*	*
2003 4/.....	12.1	11.3	8.4	6.0	3.6	2.2	2.0	1.3	*	*
2002.....	12.0	11.4	8.0	5.8	3.8	2.4	1.2	0.8	*	*
2001.....	12.6	10.8	8.0	5.9	3.4	2.1	1.7	1.0	*	*
2000.....	12.3	11.7	8.3	6.0	3.7	2.5	1.3	1.0	*	*
1999.....	13.4	11.0	8.6	5.6	3.5	2.7	1.2	0.8	*	*

White, female

2006 4/.....	2.1	1.6	1.7	0.9	0.5	0.5	*	*	*	*
2005 4/.....	2.4	1.9	1.9	0.9	0.5	*	*	*	*	*
2004 4/.....	2.4	2.1	1.5	0.8	0.4	0.5	*	*	*	*
2003 4/.....	2.6	2.1	1.3	1.1	0.6	0.5	*	*	*	*
2002.....	2.6	2.2	1.3	0.9	0.7	*	*	*	*	*
2001.....	2.4	2.0	1.0	0.9	0.7	0.5	*	*	*	*
2000.....	2.7	2.1	1.2	0.9	0.7	*	*	*	*	*
1999.....	2.8	2.0	1.2	0.7	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Human immunodeficiency virus (HIV) disease (B20-B24)

Black, both sexes

2006 4/.....	44.6	49.5	44.4	35.8	22.1	16.7	10.8	5.9	*	*
2005 4/.....	47.2	51.2	47.8	33.8	24.0	16.6	10.5	6.0	*	*
2004 4/.....	51.4	52.1	47.4	33.4	24.6	17.6	8.5	6.3	*	*
2003 4/.....	52.3	53.4	47.2	33.9	23.8	17.2	11.0	4.9	*	*
2002.....	57.3	57.4	49.7	30.2	20.7	17.5	10.5	7.0	*	*
2001.....	58.8	56.5	44.7	28.9	26.2	16.3	8.5	6.4	*	*
2000.....	57.3	58.7	46.6	29.3	22.0	15.1	11.6	6.2	*	*
1999.....	62.8	59.0	42.6	32.7	18.3	15.4	11.2	4.8	*	*

Black, male

2006 4/.....	60.6	68.2	69.1	57.1	37.3	27.2	19.4	12.5	*	*
2005 4/.....	65.8	74.8	75.4	57.8	40.7	29.7	18.7	11.6	*	*
2004 4/.....	69.9	76.9	72.4	56.1	42.0	29.6	15.2	9.6	*	*
2003 4/.....	74.0	79.5	75.7	57.6	41.3	30.0	20.0	*	*	*
2002.....	79.6	86.3	83.5	50.0	37.0	31.4	19.4	14.7	*	*
2001.....	82.8	87.5	72.2	49.9	43.5	26.4	15.5	13.0	*	*
2000.....	84.1	92.7	77.4	51.0	39.5	26.3	22.4	11.8	*	*
1999.....	94.9	96.1	70.6	57.5	30.6	27.3	21.4	*	*	*

Black, female

2006 4/.....	30.5	33.2	23.5	18.2	10.3	9.0	4.9	*	*	*
2005 4/.....	30.8	30.6	24.6	14.0	10.9	6.9	5.0	*	*	*
2004 4/.....	35.0	30.5	26.3	14.7	10.9	8.6	*	*	*	*
2003 4/.....	33.1	30.7	23.1	14.3	10.2	7.7	4.9	*	*	*
2002.....	37.6	32.4	21.0	13.9	7.9	7.2	4.5	*	*	*
2001.....	37.5	29.7	21.2	11.7	12.6	8.8	*	*	*	*
2000.....	33.7	29.2	20.4	11.5	8.2	6.9	4.4	*	*	*
1999.....	34.6	26.9	18.8	12.5	8.7	6.5	4.5	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races 3/, both sexes

2006.....	0.8	1.0	1.6	2.3	3.3	4.8	7.1	9.8	13.8	20.4
2005.....	1.1	2.4	4.2	3.9	4.5	5.9	8.3	12.0	15.8	21.2
2004.....	1.2	2.5	3.5	4.2	4.4	6.0	8.3	11.2	15.9	22.5
2003.....	1.3	2.4	3.5	3.7	4.5	5.9	8.9	12.3	14.8	22.5
2002.....	1.3	1.9	2.7	3.0	3.8	5.4	8.1	11.6	13.8	21.7
2001.....	0.9	1.5	1.9	2.4	3.9	5.6	6.9	9.4	12.3	20.1
2000.....	0.8	1.4	1.8	2.4	3.6	4.7	7.0	10.0	12.9	21.2
1999.....	1.4	2.0	2.6	3.3	4.4	6.8	8.9	12.6	16.6	27.0

All races 3/, male

2006.....	1.0	1.2	2.1	2.8	3.8	5.6	8.1	11.6	17.2	25.1
2005.....	1.5	3.3	5.8	5.2	5.3	7.0	10.0	14.1	19.6	23.6
2004.....	1.5	3.3	4.9	5.6	5.2	7.2	9.9	12.9	18.4	29.6
2003.....	1.6	3.3	4.7	4.9	5.4	7.0	10.3	13.9	18.8	26.5
2002.....	1.7	2.6	3.8	3.8	4.7	6.5	9.7	14.0	17.7	26.8
2001.....	1.2	2.1	2.4	3.0	4.5	6.2	7.5	11.4	15.9	23.6
2000.....	1.0	1.7	2.3	2.8	4.0	5.7	8.8	12.1	16.2	27.6
1999.....	1.7	2.4	3.0	3.5	5.2	8.0	10.5	15.3	20.4	31.9

All races 3/, female

2006.....	0.6	0.8	1.2	1.8	2.9	4.2	6.3	8.4	11.7	18.2
2005.....	0.7	1.5	2.5	2.7	3.8	5.0	7.0	10.5	13.4	20.1
2004.....	0.9	1.6	2.2	2.8	3.6	5.0	7.0	10.0	14.4	19.3
2003.....	0.9	1.6	2.2	2.6	3.7	5.0	7.8	11.1	12.4	20.8
2002.....	0.8	1.2	1.7	2.2	3.0	4.5	6.8	9.9	11.5	19.5
2001.....	0.6	0.9	1.5	1.8	3.3	5.1	6.5	8.0	10.1	18.7
2000.....	0.6	1.1	1.3	2.1	3.2	3.9	5.6	8.5	10.9	18.6
1999.....	1.1	1.6	2.3	3.1	3.7	5.9	7.7	10.8	14.4	25.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

White, both sexes

2006 4/.....	0.7	0.9	1.5	2.1	3.2	4.7	7.1	9.8	13.7	20.6
2005 4/.....	1.1	2.2	4.1	3.8	4.3	5.9	8.0	11.7	15.6	21.2
2004 4/.....	1.1	2.5	3.5	4.0	4.1	5.7	8.0	10.9	15.7	22.5
2003 4/.....	1.2	2.3	3.3	3.5	4.2	5.9	8.9	11.8	14.7	23.0
2002.....	1.1	1.8	2.5	2.6	3.6	5.4	7.9	11.3	13.8	22.0
2001.....	0.8	1.4	1.7	2.2	3.6	5.4	6.7	9.3	12.1	20.1
2000.....	0.7	1.3	1.6	2.3	3.2	4.6	6.8	9.8	12.7	21.0
1999.....	1.1	1.6	2.2	2.9	3.9	6.4	8.5	12.1	16.1	26.5

White, male

2006 4/.....	0.8	1.1	1.8	2.5	3.5	5.6	8.0	11.4	17.1	25.7
2005 4/.....	1.4	3.1	5.9	5.0	5.0	7.0	9.8	13.8	19.4	23.7
2004 4/.....	1.4	3.3	4.9	5.2	4.9	7.1	9.4	12.6	18.1	30.1
2003 4/.....	1.6	3.2	4.5	4.6	5.0	6.8	10.0	13.5	18.6	27.2
2002.....	1.5	2.5	3.5	3.2	4.3	6.6	9.4	13.6	17.6	27.2
2001.....	1.0	1.9	2.2	2.8	4.3	6.0	7.2	11.3	15.9	23.8
2000.....	0.9	1.5	2.0	2.7	3.6	5.5	8.6	11.6	16.1	26.9
1999.....	1.4	2.0	2.6	3.3	4.5	7.5	10.2	14.8	18.6	30.6

White, female

2006 4/.....	0.5	0.7	1.1	1.8	2.9	4.0	6.3	8.6	11.6	18.1
2005 4/.....	0.7	1.4	2.4	2.7	3.6	5.0	6.6	10.2	13.3	20.0
2004 4/.....	0.9	1.6	2.2	2.8	3.4	4.6	6.9	9.7	14.1	19.0
2003 4/.....	0.8	1.5	2.1	2.4	3.5	5.0	8.0	10.6	12.3	21.2
2002.....	0.7	1.1	1.5	2.0	2.9	4.3	6.7	9.7	11.5	19.8
2001.....	0.6	0.8	1.3	1.6	3.1	5.0	6.2	7.8	9.9	18.5
2000.....	0.6	1.0	1.1	1.9	2.8	3.8	5.4	8.6	10.8	18.7
1999.....	0.7	1.2	1.9	2.6	3.4	5.6	7.2	10.2	14.6	24.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

Black, both sexes

2006 4/.....	1.5	2.0	3.2	4.1	4.6	6.5	8.1	10.7	14.7	22.1
2005 4/.....	1.7	3.4	5.0	5.5	6.6	6.3	9.6	13.4	17.2	22.7
2004 4/.....	2.1	2.9	3.9	5.8	6.7	7.5	10.2	14.7	19.1	24.3
2003 4/.....	2.1	3.6	5.3	6.0	6.2	5.9	8.3	16.2	16.2	19.5
2002.....	2.5	3.2	4.8	6.0	5.4	5.3	9.7	15.3	11.5	21.7
2001.....	1.8	2.5	3.4	4.4	6.0	7.1	9.8	9.7	13.3	20.6
2000.....	1.7	2.6	3.5	4.0	5.9	6.1	9.9	11.6	16.1	24.1
1999.....	3.6	5.1	5.9	6.3	8.2	10.7	13.1	18.7	23.4	34.1

Black, male

2006 4/.....	2.2	2.4	4.7	5.6	6.4	6.1	9.8	15.2	17.1	20.6
2005 4/.....	2.5	4.3	6.1	7.8	8.4	8.3	10.8	18.8	20.4	23.2
2004 4/.....	2.4	4.1	5.9	8.7	9.2	7.5	14.6	18.7	21.5	29.8
2003 4/.....	2.6	4.6	7.1	7.7	8.3	8.9	13.1	16.4	22.1	22.9
2002.....	3.2	4.5	6.7	9.2	7.3	6.5	12.0	18.4	*	28.9
2001.....	2.7	3.6	4.3	5.5	7.3	9.7	10.4	10.2	16.1	*
2000.....	2.1	3.2	4.8	4.4	7.2	8.6	12.4	17.0	20.2	32.3
1999.....	3.6	5.8	6.5	5.8	10.4	12.5	15.9	21.8	41.9	45.7

Black, female

2006 4/.....	*	1.7	1.9	2.8	3.3	6.8	7.0	8.0	13.4	22.7
2005 4/.....	*	2.6	4.0	3.5	5.2	4.8	8.9	10.1	15.6	22.5
2004 4/.....	1.8	1.9	2.3	3.4	4.8	7.6	7.2	12.4	17.8	22.1
2003 4/.....	1.6	2.6	3.8	4.6	4.5	3.7	5.1	16.0	13.2	18.2
2002.....	1.9	2.0	3.1	3.4	3.9	4.4	8.1	13.4	9.9	18.9
2001.....	*	1.6	2.7	3.6	5.0	5.3	9.4	9.4	11.9	20.2
2000.....	*	2.1	2.4	3.6	4.9	4.2	8.2	8.3	14.0	21.0
1999.....	3.5	4.5	5.3	6.8	6.5	9.4	11.3	16.9	14.3	29.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

Malignant neoplasms (C00-C97)

All races 3/, both sexes

2006.....	42.3	84.1	152.1	252.7	414.6	615.1	863.3	1,144.9	1,418.8	1,606.1
2005.....	44.6	86.5	154.8	255.6	422.1	622.2	886.3	1,158.4	1,427.7	1,637.7
2004.....	44.3	86.7	155.6	261.5	427.6	637.8	892.4	1,172.1	1,424.8	1,653.3
2003.....	46.1	89.8	159.1	267.9	440.9	647.9	909.2	1,195.4	1,450.0	1,698.2
2002.....	47.5	90.6	161.5	276.8	447.0	663.8	933.6	1,207.0	1,458.3	1,723.9
2001.....	48.7	92.4	164.8	280.7	453.3	673.2	943.5	1,211.7	1,465.9	1,765.6
2000.....	48.8	93.6	166.2	289.0	463.4	690.6	951.6	1,222.5	1,505.3	1,819.4
1999.....	49.4	93.7	167.0	295.3	471.6	700.2	963.4	1,224.0	1,493.1	1,805.8

All races 3/, male

2006.....	37.1	80.7	162.1	282.3	476.3	731.6	1,045.7	1,442.3	1,901.9	2,248.7
2005.....	39.5	83.4	164.9	284.8	484.5	743.6	1,091.8	1,475.1	1,901.6	2,319.3
2004.....	39.2	86.1	168.2	289.9	492.5	756.9	1,091.6	1,503.3	1,894.3	2,349.5
2003.....	41.7	90.6	169.8	295.9	507.6	774.4	1,117.9	1,533.3	1,939.7	2,413.8
2002.....	42.8	89.8	171.7	308.6	519.7	796.8	1,157.8	1,554.1	1,951.5	2,491.1
2001.....	44.0	91.7	174.2	311.6	526.7	814.1	1,178.0	1,568.4	1,978.7	2,613.6
2000.....	44.4	93.5	173.9	321.3	535.7	834.7	1,190.4	1,586.0	2,050.4	2,710.7
1999.....	45.5	94.7	174.9	330.6	550.1	848.2	1,212.9	1,600.1	2,046.0	2,722.2

All races 3/, female

2006.....	47.6	87.5	142.6	224.8	358.2	513.3	714.8	927.2	1,116.9	1,305.5
2005.....	49.6	89.6	145.1	228.1	365.2	516.2	719.8	928.5	1,133.4	1,324.6
2004.....	49.2	87.4	143.5	234.8	368.5	534.2	732.0	934.2	1,135.5	1,340.1
2003.....	50.5	89.1	148.8	241.5	380.3	538.1	741.8	954.8	1,150.5	1,381.9
2002.....	52.2	91.4	151.8	246.8	381.1	548.9	754.4	961.9	1,158.4	1,391.1
2001.....	53.5	93.2	155.9	251.7	386.7	551.9	757.2	962.0	1,158.6	1,410.7
2000.....	53.2	93.7	158.8	258.9	398.0	567.2	763.5	969.3	1,183.8	1,456.4
1999.....	53.3	92.7	159.5	262.3	400.6	573.7	768.4	962.5	1,169.5	1,436.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms (C00-C97)

White, both sexes

2006 4/.....	40.6	80.2	146.0	245.8	408.2	612.3	866.4	1,151.9	1,427.2	1,618.1
2005 4/.....	43.0	82.0	147.8	247.2	414.2	618.5	888.9	1,164.9	1,432.7	1,645.4
2004 4/.....	42.0	82.2	148.2	253.8	421.6	633.6	891.5	1,174.1	1,429.3	1,659.7
2003 4/.....	44.0	84.8	151.3	261.2	434.5	644.8	908.5	1,195.1	1,450.7	1,701.4
2002.....	45.5	86.2	154.7	268.7	439.5	659.2	931.4	1,205.1	1,456.9	1,725.7
2001.....	46.4	87.8	158.4	272.2	447.1	665.6	938.4	1,207.5	1,458.8	1,765.2
2000.....	45.8	88.2	159.5	281.1	457.1	685.0	944.1	1,214.2	1,503.0	1,815.7
1999.....	46.8	88.5	159.9	285.8	462.2	693.1	955.3	1,213.0	1,488.4	1,798.8

White, male

2006 4/.....	35.8	77.2	154.0	271.5	464.1	721.0	1,038.6	1,439.4	1,903.8	2,258.3
2005 4/.....	38.7	79.2	156.6	271.7	470.2	731.9	1,083.4	1,472.2	1,894.9	2,322.7
2004 4/.....	37.4	81.1	158.7	278.1	479.7	745.4	1,077.9	1,490.7	1,887.6	2,348.9
2003 4/.....	39.8	85.1	159.7	285.4	494.7	763.2	1,108.1	1,518.9	1,930.1	2,412.1
2002.....	41.2	85.4	162.8	296.5	503.4	785.6	1,145.2	1,537.3	1,934.6	2,486.8
2001.....	41.9	86.5	165.1	298.2	513.9	797.6	1,160.4	1,547.8	1,950.7	2,597.6
2000.....	41.4	87.7	164.0	307.0	522.0	819.7	1,166.6	1,559.2	2,026.9	2,693.7
1999.....	43.1	88.0	164.3	316.0	532.7	831.3	1,191.9	1,571.1	2,024.1	2,696.3

White, female

2006 4/.....	45.4	83.3	138.1	220.9	356.0	515.5	723.4	937.3	1,126.1	1,317.5
2005 4/.....	47.3	84.8	139.2	223.6	362.1	517.7	728.2	937.5	1,143.0	1,333.6
2004 4/.....	46.6	83.4	137.9	230.5	367.7	534.4	738.3	942.5	1,144.7	1,348.9
2003 4/.....	48.3	84.6	143.1	237.8	378.9	540.3	745.2	961.0	1,155.2	1,386.5
2002.....	49.9	86.9	146.8	242.0	380.4	548.0	757.1	967.4	1,164.2	1,395.1
2001.....	51.0	89.0	152.0	247.4	385.4	549.7	758.6	966.4	1,161.4	1,416.7
2000.....	50.1	88.7	155.2	256.5	397.2	567.2	765.6	971.5	1,191.3	1,459.1
1999.....	50.5	89.1	155.5	257.1	397.2	572.5	767.3	961.5	1,172.1	1,438.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms (C00-C97)

Black, both sexes

2006 4/.....	59.3	123.7	224.1	358.6	561.4	771.6	1,004.1	1,283.1	1,557.0	1,678.3
2005 4/.....	60.6	129.1	231.4	374.3	573.3	782.6	1,026.2	1,296.6	1,585.5	1,744.7
2004 4/.....	64.2	129.6	235.7	377.1	570.2	803.0	1,059.1	1,355.6	1,578.2	1,774.0
2003 4/.....	66.5	138.3	243.2	382.6	589.1	802.7	1,080.1	1,385.5	1,640.8	1,853.1
2002.....	68.3	135.1	238.2	399.6	597.1	833.2	1,126.1	1,425.8	1,665.1	1,885.7
2001.....	70.4	137.8	241.5	404.7	589.4	858.4	1,156.8	1,433.0	1,716.9	1,931.7
2000.....	74.4	145.6	251.9	407.9	604.6	861.4	1,179.2	1,481.7	1,700.6	1,998.9
1999.....	73.3	144.5	255.6	429.5	633.9	879.1	1,191.7	1,502.1	1,704.2	2,008.4

Black, male

2006 4/.....	50.6	120.6	256.5	438.9	703.1	1,007.6	1,351.8	1,783.7	2,278.2	2,543.3
2005 4/.....	50.0	127.0	258.6	459.7	729.5	1,024.8	1,395.3	1,803.3	2,354.0	2,683.7
2004 4/.....	56.2	134.0	274.7	455.9	729.5	1,035.5	1,442.5	1,926.3	2,340.8	2,748.8
2003 4/.....	61.2	145.6	283.0	464.8	747.3	1,045.9	1,455.8	1,975.5	2,417.1	2,825.5
2002.....	60.4	137.1	271.5	484.4	777.8	1,079.3	1,536.2	2,047.9	2,514.3	2,976.1
2001.....	62.8	143.6	280.7	494.9	746.2	1,135.3	1,600.3	2,084.3	2,645.3	3,209.9
2000.....	70.3	151.8	293.7	512.8	769.2	1,137.3	1,654.9	2,166.0	2,683.2	3,264.8
1999.....	68.3	160.5	300.8	533.7	815.7	1,159.8	1,644.1	2,205.4	2,642.0	3,340.7

Black, female

2006 4/.....	67.0	126.3	196.7	292.4	450.9	597.3	765.2	983.5	1,175.9	1,324.4
2005 4/.....	70.0	131.0	208.4	303.9	451.6	603.0	773.5	994.4	1,179.6	1,365.8
2004 4/.....	71.2	125.9	202.6	312.3	446.1	630.3	798.9	1,017.5	1,176.3	1,387.7
2003 4/.....	71.1	132.0	209.5	314.9	465.7	622.1	826.8	1,034.8	1,234.0	1,475.3
2002.....	75.2	133.5	210.0	330.0	455.6	650.5	850.9	1,054.1	1,223.1	1,468.0
2001.....	77.1	132.8	208.1	330.6	466.1	653.9	861.4	1,044.2	1,244.0	1,457.5
2000.....	78.1	140.2	216.2	321.9	474.8	657.8	863.3	1,069.2	1,205.8	1,527.7
1999.....	77.7	130.6	217.2	344.4	491.4	671.1	891.6	1,078.5	1,239.4	1,510.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races 3/, both sexes

2006.....	0.9	1.8	3.5	5.2	6.6	8.7	10.6	13.0	14.1	16.3
2005.....	0.8	1.9	3.5	5.6	7.5	9.2	10.6	11.8	14.2	17.0
2004.....	0.9	2.0	3.8	5.1	7.5	9.4	11.0	12.7	14.9	18.0
2003.....	0.9	2.0	3.6	5.5	7.9	9.7	10.8	12.8	13.6	18.6
2002.....	0.8	2.0	3.6	5.8	8.1	8.8	12.1	13.1	14.1	18.7
2001.....	0.8	2.1	3.5	6.1	7.9	9.4	11.6	13.5	15.2	18.5
2000.....	0.9	2.0	3.8	5.7	8.1	9.9	11.5	12.6	14.0	19.4
1999.....	0.9	2.3	3.6	5.8	8.1	9.3	11.3	13.5	15.4	19.3

All races 3/, male

2006.....	1.2	2.9	5.6	8.4	10.6	13.9	16.0	19.7	21.4	21.9
2005.....	1.2	2.8	5.7	9.1	11.9	14.8	16.0	17.9	20.4	22.0
2004.....	1.2	3.2	6.2	8.6	11.9	14.9	16.4	19.3	21.7	24.9
2003.....	1.4	3.2	6.2	9.1	12.6	15.0	16.4	19.0	19.9	26.9
2002.....	1.2	3.2	5.8	9.4	12.9	13.9	18.1	19.2	20.3	26.9
2001.....	1.2	3.2	5.8	10.0	13.0	14.3	17.9	19.6	22.1	25.8
2000.....	1.4	3.1	6.0	9.5	12.7	15.2	16.9	18.7	20.2	27.1
1999.....	1.3	3.6	5.7	9.0	13.1	14.3	17.4	20.1	22.3	27.1

All races 3/, female

2006.....	0.5	0.8	1.4	2.1	2.9	4.2	6.3	8.1	9.6	13.7
2005.....	0.4	1.0	1.4	2.3	3.4	4.3	6.2	7.3	10.4	14.7
2004.....	0.5	0.9	1.5	1.7	3.6	4.7	6.7	8.0	10.8	14.9
2003.....	0.5	0.9	1.1	2.1	3.6	5.2	6.3	8.4	9.7	15.0
2002.....	0.4	0.9	1.4	2.5	3.7	4.4	7.3	8.8	10.3	15.1
2001.....	0.4	1.0	1.4	2.4	3.3	5.1	6.6	9.3	11.1	15.4
2000.....	0.4	1.0	1.6	2.2	3.9	5.4	7.2	8.3	10.4	16.3
1999.....	0.4	1.0	1.6	2.7	3.6	5.1	6.6	9.0	11.3	16.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

White, both sexes

2006 4/.....	0.8	1.7	3.2	4.8	6.2	8.3	10.4	13.0	14.3	16.9
2005 4/.....	0.7	1.7	3.1	5.1	7.1	8.7	10.5	11.7	14.2	17.3
2004 4/.....	0.7	1.7	3.4	4.6	7.0	9.0	11.0	12.7	15.0	18.6
2003 4/.....	0.8	1.8	3.2	5.2	7.3	9.3	10.4	12.6	13.4	19.0
2002.....	0.7	1.7	3.1	5.3	7.5	8.4	12.3	13.3	14.2	18.9
2001.....	0.7	1.7	3.2	5.5	7.3	8.7	11.4	13.6	15.5	19.0
2000.....	0.7	1.7	3.2	5.0	7.3	9.4	11.1	12.5	14.0	19.7
1999.....	0.7	1.8	3.0	5.0	7.5	8.9	11.0	13.2	15.5	19.8

White, male

2006 4/.....	1.1	2.6	5.2	7.8	9.8	12.9	15.3	19.5	21.3	22.4
2005 4/.....	1.0	2.4	5.1	8.1	11.2	13.8	15.3	17.3	20.0	22.1
2004 4/.....	0.9	2.6	5.4	7.8	10.9	14.0	15.8	18.8	21.5	24.8
2003 4/.....	1.3	2.7	5.5	8.5	11.3	14.1	15.2	18.2	19.2	27.2
2002.....	1.0	2.7	5.0	8.6	11.7	12.9	18.0	19.4	20.2	27.0
2001.....	1.0	2.6	5.2	8.9	11.8	13.2	17.1	19.2	22.3	26.4
2000.....	1.1	2.7	4.9	8.3	11.2	14.0	15.9	18.0	19.9	27.6
1999.....	1.1	2.7	4.7	7.6	12.0	13.5	16.8	19.3	22.0	27.3

White, female

2006 4/.....	0.5	0.7	1.3	2.0	2.8	4.2	6.3	8.2	9.9	14.3
2005 4/.....	0.5	0.9	1.3	2.2	3.3	4.2	6.5	7.6	10.6	15.1
2004 4/.....	0.4	0.8	1.4	1.5	3.4	4.6	7.1	8.2	10.9	15.7
2003 4/.....	0.3	0.9	0.9	2.1	3.7	5.1	6.4	8.5	9.8	15.4
2002.....	0.4	0.7	1.3	2.2	3.6	4.3	7.6	8.8	10.5	15.3
2001.....	0.3	0.8	1.2	2.2	3.2	4.8	6.7	9.6	11.3	15.9
2000.....	0.4	0.8	1.6	1.9	3.6	5.3	7.2	8.6	10.6	16.5
1999.....	0.4	0.8	1.3	2.5	3.3	4.9	6.4	9.0	11.7	16.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Black, both sexes

2006 4/.....	1.2	3.2	5.6	8.4	11.1	13.4	14.3	14.2	12.5	10.4
2005 4/.....	1.1	3.6	6.6	9.2	12.6	14.1	13.7	13.0	14.3	12.6
2004 4/.....	1.9	4.6	7.8	9.5	13.3	13.8	13.7	14.9	15.8	12.4
2003 4/.....	1.7	3.9	6.5	8.8	14.4	15.5	15.3	14.8	16.0	15.5
2002.....	1.5	4.2	7.1	10.1	14.7	13.3	12.5	11.3	12.6	16.0
2001.....	1.7	4.3	6.2	12.2	13.1	15.3	15.7	14.2	13.3	12.9
2000.....	1.6	4.1	8.3	12.2	15.2	14.9	16.1	14.6	15.8	15.3
1999.....	1.5	5.9	8.8	12.6	14.6	14.4	16.2	17.7	15.6	14.3

Black, male

2006 4/.....	1.7	5.1	9.1	15.0	19.8	25.5	24.9	25.1	25.3	*
2005 4/.....	2.1	6.1	11.7	17.5	22.4	26.6	25.9	26.8	28.8	21.4
2004 4/.....	2.9	7.6	13.8	17.0	23.7	26.2	27.2	25.9	27.3	26.8
2003 4/.....	1.9	7.0	11.4	16.3	27.8	27.6	30.2	26.5	30.3	25.0
2002.....	2.4	6.7	12.9	18.2	26.9	24.2	21.4	16.5	21.4	26.8
2001.....	2.4	7.0	10.8	22.2	24.8	25.4	29.3	25.1	20.9	*
2000.....	2.7	6.3	15.9	21.9	27.3	27.6	29.1	28.8	28.7	*
1999.....	2.3	10.3	14.5	21.5	25.2	24.1	26.5	33.2	28.8	24.6

Black, female

2006 4/.....	*	1.5	2.6	3.0	4.3	4.5	7.0	7.8	*	8.8
2005 4/.....	*	1.5	2.2	2.4	5.0	4.8	5.4	*	*	9.0
2004 4/.....	*	1.9	2.7	3.4	5.2	4.7	4.6	8.4	9.8	*
2003 4/.....	1.5	*	2.3	2.7	3.9	6.6	5.3	7.9	8.5	11.7
2002.....	*	1.9	2.2	3.4	5.1	5.2	6.5	8.2	8.0	11.9
2001.....	*	2.0	2.2	4.0	3.8	7.9	6.7	7.8	9.4	10.5
2000.....	*	2.1	*	4.2	5.6	5.6	7.5	6.0	9.3	14.2
1999.....	*	2.1	3.9	5.3	6.3	7.1	9.5	8.3	9.1	10.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of esophagus (C15)

All races 3/, both sexes

2006.....	0.9	2.3	4.8	8.4	13.2	17.4	22.0	26.9	27.0	25.6
2005.....	1.1	2.3	4.9	8.4	12.6	17.5	22.4	27.6	27.3	24.9
2004.....	1.0	2.3	4.7	8.1	13.2	17.5	21.3	26.0	27.7	25.8
2003.....	0.9	2.4	4.6	8.8	12.8	17.6	22.2	25.4	27.6	25.0
2002.....	1.0	2.4	4.7	8.6	12.7	18.1	22.5	25.7	27.8	24.6
2001.....	1.1	2.4	5.0	7.9	13.0	18.5	22.9	26.1	26.5	25.8
2000.....	1.0	2.4	4.9	8.8	13.2	17.9	22.1	24.5	27.2	25.9
1999.....	1.0	2.4	4.7	8.7	13.0	18.2	22.0	24.3	24.6	26.8

All races 3/, male

2006.....	1.5	4.1	8.1	15.0	23.2	30.8	38.7	47.0	51.3	48.5
2005.....	1.7	3.9	8.8	14.8	22.2	31.1	39.6	49.6	52.4	46.0
2004.....	1.6	4.0	8.1	14.1	22.9	30.5	37.0	46.8	51.7	48.5
2003.....	1.5	4.1	8.1	15.3	22.7	30.6	38.3	45.7	51.0	45.3
2002.....	1.6	4.1	8.3	15.1	22.1	31.4	40.3	46.8	50.6	43.4
2001.....	1.8	4.1	8.7	13.7	22.6	32.6	39.9	46.5	48.8	48.1
2000.....	1.8	4.2	8.7	15.2	23.0	30.7	38.0	43.9	48.8	47.8
1999.....	1.7	4.2	8.3	15.3	21.9	31.7	39.3	44.1	44.8	48.0

All races 3/, female

2006.....	0.2	0.7	1.7	2.2	4.1	5.8	8.4	12.2	11.8	14.9
2005.....	0.4	0.7	1.2	2.5	3.9	5.6	8.5	11.6	11.6	15.3
2004.....	0.3	0.7	1.4	2.4	4.3	6.1	8.7	11.1	12.9	15.7
2003.....	0.3	0.7	1.3	2.6	3.8	6.3	9.2	11.0	13.3	16.1
2002.....	0.4	0.7	1.3	2.4	4.2	6.7	8.2	10.9	14.0	16.5
2001.....	0.3	0.7	1.4	2.4	4.2	6.4	9.4	11.9	13.2	16.5
2000.....	0.2	0.6	1.4	2.8	4.4	7.1	9.6	11.0	14.5	17.1
1999.....	0.3	0.6	1.3	2.5	5.0	6.7	8.5	10.5	12.8	18.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of esophagus (C15)

White, both sexes

2006 4/.....	0.9	2.3	4.7	8.3	13.3	17.5	22.5	27.4	27.9	26.4
2005 4/.....	1.1	2.3	4.9	8.3	12.5	17.6	22.9	28.0	28.0	25.9
2004 4/.....	1.0	2.2	4.6	7.9	12.8	17.4	21.2	26.5	28.0	26.2
2003 4/.....	0.9	2.3	4.3	8.5	12.3	17.0	22.5	25.8	27.8	25.8
2002.....	0.9	2.3	4.6	8.1	12.2	17.5	22.4	25.7	28.2	25.1
2001.....	1.0	2.3	4.6	7.1	12.3	17.7	22.1	26.2	26.9	26.3
2000.....	0.9	2.3	4.5	8.1	12.4	17.4	21.8	24.0	27.2	26.6
1999.....	0.9	2.2	4.3	8.0	12.1	17.5	21.5	23.7	24.9	27.2

White, male

2006 4/.....	1.7	4.1	8.0	15.0	23.4	31.1	39.3	47.9	53.7	49.9
2005 4/.....	1.8	3.9	9.0	14.7	22.0	31.2	40.4	50.6	53.7	48.5
2004 4/.....	1.7	3.9	8.0	13.9	22.5	30.5	37.2	47.6	52.8	49.2
2003 4/.....	1.6	3.9	7.6	15.0	22.0	29.5	39.0	46.5	52.3	46.7
2002.....	1.6	4.0	8.1	14.5	21.5	30.7	40.2	47.2	51.3	44.6
2001.....	1.8	3.9	8.1	12.5	21.6	31.2	39.0	46.9	49.5	48.8
2000.....	1.8	4.1	8.0	14.1	22.0	30.0	37.5	43.2	49.7	49.3
1999.....	1.6	4.0	7.6	14.1	20.5	30.8	38.4	43.0	45.6	48.0

White, female

2006 4/.....	0.2	0.6	1.5	1.8	3.8	5.5	8.5	12.1	11.6	15.3
2005 4/.....	0.3	0.6	1.0	2.2	3.6	5.5	8.4	11.3	11.8	15.5
2004 4/.....	0.3	0.5	1.1	2.1	3.8	5.9	8.1	11.1	12.6	15.9
2003 4/.....	0.3	0.6	1.0	2.3	3.4	6.0	9.0	10.9	12.7	16.5
2002.....	0.3	0.6	1.2	2.1	3.7	5.9	7.8	10.3	14.1	16.7
2001.....	0.3	0.6	1.2	1.9	3.7	5.8	8.5	11.6	13.2	16.8
2000.....	*	0.4	1.1	2.4	3.6	6.4	9.1	10.6	13.8	17.3
1999.....	*	0.5	1.0	2.1	4.4	5.8	8.0	10.1	12.6	18.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of esophagus (C15)

Black, both sexes

2006 4/.....	0.8	3.0	7.4	11.5	16.9	21.5	22.8	25.1	23.7	18.2
2005 4/.....	1.3	3.2	6.8	11.8	17.4	21.4	23.9	30.2	22.6	17.4
2004 4/.....	1.2	4.0	6.6	12.0	20.4	23.0	26.8	27.4	27.0	21.4
2003 4/.....	1.3	4.1	8.4	14.1	20.7	26.1	24.9	26.9	29.4	21.3
2002.....	1.6	3.6	6.9	14.9	20.4	27.5	29.8	31.8	26.2	20.8
2001.....	1.6	3.7	8.6	15.9	21.9	29.6	35.8	30.7	27.6	23.6
2000.....	1.5	4.1	9.8	16.9	23.3	27.0	29.6	36.1	29.9	22.2
1999.....	1.7	4.3	9.4	17.1	23.5	28.7	31.3	35.7	23.7	23.9

Black, male

2006 4/.....	*	4.8	12.2	18.7	28.8	38.2	42.1	44.3	39.0	34.9
2005 4/.....	2.1	5.1	11.4	19.8	31.3	39.7	41.4	52.7	48.5	27.0
2004 4/.....	1.7	6.3	10.8	19.9	33.4	40.0	43.1	49.6	45.2	38.7
2003 4/.....	2.1	7.1	14.0	23.3	36.2	46.5	41.4	47.5	43.6	38.6
2002.....	2.5	5.7	11.5	26.8	34.4	45.4	54.4	54.2	48.1	37.5
2001.....	2.6	6.0	15.3	27.5	38.8	51.8	59.9	55.3	53.8	46.5
2000.....	2.4	6.3	17.4	30.5	38.5	44.8	49.8	65.7	45.5	38.1
1999.....	3.2	7.4	16.4	30.3	40.3	46.5	56.0	67.9	36.6	51.6

Black, female

2006 4/.....	*	1.4	3.3	5.6	7.6	9.2	9.5	13.7	15.6	11.3
2005 4/.....	*	1.6	2.9	5.2	6.5	7.8	11.9	16.8	8.9	13.5
2004 4/.....	*	1.9	3.1	5.5	10.2	10.4	15.8	14.3	17.4	14.6
2003 4/.....	*	1.6	3.6	6.5	8.6	11.0	13.7	14.7	22.0	14.6
2002.....	*	1.8	3.0	5.1	9.4	14.2	13.2	18.4	14.7	14.4
2001.....	*	1.7	2.9	6.3	8.6	13.1	19.7	16.1	14.3	15.2
2000.....	*	2.1	3.4	5.8	11.3	13.8	16.2	18.2	22.1	16.3
1999.....	*	1.6	3.4	6.4	10.3	15.6	14.9	16.3	17.3	13.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of stomach (C16)

All races 3/, both sexes

2006.....	1.3	2.1	3.3	5.0	7.2	11.0	15.5	22.3	29.9	35.8
2005.....	1.3	2.1	3.1	4.8	7.3	11.8	16.7	23.5	30.6	38.5
2004.....	1.4	2.1	3.5	5.3	8.0	12.1	18.2	23.4	30.3	42.2
2003.....	1.3	2.3	3.8	5.3	8.6	12.4	18.4	24.2	32.2	43.6
2002.....	1.2	2.1	3.5	5.6	8.5	12.3	18.8	25.6	33.0	47.6
2001.....	1.3	2.2	3.7	5.3	9.2	13.7	19.5	27.0	34.7	45.4
2000.....	1.4	2.3	3.6	5.6	9.7	13.8	19.6	27.1	36.9	52.7
1999.....	1.4	2.3	3.8	6.2	9.8	14.2	20.2	28.1	36.8	50.1

All races 3/, male

2006.....	1.4	2.6	4.4	7.0	10.1	15.4	21.4	31.3	44.7	50.2
2005.....	1.6	2.5	4.3	6.6	10.4	16.4	23.8	32.5	44.3	54.2
2004.....	1.7	2.6	4.6	7.5	11.2	17.0	24.9	33.6	42.5	62.9
2003.....	1.5	2.8	5.0	7.5	12.3	17.9	26.6	34.2	44.7	60.0
2002.....	1.4	2.7	4.8	7.7	11.9	18.1	27.5	36.4	47.1	67.2
2001.....	1.4	2.8	4.9	7.5	13.7	19.5	27.5	39.1	49.4	62.6
2000.....	1.7	2.9	4.8	7.8	14.1	19.5	28.3	39.4	51.4	75.6
1999.....	1.7	3.1	5.4	8.8	13.7	20.3	28.6	41.5	54.9	71.3

All races 3/, female

2006.....	1.2	1.7	2.3	3.1	4.5	7.2	10.6	15.8	20.7	29.1
2005.....	1.1	1.8	2.0	3.0	4.5	7.7	11.0	17.0	22.1	31.2
2004.....	1.2	1.6	2.4	3.3	5.1	7.8	12.8	16.0	22.7	32.9
2003.....	1.1	1.7	2.6	3.1	5.3	7.7	11.9	17.1	24.5	36.4
2002.....	1.0	1.5	2.3	3.7	5.4	7.3	11.9	18.1	24.4	39.1
2001.....	1.1	1.6	2.5	3.2	5.1	8.8	13.1	18.5	25.9	38.3
2000.....	1.2	1.7	2.4	3.5	5.8	8.9	12.7	18.5	28.3	43.4
1999.....	1.2	1.5	2.4	3.7	6.4	9.1	13.7	18.8	26.2	41.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of stomach (C16)

White, both sexes

2006 4/.....	1.0	1.8	3.0	4.4	6.1	9.5	13.5	19.6	26.1	32.4
2005 4/.....	1.1	1.7	2.7	4.2	6.2	10.0	14.7	20.5	27.0	34.8
2004 4/.....	1.2	1.7	2.9	4.5	6.9	10.2	15.9	20.9	27.3	38.8
2003 4/.....	1.1	1.8	3.1	4.6	7.6	11.0	16.3	21.4	28.8	38.7
2002.....	1.0	1.8	2.9	4.9	7.3	10.5	16.5	22.6	29.3	42.2
2001.....	1.1	1.9	3.1	4.7	7.9	11.7	17.4	24.0	31.0	41.1
2000.....	1.1	1.9	3.1	4.9	8.5	11.9	17.2	23.6	32.7	47.3
1999.....	1.1	2.0	3.1	5.2	8.3	12.8	17.7	24.8	33.0	44.8

White, male

2006 4/.....	1.1	2.2	3.9	6.2	8.6	13.4	18.8	27.8	39.2	45.8
2005 4/.....	1.3	2.0	3.7	5.8	8.6	13.7	20.7	28.7	39.1	49.0
2004 4/.....	1.4	2.1	3.9	6.5	9.5	14.4	21.3	29.9	38.0	58.4
2003 4/.....	1.2	2.3	4.1	6.6	10.6	15.5	23.5	30.3	40.3	53.5
2002.....	1.1	2.4	3.9	6.8	10.3	15.7	24.3	32.8	42.4	60.5
2001.....	1.2	2.4	4.1	6.5	11.7	16.3	24.5	35.4	44.0	57.7
2000.....	1.2	2.3	4.2	6.7	12.5	17.1	25.1	34.6	46.1	68.9
1999.....	1.2	2.7	4.4	7.6	11.5	18.4	25.2	36.9	49.0	63.4

White, female

2006 4/.....	1.0	1.3	2.0	2.6	3.9	6.0	9.1	13.5	17.8	26.1
2005 4/.....	0.9	1.5	1.8	2.6	3.9	6.7	9.7	14.5	19.5	28.2
2004 4/.....	1.0	1.3	2.0	2.7	4.4	6.5	11.5	14.2	20.7	29.9
2003 4/.....	0.9	1.4	2.2	2.7	4.8	7.0	10.5	15.0	21.7	32.1
2002.....	0.8	1.2	1.8	3.0	4.6	5.9	10.1	15.4	21.3	34.2
2001.....	1.0	1.3	2.1	2.9	4.3	7.7	11.7	16.0	23.1	34.1
2000.....	1.0	1.5	2.0	3.1	4.9	7.4	10.9	15.8	24.7	38.6
1999.....	0.9	1.2	1.9	2.9	5.4	8.0	11.7	16.3	23.5	37.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of stomach (C16)

Black, both sexes

2006 4/.....	2.3	3.6	5.3	8.3	14.7	20.7	28.4	42.1	64.4	66.9
2005 4/.....	2.3	4.0	5.2	9.0	14.4	23.5	30.1	48.6	57.3	74.3
2004 4/.....	2.7	3.9	6.9	10.6	15.8	27.3	34.0	43.8	59.9	71.3
2003 4/.....	2.4	4.3	7.4	9.6	16.7	22.3	33.1	48.8	62.4	85.8
2002.....	2.2	3.6	7.5	10.4	16.1	24.6	38.9	52.7	63.1	100.4
2001.....	2.5	3.7	7.5	9.2	17.7	27.1	34.7	51.1	69.9	88.1
2000.....	2.9	4.7	7.0	11.1	17.3	26.8	38.2	56.9	75.1	101.7
1999.....	3.2	3.9	8.5	13.2	19.1	23.6	40.3	58.6	74.3	102.4

Black, male

2006 4/.....	2.9	4.7	7.6	12.5	21.9	30.7	38.1	67.6	100.6	84.0
2005 4/.....	3.2	5.4	7.7	13.5	24.2	34.9	48.4	67.5	79.4	114.3
2004 4/.....	3.1	5.3	9.5	15.3	25.2	40.7	56.0	63.7	93.3	100.3
2003 4/.....	3.3	6.2	10.9	14.9	26.0	36.4	54.2	74.9	87.1	106.3
2002.....	2.4	5.1	11.0	15.2	24.0	36.9	59.8	73.5	90.2	137.2
2001.....	2.8	5.0	11.5	13.8	27.5	43.7	54.2	75.7	106.7	104.4
2000.....	3.9	6.6	9.6	16.5	25.9	38.5	57.2	83.2	105.4	136.1
1999.....	4.0	5.7	12.2	19.0	28.2	35.1	60.4	89.3	121.2	147.6

Black, female

2006 4/.....	1.7	2.6	3.3	4.9	9.1	13.3	21.8	26.8	45.2	60.0
2005 4/.....	1.5	2.8	3.1	5.3	6.8	15.0	17.5	37.3	45.6	58.1
2004 4/.....	2.3	2.7	4.7	6.7	8.4	17.3	19.1	32.1	42.3	59.9
2003 4/.....	1.5	2.7	4.5	5.3	9.5	11.9	18.8	33.4	49.5	77.8
2002.....	2.1	2.3	4.6	6.5	9.9	15.5	24.9	40.3	48.9	86.3
2001.....	2.2	2.5	4.1	5.4	10.1	14.8	21.7	36.3	51.1	82.1
2000.....	1.9	3.0	4.7	6.6	10.5	18.2	25.5	41.0	59.9	88.9
1999.....	2.4	2.4	5.4	8.4	12.0	15.2	27.0	40.0	51.0	85.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races 3/, both sexes

2006.....	4.0	7.6	13.5	21.8	34.8	52.0	73.9	104.4	146.8	207.5
2005.....	4.0	7.4	13.4	21.4	34.2	52.7	75.8	105.7	149.2	216.9
2004.....	3.9	7.4	13.5	22.4	35.9	53.2	78.3	111.3	152.3	225.6
2003.....	3.8	7.9	14.3	23.0	38.6	59.3	83.3	117.0	159.0	239.7
2002.....	4.3	8.2	15.0	24.6	40.7	61.6	86.2	118.9	163.2	241.4
2001.....	4.1	8.4	15.2	25.4	39.3	62.9	88.8	121.1	165.8	255.7
2000.....	4.0	8.2	14.7	25.2	42.0	62.0	92.4	126.9	174.8	270.4
1999.....	4.0	8.3	14.4	25.3	43.6	63.9	91.9	129.0	173.2	271.7

All races 3/, male

2006.....	4.2	8.5	15.1	25.6	43.1	64.9	91.9	127.3	181.3	234.6
2005.....	4.2	8.2	15.2	25.9	42.4	67.7	94.4	131.8	180.2	245.9
2004.....	4.2	8.3	16.1	27.0	43.7	66.4	98.0	139.0	183.1	254.7
2003.....	4.2	9.0	16.6	27.6	47.1	75.0	103.4	143.0	196.5	273.3
2002.....	4.9	9.1	17.5	30.1	51.6	77.8	106.8	147.7	197.4	275.1
2001.....	4.3	9.5	17.1	30.8	48.0	78.9	110.4	149.4	201.3	298.3
2000.....	4.4	9.0	17.0	30.6	51.0	77.4	115.7	159.4	213.0	315.6
1999.....	4.1	9.1	16.6	30.6	52.9	78.0	114.3	161.3	216.2	324.0

All races 3/, female

2006.....	3.8	6.7	11.9	18.2	27.2	40.8	59.2	87.6	125.3	194.9
2005.....	3.7	6.6	11.6	17.1	26.7	39.7	60.8	86.8	130.0	203.6
2004.....	3.6	6.5	11.1	18.0	28.7	41.8	62.4	91.3	133.3	212.5
2003.....	3.4	6.9	12.1	18.6	30.9	45.6	67.3	98.5	136.1	224.8
2002.....	3.6	7.4	12.6	19.5	30.8	47.6	69.7	98.6	142.3	226.8
2001.....	3.8	7.4	13.4	20.4	31.5	49.2	71.5	101.2	144.6	237.8
2000.....	3.5	7.5	12.4	20.2	33.9	48.8	74.1	104.3	152.2	251.9
1999.....	3.9	7.5	12.3	20.4	35.1	51.8	74.3	106.6	148.0	250.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of colon, rectum and anus (C18-C21)

White, both sexes

2006 4/.....	3.6	7.1	12.6	20.7	32.9	50.0	72.5	101.8	144.5	207.4
2005 4/.....	3.7	6.8	12.2	19.9	32.1	50.3	74.4	103.1	147.3	215.7
2004 4/.....	3.6	6.8	12.6	21.3	34.2	51.2	76.2	109.5	150.9	225.1
2003 4/.....	3.4	7.3	13.0	21.5	36.9	58.0	80.7	115.1	158.2	238.5
2002.....	3.9	7.5	14.2	23.3	38.9	59.9	84.6	116.9	160.3	240.7
2001.....	3.8	7.7	14.3	23.9	37.9	60.8	86.8	118.0	163.7	255.3
2000.....	3.6	7.5	13.8	23.7	40.5	60.3	90.6	125.0	172.1	269.2
1999.....	3.7	7.7	13.5	23.8	42.0	62.0	90.0	126.9	170.8	270.9

White, male

2006 4/.....	3.8	7.9	14.0	24.6	40.8	62.7	89.6	124.4	177.1	233.0
2005 4/.....	4.1	7.6	13.9	24.1	40.0	64.6	92.9	128.0	179.5	245.3
2004 4/.....	3.9	7.8	15.1	26.0	41.9	64.0	95.3	135.9	181.7	255.4
2003 4/.....	3.8	8.3	15.2	26.2	45.0	73.6	100.8	140.7	196.0	272.9
2002.....	4.5	8.4	16.5	28.7	49.4	76.0	105.4	144.9	193.3	273.6
2001.....	4.0	8.7	16.4	29.3	46.7	76.7	108.7	146.7	198.9	297.3
2000.....	3.9	8.3	16.1	28.6	49.1	75.3	113.5	157.6	208.6	313.3
1999.....	3.9	8.5	15.8	28.8	51.1	76.1	113.3	158.1	214.4	324.5

White, female

2006 4/.....	3.5	6.3	11.2	17.0	25.4	38.7	58.4	85.0	123.9	195.4
2005 4/.....	3.4	6.0	10.5	15.9	24.8	37.6	59.2	84.7	127.1	202.0
2004 4/.....	3.4	5.9	10.3	16.8	27.1	39.8	60.5	90.3	131.8	211.4
2003 4/.....	2.9	6.3	10.8	17.0	29.5	44.1	64.2	96.6	134.8	223.2
2002.....	3.3	6.5	12.0	18.2	29.3	45.7	67.6	96.9	140.1	226.4
2001.....	3.6	6.7	12.2	18.8	29.8	46.9	69.1	97.7	142.5	237.7
2000.....	3.3	6.6	11.5	19.1	32.6	47.2	72.3	102.1	150.4	251.3
1999.....	3.5	6.9	11.2	19.2	33.6	49.7	71.5	104.9	145.1	249.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of colon, rectum and anus (C18-C21)

Black, both sexes

2006 4/.....	6.9	12.6	22.4	33.8	57.3	79.1	101.6	147.8	197.8	240.8
2005 4/.....	5.7	12.6	23.8	36.5	57.5	82.9	101.3	148.3	197.6	261.7
2004 4/.....	5.9	12.2	21.9	36.1	56.7	79.9	113.2	148.6	189.1	260.8
2003 4/.....	6.6	13.2	25.5	39.4	59.7	81.2	124.6	154.7	197.7	281.5
2002.....	7.2	14.5	23.5	40.0	61.9	86.8	114.9	160.5	218.7	274.7
2001.....	6.5	14.5	24.6	43.1	58.3	90.9	120.3	172.6	211.0	286.5
2000.....	6.6	15.2	24.8	39.9	62.7	86.9	126.1	167.4	226.2	310.0
1999.....	6.6	13.3	23.4	42.0	65.7	90.5	125.4	172.3	226.8	306.1

Black, male

2006 4/.....	7.3	14.9	26.4	39.5	70.5	99.8	137.8	188.4	266.3	309.2
2005 4/.....	5.7	13.7	27.3	46.0	74.2	106.7	124.9	197.1	231.8	295.6
2004 4/.....	6.6	13.2	26.3	42.0	69.7	100.5	147.0	198.8	239.6	278.2
2003 4/.....	7.1	15.2	30.3	46.1	76.4	102.7	156.7	194.9	242.2	317.0
2002.....	8.4	15.7	27.9	47.5	79.4	109.2	138.2	209.0	285.1	328.1
2001.....	7.6	16.4	26.4	51.3	67.7	114.1	146.8	208.2	263.2	347.2
2000.....	8.0	16.4	29.2	51.4	76.4	109.3	160.1	208.9	298.4	377.1
1999.....	6.3	15.0	26.9	54.1	80.4	110.6	147.6	228.0	274.7	350.4

Black, female

2006 4/.....	6.4	10.7	19.1	29.1	47.0	63.9	76.7	123.5	161.7	212.8
2005 4/.....	5.6	11.6	20.9	28.7	44.5	65.2	85.1	119.2	179.5	248.0
2004 4/.....	5.3	11.3	18.2	31.3	46.6	64.6	90.3	118.9	162.5	254.0
2003 4/.....	6.1	11.4	21.4	33.8	46.8	65.2	103.0	130.8	174.5	267.8
2002.....	6.1	13.5	19.7	33.9	48.2	70.1	99.2	131.5	184.2	254.2
2001.....	5.6	12.8	23.1	36.3	50.9	73.8	102.7	151.4	184.4	264.0
2000.....	5.3	14.2	21.1	30.5	52.0	70.5	103.4	142.4	189.8	285.0
1999.....	6.7	11.9	20.5	32.1	54.2	75.6	110.7	138.7	203.1	289.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All races 3/, both sexes

2006.....	1.1	3.2	8.2	11.5	13.1	17.2	23.9	30.2	34.1	33.5
2005.....	1.2	3.7	8.5	10.2	12.9	16.6	24.3	30.2	34.4	32.5
2004.....	1.2	3.9	7.6	9.4	12.5	16.6	23.7	28.8	34.6	32.4
2003.....	1.2	3.7	7.4	8.5	11.7	16.5	23.5	29.6	32.1	33.4
2002.....	1.3	3.4	6.6	8.1	11.6	16.9	22.6	27.6	31.6	35.0
2001.....	1.2	3.8	5.8	7.5	11.6	16.3	21.5	27.0	32.0	34.1
2000.....	1.4	3.7	5.4	7.7	11.3	16.0	21.6	27.1	31.3	32.7
1999.....	1.4	3.4	4.9	7.1	10.9	16.3	21.8	26.1	30.9	30.6

All races 3/, male

2006.....	1.6	5.0	13.3	18.5	19.6	25.7	33.7	42.2	48.0	43.8
2005.....	1.6	5.9	13.8	16.7	19.5	24.2	34.8	42.8	48.2	44.8
2004.....	1.8	6.2	12.8	15.1	18.6	23.6	34.0	38.8	48.4	41.5
2003.....	1.8	6.0	12.2	13.0	17.3	23.3	33.4	43.7	46.1	47.5
2002.....	2.0	5.7	11.1	12.4	17.4	24.8	32.8	39.5	44.8	46.2
2001.....	1.8	6.0	9.7	11.4	17.2	23.5	30.6	39.2	43.9	50.3
2000.....	2.0	6.0	8.5	11.7	16.5	23.6	30.9	39.3	46.7	44.7
1999.....	2.1	5.5	7.6	10.6	15.6	23.7	31.9	36.6	42.9	40.1

All races 3/, female

2006.....	0.7	1.5	3.3	4.9	7.1	9.8	15.9	21.4	25.4	28.7
2005.....	0.8	1.5	3.4	4.1	6.8	9.9	15.8	21.0	25.9	26.8
2004.....	0.6	1.8	2.6	4.1	6.9	10.6	15.4	21.7	26.1	28.3
2003.....	0.7	1.6	2.8	4.3	6.6	10.7	15.6	19.6	23.6	27.2
2002.....	0.6	1.2	2.4	4.0	6.4	10.1	14.4	19.2	23.6	30.2
2001.....	0.5	1.7	2.1	3.9	6.5	10.2	14.2	18.4	24.8	27.2
2000.....	0.8	1.4	2.4	4.0	6.5	9.5	14.2	18.6	22.2	27.8
1999.....	0.8	1.3	2.3	3.8	6.6	10.1	13.8	18.7	23.9	26.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

White, both sexes

2006 4/.....	0.9	2.7	7.2	9.9	11.7	15.8	22.7	28.9	32.8	33.1
2005 4/.....	1.0	3.3	7.4	8.9	11.3	15.3	22.5	29.1	32.8	31.7
2004 4/.....	1.0	3.3	6.6	8.0	11.2	15.1	21.9	27.3	33.7	31.2
2003 4/.....	1.0	3.1	6.3	7.5	10.3	15.2	21.7	28.3	31.0	32.1
2002.....	1.1	3.0	5.5	6.8	10.6	15.8	21.1	26.7	30.7	33.8
2001.....	0.8	3.2	4.9	6.4	10.5	15.1	20.2	26.1	30.4	33.3
2000.....	1.1	3.1	4.4	6.6	10.0	14.6	20.2	25.8	30.7	32.0
1999.....	1.1	2.7	4.2	5.9	9.7	15.2	20.5	25.2	30.2	29.9

White, male

2006 4/.....	1.2	4.1	11.6	15.6	17.1	23.1	32.0	40.6	46.9	43.7
2005 4/.....	1.3	5.2	11.6	14.2	17.1	22.0	32.2	41.5	45.8	45.0
2004 4/.....	1.5	5.0	11.1	12.4	16.5	21.5	31.1	37.4	47.2	39.9
2003 4/.....	1.4	4.9	10.2	11.4	15.2	21.4	31.4	41.8	44.8	45.6
2002.....	1.6	4.9	9.0	10.4	15.6	23.0	30.4	38.0	42.9	45.1
2001.....	1.3	5.0	8.1	9.5	15.7	22.1	29.1	37.7	42.6	49.7
2000.....	1.5	4.8	6.8	9.6	14.4	21.4	28.9	38.1	45.8	43.1
1999.....	1.7	4.3	6.3	8.8	13.9	21.9	30.2	35.3	42.4	40.6

White, female

2006 4/.....	0.7	1.3	2.9	4.4	6.7	9.2	15.0	20.2	23.9	28.1
2005 4/.....	0.7	1.3	3.2	3.7	6.0	9.3	14.5	19.9	24.7	25.6
2004 4/.....	0.5	1.5	2.2	3.7	6.3	9.5	14.3	19.8	25.3	27.3
2003 4/.....	0.6	1.3	2.5	3.8	5.7	9.7	13.9	18.6	22.5	26.2
2002.....	0.5	1.1	2.0	3.4	5.9	9.3	13.5	18.6	23.2	29.0
2001.....	0.4	1.4	1.9	3.4	5.7	8.9	13.0	17.8	23.1	26.5
2000.....	0.6	1.3	2.0	3.6	5.9	8.5	13.2	17.1	21.7	27.4
1999.....	0.6	1.2	2.1	3.1	5.8	9.3	12.8	18.1	23.0	25.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Black, both sexes

2006 4/.....	1.7	5.4	14.3	22.7	20.4	22.3	26.7	30.8	36.9	31.1
2005 4/.....	1.8	5.6	14.5	18.9	22.1	21.4	29.4	29.1	39.3	29.1
2004 4/.....	1.8	7.1	12.7	17.9	16.6	23.2	28.5	33.7	35.4	36.1
2003 4/.....	1.9	6.2	14.2	14.7	18.4	18.9	27.9	30.3	32.7	41.7
2002.....	2.1	5.8	12.3	15.5	14.0	19.5	27.3	30.2	31.9	38.0
2001.....	2.3	7.1	10.7	15.2	14.3	20.0	24.0	26.9	40.9	35.0
2000.....	2.7	7.0	10.6	12.7	15.8	20.9	26.9	28.8	28.5	31.6
1999.....	2.2	6.6	8.1	12.0	14.7	21.1	28.0	26.6	33.8	34.4

Black, male

2006 4/.....	2.9	8.0	24.3	42.3	35.2	37.0	42.4	45.2	45.9	34.9
2005 4/.....	2.5	8.9	26.8	35.0	36.7	35.9	44.6	45.1	59.7	31.6
2004 4/.....	2.9	11.7	22.9	31.9	27.1	36.8	45.0	39.6	50.9	46.7
2003 4/.....	3.0	10.3	25.5	24.0	28.5	26.8	42.4	43.8	45.0	63.6
2002.....	3.3	9.8	21.7	25.3	22.8	32.2	45.1	44.6	51.2	41.8
2001.....	3.6	11.6	18.6	25.3	21.4	28.0	32.7	39.5	49.0	42.0
2000.....	3.8	12.6	17.7	22.1	25.3	34.1	41.5	36.4	45.5	48.4
1999.....	3.4	11.8	14.2	18.6	21.6	32.2	42.1	40.4	42.7	30.5

Black, female

2006 4/.....	*	3.0	5.8	6.5	9.0	11.4	15.8	22.2	32.2	29.6
2005 4/.....	*	2.6	4.2	5.6	10.8	10.6	19.0	19.5	28.6	28.1
2004 4/.....	*	3.0	4.1	6.4	8.4	13.1	17.4	30.2	27.2	31.9
2003 4/.....	*	2.7	4.5	7.0	10.5	13.0	18.2	22.2	26.3	33.2
2002.....	*	2.3	4.3	7.5	7.1	10.2	15.5	21.7	21.9	36.6
2001.....	*	3.2	3.9	6.8	8.8	14.1	18.2	19.4	36.8	32.4
2000.....	1.8	2.2	4.7	5.0	8.3	11.1	17.3	24.2	20.0	25.3
1999.....	*	2.0	2.9	6.5	9.3	12.9	18.7	18.3	29.4	35.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of pancreas (C25)

All races 3/, both sexes

2006.....	1.7	4.4	8.5	15.3	26.2	37.7	53.4	70.2	89.7	96.9
2005.....	1.9	4.5	8.7	15.7	25.1	37.6	52.6	69.6	88.4	97.4
2004.....	1.7	4.2	8.8	15.2	25.8	37.3	52.4	68.9	85.4	97.7
2003.....	1.7	4.1	8.7	14.7	24.7	36.4	53.0	67.6	86.2	95.2
2002.....	1.9	4.2	8.2	15.5	24.0	37.2	53.2	70.2	81.8	94.5
2001.....	1.8	4.0	8.6	14.9	24.8	37.0	52.8	67.2	84.2	100.1
2000.....	1.8	4.2	8.7	15.1	24.8	37.2	51.4	70.4	85.7	96.4
1999.....	1.9	4.5	8.7	15.6	25.0	38.1	52.5	67.2	82.0	99.7

All races 3/, male

2006.....	2.0	5.2	10.7	19.2	31.9	45.3	61.2	77.3	99.0	102.1
2005.....	2.2	5.2	10.6	19.6	30.6	43.9	61.6	78.7	96.5	102.8
2004.....	2.2	5.5	11.0	18.6	31.5	45.2	61.6	77.7	94.8	102.0
2003.....	2.0	5.2	10.7	18.2	29.8	42.5	60.9	76.1	99.5	95.6
2002.....	2.4	5.3	10.6	19.2	29.3	43.3	60.8	78.9	91.9	100.9
2001.....	2.0	4.9	10.7	18.5	30.3	43.6	59.0	77.2	94.9	105.1
2000.....	2.1	5.4	11.1	18.5	30.0	44.7	57.0	80.4	94.9	102.9
1999.....	2.4	5.9	11.0	19.1	29.9	44.9	60.9	75.7	92.0	107.5

All races 3/, female

2006.....	1.4	3.5	6.3	11.7	21.0	31.1	47.0	64.9	83.9	94.4
2005.....	1.6	3.8	6.9	12.0	20.0	32.1	45.3	63.0	83.4	94.9
2004.....	1.3	2.9	6.7	12.0	20.6	30.5	45.1	62.6	79.6	95.8
2003.....	1.4	3.0	6.8	11.5	20.2	31.1	46.7	61.6	78.1	95.0
2002.....	1.4	3.1	6.0	12.0	19.2	31.9	47.1	64.1	75.6	91.7
2001.....	1.6	3.1	6.5	11.5	19.8	31.3	47.8	60.1	77.8	98.1
2000.....	1.5	3.0	6.4	11.8	20.0	30.8	47.1	63.5	80.2	93.8
1999.....	1.4	3.1	6.5	12.3	20.5	32.2	46.0	61.2	76.1	96.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of pancreas (C25)

White, both sexes

2006 4/.....	1.6	4.2	8.2	14.9	25.6	37.3	52.6	70.0	89.5	96.1
2005 4/.....	1.9	4.4	8.2	15.2	24.6	37.1	52.2	68.8	87.7	96.8
2004 4/.....	1.6	4.0	8.4	14.6	25.1	36.8	51.3	68.3	85.1	97.1
2003 4/.....	1.6	3.9	8.1	14.3	23.8	36.0	52.2	67.2	85.2	94.4
2002.....	1.9	4.1	7.8	15.0	23.3	36.1	52.0	69.2	81.4	94.3
2001.....	1.7	3.8	8.3	14.4	24.7	36.3	51.7	66.3	83.3	99.0
2000.....	1.7	4.1	8.4	14.5	24.3	36.2	50.4	69.2	84.5	94.7
1999.....	1.8	4.4	8.2	14.8	24.3	37.2	51.5	65.8	80.6	98.6

White, male

2006 4/.....	1.9	5.1	10.3	18.7	31.3	45.1	60.9	77.9	100.3	103.4
2005 4/.....	2.2	5.1	10.0	18.9	30.4	43.4	62.0	78.1	96.2	102.9
2004 4/.....	2.0	5.3	10.5	17.7	30.8	44.5	60.0	77.7	95.8	102.9
2003 4/.....	1.8	4.8	10.0	17.7	28.9	42.7	60.2	76.1	99.3	96.0
2002.....	2.4	5.1	10.1	18.6	28.8	42.6	59.7	78.9	92.2	101.6
2001.....	1.9	4.7	10.5	18.2	30.3	42.8	58.0	77.0	94.5	106.5
2000.....	2.0	5.3	10.7	17.8	29.6	44.3	55.9	79.9	94.0	102.4
1999.....	2.3	5.8	10.3	17.9	29.2	44.0	60.1	74.6	90.9	107.4

White, female

2006 4/.....	1.3	3.3	6.1	11.3	20.3	30.3	45.7	64.1	82.7	92.7
2005 4/.....	1.5	3.6	6.4	11.7	19.3	31.5	44.0	61.9	82.4	94.0
2004 4/.....	1.3	2.8	6.2	11.5	19.7	30.0	44.1	61.5	78.5	94.5
2003 4/.....	1.4	2.9	6.3	11.0	19.1	30.0	45.8	60.7	76.6	93.8
2002.....	1.4	3.0	5.5	11.6	18.2	30.3	45.7	62.2	74.9	91.1
2001.....	1.6	2.9	6.3	10.8	19.6	30.6	46.5	58.7	76.5	95.9
2000.....	1.4	2.9	6.1	11.4	19.4	29.2	46.0	61.6	78.9	91.5
1999.....	1.3	3.1	6.3	11.9	19.9	31.3	44.6	59.5	74.5	95.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of pancreas (C25)

Black, both sexes

2006 4/.....	2.3	6.2	12.6	21.6	37.2	49.2	67.3	85.1	99.9	113.4
2005 4/.....	2.2	6.2	13.7	22.3	33.6	49.1	66.4	86.9	105.1	111.4
2004 4/.....	2.6	6.0	13.3	22.6	36.3	48.0	70.5	84.3	93.3	113.6
2003 4/.....	2.7	6.4	14.5	20.7	36.1	47.1	68.0	83.6	107.4	112.9
2002.....	2.5	5.6	12.3	22.4	34.4	50.4	73.7	88.6	91.8	106.6
2001.....	2.2	6.4	12.1	20.7	31.6	48.5	70.8	84.9	104.8	121.3
2000.....	2.6	5.8	13.2	22.8	33.3	50.3	69.8	93.8	104.2	119.8
1999.....	2.8	6.2	14.1	25.0	32.2	50.9	70.1	87.5	104.6	118.6

Black, male

2006 4/.....	2.7	6.6	16.8	28.7	45.8	58.6	75.7	86.8	98.6	91.1
2005 4/.....	2.3	7.2	17.3	30.1	39.6	59.2	71.5	96.1	116.6	100.4
2004 4/.....	3.4	7.6	16.1	30.0	44.8	60.6	88.1	88.7	91.1	100.3
2003 4/.....	3.5	9.0	18.6	26.2	42.8	50.0	80.8	89.0	112.2	104.3
2002.....	2.9	7.2	16.5	29.0	40.8	56.6	84.1	90.5	105.5	100.8
2001.....	2.3	7.9	15.2	24.2	38.2	60.0	78.8	95.2	109.2	99.8
2000.....	3.0	7.3	17.5	29.3	40.4	55.7	78.9	101.6	116.3	109.6
1999.....	3.4	8.3	19.5	34.1	39.5	60.0	78.4	100.7	117.7	114.8

Black, female

2006 4/.....	1.9	5.8	9.1	15.8	30.6	42.2	61.5	84.1	100.6	122.5
2005 4/.....	2.2	5.3	10.7	15.8	28.9	41.6	63.0	81.3	99.0	115.9
2004 4/.....	1.9	4.6	11.0	16.6	29.7	38.7	58.6	81.7	94.5	118.9
2003 4/.....	2.1	4.1	11.0	16.2	30.8	45.0	59.4	80.3	104.8	116.3
2002.....	2.1	4.3	8.8	17.0	29.4	45.9	66.8	87.5	84.7	108.8
2001.....	2.2	5.1	9.6	17.8	26.4	40.0	65.5	78.8	102.6	129.2
2000.....	2.3	4.4	9.4	17.4	27.8	46.3	63.7	89.1	98.1	123.6
1999.....	2.2	4.3	9.5	17.6	26.5	44.3	64.6	79.5	98.1	120.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of larynx (C32)

All races 3/, both sexes

2006.....	0.3	0.7	1.4	2.6	3.8	5.2	6.2	6.9	7.4	6.3
2005.....	0.2	0.7	1.3	2.4	3.9	5.8	6.7	6.8	7.3	6.3
2004.....	0.2	0.7	1.4	2.6	4.1	5.3	6.2	7.1	6.0	6.5
2003.....	0.3	0.7	1.4	2.6	4.6	5.8	6.6	6.6	7.4	6.5
2002.....	0.2	0.7	1.4	2.7	4.2	5.3	6.6	7.9	7.2	5.5
2001.....	0.3	0.7	1.5	2.9	4.5	5.6	7.3	7.4	6.9	6.7
2000.....	0.2	0.8	1.5	2.9	4.5	6.0	7.3	8.0	8.1	6.5
1999.....	0.3	0.8	1.5	3.1	4.6	6.1	7.4	7.2	7.4	7.0

All races 3/, male

2006.....	0.3	1.2	2.3	4.4	6.2	9.2	10.5	12.5	14.7	14.8
2005.....	0.4	1.0	2.1	4.0	6.7	10.2	11.4	12.2	14.5	15.7
2004.....	0.4	1.2	2.4	4.4	6.8	9.4	11.0	13.3	12.4	15.6
2003.....	0.5	1.2	2.4	4.2	7.9	10.3	11.4	12.6	15.4	15.3
2002.....	0.4	1.2	2.4	4.7	7.2	8.9	11.3	14.8	14.9	13.2
2001.....	0.5	1.2	2.5	4.9	7.5	9.5	12.9	14.5	13.9	18.3
2000.....	0.4	1.2	2.5	4.9	7.9	10.0	12.9	14.8	17.1	17.4
1999.....	0.4	1.3	2.6	5.4	7.8	10.4	12.6	13.2	15.8	17.8

All races 3/, female

2006.....	*	0.3	0.5	1.0	1.5	1.7	2.7	2.8	2.8	2.3
2005.....	*	0.4	0.6	0.8	1.3	1.9	3.0	2.8	2.8	1.9
2004.....	*	0.2	0.5	0.9	1.6	1.7	2.2	2.6	2.1	2.4
2003.....	*	0.3	0.4	1.0	1.6	2.0	2.8	2.3	2.5	2.7
2002.....	*	0.2	0.5	0.9	1.5	2.2	2.9	3.1	2.5	2.2
2001.....	*	0.2	0.5	1.0	1.7	2.2	2.8	2.5	2.8	1.9
2000.....	*	0.3	0.6	0.9	1.4	2.6	2.8	3.3	2.8	2.1
1999.....	*	0.2	0.4	1.0	1.7	2.5	3.3	3.0	2.6	2.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of larynx (C32)

White, both sexes

2006 4/.....	0.2	0.7	1.2	2.4	3.6	4.8	6.1	6.8	7.2	6.2
2005 4/.....	0.2	0.6	1.1	2.1	3.4	5.4	6.3	6.8	7.2	6.0
2004 4/.....	0.2	0.7	1.2	2.3	3.8	5.0	5.9	6.9	5.7	6.3
2003 4/.....	0.3	0.6	1.1	2.3	4.2	5.5	6.2	6.3	7.1	6.2
2002.....	0.2	0.6	1.2	2.5	3.9	5.0	6.4	7.5	6.9	5.6
2001.....	0.3	0.6	1.2	2.5	4.1	5.2	6.7	7.0	6.8	6.6
2000.....	0.2	0.6	1.3	2.4	3.9	5.6	6.7	7.5	8.0	6.6
1999.....	0.2	0.6	1.3	2.7	4.2	5.7	7.1	7.1	7.3	6.7

White, male

2006 4/.....	0.3	1.0	2.0	4.0	5.8	8.2	9.9	12.0	14.3	14.7
2005 4/.....	0.4	0.9	1.8	3.5	5.7	9.5	10.5	12.1	14.0	14.9
2004 4/.....	0.3	1.2	1.9	3.9	6.2	8.8	10.4	12.8	11.7	15.0
2003 4/.....	0.5	1.1	2.0	3.7	7.2	9.3	10.4	11.9	14.4	14.5
2002.....	0.3	1.0	2.0	4.1	6.3	8.3	10.7	13.8	14.6	13.3
2001.....	0.4	0.9	2.0	4.2	6.9	8.8	11.9	13.3	13.2	18.3
2000.....	0.3	1.0	2.1	4.1	6.8	9.2	11.7	13.9	16.8	17.6
1999.....	0.3	1.1	2.2	4.6	7.0	9.6	12.0	12.9	15.1	16.8

White, female

2006 4/.....	*	0.3	0.5	0.9	1.5	1.7	2.9	3.0	2.8	2.2
2005 4/.....	*	0.3	0.5	0.8	1.3	1.7	2.9	2.9	2.9	1.9
2004 4/.....	*	0.2	0.5	0.9	1.6	1.7	2.1	2.6	2.0	2.5
2003 4/.....	*	*	0.3	0.9	1.4	2.1	2.7	2.3	2.5	2.5
2002.....	*	*	0.4	0.9	1.5	2.1	2.8	2.9	2.3	2.3
2001.....	*	*	0.4	0.9	1.5	2.1	2.6	2.5	2.9	1.7
2000.....	*	0.3	0.5	0.9	1.3	2.5	2.6	3.0	2.7	2.1
1999.....	*	*	0.3	1.0	1.5	2.3	3.3	3.0	2.7	2.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of larynx (C32)

Black, both sexes

2006 4/.....	*	1.6	2.8	5.4	6.6	10.9	9.6	10.2	9.5	8.0
2005 4/.....	*	1.3	3.3	5.5	9.1	11.5	11.4	9.2	10.9	10.2
2004 4/.....	*	1.2	3.5	5.5	8.0	9.3	11.6	11.2	11.1	8.2
2003 4/.....	*	1.7	3.6	5.8	9.7	10.6	12.7	11.1	12.7	11.1
2002.....	*	1.9	3.6	6.1	9.3	9.1	11.8	15.3	11.3	*
2001.....	*	1.9	3.8	7.1	8.9	9.5	13.9	13.4	9.2	8.3
2000.....	*	2.0	4.1	7.5	10.8	10.6	15.3	15.6	11.6	6.9
1999.....	0.9	2.1	3.9	7.5	10.0	11.4	11.8	10.2	9.8	8.3

Black, male

2006 4/.....	*	2.8	5.2	9.5	12.2	21.6	19.4	22.4	21.2	18.8
2005 4/.....	*	2.1	5.0	10.3	18.5	22.1	22.1	18.3	26.0	28.8
2004 4/.....	*	2.1	6.7	11.0	14.9	18.7	22.7	24.1	25.8	23.8
2003 4/.....	*	2.7	6.8	10.7	17.5	22.7	25.3	25.6	31.7	27.1
2002.....	*	3.2	6.0	11.5	18.8	17.0	22.4	33.1	22.9	*
2001.....	*	3.6	7.0	13.0	15.9	17.5	26.3	30.7	22.5	*
2000.....	*	3.2	7.6	14.3	20.7	19.8	30.1	29.3	27.0	*
1999.....	1.6	3.9	6.9	14.4	17.8	21.0	22.7	19.9	27.9	24.6

Black, female

2006 4/.....	*	*	*	2.0	*	*	*	*	*	*
2005 4/.....	*	*	1.8	*	*	3.7	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	3.6	*	*	*	*	*
2002.....	*	*	*	*	*	*	4.7	*	*	*
2001.....	*	*	*	*	3.4	*	5.6	*	*	*
2000.....	*	*	*	*	*	3.8	5.5	7.4	*	*
1999.....	*	*	*	*	3.8	4.2	4.5	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All races 3/, both sexes

2006.....	7.1	19.9	39.4	71.8	136.5	216.9	297.1	366.8	382.3	300.5
2005.....	8.2	20.4	40.1	73.8	142.7	218.7	308.3	372.3	380.0	302.3
2004.....	8.5	20.6	40.4	77.9	144.9	226.5	311.2	372.5	375.1	297.5
2003.....	8.5	21.2	40.7	81.8	149.0	230.0	314.0	376.1	380.1	298.9
2002.....	9.2	20.4	41.6	86.3	152.9	233.4	320.9	376.9	378.5	297.2
2001.....	9.5	20.5	42.3	88.8	154.5	239.7	323.2	373.4	368.6	302.7
2000.....	9.3	21.0	43.8	92.5	159.8	245.8	325.4	368.2	374.7	302.1
1999.....	9.3	19.9	44.4	93.0	160.7	243.6	322.3	361.7	362.6	297.0

All races 3/, male

2006.....	7.3	21.6	47.5	88.3	167.8	272.4	378.8	486.0	544.1	457.7
2005.....	8.5	22.8	49.0	91.0	177.3	275.5	399.2	497.5	540.5	468.0
2004.....	9.1	23.5	49.8	95.8	179.7	285.8	400.8	508.4	530.2	466.8
2003.....	9.5	24.8	50.0	101.3	183.7	292.1	411.0	514.2	541.5	475.1
2002.....	9.6	23.6	51.4	107.0	192.7	297.5	423.0	515.5	546.8	482.2
2001.....	10.5	24.3	52.2	109.6	195.7	312.9	434.7	522.3	541.0	512.4
2000.....	10.5	25.7	53.2	115.9	202.3	322.5	440.5	516.5	558.1	521.2
1999.....	10.5	24.3	54.4	118.1	207.7	322.4	439.2	513.4	551.3	529.2

All races 3/, female

2006.....	6.9	18.3	31.6	56.2	107.8	168.4	230.7	279.5	281.3	226.9
2005.....	8.0	18.1	31.7	57.6	111.2	169.1	234.7	281.3	280.3	226.2
2004.....	7.9	17.8	31.5	61.1	113.2	174.9	239.1	274.9	279.5	221.3
2003.....	7.5	17.8	31.9	63.5	117.4	176.2	236.1	277.9	281.4	221.0
2002.....	8.8	17.3	32.2	66.8	116.8	177.9	239.3	279.1	276.1	217.0
2001.....	8.4	16.8	32.8	69.2	117.2	176.7	234.5	269.1	265.2	215.0
2000.....	8.1	16.5	34.7	70.5	121.3	180.2	234.7	264.9	266.6	212.8
1999.....	8.1	15.7	34.9	69.5	118.1	176.1	230.9	256.2	252.2	203.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

White, both sexes

2006 4/.....	7.0	19.1	37.5	70.9	136.6	220.6	303.6	375.7	388.5	306.1
2005 4/.....	8.1	19.6	38.7	72.0	142.7	222.2	315.0	380.7	385.6	305.6
2004 4/.....	8.1	19.7	38.8	76.1	145.5	229.2	318.0	378.3	378.9	302.0
2003 4/.....	8.2	19.9	39.1	80.7	149.9	233.5	319.7	381.3	384.5	302.4
2002.....	8.8	19.4	40.0	84.9	152.6	236.4	327.0	382.1	384.1	300.9
2001.....	9.1	19.3	40.9	87.9	155.9	242.0	327.4	377.2	371.2	305.4
2000.....	8.6	19.5	42.2	91.5	161.2	248.2	328.0	370.7	380.7	304.5
1999.....	8.8	18.4	42.6	92.0	160.6	244.5	324.6	362.9	365.2	298.8

White, male

2006 4/.....	7.1	20.3	44.3	84.9	164.0	271.4	380.3	490.4	548.9	464.0
2005 4/.....	8.1	21.5	46.6	86.3	173.3	275.0	401.3	503.1	543.8	469.9
2004 4/.....	8.5	21.8	46.4	91.4	175.5	284.9	401.9	508.8	530.1	472.2
2003 4/.....	8.9	22.8	46.9	97.3	180.4	291.1	412.1	513.0	541.9	478.5
2002.....	8.9	21.8	48.1	102.8	187.3	295.9	424.5	514.5	547.8	486.6
2001.....	9.7	22.2	49.0	105.6	193.4	309.8	432.8	519.4	537.4	513.9
2000.....	9.3	23.5	49.5	111.5	200.5	319.1	436.8	511.6	560.0	522.4
1999.....	9.7	21.9	50.5	114.3	203.1	317.9	435.0	508.8	548.0	531.5

White, female

2006 4/.....	6.8	17.9	30.8	57.3	111.0	175.3	240.0	290.2	287.2	231.9
2005 4/.....	8.0	17.7	30.9	58.3	114.2	175.2	243.6	290.1	286.4	229.9
2004 4/.....	7.7	17.7	31.3	61.3	117.7	179.7	249.0	282.8	285.0	225.3
2003 4/.....	7.5	17.0	31.5	64.8	121.6	182.6	244.0	286.0	287.4	224.4
2002.....	8.7	17.1	32.0	67.7	120.6	184.0	247.5	287.3	283.8	220.2
2001.....	8.4	16.4	33.1	71.0	121.3	182.5	242.1	276.4	270.7	218.1
2000.....	8.0	15.6	35.0	72.5	125.0	186.2	240.7	271.6	274.1	215.9
1999.....	7.9	15.0	35.0	70.8	121.6	180.5	236.9	260.4	257.3	205.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Black, both sexes

2006 4/.....	9.4	30.2	62.5	97.4	173.6	246.1	305.4	354.5	378.9	264.6
2005 4/.....	10.9	30.9	60.6	107.8	180.3	248.0	316.4	361.1	374.0	289.2
2004 4/.....	12.7	31.0	62.7	114.0	179.5	256.4	309.0	382.2	387.7	277.5
2003 4/.....	12.4	35.4	62.0	114.8	182.9	256.6	328.9	387.1	377.2	280.9
2002.....	14.1	32.4	64.4	120.6	193.7	267.2	330.7	391.2	366.3	272.3
2001.....	14.5	32.6	63.5	119.3	182.2	273.6	347.6	395.1	370.3	280.7
2000.....	15.7	35.5	69.3	123.1	188.7	279.9	359.1	394.9	343.3	288.1
1999.....	14.5	35.1	70.8	125.2	198.4	282.1	351.1	402.9	363.5	290.5

Black, male

2006 4/.....	10.2	36.0	82.2	139.9	252.7	365.1	452.2	543.2	595.6	442.3
2005 4/.....	12.1	38.4	79.5	156.2	262.7	363.2	469.6	544.5	597.6	495.5
2004 4/.....	14.5	40.8	89.1	161.1	271.7	367.8	467.4	600.5	627.7	481.8
2003 4/.....	15.6	45.1	86.6	167.9	267.8	376.2	495.0	618.5	628.3	502.5
2002.....	17.0	42.6	91.6	172.1	292.0	398.4	506.4	630.6	619.0	484.6
2001.....	18.3	45.0	92.4	174.1	268.7	418.8	552.7	659.8	659.7	549.2
2000.....	21.4	47.4	100.4	185.2	271.7	432.2	561.6	656.8	617.1	562.8
1999.....	18.6	48.8	103.6	186.3	301.7	429.2	556.6	658.7	665.3	574.2

Black, female

2006 4/.....	8.7	25.3	45.8	62.4	112.0	158.2	204.6	241.6	264.4	191.9
2005 4/.....	9.9	24.4	44.6	67.7	116.2	162.5	211.5	251.7	255.9	205.9
2004 4/.....	11.1	22.5	40.4	75.2	107.7	173.8	201.4	252.8	261.2	196.5
2003 4/.....	9.6	26.9	41.2	71.2	116.7	167.8	216.9	249.6	245.6	194.8
2002.....	11.5	23.6	41.5	78.3	116.8	169.8	212.8	248.1	234.7	191.0
2001.....	11.1	21.9	38.8	74.3	114.1	166.3	211.0	237.1	222.8	181.0
2000.....	10.7	25.1	42.7	72.2	123.2	167.6	224.7	237.0	205.5	185.9
1999.....	10.8	23.3	42.9	75.3	117.3	173.2	214.7	248.9	213.9	184.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant melanoma of skin (C43)

All races 3/, both sexes

2006.....	1.4	2.2	3.2	4.5	6.2	8.2	10.8	14.0	17.8	20.1
2005.....	1.5	2.2	3.5	4.6	5.9	8.1	10.9	15.1	16.2	19.8
2004.....	1.4	2.3	3.3	4.7	6.1	8.1	10.6	13.6	16.3	17.7
2003.....	1.6	2.5	3.4	4.9	6.4	7.3	10.3	13.9	15.5	17.6
2002.....	1.7	2.3	3.4	4.4	6.0	7.8	10.0	12.9	14.9	17.9
2001.....	1.8	2.6	3.6	4.6	6.1	7.8	10.7	12.4	14.7	17.8
2000.....	1.7	2.7	3.8	4.7	6.3	7.9	10.0	12.0	14.9	18.9
1999.....	1.7	2.6	3.6	4.7	6.1	8.3	9.7	12.9	14.2	16.4

All races 3/, male

2006.....	1.6	2.8	4.3	6.2	9.0	11.9	15.9	22.5	29.9	36.7
2005.....	1.9	2.7	4.6	6.4	7.9	11.9	16.2	23.5	26.0	34.9
2004.....	1.8	2.9	4.5	6.6	8.9	11.8	16.1	21.8	26.7	31.4
2003.....	1.9	3.1	4.6	6.9	8.9	10.4	15.1	22.3	24.9	30.2
2002.....	2.1	3.3	4.4	6.1	8.5	11.3	14.9	19.8	23.3	29.2
2001.....	2.2	3.2	4.7	6.3	8.8	11.9	16.3	19.6	24.0	29.9
2000.....	2.1	3.2	5.2	6.5	8.7	11.2	14.7	18.4	24.7	31.8
1999.....	2.2	3.5	4.6	6.7	8.4	12.0	14.6	19.8	22.7	27.5

All races 3/, female

2006.....	1.2	1.7	2.1	2.9	3.6	5.0	6.7	7.8	10.2	12.4
2005.....	1.1	1.8	2.4	2.9	4.2	4.8	6.6	9.0	10.1	12.9
2004.....	1.1	1.7	2.2	3.0	3.6	4.9	6.2	7.7	9.9	11.5
2003.....	1.3	1.9	2.2	3.1	4.0	4.6	6.4	7.8	9.8	12.1
2002.....	1.2	1.4	2.5	2.8	3.6	4.8	6.1	7.9	9.8	12.9
2001.....	1.4	1.9	2.6	3.0	3.7	4.3	6.4	7.3	9.2	12.8
2000.....	1.4	2.1	2.4	3.0	4.1	5.0	6.3	7.5	9.1	13.7
1999.....	1.3	1.8	2.6	2.8	4.0	5.2	5.8	8.1	9.2	11.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant melanoma of skin (C43)

White, both sexes

2006 4/.....	1.7	2.7	3.8	5.2	7.0	9.5	12.3	15.5	19.5	21.7
2005 4/.....	1.8	2.6	4.1	5.3	6.8	9.3	12.3	16.8	17.6	21.4
2004 4/.....	1.7	2.8	3.9	5.5	7.0	9.2	12.0	15.1	17.6	19.2
2003 4/.....	1.9	2.9	3.9	5.6	7.3	8.3	11.6	15.2	17.0	19.1
2002.....	2.0	2.8	4.0	5.0	6.8	8.9	11.0	14.2	16.3	19.2
2001.....	2.2	3.1	4.3	5.3	7.0	8.8	11.9	13.6	15.9	19.3
2000.....	2.0	3.1	4.4	5.3	7.2	8.9	11.0	13.1	16.0	20.3
1999.....	2.1	3.1	4.2	5.4	6.9	9.4	10.7	14.0	15.3	17.6

White, male

2006 4/.....	2.0	3.3	5.1	7.1	10.2	13.7	18.0	24.8	32.8	39.7
2005 4/.....	2.2	3.1	5.4	7.3	9.0	13.6	18.3	26.0	28.3	38.1
2004 4/.....	2.1	3.5	5.3	7.5	10.1	13.4	18.1	24.0	28.9	34.2
2003 4/.....	2.3	3.6	5.3	7.9	10.1	11.7	17.1	24.4	27.1	32.9
2002.....	2.5	3.8	5.1	6.9	9.7	12.7	16.4	21.8	25.6	31.7
2001.....	2.6	3.8	5.5	7.2	10.0	13.3	18.0	21.5	26.1	32.7
2000.....	2.5	3.8	6.0	7.3	9.9	12.5	16.2	20.1	26.6	34.7
1999.....	2.7	4.1	5.3	7.7	9.5	13.4	16.2	21.5	24.5	29.7

White, female

2006 4/.....	1.4	2.1	2.5	3.3	4.1	5.7	7.6	8.6	11.0	13.3
2005 4/.....	1.3	2.1	2.8	3.4	4.7	5.5	7.4	10.0	10.9	13.8
2004 4/.....	1.4	2.1	2.6	3.5	4.1	5.4	7.0	8.6	10.6	12.4
2003 4/.....	1.5	2.3	2.6	3.5	4.6	5.2	7.1	8.5	10.8	12.9
2002.....	1.5	1.7	2.9	3.2	4.2	5.5	6.7	8.8	10.7	13.8
2001.....	1.7	2.3	3.1	3.5	4.2	4.8	7.0	8.0	9.8	13.7
2000.....	1.6	2.5	2.8	3.4	4.6	5.7	6.9	8.2	9.7	14.5
1999.....	1.5	2.2	3.0	3.3	4.6	5.8	6.4	8.8	9.9	12.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of breast (C50)

All races 3/, both sexes

2006.....	6.9	11.6	17.0	24.7	32.7	38.0	47.1	63.1	82.7	116.7
2005.....	7.4	11.8	17.9	25.0	33.1	38.7	49.3	63.5	82.9	122.7
2004.....	7.3	11.7	18.7	26.1	33.1	40.8	49.6	64.7	82.4	124.2
2003.....	7.9	12.6	19.1	26.6	33.5	39.9	51.8	66.0	87.3	132.5
2002.....	7.9	12.9	19.7	26.2	33.7	41.3	52.4	67.1	89.8	134.6
2001.....	8.2	13.4	20.6	27.1	34.1	41.7	54.0	69.1	88.0	134.2
2000.....	8.3	13.6	20.8	28.1	35.0	43.4	55.2	70.9	90.9	147.4
1999.....	7.9	13.3	21.0	28.1	34.8	43.5	55.9	70.0	92.4	144.4

All races 3/, male

2006.....	*	*	0.3	0.4	0.8	0.8	1.2	1.8	2.6	3.0
2005.....	*	*	0.3	0.4	0.7	1.0	1.7	1.8	2.5	2.4
2004.....	*	*	0.2	0.4	0.6	0.9	1.3	2.1	2.1	3.3
2003.....	*	*	0.4	0.4	0.7	0.9	1.2	1.5	2.8	3.9
2002.....	*	*	0.3	0.4	0.7	0.8	1.3	1.9	2.7	3.4
2001.....	*	*	0.4	0.5	0.8	1.2	1.7	2.3	2.4	3.5
2000.....	*	*	0.3	0.7	0.8	1.1	1.7	2.1	2.4	4.2
1999.....	*	*	0.4	0.6	0.7	1.1	1.4	1.9	2.3	4.5

All races 3/, female

2006.....	13.8	22.7	33.0	47.5	61.9	70.4	84.4	108.0	132.9	169.9
2005.....	14.7	23.2	34.9	48.3	62.6	71.6	87.9	108.3	132.8	177.9
2004.....	14.5	23.0	36.3	50.4	62.6	75.6	88.5	109.6	131.8	178.6
2003.....	15.6	24.7	37.0	51.3	63.4	73.7	92.4	111.9	139.0	189.4
2002.....	15.6	25.3	38.3	50.4	63.7	76.2	93.2	113.2	142.7	191.5
2001.....	16.2	26.3	40.1	52.0	64.3	76.5	95.6	115.9	139.4	188.9
2000.....	16.4	26.6	40.4	53.8	66.0	79.6	97.3	118.8	143.1	205.7
1999.....	15.6	26.1	40.8	53.9	65.6	79.7	98.4	117.4	145.1	200.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of breast (C50)

White, both sexes

2006 4/.....	6.3	10.4	15.7	23.3	31.9	37.7	46.9	63.3	83.0	116.9
2005 4/.....	6.6	10.4	16.4	23.4	32.0	38.1	49.2	63.6	83.6	123.4
2004 4/.....	6.5	10.5	17.1	25.1	32.0	40.6	49.5	65.0	83.4	124.8
2003 4/.....	7.1	11.4	17.8	25.3	32.6	39.2	51.7	66.0	87.5	132.5
2002.....	7.0	11.7	18.5	25.0	33.1	40.8	52.0	67.5	89.0	135.3
2001.....	7.3	12.4	19.4	25.8	33.3	40.9	53.6	69.4	88.2	134.8
2000.....	7.5	12.4	19.7	26.9	34.4	43.3	55.5	71.1	91.7	147.3
1999.....	7.1	12.1	19.8	26.8	33.8	42.8	55.5	69.7	92.8	146.0

White, male

2006 4/.....	*	*	*	0.4	0.8	0.8	1.2	1.8	2.5	3.2
2005 4/.....	*	*	0.3	0.4	0.6	0.9	1.6	1.6	2.7	2.4
2004 4/.....	*	*	*	0.3	0.6	0.9	1.2	2.2	2.0	3.0
2003 4/.....	*	*	0.4	0.4	0.7	0.9	1.2	1.5	2.9	3.9
2002.....	*	*	0.3	0.4	0.6	0.6	1.3	1.7	2.7	3.3
2001.....	*	*	0.3	0.5	0.7	1.2	1.6	2.1	2.3	3.3
2000.....	*	*	0.3	0.6	0.7	1.1	1.7	2.2	2.3	3.9
1999.....	*	*	0.3	0.5	0.6	1.0	1.3	1.8	2.2	4.4

White, female

2006 4/.....	12.6	20.6	31.0	45.4	60.8	70.5	84.8	109.2	133.8	170.3
2005 4/.....	13.3	20.7	32.3	45.7	61.1	71.1	88.6	109.5	134.4	179.1
2004 4/.....	12.9	20.9	33.7	48.9	61.1	75.9	89.2	111.0	134.0	179.7
2003 4/.....	14.2	22.7	34.8	49.2	62.0	72.9	93.0	112.7	139.7	189.4
2002.....	13.9	23.2	36.3	48.6	63.1	76.2	93.3	114.7	142.0	192.6
2001.....	14.7	24.5	38.1	50.1	63.4	75.7	95.8	117.0	140.0	189.9
2000.....	15.0	24.4	38.7	51.8	65.4	80.2	98.6	119.7	144.9	205.5
1999.....	14.2	23.9	38.9	51.7	64.4	79.2	98.6	117.4	146.3	202.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of breast (C50)

Black, both sexes

2006 4/.....	12.5	21.3	27.7	39.5	47.4	49.8	59.1	78.1	104.6	142.2
2005 4/.....	13.6	23.1	31.8	43.6	47.7	53.8	61.5	75.5	91.5	144.3
2004 4/.....	13.7	21.2	31.8	40.0	48.9	51.2	62.8	77.0	90.1	143.5
2003 4/.....	14.1	22.6	31.2	43.4	48.1	54.6	65.1	80.3	106.4	161.6
2002.....	15.3	22.9	32.1	40.5	46.3	54.5	67.2	78.5	117.5	152.6
2001.....	14.8	22.6	33.1	42.4	47.4	58.0	70.4	83.5	105.9	152.0
2000.....	14.8	23.9	32.8	43.9	47.3	52.7	65.6	84.2	100.5	176.4
1999.....	14.6	24.2	34.7	44.4	48.6	57.8	69.9	88.0	104.9	147.3

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, female

2006 4/.....	23.5	39.6	50.4	71.3	83.6	85.5	98.3	123.7	157.7	199.2
2005 4/.....	25.4	43.0	58.4	79.1	83.9	91.8	101.5	117.3	139.5	201.8
2004 4/.....	25.6	39.5	58.1	72.4	85.8	88.1	103.7	121.6	135.7	196.9
2003 4/.....	26.3	42.0	57.3	78.3	84.7	94.1	107.4	127.2	160.5	222.4
2002.....	28.7	42.4	58.7	73.1	81.2	92.9	111.1	123.2	176.7	209.1
2001.....	27.5	41.9	60.5	76.7	83.7	99.5	115.0	130.9	157.8	205.9
2000.....	27.7	44.3	60.4	78.8	83.3	91.0	107.4	133.6	149.0	238.7
1999.....	27.5	44.9	63.3	79.6	85.2	99.5	114.1	139.3	154.7	199.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

All races 3/, both sexes

2006.....	1.7	2.1	2.4	2.5	2.7	3.1	3.4	3.6	4.1	5.2
2005.....	1.7	2.2	2.3	2.6	2.6	3.0	3.1	3.5	4.6	5.4
2004.....	1.7	2.1	2.0	2.7	2.9	3.2	3.3	3.3	4.3	5.1
2003.....	1.7	2.0	2.4	2.8	2.9	3.2	3.0	3.7	5.1	5.7
2002.....	1.7	2.1	2.4	2.7	2.8	3.3	3.6	4.1	4.2	6.2
2001.....	1.8	2.3	2.6	2.6	3.2	2.9	3.5	4.0	5.5	7.2
2000.....	1.8	2.2	2.7	3.1	3.1	3.6	3.9	4.3	5.6	7.3
1999.....	1.8	2.4	2.7	3.0	3.5	3.7	3.9	4.7	4.6	7.9

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	3.4	4.1	4.7	4.9	5.2	5.7	6.1	6.2	6.6	7.7
2005.....	3.4	4.3	4.6	5.1	5.0	5.7	5.7	6.0	7.5	7.8
2004.....	3.4	4.1	4.0	5.2	5.6	6.0	6.0	5.6	7.0	7.3
2003.....	3.3	4.0	4.6	5.5	5.5	5.9	5.4	6.3	8.3	8.3
2002.....	3.4	4.1	4.7	5.2	5.3	6.1	6.5	7.0	6.7	8.9
2001.....	3.7	4.5	5.1	5.1	6.0	5.4	6.3	6.8	8.9	10.1
2000.....	3.5	4.4	5.2	6.0	5.9	6.7	6.9	7.3	9.0	10.3
1999.....	3.6	4.6	5.2	5.8	6.7	6.8	6.9	8.0	7.2	11.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

White, both sexes

2006 4/.....	1.5	1.9	2.2	2.4	2.5	2.6	2.8	3.0	3.7	4.2
2005 4/.....	1.6	1.9	2.2	2.3	2.3	2.7	2.8	3.0	4.0	4.8
2004 4/.....	1.5	1.9	1.8	2.5	2.6	2.7	2.8	2.9	3.8	4.1
2003 4/.....	1.6	1.9	2.1	2.6	2.6	2.8	2.4	2.8	4.6	4.9
2002.....	1.6	1.8	2.2	2.5	2.4	2.7	3.0	3.6	3.5	5.1
2001.....	1.7	2.2	2.3	2.3	2.9	2.4	3.0	3.4	4.7	6.2
2000.....	1.6	2.0	2.4	2.7	2.6	2.9	3.3	3.8	5.0	6.3
1999.....	1.7	2.3	2.5	2.7	3.0	3.1	3.1	4.0	4.1	6.8

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	3.0	3.8	4.4	4.7	4.8	5.0	5.2	5.3	6.0	6.2
2005 4/.....	3.1	3.8	4.3	4.5	4.5	5.1	5.2	5.2	6.6	6.9
2004 4/.....	3.1	3.8	3.5	4.9	5.1	5.1	5.1	5.0	6.1	5.9
2003 4/.....	3.2	3.7	4.2	5.1	4.9	5.2	4.4	4.9	7.4	7.1
2002.....	3.1	3.7	4.3	4.9	4.5	5.2	5.4	6.2	5.7	7.3
2001.....	3.5	4.3	4.6	4.6	5.5	4.4	5.5	5.8	7.6	8.8
2000.....	3.1	4.0	4.7	5.3	5.0	5.5	5.9	6.5	8.0	8.8
1999.....	3.4	4.5	5.0	5.2	5.7	5.9	5.5	6.8	6.5	9.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

Black, both sexes

2006 4/.....	3.2	3.2	4.0	3.4	4.9	6.7	8.5	8.2	9.0	17.6
2005 4/.....	3.0	4.0	3.3	5.1	5.3	6.5	6.2	9.4	12.4	13.1
2004 4/.....	2.9	3.5	3.9	4.7	6.1	7.1	8.1	8.0	10.1	16.1
2003 4/.....	2.4	3.5	3.8	5.1	6.1	7.1	7.7	11.2	11.4	14.6
2002.....	3.0	3.7	4.6	4.2	6.0	8.2	10.1	9.6	12.6	17.5
2001.....	3.0	3.2	4.4	4.6	5.9	7.9	7.7	9.7	15.2	19.0
2000.....	3.1	3.7	4.9	6.2	6.4	9.4	9.7	10.8	13.6	19.7
1999.....	3.0	3.0	4.1	6.1	8.3	8.7	13.1	12.3	9.5	21.7

Black, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2006 4/.....	6.0	5.9	7.4	6.1	8.7	11.6	14.4	13.1	13.7	24.9
2005 4/.....	5.7	7.6	6.2	9.3	9.5	11.3	10.4	14.9	18.9	18.4
2004 4/.....	5.4	6.6	7.2	8.6	10.9	12.4	13.6	12.7	15.5	22.4
2003 4/.....	4.5	6.5	7.1	9.3	10.8	12.4	12.8	17.9	17.4	20.3
2002.....	5.6	7.0	8.4	7.6	10.7	14.2	16.8	15.4	19.1	24.2
2001.....	5.6	6.1	8.2	8.3	10.6	13.7	12.8	15.5	22.9	26.1
2000.....	5.9	7.0	9.1	11.2	11.4	16.3	16.2	17.4	20.4	27.0
1999.....	5.7	5.5	7.5	11.1	14.8	15.2	21.8	19.7	14.3	29.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All races 3/, both sexes

2006.....	0.4	1.0	1.9	3.8	6.2	8.8	11.6	14.3	17.6	21.6
2005.....	0.4	0.9	1.8	3.7	6.4	8.7	10.8	13.1	17.4	22.1
2004.....	0.5	0.9	1.8	3.4	5.8	8.8	11.5	13.8	17.7	22.3
2003.....	0.5	0.8	1.9	3.5	6.1	9.0	11.3	13.8	17.1	22.8
2002.....	0.4	0.9	2.0	3.5	5.8	8.4	11.8	14.2	17.0	24.6
2001.....	0.4	0.9	2.0	3.2	5.8	8.2	11.4	14.5	19.2	24.7
2000.....	0.4	0.9	1.8	3.5	6.0	8.4	11.1	14.0	18.6	24.7
1999.....	0.5	0.9	1.7	3.3	5.5	8.8	10.9	14.4	19.4	24.0

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	0.9	1.9	3.6	7.4	11.9	16.4	21.0	24.7	28.6	31.6
2005.....	0.8	1.8	3.5	7.2	12.1	16.4	19.6	22.6	28.2	32.2
2004.....	1.0	1.8	3.5	6.7	11.0	16.5	20.8	23.8	28.7	32.3
2003.....	1.0	1.6	3.7	6.8	11.7	16.8	20.4	23.5	27.6	33.0
2002.....	0.9	1.7	3.9	6.9	11.0	15.7	21.2	24.2	27.4	35.2
2001.....	0.8	1.8	4.0	6.2	11.1	15.3	20.4	24.6	30.7	35.0
2000.....	0.9	1.9	3.5	6.9	11.4	15.6	19.8	23.8	29.5	34.8
1999.....	0.9	1.8	3.3	6.3	10.5	16.2	19.5	24.5	30.7	33.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

White, both sexes

2006 4/.....	0.4	0.9	1.8	3.5	5.5	7.6	10.3	13.3	16.9	21.7
2005 4/.....	0.4	0.9	1.7	3.4	5.5	7.9	9.8	12.0	16.9	21.8
2004 4/.....	0.5	0.9	1.6	3.2	5.2	7.7	10.2	13.0	17.4	22.4
2003 4/.....	0.4	0.7	1.9	3.4	5.4	7.9	10.5	13.0	16.2	22.7
2002.....	0.4	0.8	1.9	3.2	5.2	7.2	10.8	13.1	16.3	23.8
2001.....	0.4	0.9	1.9	3.1	5.2	7.2	10.5	13.7	18.6	24.5
2000.....	0.4	0.9	1.8	3.4	5.4	7.3	9.9	13.0	17.5	24.4
1999.....	0.4	0.9	1.6	3.0	5.2	7.6	10.2	13.7	18.8	23.4

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	0.9	1.8	3.6	6.8	10.6	14.3	18.8	23.3	27.6	31.8
2005 4/.....	0.8	1.8	3.4	6.7	10.6	15.0	18.0	20.8	27.5	31.8
2004 4/.....	1.0	1.8	3.2	6.3	10.0	14.5	18.6	22.6	28.2	32.6
2003 4/.....	0.8	1.5	3.7	6.6	10.4	14.9	19.1	22.4	26.2	32.8
2002.....	0.9	1.7	3.7	6.3	10.1	13.6	19.6	22.4	26.3	34.2
2001.....	0.8	1.7	3.8	6.1	9.9	13.6	18.9	23.5	29.8	34.7
2000.....	0.8	1.9	3.5	6.7	10.3	13.6	17.8	22.2	28.0	34.3
1999.....	0.9	1.8	3.2	5.9	9.9	14.2	18.4	23.3	30.0	32.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Black, both sexes

2006 4/.....	*	1.4	2.1	6.9	13.9	20.8	25.8	27.6	30.8	25.9
2005 4/.....	*	1.3	2.5	6.1	15.1	18.0	22.7	27.6	27.4	31.3
2004 4/.....	0.7	1.0	3.0	5.2	11.9	20.2	26.4	26.6	25.5	26.2
2003 4/.....	0.9	1.1	2.5	5.3	13.9	19.9	22.4	25.5	31.2	30.0
2002.....	*	1.1	2.9	6.6	10.8	20.0	25.2	27.8	29.0	38.9
2001.....	*	1.4	2.9	4.6	11.6	18.8	24.2	26.6	31.1	32.2
2000.....	*	1.0	2.4	5.3	12.0	19.6	26.4	27.7	35.0	33.2
1999.....	0.8	*	2.2	5.8	10.5	20.9	21.8	25.7	29.8	35.4

Black, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2006 4/.....	*	2.7	3.8	12.7	24.8	36.2	43.5	44.2	47.0	36.6
2005 4/.....	*	2.5	4.7	11.2	26.9	31.4	38.3	44.0	41.9	43.9
2004 4/.....	1.4	1.9	5.5	9.4	21.1	35.3	44.4	42.3	38.9	36.6
2003 4/.....	1.6	2.1	4.7	9.6	24.8	34.7	37.4	40.7	47.6	41.7
2002.....	*	2.0	5.4	12.1	19.2	34.8	42.1	44.5	44.2	53.8
2001.....	*	2.6	5.4	8.3	20.8	32.7	40.4	42.4	47.0	44.2
2000.....	*	1.9	4.4	9.6	21.4	34.0	44.0	44.4	52.6	45.5
1999.....	1.6	*	4.1	10.5	18.8	36.4	36.2	41.2	44.5	48.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of ovary (C56)

All races 3/, both sexes

2006.....	1.3	2.8	5.1	7.7	12.4	16.3	22.0	28.3	34.8	37.8
2005.....	1.3	2.8	5.3	8.4	12.2	16.5	22.3	28.0	34.7	38.1
2004.....	1.4	2.9	5.2	7.9	11.9	17.8	22.5	28.9	36.4	37.7
2003.....	1.4	2.9	5.2	8.3	13.0	17.3	21.9	29.4	36.9	38.4
2002.....	1.4	3.2	5.5	8.3	12.9	17.3	22.3	30.3	37.8	38.1
2001.....	1.4	2.9	5.5	8.3	13.0	17.6	22.6	30.2	37.4	39.6
2000.....	1.5	3.1	5.5	8.3	12.2	17.5	21.8	30.9	36.5	40.9
1999.....	1.5	3.1	5.5	8.3	12.6	17.0	22.4	29.2	35.5	37.6

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	2.5	5.5	9.9	14.9	23.7	30.6	39.9	49.1	56.5	55.5
2005.....	2.7	5.5	10.3	16.3	23.4	30.9	40.4	48.3	56.3	55.6
2004.....	2.8	5.8	10.1	15.3	22.7	33.3	40.7	49.7	58.8	54.6
2003.....	2.8	5.7	10.2	16.2	24.8	32.3	39.5	50.4	59.5	55.4
2002.....	2.8	6.4	10.7	16.2	24.6	32.3	40.1	51.7	60.8	54.6
2001.....	2.8	5.8	10.8	16.0	24.7	32.8	40.5	51.4	59.8	56.1
2000.....	2.9	6.0	10.8	16.1	23.3	32.5	39.0	52.5	58.0	57.6
1999.....	3.0	6.2	10.8	16.0	24.0	31.6	39.9	49.5	56.2	52.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of ovary (C56)

White, both sexes

2006 4/.....	1.3	2.8	5.2	7.9	12.8	16.9	23.2	29.1	36.0	38.8
2005 4/.....	1.3	2.9	5.5	8.8	12.7	16.9	22.8	28.8	35.5	39.4
2004 4/.....	1.5	3.0	5.3	8.0	12.4	18.2	23.1	29.8	37.8	38.6
2003 4/.....	1.4	3.0	5.4	8.7	13.5	17.9	22.4	30.3	38.1	39.1
2002.....	1.4	3.3	5.8	8.6	13.5	18.0	22.9	31.1	38.8	39.0
2001.....	1.4	3.1	5.7	8.5	13.2	18.1	23.3	31.1	38.6	40.4
2000.....	1.5	3.2	5.6	8.7	12.7	18.0	22.4	32.0	37.6	42.0
1999.....	1.6	3.3	5.7	8.6	13.0	17.5	22.9	29.6	36.4	38.7

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	2.6	5.7	10.3	15.6	24.8	32.0	42.4	50.8	58.7	57.0
2005 4/.....	2.7	5.7	10.9	17.2	24.4	32.0	41.6	50.1	57.8	57.5
2004 4/.....	2.9	5.9	10.4	15.8	23.8	34.4	42.1	51.6	61.3	56.0
2003 4/.....	2.9	6.0	10.6	17.1	25.9	33.7	40.7	52.1	61.5	56.4
2002.....	2.9	6.6	11.4	16.8	25.9	33.9	41.6	53.4	62.5	55.9
2001.....	2.9	6.2	11.4	16.6	25.3	33.9	42.2	53.1	61.9	57.3
2000.....	3.0	6.4	11.1	17.0	24.5	33.7	40.3	54.5	60.0	59.0
1999.....	3.2	6.5	11.2	16.8	25.1	32.8	41.0	50.5	57.9	54.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of ovary (C56)

Black, both sexes

2006 4/.....	1.3	2.7	4.4	6.6	11.3	15.5	16.3	25.3	24.8	33.5
2005 4/.....	1.4	2.3	4.2	6.9	11.1	15.6	22.4	25.2	31.1	31.3
2004 4/.....	1.1	3.0	4.9	7.5	10.2	18.3	22.5	26.2	29.0	32.7
2003 4/.....	1.2	2.8	5.1	6.8	11.0	15.3	20.6	26.2	29.4	36.8
2002.....	1.2	2.9	4.1	7.7	10.9	14.9	20.0	26.5	32.4	34.7
2001.....	1.4	1.9	4.4	7.9	12.6	16.3	20.7	26.2	27.9	36.5
2000.....	1.5	2.2	4.8	6.4	10.9	16.8	19.6	26.1	27.4	34.4
1999.....	1.1	2.4	5.0	7.2	11.4	15.0	22.7	28.9	28.3	28.7

Black, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2006 4/.....	2.4	5.0	8.1	12.1	20.1	27.0	27.5	40.4	38.0	47.2
2005 4/.....	2.7	4.3	7.8	12.5	19.7	27.1	37.7	40.3	47.5	43.9
2004 4/.....	2.0	5.7	9.1	13.7	18.1	31.8	37.8	41.8	44.2	45.7
2003 4/.....	2.3	5.3	9.4	12.5	19.6	26.7	34.6	41.8	44.9	51.0
2002.....	2.3	5.3	7.7	13.9	19.5	25.9	33.4	42.3	49.3	48.1
2001.....	2.7	3.6	8.1	14.4	22.6	28.3	34.5	41.9	42.1	50.1
2000.....	2.9	4.0	8.8	11.6	19.4	29.2	32.6	41.9	41.2	47.2
1999.....	2.1	4.5	9.3	13.2	20.3	26.2	37.8	46.3	42.3	39.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All races 3/, both sexes

2006.....	0.8	2.0	3.7	6.5	9.6	14.8	18.2	24.0	28.9	34.3
2005.....	0.9	2.0	4.1	6.4	10.8	14.4	19.9	24.3	30.3	32.6
2004.....	0.8	2.1	4.0	6.8	11.3	14.6	19.1	24.5	29.8	32.7
2003.....	1.0	2.0	4.0	7.1	11.4	14.5	19.6	24.8	30.0	33.2
2002.....	0.9	2.4	4.3	7.2	10.9	16.0	19.6	24.6	28.7	32.5
2001.....	1.2	2.4	4.4	7.1	11.9	14.6	20.8	24.4	29.1	32.3
2000.....	1.1	2.2	4.5	7.6	10.8	15.0	19.4	24.5	29.6	33.8
1999.....	1.1	2.2	4.3	7.3	10.8	14.4	19.2	23.1	26.9	31.4

All races 3/, male

2006.....	1.2	2.8	5.4	9.5	14.0	21.6	25.0	34.5	40.8	52.1
2005.....	1.2	2.9	5.8	9.3	15.7	20.7	28.9	34.1	43.3	50.6
2004.....	1.2	2.9	5.9	9.8	16.6	20.7	27.8	34.2	44.0	47.0
2003.....	1.5	3.0	5.7	10.4	17.2	20.2	27.7	35.9	44.9	48.7
2002.....	1.3	3.4	6.4	10.6	16.1	23.0	27.6	36.6	39.9	50.1
2001.....	1.6	3.5	6.5	10.5	17.2	21.0	30.0	34.4	42.2	52.9
2000.....	1.6	3.1	6.6	11.2	15.6	21.4	27.4	34.6	44.2	55.1
1999.....	1.5	3.2	6.1	10.6	15.8	20.3	27.3	33.7	38.9	49.4

All races 3/, female

2006.....	0.4	1.2	2.1	3.6	5.6	8.8	12.6	16.4	21.5	25.9
2005.....	0.7	1.2	2.4	3.6	6.4	8.8	12.5	17.2	22.2	24.3
2004.....	0.5	1.3	2.2	3.8	6.4	9.2	12.1	17.6	21.0	26.3
2003.....	0.6	1.0	2.3	4.0	6.1	9.4	13.0	16.9	20.9	26.4
2002.....	0.5	1.5	2.3	3.9	6.2	10.0	13.2	16.2	21.9	24.9
2001.....	0.9	1.4	2.4	3.9	7.0	9.1	13.5	17.4	21.2	23.7
2000.....	0.7	1.4	2.4	4.3	6.5	9.5	13.0	17.5	21.0	25.1
1999.....	0.6	1.2	2.5	4.3	6.2	9.3	12.8	15.7	19.9	24.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of kidney and renal pelvis (C64-C65)

White, both sexes

2006 4/.....	0.8	2.0	3.7	6.7	10.0	15.3	18.7	24.9	29.6	34.9
2005 4/.....	0.9	2.0	4.2	6.5	11.0	14.8	20.6	24.9	31.1	33.1
2004 4/.....	0.8	2.2	4.2	7.0	11.7	15.1	19.4	24.9	30.6	33.7
2003 4/.....	1.0	2.0	4.2	7.2	11.6	15.0	20.2	25.4	30.8	33.8
2002.....	1.0	2.4	4.4	7.4	11.2	16.6	20.2	25.0	29.2	33.2
2001.....	1.2	2.5	4.4	7.2	12.1	15.3	21.3	24.7	29.5	32.8
2000.....	1.1	2.3	4.6	7.8	11.1	15.4	19.7	24.9	30.0	34.5
1999.....	1.1	2.1	4.3	7.5	11.2	14.8	19.4	23.4	27.3	31.5

White, male

2006 4/.....	1.2	2.8	5.4	9.5	14.4	22.0	25.7	35.7	41.8	53.4
2005 4/.....	1.2	2.8	5.9	9.5	15.7	21.3	29.6	34.8	44.2	52.0
2004 4/.....	1.1	3.0	6.2	10.2	17.2	21.3	28.2	34.5	45.4	48.6
2003 4/.....	1.5	3.1	5.9	10.7	17.5	20.8	28.2	36.5	46.2	50.0
2002.....	1.4	3.5	6.6	10.9	16.4	23.8	28.3	37.5	40.2	51.9
2001.....	1.6	3.6	6.5	10.8	17.2	21.9	30.6	34.4	42.8	54.4
2000.....	1.6	3.2	6.7	11.3	16.1	21.5	27.7	35.1	44.5	56.6
1999.....	1.5	3.1	6.0	10.7	16.4	20.6	27.6	34.1	40.1	49.8

White, female

2006 4/.....	0.4	1.3	2.0	3.9	5.8	9.3	12.9	16.9	22.0	26.2
2005 4/.....	0.7	1.2	2.6	3.7	6.7	9.1	13.1	17.6	22.9	24.3
2004 4/.....	0.5	1.3	2.3	4.0	6.7	9.6	12.2	17.8	21.3	27.0
2003 4/.....	0.6	0.9	2.5	3.8	6.2	9.7	13.7	17.4	21.3	26.6
2002.....	0.5	1.4	2.2	4.1	6.4	10.3	13.6	16.1	22.5	25.1
2001.....	0.8	1.3	2.4	3.8	7.4	9.5	13.8	17.8	21.4	23.8
2000.....	0.7	1.4	2.5	4.5	6.5	10.0	13.3	17.7	21.4	25.5
1999.....	0.6	1.1	2.6	4.4	6.3	9.7	13.0	15.8	19.8	24.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Black, both sexes

2006 4/.....	1.3	2.4	4.9	6.7	8.4	13.2	16.7	19.3	26.0	33.5
2005 4/.....	1.1	2.4	4.4	6.6	12.1	13.5	17.5	22.2	26.2	30.5
2004 4/.....	1.2	2.3	4.0	6.2	10.5	13.1	19.2	26.2	25.3	28.2
2003 4/.....	1.5	2.5	4.1	8.2	11.2	13.4	16.1	22.3	26.1	33.8
2002.....	0.9	2.8	4.4	7.0	10.6	14.3	16.8	25.8	26.7	28.2
2001.....	1.8	2.6	5.0	7.4	12.2	11.7	19.2	24.0	28.2	29.8
2000.....	1.1	2.5	4.7	7.4	10.4	14.8	18.5	23.5	29.1	29.7
1999.....	1.2	2.9	4.9	7.8	9.7	13.5	19.3	23.0	24.3	31.2

Black, male

2006 4/.....	2.1	3.8	6.8	11.1	13.4	22.0	23.1	27.3	39.7	49.1
2005 4/.....	1.9	3.6	7.1	10.0	19.6	20.0	29.1	32.2	39.3	39.0
2004 4/.....	1.6	3.2	6.1	9.2	16.1	19.6	29.8	38.2	35.9	38.7
2003 4/.....	2.5	3.6	6.5	11.5	17.0	19.5	25.9	35.6	39.1	43.8
2002.....	*	3.5	6.4	11.8	16.8	19.9	23.7	34.4	39.0	39.7
2001.....	1.9	3.5	7.8	10.5	20.5	17.0	29.3	39.0	42.5	46.5
2000.....	*	3.2	7.1	11.4	15.3	25.0	28.1	32.1	48.9	49.6
1999.....	1.8	4.1	8.0	12.5	14.0	20.0	28.2	34.2	27.9	49.2

Black, female

2006 4/.....	*	*	3.3	3.0	4.5	6.8	12.3	14.5	18.8	27.1
2005 4/.....	*	*	2.1	3.8	6.2	8.6	9.5	16.3	19.3	27.0
2004 4/.....	*	1.5	2.2	3.7	6.1	8.3	12.1	19.1	19.7	24.0
2003 4/.....	*	1.5	2.1	5.5	6.6	9.0	9.5	14.4	19.3	30.0
2002.....	*	2.3	2.6	3.1	5.7	10.2	12.1	20.6	20.3	23.8
2001.....	1.7	1.8	2.7	4.9	5.6	7.7	12.6	15.0	20.8	23.6
2000.....	*	1.8	2.6	4.2	6.5	7.3	12.2	18.2	19.1	22.3
1999.....	*	1.9	2.3	3.9	6.3	8.7	13.5	16.3	22.5	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of bladder (C67)

All races 3/, both sexes

2006.....	0.3	0.8	1.7	3.5	7.0	11.2	19.2	30.0	46.9	67.4
2005.....	0.4	0.9	1.6	3.5	6.8	11.5	18.9	30.0	46.7	68.3
2004.....	0.4	0.9	1.8	3.6	6.8	11.5	18.9	31.1	45.7	67.9
2003.....	0.3	0.8	1.8	3.5	6.9	11.8	18.5	30.0	45.1	64.8
2002.....	0.4	0.9	1.7	3.8	6.4	12.0	20.2	31.9	44.1	66.6
2001.....	0.4	0.8	1.7	3.4	7.0	12.0	19.0	29.3	44.8	68.9
2000.....	0.3	0.8	1.6	3.6	6.8	11.7	19.6	29.1	45.5	69.6
1999.....	0.3	0.7	1.9	3.8	7.0	11.8	20.2	29.7	43.8	68.5

All races 3/, male

2006.....	0.5	1.1	2.6	5.5	11.1	17.7	31.4	51.4	85.8	131.6
2005.....	0.5	1.2	2.2	5.3	10.8	18.7	31.2	50.4	86.3	132.1
2004.....	0.5	1.3	2.8	5.7	10.8	18.5	30.6	53.1	81.6	133.0
2003.....	0.4	1.4	2.7	5.5	10.9	19.1	29.6	51.7	80.3	125.1
2002.....	0.5	1.2	2.4	5.9	10.1	19.0	32.2	54.0	76.6	128.9
2001.....	0.5	1.2	2.6	5.2	11.3	19.5	31.6	50.8	81.0	134.8
2000.....	0.4	1.1	2.5	5.7	10.6	19.0	32.2	50.6	82.4	139.4
1999.....	0.4	1.1	2.9	5.9	11.0	19.3	32.9	52.3	79.5	134.8

All races 3/, female

2006.....	*	0.5	0.8	1.6	3.3	5.6	9.4	14.4	22.6	37.4
2005.....	0.3	0.6	1.0	1.7	3.1	5.2	8.9	15.2	22.1	39.0
2004.....	0.3	0.5	0.8	1.7	3.1	5.3	9.5	15.3	23.5	38.6
2003.....	0.2	0.3	0.9	1.7	3.3	5.4	9.5	14.6	23.5	38.1
2002.....	0.3	0.5	1.0	1.9	2.9	6.0	10.6	16.3	24.2	39.6
2001.....	0.3	0.4	0.9	1.7	3.2	5.6	9.0	14.3	23.2	41.4
2000.....	0.3	0.5	0.8	1.7	3.4	5.5	9.7	14.0	23.8	41.2
1999.....	0.2	0.3	0.9	1.7	3.4	5.4	10.3	13.9	22.9	41.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of bladder (C67)

White, both sexes

2006 4/.....	0.3	0.8	1.8	3.6	7.4	11.8	20.0	31.3	48.6	69.8
2005 4/.....	0.4	0.9	1.6	3.6	7.1	12.0	19.8	31.1	48.7	70.0
2004 4/.....	0.3	0.9	1.8	3.8	7.3	12.3	19.8	32.6	47.6	69.6
2003 4/.....	0.3	0.8	1.8	3.7	7.3	12.2	19.6	31.1	46.2	66.6
2002.....	0.4	0.9	1.8	3.9	6.7	12.5	20.9	32.9	45.2	67.7
2001.....	0.4	0.8	1.8	3.5	7.3	12.6	19.8	30.3	45.7	70.8
2000.....	0.3	0.8	1.7	3.8	7.2	12.3	20.2	29.7	46.4	72.0
1999.....	0.3	0.7	2.0	3.8	7.2	12.1	20.8	30.5	45.0	69.8

White, male

2006 4/.....	0.5	1.2	2.7	5.7	11.6	18.8	33.1	54.2	90.0	137.9
2005 4/.....	0.5	1.4	2.3	5.6	11.3	19.9	32.8	52.5	90.5	137.4
2004 4/.....	0.4	1.3	2.9	6.0	11.6	20.2	32.3	55.9	86.2	139.3
2003 4/.....	0.4	1.4	2.8	5.7	11.7	19.8	31.6	53.8	83.5	131.0
2002.....	0.5	1.3	2.6	6.2	10.8	20.1	33.7	56.0	79.4	134.1
2001.....	0.5	1.2	2.7	5.4	12.0	20.7	33.1	53.1	83.7	141.1
2000.....	0.4	1.2	2.7	6.0	11.4	20.2	33.6	52.7	84.6	145.7
1999.....	0.4	1.1	3.0	6.2	11.4	20.1	34.4	54.3	82.1	140.7

White, female

2006 4/.....	*	0.5	0.9	1.6	3.4	5.6	9.0	14.1	22.5	37.9
2005 4/.....	0.3	0.5	0.9	1.6	3.2	5.1	9.1	15.3	22.4	39.0
2004 4/.....	0.3	0.5	0.7	1.7	3.3	5.3	9.5	15.6	23.7	38.1
2003 4/.....	*	0.2	0.8	1.7	3.3	5.5	9.8	14.6	23.3	38.1
2002.....	0.3	0.5	1.0	1.7	2.9	5.9	10.4	16.4	24.3	38.9
2001.....	0.3	0.3	0.9	1.7	3.0	5.5	9.0	14.2	22.7	41.3
2000.....	0.3	0.5	0.8	1.8	3.4	5.5	9.6	13.6	23.7	42.1
1999.....	0.2	0.3	1.0	1.5	3.4	5.1	10.0	13.8	23.2	41.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasm of bladder (C67)

Black, both sexes

2006 4/.....	*	0.9	1.7	3.8	6.2	9.7	18.2	25.6	38.6	54.5
2005 4/.....	*	1.0	2.0	3.7	5.9	9.7	15.7	25.1	36.2	57.5
2004 4/.....	*	1.2	2.0	3.3	5.1	7.9	16.5	23.0	33.4	60.1
2003 4/.....	*	1.2	2.2	3.2	5.0	11.4	12.7	26.5	40.9	53.4
2002.....	*	*	1.6	4.0	5.2	10.9	18.6	27.0	37.2	63.3
2001.....	*	0.9	1.7	3.4	6.9	10.0	14.1	24.8	41.2	55.0
2000.....	*	0.8	1.3	3.0	5.4	9.5	17.9	28.4	44.6	49.7
1999.....	*	1.0	2.2	4.5	6.9	11.8	18.9	25.4	35.3	60.3

Black, male

2006 4/.....	*	*	2.8	5.6	10.4	12.4	24.9	32.7	56.1	92.9
2005 4/.....	*	*	2.7	4.9	9.1	13.1	25.0	38.0	56.9	101.3
2004 4/.....	*	1.7	3.0	4.4	8.0	8.2	23.0	35.0	46.6	81.5
2003 4/.....	*	1.7	2.8	4.5	6.6	18.2	16.4	41.1	61.3	80.3
2002.....	*	*	*	4.5	7.1	14.2	23.7	41.8	57.3	91.1
2001.....	*	*	2.8	4.2	9.2	12.7	20.5	34.4	57.8	78.3
2000.....	*	*	*	4.6	7.2	12.5	24.1	41.1	73.3	88.8
1999.....	*	*	3.9	5.4	9.8	15.6	23.7	38.9	59.3	86.7

Black, female

2006 4/.....	*	*	*	2.4	2.9	7.6	13.5	21.4	29.3	38.7
2005 4/.....	*	*	*	2.7	3.4	7.2	9.3	17.3	25.2	39.8
2004 4/.....	*	*	*	2.5	*	7.7	12.1	15.9	26.5	51.6
2003 4/.....	*	*	1.7	*	3.8	6.4	10.2	17.9	30.2	42.9
2002.....	*	*	*	3.7	3.7	8.5	15.2	18.1	26.7	52.6
2001.....	*	*	*	2.7	5.1	8.1	9.9	19.1	32.7	46.3
2000.....	*	*	*	*	3.9	7.3	13.8	20.8	30.1	35.2
1999.....	*	*	*	3.7	4.7	9.0	15.8	17.2	23.3	50.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes

2006.....	2.3	3.5	5.7	8.1	10.8	13.0	16.8	18.5	19.6	16.7
2005.....	2.7	3.7	5.6	8.3	10.7	14.0	18.0	20.0	20.4	15.8
2004.....	2.4	3.8	5.5	8.4	10.9	14.0	17.7	19.0	20.4	16.6
2003.....	2.9	3.9	5.8	8.5	11.2	13.8	17.3	19.4	19.6	15.2
2002.....	2.8	4.2	6.2	8.5	11.1	14.7	17.8	19.6	18.5	16.4
2001.....	2.8	4.0	6.1	8.3	11.6	14.2	16.7	20.3	20.6	16.4
2000.....	2.9	4.1	6.3	8.2	11.4	14.8	18.3	19.4	21.1	16.9
1999.....	2.8	4.3	6.1	9.2	11.7	15.7	18.3	21.0	19.4	16.0

All races 3/, male

2006.....	2.9	4.3	6.8	9.9	13.5	16.4	20.4	22.9	25.6	21.2
2005.....	3.5	4.3	7.0	10.0	13.1	17.7	21.2	25.3	25.3	19.3
2004.....	2.9	4.7	7.1	10.4	13.7	16.0	21.3	23.1	24.5	19.4
2003.....	3.4	5.1	7.4	9.7	13.7	16.4	21.3	24.0	25.4	20.0
2002.....	3.5	5.3	7.2	10.6	13.1	18.4	22.2	24.4	22.9	20.1
2001.....	3.5	4.9	7.5	10.7	13.9	17.3	21.2	25.4	26.5	20.1
2000.....	3.7	4.9	7.7	10.1	14.0	17.9	22.8	23.2	26.3	22.4
1999.....	3.4	5.2	7.4	10.6	14.7	19.5	22.3	26.1	24.5	20.7

All races 3/, female

2006.....	1.8	2.7	4.6	6.4	8.4	10.0	13.9	15.3	15.8	14.6
2005.....	1.9	3.0	4.3	6.8	8.6	10.7	15.4	16.1	17.4	14.3
2004.....	1.9	2.8	4.0	6.4	8.4	12.2	14.7	16.0	17.9	15.4
2003.....	2.4	2.8	4.3	7.3	9.0	11.6	14.1	16.2	16.0	13.1
2002.....	2.1	3.1	5.1	6.5	9.2	11.4	14.2	16.2	15.8	14.8
2001.....	2.1	3.1	4.8	6.0	9.5	11.5	13.1	16.7	17.1	14.8
2000.....	2.0	3.3	4.9	6.5	9.0	12.1	14.8	16.7	18.0	14.7
1999.....	2.3	3.4	4.9	7.9	8.9	12.4	15.1	17.5	16.5	14.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

White, both sexes

2006 4/.....	2.6	3.9	6.3	8.7	11.9	14.0	18.1	20.0	20.6	17.4
2005 4/.....	3.0	4.0	6.2	9.1	11.6	15.2	19.4	21.1	21.5	16.5
2004 4/.....	2.6	4.1	6.0	9.2	11.9	15.2	19.1	20.2	21.7	17.3
2003 4/.....	3.3	4.3	6.3	9.1	12.2	14.9	18.5	20.6	20.7	15.7
2002.....	3.1	4.7	6.7	9.1	12.0	15.9	19.0	20.7	19.3	17.1
2001.....	3.0	4.4	6.6	8.8	12.6	15.2	17.8	21.6	21.6	17.0
2000.....	3.1	4.5	6.9	8.8	12.3	16.0	19.7	20.5	21.8	17.5
1999.....	3.0	4.7	6.6	10.0	12.5	16.9	19.6	22.2	20.3	16.6

White, male

2006 4/.....	3.3	4.7	7.5	10.5	14.8	17.7	21.8	24.6	26.6	22.1
2005 4/.....	3.9	4.6	7.6	10.7	14.0	19.3	22.5	26.3	26.5	20.2
2004 4/.....	3.1	5.1	7.6	11.3	14.8	17.4	23.1	24.2	25.8	20.6
2003 4/.....	3.8	5.5	8.0	10.3	14.8	17.4	22.6	25.3	26.6	20.7
2002.....	3.9	5.9	7.9	11.2	14.0	19.9	23.6	25.6	23.7	21.2
2001.....	3.8	5.4	8.1	11.4	14.9	18.6	22.6	26.7	27.8	20.5
2000.....	4.0	5.4	8.3	10.9	15.2	19.4	24.3	24.3	27.5	23.5
1999.....	3.6	5.7	7.9	11.4	15.7	20.9	23.9	27.4	25.8	21.8

White, female

2006 4/.....	1.9	3.0	5.1	7.0	9.3	10.8	15.0	16.6	16.8	15.3
2005 4/.....	2.1	3.3	4.8	7.4	9.4	11.5	16.7	17.3	18.4	14.8
2004 4/.....	2.1	3.1	4.4	7.2	9.1	13.2	15.7	17.2	19.2	15.8
2003 4/.....	2.7	3.0	4.8	7.9	9.8	12.7	15.2	17.3	17.0	13.4
2002.....	2.3	3.5	5.5	7.1	10.2	12.4	15.3	17.2	16.7	15.4
2001.....	2.3	3.4	5.2	6.4	10.5	12.2	13.9	18.0	17.9	15.6
2000.....	2.1	3.5	5.5	6.9	9.7	13.0	16.0	17.7	18.5	15.0
1999.....	2.5	3.8	5.4	8.7	9.7	13.4	16.2	18.5	17.1	14.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Black, both sexes

2006 4/.....	1.3	1.9	3.2	5.4	5.0	7.0	9.3	7.4	12.3	9.6
2005 4/.....	1.3	2.4	3.1	5.2	5.6	7.2	9.8	12.9	10.9	9.4
2004 4/.....	1.8	2.2	3.5	4.1	6.0	7.1	9.2	10.0	9.9	10.4
2003 4/.....	1.6	2.7	3.6	5.5	5.6	8.3	9.7	10.6	10.4	10.5
2002.....	1.3	1.9	3.7	5.7	5.6	8.0	9.5	12.2	11.3	9.5
2001.....	1.6	2.6	3.4	5.4	6.4	9.0	9.4	9.4	10.8	11.7
2000.....	2.0	2.2	3.7	5.4	6.0	7.5	8.8	10.3	14.4	10.6
1999.....	1.8	2.6	4.0	5.3	6.7	8.5	7.8	12.0	11.6	9.9

Black, male

2006 4/.....	1.5	2.4	4.0	7.3	6.5	8.4	12.3	*	19.2	*
2005 4/.....	1.5	2.9	4.5	6.6	7.5	8.1	13.3	19.2	14.7	*
2004 4/.....	2.8	2.9	5.0	5.9	7.8	7.8	8.1	14.1	14.3	*
2003 4/.....	2.4	3.1	5.1	6.6	6.7	11.8	12.2	16.0	*	*
2002.....	*	2.1	3.3	8.2	7.9	9.5	11.7	15.6	17.6	*
2001.....	1.9	3.0	4.4	6.9	9.2	9.1	11.4	12.5	*	*
2000.....	2.7	2.2	5.7	6.6	7.7	7.8	12.4	11.8	*	*
1999.....	2.6	3.7	5.3	5.9	8.7	10.9	8.8	14.2	*	*

Black, female

2006 4/.....	*	1.5	2.5	3.9	3.9	5.9	7.2	7.0	8.7	9.1
2005 4/.....	*	2.0	1.9	4.0	4.1	6.5	7.4	9.1	8.9	9.8
2004 4/.....	*	1.6	2.2	2.6	4.5	6.7	9.9	7.5	7.6	11.4
2003 4/.....	*	2.3	2.2	4.6	4.7	5.7	8.0	7.3	8.5	11.7
2002.....	*	1.8	4.0	3.7	3.7	6.8	8.1	10.2	8.0	10.3
2001.....	1.3	2.3	2.6	4.1	4.3	8.8	8.1	7.5	10.2	9.7
2000.....	1.5	2.2	2.1	4.5	4.7	7.3	6.4	9.4	14.0	10.7
1999.....	*	1.6	2.8	4.8	5.2	6.7	7.2	10.6	12.5	9.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes

2006.....	4.2	6.3	11.0	18.5	32.4	51.7	81.7	119.0	162.2	186.1
2005.....	4.1	6.6	11.0	19.0	33.2	52.9	83.8	121.4	161.5	189.8
2004.....	4.1	6.7	11.6	20.2	33.1	54.5	84.2	122.7	162.6	189.4
2003.....	4.7	7.2	11.7	20.1	35.5	55.2	87.3	127.4	165.2	198.0
2002.....	4.6	7.5	12.6	21.5	36.9	59.2	90.1	125.6	168.8	197.5
2001.....	4.8	8.1	13.0	22.7	38.1	58.8	91.8	129.0	168.5	199.9
2000.....	4.8	8.3	13.4	23.4	38.2	62.2	91.6	131.4	176.0	204.6
1999.....	4.8	8.4	13.6	25.6	39.4	62.8	93.9	130.7	174.0	199.4

All races 3/, male

2006.....	5.1	7.4	14.0	23.4	41.2	65.7	103.4	157.6	219.4	244.7
2005.....	4.9	7.8	13.3	23.9	40.7	68.1	109.5	161.0	214.7	262.4
2004.....	5.0	8.1	14.4	24.9	41.7	70.0	108.7	158.6	214.5	254.9
2003.....	5.8	8.8	14.5	25.2	45.6	71.9	112.3	163.5	216.8	268.7
2002.....	5.5	9.1	15.1	26.8	45.7	76.4	116.8	164.3	224.2	269.1
2001.....	5.9	9.8	16.0	27.7	47.5	75.2	116.9	165.7	220.9	275.3
2000.....	5.7	10.3	16.1	28.4	45.8	78.3	117.7	169.8	230.6	281.8
1999.....	5.9	10.2	16.7	31.0	48.6	78.1	118.9	169.4	221.8	279.0

All races 3/, female

2006.....	3.3	5.2	8.2	13.9	24.3	39.4	64.0	90.8	126.3	158.7
2005.....	3.3	5.5	8.7	14.4	26.4	39.6	63.0	92.7	128.4	156.4
2004.....	3.2	5.3	9.0	15.7	25.2	40.9	64.5	96.8	130.6	160.0
2003.....	3.5	5.5	9.1	15.3	26.4	40.8	67.3	101.7	133.6	166.8
2002.....	3.7	6.0	10.1	16.4	28.9	44.4	68.8	98.2	135.1	166.5
2001.....	3.7	6.4	10.1	18.1	29.5	44.6	71.8	103.4	137.1	168.4
2000.....	3.9	6.5	10.8	18.8	31.3	48.4	71.0	104.7	143.8	173.1
1999.....	3.8	6.6	10.6	20.5	31.0	49.8	74.3	103.8	145.9	167.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

White, both sexes

2006 4/.....	4.1	5.9	10.9	18.2	32.0	52.4	83.5	122.3	168.0	192.2
2005 4/.....	3.9	6.3	10.5	18.7	33.3	53.6	85.6	125.4	167.1	197.3
2004 4/.....	4.0	6.5	11.2	19.9	33.1	55.0	85.0	125.4	167.2	194.9
2003 4/.....	4.6	7.0	11.3	20.1	35.8	56.1	88.7	130.3	169.5	204.5
2002.....	4.6	7.2	12.1	21.0	36.9	60.3	91.7	128.7	173.6	204.0
2001.....	4.6	7.9	12.8	22.3	38.0	59.0	93.7	131.9	172.8	205.9
2000.....	4.7	8.0	13.2	23.3	38.5	63.5	93.4	134.4	181.5	208.7
1999.....	4.7	8.3	13.4	25.3	39.2	63.8	95.6	133.1	178.4	204.1

White, male

2006 4/.....	4.9	6.9	13.7	22.9	40.7	66.4	105.6	161.6	227.7	252.0
2005 4/.....	4.7	7.4	12.7	23.5	41.1	69.4	111.8	165.6	221.9	275.3
2004 4/.....	5.0	7.8	13.9	24.8	41.9	70.4	109.7	162.3	221.1	264.0
2003 4/.....	5.7	8.7	13.9	25.1	46.3	73.6	115.2	168.7	224.3	279.5
2002.....	5.5	8.8	14.8	26.5	45.7	77.9	119.2	169.0	231.0	279.4
2001.....	5.7	9.7	15.8	27.1	47.3	75.6	119.2	169.8	226.4	284.8
2000.....	5.8	9.9	15.6	28.3	46.1	80.4	119.0	174.1	238.0	290.2
1999.....	5.6	10.0	16.5	30.6	48.6	79.4	121.0	172.1	228.5	285.8

White, female

2006 4/.....	3.3	4.9	8.1	13.6	23.9	39.9	65.1	93.1	130.3	164.1
2005 4/.....	3.0	5.3	8.3	14.1	26.1	39.5	64.0	95.7	132.7	161.4
2004 4/.....	3.0	5.2	8.6	15.2	25.0	41.4	64.6	98.3	133.7	163.8
2003 4/.....	3.4	5.4	8.8	15.3	26.1	40.7	67.0	102.6	135.8	171.3
2002.....	3.7	5.7	9.6	15.8	28.8	44.8	69.4	99.9	138.5	171.2
2001.....	3.5	6.1	9.8	17.7	29.3	44.4	73.0	105.0	140.4	172.8
2000.....	3.6	6.1	10.7	18.6	31.4	48.7	72.8	106.5	147.9	175.5
1999.....	3.8	6.6	10.3	20.2	30.5	50.3	75.5	105.7	148.8	171.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Black, both sexes

2006 4/.....	6.0	10.0	14.6	26.0	42.9	57.3	83.4	109.2	127.8	143.5
2005 4/.....	5.8	9.2	16.0	24.9	38.7	58.5	81.2	103.1	127.4	136.8
2004 4/.....	5.2	9.1	16.8	26.3	40.8	60.7	93.3	115.8	134.7	154.0
2003 4/.....	6.3	9.5	17.3	23.5	41.1	59.4	90.5	117.4	139.1	150.5
2002.....	5.7	10.5	17.3	29.8	43.9	62.3	93.2	111.8	138.2	152.3
2001.....	6.6	10.4	16.5	30.2	44.9	67.8	90.7	116.7	142.7	155.7
2000.....	6.3	12.5	17.5	29.1	44.0	62.2	90.5	114.3	131.6	175.5
1999.....	6.5	10.4	18.2	33.2	45.9	64.6	90.4	119.1	139.3	162.0

Black, male

2006 4/.....	8.0	12.2	19.2	34.4	56.0	76.2	110.7	144.5	167.7	194.8
2005 4/.....	6.8	11.9	19.9	33.1	46.7	73.4	108.2	139.0	172.1	164.5
2004 4/.....	5.9	11.3	21.4	31.5	51.2	80.4	122.7	139.2	168.6	193.7
2003 4/.....	8.0	11.9	22.5	32.0	51.1	73.4	105.1	137.8	166.8	184.5
2002.....	7.2	13.1	19.7	36.0	55.6	78.8	123.2	137.8	177.3	202.6
2001.....	8.2	11.9	21.0	37.5	56.9	87.9	114.1	141.2	191.0	206.5
2000.....	6.6	14.9	21.8	36.7	54.2	73.1	125.4	147.9	167.8	224.9
1999.....	8.4	13.3	22.5	41.2	55.2	80.7	114.6	159.1	164.8	220.3

Black, female

2006 4/.....	4.2	8.0	10.7	19.0	32.6	43.4	64.7	88.1	106.7	122.5
2005 4/.....	5.0	6.7	12.8	18.2	32.4	47.4	62.8	81.6	103.9	125.7
2004 4/.....	4.6	7.2	12.9	21.9	32.8	46.1	73.4	101.9	116.8	138.2
2003 4/.....	4.8	7.4	12.9	16.6	33.2	49.0	80.6	105.3	124.6	137.3
2002.....	4.3	8.3	15.2	24.7	34.7	50.1	73.0	96.3	117.8	133.1
2001.....	5.3	9.2	12.7	24.2	35.4	52.9	75.1	102.1	118.1	136.8
2000.....	6.0	10.4	13.9	22.9	36.0	54.1	67.3	94.0	113.4	157.1
1999.....	4.8	8.0	14.5	26.6	38.6	52.7	74.3	95.0	126.6	140.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hodgkin's disease (C81)

All races 3/, both sexes

2006.....	0.4	0.3	0.4	0.5	0.7	0.9	1.3	2.1	2.3	2.6
2005.....	0.3	0.4	0.4	0.5	0.7	1.0	1.2	2.2	2.5	2.1
2004.....	0.3	0.4	0.4	0.7	0.8	1.0	1.4	1.8	2.1	2.2
2003.....	0.3	0.3	0.4	0.6	0.8	0.9	1.5	2.1	2.3	2.4
2002.....	0.3	0.4	0.5	0.5	0.8	1.0	1.6	1.7	2.3	2.7
2001.....	0.4	0.4	0.5	0.4	0.7	1.1	1.2	2.0	2.5	2.2
2000.....	0.3	0.4	0.5	0.5	0.9	1.0	1.6	1.8	2.4	2.5
1999.....	0.3	0.4	0.5	0.7	0.9	1.3	1.8	2.0	2.3	2.4

All races 3/, male

2006.....	0.6	0.3	0.6	0.8	0.9	1.2	1.6	2.3	3.0	3.7
2005.....	0.3	0.5	0.4	0.8	0.9	1.2	1.6	2.3	2.7	3.1
2004.....	0.3	0.5	0.5	0.9	1.0	1.4	1.9	2.0	3.2	3.2
2003.....	0.5	0.4	0.5	0.7	1.1	1.1	2.0	2.4	2.9	2.6
2002.....	0.4	0.5	0.6	0.7	1.0	1.4	2.2	2.0	3.2	3.6
2001.....	0.5	0.5	0.6	0.6	0.9	1.5	1.6	2.5	2.9	2.6
2000.....	0.4	0.5	0.6	0.5	1.1	1.3	1.9	2.3	2.7	3.7
1999.....	0.4	0.5	0.6	0.9	1.2	1.6	2.3	2.3	2.8	3.8

All races 3/, female

2006.....	0.2	0.2	0.2	0.3	0.5	0.7	1.0	2.0	1.9	2.1
2005.....	0.2	0.4	0.3	0.3	0.5	0.8	1.0	2.1	2.3	1.6
2004.....	0.3	0.3	0.3	0.5	0.5	0.8	1.0	1.6	1.5	1.7
2003.....	0.2	0.2	0.4	0.5	0.6	0.6	1.2	1.8	1.9	2.3
2002.....	0.3	0.3	0.4	0.3	0.7	0.6	1.1	1.5	1.7	2.3
2001.....	0.3	0.4	0.4	0.3	0.6	0.7	0.9	1.7	2.2	2.0
2000.....	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.4	2.3	2.0
1999.....	0.3	0.3	0.4	0.5	0.5	0.9	1.5	1.8	2.0	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hodgkin's disease (C81)

White, both sexes

2006 4/.....	0.4	0.3	0.4	0.5	0.7	1.0	1.3	2.2	2.5	2.6
2005 4/.....	0.2	0.4	0.4	0.5	0.8	1.1	1.3	2.4	2.6	2.1
2004 4/.....	0.3	0.4	0.4	0.7	0.8	1.1	1.5	1.9	2.2	2.3
2003 4/.....	0.4	0.4	0.5	0.6	0.9	0.8	1.7	2.2	2.4	2.5
2002.....	0.4	0.4	0.5	0.5	0.8	1.0	1.7	1.8	2.4	2.9
2001.....	0.4	0.4	0.5	0.4	0.8	1.2	1.3	2.1	2.6	2.4
2000.....	0.3	0.4	0.5	0.5	0.9	1.0	1.6	2.0	2.5	2.6
1999.....	0.4	0.4	0.5	0.7	0.9	1.3	2.0	2.2	2.5	2.4

White, male

2006 4/.....	0.6	0.3	0.6	0.7	1.0	1.2	1.6	2.4	3.1	3.7
2005 4/.....	0.3	0.5	0.4	0.7	1.0	1.3	1.6	2.4	2.9	3.4
2004 4/.....	0.4	0.5	0.6	0.9	1.0	1.3	2.0	2.1	3.4	3.5
2003 4/.....	0.5	0.4	0.5	0.7	1.1	1.1	2.0	2.4	3.0	2.7
2002.....	0.4	0.5	0.5	0.7	0.9	1.5	2.3	2.1	3.5	3.9
2001.....	0.5	0.5	0.7	0.6	0.9	1.6	1.8	2.5	3.0	2.7
2000.....	0.4	0.5	0.6	0.5	1.1	1.4	1.9	2.6	2.8	3.9
1999.....	0.5	0.5	0.6	0.9	1.3	1.7	2.4	2.5	3.0	3.8

White, female

2006 4/.....	*	*	0.3	0.3	0.5	0.8	1.0	2.1	2.1	2.1
2005 4/.....	*	0.4	0.3	0.4	0.6	0.9	1.0	2.3	2.5	1.5
2004 4/.....	0.2	0.4	0.3	0.5	0.5	0.8	1.0	1.7	1.5	1.7
2003 4/.....	0.2	0.3	0.4	0.5	0.6	0.6	1.4	2.0	2.0	2.4
2002.....	0.3	0.3	0.5	*	0.7	0.6	1.2	1.6	1.7	2.5
2001.....	0.3	0.4	0.4	*	0.7	0.8	0.9	1.8	2.4	2.2
2000.....	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.6	2.4	2.1
1999.....	0.3	0.4	0.4	0.5	0.6	0.9	1.6	2.0	2.2	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Non-Hodgkin's lymphoma (C82-C85)

All races 3/, both sexes

2006.....	1.5	2.5	4.2	6.7	12.5	19.0	30.1	46.2	63.4	72.3
2005.....	1.6	2.4	4.3	7.6	12.3	19.2	31.5	46.9	64.4	75.2
2004.....	1.5	2.8	4.8	7.7	12.6	21.1	32.0	46.9	64.8	75.0
2003.....	1.7	3.0	4.7	8.1	14.3	21.6	32.8	49.3	65.4	78.8
2002.....	1.9	3.0	5.0	8.8	14.9	23.2	35.0	50.2	67.0	77.3
2001.....	2.0	3.4	5.5	9.6	15.6	23.0	36.0	50.9	69.8	79.6
2000.....	2.1	3.6	5.8	10.6	16.1	25.0	36.3	53.7	71.8	80.2
1999.....	2.1	3.5	6.2	10.8	16.3	25.6	38.4	54.6	72.6	77.7

All races 3/, male

2006.....	1.9	3.1	5.6	8.8	16.0	23.8	37.2	57.3	80.7	89.6
2005.....	2.2	3.1	5.7	9.7	15.0	24.5	39.9	59.0	79.4	100.5
2004.....	2.1	3.7	6.1	9.8	15.7	26.7	39.6	57.5	81.6	95.8
2003.....	2.2	3.9	6.0	10.1	18.7	27.8	41.0	60.7	81.8	103.6
2002.....	2.4	3.9	6.2	11.3	18.4	30.0	42.8	61.8	84.4	99.7
2001.....	2.7	4.4	7.4	11.9	19.1	29.2	44.7	61.3	87.3	106.6
2000.....	2.6	4.7	7.5	13.3	19.2	30.5	45.2	66.3	88.1	105.9
1999.....	2.8	4.7	7.9	13.5	19.6	30.8	46.8	68.1	85.7	104.4

All races 3/, female

2006.....	1.1	1.8	2.8	4.7	9.3	14.8	24.3	38.2	52.6	64.2
2005.....	1.1	1.7	3.0	5.7	9.9	14.5	24.7	38.1	55.2	63.5
2004.....	0.9	1.8	3.5	5.8	9.8	16.2	25.8	39.3	54.5	65.7
2003.....	1.2	2.1	3.5	6.1	10.4	16.2	26.1	41.2	55.4	67.8
2002.....	1.3	2.2	3.8	6.4	11.7	17.3	28.7	41.9	56.4	67.6
2001.....	1.3	2.5	3.8	7.5	12.4	17.7	29.1	43.6	59.3	68.3
2000.....	1.5	2.6	4.1	8.2	13.3	20.3	29.2	45.0	62.2	69.8
1999.....	1.4	2.4	4.5	8.3	13.3	21.1	31.8	45.2	65.0	66.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Non-Hodgkin's lymphoma (C82-C85)

White, both sexes

2006 4/.....	1.4	2.3	4.2	6.7	12.8	19.8	32.0	49.1	67.0	76.4
2005 4/.....	1.6	2.4	4.2	7.8	12.6	20.2	33.2	50.0	67.9	79.7
2004 4/.....	1.5	2.7	4.7	7.9	13.1	22.2	33.5	49.2	68.6	78.8
2003 4/.....	1.7	2.9	4.7	8.4	14.9	22.8	34.2	51.8	69.0	83.3
2002.....	1.8	3.0	4.9	9.0	15.6	24.5	36.5	52.9	70.4	82.1
2001.....	2.0	3.4	5.7	9.8	16.2	23.9	37.9	53.6	73.0	83.4
2000.....	2.1	3.6	5.9	11.2	16.9	26.3	38.1	56.5	75.8	84.3
1999.....	2.0	3.6	6.4	11.1	17.0	27.1	40.2	57.0	76.7	81.2

White, male

2006 4/.....	1.8	2.9	5.6	8.7	16.3	24.6	39.3	60.6	85.5	94.1
2005 4/.....	2.2	3.0	5.5	9.8	15.3	25.7	41.9	62.8	83.6	107.0
2004 4/.....	2.1	3.6	6.1	9.9	16.2	27.7	41.3	60.1	86.2	101.2
2003 4/.....	2.1	3.7	6.0	10.6	19.4	29.6	43.1	63.9	86.5	109.7
2002.....	2.3	4.0	6.2	11.6	19.1	31.6	44.5	65.0	88.6	105.6
2001.....	2.6	4.4	7.4	12.0	19.7	30.3	46.6	64.2	91.2	111.5
2000.....	2.7	4.7	7.6	13.7	20.0	32.2	46.8	69.8	92.6	111.8
1999.....	2.7	4.7	8.1	13.7	20.2	32.3	48.9	70.2	90.8	108.0

White, female

2006 4/.....	1.0	1.7	2.9	4.7	9.5	15.5	25.8	40.5	55.3	68.1
2005 4/.....	1.0	1.7	3.0	5.8	10.2	15.3	26.0	40.4	58.1	67.1
2004 4/.....	0.9	1.8	3.4	6.0	10.2	17.3	27.1	41.3	57.7	68.7
2003 4/.....	1.2	2.1	3.5	6.4	10.7	16.7	27.0	43.0	58.2	71.6
2002.....	1.3	2.1	3.7	6.5	12.3	18.2	29.9	44.2	59.3	71.8
2001.....	1.3	2.4	3.9	7.7	13.0	18.3	30.9	46.0	62.0	71.6
2000.....	1.4	2.6	4.3	8.8	14.1	21.2	31.1	47.2	65.8	73.1
1999.....	1.4	2.4	4.8	8.7	14.1	22.6	33.3	47.7	68.3	70.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Non-Hodgkin's lymphoma (C82-C85)

Black, both sexes

2006 4/.....	2.4	4.1	4.7	8.4	12.3	15.2	19.3	25.1	31.5	34.0
2005 4/.....	2.4	3.2	5.5	7.5	11.5	13.9	20.6	22.4	32.0	33.7
2004 4/.....	1.7	3.9	5.9	7.6	11.0	15.9	21.7	28.9	26.5	38.6
2003 4/.....	2.3	3.9	5.9	6.1	12.7	14.6	22.8	26.5	31.0	31.2
2002.....	2.7	3.5	6.1	9.3	12.8	17.6	25.7	26.8	35.3	30.6
2001.....	2.4	4.3	5.9	9.8	12.3	19.1	23.4	25.7	37.9	39.3
2000.....	2.3	4.5	5.5	8.8	12.6	16.6	22.7	25.9	28.2	36.3
1999.....	2.8	4.1	5.6	10.5	11.7	17.0	23.9	32.0	31.5	40.5

Black, male

2006 4/.....	3.4	5.9	7.0	11.9	16.3	20.2	23.7	30.0	34.9	42.0
2005 4/.....	2.8	4.4	8.2	10.4	14.7	17.8	29.1	25.9	37.2	40.0
2004 4/.....	2.2	5.4	7.8	10.0	14.4	19.6	28.5	31.8	34.4	41.7
2003 4/.....	3.0	5.7	7.5	7.7	16.0	17.2	24.6	28.8	34.7	34.4
2002.....	3.8	4.0	7.4	11.5	16.6	22.9	34.4	31.7	41.3	42.9
2001.....	3.4	5.1	8.7	12.2	15.5	24.6	31.6	30.7	47.4	54.5
2000.....	2.6	6.1	7.9	13.5	15.7	19.3	32.1	30.7	32.0	43.8
1999.....	3.9	5.6	7.9	13.7	15.7	23.4	28.5	47.0	28.8	59.8

Black, female

2006 4/.....	1.5	2.7	2.8	5.6	9.2	11.6	16.3	22.2	29.7	30.7
2005 4/.....	2.0	2.1	3.2	5.2	8.9	10.9	14.7	20.3	29.3	31.1
2004 4/.....	1.3	2.5	4.2	5.6	8.3	13.1	17.1	27.2	22.3	37.4
2003 4/.....	1.6	2.3	4.5	4.7	10.1	12.6	21.5	25.2	29.0	30.0
2002.....	1.7	3.1	5.0	7.6	9.8	13.7	19.9	23.9	32.2	25.9
2001.....	1.6	3.6	3.6	7.8	9.8	15.0	17.9	22.7	33.1	33.7
2000.....	2.1	3.2	3.5	5.0	10.1	14.6	16.4	23.1	26.3	33.5
1999.....	1.7	2.8	3.7	7.9	8.7	12.3	20.9	22.9	32.8	33.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Leukemia (C91-C95)

All races 3/, both sexes

2006.....	1.8	2.5	4.1	7.0	11.6	19.9	31.3	44.7	62.9	76.9
2005.....	1.7	2.7	4.1	6.5	11.8	19.8	31.3	45.7	61.0	78.7
2004.....	1.8	2.5	4.2	7.1	12.0	19.9	31.8	45.6	61.5	78.8
2003.....	2.1	2.7	4.0	7.0	12.4	19.6	32.5	47.4	62.9	80.3
2002.....	1.9	2.8	4.5	7.4	12.9	20.9	32.4	45.7	63.4	80.5
2001.....	1.9	3.0	4.6	7.6	13.1	20.6	33.3	47.7	61.3	81.4
2000.....	1.9	3.1	4.5	7.7	12.7	21.7	33.2	47.4	65.1	83.6
1999.....	1.9	3.2	4.4	8.5	13.1	21.8	33.1	46.3	62.7	82.4

All races 3/, male

2006.....	2.0	2.7	4.9	8.7	15.1	26.3	41.6	64.8	91.8	105.1
2005.....	1.8	2.9	4.4	8.3	15.4	26.1	43.9	66.0	87.9	112.4
2004.....	2.1	2.8	5.0	9.0	15.7	26.7	43.9	64.1	86.6	110.7
2003.....	2.4	3.2	4.9	8.9	15.8	26.8	44.9	66.3	88.8	111.8
2002.....	2.2	3.2	5.3	9.2	16.1	28.0	45.8	65.0	90.0	114.5
2001.....	2.1	3.4	5.2	9.2	17.4	27.6	44.9	66.2	86.9	117.5
2000.....	2.1	3.6	5.1	9.7	16.0	29.2	46.2	65.1	93.4	119.8
1999.....	2.1	3.6	5.2	10.1	16.8	29.0	45.7	64.6	87.6	119.2

All races 3/, female

2006.....	1.7	2.3	3.3	5.3	8.5	14.3	22.8	30.0	44.8	63.7
2005.....	1.6	2.5	3.7	4.9	8.5	14.4	21.1	31.0	44.3	63.3
2004.....	1.6	2.2	3.3	5.4	8.6	14.1	22.1	32.3	46.0	64.5
2003.....	1.7	2.2	3.2	5.2	9.4	13.3	22.7	33.9	47.0	66.4
2002.....	1.6	2.5	3.8	5.7	10.0	14.7	21.8	32.1	47.3	65.7
2001.....	1.7	2.7	4.0	6.0	9.3	14.6	24.1	34.8	45.9	66.3
2000.....	1.7	2.6	3.9	5.9	9.7	15.3	22.9	35.1	48.4	68.8
1999.....	1.7	2.8	3.6	7.0	9.8	15.6	23.2	33.6	48.1	67.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Leukemia (C91-C95)

White, both sexes

2006 4/.....	1.9	2.5	4.2	7.1	11.8	20.7	32.5	46.0	66.0	79.9
2005 4/.....	1.7	2.7	4.0	6.7	12.1	20.5	32.7	47.5	63.7	82.4
2004 4/.....	1.8	2.5	4.2	7.3	12.2	20.5	32.5	47.1	63.6	81.6
2003 4/.....	2.1	2.8	4.0	7.1	12.7	20.3	33.6	49.3	65.0	83.4
2002.....	2.0	2.7	4.5	7.4	13.1	21.8	33.6	47.0	65.7	83.0
2001.....	1.9	3.1	4.5	7.6	13.5	20.9	34.5	49.1	63.8	84.7
2000.....	1.9	3.0	4.5	7.8	12.8	22.6	34.4	49.2	67.6	86.0
1999.....	1.9	3.2	4.3	8.7	13.3	22.2	34.2	48.0	64.1	85.4

White, male

2006 4/.....	1.9	2.8	5.0	8.9	15.3	27.1	43.2	66.3	96.1	108.7
2005 4/.....	1.7	2.9	4.5	8.5	15.8	27.0	45.6	67.9	91.4	118.7
2004 4/.....	2.1	2.8	4.9	9.2	16.0	27.2	44.4	66.2	89.8	115.3
2003 4/.....	2.4	3.4	4.8	9.0	16.4	27.7	46.4	69.2	92.8	116.9
2002.....	2.3	3.0	5.3	9.3	16.3	29.0	47.2	67.0	93.3	119.1
2001.....	2.0	3.5	5.1	9.4	17.8	27.8	46.5	68.3	90.1	123.1
2000.....	2.2	3.5	5.0	9.8	16.1	30.5	47.4	67.5	96.7	123.6
1999.....	2.1	3.5	5.1	10.3	17.0	29.5	47.1	66.7	90.2	124.1

White, female

2006 4/.....	1.8	2.3	3.4	5.5	8.5	15.0	23.6	30.8	46.9	66.4
2005 4/.....	1.6	2.5	3.6	4.9	8.7	14.7	22.0	32.4	46.4	65.7
2004 4/.....	1.6	2.3	3.4	5.4	8.6	14.5	22.6	33.2	47.3	66.4
2003 4/.....	1.7	2.3	3.2	5.4	9.3	13.8	23.1	34.9	47.9	68.6
2002.....	1.7	2.5	3.8	5.6	10.1	15.4	22.6	32.8	48.9	67.3
2001.....	1.7	2.7	3.9	6.0	9.5	14.8	24.7	35.4	47.9	68.6
2000.....	1.6	2.5	4.0	5.9	9.8	15.7	23.9	36.3	50.2	70.8
1999.....	1.8	2.9	3.6	7.2	9.8	15.8	23.9	34.9	48.7	69.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Leukemia (C91-C95)

Black, both sexes

2006 4/.....	2.2	2.6	4.1	7.3	12.8	17.9	28.6	39.2	41.7	55.5
2005 4/.....	2.0	3.1	5.0	6.8	10.7	18.7	24.3	35.1	43.2	52.1
2004 4/.....	2.2	2.9	5.3	7.7	13.8	19.4	32.5	37.1	47.0	60.1
2003 4/.....	2.5	2.5	5.1	7.2	12.8	17.9	29.6	37.3	48.0	59.2
2002.....	1.7	3.6	5.1	9.1	13.8	17.3	28.6	39.2	46.6	64.7
2001.....	2.3	3.0	5.6	7.9	12.3	22.3	28.0	41.3	42.0	57.1
2000.....	2.4	4.5	5.2	7.9	14.1	17.9	28.4	34.7	43.8	66.0
1999.....	2.0	3.5	5.4	8.1	13.6	22.2	27.9	35.0	51.1	55.8

Black, male

2006 4/.....	3.0	2.6	5.2	9.7	16.6	26.9	38.4	58.6	61.6	84.9
2005 4/.....	2.2	3.7	5.1	8.3	14.0	24.5	34.2	52.7	66.7	62.3
2004 4/.....	2.3	3.1	7.1	9.4	17.6	28.3	48.6	48.2	58.8	79.5
2003 4/.....	3.1	2.7	6.8	10.7	14.5	24.9	39.4	50.2	58.3	77.2
2002.....	1.7	4.8	5.7	11.0	17.2	23.4	42.4	52.4	64.2	86.8
2001.....	2.7	2.8	6.6	9.9	17.1	32.3	36.4	51.6	61.0	77.2
2000.....	2.5	4.9	6.2	10.0	18.6	22.4	43.5	49.6	64.1	98.0
1999.....	2.2	4.4	6.9	10.3	17.2	29.1	38.0	51.3	61.0	77.3

Black, female

2006 4/.....	1.4	2.6	3.2	5.3	9.8	11.3	21.8	27.6	31.1	43.5
2005 4/.....	1.8	2.6	4.9	5.5	8.2	14.5	17.5	24.5	30.8	48.0
2004 4/.....	2.0	2.6	3.8	6.3	10.9	12.8	21.5	30.5	40.8	52.4
2003 4/.....	2.1	2.4	3.8	4.3	11.4	12.8	23.0	29.6	42.6	52.3
2002.....	1.6	2.6	4.7	7.6	11.2	12.8	19.3	31.3	37.4	56.3
2001.....	2.0	3.3	4.7	6.3	8.5	15.0	22.4	35.2	32.3	49.7
2000.....	2.3	4.2	4.4	6.2	10.5	14.6	18.4	25.6	33.5	54.1
1999.....	1.7	2.6	4.2	6.2	10.8	17.1	21.2	25.2	46.2	47.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All races 3/, both sexes

2006.....	0.5	1.1	2.4	4.3	7.5	11.8	19.0	25.8	33.3	33.8
2005.....	0.4	1.1	2.2	4.3	8.4	12.8	19.6	26.5	33.4	33.6
2004.....	0.5	1.0	2.3	4.6	7.7	12.3	18.9	28.3	33.9	33.1
2003.....	0.5	1.1	2.5	4.4	7.9	13.1	20.4	28.5	34.5	36.3
2002.....	0.4	1.3	2.5	4.7	8.3	14.1	21.0	27.9	35.8	36.7
2001.....	0.5	1.2	2.3	5.1	8.5	14.0	21.1	28.3	34.7	36.3
2000.....	0.5	1.2	2.6	4.5	8.5	14.4	20.4	28.3	36.4	37.8
1999.....	0.5	1.2	2.5	5.6	9.0	14.1	20.6	27.6	36.2	36.6

All races 3/, male

2006.....	0.6	1.2	2.9	5.1	9.2	14.3	22.9	33.1	43.8	45.8
2005.....	0.6	1.3	2.7	5.2	9.4	16.3	23.9	33.5	44.5	46.1
2004.....	0.5	1.1	2.7	5.2	9.3	15.3	23.2	35.0	42.8	44.6
2003.....	0.7	1.3	3.1	5.4	10.1	16.0	24.3	33.9	43.3	50.4
2002.....	0.5	1.5	3.0	5.6	10.2	16.8	25.8	35.5	46.2	50.8
2001.....	0.7	1.5	2.8	6.0	10.1	16.8	25.3	35.6	43.4	48.0
2000.....	0.6	1.4	2.9	4.9	9.5	17.1	24.2	35.7	46.1	51.8
1999.....	0.6	1.4	3.0	6.5	11.0	16.6	24.1	34.1	45.5	51.3

All races 3/, female

2006.....	0.4	0.9	1.9	3.6	6.0	9.6	15.8	20.5	26.8	28.2
2005.....	0.3	0.9	1.7	3.5	7.4	9.9	16.1	21.4	26.5	27.9
2004.....	0.4	0.9	1.8	4.0	6.2	9.8	15.5	23.5	28.4	28.0
2003.....	0.4	0.9	1.9	3.5	6.0	10.6	17.2	24.7	29.2	30.1
2002.....	0.4	1.0	2.0	4.0	6.5	11.7	17.2	22.5	29.5	30.6
2001.....	0.4	0.9	1.9	4.2	7.1	11.5	17.7	23.2	29.4	31.3
2000.....	0.4	1.0	2.3	4.2	7.6	12.1	17.3	23.1	30.7	32.2
1999.....	0.4	1.1	2.1	4.7	7.3	12.0	17.8	23.0	30.7	30.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

White, both sexes

2006 4/.....	0.5	0.8	2.0	3.8	6.6	10.9	17.7	24.8	32.4	32.8
2005 4/.....	0.4	0.9	1.9	3.7	7.7	11.7	18.3	25.5	32.7	32.9
2004 4/.....	0.4	0.9	1.9	4.0	7.1	11.3	17.4	27.0	32.5	32.0
2003 4/.....	0.5	0.9	2.1	3.9	7.3	12.1	19.1	26.9	33.1	35.0
2002.....	0.4	1.1	2.1	4.2	7.5	13.0	19.9	26.9	34.8	35.7
2001.....	0.4	1.0	2.1	4.4	7.5	12.9	19.9	27.1	33.0	35.0
2000.....	0.4	1.0	2.2	3.8	7.8	13.4	19.1	26.5	35.2	35.2
1999.....	0.4	1.1	2.1	4.7	7.9	13.2	19.2	25.7	34.9	34.8

White, male

2006 4/.....	0.6	1.0	2.5	4.5	8.0	13.4	21.4	32.0	42.8	45.0
2005 4/.....	0.5	1.0	2.3	4.4	8.9	15.2	22.5	32.2	43.9	45.8
2004 4/.....	0.5	1.0	2.3	4.6	8.7	14.2	21.8	33.8	41.2	43.5
2003 4/.....	0.6	1.1	2.6	4.8	9.4	15.1	23.4	32.9	41.9	49.7
2002.....	0.4	1.3	2.7	4.9	9.4	15.8	24.9	34.8	45.3	50.3
2001.....	0.6	1.3	2.5	5.2	9.0	15.7	24.1	34.6	41.7	47.1
2000.....	0.5	1.2	2.5	4.3	8.8	16.1	22.8	33.9	45.4	50.2
1999.....	0.4	1.3	2.6	5.6	10.0	15.8	22.5	32.4	44.3	49.7

White, female

2006 4/.....	0.4	0.7	1.6	3.1	5.3	8.6	14.6	19.5	25.8	27.1
2005 4/.....	0.3	0.7	1.4	3.1	6.6	8.6	14.9	20.5	25.6	26.9
2004 4/.....	0.3	0.8	1.5	3.3	5.6	8.7	13.8	22.0	27.0	26.8
2003 4/.....	0.3	0.7	1.6	3.1	5.4	9.5	15.6	22.6	27.7	28.5
2002.....	0.3	0.8	1.6	3.4	5.7	10.5	15.8	21.2	28.4	29.4
2001.....	*	0.7	1.6	3.7	6.1	10.4	16.5	21.7	27.8	29.9
2000.....	0.2	0.8	2.0	3.4	6.8	11.1	16.2	21.3	29.2	29.1
1999.....	0.3	1.0	1.5	3.8	5.9	10.9	16.6	21.0	29.4	28.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Black, both sexes

2006 4/.....	0.9	2.6	5.4	9.5	17.2	23.3	34.3	43.7	53.0	51.6
2005 4/.....	0.9	2.3	5.0	9.9	16.2	25.4	35.1	44.3	51.2	49.4
2004 4/.....	1.0	2.0	5.2	10.2	15.3	24.4	38.1	48.6	59.4	53.9
2003 4/.....	1.2	2.8	5.9	9.4	15.1	25.6	37.3	52.6	59.4	59.8
2002.....	1.0	2.9	5.6	10.7	16.3	26.7	37.7	44.7	54.9	55.5
2001.....	1.5	2.6	4.8	11.8	19.4	25.8	38.2	47.9	61.2	58.3
2000.....	1.1	3.1	6.3	11.7	16.5	27.3	37.9	53.3	57.9	71.6
1999.....	1.4	2.4	6.6	13.8	20.0	24.2	37.6	51.1	56.3	63.1

Black, male

2006 4/.....	*	3.0	6.4	11.6	22.4	27.4	47.1	54.6	68.5	65.2
2005 4/.....	*	3.0	6.2	12.7	17.5	30.4	42.7	58.1	67.4	61.4
2004 4/.....	*	2.1	6.1	11.3	18.1	30.8	44.3	57.8	73.9	71.5
2003 4/.....	1.5	3.2	7.7	12.2	19.7	29.3	39.4	57.5	73.1	73.0
2002.....	*	3.5	5.9	12.4	20.2	31.4	44.7	52.4	70.3	71.8
2001.....	1.6	3.4	5.3	14.5	23.4	30.2	45.1	57.1	80.3	71.5
2000.....	*	3.3	7.2	12.3	18.4	31.0	48.1	67.1	69.1	79.6
1999.....	1.9	2.6	6.7	16.1	21.6	26.2	46.8	59.4	74.1	80.9

Black, female

2006 4/.....	*	2.3	4.7	7.8	13.1	20.3	25.6	37.2	44.9	46.1
2005 4/.....	*	1.7	4.1	7.5	15.2	21.7	29.9	36.0	42.7	44.6
2004 4/.....	*	1.9	4.4	9.3	13.1	19.6	33.8	43.1	51.8	46.9
2003 4/.....	*	2.4	4.4	7.0	11.6	22.8	35.9	49.6	52.2	54.7
2002.....	*	2.4	5.3	9.3	13.3	23.1	32.9	40.1	46.9	49.3
2001.....	1.5	2.0	4.3	9.5	16.3	22.5	33.6	42.4	51.5	53.5
2000.....	*	2.8	5.5	11.2	15.0	24.6	31.1	45.0	52.2	68.7
1999.....	*	2.3	6.6	11.9	18.8	22.7	31.5	46.1	47.5	56.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes

2006.....	5.4	9.4	16.5	26.9	42.5	62.5	90.2	122.7	162.9	210.1
2005.....	5.3	9.9	16.6	27.4	43.5	62.7	90.1	121.9	167.4	212.9
2004.....	5.3	9.6	16.2	27.6	43.3	64.6	91.1	126.4	165.2	213.1
2003.....	5.3	10.3	17.7	28.1	44.4	64.6	92.9	131.5	167.7	224.0
2002.....	5.8	10.4	17.5	28.8	46.5	66.0	97.0	131.0	168.7	230.1
2001.....	5.9	10.5	18.0	30.1	46.4	67.2	98.1	133.4	170.9	234.5
2000.....	5.9	10.5	18.2	30.4	48.2	71.5	100.4	136.0	177.4	239.8
1999.....	6.7	11.4	19.1	32.5	52.5	76.1	107.8	142.3	188.5	250.4

All races 3/, male

2006.....	5.6	10.5	18.5	31.5	51.7	74.1	107.7	150.6	207.3	259.6
2005.....	5.9	11.2	19.1	32.8	52.3	74.7	109.2	148.5	208.5	263.5
2004.....	5.5	10.6	18.8	32.5	51.4	78.8	109.3	157.8	207.2	260.6
2003.....	5.9	11.9	20.1	33.0	52.9	78.3	111.3	162.7	208.2	275.0
2002.....	6.2	11.7	20.8	33.0	56.5	79.9	118.9	161.8	207.8	290.8
2001.....	6.4	12.0	21.5	34.9	55.4	79.4	120.3	162.3	211.0	295.2
2000.....	6.4	12.0	20.6	35.1	57.2	86.2	122.5	167.5	221.4	300.7
1999.....	7.6	13.2	22.0	38.2	62.6	90.6	132.8	174.0	233.0	318.9

All races 3/, female

2006.....	5.2	8.3	14.5	22.5	34.2	52.4	76.0	102.3	135.1	186.9
2005.....	4.7	8.7	14.2	22.4	35.6	52.1	74.7	102.5	141.9	189.6
2004.....	5.1	8.6	13.7	23.0	35.9	52.3	76.5	103.8	139.3	191.8
2003.....	4.8	8.7	15.5	23.6	36.7	52.8	78.1	109.2	142.9	201.4
2002.....	5.4	9.1	14.3	24.9	37.5	53.9	79.5	109.2	144.9	203.8
2001.....	5.3	9.1	14.7	25.7	38.3	56.7	80.5	113.1	146.8	209.1
2000.....	5.3	9.1	15.8	25.9	40.0	59.0	83.1	114.1	151.4	215.0
1999.....	5.8	9.7	16.3	27.1	43.3	63.8	88.2	120.3	162.5	222.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

White, both sexes

2006 4/.....	5.3	9.0	15.8	26.2	41.9	62.2	91.7	124.3	166.5	213.9
2005 4/.....	5.2	9.4	15.8	26.7	42.9	62.4	90.9	123.8	170.0	217.2
2004 4/.....	5.1	9.0	15.4	26.7	42.6	63.9	92.0	127.8	167.1	217.6
2003 4/.....	5.0	9.7	17.0	27.3	44.0	64.7	93.5	132.7	170.1	227.9
2002.....	5.5	9.9	16.8	28.1	45.7	65.6	96.9	131.7	170.0	233.0
2001.....	5.6	9.9	17.2	29.0	45.3	66.5	97.4	134.1	171.4	238.1
2000.....	5.4	9.9	17.2	29.7	47.1	71.4	99.9	135.6	179.5	243.7
1999.....	6.3	10.8	18.3	31.1	51.2	76.3	107.1	142.3	189.4	252.4

White, male

2006 4/.....	5.5	10.1	17.6	30.6	51.3	74.2	109.7	152.5	212.2	266.4
2005 4/.....	5.8	10.4	18.3	31.7	51.4	73.8	109.4	151.6	211.5	270.3
2004 4/.....	5.2	9.9	17.5	31.0	50.2	77.5	109.7	159.9	210.9	266.8
2003 4/.....	5.4	11.0	19.2	31.8	52.2	78.0	111.6	164.1	212.2	280.9
2002.....	5.7	11.1	20.0	32.0	55.1	80.2	118.8	162.2	210.9	296.5
2001.....	6.2	11.3	20.1	33.1	53.5	78.0	119.7	163.1	212.0	300.2
2000.....	5.8	11.2	19.5	34.1	54.7	86.2	121.8	165.7	225.2	307.0
1999.....	7.2	12.2	20.8	36.6	60.9	90.1	131.8	174.6	235.3	323.6

White, female

2006 4/.....	5.0	8.0	14.1	21.9	33.1	51.5	76.8	103.3	137.7	189.3
2005 4/.....	4.5	8.4	13.4	21.9	35.0	52.1	75.7	103.2	144.0	192.8
2004 4/.....	5.0	8.2	13.2	22.5	35.5	51.9	77.5	104.4	139.9	195.4
2003 4/.....	4.6	8.4	14.9	23.1	36.3	52.9	78.8	110.0	144.1	204.4
2002.....	5.4	8.7	13.7	24.3	36.9	52.7	79.1	109.9	144.9	205.5
2001.....	5.1	8.5	14.3	25.1	37.7	56.3	79.3	113.7	146.8	212.1
2000.....	5.0	8.6	14.8	25.6	40.1	58.4	82.4	114.5	152.3	218.0
1999.....	5.3	9.5	15.8	26.0	42.3	64.3	87.5	119.6	162.3	223.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Black, both sexes

2006 4/.....	7.2	13.1	24.5	38.9	57.6	77.1	96.3	127.5	149.3	186.0
2005 4/.....	7.2	14.9	25.1	39.4	56.1	76.0	97.9	127.4	168.9	183.9
2004 4/.....	7.5	14.7	25.0	40.6	57.1	84.2	99.1	133.0	167.8	186.7
2003 4/.....	7.8	16.0	26.5	41.1	57.7	77.0	102.4	139.4	166.0	206.3
2002.....	8.0	15.8	25.3	41.5	63.3	82.5	113.5	146.5	174.5	222.1
2001.....	7.7	16.7	27.8	44.1	64.1	85.2	120.3	140.7	182.3	217.1
2000.....	10.0	16.8	28.8	41.3	64.8	85.6	117.7	155.7	172.0	210.8
1999.....	10.1	18.3	29.0	48.3	73.5	85.4	128.3	157.3	193.0	242.6

Black, male

2006 4/.....	7.4	15.8	30.2	47.5	70.0	90.5	116.6	161.5	199.2	213.6
2005 4/.....	7.7	18.1	28.3	50.5	71.6	98.6	127.5	144.4	217.0	221.2
2004 4/.....	8.3	17.1	32.7	52.1	73.3	107.1	128.8	169.7	203.0	222.5
2003 4/.....	9.4	19.8	31.4	53.0	72.3	98.5	128.4	176.2	206.0	241.9
2002.....	9.6	18.4	30.9	50.2	82.0	97.2	142.9	187.4	209.4	267.0
2001.....	8.2	19.0	35.9	57.1	80.9	110.0	151.5	178.9	219.1	283.7
2000.....	11.8	20.8	33.1	51.4	89.0	102.0	149.1	211.7	208.2	253.7
1999.....	11.7	23.2	36.5	61.9	94.0	109.8	163.8	191.4	233.7	292.9

Black, female

2006 4/.....	7.1	10.7	19.7	31.9	47.9	67.2	82.4	107.1	123.0	174.7
2005 4/.....	6.7	12.1	22.4	30.3	44.1	59.2	77.7	117.3	143.6	168.8
2004 4/.....	6.8	12.5	18.5	31.1	44.4	67.3	78.9	111.3	149.3	172.5
2003 4/.....	6.4	12.6	22.3	31.3	46.3	61.0	84.8	117.5	145.1	192.4
2002.....	6.5	13.5	20.5	34.5	48.6	71.6	93.9	122.1	156.4	205.0
2001.....	7.3	14.8	20.9	33.5	50.9	66.8	99.5	117.9	163.5	192.4
2000.....	8.3	13.4	25.1	33.1	45.8	73.5	96.8	121.9	153.7	194.9
1999.....	8.8	14.0	22.7	37.2	57.4	67.3	104.6	136.7	172.9	223.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races 3/, both sexes

2006.....	0.8	1.5	2.2	3.1	5.6	9.0	16.8	28.8	46.3	78.0
2005.....	1.0	1.4	2.1	3.3	5.7	9.5	16.3	27.9	44.8	76.7
2004.....	0.8	1.4	2.0	3.5	5.8	9.3	16.7	27.3	47.2	77.7
2003.....	0.9	1.4	2.0	3.2	6.0	9.8	16.3	28.5	47.4	80.5
2002.....	1.0	1.5	2.1	3.7	5.4	9.7	16.3	28.8	46.3	78.7
2001.....	0.9	1.5	2.3	3.9	6.3	10.5	17.9	29.5	47.8	82.7
2000.....	0.9	1.5	2.3	3.8	6.6	10.6	17.4	28.9	50.6	84.9
1999.....	0.9	1.4	2.3	3.5	6.3	10.8	18.6	29.3	47.5	82.9

All races 3/, male

2006.....	0.9	1.6	2.5	3.8	6.3	11.1	21.5	38.2	60.8	92.3
2005.....	1.0	1.5	2.4	3.6	7.2	12.1	20.3	36.3	59.5	92.7
2004.....	0.9	1.4	2.3	4.0	7.1	11.7	20.5	34.8	61.3	98.9
2003.....	1.0	1.4	2.3	3.6	7.0	11.6	21.2	35.9	61.5	101.7
2002.....	1.0	1.6	2.1	4.5	6.3	12.2	20.0	37.3	59.1	101.1
2001.....	1.0	1.7	2.4	4.1	7.5	12.8	22.5	39.2	62.4	107.6
2000.....	0.9	1.6	2.6	4.1	7.5	13.3	21.2	36.8	65.3	108.9
1999.....	1.1	1.5	2.7	4.0	7.9	13.4	23.9	37.2	61.0	109.2

All races 3/, female

2006.....	0.8	1.4	1.9	2.5	4.9	7.2	13.0	21.8	37.3	71.4
2005.....	1.0	1.3	1.8	3.1	4.3	7.3	13.0	21.8	35.6	69.4
2004.....	0.7	1.3	1.7	3.0	4.6	7.1	13.7	21.9	38.5	68.1
2003.....	0.7	1.3	1.7	2.8	5.1	8.3	12.4	23.2	38.7	71.1
2002.....	0.9	1.5	2.0	3.0	4.6	7.5	13.4	22.8	38.5	68.9
2001.....	0.8	1.2	2.1	3.7	5.2	8.5	14.3	22.7	39.1	72.3
2000.....	1.0	1.3	1.9	3.5	5.7	8.2	14.4	23.4	41.9	75.2
1999.....	0.8	1.3	1.9	3.0	4.8	8.5	14.5	23.8	39.7	72.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

White, both sexes

2006 4/.....	0.8	1.4	2.2	3.2	5.4	9.2	17.2	29.5	47.8	80.0
2005 4/.....	0.9	1.3	2.1	3.3	5.6	9.7	16.6	28.7	46.3	79.6
2004 4/.....	0.7	1.4	1.9	3.4	5.7	9.4	17.1	27.7	48.5	80.2
2003 4/.....	0.9	1.3	2.0	3.1	6.1	9.8	16.8	29.5	48.7	83.0
2002.....	0.9	1.6	2.0	3.7	5.6	9.9	16.7	29.7	47.2	80.7
2001.....	0.9	1.5	2.3	3.9	6.3	10.7	18.1	30.4	48.9	83.9
2000.....	0.9	1.3	2.2	3.9	6.6	10.7	17.3	29.5	52.2	87.3
1999.....	0.9	1.3	2.2	3.5	6.2	10.7	19.0	29.7	48.7	85.3

White, male

2006 4/.....	0.8	1.5	2.6	3.8	6.2	11.3	22.2	39.5	62.5	93.7
2005 4/.....	0.9	1.4	2.4	3.7	7.1	12.4	21.0	37.5	61.9	97.0
2004 4/.....	0.9	1.4	2.3	3.9	7.0	12.0	21.0	35.4	63.2	103.2
2003 4/.....	1.0	1.3	2.3	3.5	7.1	11.8	21.8	37.4	63.3	105.5
2002.....	1.1	1.7	2.1	4.4	6.4	12.4	20.7	38.4	60.1	104.2
2001.....	1.0	1.7	2.5	4.1	7.4	13.2	23.0	40.4	63.7	109.9
2000.....	0.9	1.5	2.6	4.3	7.3	13.4	21.0	37.6	68.0	112.6
1999.....	1.1	1.4	2.6	3.9	8.0	13.3	24.6	37.7	62.9	114.3

White, female

2006 4/.....	0.7	1.3	1.8	2.6	4.7	7.4	13.1	22.0	38.5	73.6
2005 4/.....	0.9	1.1	1.8	3.0	4.1	7.3	13.0	22.1	36.5	71.6
2004 4/.....	0.6	1.3	1.5	2.9	4.6	7.0	13.8	22.1	39.3	69.9
2003 4/.....	0.7	1.2	1.7	2.8	5.3	8.0	12.7	23.9	39.8	73.0
2002.....	0.8	1.4	1.8	3.0	4.8	7.6	13.5	23.4	39.2	70.5
2001.....	0.8	1.2	2.0	3.8	5.2	8.6	14.2	23.3	40.0	73.0
2000.....	1.0	1.2	1.9	3.5	6.0	8.4	14.3	23.8	42.9	77.1
1999.....	0.7	1.2	1.8	3.0	4.6	8.3	14.7	24.0	40.4	73.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Black, both sexes

2006 4/.....	1.4	2.3	2.7	3.8	7.8	9.2	14.8	26.0	39.5	62.3
2005 4/.....	1.6	2.4	2.5	3.9	7.5	9.9	16.4	24.1	33.5	54.5
2004 4/.....	1.3	1.6	3.1	5.5	7.1	10.2	16.1	26.1	39.4	57.0
2003 4/.....	1.0	2.2	2.7	4.2	6.6	11.2	15.3	22.6	39.9	57.8
2002.....	1.7	2.0	3.0	5.2	5.8	10.6	15.0	24.4	41.9	63.6
2001.....	1.2	1.4	2.7	4.5	7.6	10.9	18.6	24.5	42.5	73.4
2000.....	1.4	2.4	3.1	4.2	7.1	10.6	20.3	26.7	37.8	64.1
1999.....	1.3	2.2	3.2	5.0	7.8	13.7	18.1	29.5	38.7	61.9

Black, male

2006 4/.....	*	2.3	2.4	4.7	8.7	12.2	18.1	32.7	54.8	80.4
2005 4/.....	1.7	2.6	2.8	4.3	9.1	10.9	18.0	28.2	42.8	54.8
2004 4/.....	*	1.7	3.5	6.6	9.3	12.4	19.4	34.6	46.6	61.6
2003 4/.....	*	2.1	2.9	5.5	8.5	12.3	21.4	26.5	53.2	66.7
2002.....	*	1.7	2.8	7.4	6.5	12.7	16.0	29.9	53.5	79.3
2001.....	*	*	2.4	4.5	9.2	13.0	22.9	31.6	58.6	88.5
2000.....	*	2.2	3.6	4.4	10.1	15.6	23.7	34.5	38.8	76.1
1999.....	*	*	3.3	6.1	8.3	16.9	21.7	36.1	42.7	63.3

Black, female

2006 4/.....	1.6	2.3	3.0	3.0	7.2	6.9	12.5	22.0	31.5	54.8
2005 4/.....	1.4	2.2	2.3	3.6	6.2	9.2	15.4	21.6	28.6	54.4
2004 4/.....	1.4	1.6	2.6	4.7	5.4	8.6	13.8	21.0	35.5	55.1
2003 4/.....	*	2.3	2.6	3.1	5.1	10.4	11.3	20.3	32.9	54.3
2002.....	2.1	2.2	3.3	3.3	5.3	9.1	14.3	21.1	35.8	57.5
2001.....	*	*	3.0	4.5	6.4	9.4	15.7	20.2	34.3	67.8
2000.....	1.3	2.5	2.7	4.0	4.7	6.9	18.0	21.9	37.4	59.7
1999.....	1.4	2.5	3.0	4.0	7.5	11.4	15.8	25.5	36.7	61.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Anemias (D50-D64)

All races 3/, both sexes

2006.....	0.5	0.5	0.5	0.8	1.2	1.4	3.0	5.1	10.5	30.0
2005.....	0.5	0.6	0.6	0.8	1.2	2.0	3.5	6.3	12.8	36.6
2004.....	0.5	0.5	0.6	0.8	1.3	1.8	3.4	5.9	12.2	38.1
2003.....	0.4	0.6	0.5	0.8	1.3	1.9	3.5	6.0	12.7	40.7
2002.....	0.4	0.6	0.6	0.8	1.2	1.9	3.7	6.4	12.9	40.3
2001.....	0.6	0.5	0.7	0.8	1.2	2.0	3.5	7.1	13.6	42.0
2000.....	0.5	0.5	0.7	0.9	1.2	2.2	3.4	6.5	13.9	42.4
1999.....	0.5	0.7	0.6	0.8	1.2	1.8	3.5	6.8	13.7	43.8

All races 3/, male

2006.....	0.5	0.5	0.5	0.8	1.1	1.5	2.8	5.4	11.6	29.6
2005.....	0.5	0.5	0.5	0.7	1.3	1.9	3.5	6.4	14.1	33.5
2004.....	0.5	0.5	0.6	1.0	1.4	1.9	3.6	6.3	12.8	36.5
2003.....	0.5	0.7	0.5	0.9	1.5	2.3	3.8	6.6	13.3	38.0
2002.....	0.4	0.5	0.6	0.8	1.3	1.7	4.3	6.6	12.4	38.3
2001.....	0.4	0.6	0.6	0.8	1.1	2.1	3.8	7.2	14.3	40.0
2000.....	0.4	0.5	0.7	1.0	1.2	2.2	3.6	6.5	14.6	40.1
1999.....	0.5	0.7	0.6	0.7	1.2	2.1	3.4	7.2	14.8	43.4

All races 3/, female

2006.....	0.5	0.6	0.6	0.8	1.2	1.3	3.2	5.0	9.8	30.2
2005.....	0.5	0.7	0.7	0.9	1.2	2.0	3.5	6.3	12.1	38.0
2004.....	0.5	0.6	0.6	0.7	1.3	1.7	3.3	5.7	11.8	38.8
2003.....	0.4	0.5	0.6	0.7	1.0	1.5	3.2	5.6	12.3	41.9
2002.....	0.5	0.7	0.7	0.8	1.1	2.1	3.2	6.2	13.2	41.2
2001.....	0.7	0.5	0.7	0.9	1.2	1.9	3.3	7.0	13.2	42.9
2000.....	0.5	0.6	0.7	0.9	1.2	2.2	3.3	6.5	13.6	43.3
1999.....	0.4	0.7	0.6	0.9	1.1	1.6	3.6	6.5	13.0	44.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Anemias (D50-D64)

White, both sexes

2006 4/.....	0.1	0.3	0.3	0.6	0.9	1.2	2.8	5.0	10.4	29.6
2005 4/.....	0.1	0.3	0.3	0.6	1.0	1.6	3.1	6.0	12.4	36.6
2004 4/.....	0.2	0.3	0.3	0.5	1.0	1.7	3.2	5.6	12.2	37.7
2003 4/.....	0.1	0.3	0.3	0.4	0.9	1.6	3.2	5.7	12.4	40.6
2002.....	0.1	0.3	0.3	0.6	0.9	1.7	3.4	5.9	12.6	39.7
2001.....	0.2	0.2	0.4	0.6	1.0	1.8	3.2	6.7	13.2	41.7
2000.....	0.1	0.2	0.4	0.7	0.9	1.9	3.2	6.2	13.8	42.0
1999.....	0.1	0.3	0.3	0.4	0.8	1.6	3.2	6.5	13.6	43.0

White, male

2006 4/.....	*	*	0.3	0.5	0.9	1.4	2.6	5.1	11.5	29.9
2005 4/.....	*	0.3	0.3	0.6	1.0	1.5	3.3	5.9	13.5	33.7
2004 4/.....	0.2	0.3	0.3	0.5	1.2	1.7	3.5	5.7	12.7	36.7
2003 4/.....	*	0.3	0.3	0.5	1.1	2.0	3.3	6.3	13.4	37.9
2002.....	*	0.3	0.3	0.6	1.0	1.6	3.7	6.3	11.9	37.2
2001.....	*	*	0.4	0.5	1.0	1.9	3.3	7.0	14.1	40.4
2000.....	*	*	0.4	0.8	0.9	2.1	3.4	6.1	14.2	39.1
1999.....	*	0.4	0.3	*	0.7	1.9	3.1	7.0	14.8	43.4

White, female

2006 4/.....	*	0.3	0.3	0.7	1.0	1.1	3.1	4.8	9.7	29.4
2005 4/.....	*	0.4	0.3	0.7	1.0	1.7	3.0	6.1	11.7	37.9
2004 4/.....	*	0.2	0.3	0.5	0.9	1.6	3.0	5.6	11.9	38.2
2003 4/.....	*	*	0.3	0.3	0.8	1.3	3.1	5.3	11.8	41.8
2002.....	*	0.3	0.3	0.5	0.8	1.8	3.1	5.6	13.0	40.8
2001.....	0.3	*	0.4	0.6	1.1	1.6	3.1	6.5	12.6	42.3
2000.....	*	0.3	0.4	0.7	0.9	1.8	3.0	6.3	13.5	43.2
1999.....	*	0.3	0.3	0.5	0.9	1.4	3.4	6.1	12.9	42.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Anemias (D50-D64)

Black, both sexes

2006 4/.....	2.9	2.6	2.4	2.9	3.4	3.0	5.5	8.0	13.5	38.9
2005 4/.....	2.9	2.5	2.9	2.5	3.7	5.3	7.3	11.0	20.1	42.5
2004 4/.....	2.4	2.5	3.0	3.9	4.2	3.7	6.2	10.7	13.4	48.2
2003 4/.....	2.5	3.3	2.7	4.1	4.4	4.3	7.1	10.9	19.8	49.3
2002.....	2.4	3.1	3.1	3.2	4.1	4.4	7.9	13.7	19.4	52.3
2001.....	3.2	3.2	2.9	3.2	2.6	3.9	7.4	12.5	21.4	48.5
2000.....	2.7	2.9	3.2	3.0	3.9	4.8	6.1	10.3	18.1	51.0
1999.....	2.8	3.2	2.9	4.2	4.3	3.6	6.6	11.4	16.5	52.9

Black, male

2006 4/.....	2.9	2.4	2.1	3.3	3.9	*	*	9.4	15.7	29.5
2005 4/.....	3.1	2.4	2.6	*	*	6.7	*	13.4	23.2	37.2
2004 4/.....	2.5	2.1	2.9	5.3	4.5	*	*	13.6	14.3	42.7
2003 4/.....	2.9	3.9	2.6	4.9	6.2	5.7	10.8	12.8	18.5	49.0
2002.....	2.3	2.6	2.6	3.0	5.1	*	11.7	12.9	23.7	56.8
2001.....	2.7	3.9	3.2	3.3	*	*	10.4	12.1	20.1	42.0
2000.....	2.2	3.2	3.4	*	*	*	*	10.9	19.4	55.4
1999.....	2.9	3.3	3.2	4.2	6.2	*	7.8	10.9	*	44.5

Black, female

2006 4/.....	3.0	2.7	2.7	2.6	3.0	*	5.7	7.2	12.3	42.8
2005 4/.....	2.7	2.6	3.2	2.9	4.0	4.2	8.2	9.6	18.5	44.6
2004 4/.....	2.4	2.8	3.1	2.7	3.9	*	6.2	8.9	12.9	50.4
2003 4/.....	2.1	2.8	2.7	3.3	3.0	*	4.7	9.8	20.5	49.4
2002.....	2.6	3.6	3.5	3.3	3.3	4.6	5.4	14.3	17.1	50.5
2001.....	3.7	2.6	2.6	3.2	*	3.9	5.4	12.8	22.1	50.9
2000.....	3.2	2.7	3.0	3.0	3.9	5.8	6.2	10.0	17.4	49.4
1999.....	2.6	3.2	2.7	4.2	*	*	5.9	11.7	16.4	56.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diabetes mellitus (E10-E14)

All races 3/, both sexes

2006.....	6.2	9.6	17.1	28.7	46.4	68.3	98.3	141.9	199.3	285.2
2005.....	5.6	9.9	17.3	29.7	47.3	71.7	104.9	151.4	211.2	312.1
2004.....	5.7	10.0	17.2	29.4	47.2	71.3	105.7	149.8	213.1	307.0
2003.....	5.8	10.3	17.9	30.4	48.9	75.0	108.8	155.3	216.5	317.5
2002.....	6.0	9.9	18.0	30.1	47.5	75.7	108.8	157.7	217.9	320.6
2001.....	5.4	9.9	17.8	29.5	48.4	75.5	108.6	155.5	218.7	321.8
2000.....	5.5	9.5	17.3	29.6	48.1	74.0	108.6	156.5	213.9	319.7
1999.....	5.5	9.9	16.5	29.5	48.9	75.6	109.2	153.5	214.8	317.2

All races 3/, male

2006.....	7.6	12.3	21.3	34.5	56.8	80.7	115.9	167.5	232.2	321.0
2005.....	7.2	12.7	21.0	36.3	56.6	82.4	121.2	174.0	244.1	344.4
2004.....	7.1	12.4	21.2	34.5	56.6	82.7	121.2	172.2	246.9	333.9
2003.....	7.4	12.6	21.3	35.4	57.8	86.8	124.4	179.0	249.7	340.7
2002.....	7.7	11.9	21.5	36.0	55.5	84.3	121.1	175.9	243.5	348.5
2001.....	6.6	12.1	21.3	34.4	54.4	84.7	121.0	171.5	242.3	343.1
2000.....	6.8	11.4	19.7	33.4	53.9	80.8	119.7	173.7	242.5	345.1
1999.....	6.4	12.3	19.4	33.5	53.9	82.5	119.4	172.3	241.3	345.7

All races 3/, female

2006.....	4.7	6.9	13.2	23.3	36.9	57.5	84.0	123.1	178.7	268.4
2005.....	4.1	7.2	13.8	23.4	38.9	62.3	91.6	134.9	190.8	297.2
2004.....	4.3	7.7	13.4	24.6	38.6	61.4	93.2	133.6	192.2	294.9
2003.....	4.2	8.1	14.7	25.7	40.9	64.8	96.2	138.4	196.2	307.2
2002.....	4.4	7.9	14.7	24.5	40.2	68.2	99.0	144.9	202.3	308.4
2001.....	4.2	7.9	14.5	24.9	42.9	67.6	98.8	144.3	204.5	312.9
2000.....	4.3	7.8	14.9	26.1	42.8	68.1	99.9	144.6	197.0	309.4
1999.....	4.6	7.5	13.8	25.8	44.5	69.7	101.2	140.4	199.3	305.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diabetes mellitus (E10-E14)

White, both sexes

2006 4/.....	5.3	8.4	14.7	25.0	40.6	60.2	89.0	130.6	185.8	270.4
2005 4/.....	4.9	8.6	15.0	25.9	41.8	63.3	94.6	139.9	198.7	296.3
2004 4/.....	5.0	8.9	14.6	25.5	41.3	63.6	94.8	137.4	199.5	289.4
2003 4/.....	5.1	8.8	15.2	26.4	42.7	66.5	98.1	143.1	202.7	300.8
2002.....	5.3	8.5	15.4	26.0	41.3	66.0	98.2	145.4	204.9	303.7
2001.....	4.8	8.6	15.3	25.3	42.3	66.8	98.0	143.0	205.9	306.0
2000.....	5.0	8.0	14.6	25.7	42.3	65.2	98.1	143.7	199.1	303.5
1999.....	4.7	8.3	13.8	24.5	41.9	66.9	97.6	141.2	200.8	298.9

White, male

2006 4/.....	6.7	10.9	18.5	30.4	50.9	73.3	107.3	158.5	223.5	312.0
2005 4/.....	6.5	11.3	18.4	32.0	51.0	74.8	111.5	164.6	235.4	336.2
2004 4/.....	6.3	11.1	18.7	30.3	51.0	75.0	111.6	162.1	238.5	323.0
2003 4/.....	6.6	11.1	18.7	31.6	51.5	79.1	115.3	170.2	241.1	333.4
2002.....	6.9	10.3	18.8	32.0	49.5	75.9	112.9	167.9	235.5	344.3
2001.....	6.1	10.5	18.6	30.4	48.8	77.6	113.5	161.8	232.1	337.3
2000.....	6.3	9.7	17.3	29.5	48.7	73.8	111.6	164.0	234.1	337.4
1999.....	5.6	10.6	16.8	28.9	47.7	75.2	111.3	163.7	233.2	333.4

White, female

2006 4/.....	3.9	5.8	10.8	19.9	31.0	48.5	73.9	109.8	162.0	250.8
2005 4/.....	3.3	5.9	11.6	20.0	33.2	53.1	80.7	121.7	175.7	278.0
2004 4/.....	3.6	6.7	10.6	20.8	32.4	53.4	81.1	119.4	175.3	274.3
2003 4/.....	3.5	6.5	11.8	21.4	34.5	55.4	84.0	123.5	179.1	286.3
2002.....	3.8	6.6	12.0	20.2	33.7	57.3	86.3	129.3	186.1	286.1
2001.....	3.4	6.7	12.0	20.4	36.3	57.4	85.4	129.7	190.1	293.0
2000.....	3.7	6.4	12.0	22.1	36.3	57.6	87.3	129.3	178.3	289.7
1999.....	3.9	6.1	11.0	20.4	36.6	59.6	86.7	125.4	181.7	285.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diabetes mellitus (E10-E14)

Black, both sexes

2006 4/.....	12.5	19.1	37.1	61.5	101.7	145.8	193.3	272.8	367.5	471.1
2005 4/.....	10.8	20.7	36.3	63.6	102.9	150.2	211.2	283.6	377.0	506.5
2004 4/.....	11.5	19.2	38.8	64.8	102.0	147.7	219.3	297.6	384.2	532.4
2003 4/.....	11.7	22.1	39.9	69.8	106.9	156.1	221.2	300.0	389.9	527.5
2002.....	12.0	21.4	40.7	67.8	105.5	161.6	221.6	300.3	392.7	537.8
2001.....	10.7	21.0	40.2	69.6	104.4	157.6	226.8	305.2	382.2	529.7
2000.....	10.1	21.9	40.7	66.3	102.4	159.2	223.7	316.5	404.9	524.9
1999.....	11.3	22.5	40.5	72.7	111.6	159.6	229.0	294.9	395.8	529.0

Black, male

2006 4/.....	15.9	23.8	45.3	71.9	120.0	163.3	220.8	297.6	373.8	463.8
2005 4/.....	12.3	25.9	42.2	78.0	118.2	161.8	236.9	310.7	391.9	464.8
2004 4/.....	13.9	24.2	44.1	74.6	115.7	168.6	238.3	309.8	389.5	505.7
2003 4/.....	14.4	25.4	44.0	77.3	121.7	170.6	232.6	306.3	387.6	453.5
2002.....	14.9	25.4	46.4	76.5	118.8	165.3	223.3	295.8	386.7	417.0
2001.....	11.7	25.5	45.6	76.7	112.5	162.1	221.5	308.1	400.5	450.5
2000.....	11.8	25.5	43.5	72.1	108.5	154.3	224.3	311.4	382.7	452.1
1999.....	13.2	27.5	44.1	78.2	113.3	163.8	216.1	282.6	374.1	506.2

Black, female

2006 4/.....	9.6	15.0	30.2	52.9	87.4	132.9	174.3	257.9	364.2	474.1
2005 4/.....	9.5	16.2	31.3	51.7	91.0	141.5	193.7	267.5	369.1	523.3
2004 4/.....	9.4	14.9	34.4	56.8	91.3	132.2	206.5	290.3	381.4	543.0
2003 4/.....	9.3	19.2	36.4	63.6	95.4	145.3	213.6	296.3	391.1	556.2
2002.....	9.4	17.9	35.8	60.6	95.1	158.9	220.4	303.0	395.9	584.1
2001.....	9.8	17.0	35.6	63.9	98.1	154.3	230.3	303.4	372.8	559.1
2000.....	8.7	18.7	38.4	61.5	97.6	162.8	223.3	319.6	416.1	552.0
1999.....	9.7	18.1	37.4	68.3	110.3	156.4	237.6	302.4	406.6	537.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nutritional deficiencies (E40-E64)

All races 3/, both sexes

2006.....	0.1	0.2	0.3	0.5	0.8	1.1	1.9	4.0	7.5	22.7
2005.....	0.1	0.2	0.4	0.5	0.6	1.2	2.7	4.7	10.0	30.8
2004.....	0.1	0.2	0.4	0.4	0.7	1.2	2.1	4.6	10.0	30.3
2003.....	0.1	0.2	0.3	0.5	0.8	1.3	2.5	5.7	10.4	35.5
2002.....	0.1	0.2	0.3	0.5	0.8	1.6	2.7	6.1	12.8	41.5
2001.....	0.1	0.2	0.3	0.5	0.8	1.4	2.8	5.9	12.8	42.9
2000.....	0.1	0.2	0.3	0.5	1.0	2.0	3.0	6.8	15.8	50.0
1999.....	0.2	0.2	0.4	0.6	0.9	1.6	3.2	6.4	15.9	53.9

All races 3/, male

2006.....	*	*	0.3	0.4	0.7	1.1	2.1	3.9	8.1	18.7
2005.....	*	0.2	0.3	0.5	0.7	1.2	2.6	4.5	9.3	27.9
2004.....	*	*	0.4	0.5	0.8	1.3	2.2	4.9	9.6	27.4
2003.....	*	*	0.3	0.5	0.7	1.4	2.7	5.6	10.9	30.6
2002.....	*	0.2	0.3	0.6	0.9	1.6	3.0	6.3	12.0	37.0
2001.....	*	0.3	0.3	0.5	0.9	1.4	2.9	5.6	13.2	35.2
2000.....	*	0.3	0.2	0.7	1.0	2.0	3.4	6.8	14.4	43.7
1999.....	0.2	0.2	0.4	0.7	1.1	1.6	3.8	6.8	16.0	48.4

All races 3/, female

2006.....	*	0.2	0.3	0.5	0.8	1.0	1.8	4.0	7.2	24.5
2005.....	*	0.2	0.4	0.4	0.5	1.2	2.8	5.0	10.4	32.1
2004.....	*	0.2	0.4	0.4	0.7	1.1	2.0	4.3	10.3	31.7
2003.....	*	0.2	0.3	0.5	0.8	1.2	2.3	5.8	10.1	37.6
2002.....	*	*	0.4	0.4	0.7	1.6	2.4	6.0	13.3	43.5
2001.....	*	0.2	0.4	0.4	0.7	1.3	2.7	6.1	12.6	46.1
2000.....	*	0.2	0.4	0.4	0.9	2.0	2.7	6.8	16.6	52.6
1999.....	0.2	0.2	0.4	0.5	0.7	1.5	2.6	6.2	15.8	56.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nutritional deficiencies (E40-E64)

White, both sexes

2006 4/.....	0.1	0.2	0.2	0.4	0.7	0.9	1.8	3.9	7.1	22.1
2005 4/.....	*	0.2	0.3	0.4	0.5	1.1	2.6	4.4	9.4	30.0
2004 4/.....	*	0.1	0.3	0.4	0.6	1.1	1.9	4.2	9.7	29.2
2003 4/.....	*	0.2	0.2	0.4	0.6	1.2	2.3	5.6	9.8	34.2
2002.....	0.1	0.2	0.3	0.4	0.7	1.4	2.5	5.9	12.3	40.2
2001.....	*	0.2	0.3	0.4	0.8	1.2	2.6	5.5	12.0	42.2
2000.....	*	0.2	0.2	0.5	0.8	1.9	2.7	6.4	15.3	48.8
1999.....	0.2	0.2	0.3	0.4	0.8	1.4	2.7	5.9	14.8	52.8

White, male

2006 4/.....	*	*	*	0.3	0.6	0.9	1.9	3.6	7.3	18.2
2005 4/.....	*	*	0.3	0.5	0.6	1.1	2.5	4.1	8.6	27.2
2004 4/.....	*	*	0.3	0.5	0.6	1.0	2.1	4.5	9.1	26.3
2003 4/.....	*	*	*	0.4	0.6	1.3	2.3	5.2	10.1	29.6
2002.....	*	*	0.3	0.4	0.7	1.4	2.8	6.1	11.0	36.3
2001.....	*	*	*	0.4	0.7	1.1	2.6	5.3	12.2	33.8
2000.....	*	0.2	*	0.5	0.8	1.8	2.9	6.2	13.9	42.8
1999.....	*	*	0.3	0.4	1.0	1.3	3.0	6.1	14.7	46.5

White, female

2006 4/.....	*	*	0.3	0.5	0.8	1.0	1.8	4.1	7.0	24.0
2005 4/.....	*	*	0.4	0.4	0.5	1.0	2.6	4.6	9.9	31.4
2004 4/.....	*	*	0.4	0.4	0.6	1.1	1.8	4.1	10.1	30.4
2003 4/.....	*	*	*	0.5	0.7	1.1	2.3	5.9	9.7	36.3
2002.....	*	*	0.4	0.4	0.7	1.3	2.2	5.7	13.1	41.9
2001.....	*	0.2	0.3	0.4	0.8	1.3	2.7	5.7	11.8	45.7
2000.....	*	*	0.3	0.4	0.9	2.0	2.5	6.5	16.2	51.3
1999.....	*	*	0.3	0.4	0.6	1.4	2.4	5.9	14.9	55.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nutritional deficiencies (E40-E64)

Black, both sexes

2006 4/.....	*	*	*	*	*	2.6	3.4	5.0	13.5	30.6
2005 4/.....	*	*	*	*	1.6	2.7	4.4	8.7	18.0	42.5
2004 4/.....	*	*	0.9	*	1.7	2.7	4.2	8.1	14.1	46.8
2003 4/.....	*	*	1.0	*	1.9	*	4.6	9.0	19.5	55.1
2002.....	*	*	*	1.5	2.0	3.5	5.1	10.5	20.7	60.6
2001.....	*	*	1.0	*	*	3.0	4.4	10.8	23.8	54.7
2000.....	*	*	*	*	2.1	3.2	6.5	12.1	22.6	66.9
1999.....	*	*	*	2.0	2.0	3.2	8.3	12.3	30.9	70.8

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	18.5	28.6
2005 4/.....	*	*	*	*	*	*	*	*	19.0	37.2
2004 4/.....	*	*	*	*	*	*	*	10.5	15.1	45.7
2003 4/.....	*	*	*	*	*	*	7.2	13.7	24.4	48.0
2002.....	*	*	*	*	*	*	*	9.6	27.5	47.2
2001.....	*	*	*	*	*	*	6.7	11.6	24.9	57.9
2000.....	*	*	*	*	*	*	9.7	15.1	22.8	63.4
1999.....	*	*	*	*	*	*	12.9	16.1	34.9	75.0

Black, female

2006 4/.....	*	*	*	*	*	*	*	*	10.9	31.4
2005 4/.....	*	*	*	*	*	*	5.0	9.1	17.4	44.6
2004 4/.....	*	*	*	*	*	*	*	6.7	13.6	47.3
2003 4/.....	*	*	*	*	*	*	*	6.2	17.0	57.9
2002.....	*	*	*	*	*	*	4.7	11.0	17.1	65.7
2001.....	*	*	*	*	*	*	*	10.3	23.3	53.5
2000.....	*	*	*	*	*	*	4.4	10.3	22.5	68.3
1999.....	*	*	*	*	*	*	5.2	10.0	29.0	69.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malnutrition (E40-E46)

All races 3/, both sexes

2006.....	0.1	0.2	0.3	0.4	0.7	1.0	1.8	3.7	7.0	20.9
2005.....	0.1	0.2	0.3	0.4	0.6	1.2	2.5	4.6	9.3	28.9
2004.....	0.1	0.2	0.4	0.4	0.7	1.2	2.0	4.4	9.6	28.3
2003.....	0.1	0.2	0.3	0.5	0.7	1.2	2.4	5.5	10.0	33.4
2002.....	0.1	0.2	0.3	0.5	0.7	1.4	2.5	5.6	12.2	38.4
2001.....	0.1	0.2	0.3	0.4	0.8	1.3	2.6	5.5	12.0	40.2
2000.....	0.1	0.2	0.3	0.5	0.9	1.9	2.8	6.4	14.6	46.3
1999.....	0.2	0.2	0.3	0.5	0.8	1.5	3.0	6.0	14.7	50.0

All races 3/, male

2006.....	*	*	0.2	0.4	0.6	1.0	1.9	3.7	7.4	17.5
2005.....	*	0.2	0.3	0.5	0.6	1.2	2.4	4.4	8.8	26.3
2004.....	*	*	0.3	0.5	0.7	1.3	2.2	4.8	9.2	25.3
2003.....	*	*	0.3	0.5	0.7	1.3	2.5	5.4	10.4	28.2
2002.....	*	*	0.3	0.5	0.8	1.4	2.8	5.7	11.4	35.1
2001.....	*	0.2	0.3	0.5	0.8	1.3	2.6	5.1	12.6	33.6
2000.....	*	0.2	*	0.6	0.9	1.8	3.3	6.4	13.1	40.7
1999.....	*	0.2	0.3	0.6	1.1	1.5	3.6	6.2	14.8	44.4

All races 3/, female

2006.....	*	0.2	0.3	0.5	0.7	1.0	1.8	3.8	6.8	22.4
2005.....	*	0.2	0.4	0.3	0.5	1.2	2.6	4.7	9.7	30.2
2004.....	*	0.2	0.4	0.3	0.7	1.1	1.9	4.2	9.9	29.6
2003.....	*	*	0.3	0.5	0.8	1.1	2.2	5.7	9.7	35.6
2002.....	*	*	0.3	0.4	0.6	1.4	2.2	5.6	12.6	39.8
2001.....	*	0.2	0.3	0.4	0.7	1.3	2.5	5.7	11.6	43.0
2000.....	*	0.2	0.3	0.3	0.9	1.9	2.4	6.5	15.5	48.6
1999.....	0.2	0.2	0.3	0.4	0.6	1.4	2.5	5.8	14.7	52.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malnutrition (E40-E46)

White, both sexes

2006 4/.....	0.1	0.2	0.2	0.4	0.7	0.9	1.7	3.7	6.6	20.3
2005 4/.....	*	0.2	0.3	0.4	0.5	1.0	2.4	4.2	8.8	28.2
2004 4/.....	*	0.1	0.3	0.4	0.6	1.0	1.9	4.1	9.3	26.9
2003 4/.....	*	0.1	0.2	0.4	0.6	1.1	2.2	5.4	9.3	32.1
2002.....	0.1	0.2	0.3	0.4	0.6	1.3	2.4	5.4	11.7	37.0
2001.....	*	0.2	0.2	0.4	0.7	1.1	2.4	5.1	11.2	39.5
2000.....	*	0.2	0.2	0.4	0.8	1.8	2.4	6.1	14.2	45.2
1999.....	0.2	0.2	0.3	0.4	0.7	1.3	2.5	5.5	13.8	48.9

White, male

2006 4/.....	*	*	*	0.3	0.6	0.8	1.8	3.4	6.7	16.9
2005 4/.....	*	*	0.3	0.5	0.5	1.0	2.3	4.0	8.1	25.7
2004 4/.....	*	*	0.3	0.4	0.6	1.0	2.0	4.3	8.7	24.1
2003 4/.....	*	*	*	0.4	0.5	1.2	2.2	4.9	9.5	27.2
2002.....	*	*	*	0.4	0.6	1.3	2.7	5.5	10.5	34.3
2001.....	*	*	*	0.4	0.7	1.0	2.3	4.7	11.7	32.2
2000.....	*	*	*	0.5	0.7	1.6	2.8	5.9	12.5	39.7
1999.....	*	*	0.3	0.4	0.9	1.2	2.8	5.6	13.4	42.2

White, female

2006 4/.....	*	*	0.2	0.5	0.7	0.9	1.7	3.9	6.5	21.8
2005 4/.....	*	*	0.3	0.4	0.5	1.0	2.4	4.4	9.2	29.4
2004 4/.....	*	*	0.3	0.4	0.6	1.1	1.7	3.9	9.7	28.2
2003 4/.....	*	*	*	0.5	0.7	1.0	2.2	5.7	9.3	34.2
2002.....	*	*	0.3	0.4	0.6	1.2	2.1	5.3	12.4	38.2
2001.....	*	*	0.3	0.4	0.8	1.2	2.5	5.4	10.9	42.6
2000.....	*	*	0.3	*	0.9	1.9	2.1	6.2	15.1	47.4
1999.....	*	*	0.3	0.4	0.5	1.3	2.3	5.5	14.0	51.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Malnutrition (E40-E46)

Black, both sexes

2006 4/.....	*	*	*	*	*	2.6	3.4	4.4	13.0	29.3
2005 4/.....	*	*	*	*	*	2.7	4.2	8.7	16.7	40.4
2004 4/.....	*	*	0.9	*	*	2.5	4.2	8.1	13.6	46.0
2003 4/.....	*	*	1.0	*	1.8	*	4.4	8.8	19.3	53.4
2002.....	*	*	*	1.5	1.9	3.0	4.4	10.1	19.9	57.6
2001.....	*	*	*	*	*	2.8	4.0	10.1	22.2	51.6
2000.....	*	*	*	*	1.9	3.1	6.5	11.2	20.9	63.8
1999.....	*	*	*	1.6	2.0	3.2	8.0	11.1	28.9	66.0

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	17.1	28.6
2005 4/.....	*	*	*	*	*	*	*	*	17.6	34.4
2004 4/.....	*	*	*	*	*	*	*	10.5	14.3	43.7
2003 4/.....	*	*	*	*	*	*	6.9	13.2	23.6	45.9
2002.....	*	*	*	*	*	*	*	9.2	25.2	46.1
2001.....	*	*	*	*	*	*	*	11.2	24.1	55.6
2000.....	*	*	*	*	*	*	9.7	14.2	22.8	61.1
1999.....	*	*	*	*	*	*	12.2	13.3	33.1	73.8

Black, female

2006 4/.....	*	*	*	*	*	*	*	*	10.9	29.6
2005 4/.....	*	*	*	*	*	*	4.8	9.1	16.3	42.8
2004 4/.....	*	*	*	*	*	*	*	6.7	13.2	46.9
2003 4/.....	*	*	*	*	*	*	*	6.2	17.0	56.3
2002.....	*	*	*	*	*	*	*	10.7	17.1	62.0
2001.....	*	*	*	*	*	*	*	9.4	21.3	50.1
2000.....	*	*	*	*	*	*	4.4	9.4	20.0	64.8
1999.....	*	*	*	*	*	*	5.2	9.7	26.8	63.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other nutritional deficiencies (E50-E64)

All races 3/, both sexes

2006.....	*	*	*	*	*	*	*	*	0.5	1.8
2005.....	*	*	*	*	*	*	*	*	0.7	1.8
2004.....	*	*	*	*	*	*	*	*	0.4	2.1
2003.....	*	*	*	*	*	*	*	*	0.5	2.1
2002.....	*	*	*	*	*	*	*	0.4	0.6	3.1
2001.....	*	*	*	*	*	*	*	0.4	0.8	2.7
2000.....	*	*	*	*	*	*	0.3	0.4	1.2	3.7
1999.....	*	*	*	*	*	*	*	0.4	1.1	3.9

All races 3/, male

2006.....	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	1.6
2004.....	*	*	*	*	*	*	*	*	*	2.1
2003.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	1.9
2001.....	*	*	*	*	*	*	*	*	*	1.6
2000.....	*	*	*	*	*	*	*	*	1.3	3.0
1999.....	*	*	*	*	*	*	*	*	1.3	3.9

All races 3/, female

2006.....	*	*	*	*	*	*	*	*	*	2.1
2005.....	*	*	*	*	*	*	*	*	0.7	1.9
2004.....	*	*	*	*	*	*	*	*	*	2.1
2003.....	*	*	*	*	*	*	*	*	*	2.0
2002.....	*	*	*	*	*	*	*	*	0.6	3.6
2001.....	*	*	*	*	*	*	*	*	1.0	3.1
2000.....	*	*	*	*	*	*	*	*	1.1	4.0
1999.....	*	*	*	*	*	*	*	*	1.1	3.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other nutritional deficiencies (E50-E64)

White, both sexes

2006 4/.....	*	*	*	*	*	*	*	*	0.5	1.9
2005 4/.....	*	*	*	*	*	*	*	*	0.6	1.8
2004 4/.....	*	*	*	*	*	*	*	*	0.4	2.2
2003 4/.....	*	*	*	*	*	*	*	*	0.5	2.2
2002.....	*	*	*	*	*	*	*	0.5	0.6	3.1
2001.....	*	*	*	*	*	*	*	0.4	0.8	2.7
2000.....	*	*	*	*	*	*	0.3	0.3	1.2	3.7
1999.....	*	*	*	*	*	*	*	0.4	1.1	3.9

White, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	1.5
2004 4/.....	*	*	*	*	*	*	*	*	*	2.2
2003 4/.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	2.0
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	1.4	3.2
1999.....	*	*	*	*	*	*	*	*	1.3	4.3

White, female

2006 4/.....	*	*	*	*	*	*	*	*	*	2.2
2005 4/.....	*	*	*	*	*	*	*	*	0.7	1.9
2004 4/.....	*	*	*	*	*	*	*	*	*	2.2
2003 4/.....	*	*	*	*	*	*	*	*	*	2.1
2002.....	*	*	*	*	*	*	*	*	0.7	3.6
2001.....	*	*	*	*	*	*	*	*	0.9	3.1
2000.....	*	*	*	*	*	*	*	*	1.0	3.9
1999.....	*	*	*	*	*	*	*	*	1.0	3.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Meningitis (G00,G03)

All races 3/, both sexes

2006.....	0.2	0.2	0.3	0.2	0.2	0.3	0.5	0.6	0.8	0.9
2005.....	0.2	0.3	0.3	0.4	0.3	0.4	0.4	0.5	0.6	0.9
2004.....	0.2	0.3	0.3	0.3	0.3	0.5	0.5	0.6	0.7	0.9
2003.....	0.2	0.3	0.3	0.3	0.4	0.6	0.5	0.7	0.6	1.1
2002.....	0.2	0.3	0.3	0.3	0.3	0.5	0.7	0.6	0.7	1.2
2001.....	0.2	0.3	0.3	0.4	0.5	0.5	0.5	0.6	1.0	0.9
2000.....	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1	1.2
1999.....	0.2	0.3	0.3	0.3	0.5	0.5	0.7	0.9	1.2	1.5

All races 3/, male

2006.....	*	0.3	0.3	0.2	*	*	0.5	*	0.9	*
2005.....	0.2	0.3	0.3	0.4	0.3	*	*	*	*	*
2004.....	0.3	0.3	0.4	0.3	0.3	*	*	*	*	*
2003.....	*	0.3	0.4	0.4	0.5	0.6	*	0.8	*	*
2002.....	0.2	0.4	0.3	0.4	*	0.6	0.9	0.7	1.0	*
2001.....	*	0.3	0.3	0.4	0.6	0.5	0.7	0.9	1.1	*
2000.....	0.2	0.3	0.4	0.4	0.5	0.6	0.9	1.3	1.1	*
1999.....	0.2	0.4	0.4	0.4	0.4	0.6	0.8	0.9	1.3	1.7

All races 3/, female

2006.....	0.2	0.2	0.2	0.3	*	0.4	0.4	0.5	0.8	0.8
2005.....	0.2	0.2	0.3	0.4	0.3	*	*	0.5	0.6	0.9
2004.....	*	*	0.3	0.3	0.3	0.6	0.6	0.6	0.6	0.7
2003.....	*	0.2	0.2	0.3	0.3	0.6	0.4	0.7	0.6	1.0
2002.....	*	0.2	0.2	*	0.3	0.4	0.6	0.6	*	1.1
2001.....	*	0.2	0.2	0.4	0.4	0.4	0.4	0.5	0.9	0.8
2000.....	*	0.2	0.3	*	*	0.4	0.5	0.6	1.1	1.0
1999.....	*	0.2	*	*	0.5	0.4	0.7	0.9	1.1	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Meningitis (G00,G03)

White, both sexes

2006 4/.....	0.2	0.2	0.3	0.2	0.2	0.3	0.5	0.5	0.8	0.9
2005 4/.....	0.2	0.3	0.2	0.4	0.3	0.3	0.4	0.5	0.6	0.9
2004 4/.....	0.2	0.2	0.2	0.2	0.3	0.4	0.4	0.6	0.6	0.8
2003 4/.....	0.1	0.2	0.2	0.3	0.3	0.5	0.4	0.7	0.6	1.0
2002.....	0.1	0.2	0.2	0.3	0.3	0.5	0.7	0.6	0.6	1.2
2001.....	0.1	0.2	0.2	0.3	0.5	0.4	0.6	0.6	0.9	0.8
2000.....	0.1	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1	1.0
1999.....	0.1	0.2	0.2	0.3	0.4	0.5	0.6	0.7	1.1	1.3

White, male

2006 4/.....	*	0.2	0.3	*	*	*	*	*	*	*
2005 4/.....	*	0.3	0.3	0.4	*	*	*	*	*	*
2004 4/.....	0.2	0.3	0.3	*	*	*	*	*	*	*
2003 4/.....	*	0.3	0.3	0.3	0.4	0.6	*	0.9	*	*
2002.....	*	0.3	*	0.3	*	0.6	0.8	*	*	*
2001.....	*	0.2	0.3	0.4	0.5	*	0.7	0.8	*	*
2000.....	*	0.3	0.3	0.4	0.5	*	0.8	1.2	*	*
1999.....	*	0.3	*	0.4	*	0.5	0.7	*	1.3	*

White, female

2006 4/.....	*	*	0.2	*	*	0.4	*	*	0.7	0.8
2005 4/.....	*	*	*	0.3	*	*	*	*	*	0.9
2004 4/.....	*	*	*	*	*	0.5	0.5	0.6	*	*
2003 4/.....	*	*	*	*	*	0.5	*	0.6	*	0.9
2002.....	*	*	*	*	*	*	0.5	0.5	*	1.2
2001.....	*	*	*	*	0.4	*	*	*	0.8	0.8
2000.....	*	*	*	*	*	*	*	0.6	1.0	1.0
1999.....	*	*	*	*	0.5	*	0.6	0.8	1.0	1.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

All races 3/, both sexes

2006.....	*	*	0.3	0.7	2.1	6.0	19.8	50.1	95.4	144.8
2005.....	*	0.1	0.2	0.8	2.3	6.8	20.3	50.2	98.7	143.7
2004.....	*	*	0.2	0.6	2.0	6.4	18.5	47.6	94.0	135.8
2003.....	*	*	0.2	0.7	2.0	6.4	19.9	49.3	93.3	138.2
2002.....	*	*	0.2	0.6	1.9	6.5	18.4	45.4	89.8	135.2
2001.....	*	*	0.2	0.7	1.8	5.8	18.1	46.4	90.9	134.2
2000.....	*	*	0.2	0.6	1.7	6.6	16.7	44.8	87.6	131.9
1999.....	*	*	0.2	0.6	1.6	5.6	16.7	41.8	82.8	124.4

All races 3/, male

2006.....	*	*	0.4	0.9	2.8	8.5	29.0	76.0	145.6	231.7
2005.....	*	*	0.3	0.9	3.2	9.6	30.0	75.0	153.6	228.6
2004.....	*	*	0.4	0.9	2.6	9.1	27.3	70.6	147.1	220.0
2003.....	*	*	0.2	0.8	2.7	8.9	29.2	74.2	140.7	222.4
2002.....	*	*	0.3	0.8	2.5	9.8	27.3	67.2	136.6	220.1
2001.....	*	*	0.3	1.0	2.7	8.2	26.5	68.8	141.7	224.4
2000.....	*	*	*	0.8	2.2	9.0	24.1	67.4	135.0	222.3
1999.....	*	*	0.3	0.7	2.3	7.7	24.8	64.4	131.3	205.6

All races 3/, female

2006.....	*	*	*	0.4	1.3	3.8	12.2	31.0	64.0	104.1
2005.....	*	*	*	0.6	1.5	4.3	12.5	32.3	64.6	104.7
2004.....	*	*	*	0.4	1.4	4.1	11.4	31.0	61.3	97.9
2003.....	*	*	0.2	0.6	1.4	4.3	12.4	31.5	64.3	101.0
2002.....	*	*	*	0.5	1.4	3.8	11.3	30.0	61.3	98.4
2001.....	*	*	*	0.4	1.0	3.7	11.5	30.7	60.4	96.4
2000.....	*	*	*	0.4	1.3	4.4	10.9	29.1	59.5	95.0
1999.....	*	*	*	0.4	0.9	3.9	10.4	26.0	54.4	91.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

White, both sexes

2006 4/.....	*	*	0.2	0.7	2.1	6.3	21.2	53.5	101.9	154.1
2005 4/.....	*	0.1	0.2	0.8	2.4	7.1	21.8	53.7	104.4	152.9
2004 4/.....	*	*	0.2	0.7	2.1	6.8	20.2	50.6	99.9	143.1
2003 4/.....	*	*	0.2	0.7	2.0	6.8	21.2	52.4	99.0	146.0
2002.....	*	*	0.2	0.6	2.1	6.9	19.6	48.3	95.7	143.3
2001.....	*	*	0.2	0.7	1.9	6.2	19.4	49.6	96.4	141.8
2000.....	*	*	0.2	0.6	1.9	7.1	17.8	47.5	92.5	139.5
1999.....	*	*	0.2	0.6	1.6	6.0	17.7	44.1	87.4	131.2

White, male

2006 4/.....	*	*	0.4	1.0	2.9	8.8	30.8	80.5	154.3	247.3
2005 4/.....	*	*	0.3	1.0	3.3	10.0	31.9	79.5	162.2	242.7
2004 4/.....	*	*	0.4	1.0	2.7	9.6	29.4	74.7	156.1	231.3
2003 4/.....	*	*	*	0.8	2.7	9.4	30.9	78.4	148.9	233.3
2002.....	*	*	0.3	0.8	2.7	10.2	28.8	71.2	145.0	232.7
2001.....	*	*	0.3	1.0	3.0	8.8	28.1	73.4	149.6	237.6
2000.....	*	*	*	0.9	2.5	9.6	25.5	70.7	141.5	235.3
1999.....	*	*	0.3	0.7	2.4	8.1	26.4	67.7	138.6	216.7

White, female

2006 4/.....	*	*	*	0.5	1.4	4.0	13.2	33.4	68.8	110.3
2005 4/.....	*	*	*	0.7	1.6	4.6	13.4	34.6	68.2	111.5
2004 4/.....	*	*	*	0.4	1.5	4.3	12.6	33.0	65.0	103.3
2003 4/.....	*	*	*	0.6	1.4	4.5	13.3	33.6	68.2	107.4
2002.....	*	*	*	0.5	1.5	3.9	12.0	31.9	65.5	104.4
2001.....	*	*	*	0.5	0.9	4.0	12.3	32.7	64.2	101.7
2000.....	*	*	*	0.4	1.4	4.8	11.7	31.2	63.3	100.6
1999.....	*	*	*	0.4	0.9	4.2	10.8	27.6	57.3	96.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Parkinson's disease (G20-G21)

Black, both sexes

2006 4/.....	*	*	*	*	1.5	4.1	11.5	25.0	37.2	51.4
2005 4/.....	*	*	*	*	*	4.3	10.9	22.7	50.0	56.7
2004 4/.....	*	*	*	*	*	3.9	7.2	22.3	39.1	61.2
2003 4/.....	*	*	*	*	1.9	4.1	10.7	22.8	38.3	52.8
2002.....	*	*	*	*	*	4.6	9.5	20.6	32.7	51.7
2001.....	*	*	*	*	*	2.7	8.9	18.9	34.9	59.0
2000.....	*	*	*	*	*	3.1	8.8	21.0	37.6	51.6
1999.....	*	*	*	*	*	3.6	8.7	19.5	32.4	52.0

Black, male

2006 4/.....	*	*	*	*	*	6.6	18.5	42.5	63.0	70.6
2005 4/.....	*	*	*	*	*	7.4	18.7	36.2	82.2	97.6
2004 4/.....	*	*	*	*	*	5.3	12.6	34.1	58.1	106.3
2003 4/.....	*	*	*	*	*	5.4	16.8	37.0	59.1	97.0
2002.....	*	*	*	*	*	7.5	16.0	30.3	55.0	89.0
2001.....	*	*	*	*	*	*	14.8	25.6	57.8	99.8
2000.....	*	*	*	*	*	*	13.7	37.8	65.8	86.5
1999.....	*	*	*	*	*	5.7	12.6	31.8	48.8	86.7

Black, female

2006 4/.....	*	*	*	*	*	*	6.8	14.5	23.5	43.5
2005 4/.....	*	*	*	*	*	*	5.6	14.7	33.0	40.1
2004 4/.....	*	*	*	*	*	*	*	15.4	29.1	43.3
2003 4/.....	*	*	*	*	*	*	6.6	14.4	27.5	35.6
2002.....	*	*	*	*	*	*	5.2	14.8	21.1	37.4
2001.....	*	*	*	*	*	*	4.9	15.0	23.3	43.8
2000.....	*	*	*	*	*	*	5.5	10.8	23.4	38.6
1999.....	*	*	*	*	*	*	6.1	12.0	24.2	39.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alzheimer's disease (G30)

All races 3/, both sexes

2006.....	*	0.1	0.4	1.1	3.4	9.6	32.9	101.2	272.6	848.3
2005.....	*	0.1	0.3	1.1	3.5	9.7	33.3	102.5	275.5	861.6
2004.....	*	0.2	0.3	0.9	3.1	10.1	31.0	99.4	261.0	818.8
2003.....	*	0.1	0.3	1.0	3.3	10.0	33.3	98.6	255.0	802.4
2002.....	*	*	0.2	0.9	3.2	9.8	30.6	95.4	245.7	752.3
2001.....	*	*	0.3	1.2	3.3	9.0	29.2	88.5	232.5	710.3
2000.....	*	*	0.3	1.1	3.2	9.1	28.9	85.2	221.1	667.7
1999.....	*	0.1	0.2	1.0	3.0	8.7	26.7	78.6	206.2	601.3

All races 3/, male

2006.....	*	*	0.4	1.0	3.4	10.3	31.6	95.1	244.5	638.6
2005.....	*	*	0.2	1.0	3.2	9.6	31.4	95.9	246.9	639.9
2004.....	*	*	0.3	0.8	3.1	9.4	29.6	89.3	236.4	619.6
2003.....	*	*	0.3	1.0	2.9	10.1	32.1	92.1	231.2	606.5
2002.....	*	*	0.2	0.8	3.1	9.2	29.5	90.9	220.6	572.0
2001.....	*	*	0.3	0.9	3.2	9.0	27.7	85.6	216.6	556.1
2000.....	*	*	0.3	0.9	2.9	9.5	29.1	79.9	206.8	522.1
1999.....	*	*	*	1.1	3.0	8.8	26.9	77.4	198.2	485.7

All races 3/, female

2006.....	*	*	0.3	1.3	3.3	9.0	34.0	105.7	290.1	946.4
2005.....	*	*	0.4	1.2	3.7	9.8	34.8	107.3	293.2	963.4
2004.....	*	0.2	0.4	1.0	3.1	10.7	32.1	106.7	276.2	908.4
2003.....	*	*	0.3	0.9	3.7	10.0	34.2	103.2	269.5	889.0
2002.....	*	*	*	1.0	3.3	10.3	31.5	98.5	260.9	830.5
2001.....	*	*	0.3	1.4	3.4	9.0	30.4	90.5	242.1	774.9
2000.....	*	*	0.3	1.2	3.5	8.7	28.8	88.9	229.5	727.0
1999.....	*	*	0.3	0.8	3.0	8.6	26.6	79.4	210.9	648.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alzheimer's disease (G30)

White, both sexes

2006 4/.....	*	0.2	0.4	1.2	3.6	10.2	34.2	104.8	282.8	878.9
2005 4/.....	*	*	0.3	1.2	3.7	10.0	34.1	105.3	284.3	891.6
2004 4/.....	*	0.2	0.4	1.0	3.3	10.4	32.0	102.3	268.5	846.8
2003 4/.....	*	*	0.3	1.1	3.6	10.3	34.5	101.8	264.4	834.0
2002.....	*	*	0.2	1.0	3.4	10.3	31.9	99.1	255.3	781.8
2001.....	*	*	0.3	1.3	3.6	9.3	30.6	90.9	240.5	738.1
2000.....	*	*	0.3	1.1	3.4	9.6	29.9	88.1	229.1	695.3
1999.....	*	*	0.2	1.0	3.3	9.1	27.6	81.3	213.6	627.9

White, male

2006 4/.....	*	*	0.4	1.1	3.6	10.9	32.6	98.3	253.6	662.5
2005 4/.....	*	*	0.2	1.1	3.5	9.8	31.9	97.5	254.3	661.6
2004 4/.....	*	*	0.4	0.9	3.3	9.7	30.2	91.6	243.9	640.9
2003 4/.....	*	*	0.3	1.1	3.0	10.3	33.1	94.9	238.5	630.3
2002.....	*	*	*	0.8	3.3	9.4	30.5	94.3	227.6	594.8
2001.....	*	*	0.3	0.9	3.4	9.2	28.8	88.1	222.7	577.2
2000.....	*	*	0.3	1.0	3.1	10.1	29.9	82.6	214.4	542.1
1999.....	*	*	*	1.1	3.3	9.2	27.6	80.1	205.6	506.9

White, female

2006 4/.....	*	*	0.3	1.4	3.5	9.7	35.5	109.7	301.2	980.5
2005 4/.....	*	*	0.4	1.4	3.9	10.2	35.8	111.1	303.1	997.5
2004 4/.....	*	0.3	0.4	1.1	3.3	11.1	33.5	110.2	283.8	939.6
2003 4/.....	*	*	0.4	1.0	4.2	10.3	35.7	106.8	280.3	924.3
2002.....	*	*	*	1.2	3.5	11.0	33.1	102.6	272.2	863.0
2001.....	*	*	0.4	1.6	3.8	9.4	32.1	92.9	251.3	805.4
2000.....	*	*	0.3	1.3	3.7	9.2	29.9	92.0	237.7	757.6
1999.....	*	*	*	0.9	3.3	9.1	27.5	82.1	218.3	676.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alzheimer's disease (G30)

Black, both sexes

2006 4/.....	*	*	*	*	3.1	7.6	32.6	90.5	226.0	663.9
2005 4/.....	*	*	*	*	2.8	10.3	37.0	100.4	241.5	689.5
2004 4/.....	*	*	*	*	2.2	10.3	30.6	96.0	236.7	657.9
2003 4/.....	*	*	*	*	1.8	10.5	30.2	91.4	207.9	583.8
2002.....	*	*	*	*	2.5	8.5	27.2	81.3	186.8	543.7
2001.....	*	*	*	*	1.9	8.7	23.3	86.8	183.6	506.7
2000.....	*	*	*	*	2.1	7.4	25.9	71.1	171.7	445.1
1999.....	*	*	*	*	*	6.9	25.3	65.0	152.0	372.4

Black, male

2006 4/.....	*	*	*	*	3.5	8.4	32.9	88.6	200.6	470.9
2005 4/.....	*	*	*	*	*	11.6	36.1	103.7	222.6	518.7
2004 4/.....	*	*	*	*	*	9.4	32.0	89.6	210.9	497.7
2003 4/.....	*	*	*	*	*	11.1	32.5	93.1	206.0	459.8
2002.....	*	*	*	*	*	9.5	29.4	82.2	189.5	412.8
2001.....	*	*	*	*	*	9.9	24.2	84.6	185.4	414.1
2000.....	*	*	*	*	*	7.8	27.7	64.7	156.0	371.3
1999.....	*	*	*	*	*	8.0	27.5	62.7	139.5	317.5

Black, female

2006 4/.....	*	*	*	*	2.8	6.9	32.3	91.6	239.4	742.8
2005 4/.....	*	*	*	*	3.5	9.3	37.7	98.4	251.5	758.5
2004 4/.....	*	*	*	*	*	11.0	29.7	99.7	250.2	721.4
2003 4/.....	*	*	*	*	*	10.1	28.6	90.3	208.9	631.9
2002.....	*	*	*	*	*	7.8	25.8	80.7	185.4	593.9
2001.....	*	*	*	*	*	7.9	22.6	88.2	182.7	541.0
2000.....	*	*	*	*	*	7.1	24.6	74.9	179.6	472.6
1999.....	*	*	*	*	*	6.0	23.9	66.4	158.2	392.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

Major cardiovascular diseases (I00-I78)

All races 3/, both sexes

2006.....	46.5	83.2	137.0	205.7	323.5	490.4	798.7	1,368.3	2,468.6	5,928.2
2005.....	47.2	84.2	140.0	208.2	336.6	515.2	848.2	1,452.7	2,635.2	6,398.5
2004.....	47.8	86.0	139.2	211.1	343.0	537.5	885.1	1,514.3	2,730.4	6,616.6
2003.....	50.3	87.7	142.8	224.5	363.7	574.0	946.1	1,627.3	2,918.8	7,138.0
2002.....	49.7	87.9	144.2	233.5	374.0	602.6	993.2	1,700.2	3,034.4	7,389.6
2001.....	48.8	85.6	144.9	234.9	385.7	619.6	1,017.5	1,759.4	3,133.9	7,673.7
2000.....	49.6	88.5	146.7	248.4	407.5	647.9	1,066.0	1,824.4	3,253.7	8,030.3
1999.....	50.0	88.3	148.7	254.8	416.0	678.9	1,111.3	1,888.7	3,356.4	8,197.1

All races 3/, male

2006.....	62.7	115.5	195.7	291.1	443.8	657.9	1,027.7	1,713.4	2,982.2	6,151.8
2005.....	64.1	116.8	199.9	294.8	462.0	679.9	1,093.3	1,820.2	3,159.0	6,615.9
2004.....	64.6	120.2	198.5	296.2	472.0	714.7	1,134.9	1,900.8	3,283.3	6,818.1
2003.....	67.7	123.1	204.5	314.7	497.9	765.9	1,225.3	2,051.8	3,517.1	7,329.3
2002.....	67.9	123.9	206.1	326.7	512.6	802.0	1,287.4	2,140.7	3,633.4	7,638.8
2001.....	66.5	121.4	204.7	327.6	526.3	821.4	1,325.6	2,223.2	3,774.2	7,936.0
2000.....	68.0	126.8	209.0	350.6	556.9	857.2	1,403.8	2,305.7	3,956.7	8,448.1
1999.....	68.9	125.7	212.5	358.4	568.4	908.7	1,461.9	2,392.7	4,078.9	8,603.2

All races 3/, female

2006.....	30.5	51.6	80.8	125.2	213.5	344.0	612.4	1,115.8	2,147.6	5,823.6
2005.....	30.5	52.4	82.5	126.6	222.2	371.4	649.7	1,185.8	2,309.9	6,298.7
2004.....	31.2	52.6	82.4	130.8	225.5	383.3	684.0	1,236.7	2,389.8	6,526.0
2003.....	33.2	53.3	83.7	139.7	241.8	407.6	722.2	1,325.1	2,552.9	7,053.5
2002.....	31.6	52.9	85.0	146.0	248.2	430.4	758.0	1,389.0	2,670.1	7,281.5
2001.....	31.5	50.7	87.6	148.1	258.3	445.9	772.7	1,434.7	2,750.2	7,563.9
2000.....	31.5	51.4	87.0	152.8	272.0	468.5	799.9	1,489.2	2,839.0	7,860.1
1999.....	31.4	52.1	87.5	158.0	278.2	482.2	837.3	1,538.3	2,933.7	8,033.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Major cardiovascular diseases (I00-I78)

White, both sexes

2006 4/.....	41.1	73.3	120.8	185.1	298.3	457.6	764.6	1,340.7	2,457.3	6,029.7
2005 4/.....	41.4	73.7	122.4	187.2	309.5	483.0	814.8	1,420.7	2,623.8	6,507.8
2004 4/.....	41.9	74.9	122.1	189.5	316.2	503.9	851.4	1,479.8	2,721.9	6,704.8
2003 4/.....	43.4	76.1	124.8	203.3	333.1	537.6	907.0	1,593.5	2,909.9	7,233.1
2002.....	43.6	76.6	127.4	211.8	344.7	567.9	957.1	1,663.0	3,023.3	7,482.7
2001.....	41.6	74.2	127.7	212.4	355.6	580.3	980.2	1,721.1	3,122.8	7,770.6
2000.....	42.6	76.5	130.6	225.6	376.3	612.7	1,027.2	1,783.3	3,243.6	8,133.9
1999.....	42.4	76.6	132.4	230.2	385.3	644.7	1,068.3	1,848.2	3,345.3	8,299.2

White, male

2006 4/.....	57.2	104.4	176.1	265.2	412.4	618.6	988.0	1,685.6	2,988.3	6,287.1
2005 4/.....	58.0	105.2	179.2	268.7	430.2	643.2	1,057.0	1,793.2	3,164.9	6,772.6
2004 4/.....	59.0	107.8	179.1	270.3	439.8	677.7	1,101.5	1,869.3	3,298.4	6,940.8
2003 4/.....	60.6	110.4	183.6	289.9	463.8	727.3	1,183.5	2,022.8	3,520.9	7,466.8
2002.....	62.0	111.9	186.7	301.4	479.1	765.2	1,253.1	2,112.9	3,642.0	7,766.6
2001.....	59.1	109.0	185.6	301.7	493.8	781.8	1,286.2	2,192.9	3,777.7	8,060.7
2000.....	60.7	114.5	190.5	323.5	523.7	822.7	1,366.4	2,273.2	3,963.6	8,585.0
1999.....	60.8	113.0	193.5	329.9	533.4	874.1	1,420.2	2,361.9	4,089.2	8,717.5

White, female

2006 4/.....	24.7	42.4	66.4	108.0	191.9	314.2	579.2	1,083.2	2,121.9	5,908.9
2005 4/.....	24.6	42.4	66.8	108.8	197.3	340.5	614.5	1,145.0	2,284.8	6,385.8
2004 4/.....	24.6	42.1	66.4	111.9	201.6	349.8	645.9	1,194.8	2,363.8	6,598.4
2003 4/.....	26.2	42.1	67.2	120.1	212.2	370.0	680.7	1,283.1	2,533.3	7,129.6
2002.....	25.2	41.6	69.3	126.1	220.5	394.1	715.8	1,341.0	2,644.0	7,359.3
2001.....	24.0	39.8	71.0	127.1	228.1	403.3	732.3	1,386.9	2,726.9	7,649.2
2000.....	24.4	39.0	72.0	132.4	240.4	429.2	755.0	1,438.7	2,815.1	7,950.6
1999.....	24.0	40.6	72.8	135.3	248.8	444.5	788.7	1,487.4	2,906.2	8,131.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Major cardiovascular diseases (I00-I78)

Black, both sexes

2006 4/.....	90.5	165.7	284.7	411.9	611.0	857.9	1,252.9	1,891.4	2,978.3	5,587.2
2005 4/.....	93.7	172.2	295.3	422.3	640.6	882.2	1,301.4	2,041.3	3,168.4	6,016.9
2004 4/.....	97.5	179.4	293.5	434.4	650.7	923.3	1,374.7	2,132.1	3,217.2	6,368.6
2003 4/.....	104.2	183.8	305.8	451.7	695.7	983.6	1,489.0	2,249.7	3,405.2	6,862.0
2002.....	99.1	184.1	302.3	468.4	697.7	1,001.3	1,510.2	2,343.9	3,555.3	7,081.7
2001.....	105.7	181.9	307.2	471.1	713.7	1,057.5	1,548.8	2,422.8	3,623.2	7,316.7
2000.....	104.9	192.0	303.5	489.3	736.9	1,061.9	1,623.2	2,496.2	3,719.8	7,518.9
1999.....	109.8	188.5	311.0	509.4	743.3	1,093.2	1,704.6	2,553.3	3,787.7	7,582.5

Black, male

2006 4/.....	112.0	216.9	391.0	571.6	846.5	1,157.4	1,641.0	2,380.2	3,493.9	5,418.9
2005 4/.....	117.6	224.6	402.0	585.8	873.4	1,156.7	1,667.3	2,508.7	3,664.5	5,707.6
2004 4/.....	118.5	234.5	393.3	594.3	890.6	1,210.4	1,729.0	2,625.1	3,677.3	6,198.0
2003 4/.....	129.8	239.7	412.9	610.0	924.3	1,273.7	1,900.2	2,789.8	4,035.4	6,547.7
2002.....	122.7	237.2	406.5	633.2	934.4	1,292.7	1,908.8	2,803.5	4,093.1	6,929.0
2001.....	129.8	237.2	405.7	627.8	934.4	1,342.7	1,988.0	2,943.9	4,284.2	7,363.9
2000.....	131.7	247.2	407.0	670.8	955.1	1,328.0	2,040.7	2,999.5	4,397.8	7,687.4
1999.....	138.3	246.5	423.5	686.4	993.1	1,400.0	2,146.5	3,046.5	4,407.7	7,921.1

Black, female

2006 4/.....	71.5	121.3	194.8	280.3	427.5	636.8	986.3	1,598.9	2,705.9	5,656.0
2005 4/.....	72.7	126.5	205.1	287.3	459.1	678.7	1,051.0	1,762.4	2,906.3	6,141.8
2004 4/.....	79.0	131.5	209.1	302.8	463.7	710.0	1,134.3	1,839.9	2,974.8	6,436.3
2003 4/.....	81.7	135.2	215.1	321.5	517.3	768.3	1,211.6	1,928.7	3,075.0	6,984.1
2002.....	78.2	138.0	213.9	333.0	512.5	785.0	1,242.7	2,069.3	3,275.3	7,140.2
2001.....	84.3	134.0	223.4	342.6	540.2	846.9	1,256.2	2,111.8	3,286.5	7,299.2
2000.....	81.2	144.0	215.2	340.5	564.6	865.5	1,346.0	2,192.8	3,378.3	7,456.2
1999.....	84.7	138.5	215.6	364.7	547.4	865.8	1,411.4	2,256.2	3,480.4	7,456.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

Diseases of heart (I00-I09,I11,I13,I20-I51)

All races 3/, both sexes

2006.....	37.3	67.4	111.0	167.7	261.3	386.9	615.9	1,031.7	1,841.0	4,480.8
2005.....	37.7	68.0	114.1	171.0	273.2	405.5	654.0	1,090.7	1,946.8	4,778.4
2004.....	38.3	69.6	113.7	173.4	278.1	423.7	679.6	1,130.0	2,007.8	4,895.9
2003.....	40.5	71.3	116.6	185.0	296.1	456.3	730.9	1,218.3	2,151.6	5,278.4
2002.....	40.2	71.3	119.2	192.8	304.4	479.6	766.1	1,278.2	2,234.3	5,446.8
2001.....	39.1	69.3	119.5	194.2	314.1	494.4	787.7	1,321.2	2,309.0	5,664.2
2000.....	39.4	71.2	120.4	205.5	330.6	515.8	826.7	1,372.3	2,392.1	5,926.1
1999.....	40.1	71.7	123.5	211.0	341.9	547.1	867.9	1,427.8	2,484.7	6,063.0

All races 3/, male

2006.....	52.4	97.4	164.1	244.4	369.3	534.4	819.7	1,341.1	2,319.5	4,819.9
2005.....	53.6	98.7	168.8	249.4	385.1	549.7	869.1	1,414.1	2,428.3	5,143.4
2004.....	54.2	101.5	167.5	252.3	393.5	580.0	899.4	1,472.2	2,509.6	5,239.3
2003.....	56.9	104.0	173.3	268.1	416.1	624.7	975.4	1,589.0	2,695.8	5,621.5
2002.....	57.7	104.7	177.0	278.2	429.1	654.3	1,025.8	1,667.3	2,786.9	5,823.5
2001.....	55.8	102.6	175.2	279.7	440.9	672.2	1,054.4	1,732.2	2,882.7	6,040.5
2000.....	57.0	107.1	178.1	298.9	463.9	702.0	1,119.7	1,787.5	3,012.3	6,430.0
1999.....	58.1	107.0	182.9	306.5	478.2	752.8	1,174.1	1,865.5	3,123.3	6,561.6

All races 3/, female

2006.....	22.4	38.1	60.0	95.4	162.6	258.0	450.1	805.2	1,541.9	4,322.1
2005.....	21.8	38.0	61.6	97.1	171.2	279.6	479.8	856.0	1,647.8	4,610.8
2004.....	22.7	38.5	62.2	99.1	173.2	287.7	502.6	884.1	1,698.7	4,741.5
2003.....	24.3	39.5	62.4	106.7	187.0	310.3	534.8	954.5	1,818.9	5,126.7
2002.....	22.9	38.8	63.9	112.5	191.2	328.8	558.5	1,003.3	1,898.3	5,283.3
2001.....	22.6	37.0	66.1	114.2	199.0	341.4	575.8	1,033.5	1,965.3	5,506.8
2000.....	22.2	36.4	65.0	118.1	209.8	356.3	596.0	1,083.0	2,026.2	5,720.9
1999.....	22.4	37.5	66.6	121.7	218.6	371.2	628.7	1,123.5	2,111.0	5,862.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of heart (I00-I09,I11,I13,I20-I51)

White, both sexes

2006 4/.....	34.1	60.9	100.6	154.3	245.0	366.0	594.8	1,017.2	1,841.6	4,576.6
2005 4/.....	34.0	61.3	102.5	156.6	255.0	385.2	634.7	1,073.2	1,946.5	4,879.3
2004 4/.....	34.4	62.4	102.3	158.8	261.2	402.0	658.9	1,112.1	2,010.3	4,977.6
2003 4/.....	35.9	63.6	104.7	170.9	275.7	432.1	706.5	1,199.6	2,153.0	5,363.5
2002.....	36.3	64.0	108.1	177.9	284.5	457.0	744.1	1,255.7	2,234.0	5,529.1
2001.....	34.2	61.8	108.6	178.5	294.7	468.4	765.0	1,297.7	2,308.3	5,752.1
2000.....	34.9	63.7	109.9	190.0	309.5	492.6	801.0	1,347.0	2,393.0	6,020.8
1999.....	34.9	64.1	112.7	193.7	321.3	524.0	839.7	1,403.0	2,484.1	6,152.7

White, male

2006 4/.....	49.2	89.8	151.8	227.9	348.6	509.3	795.9	1,326.3	2,335.5	4,947.1
2005 4/.....	49.8	91.0	155.3	231.6	364.2	527.6	848.6	1,402.3	2,445.7	5,288.4
2004 4/.....	50.5	93.4	154.7	234.6	373.5	556.2	880.1	1,458.9	2,532.6	5,348.4
2003 4/.....	52.1	95.4	159.4	251.9	394.1	599.0	949.5	1,575.7	2,710.0	5,747.2
2002.....	53.9	96.8	164.4	261.0	406.6	631.3	1,006.4	1,654.3	2,805.0	5,939.8
2001.....	50.9	94.4	163.4	261.7	420.5	647.2	1,031.2	1,716.7	2,896.4	6,157.2
2000.....	52.3	99.4	166.1	280.7	442.0	680.1	1,096.8	1,771.0	3,030.0	6,560.8
1999.....	52.4	98.8	170.3	286.1	454.8	730.0	1,148.4	1,851.7	3,144.4	6,669.0

White, female

2006 4/.....	18.8	32.0	50.3	83.4	148.5	238.3	427.9	786.4	1,529.5	4,402.6
2005 4/.....	18.0	31.8	50.8	84.4	153.5	258.6	457.9	829.6	1,633.6	4,691.0
2004 4/.....	18.2	31.5	51.0	85.9	157.2	265.2	477.1	858.3	1,685.8	4,810.4
2003 4/.....	19.6	32.1	51.1	93.1	166.1	284.6	507.6	927.7	1,809.7	5,193.6
2002.....	18.6	31.4	53.1	98.3	171.5	303.6	530.3	970.4	1,884.1	5,350.6
2001.....	17.5	29.4	54.9	99.1	178.6	311.5	549.5	1,001.0	1,952.9	5,582.5
2000.....	17.5	28.3	54.9	103.7	187.4	328.6	563.7	1,048.7	2,014.1	5,801.4
1999.....	17.4	29.7	56.4	105.8	198.3	344.3	594.5	1,088.0	2,094.3	5,945.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of heart (I00-I09,I11,I13,I20-I51)

Black, both sexes

2006 4/.....	66.5	126.1	213.2	313.8	464.1	644.1	934.7	1,382.9	2,157.4	4,056.3
2005 4/.....	69.4	128.5	224.1	326.8	494.7	662.2	958.8	1,494.7	2,282.2	4,326.7
2004 4/.....	73.8	134.9	226.0	337.4	492.5	697.0	1,020.9	1,534.2	2,297.3	4,563.8
2003 4/.....	79.0	139.6	232.4	348.4	533.8	752.7	1,115.2	1,638.3	2,445.2	4,941.2
2002.....	74.8	138.7	232.1	365.8	542.7	764.2	1,126.8	1,715.1	2,557.1	5,115.2
2001.....	79.5	137.2	232.2	369.7	546.5	809.7	1,160.0	1,780.6	2,619.7	5,275.7
2000.....	77.5	142.2	231.2	380.7	571.0	813.9	1,230.4	1,836.2	2,673.7	5,389.5
1999.....	82.9	143.0	240.2	400.6	579.9	852.1	1,289.1	1,878.6	2,749.2	5,487.4

Black, male

2006 4/.....	85.6	172.1	299.5	440.4	659.0	887.9	1,250.3	1,808.3	2,621.8	4,073.1
2005 4/.....	91.1	175.9	312.9	461.0	679.8	877.7	1,259.7	1,881.5	2,710.8	4,230.5
2004 4/.....	93.6	182.7	310.5	471.5	685.8	932.8	1,315.6	1,939.0	2,703.1	4,637.3
2003 4/.....	102.7	188.1	322.4	481.2	718.0	997.2	1,460.0	2,083.2	2,981.9	4,850.3
2002.....	96.7	186.2	319.6	503.6	740.6	1,001.5	1,448.8	2,095.3	3,039.3	5,125.7
2001.....	101.9	184.4	314.7	503.1	731.0	1,039.8	1,518.8	2,214.9	3,181.5	5,459.9
2000.....	101.3	190.8	319.1	533.0	754.7	1,033.7	1,570.9	2,245.4	3,226.1	5,633.5
1999.....	109.3	193.3	335.5	556.7	789.2	1,111.8	1,652.9	2,277.2	3,259.4	5,858.8

Black, female

2006 4/.....	49.7	86.1	140.3	209.5	312.2	464.0	717.9	1,128.4	1,912.0	4,049.4
2005 4/.....	50.2	87.1	149.0	216.0	350.4	502.4	753.0	1,264.0	2,055.7	4,365.6
2004 4/.....	56.4	93.3	154.5	227.0	341.8	521.9	820.9	1,294.4	2,083.5	4,534.7
2003 4/.....	58.2	97.4	156.3	239.2	390.1	571.3	882.6	1,373.9	2,164.0	4,976.5
2002.....	55.5	97.5	157.9	252.6	388.0	588.1	910.7	1,488.0	2,306.1	5,111.2
2001.....	59.7	96.3	162.0	260.3	401.4	639.8	921.0	1,521.3	2,333.5	5,207.3
2000.....	56.5	100.1	156.3	255.9	426.1	651.7	1,004.3	1,589.5	2,395.4	5,298.7
1999.....	59.6	99.6	159.4	273.1	415.8	659.7	1,047.8	1,638.6	2,496.4	5,348.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races 3/, both sexes

2006.....	0.2	0.4	0.4	0.9	1.5	2.5	4.1	6.8	11.0	16.7
2005.....	0.2	0.4	0.5	0.9	1.5	2.5	4.2	6.7	10.9	19.0
2004.....	0.3	0.4	0.6	0.9	1.4	2.6	4.2	7.3	10.9	17.1
2003.....	0.3	0.5	0.6	1.0	1.8	3.0	4.6	8.1	11.6	20.5
2002.....	0.2	0.5	0.7	1.3	2.1	2.8	5.0	8.4	11.4	17.8
2001.....	0.4	0.5	0.7	1.2	1.7	3.2	5.0	8.1	11.4	17.7
2000.....	0.4	0.6	0.8	1.4	2.2	3.4	5.6	8.1	11.6	16.8
1999.....	0.3	0.5	0.9	1.4	2.1	3.5	5.8	8.4	12.6	18.0

All races 3/, male

2006.....	*	0.4	0.4	0.7	1.2	1.9	3.0	5.1	7.7	12.0
2005.....	0.2	0.3	0.5	0.9	1.1	1.9	3.1	4.9	8.0	14.0
2004.....	0.2	0.4	0.5	0.7	0.9	2.1	2.8	5.7	9.5	11.5
2003.....	0.2	0.4	0.4	0.8	1.5	2.3	3.6	6.4	9.0	16.2
2002.....	0.2	0.4	0.6	1.0	1.5	1.9	4.2	5.8	8.2	11.8
2001.....	0.3	0.4	0.5	1.1	1.2	2.4	3.2	6.0	7.7	12.8
2000.....	0.3	0.5	0.6	1.1	1.8	2.2	3.6	5.8	7.8	12.2
1999.....	0.3	0.4	0.6	1.0	1.4	2.4	4.7	5.4	8.9	10.3

All races 3/, female

2006.....	0.2	0.4	0.5	1.0	1.7	3.0	5.0	8.0	13.0	18.9
2005.....	0.3	0.5	0.6	1.0	1.9	3.0	5.1	8.0	12.7	21.3
2004.....	0.3	0.4	0.7	1.1	1.9	3.0	5.3	8.3	11.8	19.5
2003.....	0.3	0.6	0.8	1.1	2.1	3.6	5.5	9.3	13.2	22.3
2002.....	0.3	0.5	0.9	1.5	2.6	3.7	5.6	10.2	13.4	20.4
2001.....	0.4	0.6	0.9	1.3	2.2	3.9	6.4	9.5	13.7	19.8
2000.....	0.5	0.7	0.9	1.6	2.5	4.5	7.1	9.7	13.9	18.7
1999.....	0.4	0.6	1.1	1.8	2.7	4.5	6.7	10.4	14.7	21.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

White, both sexes

2006 4/.....	0.2	0.3	0.4	0.8	1.3	2.5	4.2	7.1	11.6	18.0
2005 4/.....	0.2	0.3	0.4	0.8	1.4	2.5	4.4	7.0	11.5	20.4
2004 4/.....	0.2	0.3	0.6	0.7	1.4	2.6	4.4	7.6	11.6	18.2
2003 4/.....	0.2	0.4	0.5	0.8	1.6	3.1	4.7	8.6	12.4	21.8
2002.....	0.2	0.4	0.7	1.2	2.0	2.8	5.2	8.6	12.1	18.9
2001.....	0.3	0.4	0.6	1.0	1.7	3.2	5.0	8.4	12.2	18.7
2000.....	0.3	0.5	0.7	1.3	2.2	3.5	5.6	8.6	12.4	17.6
1999.....	0.3	0.5	0.8	1.4	2.0	3.6	6.0	8.8	13.1	18.9

White, male

2006 4/.....	*	0.3	0.3	0.6	1.1	1.9	3.0	5.3	8.3	12.9
2005 4/.....	*	0.3	0.4	0.8	1.1	1.9	3.3	5.2	8.3	14.7
2004 4/.....	*	0.3	0.4	0.6	0.9	2.0	3.1	6.1	10.1	12.4
2003 4/.....	*	0.3	0.3	0.7	1.3	2.3	3.7	6.9	9.8	17.3
2002.....	*	0.3	0.5	0.9	1.5	1.8	4.4	5.9	8.8	12.6
2001.....	0.3	0.4	0.4	1.0	1.2	2.3	3.3	6.3	8.0	13.3
2000.....	0.3	0.5	0.7	1.1	1.9	2.3	3.5	6.1	8.4	12.7
1999.....	0.2	0.4	0.6	1.0	1.3	2.6	4.9	5.5	9.4	11.1

White, female

2006 4/.....	*	0.3	0.4	1.0	1.6	3.0	5.2	8.4	13.6	20.4
2005 4/.....	*	0.4	0.5	0.9	1.8	3.1	5.3	8.4	13.6	23.1
2004 4/.....	0.2	0.3	0.7	0.9	1.9	3.0	5.4	8.6	12.5	20.8
2003 4/.....	0.2	0.5	0.6	1.0	1.9	3.8	5.5	9.8	14.0	23.8
2002.....	0.2	0.5	0.9	1.4	2.5	3.7	5.9	10.6	14.2	21.6
2001.....	0.3	0.4	0.7	1.1	2.3	3.9	6.4	9.9	14.7	21.0
2000.....	0.4	0.6	0.8	1.5	2.4	4.6	7.3	10.4	14.7	19.6
1999.....	0.3	0.6	0.9	1.8	2.7	4.6	6.9	11.0	15.4	22.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Black, both sexes

2006 4/.....	*	0.8	*	1.1	2.5	2.8	3.1	4.7	6.2	*
2005 4/.....	*	*	1.3	1.2	1.8	2.5	*	4.2	*	6.1
2004 4/.....	0.7	1.0	1.0	2.0	*	2.6	2.7	4.1	*	5.9
2003 4/.....	*	1.2	1.4	2.2	3.2	3.1	4.8	3.9	*	7.6
2002.....	*	0.8	1.1	2.0	2.9	3.2	*	4.8	*	*
2001.....	0.9	1.1	1.6	2.2	2.0	3.3	5.1	4.9	*	6.8
2000.....	0.9	1.0	1.1	1.9	2.2	3.4	3.6	3.7	*	8.4
1999.....	0.7	0.9	1.6	2.2	2.1	2.8	4.2	4.3	*	8.3

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, female

2006 4/.....	*	*	*	*	*	*	*	*	8.0	*
2005 4/.....	*	*	*	*	*	*	*	5.3	*	*
2004 4/.....	*	*	*	*	*	*	*	5.4	*	*
2003 4/.....	*	1.6	1.7	*	3.3	*	6.2	*	*	8.1
2002.....	*	*	*	*	3.6	*	*	5.5	*	*
2001.....	*	*	2.2	2.7	*	3.8	7.2	6.4	*	*
2000.....	*	*	*	*	*	4.4	*	*	*	*
1999.....	*	*	*	2.8	*	*	4.7	*	*	11.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart disease (I11)

All races 3/, both sexes

2006.....	4.0	6.8	10.0	11.8	14.9	18.2	23.1	35.2	64.2	194.0
2005.....	4.0	6.5	9.2	11.7	14.7	17.6	23.8	35.8	65.9	198.6
2004.....	4.0	6.8	9.0	12.2	14.9	18.8	25.1	36.7	66.1	192.8
2003.....	3.9	6.6	9.4	11.7	14.9	19.4	25.1	39.5	66.8	195.0
2002.....	3.5	5.9	8.6	11.1	14.3	18.9	24.0	37.9	66.0	187.0
2001.....	3.2	5.1	7.7	9.7	12.5	16.3	23.5	36.6	66.0	189.5
2000.....	2.9	4.7	6.6	9.3	13.2	16.5	23.0	36.3	66.1	193.1
1999.....	2.8	4.5	6.2	9.6	11.9	16.3	22.1	36.4	64.7	184.9

All races 3/, male

2006.....	5.2	9.8	13.8	16.2	20.4	24.0	27.8	40.2	63.1	143.0
2005.....	5.4	9.1	13.0	16.5	20.3	22.5	28.5	39.3	62.2	147.5
2004.....	5.2	9.2	12.4	17.0	20.4	24.7	30.5	39.1	60.9	145.4
2003.....	5.2	9.0	12.8	16.0	18.8	25.2	29.4	42.8	63.3	142.2
2002.....	4.9	8.0	11.9	15.2	19.0	23.6	27.0	41.0	62.2	134.6
2001.....	4.5	7.0	10.3	12.8	16.5	20.5	27.1	38.9	63.5	134.4
2000.....	3.9	6.3	9.4	12.1	17.4	20.7	27.2	37.3	61.8	139.1
1999.....	3.9	6.1	8.1	12.7	15.0	19.5	24.6	39.4	59.2	133.6

All races 3/, female

2006.....	2.8	3.8	6.3	7.5	9.8	13.2	19.2	31.4	64.9	217.9
2005.....	2.5	3.9	5.5	7.2	9.7	13.3	20.0	33.2	68.2	222.0
2004.....	2.7	4.4	5.8	7.6	9.9	13.5	20.7	35.0	69.3	214.1
2003.....	2.6	4.4	6.1	7.7	11.2	14.4	21.6	37.2	69.0	218.3
2002.....	2.2	3.9	5.4	7.2	10.0	14.8	21.6	35.8	68.2	209.7
2001.....	2.0	3.2	5.1	6.8	8.9	12.7	20.6	35.1	67.5	212.6
2000.....	2.0	3.2	3.8	6.6	9.3	12.9	19.6	35.7	68.6	215.1
1999.....	1.7	3.0	4.3	6.8	9.2	13.5	20.2	34.3	67.9	205.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart disease (I11)

White, both sexes

2006 4/.....	3.0	5.0	7.3	9.0	11.5	14.2	17.9	29.8	58.1	186.9
2005 4/.....	2.8	4.6	6.5	8.9	11.7	13.5	19.4	30.4	58.0	188.9
2004 4/.....	2.8	4.8	6.4	9.0	11.2	15.1	20.4	31.0	59.1	183.7
2003 4/.....	2.7	4.4	6.6	8.5	11.4	14.7	20.2	33.2	61.2	185.3
2002.....	2.4	4.3	6.0	8.1	10.5	14.4	19.3	31.9	58.1	177.3
2001.....	2.1	3.3	5.2	6.9	9.1	12.1	18.5	30.9	59.2	179.9
2000.....	2.0	3.1	4.3	6.2	9.4	12.5	17.6	30.2	58.4	181.8
1999.....	1.8	2.9	4.1	6.5	8.7	12.0	16.9	30.0	57.3	175.3

White, male

2006 4/.....	4.2	7.6	10.8	12.8	15.8	18.2	21.7	33.4	56.2	134.5
2005 4/.....	4.0	6.9	9.6	12.8	16.4	17.7	22.9	33.6	54.7	136.9
2004 4/.....	4.1	7.0	9.3	12.8	15.3	20.0	25.0	33.4	54.3	136.4
2003 4/.....	4.0	6.4	9.7	12.2	15.0	19.4	23.6	36.1	57.9	131.9
2002.....	3.7	6.1	8.9	11.6	14.0	18.1	22.0	34.5	54.8	124.8
2001.....	3.1	4.8	7.5	9.6	12.5	15.9	21.9	32.9	56.1	123.7
2000.....	2.8	4.6	6.5	8.6	12.7	16.0	21.5	31.1	53.4	129.3
1999.....	2.7	4.2	5.8	8.9	11.3	14.4	19.1	32.4	51.8	123.7

White, female

2006 4/.....	1.7	2.4	3.9	5.3	7.5	10.5	14.7	27.1	59.3	211.5
2005 4/.....	1.5	2.3	3.6	5.2	7.3	9.7	16.5	28.0	60.1	212.9
2004 4/.....	1.6	2.7	3.5	5.3	7.4	10.8	16.6	29.3	62.1	205.0
2003 4/.....	1.3	2.4	3.5	5.0	8.1	10.7	17.4	31.1	63.3	209.0
2002.....	1.2	2.4	3.3	4.7	7.1	11.1	17.1	30.0	60.1	200.1
2001.....	1.0	1.8	3.0	4.4	6.0	8.8	15.8	29.5	61.1	203.5
2000.....	1.2	1.6	2.2	4.0	6.3	9.4	14.5	29.6	61.3	203.1
1999.....	0.8	1.6	2.5	4.2	6.3	10.0	15.1	28.2	60.6	196.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart disease (I11)

Black, both sexes

2006 4/.....	11.3	20.4	31.3	37.3	48.4	56.9	75.0	96.8	145.3	299.6
2005 4/.....	12.2	20.8	30.3	37.8	46.0	57.8	69.9	100.9	168.5	338.1
2004 4/.....	12.3	22.0	31.3	41.6	49.9	55.4	76.2	103.1	161.4	314.7
2003 4/.....	12.8	23.9	32.8	41.9	48.3	63.0	77.1	113.3	145.7	332.3
2002.....	11.4	19.3	30.3	39.7	50.5	61.2	76.5	110.3	167.7	320.4
2001.....	11.5	18.6	28.4	35.7	43.6	55.8	77.8	105.1	154.7	326.1
2000.....	9.9	17.8	25.4	37.6	47.9	54.6	81.6	111.1	164.9	340.0
1999.....	10.2	17.9	24.8	38.0	41.6	57.6	79.0	113.7	161.2	305.1

Black, male

2006 4/.....	12.9	28.0	40.6	50.6	70.9	84.1	94.7	129.8	165.0	272.6
2005 4/.....	16.3	27.1	42.3	53.5	64.9	74.2	91.1	118.9	169.2	301.2
2004 4/.....	14.5	27.8	40.8	59.1	72.3	78.3	97.8	114.6	158.5	267.2
2003 4/.....	14.9	29.8	41.5	55.1	59.8	86.7	99.9	129.6	149.1	287.8
2002.....	14.6	24.4	39.9	53.1	71.1	79.0	89.8	125.4	167.4	267.0
2001.....	14.8	25.1	36.6	45.1	55.9	69.6	91.6	117.1	171.8	284.8
2000.....	12.7	21.4	35.8	47.9	63.8	70.0	95.9	117.2	178.7	273.3
1999.....	13.1	22.5	31.3	49.5	52.0	73.0	90.2	129.2	166.5	263.6

Black, female

2006 4/.....	9.9	13.7	23.4	26.4	31.0	36.9	61.5	77.1	134.9	310.7
2005 4/.....	8.6	15.3	20.2	24.9	31.3	45.7	55.4	90.1	168.0	353.0
2004 4/.....	10.3	17.0	23.2	27.2	32.5	38.5	61.5	96.2	162.9	333.5
2003 4/.....	11.0	18.7	25.5	31.0	39.4	45.5	61.8	103.6	143.9	349.6
2002.....	8.6	14.9	22.3	28.7	34.4	47.9	67.7	101.3	167.9	340.9
2001.....	8.6	13.0	21.5	28.0	33.9	45.6	68.6	97.9	145.9	341.4
2000.....	7.5	14.6	16.5	29.2	35.3	43.2	72.1	107.4	157.9	364.9
1999.....	7.6	13.8	19.3	28.7	33.4	46.2	71.6	104.4	158.6	320.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart and renal disease (I13)

All races 3/, both sexes

2006.....	0.2	0.3	0.5	0.7	1.0	1.4	2.4	4.3	8.6	22.6
2005.....	0.2	0.3	0.6	0.6	1.0	1.8	2.9	5.0	9.5	25.3
2004.....	0.3	0.5	0.5	0.7	1.3	1.6	2.6	4.5	9.4	24.8
2003.....	0.3	0.4	0.6	0.9	1.1	1.6	2.9	5.2	9.5	25.6
2002.....	0.2	0.4	0.5	0.9	1.2	1.6	2.8	4.9	9.5	23.5
2001.....	0.3	0.3	0.4	0.8	1.2	1.8	2.8	5.0	9.3	23.3
2000.....	0.3	0.3	0.4	0.8	1.0	1.6	3.0	5.1	9.8	24.3
1999.....	0.3	0.4	0.4	0.9	1.4	2.1	3.0	6.6	12.3	29.1

All races 3/, male

2006.....	0.3	0.3	0.6	0.9	1.3	1.6	2.6	4.7	10.0	23.8
2005.....	0.3	0.5	0.7	0.8	1.2	2.1	2.9	5.3	9.8	27.4
2004.....	0.3	0.5	0.7	0.7	1.7	1.7	2.8	5.1	10.4	24.3
2003.....	0.3	0.6	0.7	1.1	1.3	1.7	3.0	5.6	10.1	26.7
2002.....	0.2	0.5	0.6	1.1	1.4	1.5	3.4	5.1	9.9	23.1
2001.....	0.4	0.4	0.6	0.9	1.2	2.1	2.9	5.2	9.4	24.2
2000.....	0.3	0.4	0.5	0.9	1.0	1.7	3.4	5.0	10.8	27.8
1999.....	0.4	0.5	0.6	1.0	1.6	2.3	3.2	7.3	13.3	31.3

All races 3/, female

2006.....	*	0.3	0.4	0.4	0.8	1.3	2.3	4.0	7.8	22.1
2005.....	0.2	*	0.4	0.5	0.8	1.6	2.8	4.8	9.3	24.3
2004.....	*	0.4	0.4	0.7	0.9	1.5	2.3	4.0	8.8	25.0
2003.....	0.2	0.2	0.5	0.6	0.9	1.5	2.8	4.9	9.2	25.2
2002.....	0.2	0.2	0.4	0.7	1.0	1.7	2.4	4.7	9.2	23.7
2001.....	*	*	0.3	0.7	1.3	1.5	2.8	4.9	9.3	22.9
2000.....	0.2	0.2	0.4	0.7	1.0	1.5	2.7	5.1	9.3	22.9
1999.....	0.3	0.3	0.3	0.7	1.2	2.0	2.9	6.1	11.7	28.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart and renal disease (I13)

White, both sexes

2006 4/.....	*	0.1	0.2	0.4	0.5	0.9	1.5	3.1	7.7	20.9
2005 4/.....	*	0.1	0.2	0.3	0.5	1.1	2.0	3.9	7.9	23.9
2004 4/.....	0.1	0.1	0.2	0.3	0.7	0.9	1.8	3.5	8.1	23.0
2003 4/.....	*	0.1	0.2	0.3	0.6	0.9	2.1	4.2	8.3	23.4
2002.....	*	*	0.1	0.3	0.6	0.9	1.9	3.8	8.4	21.0
2001.....	*	*	0.2	0.4	0.7	1.1	2.0	3.9	8.3	21.0
2000.....	0.2	0.1	0.2	0.4	0.5	0.9	2.2	4.0	8.1	22.0
1999.....	*	0.1	0.2	0.4	0.7	1.2	1.9	5.2	10.4	26.1

White, male

2006 4/.....	*	*	0.3	0.5	0.6	0.8	1.6	3.3	9.0	22.5
2005 4/.....	*	*	0.3	0.4	0.5	1.1	2.1	4.2	8.2	25.5
2004 4/.....	*	*	*	0.3	1.0	1.0	2.1	3.9	9.2	22.9
2003 4/.....	*	*	*	0.5	0.8	1.1	2.2	4.9	9.3	24.3
2002.....	*	*	*	0.5	0.7	0.8	2.4	4.2	8.8	20.6
2001.....	*	*	0.3	0.4	0.7	1.3	2.0	4.1	8.7	22.7
2000.....	*	*	*	0.4	0.5	1.0	2.5	4.4	9.2	24.4
1999.....	*	*	*	0.5	0.8	1.4	2.0	5.8	12.1	28.8

White, female

2006 4/.....	*	*	*	*	0.4	0.9	1.5	2.9	6.8	20.1
2005 4/.....	*	*	*	*	0.4	1.0	1.9	3.7	7.8	23.2
2004 4/.....	*	*	*	0.3	0.4	0.8	1.5	3.2	7.4	23.0
2003 4/.....	*	*	*	*	0.5	0.7	1.9	3.7	7.8	22.9
2002.....	*	*	*	*	0.5	0.9	1.5	3.6	8.1	21.1
2001.....	*	*	*	0.3	0.8	0.8	2.0	3.7	8.0	20.3
2000.....	*	*	*	0.4	0.5	0.8	2.0	3.7	7.4	21.1
1999.....	*	*	*	0.4	0.6	1.1	1.8	4.7	9.5	25.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hypertensive heart and renal disease (I13)

Black, both sexes

2006 4/.....	1.0	1.7	2.7	3.4	5.7	6.5	11.0	16.8	21.1	41.8
2005 4/.....	1.2	1.7	3.4	3.7	5.7	8.7	10.7	16.5	28.4	42.8
2004 4/.....	1.3	2.8	2.9	3.9	6.4	7.5	9.7	15.2	26.0	48.2
2003 4/.....	1.5	2.3	3.8	5.3	5.5	7.4	11.0	16.5	24.1	53.1
2002.....	1.1	2.2	3.3	6.0	6.2	8.0	12.3	17.4	23.0	56.1
2001.....	1.4	1.9	2.3	4.4	5.7	8.4	11.8	18.4	22.8	50.1
2000.....	1.2	1.9	2.5	4.1	5.3	8.3	11.9	17.6	31.1	50.4
1999.....	2.1	2.5	2.3	5.0	7.3	10.1	15.4	21.8	35.3	65.4

Black, male

2006 4/.....	1.5	1.9	3.5	5.0	7.6	8.2	13.5	22.8	23.3	39.3
2005 4/.....	*	2.8	4.0	4.7	8.0	11.6	10.4	17.0	33.0	50.2
2004 4/.....	2.0	3.3	3.8	4.4	8.6	8.2	9.4	20.5	27.3	41.7
2003 4/.....	2.1	3.4	4.6	7.6	6.4	6.6	11.8	15.5	24.4	50.0
2002.....	*	3.0	4.7	7.6	7.3	7.2	14.0	17.0	22.9	61.1
2001.....	2.1	3.0	3.5	5.6	6.3	10.7	13.5	18.1	19.3	44.3
2000.....	1.5	2.7	2.9	5.1	5.6	9.4	13.7	13.7	33.7	63.4
1999.....	2.6	3.5	3.4	6.3	9.3	10.6	17.3	23.7	29.6	59.8

Black, female

2006 4/.....	*	1.5	1.9	2.1	4.3	5.2	9.3	13.1	19.9	42.8
2005 4/.....	*	*	2.9	2.9	3.8	6.5	10.8	16.3	26.0	39.8
2004 4/.....	*	2.4	2.2	3.5	4.7	7.0	9.9	12.1	25.3	50.8
2003 4/.....	*	*	3.1	3.5	4.8	8.0	10.4	17.1	24.0	54.3
2002.....	*	1.5	2.1	4.6	5.3	8.5	11.2	17.6	23.1	54.2
2001.....	*	*	*	3.3	5.3	6.8	10.8	18.6	24.5	52.2
2000.....	*	*	2.1	3.2	5.1	7.5	10.7	19.9	29.7	45.5
1999.....	1.6	1.6	*	4.0	5.7	9.6	14.2	20.6	38.0	67.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Ischemic heart diseases (I20-I25)

All races 3/, both sexes

2006.....	21.5	42.2	75.0	118.1	188.5	279.1	441.7	726.4	1,263.9	2,904.7
2005.....	21.6	42.9	78.3	120.9	198.6	295.4	472.3	778.6	1,359.8	3,158.1
2004.....	22.4	44.0	77.9	123.9	203.1	310.5	495.9	813.5	1,418.0	3,287.4
2003.....	23.5	45.9	80.7	132.4	218.7	335.0	538.1	886.4	1,538.2	3,610.2
2002.....	23.8	46.4	83.5	140.1	226.2	358.4	569.8	940.2	1,611.2	3,775.0
2001.....	23.4	45.6	84.8	142.8	236.6	372.4	591.8	984.3	1,685.2	3,958.2
2000.....	23.6	48.0	86.7	153.9	250.0	390.8	628.0	1,032.5	1,761.7	4,182.5
1999.....	24.2	48.5	90.1	157.2	260.0	420.1	661.9	1,081.1	1,841.7	4,318.1

All races 3/, male

2006.....	32.7	64.4	116.9	180.0	277.1	400.1	613.0	988.2	1,672.7	3,303.0
2005.....	33.0	66.2	121.3	183.8	291.3	415.2	653.7	1,054.3	1,780.2	3,576.8
2004.....	34.0	68.2	121.1	187.5	297.8	440.2	682.0	1,106.0	1,855.0	3,680.3
2003.....	35.5	71.4	126.9	200.3	319.7	476.1	744.1	1,200.1	2,013.5	4,005.2
2002.....	36.5	72.3	130.3	211.4	331.4	508.5	791.5	1,274.8	2,094.2	4,205.3
2001.....	35.8	71.6	131.1	215.5	344.4	525.7	820.3	1,337.5	2,180.0	4,395.8
2000.....	37.0	76.9	134.2	232.8	362.9	550.5	879.6	1,394.4	2,305.0	4,701.7
1999.....	37.9	77.0	140.3	237.7	376.1	595.2	924.6	1,462.5	2,400.4	4,837.5

All races 3/, female

2006.....	10.3	20.6	34.8	59.7	107.5	173.3	302.4	534.8	1,008.4	2,718.3
2005.....	10.4	20.2	37.0	61.5	114.1	190.9	325.4	578.5	1,098.7	2,965.8
2004.....	10.8	20.4	36.6	64.0	116.8	197.7	346.0	603.2	1,148.8	3,110.6
2003.....	11.6	21.1	36.5	68.4	126.9	212.6	373.0	663.1	1,247.5	3,435.5
2002.....	11.2	21.4	38.7	73.1	130.7	228.8	392.6	704.0	1,317.5	3,588.3
2001.....	11.1	20.3	40.3	74.8	138.8	240.5	410.2	737.0	1,388.7	3,775.1
2000.....	10.3	20.0	41.3	80.0	147.7	253.9	429.9	780.5	1,441.2	3,971.0
1999.....	10.7	20.8	42.1	82.1	155.0	270.2	456.7	815.9	1,514.7	4,108.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Ischemic heart diseases (I20-I25)

White, both sexes

2006 4/.....	21.0	40.1	71.3	112.1	180.9	268.6	432.1	720.3	1,267.5	2,964.3
2005 4/.....	21.0	41.1	73.8	114.2	189.8	286.3	463.7	771.1	1,364.5	3,227.0
2004 4/.....	21.6	42.0	73.4	117.5	195.2	298.7	485.8	807.0	1,423.8	3,343.9
2003 4/.....	22.4	43.5	76.1	126.4	208.2	322.6	526.3	877.2	1,542.2	3,671.7
2002.....	22.9	44.2	79.1	133.3	216.1	346.6	559.2	928.9	1,616.4	3,836.4
2001.....	22.1	43.1	80.3	135.4	225.8	357.9	580.5	972.9	1,689.9	4,022.7
2000.....	22.2	45.3	82.4	146.2	238.5	378.4	615.0	1,019.7	1,768.5	4,257.7
1999.....	22.6	46.0	85.4	149.1	249.5	408.5	647.3	1,070.6	1,849.5	4,389.2

White, male

2006 4/.....	32.6	62.2	112.7	172.6	267.5	388.4	603.3	983.8	1,689.7	3,396.0
2005 4/.....	32.7	64.1	116.2	175.7	281.5	406.5	646.8	1,053.1	1,798.9	3,687.1
2004 4/.....	33.7	66.3	116.1	180.0	289.4	428.5	673.9	1,105.8	1,880.8	3,768.8
2003 4/.....	34.4	68.8	121.6	194.2	308.7	464.4	731.5	1,195.8	2,028.8	4,106.1
2002.....	35.9	70.3	125.6	203.9	320.5	498.1	784.6	1,273.1	2,115.1	4,299.7
2001.....	34.6	69.5	126.4	207.2	333.9	512.8	810.3	1,335.1	2,201.7	4,494.0
2000.....	35.9	74.5	130.0	223.7	352.1	541.2	870.1	1,390.0	2,328.3	4,813.9
1999.....	36.2	74.7	135.0	228.2	364.4	586.0	913.4	1,463.9	2,429.7	4,933.9

White, female

2006 4/.....	9.2	18.1	30.6	53.9	100.1	161.9	289.9	523.6	1,000.8	2,761.6
2005 4/.....	9.2	18.1	32.1	54.9	104.5	179.4	312.3	562.4	1,092.2	3,015.2
2004 4/.....	9.4	17.9	31.5	57.3	107.8	183.6	331.3	588.5	1,140.0	3,152.3
2003 4/.....	10.4	18.4	31.5	61.3	115.2	197.3	358.4	646.9	1,242.2	3,479.3
2002.....	9.8	18.2	33.6	65.7	119.6	213.2	375.5	682.5	1,310.6	3,635.2
2001.....	9.5	17.0	35.2	66.8	126.1	222.0	394.4	716.3	1,380.6	3,825.5
2000.....	8.5	16.4	36.0	72.5	133.8	236.0	410.4	759.2	1,435.5	4,031.8
1999.....	8.9	17.6	37.0	73.8	143.6	253.7	435.9	794.5	1,507.0	4,170.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Ischemic heart diseases (I20-I25)

Black, both sexes

2006 4/.....	29.7	64.9	121.2	191.9	295.9	422.1	616.2	922.5	1,418.1	2,616.1
2005 4/.....	30.2	65.0	129.0	202.1	320.9	431.0	638.8	1,004.8	1,517.3	2,804.4
2004 4/.....	33.2	67.4	130.6	207.3	320.6	471.5	691.7	1,030.8	1,554.7	3,014.3
2003 4/.....	35.2	71.5	133.3	213.8	352.2	503.0	756.8	1,139.8	1,696.1	3,310.1
2002.....	35.1	72.8	136.5	230.9	362.0	522.5	779.6	1,196.0	1,768.0	3,462.9
2001.....	36.5	72.5	138.8	236.1	376.4	562.5	810.8	1,249.3	1,837.2	3,615.1
2000.....	37.1	78.2	140.3	250.9	392.9	566.1	866.5	1,296.9	1,882.8	3,684.7
1999.....	39.7	76.9	149.3	258.5	395.4	596.2	906.3	1,321.5	1,926.9	3,805.8

Black, male

2006 4/.....	41.0	91.7	178.2	281.0	431.2	594.0	850.5	1,242.2	1,785.3	2,656.7
2005 4/.....	41.2	95.4	187.9	294.0	454.6	584.9	857.4	1,296.8	1,887.0	2,785.0
2004 4/.....	44.9	96.6	189.8	297.2	454.8	641.3	922.9	1,332.6	1,860.9	3,100.5
2003 4/.....	50.2	103.2	195.6	302.7	490.5	677.1	1,027.8	1,501.3	2,138.8	3,285.3
2002.....	48.4	101.6	196.9	330.3	509.6	703.3	1,024.8	1,495.0	2,171.1	3,556.2
2001.....	50.8	101.9	199.9	333.7	515.8	743.8	1,093.0	1,582.5	2,244.8	3,775.0
2000.....	51.8	111.6	197.6	366.7	530.3	727.5	1,139.7	1,630.6	2,316.5	3,885.2
1999.....	56.8	110.4	217.6	372.5	553.5	788.8	1,188.9	1,630.3	2,312.4	4,080.1

Black, female

2006 4/.....	19.7	41.6	73.0	118.5	190.5	295.1	455.2	731.2	1,224.0	2,599.5
2005 4/.....	20.5	38.6	79.1	126.3	216.6	316.9	489.1	830.7	1,322.0	2,812.2
2004 4/.....	22.9	42.0	80.5	133.4	215.9	345.4	534.9	852.0	1,393.3	2,980.1
2003 4/.....	21.9	43.9	80.5	140.7	244.3	373.7	574.0	924.9	1,464.2	3,319.7
2002.....	23.5	47.9	85.4	149.3	246.6	388.3	615.0	1,017.3	1,558.1	3,427.2
2001.....	23.9	47.1	86.8	156.0	266.7	428.6	622.8	1,050.3	1,629.6	3,555.7
2000.....	24.1	49.2	91.4	156.0	284.5	447.0	685.1	1,095.7	1,664.3	3,610.0
1999.....	24.7	48.0	91.4	165.5	271.4	453.5	718.8	1,135.4	1,735.9	3,703.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute myocardial infarction (I21-I22)

All races 3/, both sexes

2006.....	8.5	16.5	29.2	46.7	73.9	107.0	160.4	252.0	406.4	825.8
2005.....	8.7	17.3	30.9	47.8	78.9	115.0	176.6	272.1	448.1	923.1
2004.....	8.9	17.7	31.6	50.6	82.7	124.1	187.7	291.1	476.0	992.9
2003.....	9.6	18.8	33.2	55.5	90.8	134.5	207.4	325.6	532.5	1,118.8
2002.....	9.9	19.0	34.9	58.4	96.2	148.5	227.5	352.5	572.7	1,191.3
2001.....	9.9	19.7	36.5	61.3	100.1	154.1	235.6	374.0	611.5	1,265.4
2000.....	9.9	21.1	37.8	67.0	110.9	166.8	254.7	405.3	650.1	1,338.5
1999.....	10.4	21.5	39.0	69.7	114.9	181.7	271.6	422.9	689.4	1,379.3

All races 3/, male

2006.....	13.0	24.7	44.8	70.0	107.3	151.8	219.9	331.3	527.1	936.3
2005.....	13.1	26.5	47.2	72.2	114.0	159.3	240.5	361.8	567.1	1,045.0
2004.....	13.2	27.4	48.7	76.3	120.6	173.0	254.1	386.8	608.9	1,120.5
2003.....	14.4	29.2	52.1	83.4	131.6	188.5	281.4	430.0	682.4	1,263.5
2002.....	14.8	29.2	53.7	86.6	139.0	207.6	312.0	466.1	726.6	1,366.4
2001.....	15.0	30.8	55.6	91.3	143.5	214.2	322.0	495.6	776.1	1,461.3
2000.....	15.5	33.6	57.5	99.7	158.6	231.3	349.6	535.2	835.8	1,568.3
1999.....	16.0	33.7	60.9	105.3	164.9	253.5	373.8	564.2	885.1	1,622.1

All races 3/, female

2006.....	4.1	8.4	14.3	24.7	43.4	67.8	112.0	194.0	330.9	774.1
2005.....	4.4	8.4	15.2	24.7	47.0	76.4	124.8	206.9	374.1	867.1
2004.....	4.5	8.3	15.3	26.4	48.2	81.5	134.3	222.3	394.1	935.5
2003.....	5.0	8.7	15.1	29.3	53.8	87.7	148.0	251.3	440.9	1,054.8
2002.....	5.1	9.1	17.0	32.0	57.4	97.6	159.8	272.2	479.1	1,115.3
2001.....	4.9	9.0	18.2	33.2	60.7	102.4	167.0	288.8	512.9	1,183.5
2000.....	4.4	8.9	18.9	36.5	67.7	111.5	179.9	314.9	540.6	1,244.8
1999.....	4.9	9.7	18.0	36.5	69.6	120.2	191.7	324.7	574.9	1,281.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute myocardial infarction (I21-I22)

White, both sexes

2006 4/.....	8.5	16.1	28.6	45.6	72.9	105.2	158.9	250.6	408.0	836.0
2005 4/.....	8.6	17.0	30.3	46.3	78.0	113.8	175.8	270.1	449.4	936.8
2004 4/.....	8.7	17.4	30.6	49.4	81.8	121.7	185.3	291.1	478.6	1,004.1
2003 4/.....	9.3	18.2	32.1	54.4	89.1	132.3	205.3	324.2	532.9	1,132.6
2002.....	9.7	18.4	34.2	57.3	94.3	146.4	225.0	349.5	572.7	1,201.2
2001.....	9.5	19.3	35.4	59.8	98.4	150.8	233.3	371.9	612.7	1,277.8
2000.....	9.5	20.6	36.7	65.6	108.9	164.1	251.8	401.7	652.6	1,353.5
1999.....	10.0	21.1	38.3	68.2	113.4	180.6	268.3	421.1	692.5	1,393.2

White, male

2006 4/.....	13.1	24.6	44.6	69.2	106.3	151.2	220.0	331.9	532.5	955.4
2005 4/.....	13.1	26.3	47.2	70.7	114.2	160.0	241.8	363.3	572.8	1,070.1
2004 4/.....	13.4	27.2	48.0	75.4	121.2	171.7	252.9	391.6	618.6	1,143.7
2003 4/.....	14.1	28.7	51.2	83.0	130.8	188.3	280.0	430.3	685.7	1,289.3
2002.....	14.7	29.0	53.3	86.2	138.7	207.7	313.1	467.1	730.8	1,386.1
2001.....	14.9	30.9	55.1	90.6	143.7	213.3	321.9	497.6	784.0	1,481.1
2000.....	15.3	33.6	57.2	98.8	158.3	231.1	349.3	536.3	843.5	1,592.9
1999.....	15.8	33.8	60.8	104.0	165.1	254.3	374.0	568.8	896.6	1,644.0

White, female

2006 4/.....	3.9	7.6	13.0	22.9	41.7	64.3	108.1	189.9	329.4	780.0
2005 4/.....	3.9	7.7	13.7	22.9	44.3	72.6	121.3	201.2	372.1	875.4
2004 4/.....	3.9	7.5	13.5	24.3	45.2	77.4	129.7	217.6	391.6	941.1
2003 4/.....	4.5	7.7	13.4	26.8	50.5	82.8	144.1	247.5	438.7	1,063.2
2002.....	4.6	8.0	15.4	29.7	53.3	92.5	153.2	265.3	475.8	1,120.8
2001.....	4.2	7.7	16.1	30.3	56.7	96.0	161.5	282.8	509.1	1,192.7
2000.....	3.7	7.7	16.6	34.1	63.5	105.5	173.6	307.0	539.1	1,256.2
1999.....	4.3	8.4	16.2	34.2	65.8	116.3	184.3	317.3	572.1	1,292.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Acute myocardial infarction (I21-I22)

Black, both sexes

2006 4/.....	11.1	23.1	41.6	67.0	100.9	144.9	207.1	311.5	448.2	804.0
2005 4/.....	11.8	24.2	43.3	69.9	107.8	150.2	217.0	345.4	499.1	892.1
2004 4/.....	12.4	25.0	47.9	75.3	112.7	169.2	249.7	351.1	520.9	979.9
2003 4/.....	14.4	27.5	49.5	79.4	124.3	180.3	270.2	401.4	597.0	1,094.9
2002.....	13.9	27.7	50.4	84.2	136.3	193.6	295.9	434.1	643.9	1,201.2
2001.....	14.3	27.8	53.2	89.2	135.9	211.1	304.3	452.3	671.2	1,239.9
2000.....	15.1	29.6	55.5	94.7	151.4	219.0	328.8	490.6	698.8	1,275.7
1999.....	15.4	30.0	54.9	96.9	150.2	224.0	347.5	495.1	729.6	1,318.8

Black, male

2006 4/.....	15.9	31.5	59.0	95.0	144.9	195.4	271.9	391.1	552.4	814.1
2005 4/.....	15.8	34.4	59.5	100.2	145.2	190.8	278.3	425.6	598.3	885.9
2004 4/.....	15.4	36.3	67.6	104.8	150.1	223.1	322.7	419.5	591.1	1,009.3
2003 4/.....	20.0	38.6	70.9	108.1	167.9	230.2	353.4	516.2	736.1	1,113.5
2002.....	19.0	37.1	70.4	114.5	177.6	247.8	370.2	531.4	776.4	1,280.1
2001.....	18.5	36.4	71.2	120.1	174.5	264.8	386.4	548.7	792.2	1,382.0
2000.....	20.6	41.0	72.7	131.4	196.7	270.9	419.2	592.5	856.5	1,418.5
1999.....	21.1	41.1	76.9	138.8	198.8	290.8	435.5	591.7	858.0	1,498.7

Black, female

2006 4/.....	7.0	15.8	26.9	43.9	66.7	107.7	162.5	263.8	393.2	799.9
2005 4/.....	8.4	15.3	29.6	44.9	78.6	120.0	175.1	297.6	446.6	894.6
2004 4/.....	9.7	15.2	31.2	51.0	83.6	129.2	200.1	310.5	483.8	968.3
2003 4/.....	9.5	18.0	31.5	55.7	90.3	143.3	214.0	333.2	524.2	1,087.7
2002.....	9.3	19.5	33.4	59.4	104.1	153.4	246.0	376.0	574.9	1,171.0
2001.....	10.5	20.3	37.8	63.8	105.6	171.4	249.5	394.7	609.5	1,187.2
2000.....	10.1	19.8	40.8	64.6	115.7	180.7	268.8	429.2	619.4	1,222.5
1999.....	10.4	20.5	36.2	62.6	112.1	174.5	289.2	436.8	665.9	1,251.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute ischemic heart diseases (I24)

All races 3/, both sexes

2006.....	0.3	0.6	1.1	1.5	2.5	2.9	4.4	6.4	10.0	21.5
2005.....	0.3	0.6	1.1	1.4	2.1	2.4	4.2	5.4	9.5	20.8
2004.....	0.3	0.4	0.7	1.1	1.4	1.8	2.6	4.2	6.5	14.8
2003.....	0.3	0.5	0.9	1.3	1.7	2.7	3.6	5.2	8.3	20.0
2002.....	0.3	0.6	1.2	1.4	2.2	3.0	3.9	5.7	9.0	21.1
2001.....	0.3	0.6	1.0	1.4	2.4	2.9	4.1	5.3	9.2	21.5
2000.....	0.3	0.5	1.1	1.6	2.5	3.1	4.0	5.9	9.4	21.9
1999.....	0.3	0.5	0.9	1.6	2.2	3.1	4.3	5.5	9.2	21.9

All races 3/, male

2006.....	0.4	0.8	1.6	1.9	3.6	3.6	5.9	8.0	12.1	21.8
2005.....	0.4	0.8	1.6	2.1	3.2	3.4	5.3	7.1	10.6	20.2
2004.....	0.4	0.5	1.0	1.6	2.0	2.7	3.4	5.0	8.3	16.3
2003.....	0.5	0.7	1.3	1.8	2.4	3.8	4.9	7.1	9.4	19.5
2002.....	0.4	0.8	1.8	2.1	3.0	4.3	5.1	7.0	11.3	21.4
2001.....	0.4	0.9	1.6	2.0	3.5	4.1	5.5	6.7	11.3	20.7
2000.....	0.4	0.8	1.6	2.5	3.5	4.4	5.2	7.6	11.7	20.7
1999.....	0.4	0.7	1.3	2.5	3.1	4.3	5.6	7.6	10.8	22.3

All races 3/, female

2006.....	0.2	0.3	0.6	1.0	1.5	2.2	3.3	5.3	8.8	21.4
2005.....	*	0.4	0.6	0.7	1.2	1.6	3.3	4.2	8.8	21.1
2004.....	*	0.2	0.4	0.6	0.8	1.0	1.9	3.6	5.4	14.1
2003.....	*	0.3	0.5	0.8	1.0	1.8	2.6	3.9	7.6	20.2
2002.....	*	0.3	0.7	0.7	1.4	1.9	3.0	4.7	7.6	21.0
2001.....	*	0.3	0.5	0.9	1.3	2.0	3.1	4.4	8.0	21.8
2000.....	*	*	0.5	0.8	1.5	1.9	2.9	4.8	8.1	22.4
1999.....	*	0.2	0.6	0.7	1.5	2.0	3.2	4.1	8.3	21.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute ischemic heart diseases (I24)

White, both sexes

2006 4/.....	0.2	0.5	1.0	1.4	2.3	2.7	4.2	6.2	9.8	21.6
2005 4/.....	0.2	0.5	0.9	1.2	1.9	2.3	3.9	5.1	9.2	20.5
2004 4/.....	0.3	0.3	0.6	0.9	1.3	1.6	2.4	3.9	6.3	15.0
2003 4/.....	0.3	0.4	0.8	1.1	1.6	2.6	3.4	5.0	8.1	20.2
2002.....	0.2	0.5	1.1	1.3	2.0	2.9	3.8	5.6	8.9	21.4
2001.....	0.2	0.5	0.9	1.3	2.1	2.7	4.0	5.2	9.2	21.5
2000.....	0.2	0.4	1.0	1.4	2.2	2.9	3.8	5.6	9.3	22.1
1999.....	0.2	0.3	0.8	1.4	2.1	3.0	4.0	5.4	8.9	22.4

White, male

2006 4/.....	0.3	0.7	1.5	1.8	3.4	3.5	5.5	7.6	11.6	21.8
2005 4/.....	0.3	0.7	1.4	1.8	2.8	3.2	4.9	6.6	10.3	19.7
2004 4/.....	0.4	0.4	0.8	1.4	1.8	2.3	3.1	4.6	8.3	16.2
2003 4/.....	0.4	0.6	1.2	1.5	2.2	3.6	4.8	6.7	8.9	19.7
2002.....	0.4	0.7	1.6	2.0	2.8	4.1	4.8	7.0	11.1	21.9
2001.....	0.3	0.8	1.3	1.8	3.1	3.5	5.4	6.5	11.6	20.7
2000.....	0.4	0.7	1.5	2.2	3.2	4.0	4.9	7.2	11.6	21.0
1999.....	0.3	0.5	1.2	2.2	2.8	4.1	5.2	7.6	10.3	22.9

White, female

2006 4/.....	*	0.2	0.5	0.9	1.4	2.0	3.1	5.2	8.7	21.6
2005 4/.....	*	0.2	0.4	0.6	1.0	1.6	3.1	4.0	8.5	20.8
2004 4/.....	*	*	0.3	0.5	0.8	1.0	1.9	3.4	5.1	14.4
2003 4/.....	*	0.2	0.4	0.7	0.9	1.6	2.4	3.8	7.7	20.5
2002.....	*	0.3	0.6	0.7	1.2	1.8	2.9	4.6	7.5	21.1
2001.....	*	0.2	0.4	0.8	1.2	1.9	2.9	4.3	7.7	21.7
2000.....	*	*	0.5	0.6	1.3	1.9	2.8	4.5	8.0	22.6
1999.....	*	*	0.5	0.7	1.5	2.0	3.0	3.8	8.1	22.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute ischemic heart diseases (I24)

Black, both sexes

2006 4/.....	*	1.1	2.3	2.6	4.5	4.7	7.6	9.0	14.2	20.8
2005 4/.....	*	1.3	2.8	3.2	4.5	4.1	7.3	9.7	14.8	27.3
2004 4/.....	*	1.0	1.9	2.9	3.0	3.9	4.3	8.1	9.9	15.2
2003 4/.....	0.7	1.1	1.5	2.7	3.3	4.9	6.2	8.7	11.7	19.8
2002.....	*	1.1	2.4	2.1	4.1	5.0	7.1	7.7	11.8	21.1
2001.....	0.7	1.1	2.1	2.8	5.2	5.3	6.2	6.9	10.8	24.0
2000.....	*	1.0	2.1	3.8	4.7	5.2	6.1	10.0	12.1	21.9
1999.....	*	1.5	2.2	3.3	3.9	4.3	7.3	8.7	13.9	17.2

Black, male

2006 4/.....	*	*	3.2	3.9	6.9	5.6	11.1	12.5	19.2	23.2
2005 4/.....	*	*	3.6	5.0	6.9	6.4	9.8	14.8	17.6	27.9
2004 4/.....	*	*	2.7	4.5	4.7	7.0	7.1	11.8	*	22.8
2003 4/.....	*	1.7	2.2	4.6	5.0	7.4	7.9	14.2	19.2	24.0
2002.....	*	1.9	3.6	3.1	5.7	8.0	10.7	9.2	16.0	21.4
2001.....	*	*	3.6	3.4	8.1	9.1	7.7	11.2	*	23.8
2000.....	*	*	3.6	5.8	6.6	8.3	9.0	13.2	*	*
1999.....	*	2.5	2.8	5.8	6.6	6.8	10.2	10.0	17.4	*

Black, female

2006 4/.....	*	*	*	*	*	4.0	5.3	7.0	11.6	19.7
2005 4/.....	*	*	2.2	*	*	*	5.6	6.7	13.4	27.0
2004 4/.....	*	*	*	*	*	*	*	5.9	9.1	12.2
2003 4/.....	*	*	*	*	*	*	5.1	5.4	7.7	18.2
2002.....	*	*	*	*	*	*	4.7	6.9	9.5	20.9
2001.....	*	*	*	*	*	*	5.2	*	11.9	24.0
2000.....	*	*	*	*	3.3	*	*	8.0	10.6	22.3
1999.....	*	*	*	*	*	*	5.4	8.0	12.1	16.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

Other forms of chronic ischemic heart disease (I20,I25)

All races 3/, both sexes

2006.....	12.7	25.2	44.7	70.0	112.1	169.3	276.9	468.0	847.5	2,057.4
2005.....	12.7	25.1	46.3	71.7	117.5	178.0	291.6	501.1	902.3	2,214.2
2004.....	13.2	25.9	45.6	72.2	119.0	184.6	305.6	518.2	935.6	2,279.7
2003.....	13.5	26.6	46.6	75.6	126.1	197.8	327.1	555.6	997.3	2,471.4
2002.....	13.6	26.8	47.3	80.2	127.8	206.8	338.4	582.1	1,029.5	2,562.5
2001.....	13.2	25.2	47.3	80.1	134.1	215.4	352.0	605.0	1,064.5	2,671.3
2000.....	13.4	26.5	47.9	85.2	136.6	221.0	369.4	621.2	1,102.2	2,822.1
1999.....	13.5	26.5	50.2	85.9	142.9	235.3	386.1	652.6	1,143.1	2,917.0

All races 3/, male

2006.....	19.4	38.8	70.5	108.1	166.2	244.7	387.2	649.0	1,133.5	2,344.9
2005.....	19.5	39.0	72.5	109.5	174.1	252.6	407.9	685.4	1,202.5	2,511.6
2004.....	20.4	40.3	71.4	109.5	175.2	264.4	424.5	714.2	1,237.9	2,543.6
2003.....	20.7	41.5	73.5	115.1	185.6	283.8	457.8	763.1	1,321.6	2,722.2
2002.....	21.4	42.2	74.9	122.7	189.4	296.6	474.3	801.6	1,356.2	2,817.4
2001.....	20.4	40.0	74.0	122.2	197.4	307.4	492.9	835.3	1,392.6	2,913.8
2000.....	21.1	42.5	75.1	130.6	200.8	314.8	524.7	851.5	1,457.6	3,112.7
1999.....	21.5	42.6	78.1	129.9	208.1	337.3	545.2	890.7	1,504.6	3,193.1

All races 3/, female

2006.....	6.0	11.8	19.9	34.1	62.7	103.3	187.1	335.6	668.6	1,922.9
2005.....	5.9	11.5	21.2	36.1	65.9	112.9	197.3	367.3	715.8	2,077.6
2004.....	6.1	11.9	20.9	37.0	67.8	115.1	209.8	377.3	749.3	2,161.0
2003.....	6.5	12.1	20.9	38.4	72.0	123.2	222.4	408.0	799.1	2,360.5
2002.....	6.0	11.9	21.0	40.3	72.0	129.4	229.7	427.1	830.8	2,451.9
2001.....	6.1	11.0	21.7	40.6	76.8	136.2	240.1	443.8	867.9	2,569.8
2000.....	5.8	10.9	21.8	42.7	78.5	140.5	247.0	460.8	892.5	2,703.8
1999.....	5.6	10.9	23.4	44.9	83.9	148.1	261.7	487.1	931.6	2,805.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other forms of chronic ischemic heart disease (I20,I25)

White, both sexes

2006 4/.....	12.2	23.5	41.7	65.2	105.7	160.6	269.0	463.4	849.7	2,106.6
2005 4/.....	12.2	23.6	42.6	66.7	109.9	170.2	283.9	495.8	905.8	2,269.8
2004 4/.....	12.6	24.4	42.2	67.2	112.1	175.4	298.1	512.0	939.0	2,324.8
2003 4/.....	12.9	24.9	43.2	71.0	117.6	187.7	317.6	548.0	1,001.1	2,518.8
2002.....	13.0	25.2	43.9	74.7	119.8	197.3	330.5	573.8	1,034.7	2,613.9
2001.....	12.3	23.3	44.0	74.3	125.3	204.5	343.2	595.8	1,068.1	2,723.5
2000.....	12.5	24.3	44.8	79.2	127.3	211.4	359.4	612.4	1,106.6	2,882.1
1999.....	12.3	24.6	46.3	79.4	134.0	225.0	375.0	644.2	1,148.1	2,973.6

White, male

2006 4/.....	19.2	36.9	66.7	101.7	157.9	233.6	377.8	644.3	1,145.6	2,418.9
2005 4/.....	19.2	37.1	67.6	103.3	164.5	243.3	400.0	683.2	1,215.8	2,597.3
2004 4/.....	19.9	38.7	67.2	103.3	166.4	254.5	417.9	709.7	1,254.0	2,608.9
2003 4/.....	19.9	39.5	69.2	109.6	175.8	272.6	446.7	758.8	1,334.2	2,797.1
2002.....	20.8	40.6	70.7	115.8	179.1	286.2	466.8	799.0	1,373.2	2,891.6
2001.....	19.4	37.7	69.9	114.8	187.1	296.0	483.0	831.0	1,406.1	2,992.2
2000.....	20.3	40.1	71.3	122.7	190.6	306.1	515.9	846.5	1,473.2	3,200.0
1999.....	20.1	40.4	72.9	122.0	196.5	327.6	534.1	887.5	1,522.9	3,267.1

White, female

2006 4/.....	5.2	10.2	17.1	30.0	57.0	95.6	178.7	328.5	662.7	1,960.0
2005 4/.....	5.1	10.2	18.0	31.4	59.2	105.2	187.9	357.2	711.6	2,119.0
2004 4/.....	5.3	10.2	17.7	32.5	61.8	105.2	199.7	367.4	743.3	2,196.8
2003 4/.....	5.8	10.5	17.7	33.8	63.8	112.8	211.9	395.6	795.8	2,395.5
2002.....	5.1	10.0	17.6	35.3	65.1	118.9	219.3	412.6	827.2	2,493.2
2001.....	5.2	9.0	18.7	35.8	68.2	124.1	230.0	429.2	863.8	2,611.0
2000.....	4.7	8.6	18.8	37.8	69.0	128.7	233.9	447.7	888.4	2,753.0
1999.....	4.5	9.0	20.3	38.9	76.3	135.5	248.5	473.4	926.8	2,855.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other forms of chronic ischemic heart disease (I20,I25)

Black, both sexes

2006 4/.....	18.1	40.7	77.3	122.3	190.5	272.5	401.5	602.0	955.6	1,791.4
2005 4/.....	17.9	39.5	82.8	129.0	208.5	276.7	414.4	649.7	1,003.4	1,885.0
2004 4/.....	20.3	41.4	80.8	129.1	204.8	298.4	437.7	671.6	1,023.9	2,019.1
2003 4/.....	20.1	42.8	82.2	131.7	224.6	317.7	480.4	729.7	1,087.4	2,195.3
2002.....	20.7	44.0	83.7	144.6	221.6	323.9	476.6	754.2	1,112.3	2,240.6
2001.....	21.5	43.7	83.6	144.1	235.3	346.1	500.3	790.1	1,155.2	2,351.2
2000.....	21.5	47.6	82.7	152.4	236.7	341.9	531.5	796.3	1,171.8	2,387.1
1999.....	23.8	45.4	92.2	158.4	241.2	367.9	551.4	817.7	1,183.5	2,469.8

Black, male

2006 4/.....	24.6	58.7	116.0	182.2	279.4	393.0	567.5	838.6	1,213.7	1,819.4
2005 4/.....	24.7	59.5	124.8	188.8	302.5	387.6	569.3	856.5	1,271.1	1,871.2
2004 4/.....	28.8	59.1	119.5	187.9	300.1	411.1	593.0	901.3	1,258.3	2,068.3
2003 4/.....	29.4	62.9	122.5	189.9	317.6	439.5	666.5	970.9	1,383.5	2,147.8
2002.....	29.0	62.5	122.8	212.7	326.3	447.5	643.9	954.4	1,378.6	2,254.6
2001.....	31.2	64.0	125.1	210.1	333.1	469.8	698.8	1,022.7	1,443.9	2,369.1
2000.....	30.3	69.2	121.3	229.5	326.9	448.3	711.4	1,024.9	1,444.9	2,446.0
1999.....	34.9	66.9	138.0	227.8	348.1	491.2	743.2	1,028.6	1,437.0	2,561.5

Black, female

2006 4/.....	12.4	24.9	44.5	73.0	121.2	183.5	287.4	460.4	819.3	1,779.9
2005 4/.....	11.9	22.1	47.3	79.7	135.3	194.4	308.4	526.4	862.1	1,890.6
2004 4/.....	12.8	26.0	48.2	80.7	130.5	214.6	332.3	535.6	900.4	1,999.6
2003 4/.....	11.9	25.3	48.2	83.8	152.1	227.3	354.9	586.3	932.3	2,213.8
2002.....	13.3	28.0	50.5	88.6	139.7	232.2	364.3	634.5	973.6	2,235.2
2001.....	13.0	26.1	48.2	89.9	158.2	254.8	368.1	651.2	1,008.1	2,344.5
2000.....	13.7	28.8	49.7	89.2	165.6	263.4	412.0	658.5	1,034.3	2,365.2
1999.....	14.1	26.8	53.5	101.6	157.4	276.5	424.2	690.6	1,057.9	2,435.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All races 3/, both sexes

2006.....	4.6	9.4	17.9	26.1	38.5	47.4	64.4	89.1	147.1	345.5
2005.....	4.4	9.6	18.6	26.9	37.7	46.8	64.6	96.5	161.9	376.4
2004.....	4.8	9.5	18.2	26.6	39.1	49.8	69.0	100.8	166.6	400.4
2003.....	4.7	9.9	18.6	26.6	40.5	53.3	73.3	110.7	186.2	442.9
2002.....	4.8	9.8	17.5	27.9	39.7	53.9	76.5	116.6	187.6	474.5
2001.....	4.6	8.8	17.4	27.7	40.6	55.3	76.7	121.8	197.2	493.6
2000.....	4.4	9.4	17.5	28.5	40.7	55.7	81.9	125.0	207.8	534.6
1999.....	4.7	9.1	17.4	28.6	43.4	59.6	86.8	131.6	218.2	563.4

All races 3/, male

2006.....	7.0	14.9	29.3	41.5	58.9	69.9	90.6	117.9	188.1	365.3
2005.....	7.0	15.3	30.0	42.5	57.7	68.3	91.5	127.0	204.6	390.7
2004.....	7.7	15.3	29.6	42.0	58.4	73.1	96.4	133.8	210.5	414.0
2003.....	7.4	16.0	30.5	41.8	60.6	77.1	102.8	146.6	233.1	462.1
2002.....	7.8	15.6	28.4	44.8	59.7	78.8	106.7	154.5	232.4	491.8
2001.....	7.2	14.6	28.3	43.4	61.8	79.0	107.6	160.2	247.4	502.0
2000.....	7.1	15.5	28.4	45.8	61.3	80.8	115.6	167.1	252.8	550.8
1999.....	7.6	14.9	28.0	45.0	64.6	87.3	120.8	172.7	264.3	578.6

All races 3/, female

2006.....	2.2	4.1	7.1	11.5	20.0	27.7	43.1	67.9	121.4	336.1
2005.....	1.9	4.0	7.7	12.2	19.4	28.1	42.9	74.3	135.4	369.9
2004.....	2.0	3.9	7.4	12.0	21.5	29.5	47.0	77.0	139.6	394.3
2003.....	2.0	4.0	7.2	12.4	22.2	32.6	49.7	85.2	157.6	434.4
2002.....	1.8	4.1	7.1	12.1	21.4	32.4	52.5	89.7	160.4	467.1
2001.....	2.1	3.3	7.0	12.9	21.4	34.9	52.1	94.9	167.1	490.1
2000.....	1.8	3.5	7.2	12.4	22.1	34.1	55.4	95.7	181.2	528.0
1999.....	1.8	3.5	7.3	13.4	24.1	36.0	60.2	103.1	191.3	557.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

White, both sexes

2006 4/.....	4.3	8.6	16.2	23.4	34.9	42.7	58.6	83.0	140.9	343.2
2005 4/.....	4.3	9.0	16.6	23.8	33.3	42.5	58.7	89.9	154.9	373.8
2004 4/.....	4.6	8.7	16.4	23.7	35.1	44.4	63.4	93.3	160.3	395.6
2003 4/.....	4.4	9.0	16.7	24.2	35.4	47.5	66.6	103.0	179.6	439.3
2002.....	4.4	9.1	15.6	24.7	35.3	47.3	69.7	107.8	180.5	470.6
2001.....	4.2	7.9	15.6	23.9	35.0	47.9	69.4	112.2	190.6	489.7
2000.....	4.0	8.3	15.7	24.8	35.2	49.5	73.7	116.0	198.5	529.5
1999.....	4.2	8.0	15.2	24.6	37.5	52.3	77.9	121.5	209.8	557.5

White, male

2006 4/.....	6.7	13.8	26.8	37.6	53.1	62.8	82.2	109.7	180.2	362.7
2005 4/.....	7.0	14.6	27.2	38.2	51.2	62.2	82.8	118.8	194.8	392.6
2004 4/.....	7.5	14.4	27.3	37.8	52.4	66.1	89.3	123.9	203.3	408.8
2003 4/.....	6.9	14.8	27.7	38.3	53.4	69.2	92.5	136.9	224.4	458.7
2002.....	7.4	14.7	25.7	40.0	53.6	69.4	98.0	143.3	222.2	488.0
2001.....	6.6	13.3	25.7	38.3	54.0	69.3	97.6	148.7	239.9	498.6
2000.....	6.6	14.0	25.9	40.0	54.2	72.8	104.4	154.6	241.6	545.6
1999.....	7.0	13.2	24.8	39.4	56.2	77.6	109.0	159.8	253.9	571.2

White, female

2006 4/.....	1.8	3.4	5.7	9.7	17.9	24.7	39.1	63.1	116.1	334.1
2005 4/.....	1.5	3.4	6.2	9.9	16.6	25.1	38.9	68.5	129.9	365.1
2004 4/.....	1.7	3.0	5.8	10.2	19.1	25.1	42.1	70.9	133.6	389.7
2003 4/.....	1.8	3.2	5.9	10.6	18.7	28.3	45.4	78.5	152.0	430.7
2002.....	1.5	3.5	5.6	10.2	18.4	27.8	46.6	82.5	154.9	463.0
2001.....	1.8	2.6	5.7	10.1	17.6	29.1	46.7	86.4	160.8	486.0
2000.....	1.4	2.7	5.8	10.2	17.6	29.0	49.0	88.8	172.9	522.9
1999.....	1.3	2.8	5.8	10.6	20.3	30.3	53.1	94.6	183.8	552.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerotic cardiovascular disease, so described (I25.0)

Black, both sexes

2006 4/.....	7.6	16.7	34.8	52.2	78.1	97.2	126.6	167.0	244.5	428.1
2005 4/.....	6.5	15.5	37.3	56.6	83.4	92.4	130.4	183.1	265.1	469.6
2004 4/.....	7.5	17.5	35.7	55.9	81.5	105.1	139.0	198.5	268.4	505.1
2003 4/.....	7.3	17.4	35.9	53.2	91.8	111.4	149.2	214.1	282.8	543.2
2002.....	8.1	17.0	35.9	60.5	85.5	119.5	157.5	231.9	300.4	575.5
2001.....	8.2	16.8	35.7	64.2	94.4	128.0	162.6	244.0	301.5	594.5
2000.....	7.4	19.4	36.4	65.4	94.0	119.6	176.8	239.7	340.0	646.6
1999.....	9.1	19.1	38.4	67.8	98.2	133.9	188.9	261.7	338.9	668.6

Black, male

2006 4/.....	10.5	24.8	54.4	82.0	126.0	151.8	190.7	239.0	329.3	465.6
2005 4/.....	8.9	23.1	58.1	87.6	130.3	140.2	200.8	251.7	361.7	450.8
2004 4/.....	10.7	25.1	54.0	88.3	127.8	153.1	197.1	278.0	350.8	531.5
2003 4/.....	11.4	26.4	57.8	81.3	140.0	161.3	228.9	301.3	380.2	583.9
2002.....	11.8	25.5	55.0	97.1	130.5	180.0	220.3	321.0	417.3	599.3
2001.....	12.4	25.7	56.0	97.0	144.0	183.5	241.0	330.4	388.5	609.3
2000.....	11.0	30.3	56.8	106.4	137.1	168.6	257.1	338.3	425.7	701.2
1999.....	13.7	29.9	59.7	103.8	147.9	197.3	271.0	350.0	435.1	715.9

Black, female

2006 4/.....	5.1	9.6	18.2	27.7	40.8	56.8	82.6	124.0	199.7	412.7
2005 4/.....	4.3	8.9	19.7	31.1	46.8	56.9	82.2	142.1	214.0	477.1
2004 4/.....	4.7	10.9	20.2	29.3	45.3	69.4	99.5	151.5	224.9	494.6
2003 4/.....	3.7	9.6	17.3	30.0	54.2	74.4	95.5	162.2	231.7	527.4
2002.....	4.8	9.7	19.7	30.4	50.3	74.6	115.4	178.7	239.5	566.4
2001.....	4.5	9.0	18.4	37.3	55.4	87.1	110.3	192.5	257.1	589.0
2000.....	4.2	10.0	19.0	31.8	60.0	83.5	123.4	180.3	296.8	626.3
1999.....	5.0	9.8	20.4	38.4	59.2	87.0	134.3	208.5	291.3	650.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 45-49 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85 years
years years years years years years years years years and over

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races 3/, both sexes

2006.....	8.1	15.7	26.8	43.9	73.6	121.9	212.5	379.0	700.4	1,712.0
2005.....	8.2	15.5	27.7	44.9	79.9	131.2	226.9	404.6	740.4	1,837.8
2004.....	8.4	16.4	27.4	45.7	79.9	134.8	236.6	417.4	769.0	1,879.3
2003.....	8.9	16.7	28.0	49.0	85.6	144.5	253.8	444.9	811.1	2,028.5
2002.....	8.8	17.1	29.9	52.3	88.2	152.9	261.8	465.6	841.9	2,088.0
2001.....	8.6	16.4	29.8	52.4	93.5	160.1	275.3	483.2	867.3	2,177.7
2000.....	9.0	17.0	30.3	56.7	95.9	165.3	287.4	496.2	894.4	2,287.5
1999.....	8.8	17.4	32.8	57.3	99.5	175.7	299.3	521.0	924.9	2,353.6

All races 3/, male

2006.....	12.3	23.9	41.2	66.6	107.4	174.8	296.7	531.1	945.4	1,979.5
2005.....	12.5	23.7	42.4	67.0	116.4	184.3	316.4	558.3	997.9	2,120.9
2004.....	12.7	25.0	41.8	67.5	116.8	191.3	328.2	580.4	1,027.4	2,129.6
2003.....	13.3	25.5	43.0	73.3	125.0	206.7	355.0	616.5	1,088.5	2,260.1
2002.....	13.6	26.6	46.5	78.0	129.6	217.8	367.6	647.1	1,123.8	2,325.6
2001.....	13.2	25.4	45.7	78.8	135.6	228.4	385.2	675.1	1,145.1	2,411.8
2000.....	14.0	27.0	46.7	84.9	139.5	234.0	409.1	684.5	1,204.8	2,561.9
1999.....	13.9	27.7	50.2	84.9	143.5	250.0	424.4	718.0	1,240.3	2,614.4

All races 3/, female

2006.....	3.8	7.7	12.9	22.6	42.7	75.6	144.0	267.7	547.2	1,586.8
2005.....	4.0	7.4	13.5	23.9	46.5	84.8	154.5	293.0	580.4	1,707.7
2004.....	4.1	8.0	13.6	25.0	46.3	85.6	162.8	300.3	609.8	1,766.7
2003.....	4.5	8.2	13.7	26.0	49.8	90.5	172.7	322.7	641.5	1,926.2
2002.....	4.2	7.8	13.9	28.2	50.5	97.0	177.3	337.3	670.4	1,984.9
2001.....	4.0	7.6	14.7	27.7	55.4	101.2	188.0	348.9	700.7	2,079.8
2000.....	4.0	7.3	14.6	30.3	56.4	106.3	191.6	365.1	711.3	2,175.8
1999.....	3.9	7.4	16.1	31.5	59.7	112.1	201.5	384.0	740.3	2,248.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

White, both sexes

2006 4/.....	8.0	14.9	25.5	41.8	70.8	118.0	210.4	380.4	708.8	1,763.4
2005 4/.....	8.0	14.6	26.0	42.9	76.7	127.6	225.2	405.9	750.9	1,896.0
2004 4/.....	8.1	15.7	25.8	43.5	77.0	131.0	234.7	418.8	778.7	1,929.2
2003 4/.....	8.5	16.0	26.5	46.8	82.2	140.3	251.0	445.0	821.5	2,079.5
2002.....	8.6	16.1	28.3	49.9	84.5	150.0	260.7	466.0	854.2	2,143.3
2001.....	8.1	15.4	28.4	50.5	90.2	156.6	273.8	483.6	877.5	2,233.8
2000.....	8.5	16.0	29.1	54.4	92.2	162.0	285.8	496.4	908.0	2,352.7
1999.....	8.2	16.6	31.1	54.8	96.5	172.6	297.1	522.7	938.2	2,416.1

White, male

2006 4/.....	12.5	23.1	39.9	64.1	104.8	170.8	295.6	534.6	965.4	2,056.2
2005 4/.....	12.3	22.5	40.4	65.0	113.3	181.1	317.3	564.5	1,021.0	2,204.7
2004 4/.....	12.4	24.3	40.0	65.5	114.0	188.4	328.6	585.8	1,050.7	2,200.2
2003 4/.....	12.9	24.7	41.5	71.3	122.4	203.4	354.1	622.0	1,109.8	2,338.4
2002.....	13.4	25.9	45.0	75.8	125.4	216.8	368.7	655.7	1,151.0	2,403.6
2001.....	12.8	24.4	44.2	76.5	133.1	226.8	385.4	682.4	1,166.2	2,493.6
2000.....	13.7	26.1	45.5	82.7	136.4	233.3	411.5	691.9	1,231.6	2,654.4
1999.....	13.1	27.2	48.1	82.6	140.3	249.9	425.1	727.7	1,269.0	2,695.9

White, female

2006 4/.....	3.4	6.8	11.4	20.3	39.1	70.9	139.6	265.4	546.6	1,626.0
2005 4/.....	3.6	6.8	11.8	21.5	42.6	80.1	149.1	288.6	581.7	1,753.8
2004 4/.....	3.7	7.2	11.9	22.3	42.8	80.1	157.6	296.5	609.7	1,807.0
2003 4/.....	4.0	7.3	11.8	23.3	45.1	84.5	166.6	317.1	643.8	1,964.9
2002.....	3.6	6.5	12.0	25.2	46.7	91.1	172.7	330.1	672.3	2,030.2
2001.....	3.5	6.4	13.0	25.7	50.7	95.0	183.3	342.8	703.0	2,125.0
2000.....	3.3	5.9	13.0	27.5	51.4	99.6	184.9	358.9	715.6	2,230.1
1999.....	3.2	6.2	14.5	28.3	56.1	105.2	195.4	378.8	743.0	2,303.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Black, both sexes

2006 4/.....	10.5	24.0	42.5	70.1	112.4	175.3	274.8	435.0	711.2	1,363.3
2005 4/.....	11.4	24.0	45.5	72.4	125.2	184.3	284.0	466.7	738.4	1,415.4
2004 4/.....	12.8	23.9	45.2	73.2	123.4	193.3	298.8	473.1	755.5	1,514.0
2003 4/.....	12.8	25.4	46.4	78.5	132.8	206.3	331.2	515.6	804.7	1,652.1
2002.....	12.6	27.0	47.8	84.1	136.1	204.4	319.1	522.2	811.9	1,665.1
2001.....	13.3	26.9	47.9	79.9	140.9	218.1	337.8	546.0	853.7	1,756.7
2000.....	14.1	28.1	46.3	87.0	142.8	222.3	354.7	556.6	831.8	1,740.5
1999.....	14.7	26.3	53.8	90.6	143.1	233.9	362.6	555.9	844.6	1,801.2

Black, male

2006 4/.....	14.0	33.9	61.6	100.1	153.4	241.1	376.8	599.6	884.4	1,353.8
2005 4/.....	15.8	36.4	66.7	101.3	172.1	247.4	368.4	604.8	909.5	1,420.4
2004 4/.....	18.1	34.0	65.5	99.6	172.3	258.0	395.9	623.3	907.5	1,536.8
2003 4/.....	18.0	36.5	64.7	108.6	177.6	278.2	437.5	669.6	1,003.3	1,563.9
2002.....	17.2	37.0	67.8	115.7	195.8	267.5	423.6	633.4	961.4	1,655.3
2001.....	18.8	38.3	69.2	113.1	189.2	286.4	457.8	692.3	1,055.4	1,759.8
2000.....	19.3	39.0	64.5	123.2	189.9	279.7	454.3	686.6	1,019.2	1,744.8
1999.....	21.2	37.0	78.2	124.1	200.3	293.9	472.2	678.6	1,001.9	1,845.5

Black, female

2006 4/.....	7.3	15.3	26.3	45.3	80.4	126.7	204.8	336.4	619.6	1,367.2
2005 4/.....	7.6	13.2	27.6	48.5	88.6	137.5	226.2	384.3	648.0	1,413.4
2004 4/.....	8.1	15.1	28.0	51.4	85.2	145.2	232.8	384.1	675.5	1,505.0
2003 4/.....	8.3	15.7	30.9	53.8	97.9	153.0	259.4	424.0	700.6	1,686.4
2002.....	8.5	18.4	30.8	58.2	89.4	157.6	248.9	455.8	734.1	1,668.8
2001.....	8.5	17.0	29.8	52.6	102.9	167.6	257.8	458.7	751.0	1,755.5
2000.....	9.5	18.7	30.7	57.4	105.6	179.9	288.6	478.2	737.5	1,738.9
1999.....	9.1	17.1	33.1	63.3	98.2	189.5	289.9	482.0	766.6	1,784.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other heart diseases (I26-I51)

All races 3/, both sexes

2006.....	11.4	17.7	25.0	36.3	55.5	85.7	144.6	259.0	493.3	1,342.8
2005.....	11.6	17.9	25.5	36.8	57.4	88.2	150.8	264.6	500.7	1,377.4
2004.....	11.5	17.9	25.6	35.7	57.5	90.3	151.8	268.1	503.3	1,373.9
2003.....	12.6	17.9	25.3	39.1	59.7	97.3	160.2	279.1	525.5	1,427.2
2002.....	12.4	18.1	25.9	39.4	60.6	97.9	164.5	286.7	536.3	1,443.5
2001.....	11.9	17.9	25.9	39.7	62.0	100.7	164.7	287.2	537.0	1,475.5
2000.....	12.3	17.6	25.9	40.2	64.3	103.6	167.1	290.2	542.8	1,509.4
1999.....	12.5	17.8	25.9	41.8	66.5	105.2	175.0	295.4	553.5	1,512.9

All races 3/, male

2006.....	14.0	22.6	32.4	46.6	69.5	106.9	173.2	302.8	566.0	1,338.1
2005.....	14.8	22.6	33.3	47.4	71.3	108.0	180.8	310.3	568.1	1,377.8
2004.....	14.4	23.2	32.8	46.4	72.7	111.3	181.2	316.2	573.7	1,377.7
2003.....	15.6	22.7	32.4	49.9	74.9	119.4	195.4	334.0	599.9	1,431.3
2002.....	15.9	23.5	33.6	49.4	75.7	118.8	199.8	340.7	612.4	1,448.6
2001.....	14.9	23.1	32.7	49.4	77.7	121.5	200.9	344.7	622.1	1,473.4
2000.....	15.4	23.0	33.4	52.0	80.8	126.9	205.9	345.1	626.8	1,549.1
1999.....	15.6	23.0	33.3	54.1	84.3	133.4	217.0	350.9	641.4	1,548.9

All races 3/, female

2006.....	8.9	12.9	18.0	26.6	42.7	67.2	121.3	226.9	447.9	1,345.0
2005.....	8.5	13.3	18.0	26.9	44.8	70.9	126.4	231.4	458.9	1,377.3
2004.....	8.6	12.8	18.6	25.7	43.8	72.0	128.2	233.5	459.9	1,372.1
2003.....	9.5	13.2	18.6	28.8	45.9	78.1	131.9	240.0	480.0	1,425.4
2002.....	9.0	12.8	18.5	30.1	46.8	79.8	136.3	248.6	490.0	1,441.3
2001.....	8.9	12.8	19.5	30.6	47.8	82.8	135.9	247.0	486.0	1,476.4
2000.....	9.2	12.3	18.7	29.2	49.3	83.6	136.6	252.1	493.2	1,493.2
1999.....	9.4	12.8	18.9	30.2	50.5	81.0	142.2	256.8	502.0	1,498.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other heart diseases (I26-I51)

White, both sexes

2006 4/.....	9.9	15.2	21.4	32.0	50.8	79.9	139.1	256.9	496.7	1,386.4
2005 4/.....	10.0	15.2	21.6	32.4	51.6	81.9	145.2	260.8	504.5	1,419.0
2004 4/.....	9.7	15.0	21.8	31.3	52.7	84.7	146.5	262.9	507.6	1,408.8
2003 4/.....	10.5	15.2	21.4	34.8	53.8	90.7	153.2	276.4	528.9	1,461.4
2002.....	10.7	15.0	22.1	35.0	55.3	92.4	158.5	282.4	539.1	1,475.5
2001.....	9.7	14.9	22.3	34.8	57.2	94.1	159.0	281.6	538.8	1,509.6
2000.....	10.2	14.6	22.1	35.9	59.0	97.3	160.6	284.5	545.7	1,541.6
1999.....	10.2	14.6	22.2	36.3	60.4	98.6	167.6	288.5	553.7	1,543.3

White, male

2006 4/.....	12.1	19.5	27.7	41.3	63.5	100.0	166.3	300.6	572.3	1,381.2
2005 4/.....	12.8	19.6	28.7	41.9	64.7	100.4	173.5	306.2	575.6	1,424.2
2004 4/.....	12.4	19.6	28.7	40.9	66.8	104.6	176.0	309.6	578.2	1,407.9
2003 4/.....	13.5	19.7	27.5	44.5	68.3	111.8	188.4	332.0	604.3	1,467.6
2002.....	14.1	19.9	29.2	44.1	69.9	112.5	193.0	336.5	617.5	1,482.1
2001.....	12.7	19.7	28.9	43.5	72.1	114.9	193.7	338.2	621.9	1,503.4
2000.....	13.0	19.7	28.6	46.9	74.8	119.7	199.2	339.4	630.8	1,580.5
1999.....	13.3	19.4	28.7	47.5	76.9	125.6	209.0	344.0	641.4	1,571.5

White, female

2006 4/.....	7.7	11.0	15.2	23.0	38.9	62.0	116.6	224.3	449.0	1,388.9
2005 4/.....	7.1	10.8	14.5	23.3	39.5	65.4	121.9	227.1	460.0	1,416.6
2004 4/.....	6.9	10.5	15.1	22.0	39.7	66.9	122.2	228.7	463.8	1,409.2
2003 4/.....	7.5	10.8	15.4	25.5	40.4	72.1	124.3	236.1	482.4	1,458.6
2002.....	7.3	10.2	15.2	26.2	41.8	74.7	130.3	243.7	491.0	1,472.6
2001.....	6.6	10.1	15.8	26.5	43.5	75.9	130.9	241.5	488.5	1,512.2
2000.....	7.4	9.6	15.8	25.4	44.3	77.7	129.5	245.9	495.2	1,525.8
1999.....	7.2	9.8	15.8	25.7	45.1	75.0	134.8	249.5	501.9	1,531.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Other heart diseases (I26-I51)

Black, both sexes

2006 4/.....	24.1	38.4	57.4	80.0	111.5	155.8	229.4	342.1	566.8	1,094.3
2005 4/.....	25.3	40.3	60.2	82.0	120.4	162.1	237.3	368.3	563.4	1,135.3
2004 4/.....	26.3	41.7	60.2	82.6	114.5	159.9	240.5	381.0	551.3	1,180.7
2003 4/.....	28.9	40.8	61.1	85.3	124.6	176.1	265.5	364.8	574.9	1,238.1
2002.....	26.6	43.6	60.8	87.1	121.2	169.4	255.8	386.7	594.7	1,270.4
2001.....	29.2	43.1	60.9	91.4	118.8	179.7	254.5	402.9	601.8	1,277.7
2000.....	28.5	43.3	62.0	86.2	122.8	181.5	266.8	406.9	590.7	1,306.0
1999.....	30.2	44.9	62.2	96.8	133.5	185.5	284.2	417.4	621.2	1,302.8

Black, male

2006 4/.....	30.0	49.8	76.5	102.5	147.0	199.2	288.8	409.0	645.5	1,101.8
2005 4/.....	31.8	50.1	77.6	107.6	151.1	204.4	300.1	446.6	614.5	1,088.5
2004 4/.....	31.6	54.3	75.0	109.0	149.1	202.5	284.5	469.5	653.5	1,224.9
2003 4/.....	34.7	51.0	79.5	113.5	158.3	223.6	318.0	434.1	668.1	1,220.9
2002.....	32.0	56.7	77.1	110.7	150.7	208.7	317.5	454.2	675.6	1,238.3
2001.....	33.7	53.5	73.7	117.2	151.5	212.9	318.8	494.8	743.2	1,349.1
2000.....	34.8	54.3	82.1	112.1	153.6	224.8	318.9	480.5	696.3	1,402.3
1999.....	36.2	56.1	81.9	127.0	172.7	237.8	353.1	489.1	747.3	1,454.1

Black, female

2006 4/.....	18.8	28.5	41.3	61.4	83.9	123.8	188.5	302.1	525.2	1,091.2
2005 4/.....	19.6	31.7	45.4	60.8	96.5	130.8	194.4	321.6	536.4	1,154.2
2004 4/.....	21.6	30.6	47.7	60.9	87.6	128.3	210.7	328.6	497.4	1,163.2
2003 4/.....	23.8	31.9	45.4	62.1	98.2	140.9	230.2	323.7	526.1	1,244.8
2002.....	21.8	32.3	47.0	67.7	98.2	140.2	214.4	346.3	552.6	1,282.7
2001.....	25.3	34.0	50.1	70.3	93.1	155.1	211.6	348.1	529.8	1,251.2
2000.....	22.9	33.9	44.8	64.9	98.4	149.6	232.2	362.6	537.5	1,270.2
1999.....	24.9	35.2	45.4	72.1	102.8	146.8	238.5	374.2	558.7	1,246.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute and subacute endocarditis (I33)

All races 3/, both sexes

2006.....	0.2	0.4	0.5	0.7	0.8	1.0	1.4	2.2	2.7	2.2
2005.....	0.2	0.4	0.5	0.6	0.7	1.1	1.4	2.4	2.7	3.0
2004.....	0.2	0.4	0.5	0.5	0.7	1.0	1.5	2.1	2.9	3.2
2003.....	0.3	0.4	0.5	0.6	0.9	1.0	1.5	2.3	2.7	3.0
2002.....	0.3	0.4	0.5	0.5	0.8	1.1	1.5	2.2	2.7	2.6
2001.....	0.3	0.4	0.5	0.5	0.8	1.1	1.7	2.2	3.0	2.5
2000.....	0.3	0.4	0.6	0.7	0.7	1.1	1.3	2.2	2.8	2.8
1999.....	0.3	0.3	0.4	0.7	0.7	1.1	1.5	2.1	2.7	2.9

All races 3/, male

2006.....	0.3	0.6	0.7	0.9	0.9	1.2	1.6	2.3	3.3	2.3
2005.....	0.2	0.4	0.7	0.8	0.9	1.2	1.8	3.1	3.5	5.1
2004.....	0.3	0.5	0.7	0.6	0.8	1.4	1.4	2.3	4.2	4.8
2003.....	0.4	0.6	0.7	0.8	1.1	1.3	1.6	2.9	3.7	4.1
2002.....	0.4	0.5	0.7	0.6	0.9	1.5	1.7	2.5	3.7	3.5
2001.....	0.3	0.7	0.7	0.6	1.1	1.1	2.1	3.1	4.3	3.8
2000.....	0.4	0.6	0.7	0.8	0.9	1.3	1.6	3.1	4.0	4.2
1999.....	0.4	0.5	0.7	1.0	0.8	1.2	1.7	2.8	3.9	4.7

All races 3/, female

2006.....	0.2	0.3	0.3	0.4	0.7	0.7	1.2	2.2	2.3	2.1
2005.....	*	0.3	0.3	0.4	0.6	1.0	1.1	1.9	2.2	2.1
2004.....	*	0.3	0.4	0.3	0.6	0.7	1.7	2.0	2.1	2.5
2003.....	0.2	0.3	0.3	0.4	0.8	0.9	1.4	1.9	2.1	2.5
2002.....	0.2	0.3	0.3	0.4	0.8	0.8	1.3	1.9	2.1	2.2
2001.....	0.3	0.2	0.3	0.5	0.5	1.0	1.3	1.6	2.2	2.0
2000.....	0.2	0.3	0.4	0.5	0.6	0.9	1.1	1.6	2.0	2.3
1999.....	*	*	0.2	0.5	0.7	1.0	1.4	1.7	2.0	2.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute and subacute endocarditis (I33)

White, both sexes

2006 4/.....	0.2	0.4	0.4	0.5	0.7	0.9	1.4	2.1	2.8	2.1
2005 4/.....	0.1	0.3	0.3	0.4	0.7	1.0	1.5	2.4	2.6	3.1
2004 4/.....	0.2	0.3	0.4	0.4	0.6	1.0	1.5	2.0	2.9	3.2
2003 4/.....	0.3	0.3	0.4	0.5	0.8	1.0	1.4	2.3	2.8	3.1
2002.....	0.2	0.3	0.4	0.4	0.8	1.0	1.4	2.2	2.9	2.7
2001.....	0.2	0.4	0.4	0.4	0.7	0.9	1.5	2.0	3.0	2.6
2000.....	0.3	0.3	0.4	0.5	0.7	1.1	1.3	2.2	2.7	2.7
1999.....	0.2	0.3	0.3	0.5	0.6	1.0	1.4	2.2	2.8	3.0

White, male

2006 4/.....	0.3	0.6	0.5	0.8	0.8	1.1	1.6	2.3	3.3	2.5
2005 4/.....	*	0.3	0.5	0.6	0.8	1.2	1.8	3.2	3.5	5.3
2004 4/.....	0.3	0.4	0.5	0.6	0.7	1.3	1.4	2.2	4.3	4.9
2003 4/.....	0.3	0.4	0.5	0.6	1.0	1.2	1.5	2.8	3.9	4.4
2002.....	0.3	0.4	0.6	0.4	0.9	1.3	1.7	2.7	3.9	3.6
2001.....	0.3	0.5	0.5	0.5	0.9	0.9	1.8	3.0	4.3	3.9
2000.....	0.4	0.5	0.6	0.7	0.9	1.2	1.6	3.2	4.0	4.2
1999.....	0.3	0.4	0.5	0.8	0.7	1.2	1.6	3.0	4.0	5.1

White, female

2006 4/.....	*	0.2	0.2	0.3	0.5	0.8	1.1	2.0	2.5	1.9
2005 4/.....	*	0.2	*	0.3	0.5	0.9	1.2	1.7	2.0	2.1
2004 4/.....	*	*	0.2	*	0.5	0.6	1.6	1.8	2.1	2.4
2003 4/.....	*	0.2	0.2	0.4	0.6	0.8	1.3	1.8	2.1	2.6
2002.....	*	*	*	0.3	0.7	0.7	1.2	1.8	2.3	2.3
2001.....	*	*	*	0.4	0.4	0.9	1.3	1.3	2.2	2.1
2000.....	*	*	0.3	0.4	0.5	1.0	1.0	1.6	2.0	2.1
1999.....	*	*	*	*	0.6	0.8	1.3	1.6	2.0	2.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races 3/, both sexes

2006.....	0.2	0.2	0.3	0.3	0.5	0.4	0.8	1.1	1.3	1.8
2005.....	0.2	0.3	0.3	0.3	0.4	0.7	0.8	1.1	1.3	1.9
2004.....	0.2	0.2	0.3	0.3	0.5	0.6	0.6	1.1	1.3	1.8
2003.....	0.2	0.3	0.3	0.4	0.4	0.8	0.8	0.9	1.5	1.6
2002.....	0.2	0.3	0.3	0.4	0.4	0.6	1.0	1.1	1.4	2.1
2001.....	0.2	0.3	0.3	0.4	0.5	0.5	0.8	1.2	1.3	1.9
2000.....	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.1	1.3	1.8
1999.....	0.3	0.3	0.3	0.5	0.6	0.7	0.9	1.2	1.5	2.0

All races 3/, male

2006.....	0.2	0.3	0.3	0.4	0.6	0.6	0.9	1.1	1.7	1.8
2005.....	0.3	0.4	0.4	0.3	0.5	0.7	1.0	1.4	1.5	1.9
2004.....	0.2	0.2	0.4	0.4	0.6	0.8	0.7	1.2	1.4	2.3
2003.....	0.3	0.4	0.3	0.4	0.4	1.0	1.0	1.1	1.5	1.5
2002.....	0.3	0.4	0.3	0.5	0.6	0.7	1.1	1.1	1.3	2.4
2001.....	*	0.3	0.3	0.4	0.6	0.6	0.8	1.3	1.1	1.8
2000.....	0.3	0.3	*	0.5	0.4	0.5	1.1	1.4	1.7	2.1
1999.....	0.3	0.3	0.3	0.6	0.6	0.8	0.9	1.3	1.5	2.5

All races 3/, female

2006.....	*	*	0.2	0.2	0.4	*	0.7	1.1	1.0	1.8
2005.....	*	0.3	0.3	0.3	0.4	0.6	0.6	0.9	1.1	1.9
2004.....	*	0.2	0.2	0.3	0.4	0.5	0.6	1.1	1.3	1.6
2003.....	0.2	0.2	*	0.3	0.3	0.7	0.7	0.8	1.5	1.7
2002.....	*	*	0.2	0.4	*	0.5	1.0	1.1	1.4	2.0
2001.....	0.3	0.3	0.3	0.4	*	*	0.9	1.1	1.4	2.0
2000.....	*	*	0.2	*	0.4	0.5	0.7	0.9	1.0	1.7
1999.....	0.3	0.3	0.2	0.3	0.6	0.7	0.8	1.2	1.5	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of pericardium and acute myocarditis (I30-I31,I40)

White, both sexes

2006 4/.....	0.2	0.2	0.3	0.3	0.5	0.5	0.8	1.1	1.3	1.8
2005 4/.....	0.2	0.3	0.3	0.3	0.4	0.7	0.8	1.1	1.2	1.9
2004 4/.....	0.1	0.2	0.3	0.3	0.4	0.6	0.6	1.0	1.3	1.8
2003 4/.....	0.2	0.3	0.2	0.4	0.4	0.8	0.8	1.0	1.5	1.7
2002.....	0.2	0.2	0.2	0.4	0.4	0.6	1.0	1.0	1.3	2.2
2001.....	0.2	0.2	0.2	0.3	0.5	0.4	0.9	1.2	1.3	1.9
2000.....	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.0	1.4	1.8
1999.....	0.2	0.3	0.2	0.5	0.6	0.7	0.8	1.2	1.5	2.0

White, male

2006 4/.....	*	0.3	0.3	0.4	0.7	0.6	0.9	1.2	1.8	1.8
2005 4/.....	0.2	0.3	0.4	0.3	0.4	0.7	1.0	1.3	1.5	1.9
2004 4/.....	*	*	0.3	0.4	0.6	0.6	0.7	1.2	1.4	2.2
2003 4/.....	0.2	0.3	0.3	0.4	0.5	1.0	1.1	1.1	1.6	1.7
2002.....	0.3	0.3	0.3	0.4	0.6	0.6	1.1	1.1	1.2	2.5
2001.....	*	0.2	0.3	0.4	0.6	0.5	0.9	1.3	1.1	2.0
2000.....	0.2	*	*	0.5	*	*	1.1	1.2	1.9	2.1
1999.....	0.3	0.3	*	0.6	0.6	0.7	0.9	1.3	1.5	2.5

White, female

2006 4/.....	*	*	*	*	0.4	*	0.7	1.0	1.0	1.9
2005 4/.....	*	*	0.3	*	0.4	0.6	0.6	0.9	1.1	1.8
2004 4/.....	*	*	*	0.3	*	0.5	0.6	0.9	1.3	1.6
2003 4/.....	*	*	*	0.4	*	0.7	0.6	0.8	1.4	1.7
2002.....	*	*	*	0.4	*	0.5	0.9	1.0	1.4	2.0
2001.....	*	0.3	*	0.3	*	*	0.9	1.2	1.4	1.9
2000.....	*	*	*	*	*	0.6	0.7	0.9	1.1	1.6
1999.....	*	0.3	*	*	0.5	0.7	0.8	1.1	1.4	1.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Heart failure (I50)

All races 3/, both sexes

2006.....	1.0	2.0	3.7	6.6	12.0	21.9	43.1	88.7	190.0	614.4
2005.....	1.0	1.7	3.5	6.2	11.8	21.5	43.6	88.8	190.5	618.6
2004.....	0.9	1.7	3.3	6.1	11.4	22.4	43.8	88.7	187.7	621.2
2003.....	1.0	1.6	3.3	6.4	12.1	23.6	44.7	89.9	192.7	641.1
2002.....	0.8	1.7	3.4	6.0	12.4	22.9	45.7	89.0	194.0	645.0
2001.....	0.8	1.7	3.3	6.5	13.0	23.8	45.6	92.6	199.2	674.6
2000.....	0.8	1.4	3.0	6.1	12.8	24.0	46.5	93.5	199.9	684.9
1999.....	0.8	1.5	2.9	6.5	11.7	23.5	48.3	94.3	201.6	681.9

All races 3/, male

2006.....	1.2	2.6	4.9	8.2	14.5	26.6	50.0	103.3	213.5	599.0
2005.....	1.3	2.1	4.3	7.6	14.0	24.6	50.1	102.1	213.7	603.1
2004.....	1.0	2.2	4.0	7.5	13.7	26.4	51.1	103.2	211.1	605.5
2003.....	1.3	2.1	3.9	7.9	14.7	28.3	53.1	105.3	213.9	629.8
2002.....	1.0	2.0	4.4	7.2	14.8	27.0	53.2	103.9	218.0	621.4
2001.....	0.9	2.2	4.0	8.1	15.0	28.0	54.2	107.9	226.6	646.4
2000.....	1.1	1.9	3.5	7.4	15.3	28.2	55.6	107.9	229.5	672.2
1999.....	1.0	1.9	3.6	8.0	14.2	28.0	58.9	108.4	224.9	667.0

All races 3/, female

2006.....	0.8	1.5	2.6	5.1	9.7	17.9	37.6	78.0	175.2	621.6
2005.....	0.6	1.4	2.7	4.9	9.8	18.7	38.4	79.1	176.1	625.6
2004.....	0.7	1.3	2.8	4.8	9.4	18.8	37.9	78.3	173.3	628.2
2003.....	0.7	1.2	2.7	4.9	9.7	19.6	37.8	78.9	179.8	646.1
2002.....	0.5	1.3	2.4	4.9	10.1	19.3	39.8	78.4	179.3	655.3
2001.....	0.7	1.2	2.6	5.1	11.3	20.2	38.7	81.9	182.7	686.4
2000.....	0.6	0.9	2.5	5.0	10.5	20.3	39.3	83.5	182.4	690.0
1999.....	0.6	1.1	2.1	5.1	9.5	19.6	39.9	84.5	188.0	687.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Heart failure (I50)

White, both sexes

2006 4/.....	0.8	1.5	2.8	5.6	11.0	20.5	42.2	89.0	193.3	637.2
2005 4/.....	0.7	1.2	2.7	5.3	10.5	19.8	42.3	87.6	192.4	640.6
2004 4/.....	0.7	1.4	2.5	5.1	10.3	21.5	42.9	87.9	191.0	641.5
2003 4/.....	0.7	1.3	2.6	5.6	10.7	22.4	43.2	89.5	195.3	660.4
2002.....	0.6	1.2	2.7	5.3	11.4	21.7	44.1	88.1	195.6	664.5
2001.....	0.6	1.4	2.7	5.7	12.1	22.5	44.3	92.1	201.1	695.1
2000.....	0.6	1.1	2.5	5.4	11.8	22.6	44.9	92.8	201.8	703.7
1999.....	0.6	1.1	2.3	5.6	10.9	22.0	46.8	92.5	202.5	699.2

White, male

2006 4/.....	0.9	1.9	3.7	7.2	13.4	25.3	48.3	104.2	217.8	622.6
2005 4/.....	0.8	1.5	3.4	6.5	12.5	22.7	48.4	100.6	217.9	629.3
2004 4/.....	0.8	1.6	3.1	6.2	12.2	25.2	50.7	101.7	215.6	625.5
2003 4/.....	0.9	1.7	3.0	7.0	13.1	27.0	52.1	105.9	217.1	650.8
2002.....	0.8	1.5	3.6	6.4	13.6	25.5	51.4	103.4	221.4	641.9
2001.....	0.7	1.9	3.4	7.2	14.2	26.5	52.8	107.6	228.2	666.5
2000.....	0.9	1.5	3.0	6.8	14.1	27.2	54.3	107.7	231.7	690.0
1999.....	0.8	1.4	2.9	7.1	13.3	26.2	58.1	107.2	226.3	680.0

White, female

2006 4/.....	0.6	1.0	2.0	4.1	8.8	16.3	37.1	77.6	177.8	644.0
2005 4/.....	0.5	1.0	2.1	4.2	8.6	17.3	37.2	78.0	176.5	645.8
2004 4/.....	0.6	1.1	2.0	4.0	8.6	18.1	36.5	77.8	175.7	648.7
2003 4/.....	0.4	0.9	2.1	4.2	8.5	18.3	35.9	77.6	181.9	664.7
2002.....	0.4	0.9	1.9	4.2	9.3	18.4	38.2	77.1	179.8	674.3
2001.....	0.5	0.8	2.0	4.2	10.1	18.9	37.4	81.0	184.7	707.1
2000.....	0.4	0.6	2.0	4.1	9.6	18.6	37.3	82.4	183.9	709.3
1999.....	0.4	0.8	1.7	4.2	8.8	18.4	37.8	82.1	188.5	707.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Heart failure (I50)

Black, both sexes

2006 4/.....	2.8	6.4	10.7	16.1	24.5	41.1	64.8	109.6	200.0	484.9
2005 4/.....	3.0	5.2	10.0	15.6	26.8	41.3	69.1	123.8	216.0	489.6
2004 4/.....	2.2	5.0	10.5	16.3	23.8	35.7	64.0	120.2	194.6	506.5
2003 4/.....	3.3	4.7	9.4	15.3	27.7	41.1	70.9	114.7	203.6	531.8
2002.....	2.2	5.5	9.3	13.9	24.6	38.6	71.7	118.5	215.3	528.0
2001.....	2.7	4.5	8.7	15.7	24.6	41.2	71.1	121.1	213.7	543.8
2000.....	2.2	4.2	7.6	14.0	24.3	41.2	72.6	119.8	209.5	558.4
1999.....	2.4	4.8	8.4	16.4	22.0	42.0	74.0	130.5	216.7	552.9

Black, male

2006 4/.....	3.7	8.0	14.9	18.4	31.3	49.0	82.1	126.2	223.8	463.8
2005 4/.....	4.6	6.4	12.6	20.1	32.7	48.5	82.5	152.0	224.0	423.0
2004 4/.....	2.6	6.8	12.2	21.0	30.6	45.5	69.3	148.3	220.2	503.7
2003 4/.....	3.9	5.5	11.6	19.4	35.9	51.0	79.5	124.6	227.4	513.0
2002.....	2.7	6.6	12.0	16.5	31.3	47.6	87.5	136.9	236.9	479.2
2001.....	2.6	5.0	10.4	18.9	26.4	49.6	86.2	140.8	256.8	541.2
2000.....	2.5	5.5	9.1	14.8	29.6	45.5	82.9	134.7	242.8	581.2
1999.....	2.7	6.4	11.0	19.3	27.4	51.4	83.1	146.7	238.9	603.5

Black, female

2006 4/.....	2.0	5.0	7.2	14.3	19.1	35.3	52.8	99.6	187.4	493.5
2005 4/.....	1.6	4.0	7.7	11.9	22.1	36.0	60.0	106.9	211.8	516.5
2004 4/.....	1.9	3.4	8.9	12.4	18.5	28.4	60.4	103.5	181.1	507.6
2003 4/.....	2.8	4.0	7.6	11.9	21.4	33.8	65.1	108.8	191.1	539.2
2002.....	1.7	4.5	7.0	11.7	19.4	31.8	61.2	107.6	204.1	546.7
2001.....	2.8	4.1	7.2	13.1	23.2	35.1	61.0	109.3	191.7	544.8
2000.....	1.9	3.0	6.4	13.4	20.1	38.0	65.7	110.8	192.8	549.9
1999.....	2.1	3.5	6.3	14.0	17.8	35.0	68.0	120.7	205.7	534.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All races 3/, both sexes

2006.....	10.0	15.0	20.5	28.7	42.2	62.4	99.2	167.0	299.4	724.4
2005.....	10.3	15.5	21.2	29.7	44.5	65.0	104.9	172.3	306.2	754.0
2004.....	10.2	15.5	21.4	28.8	45.0	66.3	105.9	176.1	311.3	747.7
2003.....	11.0	15.5	21.3	31.7	46.3	71.8	113.2	186.0	328.5	781.5
2002.....	11.2	15.7	21.7	32.5	46.9	73.3	116.3	194.4	338.2	793.7
2001.....	10.5	15.4	21.9	32.3	47.8	75.3	116.6	191.2	333.6	796.5
2000.....	10.9	15.5	22.1	33.1	50.4	78.0	118.4	193.4	338.9	820.0
1999.....	11.1	15.7	22.3	34.1	53.5	79.9	124.4	197.7	347.6	826.1

All races 3/, male

2006.....	12.3	19.1	26.5	37.0	53.5	78.5	120.7	196.1	347.5	735.0
2005.....	12.9	19.8	27.9	38.7	55.8	81.5	128.0	203.6	349.3	767.7
2004.....	12.9	20.2	27.8	37.9	57.6	82.7	128.0	209.5	357.0	765.2
2003.....	13.8	19.7	27.5	40.8	58.6	88.8	139.6	224.8	380.7	795.9
2002.....	14.2	20.6	28.1	41.1	59.4	89.7	143.8	233.1	389.4	821.3
2001.....	13.4	19.9	27.7	40.3	61.1	91.7	143.7	232.4	390.0	821.3
2000.....	13.6	20.2	28.9	43.4	64.2	96.8	147.6	232.6	391.5	870.7
1999.....	13.9	20.2	28.7	44.5	68.7	103.5	155.5	238.5	411.1	874.7

All races 3/, female

2006.....	7.8	11.1	14.8	20.9	31.9	48.3	81.7	145.7	269.3	719.5
2005.....	7.6	11.4	14.8	21.3	34.1	50.5	86.2	149.6	279.5	747.6
2004.....	7.6	11.0	15.3	20.3	33.4	52.0	88.0	152.2	283.2	739.9
2003.....	8.4	11.5	15.4	23.1	35.1	57.0	92.0	158.4	296.6	775.1
2002.....	8.2	11.0	15.6	24.4	35.6	59.2	94.3	167.1	307.1	781.7
2001.....	7.6	11.1	16.3	24.8	35.7	61.2	95.0	162.4	299.7	786.1
2000.....	8.2	11.0	15.6	23.4	37.8	61.9	95.4	166.0	307.9	799.3
1999.....	8.3	11.2	16.2	24.3	39.8	59.7	100.1	169.4	310.5	806.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

White, both sexes

2006 4/.....	8.7	13.2	17.9	25.5	38.6	58.0	94.8	164.8	299.3	745.3
2005 4/.....	8.9	13.4	18.2	26.4	40.1	60.3	100.7	169.7	308.2	773.4
2004 4/.....	8.6	13.2	18.6	25.4	41.3	61.7	101.4	172.0	312.3	762.4
2003 4/.....	9.4	13.4	18.2	28.4	42.0	66.5	107.8	183.7	329.3	796.1
2002.....	9.7	13.3	18.8	28.9	42.7	69.1	111.9	191.2	339.3	806.1
2001.....	8.7	12.9	18.9	28.3	44.0	70.3	112.3	186.3	333.4	809.9
2000.....	9.1	13.0	18.9	29.6	46.2	73.1	113.5	188.4	339.9	833.4
1999.....	9.2	12.9	19.4	29.8	48.2	74.9	118.6	192.7	346.9	839.0

White, male

2006 4/.....	10.7	16.7	23.2	33.0	48.7	73.0	115.4	192.9	349.4	754.3
2005 4/.....	11.5	17.3	24.4	34.5	51.0	75.8	122.3	201.1	352.7	787.7
2004 4/.....	11.2	17.4	24.7	33.7	53.3	77.5	123.2	204.5	356.9	775.3
2003 4/.....	12.0	17.3	23.6	36.5	53.7	82.6	133.8	222.2	381.7	810.7
2002.....	12.6	17.7	24.8	36.9	54.7	85.0	138.8	229.4	391.0	834.2
2001.....	11.5	17.0	24.7	35.4	56.5	86.9	138.2	226.4	388.3	831.0
2000.....	11.5	17.5	24.8	38.9	59.5	90.9	142.2	227.3	393.2	884.2
1999.....	11.9	17.3	25.1	39.0	62.4	97.5	148.4	232.5	409.5	883.9

White, female

2006 4/.....	6.7	9.6	12.7	18.4	29.2	44.6	77.7	143.8	267.7	741.1
2005 4/.....	6.3	9.4	12.0	18.6	30.0	46.6	82.9	146.5	280.4	766.9
2004 4/.....	6.0	9.0	12.7	17.5	30.3	47.7	83.5	148.2	284.6	756.5
2003 4/.....	6.8	9.5	12.9	20.5	31.1	52.3	86.5	155.9	297.0	789.7
2002.....	6.7	9.0	13.0	21.4	31.5	55.1	90.0	163.8	307.6	793.9
2001.....	5.8	8.8	13.4	21.5	32.5	55.8	91.3	158.0	300.1	801.1
2000.....	6.7	8.6	13.2	20.7	33.9	57.6	90.4	161.1	308.1	812.7
1999.....	6.5	8.6	13.7	21.0	35.2	55.1	94.9	164.7	310.0	821.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Black, both sexes

2006 4/.....	20.7	30.8	44.8	61.5	84.7	112.8	161.6	227.2	363.5	605.0
2005 4/.....	21.7	33.7	48.4	63.6	91.0	118.1	165.9	239.4	341.0	640.9
2004 4/.....	23.3	35.0	47.6	64.9	88.5	121.2	173.4	255.4	352.5	668.3
2003 4/.....	24.5	34.4	49.8	67.9	94.2	132.5	190.7	246.1	366.8	703.7
2002.....	23.6	36.5	49.5	70.6	94.5	127.7	180.3	263.4	376.5	738.5
2001.....	25.1	36.9	50.0	73.6	91.5	135.2	179.9	276.1	383.5	730.2
2000.....	25.2	37.3	52.5	69.7	96.5	137.5	191.0	282.9	377.8	742.0
1999.....	26.3	38.5	51.8	77.4	109.1	140.1	206.0	283.3	400.8	746.1

Black, male

2006 4/.....	25.7	40.2	58.8	80.9	113.3	146.7	204.2	278.8	416.2	637.2
2005 4/.....	26.2	42.0	62.5	84.2	114.3	153.5	215.1	288.8	384.1	661.9
2004 4/.....	27.9	45.0	59.8	86.2	116.2	152.2	212.4	315.7	428.3	715.3
2003 4/.....	29.7	43.0	65.2	90.9	120.3	170.1	234.2	304.9	436.3	707.9
2002.....	28.2	47.7	62.8	90.7	117.2	157.1	226.3	314.2	434.1	753.7
2001.....	30.0	45.9	60.5	95.7	121.2	158.8	227.2	347.1	479.1	805.6
2000.....	30.9	46.3	71.0	95.0	122.0	174.3	233.4	340.7	449.3	816.5
1999.....	31.8	47.7	67.7	103.6	142.8	183.3	265.3	341.0	503.1	849.5

Black, female

2006 4/.....	16.3	22.7	33.0	45.5	62.4	87.8	132.3	196.3	335.7	591.9
2005 4/.....	17.6	26.4	36.4	46.5	72.8	91.8	132.2	209.9	318.3	632.4
2004 4/.....	19.2	26.2	37.2	47.5	66.9	98.2	147.0	219.7	312.6	649.7
2003 4/.....	19.9	26.8	36.8	48.9	73.9	104.5	161.3	211.1	330.4	702.0
2002.....	19.5	26.8	38.3	54.1	76.8	105.8	149.4	233.0	346.5	732.8
2001.....	20.8	29.1	41.0	55.4	68.2	117.7	148.4	233.8	334.8	702.2
2000.....	20.3	29.5	36.8	49.0	76.5	110.4	162.9	248.1	341.8	714.3
1999.....	21.5	30.6	38.3	56.1	82.7	108.1	166.8	248.6	350.0	707.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Essential hypertension and hypertensive renal disease (I10,I12,I15)

All races 3/, both sexes

2006.....	1.2	2.1	4.0	5.5	8.8	13.3	21.0	36.6	69.7	189.4
2005.....	1.3	2.2	3.3	5.0	8.3	13.9	22.2	40.5	75.4	210.0
2004.....	1.0	2.2	3.4	4.9	8.1	13.0	21.9	38.5	71.3	198.5
2003.....	1.1	1.8	3.3	5.0	8.0	13.1	21.3	38.4	69.9	188.9
2002.....	1.0	1.8	2.9	4.5	7.3	11.8	20.7	34.9	66.7	180.4
2001.....	1.0	1.9	3.0	4.4	7.5	12.2	19.0	35.5	65.2	171.9
2000.....	1.1	1.8	3.0	4.4	7.8	11.5	19.1	33.8	63.0	162.9
1999.....	0.9	1.7	2.8	4.1	7.3	12.0	18.6	32.7	59.9	152.1

All races 3/, male

2006.....	1.5	2.6	5.1	6.7	10.6	15.1	22.8	36.6	72.2	159.2
2005.....	1.5	2.8	4.4	6.4	9.9	15.9	23.3	41.9	74.5	169.3
2004.....	1.2	2.6	4.5	5.6	9.8	14.9	22.2	40.0	72.7	166.0
2003.....	1.3	2.5	4.2	6.0	9.2	14.9	22.5	37.5	66.1	153.5
2002.....	1.3	2.3	3.5	5.5	8.2	12.7	22.2	35.9	64.8	153.4
2001.....	1.3	2.3	3.7	5.2	8.1	13.7	20.0	36.0	61.3	150.8
2000.....	1.4	2.3	3.8	5.4	9.1	11.5	20.6	34.5	62.9	142.4
1999.....	1.1	2.0	3.5	5.0	8.2	13.4	20.1	33.1	59.4	132.4

All races 3/, female

2006.....	0.9	1.6	2.8	4.4	7.1	11.6	19.6	36.7	68.1	203.5
2005.....	1.0	1.7	2.3	3.8	6.8	12.2	21.4	39.4	76.0	228.7
2004.....	0.7	1.7	2.3	4.3	6.6	11.3	21.6	37.4	70.4	213.2
2003.....	0.9	1.2	2.4	4.0	6.9	11.4	20.4	39.1	72.2	204.6
2002.....	0.7	1.3	2.3	3.7	6.5	11.0	19.5	34.2	67.9	192.1
2001.....	0.7	1.4	2.3	3.8	6.9	11.0	18.2	35.2	67.5	180.8
2000.....	0.8	1.3	2.2	3.5	6.7	11.5	17.9	33.4	63.0	171.3
1999.....	0.7	1.3	2.1	3.3	6.5	10.8	17.5	32.4	60.2	160.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Essential hypertension and hypertensive renal disease (I10,I12,I15)

White, both sexes

2006 4/.....	0.6	1.4	2.6	3.9	6.6	10.1	17.4	31.6	63.4	179.5
2005 4/.....	0.8	1.4	2.1	3.5	6.2	10.6	17.6	35.0	69.5	200.3
2004 4/.....	0.6	1.3	2.2	3.2	5.8	9.8	17.6	32.9	64.9	188.1
2003 4/.....	0.7	1.2	2.3	3.3	5.8	9.8	16.6	33.0	64.1	181.0
2002.....	0.6	1.0	1.9	3.0	5.4	9.0	16.8	29.8	60.4	170.4
2001.....	0.5	1.1	1.8	2.9	5.3	9.2	15.3	30.9	58.6	163.3
2000.....	0.6	1.0	1.8	3.0	5.5	8.5	15.3	28.9	57.4	153.2
1999.....	0.5	1.0	1.6	2.7	5.0	9.1	15.2	27.6	53.3	142.0

White, male

2006 4/.....	0.8	1.8	3.4	4.8	8.5	11.7	18.2	32.0	65.1	151.3
2005 4/.....	1.0	1.8	2.8	4.5	7.5	12.2	18.1	35.3	68.9	160.2
2004 4/.....	0.8	1.7	3.0	3.8	7.3	11.5	18.2	33.7	66.6	159.4
2003 4/.....	0.8	1.6	2.9	4.1	6.6	11.6	18.0	32.7	60.3	145.3
2002.....	0.8	1.4	2.4	3.8	6.3	9.8	18.3	30.8	57.7	145.2
2001.....	0.7	1.4	2.2	3.5	5.9	10.3	16.4	31.4	54.5	142.2
2000.....	0.9	1.4	2.5	3.8	6.6	8.8	16.8	29.9	57.5	132.5
1999.....	0.7	1.3	2.1	3.5	5.8	10.4	16.6	28.1	52.5	121.0

White, female

2006 4/.....	0.4	1.0	1.7	3.1	4.9	8.7	16.7	31.4	62.4	192.7
2005 4/.....	0.5	0.9	1.4	2.6	5.1	9.2	17.3	34.7	69.9	218.8
2004 4/.....	0.4	0.9	1.4	2.7	4.4	8.3	17.1	32.3	63.9	201.1
2003 4/.....	0.5	0.7	1.6	2.4	5.0	8.2	15.5	33.2	66.5	196.8
2002.....	0.4	0.7	1.4	2.3	4.5	8.3	15.6	29.0	62.0	181.3
2001.....	0.3	0.7	1.4	2.4	4.8	8.2	14.5	30.5	61.1	172.2
2000.....	0.4	0.6	1.1	2.2	4.5	8.3	14.1	28.2	57.4	161.6
1999.....	0.3	0.8	1.2	1.9	4.3	7.9	14.0	27.2	53.8	150.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Black, both sexes

2006 4/.....	5.1	7.7	14.9	19.1	28.8	42.9	57.1	93.7	150.0	318.3
2005 4/.....	4.6	8.5	12.8	18.4	28.6	44.6	66.4	100.7	153.2	341.0
2004 4/.....	3.6	8.3	12.4	19.7	29.6	41.8	64.6	103.9	150.0	335.6
2003 4/.....	3.9	7.1	12.0	20.5	29.0	42.9	68.9	102.1	145.4	301.1
2002.....	3.8	7.4	10.9	18.4	24.3	36.5	61.5	96.4	151.0	308.8
2001.....	4.4	7.8	12.5	18.0	27.0	40.4	56.3	89.8	148.2	288.0
2000.....	4.2	7.2	12.7	17.9	29.0	40.3	58.8	92.4	130.4	283.4
1999.....	4.0	6.3	12.8	17.0	27.0	38.0	57.2	95.0	138.7	269.4

Black, male

2006 4/.....	6.4	9.1	19.5	24.8	33.2	50.1	72.6	97.1	171.1	271.7
2005 4/.....	5.2	9.8	17.4	24.4	36.5	52.8	79.1	123.8	154.5	289.1
2004 4/.....	4.2	9.8	16.6	22.7	36.3	49.9	66.0	123.7	157.8	263.3
2003 4/.....	4.5	9.8	16.4	24.3	36.2	50.0	74.6	102.2	147.7	274.2
2002.....	4.7	9.2	13.3	21.7	25.7	40.6	67.1	105.2	164.3	274.5
2001.....	5.5	10.1	16.3	21.7	30.3	48.8	61.9	95.2	158.1	275.7
2000.....	5.3	9.0	15.6	22.6	34.4	40.1	63.9	93.1	137.4	269.9
1999.....	4.6	7.8	16.6	19.5	31.0	43.1	62.1	98.8	155.2	266.0

Black, female

2006 4/.....	3.9	6.4	11.1	14.4	25.3	37.6	46.5	91.6	138.9	337.4
2005 4/.....	4.0	7.4	9.0	13.4	22.4	38.6	57.8	86.9	152.5	362.0
2004 4/.....	3.0	7.1	8.8	17.3	24.5	35.8	63.7	92.2	145.9	364.2
2003 4/.....	3.3	4.7	8.4	17.4	23.3	37.6	65.1	102.0	144.3	311.5
2002.....	3.0	5.9	8.8	15.6	23.2	33.5	57.8	91.1	144.0	322.0
2001.....	3.4	5.8	9.3	15.0	24.5	34.2	52.5	86.5	143.1	292.6
2000.....	3.2	5.7	10.3	14.2	24.7	40.5	55.5	92.0	126.9	288.5
1999.....	3.5	5.0	9.5	15.0	24.0	34.2	54.0	92.7	130.5	270.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cerebrovascular diseases (I60-I69)

All races 3/, both sexes

2006.....	6.7	11.2	18.5	26.4	42.7	70.7	127.5	240.3	458.6	1,039.6
2005.....	6.9	11.7	18.8	25.8	42.7	73.8	133.6	253.6	497.3	1,141.8
2004.....	7.1	11.9	18.3	26.5	44.5	78.4	142.2	276.7	532.1	1,245.9
2003.....	7.4	11.8	18.6	27.8	45.9	79.7	150.7	294.1	571.1	1,370.1
2002.....	7.1	12.3	18.3	28.6	48.3	83.8	160.7	308.8	601.7	1,445.9
2001.....	7.2	12.1	18.5	28.8	49.6	85.7	164.3	320.2	622.4	1,500.2
2000.....	7.7	13.1	19.3	30.8	53.8	91.0	169.1	332.8	654.0	1,589.2
1999.....	7.5	12.5	18.4	31.7	51.4	91.1	173.5	340.6	664.0	1,614.8

All races 3/, male

2006.....	6.9	12.0	21.2	31.2	49.2	81.9	140.9	259.6	468.5	932.4
2005.....	7.0	12.1	21.5	30.1	50.0	84.6	149.5	275.8	513.0	1,023.3
2004.....	7.4	12.8	21.1	29.9	52.2	90.2	158.8	301.2	551.6	1,118.1
2003.....	7.6	12.9	21.1	31.3	53.5	91.4	170.8	322.5	595.6	1,236.0
2002.....	6.9	13.4	20.4	32.7	55.8	97.0	178.7	334.8	615.6	1,317.9
2001.....	7.4	13.2	20.5	32.8	57.2	98.5	187.9	349.0	654.6	1,380.2
2000.....	7.8	14.2	21.3	35.2	62.3	102.6	192.9	367.6	695.1	1,484.3
1999.....	7.8	13.3	20.7	35.7	60.0	102.8	197.7	376.9	706.8	1,512.8

All races 3/, female

2006.....	6.5	10.4	15.8	21.9	36.8	60.9	116.5	226.2	452.3	1,089.8
2005.....	6.7	11.3	16.2	21.8	36.0	64.3	120.7	237.6	487.5	1,196.1
2004.....	6.8	11.0	15.5	23.2	37.5	68.1	128.8	259.1	520.2	1,303.4
2003.....	7.2	10.8	16.3	24.6	38.9	69.5	134.5	273.9	556.1	1,429.4
2002.....	7.2	11.2	16.3	24.8	41.4	72.4	146.2	290.4	593.3	1,501.5
2001.....	7.1	10.9	16.5	25.1	42.7	74.7	145.6	300.0	603.2	1,550.4
2000.....	7.5	12.0	17.3	26.6	46.0	81.0	150.3	308.5	629.7	1,632.0
1999.....	7.1	11.7	16.1	28.0	43.7	81.1	154.5	315.3	639.0	1,656.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cerebrovascular diseases (I60-I69)

White, both sexes

2006 4/.....	5.3	8.9	14.5	21.3	36.3	62.4	117.8	231.6	453.0	1,050.8
2005 4/.....	5.3	9.1	14.5	21.1	36.2	65.6	123.8	244.2	491.2	1,153.3
2004 4/.....	5.7	9.1	14.1	21.8	37.5	70.0	132.6	264.8	526.2	1,257.8
2003 4/.....	5.6	9.0	14.2	22.6	38.3	70.8	140.3	283.2	565.9	1,382.2
2002.....	5.5	9.5	14.0	23.8	41.2	74.2	150.1	298.6	595.7	1,459.5
2001.....	5.6	9.3	13.8	23.9	41.7	75.2	153.2	309.7	617.1	1,511.2
2000.....	5.9	9.7	15.4	25.4	46.5	81.9	159.0	321.3	647.0	1,601.6
1999.....	5.7	9.4	14.4	26.2	43.5	82.9	161.3	329.3	658.4	1,631.6

White, male

2006 4/.....	5.6	9.6	16.4	24.5	41.0	71.7	128.9	249.8	464.4	941.3
2005 4/.....	5.5	9.5	16.5	24.2	41.8	74.3	138.0	266.4	505.3	1,033.7
2004 4/.....	6.1	9.7	16.3	24.1	43.3	80.6	147.0	288.0	546.9	1,129.3
2003 4/.....	5.9	10.1	16.0	24.7	44.5	81.8	158.5	310.0	590.4	1,247.0
2002.....	5.5	10.6	15.5	26.9	47.2	85.1	167.3	323.4	610.1	1,327.4
2001.....	5.9	10.2	15.4	26.9	48.1	86.7	174.8	337.7	648.8	1,386.2
2000.....	6.1	10.8	16.9	28.6	53.8	91.7	180.5	354.6	686.1	1,490.7
1999.....	6.1	9.8	16.2	29.5	50.6	93.2	183.6	363.2	700.5	1,522.7

White, female

2006 4/.....	4.9	8.2	12.7	18.2	31.8	54.1	108.6	218.0	445.7	1,102.2
2005 4/.....	5.2	8.6	12.5	18.2	31.1	57.8	112.1	227.7	482.4	1,208.5
2004 4/.....	5.2	8.5	11.9	19.5	32.2	60.5	120.7	247.9	513.3	1,315.7
2003 4/.....	5.3	7.9	12.3	20.5	32.5	61.1	125.4	263.8	550.7	1,442.1
2002.....	5.5	8.4	12.6	20.9	35.6	64.5	136.0	280.8	586.8	1,516.9
2001.....	5.4	8.3	12.2	21.1	35.8	65.1	135.7	289.8	598.0	1,563.5
2000.....	5.7	8.7	13.9	22.4	39.8	73.3	141.9	297.9	623.8	1,646.7
1999.....	5.3	8.9	12.7	23.0	37.0	73.9	143.7	305.5	633.6	1,675.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cerebrovascular diseases (I60-I69)

Black, both sexes

2006 4/.....	15.9	27.2	48.6	67.5	101.4	143.3	221.7	349.3	557.8	1,008.7
2005 4/.....	17.1	30.5	50.3	65.4	99.5	145.1	231.4	372.3	611.0	1,106.7
2004 4/.....	17.2	31.5	48.0	66.3	108.1	154.9	246.5	418.6	647.9	1,211.2
2003 4/.....	18.6	31.4	52.7	72.4	113.1	157.4	256.3	433.2	677.0	1,348.7
2002.....	17.4	32.1	51.2	71.1	109.9	169.6	272.5	447.2	712.5	1,383.9
2001.....	18.4	32.6	55.0	70.6	117.1	178.3	277.6	460.2	716.4	1,452.7
2000.....	19.7	37.2	51.2	77.4	114.0	174.8	281.1	475.2	769.4	1,530.0
1999.....	19.9	34.4	50.6	78.4	118.2	170.0	307.8	487.9	756.4	1,492.5

Black, male

2006 4/.....	15.6	29.0	61.1	88.9	132.2	179.3	270.4	400.0	579.8	882.0
2005 4/.....	17.4	32.0	60.5	83.6	133.4	184.9	276.7	407.2	655.2	983.5
2004 4/.....	16.9	35.4	56.4	84.7	139.4	188.3	297.8	470.9	677.2	1,061.0
2003 4/.....	18.8	34.2	62.6	90.6	141.7	184.9	307.4	503.0	727.9	1,180.3
2002.....	16.9	34.3	61.6	88.9	138.8	213.7	328.8	502.5	729.8	1,252.2
2001.....	17.8	36.5	63.9	87.6	143.4	218.5	336.3	524.6	768.1	1,349.1
2000.....	20.5	41.0	60.4	97.8	137.5	212.8	340.0	544.8	863.2	1,458.8
1999.....	20.8	38.8	61.3	92.7	149.4	204.3	366.7	559.0	818.8	1,448.3

Black, female

2006 4/.....	16.2	25.6	38.1	49.9	77.4	116.7	188.3	319.0	546.2	1,060.5
2005 4/.....	16.8	29.3	41.8	50.3	73.0	115.6	200.4	351.5	587.6	1,156.5
2004 4/.....	17.5	28.1	40.8	51.2	83.6	130.1	211.8	387.6	632.4	1,270.7
2003 4/.....	18.5	29.0	44.4	57.3	90.7	137.1	221.8	391.8	650.3	1,414.2
2002.....	17.9	30.1	42.4	56.5	87.3	136.9	234.8	414.1	703.4	1,434.3
2001.....	19.0	29.1	47.4	56.7	96.5	148.5	238.6	421.9	690.1	1,491.2
2000.....	19.0	33.9	43.4	60.6	95.4	146.7	242.0	433.2	722.2	1,556.5
1999.....	19.0	30.5	41.6	66.6	93.7	144.5	268.7	445.1	725.6	1,509.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerosis (I70)

All races 3/, both sexes

2006.....	0.1	0.2	0.4	0.9	1.7	3.2	6.1	12.6	27.2	89.9
2005.....	0.1	0.3	0.7	1.0	2.1	3.9	7.2	16.9	37.8	132.9
2004.....	0.1	0.3	0.5	1.0	2.3	4.1	8.5	17.5	38.0	137.0
2003.....	0.2	0.3	0.6	1.1	2.4	4.5	9.0	19.9	43.5	154.6
2002.....	0.1	0.3	0.6	1.4	2.4	5.2	9.3	20.5	47.8	168.1
2001.....	0.1	0.3	0.7	1.4	2.5	4.9	9.8	21.9	49.2	179.1
2000.....	0.1	0.4	0.6	1.3	2.6	5.2	10.5	22.7	52.4	188.8
1999.....	0.1	0.3	0.8	1.3	2.9	5.1	11.2	24.1	53.6	202.0

All races 3/, male

2006.....	*	0.3	0.6	1.2	2.2	3.8	7.3	14.2	29.5	85.3
2005.....	*	0.4	1.0	1.2	2.7	4.8	8.7	20.0	42.0	121.7
2004.....	*	0.4	0.7	1.2	3.0	4.9	10.3	20.2	43.5	128.2
2003.....	0.2	0.5	0.8	1.5	3.2	5.4	10.6	23.9	48.3	138.6
2002.....	0.2	0.4	0.8	1.7	3.2	6.4	11.5	24.4	52.5	156.6
2001.....	*	0.4	0.8	1.7	3.2	6.2	12.1	25.0	53.2	167.2
2000.....	0.2	0.5	0.8	1.7	3.6	6.6	13.1	27.5	59.8	176.5
1999.....	0.2	0.4	1.0	1.8	3.8	6.3	13.3	28.1	60.0	189.0

All races 3/, female

2006.....	*	*	0.2	0.5	1.3	2.6	5.1	11.4	25.7	92.1
2005.....	*	0.2	0.4	0.8	1.5	3.1	5.9	14.7	35.2	138.1
2004.....	*	*	0.3	0.8	1.7	3.4	7.1	15.5	34.6	141.0
2003.....	*	*	0.4	0.7	1.7	3.7	7.8	17.0	40.6	161.7
2002.....	*	*	0.5	1.0	1.7	4.1	7.5	17.8	45.0	173.1
2001.....	*	0.2	0.5	1.1	1.8	3.9	8.1	19.7	46.8	184.1
2000.....	*	0.2	0.4	0.9	1.7	4.0	8.4	19.4	48.1	193.8
1999.....	*	*	0.5	0.9	2.2	4.0	9.5	21.4	49.8	207.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerosis (I70)

White, both sexes

2006 4/.....	*	0.2	0.4	0.8	1.7	3.1	6.0	12.8	27.4	93.2
2005 4/.....	0.1	0.3	0.6	0.9	2.0	3.8	7.1	17.2	38.8	138.7
2004 4/.....	0.1	0.3	0.4	1.0	2.3	4.0	8.5	17.8	39.3	142.0
2003 4/.....	0.2	0.3	0.6	1.0	2.4	4.4	8.9	20.2	44.1	159.5
2002.....	0.1	0.2	0.5	1.3	2.3	5.3	9.1	20.4	48.4	174.7
2001.....	0.1	0.3	0.6	1.4	2.3	4.9	9.8	22.2	50.3	185.9
2000.....	0.1	0.3	0.5	1.2	2.4	5.2	10.4	23.1	53.0	195.2
1999.....	*	0.3	0.7	1.2	2.9	5.0	11.3	24.2	54.7	207.7

White, male

2006 4/.....	*	0.3	0.6	1.1	2.2	3.6	7.1	14.3	29.3	87.9
2005 4/.....	*	0.4	0.9	1.2	2.6	4.8	8.5	20.6	42.9	128.3
2004 4/.....	*	0.4	0.7	1.2	2.9	4.9	10.4	20.5	44.8	134.4
2003 4/.....	0.2	0.5	0.8	1.4	3.1	5.2	10.1	23.8	48.4	143.3
2002.....	*	0.4	0.6	1.7	3.0	6.5	11.3	24.0	53.1	163.2
2001.....	*	0.4	0.8	1.7	3.0	6.2	12.1	25.2	54.0	173.9
2000.....	*	0.4	0.7	1.7	3.5	6.5	13.2	28.2	60.3	181.4
1999.....	*	0.4	1.0	1.6	3.7	6.2	13.3	27.7	60.8	193.7

White, female

2006 4/.....	*	*	*	0.6	1.2	2.6	5.1	11.6	26.2	95.7
2005 4/.....	*	*	0.3	0.7	1.5	3.0	5.9	14.7	36.2	143.5
2004 4/.....	*	*	0.2	0.8	1.7	3.2	7.0	15.9	35.9	145.4
2003 4/.....	*	*	0.4	0.7	1.7	3.6	7.8	17.6	41.4	166.7
2002.....	*	*	0.4	1.0	1.6	4.2	7.3	17.7	45.5	179.7
2001.....	*	*	0.5	1.0	1.6	3.9	8.0	20.2	48.0	190.9
2000.....	*	*	0.3	0.8	1.5	4.1	8.2	19.5	48.7	200.8
1999.....	*	*	0.5	0.9	2.0	3.9	9.7	21.7	51.0	213.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Atherosclerosis (I70)

Black, both sexes

2006 4/.....	*	*	*	1.5	3.1	4.5	8.8	14.2	29.3	71.6
2005 4/.....	*	*	1.3	1.3	3.1	5.8	9.5	16.5	33.7	97.8
2004 4/.....	*	*	*	1.7	3.2	5.4	9.9	18.4	31.9	102.9
2003 4/.....	*	*	1.3	1.9	3.5	6.5	12.7	20.8	45.4	121.1
2002.....	*	*	1.5	1.9	4.2	5.2	11.5	26.8	51.3	117.0
2001.....	*	*	1.0	2.0	5.0	6.3	12.1	22.9	42.8	125.6
2000.....	*	*	1.5	2.1	4.7	5.6	12.9	23.5	53.9	138.3
1999.....	*	*	1.3	2.1	4.7	7.0	11.9	27.5	46.8	154.3

Black, male

2006 4/.....	*	*	*	2.5	3.7	6.6	11.4	16.6	39.0	75.1
2005 4/.....	*	*	*	*	4.4	7.6	12.3	17.0	40.0	75.3
2004 4/.....	*	*	*	*	4.9	6.1	12.0	23.2	40.9	78.5
2003 4/.....	*	*	*	*	4.8	8.1	18.1	28.8	59.1	104.3
2002.....	*	*	2.3	*	6.3	6.2	15.4	34.9	59.6	110.4
2001.....	*	*	*	*	5.9	7.6	15.1	30.2	49.0	113.5
2000.....	*	*	*	*	6.0	8.6	14.0	25.5	66.6	147.6
1999.....	*	*	*	3.4	5.7	8.8	15.6	37.0	56.7	164.0

Black, female

2006 4/.....	*	*	*	*	*	*	7.0	12.9	24.2	70.2
2005 4/.....	*	*	*	*	*	4.4	7.6	16.3	30.4	106.9
2004 4/.....	*	*	*	*	*	4.9	8.6	15.6	27.2	112.6
2003 4/.....	*	*	*	*	*	5.3	9.1	16.0	38.3	127.6
2002.....	*	*	*	*	*	4.4	9.0	22.0	46.9	119.5
2001.....	*	*	*	*	4.3	5.3	10.1	18.6	39.7	130.1
2000.....	*	*	*	*	3.6	*	12.2	22.2	47.6	134.8
1999.....	*	*	*	*	3.8	5.6	9.5	21.7	41.9	150.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of circulatory system (I71-I78)

All races 3/, both sexes

2006.....	1.2	2.2	3.2	5.3	9.0	16.4	28.2	47.1	72.2	128.5
2005.....	1.3	2.0	3.2	5.4	10.2	18.0	31.2	50.9	77.9	135.5
2004.....	1.2	2.1	3.4	5.2	9.9	18.3	33.0	51.7	81.1	139.3
2003.....	1.2	2.4	3.5	5.7	11.3	20.5	34.2	56.6	82.7	146.0
2002.....	1.3	2.2	3.1	6.2	11.7	22.3	36.5	57.8	83.8	148.4
2001.....	1.3	2.0	3.3	6.0	12.1	22.3	36.6	60.6	88.1	158.2
2000.....	1.3	2.1	3.5	6.4	12.7	24.5	40.6	62.9	92.2	163.2
1999.....	1.4	2.1	3.3	6.7	12.5	23.5	40.1	63.5	94.3	165.2

All races 3/, male

2006.....	1.8	3.1	4.5	7.6	12.4	22.7	37.0	61.9	92.5	155.0
2005.....	1.8	2.9	4.4	7.8	14.2	24.8	42.7	68.4	101.2	158.2
2004.....	1.7	3.0	4.8	7.2	13.6	24.7	44.1	67.1	105.8	166.5
2003.....	1.7	3.2	5.1	7.9	15.9	29.4	46.1	78.9	111.3	179.7
2002.....	1.8	3.2	4.4	8.6	16.3	31.7	49.2	78.3	113.7	187.4
2001.....	1.8	2.8	4.4	8.2	16.8	30.8	51.2	81.0	122.5	197.3
2000.....	1.7	2.8	4.9	9.3	18.0	34.6	57.6	88.6	126.6	214.9
1999.....	1.6	3.0	4.4	9.4	18.2	33.5	56.7	89.1	129.3	207.5

All races 3/, female

2006.....	0.7	1.4	1.9	3.0	5.8	10.8	21.0	36.3	59.5	116.1
2005.....	0.9	1.1	2.0	3.2	6.6	12.1	21.9	38.2	63.4	125.0
2004.....	0.8	1.2	2.1	3.3	6.6	12.8	24.0	40.6	65.9	127.0
2003.....	0.8	1.6	2.1	3.7	7.2	12.8	24.6	40.7	65.2	131.1
2002.....	0.7	1.4	2.0	3.9	7.5	14.2	26.4	43.4	65.6	131.5
2001.....	0.9	1.2	2.2	3.9	7.8	15.0	25.1	46.3	67.4	141.8
2000.....	0.9	1.5	2.1	3.7	7.9	15.7	27.2	45.0	71.9	142.2
1999.....	1.1	1.3	2.1	4.1	7.3	15.0	27.2	45.7	73.7	148.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of circulatory system (I71-I78)

White, both sexes

2006 4/.....	1.0	2.0	2.7	4.8	8.7	16.0	28.6	47.5	72.0	129.7
2005 4/.....	1.2	1.7	2.8	5.0	10.0	17.7	31.6	51.1	77.9	136.1
2004 4/.....	1.1	1.8	3.0	4.8	9.3	18.1	33.8	52.1	81.2	139.3
2003 4/.....	1.1	2.0	3.1	5.5	11.0	20.6	34.7	57.5	82.9	146.9
2002.....	1.1	1.9	2.7	5.8	11.4	22.4	37.0	58.7	84.8	148.9
2001.....	1.1	1.8	2.9	5.6	11.7	22.6	36.8	60.6	88.4	158.2
2000.....	1.0	1.8	3.0	6.0	12.3	24.5	41.4	63.0	93.0	163.0
1999.....	1.2	1.9	3.0	6.3	12.6	23.7	40.8	64.1	94.8	165.1

White, male

2006 4/.....	1.6	2.8	3.9	7.0	12.2	22.3	37.9	63.3	94.0	159.4
2005 4/.....	1.5	2.4	3.8	7.2	14.1	24.3	43.9	68.5	102.1	162.0
2004 4/.....	1.5	2.6	4.3	6.6	12.8	24.5	45.7	68.2	107.5	169.3
2003 4/.....	1.5	2.8	4.6	7.7	15.4	29.7	47.4	80.5	111.9	184.1
2002.....	1.6	2.7	3.8	7.9	16.0	32.5	49.8	80.4	116.1	190.9
2001.....	1.4	2.5	3.8	7.9	16.4	31.5	51.7	82.0	124.0	201.3
2000.....	1.3	2.5	4.4	8.9	17.8	35.5	59.1	89.4	129.7	219.6
1999.....	1.5	2.7	4.0	9.2	18.6	34.4	58.4	91.1	131.1	211.1

White, female

2006 4/.....	0.5	1.1	1.5	2.7	5.5	10.4	20.9	35.7	58.0	115.7
2005 4/.....	0.8	0.9	1.8	2.9	6.2	11.8	21.3	38.3	62.7	124.2
2004 4/.....	0.7	1.1	1.8	3.1	6.1	12.4	23.9	40.4	64.8	125.7
2003 4/.....	0.7	1.3	1.7	3.4	6.9	12.6	24.4	40.9	65.0	130.5
2002.....	0.6	1.0	1.7	3.7	7.2	13.5	26.5	43.1	65.6	130.7
2001.....	0.7	1.1	2.0	3.4	7.4	14.7	24.7	45.5	67.0	140.1
2000.....	0.7	1.1	1.8	3.3	7.2	15.0	27.1	44.5	71.2	140.1
1999.....	0.9	1.0	1.9	3.6	7.1	14.5	26.9	45.1	73.4	146.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of circulatory system (I71-I78)

Black, both sexes

2006 4/.....	2.7	4.4	7.2	10.1	13.6	23.2	30.6	51.3	83.8	132.3
2005 4/.....	2.6	4.3	6.7	10.4	14.7	24.5	35.2	57.0	88.4	144.6
2004 4/.....	2.6	4.4	6.4	9.4	17.4	24.3	32.7	56.9	90.1	155.1
2003 4/.....	2.6	5.3	7.3	8.5	16.3	24.0	35.9	55.3	92.1	150.0
2002.....	2.8	5.3	6.7	11.2	16.5	25.7	37.8	58.4	83.5	156.8
2001.....	3.2	4.0	6.5	10.7	18.0	22.9	42.8	69.3	96.2	174.7
2000.....	3.2	4.5	6.8	11.2	18.3	27.3	39.9	68.9	92.3	177.7
1999.....	2.8	4.5	6.1	11.4	13.5	26.1	38.6	64.3	96.5	178.9

Black, male

2006 4/.....	3.9	6.1	10.0	15.0	18.4	33.5	36.3	58.2	82.1	117.1
2005 4/.....	3.7	6.7	9.5	15.1	19.3	33.7	39.5	79.1	103.9	129.2
2004 4/.....	3.6	6.3	8.9	13.7	24.3	33.4	37.6	68.2	98.3	158.0
2003 4/.....	3.5	6.8	9.9	11.2	23.5	33.5	40.1	72.6	118.9	138.7
2002.....	4.1	6.7	9.8	16.5	23.0	30.7	48.7	65.7	100.1	166.2
2001.....	4.5	5.5	9.4	13.1	23.8	28.0	55.9	79.0	127.6	165.7
2000.....	4.2	5.5	10.0	15.1	22.6	32.8	51.8	90.7	104.5	177.6
1999.....	3.1	6.1	8.6	14.1	17.8	31.9	49.2	74.6	117.7	184.0

Black, female

2006 4/.....	1.7	2.9	4.7	6.0	9.9	15.6	26.6	47.1	84.6	138.5
2005 4/.....	1.5	2.3	4.4	6.5	11.2	17.6	32.2	43.7	80.1	150.8
2004 4/.....	1.8	2.7	4.2	5.8	11.9	17.4	29.4	50.1	85.8	154.0
2003 4/.....	1.8	4.0	5.1	6.2	10.7	17.0	33.0	45.0	78.1	154.3
2002.....	1.7	4.0	4.0	6.8	11.5	22.0	30.5	54.1	74.8	153.2
2001.....	2.0	2.7	4.1	8.7	13.4	19.1	34.1	63.5	80.1	178.1
2000.....	2.3	3.7	4.1	8.0	14.9	23.2	32.0	55.8	86.2	177.7
1999.....	2.5	3.2	4.0	9.1	10.1	21.7	31.5	58.1	86.0	177.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Aortic aneurysm and dissection (I71)

All races 3/, both sexes

2006.....	1.0	1.6	2.1	3.4	5.6	10.6	18.0	29.5	43.2	62.7
2005.....	0.9	1.4	2.2	3.3	6.8	11.7	19.8	31.6	46.6	64.9
2004.....	0.8	1.4	2.2	3.1	6.4	11.4	21.1	32.6	48.8	64.8
2003.....	0.8	1.7	2.4	3.7	7.3	13.5	22.7	36.6	50.7	70.0
2002.....	0.9	1.5	2.0	4.0	7.4	14.7	23.9	37.5	51.5	69.4
2001.....	0.9	1.4	2.0	3.7	7.8	15.0	23.8	39.0	55.4	77.5
2000.....	0.9	1.3	2.2	4.3	8.4	16.3	27.8	41.0	56.5	79.2
1999.....	0.9	1.4	2.1	4.4	8.1	16.5	27.3	42.2	58.2	80.3

All races 3/, male

2006.....	1.5	2.4	3.3	5.3	8.4	15.2	24.8	41.1	59.0	87.6
2005.....	1.5	2.3	3.3	5.3	10.0	17.1	29.0	43.8	65.4	91.0
2004.....	1.3	2.1	3.3	4.9	9.7	17.0	30.6	44.2	70.3	95.0
2003.....	1.3	2.5	3.7	5.8	11.4	20.2	33.0	53.8	74.2	105.5
2002.....	1.4	2.4	3.2	6.3	11.5	23.0	34.9	55.0	74.5	106.6
2001.....	1.4	2.1	3.2	5.8	12.0	22.2	36.0	56.3	82.5	114.8
2000.....	1.3	1.9	3.5	6.7	13.1	24.9	41.8	61.9	84.9	127.3
1999.....	1.2	2.2	3.3	6.8	13.0	25.3	41.0	63.3	88.6	125.3

All races 3/, female

2006.....	0.4	0.8	1.0	1.6	3.1	6.5	12.5	21.0	33.3	51.0
2005.....	0.4	0.6	1.1	1.5	3.8	7.0	12.4	22.7	35.0	52.9
2004.....	0.4	0.7	1.0	1.5	3.3	6.6	13.5	24.3	35.6	51.2
2003.....	0.4	0.9	1.1	1.7	3.6	7.8	14.5	24.3	36.3	54.2
2002.....	0.3	0.7	1.0	1.8	3.7	7.5	15.1	25.1	37.6	53.3
2001.....	0.3	0.6	0.9	1.8	3.9	8.8	14.0	26.9	39.2	61.9
2000.....	0.5	0.8	1.0	2.1	4.1	8.9	16.7	26.5	39.7	59.6
1999.....	0.5	0.7	1.0	2.1	3.7	9.0	16.6	27.5	40.4	62.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Aortic aneurysm and dissection (I71)

White, both sexes

2006 4/.....	0.8	1.4	1.8	3.1	5.6	10.5	18.5	30.4	44.1	64.3
2005 4/.....	0.8	1.2	1.8	3.2	6.7	11.7	20.6	32.5	47.6	66.2
2004 4/.....	0.7	1.2	1.9	2.8	6.1	11.7	22.1	33.7	49.8	66.8
2003 4/.....	0.7	1.4	2.1	3.6	7.3	14.0	23.6	38.1	52.2	72.4
2002.....	0.7	1.2	1.7	3.8	7.5	15.3	24.8	39.0	53.1	71.2
2001.....	0.7	1.2	1.7	3.6	7.7	15.5	24.6	39.8	57.0	79.7
2000.....	0.7	1.2	2.0	4.1	8.4	16.7	29.0	42.0	58.1	80.8
1999.....	0.7	1.2	1.9	4.2	8.5	17.1	28.4	43.5	60.0	82.2

White, male

2006 4/.....	1.3	2.2	2.9	5.0	8.5	15.2	25.8	42.8	61.0	90.4
2005 4/.....	1.2	1.9	2.8	5.1	10.0	17.2	30.6	45.1	67.1	94.2
2004 4/.....	1.1	1.8	3.0	4.5	9.3	17.4	32.1	45.8	72.1	98.7
2003 4/.....	1.2	2.1	3.4	5.7	11.4	21.1	34.6	56.2	76.5	110.0
2002.....	1.2	2.0	2.7	5.9	11.6	24.3	36.1	57.5	77.7	111.8
2001.....	1.1	1.9	2.8	5.6	11.9	23.2	37.3	57.7	85.5	120.3
2000.....	1.0	1.8	3.2	6.5	13.3	26.1	43.8	63.6	88.6	132.8
1999.....	1.0	1.9	2.9	6.7	13.6	26.5	43.0	66.3	91.5	129.8

White, female

2006 4/.....	0.3	0.6	0.7	1.4	2.9	6.3	12.5	21.2	33.4	52.0
2005 4/.....	0.3	0.4	0.9	1.4	3.5	6.9	12.3	23.1	35.3	53.3
2004 4/.....	0.3	0.5	0.9	1.2	3.2	6.7	13.9	24.8	35.9	52.4
2003 4/.....	0.3	0.7	0.8	1.5	3.5	7.9	14.6	25.0	37.2	55.7
2002.....	0.3	0.5	0.8	1.7	3.7	7.4	15.5	25.8	38.1	53.5
2001.....	0.3	0.5	0.7	1.6	3.9	8.8	14.3	27.1	39.8	62.6
2000.....	0.3	0.5	0.9	1.9	3.9	8.5	17.1	26.8	40.0	59.6
1999.....	0.4	0.5	0.9	1.8	3.7	8.9	16.7	27.5	41.3	63.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Aortic aneurysm and dissection (I71)

Black, both sexes

2006 4/.....	2.2	3.1	5.1	6.0	7.5	13.0	16.0	23.6	34.1	47.2
2005 4/.....	1.8	3.3	4.9	5.6	8.8	13.3	15.9	25.6	38.6	50.2
2004 4/.....	1.8	3.2	4.0	5.8	10.0	11.6	14.9	26.1	40.6	43.7
2003 4/.....	1.8	3.8	4.7	5.7	9.0	11.7	17.9	25.9	36.0	46.4
2002.....	2.1	3.8	4.7	6.8	8.3	11.6	18.4	24.9	37.4	50.5
2001.....	2.1	2.7	4.4	6.2	9.3	11.6	18.4	34.4	42.5	55.3
2000.....	2.3	2.9	3.7	7.3	9.3	14.3	20.3	33.8	40.4	62.3
1999.....	2.0	3.5	4.0	6.5	6.5	14.1	19.8	29.6	40.5	60.3

Black, male

2006 4/.....	3.1	4.5	7.5	9.2	10.2	18.5	19.7	26.4	34.9	50.9
2005 4/.....	3.1	5.3	7.1	8.8	12.0	18.8	17.1	34.0	47.1	51.1
2004 4/.....	2.8	5.1	6.0	8.8	15.9	17.9	18.5	29.6	48.1	56.6
2003 4/.....	2.8	5.8	7.1	8.0	14.7	16.3	20.4	32.9	49.5	51.1
2002.....	3.5	5.7	7.7	11.3	13.1	14.7	24.7	31.2	41.3	49.3
2001.....	3.4	3.9	6.9	8.6	14.6	15.0	26.6	43.2	54.6	48.8
2000.....	3.5	3.6	6.0	10.4	12.8	17.7	27.7	47.7	45.5	69.2
1999.....	2.7	5.4	6.4	9.0	9.3	19.2	25.4	29.9	54.1	76.2

Black, female

2006 4/.....	1.4	1.9	3.0	3.4	5.4	9.0	13.5	22.0	33.6	45.7
2005 4/.....	*	1.6	3.1	3.0	6.4	9.2	15.2	20.5	34.1	49.9
2004 4/.....	*	1.6	2.4	3.3	5.4	6.8	12.5	24.0	36.7	38.6
2003 4/.....	*	2.1	2.7	3.8	4.7	8.4	16.2	21.7	29.0	44.6
2002.....	*	2.1	2.2	3.2	4.5	9.3	14.1	21.1	35.4	50.9
2001.....	*	1.7	2.2	4.2	5.1	9.0	13.0	29.1	36.4	57.7
2000.....	*	2.3	*	4.7	6.5	11.7	15.3	25.3	37.8	59.7
1999.....	1.4	2.0	*	4.4	4.3	10.4	16.0	29.5	33.7	54.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races 3/, both sexes

2006.....	0.3	0.7	1.0	1.9	3.3	5.8	10.2	17.6	29.1	65.8
2005.....	0.4	0.5	1.0	2.1	3.5	6.3	11.4	19.3	31.3	70.6
2004.....	0.4	0.7	1.2	2.1	3.5	6.9	11.9	19.1	32.3	74.5
2003.....	0.4	0.7	1.2	2.0	4.0	7.0	11.4	20.0	32.0	76.0
2002.....	0.4	0.7	1.1	2.2	4.3	7.7	12.6	20.4	32.3	79.0
2001.....	0.5	0.6	1.2	2.2	4.3	7.3	12.9	21.5	32.7	80.7
2000.....	0.4	0.8	1.3	2.1	4.3	8.2	12.8	21.8	35.8	84.0
1999.....	0.5	0.7	1.2	2.3	4.4	7.0	12.9	21.3	36.1	84.9

All races 3/, male

2006.....	0.3	0.7	1.2	2.3	4.0	7.5	12.1	20.8	33.5	67.3
2005.....	0.3	0.6	1.1	2.5	4.2	7.7	13.7	24.6	35.8	67.2
2004.....	0.4	0.8	1.4	2.3	3.9	7.7	13.6	22.9	35.5	71.6
2003.....	0.3	0.7	1.3	2.0	4.4	9.3	13.0	25.1	37.1	74.2
2002.....	0.4	0.8	1.2	2.3	4.9	8.7	14.2	23.4	39.2	80.8
2001.....	0.4	0.7	1.2	2.4	4.8	8.5	15.2	24.7	40.0	82.5
2000.....	0.3	0.8	1.4	2.6	4.9	9.7	15.7	26.6	41.7	87.6
1999.....	0.4	0.8	1.2	2.6	5.2	8.1	15.7	25.9	40.8	82.2

All races 3/, female

2006.....	0.2	0.6	0.9	1.5	2.7	4.3	8.6	15.3	26.2	65.0
2005.....	0.5	0.5	0.9	1.7	2.8	5.2	9.6	15.5	28.4	72.2
2004.....	0.5	0.6	1.0	1.8	3.2	6.2	10.5	16.3	30.3	75.8
2003.....	0.5	0.7	1.0	1.9	3.6	5.0	10.1	16.3	28.9	76.9
2002.....	0.4	0.6	1.0	2.1	3.7	6.7	11.3	18.3	28.1	78.2
2001.....	0.5	0.6	1.3	2.0	3.9	6.3	11.1	19.3	28.3	79.9
2000.....	0.5	0.7	1.1	1.6	3.7	6.9	10.5	18.5	32.2	82.6
1999.....	0.5	0.6	1.2	2.1	3.6	6.0	10.6	18.2	33.3	86.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

White, both sexes

2006 4/.....	0.2	0.6	0.9	1.7	3.1	5.5	10.0	17.1	27.8	65.4
2005 4/.....	0.4	0.5	1.0	1.8	3.3	6.0	11.0	18.6	30.3	69.9
2004 4/.....	0.4	0.7	1.1	1.9	3.2	6.4	11.7	18.4	31.4	72.5
2003 4/.....	0.3	0.6	1.0	1.9	3.7	6.6	11.1	19.5	30.7	74.6
2002.....	0.4	0.6	1.0	2.0	3.9	7.1	12.2	19.7	31.7	77.8
2001.....	0.4	0.6	1.2	2.1	4.0	7.1	12.2	20.8	31.4	78.5
2000.....	0.3	0.7	1.0	1.9	3.9	7.8	12.4	21.0	34.9	82.3
1999.....	0.5	0.7	1.1	2.1	4.1	6.6	12.5	20.6	34.8	82.9

White, male

2006 4/.....	0.2	0.7	1.1	2.0	3.7	7.1	12.1	20.5	33.0	69.0
2005 4/.....	0.3	0.5	1.0	2.1	4.1	7.1	13.3	23.4	35.0	67.8
2004 4/.....	0.4	0.8	1.3	2.1	3.6	7.1	13.6	22.3	35.4	70.7
2003 4/.....	0.3	0.7	1.2	2.0	4.0	8.7	12.8	24.4	35.4	74.1
2002.....	0.4	0.8	1.1	2.0	4.4	8.2	13.7	23.0	38.4	79.1
2001.....	0.3	0.6	1.1	2.3	4.4	8.3	14.4	24.2	38.5	80.9
2000.....	0.3	0.7	1.1	2.4	4.5	9.4	15.3	25.9	41.1	86.9
1999.....	0.4	0.8	1.1	2.4	4.9	7.9	15.3	24.8	39.6	81.3

White, female

2006 4/.....	0.3	0.5	0.8	1.4	2.6	4.1	8.3	14.5	24.6	63.7
2005 4/.....	0.5	0.5	0.9	1.5	2.6	4.9	9.0	15.1	27.4	70.8
2004 4/.....	0.4	0.5	0.9	1.8	2.8	5.7	10.1	15.6	28.9	73.3
2003 4/.....	0.4	0.6	0.9	1.9	3.4	4.8	9.8	15.9	27.8	74.8
2002.....	0.3	0.5	0.9	2.0	3.4	6.1	11.0	17.3	27.5	77.2
2001.....	0.5	0.6	1.2	1.8	3.5	6.0	10.4	18.4	27.1	77.5
2000.....	0.4	0.6	0.9	1.4	3.3	6.5	10.0	17.6	31.2	80.4
1999.....	0.5	0.5	1.0	1.8	3.4	5.6	10.2	17.6	32.0	83.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

Black, both sexes

2006 4/.....	*	1.3	2.1	4.0	6.1	10.2	14.5	27.6	49.7	85.1
2005 4/.....	0.7	1.0	1.8	4.7	5.9	11.2	19.3	31.4	49.8	94.3
2004 4/.....	0.8	1.1	2.3	3.6	7.4	12.7	17.8	30.8	49.5	111.4
2003 4/.....	0.8	1.5	2.6	2.8	7.3	12.3	18.0	29.4	56.1	103.6
2002.....	0.7	1.5	2.0	4.4	8.3	14.1	19.4	33.5	46.1	106.3
2001.....	1.1	1.3	2.1	4.5	8.7	11.3	24.4	34.9	53.6	119.4
2000.....	0.9	1.7	3.1	3.9	9.0	13.0	19.6	35.2	52.0	115.4
1999.....	0.8	1.0	2.1	4.9	7.0	11.9	18.8	34.6	56.1	118.6

Black, male

2006 4/.....	*	1.6	2.6	5.8	8.1	15.0	16.6	31.8	47.2	66.1
2005 4/.....	*	*	2.4	6.2	7.3	15.0	22.5	45.1	56.9	78.1
2004 4/.....	*	*	2.9	4.9	8.4	15.5	19.1	38.7	50.2	101.3
2003 4/.....	*	*	2.8	3.2	8.9	17.2	19.7	39.7	69.4	87.6
2002.....	*	*	2.1	5.2	9.9	16.0	24.0	34.4	58.8	116.9
2001.....	*	*	2.5	4.5	9.2	13.0	29.3	35.8	73.0	116.9
2000.....	*	1.9	4.0	4.8	9.7	15.1	24.1	43.0	59.0	108.4
1999.....	*	*	*	5.1	8.5	12.7	23.7	44.6	63.7	107.8

Black, female

2006 4/.....	*	*	1.7	2.6	4.5	6.6	13.1	25.2	51.0	92.9
2005 4/.....	*	*	*	3.4	4.8	8.5	17.1	23.2	46.0	100.9
2004 4/.....	*	*	1.8	2.5	6.6	10.6	16.9	26.1	49.1	115.4
2003 4/.....	*	1.8	2.4	2.4	6.0	8.6	16.8	23.3	49.1	109.8
2002.....	*	1.9	1.8	3.7	7.0	12.8	16.4	32.9	39.4	102.3
2001.....	*	*	1.8	4.5	8.3	10.1	21.1	34.4	43.7	120.4
2000.....	*	*	2.4	3.2	8.3	11.5	16.6	30.5	48.4	118.0
1999.....	*	*	*	4.7	5.8	11.4	15.5	28.6	52.3	122.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes

2006.....	0.7	1.1	1.4	1.6	1.9	2.4	3.5	5.6	8.7	18.4
2005.....	0.9	1.3	1.6	1.9	2.4	3.3	4.4	7.0	10.8	23.0
2004.....	0.9	1.3	1.4	1.9	2.5	3.1	4.3	7.1	11.5	23.3
2003.....	0.9	1.3	1.5	1.7	2.5	3.3	4.9	7.0	11.9	24.0
2002.....	0.8	1.2	1.5	2.0	2.5	3.4	4.3	7.4	13.2	24.4
2001.....	0.8	1.1	1.3	1.9	2.6	3.8	5.4	7.7	11.9	26.1
2000.....	0.9	1.2	1.5	1.7	2.6	3.4	5.3	7.7	11.9	25.6
1999.....	0.8	1.0	1.0	1.6	2.2	3.3	4.3	6.8	12.7	23.4

All races 3/, male

2006.....	0.8	1.3	1.8	1.9	2.3	2.4	3.5	6.1	8.3	16.2
2005.....	1.0	1.6	2.1	2.3	3.0	4.0	5.0	6.6	11.1	19.2
2004.....	1.0	1.4	1.8	2.2	3.0	3.3	4.6	7.9	12.2	19.6
2003.....	1.0	1.4	1.8	2.2	2.7	3.5	4.9	6.9	10.9	19.1
2002.....	0.8	1.3	1.6	2.3	2.6	3.6	4.2	7.0	13.3	20.5
2001.....	0.9	1.2	1.5	2.2	2.9	3.9	5.9	8.3	12.1	22.0
2000.....	1.1	1.5	1.7	1.9	2.7	3.9	5.6	8.0	12.4	22.2
1999.....	0.9	1.1	1.2	1.7	2.3	3.7	4.8	7.5	13.2	20.8

All races 3/, female

2006.....	0.7	1.0	0.9	1.2	1.6	2.5	3.6	5.3	9.0	19.5
2005.....	0.9	1.0	1.2	1.5	1.8	2.7	4.0	7.2	10.7	24.8
2004.....	0.8	1.1	1.1	1.6	2.0	2.9	4.0	6.5	11.0	25.0
2003.....	0.9	1.1	1.2	1.3	2.3	3.0	5.0	7.0	12.5	26.2
2002.....	0.8	1.1	1.4	1.7	2.5	3.2	4.4	7.7	13.1	26.2
2001.....	0.8	0.9	1.0	1.7	2.3	3.7	5.0	7.3	11.7	27.8
2000.....	0.7	1.0	1.3	1.5	2.6	3.0	5.0	7.5	11.6	26.9
1999.....	0.6	1.0	0.9	1.5	2.1	2.9	3.9	6.3	12.4	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other disorders of circulatory system (I80-I99)

White, both sexes

2006 4/.....	0.6	1.0	1.2	1.5	1.7	2.3	3.4	5.5	8.4	18.1
2005 4/.....	0.7	1.2	1.4	1.7	2.1	3.2	4.2	6.7	10.5	23.5
2004 4/.....	0.7	1.1	1.2	1.8	2.3	3.0	3.9	6.8	11.3	23.2
2003 4/.....	0.8	1.1	1.4	1.6	2.2	3.0	4.7	6.5	11.6	23.6
2002.....	0.6	1.0	1.4	1.9	2.2	3.2	4.1	7.1	12.7	24.3
2001.....	0.7	0.9	1.1	1.8	2.3	3.6	5.1	7.2	11.8	26.0
2000.....	0.8	1.1	1.4	1.5	2.4	2.9	5.0	7.3	11.8	25.5
1999.....	0.6	0.9	0.9	1.6	2.1	2.9	4.1	6.5	12.4	23.2

White, male

2006 4/.....	0.6	1.1	1.5	1.9	1.9	2.2	3.4	6.0	7.6	16.0
2005 4/.....	0.8	1.4	1.9	2.1	2.6	3.7	4.6	6.4	10.5	19.1
2004 4/.....	0.8	1.4	1.5	2.1	2.7	3.1	4.3	7.6	11.9	19.7
2003 4/.....	0.9	1.2	1.6	2.1	2.3	3.2	4.6	6.4	10.2	18.5
2002.....	0.6	1.1	1.5	2.1	2.3	3.3	4.0	6.7	12.4	19.9
2001.....	0.7	1.2	1.3	2.0	2.5	3.6	5.6	7.8	12.0	21.7
2000.....	1.0	1.3	1.6	1.7	2.4	3.4	5.5	7.8	12.3	22.0
1999.....	0.8	0.9	1.0	1.7	2.3	3.2	4.7	7.0	12.8	20.7

White, female

2006 4/.....	0.5	0.8	0.8	1.1	1.4	2.3	3.5	5.2	8.9	19.1
2005 4/.....	0.7	1.0	1.0	1.4	1.6	2.7	3.9	6.9	10.5	25.5
2004 4/.....	0.6	0.9	0.9	1.4	1.8	2.8	3.6	6.2	11.0	24.8
2003 4/.....	0.7	1.1	1.2	1.1	2.2	2.9	4.7	6.5	12.4	25.8
2002.....	0.7	1.0	1.2	1.6	2.1	3.0	4.2	7.4	12.9	26.3
2001.....	0.7	0.7	1.0	1.5	2.1	3.5	4.8	6.8	11.7	27.9
2000.....	0.5	0.9	1.1	1.4	2.4	2.5	4.7	7.0	11.5	26.9
1999.....	0.5	0.9	0.9	1.5	1.9	2.7	3.7	6.1	12.2	24.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other disorders of circulatory system (I80-I99)

Black, both sexes

2006 4/.....	1.9	2.7	3.0	2.7	4.4	4.2	5.8	8.0	14.9	25.7
2005 4/.....	2.3	2.6	3.5	4.0	5.2	5.5	7.6	11.5	17.5	24.3
2004 4/.....	2.5	2.4	3.6	3.6	5.0	4.9	8.9	12.9	14.6	28.8
2003 4/.....	2.3	2.6	2.5	3.6	5.2	6.2	9.3	13.8	17.8	33.6
2002.....	1.8	2.4	2.9	4.2	5.9	6.4	7.2	12.9	22.2	30.0
2001.....	1.8	2.1	2.4	3.9	6.1	6.9	9.0	14.1	14.9	30.1
2000.....	1.8	2.5	3.2	3.9	5.1	8.7	9.1	14.6	14.1	30.0
1999.....	1.9	2.4	2.3	2.1	3.9	6.9	6.4	11.4	17.6	28.4

Black, male

2006 4/.....	1.8	2.6	4.7	2.9	6.2	4.7	*	*	17.1	22.3
2005 4/.....	2.4	3.6	4.1	4.8	6.9	7.8	10.1	12.1	22.5	25.1
2004 4/.....	2.5	2.1	5.0	3.6	6.2	5.8	8.7	14.1	16.5	24.8
2003 4/.....	1.8	3.4	3.4	3.6	7.3	7.6	10.2	15.5	20.7	32.3
2002.....	2.1	3.0	2.9	5.4	5.9	7.7	7.7	12.4	28.3	32.2
2001.....	2.2	1.7	3.1	4.7	8.5	7.9	10.4	16.3	*	29.5
2000.....	2.0	3.1	3.7	4.8	5.6	9.4	8.7	13.2	*	30.0
1999.....	2.6	2.7	3.0	*	*	9.3	6.8	14.7	20.9	*

Black, female

2006 4/.....	1.9	2.8	1.6	2.6	3.0	3.8	5.9	8.0	13.7	27.1
2005 4/.....	2.2	1.8	3.0	3.3	3.8	3.7	5.8	11.2	14.8	24.0
2004 4/.....	2.6	2.6	2.4	3.7	4.1	4.1	9.0	12.1	13.6	30.3
2003 4/.....	2.7	1.9	1.8	3.6	3.6	5.1	8.6	12.8	16.2	34.0
2002.....	1.6	1.8	2.9	3.2	5.9	5.4	6.9	13.2	19.1	29.2
2001.....	1.5	2.4	*	3.2	4.2	6.2	8.1	12.8	14.7	30.3
2000.....	1.7	1.9	2.8	3.2	4.7	8.3	9.3	15.4	14.4	30.0
1999.....	*	2.1	*	*	4.2	5.0	6.1	9.4	16.0	30.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza and pneumonia (J10-J18)

All races 3/, both sexes

2006.....	2.4	3.6	5.7	7.6	13.2	23.1	42.9	86.2	181.8	502.5
2005.....	2.7	4.2	6.1	8.8	14.6	24.5	48.6	96.1	202.7	593.9
2004.....	2.5	3.6	5.6	8.3	14.2	23.8	47.2	94.7	198.7	582.6
2003.....	2.7	4.2	6.4	8.2	15.2	26.6	49.4	103.3	216.8	666.1
2002.....	2.6	4.2	5.5	8.5	14.8	25.3	50.9	106.5	227.3	696.6
2001.....	2.6	3.9	5.4	8.1	13.9	24.4	49.3	101.1	216.9	685.6
2000.....	3.0	3.9	5.6	8.9	15.6	26.6	52.6	106.7	240.8	744.1
1999.....	2.7	4.1	5.3	8.4	14.3	25.0	50.5	105.9	233.8	751.8

All races 3/, male

2006.....	2.7	4.2	7.1	9.4	16.4	28.1	52.5	108.1	226.8	586.4
2005.....	3.1	5.1	7.5	10.7	17.5	29.6	59.0	119.3	253.7	673.5
2004.....	2.7	4.3	7.1	9.9	17.2	29.8	58.6	118.3	252.3	677.2
2003.....	3.1	4.9	8.1	9.6	17.9	33.3	62.1	130.1	272.3	757.8
2002.....	3.1	4.9	6.7	10.2	18.0	31.2	63.9	133.9	287.7	805.8
2001.....	3.0	4.9	6.7	10.0	17.0	30.3	61.6	130.7	274.7	808.4
2000.....	3.8	4.7	7.0	10.9	18.9	32.9	65.7	134.1	308.8	892.0
1999.....	3.1	5.2	6.8	9.7	16.8	31.7	64.2	133.5	301.7	894.7

All races 3/, female

2006.....	2.2	3.1	4.4	6.0	10.3	18.8	35.0	70.2	153.7	463.3
2005.....	2.3	3.4	4.9	6.9	12.0	20.1	40.2	79.3	171.1	557.4
2004.....	2.3	3.0	4.2	6.8	11.4	18.6	37.9	77.7	165.7	540.0
2003.....	2.3	3.5	4.8	6.8	12.7	20.7	39.2	84.2	182.9	625.6
2002.....	2.2	3.4	4.3	6.8	11.8	20.1	40.6	87.2	190.6	649.2
2001.....	2.2	3.0	4.1	6.4	11.1	19.2	39.5	80.5	182.3	634.2
2000.....	2.2	3.1	4.3	7.0	12.5	21.2	42.2	87.6	200.7	683.9
1999.....	2.3	3.0	3.9	7.2	12.0	19.2	39.7	86.7	194.1	694.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza and pneumonia (J10-J18)

White, both sexes

2006 4/.....	2.3	3.2	5.2	7.2	12.7	21.7	42.5	85.6	181.5	509.3
2005 4/.....	2.4	3.8	5.5	8.0	13.7	23.3	47.0	95.1	204.0	606.4
2004 4/.....	2.2	3.2	5.0	7.7	13.3	22.3	46.2	93.1	198.3	590.7
2003 4/.....	2.3	3.7	5.7	7.4	14.0	25.2	47.9	102.4	218.4	679.3
2002.....	2.3	3.8	5.0	7.7	13.8	24.5	49.9	105.4	228.0	709.1
2001.....	2.2	3.3	4.8	7.5	13.0	23.3	48.1	98.6	216.6	695.0
2000.....	2.5	3.4	4.9	8.1	14.3	24.5	50.3	105.6	240.7	758.3
1999.....	2.1	3.4	4.6	7.7	13.1	24.0	48.8	105.5	235.5	764.3

White, male

2006 4/.....	2.6	3.7	6.3	8.9	15.4	26.2	51.3	106.5	225.6	591.4
2005 4/.....	2.7	4.5	6.6	9.4	16.1	27.6	56.5	117.7	253.9	684.4
2004 4/.....	2.4	3.8	6.2	9.1	15.9	27.1	56.7	114.6	250.8	685.5
2003 4/.....	2.6	4.3	7.2	8.5	15.7	30.8	59.2	127.2	271.4	767.1
2002.....	2.7	4.4	6.0	9.1	16.5	29.8	62.4	131.4	286.5	815.9
2001.....	2.5	4.2	5.7	9.0	15.4	28.3	59.3	125.5	272.3	814.4
2000.....	3.2	3.9	5.9	9.5	17.2	29.7	62.2	131.8	306.4	906.1
1999.....	2.3	4.2	5.7	8.6	15.1	30.1	60.9	132.2	301.9	905.7

White, female

2006 4/.....	2.0	2.8	4.2	5.6	10.1	17.6	35.3	70.0	153.6	470.7
2005 4/.....	2.1	3.2	4.4	6.6	11.4	19.5	39.2	78.4	172.8	570.4
2004 4/.....	2.1	2.7	3.8	6.2	10.9	18.1	37.5	77.3	165.6	548.0
2003 4/.....	2.1	3.1	4.3	6.3	12.4	20.2	38.6	84.5	185.7	640.4
2002.....	1.9	3.2	3.9	6.3	11.3	19.8	39.7	86.8	192.1	662.6
2001.....	1.8	2.5	4.0	6.0	10.7	19.0	39.0	79.6	182.9	645.0
2000.....	1.8	2.8	4.0	6.7	11.6	20.0	40.8	87.1	201.7	698.3
1999.....	1.9	2.6	3.6	6.7	11.2	18.7	39.1	86.8	196.4	707.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza and pneumonia (J10-J18)

Black, both sexes

2006 4/.....	4.1	7.2	10.8	12.9	20.3	39.7	51.0	99.4	196.9	434.6
2005 4/.....	4.9	7.9	12.1	16.8	24.5	39.2	68.2	115.8	199.3	471.2
2004 4/.....	5.0	7.0	11.6	15.5	23.8	39.8	62.4	117.6	215.1	505.1
2003 4/.....	5.5	8.5	12.6	15.7	27.1	41.6	70.1	122.5	205.6	525.4
2002.....	5.2	7.5	10.1	17.2	26.3	35.5	68.9	123.3	231.6	575.2
2001.....	5.8	9.0	10.7	15.5	23.3	34.3	65.7	133.7	224.0	581.3
2000.....	6.1	8.3	12.2	17.8	27.2	46.0	77.5	124.4	245.4	600.9
1999.....	6.9	9.7	11.8	16.2	26.6	36.8	75.2	118.7	219.9	634.8

Black, male

2006 4/.....	4.5	8.3	15.4	16.1	29.3	50.1	68.9	139.2	245.1	504.9
2005 4/.....	6.1	10.5	15.9	23.8	33.3	53.7	90.5	144.4	259.1	531.7
2004 4/.....	5.9	8.5	15.8	19.1	33.1	61.3	87.4	169.7	297.7	598.0
2003 4/.....	7.0	10.4	16.7	21.3	39.7	61.6	100.5	177.6	285.0	641.2
2002.....	6.2	9.9	12.9	23.7	36.0	48.1	89.8	173.2	321.7	689.4
2001.....	7.0	11.4	15.7	20.8	33.6	50.1	94.9	203.5	309.0	714.8
2000.....	8.3	11.3	17.2	25.9	35.2	64.8	107.6	172.5	343.9	755.4
1999.....	9.0	13.4	17.2	21.7	34.2	51.9	109.6	160.0	312.2	803.8

Black, female

2006 4/.....	3.8	6.2	6.9	10.3	13.2	32.0	38.7	75.5	171.4	405.8
2005 4/.....	3.9	5.7	9.0	11.0	17.7	28.4	53.0	98.7	167.7	446.8
2004 4/.....	4.2	5.7	8.0	12.6	16.6	23.9	45.5	86.8	171.6	468.2
2003 4/.....	4.1	7.0	9.1	11.1	17.3	26.9	49.6	89.8	164.0	480.4
2002.....	4.3	5.5	7.7	12.0	18.7	26.1	54.9	93.6	184.6	531.5
2001.....	4.8	6.9	6.5	11.2	15.2	22.7	46.2	92.1	180.7	531.7
2000.....	4.2	5.8	7.9	11.2	20.9	32.1	57.5	95.4	195.7	543.4
1999.....	5.1	6.6	7.1	11.8	20.6	25.6	52.4	93.8	174.1	571.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza (J10-J11)

All races 3/, both sexes

2006.....	*	*	*	0.1	0.2	0.3	0.6	1.3	2.5	7.0
2005.....	*	*	0.1	0.1	0.3	0.4	1.1	2.4	5.4	19.3
2004.....	*	*	*	0.1	0.3	0.3	0.8	1.6	3.2	11.4
2003.....	*	0.1	0.2	0.3	0.3	0.7	1.3	2.7	4.7	17.0
2002.....	*	*	*	*	*	*	0.4	0.8	2.2	8.8
2001.....	*	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	0.1	0.1	0.3	0.5	0.8	1.4	2.5	6.1	20.2
1999.....	*	*	*	0.2	0.4	0.6	1.2	2.1	6.1	21.2

All races 3/, male

2006.....	*	*	*	*	*	*	0.7	1.3	3.0	8.1
2005.....	*	*	*	*	0.4	*	1.1	2.5	5.3	18.2
2004.....	*	*	*	*	*	*	1.1	1.8	3.3	11.1
2003.....	*	*	*	0.3	*	0.8	1.3	2.9	4.9	17.0
2002.....	*	*	*	*	*	*	*	0.9	2.6	8.9
2001.....	*	*	*	*	*	*	*	*	*	2.2
2000.....	*	*	*	0.3	0.5	1.0	1.6	2.7	6.3	20.7
1999.....	*	*	*	*	0.5	0.6	1.2	2.4	7.3	19.9

All races 3/, female

2006.....	*	*	*	*	*	*	0.5	1.3	2.2	6.5
2005.....	*	*	*	*	*	0.5	1.1	2.3	5.5	19.8
2004.....	*	*	*	*	*	*	0.6	1.4	3.1	11.5
2003.....	*	*	*	0.3	0.3	0.7	1.3	2.5	4.7	17.1
2002.....	*	*	*	*	*	*	*	0.7	2.0	8.7
2001.....	*	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	*	*	*	0.5	0.6	1.2	2.5	6.0	19.9
1999.....	*	*	*	*	*	0.6	1.1	1.9	5.4	21.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Influenza (J10-J11)

White, both sexes

2006 4/.....	*	*	*	0.1	0.2	0.3	0.7	1.4	2.7	7.5
2005 4/.....	*	*	0.1	0.2	0.3	0.5	1.2	2.5	5.8	20.8
2004 4/.....	*	*	*	0.2	0.3	0.3	0.9	1.5	3.4	12.3
2003 4/.....	*	0.1	0.2	0.3	0.3	0.8	1.4	2.9	5.1	18.2
2002.....	*	*	*	*	*	*	0.4	0.9	2.4	9.4
2001.....	*	*	*	*	*	*	*	*	0.7	2.3
2000.....	*	*	0.1	0.3	0.5	0.8	1.4	2.7	6.5	21.4
1999.....	*	*	*	0.2	0.4	0.7	1.2	2.2	6.4	22.8

White, male

2006 4/.....	*	*	*	*	*	*	0.8	1.3	3.1	8.6
2005 4/.....	*	*	*	*	*	*	1.2	2.7	5.6	19.2
2004 4/.....	*	*	*	*	*	*	1.1	1.7	3.6	12.1
2003 4/.....	*	*	*	0.3	*	0.9	1.4	3.1	5.2	18.3
2002.....	*	*	*	*	*	*	*	1.0	2.8	9.7
2001.....	*	*	*	*	*	*	*	*	*	2.3
2000.....	*	*	*	*	0.5	1.0	1.7	2.7	6.7	21.6
1999.....	*	*	*	*	0.5	0.6	1.2	2.6	7.5	21.7

White, female

2006 4/.....	*	*	*	*	*	*	0.5	1.4	2.4	7.1
2005 4/.....	*	*	*	*	*	0.5	1.2	2.4	5.9	21.5
2004 4/.....	*	*	*	*	*	*	0.7	1.4	3.3	12.4
2003 4/.....	*	*	*	*	0.4	0.7	1.4	2.8	5.0	18.1
2002.....	*	*	*	*	*	*	*	0.8	2.2	9.3
2001.....	*	*	*	*	*	*	*	*	0.7	2.3
2000.....	*	*	*	*	0.6	0.7	1.2	2.7	6.3	21.3
1999.....	*	*	*	*	*	0.7	1.1	2.0	5.7	23.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonia (J12-J18)

All races 3/, both sexes

2006.....	2.4	3.6	5.6	7.5	13.0	22.8	42.3	85.0	179.3	495.5
2005.....	2.6	4.2	6.0	8.6	14.3	24.1	47.5	93.7	197.3	574.7
2004.....	2.5	3.6	5.6	8.2	13.9	23.5	46.4	93.1	195.6	571.2
2003.....	2.6	4.1	6.2	7.9	14.9	25.8	48.1	100.6	212.1	649.1
2002.....	2.6	4.1	5.4	8.4	14.6	25.1	50.5	105.7	225.1	687.8
2001.....	2.6	3.9	5.3	8.1	13.8	24.3	49.2	100.9	216.1	683.4
2000.....	2.9	3.8	5.5	8.6	15.1	25.8	51.2	104.1	234.7	723.9
1999.....	2.7	4.0	5.2	8.2	13.9	24.3	49.3	103.8	227.7	730.6

All races 3/, male

2006.....	2.7	4.2	7.0	9.3	16.3	27.8	51.8	106.9	223.8	578.3
2005.....	3.0	5.0	7.4	10.5	17.1	29.2	57.9	116.8	248.4	655.3
2004.....	2.7	4.2	7.0	9.7	17.0	29.4	57.6	116.5	249.1	666.1
2003.....	3.0	4.8	7.9	9.3	17.6	32.4	60.8	127.2	267.5	740.9
2002.....	3.0	4.9	6.7	10.1	17.8	30.9	63.4	133.0	285.0	796.9
2001.....	3.0	4.8	6.6	9.9	16.9	30.3	61.4	130.5	273.9	806.2
2000.....	3.7	4.6	6.8	10.6	18.4	31.9	64.1	131.5	302.6	871.3
1999.....	3.1	5.1	6.7	9.5	16.3	31.1	63.0	131.1	294.5	874.7

All races 3/, female

2006.....	2.1	3.1	4.3	5.9	10.1	18.5	34.5	69.0	151.5	456.7
2005.....	2.3	3.4	4.7	6.8	11.7	19.6	39.1	77.0	165.6	537.6
2004.....	2.3	2.9	4.1	6.7	11.1	18.4	37.3	76.4	162.6	528.5
2003.....	2.2	3.4	4.7	6.5	12.4	20.1	37.9	81.7	178.2	608.5
2002.....	2.2	3.4	4.3	6.8	11.8	20.0	40.2	86.5	188.6	640.5
2001.....	2.2	2.9	4.1	6.3	11.0	19.1	39.4	80.2	181.5	632.0
2000.....	2.1	3.0	4.2	6.7	12.0	20.5	41.0	85.1	194.6	663.9
1999.....	2.3	2.9	3.9	7.0	11.7	18.6	38.6	84.8	188.7	672.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonia (J12-J18)

White, both sexes

2006 4/.....	2.2	3.2	5.1	7.1	12.5	21.4	41.9	84.2	178.8	501.7
2005 4/.....	2.4	3.8	5.4	7.8	13.4	22.8	45.8	92.6	198.2	585.6
2004 4/.....	2.2	3.2	4.9	7.5	13.1	22.0	45.3	91.5	194.9	578.4
2003 4/.....	2.3	3.6	5.6	7.1	13.7	24.4	46.5	99.5	213.3	661.1
2002.....	2.3	3.8	4.9	7.6	13.7	24.3	49.5	104.5	225.6	699.6
2001.....	2.1	3.3	4.8	7.4	12.9	23.2	48.0	98.3	215.8	692.7
2000.....	2.4	3.2	4.8	7.8	13.8	23.7	48.9	102.9	234.3	737.0
1999.....	2.1	3.3	4.6	7.4	12.6	23.4	47.6	103.3	229.2	741.6

White, male

2006 4/.....	2.5	3.7	6.2	8.8	15.3	25.8	50.5	105.2	222.6	582.8
2005 4/.....	2.7	4.4	6.5	9.2	15.8	27.2	55.3	115.1	248.3	665.2
2004 4/.....	2.3	3.8	6.2	9.0	15.7	26.7	55.6	112.9	247.3	673.4
2003 4/.....	2.5	4.2	7.0	8.2	15.4	29.9	57.8	124.1	266.2	748.8
2002.....	2.7	4.3	6.0	9.0	16.3	29.6	61.9	130.4	283.7	806.2
2001.....	2.5	4.1	5.6	9.0	15.3	28.3	59.2	125.3	271.5	812.1
2000.....	3.1	3.8	5.7	9.2	16.7	28.6	60.6	129.1	299.7	884.4
1999.....	2.2	4.2	5.6	8.4	14.6	29.5	59.7	129.6	294.3	884.0

White, female

2006 4/.....	1.9	2.8	4.1	5.5	9.9	17.4	34.8	68.5	151.2	463.7
2005 4/.....	2.1	3.1	4.3	6.5	11.1	19.0	38.0	75.9	166.9	548.9
2004 4/.....	2.1	2.6	3.7	6.1	10.6	17.8	36.9	75.9	162.3	535.6
2003 4/.....	2.0	3.0	4.2	6.0	12.1	19.5	37.2	81.7	180.7	622.3
2002.....	1.9	3.2	3.9	6.2	11.3	19.7	39.4	86.0	190.0	653.3
2001.....	1.8	2.4	3.9	5.9	10.6	18.8	38.9	79.3	182.2	642.8
2000.....	1.8	2.7	3.9	6.4	11.1	19.4	39.6	84.5	195.4	677.1
1999.....	1.9	2.5	3.5	6.5	10.8	18.0	37.9	84.8	190.7	684.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonia (J12-J18)

Black, both sexes

2006 4/.....	4.1	7.1	10.7	12.9	20.1	39.5	50.9	99.0	195.0	432.7
2005 4/.....	4.9	7.8	12.0	16.7	24.1	39.0	67.8	114.9	197.1	467.2
2004 4/.....	4.9	6.9	11.5	15.5	23.5	39.5	61.9	115.6	214.1	502.5
2003 4/.....	5.4	8.3	12.4	15.4	27.0	41.0	69.6	121.7	203.6	519.6
2002.....	5.2	7.5	10.1	17.2	26.1	35.0	68.6	123.2	230.8	572.8
2001.....	5.8	9.0	10.7	15.4	23.1	34.1	65.7	133.7	222.9	579.7
2000.....	6.0	8.1	12.1	17.7	26.9	45.4	76.0	123.5	243.1	591.5
1999.....	6.8	9.7	11.6	16.0	26.4	36.5	73.9	117.8	217.0	629.1

Black, male

2006 4/.....	4.5	8.3	15.3	16.1	29.2	49.9	68.9	138.7	241.6	502.2
2005 4/.....	6.1	10.2	15.7	23.6	32.5	53.7	89.8	143.9	257.7	524.3
2004 4/.....	5.9	8.4	15.8	19.1	32.9	61.1	86.4	165.6	297.0	597.1
2003 4/.....	7.0	10.0	16.5	20.8	39.5	60.8	100.2	176.6	283.5	638.1
2002.....	6.2	9.9	12.9	23.5	35.6	47.4	89.8	173.2	320.2	687.2
2001.....	7.0	11.4	15.7	20.6	33.2	49.6	94.9	203.5	309.0	712.6
2000.....	8.1	11.0	17.0	25.5	35.0	64.3	106.6	170.6	343.1	742.7
1999.....	8.9	13.3	17.1	21.7	33.9	51.4	108.2	159.6	307.8	800.3

Black, female

2006 4/.....	3.7	6.0	6.9	10.3	13.1	31.8	38.5	75.3	170.4	404.3
2005 4/.....	3.8	5.7	8.8	10.9	17.6	28.0	52.8	97.6	165.1	444.1
2004 4/.....	4.1	5.7	7.9	12.6	16.2	23.6	45.3	86.0	170.5	465.0
2003 4/.....	4.1	6.7	8.9	11.0	17.1	26.3	49.0	89.0	161.7	473.5
2002.....	4.3	5.5	7.7	12.0	18.6	25.9	54.4	93.3	184.2	529.0
2001.....	4.7	6.9	6.5	11.2	15.2	22.7	46.2	92.1	179.1	530.5
2000.....	4.2	5.6	7.9	11.2	20.4	31.5	55.7	95.1	192.8	535.3
1999.....	5.0	6.6	7.0	11.4	20.5	25.4	51.1	92.7	172.0	565.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute lower respiratory infections (J20-J22)

All races 3/, both sexes

2006.....	*	*	*	*	*	*	*	0.3	0.6	2.0
2005.....	*	*	*	*	*	*	*	0.4	0.9	3.3
2004.....	*	*	*	*	*	*	*	0.3	0.6	2.6
2003.....	*	*	*	*	*	*	*	0.5	0.8	3.5
2002.....	*	*	*	*	*	*	0.3	0.3	0.8	3.3
2001.....	*	*	*	*	*	*	*	0.3	0.7	3.3
2000.....	*	*	*	*	*	*	0.3	0.6	0.8	4.5
1999.....	*	*	*	*	*	*	0.2	0.6	1.3	6.0

All races 3/, male

2006.....	*	*	*	*	*	*	*	*	*	1.8
2005.....	*	*	*	*	*	*	*	*	1.3	3.1
2004.....	*	*	*	*	*	*	*	*	*	2.5
2003.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	2.8
2001.....	*	*	*	*	*	*	*	*	*	2.3
2000.....	*	*	*	*	*	*	*	0.7	*	3.8
1999.....	*	*	*	*	*	*	*	0.7	1.6	4.0

All races 3/, female

2006.....	*	*	*	*	*	*	*	*	0.7	2.1
2005.....	*	*	*	*	*	*	*	*	0.6	3.5
2004.....	*	*	*	*	*	*	*	*	0.7	2.7
2003.....	*	*	*	*	*	*	*	*	0.8	3.9
2002.....	*	*	*	*	*	*	*	*	0.9	3.5
2001.....	*	*	*	*	*	*	*	*	0.7	3.7
2000.....	*	*	*	*	*	*	*	0.5	0.9	4.8
1999.....	*	*	*	*	*	*	*	0.5	1.1	6.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other acute lower respiratory infections (J20-J22)

White, both sexes

2006 4/.....	*	*	*	*	*	*	*	0.3	0.7	2.2
2005 4/.....	*	*	*	*	*	*	*	0.4	0.9	3.5
2004 4/.....	*	*	*	*	*	*	*	*	0.7	2.8
2003 4/.....	*	*	*	*	*	*	*	0.5	0.9	3.7
2002.....	*	*	*	*	*	*	*	*	0.8	3.5
2001.....	*	*	*	*	*	*	*	0.3	0.7	3.5
2000.....	*	*	*	*	*	*	0.3	0.6	0.9	4.7
1999.....	*	*	*	*	*	*	*	0.6	1.3	6.3

White, male

2006 4/.....	*	*	*	*	*	*	*	*	*	1.9
2005 4/.....	*	*	*	*	*	*	*	*	1.4	3.3
2004 4/.....	*	*	*	*	*	*	*	*	*	2.6
2003 4/.....	*	*	*	*	*	*	*	*	*	2.7
2002.....	*	*	*	*	*	*	*	*	*	2.9
2001.....	*	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	*	*	*	*	0.8	*	3.8
1999.....	*	*	*	*	*	*	*	0.8	1.6	4.0

White, female

2006 4/.....	*	*	*	*	*	*	*	*	0.8	2.3
2005 4/.....	*	*	*	*	*	*	*	*	*	3.6
2004 4/.....	*	*	*	*	*	*	*	*	0.7	2.8
2003 4/.....	*	*	*	*	*	*	*	*	0.8	4.2
2002.....	*	*	*	*	*	*	*	*	0.9	3.7
2001.....	*	*	*	*	*	*	*	*	0.8	3.9
2000.....	*	*	*	*	*	*	*	*	1.0	5.1
1999.....	*	*	*	*	*	*	*	0.5	1.1	7.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute bronchitis and bronchiolitis (J20-J21)

All races 3/, both sexes

2006.....	*	*	*	*	*	*	*	0.4	1.1
2005.....	*	*	*	*	*	*	0.3	0.5	1.9
2004.....	*	*	*	*	*	*	*	*	1.4
2003.....	*	*	*	*	*	*	*	0.5	2.1
2002.....	*	*	*	*	*	*	*	0.5	2.0
2001.....	*	*	*	*	*	*	*	0.4	2.1
2000.....	*	*	*	*	*	*	0.4	0.5	2.6
1999.....	*	*	*	*	*	*	0.4	0.8	3.6

All races 3/, male

2006.....	*	*	*	*	*	*	*	*	1.2
2005.....	*	*	*	*	*	*	*	*	1.7
2004.....	*	*	*	*	*	*	*	*	1.8
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	2.4
1999.....	*	*	*	*	*	*	*	*	2.6

All races 3/, female

2006.....	*	*	*	*	*	*	*	*	1.1
2005.....	*	*	*	*	*	*	*	*	2.0
2004.....	*	*	*	*	*	*	*	*	1.3
2003.....	*	*	*	*	*	*	*	*	2.5
2002.....	*	*	*	*	*	*	*	*	2.2
2001.....	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	*	*	*	*	*	2.7
1999.....	*	*	*	*	*	*	*	0.7	4.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute bronchitis and bronchiolitis (J20-J21)

White, both sexes

2006 4/.....	*	*	*	*	*	*	*	*	0.5	1.2
2005 4/.....	*	*	*	*	*	*	*	0.3	0.6	2.0
2004 4/.....	*	*	*	*	*	*	*	*	*	1.5
2003 4/.....	*	*	*	*	*	*	*	*	0.5	2.3
2002.....	*	*	*	*	*	*	*	*	0.5	2.1
2001.....	*	*	*	*	*	*	*	*	0.5	2.1
2000.....	*	*	*	*	*	*	*	0.4	0.5	2.7
1999.....	*	*	*	*	*	*	*	0.4	0.9	3.8

White, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	1.7
2004 4/.....	*	*	*	*	*	*	*	*	*	1.8
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	2.4
1999.....	*	*	*	*	*	*	*	*	*	2.6

White, female

2006 4/.....	*	*	*	*	*	*	*	*	*	1.1
2005 4/.....	*	*	*	*	*	*	*	*	*	2.1
2004 4/.....	*	*	*	*	*	*	*	*	*	1.3
2003 4/.....	*	*	*	*	*	*	*	*	*	2.7
2002.....	*	*	*	*	*	*	*	*	*	2.3
2001.....	*	*	*	*	*	*	*	*	*	2.5
2000.....	*	*	*	*	*	*	*	*	*	2.9
1999.....	*	*	*	*	*	*	*	*	0.7	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic lower respiratory diseases (J40-J47)

All races 3/, both sexes

2006.....	2.8	5.8	12.7	25.8	57.4	111.8	194.8	308.5	434.8	589.1
2005.....	2.8	6.5	12.6	26.9	62.1	118.6	210.3	330.6	457.8	637.2
2004.....	2.6	5.6	11.7	25.5	60.0	113.7	200.6	315.7	434.7	601.7
2003.....	2.8	5.9	11.9	27.5	63.9	121.1	210.8	330.0	455.9	635.1
2002.....	2.9	5.8	12.0	28.2	60.8	119.0	211.5	330.1	465.7	637.6
2001.....	2.7	5.4	11.9	29.4	62.9	124.2	215.4	329.2	452.9	644.7
2000.....	2.8	5.8	11.8	29.6	62.5	124.1	218.2	327.8	473.4	648.6
1999.....	2.6	5.5	12.0	31.9	66.6	130.4	227.6	340.2	484.5	646.0

All races 3/, male

2006.....	2.6	5.5	13.4	27.4	62.0	119.8	218.0	364.6	541.3	760.8
2005.....	2.7	6.4	13.6	28.5	67.6	132.9	240.0	393.6	564.7	819.9
2004.....	2.6	5.6	12.5	26.9	64.6	126.1	228.2	378.8	543.4	811.1
2003.....	2.5	6.3	12.3	29.4	69.2	132.6	244.0	395.8	569.1	865.9
2002.....	3.0	5.9	12.8	29.5	65.9	131.5	246.0	405.6	595.9	894.8
2001.....	2.5	5.8	12.3	31.4	67.0	137.1	252.9	406.7	583.5	916.9
2000.....	2.6	6.2	12.1	31.8	68.0	139.7	257.8	404.2	628.5	967.9
1999.....	2.4	5.7	12.1	34.3	74.6	151.0	278.3	432.4	648.3	996.7

All races 3/, female

2006.....	2.9	6.1	12.1	24.2	53.3	104.8	175.8	267.5	368.3	508.7
2005.....	2.8	6.6	11.7	25.5	57.1	106.1	186.2	284.8	391.5	553.3
2004.....	2.7	5.5	10.9	24.2	55.8	102.9	178.4	270.4	367.7	507.4
2003.....	3.1	5.4	11.5	25.7	59.0	111.2	184.2	283.1	386.7	533.0
2002.....	2.8	5.6	11.2	26.9	56.1	108.3	183.8	276.8	386.5	526.0
2001.....	2.9	5.1	11.5	27.5	59.1	113.1	185.6	275.0	374.6	530.8
2000.....	2.9	5.5	11.5	27.5	57.5	110.7	187.0	274.6	381.9	518.5
1999.....	2.9	5.3	11.9	29.7	59.3	112.8	188.0	276.1	388.7	504.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic lower respiratory diseases (J40-J47)

White, both sexes

2006 4/.....	2.5	5.8	12.7	26.7	60.4	119.4	209.5	326.4	457.6	618.3
2005 4/.....	2.5	6.2	12.4	27.7	65.6	126.2	224.5	349.4	481.5	667.6
2004 4/.....	2.4	5.2	11.6	26.1	63.2	121.7	214.3	333.3	457.6	630.1
2003 4/.....	2.5	5.6	11.7	28.0	67.3	129.5	223.2	347.2	478.9	664.5
2002.....	2.5	5.5	11.7	28.9	63.8	126.8	223.3	346.9	486.5	666.2
2001.....	2.4	5.1	11.7	30.2	66.2	131.3	227.1	346.1	473.5	671.8
2000.....	2.3	5.3	11.4	30.1	65.5	130.5	229.6	343.1	495.5	675.6
1999.....	2.2	4.9	11.8	32.6	69.6	137.2	237.9	353.4	504.9	672.9

White, male

2006 4/.....	2.3	5.3	13.1	27.9	63.3	125.0	229.8	378.5	559.5	788.6
2005 4/.....	2.4	6.0	13.2	28.5	70.2	138.6	251.2	406.6	583.5	846.1
2004 4/.....	2.4	5.3	12.3	26.6	66.5	132.4	238.5	392.8	563.4	837.7
2003 4/.....	2.2	6.1	12.0	29.4	71.5	139.3	252.4	406.9	585.6	894.4
2002.....	2.5	5.6	12.4	29.6	67.5	137.1	254.3	417.4	612.9	923.4
2001.....	2.2	5.6	11.9	31.8	69.1	142.0	260.8	418.9	600.0	943.6
2000.....	2.2	5.7	11.5	31.8	69.8	143.1	265.8	415.5	648.5	997.4
1999.....	2.0	5.0	11.6	34.5	76.3	155.7	285.6	441.3	664.4	1,029.4

White, female

2006 4/.....	2.6	6.2	12.3	25.5	57.6	114.5	192.6	287.6	393.2	538.4
2005 4/.....	2.6	6.3	11.7	26.9	61.4	115.1	202.4	307.0	417.6	585.4
2004 4/.....	2.5	5.1	11.0	25.6	60.1	112.2	194.4	289.7	391.9	536.5
2003 4/.....	2.8	5.1	11.4	26.7	63.5	120.8	199.3	304.0	413.2	562.8
2002.....	2.5	5.4	11.0	28.3	60.4	117.6	198.0	296.4	408.9	554.5
2001.....	2.6	4.6	11.6	28.6	63.5	122.0	199.7	294.5	397.1	557.9
2000.....	2.4	4.9	11.3	28.4	61.5	119.4	200.5	292.1	404.4	544.8
1999.....	2.5	4.7	12.0	30.8	63.5	121.1	200.0	291.6	410.7	529.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic lower respiratory diseases (J40-J47)

Black, both sexes

2006 4/.....	5.2	7.7	17.1	27.7	51.9	83.5	122.6	199.9	262.7	331.8
2005 4/.....	5.0	10.5	17.9	30.4	52.4	92.0	142.8	220.8	271.1	366.7
2004 4/.....	4.8	9.6	16.3	27.9	52.1	81.7	128.1	204.6	249.3	333.9
2003 4/.....	5.7	9.1	16.7	31.3	52.7	86.5	146.7	211.9	271.9	353.0
2002.....	6.5	9.1	17.4	30.9	53.4	86.1	148.1	215.9	298.5	368.2
2001.....	5.1	9.1	16.6	31.6	52.2	94.3	149.6	206.9	276.5	375.8
2000.....	6.1	11.0	18.2	33.0	57.4	97.3	151.2	211.6	271.3	373.8
1999.....	6.1	11.2	17.2	36.1	57.6	100.0	163.3	242.4	295.3	376.2

Black, male

2006 4/.....	5.0	7.8	19.9	33.6	70.2	112.0	169.2	287.7	417.6	512.0
2005 4/.....	4.9	11.0	20.1	37.5	66.0	126.0	198.9	345.5	422.8	594.9
2004 4/.....	4.9	9.9	18.9	37.7	67.6	109.0	182.3	300.7	392.4	566.3
2003 4/.....	5.1	9.8	17.7	38.0	67.7	112.0	212.5	326.8	471.8	601.6
2002.....	6.9	9.9	19.9	38.1	71.1	116.2	214.0	340.8	477.6	645.4
2001.....	5.1	8.9	18.8	37.1	66.3	127.1	222.2	329.9	468.7	671.7
2000.....	5.9	11.4	19.6	39.5	73.5	138.4	226.3	326.5	448.5	693.1
1999.....	6.2	11.9	19.7	43.2	79.3	139.4	243.5	387.0	521.4	701.9

Black, female

2006 4/.....	5.4	7.5	14.7	22.8	37.7	62.5	90.7	147.3	180.9	258.1
2005 4/.....	5.1	10.1	16.1	24.6	41.8	66.8	104.3	146.4	191.0	274.6
2004 4/.....	4.8	9.3	14.2	19.9	40.1	61.3	91.4	147.7	173.9	241.8
2003 4/.....	6.2	8.6	15.8	25.7	41.1	67.6	102.4	143.5	167.1	256.4
2002.....	6.1	8.5	15.3	25.1	39.6	63.8	104.0	141.3	205.3	262.0
2001.....	5.1	9.2	14.7	27.1	41.1	70.0	101.3	133.4	178.7	266.1
2000.....	6.2	10.6	16.9	27.7	44.6	67.0	101.2	142.4	182.1	255.0
1999.....	6.1	10.7	15.0	30.2	40.6	70.8	110.0	155.3	183.2	254.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Bronchitis, chronic and unspecified (J40-J42)

All races 3/, both sexes

2006.....	*	*	*	0.1	0.2	0.3	0.6	1.1	2.1	6.0
2005.....	*	*	*	0.2	0.3	0.5	0.9	1.1	2.2	7.4
2004.....	*	*	0.1	0.2	0.3	0.4	0.7	1.2	2.0	6.5
2003.....	*	*	*	0.1	0.2	0.4	0.7	1.4	2.6	7.3
2002.....	*	*	0.1	0.1	0.3	0.5	1.0	1.8	2.9	8.1
2001.....	*	0.1	0.1	0.2	0.3	0.6	0.9	1.7	2.6	9.1
2000.....	*	*	*	0.1	0.4	0.7	1.0	2.1	3.6	11.8
1999.....	*	*	0.1	0.3	0.6	0.7	1.2	1.9	4.1	10.7

All races 3/, male

2006.....	*	*	*	*	*	*	0.6	1.1	2.2	5.9
2005.....	*	*	*	*	0.4	*	0.9	1.4	2.5	7.3
2004.....	*	*	*	*	*	0.5	0.8	1.2	2.2	7.0
2003.....	*	*	*	*	*	*	0.8	1.9	2.6	7.0
2002.....	*	*	*	*	*	0.7	1.1	2.2	2.9	8.8
2001.....	*	*	*	*	0.4	0.8	1.1	2.2	2.5	9.2
2000.....	*	*	*	*	0.4	0.9	1.0	2.8	4.9	12.9
1999.....	*	*	*	0.4	0.5	0.8	1.5	2.4	4.7	11.9

All races 3/, female

2006.....	*	*	*	*	*	*	0.6	1.1	2.0	6.0
2005.....	*	*	*	0.2	*	0.5	0.8	0.8	2.0	7.4
2004.....	*	*	*	*	*	*	0.6	1.2	1.9	6.3
2003.....	*	*	*	*	*	0.4	0.7	1.1	2.5	7.5
2002.....	*	*	*	*	*	*	0.9	1.4	2.9	7.8
2001.....	*	*	*	*	*	0.4	0.7	1.4	2.6	9.1
2000.....	*	*	*	*	0.4	0.5	1.1	1.6	2.8	11.4
1999.....	*	*	*	*	0.6	0.5	1.0	1.5	3.8	10.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Bronchitis, chronic and unspecified (J40-J42)

White, both sexes

2006 4/.....	*	*	*	*	0.2	0.4	0.7	1.1	2.1	6.2
2005 4/.....	*	*	*	0.1	0.3	0.5	0.9	1.1	2.2	7.9
2004 4/.....	*	*	*	0.2	0.3	0.4	0.7	1.2	2.1	6.7
2003 4/.....	*	*	*	*	0.2	0.5	0.7	1.5	2.6	7.5
2002.....	*	*	*	*	0.3	0.6	1.1	1.9	3.0	8.6
2001.....	*	*	0.1	0.2	0.3	0.6	0.9	1.8	2.7	9.6
2000.....	*	*	*	*	0.5	0.7	1.0	2.2	3.7	12.7
1999.....	*	*	*	0.3	0.6	0.7	1.2	1.9	4.1	11.2

White, male

2006 4/.....	*	*	*	*	*	*	0.6	1.0	2.2	6.3
2005 4/.....	*	*	*	*	0.4	*	0.9	1.4	2.5	7.4
2004 4/.....	*	*	*	*	*	0.6	0.8	1.2	2.4	7.1
2003 4/.....	*	*	*	*	*	*	0.7	1.9	2.6	7.3
2002.....	*	*	*	*	*	0.7	1.2	2.4	2.9	9.3
2001.....	*	*	*	*	*	0.8	1.0	2.2	2.5	9.4
2000.....	*	*	*	*	0.5	1.0	1.0	2.8	5.2	13.8
1999.....	*	*	*	0.4	0.5	0.9	1.5	2.4	4.4	12.4

White, female

2006 4/.....	*	*	*	*	*	*	0.7	1.2	2.0	6.2
2005 4/.....	*	*	*	*	*	0.6	0.9	0.9	2.1	8.1
2004 4/.....	*	*	*	*	*	*	0.6	1.2	2.0	6.5
2003 4/.....	*	*	*	*	*	0.5	0.8	1.2	2.6	7.6
2002.....	*	*	*	*	*	*	1.0	1.6	3.0	8.3
2001.....	*	*	*	*	*	*	0.7	1.5	2.8	9.7
2000.....	*	*	*	*	0.5	0.5	1.1	1.7	2.8	12.2
1999.....	*	*	*	*	0.7	0.5	1.0	1.5	3.9	10.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Emphysema (J43)

All races 3/, both sexes

2006.....	0.3	0.7	1.4	3.2	7.1	13.2	22.3	33.1	41.6	45.3
2005.....	0.2	0.8	1.4	3.7	8.0	15.3	25.8	37.8	46.0	51.9
2004.....	0.2	0.6	1.5	3.4	8.1	15.8	26.1	37.5	45.7	49.9
2003.....	0.3	0.7	1.5	4.0	9.3	16.6	28.5	41.9	51.6	54.6
2002.....	0.3	0.7	1.6	4.3	9.1	17.4	31.3	43.2	54.7	59.3
2001.....	0.2	0.7	1.7	5.0	10.1	19.8	33.2	46.9	55.5	62.1
2000.....	0.3	0.8	1.8	5.0	10.4	20.7	35.5	47.5	60.4	63.3
1999.....	0.4	0.7	2.0	5.5	12.1	23.5	38.4	51.3	65.3	62.1

All races 3/, male

2006.....	0.4	0.7	1.9	3.5	7.9	15.1	25.2	40.3	52.8	63.7
2005.....	0.3	1.1	1.6	4.4	9.7	18.4	30.7	47.5	59.5	71.5
2004.....	0.3	0.8	1.6	3.8	9.5	18.4	31.6	46.5	60.5	74.4
2003.....	0.3	1.0	1.9	4.4	10.3	18.3	34.4	51.9	67.5	79.2
2002.....	0.5	0.9	1.9	4.6	10.7	19.7	37.8	56.1	74.9	90.0
2001.....	0.3	0.9	2.0	5.8	10.8	22.4	40.3	59.2	74.9	101.2
2000.....	0.3	1.0	2.1	5.6	11.2	24.5	42.6	60.4	83.6	104.2
1999.....	0.4	0.9	2.2	6.2	13.6	28.0	48.9	67.3	91.5	105.4

All races 3/, female

2006.....	0.3	0.7	1.0	2.8	6.4	11.5	19.9	27.8	34.6	36.7
2005.....	*	0.5	1.1	3.0	6.5	12.6	21.9	30.8	37.6	42.8
2004.....	0.2	0.4	1.3	3.1	6.8	13.5	21.7	31.1	36.6	38.9
2003.....	0.3	0.5	1.2	3.6	8.4	15.2	23.7	34.8	41.9	43.7
2002.....	0.2	0.6	1.3	4.0	7.7	15.4	26.1	34.0	42.4	46.0
2001.....	*	0.5	1.4	4.2	9.5	17.6	27.5	38.3	43.8	45.8
2000.....	0.2	0.6	1.5	4.5	9.7	17.5	29.9	38.5	46.8	46.7
1999.....	0.3	0.5	1.7	4.9	10.8	19.8	30.2	40.2	49.9	44.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Emphysema (J43)

White, both sexes

2006 4/.....	0.3	0.7	1.5	3.4	7.5	14.2	24.3	35.2	43.9	47.8
2005 4/.....	0.2	0.8	1.3	3.9	8.6	16.4	27.9	39.9	48.2	54.5
2004 4/.....	0.2	0.6	1.4	3.6	8.6	17.2	28.2	40.1	48.4	52.5
2003 4/.....	0.3	0.7	1.6	4.2	9.9	18.0	30.5	44.3	54.2	57.0
2002.....	0.3	0.7	1.7	4.6	9.8	18.6	33.3	45.8	57.4	62.3
2001.....	0.2	0.7	1.7	5.1	10.9	21.5	35.2	49.8	58.7	65.1
2000.....	0.2	0.7	1.8	5.3	11.2	22.0	37.9	50.2	63.6	65.8
1999.....	0.3	0.6	2.0	5.8	13.0	25.4	40.9	53.8	68.8	64.7

White, male

2006 4/.....	0.4	0.7	1.9	3.6	7.9	15.8	27.1	41.7	54.6	66.5
2005 4/.....	0.3	1.1	1.5	4.5	10.1	19.2	32.4	49.3	60.6	73.6
2004 4/.....	0.3	0.8	1.6	3.8	9.8	19.4	33.4	48.5	63.2	77.1
2003 4/.....	0.3	1.0	1.9	4.5	10.6	19.5	35.9	53.2	69.6	80.4
2002.....	0.4	0.9	1.9	4.8	11.1	20.2	39.5	58.2	77.4	93.0
2001.....	0.3	1.0	1.9	5.8	11.3	23.9	41.6	61.7	78.0	105.1
2000.....	0.2	0.9	2.0	5.8	11.8	25.0	44.6	63.1	87.0	105.5
1999.....	0.3	0.8	2.1	6.4	14.2	29.4	50.6	69.6	94.4	109.2

White, female

2006 4/.....	0.3	0.7	1.2	3.2	7.1	12.7	21.9	30.3	37.1	39.1
2005 4/.....	*	0.5	1.2	3.3	7.2	13.8	24.2	33.0	40.5	45.7
2004 4/.....	*	0.4	1.3	3.4	7.5	15.2	24.0	33.9	39.2	41.4
2003 4/.....	0.3	0.5	1.4	3.9	9.2	16.7	26.0	37.8	44.7	46.6
2002.....	*	0.6	1.5	4.4	8.6	17.3	28.3	37.0	45.1	48.9
2001.....	*	0.5	1.4	4.5	10.6	19.4	30.0	41.3	47.0	48.4
2000.....	0.2	0.6	1.5	4.8	10.7	19.3	32.5	41.1	49.8	49.7
1999.....	0.3	0.5	1.9	5.3	11.9	21.9	33.1	42.6	53.7	46.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Emphysema (J43)

Black, both sexes

2006 4/.....	*	0.7	1.3	2.6	6.7	8.9	11.5	20.4	24.4	23.9
2005 4/.....	*	0.9	1.9	3.5	6.0	11.6	13.6	26.2	26.2	27.3
2004 4/.....	*	*	2.1	3.1	6.5	8.4	13.7	20.7	24.3	27.6
2003 4/.....	*	1.0	1.3	3.6	7.2	10.2	16.4	25.2	30.5	30.9
2002.....	0.8	1.0	1.5	3.3	5.9	11.9	20.9	23.5	34.8	32.1
2001.....	*	*	2.3	4.9	6.4	11.6	20.1	21.9	26.3	35.3
2000.....	*	1.6	2.5	4.7	7.7	14.9	19.5	24.5	30.8	37.2
1999.....	0.8	1.4	2.4	4.6	8.0	14.0	20.8	31.4	31.2	37.6

Black, male

2006 4/.....	*	*	2.5	3.9	11.0	13.4	14.5	32.2	41.1	40.2
2005 4/.....	*	*	2.8	5.6	9.8	17.8	20.2	39.8	45.6	51.1
2004 4/.....	*	*	3.0	4.9	9.7	13.8	21.4	34.6	39.5	56.6
2003 4/.....	*	*	2.0	5.2	10.2	13.3	24.6	43.4	50.2	70.9
2002.....	*	*	2.4	4.5	9.3	20.4	30.7	41.3	61.1	64.3
2001.....	*	*	2.9	6.9	9.6	15.8	31.6	34.4	49.0	68.1
2000.....	*	2.4	3.5	5.9	10.8	25.2	30.1	35.0	53.1	87.6
1999.....	*	2.0	4.1	6.4	11.9	22.8	37.0	48.9	65.4	72.6

Black, female

2006 4/.....	*	*	*	*	3.3	5.5	9.5	13.4	15.6	17.2
2005 4/.....	*	*	*	*	3.0	7.1	9.1	18.1	16.0	17.6
2004 4/.....	*	*	*	*	3.9	4.3	8.6	12.4	16.3	16.1
2003 4/.....	*	*	*	2.3	4.8	7.9	10.9	14.4	20.1	15.4
2002.....	*	*	*	*	3.3	5.6	14.3	12.9	21.1	19.7
2001.....	*	*	*	3.3	4.0	8.4	12.3	14.4	14.7	23.2
2000.....	*	*	*	3.6	5.2	7.3	12.4	18.2	19.5	18.5
1999.....	*	*	*	3.0	5.0	7.5	10.1	20.9	14.3	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Asthma (J45-J46)

All races 3/, both sexes

2006.....	1.0	1.2	1.4	1.4	1.8	2.0	2.7	3.6	6.4	12.3
2005.....	1.0	1.4	1.4	1.5	1.9	2.3	2.9	4.1	7.1	14.5
2004.....	0.9	1.3	1.4	1.7	1.9	2.5	3.0	4.2	6.6	14.1
2003.....	1.1	1.5	1.7	1.9	2.2	2.7	3.1	4.9	7.2	14.2
2002.....	1.2	1.4	1.6	1.9	2.1	2.7	3.8	5.4	7.7	14.7
2001.....	1.1	1.3	1.8	1.9	2.6	3.0	3.9	5.2	8.1	16.1
2000.....	1.2	1.6	1.7	2.0	2.5	3.4	4.6	5.7	8.7	16.7
1999.....	1.1	1.6	1.7	2.6	2.2	3.7	4.7	6.2	9.6	17.3

All races 3/, male

2006.....	0.8	0.9	1.1	1.0	1.2	1.4	2.1	2.9	4.4	9.1
2005.....	0.8	0.9	1.2	1.1	1.2	1.7	1.9	3.7	4.8	8.7
2004.....	0.8	1.1	1.0	1.4	1.3	2.0	2.3	2.8	4.4	11.1
2003.....	0.9	1.2	1.2	1.5	1.7	1.8	2.4	3.7	5.7	10.7
2002.....	0.9	1.1	1.3	1.5	1.7	2.0	2.8	4.3	6.2	10.9
2001.....	0.9	1.0	1.3	1.2	1.9	2.4	3.0	3.8	5.6	12.1
2000.....	1.0	1.3	1.3	1.6	1.8	2.7	3.4	3.9	6.7	12.9
1999.....	0.8	1.0	1.2	1.9	1.7	2.6	3.9	4.8	7.2	12.7

All races 3/, female

2006.....	1.2	1.6	1.7	1.7	2.3	2.6	3.2	4.2	7.6	13.9
2005.....	1.2	1.8	1.7	1.9	2.6	2.8	3.7	4.4	8.6	17.2
2004.....	1.1	1.5	1.8	2.0	2.5	2.9	3.6	5.1	7.9	15.4
2003.....	1.4	1.7	2.1	2.2	2.6	3.5	3.7	5.7	8.0	15.8
2002.....	1.4	1.8	1.9	2.3	2.5	3.3	4.5	6.2	8.7	16.3
2001.....	1.3	1.7	2.2	2.5	3.3	3.6	4.7	6.1	9.6	17.8
2000.....	1.4	1.9	2.1	2.4	3.1	4.0	5.5	6.9	9.8	18.2
1999.....	1.3	2.2	2.2	3.3	2.7	4.6	5.4	7.2	11.1	19.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Asthma (J45-J46)

White, both sexes

2006 4/.....	0.8	0.9	1.0	1.0	1.4	1.7	2.3	3.4	6.0	12.1
2005 4/.....	0.8	1.0	1.1	1.1	1.6	1.8	2.3	3.8	6.4	14.5
2004 4/.....	0.7	0.8	1.0	1.3	1.4	2.1	2.8	3.6	6.2	13.7
2003 4/.....	0.8	1.1	1.1	1.5	1.8	2.3	2.6	4.4	6.7	14.3
2002.....	0.8	1.1	1.1	1.4	1.8	2.2	3.2	5.1	7.5	14.7
2001.....	0.8	1.0	1.4	1.4	1.9	2.6	3.4	4.7	7.8	15.7
2000.....	0.9	1.1	1.2	1.4	2.0	2.8	4.0	5.2	8.4	15.9
1999.....	0.8	1.2	1.3	2.0	1.8	3.1	4.0	5.8	9.4	16.6

White, male

2006 4/.....	0.6	0.6	0.7	0.6	0.8	1.1	1.6	2.7	3.9	9.1
2005 4/.....	0.6	0.6	1.0	0.8	0.9	1.3	1.4	3.1	4.1	8.6
2004 4/.....	0.6	0.6	0.7	0.9	0.8	1.6	1.9	2.2	3.8	10.3
2003 4/.....	0.6	0.9	0.8	1.2	1.2	1.5	2.0	3.0	5.4	10.6
2002.....	0.6	0.8	0.9	1.0	1.4	1.5	2.3	3.8	5.8	10.7
2001.....	0.6	0.7	0.9	1.0	1.2	2.0	2.4	3.3	5.1	11.3
2000.....	0.7	1.0	1.0	1.0	1.4	2.0	3.0	3.7	6.3	11.4
1999.....	0.5	0.7	0.9	1.5	1.2	2.0	3.0	4.2	7.0	10.8

White, female

2006 4/.....	1.0	1.2	1.3	1.4	1.8	2.3	2.9	3.9	7.4	13.6
2005 4/.....	0.9	1.3	1.2	1.4	2.2	2.2	3.0	4.3	7.9	17.2
2004 4/.....	0.8	1.0	1.4	1.6	2.1	2.6	3.5	4.7	7.7	15.2
2003 4/.....	1.0	1.3	1.5	1.7	2.3	3.0	3.1	5.4	7.5	15.9
2002.....	1.0	1.3	1.3	1.7	2.2	2.8	3.9	6.1	8.6	16.4
2001.....	1.0	1.2	1.8	1.9	2.5	3.2	4.2	5.7	9.4	17.6
2000.....	1.0	1.2	1.5	1.8	2.6	3.4	4.8	6.3	9.7	17.7
1999.....	1.1	1.7	1.7	2.6	2.2	4.0	4.8	6.9	10.8	18.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Asthma (J45-J46)

Black, both sexes

2006 4/.....	2.7	3.6	4.3	4.4	6.1	4.9	6.3	5.0	7.3	12.5
2005 4/.....	2.8	4.4	3.9	5.4	5.3	6.0	8.7	7.9	12.6	11.8
2004 4/.....	2.9	4.8	4.1	5.1	6.3	5.7	5.6	9.0	9.4	15.5
2003 4/.....	3.4	4.1	5.7	5.2	5.0	6.6	7.5	8.2	10.4	12.0
2002.....	4.0	4.1	5.7	7.1	5.6	6.1	10.1	7.4	8.9	11.9
2001.....	3.1	4.3	4.9	5.9	7.9	6.3	9.4	10.2	11.1	16.3
2000.....	3.6	5.4	5.3	6.6	7.2	8.8	9.7	10.7	9.9	22.2
1999.....	3.3	4.8	5.8	7.3	6.2	8.4	11.1	9.8	11.3	20.4

Black, male

2006 4/.....	2.6	2.9	4.2	4.1	5.7	*	*	*	*	*
2005 4/.....	2.5	3.3	2.5	4.3	*	4.8	6.3	9.8	*	*
2004 4/.....	2.7	4.7	2.9	4.8	5.8	4.8	6.5	*	*	*
2003 4/.....	2.7	3.5	4.4	3.8	4.8	4.9	*	9.1	*	*
2002.....	3.7	3.4	4.6	6.0	5.3	6.2	8.7	*	*	*
2001.....	2.9	3.8	4.5	3.6	5.9	*	9.8	*	*	*
2000.....	3.1	4.0	3.5	6.3	6.0	8.6	7.0	*	*	25.4
1999.....	2.9	3.6	4.2	5.1	5.7	6.8	12.2	9.5	*	*

Black, female

2006 4/.....	2.7	4.2	4.4	4.6	6.5	5.5	6.8	5.9	8.3	15.0
2005 4/.....	3.1	5.3	5.1	6.4	6.7	6.9	10.4	6.7	13.0	13.5
2004 4/.....	3.1	4.9	5.1	5.4	6.7	6.3	5.1	9.7	9.1	16.9
2003 4/.....	4.1	4.7	6.7	6.3	5.1	7.9	8.6	7.6	11.6	13.8
2002.....	4.2	4.7	6.6	8.1	5.7	5.9	11.0	8.0	9.5	12.3
2001.....	3.3	4.8	5.3	7.8	9.4	7.5	9.2	11.4	11.9	16.8
2000.....	4.0	6.6	6.9	6.9	8.2	9.0	11.5	14.2	10.2	21.0
1999.....	3.6	5.9	7.1	9.1	6.5	9.6	10.4	10.0	13.8	21.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic lower respiratory diseases (J44,J47)

All races 3/, both sexes

2006.....	1.4	3.8	9.8	21.1	48.3	96.3	169.1	270.7	384.8	525.4
2005.....	1.5	4.3	9.7	21.5	51.8	100.6	180.7	287.5	402.5	563.5
2004.....	1.4	3.6	8.8	20.2	49.7	95.1	170.8	272.9	380.4	531.1
2003.....	1.4	3.6	8.6	21.5	52.2	101.4	178.5	281.8	394.6	558.9
2002.....	1.4	3.5	8.7	21.8	49.3	98.4	175.4	279.8	400.3	555.5
2001.....	1.3	3.3	8.4	22.4	49.8	100.7	177.4	275.4	386.8	557.4
2000.....	1.2	3.4	8.2	22.4	49.1	99.2	177.1	272.5	400.7	556.7
1999.....	1.2	3.1	8.2	23.5	51.6	102.5	183.2	280.8	405.5	555.8

All races 3/, male

2006.....	1.3	3.9	10.3	22.8	52.7	103.1	190.1	320.3	481.9	682.2
2005.....	1.4	4.3	10.7	22.8	56.3	112.4	206.5	341.0	497.9	732.4
2004.....	1.5	3.6	9.8	21.5	53.5	105.2	193.5	328.2	476.3	718.6
2003.....	1.3	4.1	9.1	23.3	57.0	112.2	206.5	338.3	493.3	769.0
2002.....	1.6	3.8	9.5	23.3	53.2	109.1	204.3	343.0	511.9	785.1
2001.....	1.3	3.8	9.0	24.2	54.0	111.6	208.6	341.5	500.4	794.5
2000.....	1.2	3.8	8.7	24.4	54.5	111.6	210.9	337.1	533.3	838.0
1999.....	1.2	3.7	8.6	25.8	58.7	119.6	223.9	357.9	544.9	866.7

All races 3/, female

2006.....	1.4	3.8	9.3	19.6	44.3	90.3	152.0	234.3	324.2	452.1
2005.....	1.5	4.3	8.8	20.4	47.7	90.2	159.8	248.7	343.2	485.9
2004.....	1.4	3.6	7.8	18.9	46.2	86.2	152.5	233.1	321.3	446.8
2003.....	1.4	3.2	8.0	19.8	47.8	92.1	156.0	241.5	334.2	466.0
2002.....	1.3	3.3	7.9	20.4	45.7	89.3	152.3	235.2	332.4	455.8
2001.....	1.4	2.8	7.8	20.7	46.1	91.4	152.7	229.1	318.6	458.2
2000.....	1.2	2.9	7.8	20.5	44.3	88.7	150.4	227.6	322.5	442.2
1999.....	1.1	2.6	7.8	21.4	45.2	87.9	151.4	227.2	324.0	430.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic lower respiratory diseases (J44,J47)

White, both sexes

2006 4/.....	1.3	4.1	10.1	22.2	51.3	103.2	182.3	286.7	405.6	552.1
2005 4/.....	1.5	4.4	9.9	22.5	55.1	107.6	193.4	304.5	424.6	590.7
2004 4/.....	1.5	3.7	9.1	21.1	52.9	102.0	182.6	288.3	400.9	557.2
2003 4/.....	1.3	3.8	8.9	22.3	55.5	108.7	189.4	297.0	415.4	585.8
2002.....	1.4	3.6	8.8	22.9	51.9	105.4	185.7	294.0	418.6	580.7
2001.....	1.4	3.3	8.6	23.4	53.1	106.6	187.6	289.8	404.4	581.3
2000.....	1.2	3.4	8.4	23.2	51.8	105.0	186.7	285.5	419.7	581.2
1999.....	1.1	3.0	8.4	24.4	54.2	108.1	191.8	292.0	422.6	580.3

White, male

2006 4/.....	1.3	4.0	10.5	23.5	54.4	107.8	200.4	333.1	498.7	706.7
2005 4/.....	1.4	4.3	10.6	23.1	58.7	117.8	216.4	352.8	516.3	756.5
2004 4/.....	1.5	3.7	9.9	21.7	55.6	110.7	202.4	340.9	494.0	743.2
2003 4/.....	1.2	4.2	9.3	23.6	59.4	117.8	213.8	348.8	507.9	796.0
2002.....	1.6	3.8	9.5	23.7	54.7	114.7	211.4	353.0	526.8	810.4
2001.....	1.3	3.9	9.0	24.8	56.2	115.4	215.7	351.7	514.4	817.8
2000.....	1.2	3.8	8.5	24.9	56.2	115.1	217.3	345.8	550.1	866.7
1999.....	1.1	3.5	8.6	26.2	60.3	123.4	230.5	365.1	558.5	897.0

White, female

2006 4/.....	1.3	4.2	9.7	20.9	48.4	99.1	167.2	252.1	346.8	479.5
2005 4/.....	1.5	4.5	9.2	22.0	51.8	98.5	174.4	268.8	367.2	514.4
2004 4/.....	1.4	3.6	8.2	20.5	50.3	94.2	166.3	249.9	343.0	473.4
2003 4/.....	1.5	3.3	8.5	21.0	51.8	100.6	169.5	259.5	358.4	492.7
2002.....	1.3	3.5	8.2	22.1	49.3	97.1	164.8	251.8	352.3	480.9
2001.....	1.4	2.8	8.2	22.1	50.2	98.9	164.8	246.0	337.9	482.3
2000.....	1.1	3.0	8.2	21.6	47.7	96.1	162.1	243.0	342.1	465.2
1999.....	1.1	2.6	8.3	22.7	48.7	94.7	161.1	240.6	342.4	453.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic lower respiratory diseases (J44,J47)

Black, both sexes

2006 4/.....	1.9	3.3	11.3	20.4	39.1	69.5	104.1	173.1	229.1	291.9
2005 4/.....	1.8	5.1	12.0	21.2	40.8	73.9	119.9	186.1	230.8	326.1
2004 4/.....	1.5	4.1	9.8	19.5	39.0	67.5	108.2	174.2	214.9	285.9
2003 4/.....	1.8	3.8	9.4	22.1	40.4	69.5	122.2	178.0	228.7	304.6
2002.....	1.7	3.9	10.1	20.3	41.8	67.8	116.9	184.3	252.5	320.1
2001.....	1.6	4.1	9.4	20.7	37.6	75.7	119.1	173.3	238.4	320.9
2000.....	1.9	3.9	10.1	21.7	42.2	72.8	120.7	174.7	227.9	310.6
1999.....	2.0	4.8	8.5	23.9	43.1	77.1	129.9	199.1	248.8	313.7

Black, male

2006 4/.....	1.7	3.8	13.1	25.1	53.4	94.4	148.3	249.7	369.0	463.8
2005 4/.....	1.7	6.3	14.7	27.3	52.4	102.7	171.4	294.1	363.1	532.6
2004 4/.....	1.5	3.9	12.3	28.1	51.6	90.1	153.8	257.5	342.9	492.7
2003 4/.....	1.9	4.5	11.0	28.8	52.4	93.8	181.3	273.0	410.5	520.3
2002.....	2.0	4.8	12.7	27.5	56.2	88.8	174.3	291.6	407.3	565.0
2001.....	1.6	4.4	11.3	26.6	50.4	105.7	179.4	285.3	408.5	583.2
2000.....	1.7	5.0	12.5	27.3	56.1	104.1	187.9	284.9	383.6	576.6
1999.....	2.5	6.0	11.0	31.2	61.3	109.6	192.3	325.8	441.2	607.0

Black, female

2006 4/.....	2.1	2.8	9.9	16.5	28.0	51.1	73.8	127.2	155.2	221.5
2005 4/.....	1.8	4.1	9.8	16.1	31.7	52.5	84.6	121.6	161.0	242.7
2004 4/.....	1.4	4.2	7.6	12.4	29.1	50.7	77.3	124.8	147.4	204.0
2003 4/.....	1.7	3.1	8.0	16.6	31.0	51.5	82.4	121.5	133.5	220.8
2002.....	1.4	3.1	8.0	14.4	30.5	52.2	78.4	120.2	171.9	226.3
2001.....	1.5	3.8	7.7	15.8	27.5	53.5	78.9	106.5	151.7	223.5
2000.....	1.9	2.8	8.0	17.0	31.2	49.7	76.1	108.2	149.4	211.6
1999.....	1.5	3.7	6.5	17.9	28.8	53.1	88.4	122.7	153.4	204.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonitis due to solids and liquids (J69)

All races 3/, both sexes

2006.....	0.5	0.8	1.2	2.0	3.7	6.3	12.8	26.5	58.7	155.8
2005.....	0.4	0.8	1.5	2.0	3.7	6.4	12.9	28.5	61.1	163.9
2004.....	0.5	0.8	1.2	1.9	3.4	6.0	13.7	28.4	62.2	164.9
2003.....	0.4	0.9	1.1	2.0	3.8	6.3	13.3	30.1	65.4	176.2
2002.....	0.5	0.7	1.1	1.8	3.4	6.5	13.4	31.3	67.3	186.0
2001.....	0.5	0.7	1.3	1.8	3.6	6.2	14.1	30.4	68.0	189.4
2000.....	0.5	0.8	1.2	1.9	3.2	6.7	14.2	30.4	65.7	187.6
1999.....	0.5	0.7	0.9	2.0	3.1	6.1	13.2	28.2	60.7	175.6

All races 3/, male

2006.....	0.6	1.0	1.6	2.5	4.9	8.7	17.5	37.3	85.7	224.8
2005.....	0.4	0.9	1.8	2.4	4.9	8.4	17.5	38.3	86.2	229.3
2004.....	0.6	1.0	1.5	2.3	4.5	8.1	18.7	39.0	87.4	237.2
2003.....	0.4	1.0	1.4	2.6	4.8	8.6	17.8	41.5	92.3	246.2
2002.....	0.7	0.9	1.5	2.2	4.2	8.6	18.2	43.4	95.1	268.5
2001.....	0.7	0.8	1.7	2.3	4.2	8.2	19.1	41.0	98.1	270.3
2000.....	0.6	1.0	1.5	2.4	3.9	9.2	20.2	42.4	95.2	276.0
1999.....	0.6	0.9	1.0	2.4	4.2	8.3	18.4	41.6	90.1	258.4

All races 3/, female

2006.....	0.4	0.7	0.9	1.5	2.6	4.3	8.9	18.6	41.9	123.5
2005.....	0.4	0.7	1.1	1.6	2.7	4.6	9.1	21.3	45.5	133.9
2004.....	0.3	0.5	0.8	1.5	2.5	4.2	9.8	20.9	46.6	132.4
2003.....	0.4	0.7	0.8	1.5	2.9	4.3	9.7	22.1	49.0	145.3
2002.....	0.4	0.5	0.8	1.4	2.8	4.8	9.6	22.8	50.4	150.2
2001.....	0.4	0.6	0.9	1.4	3.0	4.4	10.2	23.0	49.9	155.6
2000.....	0.4	0.6	0.9	1.5	2.5	4.6	9.4	22.1	48.2	151.6
1999.....	0.4	0.5	0.8	1.6	2.0	4.2	9.2	18.8	43.5	142.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonitis due to solids and liquids (J69)

White, both sexes

2006 4/.....	0.5	0.8	1.2	1.9	3.7	6.2	12.4	26.3	60.1	160.0
2005 4/.....	0.5	0.8	1.4	2.0	3.5	6.0	12.7	27.9	61.4	166.5
2004 4/.....	0.4	0.7	1.1	1.8	3.1	5.9	13.4	28.2	62.9	167.8
2003 4/.....	0.4	0.8	1.1	2.0	3.3	5.9	13.1	29.7	65.0	178.7
2002.....	0.5	0.7	1.0	1.8	3.3	6.1	13.1	30.9	68.1	189.5
2001.....	0.5	0.7	1.2	1.8	3.5	6.0	13.6	30.2	68.1	190.8
2000.....	0.5	0.7	1.1	1.9	2.9	6.4	13.4	30.1	65.7	191.0
1999.....	0.4	0.6	0.8	1.8	2.9	5.8	12.5	27.6	60.3	178.0

White, male

2006 4/.....	0.6	1.0	1.6	2.4	4.9	8.3	17.0	36.4	87.5	231.8
2005 4/.....	0.5	0.9	1.7	2.4	4.5	7.8	17.1	37.1	86.1	234.2
2004 4/.....	0.5	1.0	1.4	2.2	3.9	7.8	17.9	38.3	88.4	241.7
2003 4/.....	0.4	1.0	1.4	2.5	4.2	8.2	17.2	40.6	91.7	250.8
2002.....	0.6	0.8	1.3	2.1	4.1	8.0	17.6	42.5	96.5	274.6
2001.....	0.7	0.8	1.5	2.3	4.0	8.0	18.4	40.6	98.7	275.4
2000.....	0.7	0.9	1.3	2.3	3.6	8.5	18.7	41.4	96.2	284.0
1999.....	0.5	0.8	0.9	2.2	3.9	7.7	17.1	40.4	88.9	263.6

White, female

2006 4/.....	0.3	0.6	0.9	1.4	2.7	4.3	8.6	18.8	42.7	126.3
2005 4/.....	0.4	0.7	1.1	1.6	2.6	4.5	9.0	21.1	45.9	135.3
2004 4/.....	0.3	0.5	0.8	1.5	2.3	4.2	9.7	20.8	47.0	134.5
2003 4/.....	0.4	0.7	0.8	1.5	2.5	3.9	9.7	21.8	48.6	146.7
2002.....	0.4	0.5	0.7	1.5	2.6	4.5	9.5	22.6	50.7	152.5
2001.....	0.4	0.5	0.8	1.3	3.0	4.2	9.7	22.8	49.6	155.3
2000.....	0.4	0.5	0.9	1.4	2.3	4.6	9.1	22.2	47.5	153.2
1999.....	0.4	0.4	0.8	1.5	2.0	4.1	8.8	18.6	43.4	143.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Pneumonitis due to solids and liquids (J69)

Black, both sexes

2006 4/.....	0.9	1.2	1.7	2.9	4.9	9.2	19.8	32.8	55.1	128.4
2005 4/.....	*	1.2	2.4	3.1	6.4	10.4	17.0	40.9	69.2	153.1
2004 4/.....	0.9	1.0	1.7	3.0	7.3	8.8	20.4	34.7	64.9	144.7
2003 4/.....	0.9	1.3	1.2	2.7	8.5	11.0	17.2	39.8	76.2	163.4
2002.....	0.9	0.8	2.1	2.4	5.5	11.7	18.8	41.9	67.8	165.1
2001.....	0.7	1.0	2.4	2.4	6.0	9.3	22.3	38.6	76.1	185.5
2000.....	*	1.7	2.2	2.9	5.8	11.3	25.1	38.9	71.4	159.8
1999.....	1.0	1.2	1.8	3.9	5.0	10.1	23.7	40.5	71.1	167.1

Black, male

2006 4/.....	*	*	2.4	3.7	7.6	15.0	28.0	55.0	84.9	173.4
2005 4/.....	*	1.5	3.4	3.6	9.8	15.4	25.6	62.1	106.0	201.7
2004 4/.....	*	*	2.2	4.4	11.0	13.6	31.4	51.4	93.3	202.7
2003 4/.....	*	1.7	*	3.8	11.6	14.5	26.3	59.3	107.8	212.7
2002.....	*	*	3.1	3.7	6.3	16.7	28.4	62.5	91.7	233.7
2001.....	*	*	3.4	3.1	8.1	12.5	31.0	56.2	108.3	243.9
2000.....	*	2.4	3.1	3.3	7.2	17.4	41.1	62.4	96.1	209.9
1999.....	*	*	*	5.1	8.1	15.6	37.3	65.1	109.9	228.5

Black, female

2006 4/.....	*	*	*	2.3	2.8	4.8	14.2	19.6	39.4	110.0
2005 4/.....	*	*	*	2.7	3.7	6.7	11.0	28.3	49.7	133.5
2004 4/.....	*	*	*	*	4.4	5.2	13.0	24.8	49.9	121.7
2003 4/.....	*	*	*	*	6.0	8.4	11.1	28.2	59.6	144.2
2002.....	*	*	*	*	4.8	8.0	12.3	29.6	55.3	138.8
2001.....	*	*	*	*	4.3	6.9	16.6	28.0	59.7	163.8
2000.....	*	*	*	*	4.6	6.7	14.4	24.8	59.0	141.2
1999.....	*	*	*	2.9	*	6.0	14.6	25.7	51.9	144.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races 3/, both sexes

2006.....	1.5	2.3	4.0	6.7	11.9	21.4	36.6	59.1	91.9	136.6
2005.....	1.4	2.4	4.1	6.6	11.8	21.5	37.5	59.4	89.7	136.5
2004.....	1.5	2.6	4.0	7.3	12.4	21.7	39.5	61.4	91.3	139.4
2003.....	1.3	2.6	4.0	6.6	11.6	21.0	36.2	57.3	83.8	138.1
2002.....	1.5	2.5	4.1	7.0	11.9	20.6	37.6	56.8	83.7	133.5
2001.....	1.5	2.6	4.4	7.3	12.6	22.8	38.0	61.8	89.6	142.7
2000.....	1.7	2.6	4.6	7.8	12.7	22.9	36.9	60.5	89.2	141.7
1999.....	1.6	2.7	4.1	6.9	12.7	22.5	36.8	59.5	87.6	147.6

All races 3/, male

2006.....	1.5	2.4	4.3	7.7	14.0	26.0	46.3	75.7	118.0	163.7
2005.....	1.4	2.7	4.4	7.3	14.1	25.9	46.0	74.5	113.9	157.7
2004.....	1.5	2.7	4.5	8.1	14.3	26.6	49.4	77.3	118.1	168.8
2003.....	1.4	2.8	4.3	7.5	13.8	25.9	43.8	71.8	105.5	165.0
2002.....	1.6	2.5	4.3	7.9	13.9	24.4	47.8	70.5	104.2	156.6
2001.....	1.8	2.9	4.9	7.7	15.0	27.1	47.0	76.2	110.0	169.1
2000.....	1.8	2.8	5.0	8.8	14.6	26.8	45.7	77.3	112.4	179.7
1999.....	1.7	2.8	4.4	7.4	14.7	27.3	46.4	74.9	115.0	182.3

All races 3/, female

2006.....	1.5	2.3	3.8	5.7	10.0	17.3	28.7	47.0	75.6	124.0
2005.....	1.4	2.1	3.8	5.9	9.8	17.7	30.6	48.5	74.6	126.7
2004.....	1.5	2.6	3.4	6.6	10.7	17.4	31.6	50.0	74.8	126.1
2003.....	1.3	2.5	3.7	5.7	9.6	16.8	30.0	47.0	70.5	126.2
2002.....	1.4	2.4	3.9	6.2	10.2	17.3	29.4	47.0	71.1	123.4
2001.....	1.3	2.3	4.0	6.8	10.5	19.1	30.8	51.7	77.3	131.6
2000.....	1.6	2.4	4.2	6.8	10.8	19.6	29.9	48.8	75.6	126.2
1999.....	1.4	2.6	3.8	6.5	10.9	18.4	29.3	48.8	71.6	133.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

White, both sexes

2006 4/.....	1.3	2.1	3.7	6.3	11.7	21.7	38.0	61.3	95.2	141.3
2005 4/.....	1.3	2.1	3.7	6.2	11.6	21.6	38.4	60.7	92.8	140.2
2004 4/.....	1.3	2.2	3.6	7.1	12.2	22.0	40.9	63.6	94.1	144.1
2003 4/.....	1.1	2.2	3.7	6.3	11.5	21.5	37.4	58.5	87.4	143.2
2002.....	1.3	2.1	3.8	6.7	11.6	21.0	38.2	58.7	85.5	137.9
2001.....	1.3	2.3	3.9	7.0	12.4	23.0	38.7	63.5	92.3	146.0
2000.....	1.6	2.2	4.3	7.5	12.4	22.8	37.7	62.0	91.2	145.9
1999.....	1.3	2.3	3.8	6.4	12.6	22.6	37.0	60.6	89.5	150.7

White, male

2006 4/.....	1.3	2.2	4.0	7.6	13.7	26.5	47.7	78.4	122.1	168.5
2005 4/.....	1.2	2.4	4.0	6.8	13.6	26.3	47.2	75.5	117.0	160.0
2004 4/.....	1.3	2.2	4.3	7.8	14.2	27.0	50.8	79.7	121.5	174.3
2003 4/.....	1.2	2.5	4.1	7.1	13.7	26.4	45.2	72.7	110.0	170.5
2002.....	1.4	2.1	4.0	7.6	13.4	25.3	48.3	72.5	106.8	162.0
2001.....	1.7	2.5	4.4	7.6	14.5	27.5	48.3	78.4	113.3	173.5
2000.....	1.7	2.4	4.6	8.5	14.4	26.8	46.3	78.9	115.0	184.4
1999.....	1.5	2.4	4.2	6.7	14.7	27.7	47.0	76.5	117.6	185.7

White, female

2006 4/.....	1.4	1.9	3.3	5.0	9.8	17.3	29.9	48.5	78.2	128.6
2005 4/.....	1.3	1.8	3.3	5.7	9.7	17.4	31.1	49.8	77.5	131.1
2004 4/.....	1.4	2.2	3.0	6.3	10.4	17.5	32.7	51.8	77.2	130.5
2003 4/.....	1.0	2.0	3.3	5.4	9.6	17.2	30.9	48.2	73.6	131.0
2002.....	1.2	2.0	3.6	5.9	10.0	17.2	30.0	48.7	72.5	127.4
2001.....	1.0	2.0	3.4	6.4	10.4	19.1	31.0	52.9	79.6	134.4
2000.....	1.5	2.1	3.9	6.6	10.6	19.3	30.7	50.1	77.1	130.3
1999.....	1.1	2.2	3.5	6.2	10.7	18.2	29.1	49.5	72.9	136.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Black, both sexes

2006 4/.....	2.8	4.9	7.0	10.9	16.0	20.1	32.2	44.2	61.8	95.2
2005 4/.....	2.8	4.8	7.7	9.9	15.3	24.7	34.4	52.6	60.9	102.6
2004 4/.....	3.0	5.5	7.1	11.1	16.5	23.2	32.8	46.2	63.4	92.8
2003 4/.....	3.1	5.7	7.3	10.5	14.4	20.2	29.1	51.4	50.0	89.6
2002.....	2.9	5.5	6.9	11.1	16.8	20.7	33.9	40.5	68.3	90.3
2001.....	3.3	5.7	9.3	10.9	17.0	24.2	36.1	50.2	66.9	109.6
2000.....	3.0	6.0	7.8	11.6	15.5	28.1	33.4	49.8	71.2	98.2
1999.....	3.4	5.8	7.1	11.7	15.4	25.0	38.0	52.8	66.2	111.6

Black, male

2006 4/.....	2.9	4.8	6.6	10.2	19.8	24.1	44.0	57.3	80.1	116.2
2005 4/.....	2.8	5.3	8.4	12.1	20.0	28.3	40.8	70.2	82.2	132.0
2004 4/.....	3.3	6.3	7.2	12.8	18.7	29.3	41.4	61.4	81.1	115.2
2003 4/.....	2.5	5.5	7.3	12.6	18.5	25.6	37.8	70.8	65.0	113.6
2002.....	3.4	5.7	7.6	12.4	21.2	22.2	44.4	53.3	86.4	100.8
2001.....	3.2	6.2	9.4	10.5	21.6	28.0	39.4	60.4	85.1	129.3
2000.....	3.3	7.2	8.6	13.3	17.6	32.5	40.5	67.6	82.6	132.6
1999.....	3.3	6.6	7.8	13.7	17.8	28.3	44.1	66.5	83.7	145.3

Black, female

2006 4/.....	2.8	5.0	7.2	11.5	13.1	17.1	24.1	36.4	52.1	86.6
2005 4/.....	2.8	4.4	7.2	8.1	11.6	22.0	30.1	42.1	49.7	90.8
2004 4/.....	2.7	4.9	7.0	9.7	14.9	18.7	27.0	37.2	54.1	83.9
2003 4/.....	3.6	5.8	7.3	8.8	11.1	16.3	23.3	39.9	42.2	80.2
2002.....	2.3	5.3	6.2	10.0	13.3	19.6	26.9	32.9	58.9	86.3
2001.....	3.3	5.3	9.1	11.3	13.4	21.4	33.9	44.1	57.6	102.3
2000.....	2.6	5.0	7.1	10.3	13.9	24.8	28.6	39.0	65.4	85.4
1999.....	3.5	5.1	6.6	10.1	13.5	22.5	34.0	44.6	57.5	99.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Peptic ulcer (K25-K28)

All races 3/, both sexes

2006.....	0.2	0.5	0.8	1.0	1.5	2.1	3.5	5.3	9.6	20.7
2005.....	0.3	0.5	0.8	0.9	1.6	2.2	3.8	5.8	10.8	21.4
2004.....	0.3	0.5	0.8	1.0	1.7	2.5	3.9	6.8	11.6	24.4
2003.....	0.3	0.6	0.8	1.2	1.7	2.3	4.2	7.3	13.6	26.4
2002.....	0.3	0.6	0.8	1.4	1.7	2.7	4.7	8.0	13.2	28.3
2001.....	0.4	0.7	0.8	1.3	2.1	3.1	4.8	8.3	16.0	33.4
2000.....	0.3	0.6	1.0	1.5	2.2	3.1	5.1	8.7	15.6	35.6
1999.....	0.4	0.6	0.8	1.4	2.2	3.3	5.9	9.1	17.1	34.6

All races 3/, male

2006.....	*	0.8	1.0	1.4	1.9	2.6	4.0	6.3	11.0	22.0
2005.....	0.5	0.8	1.2	1.3	2.2	2.6	4.8	7.3	13.1	22.4
2004.....	0.3	0.7	1.0	1.2	2.2	2.9	4.5	8.3	12.9	24.6
2003.....	0.4	0.8	1.2	1.4	2.3	3.1	5.4	8.8	16.2	28.9
2002.....	0.4	0.7	1.2	1.8	2.1	3.2	6.1	9.3	16.7	31.6
2001.....	0.5	0.9	1.1	1.8	2.8	4.1	5.6	10.3	17.3	34.9
2000.....	0.4	0.9	1.5	2.1	2.9	3.9	6.9	11.2	17.8	40.7
1999.....	0.4	0.7	1.2	1.8	3.0	4.5	7.9	11.0	21.1	37.5

All races 3/, female

2006.....	0.2	0.3	0.5	0.6	1.1	1.7	3.1	4.6	8.8	20.1
2005.....	0.2	0.3	0.5	0.6	1.1	1.8	3.0	4.7	9.4	21.0
2004.....	0.2	0.4	0.5	0.7	1.3	2.2	3.4	5.8	10.8	24.3
2003.....	0.2	0.4	0.5	1.0	1.1	1.7	3.2	6.2	11.9	25.3
2002.....	0.3	0.4	0.5	1.0	1.4	2.2	3.6	7.0	11.0	26.9
2001.....	0.2	0.4	0.6	0.8	1.5	2.2	4.2	6.9	15.2	32.7
2000.....	0.2	0.4	0.5	0.9	1.6	2.5	3.6	6.9	14.4	33.5
1999.....	0.3	0.5	0.5	1.1	1.5	2.3	4.3	7.7	14.7	33.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Peptic ulcer (K25-K28)

White, both sexes

2006 4/.....	0.2	0.5	0.7	0.9	1.4	2.0	3.4	5.2	9.9	21.2
2005 4/.....	0.3	0.4	0.8	0.9	1.6	2.2	3.7	5.7	10.9	21.6
2004 4/.....	0.2	0.4	0.7	0.9	1.7	2.5	3.8	6.8	11.7	25.1
2003 4/.....	0.3	0.5	0.7	1.1	1.6	2.4	4.2	7.2	13.4	26.9
2002.....	0.2	0.5	0.8	1.3	1.6	2.6	4.8	7.9	13.6	29.1
2001.....	0.3	0.6	0.7	1.1	2.0	3.0	4.8	8.3	16.2	34.4
2000.....	0.3	0.6	0.9	1.4	2.2	3.0	5.1	8.5	16.0	36.3
1999.....	0.3	0.5	0.7	1.4	2.2	3.1	5.8	9.1	17.1	35.3

White, male

2006 4/.....	*	0.7	0.9	1.4	1.8	2.5	3.6	6.0	11.3	22.0
2005 4/.....	0.4	0.6	1.1	1.2	2.1	2.5	4.5	7.0	13.5	22.5
2004 4/.....	0.3	0.6	1.0	1.2	2.1	2.9	4.3	8.1	12.6	25.5
2003 4/.....	0.3	0.7	1.0	1.3	2.2	3.1	5.3	8.6	15.6	29.0
2002.....	0.3	0.6	1.0	1.6	2.0	3.1	6.0	9.2	17.1	32.5
2001.....	0.5	0.8	0.9	1.5	2.7	3.8	5.3	10.1	17.3	35.7
2000.....	0.4	0.8	1.3	2.0	2.9	3.8	6.7	10.9	17.8	40.7
1999.....	0.3	0.6	1.1	1.6	2.9	4.1	7.8	11.1	21.1	38.4

White, female

2006 4/.....	*	0.3	0.5	0.5	1.1	1.6	3.2	4.6	9.0	20.8
2005 4/.....	*	0.2	0.5	0.7	1.1	1.9	3.0	4.7	9.3	21.2
2004 4/.....	*	0.3	0.5	0.7	1.3	2.1	3.3	5.8	11.1	25.0
2003 4/.....	0.2	0.3	0.5	0.9	1.0	1.7	3.3	6.2	12.1	26.0
2002.....	*	0.4	0.5	1.0	1.3	2.1	3.8	7.0	11.5	27.6
2001.....	*	0.4	0.6	0.8	1.4	2.2	4.4	7.0	15.6	33.9
2000.....	*	0.3	0.5	0.9	1.6	2.4	3.8	6.9	14.9	34.5
1999.....	0.3	0.4	0.4	1.1	1.5	2.2	4.2	7.7	14.8	34.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Peptic ulcer (K25-K28)

Black, both sexes

2006 4/.....	*	1.0	1.6	1.7	2.6	2.9	4.8	7.0	7.3	14.8
2005 4/.....	0.7	1.1	1.4	1.2	2.2	2.7	5.4	5.8	9.7	17.4
2004 4/.....	*	1.3	1.3	1.6	2.6	3.5	5.1	6.6	9.4	16.1
2003 4/.....	*	1.2	1.8	2.1	2.4	2.1	4.5	8.7	14.5	16.9
2002.....	0.9	0.9	1.6	2.3	2.9	3.2	5.2	8.8	6.8	19.0
2001.....	*	1.1	1.6	2.5	3.0	3.6	5.5	7.8	11.1	20.9
2000.....	*	1.4	2.0	2.6	2.7	4.1	5.7	9.4	11.0	26.6
1999.....	1.0	1.2	1.8	2.4	3.2	5.0	6.5	7.7	16.2	25.5

Black, male

2006 4/.....	*	1.8	2.8	2.5	*	*	8.3	9.8	*	*
2005 4/.....	*	1.9	2.5	*	*	*	7.9	9.4	*	20.5
2004 4/.....	*	2.1	*	*	3.9	*	*	10.0	*	*
2003 4/.....	*	1.6	3.1	3.1	3.9	*	7.2	11.4	23.6	*
2002.....	*	*	2.8	3.6	4.2	*	8.0	11.9	*	*
2001.....	*	1.8	2.7	4.1	4.5	5.3	9.1	10.7	*	23.8
2000.....	*	*	3.3	3.5	*	*	10.0	12.8	*	31.1
1999.....	1.5	1.9	2.8	3.7	5.3	7.5	8.8	*	18.3	28.1

Black, female

2006 4/.....	*	*	*	*	*	*	*	5.4	*	13.9
2005 4/.....	*	*	*	*	*	*	*	*	10.4	16.1
2004 4/.....	*	*	*	*	*	3.6	4.4	*	7.6	16.5
2003 4/.....	*	*	*	*	*	*	*	7.1	9.7	15.8
2002.....	*	*	*	*	*	*	*	6.9	*	18.5
2001.....	*	*	*	*	*	*	*	6.1	9.0	19.8
2000.....	*	*	*	*	*	*	*	7.4	8.5	24.9
1999.....	*	*	*	*	*	*	5.0	6.9	15.1	24.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of appendix (K35-K38)

All races 3/, both sexes

2006.....	*	*	*	0.2	0.2	0.4	0.5	0.6	1.0	1.7
2005.....	*	*	0.1	0.2	0.2	0.4	0.4	0.8	0.9	2.1
2004.....	*	*	0.2	0.1	0.2	0.3	0.6	0.8	1.4	1.7
2003.....	*	0.1	0.1	*	0.2	0.3	0.4	0.8	1.4	1.7
2002.....	*	0.1	0.2	0.2	0.3	0.3	0.7	0.8	1.1	2.0
2001.....	*	0.1	*	0.2	0.2	0.4	0.6	0.5	1.0	2.1
2000.....	*	*	0.1	0.2	0.3	0.3	0.5	0.7	1.4	1.6
1999.....	*	0.1	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.5

All races 3/, male

2006.....	*	*	*	0.2	*	0.5	0.8	1.1	1.5	2.3
2005.....	*	*	*	0.3	*	0.6	0.7	0.9	1.3	2.8
2004.....	*	*	0.2	*	*	*	0.7	1.1	1.7	2.5
2003.....	*	*	*	*	*	0.4	0.7	0.9	2.3	2.4
2002.....	*	*	0.2	*	0.5	0.5	0.8	1.1	1.6	2.2
2001.....	*	*	*	*	*	0.5	0.9	0.7	1.6	2.8
2000.....	*	*	*	0.3	0.4	*	0.6	1.0	1.6	2.0
1999.....	*	*	*	*	*	*	0.5	0.9	*	2.9

All races 3/, female

2006.....	*	*	*	*	*	*	*	*	0.7	1.5
2005.....	*	*	*	*	*	*	*	0.7	0.7	1.8
2004.....	*	*	*	*	*	*	0.5	0.6	1.2	1.4
2003.....	*	*	*	*	*	*	*	0.6	0.9	1.3
2002.....	*	*	*	*	*	*	0.5	0.6	0.8	2.0
2001.....	*	*	*	*	*	*	0.4	*	0.7	1.8
2000.....	*	*	*	*	*	*	0.4	0.5	1.2	1.5
1999.....	*	*	*	*	*	*	*	*	1.1	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Diseases of appendix (K35-K38)

White, both sexes

2006 4/.....	*	*	*	0.2	*	0.4	0.5	0.6	1.0	1.8
2005 4/.....	*	*	*	0.2	0.2	0.4	0.4	0.7	1.0	2.2
2004 4/.....	*	*	0.1	0.2	0.2	0.3	0.6	0.8	1.3	1.8
2003 4/.....	*	*	*	*	*	0.3	0.4	0.7	1.5	1.7
2002.....	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.2	2.1
2001.....	*	*	*	*	*	0.3	0.7	0.5	0.9	2.1
2000.....	*	*	*	0.2	0.3	0.2	0.4	0.6	1.3	1.7
1999.....	*	*	0.2	*	*	0.3	0.4	0.6	1.0	1.5

White, male

2006 4/.....	*	*	*	*	*	0.6	0.7	1.0	1.4	2.3
2005 4/.....	*	*	*	*	*	0.6	0.7	0.9	1.4	2.9
2004 4/.....	*	*	*	*	*	*	0.7	1.1	1.7	2.3
2003 4/.....	*	*	*	*	*	*	0.7	0.8	2.4	2.5
2002.....	*	*	*	*	0.5	*	0.7	1.2	1.6	2.0
2001.....	*	*	*	*	*	*	1.0	0.7	1.5	2.5
2000.....	*	*	*	*	0.5	*	*	0.9	1.5	2.0
1999.....	*	*	*	*	*	*	*	0.9	*	3.1

White, female

2006 4/.....	*	*	*	*	*	*	*	*	0.7	1.5
2005 4/.....	*	*	*	*	*	*	*	0.6	0.7	1.9
2004 4/.....	*	*	*	*	*	*	0.5	0.6	1.0	1.6
2003 4/.....	*	*	*	*	*	*	*	0.6	0.9	1.3
2002.....	*	*	*	*	*	*	0.5	0.6	0.9	2.1
2001.....	*	*	*	*	*	*	*	*	*	1.9
2000.....	*	*	*	*	*	*	*	*	1.2	1.6
1999.....	*	*	*	*	*	*	*	*	1.1	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hernia (K40-K46)

All races 3/, both sexes

2006.....	0.1	0.1	0.3	0.5	0.7	0.9	1.7	2.8	5.4	12.1
2005.....	0.1	0.1	0.3	0.4	0.9	0.9	1.4	2.5	4.8	12.1
2004.....	0.1	0.2	0.2	0.4	0.5	0.9	1.8	2.9	5.3	11.0
2003.....	0.1	0.2	0.3	0.5	0.7	1.0	1.6	2.7	4.8	11.9
2002.....	0.1	0.2	0.2	0.4	0.6	0.9	1.6	2.8	5.3	12.8
2001.....	0.1	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.6
2000.....	*	0.1	0.3	0.4	0.6	0.9	1.4	3.0	4.9	13.1
1999.....	0.1	*	0.2	0.3	0.7	0.8	1.6	2.5	5.4	12.7

All races 3/, male

2006.....	*	*	0.3	0.4	0.8	0.8	1.9	2.9	5.7	13.5
2005.....	*	*	0.3	0.4	0.9	0.7	1.3	2.4	4.6	12.7
2004.....	*	0.2	0.3	0.4	0.5	0.8	2.2	3.2	4.7	11.3
2003.....	*	0.2	0.4	0.4	0.7	1.1	1.6	3.0	5.2	12.3
2002.....	*	*	*	0.4	0.5	0.9	1.6	2.7	5.2	11.7
2001.....	*	*	0.3	0.3	0.8	0.9	1.4	2.3	5.3	12.0
2000.....	*	*	0.4	0.3	0.6	0.9	1.4	3.0	5.7	12.6
1999.....	*	*	0.2	0.3	0.8	0.8	1.5	2.5	6.5	13.2

All races 3/, female

2006.....	*	*	0.3	0.5	0.7	1.0	1.5	2.7	5.1	11.4
2005.....	*	*	0.2	0.4	0.8	1.1	1.4	2.5	5.0	11.8
2004.....	*	0.2	0.2	0.4	0.5	1.0	1.4	2.7	5.6	10.8
2003.....	*	0.2	0.3	0.6	0.7	1.0	1.5	2.5	4.6	11.8
2002.....	*	*	0.3	0.4	0.7	1.0	1.5	2.9	5.3	13.3
2001.....	*	*	0.3	0.5	0.6	0.8	1.6	2.6	4.8	12.8
2000.....	*	*	0.3	0.4	0.6	0.9	1.5	3.0	4.4	13.3
1999.....	*	*	*	*	0.6	0.8	1.6	2.6	4.8	12.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Hernia (K40-K46)

White, both sexes

2006 4/.....	0.1	0.1	0.3	0.4	0.8	0.9	1.6	2.8	5.3	12.6
2005 4/.....	*	0.1	0.3	0.4	0.9	0.9	1.4	2.5	5.0	12.5
2004 4/.....	*	0.2	0.2	0.4	0.5	0.9	1.8	3.1	5.4	11.2
2003 4/.....	*	0.2	0.3	0.4	0.7	1.1	1.6	2.8	4.7	12.3
2002.....	*	0.1	0.2	0.4	0.6	0.9	1.6	3.0	5.4	13.4
2001.....	*	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.8
2000.....	*	0.1	0.3	0.4	0.6	0.8	1.5	3.0	5.0	13.3
1999.....	*	*	0.2	0.3	0.7	0.7	1.6	2.5	5.5	13.0

White, male

2006 4/.....	*	*	0.3	0.3	0.8	0.8	1.9	2.9	5.6	14.0
2005 4/.....	*	*	0.3	0.4	0.9	0.7	1.4	2.4	4.6	12.8
2004 4/.....	*	*	0.3	0.4	0.5	0.7	2.2	3.3	4.7	11.4
2003 4/.....	*	*	0.4	0.4	0.7	1.1	1.6	3.1	5.1	12.5
2002.....	*	*	*	0.4	0.5	0.8	1.6	2.9	5.4	11.9
2001.....	*	*	0.3	*	0.7	0.9	1.4	2.3	5.2	11.8
2000.....	*	*	0.3	*	0.6	0.8	1.5	3.1	5.7	12.7
1999.....	*	*	*	0.4	0.8	0.8	1.6	2.5	6.6	12.9

White, female

2006 4/.....	*	*	0.2	0.5	0.7	1.0	1.4	2.7	5.2	11.9
2005 4/.....	*	*	*	0.4	0.8	1.1	1.4	2.6	5.2	12.3
2004 4/.....	*	*	*	0.4	0.5	1.1	1.4	2.9	5.8	11.2
2003 4/.....	*	*	*	0.5	0.7	1.0	1.6	2.6	4.5	12.2
2002.....	*	*	*	0.4	0.6	0.9	1.6	3.0	5.4	14.0
2001.....	*	*	0.3	0.5	0.6	0.8	1.6	2.6	5.0	13.2
2000.....	*	*	0.3	0.5	0.6	0.8	1.5	2.9	4.6	13.6
1999.....	*	*	*	*	0.5	0.7	1.5	2.5	4.9	13.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic liver disease and cirrhosis (K70,K73-K74)

All races 3/, both sexes

2006.....	8.2	14.8	21.2	22.8	22.9	24.8	27.4	29.6	28.2	19.4
2005.....	8.6	15.0	20.7	23.0	24.1	25.4	29.3	29.8	27.9	19.7
2004.....	8.7	15.1	21.3	21.7	23.7	26.2	29.5	30.1	27.0	19.7
2003.....	9.3	15.9	21.1	22.0	24.4	28.0	31.2	31.0	28.5	20.1
2002.....	9.7	16.0	20.3	21.7	24.5	27.9	31.2	32.5	29.9	21.4
2001.....	9.8	17.1	20.1	21.1	24.7	28.9	31.1	31.3	28.7	22.2
2000.....	10.2	16.8	18.7	21.9	26.2	28.9	30.8	31.9	29.6	23.1
1999.....	9.9	16.4	18.6	21.4	26.5	29.7	31.6	32.6	30.8	23.2

All races 3/, male

2006.....	10.8	20.7	31.8	33.7	33.1	34.1	35.5	36.9	34.4	26.1
2005.....	11.5	21.3	31.3	34.5	35.3	33.9	38.1	37.2	35.1	27.6
2004.....	12.2	21.5	31.9	32.3	33.4	35.7	38.7	38.6	34.0	25.5
2003.....	12.8	23.4	32.3	32.4	35.6	38.4	40.0	38.1	35.0	27.5
2002.....	12.9	23.3	31.0	32.0	34.5	37.2	40.0	39.4	36.9	28.4
2001.....	13.2	25.2	30.1	31.5	35.3	39.7	40.8	37.6	35.3	32.0
2000.....	14.4	25.4	28.6	31.7	37.7	39.3	40.7	40.7	34.6	34.3
1999.....	14.5	25.3	28.8	32.5	37.2	41.8	41.5	40.1	38.7	33.1

All races 3/, female

2006.....	5.6	9.0	11.1	12.5	13.6	16.7	20.9	24.2	24.3	16.2
2005.....	5.7	8.9	10.6	12.2	13.9	17.9	22.2	24.4	23.4	16.0
2004.....	5.4	9.0	11.1	11.8	14.9	18.0	22.1	24.0	22.7	17.1
2003.....	5.9	8.6	10.3	12.2	14.2	19.0	24.1	26.0	24.5	16.8
2002.....	6.5	8.9	10.1	12.1	15.3	19.8	24.1	27.6	25.6	18.4
2001.....	6.4	9.3	10.5	11.4	15.2	19.6	23.5	26.9	24.7	18.1
2000.....	6.1	8.4	9.1	12.7	15.7	19.9	23.1	25.8	26.6	18.5
1999.....	5.4	7.8	8.8	11.1	16.8	19.4	23.9	27.4	26.2	19.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic liver disease and cirrhosis (K70,K73-K74)

White, both sexes

2006 4/.....	8.5	15.5	21.9	23.3	23.6	25.7	29.1	30.8	29.6	20.2
2005 4/.....	9.0	15.4	21.0	23.5	24.2	26.5	30.9	30.8	28.8	20.2
2004 4/.....	9.0	15.6	21.6	21.6	24.0	27.1	31.1	31.8	28.1	20.3
2003 4/.....	9.7	16.2	21.0	22.0	24.6	28.9	32.8	32.9	29.8	20.9
2002.....	10.0	16.1	20.2	21.8	24.8	28.9	32.7	34.2	30.9	22.3
2001.....	9.8	17.2	19.6	20.7	25.3	29.7	32.3	32.6	30.0	23.2
2000.....	10.1	16.9	18.1	21.6	26.5	29.4	31.8	33.4	30.5	23.9
1999.....	9.7	15.8	18.3	20.8	26.3	30.2	32.6	33.7	32.0	23.8

White, male

2006 4/.....	11.2	21.7	32.6	34.4	34.0	34.9	37.3	38.0	36.3	27.2
2005 4/.....	12.1	21.8	31.9	35.0	35.2	35.0	39.8	38.1	35.9	28.5
2004 4/.....	12.6	22.0	32.5	32.0	33.7	36.4	40.3	40.5	35.4	26.7
2003 4/.....	13.4	24.0	32.0	32.2	35.9	39.5	41.8	40.1	36.2	28.4
2002.....	13.5	23.8	30.9	32.1	34.7	38.2	41.6	41.6	38.1	29.6
2001.....	13.2	25.5	29.3	30.6	35.6	40.2	42.1	39.1	37.2	33.5
2000.....	14.4	26.0	27.8	31.5	38.3	40.2	41.7	42.3	35.9	35.5
1999.....	14.2	24.6	28.4	31.5	37.4	42.2	42.5	41.1	40.3	34.2

White, female

2006 4/.....	5.8	9.3	11.3	12.7	13.9	17.5	22.2	25.4	25.4	16.9
2005 4/.....	5.9	9.1	10.3	12.4	13.9	19.0	23.5	25.5	24.3	16.4
2004 4/.....	5.4	9.2	11.0	11.6	15.0	18.8	23.5	25.5	23.6	17.4
2003 4/.....	5.9	8.4	10.2	12.2	14.2	19.6	25.4	27.8	25.9	17.5
2002.....	6.5	8.5	9.8	11.9	15.6	20.7	25.4	28.9	26.5	19.1
2001.....	6.3	8.9	10.0	11.3	15.8	20.4	24.3	28.0	25.7	18.9
2000.....	5.9	7.8	8.7	12.2	15.6	20.0	23.9	27.2	27.3	19.2
1999.....	5.1	7.2	8.5	10.7	16.2	19.8	24.7	28.4	27.2	19.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic liver disease and cirrhosis (K70,K73-K74)

Black, both sexes

2006 4/.....	6.5	11.0	20.8	23.1	20.4	21.0	16.2	19.4	12.1	10.6
2005 4/.....	6.2	14.2	21.6	24.6	25.4	20.0	20.4	19.9	18.0	12.3
2004 4/.....	7.7	14.4	23.0	25.7	24.7	22.6	20.3	13.9	15.1	12.1
2003 4/.....	7.7	15.0	25.6	26.4	25.0	25.4	21.7	14.1	14.7	12.3
2002.....	7.9	17.4	25.0	24.6	25.5	22.9	20.0	20.1	16.0	12.2
2001.....	10.6	19.2	26.8	27.7	23.6	26.9	22.6	21.0	14.6	10.4
2000.....	10.9	19.2	25.9	25.9	25.9	27.5	24.7	18.3	17.5	13.8
1999.....	11.9	22.5	22.9	28.2	30.7	27.7	25.3	23.2	18.8	17.2

Black, male

2006 4/.....	8.9	15.2	31.5	35.9	29.7	34.4	22.5	31.8	13.7	*
2005 4/.....	8.1	20.8	31.7	38.3	40.5	30.9	31.6	32.6	23.2	*
2004 4/.....	10.5	21.3	34.3	40.9	36.4	35.1	32.0	20.0	21.5	*
2003 4/.....	9.7	21.4	41.5	42.2	38.9	36.7	28.9	19.6	22.1	21.9
2002.....	9.8	23.4	38.2	36.9	39.0	34.9	29.0	23.0	21.4	*
2001.....	14.5	28.2	42.2	43.8	38.4	43.2	33.7	26.9	*	*
2000.....	14.6	26.4	40.7	38.5	37.3	38.0	37.4	27.4	19.4	*
1999.....	17.1	34.5	36.6	44.1	42.4	43.4	37.3	31.8	25.3	*

Black, female

2006 4/.....	4.3	7.4	11.8	12.6	13.2	11.1	11.8	12.1	11.2	8.8
2005 4/.....	4.5	8.5	13.1	13.2	13.6	12.0	12.8	12.3	15.2	10.9
2004 4/.....	5.2	8.5	13.5	13.1	15.6	13.3	12.3	10.2	11.7	11.4
2003 4/.....	5.9	9.3	12.2	13.4	14.1	17.0	16.8	10.9	10.8	8.5
2002.....	6.3	12.2	13.7	14.5	14.9	14.1	13.9	18.4	13.1	10.7
2001.....	7.1	11.3	13.7	14.5	12.0	14.8	15.2	17.5	14.7	9.7
2000.....	7.7	13.0	13.3	15.5	17.0	19.8	16.2	12.8	16.6	11.6
1999.....	7.3	12.2	11.4	15.2	21.5	16.2	17.3	18.0	15.6	15.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcoholic liver disease (K70)

All races 3/, both sexes

2006.....	5.5	9.2	12.0	12.5	11.6	10.1	8.6	7.0	4.5	2.2
2005.....	5.7	9.3	12.0	12.4	12.0	10.3	9.4	6.4	4.8	1.8
2004.....	5.8	9.1	12.2	11.8	11.5	10.8	9.0	6.5	4.1	2.1
2003.....	5.9	9.4	11.8	11.6	11.6	11.0	8.3	6.6	4.4	1.8
2002.....	5.9	9.4	11.4	11.4	11.8	10.7	9.5	6.7	4.4	2.1
2001.....	6.1	10.2	11.3	11.1	12.0	11.1	9.0	6.9	3.9	2.4
2000.....	6.6	10.0	10.7	11.7	12.9	11.4	9.1	7.2	4.4	2.2
1999.....	6.4	10.0	10.9	11.5	12.6	11.6	9.4	7.3	5.2	2.8

All races 3/, male

2006.....	7.1	12.9	18.4	19.4	18.1	16.9	14.1	12.0	7.5	4.7
2005.....	7.5	13.2	18.5	19.5	19.4	16.5	15.0	11.2	8.7	4.2
2004.....	8.0	12.9	18.6	18.2	18.3	17.8	15.0	11.5	8.0	4.6
2003.....	8.0	14.2	18.3	18.1	18.6	17.7	14.2	11.3	8.2	3.6
2002.....	8.0	13.8	17.7	17.6	19.2	17.4	15.4	10.9	8.0	4.7
2001.....	8.4	15.3	17.0	18.0	19.0	17.9	15.2	12.1	7.0	5.5
2000.....	9.4	15.3	16.9	18.0	20.3	18.7	15.7	13.8	7.7	5.3
1999.....	9.4	15.5	17.2	18.7	20.0	19.7	16.2	12.1	9.7	6.1

All races 3/, female

2006.....	3.8	5.6	5.9	6.0	5.6	4.2	4.1	3.4	2.6	1.1
2005.....	3.9	5.4	5.8	5.7	5.3	4.9	4.8	3.0	2.4	0.7
2004.....	3.6	5.5	6.0	5.7	5.4	4.8	4.1	2.9	1.7	1.0
2003.....	3.8	4.8	5.5	5.5	5.3	5.3	3.6	3.3	2.1	1.0
2002.....	3.9	5.1	5.4	5.6	5.1	4.9	4.8	3.7	2.2	1.0
2001.....	3.9	5.3	5.8	4.7	5.6	5.3	4.0	3.3	2.1	1.0
2000.....	3.9	4.8	4.8	5.8	6.2	5.1	3.9	2.6	2.4	1.0
1999.....	3.5	4.6	4.8	4.9	5.9	4.6	4.0	3.9	2.7	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcoholic liver disease (K70)

White, both sexes

2006 4/.....	5.8	9.7	12.6	13.0	12.1	10.4	8.9	7.3	4.7	2.4
2005 4/.....	5.9	9.6	12.5	12.9	12.1	10.7	9.7	6.6	4.9	1.9
2004 4/.....	6.1	9.5	12.6	11.8	11.7	11.1	9.3	6.8	4.1	2.2
2003 4/.....	6.2	9.7	11.8	11.8	11.6	11.2	8.6	7.0	4.5	1.9
2002.....	6.2	9.6	11.5	11.4	12.0	11.0	9.8	6.8	4.5	2.1
2001.....	6.2	10.5	11.0	11.0	12.2	11.2	9.2	7.1	4.0	2.3
2000.....	6.6	10.0	10.6	11.5	13.0	11.6	9.3	7.3	4.3	2.3
1999.....	6.2	9.8	10.8	11.2	12.4	11.5	9.4	7.2	5.3	2.7

White, male

2006 4/.....	7.6	13.5	19.3	20.0	19.0	17.1	14.7	12.4	7.9	4.9
2005 4/.....	7.8	13.7	19.2	20.3	19.5	17.0	15.4	11.3	8.9	4.4
2004 4/.....	8.5	13.3	19.4	18.1	18.4	18.1	15.3	12.0	8.0	4.8
2003 4/.....	8.4	14.6	18.4	18.3	18.7	17.9	14.6	11.8	8.2	3.7
2002.....	8.3	14.3	17.9	17.9	19.4	17.7	15.8	11.1	8.2	4.5
2001.....	8.5	15.9	16.8	17.6	19.4	17.8	15.4	12.5	7.2	5.4
2000.....	9.4	15.6	16.8	18.0	20.6	19.0	15.9	14.1	7.8	5.4
1999.....	9.2	15.4	17.1	18.2	20.0	19.5	16.2	11.9	9.8	5.8

White, female

2006 4/.....	3.9	5.9	6.1	6.2	5.7	4.4	4.1	3.5	2.6	1.1
2005 4/.....	3.9	5.6	6.0	5.8	5.2	5.1	5.0	3.1	2.4	0.8
2004 4/.....	3.7	5.7	5.9	5.7	5.4	4.9	4.4	3.0	1.7	1.0
2003 4/.....	3.9	4.9	5.4	5.5	5.1	5.3	3.7	3.5	2.2	1.1
2002.....	4.0	5.0	5.2	5.3	5.1	5.0	4.9	3.7	2.2	1.1
2001.....	3.9	5.2	5.4	4.6	5.7	5.4	4.2	3.3	2.1	1.0
2000.....	3.8	4.4	4.5	5.3	6.0	5.1	4.0	2.6	2.3	1.0
1999.....	3.3	4.3	4.5	4.7	5.5	4.6	4.0	3.9	2.6	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcoholic liver disease (K70)

Black, both sexes

2006 4/.....	3.5	6.0	9.4	10.0	8.6	8.9	6.5	5.2	*	*
2005 4/.....	3.9	7.8	10.1	11.0	11.9	9.0	8.4	5.3	*	*
2004 4/.....	4.1	7.5	10.8	12.9	12.0	10.8	6.5	*	*	*
2003 4/.....	4.2	7.6	12.7	12.2	12.9	11.2	6.9	*	*	*
2002.....	4.2	8.2	12.1	12.5	12.0	9.6	7.4	6.2	*	*
2001.....	5.9	9.0	14.7	14.1	11.5	13.0	7.8	5.7	*	*
2000.....	6.9	11.0	13.3	13.9	12.6	11.8	8.4	6.2	*	*
1999.....	7.2	12.8	12.8	14.8	15.6	12.9	10.4	8.0	*	*

Black, male

2006 4/.....	4.5	8.8	14.1	16.3	12.7	16.9	10.2	9.8	*	*
2005 4/.....	4.6	11.7	15.0	16.8	20.5	15.4	15.2	12.1	*	*
2004 4/.....	5.6	10.7	15.6	20.8	19.4	19.4	13.6	*	*	*
2003 4/.....	5.4	12.0	20.6	19.5	20.8	18.7	12.5	*	*	*
2002.....	5.6	11.5	18.7	17.4	20.4	16.5	13.0	10.6	*	*
2001.....	8.2	12.9	22.3	23.5	18.7	22.9	15.1	*	*	*
2000.....	9.6	15.0	20.6	20.4	18.0	19.3	16.7	11.3	*	*
1999.....	10.1	19.2	20.2	24.9	23.3	24.1	18.7	14.2	*	*

Black, female

2006 4/.....	2.6	3.5	5.5	4.7	5.4	*	*	*	*	*
2005 4/.....	3.3	4.3	5.9	6.2	5.1	4.2	*	*	*	*
2004 4/.....	2.8	4.6	6.8	6.3	6.1	4.5	*	*	*	*
2003 4/.....	3.0	3.7	6.0	6.1	6.8	5.7	*	*	*	*
2002.....	3.0	5.3	6.6	8.4	5.4	4.4	*	*	*	*
2001.....	3.9	5.5	8.2	6.4	5.8	5.6	*	*	*	*
2000.....	4.5	7.5	7.1	8.5	8.3	6.3	*	*	*	*
1999.....	4.7	7.2	6.6	6.5	9.5	4.6	5.0	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic liver disease and cirrhosis (K73-K74)

All races 3/, both sexes

2006.....	2.7	5.6	9.2	10.3	11.4	14.7	18.9	22.5	23.7	17.2
2005.....	2.9	5.7	8.7	10.6	12.0	15.0	20.0	23.3	23.1	17.8
2004.....	2.9	6.0	9.1	9.9	12.2	15.4	20.5	23.6	22.9	17.6
2003.....	3.4	6.5	9.3	10.4	12.7	17.0	22.9	24.5	24.1	18.2
2002.....	3.7	6.6	8.9	10.3	12.6	17.2	21.7	25.8	25.5	19.3
2001.....	3.6	6.9	8.8	10.0	12.8	17.7	22.2	24.4	24.7	19.9
2000.....	3.6	6.8	7.9	10.2	13.3	17.5	21.7	24.7	25.2	20.9
1999.....	3.5	6.4	7.7	9.9	13.9	18.1	22.3	25.4	25.6	20.4

All races 3/, male

2006.....	3.7	7.8	13.4	14.4	15.0	17.2	21.3	24.9	26.9	21.4
2005.....	4.0	8.1	12.8	14.9	15.9	17.4	23.1	26.0	26.5	23.3
2004.....	4.1	8.6	13.3	14.1	15.2	17.9	23.8	27.1	26.0	21.0
2003.....	4.8	9.2	14.0	14.3	17.0	20.7	25.8	26.8	26.8	23.9
2002.....	4.9	9.5	13.3	14.4	15.3	19.8	24.6	28.6	28.9	23.7
2001.....	4.8	10.0	13.1	13.5	16.3	21.8	25.6	25.5	28.4	26.5
2000.....	5.0	10.2	11.6	13.7	17.4	20.6	25.0	27.0	26.8	29.0
1999.....	5.2	9.8	11.6	13.8	17.3	22.0	25.3	28.0	29.0	27.0

All races 3/, female

2006.....	1.8	3.4	5.1	6.5	8.1	12.5	16.8	20.7	21.7	15.2
2005.....	1.9	3.5	4.7	6.5	8.5	13.0	17.4	21.4	21.0	15.3
2004.....	1.8	3.5	5.1	6.0	9.5	13.2	18.0	21.1	21.0	16.1
2003.....	2.0	3.8	4.8	6.6	8.9	13.8	20.5	22.8	22.5	15.8
2002.....	2.6	3.8	4.7	6.5	10.2	14.9	19.4	23.9	23.4	17.4
2001.....	2.5	4.0	4.7	6.7	9.6	14.2	19.5	23.6	22.6	17.1
2000.....	2.2	3.6	4.3	6.9	9.5	14.8	19.2	23.2	24.2	17.5
1999.....	1.8	3.2	4.0	6.2	10.8	14.8	19.9	23.5	23.6	17.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic liver disease and cirrhosis (K73-K74)

White, both sexes

2006 4/.....	2.8	5.8	9.3	10.3	11.5	15.3	20.1	23.4	24.9	17.8
2005 4/.....	3.1	5.8	8.5	10.6	12.0	15.8	21.2	24.2	23.8	18.3
2004 4/.....	2.9	6.1	9.0	9.8	12.3	16.0	21.7	25.0	24.0	18.1
2003 4/.....	3.5	6.5	9.2	10.2	13.0	17.7	24.2	26.0	25.3	19.0
2002.....	3.9	6.5	8.7	10.4	12.8	18.0	22.8	27.4	26.4	20.2
2001.....	3.6	6.6	8.5	9.7	13.1	18.5	23.1	25.5	26.0	20.9
2000.....	3.6	6.9	7.5	10.1	13.5	17.8	22.5	26.1	26.2	21.6
1999.....	3.4	6.0	7.6	9.6	13.9	18.7	23.2	26.5	26.8	21.1

White, male

2006 4/.....	3.7	8.2	13.3	14.3	15.0	17.8	22.6	25.6	28.4	22.3
2005 4/.....	4.2	8.1	12.7	14.7	15.7	18.0	24.5	26.7	26.9	24.2
2004 4/.....	4.1	8.6	13.1	13.9	15.3	18.3	24.9	28.5	27.3	22.0
2003 4/.....	5.0	9.4	13.6	13.8	17.2	21.5	27.2	28.2	27.9	24.7
2002.....	5.2	9.5	13.0	14.3	15.3	20.5	25.8	30.5	30.0	25.1
2001.....	4.7	9.6	12.5	13.0	16.2	22.4	26.7	26.6	30.0	28.1
2000.....	5.0	10.3	11.0	13.5	17.6	21.2	25.8	28.2	28.1	30.1
1999.....	5.1	9.2	11.3	13.4	17.3	22.8	26.3	29.1	30.5	28.4

White, female

2006 4/.....	1.9	3.3	5.3	6.5	8.3	13.1	18.1	21.8	22.7	15.7
2005 4/.....	2.0	3.5	4.4	6.6	8.7	13.9	18.5	22.3	21.9	15.6
2004 4/.....	1.7	3.5	5.0	5.9	9.6	13.9	19.1	22.4	21.9	16.4
2003 4/.....	1.9	3.6	4.8	6.7	9.2	14.3	21.7	24.3	23.7	16.5
2002.....	2.5	3.5	4.6	6.6	10.5	15.7	20.4	25.1	24.3	18.1
2001.....	2.4	3.8	4.6	6.6	10.2	15.1	20.2	24.7	23.5	17.9
2000.....	2.1	3.4	4.2	6.9	9.6	14.8	19.9	24.6	25.1	18.2
1999.....	1.8	2.9	3.9	6.0	10.7	15.1	20.7	24.6	24.6	18.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other chronic liver disease and cirrhosis (K73-K74)

Black, both sexes

2006 4/.....	3.0	5.0	11.3	13.2	11.8	12.1	9.6	14.2	9.5	9.9
2005 4/.....	2.3	6.5	11.5	13.6	13.5	11.0	12.1	14.5	14.6	10.7
2004 4/.....	3.6	7.0	12.2	12.8	12.8	11.8	13.7	11.2	11.4	11.3
2003 4/.....	3.6	7.4	13.0	14.2	12.1	14.2	14.8	11.1	10.9	10.8
2002.....	3.7	9.2	12.8	12.1	13.5	13.4	12.6	13.9	12.0	9.8
2001.....	4.7	10.2	12.1	13.6	12.2	13.9	14.8	15.3	11.6	7.7
2000.....	4.1	8.2	12.6	12.0	13.3	15.7	16.3	12.1	13.3	12.2
1999.....	4.7	9.8	10.1	13.4	15.1	14.8	14.9	15.2	13.3	12.8

Black, male

2006 4/.....	4.5	6.4	17.4	19.6	17.0	17.6	12.3	21.9	*	*
2005 4/.....	3.5	9.1	16.7	21.6	20.0	15.4	16.4	20.6	19.0	*
2004 4/.....	4.9	10.6	18.7	20.1	17.0	15.7	18.5	15.9	14.3	*
2003 4/.....	4.3	9.5	20.9	22.7	18.1	18.0	16.4	14.6	*	*
2002.....	4.2	11.9	19.5	19.5	18.6	18.4	16.0	12.4	*	*
2001.....	6.4	15.3	19.9	20.3	19.7	20.3	18.5	18.1	*	*
2000.....	5.0	11.4	20.1	18.1	19.3	18.7	20.7	16.1	*	*
1999.....	7.0	15.3	16.4	19.2	19.1	19.2	18.7	17.6	*	*

Black, female

2006 4/.....	1.7	3.8	6.2	7.8	7.8	8.1	7.8	9.6	9.4	8.8
2005 4/.....	*	4.2	7.1	7.1	8.5	7.8	9.1	10.9	12.2	10.1
2004 4/.....	2.4	3.8	6.7	6.8	9.5	8.8	10.5	8.4	9.8	11.0
2003 4/.....	2.9	5.6	6.2	7.3	7.4	11.3	13.7	9.0	10.1	8.1
2002.....	3.3	6.9	7.1	6.1	9.4	9.6	10.3	14.8	10.7	9.9
2001.....	3.2	5.8	5.5	8.1	6.2	9.2	12.3	13.6	13.1	*
2000.....	3.2	5.5	6.1	7.0	8.7	13.4	13.3	9.7	14.0	11.2
1999.....	2.6	5.0	4.8	8.7	12.0	11.5	12.4	13.7	12.1	13.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races 3/, both sexes

2006.....	0.1	0.2	0.4	0.6	1.2	1.8	3.3	6.2	9.9	21.8
2005.....	0.1	0.2	0.4	0.5	0.9	1.7	3.3	5.8	11.1	22.9
2004.....	0.2	0.3	0.4	0.6	1.2	1.8	3.2	5.8	10.4	23.8
2003.....	0.1	0.2	0.4	0.5	0.9	2.0	3.4	5.9	11.3	21.8
2002.....	0.2	0.2	0.4	0.6	1.1	2.0	3.6	5.7	10.8	22.9
2001.....	0.1	0.3	0.4	0.7	1.3	2.0	3.8	5.8	10.0	23.7
2000.....	0.1	0.2	0.4	0.6	1.0	1.9	3.0	5.7	10.3	24.8
1999.....	0.2	0.3	0.4	0.7	1.0	1.8	3.4	5.7	10.9	23.9

All races 3/, male

2006.....	*	0.2	0.5	0.6	1.2	2.3	4.0	7.6	12.5	21.7
2005.....	*	*	0.5	0.5	1.2	2.3	4.3	7.1	12.7	24.3
2004.....	*	0.3	0.4	0.7	1.4	2.2	3.8	7.0	12.5	25.5
2003.....	*	0.2	0.5	0.6	1.1	2.2	4.3	7.5	13.9	22.7
2002.....	0.2	0.3	0.5	0.8	1.1	2.3	4.0	7.5	13.5	24.1
2001.....	*	0.4	0.4	0.9	1.5	2.2	4.4	7.6	12.1	26.9
2000.....	*	0.2	0.3	0.7	1.2	2.3	3.6	7.4	12.9	27.6
1999.....	*	0.3	0.4	0.9	1.3	2.0	4.2	6.9	13.3	27.7

All races 3/, female

2006.....	*	0.2	0.3	0.5	1.2	1.3	2.7	5.2	8.3	21.9
2005.....	*	0.2	0.3	0.6	0.7	1.2	2.4	4.8	10.1	22.3
2004.....	*	0.3	0.4	0.5	1.0	1.5	2.8	5.0	9.0	23.0
2003.....	*	0.2	0.4	0.4	0.7	1.8	2.8	4.8	9.8	21.4
2002.....	*	*	0.3	0.4	1.0	1.8	3.2	4.4	9.1	22.4
2001.....	*	0.3	0.3	0.5	1.1	1.9	3.3	4.6	8.7	22.4
2000.....	*	0.2	0.4	0.5	0.8	1.6	2.5	4.5	8.7	23.7
1999.....	*	0.3	0.5	0.6	0.7	1.6	2.8	4.8	9.5	22.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

White, both sexes

2006 4/.....	*	0.2	0.3	0.5	1.1	1.6	3.0	6.1	10.0	22.3
2005 4/.....	*	0.1	0.3	0.5	0.9	1.5	3.1	5.7	11.1	23.1
2004 4/.....	0.2	0.2	0.3	0.5	1.1	1.8	3.0	5.7	10.3	24.2
2003 4/.....	*	0.2	0.3	0.4	0.8	1.8	3.6	5.9	11.3	21.9
2002.....	0.2	0.2	0.3	0.6	1.0	2.0	3.5	5.5	10.6	23.5
2001.....	*	0.3	0.3	0.6	1.3	1.8	3.7	5.7	9.8	23.9
2000.....	0.1	0.2	0.3	0.6	0.9	1.8	2.7	5.6	10.1	25.6
1999.....	0.1	0.3	0.4	0.6	0.9	1.6	3.4	5.7	11.0	24.4

White, male

2006 4/.....	*	*	0.4	0.5	1.1	2.1	3.8	7.5	12.7	22.8
2005 4/.....	*	*	0.4	0.5	1.1	2.1	4.1	7.1	13.1	24.8
2004 4/.....	*	0.2	0.3	0.6	1.4	2.2	3.7	6.9	12.7	26.7
2003 4/.....	*	*	0.4	0.5	0.9	2.1	4.4	7.5	13.9	23.0
2002.....	*	0.3	0.5	0.8	1.1	2.3	4.0	7.4	13.3	24.7
2001.....	*	0.3	0.4	0.8	1.4	1.9	4.4	7.6	12.0	26.7
2000.....	*	*	0.3	0.7	1.1	2.3	3.4	7.4	13.2	28.9
1999.....	*	0.3	0.3	0.7	1.2	1.9	4.2	7.0	13.8	28.1

White, female

2006 4/.....	*	*	0.3	0.5	1.2	1.1	2.4	5.0	8.3	22.0
2005 4/.....	*	*	0.2	0.5	0.7	0.9	2.3	4.7	9.8	22.4
2004 4/.....	*	0.2	0.3	0.5	0.8	1.4	2.4	4.8	8.8	23.1
2003 4/.....	*	*	0.3	0.3	0.7	1.6	2.8	4.7	9.8	21.4
2002.....	*	*	0.3	0.4	0.9	1.7	3.1	4.1	9.0	23.0
2001.....	*	0.3	*	0.4	1.1	1.8	3.1	4.4	8.4	22.8
2000.....	*	*	0.3	0.4	0.7	1.4	2.2	4.4	8.3	24.3
1999.....	*	0.3	0.4	0.6	0.6	1.4	2.7	4.8	9.3	22.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

Black, both sexes

2006 4/.....	*	*	0.9	1.1	1.9	3.4	4.6	9.0	9.0	17.1
2005 4/.....	*	*	*	*	*	3.6	4.6	7.0	12.4	20.6
2004 4/.....	*	*	1.0	1.3	2.0	2.3	5.1	7.1	12.1	19.5
2003 4/.....	*	*	1.0	*	*	3.0	2.8	6.0	10.9	20.7
2002.....	*	*	*	*	*	2.3	4.6	7.7	12.8	14.8
2001.....	*	*	*	1.4	2.0	3.5	6.2	6.1	11.6	19.7
2000.....	*	*	*	*	*	3.1	4.9	6.6	11.0	17.8
1999.....	*	*	*	*	2.0	3.2	4.1	4.6	11.3	15.3

Black, male

2006 4/.....	*	*	*	*	*	*	*	10.3	*	*
2005 4/.....	*	*	*	*	*	*	6.3	9.8	*	19.5
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	20.9
2002.....	*	*	*	*	*	*	*	*	16.0	*
2001.....	*	*	*	*	*	*	*	*	*	27.2
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, female

2006 4/.....	*	*	*	*	*	*	5.3	8.3	9.4	19.7
2005 4/.....	*	*	*	*	*	*	*	5.3	13.4	21.0
2004 4/.....	*	*	*	*	*	*	5.5	6.7	12.9	21.7
2003 4/.....	*	*	*	*	*	*	*	*	9.3	20.7
2002.....	*	*	*	*	*	*	5.4	7.7	11.1	16.0
2001.....	*	*	*	*	*	*	6.3	6.9	13.1	16.8
2000.....	*	*	*	*	*	*	4.9	6.3	11.5	19.3
1999.....	*	*	*	*	*	*	*	*	12.1	15.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races 3/, both sexes

2006.....	2.3	3.8	6.7	10.7	18.1	30.5	50.3	86.4	144.0	290.5
2005.....	2.1	3.8	5.8	10.6	17.7	29.2	51.3	85.6	142.8	288.3
2004.....	2.2	3.7	6.4	10.6	17.6	29.6	49.0	85.4	139.0	286.6
2003.....	2.3	3.8	6.1	10.4	17.9	30.7	50.6	84.5	143.9	293.1
2002.....	2.1	3.7	5.9	10.0	16.8	29.4	50.0	85.4	142.2	288.6
2001.....	2.1	3.6	5.8	9.9	16.9	31.7	49.3	82.2	135.8	287.7
2000.....	2.1	3.4	5.5	9.4	16.9	29.5	47.2	79.5	132.8	277.8
1999.....	2.0	3.1	5.2	9.0	15.8	28.2	46.7	77.7	127.4	268.9

All races 3/, male

2006.....	2.7	4.7	8.0	12.4	21.0	33.8	59.4	105.1	182.1	377.1
2005.....	2.5	4.8	7.2	12.4	21.1	31.4	59.1	102.7	183.8	376.5
2004.....	2.6	4.4	7.7	11.8	19.6	33.3	56.2	102.2	181.2	372.2
2003.....	2.7	4.5	7.6	12.2	19.6	34.6	58.1	102.9	187.9	387.6
2002.....	2.3	4.3	6.9	11.0	18.6	33.6	59.8	105.7	182.7	385.5
2001.....	2.3	4.4	6.7	10.9	18.2	36.5	57.0	100.5	176.8	393.7
2000.....	2.5	4.4	6.4	10.7	18.6	32.8	54.3	99.8	173.0	380.7
1999.....	2.3	3.8	5.8	9.9	17.8	31.0	54.9	99.2	168.9	371.1

All races 3/, female

2006.....	1.9	2.9	5.5	9.0	15.5	27.6	42.8	72.7	120.2	250.0
2005.....	1.7	2.9	4.5	8.8	14.7	27.3	44.9	73.2	117.2	247.8
2004.....	1.8	3.1	5.2	9.4	15.9	26.4	43.2	73.3	113.0	248.1
2003.....	1.9	3.2	4.7	8.6	16.4	27.4	44.7	71.4	117.1	251.4
2002.....	1.9	3.1	4.9	9.2	15.2	25.8	42.1	71.1	117.6	246.6
2001.....	1.9	2.9	4.9	8.9	15.7	27.6	43.2	69.4	111.2	243.3
2000.....	1.6	2.5	4.6	8.2	15.5	26.6	41.6	65.3	109.0	235.9
1999.....	1.7	2.4	4.5	8.1	14.0	25.8	40.3	62.8	103.1	227.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

White, both sexes

2006 4/.....	1.5	2.8	4.7	8.1	14.2	24.5	43.1	79.2	136.5	281.5
2005 4/.....	1.4	2.7	4.3	7.7	13.8	23.9	44.3	78.7	135.3	280.3
2004 4/.....	1.6	2.5	4.5	7.8	13.6	24.3	42.9	78.2	132.3	278.2
2003 4/.....	1.5	2.6	4.3	7.8	14.1	25.2	43.5	77.6	136.0	284.8
2002.....	1.4	2.6	3.9	7.4	13.1	24.4	43.2	78.1	135.5	278.8
2001.....	1.4	2.4	4.1	7.3	12.8	25.8	43.1	75.1	127.9	278.2
2000.....	1.5	2.2	3.8	6.8	13.2	23.8	40.6	73.0	125.3	268.7
1999.....	1.3	2.0	3.5	6.6	12.2	23.0	40.4	70.9	118.5	255.7

White, male

2006 4/.....	1.9	3.6	5.7	9.4	17.2	27.8	52.4	98.0	175.3	373.8
2005 4/.....	1.7	3.4	5.5	8.9	16.8	27.1	52.7	95.9	178.1	373.7
2004 4/.....	2.0	2.9	5.3	9.0	15.2	27.6	50.4	95.6	176.4	366.7
2003 4/.....	1.9	3.1	5.1	9.2	15.5	29.1	50.5	96.9	179.7	383.6
2002.....	1.6	3.0	4.6	8.0	14.3	28.4	53.0	99.0	177.9	377.4
2001.....	1.6	2.8	4.9	8.2	14.4	30.5	51.3	94.6	169.0	388.9
2000.....	1.8	2.9	4.4	7.9	14.9	27.8	48.7	93.4	166.7	376.2
1999.....	1.5	2.4	4.0	7.8	14.0	26.0	48.9	91.6	159.6	360.8

White, female

2006 4/.....	1.1	2.1	3.7	6.9	11.4	21.6	35.3	65.2	112.0	238.1
2005 4/.....	1.1	2.0	3.2	6.5	11.0	21.0	37.4	66.0	108.5	237.3
2004 4/.....	1.1	2.1	3.7	6.5	12.0	21.5	36.7	65.4	104.9	238.3
2003 4/.....	1.1	2.1	3.4	6.5	12.8	21.8	37.7	63.6	109.1	241.0
2002.....	1.3	2.3	3.3	6.7	12.0	20.8	35.1	63.2	109.4	235.9
2001.....	1.2	1.9	3.3	6.3	11.4	21.7	36.4	61.3	103.1	231.9
2000.....	1.1	1.7	3.2	5.8	11.6	20.3	34.1	58.6	100.7	225.0
1999.....	1.1	1.6	3.1	5.5	10.6	20.4	33.6	56.4	94.3	213.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Black, both sexes

2006 4/.....	7.6	11.0	22.4	32.6	55.5	86.5	122.4	177.9	254.6	442.6
2005 4/.....	7.3	12.1	18.2	35.8	55.7	79.4	124.8	175.9	254.9	424.4
2004 4/.....	6.9	12.6	21.5	35.6	56.7	79.2	115.9	175.4	242.9	430.9
2003 4/.....	7.9	12.6	21.0	34.3	54.2	83.8	127.8	171.0	260.7	440.5
2002.....	6.7	11.7	21.6	33.7	50.4	79.2	126.6	178.0	241.5	441.6
2001.....	6.7	12.5	19.8	32.5	53.1	88.6	119.8	172.3	248.6	429.6
2000.....	6.3	12.7	19.7	32.3	50.7	83.5	119.7	165.4	242.3	406.4
1999.....	7.0	11.8	19.6	29.1	47.1	77.8	117.0	163.7	246.2	433.3

Black, male

2006 4/.....	9.0	13.6	27.7	40.4	64.5	98.9	135.6	210.3	312.8	488.8
2005 4/.....	8.5	15.9	22.6	45.8	68.3	79.4	133.5	208.8	303.4	482.4
2004 4/.....	7.1	15.4	27.8	39.7	64.1	92.1	129.5	199.3	285.5	512.6
2003 4/.....	8.8	14.8	29.3	43.3	62.7	94.1	155.7	191.7	333.7	518.2
2002.....	7.6	14.7	26.8	40.0	61.2	89.3	149.6	207.6	281.2	561.8
2001.....	7.0	16.5	22.8	37.7	55.5	101.1	130.9	189.1	309.8	511.7
2000.....	7.9	17.4	24.1	37.4	54.2	86.9	127.7	198.9	290.0	490.1
1999.....	8.2	15.5	23.4	29.2	54.5	81.8	130.6	208.0	327.0	526.1

Black, female

2006 4/.....	6.4	8.7	17.9	26.2	48.5	77.4	113.3	158.6	223.9	423.7
2005 4/.....	6.3	8.8	14.5	27.5	45.9	79.5	118.8	156.3	229.2	401.0
2004 4/.....	6.6	10.1	16.2	32.3	51.0	69.6	106.8	161.2	220.4	398.5
2003 4/.....	7.2	10.7	13.9	26.9	47.5	76.2	109.0	158.7	222.4	410.4
2002.....	5.9	9.1	17.2	28.5	42.0	71.8	111.1	160.3	220.8	395.5
2001.....	6.4	9.1	17.1	28.2	51.2	79.4	112.3	162.3	217.5	399.1
2000.....	4.8	8.6	16.1	28.2	47.9	81.0	114.3	145.3	218.2	375.2
1999.....	5.9	8.6	16.4	29.1	41.3	74.8	108.0	137.0	206.1	398.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All races 3/, both sexes

2006.....	*	*	*	*	*	*	*	0.4	0.6
2005.....	*	*	*	*	*	*	0.3	*	0.8
2004.....	*	*	*	*	*	*	0.3	0.4	0.8
2003.....	*	*	*	*	*	*	*	0.6	0.6
2002.....	*	*	*	*	*	*	*	0.6	0.9
2001.....	*	*	*	*	*	*	0.4	0.4	0.9
2000.....	*	*	*	*	*	*	*	0.7	1.2
1999.....	*	*	*	*	*	*	0.3	0.5	0.8

All races 3/, male

2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	1.8
1999.....	*	*	*	*	*	*	*	*	*

All races 3/, female

2006.....	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	0.7
2004.....	*	*	*	*	*	*	*	*	0.8
2003.....	*	*	*	*	*	*	*	*	0.6
2002.....	*	*	*	*	*	*	*	0.6	0.9
2001.....	*	*	*	*	*	*	*	*	0.7
2000.....	*	*	*	*	*	*	*	0.7	0.9
1999.....	*	*	*	*	*	*	0.5	*	0.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All races 3/, both sexes

2006.....	*	0.1	0.3	0.4	0.6	1.0	1.8	3.2	6.0	13.4
2005.....	*	*	0.2	0.3	0.4	0.5	0.9	1.6	2.4	5.4
2004.....	*	*	0.1	0.2	0.3	0.4	0.6	1.0	1.6	2.8
2003.....	*	*	0.2	0.1	0.2	0.4	0.7	1.0	1.6	2.8
2002.....	*	0.1	0.1	0.2	*	0.3	0.6	1.2	1.7	2.8
2001.....	*	*	0.1	*	0.2	0.5	0.7	0.9	1.5	2.8
2000.....	0.1	*	0.1	0.2	0.3	0.5	0.7	1.1	1.9	2.8
1999.....	*	*	0.2	0.2	0.3	0.4	0.7	1.4	2.0	3.2

All races 3/, male

2006.....	*	*	0.3	0.4	0.8	1.3	1.8	4.3	7.7	17.2
2005.....	*	*	0.2	0.3	0.4	0.6	1.1	1.9	3.2	7.5
2004.....	*	*	*	0.2	0.3	0.4	0.7	1.3	2.0	4.0
2003.....	*	*	0.2	*	*	0.5	0.6	1.3	2.0	4.1
2002.....	*	*	*	0.3	*	*	0.9	1.6	2.1	3.8
2001.....	*	*	*	*	*	0.5	1.0	1.3	2.3	3.7
2000.....	*	*	*	*	*	0.5	0.8	1.5	2.1	4.5
1999.....	*	*	*	*	*	0.5	0.6	1.9	2.7	5.0

All races 3/, female

2006.....	*	*	0.3	0.4	0.4	0.8	1.7	2.3	5.0	11.6
2005.....	*	*	*	*	0.3	0.5	0.8	1.4	1.9	4.4
2004.....	*	*	*	*	*	*	0.5	0.8	1.4	2.2
2003.....	*	*	*	*	*	*	0.8	0.9	1.4	2.2
2002.....	*	*	*	*	*	*	0.4	0.8	1.5	2.3
2001.....	*	*	*	*	*	0.5	0.5	0.7	1.0	2.4
2000.....	*	*	*	*	*	*	0.6	0.8	1.8	2.1
1999.....	*	*	*	*	*	0.4	0.7	1.1	1.5	2.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

White, both sexes

2006 4/.....	*	*	0.2	0.3	0.5	0.8	1.6	2.9	5.5	12.5
2005 4/.....	*	*	*	0.2	0.3	0.5	0.8	1.5	2.3	5.2
2004 4/.....	*	*	0.1	0.2	0.2	0.4	0.6	1.0	1.6	2.7
2003 4/.....	*	*	*	0.2	0.2	0.4	0.6	0.9	1.5	2.6
2002.....	*	*	*	0.2	*	0.3	0.7	1.0	1.7	2.6
2001.....	*	*	*	*	*	0.4	0.7	0.9	1.5	2.9
2000.....	*	*	*	0.2	*	0.4	0.7	1.0	1.9	2.4
1999.....	*	*	*	*	0.3	0.3	0.6	1.4	1.8	2.9

White, male

2006 4/.....	*	*	*	*	0.7	0.9	1.7	3.9	7.1	15.9
2005 4/.....	*	*	*	*	*	0.5	1.0	1.7	3.1	7.3
2004 4/.....	*	*	*	*	*	*	0.6	1.2	2.0	3.7
2003 4/.....	*	*	*	*	*	0.5	*	1.1	1.9	3.8
2002.....	*	*	*	*	*	*	0.9	1.5	2.1	3.8
2001.....	*	*	*	*	*	*	1.1	1.2	2.3	4.0
2000.....	*	*	*	*	*	*	0.8	1.4	2.2	4.0
1999.....	*	*	*	*	*	*	0.6	1.9	2.6	4.6

White, female

2006 4/.....	*	*	*	0.3	*	0.6	1.4	2.2	4.4	10.9
2005 4/.....	*	*	*	*	*	*	0.6	1.3	1.8	4.2
2004 4/.....	*	*	*	*	*	*	0.5	0.8	1.3	2.3
2003 4/.....	*	*	*	*	*	*	0.6	0.8	1.3	2.1
2002.....	*	*	*	*	*	*	0.5	0.7	1.4	2.0
2001.....	*	*	*	*	*	*	0.5	0.7	0.9	2.4
2000.....	*	*	*	*	*	*	0.6	0.7	1.7	1.8
1999.....	*	*	*	*	*	*	0.7	1.0	1.4	2.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Renal failure (N17-N19)

All races 3/, both sexes

2006.....	2.2	3.7	6.4	10.2	17.5	29.3	48.4	83.0	137.5	276.4
2005.....	2.0	3.8	5.6	10.3	17.3	28.6	50.2	83.7	140.0	282.1
2004.....	2.1	3.6	6.2	10.3	17.3	29.1	48.2	84.0	136.8	282.9
2003.....	2.2	3.7	5.9	10.2	17.6	30.2	49.7	83.2	141.7	289.7
2002.....	2.0	3.6	5.7	9.8	16.5	29.0	49.1	83.9	139.8	284.8
2001.....	2.0	3.5	5.7	9.7	16.6	31.1	48.4	80.9	133.8	284.0
2000.....	2.0	3.3	5.3	9.2	16.6	28.9	46.2	78.1	130.0	273.7
1999.....	1.9	3.0	4.9	8.7	15.3	27.7	45.7	75.7	124.8	264.8

All races 3/, male

2006.....	2.6	4.5	7.7	12.0	20.2	32.4	57.4	100.4	174.0	358.8
2005.....	2.3	4.7	6.9	12.0	20.5	30.7	57.8	100.4	180.3	368.0
2004.....	2.5	4.3	7.4	11.6	19.1	32.8	55.3	100.5	178.4	367.4
2003.....	2.6	4.3	7.4	12.0	19.3	33.9	57.2	101.3	185.2	383.1
2002.....	2.2	4.2	6.7	10.6	18.4	33.1	58.8	103.9	180.0	380.7
2001.....	2.2	4.2	6.5	10.7	17.9	35.9	55.8	98.9	173.9	388.7
2000.....	2.4	4.2	6.1	10.4	18.1	32.1	53.3	97.9	170.1	374.2
1999.....	2.2	3.6	5.6	9.7	17.3	30.5	53.9	96.7	165.1	364.6

All races 3/, female

2006.....	1.8	2.8	5.2	8.6	15.1	26.6	41.0	70.2	114.8	237.8
2005.....	1.7	2.8	4.4	8.6	14.3	26.7	44.0	71.6	115.0	242.6
2004.....	1.8	3.0	5.0	9.2	15.6	25.8	42.5	72.1	111.2	244.9
2003.....	1.8	3.2	4.6	8.5	16.0	27.0	43.6	70.2	115.0	248.4
2002.....	1.8	2.9	4.8	9.0	14.9	25.4	41.4	69.9	115.3	243.2
2001.....	1.8	2.8	4.8	8.8	15.5	27.0	42.5	68.3	109.8	240.1
2000.....	1.6	2.4	4.5	8.0	15.2	26.1	40.6	64.4	106.4	232.8
1999.....	1.6	2.3	4.3	7.9	13.5	25.3	39.3	61.1	101.2	224.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Renal failure (N17-N19)

White, both sexes

2006 4/.....	1.4	2.8	4.5	7.8	13.7	23.7	41.4	76.0	130.6	268.2
2005 4/.....	1.3	2.6	4.2	7.4	13.3	23.3	43.3	76.8	132.7	274.3
2004 4/.....	1.5	2.5	4.3	7.6	13.3	23.8	42.1	76.8	130.1	274.7
2003 4/.....	1.4	2.5	4.1	7.6	13.8	24.7	42.7	76.4	133.9	281.5
2002.....	1.4	2.5	3.8	7.1	12.9	23.9	42.3	76.8	133.2	275.3
2001.....	1.4	2.3	4.0	7.1	12.6	25.2	42.1	73.8	126.0	274.6
2000.....	1.4	2.1	3.7	6.6	12.8	23.3	39.6	71.7	122.6	265.0
1999.....	1.2	1.8	3.4	6.4	11.9	22.5	39.4	69.0	116.2	251.9

White, male

2006 4/.....	1.8	3.5	5.5	9.1	16.5	26.8	50.6	93.7	167.8	356.8
2005 4/.....	1.6	3.4	5.3	8.6	16.3	26.5	51.4	93.7	174.6	365.4
2004 4/.....	1.9	2.9	5.1	8.8	14.9	27.1	49.5	94.0	173.7	362.3
2003 4/.....	1.8	3.0	4.9	9.0	15.3	28.4	49.7	95.6	177.3	379.3
2002.....	1.5	2.9	4.5	7.7	14.0	27.9	52.0	97.3	175.3	372.7
2001.....	1.6	2.7	4.7	8.0	14.1	30.0	50.1	93.0	166.1	383.9
2000.....	1.7	2.7	4.2	7.6	14.5	27.2	47.7	91.6	163.8	370.4
1999.....	1.4	2.2	3.8	7.6	13.6	25.5	48.0	89.1	156.2	354.9

White, female

2006 4/.....	1.0	2.0	3.5	6.5	11.1	20.9	33.8	62.8	107.1	226.6
2005 4/.....	1.0	1.9	3.2	6.3	10.5	20.6	36.7	64.3	106.4	232.4
2004 4/.....	1.1	2.1	3.5	6.4	11.8	21.0	36.0	64.2	103.1	235.2
2003 4/.....	1.0	2.1	3.3	6.4	12.4	21.4	36.9	62.4	107.1	238.2
2002.....	1.2	2.1	3.2	6.6	11.8	20.3	34.4	62.1	107.4	232.9
2001.....	1.1	1.9	3.3	6.2	11.2	21.1	35.6	60.2	101.8	228.8
2000.....	1.1	1.6	3.1	5.6	11.3	19.9	33.2	57.8	98.1	222.1
1999.....	1.1	1.5	2.9	5.3	10.3	19.9	32.5	54.8	92.5	210.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Renal failure (N17-N19)

Black, both sexes

2006 4/.....	7.4	10.6	21.3	31.3	53.6	82.6	118.0	172.2	242.1	418.7
2005 4/.....	7.0	11.9	17.6	34.9	54.9	78.4	122.1	172.7	250.0	413.7
2004 4/.....	6.6	12.2	21.3	35.2	56.1	78.0	114.8	173.3	239.6	426.4
2003 4/.....	7.8	12.4	20.5	34.2	53.6	82.9	126.1	168.8	257.4	436.4
2002.....	6.6	11.2	21.1	33.3	50.1	78.8	125.7	175.4	238.1	434.2
2001.....	6.5	12.3	19.3	32.2	52.5	87.1	119.0	170.6	245.9	425.6
2000.....	6.1	12.3	18.9	32.0	50.1	82.1	118.6	163.1	238.9	400.4
1999.....	6.7	11.7	18.9	28.5	46.1	76.5	115.7	161.0	241.6	425.3

Black, male

2006 4/.....	8.9	13.1	26.4	38.3	61.8	93.7	132.3	202.3	297.1	454.9
2005 4/.....	8.0	15.6	21.9	44.7	67.1	78.7	131.2	203.4	297.1	469.4
2004 4/.....	6.8	15.0	27.6	39.1	63.0	91.1	127.9	196.5	282.7	506.7
2003 4/.....	8.6	14.4	28.7	43.3	61.9	92.6	153.7	188.1	328.5	513.0
2002.....	7.4	14.4	26.1	39.4	61.2	88.8	148.2	203.5	278.2	555.3
2001.....	6.8	16.1	22.1	37.2	54.5	99.1	129.9	187.7	307.4	508.3
2000.....	7.7	17.1	22.7	37.0	53.2	84.8	127.0	196.6	287.5	480.9
1999.....	8.0	15.3	22.7	28.7	53.5	80.5	128.9	204.7	320.9	513.2

Black, female

2006 4/.....	6.1	8.4	17.0	25.4	47.2	74.4	108.2	154.3	213.0	403.9
2005 4/.....	6.1	8.6	13.9	26.8	45.3	78.3	115.8	154.4	225.2	391.2
2004 4/.....	6.3	9.8	16.0	31.9	50.7	68.4	105.9	159.6	217.0	394.6
2003 4/.....	7.0	10.7	13.7	26.7	47.1	75.7	107.4	157.4	220.1	406.7
2002.....	5.9	8.5	16.8	28.2	41.5	71.4	110.7	158.6	217.2	387.7
2001.....	6.4	9.0	16.9	28.1	50.9	78.3	111.7	160.3	214.6	394.9
2000.....	4.7	8.2	15.7	27.8	47.6	80.1	113.0	143.0	214.4	370.4
1999.....	5.6	8.5	15.8	28.4	40.3	73.5	106.9	134.7	202.2	392.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Infections of kidney (N10-N12,N13.6,N15.1)

All races 3/, both sexes

2006.....	*	0.1	0.1	0.1	0.2	0.4	0.6	1.0	1.6	4.7
2005.....	*	0.1	0.1	0.1	0.2	0.4	0.7	1.3	2.5	5.2
2004.....	0.1	0.1	0.1	0.2	0.2	0.4	0.7	1.3	2.6	5.2
2003.....	0.1	0.1	0.2	0.2	0.2	0.6	0.6	1.3	2.8	5.9
2002.....	*	*	0.1	0.2	0.3	0.5	0.7	1.6	2.3	6.3
2001.....	*	0.1	0.1	0.2	0.2	0.4	0.6	1.7	2.4	6.0
2000.....	*	0.1	0.1	0.2	0.3	0.4	0.7	1.5	2.7	7.5
1999.....	*	0.1	0.2	0.2	0.3	0.5	0.9	1.5	2.7	7.5

All races 3/, male

2006.....	*	*	*	*	*	*	0.7	*	1.2	3.7
2005.....	*	*	*	*	*	*	0.7	0.9	2.5	3.6
2004.....	*	*	*	0.3	*	*	*	1.0	2.2	4.5
2003.....	*	*	*	*	*	*	*	0.8	2.4	5.4
2002.....	*	*	*	*	*	*	0.7	1.4	1.4	5.3
2001.....	*	*	*	*	*	*	0.5	1.2	2.1	4.3
2000.....	*	*	*	*	*	*	*	1.1	2.2	5.6
1999.....	*	*	*	*	*	*	0.6	1.2	2.6	6.8

All races 3/, female

2006.....	*	0.2	*	*	*	0.4	0.6	1.3	1.9	5.2
2005.....	*	*	*	*	*	0.5	0.8	1.6	2.6	5.9
2004.....	*	*	*	*	*	0.5	0.9	1.5	2.8	5.5
2003.....	*	*	*	0.3	*	0.8	0.7	1.6	3.1	6.1
2002.....	*	*	*	*	0.4	0.6	0.7	1.7	2.8	6.7
2001.....	*	*	0.2	*	*	*	0.6	2.1	2.6	6.7
2000.....	*	*	*	0.3	0.4	0.5	1.1	1.8	3.0	8.3
1999.....	*	*	0.3	*	*	0.6	1.1	1.7	2.8	7.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Infections of kidney (N10-N12,N13.6,N15.1)

White, both sexes

2006 4/.....	*	0.1	0.2	0.2	0.2	0.4	0.6	0.9	1.7	4.9
2005 4/.....	*	0.1	*	*	0.2	0.4	0.7	1.2	2.6	5.2
2004 4/.....	0.1	*	*	0.2	0.2	0.4	0.7	1.2	2.5	5.2
2003 4/.....	0.1	0.1	0.2	0.2	0.2	0.6	0.6	1.3	2.7	6.0
2002.....	*	*	*	0.2	0.2	0.5	0.8	1.6	2.3	6.4
2001.....	*	*	*	*	*	0.3	0.6	1.7	2.2	6.1
2000.....	*	*	*	0.2	0.3	0.4	0.7	1.4	2.5	7.6
1999.....	*	*	0.2	0.2	0.2	0.5	0.9	1.5	2.8	7.7

White, male

2006 4/.....	*	*	*	*	*	*	0.7	*	1.2	3.7
2005 4/.....	*	*	*	*	*	*	0.6	0.9	2.6	3.5
2004 4/.....	*	*	*	*	*	*	*	0.9	2.4	4.7
2003 4/.....	*	*	*	*	*	*	*	0.8	2.2	5.8
2002.....	*	*	*	*	*	*	0.8	1.4	1.4	5.4
2001.....	*	*	*	*	*	*	*	1.1	2.0	4.3
2000.....	*	*	*	*	*	*	*	1.1	2.1	5.9
1999.....	*	*	*	*	*	*	*	1.3	2.6	7.2

White, female

2006 4/.....	*	*	*	*	*	0.5	*	1.2	2.0	5.4
2005 4/.....	*	*	*	*	*	0.5	0.7	1.5	2.7	6.0
2004 4/.....	*	*	*	*	*	0.5	1.0	1.4	2.7	5.5
2003 4/.....	*	*	*	*	*	0.8	0.7	1.6	3.1	6.0
2002.....	*	*	*	*	*	0.6	0.8	1.7	2.9	6.8
2001.....	*	*	*	*	*	*	0.7	2.2	2.4	6.9
2000.....	*	*	*	*	*	0.5	1.0	1.7	2.8	8.3
1999.....	*	*	*	*	*	0.6	1.1	1.7	2.9	7.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes

2006.....	1.1	1.5	1.6	2.2	2.4	1.9	2.0	2.7	3.5	5.6
2005.....	1.2	1.5	1.8	2.3	2.4	1.7	2.0	2.4	4.0	6.4
2004.....	1.2	1.5	1.8	2.2	2.2	2.3	2.3	3.1	4.3	6.1
2003.....	1.4	1.5	1.8	2.4	2.4	2.3	2.2	3.0	4.4	6.8
2002.....	1.3	1.6	1.9	2.3	2.4	2.3	2.5	3.1	4.8	6.6
2001.....	1.4	1.5	1.8	2.1	2.5	2.2	2.7	3.3	4.3	7.2
2000.....	1.3	1.4	1.8	2.2	2.5	2.6	2.4	3.3	5.0	7.5
1999.....	1.2	1.7	1.8	2.3	2.4	2.6	2.5	3.4	5.0	7.8

All races 3/, male

2006.....	1.2	1.7	1.9	2.5	2.7	1.9	2.3	3.0	4.5	6.8
2005.....	1.3	1.5	1.9	2.5	2.6	1.6	2.0	2.8	4.9	6.9
2004.....	1.3	1.7	2.0	2.3	2.2	2.3	2.7	3.7	3.8	6.8
2003.....	1.5	1.6	1.8	2.5	2.6	2.4	2.4	2.8	4.5	6.7
2002.....	1.4	1.9	2.0	2.3	2.7	2.4	2.7	3.4	4.8	7.0
2001.....	1.6	1.5	1.9	2.1	2.5	2.5	3.1	3.6	4.8	7.7
2000.....	1.5	1.6	2.2	2.0	2.6	2.7	2.6	3.3	5.3	10.6
1999.....	1.3	1.9	2.0	2.5	2.5	2.4	2.6	3.4	5.7	8.5

All races 3/, female

2006.....	1.0	1.3	1.3	1.9	2.1	1.8	1.8	2.4	2.8	5.1
2005.....	1.2	1.4	1.8	2.1	2.3	1.7	2.0	2.2	3.5	6.2
2004.....	1.2	1.3	1.7	2.0	2.1	2.3	2.0	2.7	4.6	5.8
2003.....	1.3	1.3	1.8	2.3	2.3	2.2	2.1	3.2	4.4	6.8
2002.....	1.2	1.4	1.8	2.3	2.2	2.3	2.3	2.9	4.9	6.5
2001.....	1.3	1.4	1.8	2.1	2.6	2.0	2.4	3.1	4.0	7.0
2000.....	1.1	1.3	1.5	2.4	2.5	2.5	2.3	3.3	4.8	6.3
1999.....	1.2	1.6	1.7	2.2	2.3	2.8	2.3	3.4	4.5	7.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, both sexes

2006 4/.....	1.1	1.5	1.6	2.3	2.5	1.9	2.1	2.8	3.5	5.7
2005 4/.....	1.3	1.5	2.0	2.4	2.6	1.8	2.2	2.5	4.3	6.5
2004 4/.....	1.3	1.6	1.9	2.3	2.3	2.4	2.5	3.1	4.4	6.3
2003 4/.....	1.5	1.5	1.9	2.5	2.5	2.3	2.3	3.1	4.5	6.9
2002.....	1.3	1.7	2.0	2.4	2.5	2.5	2.5	3.2	4.8	6.9
2001.....	1.5	1.4	1.9	2.2	2.7	2.3	2.7	3.4	4.4	7.4
2000.....	1.3	1.5	2.0	2.2	2.6	2.6	2.4	3.4	5.2	7.8
1999.....	1.3	1.8	1.8	2.4	2.4	2.7	2.4	3.3	5.1	8.0

White, male

2006 4/.....	1.2	1.7	1.9	2.6	2.8	2.0	2.3	3.0	4.6	6.8
2005 4/.....	1.3	1.6	2.0	2.6	2.7	1.8	2.2	2.8	5.2	6.9
2004 4/.....	1.4	1.8	2.0	2.4	2.3	2.4	2.9	3.6	3.9	7.2
2003 4/.....	1.6	1.5	2.0	2.6	2.6	2.3	2.3	2.8	4.3	6.9
2002.....	1.4	1.8	2.0	2.3	2.8	2.6	2.7	3.4	4.6	7.3
2001.....	1.6	1.5	1.9	2.2	2.6	2.7	3.2	3.9	4.6	7.5
2000.....	1.4	1.6	2.3	2.0	2.8	2.6	2.6	3.4	5.6	11.1
1999.....	1.3	1.9	2.0	2.6	2.6	2.5	2.6	3.0	5.8	8.6

White, female

2006 4/.....	1.0	1.4	1.3	2.0	2.2	1.9	1.9	2.6	2.7	5.2
2005 4/.....	1.3	1.4	1.9	2.2	2.5	1.8	2.2	2.2	3.7	6.3
2004 4/.....	1.2	1.3	1.9	2.2	2.3	2.5	2.2	2.8	4.7	6.0
2003 4/.....	1.3	1.4	1.9	2.4	2.4	2.2	2.2	3.3	4.7	6.9
2002.....	1.2	1.5	2.0	2.4	2.2	2.4	2.4	3.0	5.0	6.7
2001.....	1.3	1.4	1.9	2.2	2.8	2.1	2.3	3.1	4.2	7.3
2000.....	1.1	1.3	1.7	2.4	2.4	2.7	2.3	3.4	4.9	6.5
1999.....	1.2	1.7	1.7	2.3	2.3	2.9	2.3	3.6	4.6	7.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2006.....	4.2	4.9	5.6	5.9	7.2	9.5	15.5	28.7	58.6	236.1
2005.....	5.2	6.4	7.0	6.9	7.7	10.4	15.7	27.6	54.0	224.4
2004.....	4.9	5.7	5.8	6.3	7.9	10.2	15.4	28.0	55.5	221.5
2003.....	5.9	7.3	6.8	6.8	8.2	10.6	17.3	26.7	55.7	224.1
2002.....	6.0	6.4	6.9	7.0	8.3	10.4	16.1	28.3	53.3	208.0
2001.....	7.7	9.1	8.8	8.4	9.9	12.7	17.8	29.8	54.3	206.4
2000.....	7.6	8.6	8.0	8.6	10.2	14.2	19.0	30.1	56.5	208.1
1999.....	5.1	5.9	5.8	6.9	8.9	11.9	17.6	26.5	50.5	187.7

All races 3/, male

2006.....	5.5	6.3	7.2	7.7	9.0	11.9	18.5	32.3	61.3	191.1
2005.....	6.4	8.2	9.3	9.2	9.6	12.8	18.8	31.0	54.8	183.9
2004.....	6.5	7.4	7.7	8.3	10.1	12.6	17.9	30.6	58.1	183.6
2003.....	7.4	9.4	8.9	8.9	10.6	13.7	21.3	31.2	58.0	177.9
2002.....	7.6	8.5	9.0	9.0	11.2	12.7	19.7	32.9	60.6	175.5
2001.....	9.9	11.9	11.8	11.0	13.2	15.5	20.9	33.2	60.6	175.0
2000.....	10.2	11.7	11.2	11.4	13.4	18.9	22.4	36.1	62.3	182.7
1999.....	6.7	8.2	7.9	9.1	11.9	15.1	21.8	31.5	54.4	166.3

All races 3/, female

2006.....	3.0	3.5	4.1	4.2	5.6	7.3	13.1	26.0	56.9	257.2
2005.....	4.1	4.6	4.8	4.7	6.0	8.3	13.2	25.1	53.5	243.0
2004.....	3.3	4.1	4.0	4.5	5.9	8.1	13.4	26.1	53.8	238.5
2003.....	4.4	5.1	4.8	4.9	6.0	7.9	14.2	23.5	54.3	244.6
2002.....	4.5	4.4	5.0	5.0	5.6	8.4	13.3	25.1	48.9	222.1
2001.....	5.5	6.3	5.9	5.9	6.9	10.2	15.4	27.4	50.4	219.5
2000.....	5.0	5.6	5.0	5.9	7.3	10.1	16.4	26.0	53.0	218.5
1999.....	3.4	3.7	3.9	4.9	6.2	9.2	14.2	23.0	48.3	196.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, both sexes

2006 4/.....	4.0	4.7	5.2	5.6	7.0	9.2	15.0	28.3	58.3	241.1
2005 4/.....	5.0	5.9	6.6	6.3	7.3	9.7	15.0	27.2	53.7	228.9
2004 4/.....	4.7	5.6	5.4	5.8	7.4	9.8	14.7	27.6	55.1	224.4
2003 4/.....	5.6	6.6	6.2	6.3	7.6	10.1	16.6	26.5	56.1	228.3
2002.....	5.7	5.8	6.1	6.4	7.9	9.8	15.9	27.9	53.3	211.6
2001.....	7.6	8.8	8.3	7.9	9.4	11.8	17.2	29.5	53.9	210.0
2000.....	7.4	8.2	7.3	7.9	9.5	13.0	18.5	29.5	56.0	212.2
1999.....	4.4	5.2	5.0	6.2	8.2	11.1	17.0	25.5	49.5	190.2

White, male

2006 4/.....	5.1	5.9	6.6	7.3	8.8	11.5	17.2	31.5	60.5	193.8
2005 4/.....	6.3	7.8	8.7	8.3	8.9	11.8	17.9	30.4	53.6	185.6
2004 4/.....	6.3	7.2	6.9	7.5	9.3	11.9	17.4	30.1	57.8	186.0
2003 4/.....	7.0	8.6	8.2	8.1	9.7	12.8	20.4	31.0	57.8	180.0
2002.....	7.2	7.6	7.7	8.1	10.7	11.9	19.2	31.6	60.0	178.2
2001.....	10.0	11.6	11.2	10.1	12.2	14.3	20.3	33.2	60.6	177.0
2000.....	10.1	11.2	10.1	10.2	12.4	17.1	21.5	35.4	62.0	186.8
1999.....	5.9	7.0	6.7	8.0	10.9	13.9	21.0	30.3	52.6	167.3

White, female

2006 4/.....	2.8	3.5	3.8	3.9	5.3	7.1	13.2	25.8	56.9	263.3
2005 4/.....	3.8	4.1	4.5	4.3	5.8	7.9	12.6	24.9	53.7	248.8
2004 4/.....	3.1	4.0	3.8	4.2	5.6	7.9	12.5	25.8	53.5	241.7
2003 4/.....	4.2	4.6	4.2	4.7	5.6	7.7	13.5	23.3	55.0	249.7
2002.....	4.2	4.0	4.4	4.7	5.3	7.9	13.1	25.3	49.2	226.1
2001.....	5.3	6.0	5.5	5.7	6.7	9.7	14.6	26.8	49.9	223.8
2000.....	4.6	5.2	4.6	5.6	6.7	9.4	16.0	25.4	52.4	222.5
1999.....	2.9	3.4	3.4	4.5	5.7	8.6	13.8	22.1	47.6	199.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, both sexes

2006 4/.....	6.9	7.6	10.2	10.3	11.1	14.9	24.4	41.0	75.7	223.9
2005 4/.....	8.0	11.2	12.4	13.9	14.2	19.2	26.2	38.8	70.9	215.1
2004 4/.....	7.6	8.6	11.0	12.5	14.6	16.1	25.9	39.9	72.0	220.8
2003 4/.....	9.7	13.4	13.4	13.0	15.7	17.9	30.3	34.9	60.9	210.9
2002.....	9.5	12.1	15.7	13.8	13.6	17.8	23.1	39.5	64.9	196.0
2001.....	9.8	13.2	14.6	14.3	15.4	21.7	27.9	38.6	66.6	191.6
2000.....	11.2	13.8	15.9	16.0	19.0	28.3	29.7	44.2	73.4	186.4
1999.....	10.3	13.1	13.3	14.9	17.0	22.3	27.2	42.7	72.5	182.4

Black, male

2006 4/.....	9.0	10.8	14.2	14.0	12.9	20.6	36.3	52.8	92.4	193.9
2005 4/.....	9.2	13.8	17.2	19.5	20.0	26.4	33.5	49.2	85.7	199.9
2004 4/.....	9.7	11.8	16.3	18.2	20.2	22.3	28.2	45.0	81.1	181.8
2003 4/.....	12.5	17.5	17.8	18.5	22.8	25.9	38.8	41.5	70.1	193.9
2002.....	12.4	17.1	22.1	19.5	20.0	22.9	31.0	57.0	87.1	181.2
2001.....	11.7	17.3	18.9	21.7	25.4	29.5	31.6	40.9	70.6	184.9
2000.....	13.5	18.2	23.6	25.0	25.5	41.4	38.1	54.8	83.5	161.5
1999.....	13.6	19.7	19.1	22.4	24.0	30.1	35.6	54.1	88.1	184.0

Black, female

2006 4/.....	5.0	4.8	6.9	7.3	9.6	10.7	16.3	34.0	66.9	236.2
2005 4/.....	6.9	8.9	8.4	9.3	9.6	13.9	21.2	32.5	63.1	221.3
2004 4/.....	5.9	5.9	6.6	7.7	10.2	11.5	24.4	36.9	67.3	236.3
2003 4/.....	7.3	9.9	9.7	8.4	10.2	12.1	24.6	30.9	56.1	217.5
2002.....	7.0	7.7	10.2	9.0	8.7	14.1	17.7	29.1	53.3	201.7
2001.....	8.1	9.7	11.0	8.2	7.5	16.0	25.3	37.2	64.6	194.1
2000.....	9.1	9.9	9.4	8.6	13.9	18.6	24.2	37.9	68.4	195.7
1999.....	7.3	7.3	8.5	8.9	11.5	16.5	21.6	35.8	64.8	181.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races 3/, both sexes

2006.....	23.0	33.3	47.6	61.8	85.0	118.6	191.7	335.2	646.8	1,724.2
2005.....	22.8	33.6	47.8	60.3	81.9	115.6	182.4	313.2	593.5	1,559.1
2004.....	21.9	33.6	45.0	58.1	78.7	111.5	176.4	300.6	554.8	1,461.6
2003.....	23.1	33.3	45.3	56.6	80.9	114.5	178.3	303.7	559.8	1,508.4
2002.....	23.5	33.6	42.6	56.2	77.1	110.4	173.8	297.8	555.5	1,493.0
2001.....	21.7	32.1	40.9	53.2	76.4	112.0	172.3	295.5	539.3	1,476.7
2000.....	22.1	31.0	40.4	53.1	74.1	106.0	168.2	288.9	537.8	1,445.7
1999.....	21.6	29.9	38.4	51.0	72.1	107.7	167.8	281.3	510.8	1,410.7

All races 3/, male

2006.....	27.2	39.7	57.9	73.0	96.3	131.0	207.6	349.9	652.5	1,455.0
2005.....	27.6	40.7	58.5	71.9	93.1	125.3	194.2	327.1	595.6	1,305.3
2004.....	26.0	40.6	56.0	68.3	88.8	123.1	187.2	315.5	549.4	1,236.6
2003.....	27.8	40.3	55.2	65.6	91.0	125.7	191.0	316.5	563.7	1,258.1
2002.....	28.2	41.0	52.4	64.4	87.9	122.8	186.8	309.6	555.2	1,262.9
2001.....	25.7	38.8	49.1	59.9	85.8	124.3	186.5	307.6	539.6	1,271.6
2000.....	26.3	38.5	47.6	60.9	83.1	117.6	183.1	303.4	537.6	1,266.2
1999.....	26.6	36.3	45.4	59.5	80.9	117.1	184.3	298.1	528.1	1,225.4

All races 3/, female

2006.....	19.0	27.0	37.8	51.3	74.7	107.8	178.8	324.4	643.2	1,850.2
2005.....	17.9	26.7	37.6	49.4	71.7	107.2	172.9	303.1	592.2	1,675.6
2004.....	17.9	26.7	34.6	48.4	69.5	101.4	167.6	289.9	558.0	1,562.8
2003.....	18.5	26.5	36.0	48.1	71.7	104.8	168.0	294.7	557.4	1,619.1
2002.....	18.9	26.4	33.2	48.6	67.3	99.6	163.5	289.5	555.6	1,592.8
2001.....	17.7	25.6	33.0	46.9	67.9	101.4	161.0	287.0	539.1	1,562.6
2000.....	17.9	23.7	33.4	45.8	65.9	96.1	156.5	278.8	537.8	1,518.8
1999.....	16.6	23.7	31.7	43.1	64.2	99.5	154.8	269.7	500.7	1,485.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

White, both sexes

2006 4/.....	21.5	31.2	44.8	58.2	82.7	116.0	189.3	336.3	650.6	1,757.7
2005 4/.....	21.4	31.1	44.7	57.4	78.8	112.6	179.6	311.2	596.4	1,586.5
2004 4/.....	20.4	31.0	42.0	54.8	75.6	108.1	173.5	298.9	558.1	1,486.9
2003 4/.....	21.4	30.8	41.6	53.5	77.1	111.9	175.4	301.8	562.0	1,534.2
2002.....	21.6	30.9	39.2	53.1	73.4	107.7	170.2	295.6	556.2	1,517.6
2001.....	19.3	28.7	37.3	49.5	72.6	108.9	169.2	292.6	541.2	1,498.0
2000.....	19.8	27.4	37.1	49.3	70.1	102.5	164.5	285.7	537.2	1,464.6
1999.....	19.1	26.3	34.8	47.4	68.6	103.2	163.6	277.1	512.1	1,434.0

White, male

2006 4/.....	26.0	38.1	55.1	68.7	93.1	128.5	203.2	347.8	653.1	1,481.8
2005 4/.....	26.8	38.6	55.5	68.5	89.5	120.8	190.8	322.7	594.7	1,328.2
2004 4/.....	25.2	38.4	52.7	64.4	85.2	118.1	183.1	309.7	550.1	1,251.8
2003 4/.....	26.4	38.0	50.8	62.0	86.2	121.6	186.0	311.4	561.0	1,279.1
2002.....	26.8	38.2	48.7	60.6	83.4	118.6	181.2	304.6	553.0	1,274.7
2001.....	23.4	35.3	45.1	55.3	81.8	119.9	182.1	302.0	540.6	1,282.9
2000.....	24.0	35.0	44.1	56.8	77.5	112.7	177.5	298.2	533.4	1,274.8
1999.....	24.3	32.5	41.6	55.4	76.8	111.8	179.1	290.9	525.5	1,231.6

White, female

2006 4/.....	17.0	24.4	34.6	48.0	73.0	104.9	177.9	327.7	649.1	1,887.2
2005 4/.....	15.9	23.7	34.2	46.7	68.9	105.3	170.4	302.7	597.5	1,705.4
2004 4/.....	15.7	23.6	31.6	45.5	66.7	99.1	165.6	291.0	563.1	1,592.9
2003 4/.....	16.3	23.6	32.6	45.3	68.6	103.4	166.8	294.9	562.7	1,647.2
2002.....	16.3	23.6	29.9	46.0	64.2	98.1	161.2	289.2	558.2	1,623.1
2001.....	15.3	22.3	29.7	44.0	64.2	99.2	158.8	285.9	541.6	1,588.0
2000.....	15.5	19.8	30.3	42.2	63.2	93.6	154.2	277.0	539.4	1,541.7
1999.....	13.8	20.2	28.3	39.9	61.0	95.7	151.2	267.4	504.2	1,515.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Black, both sexes

2006 4/.....	37.3	55.0	80.9	106.2	126.5	165.6	256.3	400.1	732.2	1,636.8
2005 4/.....	36.6	58.1	80.1	98.8	128.6	164.8	247.6	401.0	680.4	1,501.2
2004 4/.....	35.5	59.5	78.0	99.9	125.7	165.8	244.3	385.2	624.6	1,405.8
2003 4/.....	38.7	58.2	83.5	96.1	131.7	161.7	241.2	382.2	626.5	1,435.7
2002.....	41.6	60.3	78.4	96.1	123.7	155.7	242.8	369.7	642.1	1,425.4
2001.....	41.2	63.0	79.4	95.6	122.1	163.6	237.0	380.4	612.1	1,418.0
2000.....	41.9	64.3	75.6	95.5	122.2	157.2	233.9	372.5	618.9	1,388.9
1999.....	42.5	61.8	75.4	92.1	117.2	167.2	239.7	369.5	556.5	1,286.2

Black, male

2006 4/.....	40.1	61.6	95.3	127.3	154.1	189.1	303.3	459.1	803.0	1,411.9
2005 4/.....	39.8	65.7	94.2	119.4	151.1	194.4	274.5	447.9	743.7	1,241.0
2004 4/.....	37.6	66.9	95.4	121.7	147.6	200.1	280.0	464.5	659.3	1,291.5
2003 4/.....	42.0	65.9	102.4	113.0	158.7	191.6	289.7	441.8	710.2	1,224.1
2002.....	44.2	70.9	94.6	116.3	146.1	185.2	288.4	419.3	693.1	1,319.8
2001.....	47.0	74.8	95.9	114.8	141.6	196.9	273.3	445.6	659.7	1,313.9
2000.....	48.0	74.7	89.1	114.4	150.1	188.4	279.8	428.6	692.9	1,325.1
1999.....	47.6	73.3	89.5	111.0	134.9	189.5	274.1	434.1	635.7	1,291.3

Black, female

2006 4/.....	34.7	49.2	68.7	88.8	105.1	148.2	224.0	364.8	694.8	1,728.8
2005 4/.....	33.8	51.4	68.1	81.8	111.1	142.8	229.2	373.1	646.9	1,606.2
2004 4/.....	33.7	53.1	63.3	82.0	108.5	140.3	220.1	338.3	606.3	1,451.1
2003 4/.....	35.7	51.5	67.6	82.2	110.7	139.6	208.5	346.7	582.6	1,517.9
2002.....	39.3	51.0	64.6	79.6	106.1	133.8	212.2	340.0	615.5	1,465.9
2001.....	36.2	52.9	65.3	79.8	106.7	138.9	212.8	341.4	587.9	1,456.6
2000.....	36.5	55.2	64.0	79.9	100.2	134.2	203.3	338.6	581.7	1,412.7
1999.....	37.9	52.0	63.5	76.6	103.3	150.6	216.9	330.7	517.3	1,284.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes

2006.....	43.3	47.0	43.7	37.1	35.0	39.6	50.5	81.3	136.0	274.9
2005.....	41.6	44.6	41.6	35.5	36.0	40.1	53.7	82.7	137.0	279.5
2004.....	39.7	42.6	38.6	32.8	33.7	38.4	50.5	79.6	135.9	276.7
2003.....	40.4	40.7	36.6	32.8	32.9	38.1	50.9	78.8	133.5	278.9
2002.....	39.4	38.7	34.3	30.4	32.6	38.3	50.8	80.0	131.1	275.4
2001.....	36.7	36.0	31.9	29.3	31.5	36.8	49.3	80.4	130.4	276.4
2000.....	35.4	34.2	30.8	29.9	32.2	35.4	48.8	75.5	124.6	273.5
1999.....	35.4	34.1	29.2	30.1	31.2	38.7	50.9	78.6	133.3	282.4

All races 3/, male

2006.....	60.8	66.8	63.0	52.9	49.6	55.1	66.4	107.3	176.5	337.1
2005.....	58.7	63.5	60.7	51.3	51.4	54.6	72.6	110.0	180.8	344.2
2004.....	56.0	60.5	56.5	46.9	46.8	52.8	67.0	104.0	179.7	345.5
2003.....	58.0	58.9	52.3	47.2	47.0	52.3	66.8	104.9	175.2	352.5
2002.....	56.8	56.6	49.6	44.8	45.7	53.4	68.9	106.4	170.1	346.6
2001.....	54.1	54.0	46.9	41.8	44.5	49.9	65.6	108.3	173.2	358.5
2000.....	53.1	51.6	46.0	43.1	45.3	48.1	64.6	99.8	164.8	359.7
1999.....	53.1	51.7	43.6	43.6	44.5	53.1	68.2	102.8	181.8	375.4

All races 3/, female

2006.....	25.8	27.8	25.2	22.2	21.7	26.0	37.5	62.3	110.7	245.8
2005.....	24.6	26.0	23.3	20.7	22.0	27.4	38.5	62.8	109.7	249.8
2004.....	23.6	25.2	21.4	19.5	21.7	25.8	37.2	62.1	109.0	245.8
2003.....	23.0	23.0	21.6	19.3	20.1	25.8	38.1	60.3	108.1	246.4
2002.....	22.4	21.3	19.6	16.8	20.7	25.3	36.3	61.4	107.3	244.6
2001.....	19.6	18.5	17.5	17.6	19.6	25.5	36.3	60.9	104.8	242.0
2000.....	18.0	17.3	16.1	17.5	20.4	24.5	36.4	58.5	100.9	238.4
1999.....	18.1	17.0	15.5	17.6	19.2	26.4	37.3	61.8	105.0	244.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

White, both sexes

2006 4/.....	44.9	47.5	43.3	36.3	34.4	39.5	50.9	83.0	140.9	288.6
2005 4/.....	43.2	45.4	40.7	34.7	35.3	39.8	53.8	83.1	140.0	290.9
2004 4/.....	41.3	43.2	37.9	32.0	33.3	38.2	50.5	80.6	140.2	289.9
2003 4/.....	41.5	41.0	35.7	32.1	32.5	37.3	50.9	80.2	136.5	291.1
2002.....	40.5	38.5	33.6	29.6	32.1	37.3	51.0	81.1	134.6	286.1
2001.....	36.8	34.9	31.1	28.5	30.5	36.4	48.9	81.2	133.3	285.5
2000.....	35.2	33.2	29.2	28.9	31.8	34.5	48.1	75.7	127.3	282.8
1999.....	35.2	32.8	27.8	28.9	29.5	37.7	49.5	78.0	135.0	290.4

White, male

2006 4/.....	62.6	66.7	61.2	50.9	48.0	54.7	66.2	107.9	182.9	352.4
2005 4/.....	60.6	63.8	58.6	49.2	49.2	53.7	71.7	110.0	182.9	356.6
2004 4/.....	57.9	61.0	54.7	44.8	45.4	52.4	66.5	104.6	183.9	359.4
2003 4/.....	59.6	58.6	50.6	45.6	46.1	50.8	66.9	105.8	177.3	365.7
2002.....	58.2	56.4	47.8	42.8	44.5	50.9	68.6	106.5	173.0	356.6
2001.....	54.2	51.8	45.1	40.0	42.6	48.9	64.6	108.2	175.2	368.1
2000.....	52.5	49.8	43.2	40.8	43.8	45.9	63.3	98.7	166.4	369.8
1999.....	52.5	49.1	41.1	41.1	42.0	51.1	65.7	100.7	183.4	383.6

White, female

2006 4/.....	26.9	28.4	25.8	22.3	21.7	26.0	38.3	64.3	114.4	258.7
2005 4/.....	25.6	26.9	23.0	20.8	22.4	27.5	39.0	63.2	113.1	260.7
2004 4/.....	24.5	25.5	21.5	19.7	22.0	25.6	37.3	63.0	113.0	258.6
2003 4/.....	23.4	23.4	21.1	19.2	19.9	25.4	37.9	61.7	111.3	258.0
2002.....	22.8	20.7	19.7	16.9	20.7	25.3	36.7	62.9	111.2	255.4
2001.....	19.3	18.1	17.4	17.6	19.3	25.4	36.1	62.0	107.9	251.0
2000.....	17.8	16.8	15.5	17.5	20.7	24.6	35.8	59.6	104.0	247.4
1999.....	17.9	16.6	14.9	17.4	18.0	26.1	36.6	62.0	106.5	253.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Black, both sexes

2006 4/.....	44.2	53.6	56.7	51.2	44.0	45.3	51.4	71.7	93.7	150.5
2005 4/.....	41.1	49.5	57.2	49.7	45.6	47.5	58.9	87.4	110.9	174.8
2004 4/.....	38.4	49.0	50.7	45.2	41.7	43.8	53.8	77.9	100.5	152.3
2003 4/.....	41.7	47.9	49.1	43.7	40.4	47.7	53.8	67.7	105.1	160.2
2002.....	40.0	48.3	45.1	42.0	40.6	47.3	51.9	74.9	96.8	173.4
2001.....	44.3	50.1	44.8	40.6	43.7	45.5	58.0	78.2	104.5	183.6
2000.....	43.8	47.0	47.0	42.0	39.4	46.1	60.2	75.3	100.0	181.1
1999.....	43.6	50.4	44.4	43.4	46.3	50.7	66.6	90.3	124.2	207.6

Black, male

2006 4/.....	65.5	81.1	91.2	81.9	72.1	69.4	77.5	109.6	125.3	199.3
2005 4/.....	60.1	75.7	90.9	80.3	79.6	73.2	90.5	130.5	167.1	225.0
2004 4/.....	56.5	72.2	82.2	73.2	66.7	67.4	82.2	116.0	157.1	213.6
2003 4/.....	60.8	73.9	75.2	70.1	64.8	74.4	74.2	105.0	152.8	225.2
2002.....	57.9	72.0	72.6	71.2	65.3	75.8	78.4	117.6	149.8	260.5
2001.....	66.6	81.7	71.8	65.9	69.8	69.1	82.5	119.4	164.5	255.3
2000.....	68.8	74.5	76.5	71.5	65.4	74.2	86.3	119.5	156.0	275.6
1999.....	69.0	82.7	73.2	71.5	70.6	77.9	101.8	135.3	177.0	316.4

Black, female

2006 4/.....	25.5	29.6	27.6	25.9	22.2	27.5	33.4	49.0	77.0	130.5
2005 4/.....	24.3	26.7	28.7	24.5	19.1	28.4	37.2	61.6	81.2	154.5
2004 4/.....	22.6	28.8	24.2	22.0	22.3	26.3	34.5	55.3	70.7	128.0
2003 4/.....	24.8	25.3	27.0	22.0	21.4	28.0	40.1	45.6	80.1	134.9
2002.....	24.2	27.7	21.8	18.1	21.2	26.1	34.1	49.4	69.2	140.1
2001.....	24.5	22.7	21.8	19.9	23.2	28.0	41.7	53.5	74.0	157.0
2000.....	21.7	23.1	21.9	17.8	18.8	25.3	42.8	48.7	71.8	145.9
1999.....	21.3	22.5	20.1	20.4	27.3	30.6	43.2	63.2	98.1	166.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Transport accidents (V01-V99,Y85)

All races 3/, both sexes

2006.....	16.8	17.2	16.4	15.6	16.1	16.2	17.5	21.6	26.3	24.4
2005.....	17.0	16.8	16.6	16.2	16.3	17.2	18.9	23.0	26.0	26.1
2004.....	16.4	17.1	16.3	15.2	15.9	17.0	18.1	23.6	28.1	27.2
2003.....	17.5	16.8	16.3	15.9	16.0	16.2	19.0	23.9	29.4	29.7
2002.....	17.1	16.9	15.8	14.9	16.7	16.7	20.2	25.0	29.7	28.8
2001.....	17.3	16.3	15.6	15.2	15.4	16.4	19.5	26.2	30.4	28.9
2000.....	16.8	16.2	15.5	15.7	16.4	16.8	19.4	25.1	30.0	31.5
1999.....	16.6	15.9	14.9	16.3	15.7	18.1	20.9	25.9	32.1	32.0

All races 3/, male

2006.....	24.2	25.7	24.5	22.9	22.9	22.9	23.9	30.1	37.5	42.9
2005.....	24.8	24.9	25.0	23.7	23.2	23.9	27.0	32.5	38.0	46.0
2004.....	24.1	24.9	24.4	22.2	22.4	23.2	24.3	32.3	42.6	46.5
2003.....	25.5	24.9	23.6	23.2	22.7	22.5	25.0	32.9	42.1	52.0
2002.....	24.5	25.1	22.7	22.2	22.8	23.4	27.5	34.5	42.6	53.0
2001.....	25.4	24.0	22.5	21.4	22.0	22.1	26.8	36.2	45.5	52.7
2000.....	24.8	23.6	22.3	22.4	22.6	22.6	26.3	34.5	44.0	59.8
1999.....	24.0	23.1	21.5	23.0	21.8	24.8	27.5	36.2	48.3	63.5

All races 3/, female

2006.....	9.4	8.9	8.6	8.6	9.9	10.2	12.3	15.3	19.3	15.7
2005.....	9.2	8.9	8.6	9.0	10.0	11.3	12.4	16.1	18.5	17.0
2004.....	8.9	9.4	8.6	8.7	10.0	11.5	13.1	17.3	19.1	18.5
2003.....	9.5	9.0	9.2	9.0	9.9	10.7	14.2	17.5	21.7	19.9
2002.....	9.9	8.8	9.2	7.9	11.1	10.9	14.3	18.3	21.8	18.3
2001.....	9.4	8.8	8.9	9.4	9.3	11.6	13.8	19.2	21.3	18.9
2000.....	9.0	8.9	8.9	9.5	10.7	11.7	13.9	18.5	21.7	19.9
1999.....	9.4	8.9	8.6	10.0	10.2	12.4	15.7	18.8	22.6	19.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Transport accidents (V01-V99,Y85)

White, both sexes

2006 4/.....	17.3	17.2	16.7	15.4	15.9	16.4	17.7	21.6	26.8	25.3
2005 4/.....	17.5	17.2	16.6	16.4	16.2	17.4	19.0	23.0	26.8	27.1
2004 4/.....	16.8	17.2	16.0	15.2	15.9	17.1	18.2	23.6	29.0	28.4
2003 4/.....	17.7	16.9	16.1	15.7	15.8	16.0	19.2	23.9	30.0	30.5
2002.....	17.5	16.9	15.9	14.8	16.7	16.5	20.5	25.5	30.3	29.6
2001.....	17.5	15.9	15.7	15.1	15.2	16.4	19.5	26.6	31.2	29.9
2000.....	16.8	16.0	15.1	15.6	16.4	16.6	19.2	25.3	30.7	32.2
1999.....	16.7	15.6	14.6	16.1	15.4	18.0	20.8	26.0	32.4	32.3

White, male

2006 4/.....	24.8	25.6	24.7	22.7	22.4	23.0	23.9	29.8	38.4	44.0
2005 4/.....	25.3	25.3	24.8	23.8	23.0	23.7	26.9	32.1	39.0	47.5
2004 4/.....	24.4	25.0	23.7	21.9	22.0	23.2	24.1	32.2	43.2	48.6
2003 4/.....	25.8	24.9	23.3	22.8	22.6	22.2	25.4	32.7	42.6	52.8
2002.....	24.9	25.3	22.5	21.9	22.8	22.7	27.5	34.4	42.8	53.8
2001.....	25.4	23.2	22.6	21.0	21.5	21.7	26.6	36.2	46.6	54.1
2000.....	24.6	23.4	21.7	21.8	22.2	21.9	25.8	34.3	44.5	61.6
1999.....	24.0	22.5	21.0	22.5	21.1	24.6	27.1	35.9	48.4	63.7

White, female

2006 4/.....	9.6	8.9	8.8	8.5	9.9	10.4	12.6	15.5	19.5	16.5
2005 4/.....	9.5	9.1	8.5	9.2	10.0	11.7	12.5	16.2	19.2	17.7
2004 4/.....	9.0	9.5	8.5	8.8	10.3	11.6	13.4	17.3	20.1	19.3
2003 4/.....	9.6	9.1	9.1	8.8	9.6	10.5	14.0	17.6	22.2	20.6
2002.....	10.0	8.6	9.3	8.1	11.0	11.0	14.8	19.1	22.6	19.1
2001.....	9.5	8.7	8.9	9.5	9.3	11.6	13.8	19.8	21.8	19.7
2000.....	9.0	8.7	8.6	9.6	11.1	12.0	13.9	19.0	22.6	20.3
1999.....	9.4	8.8	8.4	9.9	10.1	12.3	15.8	19.1	23.0	19.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Transport accidents (V01-V99,Y85)

Black, both sexes

2006 4/.....	17.1	18.7	16.9	18.9	18.3	15.5	15.4	20.1	19.9	15.6
2005 4/.....	16.8	16.9	18.9	16.9	17.3	17.2	18.8	22.4	16.3	16.6
2004 4/.....	16.7	19.1	19.9	17.2	17.4	16.6	16.2	23.0	19.6	14.7
2003 4/.....	18.7	18.1	18.2	18.9	17.8	18.5	17.9	22.8	21.8	21.0
2002.....	17.1	18.9	16.5	16.8	17.4	17.3	16.9	20.1	24.1	18.4
2001.....	19.4	20.5	16.8	17.8	18.6	18.6	19.0	22.1	19.8	17.5
2000.....	19.3	19.4	19.3	18.1	16.4	18.2	21.3	19.4	20.9	21.6
1999.....	18.0	19.6	18.3	18.7	17.7	18.0	21.0	24.6	28.3	27.7

Black, male

2006 4/.....	26.0	29.5	27.4	29.2	29.5	25.3	25.2	33.6	28.1	36.6
2005 4/.....	26.2	25.5	30.8	27.4	27.8	28.8	30.4	38.9	27.4	33.5
2004 4/.....	25.8	29.6	32.6	27.1	28.6	27.1	26.2	36.9	40.9	28.8
2003 4/.....	28.3	29.1	28.9	30.3	27.6	28.6	24.0	37.4	37.7	42.7
2002.....	25.2	29.1	26.3	29.0	27.3	28.7	28.4	38.1	44.3	46.1
2001.....	30.6	33.8	25.7	28.6	30.9	28.2	26.3	39.0	31.3	35.2
2000.....	31.0	29.3	30.0	31.0	27.5	31.2	33.4	34.5	39.6	45.0
1999.....	28.3	31.2	29.1	29.3	28.4	27.3	31.5	40.4	48.8	68.0

Black, female

2006 4/.....	9.2	9.3	8.0	10.3	9.6	8.3	8.7	12.1	15.6	*
2005 4/.....	8.5	9.4	8.8	8.3	9.1	8.6	10.8	12.5	10.4	9.8
2004 4/.....	8.6	10.0	9.1	9.1	8.7	8.8	9.4	14.8	8.3	9.1
2003 4/.....	10.3	8.4	9.2	9.6	10.2	11.0	13.7	14.1	13.5	12.6
2002.....	10.0	10.0	8.2	6.8	9.6	8.9	9.2	9.3	13.5	*
2001.....	9.4	8.9	9.3	8.9	9.0	11.5	14.1	11.9	13.9	10.9
2000.....	9.0	10.8	10.2	7.6	7.7	8.6	13.3	10.3	11.5	12.9
1999.....	9.0	9.6	9.1	10.1	9.3	11.2	14.0	15.2	18.1	12.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes

2006.....	15.4	15.7	14.8	14.0	14.7	14.9	16.1	20.2	25.0	23.4
2005.....	15.6	15.3	14.8	14.6	14.8	15.8	17.8	21.6	24.6	25.1
2004.....	15.1	15.5	14.6	13.9	14.3	15.5	16.8	22.7	26.9	26.1
2003.....	16.0	15.1	14.6	14.1	14.3	14.8	17.7	22.5	28.2	28.8
2002.....	15.7	15.2	14.4	13.4	15.2	15.3	18.9	23.8	28.4	28.0
2001.....	15.5	14.5	13.9	13.7	13.7	14.8	18.0	24.6	28.9	27.8
2000.....	15.1	14.6	13.8	14.2	14.7	15.1	18.1	23.8	28.5	30.4
1999.....	14.8	14.0	13.1	14.4	13.9	16.4	19.3	24.1	30.1	30.4

All races 3/, male

2006.....	21.9	23.2	22.0	20.5	20.4	20.8	21.5	27.8	35.0	41.0
2005.....	22.6	22.2	22.1	21.0	20.7	21.5	25.2	30.1	35.7	44.1
2004.....	21.9	22.3	21.6	19.9	19.6	20.8	22.0	30.8	40.4	44.2
2003.....	23.1	22.0	20.8	20.3	19.6	20.1	22.8	30.5	40.0	50.0
2002.....	22.1	22.2	20.3	19.6	20.2	20.8	25.0	32.3	39.9	51.7
2001.....	22.5	21.2	19.7	18.9	19.2	19.2	24.2	33.6	43.1	51.1
2000.....	21.9	21.1	19.6	19.9	19.7	19.8	23.9	32.2	41.4	57.5
1999.....	21.0	20.0	18.6	19.8	18.7	22.0	24.7	33.6	45.2	59.8

All races 3/, female

2006.....	8.9	8.4	8.0	7.9	9.4	9.6	11.7	14.6	18.7	15.2
2005.....	8.7	8.5	7.9	8.5	9.4	10.7	11.8	15.5	17.8	16.4
2004.....	8.3	8.8	8.0	8.2	9.5	10.9	12.6	16.9	18.6	17.9
2003.....	9.0	8.3	8.7	8.3	9.4	10.3	13.7	16.9	21.0	19.5
2002.....	9.4	8.4	8.7	7.5	10.6	10.6	14.1	17.7	21.4	17.7
2001.....	8.7	8.0	8.3	8.8	8.8	11.0	13.1	18.2	20.4	18.0
2000.....	8.4	8.3	8.2	9.0	10.2	11.0	13.5	17.9	20.9	19.3
1999.....	8.6	8.1	8.0	9.4	9.5	11.6	15.0	17.5	21.3	18.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, both sexes

2006 4/.....	15.8	15.7	15.1	13.9	14.4	15.0	16.3	20.3	25.4	24.4
2005 4/.....	16.1	15.7	14.9	14.7	14.7	15.9	17.8	21.6	25.5	26.1
2004 4/.....	15.4	15.6	14.3	13.9	14.3	15.6	16.8	22.8	27.7	27.3
2003 4/.....	16.2	15.2	14.4	13.8	14.0	14.6	17.8	22.6	28.8	29.5
2002.....	16.0	15.2	14.3	13.3	15.1	15.1	19.2	24.3	29.0	28.8
2001.....	15.6	14.2	13.9	13.6	13.5	14.7	17.9	25.0	29.7	28.8
2000.....	15.0	14.5	13.4	14.1	14.7	14.9	17.9	24.0	29.2	31.1
1999.....	14.8	13.7	12.9	14.1	13.5	16.3	19.2	24.3	30.6	30.7

White, male

2006 4/.....	22.4	23.1	22.2	20.2	19.9	20.8	21.4	27.4	35.6	42.2
2005 4/.....	23.1	22.7	22.1	21.0	20.5	21.4	25.0	29.7	36.7	45.6
2004 4/.....	22.2	22.3	20.9	19.6	19.2	20.7	21.7	30.8	40.8	46.3
2003 4/.....	23.2	22.1	20.5	19.7	19.4	19.9	23.0	30.3	40.6	50.7
2002.....	22.5	22.3	20.0	19.3	20.0	20.1	25.0	32.4	40.1	52.4
2001.....	22.5	20.5	19.7	18.7	18.5	19.0	23.9	33.6	44.3	52.6
2000.....	21.6	20.9	18.9	19.3	19.2	19.1	23.4	32.0	41.7	59.2
1999.....	20.9	19.4	18.1	19.2	18.1	21.8	24.4	33.3	45.5	60.3

White, female

2006 4/.....	9.1	8.3	8.2	7.7	9.2	9.8	12.0	14.9	19.0	16.0
2005 4/.....	9.0	8.7	7.8	8.7	9.3	11.0	11.9	15.5	18.5	17.1
2004 4/.....	8.4	8.9	7.9	8.3	9.8	11.1	12.8	16.9	19.6	18.7
2003 4/.....	9.0	8.4	8.6	8.1	9.1	10.0	13.5	17.1	21.5	20.1
2002.....	9.5	8.2	8.8	7.6	10.5	10.7	14.5	18.5	22.2	18.5
2001.....	8.8	8.0	8.2	8.8	8.8	11.0	13.0	19.0	20.9	18.9
2000.....	8.4	8.1	7.9	9.1	10.6	11.3	13.5	18.4	21.8	19.7
1999.....	8.6	8.0	7.7	9.3	9.3	11.6	15.1	18.0	21.8	18.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2006 4/.....	15.8	17.2	15.5	17.3	17.6	14.6	14.8	18.8	18.9	14.0
2005 4/.....	15.6	15.0	16.7	15.9	16.2	15.9	18.0	21.4	15.0	15.8
2004 4/.....	15.4	17.3	18.3	15.9	15.5	15.6	15.3	20.8	19.3	14.7
2003 4/.....	17.5	16.2	16.8	17.5	16.4	16.9	17.2	21.3	20.6	20.7
2002.....	15.9	16.9	15.1	15.0	16.2	15.9	15.8	18.4	22.5	18.1
2001.....	17.6	18.1	15.3	15.9	16.9	16.5	18.3	20.1	19.0	16.6
2000.....	17.9	17.6	18.3	16.7	15.4	16.6	20.1	18.1	20.0	20.3
1999.....	16.6	17.8	16.3	17.1	15.8	16.6	19.6	22.9	26.0	25.5

Black, male

2006 4/.....	24.1	26.6	24.7	26.5	27.7	23.7	23.7	31.3	27.4	33.1
2005 4/.....	24.3	22.4	26.8	25.5	26.0	25.7	29.4	36.7	26.0	32.5
2004 4/.....	23.5	26.7	29.7	24.5	25.0	25.4	24.3	33.2	40.2	28.8
2003 4/.....	26.4	25.8	26.0	27.6	24.5	25.1	23.0	35.1	34.7	41.7
2002.....	23.0	25.6	23.6	25.3	24.9	25.4	25.7	33.5	40.5	45.0
2001.....	27.5	29.9	23.2	24.7	28.5	23.9	24.6	36.2	30.5	35.2
2000.....	28.4	26.3	28.1	28.2	25.7	27.8	31.4	32.1	38.8	43.8
1999.....	25.6	28.2	25.5	26.3	24.6	24.9	28.5	38.9	45.3	60.9

Black, female

2006 4/.....	8.6	9.0	7.7	9.7	9.6	8.0	8.7	11.2	14.5	*
2005 4/.....	7.9	8.5	8.1	8.0	8.6	8.6	10.2	12.3	9.3	9.0
2004 4/.....	8.3	9.1	8.8	8.8	8.0	8.3	9.2	13.5	8.3	9.1
2003 4/.....	9.7	7.9	9.1	9.2	10.1	10.8	13.3	13.0	13.2	12.6
2002.....	9.6	9.4	8.0	6.6	9.4	8.9	9.2	9.3	13.1	*
2001.....	8.9	8.0	8.7	8.7	7.8	11.1	14.1	10.5	13.1	9.7
2000.....	8.6	10.0	9.9	7.3	7.4	8.3	12.7	9.7	10.6	11.6
1999.....	8.7	8.9	8.5	9.6	8.8	10.4	13.7	13.2	16.4	12.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2006.....	0.5	0.5	0.6	0.5	0.4	0.5	0.5	0.6	0.6	0.5
2005.....	0.5	0.5	0.7	0.5	0.5	0.4	0.4	0.6	0.6	0.5
2004.....	0.5	0.6	0.5	0.4	0.5	0.4	0.6	0.4	0.6	0.6
2003.....	0.6	0.6	0.5	0.6	0.5	0.4	0.5	0.6	0.7	0.5
2002.....	0.5	0.6	0.5	0.4	0.4	0.3	0.4	0.5	0.6	*
2001.....	0.6	0.7	0.6	0.5	0.6	0.7	0.6	0.8	0.8	0.7
2000.....	0.7	0.6	0.6	0.5	0.5	0.5	0.5	0.7	0.8	0.8
1999.....	0.9	0.9	0.7	0.8	0.8	0.8	0.7	1.3	1.4	1.3

All races 3/, male

2006.....	0.9	0.9	1.0	0.8	0.7	0.7	1.0	1.0	1.1	*
2005.....	0.8	0.9	1.2	0.7	0.8	0.7	0.6	1.1	1.1	*
2004.....	0.8	0.9	0.8	0.7	0.7	0.6	0.8	0.6	1.2	*
2003.....	0.9	1.1	0.8	0.9	0.8	0.8	0.7	0.8	1.3	*
2002.....	0.9	1.0	0.8	0.7	0.7	0.6	0.8	0.8	1.2	*
2001.....	1.0	1.0	0.9	0.8	0.9	1.1	0.9	1.2	*	*
2000.....	1.2	0.9	0.9	0.7	1.0	0.9	0.9	1.1	1.3	*
1999.....	1.5	1.4	1.2	1.2	1.1	1.2	1.2	1.7	2.1	2.9

All races 3/, female

2006.....	*	0.2	0.2	0.3	*	*	*	*	*	*
2005.....	0.2	*	0.4	*	*	*	*	*	*	*
2004.....	0.2	0.2	*	*	*	*	*	*	*	*
2003.....	0.3	0.2	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.2	0.4	0.2	*	*	*	*	0.5	0.7	*
2000.....	0.3	0.3	0.4	*	*	*	*	*	*	*
1999.....	0.4	0.4	0.3	0.4	0.4	0.5	0.4	0.9	1.0	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, both sexes

2006 4/.....	0.5	0.5	0.6	0.5	0.4	0.5	0.5	0.6	0.6	0.5
2005 4/.....	0.5	0.5	0.6	0.5	0.4	0.4	0.3	0.6	0.7	0.5
2004 4/.....	0.5	0.5	0.4	0.4	0.4	0.4	0.6	0.3	0.7	0.6
2003 4/.....	0.6	0.6	0.5	0.5	0.5	0.4	0.4	0.6	0.7	0.5
2002.....	0.5	0.5	0.5	0.4	0.4	0.3	0.4	0.4	0.6	*
2001.....	0.6	0.6	0.6	0.5	0.5	0.6	0.6	0.8	0.7	0.6
2000.....	0.7	0.6	0.6	0.4	0.6	0.5	0.5	0.7	0.8	0.8
1999.....	0.9	0.9	0.7	0.8	0.8	0.8	0.7	1.2	1.4	1.3

White, male

2006 4/.....	0.9	0.8	0.9	0.8	0.6	0.7	1.0	1.0	1.2	*
2005 4/.....	0.7	0.8	0.9	0.8	0.6	0.5	*	1.0	1.1	*
2004 4/.....	0.8	0.8	0.7	0.7	0.7	0.6	0.8	*	1.4	*
2003 4/.....	0.9	0.9	0.8	0.9	0.8	0.7	0.7	0.8	1.2	*
2002.....	0.8	0.9	0.8	0.6	0.7	0.5	0.7	*	1.3	*
2001.....	0.9	0.9	0.9	0.8	0.9	1.0	0.9	1.3	*	*
2000.....	1.2	0.8	0.8	0.6	1.0	0.8	0.9	1.1	1.3	*
1999.....	1.5	1.3	1.2	1.3	1.1	1.2	1.1	1.7	2.0	2.8

White, female

2006 4/.....	*	0.2	0.2	0.3	*	*	*	*	*	*
2005 4/.....	0.2	*	0.3	*	*	*	*	*	*	*
2004 4/.....	0.2	*	*	*	*	*	*	*	*	*
2003 4/.....	0.2	0.2	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	0.2	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	0.3	0.4	*	*	*	*	*	*	*
1999.....	0.4	0.5	0.3	0.4	0.4	0.5	*	0.9	1.0	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes

2006.....	0.9	0.9	0.9	1.0	1.0	0.8	0.9	0.8	0.7	0.4
2005.....	0.8	1.0	1.0	1.1	1.0	1.0	0.7	0.7	0.7	0.5
2004.....	0.8	1.0	1.2	0.9	1.1	1.0	0.8	0.5	0.5	0.5
2003.....	0.9	1.1	1.1	1.2	1.2	0.9	0.8	0.8	0.5	*
2002.....	0.9	1.1	1.0	1.1	1.1	1.0	0.8	0.8	0.7	0.4
2001.....	1.2	1.1	1.2	1.0	1.1	1.0	0.9	0.8	0.7	0.5
2000.....	1.0	0.9	1.0	1.0	1.1	1.2	0.7	0.6	0.7	*
1999.....	0.9	1.0	1.0	1.1	1.1	0.9	0.8	0.5	0.6	*

All races 3/, male

2006.....	1.4	1.6	1.5	1.6	1.7	1.4	1.4	1.3	1.4	*
2005.....	1.4	1.7	1.8	1.9	1.7	1.7	1.3	1.3	1.2	*
2004.....	1.3	1.7	2.0	1.5	2.1	1.8	1.5	0.9	1.0	*
2003.....	1.6	1.7	2.0	2.0	2.3	1.7	1.5	1.6	*	*
2002.....	1.5	1.9	1.6	1.9	1.9	2.0	1.7	1.4	1.5	*
2001.....	1.8	1.8	1.9	1.7	1.9	1.8	1.7	1.3	1.4	*
2000.....	1.7	1.6	1.8	1.8	1.9	2.0	1.4	1.2	1.4	*
1999.....	1.5	1.7	1.8	2.0	1.9	1.6	1.6	0.9	*	*

All races 3/, female

2006.....	0.4	0.3	0.3	0.4	0.4	*	*	*	*	*
2005.....	0.3	0.3	0.3	0.3	0.4	0.4	*	*	*	*
2004.....	0.3	0.4	0.4	0.3	*	*	*	*	*	*
2003.....	0.3	0.4	0.3	0.4	*	*	*	*	*	*
2002.....	0.3	0.3	0.3	0.3	0.4	*	*	*	*	*
2001.....	0.5	0.4	0.4	0.4	*	*	*	0.5	*	*
2000.....	0.3	0.3	0.3	*	0.4	0.4	*	*	*	*
1999.....	0.4	0.3	0.3	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

White, both sexes

2006 4/.....	1.0	1.0	1.0	1.0	1.1	0.9	0.9	0.8	0.8	*
2005 4/.....	0.9	1.0	1.1	1.2	1.1	1.1	0.8	0.8	0.7	0.5
2004 4/.....	0.9	1.1	1.3	0.9	1.2	1.1	0.8	0.5	0.6	0.6
2003 4/.....	1.0	1.1	1.2	1.3	1.3	0.9	0.9	0.8	0.5	*
2002.....	1.0	1.2	1.0	1.1	1.2	1.0	0.9	0.8	0.7	*
2001.....	1.3	1.1	1.2	1.0	1.2	1.0	1.0	0.8	0.7	*
2000.....	1.1	1.0	1.1	1.1	1.2	1.2	0.8	0.6	0.7	*
1999.....	1.0	1.1	1.1	1.1	1.1	0.9	0.9	0.5	0.5	*

White, male

2006 4/.....	1.5	1.7	1.6	1.7	1.9	1.5	1.5	1.3	1.6	*
2005 4/.....	1.6	1.8	1.8	2.1	1.9	1.8	1.4	1.4	1.2	*
2004 4/.....	1.4	1.9	2.1	1.6	2.1	1.9	1.6	0.9	1.0	*
2003 4/.....	1.7	1.9	2.1	2.2	2.5	1.6	1.7	1.6	*	*
2002.....	1.6	2.1	1.7	1.9	2.1	2.0	1.8	1.3	1.5	*
2001.....	2.0	1.8	2.0	1.6	2.1	1.8	1.8	1.3	1.5	*
2000.....	1.9	1.6	2.0	1.9	2.0	2.0	1.5	1.2	1.5	*
1999.....	1.5	1.8	1.8	2.0	1.9	1.6	1.6	0.9	*	*

White, female

2006 4/.....	0.4	0.3	0.4	0.4	0.4	*	*	*	*	*
2005 4/.....	0.3	0.3	0.4	0.3	0.4	0.4	*	*	*	*
2004 4/.....	0.4	0.4	0.4	0.3	*	*	*	*	*	*
2003 4/.....	0.4	0.4	0.4	0.4	*	*	*	*	*	*
2002.....	0.3	0.2	0.4	0.4	0.4	*	*	*	*	*
2001.....	0.6	0.4	0.5	0.5	*	*	*	*	*	*
2000.....	0.4	0.3	0.3	*	0.4	*	*	*	*	*
1999.....	0.4	0.3	0.3	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes

2006.....	26.5	29.9	27.4	21.6	18.9	23.4	33.0	59.8	109.7	250.5
2005.....	24.6	27.8	25.0	19.4	19.8	22.9	34.8	59.7	111.0	253.4
2004.....	23.3	25.5	22.3	17.6	17.8	21.4	32.4	56.0	107.8	249.5
2003.....	22.9	23.9	20.3	17.0	16.9	21.9	31.9	54.9	104.1	249.2
2002.....	22.3	21.8	18.4	15.5	16.0	21.7	30.6	55.0	101.4	246.7
2001.....	19.4	19.7	16.3	14.1	16.1	20.3	29.8	54.3	100.1	247.5
2000.....	18.6	18.0	15.3	14.2	15.9	18.6	29.4	50.4	94.6	242.0
1999.....	18.8	18.2	14.3	13.9	15.5	20.6	30.0	52.7	101.3	250.3

All races 3/, male

2006.....	36.6	41.1	38.5	30.0	26.8	32.2	42.5	77.2	139.0	294.3
2005.....	33.9	38.7	35.7	27.5	28.3	30.7	45.6	77.5	142.8	298.3
2004.....	31.9	35.6	32.1	24.7	24.5	29.6	42.8	71.7	137.0	299.0
2003.....	32.4	34.1	28.7	24.0	24.3	29.7	41.8	72.0	133.0	300.5
2002.....	32.3	31.5	26.8	22.6	22.9	30.0	41.4	71.9	127.5	293.6
2001.....	28.7	30.0	24.3	20.5	22.5	27.8	38.8	72.1	127.8	305.8
2000.....	28.3	28.0	23.7	20.7	22.7	25.5	38.4	65.3	120.8	299.8
1999.....	29.1	28.6	22.0	20.6	22.7	28.3	40.7	66.6	133.5	311.9

All races 3/, female

2006.....	16.4	18.9	16.7	13.6	11.8	15.8	25.3	47.0	91.4	230.1
2005.....	15.3	17.1	14.7	11.7	12.0	16.0	26.1	46.7	91.3	232.8
2004.....	14.7	15.8	12.8	10.8	11.7	14.3	24.1	44.8	89.9	227.3
2003.....	13.5	14.1	12.3	10.4	10.2	15.1	23.9	42.7	86.4	226.5
2002.....	12.5	12.5	10.4	8.9	9.7	14.5	21.9	43.1	85.5	226.3
2001.....	10.2	9.7	8.6	8.2	10.3	13.9	22.6	41.8	83.5	223.1
2000.....	9.0	8.4	7.3	8.0	9.7	12.8	22.4	40.0	79.2	218.4
1999.....	8.7	8.1	6.9	7.6	9.0	14.0	21.7	43.0	82.4	225.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nontransport accidents (W00-X59,Y86)

White, both sexes

2006 4/.....	27.6	30.3	26.6	20.9	18.5	23.1	33.2	61.4	114.1	263.4
2005 4/.....	25.7	28.1	24.1	18.3	19.1	22.4	34.8	60.2	113.2	263.9
2004 4/.....	24.5	26.0	21.9	16.8	17.4	21.2	32.3	57.0	111.2	261.5
2003 4/.....	23.8	24.0	19.5	16.5	16.7	21.4	31.8	56.2	106.5	260.6
2002.....	23.1	21.6	17.7	14.7	15.5	20.8	30.5	55.6	104.3	256.5
2001.....	19.3	19.0	15.4	13.4	15.3	20.0	29.4	54.6	102.1	255.7
2000.....	18.3	17.2	14.1	13.3	15.3	17.9	28.9	50.4	96.5	250.6
1999.....	18.5	17.1	13.2	12.9	14.2	19.7	28.7	51.9	102.6	258.1

White, male

2006 4/.....	37.8	41.1	36.5	28.2	25.6	31.7	42.3	78.2	144.5	308.4
2005 4/.....	35.2	38.5	33.8	25.4	26.2	29.9	44.8	77.9	143.9	309.1
2004 4/.....	33.5	36.0	30.9	22.9	23.4	29.2	42.4	72.5	140.7	310.8
2003 4/.....	33.8	33.8	27.2	22.8	23.4	28.6	41.5	73.1	134.7	312.9
2002.....	33.3	31.1	25.2	20.9	21.7	28.2	41.1	72.1	130.1	302.8
2001.....	28.7	28.6	22.4	19.0	21.1	27.1	38.1	72.0	128.6	314.0
2000.....	27.9	26.4	21.5	19.0	21.6	24.0	37.5	64.4	122.0	308.2
1999.....	28.5	26.6	20.0	18.6	20.9	26.5	38.6	64.8	135.0	319.8

White, female

2006 4/.....	17.3	19.5	17.0	13.8	11.8	15.6	25.7	48.8	94.9	242.2
2005 4/.....	16.1	17.8	14.5	11.6	12.5	15.8	26.5	47.1	94.0	243.0
2004 4/.....	15.5	16.0	13.1	10.9	11.7	14.1	24.0	45.7	92.9	239.3
2003 4/.....	13.8	14.3	12.0	10.4	10.4	14.9	23.9	44.0	89.1	237.4
2002.....	12.8	12.1	10.4	8.8	9.7	14.3	21.9	43.8	88.5	236.3
2001.....	9.7	9.4	8.5	8.1	10.0	13.8	22.3	42.2	86.1	231.2
2000.....	8.7	8.1	6.9	7.9	9.5	12.6	21.9	40.6	81.4	227.2
1999.....	8.5	7.8	6.5	7.4	8.0	13.7	20.8	42.9	83.5	233.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Nontransport accidents (W00-X59,Y86)

Black, both sexes

2006 4/.....	27.2	34.9	39.8	32.3	25.7	29.8	36.0	51.6	73.8	134.9
2005 4/.....	24.3	32.6	38.3	32.8	28.3	30.3	40.1	65.0	94.7	158.2
2004 4/.....	21.8	29.9	30.9	27.9	24.3	27.2	37.6	54.8	81.0	137.6
2003 4/.....	22.9	29.9	30.9	24.8	22.6	29.3	36.0	44.9	83.3	139.2
2002.....	22.9	29.4	28.6	25.2	23.2	29.9	35.0	54.8	72.7	155.0
2001.....	24.9	29.6	28.0	22.8	25.1	26.9	39.0	56.1	84.8	166.1
2000.....	24.5	27.6	27.7	23.9	22.9	27.8	38.8	56.0	79.1	159.5
1999.....	25.6	30.7	26.2	24.6	28.6	32.7	45.6	65.7	95.9	179.8

Black, male

2006 4/.....	39.6	51.7	63.7	52.6	42.6	44.1	52.3	76.1	97.2	162.6
2005 4/.....	34.0	50.2	60.1	52.9	51.8	44.4	60.1	91.6	139.8	191.5
2004 4/.....	30.7	42.6	49.6	46.2	38.1	40.2	56.0	79.2	116.2	184.8
2003 4/.....	32.4	44.8	46.3	39.8	37.2	45.8	50.3	67.6	115.2	182.5
2002.....	32.7	42.9	46.3	42.1	38.0	47.1	50.1	79.5	105.5	214.4
2001.....	36.0	47.8	46.1	37.2	38.8	40.9	56.2	80.4	133.2	220.1
2000.....	37.8	45.3	46.5	40.5	37.9	42.9	52.8	85.1	116.3	230.6
1999.....	40.7	51.5	44.1	42.2	42.2	50.6	70.2	95.0	128.2	248.4

Black, female

2006 4/.....	16.2	20.2	19.6	15.6	12.5	19.2	24.7	37.0	61.5	123.6
2005 4/.....	15.8	17.3	19.8	16.2	10.1	19.7	26.4	49.1	70.8	144.8
2004 4/.....	13.9	18.9	15.1	12.9	13.5	17.4	25.0	40.4	62.4	118.9
2003 4/.....	14.5	16.8	17.8	12.5	11.1	17.0	26.4	31.5	66.5	122.3
2002.....	14.2	17.7	13.6	11.2	11.6	17.2	24.9	40.1	55.7	132.3
2001.....	15.2	13.8	12.5	11.0	14.2	16.5	27.6	41.6	60.1	146.1
2000.....	12.7	12.3	11.7	10.3	11.1	16.7	29.5	38.4	60.3	133.1
1999.....	12.3	12.9	11.0	10.2	18.0	19.4	29.3	48.1	79.9	154.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Falls (W00-W19)

All races 3/, both sexes

2006.....	1.8	2.5	3.5	4.4	6.0	9.5	16.5	33.5	66.2	151.6
2005.....	1.7	2.4	3.2	3.9	5.9	8.8	16.7	31.9	63.6	147.7
2004.....	1.8	2.6	3.2	4.1	5.8	8.9	16.1	30.3	61.8	143.3
2003.....	1.7	2.3	2.8	3.7	5.3	8.1	14.7	28.7	57.5	135.9
2002.....	1.8	2.3	2.8	3.4	4.9	7.9	13.9	27.1	54.0	130.4
2001.....	1.6	2.5	2.7	3.3	4.9	7.8	12.4	25.2	49.9	121.5
2000.....	1.6	2.0	2.6	3.3	4.6	6.2	12.0	22.1	44.5	112.6
1999.....	1.7	2.3	2.2	3.0	4.6	6.9	11.3	22.3	45.5	110.2

All races 3/, male

2006.....	2.8	4.0	5.5	6.4	8.5	13.3	20.6	42.4	83.7	181.1
2005.....	2.6	3.7	5.0	5.9	8.5	12.1	21.4	41.0	82.1	174.8
2004.....	2.9	4.1	4.9	6.2	8.1	12.4	21.6	38.4	78.7	174.4
2003.....	2.7	3.6	4.4	5.3	7.3	10.7	18.9	36.9	73.8	165.1
2002.....	2.9	3.6	4.3	5.2	7.1	10.7	18.9	35.1	67.9	158.9
2001.....	2.8	4.2	4.4	5.0	7.1	10.6	16.0	34.0	64.1	155.7
2000.....	2.7	3.4	4.2	5.0	6.9	8.6	15.7	29.4	57.3	145.4
1999.....	2.8	3.8	3.6	4.7	7.0	9.8	15.3	27.9	61.6	142.5

All races 3/, female

2006.....	0.7	1.1	1.5	2.4	3.7	6.2	13.1	27.0	55.3	137.8
2005.....	0.7	1.2	1.4	2.1	3.6	6.0	13.0	25.4	52.2	135.2
2004.....	0.7	1.1	1.5	2.0	3.6	5.8	11.7	24.5	51.4	129.2
2003.....	0.6	1.1	1.3	2.1	3.5	5.8	11.3	22.8	47.5	122.9
2002.....	0.7	1.0	1.4	1.8	3.0	5.5	9.9	21.4	45.5	118.1
2001.....	0.6	0.9	1.2	1.6	2.9	5.4	9.5	19.0	41.3	107.2
2000.....	0.6	0.8	1.2	1.8	2.6	4.2	9.1	17.1	36.9	99.2
1999.....	0.6	0.8	0.9	1.5	2.4	4.5	8.1	18.4	36.0	97.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Falls (W00-W19)

White, both sexes

2006 4/.....	1.9	2.6	3.5	4.4	6.1	9.7	17.3	35.4	69.9	161.1
2005 4/.....	1.8	2.5	3.2	4.0	6.0	9.1	17.6	33.4	66.6	156.7
2004 4/.....	1.9	2.6	3.2	4.0	5.8	9.2	16.7	32.0	65.3	152.6
2003 4/.....	1.7	2.3	2.7	3.7	5.4	8.2	15.2	30.1	60.0	144.4
2002.....	1.8	2.3	2.8	3.4	5.1	8.1	14.4	28.3	57.1	138.0
2001.....	1.7	2.5	2.8	3.2	4.9	8.0	12.8	26.3	52.4	128.0
2000.....	1.7	2.1	2.5	3.2	4.6	6.2	12.3	23.1	46.6	119.1
1999.....	1.7	2.2	2.1	2.9	4.3	6.9	11.4	23.0	47.7	115.9

White, male

2006 4/.....	3.0	4.1	5.4	6.4	8.5	13.5	21.4	44.0	88.0	191.2
2005 4/.....	2.8	3.7	5.0	5.9	8.4	12.4	22.4	43.1	84.6	183.8
2004 4/.....	3.1	4.2	4.8	6.1	7.9	13.0	22.0	40.4	82.7	183.7
2003 4/.....	2.8	3.6	4.1	5.3	7.3	10.6	19.5	38.6	76.4	175.1
2002.....	3.0	3.7	4.2	5.1	7.2	10.8	19.3	36.4	71.2	166.4
2001.....	2.8	4.1	4.4	4.9	7.0	10.8	16.5	35.1	66.6	162.3
2000.....	2.7	3.5	3.9	4.8	6.9	8.4	16.2	30.3	59.7	152.8
1999.....	2.7	3.7	3.3	4.4	6.6	9.6	15.3	28.3	64.2	149.1

White, female

2006 4/.....	0.7	1.1	1.6	2.5	3.8	6.3	14.0	29.1	58.4	147.0
2005 4/.....	0.8	1.3	1.4	2.1	3.7	6.2	13.6	26.3	55.3	144.2
2004 4/.....	0.7	1.1	1.6	2.1	3.8	5.8	12.3	25.9	54.6	138.6
2003 4/.....	0.6	1.0	1.3	2.1	3.6	6.1	11.6	24.0	49.9	130.8
2002.....	0.7	1.0	1.5	1.8	3.1	5.8	10.3	22.6	48.5	125.6
2001.....	0.5	0.9	1.2	1.6	2.9	5.6	9.9	20.0	43.9	113.6
2000.....	0.6	0.7	1.2	1.8	2.6	4.3	9.3	18.1	38.8	105.4
1999.....	0.6	0.8	0.9	1.5	2.2	4.6	8.3	19.3	37.9	102.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Falls (W00-W19)

Black, both sexes

2006 4/.....	1.2	2.6	3.8	4.5	5.0	8.9	10.5	16.4	32.9	54.0
2005 4/.....	1.0	2.4	3.9	4.1	5.7	7.2	10.9	20.5	32.5	52.7
2004 4/.....	1.7	2.6	3.5	5.1	6.3	7.9	12.8	16.8	25.5	46.5
2003 4/.....	1.5	3.0	3.6	4.3	5.5	7.6	10.6	13.4	28.2	44.1
2002.....	1.6	2.5	3.2	3.8	4.7	6.1	10.9	14.8	18.1	49.3
2001.....	2.0	2.6	3.3	3.9	4.9	7.9	9.7	16.3	21.9	44.5
2000.....	1.6	2.0	3.6	4.5	5.1	7.1	9.7	11.6	20.3	38.8
1999.....	1.7	3.2	3.6	4.5	6.7	9.0	9.5	13.4	23.1	45.3

Black, male

2006 4/.....	2.2	4.3	6.4	7.6	8.1	12.9	16.3	25.5	45.9	67.9
2005 4/.....	1.7	4.4	6.4	6.5	9.8	10.9	14.5	23.7	52.7	63.2
2004 4/.....	2.6	4.1	6.0	8.8	11.8	9.7	20.7	23.7	38.0	71.5
2003 4/.....	2.6	4.8	5.8	6.5	8.1	12.8	12.2	18.3	35.4	56.3
2002.....	2.6	4.1	5.6	6.3	7.3	9.2	15.4	22.0	28.3	77.2
2001.....	3.3	4.9	6.0	6.4	7.7	11.9	13.5	24.2	36.1	68.1
2000.....	2.6	3.4	6.3	7.7	8.7	10.7	12.4	18.4	26.1	61.1
1999.....	3.3	5.6	6.0	7.6	10.2	14.8	14.6	20.4	31.4	58.6

Black, female

2006 4/.....	*	*	*	2.0	*	5.9	6.6	11.0	26.0	48.3
2005 4/.....	*	*	1.8	2.1	*	4.4	8.4	18.7	21.9	48.4
2004 4/.....	*	*	*	*	*	6.7	7.5	12.7	18.9	36.6
2003 4/.....	*	1.4	1.7	2.5	3.5	3.7	9.5	10.6	24.4	39.3
2002.....	*	*	*	*	*	3.7	7.8	10.4	12.7	38.6
2001.....	*	*	*	*	*	4.9	7.2	11.6	14.7	35.8
2000.....	*	*	*	*	*	4.4	8.0	7.4	17.4	30.5
1999.....	*	*	*	*	4.0	4.6	6.1	9.2	19.0	40.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental drowning and submersion (W65-W74)

All races 3/, both sexes

2006.....	1.1	1.2	1.0	1.0	0.9	1.1	1.2	1.6	1.4	1.4
2005.....	1.3	1.2	1.1	0.9	0.9	0.9	1.2	1.4	1.4	1.8
2004.....	1.0	1.1	1.0	0.9	0.9	1.1	1.0	1.0	1.5	1.6
2003.....	1.1	1.0	1.0	1.0	0.9	1.0	1.2	1.3	1.3	1.8
2002.....	1.1	1.0	0.9	0.8	0.8	0.8	1.0	1.2	1.5	1.7
2001.....	1.1	1.0	0.8	0.8	0.9	0.8	0.9	1.1	1.4	1.7
2000.....	1.0	0.9	1.0	0.9	0.9	0.8	1.1	1.0	1.7	1.5
1999.....	1.0	1.0	1.0	0.9	0.9	0.9	1.1	1.2	1.4	1.6

All races 3/, male

2006.....	1.8	2.0	1.6	1.6	1.4	1.6	1.9	2.5	2.3	2.4
2005.....	2.0	2.0	1.7	1.5	1.5	1.4	1.8	2.1	2.5	3.9
2004.....	1.7	1.8	1.7	1.5	1.5	1.7	1.6	1.7	2.8	2.9
2003.....	1.8	1.6	1.6	1.6	1.5	1.6	1.9	1.8	2.3	3.3
2002.....	1.9	1.6	1.5	1.3	1.2	1.4	1.8	1.8	2.5	3.6
2001.....	1.7	1.8	1.3	1.2	1.4	1.4	1.6	1.8	2.4	3.5
2000.....	1.7	1.6	1.7	1.6	1.5	1.4	1.8	1.2	2.7	2.8
1999.....	1.7	1.7	1.8	1.5	1.5	1.5	1.7	2.1	2.7	2.3

All races 3/, female

2006.....	0.5	0.5	0.4	0.5	0.4	0.7	0.6	0.9	0.8	0.9
2005.....	0.5	0.5	0.4	0.3	0.3	0.5	0.7	0.8	0.7	0.9
2004.....	0.4	0.4	0.5	0.4	0.5	0.5	0.5	0.5	0.7	1.0
2003.....	0.3	0.3	0.4	0.3	0.4	0.6	0.6	0.8	0.6	1.1
2002.....	0.4	0.4	0.3	0.3	0.4	*	0.4	0.7	0.9	0.9
2001.....	0.4	0.3	0.3	0.4	0.4	*	*	0.7	0.8	1.0
2000.....	0.3	0.3	0.3	*	0.4	*	0.6	0.8	1.1	1.0
1999.....	0.3	0.3	0.3	0.4	0.4	0.4	0.7	0.6	0.7	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental drowning and submersion (W65-W74)

White, both sexes

2006 4/.....	1.2	1.3	1.0	1.0	0.9	1.1	1.1	1.5	1.5	1.4
2005 4/.....	1.3	1.2	1.0	0.9	0.9	0.9	1.2	1.3	1.2	1.7
2004 4/.....	1.0	1.1	1.0	0.9	0.9	1.1	0.9	1.0	1.4	1.5
2003 4/.....	1.0	1.0	1.0	1.0	0.9	1.1	1.1	1.2	1.3	1.6
2002.....	1.2	0.9	0.8	0.8	0.8	0.8	0.9	1.1	1.4	1.6
2001.....	1.0	1.0	0.8	0.7	0.9	0.8	0.9	1.1	1.3	1.7
2000.....	1.0	0.9	0.9	0.9	0.9	0.8	1.1	0.9	1.8	1.5
1999.....	1.0	1.1	1.0	0.9	0.9	0.9	1.0	1.1	1.4	1.6

White, male

2006 4/.....	1.8	2.0	1.6	1.6	1.3	1.7	1.9	2.5	2.5	2.6
2005 4/.....	2.0	2.0	1.6	1.5	1.4	1.3	1.7	2.1	2.3	3.9
2004 4/.....	1.7	1.8	1.6	1.4	1.4	1.8	1.4	1.6	2.8	2.8
2003 4/.....	1.8	1.5	1.5	1.6	1.4	1.7	1.9	1.8	2.4	3.3
2002.....	1.9	1.4	1.4	1.3	1.2	1.4	1.6	1.8	2.4	3.7
2001.....	1.6	1.7	1.3	1.1	1.3	1.4	1.6	1.8	2.3	3.5
2000.....	1.6	1.5	1.5	1.6	1.6	1.4	1.7	1.1	3.0	2.7
1999.....	1.7	1.8	1.7	1.4	1.4	1.4	1.5	1.9	2.7	2.4

White, female

2006 4/.....	0.5	0.5	0.4	0.5	0.5	0.7	0.5	0.8	0.8	0.8
2005 4/.....	0.6	0.5	0.5	0.3	0.4	0.4	0.7	0.7	*	0.8
2004 4/.....	0.3	0.4	0.4	0.4	0.5	*	*	*	*	0.9
2003 4/.....	0.3	0.4	0.4	0.4	0.4	0.6	0.6	0.8	0.7	0.9
2002.....	0.4	0.4	0.3	0.3	0.4	*	*	0.6	0.8	0.7
2001.....	0.4	0.3	0.3	0.4	0.5	*	*	0.7	0.7	1.0
2000.....	0.3	0.3	0.3	*	*	*	0.6	0.7	1.0	1.0
1999.....	0.4	0.4	0.3	0.4	*	*	0.7	0.5	0.7	1.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental exposure to smoke, fire and flames (X00-X09)

All races 3/, both sexes

2006.....	0.7	1.1	1.1	1.4	1.5	2.1	2.2	3.2	3.8	4.3
2005.....	0.9	1.2	1.1	1.1	1.6	1.9	2.3	3.1	4.2	5.0
2004.....	0.7	1.1	1.2	1.2	1.6	1.8	2.0	2.9	4.4	5.6
2003.....	1.0	1.2	1.3	1.3	1.6	2.0	2.6	3.0	4.3	5.4
2002.....	0.9	1.0	1.3	1.2	1.4	1.8	2.4	2.8	4.2	4.8
2001.....	1.1	1.0	1.1	1.5	1.5	2.0	2.5	3.0	3.8	5.7
2000.....	1.0	1.1	1.3	1.4	1.7	2.0	2.4	3.5	4.3	6.2
1999.....	1.0	1.0	1.0	1.2	1.5	2.0	2.5	3.5	4.7	5.9

All races 3/, male

2006.....	0.9	1.4	1.5	1.9	1.9	2.7	2.5	4.0	5.1	6.3
2005.....	1.2	1.5	1.6	1.5	2.0	2.5	2.8	3.9	5.6	7.0
2004.....	1.0	1.5	1.8	1.6	2.1	2.6	2.3	3.5	5.9	7.2
2003.....	1.2	1.7	1.6	1.8	2.0	2.5	3.4	4.0	5.1	7.4
2002.....	1.3	1.4	1.6	1.5	2.0	2.6	3.2	3.5	6.1	7.0
2001.....	1.6	1.5	1.5	2.1	1.8	2.5	3.0	3.5	5.1	8.3
2000.....	1.2	1.5	1.8	1.9	2.2	2.4	3.2	4.5	5.9	9.4
1999.....	1.5	1.5	1.3	1.6	1.9	2.5	3.4	4.3	6.6	9.3

All races 3/, female

2006.....	0.5	0.8	0.8	1.0	1.1	1.5	2.0	2.6	2.9	3.4
2005.....	0.5	0.9	0.7	0.8	1.2	1.4	1.9	2.5	3.4	4.0
2004.....	0.5	0.8	0.7	0.9	1.3	1.2	1.6	2.5	3.5	4.9
2003.....	0.7	0.8	1.1	0.8	1.1	1.6	1.9	2.3	3.9	4.5
2002.....	0.6	0.6	0.9	0.9	0.8	1.1	1.8	2.4	3.0	3.9
2001.....	0.7	0.6	0.7	1.0	1.2	1.5	2.1	2.7	3.1	4.6
2000.....	0.7	0.7	0.7	0.9	1.2	1.6	1.8	2.8	3.4	4.9
1999.....	0.5	0.5	0.7	0.9	1.2	1.6	1.8	3.0	3.6	4.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental exposure to smoke, fire and flames (X00-X09)

White, both sexes

2006 4/.....	0.7	1.0	1.1	1.3	1.4	1.9	2.1	2.8	3.3	3.7
2005 4/.....	0.8	1.2	1.0	0.8	1.4	1.7	2.1	2.7	3.8	4.2
2004 4/.....	0.7	1.0	1.1	1.1	1.4	1.6	1.6	2.5	3.9	4.9
2003 4/.....	1.0	1.2	1.2	1.2	1.4	1.8	2.2	2.6	3.7	4.7
2002.....	0.9	0.9	1.1	1.1	1.1	1.6	2.1	2.5	3.7	4.0
2001.....	1.0	0.9	1.0	1.3	1.3	1.7	2.1	2.6	3.2	4.8
2000.....	0.9	0.9	1.2	1.1	1.4	1.7	2.0	2.8	3.5	5.1
1999.....	0.9	0.9	0.9	1.1	1.2	1.6	2.0	2.8	3.9	4.7

White, male

2006 4/.....	0.9	1.2	1.4	1.8	1.8	2.4	2.4	3.3	4.7	6.0
2005 4/.....	1.1	1.5	1.4	1.1	1.7	2.1	2.6	3.4	5.1	6.0
2004 4/.....	0.9	1.2	1.6	1.3	1.9	2.2	1.9	2.9	5.4	6.5
2003 4/.....	1.3	1.5	1.5	1.6	1.9	2.2	3.0	3.2	4.2	6.7
2002.....	1.3	1.3	1.4	1.4	1.6	2.2	2.7	3.1	5.8	5.8
2001.....	1.4	1.2	1.2	1.8	1.5	2.1	2.4	2.8	4.4	7.0
2000.....	1.1	1.3	1.7	1.6	1.8	2.0	2.7	3.6	4.8	8.2
1999.....	1.4	1.3	1.2	1.4	1.7	2.0	2.6	3.3	5.8	8.0

White, female

2006 4/.....	0.5	0.8	0.7	1.0	1.0	1.4	1.8	2.3	2.4	2.7
2005 4/.....	0.6	0.9	0.7	0.6	1.2	1.3	1.8	2.2	2.9	3.4
2004 4/.....	0.4	0.7	0.7	0.9	1.0	1.1	1.3	2.2	2.9	4.2
2003 4/.....	0.7	0.8	0.9	0.8	1.0	1.5	1.6	2.2	3.4	3.8
2002.....	0.6	0.6	0.9	0.8	0.6	1.0	1.6	2.1	2.5	3.2
2001.....	0.6	0.5	0.7	0.8	1.1	1.3	1.9	2.4	2.5	3.8
2000.....	0.6	0.6	0.7	0.7	1.1	1.4	1.5	2.3	2.7	3.9
1999.....	0.5	0.5	0.6	0.9	0.8	1.3	1.5	2.5	2.8	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental exposure to smoke, fire and flames (X00-X09)

Black, both sexes

2006 4/.....	1.0	1.9	1.9	2.7	2.6	4.3	4.1	9.0	9.9	12.2
2005 4/.....	1.2	1.7	2.2	3.7	3.2	4.4	4.2	8.0	11.2	14.4
2004 4/.....	1.4	2.6	2.2	2.8	4.2	4.0	5.9	8.0	12.1	14.9
2003 4/.....	1.4	2.3	2.9	2.8	2.9	4.4	6.4	8.7	12.4	14.9
2002.....	1.1	1.6	2.8	2.6	4.2	4.0	6.3	7.2	10.7	15.7
2001.....	2.1	2.3	2.5	3.7	3.3	5.2	7.3	8.5	11.6	17.8
2000.....	2.0	2.2	2.6	3.9	4.5	5.0	7.1	11.4	16.1	20.3
1999.....	1.7	2.0	2.3	2.6	4.3	5.7	8.3	12.0	15.3	21.7

Black, male

2006 4/.....	*	2.5	2.6	3.8	4.1	6.6	*	14.8	*	*
2005 4/.....	1.9	2.3	3.8	5.4	5.3	6.7	*	12.1	15.5	22.3
2004 4/.....	2.0	3.9	3.7	4.2	4.7	7.3	7.8	12.3	15.1	*
2003 4/.....	1.5	3.3	3.2	4.6	3.9	6.4	8.9	15.5	17.0	*
2002.....	1.8	2.1	4.0	3.4	6.3	6.7	9.3	9.2	*	25.7
2001.....	2.9	3.7	4.2	5.0	4.9	7.4	11.1	12.1	*	27.2
2000.....	2.7	3.2	4.0	5.8	7.0	7.5	10.4	16.5	21.9	27.7
1999.....	2.5	3.0	3.3	4.1	4.2	7.8	13.2	17.1	20.1	29.3

Black, female

2006 4/.....	*	1.4	*	*	*	*	*	5.6	8.7	11.7
2005 4/.....	*	*	*	2.3	*	*	*	5.6	8.9	11.3
2004 4/.....	*	1.5	*	*	3.8	*	4.6	5.4	10.6	13.4
2003 4/.....	*	1.4	2.6	*	*	*	4.7	*	10.1	13.4
2002.....	*	*	*	*	*	*	*	6.0	9.2	11.9
2001.....	1.3	*	*	2.7	*	*	4.7	6.4	10.2	14.3
2000.....	1.3	*	*	*	*	*	4.9	8.3	13.2	17.6
1999.....	*	*	*	*	4.3	4.2	5.0	8.9	13.0	18.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes

2006.....	19.3	20.7	17.1	9.8	4.6	2.8	2.2	2.4	2.9	3.8
2005.....	17.1	18.2	14.4	8.3	4.4	2.6	2.0	2.3	2.6	3.5
2004.....	16.3	16.7	12.0	6.7	3.7	2.4	2.1	2.4	2.5	3.4
2003.....	15.5	15.4	10.9	6.1	3.4	2.3	2.0	2.0	2.5	3.5
2002.....	14.8	13.8	9.2	5.0	3.0	2.3	1.9	2.1	2.7	3.9
2001.....	12.0	10.9	7.0	3.7	2.4	1.7	1.5	2.1	2.4	3.5
2000.....	11.2	9.9	6.1	3.3	2.2	1.5	1.5	1.6	2.6	4.3
1999.....	11.1	9.8	5.4	3.4	2.0	1.7	1.6	1.9	2.6	3.4

All races 3/, male

2006.....	25.4	26.9	22.5	12.7	6.1	3.0	2.7	3.0	2.7	4.6
2005.....	22.1	24.0	19.1	10.9	5.7	3.0	2.3	3.2	2.8	4.6
2004.....	20.9	21.6	16.3	8.4	4.2	2.7	2.5	2.6	2.9	4.2
2003.....	20.7	20.8	14.7	7.6	4.5	3.0	2.2	2.3	3.5	4.0
2002.....	20.3	18.7	12.8	6.8	3.6	2.8	2.1	2.3	3.2	5.1
2001.....	16.8	15.7	9.9	5.0	3.0	2.0	2.1	2.7	2.8	3.8
2000.....	16.4	14.7	8.9	4.0	2.7	1.8	1.7	1.6	2.9	6.2
1999.....	16.6	15.1	7.9	4.7	3.0	2.2	1.9	2.1	2.9	4.4

All races 3/, female

2006.....	13.4	14.7	11.9	7.1	3.4	2.6	1.9	1.9	3.0	3.4
2005.....	12.0	12.6	10.0	5.8	3.2	2.3	1.7	1.7	2.4	3.0
2004.....	11.7	11.9	7.9	5.1	3.2	2.1	1.8	2.2	2.3	3.0
2003.....	10.3	10.2	7.3	4.6	2.5	1.8	1.8	1.7	2.0	3.3
2002.....	9.4	9.1	5.8	3.4	2.4	1.8	1.7	1.9	2.3	3.4
2001.....	7.3	6.2	4.1	2.6	1.9	1.4	1.0	1.6	2.2	3.3
2000.....	6.0	5.3	3.4	2.7	1.8	1.3	1.3	1.6	2.4	3.6
1999.....	5.7	4.7	3.1	2.2	1.2	1.2	1.4	1.7	2.4	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental poisoning and exposure to noxious substances (X40-X49)

White, both sexes

2006 4/.....	20.3	21.1	16.6	9.3	4.2	2.5	2.1	2.4	2.9	3.8
2005 4/.....	18.0	18.7	13.8	7.6	4.1	2.4	1.9	2.3	2.5	3.6
2004 4/.....	17.3	17.1	11.7	6.2	3.5	2.3	2.1	2.3	2.4	3.4
2003 4/.....	16.3	15.7	10.6	5.8	3.4	2.0	2.0	2.0	2.7	3.5
2002.....	15.4	13.7	8.7	4.6	2.7	2.2	1.9	2.1	2.6	3.9
2001.....	12.1	10.5	6.1	3.4	2.1	1.6	1.4	2.0	2.3	3.3
2000.....	11.0	9.4	5.3	3.0	2.1	1.5	1.4	1.7	2.6	4.1
1999.....	10.8	9.1	4.8	3.1	1.7	1.5	1.4	1.9	2.6	3.3

White, male

2006 4/.....	26.3	27.0	21.0	11.3	5.1	2.5	2.3	2.9	2.8	4.7
2005 4/.....	23.3	24.2	17.9	9.4	4.9	2.3	2.0	3.1	2.6	4.8
2004 4/.....	22.2	22.0	15.4	7.3	3.6	2.3	2.4	2.4	2.7	4.3
2003 4/.....	21.8	20.9	13.8	7.0	4.2	2.3	2.2	2.3	3.6	3.7
2002.....	21.0	18.3	11.7	5.8	2.9	2.6	2.0	2.3	3.0	5.2
2001.....	17.1	14.9	8.5	4.3	2.3	1.9	2.0	2.4	2.7	3.7
2000.....	16.2	13.7	7.6	3.3	2.4	1.6	1.6	1.7	2.9	5.9
1999.....	16.2	13.8	6.6	4.0	2.3	2.0	1.7	2.1	3.0	4.5

White, female

2006 4/.....	14.1	15.3	12.2	7.3	3.5	2.6	2.0	2.0	3.0	3.3
2005 4/.....	12.6	13.3	9.9	5.9	3.4	2.4	1.8	1.7	2.5	3.0
2004 4/.....	12.5	12.2	8.2	5.2	3.4	2.2	1.8	2.3	2.2	3.0
2003 4/.....	10.8	10.5	7.3	4.7	2.6	1.7	1.8	1.8	2.1	3.4
2002.....	9.8	9.0	5.8	3.4	2.5	1.9	1.8	1.9	2.5	3.4
2001.....	7.1	6.1	3.8	2.6	1.9	1.4	1.0	1.7	2.0	3.1
2000.....	5.8	5.1	3.1	2.7	1.8	1.4	1.3	1.7	2.5	3.4
1999.....	5.4	4.5	3.0	2.3	1.2	1.1	1.2	1.8	2.4	2.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Accidental poisoning and exposure to noxious substances (X40-X49)

Black, both sexes

2006 4/.....	20.3	24.3	26.9	17.7	9.3	5.4	3.5	*	*	5.4
2005 4/.....	17.0	20.7	23.1	16.3	8.1	5.0	3.7	*	*	*
2004 4/.....	14.6	19.6	17.9	12.6	6.0	4.1	3.1	*	*	*
2003 4/.....	15.2	18.8	16.9	9.6	4.8	5.5	*	*	*	*
2002.....	15.7	19.4	16.1	10.3	6.0	3.0	*	*	*	*
2001.....	15.2	17.7	15.4	7.4	5.4	2.6	*	*	*	6.4
2000.....	15.3	16.5	14.2	6.8	3.9	2.7	*	*	*	8.1
1999.....	16.1	17.9	12.3	6.5	5.5	2.9	3.2	*	*	*

Black, male

2006 4/.....	27.8	34.5	41.7	29.5	17.0	8.4	6.5	*	*	*
2005 4/.....	22.0	30.3	34.6	26.9	15.1	9.3	6.6	*	*	*
2004 4/.....	19.1	25.8	28.7	20.5	10.1	7.3	*	*	*	*
2003 4/.....	20.5	26.7	25.7	15.8	8.7	10.3	*	*	*	*
2002.....	21.6	27.2	25.6	17.7	11.3	5.5	*	*	*	*
2001.....	20.3	28.0	24.8	12.7	9.6	*	*	*	*	*
2000.....	22.8	26.3	23.1	11.5	7.0	*	*	*	*	*
1999.....	24.2	29.7	20.9	11.9	10.0	*	*	*	*	*

Black, female

2006 4/.....	13.6	15.3	14.4	8.0	3.3	*	*	*	*	*
2005 4/.....	12.6	12.4	13.4	7.6	*	*	*	*	*	*
2004 4/.....	10.6	14.2	8.9	6.1	*	*	*	*	*	*
2003 4/.....	10.5	11.9	9.5	4.5	*	*	*	*	*	*
2002.....	10.5	12.5	8.1	4.3	*	*	*	*	*	*
2001.....	10.7	8.9	7.5	3.1	*	*	*	*	*	*
2000.....	8.6	8.0	6.6	3.0	*	*	*	*	*	*
1999.....	9.1	7.8	4.9	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes

2006.....	3.3	4.0	4.5	4.7	5.8	7.7	10.7	18.8	35.2	89.3
2005.....	3.5	4.4	4.9	5.0	6.8	8.4	12.3	20.7	38.9	95.1
2004.....	3.2	3.9	4.5	4.5	5.7	7.0	11.0	19.2	37.4	95.5
2003.....	3.4	3.8	4.0	4.7	5.5	8.1	11.2	19.8	38.1	102.5
2002.....	3.3	3.5	4.0	4.8	5.7	8.6	11.1	21.6	38.8	105.6
2001.....	3.2	3.9	4.5	4.6	6.3	7.8	12.3	22.5	42.2	115.0
2000.....	3.4	3.9	4.1	5.0	6.3	7.8	12.2	22.0	41.4	117.1
1999.....	3.7	3.8	4.4	5.1	6.3	8.9	13.3	23.6	46.9	128.9

All races 3/, male

2006.....	5.4	6.5	7.1	7.1	8.6	11.2	14.5	24.5	44.8	99.4
2005.....	5.6	7.1	7.8	7.4	10.2	11.5	16.6	26.8	49.0	107.1
2004.....	5.0	6.3	7.0	6.7	8.4	9.8	14.3	25.1	46.1	109.8
2003.....	5.5	6.0	6.1	7.1	8.7	11.4	15.1	26.5	47.6	120.2
2002.....	5.4	5.8	6.2	7.4	8.7	12.0	15.0	28.6	47.2	118.6
2001.....	5.3	6.3	6.8	6.7	8.9	10.9	15.9	29.3	52.7	134.0
2000.....	5.6	6.5	6.7	7.7	9.1	10.8	15.6	28.1	51.6	135.2
1999.....	6.1	6.0	7.2	7.8	9.1	12.0	18.0	29.9	59.3	152.7

All races 3/, female

2006.....	1.3	1.6	2.0	2.5	3.2	4.7	7.7	14.6	29.2	84.5
2005.....	1.5	1.8	2.1	2.7	3.6	5.7	8.7	16.3	32.6	89.6
2004.....	1.4	1.5	2.1	2.3	3.2	4.6	8.4	15.0	32.1	89.1
2003.....	1.4	1.6	2.1	2.5	2.7	5.3	8.2	15.0	32.3	94.7
2002.....	1.3	1.3	1.9	2.4	3.0	5.7	8.1	16.6	33.8	100.0
2001.....	1.2	1.6	2.2	2.5	3.8	5.2	9.5	17.7	35.9	107.0
2000.....	1.3	1.4	1.7	2.4	3.8	5.2	9.6	17.8	35.4	109.8
1999.....	1.4	1.7	1.7	2.5	3.7	6.3	9.6	19.3	39.6	119.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

White, both sexes

2006 4/.....	3.4	4.0	4.4	4.7	5.7	7.6	10.4	18.9	36.4	93.2
2005 4/.....	3.5	4.3	4.7	4.8	6.5	8.2	11.7	20.3	38.8	97.4
2004 4/.....	3.4	4.0	4.5	4.4	5.7	6.8	10.8	19.1	38.0	98.8
2003 4/.....	3.5	3.7	3.9	4.6	5.4	8.0	11.1	20.0	38.4	106.3
2002.....	3.4	3.5	4.0	4.6	5.6	7.9	11.0	21.4	39.1	108.8
2001.....	3.2	3.8	4.4	4.5	5.9	7.7	11.9	22.3	42.5	117.7
2000.....	3.4	3.7	4.0	4.7	6.2	7.4	11.8	21.7	41.9	120.5
1999.....	3.7	3.6	4.2	4.7	5.8	8.6	12.7	22.8	46.8	132.4

White, male

2006 4/.....	5.4	6.5	6.8	6.9	8.6	11.1	14.1	24.7	46.2	103.5
2005 4/.....	5.6	6.8	7.4	7.0	9.5	11.4	15.5	25.9	48.6	110.0
2004 4/.....	5.2	6.4	7.0	6.5	8.5	9.5	14.2	24.8	46.6	112.9
2003 4/.....	5.7	5.8	5.9	6.9	8.2	11.4	14.7	26.6	47.3	123.9
2002.....	5.6	5.9	6.1	6.9	8.5	10.9	14.9	28.1	47.2	121.2
2001.....	5.2	6.2	6.6	6.5	8.5	10.7	15.3	29.1	51.9	137.0
2000.....	5.6	6.1	6.4	7.3	8.7	10.1	15.1	27.1	51.2	137.6
1999.....	6.0	5.5	6.8	7.1	8.5	11.3	17.0	28.7	59.0	155.2

White, female

2006 4/.....	1.3	1.6	2.0	2.5	3.1	4.5	7.4	14.5	30.2	88.3
2005 4/.....	1.4	1.7	1.9	2.6	3.7	5.3	8.6	16.1	32.6	91.6
2004 4/.....	1.5	1.6	2.2	2.4	3.1	4.4	8.1	14.9	32.6	92.5
2003 4/.....	1.3	1.6	1.9	2.4	2.7	5.0	8.1	15.3	32.9	98.5
2002.....	1.2	1.1	1.9	2.4	2.9	5.3	7.8	16.5	34.2	103.4
2001.....	1.1	1.5	2.3	2.6	3.5	5.1	9.1	17.4	36.8	109.7
2000.....	1.3	1.4	1.6	2.3	3.8	5.1	9.1	17.9	36.4	113.6
1999.....	1.5	1.6	1.7	2.4	3.3	6.2	9.3	18.7	39.6	123.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Black, both sexes

2006 4/.....	3.6	4.9	5.9	6.1	7.4	10.0	16.0	22.1	28.2	61.7
2005 4/.....	3.9	6.4	7.6	7.6	10.1	12.4	19.3	31.9	46.6	86.1
2004 4/.....	2.9	4.0	5.5	6.3	6.9	10.1	13.7	25.9	37.4	70.2
2003 4/.....	3.7	4.7	6.2	6.7	8.0	11.2	14.7	19.6	39.6	73.2
2002.....	3.3	4.3	5.1	7.3	7.3	15.9	14.3	27.8	39.0	83.1
2001.....	3.9	5.4	5.8	5.9	10.4	10.4	18.7	27.3	43.9	96.1
2000.....	3.9	5.7	5.6	7.9	8.6	12.3	18.1	30.2	39.2	90.1
1999.....	4.7	6.6	6.6	9.3	10.9	13.9	22.9	36.2	53.7	105.5

Black, male

2006 4/.....	6.2	8.2	10.4	9.4	11.0	14.3	22.8	29.1	36.3	75.1
2005 4/.....	6.0	10.8	12.1	12.1	18.9	15.4	29.1	48.3	63.2	97.6
2004 4/.....	4.9	6.6	8.7	10.6	9.9	14.5	18.8	35.0	53.8	87.4
2003 4/.....	5.6	7.9	9.2	10.4	13.9	15.3	22.3	29.2	56.8	91.8
2002.....	4.3	6.4	8.6	12.7	11.3	23.7	18.7	40.9	53.5	106.1
2001.....	6.6	8.6	9.5	9.9	14.6	15.8	25.2	35.3	72.2	114.6
2000.....	6.5	10.1	10.0	13.8	13.9	18.2	23.1	46.8	64.1	124.5
1999.....	8.1	11.1	11.0	15.4	16.1	20.8	34.9	50.3	69.8	153.5

Black, female

2006 4/.....	1.4	2.1	2.1	3.4	4.7	6.8	11.4	17.9	23.9	56.3
2005 4/.....	2.0	2.6	3.8	3.9	3.3	10.2	12.6	22.1	37.8	81.4
2004 4/.....	*	1.7	2.7	2.7	4.5	6.8	10.3	20.5	28.7	63.4
2003 4/.....	2.0	2.0	3.7	3.7	3.5	8.2	9.5	13.8	30.6	66.0
2002.....	2.4	2.5	2.1	2.9	4.2	10.2	11.4	20.0	31.4	74.3
2001.....	1.6	2.7	2.7	2.7	7.0	6.4	14.3	22.5	29.4	89.3
2000.....	1.5	1.8	*	3.0	4.4	7.9	14.9	20.2	26.7	77.3
1999.....	1.6	2.7	2.8	4.3	6.8	8.9	14.9	27.7	45.8	87.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All races 3/, both sexes

2006.....	15.5	17.1	17.3	15.3	13.4	12.6	12.6	15.6	16.3	15.9
2005.....	16.0	16.7	16.1	14.3	13.3	11.8	13.5	15.4	18.7	16.9
2004.....	16.0	16.9	16.2	14.2	13.2	11.4	13.5	16.0	16.7	16.4
2003.....	15.9	16.0	15.7	14.7	12.6	12.0	13.6	15.6	17.7	16.9
2002.....	16.1	16.3	15.1	14.6	12.3	12.5	14.6	16.6	19.3	18.0
2001 5/.....	15.2	15.7	14.6	14.0	12.0	12.7	13.9	16.4	18.9	17.5
2000.....	15.4	14.9	13.9	12.7	11.4	11.3	13.8	16.5	19.3	19.6
1999.....	14.3	14.3	13.4	12.8	11.4	12.4	14.5	17.4	19.2	19.3

All races 3/, male

2006.....	23.6	26.0	26.4	23.9	21.1	22.0	23.6	31.6	35.7	43.2
2005.....	24.6	25.4	25.0	22.4	22.0	20.6	25.4	31.9	41.4	45.0
2004.....	24.2	25.2	24.4	22.3	21.7	20.2	25.4	32.4	38.3	45.0
2003.....	24.5	24.4	24.4	23.1	21.1	21.2	26.0	31.8	40.1	47.8
2002.....	24.7	24.9	23.8	23.5	20.4	22.0	27.9	34.3	44.0	50.7
2001 5/.....	23.6	24.3	22.4	22.3	19.5	22.7	26.7	33.8	44.2	51.1
2000.....	23.9	23.3	21.4	20.1	18.7	19.8	26.1	34.9	44.8	57.5
1999.....	22.1	22.3	20.6	20.3	19.1	22.1	27.7	35.6	44.1	57.1

All races 3/, female

2006.....	7.4	8.4	8.4	7.2	6.3	4.4	3.7	3.8	4.2	3.1
2005.....	7.4	8.2	7.7	6.6	5.5	4.2	3.8	3.5	4.7	4.0
2004.....	7.9	8.9	8.4	6.6	5.5	3.7	3.9	4.3	3.5	3.6
2003.....	7.4	7.9	7.5	6.7	4.9	4.0	3.6	4.0	3.9	3.3
2002.....	7.7	8.0	6.7	6.2	5.0	4.3	3.9	4.1	4.4	3.8
2001 5/.....	6.9	7.4	7.0	6.2	5.2	4.1	3.7	4.3	3.8	3.4
2000.....	7.0	6.7	6.8	5.8	4.9	4.0	4.1	3.8	4.2	4.2
1999.....	6.7	6.7	6.5	5.8	4.4	4.2	4.1	4.8	4.6	4.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

White, both sexes

2006 4/.....	17.5	19.3	19.5	17.1	14.8	13.8	13.8	16.9	17.6	16.9
2005 4/.....	18.0	18.9	18.1	15.9	14.7	13.0	14.7	16.7	20.0	17.9
2004 4/.....	18.1	19.0	18.3	15.7	14.4	12.4	14.5	17.3	17.7	17.5
2003 4/.....	17.9	18.1	17.7	16.2	13.8	13.0	14.7	16.7	18.9	18.1
2002.....	18.2	18.4	16.8	16.1	13.6	13.7	15.7	17.6	20.9	19.1
2001 5/.....	16.9	17.6	16.2	15.3	13.1	13.8	14.9	17.6	20.0	18.6
2000.....	17.0	16.5	15.3	13.9	12.6	12.3	14.7	17.6	20.4	21.0
1999.....	16.0	15.9	14.8	14.0	12.3	13.3	15.6	18.6	20.4	20.7

White, male

2006 4/.....	26.3	29.0	29.5	26.4	23.0	23.8	25.9	34.1	38.7	46.1
2005 4/.....	27.3	28.5	27.7	24.8	24.1	22.6	27.6	34.2	44.3	48.2
2004 4/.....	27.0	27.8	27.3	24.1	23.6	21.9	27.1	34.8	40.4	48.4
2003 4/.....	27.3	27.3	27.2	25.4	22.9	22.9	27.9	34.0	42.9	51.4
2002.....	27.6	27.6	26.4	25.7	22.5	23.8	30.1	36.3	47.2	53.9
2001 5/.....	26.1	27.1	24.7	24.3	21.0	24.5	28.3	36.0	46.8	55.0
2000.....	26.2	25.6	23.2	21.8	20.4	21.3	27.6	37.1	47.7	61.6
1999.....	24.5	24.4	22.3	21.9	20.5	23.6	29.5	37.9	46.8	61.5

White, female

2006 4/.....	8.5	9.6	9.7	8.3	7.0	4.8	3.8	4.1	4.3	3.2
2005 4/.....	8.5	9.4	8.7	7.4	6.0	4.4	4.1	3.7	4.8	3.9
2004 4/.....	9.1	10.2	9.5	7.6	5.8	4.1	4.1	4.5	3.6	3.6
2003 4/.....	8.4	9.0	8.3	7.4	5.5	4.3	3.9	4.2	4.1	3.3
2002.....	8.7	9.3	7.4	6.9	5.3	4.8	3.9	4.2	4.7	4.0
2001 5/.....	7.8	8.1	7.9	6.8	5.8	4.3	4.0	4.6	3.8	3.4
2000.....	7.9	7.5	7.5	6.4	5.5	4.4	4.4	3.9	4.2	4.5
1999.....	7.5	7.4	7.4	6.4	4.8	4.4	4.5	5.0	4.8	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes

2006.....	6.7	7.8	8.3	8.1	8.1	8.5	9.4	11.8	12.2	10.9
2005.....	7.0	8.0	8.3	8.1	8.2	8.2	9.9	11.9	13.6	11.2
2004.....	7.1	8.1	8.0	7.8	8.3	8.0	9.8	12.2	12.3	11.1
2003.....	7.2	7.8	8.3	8.4	8.2	8.5	10.1	11.9	13.1	11.9
2002.....	7.4	7.8	7.9	8.6	8.1	8.8	10.7	12.4	14.3	12.0
2001.....	7.0	7.5	8.0	8.4	8.0	9.0	10.3	12.4	14.1	12.2
2000.....	7.4	7.4	7.8	7.4	7.7	8.3	10.3	12.4	14.0	13.2
1999.....	6.7	7.4	7.2	8.1	7.5	8.7	10.9	12.9	13.5	12.6

All races 3/, male

2006.....	11.2	13.1	14.3	14.3	14.4	16.3	19.2	26.4	29.7	32.7
2005.....	11.8	13.7	14.6	14.1	15.3	15.5	20.2	26.6	33.3	33.8
2004.....	11.7	13.4	13.6	13.8	14.9	15.5	20.0	26.7	30.7	34.4
2003.....	12.1	13.2	14.5	14.6	15.3	16.5	20.7	26.3	32.4	36.6
2002.....	12.6	13.3	14.0	15.4	14.8	17.0	22.5	27.8	35.9	37.8
2001.....	12.0	12.8	13.8	14.9	14.2	17.3	21.4	27.7	35.5	38.9
2000.....	12.6	12.8	13.3	13.1	13.9	15.9	21.3	28.1	35.4	42.9
1999.....	11.2	12.6	12.4	14.0	14.0	16.9	22.4	29.1	34.6	41.6

All races 3/, female

2006.....	2.3	2.7	2.6	2.3	2.4	1.8	1.4	1.1	1.3	0.7
2005.....	2.3	2.5	2.4	2.3	1.8	1.7	1.6	1.3	1.3	0.8
2004.....	2.6	2.8	2.7	2.1	2.3	1.4	1.7	1.7	1.0	*
2003.....	2.3	2.6	2.3	2.5	1.7	1.6	1.6	1.6	1.3	0.9
2002.....	2.3	2.4	2.1	2.3	2.0	1.8	1.3	1.5	1.2	0.8
2001.....	2.2	2.3	2.4	2.4	2.3	1.9	1.4	1.7	1.3	1.0
2000.....	2.4	2.2	2.5	2.1	2.1	1.8	1.6	1.4	1.4	1.1
1999.....	2.2	2.4	2.1	2.6	1.6	1.7	1.8	1.6	1.2	0.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

White, both sexes

2006 4/.....	7.7	9.0	9.5	9.3	9.0	9.5	10.5	13.0	13.4	11.7
2005 4/.....	8.0	9.2	9.5	9.2	9.2	9.2	10.9	13.1	14.7	12.2
2004 4/.....	8.1	9.2	9.2	8.7	9.3	8.8	10.8	13.3	13.3	12.1
2003 4/.....	8.3	9.1	9.4	9.5	9.1	9.5	11.1	12.9	14.1	12.9
2002.....	8.5	8.9	8.9	9.7	9.0	9.9	11.7	13.3	15.6	12.9
2001.....	7.9	8.5	9.0	9.4	8.9	10.0	11.1	13.4	15.0	13.2
2000.....	8.4	8.3	8.7	8.2	8.5	9.2	11.1	13.4	15.2	14.3
1999.....	7.5	8.3	8.0	8.9	8.2	9.5	11.8	13.9	14.4	13.6

White, male

2006 4/.....	12.6	15.0	16.2	16.1	15.8	17.9	21.2	28.6	32.3	35.1
2005 4/.....	13.3	15.5	16.3	15.9	17.0	17.4	22.0	28.7	36.0	36.8
2004 4/.....	13.1	15.1	15.5	15.1	16.6	16.9	21.6	28.7	32.9	37.5
2003 4/.....	13.8	15.1	16.2	16.4	16.8	18.3	22.5	28.3	34.8	39.9
2002.....	14.3	15.0	15.5	16.9	16.3	18.8	24.4	29.6	38.8	40.5
2001.....	13.3	14.5	15.4	16.3	15.7	19.0	22.9	29.7	37.8	42.1
2000.....	14.0	14.2	14.7	14.2	15.3	17.3	22.7	30.4	38.1	46.5
1999.....	12.5	14.0	13.7	15.1	15.2	18.3	24.0	31.2	36.8	44.9

White, female

2006 4/.....	2.7	3.1	3.0	2.7	2.7	2.0	1.6	1.3	1.4	0.7
2005 4/.....	2.7	2.9	2.8	2.7	2.0	1.9	1.7	1.5	1.4	0.9
2004 4/.....	3.0	3.3	3.1	2.5	2.6	1.7	1.9	2.0	1.1	*
2003 4/.....	2.7	3.1	2.6	2.9	2.0	1.8	1.8	1.8	1.4	0.9
2002.....	2.7	2.8	2.4	2.7	2.3	2.1	1.4	1.6	1.3	0.9
2001.....	2.4	2.6	2.8	2.7	2.7	2.0	1.5	1.8	1.3	1.1
2000.....	2.7	2.5	2.8	2.4	2.3	2.1	1.8	1.5	1.5	1.2
1999.....	2.5	2.7	2.5	3.0	1.8	1.8	2.1	1.8	1.3	1.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All races 3/, both sexes

2006.....	8.7	9.3	9.0	7.2	5.3	4.0	3.2	3.8	4.1	5.0
2005.....	9.0	8.7	7.8	6.2	5.1	3.7	3.6	3.5	5.2	5.7
2004.....	8.9	8.9	8.2	6.5	4.9	3.4	3.6	3.9	4.4	5.4
2003.....	8.7	8.2	7.5	6.2	4.5	3.4	3.5	3.7	4.6	5.1
2002.....	8.7	8.5	7.2	5.9	4.2	3.7	3.9	4.2	5.0	6.0
2001 5/.....	8.2	8.2	6.6	5.6	4.0	3.7	3.6	4.0	4.8	5.3
2000.....	7.9	7.4	6.2	5.2	3.7	3.0	3.5	4.2	5.3	6.5
1999.....	7.7	6.9	6.2	4.7	3.9	3.8	3.6	4.5	5.7	6.7

All races 3/, male

2006.....	12.4	12.9	12.2	9.6	6.8	5.7	4.4	5.3	6.0	10.5
2005.....	12.8	11.8	10.4	8.3	6.7	5.1	5.2	5.3	8.1	11.2
2004.....	12.5	11.8	10.8	8.6	6.8	4.7	5.4	5.7	7.6	10.7
2003.....	12.4	11.2	9.9	8.5	5.8	4.7	5.4	5.5	7.7	11.1
2002.....	12.1	11.6	9.9	8.2	5.7	5.0	5.5	6.5	8.1	12.9
2001 5/.....	11.6	11.5	8.6	7.5	5.3	5.4	5.4	6.1	8.7	12.2
2000.....	11.3	10.6	8.1	6.9	4.8	3.9	4.8	6.8	9.4	14.6
1999.....	10.8	9.6	8.1	6.4	5.1	5.2	5.3	6.6	9.5	15.5

All races 3/, female

2006.....	5.1	5.7	5.9	4.9	4.0	2.6	2.3	2.7	2.9	2.4
2005.....	5.1	5.7	5.3	4.3	3.6	2.4	2.2	2.2	3.4	3.1
2004.....	5.3	6.0	5.7	4.5	3.2	2.2	2.2	2.5	2.5	3.0
2003.....	5.1	5.3	5.1	4.1	3.2	2.4	2.0	2.4	2.6	2.4
2002.....	5.4	5.6	4.6	3.9	2.9	2.5	2.6	2.6	3.1	3.0
2001 5/.....	4.8	5.1	4.6	3.8	2.8	2.2	2.3	2.6	2.4	2.4
2000.....	4.6	4.4	4.3	3.7	2.8	2.1	2.5	2.4	2.8	3.2
1999.....	4.5	4.2	4.4	3.2	2.8	2.5	2.3	3.1	3.4	3.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

White, both sexes

2006 4/.....	9.8	10.3	10.0	7.9	5.7	4.2	3.3	3.9	4.2	5.3
2005 4/.....	9.9	9.7	8.6	6.8	5.5	3.8	3.8	3.6	5.3	5.6
2004 4/.....	10.0	9.7	9.1	7.0	5.1	3.6	3.7	4.0	4.4	5.5
2003 4/.....	9.6	9.1	8.3	6.7	4.7	3.5	3.6	3.8	4.8	5.2
2002.....	9.7	9.6	7.9	6.4	4.6	3.8	4.0	4.3	5.3	6.2
2001 5/.....	9.1	9.1	7.2	5.9	4.2	3.8	3.7	4.2	5.0	5.5
2000.....	8.7	8.2	6.6	5.7	4.1	3.1	3.6	4.2	5.3	6.7
1999.....	8.4	7.6	6.7	5.1	4.1	3.8	3.8	4.7	5.9	7.1

White, male

2006 4/.....	13.8	14.1	13.3	10.3	7.2	5.9	4.7	5.4	6.3	11.0
2005 4/.....	14.0	13.0	11.4	8.9	7.2	5.3	5.6	5.5	8.4	11.4
2004 4/.....	13.9	12.6	11.8	9.0	7.1	4.9	5.4	6.1	7.5	10.9
2003 4/.....	13.5	12.2	11.0	9.0	6.1	4.7	5.4	5.6	8.1	11.5
2002.....	13.4	12.6	10.9	8.7	6.2	5.0	5.8	6.7	8.4	13.4
2001 5/.....	12.7	12.6	9.3	7.9	5.3	5.5	5.3	6.3	9.0	12.9
2000.....	12.2	11.4	8.5	7.6	5.2	4.0	4.9	6.7	9.6	15.2
1999.....	11.9	10.4	8.6	6.7	5.2	5.3	5.5	6.7	10.0	16.6

White, female

2006 4/.....	5.8	6.5	6.7	5.6	4.4	2.8	2.2	2.8	2.9	2.6
2005 4/.....	5.8	6.5	5.9	4.8	3.9	2.5	2.3	2.2	3.4	3.0
2004 4/.....	6.0	6.8	6.4	5.1	3.3	2.5	2.2	2.5	2.5	3.0
2003 4/.....	5.7	5.9	5.7	4.5	3.5	2.5	2.2	2.4	2.8	2.4
2002.....	6.0	6.5	5.0	4.2	3.0	2.7	2.5	2.6	3.4	3.1
2001 5/.....	5.4	5.6	5.1	4.1	3.1	2.3	2.5	2.8	2.5	2.3
2000.....	5.1	5.0	4.8	4.0	3.1	2.3	2.6	2.5	2.7	3.3
1999.....	4.9	4.7	4.9	3.5	3.0	2.6	2.4	3.3	3.5	3.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All races 3/, both sexes

2006.....	6.3	5.6	4.5	3.4	2.9	2.0	2.2	2.0	2.3	1.9
2005.....	6.7	5.4	4.2	2.9	2.7	2.3	2.4	2.2	2.1	2.1
2004.....	6.3	5.3	4.2	3.2	2.8	2.6	2.2	2.2	2.2	2.1
2003.....	6.4	5.6	4.2	3.1	2.4	2.3	2.4	2.5	2.4	2.2
2002.....	6.5	5.4	4.0	3.5	2.7	2.4	2.2	2.3	2.4	2.1
2001 5/.....	8.6	7.0	5.5	4.4	3.5	3.3	2.5	2.5	2.5	2.4
2000.....	6.6	5.3	4.0	3.4	2.6	2.4	2.4	2.3	2.6	2.4
1999.....	6.6	5.1	4.0	3.2	2.8	2.7	2.4	2.5	2.4	2.4

All races 3/, male

2006.....	9.4	8.5	6.7	5.2	4.4	2.9	3.2	2.9	2.3	2.3
2005.....	10.0	8.3	6.7	4.5	4.2	3.1	3.5	2.8	2.4	2.7
2004.....	9.0	7.8	6.7	4.9	4.1	3.8	3.3	2.6	2.7	2.7
2003.....	9.5	8.3	6.4	4.6	3.7	3.4	3.4	3.5	3.0	3.2
2002.....	9.6	8.0	6.0	5.3	4.2	3.4	3.2	3.3	2.9	3.0
2001 5/.....	12.6	10.6	8.3	6.7	5.0	4.6	3.3	3.2	3.0	3.2
2000.....	9.7	7.8	5.9	5.2	3.9	3.3	3.4	2.8	3.8	3.3
1999.....	9.5	7.7	5.8	5.1	4.3	4.2	3.1	3.4	3.4	3.4

All races 3/, female

2006.....	3.3	2.8	2.5	1.7	1.6	1.3	1.4	1.4	2.3	1.8
2005.....	3.4	2.6	1.8	1.5	1.4	1.6	1.5	1.8	2.0	1.9
2004.....	3.6	3.0	1.9	1.6	1.6	1.6	1.3	1.9	2.0	1.9
2003.....	3.4	3.0	2.1	1.7	1.3	1.4	1.6	1.8	2.1	1.8
2002.....	3.5	2.9	2.2	1.8	1.4	1.5	1.4	1.6	2.1	1.7
2001 5/.....	4.7	3.6	2.7	2.3	2.2	2.1	1.9	1.9	2.2	2.0
2000.....	3.6	2.8	2.1	1.7	1.5	1.7	1.6	2.0	1.9	2.0
1999.....	3.6	2.5	2.2	1.4	1.5	1.5	1.9	1.9	1.9	2.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

White, both sexes

2006 4/.....	4.3	3.9	3.2	2.5	2.4	1.5	1.9	1.7	2.2	1.7
2005 4/.....	4.4	3.8	2.9	2.3	2.1	1.8	1.9	2.0	1.9	1.9
2004 4/.....	4.3	3.8	3.0	2.4	2.3	2.2	1.8	1.8	2.0	1.9
2003 4/.....	4.2	3.9	3.1	2.4	2.0	1.9	1.9	2.2	2.1	2.0
2002.....	4.5	3.8	2.9	2.9	2.2	1.9	1.9	1.8	2.1	1.8
2001 5/.....	6.8	5.6	4.4	3.8	3.0	2.7	2.2	2.1	2.2	2.2
2000.....	4.4	3.7	2.9	2.7	2.0	1.8	2.0	2.1	2.0	2.2
1999.....	4.5	3.6	3.0	2.6	2.3	2.3	2.0	2.0	2.1	2.1

White, male

2006 4/.....	6.1	5.7	4.5	3.5	3.5	1.9	2.4	2.3	2.1	2.0
2005 4/.....	6.2	5.6	4.5	3.3	3.3	2.2	2.7	2.2	2.0	1.9
2004 4/.....	5.8	5.2	4.6	3.6	3.2	3.0	2.6	2.0	2.2	2.3
2003 4/.....	5.8	5.4	4.5	3.3	2.9	2.7	2.6	2.9	2.5	3.0
2002.....	6.5	5.1	4.0	4.1	3.2	2.7	2.6	2.4	2.3	2.7
2001 5/.....	9.8	8.2	6.5	5.5	4.2	3.7	2.6	2.7	2.6	2.8
2000.....	6.4	5.1	4.0	4.0	2.8	2.2	2.6	2.4	2.5	2.8
1999.....	6.3	5.3	4.3	3.8	3.3	3.4	2.5	2.6	2.9	2.7

White, female

2006 4/.....	2.4	2.2	2.0	1.5	1.3	1.2	1.4	1.3	2.3	1.6
2005 4/.....	2.6	2.0	1.4	1.3	1.1	1.4	1.3	1.8	1.8	1.9
2004 4/.....	2.9	2.3	1.4	1.3	1.4	1.6	1.1	1.6	1.9	1.8
2003 4/.....	2.7	2.4	1.7	1.6	1.0	1.2	1.3	1.6	1.8	1.6
2002.....	2.5	2.5	1.8	1.7	1.3	1.3	1.4	1.4	1.9	1.5
2001 5/.....	3.8	3.0	2.4	2.1	1.9	1.9	1.8	1.7	1.9	2.0
2000.....	2.5	2.3	1.8	1.4	1.3	1.4	1.5	1.8	1.7	1.9
1999.....	2.6	1.9	1.7	1.3	1.3	1.4	1.7	1.6	1.6	1.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

Black, both sexes

2006 4/.....	20.5	17.1	14.5	10.7	8.0	6.6	5.4	5.0	*	*
2005 4/.....	21.6	17.2	13.9	8.3	7.8	7.1	7.1	5.8	5.6	*
2004 4/.....	20.2	16.8	13.4	9.4	7.9	5.7	6.4	7.1	5.2	*
2003 4/.....	21.4	17.7	12.7	9.1	6.8	5.7	6.9	6.6	6.6	*
2002.....	20.6	17.6	13.0	8.9	7.5	5.8	5.6	8.2	6.0	*
2001 5/.....	21.7	18.1	13.6	9.9	7.5	7.7	6.5	6.9	6.2	*
2000.....	21.8	16.8	12.4	9.1	7.4	8.2	6.8	5.3	9.6	*
1999.....	21.2	16.2	11.2	8.8	7.4	6.6	6.0	8.2	6.1	*

Black, male

2006 4/.....	33.6	29.3	24.5	19.3	13.6	12.4	10.2	10.3	*	*
2005 4/.....	36.7	29.3	24.1	15.3	12.9	12.6	12.7	11.2	*	*
2004 4/.....	32.9	27.8	23.8	16.5	13.1	11.6	11.3	10.5	*	*
2003 4/.....	36.8	30.0	21.8	17.3	11.4	10.8	11.5	12.3	*	*
2002.....	32.4	30.8	22.5	16.8	13.5	10.0	11.3	15.2	*	*
2001 5/.....	34.2	30.0	23.1	17.4	12.0	13.2	11.8	11.6	*	*
2000.....	34.1	28.7	22.4	16.0	13.0	15.1	12.4	*	19.4	*
1999.....	34.1	27.3	18.5	17.0	12.7	13.0	10.2	13.8	*	*

Black, female

2006 4/.....	8.9	6.4	6.1	3.6	3.6	*	*	*	*	*
2005 4/.....	8.3	6.7	5.2	2.5	3.8	*	*	*	*	*
2004 4/.....	8.9	7.2	4.6	3.6	3.8	*	*	*	*	*
2003 4/.....	7.9	7.0	5.0	2.3	3.2	*	*	*	*	*
2002.....	10.1	6.1	4.9	*	*	*	*	*	*	*
2001 5/.....	10.6	7.7	5.6	3.8	4.0	*	*	*	*	*
2000.....	10.9	6.5	3.8	3.5	*	*	*	*	*	*
1999.....	9.8	6.7	5.0	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All races 3/, both sexes

2006.....	3.9	3.1	2.4	1.6	1.4	0.9	0.8	0.7	0.6	0.4
2005.....	4.1	3.0	2.1	1.4	1.3	1.1	1.1	0.8	0.6	0.5
2004.....	3.8	2.9	2.2	1.7	1.5	1.2	0.8	0.8	0.7	0.6
2003.....	3.8	3.1	2.2	1.6	1.2	0.9	0.9	0.7	0.6	0.5
2002.....	3.9	2.8	2.2	1.8	1.3	1.0	0.8	1.0	0.8	*
2001.....	3.7	2.7	2.1	1.5	1.4	1.2	0.9	0.8	0.7	*
2000.....	3.8	2.9	2.1	1.7	1.2	1.0	1.0	0.7	0.5	0.7
1999.....	3.7	2.8	2.0	1.6	1.6	1.1	1.0	0.8	0.7	0.6

All races 3/, male

2006.....	6.2	4.9	3.6	2.3	2.1	1.2	1.3	1.0	*	*
2005.....	6.6	4.7	3.4	2.2	2.0	1.4	1.7	1.0	*	*
2004.....	5.9	4.4	3.5	2.7	2.2	1.8	1.4	1.0	*	*
2003.....	5.9	5.0	3.5	2.4	1.8	1.4	1.4	0.9	*	*
2002.....	6.2	4.2	3.4	2.7	1.9	1.4	1.0	1.4	1.0	*
2001.....	5.6	4.2	3.2	2.3	2.0	1.8	1.1	1.2	*	*
2000.....	5.8	4.5	3.3	2.6	1.8	1.3	1.3	0.8	*	*
1999.....	5.5	4.4	3.0	2.6	2.5	1.5	1.4	1.1	*	*

All races 3/, female

2006.....	1.6	1.3	1.2	0.9	0.7	0.6	0.4	*	0.7	*
2005.....	1.7	1.4	0.8	0.6	0.6	0.8	0.6	0.7	0.6	*
2004.....	1.6	1.4	0.9	0.7	0.9	0.7	*	0.7	0.8	*
2003.....	1.7	1.4	1.0	0.8	0.6	0.4	0.5	0.6	0.7	*
2002.....	1.7	1.5	1.1	0.9	0.7	0.6	0.6	0.6	0.7	*
2001.....	1.7	1.1	1.0	0.7	0.8	0.7	0.7	0.5	0.7	*
2000.....	1.7	1.3	1.0	0.9	0.7	0.8	0.7	0.7	*	*
1999.....	1.8	1.3	1.1	0.6	0.7	0.7	0.7	0.6	0.6	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

White, both sexes

2006 4/.....	2.5	2.0	1.5	1.2	1.1	0.6	0.6	0.5	0.7	*
2005 4/.....	2.6	2.1	1.5	1.1	1.0	0.9	0.9	0.8	0.5	*
2004 4/.....	2.5	2.0	1.5	1.3	1.2	1.1	0.7	0.7	0.7	0.5
2003 4/.....	2.4	2.2	1.7	1.2	1.0	0.7	0.8	0.6	0.6	0.5
2002.....	2.6	1.9	1.6	1.5	0.9	0.8	0.7	0.8	0.8	*
2001.....	2.6	1.9	1.5	1.2	1.2	1.0	0.7	0.7	0.7	*
2000.....	2.5	2.0	1.5	1.4	1.0	0.8	0.8	0.7	*	0.6
1999.....	2.5	1.9	1.5	1.2	1.2	1.0	0.9	0.7	0.7	0.6

White, male

2006 4/.....	3.9	2.8	2.2	1.6	1.7	0.6	0.9	*	*	*
2005 4/.....	3.9	3.0	2.3	1.6	1.6	1.1	1.2	0.9	*	*
2004 4/.....	3.5	2.9	2.3	1.9	1.7	1.4	1.0	0.9	*	*
2003 4/.....	3.4	3.1	2.5	1.8	1.4	1.1	1.1	0.7	*	*
2002.....	4.0	2.6	2.4	2.1	1.4	1.0	0.8	1.2	*	*
2001.....	3.7	2.8	2.3	1.6	1.7	1.3	0.8	1.0	*	*
2000.....	3.7	3.0	2.2	2.1	1.4	1.0	0.9	0.7	*	*
1999.....	3.6	2.8	2.3	1.9	1.8	1.3	1.1	0.7	*	*

White, female

2006 4/.....	1.2	1.2	0.9	0.8	0.6	0.6	*	*	0.8	*
2005 4/.....	1.4	1.1	0.7	0.6	0.5	0.7	0.6	0.8	*	*
2004 4/.....	1.5	1.1	0.8	0.7	0.8	0.8	*	0.6	0.9	*
2003 4/.....	1.3	1.2	0.9	0.8	0.5	*	0.5	0.5	0.7	*
2002.....	1.3	1.3	0.9	0.9	0.5	0.5	0.6	0.6	0.7	*
2001.....	1.4	0.9	0.8	0.8	0.7	0.8	0.7	0.5	0.7	*
2000.....	1.3	1.1	0.9	0.8	0.7	0.7	0.7	0.7	*	*
1999.....	1.4	1.0	0.7	0.6	0.7	0.7	0.7	0.6	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, both sexes

2006 4/.....	13.7	10.3	8.4	4.5	3.7	3.3	*	*	*	*
2005 4/.....	14.4	9.9	6.9	3.7	3.3	2.8	3.3	*	*	*
2004 4/.....	12.8	9.4	7.2	5.2	4.2	2.6	*	*	*	*
2003 4/.....	13.7	10.2	6.8	4.6	3.1	2.4	*	*	*	*
2002.....	13.1	9.6	7.1	4.2	4.0	2.8	*	*	*	*
2001.....	11.6	8.4	6.5	4.3	3.0	2.9	*	*	*	*
2000.....	12.6	9.4	7.0	4.2	2.6	2.8	2.7	*	*	*
1999.....	12.2	9.7	5.9	4.5	4.2	2.2	*	*	*	*

Black, male

2006 4/.....	23.9	19.6	15.0	8.4	7.1	6.3	*	*	*	*
2005 4/.....	26.1	17.9	12.6	7.3	5.6	5.2	7.0	*	*	*
2004 4/.....	23.4	16.5	13.7	9.6	7.1	5.6	*	*	*	*
2003 4/.....	24.3	19.1	11.9	9.1	5.2	*	*	*	*	*
2002.....	22.7	17.4	12.3	8.3	6.5	5.2	*	*	*	*
2001.....	20.3	15.3	11.8	8.6	5.1	6.4	*	*	*	*
2000.....	21.6	16.7	13.1	7.1	5.4	5.2	*	*	*	*
1999.....	20.5	17.3	9.5	9.3	8.3	*	*	*	*	*

Black, female

2006 4/.....	4.6	2.2	2.9	*	*	*	*	*	*	*
2005 4/.....	4.2	3.0	2.1	*	*	*	*	*	*	*
2004 4/.....	3.5	3.2	1.7	*	*	*	*	*	*	*
2003 4/.....	4.4	2.5	2.5	*	*	*	*	*	*	*
2002.....	4.6	2.8	2.8	*	*	*	*	*	*	*
2001.....	3.9	2.4	1.9	*	*	*	*	*	*	*
2000.....	4.7	3.2	*	*	*	*	*	*	*	*
1999.....	4.8	3.2	2.8	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes

2006.....	2.4	2.5	2.2	1.8	1.6	1.2	1.4	1.4	1.7	1.5
2005.....	2.5	2.4	2.1	1.6	1.4	1.2	1.4	1.4	1.5	1.7
2004.....	2.5	2.5	2.1	1.5	1.3	1.4	1.4	1.4	1.6	1.5
2003.....	2.6	2.5	1.9	1.5	1.3	1.5	1.5	1.8	1.8	1.7
2002.....	2.6	2.6	1.8	1.7	1.5	1.4	1.4	1.3	1.6	1.7
2001 5/.....	4.9	4.4	3.4	2.9	2.1	2.1	1.6	1.6	1.8	2.0
2000.....	2.9	2.4	1.8	1.6	1.4	1.4	1.4	1.6	2.1	1.7
1999.....	2.9	2.3	1.9	1.6	1.3	1.6	1.4	1.7	1.7	1.8

All races 3/, male

2006.....	3.2	3.6	3.1	2.9	2.3	1.7	1.9	2.0	1.8	1.8
2005.....	3.4	3.6	3.2	2.3	2.1	1.7	1.9	1.8	1.8	2.1
2004.....	3.1	3.4	3.2	2.2	2.0	2.0	1.8	1.6	2.2	1.7
2003.....	3.6	3.3	2.9	2.2	1.9	2.1	2.0	2.7	2.6	2.5
2002.....	3.4	3.8	2.6	2.6	2.3	2.0	2.2	1.8	1.9	2.3
2001 5/.....	7.0	6.4	5.0	4.4	3.0	2.8	2.2	2.0	2.1	2.7
2000.....	3.9	3.3	2.6	2.6	2.0	2.0	2.1	2.0	3.0	2.5
1999.....	4.0	3.3	2.8	2.5	1.8	2.6	1.7	2.2	2.6	2.6

All races 3/, female

2006.....	1.6	1.5	1.3	0.9	0.9	0.7	1.0	0.9	1.6	1.4
2005.....	1.7	1.2	1.0	0.9	0.8	0.8	1.0	1.1	1.3	1.5
2004.....	1.9	1.6	1.0	0.8	0.8	0.8	1.0	1.2	1.2	1.5
2003.....	1.6	1.6	1.0	0.9	0.7	1.0	1.0	1.1	1.3	1.4
2002.....	1.8	1.4	1.1	1.0	0.7	0.9	0.8	0.9	1.4	1.4
2001 5/.....	2.9	2.5	1.8	1.5	1.4	1.4	1.2	1.4	1.6	1.7
2000.....	1.8	1.5	1.0	0.8	0.8	0.9	0.9	1.3	1.5	1.4
1999.....	1.8	1.3	1.1	0.9	0.8	0.8	1.2	1.3	1.2	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

White, both sexes

2006 4/.....	1.7	1.9	1.7	1.3	1.3	0.9	1.2	1.2	1.5	1.4
2005 4/.....	1.8	1.7	1.4	1.2	1.1	0.9	1.1	1.1	1.3	1.5
2004 4/.....	1.8	1.8	1.5	1.2	1.0	1.2	1.1	1.0	1.3	1.4
2003 4/.....	1.9	1.7	1.4	1.2	1.0	1.2	1.1	1.5	1.5	1.6
2002.....	1.9	1.8	1.3	1.4	1.3	1.2	1.2	1.0	1.3	1.4
2001 5/.....	4.2	3.7	2.9	2.6	1.8	1.7	1.4	1.4	1.5	1.8
2000.....	1.9	1.7	1.4	1.2	1.0	1.0	1.1	1.4	1.6	1.5
1999.....	2.0	1.7	1.5	1.3	1.0	1.4	1.2	1.4	1.4	1.5

White, male

2006 4/.....	2.2	2.9	2.3	1.9	1.8	1.3	1.5	1.7	1.6	1.7
2005 4/.....	2.4	2.5	2.2	1.7	1.6	1.1	1.5	1.3	1.6	1.6
2004 4/.....	2.3	2.3	2.3	1.7	1.4	1.6	1.5	1.1	1.8	1.5
2003 4/.....	2.4	2.3	2.0	1.5	1.5	1.6	1.5	2.2	2.2	2.3
2002.....	2.6	2.5	1.6	1.9	1.9	1.7	1.8	1.2	1.5	2.0
2001 5/.....	6.0	5.3	4.2	3.9	2.5	2.4	1.8	1.7	2.0	2.3
2000.....	2.7	2.1	1.8	1.9	1.5	1.3	1.6	1.7	2.2	2.1
1999.....	2.8	2.4	2.1	1.9	1.5	2.1	1.3	1.8	2.2	1.9

White, female

2006 4/.....	1.2	1.0	1.0	0.7	0.7	0.6	1.0	0.8	1.5	1.2
2005 4/.....	1.2	0.9	0.7	0.8	0.6	0.7	0.8	1.0	1.2	1.5
2004 4/.....	1.4	1.2	0.7	0.7	0.6	0.8	0.7	1.0	1.0	1.3
2003 4/.....	1.3	1.2	0.9	0.8	0.5	0.9	0.8	1.0	1.1	1.2
2002.....	1.2	1.2	0.9	0.9	0.7	0.7	0.8	0.8	1.2	1.2
2001 5/.....	2.3	2.0	1.6	1.4	1.1	1.1	1.1	1.2	1.2	1.6
2000.....	1.2	1.2	0.9	0.6	0.6	0.7	0.7	1.1	1.2	1.3
1999.....	1.2	0.9	1.0	0.8	0.6	0.8	1.0	1.0	1.0	1.4

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

Black, both sexes

2006 4/.....	6.8	6.8	6.1	6.2	4.3	3.3	3.1	*	*	*
2005 4/.....	7.2	7.3	7.0	4.6	4.5	4.3	3.7	4.5	*	*
2004 4/.....	7.3	7.4	6.2	4.2	3.7	3.1	4.3	5.2	*	*
2003 4/.....	7.7	7.5	5.9	4.5	3.6	3.3	4.4	4.8	5.1	*
2002.....	7.5	8.0	5.9	4.6	3.5	3.1	3.8	5.7	*	*
2001 5/.....	10.1	9.7	7.2	5.6	4.6	4.8	4.0	4.7	*	*
2000.....	9.2	7.4	5.3	5.0	4.7	5.4	4.1	4.1	6.8	*
1999.....	9.0	6.5	5.3	4.3	3.2	4.4	3.4	5.2	*	*

Black, male

2006 4/.....	9.7	9.7	9.5	10.9	6.5	6.1	*	*	*	*
2005 4/.....	10.6	11.5	11.5	8.0	7.3	7.4	*	*	*	*
2004 4/.....	9.5	11.3	10.1	6.9	6.0	6.1	*	*	*	*
2003 4/.....	12.5	10.9	9.9	8.1	6.2	6.2	6.9	9.1	*	*
2002.....	9.7	13.4	10.3	8.5	6.9	*	7.3	10.1	*	*
2001 5/.....	13.9	14.7	11.3	8.8	6.9	6.9	7.1	*	*	*
2000.....	12.6	12.0	9.3	8.9	7.7	9.9	6.7	*	*	*
1999.....	13.6	10.0	9.0	7.6	4.5	8.6	*	*	*	*

Black, female

2006 4/.....	4.3	4.2	3.2	2.3	*	*	*	*	*	*
2005 4/.....	4.2	3.7	3.1	*	*	*	*	*	*	*
2004 4/.....	5.4	4.0	2.9	*	*	*	*	*	*	*
2003 4/.....	3.5	4.5	2.5	*	*	*	*	*	*	*
2002.....	5.5	3.4	2.1	*	*	*	*	*	*	*
2001 5/.....	6.8	5.3	3.7	3.1	*	*	*	*	*	*
2000.....	6.2	3.3	2.0	*	*	*	*	*	*	*
1999.....	5.0	3.5	2.2	*	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2006.....	3.2	3.5	2.8	1.8	1.2	0.9	0.8	1.0	1.1	1.4
2005.....	3.1	3.2	2.9	1.6	1.1	0.9	0.7	0.8	0.9	1.5
2004.....	3.4	3.6	2.6	1.7	1.0	0.9	0.8	0.7	1.3	1.6
2003.....	3.9	3.6	2.6	1.6	1.0	0.8	0.7	0.6	0.9	1.1
2002.....	3.4	3.3	2.3	1.4	1.0	0.8	0.7	0.6	1.0	1.4
2001.....	3.2	3.0	2.0	1.1	0.7	0.6	0.5	0.7	1.1	1.0
2000.....	3.1	2.7	1.7	1.1	0.7	0.6	0.6	0.6	0.7	1.3
1999.....	2.9	2.7	1.6	1.0	0.8	0.8	0.5	0.7	0.9	1.4

All races 3/, male

2006.....	3.8	4.2	3.5	2.3	1.4	1.2	0.9	1.2	1.7	2.2
2005.....	3.6	3.8	3.5	2.1	1.1	1.1	1.0	1.0	*	1.6
2004.....	4.1	4.5	3.3	2.0	1.1	1.2	0.9	0.8	1.3	1.9
2003.....	4.9	4.6	3.4	2.0	1.2	0.8	0.7	0.8	1.1	1.5
2002.....	4.3	4.1	2.9	1.8	1.3	1.0	1.0	0.8	1.1	2.2
2001.....	4.4	4.2	2.5	1.5	0.8	0.9	0.6	0.9	1.5	1.5
2000.....	4.2	3.8	2.3	1.6	0.9	0.9	0.8	1.0	1.1	1.7
1999.....	4.0	3.7	2.1	1.5	1.1	1.1	0.6	1.0	1.3	2.1

All races 3/, female

2006.....	2.6	2.7	2.0	1.5	0.9	0.7	0.7	0.8	0.8	1.0
2005.....	2.6	2.7	2.4	1.2	1.1	0.8	0.5	0.6	0.9	1.4
2004.....	2.7	2.6	2.0	1.3	0.8	0.7	0.6	0.6	1.2	1.6
2003.....	2.9	2.6	1.8	1.1	0.9	0.7	0.6	*	0.8	0.9
2002.....	2.6	2.6	1.7	1.0	0.7	0.7	0.5	0.5	0.9	1.1
2001.....	2.0	1.8	1.5	0.6	0.5	*	*	0.6	0.8	0.8
2000.....	1.9	1.7	1.1	0.7	0.5	*	*	*	*	1.2
1999.....	1.9	1.7	1.0	0.6	0.6	0.5	*	0.5	0.7	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

White, both sexes

2006 4/.....	3.3	3.5	2.7	1.7	1.2	0.9	0.7	0.8	1.1	1.1
2005 4/.....	3.1	3.3	2.9	1.6	1.1	0.9	0.7	0.7	0.9	1.5
2004 4/.....	3.5	3.7	2.6	1.7	1.0	0.8	0.8	0.6	1.3	1.6
2003 4/.....	4.0	3.6	2.5	1.5	0.9	0.7	0.6	0.6	0.9	1.2
2002.....	3.5	3.3	2.2	1.4	0.9	0.8	0.7	0.6	0.9	1.4
2001.....	3.2	2.9	1.9	1.1	0.6	0.6	0.5	0.7	1.0	1.0
2000.....	2.9	2.6	1.6	1.0	0.7	0.5	0.5	0.6	0.8	1.3
1999.....	2.8	2.5	1.4	1.0	0.8	0.8	0.4	0.7	0.9	1.4

White, male

2006 4/.....	3.8	4.1	3.3	2.0	1.4	1.1	0.7	0.8	1.6	2.0
2005 4/.....	3.5	3.7	3.3	2.0	0.9	1.1	1.0	0.8	*	1.7
2004 4/.....	4.1	4.5	3.1	2.0	1.0	1.0	0.9	*	1.4	1.9
2003 4/.....	5.0	4.5	3.2	1.8	1.0	0.7	0.7	0.9	1.1	1.5
2002.....	4.3	4.0	2.5	1.8	1.2	0.9	0.8	*	1.1	2.3
2001.....	4.2	4.0	2.3	1.4	0.7	0.8	0.6	0.9	1.5	*
2000.....	4.0	3.5	2.1	1.3	0.9	0.8	0.7	1.0	*	*
1999.....	3.8	3.2	1.8	1.3	1.0	1.1	*	1.0	1.3	2.1

White, female

2006 4/.....	2.9	2.9	2.1	1.5	1.0	0.7	0.7	0.8	0.7	0.7
2005 4/.....	2.6	2.9	2.5	1.3	1.2	0.8	0.5	0.6	0.8	1.3
2004 4/.....	2.9	2.9	2.0	1.4	0.9	0.6	0.6	0.6	1.2	1.5
2003 4/.....	3.1	2.8	1.9	1.2	0.8	0.7	0.5	*	0.8	1.0
2002.....	2.6	2.6	1.9	1.1	0.7	0.7	0.5	*	0.7	1.1
2001.....	2.1	1.9	1.6	0.7	0.6	*	*	0.6	0.7	0.8
2000.....	1.9	1.7	1.1	0.7	0.5	*	*	*	*	1.1
1999.....	1.7	1.8	1.0	0.7	0.7	0.5	*	*	0.8	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2006.....	3.2	3.4	2.7	1.8	1.1	0.9	0.8	0.9	1.1	1.4
2005.....	3.0	3.2	2.8	1.6	1.0	0.9	0.7	0.7	0.9	1.4
2004.....	3.3	3.5	2.5	1.6	0.9	0.8	0.7	0.6	1.2	1.6
2003.....	3.8	3.5	2.5	1.5	1.0	0.7	0.6	0.6	0.8	1.1
2002.....	3.3	3.2	2.2	1.3	0.9	0.8	0.6	0.6	0.9	1.4
2001.....	3.1	2.9	1.9	1.0	0.6	0.5	0.4	0.6	1.0	1.0
2000.....	3.0	2.7	1.6	1.1	0.7	0.5	0.5	0.5	0.7	1.3
1999.....	2.8	2.6	1.5	1.0	0.7	0.7	0.4	0.6	0.8	1.3

All races 3/, male

2006.....	3.8	4.2	3.5	2.2	1.3	1.1	0.8	1.2	1.7	2.2
2005.....	3.5	3.7	3.3	2.0	1.0	1.1	0.9	0.9	*	1.5
2004.....	3.9	4.5	3.1	1.9	1.0	1.0	0.8	0.7	1.1	1.8
2003.....	4.8	4.5	3.3	1.8	1.1	0.8	0.6	0.7	*	1.4
2002.....	4.1	4.0	2.8	1.7	1.3	0.9	0.8	0.7	1.0	2.2
2001.....	4.2	4.1	2.4	1.4	0.7	0.8	0.6	0.7	1.3	*
2000.....	4.1	3.7	2.1	1.5	0.9	0.7	0.7	0.8	*	*
1999.....	3.8	3.5	2.0	1.4	0.9	0.9	*	0.9	*	1.8

All races 3/, female

2006.....	2.6	2.7	2.0	1.4	0.9	0.6	0.7	0.8	0.8	1.0
2005.....	2.6	2.7	2.4	1.1	1.0	0.7	0.5	0.6	0.9	1.4
2004.....	2.6	2.6	2.0	1.3	0.8	0.6	0.6	0.6	1.2	1.5
2003.....	2.8	2.5	1.8	1.1	0.8	0.7	0.6	*	0.8	0.9
2002.....	2.6	2.5	1.7	0.9	0.7	0.6	0.4	0.5	0.8	1.1
2001.....	2.0	1.7	1.5	0.6	0.5	*	*	0.6	0.7	0.8
2000.....	1.9	1.6	1.1	0.7	0.4	*	*	*	*	1.2
1999.....	1.8	1.7	1.0	0.6	0.5	0.5	*	*	0.7	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

White, both sexes

2006 4/.....	3.3	3.4	2.7	1.7	1.1	0.9	0.7	0.8	1.0	1.1
2005 4/.....	3.0	3.3	2.8	1.6	1.0	0.9	0.6	0.7	0.8	1.4
2004 4/.....	3.4	3.6	2.5	1.6	0.9	0.7	0.7	0.6	1.2	1.6
2003 4/.....	3.9	3.5	2.5	1.4	0.9	0.7	0.5	0.5	0.8	1.2
2002.....	3.4	3.2	2.1	1.3	0.9	0.7	0.6	0.5	0.8	1.4
2001.....	3.1	2.9	1.8	1.0	0.6	0.5	0.4	0.6	0.9	1.0
2000.....	2.8	2.5	1.5	0.9	0.6	0.5	0.5	0.5	0.7	1.2
1999.....	2.6	2.4	1.3	0.9	0.7	0.7	0.4	0.6	0.8	1.3

White, male

2006 4/.....	3.7	4.0	3.2	1.9	1.3	1.1	0.6	0.8	1.5	2.0
2005 4/.....	3.4	3.7	3.1	1.9	0.8	1.0	0.8	0.7	*	1.6
2004 4/.....	4.0	4.4	3.0	1.9	0.9	0.8	0.8	*	1.1	1.8
2003 4/.....	4.8	4.4	3.1	1.7	0.9	0.6	*	0.7	*	*
2002.....	4.2	3.9	2.4	1.7	1.2	0.8	0.7	*	*	2.3
2001.....	4.1	3.9	2.2	1.3	0.6	0.7	*	0.7	1.4	*
2000.....	3.8	3.5	2.0	1.2	0.8	0.6	0.6	0.8	*	*
1999.....	3.6	3.1	1.7	1.2	0.9	0.9	*	0.8	*	*

White, female

2006 4/.....	2.9	2.9	2.1	1.4	1.0	0.7	0.7	0.8	0.7	0.7
2005 4/.....	2.6	2.9	2.5	1.2	1.1	0.8	0.5	0.6	0.8	1.3
2004 4/.....	2.8	2.8	2.0	1.4	0.8	0.5	0.6	0.6	1.2	1.5
2003 4/.....	3.0	2.7	1.9	1.2	0.8	0.7	0.5	*	0.8	1.0
2002.....	2.6	2.5	1.9	1.0	0.7	0.6	0.5	*	0.7	1.1
2001.....	2.1	1.8	1.5	0.7	0.6	*	*	0.6	*	0.8
2000.....	1.8	1.6	1.1	0.7	0.4	*	*	*	*	1.1
1999.....	1.7	1.7	1.0	0.7	0.6	0.5	*	*	0.8	1.1

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes

2006.....	0.4	0.5	0.8	0.9	1.3	2.0	3.1	4.7	6.1	8.3
2005.....	0.4	0.5	0.8	1.0	1.4	2.0	3.3	4.7	7.2	9.1
2004.....	0.4	0.6	0.8	1.2	1.6	2.6	3.5	5.0	7.6	10.6
2003.....	0.5	0.7	0.7	1.1	1.7	2.2	3.7	5.3	7.3	11.2
2002.....	0.4	0.5	0.8	1.1	1.6	2.3	3.9	5.1	7.7	11.3
2001.....	0.4	0.7	0.8	1.3	1.9	2.6	3.6	5.9	8.3	12.8
2000.....	0.4	0.6	0.9	1.2	2.0	2.8	4.3	5.8	8.0	13.6
1999.....	0.4	0.5	0.7	1.1	1.7	2.4	4.0	5.5	8.4	12.7

All races 3/, male

2006.....	0.3	0.5	0.9	1.0	1.2	2.0	2.9	5.3	7.1	8.8
2005.....	0.4	0.5	0.8	1.1	1.7	2.3	3.7	4.6	8.9	10.8
2004.....	0.4	0.6	0.8	1.5	1.7	2.5	4.3	5.8	9.6	11.9
2003.....	0.5	0.7	0.7	1.3	1.6	2.3	4.2	5.6	8.8	12.9
2002.....	0.4	0.6	0.7	1.2	1.5	2.5	3.8	5.4	7.7	12.3
2001.....	0.4	0.7	0.8	1.3	2.0	2.6	3.8	6.3	9.9	14.5
2000.....	0.4	0.6	1.0	1.2	2.0	2.8	5.3	7.0	10.4	16.0
1999.....	0.4	0.4	0.7	1.2	1.7	2.3	4.5	6.4	10.1	16.5

All races 3/, female

2006.....	0.4	0.5	0.7	0.9	1.4	2.0	3.3	4.2	5.4	8.1
2005.....	0.4	0.6	0.9	1.0	1.2	1.8	3.0	4.8	6.2	8.3
2004.....	0.5	0.6	0.8	1.0	1.5	2.7	2.9	4.3	6.3	9.9
2003.....	0.4	0.6	0.6	0.9	1.7	2.1	3.3	5.1	6.4	10.5
2002.....	0.4	0.5	0.9	1.0	1.7	2.2	3.9	5.0	7.7	10.9
2001.....	0.5	0.6	0.8	1.2	1.7	2.6	3.5	5.7	7.3	12.1
2000.....	0.4	0.7	0.9	1.1	2.0	2.7	3.6	4.9	6.7	12.7
1999.....	0.4	0.5	0.7	0.9	1.6	2.5	3.6	4.8	7.4	11.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Complications of medical and surgical care (Y40-Y84,Y88)

White, both sexes

2006 4/.....	0.3	0.5	0.7	0.8	1.2	1.9	3.1	4.5	5.9	8.3
2005 4/.....	0.4	0.5	0.7	0.9	1.3	1.8	3.0	4.4	7.3	9.2
2004 4/.....	0.4	0.6	0.8	1.1	1.5	2.4	3.4	4.8	7.3	10.4
2003 4/.....	0.4	0.6	0.6	1.0	1.6	2.1	3.6	5.1	7.4	11.0
2002.....	0.4	0.5	0.7	1.0	1.5	2.2	3.8	5.0	7.6	11.2
2001.....	0.4	0.6	0.7	1.1	1.6	2.4	3.2	5.8	8.2	12.1
2000.....	0.3	0.6	0.9	1.0	1.8	2.6	4.1	5.4	8.0	13.1
1999.....	0.3	0.4	0.6	0.9	1.5	2.3	3.8	5.2	8.0	12.1

White, male

2006 4/.....	0.3	0.4	0.8	0.8	1.1	2.0	2.8	5.1	7.2	8.8
2005 4/.....	0.4	0.4	0.7	0.9	1.5	2.3	3.4	4.3	8.8	11.2
2004 4/.....	0.3	0.6	0.8	1.3	1.7	2.5	4.2	5.8	9.5	11.8
2003 4/.....	0.6	0.6	0.5	1.2	1.6	2.3	4.1	5.5	9.1	12.8
2002.....	0.4	0.5	0.6	1.0	1.5	2.2	3.7	5.3	7.8	11.8
2001.....	0.3	0.6	0.6	1.1	1.7	2.5	3.2	6.2	10.0	13.6
2000.....	0.4	0.5	1.0	1.1	1.8	2.5	5.0	6.8	10.6	15.8
1999.....	0.3	0.3	0.6	1.0	1.6	2.3	4.2	6.2	9.7	15.6

White, female

2006 4/.....	0.3	0.5	0.5	0.7	1.3	1.8	3.3	4.0	5.0	8.0
2005 4/.....	0.4	0.5	0.7	1.0	1.1	1.5	2.6	4.5	6.3	8.3
2004 4/.....	0.4	0.5	0.7	0.9	1.4	2.4	2.7	4.0	6.0	9.8
2003 4/.....	0.3	0.5	0.6	0.8	1.6	2.0	3.1	4.7	6.3	10.2
2002.....	0.3	0.4	0.7	0.9	1.5	2.1	3.9	4.8	7.5	10.9
2001.....	0.4	0.5	0.7	1.1	1.6	2.4	3.2	5.6	7.0	11.5
2000.....	0.3	0.6	0.9	0.9	1.8	2.7	3.3	4.5	6.4	12.0
1999.....	0.4	0.5	0.6	0.7	1.5	2.4	3.4	4.5	7.1	10.7

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Complications of medical and surgical care (Y40-Y84,Y88)

Black, both sexes

2006 4/.....	0.7	1.0	1.7	2.3	3.1	3.9	4.3	8.2	9.5	10.4
2005 4/.....	*	1.0	2.1	2.2	2.9	4.5	6.4	9.4	8.5	9.1
2004 4/.....	0.9	1.0	1.4	2.7	2.5	4.5	5.5	7.6	11.9	13.8
2003 4/.....	*	1.3	1.6	2.5	3.0	3.6	5.8	9.5	7.6	15.2
2002.....	1.0	1.2	2.1	2.6	2.8	4.0	5.2	7.6	9.4	14.0
2001.....	1.0	1.6	1.5	2.7	4.2	4.6	8.5	8.0	11.1	23.3
2000.....	*	1.1	1.7	2.6	4.7	4.6	7.9	10.5	10.4	21.0
1999.....	0.8	0.9	1.8	2.6	2.6	3.5	7.3	9.5	12.4	20.1

Black, male

2006 4/.....	*	*	*	*	*	*	*	9.4	*	*
2005 4/.....	*	*	2.2	2.6	3.8	*	*	9.4	*	*
2004 4/.....	*	*	*	3.4	*	*	*	*	*	*
2003 4/.....	*	*	2.1	2.8	*	*	*	*	*	*
2002.....	*	*	*	*	*	5.0	*	*	*	*
2001.....	*	1.7	*	3.4	5.1	5.3	11.1	*	*	29.5
2000.....	*	*	*	*	4.6	6.2	9.4	10.4	*	*
1999.....	*	*	*	*	*	*	9.2	10.4	*	23.4

Black, female

2006 4/.....	*	*	1.9	2.6	2.9	4.2	*	7.5	9.4	10.2
2005 4/.....	*	*	2.1	*	*	4.8	6.7	9.3	7.4	9.0
2004 4/.....	*	*	1.7	*	*	5.4	5.1	7.5	11.0	13.4
2003 4/.....	*	*	*	*	3.0	3.8	6.4	10.6	7.7	15.4
2002.....	*	*	2.5	2.4	3.1	*	4.9	8.2	9.9	11.5
2001.....	*	*	*	*	3.5	4.1	6.7	8.0	11.0	21.0
2000.....	*	*	*	3.1	4.7	*	6.9	10.5	10.6	21.9
1999.....	*	*	*	*	*	*	6.1	8.9	10.8	18.8

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 6/

All races 3/, both sexes

2006.....	10.1	15.7	19.7	19.9	18.6	16.0	13.2	11.0	8.1	4.9
2005.....	10.1	15.7	19.7	19.7	19.3	16.3	13.7	10.1	8.1	4.6
2004.....	10.0	15.5	19.5	18.9	18.5	17.0	14.1	10.7	7.3	4.3
2003.....	10.6	16.0	18.9	18.3	18.8	16.8	13.2	10.3	7.6	4.1
2002.....	10.6	15.6	18.2	18.2	18.6	16.5	14.4	10.9	7.2	4.4
2001.....	10.6	16.5	17.9	17.6	18.9	17.3	13.7	11.0	7.5	5.0
2000.....	11.1	15.9	16.7	18.2	19.3	17.2	14.3	11.2	7.9	5.4
1999.....	10.9	16.0	16.9	18.1	19.3	17.6	14.1	11.9	8.5	5.5

All races 3/, male

2006.....	14.1	22.8	30.6	31.8	30.1	27.1	22.2	19.1	14.1	9.5
2005.....	14.4	22.9	30.9	31.7	31.3	27.0	22.6	18.4	14.9	9.3
2004.....	14.6	22.7	30.6	30.0	30.4	28.1	24.5	19.4	13.8	9.0
2003.....	15.2	24.5	29.9	29.1	30.6	27.6	22.8	18.0	15.0	7.8
2002.....	15.3	23.7	29.2	29.2	31.0	27.9	23.8	18.7	13.4	9.7
2001.....	15.3	25.4	28.0	28.7	31.0	28.5	23.8	19.4	13.8	11.6
2000.....	16.4	25.0	26.9	28.8	31.4	28.9	25.0	21.1	14.9	11.9
1999.....	16.6	25.2	27.4	29.7	31.5	29.9	24.5	21.2	16.6	11.8

All races 3/, female

2006.....	6.2	8.7	9.1	8.7	8.2	6.4	5.9	5.0	4.3	2.7
2005.....	5.8	8.6	8.9	8.4	8.4	7.1	6.5	4.0	3.8	2.5
2004.....	5.3	8.4	8.8	8.3	7.8	7.3	5.8	4.4	3.3	2.3
2003.....	6.0	7.7	8.3	8.1	8.1	7.5	5.4	4.9	3.1	2.5
2002.....	6.1	7.7	7.7	7.9	7.2	6.7	6.9	5.3	3.4	2.1
2001.....	6.0	7.9	8.3	7.2	8.0	7.6	5.7	5.1	3.8	2.3
2000.....	5.8	7.1	7.0	8.3	8.3	7.2	5.9	4.3	3.8	2.7
1999.....	5.3	7.1	6.8	7.3	8.3	7.0	6.0	5.5	3.8	3.0

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 6/

White, both sexes

2006 4/.....	10.5	16.4	20.3	20.1	19.0	16.1	13.3	11.4	8.3	5.0
2005 4/.....	10.4	16.1	20.0	20.2	19.1	16.6	13.8	10.1	8.0	4.6
2004 4/.....	10.4	16.0	19.8	18.7	18.2	16.6	14.0	10.9	7.5	4.5
2003 4/.....	10.8	16.2	18.7	18.2	18.2	17.0	13.2	10.6	7.6	4.2
2002.....	10.9	15.8	18.2	17.7	17.9	16.5	14.2	10.9	7.1	4.3
2001.....	10.4	16.5	17.1	16.8	18.6	16.7	13.8	10.9	7.7	4.8
2000.....	10.8	15.5	16.1	17.4	18.7	16.9	14.2	11.1	7.9	5.3
1999.....	10.5	15.3	16.2	17.1	18.5	17.0	13.6	11.6	8.3	5.3

White, male

2006 4/.....	14.7	23.7	31.4	31.7	30.4	26.8	22.1	19.7	14.4	9.7
2005 4/.....	14.9	23.4	31.2	32.3	30.8	27.0	22.4	18.2	14.8	9.4
2004 4/.....	15.2	23.3	31.1	29.5	29.5	27.1	23.8	19.5	14.2	9.3
2003 4/.....	15.6	24.7	29.3	28.9	29.2	27.6	22.7	18.2	14.6	7.8
2002.....	15.8	24.0	29.1	28.4	29.8	27.5	23.0	18.7	13.2	9.3
2001.....	14.9	25.4	26.8	27.0	30.2	27.1	23.5	19.2	14.0	11.1
2000.....	16.0	24.6	26.0	27.6	30.3	28.1	24.5	20.8	14.9	11.6
1999.....	16.1	24.2	26.3	27.7	30.4	28.4	23.4	20.4	16.4	11.4

White, female

2006 4/.....	6.3	9.1	9.4	8.9	8.4	6.6	6.1	5.2	4.4	2.8
2005 4/.....	5.8	8.9	9.0	8.5	8.2	7.3	6.6	4.1	3.8	2.5
2004 4/.....	5.5	8.7	8.8	8.3	7.7	7.4	6.0	4.6	3.3	2.3
2003 4/.....	6.0	7.8	8.3	8.0	8.0	7.5	5.4	5.1	3.3	2.5
2002.....	6.1	7.7	7.4	7.5	6.9	6.8	7.1	5.3	3.4	2.2
2001.....	5.9	7.6	7.6	7.0	8.0	7.6	5.8	5.1	3.9	2.2
2000.....	5.6	6.5	6.5	7.7	8.0	7.1	5.9	4.4	3.7	2.7
1999.....	4.8	6.5	6.3	7.0	7.6	7.0	5.9	5.4	3.6	2.9

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 6/

Black, both sexes

2006 4/.....	7.3	12.0	17.9	21.1	19.3	17.5	13.4	9.4	6.6	*
2005 4/.....	8.1	14.6	19.6	20.3	23.5	17.0	15.9	10.7	8.5	5.6
2004 4/.....	7.5	14.2	20.6	23.2	23.9	22.4	17.1	8.6	6.2	*
2003 4/.....	8.7	15.1	22.6	22.4	26.7	19.0	14.3	9.0	8.6	*
2002.....	9.0	15.8	21.8	25.3	25.3	18.7	17.7	12.5	9.2	*
2001.....	12.2	18.4	27.4	26.9	24.6	25.3	14.7	13.5	7.3	*
2000.....	13.0	21.6	24.1	26.6	26.1	23.9	17.6	13.5	8.2	6.3
1999.....	12.9	22.8	24.1	29.3	28.9	24.9	21.2	16.4	11.6	7.7

Black, male

2006 4/.....	9.9	18.1	28.9	37.2	33.9	33.7	25.5	18.3	*	*
2005 4/.....	11.1	23.1	31.7	33.6	41.3	31.1	30.4	24.1	15.5	*
2004 4/.....	11.5	22.2	33.0	40.6	42.6	42.9	35.0	17.3	*	*
2003 4/.....	12.6	24.7	39.4	37.2	49.2	33.0	27.6	17.3	21.4	*
2002.....	12.5	24.4	35.3	42.0	45.5	36.4	35.1	23.4	19.1	*
2001.....	18.2	28.9	43.9	47.9	44.3	47.8	29.6	26.0	*	*
2000.....	19.3	32.9	39.9	43.6	44.1	42.9	36.1	28.4	16.9	*
1999.....	18.9	35.9	39.9	52.9	48.6	48.3	41.4	31.3	22.7	*

Black, female

2006 4/.....	5.0	6.6	8.6	7.9	7.8	5.5	5.1	*	*	*
2005 4/.....	5.4	7.3	9.4	9.3	9.6	6.5	6.1	*	*	*
2004 4/.....	4.0	7.3	10.1	8.8	9.3	7.2	5.1	*	*	*
2003 4/.....	5.3	6.7	8.4	10.3	9.2	8.6	5.3	*	*	*
2002.....	5.8	8.4	10.3	11.6	9.4	5.6	6.0	6.0	*	*
2001.....	6.9	9.2	13.4	9.6	9.1	8.6	4.7	6.1	*	*
2000.....	7.4	11.7	10.7	12.7	11.9	9.8	5.3	*	*	*
1999.....	7.6	11.6	10.8	10.1	13.5	7.5	7.9	7.4	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

All races 3/, both sexes

2006.....	26.2	28.4	24.8	15.2	8.1	5.7	4.9	5.5	6.8	8.1
2005.....	24.0	25.8	21.2	12.8	7.8	5.0	4.4	4.9	6.1	7.6
2004.....	23.2	24.4	18.5	11.0	6.6	4.4	4.1	4.7	4.9	6.3
2003.....	22.8	22.7	17.0	9.8	5.7	4.5	3.7	3.9	4.6	6.0
2002.....	21.8	20.9	14.5	8.2	5.1	3.8	3.3	3.6	4.1	5.7
2001.....	18.5	17.6	11.6	6.3	4.1	3.1	2.9	3.4	3.6	5.1
2000.....	17.5	15.8	10.3	5.9	3.6	2.6	2.7	3.2	3.9	5.7
1999.....	17.4	15.5	9.3	5.8	3.8	3.2	2.9	3.6	4.2	4.8

All races 3/, male

2006.....	32.9	35.0	31.3	18.7	9.5	6.0	5.5	6.2	8.0	10.7
2005.....	29.9	31.9	26.6	16.0	9.0	5.4	5.2	6.6	6.9	9.9
2004.....	28.3	30.1	23.6	13.0	7.6	4.8	4.5	5.2	5.6	8.6
2003.....	29.1	29.2	21.7	11.5	6.6	5.2	3.9	4.7	5.3	7.8
2002.....	28.0	26.4	19.1	10.0	5.7	4.1	3.4	3.9	4.7	7.3
2001.....	24.6	23.9	15.0	7.7	4.3	3.7	3.3	3.8	3.9	5.7
2000.....	23.8	21.9	13.8	6.7	3.9	2.9	2.8	3.6	4.3	7.5
1999.....	24.3	21.9	12.2	7.1	4.7	3.7	3.1	3.5	4.4	6.2

All races 3/, female

2006.....	19.6	21.9	18.6	12.0	6.9	5.4	4.3	5.0	6.0	6.8
2005.....	18.2	19.8	16.1	9.8	6.6	4.8	3.8	3.7	5.5	6.6
2004.....	18.2	19.0	13.7	9.1	5.7	4.1	3.7	4.4	4.4	5.2
2003.....	16.6	16.5	12.4	8.2	4.9	4.0	3.4	3.3	4.2	5.2
2002.....	15.8	15.6	10.2	6.6	4.6	3.6	3.3	3.3	3.8	4.9
2001.....	12.6	11.6	8.3	5.1	3.9	2.6	2.6	3.2	3.4	4.9
2000.....	11.4	9.8	7.0	5.2	3.4	2.3	2.7	2.9	3.8	4.9
1999.....	10.6	9.2	6.6	4.6	2.9	2.7	2.7	3.7	4.0	4.3

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

White, both sexes

2006 4/.....	27.5	28.9	24.3	14.6	7.6	5.6	4.8	5.6	7.0	8.3
2005 4/.....	25.4	26.5	20.5	12.2	7.8	5.1	4.6	5.0	6.3	8.0
2004 4/.....	24.8	24.9	18.2	10.6	6.4	4.3	4.3	4.9	5.0	6.4
2003 4/.....	23.9	23.1	16.5	9.5	5.6	4.4	3.8	4.2	4.9	6.3
2002.....	22.6	20.8	13.8	7.9	4.9	3.9	3.5	3.7	4.3	5.8
2001.....	18.7	16.9	10.8	6.1	3.9	3.1	3.0	3.5	3.7	5.1
2000.....	17.3	15.0	9.5	5.7	3.6	2.5	2.8	3.4	4.1	5.6
1999.....	16.9	14.5	8.6	5.5	3.5	3.1	2.9	3.8	4.4	4.8

White, male

2006 4/.....	34.1	34.9	29.4	16.8	8.1	5.4	5.0	6.2	8.5	11.2
2005 4/.....	31.4	32.0	24.8	14.4	8.4	5.0	5.2	6.7	6.9	10.6
2004 4/.....	29.9	30.4	22.1	11.6	6.8	4.2	4.6	5.0	5.7	8.7
2003 4/.....	30.3	29.2	20.4	10.7	6.2	4.7	3.9	4.9	5.5	7.8
2002.....	29.0	25.9	17.3	9.0	4.9	3.8	3.4	4.0	4.6	7.6
2001.....	24.8	22.3	13.3	6.9	3.6	3.6	3.2	3.6	4.1	5.6
2000.....	23.4	20.4	12.1	5.9	3.6	2.7	2.7	3.8	4.4	7.2
1999.....	23.5	20.0	10.6	6.2	4.0	3.5	3.1	3.5	4.6	6.4

White, female

2006 4/.....	20.8	23.0	19.2	12.5	7.3	5.8	4.6	5.2	6.1	6.9
2005 4/.....	19.2	21.0	16.2	10.2	7.2	5.1	4.1	3.8	5.9	6.8
2004 4/.....	19.6	19.5	14.3	9.6	6.1	4.5	4.1	4.8	4.5	5.3
2003 4/.....	17.3	17.0	12.7	8.5	5.2	4.2	3.6	3.6	4.4	5.6
2002.....	16.3	15.8	10.4	7.0	4.9	3.9	3.5	3.5	4.2	4.9
2001.....	12.7	11.6	8.4	5.3	4.2	2.6	2.8	3.5	3.5	4.9
2000.....	11.3	9.7	6.9	5.5	3.7	2.4	3.0	3.0	3.9	4.9
1999.....	10.1	9.1	6.6	4.8	3.2	2.8	2.7	4.0	4.2	4.2

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

Black, both sexes

2006 4/.....	27.3	33.5	36.7	25.3	14.9	7.4	6.3	4.7	*	6.5
2005 4/.....	23.8	29.3	32.8	22.4	10.4	5.6	4.6	4.3	*	*
2004 4/.....	21.0	29.0	27.0	18.0	9.4	5.9	3.0	3.7	*	*
2003 4/.....	23.6	28.2	25.4	14.6	7.8	6.4	3.2	*	*	*
2002.....	24.1	28.2	25.0	13.6	8.9	3.7	*	*	*	*
2001.....	23.3	28.0	21.7	10.2	7.1	3.5	3.0	*	*	6.1
2000.....	23.9	26.0	21.0	9.6	4.8	3.6	*	*	*	7.5
1999.....	25.7	27.8	18.4	9.8	6.5	3.9	3.0	*	*	*

Black, male

2006 4/.....	37.2	47.1	56.4	41.2	25.8	13.1	11.7	*	*	*
2005 4/.....	30.2	41.3	48.4	36.0	19.1	10.0	7.6	*	*	*
2004 4/.....	26.9	37.1	41.6	29.7	16.1	10.9	*	*	*	*
2003 4/.....	31.1	39.3	38.5	23.3	13.1	11.6	*	*	*	*
2002.....	31.1	38.7	39.9	22.8	16.0	7.0	*	*	*	*
2001.....	31.7	43.3	34.4	17.1	12.6	5.3	*	*	*	*
2000.....	33.8	40.7	33.6	16.0	7.9	6.0	*	*	*	*
1999.....	36.8	44.3	30.2	17.6	12.1	6.5	*	*	*	*

Black, female

2006 4/.....	18.6	21.7	19.9	12.3	6.3	*	*	*	*	7.3
2005 4/.....	18.1	18.9	19.6	11.2	3.5	*	*	*	*	*
2004 4/.....	15.9	22.0	14.6	8.4	4.2	*	*	*	*	*
2003 4/.....	17.0	18.5	14.3	7.5	3.6	*	*	*	*	*
2002.....	17.9	19.1	12.4	6.1	3.3	*	*	*	*	*
2001.....	15.9	14.8	10.8	4.6	*	*	*	*	*	*
2000.....	15.1	13.3	10.2	4.5	*	*	*	*	*	*
1999.....	15.9	13.5	8.5	3.3	*	*	*	*	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

All races 3/, both sexes

2006.....	11.1	11.3	11.0	9.9	9.7	9.7	10.4	12.8	13.1	11.5
2005.....	11.6	11.4	10.9	9.8	9.9	9.5	11.3	13.0	14.5	12.0
2004.....	11.4	11.3	10.7	9.7	10.0	9.5	11.0	13.3	13.4	11.9
2003.....	11.5	11.4	10.9	10.4	9.6	9.7	11.3	12.9	14.2	12.5
2002.....	11.9	11.1	10.5	10.7	9.6	10.1	11.8	13.6	15.5	12.5
2001.....	11.3	10.6	10.4	10.4	9.7	10.5	11.4	13.7	15.3	12.8
2000.....	11.8	10.7	10.3	9.6	9.1	9.7	11.5	13.4	14.7	14.2
1999.....	10.9	10.7	9.6	9.9	9.4	10.0	12.2	14.0	14.5	13.5

All races 3/, male

2006.....	18.3	18.7	18.4	17.0	17.0	18.0	20.8	28.0	30.7	33.6
2005.....	19.2	19.1	18.7	17.0	17.9	17.4	22.6	28.1	34.6	35.4
2004.....	18.5	18.4	18.0	16.9	17.4	17.8	22.1	28.2	32.0	35.9
2003.....	18.9	18.9	18.6	17.8	17.6	18.5	22.5	27.8	33.8	37.8
2002.....	19.7	18.4	18.0	18.6	17.1	18.8	24.1	29.8	37.6	38.9
2001.....	18.6	17.9	17.6	17.9	16.7	19.6	22.9	29.9	37.3	40.2
2000.....	19.4	17.9	17.3	16.5	16.1	17.8	23.1	29.6	36.6	44.7
1999.....	17.7	17.8	16.1	17.1	17.1	18.9	24.5	30.8	36.2	43.3

All races 3/, female

2006.....	4.1	4.1	3.9	3.2	3.1	2.4	1.9	1.6	2.1	1.1
2005.....	4.1	4.0	3.3	3.0	2.6	2.7	2.3	2.1	2.0	1.3
2004.....	4.4	4.3	3.7	3.0	3.2	2.3	2.1	2.6	1.9	1.0
2003.....	4.2	4.1	3.5	3.4	2.4	2.1	2.3	2.3	2.2	1.3
2002.....	4.2	4.0	3.2	3.3	2.8	2.5	2.0	2.2	2.0	1.1
2001.....	4.1	3.6	3.5	3.3	3.2	2.6	2.2	2.3	2.2	1.3
2000.....	4.3	3.7	3.5	3.1	2.8	2.7	2.4	2.2	1.9	1.7
1999.....	4.2	3.7	3.4	3.2	2.4	2.4	2.5	2.3	1.9	1.5

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

White, both sexes

2006 4/.....	10.7	11.5	11.4	10.8	10.4	10.5	11.3	13.8	14.3	12.2
2005 4/.....	11.1	11.7	11.4	10.7	10.7	10.3	12.2	14.2	15.6	13.0
2004 4/.....	11.1	11.6	11.3	10.2	10.7	10.3	11.8	14.3	14.4	12.8
2003 4/.....	11.2	11.6	11.5	11.1	10.4	10.5	12.1	13.9	15.1	13.5
2002.....	11.6	11.3	10.8	11.5	10.2	11.0	12.8	14.4	16.7	13.4
2001.....	11.0	10.9	10.9	10.9	10.5	11.3	12.1	14.6	16.1	13.8
2000.....	11.5	10.7	10.6	10.0	9.8	10.4	12.1	14.5	15.7	15.2
1999.....	10.6	10.6	10.0	10.4	9.8	10.7	12.9	14.9	15.4	14.5

White, male

2006 4/.....	17.3	18.5	18.9	18.2	18.0	19.2	22.5	30.0	33.3	35.8
2005 4/.....	18.0	19.2	19.4	18.2	19.2	18.9	24.0	30.0	37.2	38.0
2004 4/.....	17.6	18.6	18.7	17.4	18.6	18.9	23.4	30.1	34.2	38.9
2003 4/.....	18.1	18.9	19.4	18.9	18.9	20.0	24.1	29.7	36.0	41.0
2002.....	19.1	18.4	18.5	19.7	18.1	20.3	25.8	31.2	40.3	41.7
2001.....	18.0	18.2	18.3	18.6	18.0	20.9	24.1	31.8	39.2	43.3
2000.....	18.7	17.8	17.6	17.0	17.0	18.9	24.1	31.9	38.8	48.4
1999.....	17.1	17.6	16.7	17.5	17.7	20.1	25.8	32.6	38.2	46.8

White, female

2006 4/.....	4.0	4.5	4.0	3.6	3.4	2.7	2.1	1.8	2.3	1.1
2005 4/.....	4.2	4.2	3.6	3.4	2.7	2.8	2.4	2.4	2.1	1.4
2004 4/.....	4.6	4.5	4.0	3.2	3.4	2.6	2.3	2.7	2.0	1.1
2003 4/.....	4.2	4.4	3.6	3.7	2.5	2.2	2.3	2.4	2.2	1.3
2002.....	4.1	4.2	3.3	3.6	2.9	2.8	2.1	2.3	2.2	1.2
2001.....	4.0	3.7	3.7	3.6	3.6	2.9	2.3	2.4	2.2	1.4
2000.....	4.1	3.7	3.7	3.3	3.1	2.9	2.6	2.2	1.9	1.8
1999.....	4.1	3.8	3.4	3.6	2.5	2.5	2.7	2.4	1.9	1.6

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Black, both sexes

2006 4/.....	17.2	13.2	11.3	6.6	6.9	6.5	5.3	6.0	*	6.2
2005 4/.....	18.1	12.9	10.1	6.0	6.2	5.7	7.6	5.3	6.3	*
2004 4/.....	15.8	12.1	9.9	8.8	7.0	6.0	6.5	7.1	5.9	*
2003 4/.....	16.6	12.4	10.2	7.3	6.3	5.9	6.2	6.0	6.3	*
2002.....	16.6	13.0	10.3	7.3	7.0	5.1	6.7	8.8	*	*
2001.....	15.4	11.3	9.6	8.1	5.5	5.9	7.4	6.9	8.4	*
2000.....	16.5	13.2	10.6	7.4	5.8	6.2	7.7	4.6	6.8	*
1999.....	15.6	13.6	8.6	8.5	7.9	6.4	6.9	8.4	6.1	*

Black, male

2006 4/.....	30.9	25.1	20.4	12.8	13.4	13.4	11.4	15.7	*	*
2005 4/.....	33.3	23.5	18.8	11.8	11.1	10.2	16.1	14.3	14.7	*
2004 4/.....	29.2	21.6	18.6	16.5	12.3	13.3	15.9	15.9	15.8	*
2003 4/.....	29.4	23.4	18.2	13.7	11.8	11.3	12.2	13.2	14.8	*
2002.....	29.1	24.1	18.1	14.4	12.3	10.2	14.7	20.7	*	*
2001.....	26.8	20.4	17.4	16.3	10.4	12.2	15.8	16.3	20.9	*
2000.....	28.5	23.4	19.4	13.7	11.2	12.7	18.1	9.5	16.9	*
1999.....	26.9	24.4	15.2	17.3	15.5	12.7	15.6	19.5	*	*

Worktable 291R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second-Edition 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Black, female

2006 4/.....	5.2	2.9	3.7	*	*	*	*	*	*	*
2005 4/.....	4.8	3.7	2.7	*	*	*	*	*	*	*
2004 4/.....	4.0	4.0	2.5	2.5	*	*	*	*	*	*
2003 4/.....	5.4	2.8	3.4	*	*	*	*	*	*	*
2002.....	5.6	3.3	3.7	*	*	*	*	*	*	*
2001.....	5.4	3.5	2.9	*	*	*	*	*	*	*
2000.....	5.9	4.4	3.1	*	*	*	*	*	*	*
1999.....	5.6	4.3	3.1	*	*	*	*	*	*	*

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Includes races other than white and black.

4/ Multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2006, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality."

5/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to Appendix from Vital Statistics of the United States: Mortality, 2001."

6/ Tabular data in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.