



Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Certain other intestinal infections (A04,A07-A09)

2007.....	2.1	2.1	2.2	1.4	1.4	1.3	2.2	2.1	2.2	2.3	2.2	2.3	1.6	1.6	1.6
2006.....	2.1	2.1	2.1	1.5	1.4	1.5	2.2	2.1	2.1	2.3	2.2	2.3	1.4	1.4	1.4
2005.....	1.8	1.9	1.8	1.1	0.9	1.2	1.9	1.9	1.9	2.0	2.0	2.0	1.2	1.3	1.2
2004.....	1.5	1.4	1.5	0.9	0.8	0.9	1.5	1.4	1.5	1.6	1.5	1.6	1.0	1.0	0.9
2003.....	1.0	1.0	1.1	0.6	0.7	0.5	1.1	1.0	1.1	1.1	1.0	1.1	0.7	0.7	0.8
2002.....	0.8	0.8	0.8	0.5	0.5	0.5	0.9	0.9	0.8	0.9	0.9	0.9	0.6	0.7	0.6
2001.....	0.5	0.5	0.5	0.4	*	0.4	0.5	0.5	0.5	0.6	0.6	0.6	0.4	0.5	0.4
2000.....	0.5	0.5	0.5	0.3	0.4	0.3	0.5	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.3
1999.....	0.4	0.4	0.4	0.2	*	*	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3

Tuberculosis (A16-A19)

2007.....	0.2	0.2	0.1	0.4	0.6	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.4	0.6	0.2
2006.....	0.2	0.3	0.1	0.4	0.6	0.3	0.2	0.3	0.1	0.1	0.1	0.1	0.5	0.9	0.3
2005.....	0.2	0.3	0.1	0.4	0.6	0.2	0.2	0.3	0.1	0.1	0.1	0.1	0.5	0.8	0.3
2004.....	0.2	0.3	0.1	0.5	0.8	0.3	0.2	0.3	0.1	0.1	0.2	0.1	0.7	1.0	0.4
2003.....	0.2	0.3	0.1	0.4	0.6	0.3	0.2	0.3	0.1	0.1	0.2	0.1	0.6	1.0	0.3
2002.....	0.3	0.4	0.2	0.5	0.6	0.4	0.3	0.4	0.2	0.1	0.2	0.1	0.8	1.1	0.5
2001.....	0.3	0.4	0.2	0.5	0.8	0.3	0.2	0.3	0.2	0.1	0.2	0.1	0.8	1.2	0.5
2000.....	0.3	0.4	0.2	0.5	0.7	0.4	0.2	0.4	0.2	0.1	0.2	0.1	0.9	1.5	0.6
1999.....	0.3	0.5	0.2	0.8	1.1	0.6	0.3	0.5	0.2	0.2	0.3	0.1	1.0	1.5	0.6

Respiratory tuberculosis (A16)

2007.....	0.1	0.2	0.1	0.3	0.5	0.2	0.1	0.2	0.1	0.1	0.1	0.0	0.3	0.5	0.2
2006.....	0.2	0.3	0.1	0.3	0.4	0.2	0.1	0.2	0.1	0.1	0.1	0.0	0.4	0.7	0.2
2005.....	0.1	0.2	0.1	0.2	0.3	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.4	0.6	0.2
2004.....	0.2	0.2	0.1	0.4	0.6	0.3	0.1	0.2	0.1	0.1	0.1	0.1	0.5	0.8	0.3
2003.....	0.2	0.3	0.1	0.4	0.6	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.5	0.8	0.3
2002.....	0.2	0.3	0.1	0.4	0.5	0.3	0.2	0.3	0.1	0.1	0.2	0.1	0.6	0.9	0.4
2001.....	0.2	0.3	0.1	0.4	0.7	0.2	0.2	0.3	0.1	0.1	0.2	0.1	0.6	1.0	0.3
2000.....	0.2	0.3	0.1	0.4	0.7	0.2	0.2	0.3	0.1	0.1	0.2	0.1	0.8	1.3	0.4
1999.....	0.3	0.4	0.2	0.6	0.8	0.5	0.3	0.4	0.1	0.2	0.3	0.1	0.8	1.2	0.5









Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)</b>															
2007.....	1.8	2.2	1.6	1.7	2.0	1.6	1.8	2.2	1.6	1.8	2.1	1.5	2.4	3.0	2.1
2006.....	1.9	2.3	1.6	1.5	1.9	1.2	1.9	2.3	1.7	1.8	2.2	1.6	2.7	3.2	2.3
2005.....	2.5	3.1	2.0	2.9	3.4	2.4	2.5	3.0	2.0	2.4	2.9	1.9	3.2	4.0	2.6
2004.....	2.5	3.1	2.0	2.9	3.7	2.3	2.5	3.0	2.0	2.4	2.9	1.9	3.3	4.2	2.6
2003.....	2.5	3.1	2.1	2.9	3.7	2.2	2.5	3.0	2.1	2.4	2.8	2.0	3.2	4.1	2.6
2002.....	2.3	2.9	1.9	2.5	3.2	1.9	2.3	2.8	1.9	2.2	2.6	1.8	3.2	4.1	2.5
2001.....	2.0	2.4	1.7	2.0	2.4	1.5	2.0	2.4	1.7	1.9	2.2	1.6	2.9	3.4	2.5
2000.....	2.0	2.4	1.7	2.1	2.5	1.7	2.0	2.4	1.7	1.8	2.2	1.6	3.0	3.8	2.5
1999.....	2.6	3.1	2.3	2.6	2.8	2.5	2.6	3.1	2.3	2.4	2.8	2.0	4.7	5.6	4.1
<b>Malignant neoplasms (C00-C97)</b>															
2007.....	178.4	217.5	151.3	116.2	141.4	98.6	183.1	223.2	155.3	182.3	220.8	155.3	220.0	288.6	178.4
2006.....	180.7	220.1	153.6	118.0	143.4	100.4	185.3	225.6	157.6	184.6	223.4	157.6	221.7	290.8	179.5
2005.....	183.8	225.1	155.6	122.8	152.7	101.9	188.2	230.2	159.4	187.0	227.3	159.1	226.8	299.0	183.0
2004.....	185.8	227.7	157.4	121.9	151.2	101.4	190.1	232.7	161.2	188.6	229.2	160.9	231.1	306.4	185.7
2003.....	190.1	233.3	160.9	126.6	156.5	105.9	194.1	238.1	164.4	192.4	234.6	163.8	237.3	314.2	190.8
2002.....	193.5	238.9	163.1	128.4	161.4	106.1	197.4	243.4	166.6	195.6	239.6	165.9	242.5	324.4	193.4
2001.....	196.0	243.7	164.7	132.3	168.2	108.6	199.7	247.9	168.0	197.4	243.1	167.2	246.7	335.3	194.4
2000.....	199.6	248.9	167.6	134.9	171.7	110.8	203.1	253.0	170.7	200.6	247.7	170.0	252.0	344.8	196.9
1999.....	200.8	251.9	167.6	134.8	170.6	110.9	204.2	256.0	170.6	201.2	250.4	169.4	255.9	350.3	199.8
<b>Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)</b>															
2007.....	2.5	3.9	1.4	1.5	2.3	0.8	2.6	4.0	1.5	2.5	3.8	1.4	3.4	5.9	1.6
2006.....	2.5	3.8	1.4	1.5	2.4	0.7	2.5	3.9	1.4	2.5	3.7	1.4	3.3	5.8	1.5
2005.....	2.5	3.9	1.4	1.4	2.3	0.7	2.6	4.0	1.5	2.5	3.7	1.5	3.6	6.6	1.4
2004.....	2.6	4.0	1.5	1.6	2.6	0.8	2.7	4.1	1.5	2.5	3.8	1.5	3.9	7.0	1.7
2003.....	2.6	4.0	1.5	1.7	2.7	0.9	2.7	4.2	1.5	2.6	3.9	1.5	3.9	7.0	1.7
2002.....	2.7	4.1	1.5	1.5	2.6	0.7	2.8	4.1	1.6	2.6	3.9	1.5	3.7	6.4	1.7
2001.....	2.7	4.1	1.6	1.8	2.9	1.0	2.8	4.2	1.6	2.6	3.9	1.6	3.9	6.8	1.8
2000.....	2.7	4.1	1.6	1.8	3.1	0.8	2.8	4.1	1.6	2.6	3.8	1.6	4.2	7.4	1.8
1999.....	2.7	4.1	1.6	1.9	3.0	1.0	2.8	4.2	1.6	2.6	3.9	1.6	4.4	7.6	2.1

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Malignant neoplasm of esophagus (C15)</b>															
2007.....	4.3	7.7	1.6	2.0	3.7	0.7	4.5	8.0	1.6	4.5	8.2	1.6	4.8	8.6	2.3
2006.....	4.4	7.8	1.7	2.3	4.0	0.9	4.6	8.1	1.7	4.6	8.3	1.7	4.9	8.5	2.4
2005.....	4.4	7.9	1.7	2.3	4.2	0.8	4.6	8.2	1.7	4.6	8.3	1.6	5.1	9.0	2.4
2004.....	4.3	7.7	1.7	2.2	4.0	0.9	4.5	8.0	1.8	4.5	8.0	1.7	5.5	9.1	2.9
2003.....	4.4	7.7	1.7	2.2	3.8	1.0	4.5	8.0	1.8	4.5	8.0	1.7	5.8	9.9	2.9
2002.....	4.4	7.8	1.7	2.3	4.2	0.9	4.5	8.0	1.8	4.5	8.0	1.7	6.0	10.3	3.0
2001.....	4.4	7.8	1.8	2.4	4.4	1.0	4.5	8.0	1.8	4.4	7.9	1.7	6.5	11.5	3.0
2000.....	4.4	7.7	1.8	2.5	4.6	1.1	4.5	7.9	1.9	4.4	7.8	1.7	6.6	11.2	3.3
1999.....	4.4	7.7	1.8	2.5	4.7	1.0	4.5	7.9	1.8	4.3	7.6	1.7	6.6	11.6	3.1
<b>Malignant neoplasm of stomach (C16)</b>															
2007.....	3.6	5.0	2.6	5.5	7.4	4.1	3.4	4.8	2.4	2.9	4.1	2.0	7.0	10.5	4.7
2006.....	3.7	5.0	2.7	5.5	6.9	4.5	3.5	4.9	2.5	3.0	4.2	2.0	6.9	10.0	5.0
2005.....	3.8	5.2	2.7	5.9	8.0	4.3	3.6	5.0	2.6	3.0	4.2	2.2	7.3	10.8	5.0
2004.....	4.0	5.5	2.8	6.6	8.7	5.0	3.8	5.3	2.7	3.2	4.5	2.2	7.6	11.2	5.3
2003.....	4.1	5.7	3.0	6.5	8.5	5.0	3.9	5.5	2.8	3.4	4.7	2.4	7.9	11.5	5.5
2002.....	4.2	5.9	3.0	6.6	8.8	5.0	4.1	5.7	2.8	3.5	4.9	2.3	8.5	12.1	6.2
2001.....	4.4	6.1	3.1	6.9	9.2	5.3	4.2	5.9	2.9	3.6	5.0	2.5	8.4	12.1	5.9
2000.....	4.6	6.4	3.3	7.0	9.8	5.1	4.4	6.2	3.1	3.7	5.3	2.6	9.1	12.9	6.6
1999.....	4.6	6.6	3.3	7.1	9.3	5.3	4.5	6.4	3.1	3.8	5.5	2.6	9.3	13.7	6.5
<b>Malignant neoplasms of colon, rectum and anus (C18-C21)</b>															
2007.....	16.9	20.1	14.4	12.0	14.5	10.1	17.2	20.5	14.7	16.7	19.9	14.2	24.0	29.6	20.2
2006.....	17.2	20.5	14.7	12.6	15.1	10.7	17.5	20.9	15.0	17.0	20.2	14.5	24.7	31.2	20.6
2005.....	17.5	20.9	14.8	12.4	15.2	10.4	17.8	21.3	15.1	17.2	20.7	14.5	25.2	30.7	21.6
2004.....	18.0	21.5	15.3	12.6	15.6	10.2	18.3	21.9	15.7	17.9	21.4	15.2	25.1	30.6	21.6
2003.....	19.1	22.9	16.2	13.4	16.8	10.8	19.4	23.2	16.5	18.8	22.6	16.0	26.8	32.5	23.1
2002.....	19.7	23.7	16.7	13.7	16.9	11.2	20.0	24.1	17.0	19.5	23.5	16.5	27.2	33.6	23.1
2001.....	20.1	24.2	17.2	14.1	17.5	11.8	20.5	24.6	17.5	19.9	24.0	16.9	28.0	33.6	24.5
2000.....	20.8	25.1	17.7	14.1	17.8	11.5	21.1	25.5	18.0	20.5	24.8	17.4	28.7	35.5	24.3
1999.....	20.9	25.3	17.8	14.3	17.7	11.9	21.2	25.7	18.1	20.7	25.2	17.5	28.7	34.5	25.0



Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of liver and intrahepatic bile ducts (C22)															
2007.....	5.4	7.9	3.2	7.8	11.1	5.1	5.2	7.7	3.1	4.7	6.9	2.9	7.2	11.8	3.9
2006.....	5.3	7.7	3.3	7.9	11.2	5.1	5.1	7.4	3.1	4.6	6.6	2.9	7.1	11.3	3.9
2005.....	5.2	7.7	3.2	8.3	11.7	5.4	5.0	7.4	3.0	4.5	6.6	2.8	7.1	11.4	3.8
2004.....	5.1	7.4	3.2	7.6	10.9	4.9	4.9	7.1	3.1	4.4	6.3	2.8	7.0	10.9	4.2
2003.....	5.0	7.4	3.1	7.8	10.8	5.2	4.8	7.1	3.0	4.3	6.3	2.7	6.7	10.3	4.0
2002.....	4.9	7.1	3.0	7.5	10.7	4.9	4.7	6.9	2.9	4.2	6.2	2.7	6.4	10.0	3.7
2001.....	4.7	6.9	3.0	7.3	10.4	4.9	4.6	6.7	2.9	4.1	6.0	2.6	6.3	9.2	4.1
2000.....	4.7	6.8	3.0	7.5	10.7	4.9	4.5	6.6	2.9	4.0	5.8	2.6	6.2	9.7	3.6
1999.....	4.5	6.5	3.0	7.2	10.0	5.0	4.4	6.3	2.8	3.9	5.6	2.6	6.0	8.8	3.9
Malignant neoplasm of pancreas (C25)															
2007.....	10.8	12.5	9.4	7.9	8.4	7.4	11.0	12.8	9.6	10.8	12.7	9.3	14.3	15.9	12.9
2006.....	10.8	12.3	9.5	8.0	8.9	7.3	11.0	12.6	9.6	10.8	12.6	9.4	13.8	15.1	12.7
2005.....	10.8	12.3	9.5	8.6	9.4	7.8	10.9	12.5	9.6	10.7	12.4	9.4	13.9	15.6	12.5
2004.....	10.6	12.3	9.3	8.2	9.1	7.4	10.8	12.5	9.4	10.6	12.3	9.2	13.8	15.7	12.2
2003.....	10.5	12.0	9.3	8.2	9.2	7.4	10.6	12.2	9.4	10.4	12.0	9.1	13.9	15.6	12.6
2002.....	10.5	12.1	9.2	7.9	8.4	7.4	10.7	12.3	9.3	10.5	12.2	9.0	13.8	15.5	12.5
2001.....	10.5	12.1	9.3	8.4	9.0	7.8	10.7	12.3	9.4	10.5	12.2	9.2	13.8	15.1	12.6
2000.....	10.6	12.2	9.3	8.4	9.4	7.5	10.7	12.3	9.4	10.5	12.2	9.1	14.3	16.0	12.9
1999.....	10.6	12.3	9.3	7.9	8.8	7.2	10.8	12.4	9.4	10.5	12.2	9.2	14.3	16.7	12.5
Malignant neoplasm of larynx (C32)															
2007.....	1.1	2.1	0.4	0.7	1.5	*	1.2	2.1	0.4	1.1	1.9	0.4	2.2	4.4	0.6
2006.....	1.2	2.2	0.5	0.8	1.8	*	1.3	2.2	0.5	1.2	2.1	0.5	2.2	4.3	0.7
2005.....	1.2	2.2	0.5	0.9	1.9	*	1.3	2.3	0.5	1.2	2.1	0.5	2.3	4.6	0.8
2004.....	1.2	2.2	0.4	1.0	2.1	*	1.2	2.2	0.5	1.2	2.1	0.4	2.3	4.6	0.7
2003.....	1.3	2.4	0.5	0.9	1.9	*	1.3	2.4	0.5	1.2	2.2	0.5	2.5	5.2	0.7
2002.....	1.3	2.3	0.5	0.9	1.7	0.3	1.3	2.4	0.5	1.2	2.2	0.5	2.5	4.9	0.9
2001.....	1.3	2.5	0.5	1.0	1.9	0.2	1.4	2.5	0.5	1.2	2.3	0.5	2.6	5.0	0.9
2000.....	1.4	2.5	0.5	0.9	1.9	*	1.4	2.6	0.5	1.3	2.3	0.5	2.8	5.5	1.0
1999.....	1.4	2.5	0.5	1.2	2.5	0.3	1.4	2.5	0.5	1.3	2.3	0.5	2.6	5.2	0.8

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Malignant neoplasms of trachea, bronchus and lung (C33-C34)</b>															
2007.....	50.6	65.1	40.0	20.9	29.6	14.4	53.0	67.8	42.0	53.9	67.7	43.5	56.8	84.0	39.0
2006.....	51.5	67.0	40.0	20.7	30.3	13.6	53.9	69.8	42.1	54.7	69.7	43.5	57.9	85.4	39.8
2005.....	52.6	69.0	40.5	22.4	33.3	14.4	54.8	71.6	42.5	55.5	71.4	43.7	59.6	88.1	40.9
2004.....	53.2	70.1	40.9	22.4	33.4	14.3	55.4	72.6	42.8	56.0	72.1	44.1	60.9	91.7	40.6
2003.....	54.1	71.7	41.3	23.2	34.5	14.8	56.2	74.1	43.1	56.7	73.6	44.3	62.0	94.2	40.9
2002.....	54.9	73.2	41.6	23.7	36.2	14.6	56.9	75.6	43.3	57.5	75.0	44.6	62.9	96.6	40.8
2001.....	55.3	75.2	41.0	23.8	36.8	14.5	57.2	77.4	42.7	57.7	76.6	44.0	63.4	100.5	39.4
2000.....	56.1	76.7	41.3	24.8	39.0	14.7	57.9	78.9	42.9	58.2	77.9	44.1	64.9	102.6	40.4
1999.....	55.5	76.9	40.2	25.0	39.6	14.6	57.2	78.8	41.7	57.2	77.6	42.7	65.7	104.3	40.6
<b>Malignant melanoma of skin (C43)</b>															
2007.....	2.7	4.0	1.7	0.8	1.0	0.6	2.8	4.2	1.8	3.3	4.9	2.1	0.5	0.5	0.4
2006.....	2.7	4.0	1.7	0.8	1.0	0.7	2.9	4.3	1.8	3.3	4.9	2.1	0.4	0.6	0.3
2005.....	2.7	4.0	1.8	0.7	1.0	0.5	2.9	4.2	1.9	3.3	4.8	2.2	0.5	0.5	0.4
2004.....	2.7	3.9	1.7	0.6	0.7	0.5	2.8	4.2	1.8	3.3	4.8	2.1	0.4	0.5	0.4
2003.....	2.7	3.9	1.8	0.8	1.0	0.6	2.8	4.1	1.8	3.2	4.7	2.1	0.5	0.4	0.4
2002.....	2.6	3.8	1.7	0.8	0.8	0.7	2.7	4.0	1.8	3.1	4.5	2.1	0.4	0.5	0.3
2001.....	2.7	3.9	1.8	0.8	0.9	0.7	2.8	4.1	1.8	3.2	4.7	2.1	0.4	0.3	0.5
2000.....	2.7	3.8	1.8	0.8	1.1	0.5	2.8	4.0	1.9	3.2	4.5	2.2	0.6	0.6	0.5
1999.....	2.6	3.8	1.7	0.8	1.0	0.6	2.8	4.0	1.8	3.1	4.5	2.1	0.4	0.5	0.3
<b>Malignant neoplasm of breast (C50)</b>															
2007.....	12.9	0.3	22.9	7.9	*	14.5	13.2	0.3	23.6	12.8	0.3	23.0	19.0	0.4	32.2
2006.....	13.2	0.3	23.5	8.2	*	15.0	13.5	0.3	24.1	13.1	0.3	23.5	19.0	0.5	32.2
2005.....	13.5	0.3	24.1	8.2	*	15.0	13.9	0.3	24.8	13.5	0.3	24.0	19.7	0.5	33.5
2004.....	13.8	0.3	24.4	8.7	*	15.6	14.1	0.3	25.1	13.7	0.3	24.5	19.4	0.5	32.9
2003.....	14.2	0.3	25.3	8.9	*	16.1	14.6	0.3	25.9	14.2	0.3	25.2	20.5	0.5	34.6
2002.....	14.5	0.3	25.6	8.6	*	15.5	14.8	0.3	26.2	14.4	0.3	25.6	20.5	0.6	34.6
2001.....	14.8	0.3	26.0	9.1	*	16.3	15.1	0.4	26.6	14.7	0.3	26.0	20.8	0.6	35.1
2000.....	15.3	0.4	26.8	9.5	*	16.9	15.6	0.4	27.3	15.2	0.4	26.8	20.9	0.6	35.1
1999.....	15.2	0.3	26.6	9.2	*	16.4	15.5	0.3	27.2	15.1	0.3	26.6	21.2	0.7	35.7

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasm of cervix uteri (C53)															
2007.....	1.3	...	2.4	1.5	...	3.0	1.2	...	2.4	1.1	...	2.1	2.5	...	4.4
2006.....	1.3	...	2.4	1.6	...	3.1	1.3	...	2.4	1.1	...	2.1	2.5	...	4.4
2005.....	1.3	...	2.4	1.6	...	3.1	1.3	...	2.4	1.1	...	2.1	2.6	...	4.5
2004.....	1.3	...	2.4	1.7	...	3.1	1.3	...	2.4	1.1	...	2.1	2.7	...	4.6
2003.....	1.3	...	2.5	1.6	...	3.0	1.3	...	2.4	1.1	...	2.2	2.7	...	4.8
2002.....	1.3	...	2.6	1.7	...	3.1	1.3	...	2.5	1.1	...	2.2	2.9	...	5.0
2001.....	1.4	...	2.7	1.8	...	3.4	1.4	...	2.6	1.2	...	2.3	2.8	...	4.9
2000.....	1.5	...	2.8	2.2	...	4.0	1.5	...	2.7	1.2	...	2.3	3.2	...	5.5
1999.....	1.5	...	2.8	1.9	...	3.6	1.5	...	2.8	1.3	...	2.5	3.2	...	5.4
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)															
2007.....	2.3	...	4.2	1.6	...	2.9	2.4	...	4.3	2.2	...	3.9	4.6	...	7.7
2006.....	2.4	...	4.2	1.7	...	3.1	2.4	...	4.3	2.2	...	4.0	4.4	...	7.4
2005.....	2.3	...	4.1	1.8	...	3.2	2.3	...	4.1	2.2	...	3.9	4.3	...	7.2
2004.....	2.3	...	4.1	1.6	...	2.9	2.4	...	4.2	2.2	...	3.9	4.2	...	7.0
2003.....	2.3	...	4.1	1.7	...	3.0	2.4	...	4.2	2.2	...	3.9	4.3	...	7.1
2002.....	2.4	...	4.2	1.9	...	3.3	2.4	...	4.2	2.2	...	3.9	4.5	...	7.4
2001.....	2.4	...	4.2	1.9	...	3.3	2.4	...	4.2	2.3	...	4.0	4.3	...	7.0
2000.....	2.4	...	4.1	1.8	...	3.1	2.4	...	4.2	2.2	...	3.9	4.5	...	7.4
1999.....	2.4	...	4.1	1.8	...	3.1	2.4	...	4.1	2.3	...	3.9	4.2	...	6.9
Malignant neoplasm of ovary (C56)															
2007.....	4.6	...	8.2	3.2	...	5.7	4.7	...	8.4	4.9	...	8.9	4.0	...	6.7
2006.....	4.8	...	8.5	3.4	...	6.0	4.9	...	8.7	5.1	...	9.2	4.1	...	6.9
2005.....	4.8	...	8.6	3.3	...	6.0	4.9	...	8.8	5.1	...	9.2	4.4	...	7.4
2004.....	4.9	...	8.8	3.3	...	5.9	5.1	...	9.0	5.2	...	9.3	4.6	...	7.8
2003.....	5.0	...	8.8	3.3	...	5.9	5.1	...	9.1	5.3	...	9.4	4.5	...	7.5
2002.....	5.1	...	9.0	3.3	...	6.0	5.2	...	9.2	5.4	...	9.6	4.5	...	7.5
2001.....	5.1	...	9.0	3.5	...	6.1	5.2	...	9.2	5.4	...	9.5	4.6	...	7.6
2000.....	5.1	...	8.9	3.7	...	6.4	5.2	...	9.1	5.3	...	9.5	4.4	...	7.3
1999.....	5.0	...	8.8	3.1	...	5.5	5.1	...	9.0	5.2	...	9.3	4.4	...	7.4

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasm of prostate (C61)															
2007.....	9.2	23.5	...	7.2	17.7	...	9.3	23.9	...	8.6	21.9	...	18.9	52.6	...
2006.....	9.2	23.5	...	7.3	18.1	...	9.3	23.8	...	8.5	21.9	...	18.5	51.3	...
2005.....	9.5	24.5	...	7.4	18.5	...	9.6	24.9	...	8.9	22.8	...	19.4	54.1	...
2004.....	9.8	25.4	...	7.6	19.1	...	9.9	25.7	...	9.1	23.6	...	20.0	56.4	...
2003.....	10.1	26.5	...	7.9	20.2	...	10.2	26.8	...	9.4	24.6	...	20.7	58.2	...
2002.....	10.6	27.9	...	8.4	21.6	...	10.7	28.2	...	9.8	25.8	...	22.2	62.8	...
2001.....	10.9	29.1	...	9.1	23.5	...	11.0	29.3	...	10.0	26.7	...	23.3	66.8	...
2000.....	11.3	30.4	...	8.3	21.6	...	11.4	30.7	...	10.3	28.0	...	24.1	68.8	...
1999.....	11.6	31.3	...	8.8	23.0	...	11.7	31.7	...	10.7	28.9	...	24.5	69.7	...
Malignant neoplasms of kidney and renal pelvis (C64-C65)															
2007.....	4.0	5.8	2.6	3.7	5.5	2.2	4.0	5.8	2.7	4.1	5.9	2.7	4.1	6.1	2.8
2006.....	4.0	5.7	2.6	3.2	4.7	2.1	4.1	5.8	2.7	4.1	5.9	2.7	3.9	6.0	2.5
2005.....	4.1	5.9	2.7	3.7	5.1	2.6	4.1	5.9	2.7	4.2	6.0	2.8	4.1	6.1	2.6
2004.....	4.1	5.9	2.7	3.7	5.5	2.2	4.1	5.9	2.8	4.2	6.1	2.8	4.1	5.9	2.8
2003.....	4.2	6.0	2.7	3.7	5.0	2.6	4.2	6.1	2.7	4.3	6.2	2.8	4.1	6.1	2.8
2002.....	4.2	6.1	2.8	3.6	5.3	2.3	4.2	6.2	2.8	4.4	6.3	2.9	4.1	5.9	2.9
2001.....	4.3	6.2	2.8	3.7	5.6	2.3	4.3	6.2	2.8	4.4	6.3	2.9	4.3	6.5	2.9
2000.....	4.2	6.2	2.8	3.5	5.3	2.2	4.3	6.2	2.8	4.3	6.3	2.9	4.2	6.3	2.8
1999.....	4.1	5.9	2.7	3.6	5.1	2.4	4.1	5.9	2.7	4.1	6.0	2.7	4.2	6.2	2.8
Malignant neoplasm of bladder (C67)															
2007.....	4.4	7.6	2.2	2.3	4.0	1.1	4.5	7.8	2.3	4.7	8.2	2.3	3.8	5.5	2.8
2006.....	4.3	7.5	2.2	2.2	3.5	1.3	4.5	7.8	2.2	4.7	8.2	2.3	3.8	5.6	2.8
2005.....	4.3	7.5	2.2	2.4	4.2	1.1	4.5	7.7	2.3	4.7	8.2	2.3	3.7	5.7	2.6
2004.....	4.4	7.6	2.2	2.2	3.5	1.4	4.5	7.8	2.3	4.7	8.2	2.3	3.6	5.0	2.8
2003.....	4.3	7.3	2.2	2.4	4.0	1.4	4.4	7.5	2.3	4.5	7.9	2.3	3.8	5.5	2.7
2002.....	4.4	7.5	2.4	2.5	4.2	1.5	4.5	7.7	2.4	4.6	8.0	2.4	4.0	5.5	3.1
2001.....	4.3	7.5	2.3	2.4	4.0	1.4	4.4	7.7	2.3	4.6	8.1	2.3	3.8	5.1	2.9
2000.....	4.3	7.6	2.3	2.3	4.1	1.2	4.4	7.8	2.3	4.6	8.1	2.3	3.8	5.6	2.7
1999.....	4.4	7.6	2.3	2.3	4.1	1.1	4.4	7.8	2.3	4.6	8.1	2.3	4.0	5.8	3.0

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)</b>															
2007.....	4.2	5.1	3.5	2.8	3.1	2.5	4.4	5.3	3.6	4.8	5.8	3.9	2.4	2.9	2.1
2006.....	4.2	5.1	3.4	2.6	2.9	2.2	4.3	5.3	3.5	4.7	5.8	3.8	2.4	3.0	2.0
2005.....	4.3	5.3	3.5	2.9	3.5	2.4	4.5	5.5	3.6	4.9	5.9	4.0	2.6	3.3	2.0
2004.....	4.3	5.2	3.5	2.7	3.1	2.3	4.4	5.4	3.6	4.8	5.8	3.9	2.5	3.1	2.0
2003.....	4.4	5.3	3.6	2.8	3.3	2.4	4.5	5.5	3.7	4.9	5.9	4.1	2.7	3.4	2.2
2002.....	4.4	5.5	3.6	2.9	3.4	2.4	4.6	5.6	3.7	4.9	6.1	4.0	2.6	3.2	2.2
2001.....	4.5	5.5	3.6	2.9	3.4	2.4	4.6	5.7	3.7	4.9	6.1	4.0	2.6	3.2	2.2
2000.....	4.5	5.6	3.7	3.0	3.7	2.6	4.6	5.7	3.8	5.0	6.2	4.0	2.7	3.3	2.3
1999.....	4.6	5.6	3.8	2.9	3.2	2.6	4.8	5.8	3.9	5.1	6.3	4.2	2.8	3.3	2.3
<b>Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)</b>															
2007.....	17.6	22.8	13.7	13.0	15.5	11.1	17.8	23.3	13.8	18.1	23.7	13.9	17.9	22.5	14.9
2006.....	17.9	23.0	14.1	12.9	15.6	10.9	18.2	23.5	14.3	18.5	23.9	14.5	18.4	24.1	14.5
2005.....	18.2	23.5	14.3	13.7	16.9	11.3	18.4	23.9	14.4	18.7	24.4	14.6	17.9	23.0	14.5
2004.....	18.4	23.6	14.6	13.2	16.2	10.7	18.7	24.0	14.8	18.9	24.4	14.9	19.2	23.9	16.0
2003.....	19.1	24.5	15.1	13.8	16.3	11.9	19.3	24.9	15.2	19.6	25.5	15.3	19.2	23.4	16.4
2002.....	19.5	25.1	15.5	14.5	17.4	12.4	19.8	25.5	15.7	20.1	26.0	15.8	19.6	24.8	16.2
2001.....	20.0	25.5	16.0	14.9	19.3	11.6	20.2	25.8	16.2	20.4	26.1	16.3	20.3	25.7	16.8
2000.....	20.4	26.0	16.5	15.5	19.3	12.7	20.6	26.3	16.6	20.9	26.7	16.8	20.2	25.4	16.9
1999.....	20.6	26.1	16.7	15.4	17.6	13.5	20.8	26.5	16.8	21.0	26.8	16.9	20.6	26.0	17.0
<b>Hodgkin's disease (C81)</b>															
2007.....	0.4	0.5	0.3	0.4	0.5	0.4	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.5	0.3
2006.....	0.4	0.6	0.3	0.4	0.4	0.3	0.5	0.6	0.3	0.5	0.6	0.4	0.4	0.6	0.3
2005.....	0.4	0.5	0.4	0.5	0.7	0.4	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3
2004.....	0.4	0.5	0.3	0.4	0.5	0.3	0.4	0.6	0.4	0.4	0.6	0.4	0.4	0.5	0.4
2003.....	0.5	0.5	0.4	0.4	0.5	0.3	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.5	0.3
2002.....	0.5	0.6	0.4	0.4	0.5	0.4	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.6	0.3
2001.....	0.5	0.6	0.4	0.5	0.7	0.3	0.5	0.6	0.4	0.5	0.6	0.4	0.4	0.5	0.4
2000.....	0.5	0.6	0.4	0.5	0.7	0.3	0.5	0.5	0.4	0.5	0.6	0.4	0.3	0.4	0.3
1999.....	0.5	0.6	0.4	0.6	0.8	0.5	0.5	0.6	0.4	0.5	0.7	0.4	0.4	0.6	0.3

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Non-Hodgkin's lymphoma (C82-C85)</b>															
2007.....	6.5	8.3	5.2	5.1	6.2	4.4	6.6	8.4	5.2	6.9	8.7	5.5	4.6	5.9	3.7
2006.....	6.7	8.4	5.4	5.0	6.1	4.2	6.8	8.5	5.5	7.1	8.8	5.7	5.0	6.4	4.0
2005.....	6.9	8.7	5.5	5.3	6.3	4.6	7.0	8.9	5.6	7.3	9.3	5.8	4.8	6.1	3.8
2004.....	7.0	8.8	5.7	5.2	6.4	4.2	7.1	9.0	5.8	7.5	9.3	6.1	5.0	6.2	4.1
2003.....	7.3	9.3	5.9	5.4	6.4	4.7	7.4	9.4	6.0	7.8	9.9	6.2	5.0	5.9	4.3
2002.....	7.6	9.6	6.2	5.8	6.6	5.2	7.7	9.7	6.2	8.0	10.1	6.5	5.4	6.9	4.4
2001.....	7.9	9.9	6.4	5.9	7.5	4.6	8.0	10.0	6.5	8.3	10.4	6.8	5.7	7.4	4.5
2000.....	8.2	10.2	6.7	6.3	7.7	5.3	8.3	10.3	6.8	8.7	10.8	7.1	5.3	6.7	4.3
1999.....	8.3	10.3	6.9	6.2	7.1	5.5	8.4	10.5	6.9	8.8	10.8	7.3	5.7	7.6	4.4
<b>Leukemia (C91-C95)</b>															
2007.....	7.0	9.4	5.3	4.5	5.5	3.8	7.1	9.6	5.3	7.3	10.0	5.4	6.2	8.0	5.0
2006.....	7.1	9.5	5.4	4.7	5.7	3.9	7.3	9.8	5.5	7.5	10.1	5.7	6.3	8.8	4.7
2005.....	7.1	9.6	5.4	4.9	6.5	3.8	7.2	9.8	5.5	7.5	10.1	5.6	6.0	8.0	4.8
2004.....	7.2	9.6	5.5	4.7	5.9	3.7	7.3	9.8	5.6	7.5	10.1	5.7	6.7	8.7	5.4
2003.....	7.4	9.9	5.6	5.0	6.2	4.2	7.5	10.0	5.7	7.7	10.4	5.8	6.6	8.4	5.3
2002.....	7.5	10.0	5.7	4.9	6.0	4.1	7.6	10.2	5.7	7.8	10.5	5.9	6.7	8.8	5.4
2001.....	7.6	10.1	5.9	5.2	6.7	4.2	7.7	10.2	5.9	7.9	10.5	6.1	6.7	8.8	5.4
2000.....	7.7	10.3	5.9	5.4	6.9	4.2	7.8	10.4	6.0	8.0	10.7	6.1	6.8	9.1	5.3
1999.....	7.6	10.2	6.0	5.2	6.3	4.5	7.7	10.3	6.0	7.9	10.6	6.1	6.8	8.8	5.4
<b>Multiple myeloma and immunoproliferative neoplasms (C88,C90)</b>															
2007.....	3.6	4.6	2.9	2.9	3.4	2.5	3.7	4.7	2.9	3.4	4.4	2.6	6.7	8.0	5.9
2006.....	3.6	4.5	2.9	2.9	3.3	2.5	3.7	4.6	3.0	3.4	4.3	2.7	6.6	8.3	5.5
2005.....	3.7	4.7	3.0	2.9	3.4	2.5	3.7	4.7	3.0	3.5	4.5	2.8	6.6	8.2	5.6
2004.....	3.7	4.6	3.1	2.9	3.4	2.5	3.7	4.6	3.1	3.5	4.4	2.8	7.0	8.4	6.1
2003.....	3.8	4.8	3.2	3.0	3.3	2.8	3.9	4.9	3.2	3.6	4.6	2.9	7.3	8.5	6.4
2002.....	4.0	5.0	3.3	3.4	4.3	2.7	4.0	5.0	3.3	3.7	4.8	3.0	7.0	8.4	6.1
2001.....	4.0	4.9	3.3	3.3	4.3	2.6	4.0	4.9	3.4	3.7	4.6	3.1	7.4	8.9	6.4
2000.....	4.0	4.9	3.4	3.3	3.8	2.9	4.0	4.9	3.4	3.7	4.6	3.1	7.7	9.1	6.9
1999.....	4.0	4.9	3.4	3.2	3.4	3.0	4.1	5.0	3.4	3.7	4.7	3.1	7.6	9.0	6.8

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year

	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)
2007..... 0.0 0.0 0.0 \* \* \* 0.0 0.0 0.0 0.0 0.0 0.0 \* \* \*
2006..... 0.0 0.0 0.0 \* \* \* 0.0 0.0 0.0 0.0 0.0 0.0 \* \* \*
2005..... 0.0 0.0 0.0 \* \* \* 0.0 0.0 \* 0.0 0.0 0.0 \* \* \*
2004..... 0.0 0.0 0.0 \* \* \* 0.0 0.0 0.0 0.0 0.0 0.0 \* \* \*
2003..... 0.0 0.0 0.0 \* \* \* 0.0 0.0 0.0 0.0 0.0 0.0 \* \* \*
2002..... 0.0 0.0 0.0 \* \* \* 0.0 0.0 0.0 0.0 0.0 0.0 \* \* \*
2001..... 0.0 0.0 0.0 \* \* \* 0.0 0.0 0.0 0.0 0.0 0.0 \* \* \*
2000..... 0.0 0.0 0.0 \* \* \* 0.0 0.1 0.0 0.0 0.0 0.0 \* \* \*
1999..... 0.0 0.0 0.0 \* \* \* 0.0 0.0 0.0 0.0 0.0 0.0 \* \* \*

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)
2007..... 20.0 24.3 16.8 13.9 15.9 12.3 20.4 24.9 17.1 20.5 25.1 17.1 22.5 27.5 19.1
2006..... 20.3 24.5 17.3 14.6 16.7 13.1 20.8 25.0 17.6 20.8 25.2 17.5 23.1 28.0 19.9
2005..... 20.6 24.8 17.5 14.8 17.5 12.8 21.0 25.3 17.9 21.1 25.3 17.9 23.6 28.9 20.0
2004..... 20.8 25.0 17.6 14.5 16.6 12.9 21.2 25.6 17.9 21.2 25.6 17.9 24.2 30.2 20.2
2003..... 21.4 25.7 18.2 15.8 18.5 13.8 21.7 26.1 18.4 21.7 26.1 18.4 24.7 30.4 20.8
2002..... 21.8 26.5 18.5 16.2 19.1 14.0 22.2 26.9 18.7 22.1 26.8 18.6 26.1 31.8 22.3
2001..... 22.2 26.7 18.9 16.6 19.3 14.6 22.5 27.2 19.1 22.3 26.9 19.0 26.6 33.2 22.2
2000..... 22.8 27.5 19.4 17.3 20.2 15.4 23.0 27.8 19.6 22.9 27.7 19.5 26.8 33.4 22.4
1999..... 24.2 29.3 20.6 17.9 20.8 15.8 24.5 29.8 20.8 24.3 29.5 20.6 28.9 35.7 24.2

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)
2007..... 4.5 5.6 3.7 3.0 3.1 2.8 4.6 5.8 3.8 4.7 5.9 3.8 4.2 5.3 3.5
2006..... 4.5 5.6 3.9 3.2 3.7 2.8 4.6 5.7 3.9 4.7 5.8 3.9 4.5 5.4 3.9
2005..... 4.5 5.5 3.8 2.6 3.2 2.2 4.6 5.7 3.9 4.7 5.9 3.9 4.2 4.6 4.0
2004..... 4.5 5.7 3.8 3.1 3.6 2.7 4.6 5.8 3.9 4.7 5.9 3.9 4.5 5.3 4.0
2003..... 4.6 5.7 3.9 3.3 3.7 2.9 4.7 5.8 4.0 4.8 6.0 4.0 4.3 5.1 3.8
2002..... 4.6 5.7 3.9 3.2 3.7 2.8 4.7 5.8 4.0 4.8 5.9 4.0 4.5 5.2 4.1
2001..... 4.9 6.0 4.1 3.3 4.0 2.8 4.9 6.1 4.2 5.0 6.3 4.2 4.7 5.7 4.1
2000..... 4.9 6.0 4.2 3.5 3.8 3.3 5.0 6.1 4.2 5.0 6.2 4.3 4.6 5.5 4.1
1999..... 4.9 6.1 4.1 3.3 3.4 3.2 4.9 6.2 4.1 5.0 6.3 4.1 4.9 5.5 4.5

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year

	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Anemias (D50-D64)

2007.....	1.5	1.5	1.5	1.0	1.1	0.8	1.6	1.6	1.5	1.3	1.3	1.3	3.1	3.2	2.9
2006.....	1.3	1.3	1.3	0.9	0.8	1.0	1.3	1.3	1.3	1.1	1.1	1.1	2.9	2.8	2.9
2005.....	1.5	1.5	1.5	0.9	0.9	0.9	1.6	1.5	1.6	1.3	1.2	1.3	3.4	3.5	3.3
2004.....	1.5	1.5	1.5	1.0	0.9	1.0	1.6	1.6	1.5	1.3	1.3	1.3	3.4	3.6	3.2
2003.....	1.6	1.6	1.5	1.1	1.2	1.1	1.6	1.6	1.5	1.3	1.3	1.3	3.6	4.0	3.2
2002.....	1.6	1.6	1.6	1.3	1.6	1.1	1.6	1.6	1.6	1.3	1.3	1.4	3.7	4.0	3.5
2001.....	1.6	1.6	1.6	1.2	1.1	1.2	1.7	1.6	1.7	1.4	1.4	1.4	3.6	3.7	3.4
2000.....	1.6	1.6	1.6	0.9	0.9	1.0	1.7	1.6	1.7	1.4	1.4	1.4	3.5	3.5	3.5
1999.....	1.6	1.7	1.6	1.2	1.4	1.1	1.7	1.7	1.6	1.4	1.4	1.4	3.7	3.8	3.5

Diabetes mellitus (E10-E14)

2007.....	22.5	26.4	19.5	28.9	31.9	26.2	22.1	26.0	19.1	19.8	23.9	16.4	43.7	46.9	41.0
2006.....	23.3	27.4	20.1	29.9	33.7	26.8	22.9	27.0	19.7	20.4	24.7	17.0	45.9	50.6	42.4
2005.....	24.6	28.4	21.6	33.6	37.3	30.5	24.0	27.9	21.1	21.5	25.6	18.4	47.7	51.7	44.6
2004.....	24.5	28.2	21.7	32.1	34.3	30.2	24.0	27.7	21.1	21.5	25.5	18.4	48.7	52.1	46.0
2003.....	25.3	28.9	22.5	35.0	38.3	32.4	24.7	28.3	21.9	22.1	26.1	19.0	50.0	51.7	48.3
2002.....	25.4	28.6	23.0	35.6	38.1	33.6	24.8	28.0	22.3	22.2	25.9	19.3	50.3	50.3	49.4
2001.....	25.3	28.1	23.1	36.7	37.8	35.5	24.6	27.5	22.4	22.1	25.4	19.5	49.8	50.5	48.7
2000.....	25.0	27.8	23.0	36.9	39.6	34.6	24.4	27.1	22.4	21.8	25.1	19.4	50.2	49.4	50.2
1999.....	25.0	27.8	23.0	37.8	39.1	36.6	24.3	27.1	22.3	21.6	25.0	19.2	50.4	50.0	50.1

Nutritional deficiencies (E40-E64)

2007.....	0.9	0.9	0.9	0.5	0.5	0.6	0.9	0.9	0.9	0.8	0.8	0.8	1.5	1.7	1.4
2006.....	0.8	0.7	0.8	0.6	0.6	0.7	0.8	0.8	0.8	0.8	0.7	0.8	1.3	1.6	1.1
2005.....	1.0	1.0	1.1	0.8	0.8	0.8	1.0	1.0	1.1	1.0	0.9	1.0	1.7	1.6	1.7
2004.....	1.0	0.9	1.0	0.6	0.5	0.7	1.0	1.0	1.0	1.0	0.9	1.0	1.7	1.9	1.5
2003.....	1.1	1.1	1.2	0.8	0.9	0.8	1.1	1.1	1.2	1.1	1.0	1.1	1.9	2.1	1.7
2002.....	1.3	1.2	1.3	1.0	1.0	1.0	1.3	1.2	1.3	1.2	1.2	1.3	2.2	2.1	2.1
2001.....	1.3	1.2	1.4	1.1	1.2	1.1	1.3	1.2	1.4	1.3	1.1	1.3	2.1	2.4	1.9
2000.....	1.5	1.4	1.6	1.2	1.2	1.2	1.5	1.4	1.6	1.5	1.3	1.6	2.4	2.7	2.2
1999.....	1.6	1.5	1.6	1.3	1.3	1.3	1.6	1.5	1.6	1.5	1.4	1.5	2.7	3.2	2.4



Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Malnutrition (E40-E46)</b>															
2007.....	0.8	0.8	0.8	0.5	0.5	0.6	0.8	0.8	0.8	0.8	0.8	0.8	1.4	1.6	1.3
2006.....	0.7	0.7	0.8	0.6	0.5	0.7	0.7	0.7	0.8	0.7	0.6	0.7	1.2	1.5	1.1
2005.....	1.0	0.9	1.0	0.8	0.8	0.7	1.0	0.9	1.0	0.9	0.8	1.0	1.6	1.5	1.7
2004.....	0.9	0.9	1.0	0.6	0.5	0.7	1.0	0.9	1.0	0.9	0.8	0.9	1.6	1.8	1.5
2003.....	1.1	1.0	1.1	0.8	0.7	0.8	1.1	1.0	1.1	1.0	0.9	1.1	1.9	2.1	1.7
2002.....	1.2	1.1	1.2	1.0	1.0	0.9	1.2	1.2	1.2	1.1	1.1	1.2	2.0	2.0	2.0
2001.....	1.2	1.1	1.3	1.1	1.1	1.1	1.2	1.1	1.3	1.2	1.0	1.2	1.9	2.2	1.7
2000.....	1.4	1.3	1.5	1.1	1.1	1.2	1.4	1.3	1.5	1.4	1.2	1.4	2.3	2.5	2.1
1999.....	1.5	1.4	1.5	1.2	1.2	1.2	1.5	1.4	1.5	1.4	1.3	1.4	2.5	3.0	2.2
<b>Other nutritional deficiencies (E50-E64)</b>															
2007.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	*	*
2006.....	0.1	0.0	0.0	*	*	*	0.1	0.0	0.0	0.1	0.0	0.0	*	*	*
2005.....	0.1	0.0	0.1	*	*	*	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
2004.....	0.1	0.0	0.0	*	*	*	0.1	0.1	0.0	0.1	0.0	0.1	*	*	*
2003.....	0.1	0.1	0.0	*	*	*	0.1	0.1	0.0	0.1	0.1	0.0	*	*	*
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.2	*	0.2
<b>Meningitis (G00,G03)</b>															
2007.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.3
2006.....	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.3
2005.....	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.5	0.3
2004.....	0.3	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.5	0.5	0.4
2003.....	0.3	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.6	0.6	0.6
2002.....	0.2	0.3	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.2	0.2	0.2	0.6	0.8	0.4
2001.....	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.6	0.8	0.4
2000.....	0.3	0.3	0.2	0.2	0.3	0.1	0.3	0.3	0.2	0.2	0.3	0.2	0.6	0.8	0.6
1999.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.7	0.9	0.6

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Parkinson's disease (G20-G21)</b>															
2007.....	6.4	9.6	4.3	4.1	5.8	2.9	6.5	9.9	4.4	6.9	10.5	4.7	3.2	5.0	2.2
2006.....	6.3	9.6	4.3	4.0	6.1	2.6	6.5	9.8	4.4	6.9	10.5	4.7	2.8	4.4	1.9
2005.....	6.4	9.8	4.4	4.3	6.3	3.1	6.5	10.0	4.4	7.0	10.6	4.7	3.0	5.0	1.9
2004.....	6.1	9.3	4.1	3.8	5.7	2.6	6.2	9.5	4.2	6.6	10.1	4.4	2.7	4.3	1.9
2003.....	6.2	9.3	4.2	3.6	5.5	2.4	6.3	9.5	4.3	6.7	10.1	4.6	2.8	4.5	1.9
2002.....	5.9	9.0	4.1	3.5	5.1	2.6	6.0	9.2	4.1	6.4	9.7	4.4	2.5	4.0	1.7
2001.....	5.9	9.1	4.0	3.8	5.6	2.7	5.9	9.3	4.1	6.3	9.9	4.3	2.5	4.0	1.8
2000.....	5.7	8.8	3.9	3.6	5.4	2.6	5.8	9.0	4.0	6.1	9.5	4.3	2.5	4.2	1.6
1999.....	5.4	8.4	3.7	3.5	5.1	2.5	5.4	8.5	3.7	5.8	9.0	3.9	2.4	3.8	1.7
<b>Alzheimer's disease (G30)</b>															
2007.....	22.7	18.5	24.9	13.5	11.3	14.7	23.2	18.9	25.5	24.1	19.6	26.6	19.4	16.0	20.9
2006.....	22.6	18.4	24.7	14.0	12.2	14.9	23.0	18.8	25.2	24.0	19.5	26.3	18.6	14.9	20.2
2005.....	22.9	18.5	25.1	13.8	12.5	14.5	23.3	18.8	25.7	24.2	19.4	26.7	19.7	16.7	20.9
2004.....	21.8	17.7	23.8	13.0	10.8	14.1	22.2	18.1	24.3	23.0	18.7	25.2	18.7	15.5	20.1
2003.....	21.4	17.5	23.3	13.4	12.2	14.1	21.7	17.8	23.7	22.6	18.4	24.7	16.9	14.9	17.6
2002.....	20.2	16.6	22.0	11.6	9.7	12.6	20.6	17.0	22.4	21.5	17.7	23.4	15.3	13.2	16.1
2001.....	19.1	16.1	20.5	10.7	9.2	11.5	19.4	16.4	20.8	20.2	17.0	21.7	14.8	13.3	15.3
2000.....	18.1	15.2	19.3	10.4	9.1	10.9	18.3	15.5	19.6	19.1	16.1	20.4	13.2	11.6	13.9
1999.....	16.5	14.4	17.4	9.6	8.1	10.3	16.7	14.6	17.6	17.4	15.2	18.4	11.4	10.3	11.8
<b>Major cardiovascular diseases (I00-I78)</b>															
2007.....	249.9	298.9	210.4	181.0	213.2	153.9	254.6	305.0	214.1	248.2	298.2	207.6	341.4	411.5	289.5
2006.....	261.2	312.0	220.4	190.3	224.6	162.0	265.7	317.9	224.0	258.9	310.6	217.2	354.9	428.3	301.2
2005.....	277.3	329.5	235.6	207.2	246.6	175.4	281.5	334.8	239.0	274.5	327.8	231.7	373.9	442.6	322.5
2004.....	286.5	340.0	243.8	210.2	250.9	177.3	290.7	345.3	247.4	283.3	338.1	239.5	387.8	458.3	336.2
2003.....	306.1	363.4	260.7	228.1	266.3	196.7	310.1	368.6	263.8	302.2	360.9	255.3	411.8	485.9	357.4
2002.....	317.4	377.2	270.5	236.3	280.1	201.7	321.1	382.0	273.4	313.4	374.8	264.9	421.3	493.6	368.4
2001.....	326.5	388.1	278.8	252.0	298.0	216.3	329.8	392.2	281.3	321.6	384.4	272.5	432.5	511.0	377.0
2000.....	339.7	407.3	288.4	258.2	307.0	220.5	342.9	411.4	290.9	334.9	404.1	281.9	443.9	523.4	388.2
1999.....	349.3	419.0	296.5	267.7	320.6	227.6	352.3	422.7	298.9	344.2	414.8	290.1	452.1	536.9	392.9

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Diseases of heart (I00-I09,I11,I13,I20-I51)</b>															
2007.....	190.9	237.7	154.0	136.0	165.0	111.8	194.8	243.1	156.8	191.4	239.8	153.0	251.9	311.8	208.2
2006.....	200.2	248.5	162.2	144.1	175.2	118.9	203.9	253.6	164.9	200.3	250.0	160.9	262.3	326.5	216.1
2005.....	211.1	260.9	172.3	157.3	192.4	129.1	214.5	265.5	174.8	210.7	262.2	170.3	275.6	335.1	231.9
2004.....	217.0	267.9	177.3	158.4	193.9	130.0	220.4	272.4	179.9	216.3	268.7	175.1	284.7	347.3	240.0
2003.....	232.3	286.6	190.3	173.2	206.8	145.8	235.5	291.0	192.5	230.9	286.9	187.1	304.1	369.2	257.3
2002.....	240.8	297.4	197.2	180.5	219.8	149.7	243.7	301.4	199.3	239.2	297.7	193.7	312.1	375.5	266.4
2001.....	247.8	305.4	203.9	192.2	232.6	161.0	250.4	309.0	205.7	245.6	304.8	200.0	320.4	388.7	272.9
2000.....	257.6	320.0	210.9	196.0	238.2	163.7	260.2	323.6	212.7	255.5	319.9	206.8	328.4	396.6	281.0
1999.....	266.5	331.0	218.1	205.8	249.2	172.3	268.8	334.3	219.7	263.8	330.0	213.9	337.4	410.3	286.7
<b>Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)</b>															
2007.....	1.0	0.8	1.2	0.7	0.5	0.8	1.0	0.8	1.2	1.0	0.8	1.2	0.8	0.7	0.9
2006.....	1.1	0.8	1.2	0.8	0.6	0.9	1.1	0.8	1.2	1.1	0.8	1.3	0.8	0.7	1.0
2005.....	1.1	0.8	1.3	0.8	0.4	1.0	1.1	0.9	1.3	1.1	0.9	1.3	0.8	0.7	0.9
2004.....	1.1	0.8	1.3	0.8	0.7	0.9	1.1	0.8	1.3	1.1	0.8	1.3	0.8	0.6	1.0
2003.....	1.2	1.0	1.4	0.8	0.6	1.0	1.2	1.0	1.4	1.3	1.0	1.4	1.1	0.9	1.2
2002.....	1.2	0.9	1.5	1.1	0.6	1.4	1.2	0.9	1.5	1.3	0.9	1.5	0.9	0.8	1.0
2001.....	1.2	0.9	1.5	0.9	0.7	1.1	1.2	0.9	1.5	1.2	0.9	1.4	1.1	0.7	1.4
2000.....	1.3	1.0	1.5	0.9	0.7	1.1	1.3	1.0	1.6	1.3	1.0	1.6	1.0	0.7	1.2
1999.....	1.3	0.9	1.6	1.1	0.4	1.5	1.3	0.9	1.6	1.4	1.0	1.6	1.0	0.8	1.2
<b>Hypertensive heart disease (I11)</b>															
2007.....	9.5	10.3	8.4	7.4	8.2	6.4	9.6	10.5	8.6	8.1	8.6	7.3	23.8	28.9	19.7
2006.....	9.4	10.1	8.3	7.4	8.1	6.6	9.5	10.3	8.4	8.0	8.4	7.2	23.5	29.1	19.2
2005.....	9.4	10.1	8.5	7.2	7.9	6.4	9.6	10.2	8.6	8.0	8.4	7.2	24.4	28.9	20.8
2004.....	9.5	10.1	8.5	7.8	9.1	6.7	9.6	10.2	8.6	7.9	8.3	7.2	24.6	28.8	21.0
2003.....	9.6	10.1	8.8	8.9	10.3	7.5	9.6	10.0	8.8	7.9	8.1	7.3	25.4	29.1	22.2
2002.....	9.1	9.5	8.3	8.4	9.7	7.1	9.2	9.5	8.4	7.5	7.6	7.0	24.7	28.2	21.5
2001.....	8.7	8.9	8.1	7.7	8.8	6.7	8.7	8.9	8.2	7.2	7.1	6.8	23.4	26.5	20.7
2000.....	8.6	8.7	8.1	7.0	7.5	6.3	8.7	8.7	8.1	7.1	7.0	6.7	24.0	26.6	21.5
1999.....	8.3	8.3	7.9	6.9	7.1	6.6	8.4	8.4	7.9	6.8	6.7	6.5	23.2	26.0	20.7

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Hypertensive heart and renal disease (I13)</b>															
2007.....	0.9	1.0	0.9	0.9	1.0	0.8	0.9	1.0	0.9	0.7	0.8	0.7	2.9	3.5	2.4
2006.....	0.9	1.0	0.8	0.7	0.8	0.6	0.9	1.0	0.8	0.7	0.8	0.6	3.0	3.6	2.5
2005.....	1.0	1.1	0.9	1.0	1.1	1.0	1.0	1.1	1.0	0.8	0.8	0.7	3.3	3.9	2.8
2004.....	1.0	1.1	0.9	1.0	1.2	0.9	1.0	1.1	0.9	0.7	0.8	0.7	3.3	3.7	3.0
2003.....	1.1	1.1	1.0	1.0	1.2	0.9	1.1	1.2	1.0	0.8	0.9	0.7	3.5	3.8	3.2
2002.....	1.0	1.0	0.9	1.1	1.2	1.0	1.0	1.0	0.9	0.7	0.7	0.7	3.6	3.9	3.2
2001.....	1.0	1.1	1.0	1.1	1.3	1.0	1.0	1.0	0.9	0.7	0.8	0.7	3.4	3.6	3.1
2000.....	1.0	1.1	0.9	1.0	1.2	0.9	1.0	1.1	0.9	0.8	0.8	0.7	3.4	3.7	3.1
1999.....	1.2	1.4	1.1	1.1	1.2	1.0	1.2	1.4	1.1	0.9	1.0	0.8	4.2	4.5	3.9
<b>Ischemic heart diseases (I20-I25)</b>															
2007.....	126.0	165.4	95.7	97.8	122.3	77.8	128.0	168.6	96.9	127.4	168.8	95.2	153.1	194.9	123.5
2006.....	134.9	176.5	103.1	106.4	132.8	85.4	136.8	179.5	104.2	136.0	179.3	102.4	164.1	209.8	132.0
2005.....	144.4	187.4	111.7	118.0	148.4	94.1	146.0	189.9	112.7	145.2	190.2	110.7	173.7	216.9	142.8
2004.....	150.2	194.1	116.7	119.2	149.6	95.3	151.9	196.8	117.8	150.9	196.8	115.6	182.0	226.7	150.5
2003.....	162.9	209.9	127.2	130.0	158.0	107.4	164.6	212.7	128.1	163.3	212.3	125.7	197.2	246.1	162.7
2002.....	170.8	220.4	133.6	138.3	172.2	112.2	172.3	222.7	134.4	171.0	222.7	131.7	205.1	253.2	171.4
2001.....	177.8	228.5	139.9	149.9	185.8	122.6	179.0	230.5	140.5	177.5	230.2	137.7	213.5	264.4	178.4
2000.....	186.8	241.4	146.5	153.2	190.1	125.1	188.0	243.5	147.2	186.6	243.5	144.3	220.4	271.9	185.0
1999.....	194.6	251.2	152.9	162.2	200.1	133.1	195.7	253.1	153.5	194.3	253.0	150.8	226.6	280.3	189.7
<b>Acute myocardial infarction (I21-I22)</b>															
2007.....	41.4	54.0	31.5	32.2	40.4	25.5	42.2	55.1	31.9	42.1	55.5	31.3	50.5	62.2	42.0
2006.....	45.0	58.8	34.2	35.0	43.4	28.4	45.8	60.0	34.6	45.7	60.4	34.0	54.2	67.6	44.7
2005.....	49.1	63.6	37.8	40.4	51.1	32.1	49.7	64.5	38.1	49.7	65.1	37.4	58.5	70.9	49.5
2004.....	52.2	67.8	40.2	40.9	51.2	32.7	53.0	68.9	40.7	52.9	69.4	39.9	62.9	76.0	53.5
2003.....	57.9	75.0	44.8	45.4	54.8	37.9	58.7	76.3	45.2	58.5	76.6	44.3	69.6	85.5	58.4
2002.....	62.1	80.6	48.1	49.5	61.6	40.2	62.8	81.7	48.5	62.5	82.1	47.4	75.0	90.9	64.1
2001.....	65.4	84.9	50.9	52.8	66.2	43.0	66.1	86.0	51.2	65.8	86.5	50.1	77.8	94.3	66.8
2000.....	69.9	91.3	54.1	54.1	67.6	43.9	70.6	92.5	54.5	70.4	93.0	53.4	82.0	100.1	70.0
1999.....	73.2	96.1	56.3	57.2	71.1	46.6	73.9	97.3	56.7	73.8	98.0	55.7	84.1	103.3	71.3

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Other acute ischemic heart diseases (I24)</b>															
2007.....	1.3	1.6	1.0	0.5	0.7	0.4	1.3	1.7	1.0	1.3	1.6	1.0	1.8	2.5	1.4
2006.....	1.2	1.5	1.0	0.5	0.5	0.5	1.3	1.6	1.1	1.2	1.5	1.0	1.9	2.5	1.5
2005.....	1.2	1.4	0.9	0.5	0.5	0.4	1.2	1.5	1.0	1.1	1.4	0.9	2.0	2.6	1.6
2004.....	0.8	1.0	0.6	0.2	0.2	*	0.9	1.1	0.7	0.8	1.0	0.6	1.4	2.1	1.0
2003.....	1.1	1.3	0.9	0.3	0.2	0.3	1.1	1.4	0.9	1.1	1.3	0.9	1.7	2.4	1.2
2002.....	1.2	1.5	0.9	0.4	0.5	0.3	1.2	1.5	1.0	1.2	1.5	0.9	1.7	2.3	1.3
2001.....	1.2	1.5	0.9	0.4	0.7	0.2	1.2	1.5	1.0	1.2	1.5	0.9	1.8	2.4	1.4
2000.....	1.2	1.5	0.9	0.5	0.7	0.4	1.2	1.6	1.0	1.2	1.5	0.9	1.8	2.5	1.4
1999.....	1.2	1.5	0.9	0.4	0.4	0.4	1.2	1.6	1.0	1.2	1.5	0.9	1.8	2.5	1.3
<b>Other forms of chronic ischemic heart disease (I20,I25)</b>															
2007.....	83.3	109.8	63.2	65.1	81.1	51.9	84.5	111.8	63.9	84.0	111.7	62.8	100.8	130.2	80.0
2006.....	88.7	116.1	67.9	70.8	88.9	56.4	89.8	117.9	68.5	89.0	117.4	67.4	108.1	139.7	85.9
2005.....	94.2	122.4	73.0	77.2	96.7	61.6	95.1	123.9	73.6	94.4	123.7	72.4	113.2	143.4	91.7
2004.....	97.1	125.3	75.8	78.1	98.2	62.4	98.0	126.8	76.4	97.2	126.4	75.1	117.6	148.6	96.0
2003.....	103.8	133.5	81.5	84.3	102.9	69.3	104.7	135.0	82.0	103.7	134.4	80.5	125.9	158.1	103.1
2002.....	107.6	138.3	84.5	88.4	110.0	71.8	108.3	139.5	84.9	107.3	139.1	83.4	128.5	160.0	106.1
2001.....	111.2	142.1	88.1	96.6	118.9	79.4	111.7	143.0	88.3	110.5	142.3	86.7	134.0	167.8	110.2
2000.....	115.7	148.6	91.5	98.7	121.9	80.8	116.1	149.4	91.7	115.1	149.0	90.0	136.5	169.3	113.7
1999.....	120.2	153.6	95.6	104.6	128.5	86.2	120.5	154.2	95.8	119.4	153.5	94.1	140.7	174.5	117.0
<b>Atherosclerotic cardiovascular disease, so described (I25.0)</b>															
2007.....	18.2	24.3	13.1	14.1	19.4	9.6	18.5	24.6	13.3	17.4	23.2	12.5	30.8	43.0	21.9
2006.....	19.3	25.6	14.0	16.0	21.9	11.0	19.5	25.8	14.1	18.2	24.1	13.2	33.2	46.6	23.7
2005.....	20.3	26.6	15.0	17.3	24.4	11.6	20.4	26.6	15.1	19.1	24.9	14.1	34.9	47.5	25.7
2004.....	21.1	27.5	15.7	17.5	24.6	11.8	21.2	27.6	15.9	19.9	25.9	14.8	36.5	49.5	27.3
2003.....	22.7	29.6	17.1	19.1	26.7	13.0	22.9	29.6	17.3	21.5	27.6	16.2	38.6	53.1	28.4
2002.....	23.5	30.4	17.8	19.3	26.2	13.9	23.6	30.4	17.9	22.1	28.4	16.8	40.6	55.2	30.3
2001.....	24.1	30.8	18.5	21.0	28.5	15.1	24.1	30.8	18.6	22.5	28.6	17.3	42.1	56.4	32.0
2000.....	25.2	32.3	19.4	21.3	29.0	15.4	25.2	32.2	19.5	23.5	30.0	18.2	43.9	59.2	33.3
1999.....	26.4	33.5	20.6	23.2	31.1	17.0	26.4	33.4	20.7	24.6	30.9	19.3	46.0	61.5	35.1

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)															
2007.....	65.1	85.5	50.1	50.9	61.8	42.2	66.0	87.2	50.6	66.5	88.4	50.4	70.0	87.2	58.1
2006.....	69.4	90.6	53.9	54.8	67.0	45.5	70.3	92.1	54.4	70.8	93.3	54.1	74.8	93.1	62.2
2005.....	73.9	95.8	58.0	59.9	72.4	50.1	74.7	97.3	58.4	75.3	98.7	58.2	78.3	95.9	66.0
2004.....	76.0	97.8	60.1	60.6	73.6	50.6	76.8	99.2	60.6	77.3	100.5	60.2	81.2	99.1	68.7
2003.....	81.1	104.0	64.4	65.2	76.3	56.3	81.9	105.4	64.7	82.2	106.8	64.2	87.3	105.0	74.7
2002.....	84.1	108.0	66.7	69.1	83.9	57.9	84.7	109.1	66.9	85.2	110.7	66.6	87.9	104.8	75.8
2001.....	87.1	111.3	69.6	75.6	90.5	64.3	87.6	112.2	69.7	88.0	113.7	69.4	91.8	111.4	78.2
2000.....	90.5	116.3	72.0	77.3	92.9	65.4	90.9	117.2	72.2	91.5	119.0	71.8	92.6	110.1	80.4
1999.....	93.8	120.1	75.0	81.4	97.4	69.2	94.1	120.9	75.1	94.8	122.6	74.8	94.7	113.0	81.9
Other heart diseases (I26-I51)															
2007.....	53.4	60.2	48.0	29.3	33.1	26.0	55.1	62.2	49.4	54.1	60.8	48.7	71.3	83.9	61.8
2006.....	53.9	60.0	48.7	28.8	32.8	25.4	55.5	62.0	50.2	54.6	60.6	49.5	70.8	83.4	61.5
2005.....	55.2	61.5	49.9	30.2	34.5	26.6	56.8	63.3	51.3	55.6	61.9	50.3	73.4	84.8	64.7
2004.....	55.2	61.7	49.9	29.5	33.3	26.3	56.8	63.5	51.3	55.6	61.9	50.3	74.0	87.4	64.5
2003.....	57.6	64.5	52.0	32.4	36.6	29.0	59.0	66.2	53.2	57.7	64.6	52.0	77.0	89.3	67.9
2002.....	58.5	65.5	52.8	31.5	36.1	27.9	60.0	67.2	54.1	58.6	65.7	52.9	77.8	89.3	69.2
2001.....	59.1	66.1	53.4	32.6	36.2	29.6	60.4	67.6	54.6	58.9	65.8	53.3	79.1	93.3	69.3
2000.....	60.0	67.9	53.8	33.9	38.7	30.2	61.2	69.3	54.9	59.7	67.6	53.5	79.7	93.7	70.1
1999.....	60.9	69.2	54.5	34.5	40.4	30.0	62.1	70.5	55.5	60.4	68.4	54.1	82.3	98.8	71.1
Acute and subacute endocarditis (I33)															
2007.....	0.4	0.5	0.3	0.3	0.3	0.2	0.4	0.5	0.3	0.4	0.5	0.3	0.7	1.1	0.4
2006.....	0.4	0.5	0.3	0.3	0.4	0.3	0.4	0.5	0.3	0.4	0.4	0.3	0.7	0.9	0.6
2005.....	0.4	0.5	0.3	0.3	0.5	0.2	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.9	0.6
2004.....	0.4	0.5	0.3	0.4	0.6	0.2	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6
2003.....	0.4	0.5	0.3	0.4	0.4	0.3	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6
2002.....	0.4	0.5	0.3	0.3	0.5	0.2	0.4	0.5	0.3	0.4	0.5	0.3	0.7	0.8	0.6
2001.....	0.4	0.5	0.3	0.3	0.3	0.2	0.4	0.6	0.3	0.4	0.5	0.3	0.8	1.1	0.6
2000.....	0.4	0.5	0.3	0.4	0.4	0.3	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.7
1999.....	0.4	0.5	0.3	0.3	0.6	0.2	0.4	0.5	0.3	0.4	0.5	0.3	0.8	0.9	0.6

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Diseases of pericardium and acute myocarditis (I30-I31,I40)</b>															
2007.....	0.3	0.3	0.2	0.2	0.2	0.1	0.3	0.4	0.2	0.3	0.3	0.2	0.4	0.5	0.4
2006.....	0.3	0.3	0.2	0.2	0.2	0.1	0.3	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.3
2005.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.2	0.5	0.5	0.5
2004.....	0.3	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.5	0.6	0.4
2003.....	0.3	0.3	0.3	0.2	0.2	0.2	0.3	0.4	0.3	0.3	0.3	0.2	0.5	0.5	0.5
2002.....	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.2	0.5	0.5	0.5
2001.....	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.5	0.5
2000.....	0.3	0.3	0.2	0.2	0.3	0.2	0.3	0.3	0.2	0.2	0.3	0.2	0.4	0.5	0.4
1999.....	0.3	0.4	0.3	0.3	0.2	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.5	0.6	0.5
<b>Heart failure (I50)</b>															
2007.....	17.3	18.6	16.2	9.6	10.4	9.0	17.8	19.2	16.6	17.8	19.2	16.7	20.5	22.9	18.8
2006.....	18.9	20.0	17.9	10.0	10.6	9.4	19.4	20.6	18.4	19.5	20.7	18.6	21.3	23.6	19.6
2005.....	18.9	19.9	18.0	10.1	10.7	9.6	19.4	20.4	18.5	19.4	20.5	18.5	22.1	23.5	20.7
2004.....	18.9	19.9	18.0	9.7	10.2	9.2	19.4	20.5	18.4	19.5	20.5	18.6	21.3	23.7	19.7
2003.....	19.4	20.6	18.4	10.8	11.7	10.1	19.8	21.0	18.8	19.8	21.1	18.8	22.2	23.9	20.9
2002.....	19.5	20.4	18.6	10.2	10.7	9.8	19.9	20.9	18.9	19.9	21.0	19.0	22.1	23.7	20.7
2001.....	20.2	21.2	19.2	10.7	10.9	10.4	20.5	21.6	19.6	20.6	21.7	19.7	22.4	24.7	20.9
2000.....	20.3	21.5	19.2	10.9	11.6	10.4	20.6	22.0	19.6	20.7	22.0	19.6	22.4	24.6	21.0
1999.....	20.3	21.4	19.3	11.1	11.7	10.6	20.6	21.8	19.6	20.6	21.7	19.6	22.8	25.5	21.1
<b>All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)</b>															
2007.....	35.5	40.7	31.2	19.2	22.1	16.7	36.7	42.1	32.2	35.7	40.8	31.4	49.6	59.5	42.2
2006.....	34.3	39.2	30.2	18.3	21.6	15.6	35.4	40.5	31.2	34.4	39.1	30.4	48.4	58.6	41.0
2005.....	35.6	40.7	31.3	19.6	23.1	16.6	36.7	42.0	32.3	35.5	40.6	31.3	50.1	59.9	42.8
2004.....	35.7	41.0	31.4	19.2	22.3	16.6	36.8	42.2	32.3	35.5	40.6	31.2	51.5	62.2	43.9
2003.....	37.5	43.1	33.0	21.1	24.3	18.4	38.5	44.2	33.8	37.2	42.6	32.6	53.6	63.9	45.9
2002.....	38.4	44.2	33.7	20.7	24.7	17.7	39.4	45.5	34.6	38.1	43.9	33.4	54.6	64.2	47.5
2001.....	38.2	44.1	33.6	21.4	24.7	18.7	39.1	45.2	34.3	37.6	43.4	33.1	55.3	66.9	47.3
2000.....	39.0	45.5	34.0	22.4	26.5	19.3	39.9	46.5	34.7	38.4	44.7	33.5	56.1	67.6	48.0
1999.....	39.9	46.9	34.6	22.7	27.9	18.9	40.8	47.8	35.3	39.1	45.8	34.0	58.2	71.7	48.9

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year All origins 1/ Both sexes Male Female Hispanic Both sexes Male Female Non-Hispanic 2/ Both sexes Male Female Non-Hispanic white 3/ Both sexes Male Female Non-Hispanic black 3/ Both sexes Male Female

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Table with 17 columns (Year, All origins, Hispanic, Non-Hispanic, Non-Hispanic white, Non-Hispanic black) and 17 rows (2007-1999) showing death rates for Essential hypertension and hypertensive renal disease.

Cerebrovascular diseases (I60-I69)

Table with 17 columns (Year, All origins, Hispanic, Non-Hispanic, Non-Hispanic white, Non-Hispanic black) and 17 rows (2007-1999) showing death rates for Cerebrovascular diseases.

Atherosclerosis (I70)

Table with 17 columns (Year, All origins, Hispanic, Non-Hispanic, Non-Hispanic white, Non-Hispanic black) and 17 rows (2007-1999) showing death rates for Atherosclerosis.



Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Other diseases of circulatory system (I71-I78)</b>															
2007.....	6.9	8.8	5.5	4.0	4.9	3.2	7.1	9.1	5.6	7.0	9.0	5.5	8.7	10.8	7.3
2006.....	7.2	9.4	5.6	4.2	5.6	3.1	7.4	9.6	5.8	7.3	9.6	5.6	9.1	10.5	7.9
2005.....	7.8	10.0	6.1	4.7	6.6	3.3	7.9	10.3	6.2	7.8	10.2	6.1	9.8	11.8	8.2
2004.....	7.9	10.2	6.3	4.6	6.0	3.4	8.1	10.5	6.5	8.1	10.5	6.3	9.9	11.8	8.4
2003.....	8.4	11.2	6.4	5.3	7.2	3.9	8.6	11.5	6.6	8.6	11.5	6.4	10.0	12.0	8.5
2002.....	8.7	11.5	6.6	5.2	6.6	4.1	8.8	11.8	6.7	8.8	11.9	6.6	10.1	12.2	8.7
2001.....	9.0	12.0	6.9	5.0	6.5	3.9	9.2	12.2	7.0	9.1	12.3	6.9	10.9	12.9	9.5
2000.....	9.4	12.9	7.1	5.8	7.7	4.5	9.6	13.1	7.1	9.5	13.3	6.9	11.1	13.2	9.6
1999.....	9.5	12.8	7.1	5.6	7.9	4.2	9.6	13.0	7.3	9.6	13.2	7.1	10.6	12.5	9.3
<b>Aortic aneurysm and dissection (I71)</b>															
2007.....	4.1	5.7	2.9	2.1	2.9	1.4	4.3	6.0	3.0	4.3	6.0	3.0	4.3	5.6	3.3
2006.....	4.3	6.0	3.0	2.2	3.3	1.3	4.4	6.3	3.1	4.4	6.3	3.0	4.4	5.6	3.5
2005.....	4.6	6.5	3.1	2.3	3.6	1.4	4.7	6.7	3.2	4.7	6.8	3.2	4.7	6.1	3.6
2004.....	4.6	6.7	3.1	2.4	3.5	1.5	4.8	6.9	3.3	4.8	7.0	3.3	4.5	6.2	3.4
2003.....	5.1	7.5	3.3	3.0	4.6	1.9	5.2	7.7	3.4	5.3	7.9	3.4	4.7	6.2	3.5
2002.....	5.2	7.7	3.3	2.5	3.8	1.6	5.3	7.9	3.4	5.4	8.1	3.5	4.7	6.1	3.6
2001.....	5.4	8.0	3.6	2.5	4.1	1.5	5.5	8.2	3.7	5.6	8.4	3.7	5.0	6.5	4.0
2000.....	5.7	8.7	3.7	3.2	4.8	2.1	5.8	8.9	3.7	5.9	9.2	3.7	5.3	6.9	4.2
1999.....	5.8	8.8	3.7	2.8	4.5	1.7	5.9	9.0	3.8	6.1	9.3	3.8	5.0	6.5	3.9
<b>Other diseases of arteries, arterioles and capillaries (I72-I78)</b>															
2007.....	2.8	3.1	2.5	1.9	2.0	1.8	2.8	3.1	2.6	2.7	3.0	2.5	4.5	5.2	4.0
2006.....	2.9	3.3	2.7	2.0	2.3	1.8	3.0	3.4	2.7	2.9	3.3	2.6	4.6	4.9	4.4
2005.....	3.2	3.5	2.9	2.4	3.0	1.9	3.2	3.5	3.0	3.1	3.4	2.9	5.1	5.7	4.6
2004.....	3.3	3.5	3.1	2.2	2.4	2.0	3.4	3.6	3.2	3.2	3.5	3.0	5.3	5.7	5.1
2003.....	3.3	3.7	3.1	2.2	2.6	2.0	3.4	3.7	3.2	3.3	3.6	3.0	5.3	5.8	4.9
2002.....	3.5	3.8	3.3	2.7	2.9	2.6	3.5	3.9	3.3	3.4	3.7	3.2	5.5	6.1	5.1
2001.....	3.6	3.9	3.3	2.4	2.4	2.4	3.6	4.0	3.3	3.4	3.8	3.2	5.9	6.5	5.6
2000.....	3.7	4.2	3.4	2.6	2.9	2.4	3.7	4.2	3.4	3.6	4.1	3.2	5.8	6.3	5.4
1999.....	3.7	4.0	3.4	2.8	3.4	2.4	3.7	4.0	3.4	3.6	3.9	3.3	5.7	6.0	5.4

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year

	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Other disorders of circulatory system (I80-I99)

2007.....	1.3	1.4	1.2	0.9	0.9	0.8	1.3	1.4	1.2	1.3	1.3	1.1	2.2	2.5	2.0
2006.....	1.3	1.3	1.2	0.8	0.9	0.8	1.3	1.3	1.2	1.2	1.3	1.2	2.4	2.5	2.2
2005.....	1.6	1.6	1.5	1.2	1.2	1.3	1.6	1.7	1.5	1.5	1.6	1.4	2.7	3.1	2.4
2004.....	1.5	1.6	1.5	1.0	0.9	1.0	1.6	1.7	1.5	1.5	1.6	1.4	2.8	2.9	2.6
2003.....	1.6	1.6	1.6	1.1	1.2	1.1	1.6	1.6	1.6	1.5	1.5	1.5	3.0	3.2	2.8
2002.....	1.6	1.6	1.6	1.1	1.0	1.2	1.7	1.6	1.6	1.6	1.5	1.6	2.9	3.2	2.7
2001.....	1.6	1.7	1.6	1.4	1.5	1.3	1.7	1.7	1.6	1.6	1.6	1.5	2.8	3.1	2.5
2000.....	1.7	1.7	1.6	1.4	1.6	1.3	1.7	1.7	1.6	1.6	1.6	1.5	2.9	3.1	2.8
1999.....	1.5	1.5	1.4	1.2	1.3	1.2	1.5	1.5	1.4	1.4	1.4	1.3	2.5	2.7	2.2

Influenza and pneumonia (J09-J18)

2007.....	16.2	19.3	14.2	13.1	15.6	11.3	16.4	19.6	14.4	16.2	19.1	14.3	18.8	24.0	15.5
2006.....	17.8	21.2	15.5	15.0	17.4	13.3	17.9	21.4	15.6	17.8	21.1	15.6	19.9	24.7	16.9
2005.....	20.3	23.9	17.9	16.8	19.6	14.8	20.5	24.1	18.1	20.4	23.8	18.1	22.1	27.3	18.7
2004.....	19.8	23.7	17.3	17.1	19.8	15.2	19.9	23.8	17.3	19.6	23.4	17.2	22.5	29.4	18.4
2003.....	22.0	26.1	19.4	18.4	21.6	16.2	22.2	26.3	19.6	22.0	25.8	19.6	23.6	31.2	19.0
2002.....	22.6	27.0	19.9	19.2	23.3	16.5	22.7	27.1	20.0	22.6	26.8	20.0	24.3	31.2	20.3
2001.....	22.0	26.6	19.2	20.5	24.6	17.8	22.0	26.6	19.2	21.7	26.0	19.1	24.3	32.5	19.6
2000.....	23.7	28.9	20.7	20.6	25.6	17.4	23.8	28.9	20.8	23.5	28.3	20.7	25.9	34.3	21.0
1999.....	23.5	28.5	20.6	18.8	22.7	16.2	23.6	28.7	20.7	23.4	28.2	20.7	25.5	33.6	20.9

Influenza (J09-J11)

2007.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.0	*	*
2006.....	0.3	0.3	0.2	0.1	0.1	*	0.3	0.3	0.3	0.3	0.3	0.3	0.1	0.2	*
2005.....	0.6	0.6	0.6	0.2	0.3	0.2	0.6	0.6	0.6	0.6	0.6	0.6	0.2	0.3	0.2
2004.....	0.4	0.4	0.3	0.1	*	*	0.4	0.4	0.3	0.4	0.4	0.4	0.2	0.2	0.1
2003.....	0.6	0.6	0.6	0.3	0.4	0.2	0.6	0.6	0.6	0.7	0.7	0.6	0.3	0.3	0.3
2002.....	0.2	0.3	0.2	*	*	*	0.2	0.3	0.2	0.3	0.3	0.2	0.1	*	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.6	0.7	0.6	0.3	0.4	0.3	0.6	0.7	0.6	0.7	0.7	0.7	0.3	0.4	0.3
1999.....	0.6	0.6	0.6	0.3	0.3	0.3	0.6	0.6	0.6	0.6	0.7	0.6	0.3	0.3	0.3

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Pneumonia (J12-J18)</b>															
2007.....	16.1	19.2	14.1	13.0	15.5	11.3	16.3	19.5	14.2	16.0	19.0	14.2	18.7	24.0	15.4
2006.....	17.5	20.9	15.3	14.9	17.3	13.2	17.7	21.1	15.4	17.5	20.8	15.3	19.7	24.6	16.8
2005.....	19.7	23.3	17.3	16.6	19.3	14.5	19.9	23.5	17.5	19.7	23.1	17.5	21.8	27.0	18.5
2004.....	19.4	23.3	16.9	17.0	19.7	15.1	19.5	23.5	17.0	19.2	23.0	16.8	22.3	29.2	18.2
2003.....	21.4	25.5	18.8	18.1	21.3	15.9	21.6	25.6	18.9	21.4	25.1	19.0	23.2	30.9	18.6
2002.....	22.4	26.7	19.6	19.2	23.2	16.4	22.5	26.9	19.8	22.4	26.5	19.8	24.2	31.1	20.2
2001.....	21.9	26.5	19.1	20.4	24.5	17.8	21.9	26.5	19.1	21.6	25.9	19.0	24.3	32.4	19.5
2000.....	23.1	28.2	20.1	20.3	25.3	17.1	23.1	28.2	20.1	22.8	27.6	20.0	25.6	33.9	20.6
1999.....	22.9	27.9	20.0	18.5	22.5	15.9	23.0	28.0	20.1	22.8	27.5	20.0	25.3	33.4	20.6
<b>Other acute lower respiratory infections (J20-J22,U04)</b>															
2007.....	0.1	0.1	0.1	0.0	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2006.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2005.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
2004.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2002.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
2001.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.1
1999.....	0.2	0.2	0.2	0.1	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
<b>Acute bronchitis and bronchiolitis (J20-J21)</b>															
2007.....	0.1	0.1	0.0	0.0	*	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	*	*
2006.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2005.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2004.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2003.....	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2002.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*
2001.....	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*
2000.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1
1999.....	0.1	0.1	0.1	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified acute lower respiratory infections (J22, U04)															
2007.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
2006.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2005.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2004.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2003.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2002.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2001.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.0	0.0	0.0	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	0.0	0.0	0.1	0.0	0.0	*	*	*
1999.....	0.1	0.0	0.1	*	*	*	0.1	0.0	0.1	0.1	0.0	0.1	*	*	*
Chronic lower respiratory diseases (J40-J47)															
2007.....	40.8	48.0	36.0	17.5	22.8	13.9	42.4	49.9	37.6	44.9	51.8	40.4	28.7	40.4	21.9
2006.....	40.5	47.6	35.9	17.3	21.9	14.3	42.0	49.3	37.4	44.4	51.2	40.2	28.6	40.3	21.7
2005.....	43.2	51.2	38.1	19.3	25.1	15.4	44.8	52.9	39.6	47.2	54.7	42.5	31.1	44.8	23.2
2004.....	41.1	49.5	36.0	18.4	23.8	14.9	42.6	51.0	37.3	44.9	52.8	40.0	28.7	41.6	21.2
2003.....	43.3	52.3	37.8	20.2	27.1	15.8	44.7	53.7	39.1	47.0	55.4	41.8	30.5	45.1	22.3
2002.....	43.5	53.5	37.4	20.6	27.2	16.2	44.8	54.9	38.6	46.9	56.5	41.2	31.5	46.8	22.9
2001.....	43.7	54.0	37.6	20.7	27.6	16.5	44.9	55.4	38.7	47.0	56.9	41.1	31.3	47.0	22.6
2000.....	44.2	55.8	37.4	21.1	28.8	16.3	45.3	57.1	38.5	47.2	58.5	40.7	32.0	47.9	23.1
1999.....	45.4	58.7	37.7	23.4	33.0	17.7	46.5	59.9	38.6	48.3	61.3	40.8	33.9	52.0	23.7
Bronchitis, chronic and unspecified (J40-J42)															
2007.....	0.2	0.2	0.2	0.2	0.2	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2006.....	0.2	0.2	0.2	0.2	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.2
2005.....	0.3	0.3	0.3	0.2	*	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.1
2004.....	0.2	0.3	0.2	0.2	*	*	0.2	0.3	0.2	0.3	0.3	0.2	0.2	0.2	0.2
2003.....	0.3	0.3	0.3	0.2	0.3	0.2	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
2002.....	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.4	0.3	0.3	0.4	0.3	0.2	0.3	0.2
2001.....	0.3	0.4	0.3	0.2	*	0.2	0.3	0.4	0.3	0.4	0.4	0.3	0.2	0.3	0.2
2000.....	0.4	0.5	0.4	0.4	0.5	0.3	0.4	0.5	0.4	0.4	0.5	0.4	0.3	0.3	0.3
1999.....	0.4	0.5	0.4	0.4	*	0.4	0.4	0.5	0.4	0.4	0.5	0.4	0.4	0.5	0.3

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Emphysema (J43)</b>															
2007.....	4.1	5.1	3.4	1.4	2.1	0.9	4.3	5.3	3.6	4.6	5.5	3.9	2.7	4.4	1.7
2006.....	4.1	5.0	3.5	1.5	2.1	1.0	4.3	5.2	3.7	4.6	5.4	4.0	2.6	4.1	1.7
2005.....	4.6	5.8	3.8	1.7	2.4	1.1	4.8	6.1	4.0	5.1	6.3	4.4	3.1	5.0	1.9
2004.....	4.6	5.8	3.8	1.6	2.2	1.2	4.8	6.1	4.0	5.1	6.3	4.3	2.8	4.7	1.7
2003.....	5.1	6.3	4.3	2.0	2.9	1.4	5.3	6.6	4.5	5.6	6.8	4.9	3.3	5.4	2.1
2002.....	5.4	7.0	4.4	2.0	2.8	1.5	5.6	7.2	4.6	6.0	7.5	5.0	3.5	5.9	2.1
2001.....	5.8	7.4	4.7	2.1	3.3	1.4	6.0	7.7	4.9	6.4	8.0	5.3	3.4	5.4	2.2
2000.....	6.0	7.8	4.9	2.1	3.2	1.3	6.2	8.1	5.1	6.6	8.3	5.5	3.8	6.2	2.4
1999.....	6.5	8.6	5.2	2.4	3.8	1.5	6.7	8.8	5.3	7.1	9.2	5.8	4.0	6.8	2.3
<b>Asthma (J45-J46)</b>															
2007.....	1.1	0.9	1.2	0.8	0.7	0.9	1.1	0.9	1.3	0.9	0.7	1.1	2.6	2.5	2.6
2006.....	1.2	0.9	1.3	1.0	0.7	1.2	1.2	1.0	1.4	0.9	0.7	1.1	2.8	2.6	2.9
2005.....	1.3	1.0	1.5	1.0	0.9	1.1	1.3	1.0	1.6	1.0	0.7	1.3	3.1	2.7	3.4
2004.....	1.3	1.0	1.4	1.1	0.9	1.4	1.3	1.1	1.5	1.0	0.8	1.2	3.1	3.1	3.1
2003.....	1.4	1.1	1.6	1.3	1.1	1.4	1.4	1.2	1.6	1.1	0.9	1.3	3.2	3.0	3.4
2002.....	1.5	1.2	1.7	1.3	1.0	1.5	1.5	1.2	1.7	1.2	0.9	1.4	3.5	3.4	3.5
2001.....	1.5	1.2	1.7	1.4	1.0	1.7	1.5	1.2	1.8	1.2	0.9	1.4	3.6	3.3	3.9
2000.....	1.6	1.3	1.8	1.6	1.5	1.7	1.6	1.3	1.8	1.3	1.0	1.5	3.9	3.5	4.2
1999.....	1.7	1.3	2.0	1.8	1.3	2.2	1.7	1.3	1.9	1.4	1.0	1.6	4.0	3.6	4.2
<b>Other chronic lower respiratory diseases (J44,J47)</b>															
2007.....	35.3	41.9	31.2	15.1	19.7	11.9	36.8	43.4	32.5	39.2	45.4	35.2	23.2	33.3	17.4
2006.....	34.9	41.4	30.8	14.7	18.9	11.9	36.3	42.9	32.1	38.6	44.8	34.8	23.0	33.4	16.9
2005.....	37.0	44.1	32.5	16.4	21.6	12.9	38.4	45.5	33.8	40.8	47.4	36.6	24.7	36.7	17.7
2004.....	35.0	42.3	30.5	15.4	20.4	12.1	36.2	43.6	31.6	38.4	45.4	34.2	22.5	33.5	16.2
2003.....	36.5	44.5	31.7	16.7	22.8	12.8	37.7	45.7	32.8	39.9	47.4	35.4	23.8	36.5	16.6
2002.....	36.2	44.9	31.1	16.9	23.1	12.9	37.3	46.0	32.1	39.4	47.7	34.5	24.3	37.3	17.1
2001.....	36.1	45.0	30.8	16.9	23.1	13.2	37.1	46.1	31.7	39.0	47.7	34.0	24.0	38.0	16.3
2000.....	36.1	46.1	30.3	17.1	23.6	13.0	37.0	47.2	31.1	38.9	48.7	33.3	23.9	37.8	16.2
1999.....	36.8	48.3	30.2	18.8	27.5	13.6	37.7	49.2	31.0	39.4	50.6	32.9	25.5	41.1	16.9

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Pneumoconioses and chemical effects (J60-J66,J68)</b>															
2007.....	0.3	0.7	0.0	0.1	0.3	*	0.3	0.7	0.0	0.3	0.8	0.0	0.2	0.4	*
2006.....	0.3	0.7	0.0	*	*	*	0.3	0.8	0.0	0.3	0.8	0.0	0.2	0.5	*
2005.....	0.3	0.8	0.0	0.1	*	*	0.3	0.9	0.0	0.4	0.9	0.0	0.2	0.5	*
2004.....	0.3	0.9	0.0	0.1	0.4	*	0.4	0.9	0.0	0.4	1.0	0.0	0.2	0.7	*
2003.....	0.4	1.0	0.0	0.1	*	*	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.5	*
2002.....	0.4	1.0	0.0	0.2	0.5	*	0.4	1.0	0.0	0.4	1.0	0.0	0.2	0.6	*
2001.....	0.4	1.0	0.0	0.2	0.4	*	0.4	1.0	0.0	0.4	1.1	0.0	0.2	0.7	*
2000.....	0.4	1.1	0.0	0.2	0.5	*	0.4	1.1	0.0	0.4	1.2	0.0	0.3	0.7	*
1999.....	0.4	1.0	0.0	*	*	*	0.4	1.1	0.0	0.4	1.1	0.0	0.3	0.8	*
<b>Pneumonitis due to solids and liquids (J69)</b>															
2007.....	5.2	7.3	4.0	2.7	3.5	2.1	5.4	7.6	4.1	5.5	7.7	4.2	5.4	7.7	4.1
2006.....	5.3	7.5	4.0	3.0	3.6	2.5	5.5	7.7	4.1	5.5	7.8	4.1	5.6	8.2	4.1
2005.....	5.6	7.6	4.4	3.4	4.6	2.6	5.7	7.8	4.5	5.7	7.7	4.5	6.5	9.2	4.9
2004.....	5.6	7.7	4.3	3.3	4.1	2.8	5.7	7.9	4.4	5.7	7.9	4.4	6.2	8.8	4.7
2003.....	5.9	8.0	4.6	3.5	4.3	2.9	6.0	8.2	4.7	6.0	8.2	4.7	6.8	9.3	5.4
2002.....	6.1	8.4	4.7	3.9	5.3	3.1	6.2	8.6	4.8	6.2	8.7	4.8	6.7	9.4	5.2
2001.....	6.1	8.5	4.8	4.4	6.0	3.5	6.2	8.6	4.9	6.2	8.7	4.8	7.1	9.5	5.8
2000.....	6.1	8.6	4.6	4.1	5.5	3.3	6.1	8.7	4.7	6.1	8.7	4.7	6.8	9.5	5.3
1999.....	5.6	8.1	4.3	3.7	4.7	3.1	5.7	8.2	4.3	5.7	8.1	4.3	6.8	10.0	5.2
<b>Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)</b>															
2007.....	9.0	11.1	7.6	7.7	9.0	6.6	9.1	11.2	7.7	9.2	11.3	7.8	8.8	10.5	7.6
2006.....	8.9	10.9	7.6	7.3	8.3	6.6	9.1	11.0	7.7	9.2	11.2	7.7	8.4	10.1	7.4
2005.....	8.9	10.6	7.7	7.9	9.0	7.1	9.0	10.7	7.7	9.0	10.8	7.8	8.9	10.9	7.6
2004.....	9.2	11.1	7.8	8.1	9.0	7.4	9.3	11.3	7.9	9.3	11.4	8.0	8.7	10.6	7.5
2003.....	8.7	10.5	7.5	7.6	8.7	6.8	8.8	10.6	7.6	8.9	10.7	7.6	8.2	10.2	7.0
2002.....	8.7	10.4	7.6	7.5	8.4	6.7	8.7	10.5	7.6	8.8	10.5	7.6	8.6	10.1	7.5
2001.....	9.3	11.0	8.1	8.2	9.8	7.1	9.3	11.1	8.1	9.3	11.2	8.1	9.6	10.9	8.8
2000.....	9.2	11.3	7.9	8.3	9.5	7.3	9.3	11.3	8.0	9.3	11.4	8.0	9.3	11.4	8.1
1999.....	9.2	11.2	7.8	8.3	10.0	7.1	9.2	11.2	7.9	9.2	11.2	7.8	9.6	11.3	8.5

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Peptic ulcer (K25-K28)</b>															
2007.....	1.0	1.1	0.8	0.6	0.8	0.5	1.0	1.1	0.8	1.0	1.1	0.8	1.0	1.4	0.8
2006.....	1.1	1.2	0.9	0.9	1.1	0.7	1.1	1.2	0.9	1.0	1.2	1.0	1.3	1.7	0.9
2005.....	1.1	1.4	0.9	0.9	1.0	0.8	1.2	1.4	1.0	1.1	1.3	1.0	1.3	1.8	0.9
2004.....	1.2	1.4	1.1	0.9	1.1	0.7	1.2	1.4	1.1	1.2	1.4	1.1	1.3	1.7	1.1
2003.....	1.3	1.6	1.1	1.1	1.4	0.9	1.3	1.6	1.1	1.3	1.5	1.2	1.5	2.1	1.0
2002.....	1.4	1.7	1.2	1.2	1.6	0.8	1.4	1.7	1.2	1.4	1.7	1.2	1.4	2.0	1.1
2001.....	1.6	1.8	1.4	1.2	1.6	0.8	1.6	1.8	1.4	1.6	1.8	1.4	1.6	2.3	1.1
2000.....	1.6	2.0	1.4	1.4	1.6	1.2	1.7	2.1	1.4	1.6	2.0	1.4	1.7	2.4	1.3
1999.....	1.7	2.0	1.4	1.3	1.8	0.9	1.7	2.1	1.4	1.7	2.0	1.4	1.9	2.5	1.4
<b>Diseases of appendix (K35-K38)</b>															
2007.....	0.1	0.2	0.1	0.1	*	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
2006.....	0.1	0.2	0.1	0.1	0.2	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1
2005.....	0.1	0.2	0.1	0.1	*	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2	0.1
2004.....	0.1	0.2	0.1	0.1	0.1	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.1
2003.....	0.1	0.2	0.1	0.1	0.2	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.3	0.2
2002.....	0.2	0.2	0.1	0.1	*	*	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.1
2001.....	0.1	0.2	0.1	0.1	*	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.2
2000.....	0.1	0.2	0.1	0.1	*	*	0.1	0.2	0.1	0.1	0.2	0.1	0.3	0.4	0.2
1999.....	0.1	0.2	0.1	0.1	0.2	*	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.4	0.1
<b>Hernia (K40-K46)</b>															
2007.....	0.5	0.5	0.5	0.4	0.3	0.5	0.5	0.6	0.5	0.5	0.6	0.5	0.6	0.8	0.5
2006.....	0.5	0.6	0.5	0.4	0.5	0.4	0.6	0.6	0.5	0.6	0.6	0.5	0.6	0.7	0.6
2005.....	0.5	0.5	0.5	0.4	0.3	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.5
2004.....	0.5	0.5	0.5	0.4	0.3	0.5	0.5	0.5	0.5	0.5	0.6	0.5	0.6	0.6	0.5
2003.....	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.6	0.5	0.5	0.6	0.5	0.6	0.7	0.6
2002.....	0.5	0.5	0.6	0.5	0.4	0.5	0.5	0.5	0.6	0.6	0.5	0.6	0.5	0.6	0.5
2001.....	0.5	0.5	0.5	0.6	0.4	0.7	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.8	0.5
2000.....	0.5	0.5	0.5	0.6	0.4	0.6	0.5	0.5	0.5	0.5	0.6	0.5	0.6	0.6	0.6
1999.....	0.5	0.6	0.5	0.5	0.4	0.5	0.5	0.6	0.5	0.5	0.6	0.5	0.6	0.6	0.6

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Chronic liver disease and cirrhosis (K70,K73-K74)</b>															
2007.....	9.1	12.7	5.9	13.8	19.9	8.1	8.7	12.0	5.7	8.9	12.2	5.9	7.5	11.3	4.6
2006.....	8.8	12.1	5.8	13.3	18.9	8.1	8.4	11.4	5.6	8.6	11.7	5.8	7.1	10.4	4.5
2005.....	9.0	12.4	5.8	13.9	20.4	7.9	8.5	11.7	5.7	8.7	11.8	5.8	7.8	11.7	4.9
2004.....	9.0	12.5	5.8	14.0	19.7	8.6	8.5	11.8	5.7	8.7	11.9	5.8	8.1	12.0	4.9
2003.....	9.3	13.0	6.0	14.7	20.9	9.0	8.9	12.3	5.8	9.0	12.4	5.9	8.5	12.6	5.3
2002.....	9.4	12.9	6.3	15.4	22.3	9.0	8.9	12.0	6.1	9.0	12.2	6.1	8.6	12.1	5.8
2001.....	9.5	13.2	6.2	15.8	23.7	8.7	9.0	12.3	6.1	9.0	12.2	6.1	9.4	14.1	5.7
2000.....	9.5	13.4	6.2	16.5	24.7	9.3	9.0	12.5	6.0	9.0	12.4	5.9	9.6	13.7	6.4
1999.....	9.6	13.5	6.1	16.1	23.8	9.1	9.1	12.7	5.9	9.0	12.5	5.8	10.2	15.0	6.4
<b>Alcoholic liver disease (K70)</b>															
2007.....	4.5	6.8	2.3	6.9	12.0	2.2	4.2	6.3	2.4	4.4	6.5	2.4	3.5	5.6	1.8
2006.....	4.1	6.3	2.2	6.5	11.2	2.2	3.9	5.7	2.2	4.0	5.9	2.3	3.0	4.7	1.8
2005.....	4.2	6.4	2.2	6.4	11.3	2.0	3.9	5.9	2.2	4.0	5.9	2.3	3.5	5.5	2.0
2004.....	4.2	6.3	2.2	6.3	10.9	2.0	3.9	5.9	2.2	4.0	6.0	2.2	3.6	5.7	2.0
2003.....	4.2	6.4	2.1	6.7	11.6	2.2	3.9	5.9	2.1	4.0	6.0	2.1	3.9	6.1	2.1
2002.....	4.2	6.3	2.2	6.9	12.1	2.1	3.9	5.8	2.2	3.9	5.8	2.2	3.9	5.9	2.3
2001.....	4.3	6.6	2.2	7.6	13.6	2.3	4.0	6.0	2.2	3.9	5.9	2.2	4.5	7.1	2.4
2000.....	4.3	6.8	2.2	7.8	14.4	2.1	4.1	6.2	2.1	4.0	6.1	2.0	4.7	7.1	2.9
1999.....	4.3	6.9	2.1	7.6	13.5	2.4	4.1	6.3	2.1	4.0	6.2	2.0	5.2	8.2	2.8
<b>Other chronic liver disease and cirrhosis (K73-K74)</b>															
2007.....	4.6	5.9	3.5	6.9	7.9	5.9	4.4	5.7	3.4	4.5	5.8	3.4	4.1	5.7	2.8
2006.....	4.6	5.9	3.6	6.8	7.7	5.9	4.5	5.7	3.4	4.6	5.8	3.5	4.0	5.7	2.8
2005.....	4.7	6.0	3.6	7.5	9.1	5.9	4.5	5.8	3.5	4.6	5.8	3.6	4.3	6.2	2.9
2004.....	4.8	6.1	3.7	7.7	8.8	6.6	4.6	5.9	3.5	4.7	5.9	3.5	4.4	6.3	3.0
2003.....	5.2	6.5	3.9	8.0	9.3	6.7	4.9	6.3	3.7	5.0	6.4	3.8	4.6	6.4	3.2
2002.....	5.2	6.5	4.1	8.5	10.2	6.9	5.0	6.2	3.9	5.1	6.3	4.0	4.7	6.2	3.5
2001.....	5.2	6.6	4.0	8.2	10.1	6.4	5.0	6.3	3.9	5.0	6.3	4.0	5.0	7.0	3.2
2000.....	5.2	6.6	4.0	8.7	10.3	7.2	5.0	6.3	3.8	5.0	6.3	3.8	4.9	6.6	3.5
1999.....	5.2	6.7	4.0	8.5	10.3	6.7	5.0	6.4	3.8	5.0	6.4	3.8	5.0	6.8	3.6



Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year All origins 1/ Both sexes Male Female Hispanic Both sexes Male Female Non-Hispanic 2/ Both sexes Male Female Non-Hispanic white 3/ Both sexes Male Female Non-Hispanic black 3/ Both sexes Male Female

Cholelithiasis and other disorders of gallbladder (K80-K82)

Table with 16 columns (Year, All origins, Hispanic, Non-Hispanic, Non-Hispanic white, Non-Hispanic black) and 16 rows (2007-1999) showing death rates for cholelithiasis.

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Table with 16 columns (Year, All origins, Hispanic, Non-Hispanic, Non-Hispanic white, Non-Hispanic black) and 16 rows (2007-1999) showing death rates for nephritis.

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

Table with 16 columns (Year, All origins, Hispanic, Non-Hispanic, Non-Hispanic white, Non-Hispanic black) and 16 rows (2007-1999) showing death rates for acute nephritic syndrome.

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)															
2007.....	0.9	1.2	0.8	0.8	1.0	0.6	0.9	1.2	0.8	0.8	1.1	0.7	1.9	2.3	1.7
2006.....	0.6	0.7	0.5	0.5	0.6	0.4	0.6	0.7	0.5	0.5	0.6	0.4	1.3	1.7	1.1
2005.....	0.3	0.4	0.2	0.3	0.3	0.3	0.3	0.4	0.2	0.2	0.3	0.2	0.6	0.8	0.5
2004.....	0.2	0.2	0.2	0.1	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.3	0.4	0.3
2003.....	0.2	0.2	0.1	0.2	*	*	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.4	0.2
2002.....	0.2	0.2	0.2	0.2	*	*	0.2	0.2	0.2	0.2	0.2	0.1	0.3	0.4	0.3
2001.....	0.2	0.2	0.2	0.2	*	*	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.3	0.2
2000.....	0.2	0.3	0.2	0.2	*	*	0.2	0.3	0.2	0.2	0.2	0.1	0.4	0.5	0.3
1999.....	0.2	0.3	0.2	0.2	*	*	0.2	0.3	0.2	0.2	0.2	0.2	0.4	0.5	0.3
Renal failure (N17-N19)															
2007.....	13.6	16.5	11.6	11.4	13.4	10.0	13.7	16.7	11.7	12.2	15.3	10.3	28.0	32.0	25.4
2006.....	13.9	17.0	11.8	12.0	13.9	10.6	13.9	17.2	11.9	12.4	15.6	10.3	29.5	34.1	26.6
2005.....	14.0	17.1	12.0	11.6	14.0	10.0	14.1	17.3	12.1	12.6	15.8	10.5	29.6	33.9	26.8
2004.....	13.9	17.0	12.0	11.7	13.7	10.3	14.0	17.2	12.1	12.5	15.6	10.5	29.6	34.4	26.7
2003.....	14.2	17.5	12.1	12.4	14.6	10.9	14.3	17.7	12.2	12.6	16.1	10.6	30.6	36.2	27.0
2002.....	13.9	17.3	11.9	11.4	13.4	9.9	14.0	17.4	11.9	12.5	15.9	10.4	29.7	35.3	26.3
2001.....	13.7	17.1	11.8	12.1	14.2	10.8	13.8	17.2	11.8	12.2	15.7	10.1	29.6	33.9	27.1
2000.....	13.2	16.6	11.3	11.6	13.6	10.2	13.3	16.7	11.3	11.8	15.2	9.7	28.6	33.0	25.9
1999.....	12.8	16.1	10.8	11.7	14.9	9.6	12.8	16.1	10.9	11.2	14.5	9.3	28.2	33.5	25.3
Other disorders of kidney (N25,N27)															
2007.....	0.0	*	*	*	*	*	0.0	*	*	*	*	*	*	*	*
2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	0.0	*	*	0.0	*	*	*	*	*
2004.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	0.0	*	*	0.0	*	*	*	*	*
2002.....	0.0	*	0.0	*	*	*	0.0	*	0.0	0.0	*	0.0	*	*	*
2001.....	0.0	*	0.0	*	*	*	0.0	*	0.0	*	*	*	*	*	*
2000.....	0.0	0.0	0.0	*	*	*	0.0	*	0.0	0.0	*	*	*	*	*
1999.....	0.0	*	0.0	*	*	*	0.0	*	0.0	*	*	*	*	*	*

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Infections of kidney (N10-N12,N13.6,N15.1)</b>															
2007.....	0.2	0.2	0.2	0.2	*	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2006.....	0.2	0.2	0.2	0.2	*	0.2	0.2	0.1	0.2	0.2	0.2	0.3	0.2	0.2	0.2
2005.....	0.2	0.2	0.3	0.2	0.3	0.2	0.2	0.2	0.3	0.2	0.2	0.3	0.3	0.3	0.4
2004.....	0.2	0.2	0.3	0.3	*	0.3	0.2	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.4
2003.....	0.3	0.2	0.3	0.3	*	0.3	0.3	0.2	0.3	0.3	0.2	0.3	0.4	0.3	0.4
2002.....	0.3	0.2	0.3	0.3	*	0.3	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.3	0.3
2001.....	0.3	0.2	0.3	0.2	*	0.2	0.3	0.2	0.3	0.3	0.2	0.3	0.3	0.3	0.3
2000.....	0.3	0.2	0.3	0.3	*	0.4	0.3	0.2	0.3	0.3	0.2	0.3	0.4	0.3	0.5
1999.....	0.3	0.2	0.3	0.4	0.4	0.4	0.3	0.2	0.3	0.3	0.2	0.4	0.4	0.3	0.4
<b>Hyperplasia of prostate (N40)</b>															
2007.....	0.1	0.4	...	0.2	0.4	...	0.1	0.4	...	0.1	0.4	...	0.2	0.5	...
2006.....	0.2	0.5	...	0.1	0.4	...	0.2	0.5	...	0.2	0.5	...	0.2	0.4	...
2005.....	0.2	0.5	...	*	*	...	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2004.....	0.2	0.5	...	*	*	...	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2003.....	0.2	0.5	...	0.2	0.4	...	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2002.....	0.1	0.4	...	*	*	...	0.1	0.4	...	0.1	0.4	...	0.2	0.5	...
2001.....	0.1	0.5	...	*	*	...	0.2	0.5	...	0.2	0.5	...	0.1	0.4	...
2000.....	0.2	0.5	...	*	*	...	0.2	0.5	...	0.2	0.5	...	0.2	0.5	...
1999.....	0.2	0.5	...	*	*	...	0.2	0.5	...	0.2	0.5	...	0.2	0.6	...
<b>Inflammatory diseases of female pelvic organs (N70-N76)</b>															
2007.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	*	...	*
2006.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	*	...	*
2005.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	*	...	*
2004.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.0	*	...	*
2003.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	*	...	*
2002.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.0	0.1	...	0.1
2001.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
2000.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.1	0.1	...	0.1
1999.....	0.0	...	0.1	*	...	*	0.0	...	0.1	0.0	...	0.0	0.1	...	0.2

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Pregnancy, childbirth and the puerperium (000-099)</b>															
2007.....	0.3	...	0.5	0.3	...	0.6	0.3	...	0.5	0.2	...	0.4	0.6	...	1.2
2006.....	0.2	...	0.5	0.3	...	0.6	0.2	...	0.5	0.2	...	0.3	0.7	...	1.3
2005.....	0.3	...	0.5	0.3	...	0.6	0.3	...	0.5	0.2	...	0.3	0.7	...	1.3
2004.....	0.2	...	0.5	0.2	...	0.5	0.2	...	0.5	0.1	...	0.3	0.6	...	1.2
2003.....	0.2	...	0.4	0.2	...	0.5	0.2	...	0.3	0.1	...	0.2	0.5	...	1.0
2002.....	0.1	...	0.3	0.2	...	0.3	0.1	...	0.2	0.1	...	0.2	0.4	...	0.8
2001.....	0.1	...	0.3	0.2	...	0.4	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
2000.....	0.1	...	0.3	0.2	...	0.4	0.1	...	0.3	0.1	...	0.2	0.4	...	0.7
1999.....	0.2	...	0.3	0.2	...	0.4	0.1	...	0.3	0.1	...	0.2	0.4	...	0.8
<b>Pregnancy with abortive outcome (000-007)</b>															
2007.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2006.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2005.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2004.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2003.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2002.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2001.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
2000.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
1999.....	0.0	...	0.0	*	...	*	0.0	...	0.0	*	...	*	*	...	*
<b>Other complications of pregnancy, childbirth and the puerperium (010-099)</b>															
2007.....	0.3	...	0.5	0.2	...	0.5	0.3	...	0.5	0.2	...	0.4	0.6	...	1.1
2006.....	0.2	...	0.5	0.3	...	0.6	0.2	...	0.5	0.2	...	0.3	0.6	...	1.2
2005.....	0.2	...	0.5	0.3	...	0.6	0.2	...	0.5	0.2	...	0.3	0.6	...	1.3
2004.....	0.2	...	0.5	0.2	...	0.5	0.2	...	0.4	0.1	...	0.3	0.6	...	1.1
2003.....	0.2	...	0.3	0.2	...	0.5	0.2	...	0.3	0.1	...	0.2	0.5	...	0.9
2002.....	0.1	...	0.2	0.2	...	0.3	0.1	...	0.2	0.1	...	0.1	0.4	...	0.7
2001.....	0.1	...	0.3	0.2	...	0.4	0.1	...	0.2	0.1	...	0.2	0.4	...	0.7
2000.....	0.1	...	0.3	0.2	...	0.4	0.1	...	0.2	0.1	...	0.2	0.3	...	0.6
1999.....	0.1	...	0.3	0.2	...	0.4	0.1	...	0.2	0.1	...	0.1	0.4	...	0.7

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year

	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Certain conditions originating in the perinatal period (P00-P96)

2007.....	4.7	5.2	4.2	3.9	4.3	3.6	4.9	5.4	4.4	3.6	4.0	3.2	10.2	11.2	9.1
2006.....	4.8	5.3	4.3	4.0	4.4	3.6	5.0	5.5	4.5	3.6	4.0	3.2	10.7	11.5	9.8
2005.....	4.9	5.4	4.4	4.2	4.7	3.6	5.0	5.5	4.5	3.7	4.1	3.3	10.6	11.7	9.5
2004.....	4.8	5.4	4.2	4.1	4.6	3.6	4.9	5.5	4.4	3.6	4.1	3.2	10.3	11.4	9.3
2003.....	4.9	5.5	4.3	4.2	4.6	3.8	5.1	5.7	4.4	3.8	4.3	3.3	10.3	11.5	9.0
2002.....	4.9	5.4	4.3	4.0	4.4	3.6	5.0	5.6	4.5	3.8	4.2	3.3	10.2	11.1	9.3
2001.....	4.7	5.3	4.2	3.8	4.2	3.3	4.9	5.4	4.4	3.6	4.0	3.2	10.7	11.8	9.5
2000.....	5.1	5.6	4.5	3.8	4.2	3.5	5.3	5.9	4.7	3.8	4.3	3.4	11.7	13.0	10.3
1999.....	5.2	5.7	4.6	3.9	4.3	3.6	5.4	5.9	4.8	3.8	4.3	3.4	12.1	13.5	10.8

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

2007.....	3.4	3.6	3.2	3.0	3.2	2.9	3.4	3.6	3.2	3.3	3.4	3.1	4.1	4.2	3.9
2006.....	3.5	3.7	3.2	3.2	3.5	3.0	3.5	3.7	3.2	3.4	3.6	3.1	4.2	4.5	3.9
2005.....	3.5	3.7	3.3	3.3	3.4	3.2	3.5	3.7	3.3	3.4	3.6	3.2	3.9	4.2	3.8
2004.....	3.6	3.7	3.4	3.2	3.3	3.1	3.6	3.8	3.4	3.5	3.7	3.4	4.1	4.5	3.7
2003.....	3.6	3.8	3.4	3.4	3.5	3.2	3.6	3.8	3.4	3.5	3.6	3.4	4.0	4.5	3.7
2002.....	3.7	3.8	3.5	3.4	3.6	3.2	3.7	3.8	3.5	3.6	3.8	3.4	4.2	4.5	4.0
2001.....	3.6	3.8	3.4	3.4	3.8	3.0	3.6	3.8	3.4	3.5	3.8	3.3	4.2	4.4	4.1
2000.....	3.8	4.1	3.6	3.6	3.9	3.3	3.8	4.1	3.6	3.7	3.9	3.5	4.5	4.9	4.2
1999.....	3.8	3.9	3.6	3.4	3.4	3.4	3.8	4.0	3.6	3.7	3.9	3.5	4.5	4.8	4.2

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

2007.....	10.4	10.5	9.8	5.8	6.6	4.9	10.7	10.9	10.2	10.4	10.5	9.9	14.3	15.6	12.8
2006.....	10.1	10.5	9.3	5.5	6.1	4.7	10.4	10.9	9.7	10.2	10.5	9.5	13.7	15.7	11.8
2005.....	10.4	11.1	9.4	5.7	6.9	4.6	10.7	11.4	9.7	10.4	10.9	9.4	14.6	16.9	12.5
2004.....	10.0	10.7	9.0	5.3	6.1	4.5	10.3	11.0	9.4	9.9	10.5	9.0	14.2	16.1	12.3
2003.....	10.6	11.4	9.5	5.6	6.6	4.5	11.0	11.8	9.9	10.6	11.3	9.6	15.0	17.6	12.8
2002.....	10.3	11.4	9.0	5.5	6.6	4.5	10.7	11.8	9.3	10.2	11.1	9.0	14.8	18.2	12.0
2001.....	11.3	12.7	9.8	7.9	9.9	5.8	11.5	12.8	10.0	11.0	12.2	9.7	15.4	18.0	13.1
2000.....	11.4	13.1	9.7	7.8	10.3	5.5	11.6	13.2	10.0	11.0	12.5	9.4	16.7	20.0	13.9
1999.....	9.7	11.0	8.4	5.2	6.2	4.3	10.0	11.3	8.7	9.3	10.4	8.1	15.6	19.3	12.7

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Table with columns: Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004) and year; All origins 1/ (Both sexes, Male, Female); Hispanic (Both sexes, Male, Female); Non-Hispanic 2/ (Both sexes, Male, Female); Non-Hispanic white 3/ (Both sexes, Male, Female); Non-Hispanic black 3/ (Both sexes, Male, Female). Rows include: All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98); Accidents (unintentional injuries) (V01-X59, Y85-Y86); Transport accidents (V01-V99, Y85).

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year

Table with 16 columns: Cause of death, All origins 1/ (Both sexes, Male, Female), Hispanic (Both sexes, Male, Female), Non-Hispanic 2/ (Both sexes, Male, Female), Non-Hispanic white 3/ (Both sexes, Male, Female), Non-Hispanic black 3/ (Both sexes, Male, Female). Rows include Motor vehicle accidents (V02-V04, V09.0, V09.2, V12-V14, V19.0-V19.2, V19.4-V19.6, V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1, V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0, V89.2) for years 2007-1999.

Other land transport accidents (V01, V05-V06, V09.1, V09.3-V09.9, V10-V11, V15-V18, V19.3, V19.8-V19.9, V80.0-V80.2, V80.6-V80.9, V81.2-V81.9, V82.2-V82.9, V87.9, V88.9, V89.1, V89.3, V89.9)

Table with 16 columns: Cause of death, All origins 1/ (Both sexes, Male, Female), Hispanic (Both sexes, Male, Female), Non-Hispanic 2/ (Both sexes, Male, Female), Non-Hispanic white 3/ (Both sexes, Male, Female), Non-Hispanic black 3/ (Both sexes, Male, Female). Rows include Other land transport accidents for years 2007-1999.

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99, Y85)

Table with 16 columns: Cause of death, All origins 1/ (Both sexes, Male, Female), Hispanic (Both sexes, Male, Female), Non-Hispanic 2/ (Both sexes, Male, Female), Non-Hispanic white 3/ (Both sexes, Male, Female), Non-Hispanic black 3/ (Both sexes, Male, Female). Rows include Water, air and space, and other and unspecified transport accidents and their sequelae for years 2007-1999.

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Nontransport accidents (W00-X59,Y86)</b>															
2007.....	24.6	32.7	17.2	16.1	23.2	8.9	25.7	34.0	18.2	27.0	35.6	19.2	22.2	31.5	14.7
2006.....	23.8	32.1	16.2	16.2	23.4	8.9	24.7	33.2	17.0	25.7	34.2	17.9	23.3	34.2	14.4
2005.....	22.9	30.9	15.7	16.0	23.0	8.9	23.6	31.7	16.4	24.3	32.4	17.0	23.7	34.5	15.1
2004.....	21.5	29.0	14.9	14.6	20.8	8.5	22.2	29.9	15.5	23.0	30.7	16.1	21.1	30.6	13.6
2003.....	20.9	28.4	14.3	14.7	21.7	7.9	21.4	28.9	14.8	22.0	29.5	15.3	20.6	29.9	13.3
2002.....	20.2	27.6	13.5	14.7	22.1	7.7	20.6	28.0	14.0	20.9	28.2	14.3	21.3	31.7	13.2
2001.....	19.1	26.4	12.7	14.0	20.1	7.9	19.4	26.8	13.1	19.5	26.7	13.2	21.4	31.8	13.3
2000.....	18.4	25.6	12.0	14.4	21.4	7.8	18.6	25.8	12.3	18.4	25.5	12.3	21.3	32.5	12.7
1999.....	18.7	26.3	12.1	15.2	23.3	7.6	18.7	26.2	12.4	18.4	25.5	12.3	23.3	35.3	13.8
<b>Falls (W00-W19)</b>															
2007.....	7.0	9.0	5.5	5.0	6.4	3.7	7.1	9.1	5.6	7.6	9.6	6.0	3.5	5.0	2.4
2006.....	6.6	8.6	5.1	5.1	6.9	3.4	6.7	8.6	5.3	7.0	9.0	5.6	3.7	5.5	2.5
2005.....	6.4	8.3	4.9	5.0	6.6	3.5	6.4	8.3	5.1	6.8	8.7	5.3	3.7	5.2	2.6
2004.....	6.3	8.2	4.8	4.7	6.5	3.2	6.3	8.2	4.9	6.7	8.6	5.2	3.6	5.6	2.3
2003.....	5.9	7.6	4.5	4.2	5.8	2.9	5.9	7.7	4.6	6.2	8.0	4.9	3.4	4.7	2.4
2002.....	5.6	7.4	4.3	4.3	6.1	2.9	5.6	7.4	4.4	5.9	7.6	4.6	3.1	4.9	2.0
2001.....	5.3	7.2	3.9	4.1	5.4	2.8	5.3	7.3	3.9	5.6	7.5	4.2	3.3	5.2	2.0
2000.....	4.8	6.6	3.6	4.2	5.9	2.8	4.8	6.5	3.6	5.0	6.7	3.8	3.0	4.6	1.9
1999.....	4.8	6.6	3.5	4.1	6.0	2.6	4.8	6.6	3.6	5.0	6.7	3.7	3.5	5.3	2.2
<b>Accidental discharge of firearms (W32-W34)</b>															
2007.....	0.2	0.4	0.0	0.1	0.2	*	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.6	*
2006.....	0.2	0.4	0.0	0.1	0.2	*	0.2	0.4	0.1	0.2	0.4	0.0	0.3	0.6	*
2005.....	0.3	0.5	0.1	0.2	0.4	*	0.3	0.5	0.1	0.3	0.4	0.1	0.4	0.7	*
2004.....	0.2	0.4	0.0	0.1	0.2	*	0.2	0.4	0.1	0.2	0.4	0.1	0.3	0.6	*
2003.....	0.2	0.5	0.1	0.2	0.3	*	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.6	*
2002.....	0.3	0.4	0.1	0.2	0.2	*	0.3	0.5	0.1	0.3	0.5	0.1	0.4	0.8	*
2001.....	0.3	0.5	0.1	0.2	0.4	*	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.6	*
2000.....	0.3	0.5	0.1	0.2	0.2	*	0.3	0.5	0.1	0.3	0.5	0.1	0.3	0.7	0.1
1999.....	0.3	0.5	0.1	0.2	0.3	*	0.3	0.5	0.1	0.3	0.5	0.1	0.4	0.8	0.1



Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
<b>Accidental drowning and submersion (W65-W74)</b>															
2007.....	1.1	1.8	0.5	1.0	1.7	0.3	1.1	1.8	0.5	1.1	1.7	0.6	1.3	2.1	0.4
2006.....	1.2	1.9	0.5	1.0	1.7	0.4	1.2	1.9	0.5	1.2	1.8	0.6	1.3	2.3	0.4
2005.....	1.2	1.9	0.5	1.1	1.8	0.4	1.2	1.9	0.5	1.2	1.8	0.5	1.4	2.6	0.4
2004.....	1.1	1.8	0.5	1.0	1.6	0.3	1.1	1.8	0.5	1.1	1.7	0.5	1.4	2.3	0.5
2003.....	1.1	1.8	0.5	1.1	1.8	0.3	1.1	1.8	0.5	1.1	1.7	0.5	1.2	2.2	0.4
2002.....	1.2	2.0	0.5	1.1	1.9	0.3	1.2	1.9	0.5	1.1	1.7	0.5	1.6	2.8	0.5
2001.....	1.1	1.8	0.5	1.0	1.6	0.3	1.2	1.8	0.5	1.1	1.7	0.5	1.5	2.7	0.5
2000.....	1.2	2.0	0.5	1.1	1.7	0.4	1.3	2.0	0.5	1.2	1.8	0.5	1.6	2.7	0.6
1999.....	1.3	2.0	0.5	1.2	2.0	0.3	1.2	2.0	0.5	1.1	1.8	0.5	1.7	3.0	0.5
<b>Accidental exposure to smoke, fire and flames (X00-X09)</b>															
2007.....	1.1	1.3	0.8	0.6	0.9	0.4	1.1	1.4	0.9	1.0	1.2	0.8	2.3	3.1	1.7
2006.....	1.0	1.3	0.8	0.6	0.8	0.4	1.1	1.3	0.9	0.9	1.2	0.8	2.1	2.7	1.7
2005.....	1.1	1.4	0.8	0.7	0.8	0.6	1.1	1.4	0.8	1.0	1.3	0.7	2.3	3.1	1.6
2004.....	1.1	1.4	0.8	0.6	0.8	0.5	1.2	1.5	0.9	1.0	1.3	0.8	2.5	3.4	1.8
2003.....	1.2	1.4	0.9	0.6	0.9	0.4	1.2	1.5	1.0	1.1	1.4	0.9	2.5	3.3	1.9
2002.....	1.1	1.4	0.8	0.8	1.0	0.5	1.2	1.5	0.8	1.0	1.3	0.7	2.5	3.3	1.8
2001.....	1.2	1.5	0.9	0.6	0.8	0.5	1.2	1.6	0.9	1.0	1.4	0.8	2.7	3.8	1.9
2000.....	1.2	1.6	0.9	0.7	0.9	0.5	1.3	1.7	0.9	1.0	1.4	0.8	3.1	4.2	2.3
1999.....	1.2	1.6	0.9	0.7	1.1	0.4	1.3	1.6	0.9	1.1	1.4	0.8	3.1	4.1	2.3
<b>Accidental poisoning and exposure to noxious substances (X40-X49)</b>															
2007.....	9.8	13.0	6.6	5.8	8.9	2.5	10.5	13.8	7.2	11.6	15.2	8.0	8.8	12.5	5.7
2006.....	9.1	12.4	5.9	5.7	8.5	2.7	9.7	13.2	6.3	10.5	14.1	6.9	9.7	14.5	5.6
2005.....	7.9	10.7	5.1	5.2	7.9	2.3	8.4	11.3	5.5	9.0	12.0	6.0	8.3	12.1	5.1
2004.....	7.1	9.5	4.7	4.6	6.9	2.3	7.5	10.0	5.0	8.1	10.8	5.5	7.1	10.1	4.6
2003.....	6.7	9.1	4.3	5.0	7.6	2.2	6.9	9.4	4.5	7.4	9.9	4.8	6.9	10.1	4.2
2002.....	6.1	8.4	3.7	4.7	7.3	2.0	6.3	8.6	4.0	6.6	9.0	4.2	6.9	10.3	4.0
2001.....	4.9	7.0	2.9	3.9	6.1	1.6	5.0	7.1	3.0	5.1	7.2	3.1	6.4	9.6	3.8
2000.....	4.5	6.6	2.5	4.1	6.6	1.5	4.5	6.5	2.6	4.6	6.5	2.6	6.1	9.4	3.4
1999.....	4.4	6.4	2.3	4.9	8.1	1.6	4.2	6.2	2.4	4.2	6.0	2.3	6.3	9.8	3.2

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year

	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

2007.....	5.4	7.2	3.7	3.5	5.2	1.9	5.5	7.4	3.9	5.5	7.4	3.8	6.0	8.2	4.4
2006.....	5.6	7.6	3.8	3.7	5.3	2.1	5.8	7.8	4.0	5.8	7.9	4.0	6.1	8.5	4.1
2005.....	6.1	8.2	4.2	3.8	5.6	2.2	6.2	8.4	4.4	6.2	8.3	4.3	7.6	10.7	5.3
2004.....	5.7	7.6	4.0	3.6	4.9	2.2	5.8	7.9	4.1	5.9	7.9	4.1	6.1	8.7	4.3
2003.....	5.8	7.9	4.1	3.6	5.4	2.0	6.0	8.0	4.2	6.0	8.0	4.2	6.3	9.1	4.3
2002.....	5.9	7.9	4.2	3.6	5.5	2.0	6.1	8.1	4.3	6.1	8.1	4.3	6.8	9.6	4.8
2001.....	6.3	8.4	4.5	4.2	5.8	2.6	6.4	8.5	4.6	6.3	8.5	4.6	7.1	10.0	5.0
2000.....	6.3	8.5	4.4	4.2	6.0	2.5	6.4	8.6	4.5	6.3	8.5	4.5	7.2	11.0	4.6
1999.....	6.7	9.1	4.8	4.1	5.8	2.7	6.8	9.2	4.9	6.8	9.1	4.9	8.3	12.4	5.5

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

2007.....	11.3	18.4	4.7	6.0	10.1	1.9	12.0	19.7	5.0	13.5	21.9	5.7	5.1	9.0	1.8
2006.....	10.9	18.0	4.5	5.3	8.8	1.8	11.7	19.3	4.9	13.2	21.4	5.6	5.2	9.6	1.4
2005.....	10.9	18.0	4.4	5.6	9.4	1.8	11.6	19.1	4.7	12.9	21.2	5.3	5.4	9.4	1.9
2004.....	10.9	18.0	4.5	5.9	9.8	2.0	11.6	19.1	4.9	12.9	21.0	5.4	5.5	9.8	1.8
2003.....	10.8	18.0	4.2	5.6	9.7	1.7	11.4	19.1	4.5	12.7	21.0	5.0	5.4	9.4	1.9
2002.....	10.9	18.4	4.2	5.7	9.9	1.8	11.6	19.5	4.5	12.9	21.4	5.1	5.4	10.0	1.6
2001 4/.....	10.7	18.2	4.0	5.7	10.1	1.6	11.3	19.2	4.3	12.5	21.0	4.9	5.5	9.9	1.8
2000.....	10.4	17.7	4.0	5.9	10.3	1.7	10.9	18.6	4.2	12.0	20.2	4.7	5.6	10.2	1.8
1999.....	10.5	17.8	4.0	5.9	10.3	1.9	11.0	18.7	4.2	12.0	20.2	4.7	5.7	10.5	1.7

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

2007.....	5.6	10.3	1.4	2.4	4.4	0.4	6.1	11.2	1.5	6.9	12.6	1.8	2.6	5.0	0.5
2006.....	5.5	10.2	1.4	2.0	3.8	0.3	6.0	11.0	1.5	6.8	12.3	1.8	2.7	5.4	0.5
2005.....	5.7	10.5	1.4	2.2	4.0	0.3	6.1	11.3	1.5	6.9	12.6	1.7	2.8	5.4	0.5
2004.....	5.6	10.3	1.5	2.4	4.4	0.5	6.0	11.1	1.6	6.8	12.3	1.9	2.8	5.6	0.5
2003.....	5.8	10.7	1.4	2.4	4.5	0.3	6.2	11.5	1.5	6.9	12.8	1.8	2.8	5.5	0.6
2002.....	5.9	11.1	1.4	2.5	4.8	0.4	6.3	11.8	1.5	7.0	13.0	1.8	3.0	6.0	0.5
2001.....	5.9	11.0	1.4	2.5	4.8	0.4	6.3	11.7	1.6	7.0	12.9	1.8	3.1	6.1	0.7
2000.....	5.9	11.0	1.5	2.7	5.1	0.5	6.3	11.6	1.6	6.9	12.8	1.8	3.2	6.1	0.7
1999.....	6.0	11.1	1.5	2.8	5.3	0.6	6.3	11.8	1.6	6.9	12.8	1.8	3.3	6.7	0.6

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

2007.....	5.6	8.1	3.3	3.6	5.7	1.5	6.0	8.5	3.5	6.6	9.4	3.9	2.5	4.0	1.2
2006.....	5.4	7.8	3.1	3.3	5.0	1.5	5.7	8.3	3.3	6.4	9.1	3.8	2.5	4.1	1.0
2005.....	5.2	7.5	3.0	3.5	5.4	1.5	5.5	7.9	3.2	6.1	8.6	3.6	2.6	4.0	1.4
2004.....	5.3	7.6	3.1	3.5	5.4	1.5	5.6	8.0	3.3	6.1	8.7	3.6	2.6	4.2	1.3
2003.....	5.0	7.3	2.8	3.2	5.2	1.3	5.2	7.6	3.0	5.7	8.3	3.3	2.6	3.9	1.3
2002.....	5.0	7.4	2.8	3.2	5.1	1.4	5.3	7.7	3.0	5.8	8.4	3.3	2.4	3.9	1.1
2001 4/.....	4.8	7.1	2.6	3.2	5.3	1.2	5.0	7.4	2.8	5.5	8.1	3.1	2.4	3.8	1.1
2000.....	4.5	6.7	2.5	3.1	5.2	1.2	4.7	6.9	2.6	5.1	7.4	2.9	2.5	4.0	1.1
1999.....	4.5	6.6	2.5	3.1	5.0	1.3	4.7	6.9	2.6	5.0	7.4	2.9	2.4	3.9	1.1

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

2007.....	6.1	9.6	2.5	6.9	11.2	2.3	5.9	9.2	2.5	2.8	3.7	1.8	21.8	38.5	6.3
2006.....	6.2	9.7	2.5	7.3	11.7	2.3	6.0	9.3	2.6	2.7	3.6	1.8	22.4	39.2	6.7
2005.....	6.1	9.6	2.5	7.5	12.1	2.4	5.8	9.1	2.5	2.7	3.5	1.8	21.8	38.5	6.2
2004.....	5.9	9.2	2.5	7.2	11.5	2.3	5.6	8.6	2.6	2.7	3.6	1.8	20.7	36.2	6.5
2003.....	6.0	9.4	2.6	7.7	12.1	2.7	5.7	8.9	2.6	2.7	3.6	1.8	21.7	37.9	6.6
2002.....	6.1	9.4	2.8	7.3	11.6	2.5	5.8	8.9	2.8	2.8	3.7	1.9	21.6	37.4	7.0
2001 4/.....	7.1	10.8	3.3	8.3	12.9	3.1	6.8	10.3	3.3	4.0	5.6	2.5	21.7	37.2	7.5
2000.....	5.9	9.0	2.8	7.5	11.8	2.8	5.6	8.5	2.7	2.8	3.6	1.9	21.1	36.4	7.2
1999.....	6.0	9.1	2.9	7.6	12.2	2.6	5.7	8.5	2.9	2.9	3.8	2.0	20.7	35.2	7.5

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

2007.....	4.2	7.1	1.2	4.7	7.8	1.1	4.1	6.9	1.3	1.6	2.2	0.9	16.9	31.2	3.3
2006.....	4.3	7.2	1.3	4.9	8.3	1.1	4.1	6.9	1.3	1.4	2.1	0.8	17.4	32.0	3.6
2005.....	4.2	7.0	1.2	5.1	8.5	1.2	4.0	6.7	1.2	1.4	2.1	0.9	16.6	30.9	3.1
2004.....	4.0	6.6	1.1	4.8	8.1	1.0	3.7	6.3	1.2	1.5	2.1	0.8	15.6	29.0	3.0
2003.....	4.1	6.8	1.2	5.1	8.5	1.2	3.9	6.5	1.2	1.5	2.1	0.8	16.3	30.1	3.3
2002.....	4.1	6.8	1.3	4.8	8.1	1.2	3.9	6.5	1.3	1.6	2.2	0.9	16.1	29.5	3.6
2001.....	3.9	6.6	1.3	4.9	8.2	1.2	3.7	6.2	1.3	1.6	2.2	0.9	15.2	28.2	3.1
2000.....	3.8	6.3	1.2	4.8	8.0	1.3	3.6	6.0	1.3	1.5	2.0	0.9	15.0	27.8	3.2
1999.....	3.8	6.3	1.3	5.0	8.2	1.4	3.6	6.0	1.3	1.5	2.2	0.9	14.8	26.9	3.6

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year

	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

2007.....	1.9	2.5	1.3	2.3	3.3	1.2	1.8	2.3	1.3	1.2	1.5	0.9	5.0	7.3	3.0
2006.....	1.9	2.6	1.3	2.3	3.4	1.2	1.9	2.4	1.3	1.3	1.6	0.9	5.0	7.2	3.1
2005.....	1.9	2.6	1.3	2.4	3.6	1.2	1.8	2.4	1.3	1.2	1.5	0.9	5.2	7.6	3.1
2004.....	2.0	2.5	1.4	2.4	3.4	1.3	1.8	2.3	1.4	1.3	1.5	1.0	5.1	7.2	3.5
2003.....	2.0	2.6	1.4	2.6	3.7	1.5	1.9	2.4	1.4	1.2	1.5	1.0	5.4	7.9	3.3
2002.....	2.0	2.6	1.4	2.4	3.5	1.4	1.9	2.4	1.4	1.3	1.6	1.0	5.5	7.9	3.5
2001 4/.....	3.1	4.3	2.0	3.3	4.7	1.9	3.1	4.1	2.0	2.4	3.3	1.5	6.5	9.0	4.4
2000.....	2.1	2.7	1.5	2.7	3.9	1.6	2.0	2.6	1.5	1.3	1.6	1.0	6.1	8.6	4.0
1999.....	2.2	2.8	1.6	2.6	4.0	1.3	2.1	2.6	1.6	1.4	1.7	1.1	5.9	8.2	3.9

Legal intervention (Y35,Y89.0)

2007.....	0.1	0.3	*	0.2	0.4	*	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2006.....	0.2	0.3	*	0.2	0.4	*	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2005.....	0.1	0.3	*	0.2	0.3	*	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2004.....	0.1	0.3	*	0.2	0.3	*	0.1	0.3	*	0.1	0.2	*	0.2	0.5	*
2003.....	0.1	0.3	*	0.2	0.3	*	0.1	0.2	*	0.1	0.2	*	0.3	0.6	*
2002.....	0.1	0.3	*	0.2	0.3	*	0.1	0.3	*	0.1	0.2	*	0.3	0.6	*
2001.....	0.1	0.3	*	0.1	0.3	*	0.1	0.3	*	0.1	0.2	*	0.2	0.4	*
2000.....	0.1	0.2	*	0.1	0.2	*	0.1	0.2	*	0.1	0.2	*	0.3	0.6	*
1999.....	0.1	0.3	*	0.2	0.3	*	0.1	0.3	*	0.1	0.2	*	0.4	0.7	*

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

2007.....	1.8	2.2	1.3	0.8	1.1	0.5	1.9	2.4	1.4	2.0	2.5	1.6	2.0	3.0	1.2
2006.....	1.7	2.2	1.2	0.8	1.1	0.5	1.9	2.4	1.4	1.9	2.4	1.5	2.0	3.0	1.1
2005.....	1.6	1.9	1.2	0.8	1.1	0.6	1.7	2.1	1.3	1.8	2.1	1.4	1.9	2.6	1.2
2004.....	1.7	2.2	1.2	0.9	1.2	0.5	1.8	2.3	1.3	1.9	2.4	1.4	1.9	2.8	1.2
2003.....	1.7	2.3	1.2	0.9	1.4	0.4	1.8	2.4	1.3	2.0	2.5	1.4	1.9	2.9	1.0
2002.....	1.7	2.2	1.2	0.8	1.3	0.4	1.8	2.3	1.3	1.8	2.3	1.3	2.1	3.1	1.3
2001.....	1.5	2.0	0.9	0.9	1.3	0.4	1.5	2.1	1.0	1.6	2.1	1.1	1.8	2.9	0.9
2000.....	1.3	1.9	0.8	0.6	0.9	0.3	1.4	2.0	0.9	1.4	1.9	0.9	1.9	3.1	1.0
1999.....	1.4	2.0	0.9	0.7	1.2	0.3	1.5	2.0	0.9	1.4	1.9	0.9	2.0	3.3	1.0

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year

	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female

Discharge of firearms, undetermined intent (Y22-Y24)

2007.....	0.1	0.1	0.0	0.0	0.1	*	0.1	0.1	0.0	0.1	0.2	0.0	0.1	0.2	*
2006.....	0.1	0.1	0.0	0.1	0.1	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2005.....	0.1	0.1	0.0	0.0	0.1	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2004.....	0.1	0.1	0.0	0.0	*	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.1	*
2003.....	0.1	0.1	0.0	0.1	0.1	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2002.....	0.1	0.1	0.0	0.1	0.1	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2001.....	0.1	0.1	0.0	0.1	*	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.2	*
2000.....	0.1	0.1	0.0	0.0	*	*	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.3	*
1999.....	0.1	0.2	0.0	0.1	0.2	*	0.1	0.2	0.0	0.1	0.2	0.0	0.2	0.3	*

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

2007.....	1.7	2.1	1.3	0.7	0.9	0.5	1.8	2.3	1.4	1.9	2.4	1.5	1.9	2.8	1.1
2006.....	1.6	2.1	1.2	0.8	1.0	0.5	1.8	2.3	1.3	1.8	2.2	1.4	1.8	2.8	1.1
2005.....	1.5	1.8	1.2	0.8	1.0	0.5	1.6	2.0	1.3	1.7	2.0	1.4	1.8	2.5	1.2
2004.....	1.6	2.0	1.2	0.8	1.1	0.5	1.7	2.2	1.3	1.8	2.3	1.4	1.8	2.7	1.1
2003.....	1.7	2.2	1.2	0.8	1.3	0.4	1.8	2.3	1.3	1.9	2.4	1.4	1.9	2.8	1.0
2002.....	1.6	2.0	1.1	0.8	1.2	0.4	1.7	2.2	1.2	1.7	2.2	1.3	2.0	2.9	1.2
2001.....	1.4	1.9	0.9	0.8	1.2	0.4	1.5	2.0	1.0	1.5	1.9	1.0	1.7	2.8	0.9
2000.....	1.3	1.8	0.8	0.6	0.9	0.3	1.4	1.9	0.9	1.3	1.8	0.9	1.8	2.8	0.9
1999.....	1.3	1.8	0.8	0.6	1.0	0.3	1.3	1.8	0.9	1.3	1.7	0.9	1.9	3.0	0.9

Operations of war and their sequelae (Y36,Y89.1)

2007.....	0.0	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2006.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2005.....	0.0	0.0	*	*	*	*	0.0	0.0	*	*	*	*	*	*	*
2004.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	0.0	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	0.0	*	*	*	*	0.0	0.0	*	0.0	*	*	*	*	*

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year

Complications of medical and surgical care (Y40-Y84,Y88)

Table with 17 columns: Cause of death, All origins 1/ (Both sexes, Male, Female), Hispanic (Both sexes, Male, Female), Non-Hispanic 2/ (Both sexes, Male, Female), Non-Hispanic white 3/ (Both sexes, Male, Female), Non-Hispanic black 3/ (Both sexes, Male, Female). Rows for years 2007-1999.

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 5/

Table with 17 columns: Cause of death, All origins 1/ (Both sexes, Male, Female), Hispanic (Both sexes, Male, Female), Non-Hispanic 2/ (Both sexes, Male, Female), Non-Hispanic white 3/ (Both sexes, Male, Female), Non-Hispanic black 3/ (Both sexes, Male, Female). Rows for years 2007-1999.

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 5/

Table with 17 columns: Cause of death, All origins 1/ (Both sexes, Male, Female), Hispanic (Both sexes, Male, Female), Non-Hispanic 2/ (Both sexes, Male, Female), Non-Hispanic white 3/ (Both sexes, Male, Female), Non-Hispanic black 3/ (Both sexes, Male, Female). Rows for years 2007-1999.

Worktable Orig293. Age-adjusted death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Age-adjusted death rates per 100,000 U.S. standard population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Race categories are consistent with the 1977 Office of Management and Budget standards. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data--Mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision,) Second Edition, 2004) and year	All origins 1/			Hispanic			Non-Hispanic 2/			Non-Hispanic white 3/			Non-Hispanic black 3/		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 5/															
2007.....	10.2	18.2	2.7	7.4	12.9	1.5	10.6	18.8	2.9	8.8	15.4	2.8	20.2	37.6	4.0
2006.....	10.2	18.1	2.7	7.3	12.7	1.5	10.5	18.7	2.9	8.6	15.0	2.7	20.8	38.8	4.1
2005.....	10.2	18.3	2.7	7.6	13.3	1.6	10.5	18.8	2.8	8.8	15.3	2.7	20.0	37.7	3.7
2004.....	10.0	17.7	2.7	7.5	13.1	1.5	10.2	18.2	2.9	8.7	15.1	2.8	19.0	35.7	3.7
2003.....	10.3	18.4	2.7	7.8	13.6	1.6	10.5	18.8	2.8	8.8	15.6	2.7	19.7	36.8	3.9
2002.....	10.4	18.6	2.8	7.6	13.4	1.6	10.7	19.1	3.0	9.0	16.0	2.8	19.8	37.0	4.2
2001.....	10.3	18.5	2.8	7.8	13.7	1.7	10.5	18.8	3.0	9.1	16.0	2.8	18.9	35.4	3.9
2000.....	10.2	18.1	2.8	7.8	13.6	1.8	10.3	18.4	3.0	8.8	15.5	2.8	18.9	35.2	4.1
1999.....	10.3	18.4	2.9	8.2	14.2	2.0	10.5	18.7	3.0	8.9	15.8	2.8	19.0	35.2	4.4

\* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for origin not stated are included in "All origins" but not distributed among specified origins.

2/ Includes races other than white and black.

3/ Multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2007, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2006, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality."

4/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see http://www.cdc.gov/nchs and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

5/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.